

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
31 January 2002 (31.01.2002)


PCT

(10) International Publication Number
WO 02/07669 A2(51) International Patent Classification⁷: **A61K**(21) International Application Number: **PCT/IL01/00629**(22) International Filing Date: **10 July 2001 (10.07.2001)**(25) Filing Language: **English**(26) Publication Language: **English**(30) Priority Data:
09/614,271 **12 July 2000 (12.07.2000)** **US**(71) Applicant (for all designated States except US):
D-PHARM LTD. [IL/IL]; Kiryat Weizmann Science
Park, 76326 Rehovot (IL).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **KOZAK, Alexander**
[IL/IL]; 8 Naphtali Ben Ephraim Street, 76555 Rehovot
(IL).(74) Agent: **WALKER, Naomi**; Kiryat Weizmann Science
Park, P.O. Box 2313, 76123 Rehovot (IL).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.(54) Title: **PHOSPHOLIPID DERIVATIVES OF VALPROIC ACID AND MIXTURES THEREOF**

(57) Abstract: The present invention relates to compounds, which are phospholipid derivatives of valproic acid, to compositions comprising said compounds and their use for treating epilepsy, migraine, bipolar disorders and pain.

WO 02/07669 A2

PHOSPHOLIPID DERIVATIVES OF VALPROIC ACID AND MIXTURES THEREOF

5

10

FIELD OF THE INVENTION

The present invention relates to compounds, which are phospholipid
15 derivatives of valproic acid, to compositions comprising said compounds and
their use for treating epilepsy, migraine, bipolar disorders and pain.

BACKGROUND OF THE INVENTION

Epilepsy is a neurological disease that is characterized by paroxysmal
20 transient disturbances of the electrical activity of the brain. Epileptic seizures
may be partial or focal seizures that are restricted to a particular locus within
the brain, or generalized seizures which can result in abnormal activity
throughout the brain. The disturbances of brain function during an epileptic
attack may be manifested as psychic or sensory incidents such as amnesia,
25 hallucinations, déjà vu states etc., as abnormal motor phenomenon such as
spasms or whole body convulsions or as loss of consciousness. In extreme
cases, epilepsy can degenerate into status epilepticus which may be fatal
(DeLorenzo et al.- J. Clin. Neurophysiol. (1995) 12: 316-325).

Valproic acid (VPA) and its sodium salt (sodium valproate, NaVPA) are
30 among the most prescribed anti-epileptic drugs. These drugs are also effective
in the treatment of bipolar disorders and in prophylaxis of migraines.

Although the clinical usefulness of valproic acid is well established, this
compound suffers major drawbacks. Treatment with VPA is associated with

adverse side effects such as gastro-intestinal irritation, bone marrow suppression (especially manifested as aplastic anemia and thrombocytopenia), and hepatic dysfunction. VPA has also been reported to have teratogenic effects and patients treated with VPA may experience nausea, vomiting, dizziness, confusion or sedation.

Another drawback of valproic acid is its short half-life due to rapid clearance of the drug. As a result plasma levels of VPA fluctuate during chronic treatment and the drug has to be administered several times a day even as a sustained release formulation. In addition, valproic acid, which is a liquid, is less desirable for use as an oral dosage form. The sodium valproate, on the other hand, is solid, but being hygroscopic is characterized by poor stability.

Efforts have been made in order to overcome the VPA-induced side effects and the disadvantageous physical and pharmacokinetic properties of the drug. Most approaches that have been taken involve modification of the VPA molecule. However, although some of the modified drugs were devoid of adverse side effects, in many cases they also lost the therapeutic effect or were much less potent.

Mergen et al (J. Pharm. Pharmacol. (1991), 43: 815-816) describe conjugates of valproic acid with 1,3-dipalmitoylglycerol, 1,2-dipalmitoylglycerol or 1,3-diaminopalmitoyl-propan-2-ol. According to the Mergen et al.'s publication, only the latter compound was found to have antiepileptic activity, while both conjugates of VPA with the diglycerides were inactive.

Hadad et al. (Biopharmaceutics & Drug Disposition (1993), 14: 51-59) investigated the anticonvulsant activity of 1,4-butanediol divalproate, glyceryl trivalproate and valpromide in comparison to valproic acid. Their study demonstrated that only 1,4-butanediol divalproate, in one of the model systems tested, had a better protective index value than VPA.

WO 02/07669

PCT/IL01/00629

U.S. Patent No. 4,654,370 to Marriott and Paris discloses glycerides esterified with one or two moles of valproic acid. These compounds have been found to have the same useful therapeutic effect as valproic acid alone but without causing gastric irritation.

5

U.S. Patents Nos. 4,988,731 and 5,212,326 both to Meade, disclose oligomers having 1:1 molar ratio of sodium valproate and valproic acid which have physiological properties similar to those of valproic acid or sodium valproate but show superior stability characteristics.

10

U.S. Patent No. 4,558,070 to Bauer and Shada discloses a stable complex between valproic acid and potassium, cesium or rubidium which may be formed by combining four moles of valproic acid with one mole of the alkali metal ion. The alkali metal salts of valproic acid were reported to have improved stability.

15

Despite continuous efforts in the field, it is still an unmet need to provide an anti-epileptic medication with improved pharmacokinetic properties and overall superior therapeutic index.

20

SUMMARY OF THE INVENTION

A first aspect of the present invention provides a pharmaceutical composition comprising a therapeutically effective amount of valproic acid or a pharmaceutically acceptable derivative thereof covalently bonded to a phospholipid moiety; and a pharmaceutically acceptable carrier. In preferred embodiments, the phospholipid moiety may be selected from plasmalogens, phosphatidic acids and phosphoglycerides. More preferred are compounds, wherein said phospholipid moiety may be lysophosphatidyl-ethanolamine, N-mono-(C₁₋₄)-alkyl, N,N-di-(C₁₋₄)-alkyl and quaternary derivatives of the amines thereof.

25

Most preferred embodiments may be compositions comprising phospholipid derivatives of valproic acid (hereinafter

30

referred to as DP-VPA) wherein valproic acid is covalently linked as an ester at the sn-2 position of a phospholipid moiety.

Currently the most preferred DP-VPA compounds are 1-Palmitoyl-2-valproyl-sn-glycero-3-phosphocholine, also referred to as 1-hexadecanoyl-sn-glycero-3-phosphorylcholine (hereinafter denoted as C₁₆-DP-VPA) and 1-Stearoyl-2-valproyl-sn-glycero-3-phosphocholine, also referred to as 1-octadecanoyl-sn-glycero-3-phosphorylcholine (hereinafter denoted as C₁₈-DP-VPA).

According to preferred embodiments, the pharmaceutical compositions may comprise a mixture of DP-VPA compounds, more preferably a mixture of C₁₆-DP-VPA and C₁₈-DP-VPA (hereinafter denoted as C₁₆/C₁₈-DP-VPA).

In one preferred embodiment the ratio of C₁₆-DP-VPA to C₁₈-DP-VPA in the C₁₆/C₁₈-DP-VPA mixture can be from around 1:20 to around 1:2 by weight. Most preferred are mixtures wherein the ratio of C₁₆-DP-VPA to C₁₈-DP-VPA can be from around 1:5 to around 1:7 w/w (equivalent to 15±5% C₁₆-DP-VPA: 85±5% C₁₈-DP-VPA (w/w)).

A second aspect of the present invention provides a use of valproic acid or pharmaceutically acceptable derivative in the form of a composition as defined above in the manufacture of a medicament for reducing side effects of valproic acid or pharmaceutically acceptable derivative thereof.

A third aspect of the present invention provides a use of valproic acid or pharmaceutically acceptable derivative in the form of a composition as defined above in the manufacture of a medicament for reducing interactions of valproic acid or pharmaceutically acceptable derivative thereof with other therapeutic agents.

A fourth aspect of the present invention provides a use of valproic acid or pharmaceutically acceptable derivative in the form of a composition as defined above in the manufacture of a medicament for increasing valproic acid or pharmaceutically acceptable derivative thereof efficacy.

A fifth aspect of the present invention provides a compound comprising valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to a phospholipid moiety.

The compounds and compositions of the invention are useful for the treatment of central nervous system disorders including, but not limited to, epilepsy, migraines, chronic pain and bipolar disorders.

Thus, according to a sixth aspect of the present invention, there is provided a use of a pharmaceutical composition comprising a therapeutically effective amount of valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to a phospholipid moiety; a pharmaceutically acceptable carrier; said composition providing a therapeutic effect in the manufacture of a medicament for the treatment of a central nervous system disorder in a mammal.

Further embodiments will become apparent to those skilled in the art upon further review of the following disclosure, including the detailed descriptions of specific embodiments of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 depicts a typical HPLC chromatogram of a C₁₆/C₁₈-DP-VPA composition, where the C₁₆-DP-VPA to C₁₈-DP-VPA ratio is 15%:85%, by weight

- 5 Figs. 2A-B depict plasma concentrations of C₁₆-DP-VPA (2A) and C₁₈-DP-VPA (2B) measured at different time points following a single oral administration to human subjects of 0.625 g of C₁₆/C₁₈-DP-VPA (C₁₆/C₁₈ ratio = 13%: 87% w/w).

10 **DETAILED DESCRIPTION THE INVENTION**

The present invention relates to phospholipid derivatives of valproic acid, to pharmaceutical compositions comprising these compounds and mixtures thereof, and to their use in the treatment of neurological disorders.

- DP-VPA molecules are disclosed in US Patent Application No. 15 08/479,959 and International Patent Publication WO 94/22483, the disclosure of which is herein incorporated by reference.


- Specifically disclosed in the abovementioned applications is a molecule, referred to as "TVA 16", which is a 1:1 ester of valproic acid with 1-hexadecanoyl-sn-glycero-3-phosphorylcholine. TVA 16 was shown to have 20 significant anticonvulsant activity and to be more potent than sodium valproate. In the present application, the 1:1 ester of valproic acid with 1-hexadecanoyl-sn-glycero-3-phosphorylcholine, or in its chemical name 1-Palmitoyl-2-valproyl-sn-glycero-3-phosphocholine, is hereinafter referred to as C₁₆-DP-VPA.

- 25 Another embodiment, disclosed in the present application, is a 1:1 ester of valproic acid with 1-octadecanoyl-sn-glycero-3-phosphorylcholine, or in its chemical name 1-Stearoyl-2-valproyl-sn-glycero-3-phosphocholine. This molecule is hereinafter referred to as C₁₈-DP-VPA.

- The DP-VPA compounds of the invention include conjugates of valproic 30 acid, or a pharmaceutically acceptable derivative thereof, with any phospholipid, preferably a phosphoglyceride. Suitable phospholipids include,

but are not limited to, plasmalogens, phosphatidic acids and phospho-ester derivatives thereof. Preferred phospholipid moieties, in accordance with the invention, include lysophosphatidyl-ethanolamine, N-mono-(C₁₋₄)-alkyl, N,N-di-(C₁₋₄)-alkyl and quaternary derivatives of the amines thereof. Currently, the most preferred phospholipid in the compounds of the invention is phosphatidylcholine.

The choice of the fatty acid residue at position sn-1 of a glycerophospholipid moiety is specifically discussed below in connection with the preferred compounds in accordance with the invention. However, it should be appreciated that VPA or its pharmaceutically acceptable derivative may be covalently linked to the phospholipid moiety at positions sn-1, sn-2 or linked to the phospholipid head group at position sn-3. Accordingly, it is possible that VPA or its derivative is released by cleavage by the respective phospholipases, PLA₁, PLA₂, PLC and PLD as depicted in the following scheme 1.



X= H or a polar head group.

In preferred embodiments of the invention, VPA, or its pharmaceutically acceptable derivative, is joined to the phospholipid via an ester linkage at position sn-2, thus enabling release of VPA by phospholipases A₂. In

particularly preferred embodiments, VPA is covalently linked by an ester bond to the sn-2 position of phosphatidylcholine.

The term "pharmaceutically acceptable derivatives of VPA" as used in the specification refers to pharmaceutically acceptable analogs of valproic acid having similar therapeutic activity. This includes derivatives of VPA with saturated or unsaturated carbon chains having one or more double and/or triple bonds. Pharmaceutically acceptable substituent on the carbon atoms of the molecules are also allowed and may include, for example, halogen atoms or lower alkyl groups comprising 1-5 carbon atoms. Amides of VPA and its analogs as mentioned above are also included within the scope of the invention. Furthermore, for those compounds having a chiral centre of asymmetry, the compounds of the present invention include optically active isomers, racemates or preferred mixtures thereof.

It should be appreciated that within the scope of the invention are also pharmaceutically acceptable salts of the DP-VPA compounds. The term "pharmaceutically acceptable salts" means non-toxic salts of the compounds of the invention including, but not limited to, sodium, potassium, calcium, magnesium, ammonium, alkyl ammonium or amine derived salts.

Although both C₁₈-DP-VPA and C₁₆-DP-VPA are potent anticonvulsant agents having similar efficacies, it has now been unexpectedly found that the two compounds differ in their pharmacokinetic profiles. The C₁₆-DP-VPA compound, following a single oral administration, exhibits a significantly prolonged half-life in plasma in comparison to the half-life of C₁₈-DP-VPA. However, the plasma concentrations of C₁₈-DP-VPA were found to reach peak levels at a later time point in comparison to the peak levels of the C₁₆-DP-VPA molecule.

It is now disclosed, for the first time, that mixtures of C₁₆-DP-VPA and C₁₈-DP-VPA (herein referred to as C₁₆/C₁₈-DP-VPA) offer an advantage by exhibiting higher neuroprotective values and prolonged therapeutic effect in

comparison to the effect of compositions comprising either C₁₆-DP-VPA or C₁₈-DP-VPA alone.

Preferred compositions, in accordance with the invention, are those wherein the C₁₆-DP-VPA to C₁₈-DP-VPA ratio in the C₁₆/C₁₈-DP-VPA mixture is from around 1:20 to around 1:2 (by weight). Most preferred are mixtures wherein the ratio of C₁₆-DP-VPA to C₁₈-DP-VPA is around 15±5% : 85±5% (w/w).

Without wishing to be limited to a single mechanism or theory, it is suggested that the length of the alkyl moiety esterified at position sn-1 of the phospholipid may determine the lipophilicity of the DP-VPA molecule, and thus also its transport across cellular membranes.

Alternatively, again without wishing to be limited to a single mechanism or theory, the fatty acid residue at position sn-1 may determine the properties of the DP-VPA conjugate as a substrate for phospholipases, thus affecting the regulated release of valproic acid, for example, by elevated activity of phospholipase A₂ (PLA₂) at the diseased site. Phospholipases A₂ are a family of esterases that hydrolyze the sn-2 ester bonds in phosphoglyceride molecules. It has been shown that in disorders such as epilepsy, PLA₂ activation coincides with epileptic seizures (Flynn and Wecker (1987) J. Neurochem. 48: 1178-84; Bazan, et al. (1986) Adv. Neurol. 44: 879-902).

Also associated with elevated phospholipase A₂ activity, are bipolar disorders and some types of pain and migraine that are associated with inflammatory processes (Horrobin and Bennett (1999) Prostaglandins Leukot Essent Fatty Acids 60: 141-167).

The compounds according to the invention, being hydrophobic in nature, may penetrate biological membranes and barriers, thus facilitating the transport of the drug into cells or organs, for example, into the brain where their effect is needed.

It may be envisaged that regulated release of the valproic acid moiety at the diseased target site may even further improve the therapeutic index of the drug, as the efficacy the drug is expected to increase while potential side effects

and toxicity are reduced. Valproic acid may be released by cleavage of the DP-VPA compound at position sn-2 of the phospholipid by phospholipase A₂ or any other lipase or esterase. However, it may not be excluded that the active drug may be different from the original parent drug molecule, VPA, with a chemical group(s) being removed from or added to its structure while being released from its intracellular transporting adjuvant or as a result of physiological phospholipid metabolism.

It is important to note that the conjugate of the invention, namely the valproic acid or its pharmaceutically acceptable derivative covalently linked to the phospholipid moiety, may be active *per se*. Alternatively, the covalent bond of the lipid-drug conjugate may, under certain circumstances, be cleaved to release the pharmacologically active drug. In the latter case, the compound of the invention may be regarded as a prodrug, in the sense that the therapeutic agent is released from its transporting adjuvant.

Irrespective of the exact mechanism of action, it is evident that the compounds of the invention have an improved therapeutic profile and are more effective comparing to VPA in at least two aspects: (i) increased efficacy, and (ii) decreased side effects.

The DP-VPA compounds were found to be effective at much lower equivalent molar doses compared to the doses currently used for VPA. The reduced therapeutic doses in turn reduce the toxicological risk, accompanying side effects and also reduce the risk of undesirable interactions with other drugs. In addition, the DP-VPA molecules have been found to exhibit significantly improved pharmacokinetic properties compared to VPA (e.g. substantially increased half-life in serum and in brain tissue). Thus, the DP-VPA molecules represent a class of superior anti-epileptic drugs.

Furthermore, the preferred pharmaceutical compositions in accordance with the invention, i.e. compositions comprising a mixture of both C₁₆-DP-VPA and C₁₈-DP-VPA, may conveniently be prepared from natural sources.

It would be highly advantageous to have DP-VPA molecules that can be obtained from starting material which can be derived from natural sources by a

relatively simple procedure. Such starting material, which is readily available, is lyso-lecithin obtained from egg or soybean lecithins. The soybean, being a non-animal source, is the preferred starting material in the preparation of medicaments for human use. In typical hydrogenated preparations derived from soybean the content of 1-palmitoyl-lysolecithin is around 8-18% and the content of 1-stearoyl-lysolecithin is around 80- 90% (by weight).

Pure C₁₆-DP-VPA and C₁₈-DP-VPA molecules may be chemically synthesized *de novo*. Alternatively, pure C₁₆-DP-VPA and C₁₈-DP-VPA compounds may be prepared by using starting materials obtainable from natural sources, i.e C₁₆-lyso-lecithin and C₁₈-lyso-lecithin may be isolated and purified, for example, from eggs or soybeans, and then acylated by VPA (= semi-natural preparation).

C₁₆-DP-VPA and C₁₈-DP-VPA, in particular when used as mixtures within a preferred range of ratios in accordance with the present invention, have been shown, to exhibit significantly improved therapeutic properties. The advantageous properties were exemplified by the substantially increased half-life of DP-VPA in serum and the high efficacy of the drug. The improved residency time in the serum may facilitate the attainment of steady-state drug levels with reduced fluctuation around the therapeutic blood level and reduction in the frequency of drug administration to once or twice per day.

The compositions of the invention can be administered orally, parenterally, (for example by intravenous drip or intraperitoneal, subcutaneous, or intramuscular injection), topically, (for example by nasal application or inhalation) or rectally. Oral administration is a currently more preferred route of administration.

Suitable formulations for administration of the compounds of the invention, whether used separately or as a mixture, include, but are not limited

to, powders, granules, emulsions, suspensions or solutions in water or non-aqueous media, in the dosage form of tablets, capsules, syrups or solutions.

For oral administration, DP-VPA amounts of from about 0.5 to 20 mg/kg body weight per day are useful, preferably 1 to 8 mg/kg body weight per day.

- 5 Dosing will be dependent on the severity of the symptoms and on the responsiveness of the subject to the DP-VPA drug. A physician or other persons of ordinary skill in the art can easily determine optimum dosages and dosage form as well as dosage regimen and means of administration.


- 10 The invention will now be illustrated by the following non-limiting examples.

EXAMPLES:

15 **EXAMPLE 1: Synthesis of DP-VPA**

- The synthesis of DP-VPA is a two-stage process. The first stage is aimed at obtaining valproic anhydride by heating of valproic acid in a solution of acetic anhydride under catalysis of pyridine. In the second stage, DP-VPA is prepared by interaction of valproic anhydride with lyso-lecithin. This reaction
20 is conducted in a solution of valproic anhydride by catalysis of 4-dimethylaminopyridine at 90-100°C.

- Extraction and purification of the product obtained are carried out in four stages. The first stage of purification is performed by extraction of the un-reacted valproic anhydride, valproic acid and catalyst (4-aminopyridine) in
25 acetone. The crude product obtained is precipitated and separated from solution at the second stage. The solid product obtained is washed from the remaining compounds at the third stage. Finally, the product is re-crystallized several times from an acetone/ethanol solution and the residual solvents are removed under vacuum. Yield of the product is around 80%.

Scheme of synthesis of C₁₈-DP-VPA

40 The 1-Palmitoyl-2-valproyl-sn-glycero-3-phosphocholine (C₁₆-DP-VPA) and 1-Stearoyl-2-valproyl-sn-glycero-3-phosphocholine (C₁₈-DP-VPA) compounds were prepared using, respectively, lyso-stearic- and lyso-palmitic-phosphatidylcholines.

WO 02/07669

PCT/IL01/00629

The lyso-stearic- and lyso-palmitic-phosphatidylcholines may be purified from a natural source (e.g. egg or soybean) by means and procedures well known in the art (F. Gunstone (1999) Fatty Acid and Lipid Chemistry, pp. 87-99, Aspen Publishers, Inc.).

- 5 Alternatively these starting materials may be obtained by chemical synthesis procedures as known in the art.

EXAMPLE 2: Synthesis of C₁₆/C₁₈-DP-VPA

- 10 Mixtures of C₁₆/C₁₈-DP-VPA were prepared by the same procedure as described in Example 1 for the preparation of DP-VPA. The difference was that in the case of the mixture compositions the interaction of valproic anhydride was with lyso-lecithin, which was obtained from soybean and saturated by hydrogenation (S VPC-3 from Lipoid GmbH, Ludwigshafen, Germany).

15

EXAMPLE 3: Analysis of DP-VPA compounds

- C₁₆/C₁₈-DP-VPA mixtures synthesized as described above in Example 2 were subjected to analytical assays for characterization and proof of structure. Analytical results of a product containing 1-Palmitoyl-2-valproyl-sn-glycero-3-phosphocholine (C₁₆-DP-VPA) and 1-Stearoyl-2-valproyl-sn-glycero-3-phosphocholine (C₁₈-DP-VPA), at a ratio of 13%:87% (by weight) are given below.

20

Mass spectroscopy

- The mass of the protonated DP-VPA molecules as determined by ESI (+) is 25 622.4 to 622.8 for C₁₆-DP-VPA, and 650.4 to 650.8 for C₁₈-DP-VPA.

 This agrees well with the calculated molecular weight values.

Elementary analysis

- Calculated for M.H₂O: C 60.93%, H 10.25%, N 2.11%, P 4.66% (M is corrected for the content of 1-palmitoyl-2-valproyl-sn-glycero-3-phosphocholine).
30 The average values found were C 60.72%, H 10.58%, N 2.09%, P 4.56%.

These values agree well with the calculated values.

Thin layer chromatography (TLC) analysis

A TLC is performed on silica gel 60 F₂₅₄ on aluminium sheet with a chloroform:methanol:water (65:35:5, v/v) fluent. The spray reagent for
5 detection is a mixture of 4-methoxybenzaldehyde (5ml), sulphuric acid 95-98% (5ml), ethanol (100ml) and glacial acetic acid (1ml). The sheet is sprayed with this reagent and then heated with hot air at 120-150°C.

The TLC analysis results show that there is one spot at an R_f of 0.58 to 0.60.

Analytical NMR data

10 The typical NMR data given below are for proton, carbon-13 and phosphorus-31.

¹H NMR(CDCl₃), δ (ppm): 0.84-0.90 (m, 9H), 1.24-1.27 (broad s) + 1.31-1.41 (m) (both 34H), 1.50-1.59 (m, 4H), 2.21-2.28 (t, 2H), 2.29-2.37 (m, 1H), 3.35 (s, 9H), 3.77-3.78 (broad s, 2H), 3.88-3.96 (m, 2H), 4.06-4.14 (m, 1H), 4.30 (broad s, 2H), 4.40-4.46 (d, 1H), 5.18 (m, 1H).

15 ¹³C NMR(CDCl₃), δ (ppm): 8.62 (CH₃), 22.65 (CH₃), 62.99 [(CH₃)₃N], 29.17, 30.93, 33.15, 39.92, 43.07, 53.86, 67.97, 71.89, 74.99, 78.91 (CH₂ and CH), 182.21 (CO), 184.43 (CO).

³¹P NMR(CDCl₃), δ (ppm): -0.29 (respectively to H₃PO₄ in D₂O).

20 HPLC analysis

DP-VPA is analysed by HPLC using the following conditions:

Instrument:	Liquid chromatograph equipped with integrating device
Column:	Zorbax Eclipse XDB C18, 5μ, 4.6 x 250 mm
Mobile phase:	Methanol – Acetonitrile – Water (85:15:5 v/v)
Flow rate:	1.0 mL/min
Detection:	UV @ 220 nm
Injection volume	20 μL

25 Typical retention times are given in the following table:

Name of the compound	Typical Retention time, min
1-Palmitoyl-lysolecithine (potential impurity)	6
1-Stearoyl-lysolecithine (potential impurity)	8
1-Palmitoyl-DP-VPA	12
1-Stearoyl-DP-VPA	18

A typical HPLC chromatogram of C₁₆-DP-VPA /C₁₈-DP-VPA mixture at a ratio of 15%:85% (by weight) which was analysed as described above is depicted in Figure 1.

5

EXAMPLE 4: Toxicology and safety studies in humans

A Phase I safety and tolerability clinical trial has been carried out using C₁₆/C₁₈-DP-VPA mixture (C₁₆-DP-VPA /C₁₈-DP-VPA at a ratio of 15%:85% w/w) in solution of 5% Poloxamer F-127 + 0.5% Tween-80 and the oral route of administration.

10

The design of this study was double blind, placebo randomized. It was divided into a single administration part and a part in which there was repeated daily administration for 7 consecutive days. In each part the doses were increased at intervals of 7 days if the previous dose was well tolerated.

Five doses ranging from 0.3125 g to 5 g of DP-VPA were administered as single doses in the first part of the study and 3 doses (0.3125 g, 0.625 g and 1.25 g) were administered as repeated doses in the second part of the study. Fifty six subjects in total were enrolled in this study: 29 in the first part and 27 in the second part.

In the first part, nausea and vomiting were the most commonly reported adverse effects, the incidence of which was highest in the highest dose group (DP-VPA 5 g) with 3 of the 6 subjects reporting it. Two of the 6 subjects in the second highest dose group (DP-VPA 2.5 g) reported nausea and vomiting. Headache, diarrhoea, abdominal pain and dizziness were also reported but in the investigator's opinion they were probably not study drug related.

25

With regard to the laboratory results, vital signs and ECG parameters, no trends were seen and all results remained within acceptable parameters for all subjects.

In the second part, fewer adverse effects occurred and only 3 (burning stomach and abdominal pain) were considered to be probably related to the

30

drug as they occurred soon after administration. There were no incidences of nausea and the 1 episode of vomiting occurred more than 24 h after administration and was not related to the study drug.

From the point of view of the vital signs, ECGs, laboratory tests, urinalysis and physical examinations the tolerability was very good.

Conclusions: In both the single and repeated administration parts of this study the clinical and biological tolerability was found to be very good at DP-VPA doses of up to 2.5 g. It can be concluded that the DP-VPA's toxicological profile is significantly improved compared to that for the parent drug VPA.

EXAMPLE 5: Pharmacokinetics studies in humans

In order to assess the pharmacokinetic properties of the C₁₆-DP-VPA and C₁₈-DP-VPA compounds, the plasma levels of these compounds were monitored in human subjects.

Healthy male volunteers, 18-40 year old, (7 individuals for each dose tested) received, by a single oral administration, 0.3125 g, 0.625 g or 1.25 g of C₁₆ / C₁₈-DP-VPA mixture at a ratio of C₁₆ / C₁₈ =13% : 87% (w/w). Blood samples, 10 ml each, were drawn from each individual at the time points after the administration of the drug, as indicated. The samples were centrifuged at 4° C at 1100 g for 10 minutes immediately after collection. Plasma levels of C₁₆-DP-VPA and C₁₈-DP-VPA were determined using a LC-MS/MS technique. The plasma concentration profiles of C₁₆-DP-VPA and C₁₈-DP-VPA as monitored for 24 hours following a single oral administration 0.625 g of C₁₆/C₁₈-DP-VPA are shown, respectively, in Figures 2A and 2B.

As can be seen from the results of the human study, the C₁₆-DP-VPA and C₁₈-DP-VPA compounds have different kinetic profiles. While the peak concentration of C₁₆-DP-VPA in the plasma was reached at 6 hours after the administering of the drug, the peak of C₁₈-DP-VPA was reached two hours later, at 8 hours post administration.

The terminal plasma half-life ($t_{1/2}$) for the C₁₆-DP-VPA and C₁₈-DP-VPA compounds was calculated from their plasma concentration-time profiles. It was found that the calculated $t_{1/2}$ values for the two compounds were significantly different; $t_{1/2}$ for C₁₆-DP-VPA was 14.0 ± 0.6 hours compared to 8.3 $\pm$ 1.3 hours for C₁₈-DP-VPA.

The $t_{1/2}$ measured for the global DP-VPA was 10.6 ± 1.2 hours, a value that combines the pharmacokinetic profiles of both the C₁₆-DP-VPA and C₁₈-DP-VPA compounds.

Plasma concentration ($\mu\text{g/ml}$) of C₁₆-DP-VPA and of C₁₈-DP-VPA measured in samples collected at various time points following a single oral administration of 0.625 g C₁₆/C₁₈-DP-VPA are summarized in Table 1.

Table 1: C₁₆-DP-VPA and of C₁₈-DP-VPA concentrations in humans plasma following a single oral administration of C₁₆/C₁₈-DP-VPA.

Time (hours)	C ₁₆ -DP-VPA ($\mu\text{g/ml}$)	C ₁₈ -DP-VPA ($\mu\text{g/ml}$)	Ratio C ₁₈ / C ₁₆
1	0.013 ± 0.034	0.000 ± 0.000	
3	0.167 ± 0.094	0.058 ± 0.043	0.35
6	0.397 ± 0.172	0.818 ± 0.292	2.1
8	0.385 ± 0.140	1.175 ± 0.137	3.1
10	0.368 ± 0.182	0.949 ± 0.254	2.6
12	0.318 ± 0.150	0.698 ± 0.163	2.2

As can be seen from the results in Table 1, the observed ratios of C₁₈-DP-VPA / C₁₆-DP-VPA were different from the expected ratio of 6.7 (C₁₈-DP-VPA : C₁₆-DP-VPA ratio 87%: 13% (by weight)).

Surprisingly, the presentation of the C₁₆-DP-VPA compound in the plasma was found to be higher than its proportion in the administered mixture.

This phenomenon was more pronounced at the shorter time points, i.e. less than 6 hours following the administration of C₁₆/C₁₈-DP-VPA.

The highest ratio of C₁₈/ C₁₆, i.e. 3.1 (which is still below the expected ratio of 6.7) was reached at 8 hours following the administration of C₁₆/C₁₈-DP-VPA. At this point the levels of C₁₈-DP-VPA reach their peak concentration in the plasma.

These observations indicate that the C₁₆-DP-VPA and C₁₈-DP-VPA compounds demonstrate different pharmacokinetic profiles.

10 **EXAMPLE 6: Anti-convulsive effect of C₁₆-DP-VPA; C₁₈-DP-VPA and C₁₆/ C₁₈-DP-VPA mixtures (efficacy study)**

The anti-epileptic efficacies of C₁₆-DP-VPA, C₁₈-DP-VPA and mixtures comprising both C₁₆-DP-VPA and C₁₈-DP-VPA were evaluated in mice. The protective effect of the compounds was compared at different time points following chemical seizure induction by pentylenetetrazol (PTZ).

15 The pentylenetetrazol (PTZ) induced seizure model in mice is an established animal model system for epilepsy. Subcutaneous injection of PTZ into control animals results in the following sequence of events: myoclonic jerks within 1-2 mins, followed by clonic and clonic-tonic seizures each lasting approximately 5-10 secs with the severity of seizures increasing with time for up to 30 mins. 80% of the first seizures are observed within 5 mins, and 100% within 20 mins. The second seizures usually follow within 6-10 mins and subsequent seizures (if any) every 6 mins.

CD-1 mice (25-30 g) were pre-treated by subcutaneous (s.c.) injection of either C₁₆-DP-VPA or C₁₈-DP-VPA or a mixture thereof. The amounts used were equivalent to 40 mg/kg VPA. After different times as indicated, 1, 2 or 4 hours, a convulsive dose of pentylenetetrazol (85-100 mg/kg) was subcutaneously injected into the mice. The animals were monitored for one hour following the PTZ administration for the occurrence of episodes of clonic spasms persisting for at least 5 sec. The protection effect was calculated as the

25
30

number of animals that did not experience a second seizure divided by the total number of animals tested.

In Tables 2 A-D are shown the results of PTZ-induced seizures test as described above, wherein the tested compositions were as follows:

5 Table A - C₁₆-DP-VPA 100%

Table B - C₁₆-DP-VPA/C₁₈-DP-VPA ratio 50/50 (by weight)

Table C - C₁₆-DP-VPA/C₁₈-DP-VPA ratio 10/90 (by weight)

Table D - C₁₈-DP-VPA 100%

n= the number of animals experiencing a seizure;

10 N= the total number of animals in the assay.

15 **Tables 2 A-D: PTZ-induced seizures test in mice**

A) C₁₆-DP-VPA -100%

Time (h)	Second seizure (n/N)	% Protection
1	4/8	50
2	6/8	25
4	5/8	38

20 **B) C₁₆-DP-VPA/C₁₈-DP-VPA :50/50**

Time (h)	Second seizure (n/N)	% Protection
1	6/7	14
2	6/7	14
4	5/8	38

25 **C) C₁₆-DP-VPA/C₁₈-DP-VPA :10/90**

Time (h)	Second seizure (n/N)	% Protection
1	5/7	29
2	6/7	14
4	2/8	75

D) C₁₈-DP-VPA -100%

Time (h)	Second seizure (n/N)	% Protection
1	5/6	17
2	6/7	14
4	1/7	86

5 As can be seen in Table 2, the C₁₈-DP-VPA compound is more effective in prevention of second seizures, showing 86% protection compared to a maximum of 50% protection obtained by 100% C₁₆-DP-VPA.

The C₁₆-DP-VPA/C₁₈-DP-VPA mixtures demonstrated intermediate behaviour, with the effect of the C₁₆-DP-VPA/C₁₈-DP-VPA at a ratio of 10/90 (Table C) being closer to that obtained with the pure C₁₈-DP-VPA compound. However, protection was more apparent at 1 hour with the 10/90 mixture than with 100% C₁₈-DP-VPA.

It is also important to note the different kinetics of the protective effects by the two DP-VPA compounds. The peak activity of C₁₈-DP-VPA was obtained 4 hours after the administration of PTZ, while the pure C₁₆-DP-VPA reached its maximal therapeutic effect already within 1 hour.

Conclusion: although C₁₆-DP-VPA and C₁₈-DP-VPA have similar potency as anti-convulsant (i.e. similar ED₅₀ values), their therapeutic profiles are different. The C₁₆-DP-VPA compound is more effective in preventing seizures in the first hour after induction of seizures by subcutaneous administration of PTZ. The C₁₈-DP-VPA compound, on the other hand, shows maximal anti-convulsant activity at the later times, i.e. at four hours after the injection of PTZ (see Table A in comparison to Table D).

It is important to note that the tested C₁₆-DP-VPA/C₁₈-DP-VPA mixtures, in particular the C₁₆/C₁₈ mixture at a ratio of 10:90, behave in a similar way to the 100% C₁₈-DP-VPA compound, namely demonstrating high maximal seizure protection, but over a more prolonged time.

EXAMPLE 7: Suitable formulations for administration of DP-VPA

The DP-VPA compounds and compositions of the present invention can be administered to a subject in a number of ways, which are well known in the art. For example, administration may be done orally, parenterally, (for example
5 by intravenous drip or intraperitoneal, subcutaneous, or intramuscular injection), topically, (for example by nasal application or inhalation) or rectally.

Formulations for topical administration may include, but are not limited to drops, liquids, sprays, powders, suppositories, creams, gels and ointments. Conventional pharmaceutical carriers, aqueous, powder or oily bases,
10 thickeners and the like may be necessary or desirable.

Formulations for parenteral administration may include, but are not limited to, sterile aqueous solutions which may also contain buffers, diluents and other suitable additives.

Compositions for oral administration may be formulated as powders or
15 granules, suspensions or solutions in water or non-aqueous media, in the dosage form of tablets, capsules, syrups or solutions. The formulation may be designed so to enable modified controlled release of the active agent. The capsules and tablets may be coated so to afford site-specific delivery to different parts of the gastrointestinal tract. Thickeners, diluents, flavorings,
20 dispersing aids, emulsifiers or binders may be desirable.

Suitable pharmaceutical excipients that may be included in the formulations include, but are not limited to, phospholipids, triglycerides, propylene glycols, polyethylene glycols, poloxamers, surfactant and co-surfactants.

25 Dosing is dependent on the severity of the symptoms and on the responsiveness of the subject to the DP-VPA drug. Persons of ordinary skill in the art can easily determine optimum dosages and dosage form as well as dosage regimen and means of administration.

While the present invention has been particularly described, persons skilled in the art will appreciate that many variations and modifications can be made.

Therefore, the invention is not to be construed as restricted to the particularly described embodiments, rather the scope, spirit and concept of the invention will

5 be more readily understood by reference to the claims which follow.

WO 02/07669

PCT/IL01/00629

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising a therapeutically effective amount of valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to a phospholipid moiety; and a pharmaceutically acceptable carrier.
2. The pharmaceutical composition according to claim 1, wherein said phospholipid moiety is selected from the group consisting of plasmalogens, phosphatidic acids and phospho-esters derivatives thereof.
3. The pharmaceutical composition according to claim 1, wherein said phospholipid moiety is selected from the group consisting of lysophosphatidyl-ethanolamine, N-mono-(C₁₋₄)-alkyl, N,N-di-(C₁₋₄)-alkyl and quaternary derivatives of the amines thereof.
4. The pharmaceutical composition according to claim 3, wherein said quaternary derivative of lysophosphatidyl-ethanolamine is lysophosphatidylcholine.
5. The pharmaceutical composition according to claim 1, wherein said valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said phospholipid moiety is 1-palmitoyl-2-valproyl-sn-glycero-3-phosphocholine.
6. The pharmaceutical composition according to claim 1, wherein said valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said phospholipid moiety is 1-stearoyl-2-valproyl-sn-glycero-3-phosphocholine.

7. The pharmaceutical composition according to claim 1 wherein said valproic acid or pharmaceutically acceptable derivative thereof is covalently linked to said phospholipid moiety at a position selected from the group consisting of sn-1, sn-2, and sn-3.

8. The pharmaceutical composition according to claim 1 wherein said valproic acid or pharmaceutically acceptable derivative thereof is covalently linked to said phospholipid moiety at a position sn-2.

9. The pharmaceutical composition according to claim 8 wherein said valproic acid or pharmaceutically acceptable derivative thereof is released by cleavage by phospholipase A₂ at said sn-2 position.

10. The pharmaceutical composition according to claim 1, wherein said composition is in a form selected from the group consisting of solutions, suspensions, emulsions, syrups, capsules, tablets and suppositories.

11. The pharmaceutical composition of claim 1, which is suitable for oral administration, intravenous administration, or rectal administration.

12. The pharmaceutical composition of claim 1, further comprising a second valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to a different phospholipid moiety;

wherein said valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said phospholipid moiety and said second valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said different phospholipid moiety are in a ratio of from about 1:20 to about 1:2 by weight.

WO 02/07669

PCT/IL01/00629

13. The pharmaceutical composition of claim 1, further comprising a second valproic acid or said pharmaceutically acceptable derivative thereof covalently bonded to a different phospholipid moiety;

wherein said valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said phospholipid moiety and said second valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said different phospholipid moiety are in a ratio of from about 1:5 to about 1:7 by weight.

14. The pharmaceutical composition of claim 1, further comprising a second valproic acid or a pharmaceutically acceptable derivative thereof covalently bonded to a different phospholipid moiety.

15. The pharmaceutical composition of claim 14, wherein said valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said phospholipid moiety is 1-palmitoyl-2-valproyl-sn-glycero-3-phosphocholine.

16. The pharmaceutical composition of claim 14, wherein said second valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said different phospholipid moiety is 1-stearoyl-2-valproyl-sn-glycero-3-phosphocholine.

17. The pharmaceutical composition of claim 14, wherein the total amount of said first valproic acid or said pharmaceutically acceptable derivative thereof covalently bonded to said phospholipid moiety and said second valproic acid or said pharmaceutically acceptable derivative thereof covalently bonded to said different phospholipid moiety is from about 35 mg to about 2500 mg.

18. The pharmaceutical composition of claim 14, wherein the total amount

WO 02/07669

PCT/IL01/00629

of said first valproic acid or said pharmaceutically acceptable derivative thereof covalently bonded to said phospholipid moiety and said second valproic acid or said pharmaceutically acceptable derivative thereof covalently bonded to said different phospholipid moiety is from about 70 mg to about 560 mg.

19. The pharmaceutical composition of claim 14, wherein said combination exhibits greater therapeutic effects, prolonged therapeutic effects, or greater and prolonged therapeutic effects, on central nervous system disorders than a composition comprising only said valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to a phospholipid moiety, or only said second valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said different phospholipid moiety.

20. The pharmaceutical composition of claim 14, wherein said valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said phospholipid moiety is 1-palmitoyl-2-valproyl-sn-glycero-3-phosphocholine and said second valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to said different phospholipid moiety is 1-stearoyl-2-valproyl-sn-glycero-3-phosphocholine.

21. The pharmaceutical composition of claim 20, wherein 1-palmitoyl-2-valproyl-sn-glycero-3-phosphatidyl-choline and 1-stearoyl-2-valproyl-sn-glycero-3-phosphatidyl-choline are in a ratio of from about 1:20 to about 1:2 by weight.

22. The pharmaceutical composition of claim 20, wherein 1-palmitoyl-2-valproyl-sn-glycero-3-phosphocholine and 1-stearoyl-2-valproyl-sn-glycero-3-phosphocholine are in a ratio of from about 1:20 to about 1:2 by weight.

WO 02/07669

PCT/IL01/00629

23. The pharmaceutical composition of claim 20, which is suitable for oral administration, intravenous administration, or rectal administration.

24. Use of valproic acid or pharmaceutically acceptable derivative in the form of a composition as claimed in claim 1 in the manufacture of a medicament for reducing side effects of valproic acid or pharmaceutically acceptable derivative thereof.

25. Use of valproic acid or pharmaceutically acceptable derivative in the form of a composition as claimed in claim 1 in the manufacture of a medicament for reducing interactions of valproic acid or pharmaceutically acceptable derivative thereof with other therapeutic agents..

26. Use of valproic acid or pharmaceutically acceptable derivative in the form of a composition as claimed in claim 1 in the manufacture of a medicament for increasing valproic acid or pharmaceutically acceptable derivative thereof efficacy.

27. A compound comprising valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to a phospholipid moiety.

28. The compound of claim 27 wherein said phospholipid moiety is 1-stearoyl-sn-glycero-3-phosphorylcholine.

29. The compound of claim 27 wherein said phospholipid moiety is 1-palmitoyl-sn-glycero-3-phosphorylcholine.

30. Use of a pharmaceutical composition comprising a therapeutically effective amount of valproic acid or pharmaceutically acceptable derivative thereof covalently bonded to a phospholipid moiety; a pharmaceutically acceptable carrier; said composition providing a therapeutic effect in the manufacture of a medicament for the treatment of a central nervous system disorder in a mammal

31. A use according to claim 30, wherein the central nervous system disorder is selected from epilepsy, migraine, bipolar disorders and pain.

32. A use according to claim 30, wherein the central nervous system disorder is epilepsy.

33. A use according to claim 30, wherein said said phospholipid moiety is 1-stearoyl-sn-glycero-3-phosphorylcholine.

34. A use according to claim 30, wherein said said phospholipid moiety is 1-palmitoyl-sn-glycero-3-phosphorylcholine.