

- 1 -

REPLACES SHEET		80
☐	Abstract (pages 1-)	to page(s)
☒	Claims (pages 1-)	to page(s) 71.....
☐	Drawing (pages 1-)	to page(s)
☐	Material (pages 1-)	to page(s)
DATE OF RECEIPT		

" NEW BENZIMIDAZOLONE DERIVATIVES "

The present invention relates to novel pharmacologically active benzimidazolone derivatives and acid addition salts thereof, to processes for their preparation and to pharmaceutical compositions containing them. The new compounds possess central serotonergic activity and are useful in the treatment of central nervous system (CNS) disorders.

It is known that 1 A and 2 serotonergic receptors (5-HT_{1A} and 5-HT₂) seem to be important for many functions in the animal body. For instance, altered function of these receptors is involved in the genesis and/or treatment of anxiety, depression, psychoses, abnormality of sleep and feeding, organic mental diseases and alteration of blood pressure. In spite of the clear involvement of 5-HT_{1A} receptors in such a huge amount of pathological events, it is not clear why, for example, some compounds acting upon 5-HT_{1A} receptors exert in humans a preferential anxiolytic effects, while others exert a preferential hypotensive action. The same holds for 5-HT₂ antagonists. This is probably due to heterogeneous characteristics, so far unknown, of 5-HT_{1A} and 5-HT₂ receptors. Therefore, there is the possibility that compounds acting on 5-HT_{1A} and 5-HT₂ receptors may exert a wide range of therapeutic effects in humans.

GB 2023594 describes a class of 1-substituted alkyl-4-(3-

trifluoromethylthiophenyl)-piperazines which may contain, as a substituent of the alkyl group, an optionally 3-substituted benzimidazolone groupment. The above compounds were found to exert activity in the central nervous system, which was showed by behavioural tests in mice.

The benzimidazolone groupment was also used as a generic substituent in the preparation of structurally different classes of compounds endowed with activity on the central nervous system; examples may be found in BE 904,945, US 4,954,503 and EP 200,322.


US 3,472,854 describes (benzimidazolyl)-lower alkyl substituted piperazines useful, among other indications, as tranquilisers and sedatives.

EP 0376607 describes piperazinybutyl indole derivatives, including 2-indolones, which have been found to possess central serotonergic activity with preference for the 5-HT_{1A} receptor subtype.

We have now synthetised, and this is the object of the present invention, a novel class of structurally distinct compounds showing affinity for the 5-HT_{1A} and 5-HT₂ receptors. These new compounds may be useful in the treatment of CNS diseases such as affective disorders, (for example depression and bipolar disorders), anxiety, sleep and sexual disorders, psychosis, schizophrenia, personality disorders, mental organic disorders and mental disorders in childhood, aggressiveness, age associated memory impairment. Moreover they may be used for cardiovascular disorders such

as hypertension and thrombosis.

According to the present invention we provide compounds of general formula (I)



(I)

where

R₁ and R₂ may be at the same time or not a hydrogen atom, halogen, trifluoromethyl, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ acyl, carboxyl, C₁₋₆ alkoxycarbonyl, hydroxy, nitro, amino optionally C₁₋₄ alkyl N-mono or di-substituted, C₁₋₆ acylamino, C₁₋₆ alkoxycarbonylamino, carbamoyl optionally C₁₋₄ alkyl N-mono or di-substituted, cyano, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, amino sulphonyl optionally C₁₋₄ alkyl N-mono or di-substituted, C₁₋₄ alkyl N-mono or di-substituted aminosulphonylamino, aminosulphonyl-

- amino;
- R_3 is hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl or C_2-C_6 alkynyl;
- A is -CO- or -CONH- or it is absent;
- B is a straight or branched, saturated or unsaturated C_{2-6} alkyl;
- m and n are both independently an integer from 1 to 3;
- R_4 is an aryl, aralkyl, a heteroaryl or heteroaralkyl group, each group being optionally substituted by one or more substituents selected from halogen, trifluoromethyl, cyano, C_{1-3} alkoxy, C_{1-4} alkyl and acid addition salts thereof.

For pharmaceutical use the compounds of general formula (I) may be used as such or in the form of tautomers or of physiologically acceptable acid addition salts thereof. The term "acid addition salts" includes salts either with inorganic or organic acids. Physiologically acceptable organic acids which may be used in salt formation include, for example, maleic, citric, tartaric, fumaric, methansulphonic, acetic, benzoic, succinic, gluconic, isethionic, glycinic, lactic, malic, mucoic, glutammic, sulphamic and ascorbic acid; suitable inorganic acids include hydrochloric, hydrobromic, nitric, sulfuric or phosphoric acid.

Some of the compounds of formula (I) according to the present invention contain chiral or prochiral centres and thus may exist in different stereoisomeric forms including

enantiomers of (+) and (-) type or mixtures of them. The present invention includes in its scope both the individual isomers and the mixtures thereof.

It has to be understood that, when mixtures of optical isomers are present, they may be separated according to the classic resolution methods based on their different physico-chemical properties, e.g. by fractional crystallization of their acid addition salts with a suitable optically active acid or by the chromatographic separation with a suitable mixture of solvents.

When in the compounds of formula (I) R_1 , R_2 and R_3 represent C_{1-6} alkyl group, such groups refer to an alkyl group having a straight or branched chain. Typical groups of that kind include methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, n-hexyl, 2-methylpentyl and the like. The term halogen means fluoro, chloro, bromo and iodo. Preferred halogens are fluoro, chloro and bromo and particularly fluoro and chloro. When R_1 and R_2 represent a C_{1-6} alkoxy group, they may, for example, be methoxy, ethoxy, propoxy, butoxy, pentoxy, hexyloxy. When R_1 and R_2 represent a C_{1-6} acyloxy group it may, for example, be acetoxo, propionyloxy. When B is a straight or branched, saturated or unsaturated C_{2-6} alkyl group, it may, for example, ethyl, propyl, butyl, hexyl, 2-methylpropyl, 2-butenyl.

When R_4 is aryl, aralkyl it may, for example, be phenyl, benzyl or naphthyl respectively, each group being optionally


substituted by one or more substituents selected from fluoro, chloro, methoxy, methyl, trifluoromethyl, ethyl, ethoxy. When R_4 is heteroaryl or heteroalkyl, it may, for example, be 1,2-benzisothiazole, benzodioxane, pyrimidine or 1,2-benzioxazole.

When m and n are one of integers from 1 to 3, they may, for example, form a saturated 5-7-membered heterocyclic ring, such as piperazine, imidazolidine, diazepine.

Preferred compounds according to the present invention are those wherein A is absent, B is a straight, saturated C_{2-4} alkyl, m and n are the integer 2. R_4 is a substituted phenyl ring wherein the substituents are selected from methoxy, chloro and trifluoromethyl.


The compounds of the general formula (I) may conveniently be prepared by a variety of synthetic routes using conventional methods. According to a further feature of the invention we provide processes for the preparation of compounds of formula I as hereinbefore described in which either

a) a compound of general formula (II)



(II)


wherein G is R₃, or it is a protecting group selected from alkoxycarbonyl, aryloxycarbonyl, arylalkenyl, alkylalkenyl group, preferably ethoxycarbonyl, Á-methylvinyl, Á-phenylvinyl, A is absent, R₁, R₂, R₃ and B are as hereinbefore defined and X is a leaving group such as halogen, methansulfonate or 4-methylbenzensulfonate, is reacted with a compound of formula (III)



wherein R₄, m and n are as hereinbefore defined. The reaction may be conveniently carried out in solvents such as alcohols, ketones, benzene, ethyl acetate, acetonitrile, dioxane, chloroform, dimethylformamide at a temperature ranging from 0°C to 150°C, preferably at 5°C or at the boiling point of the same solvent. The presence of an acid acceptor such as sodium carbonate, triethylamine and the like may be useful. When G represents an alkoxy- or aryloxy-protecting group, it may be either conveniently removed during the process or it may be cleared by subsequent treatment with aqueous alkali such as diluted sodium hydroxyde, diluted potassium hydroxyde, sodium or potassium carbonate, sodium or potassium hydrogencarbonates; in the


case G is arylalkenyl or alkylalkenyl group it may be removed by subsequent treatment with acids such as aqueous hydrochloric or sulphuric acid; in every instance choice products of general formula (I) in which R_3 is H are obtained.

The compounds of general formula (II), used as starting materials in the above mentioned process, may be prepared by reacting a compound of general formula (IV)



wherein R_1 , R_2 and G are as hereinbefore defined, with an alkyldihalide or a haloalkanole in the presence of a strong base, such as sodium hydride in an aprotic solvent such as tetrahydrofuran or dimethylformamide, or solid potassium hydroxide in dimethylformamide at a temperature ranging from 20°C and 100°C, or in the presence of aqueous alkali such as sodium or potassium hydroxide in the presence of an organic solvent unsoluble with water, such as methylene chloride, benzene or toluene and in the presence of a catalytic amount of an phase transfer catalyst such as ammonium quaternary salt, at a temperature ranging from 20°C and the boiling point of the same solvent. When a haloalkanol is used, the hydroxyl group of the obtained product is changed into methansulfonate or 4-methylbenzensulfonate by the treatment with methansulfonylchloride or with 4-methylbenzensulfonylchloride to give the compound of general formula III. The compounds of general formula IV may in turn be prepared by methods known in the literature such as for example those exemplified in J. Org. Chem. 38, 3498-502 (1973); or

b) a compound of general formula (V)




wherein R_1 , R_2 , R_4 , A , B , m and n are as hereinbefore defined, is reacted with a carbonyl derivative of formula (VI)



wherein Y and Y' are leaving groups identical or different from each other such as halogen, halogenated alkoxy, alkoxy, aryloxy or heterocycle. Preferred groups are chlorine, trichloromethoxy, methoxy, ethoxy or imidazolyl. The reaction may be generally carried out in an aprotic solvent such as tetrahydrofuran, methylene chloride, chloroform, acetone, acetonitrile, benzene, toluene, ethylacetate, carbon tetrachloride or dimethylformamide, optionally in the presence of an acid acceptor, such as triethylamine,


pyridine, sodium or potassium carbonate at a temperature between 0°C and 100°C, preferably at room temperature.

Compounds of general formula V, used as starting materials in the above described process may be prepared by reducing a compound of general formula VII




wherein R_1 , R_2 , R_4 , A, B, m and n are as hereinbefore defined, with hydrogen or a hydrogen donor such as ammonium formate, cyclohexene, cyclohexadiene or hydrazine. The reduction is preferably carried out with hydrogen in the presence of a suitable catalyst, preferably 5% or 10% Pd on charcoal or Raney nickel in the presence of a suitable solvent such as methanol, ethanol, toluene, water or a mixture of them. The reaction is preferably carried out at room pressure and temperature. The same reduction may be conveniently carried out with iron in acidic medium, for example hydrochloric acid, optionally in the presence of $FeCl_3$, or with Zn in acetic or hydrochloric acid, with $SnCl_2$

in hydrochloric acid or with other reducing agents such as titanium trichloride, ferrous sulphate, hydrogen sulphide or its salts, sodium hydrosulphide. When A is absent, the compounds of formula VII may be conveniently prepared by reacting a compound of general formula (VIII)




with a compound of general formula (IX)



wherein R₁, R₂, R₄, B, m and n are as hereinbefore defined and Hal is a leaving group such as halogen, preferably chlorine. The reaction may be conveniently carried out with inert solvents such as butanol, isopropanol, ethanol and like or without solvents at a temperature between 50°C and 200°C.


The compounds of general formula (IX) may be, in turn, conveniently prepared for example by reducing the

corresponding nitrile of general formula (X)




wherein R_4 , m and n are as hereinbefore defined and B contains a carbon atom less in comparison with the above defined. The reaction may be conveniently carried by catalytic hydrogenation in the presence of ammonia or of acids, such as hydrochloric acid in the presence of a catalyst such as Ni-Raney, platinum dioxide and like. Alternatively the nitriles of general formula (X) may be reduced with metal hydride such as lithium aluminium hydride or with diborane.


When A represents a carbonyl group CO , the compounds of general formula (VII) may be prepared by reacting a compound of formula (XI)




with a compound of formula (XII)




wherein R_1 , R_2 , R_4 , B, m, n and Hal are as hereinbefore defined and A is carbonyl group. The reaction is carried out in an aprotic solvent such as tetrahydrofuran, acetonitrile, chloroform, toluene, chlorobenzene or without solvents and, optionally, in the presence of an acid acceptor, preferably in pyridine at a temperature between 20°C and 100°C, preferably between 20°C and 80°C. Compounds of formula (XII) may be prepared by known methods which are well known to anyone skilled in the art. When A represents a carboxyamidic group -CONH-, the compounds of general formula (VII) may be prepared by reacting a compound of general formula (XIII)



with a compound of formula (IX), wherein R_1 and R_2 are as above described. The reaction may be conveniently carried out in an aprotic solvent such as tetrahydrofurane, chloroform, toluene, benzene, cyclohexane at a temperature between 0°C and 80°C , preferably between 5°C and 30°C ; or
c) when it is desired to prepare compounds of formula (I) wherein A is absent or it represents a carbonyl group, a compound of general formula (XIV)



wherein R_1 , R_2 and R_3 are as hereinbefore defined and M is a metal atom, such as sodium, potassium or lithium, preferably sodium, is reacted with a compound of formula XV




wherein Hal, B, m, n, A and R_4 are as above described.

The reaction is preferably carried out in a polar aprotic solvent, such as dimethylformamide, tetrahydrofurane or pyridine at a temperature ranging from 0°C to 100°C ,

preferably at room temperature.

Compound of formula XIV is generated "in situ" from the corresponding hydrogen compounds by means of sodium, potassium, sodium hydride, potassium hydride, potassium hydroxide, sodium hydroxide, potassium tert-butyrate, butyllithium, lithium diisopropylamide, preferably sodium hydride; in case sodium or potassium hydroxide in aqueous concentrated solution are used, the reaction may be conveniently carried out in the presence of an inorganic insoluble solvent such as methylene chloride, in the presence of a phase transfer catalyst, such as a suitable ammonium quaternary salt at a temperature between 20°C and 50°C. Compounds of general formula (XV) wherein A is absent or represents a carbonyl group may be prepared from suitable starting compounds by methods which are well known to anyone skilled in the art, or

d) when it is desired to prepare compounds of formula (I) wherein A is a -CONH- group, a compound of formula XVI



wherein R_1 , R_2 , and R_3 are as above defined, L represents a leaving group such as halogen or alkoxy, preferably chlorine, methoxy or ethoxy, is reacted with a compound of

formula IX. The reaction is carried out in an inert aprotic solvent such as tetrahydrofuran, methylene chloride, ethyl acetate, acetonitrile, acetone, benzene, optionally in the presence of an organic or inorganic acid acceptor such as triethylamine, pyridine, sodium or potassium carbonate at a temperature ranging from -10°C to the boiling point of the selected solvent, preferably at room temperature. Compounds of general formula XVI can be prepared with known methods from suitable starting compounds. Examples of compounds of general formula XVI may be found in EP 309,423, US 4,061,861 and in J. Org. Chem. 38, 3498-502 (1973).

It has to be understood that compounds of general formula (I) containing an R_1 , R_2 , R_3 and R_4 group which may give rise to another R_1 , R_2 , R_3 and R_4 group, are useful novel intermediates. Some of these transformations may also occur in the intermediates for compounds of general formula (I).

Some examples of such conversions, which obviously are not exhaustive of all possibilities, are:

- 1) a nitro group may be transformed into an amino group by reduction.
- 2) an amino group may be transformed into a C_{1-6} acylamino group by acylation with a suitable carboxylic acid derivative.
- 3) an amino group may be transformed into a C_{1-4} alkyl N-mono or di-substituted group by alkylation.
- 4) an amino group may be transformed into a C_{1-6}

alkoxy carbonyl amino group by reaction with a suitable reactive C₁₋₆ alkyl carbonic acid monoester derivative.

- 5) a carboxyl group may be transformed into a C₁₋₆ alkoxy carbonyl group, or into a carbamoyl group optionally C₁₋₄ alkyl N-mono or di-substituted by reaction of a suitable reactive carboxylic acid derivative with appropriate alcohols and amines.
- 6) a carbamoyl group may be transformed into a cyano group by dehydration.
- 7) a C₁₋₆ alkyl thio or a C₁₋₆ alkyl sulphinyl group may be transformed into a C₁₋₆ alkyl sulphinyl or a C₁₋₆ alkylsulphonyl group by oxidation.
- 8) an aromatic hydrogen group may be transformed into a nitro group by nitration.
- 9) a hydrogen group may be transformed into a halogen group by halogenation.
- 10) A product of general formula I where R₃ is H, may be transformed in a product of formula I where R₃ is C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl by alkylation with a suitable alkyl halide in the presence of a strong base such as sodium or potassium hydroxide, sodium or potassium hydride, potassium t-butilate in an aprotic solvent such as dimethylformamide or tetrahydrofuran at a temperature between 20°C and 100°C. When aqueous concentrated solutions of sodium or potassium

hydroxide are used, the reaction may be conveniently carried out in the presence of an unsoluble organic solvent, such as methylene chloride in the presence of phase transfer catalyst such as a suitable ammonium quaternary salt at a temperature between 20°C and 50°C.

- 11) a tertiary amino group may be transformed into a quaternary ammonium derivative by reaction with a suitable alkylating agent such as methyl bromide or methyl iodide.

These transformations are well known to any expert of the branch.

The compounds of the general formula (I) prepared according to the above methods may optionally be converted by inorganic or organic acids into non-toxic, physiologically acceptable acid addition salts, for example by conventional methods such as by reacting the compounds as bases with a solution of the corresponding acid in a suitable solvent. Examples of non-toxic physiologically acceptable acid addition salts are those formed with hydrochloric, nitric, sulfuric, maleic, fumaric, citric, tartaric, methansulphonic, acetic, benzoic, succinic, gluconic, lactic, glycinic, malic, mucoic, glutammic, isethionic, phosphoric, ascorbic or sulphamic acid. Particularly preferred acids are hydrochloric, maleic and fumaric acid.

Particularly preferred compounds according to the present invention are:

1-[2-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)ethyl]-2,3-dihydro-1H-benzimidazol-2-one (Compound 3)

1-[4-(4-(3-chloro-phenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one (Compound 4)

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one (Compound 8)

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-3-methyl-2,3-dihydro-1H-benzimidazol-2-one (Compound 9)

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-3-isopropyl-2,3-dihydro-1H-benzimidazol-2-one (Compound 15)

1-[3-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)propyl]-2,3-dihydro-1H-benzimidazol-2-one (Compound 18)

6-methoxy-1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one (Compound 25)

1-[4-(4-(1-naphthyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one (Compound 30)

As already mentioned hereinbefore, the new compounds of formula (I), according to the present invention, show interesting pharmacological properties owing to their

activity on CNS serotonergic receptors, particularly 5-HT_{1A} and 5-HT₂ receptor subtypes. Therefore the new compounds are commercially useful in the prevention and in the treatment of disorders wherein the altered functionality of 5-HT_{1A} and 5-HT₂ receptors, as above mentioned, is involved.

The biochemical and pharmacological profile of the compounds object of the present invention was assessed by evaluating their affinity for 5-HT_{1A} and 5-HT₂ receptors and their efficacy was established: a) in inducing the well-known behavioural syndrome due to the stimulation of 5-HT_{1A} receptors and b) by evaluating the antagonism towards the behavioural syndrome induced by quipazine stimulating the 5-HT₂ receptors.

RECEPTOR BINDING STUDIES

Receptor binding studies on 5-HT_{1A} and 5-HT₂ receptors were carried out to determine the affinity of the test compounds.

5-HT_{1A} RECEPTOR

- Tissue preparation

Rats (male Sprague Dawley, 200-250 g) were used. The Hippocampi of these animals were homogenized in 10 volumes of ice cold TRIS buffer (pH 7.4). The homogenate was diluted 1:400 (w:v) in the same buffer to have a final protein concentration of about 200 µg/mL, filtered and incubated at 37°C for 10 min, before use.

- Binding assay

Displacement experiments were performed by incubating the homogenate (980 µL) in the presence of [³H]-8OH-DPAT (1.0-

1,5 nM) (10 μ L) and of different concentrations of the test compounds dissolved in the test buffer (10 μ L), at 30°C for 15 min (final volume: 1 mL).

Non specific binding was determined in the presence of 100 μ M 5-HT (10 μ L). The separation of [3 H]-8-OH-DPAT, free from that bound to the receptor, was carried out by the filtration technique (GF/B filters, Whatman). The radioactivity present was counted by liquid scintillation spectrometry.

- Data analysis

The affinity values (K_i) for the compounds were obtained by a non linear least squares regression analysis on the basis of a one binding site model. The values were corrected on the basis of the radioligand occupancy on the receptors according to the equation: $K_i = IC_{50}/(1 + [C]/K_D)$, where $[C]$ and K_D represent the concentration and the dissociation constant, respectively, of the radioligand used ([3 H]-8-OH-DPAT).

5-HT₂ RECEPTOR

- Tissue preparation

Rats (male Sprague Dawley, 200-250 g) were used. Cerebral cortices were homogenized in 10 volumes of ice cold 0.32 M sucrose. After the centrifugation of the homogenate (1,000 x g for 10 min) the supernatant was then recentrifuged at 48,000 x g for 15 min. The resulting pellet was resuspended in 10 volumes of 50 mM TRIS buffer (pH 7.4), incubated at 37°C for 10 min and recentrifuged at 48,000 x g for 15 min.

The residue was then resuspended in 10 volumes of 50 mM TRIS buffer (pH 7.4).

- Binding assay

The tissue was diluted 1:100 (w:v) in 50 mM TRIS buffer (pH 7.4) to have a final protein concentration of about 200 $\mu\text{g/mL}$.

Displacement experiments were performed by incubating the homogenate (980 μL) in the presence of [^3H]-Ketanserine (0.5-1.0 nM) (10 μL) and of different concentrations of the test compounds dissolved in the assay buffer (10 μL), at 37°C for 10 min (final volume: 1 mL).

Non specific binding was determined in the presence of 100 μM Methysergide (10 μL). The separation of [^3H]-Ketanserine free from that bound to the receptor was carried by the filtration technique (GF/B filters, Whatman). The radioactivity present was counted by liquid scintillation spectrometry.

- Data analysis

The affinity values (K_i) for the compounds were obtained by non linear least squares regression analysis on the basis of a one binding site model. These values were corrected on the basis of the radioligand occupancy on the receptors according to the equation: $K_i = \text{IC}_{50} / (1 + [\text{C}] / K_D)$, where $[\text{C}]$ and K_D represent the concentration and the dissociation constant, respectively, of the radioligand used ([^3H]-Ketanserine).

The results of some of the compounds of the present invention on the affinity to the 5-HT_{1A} and 5-HT₂ receptors are reported in Table 1.

TABLE 1 - AFFINITY FOR 5-HT_{1A} AND 5-HT₂ RECEPTORS

Compound	K _i (nM)	
	5-HT _{1A}	5-HT ₂
1	13	36
3	50	133
4	74	1.4
8	13	9.5
9	23	14
10	5	30
15	80	20
17	30	8
18	25	4
22	15	15
24	12	65
25	100	25
30	10	7

Animal studies

Behavioural syndrome

This syndrome, which relates to the stimulation of 5-HT_{1A} receptors and has been described by Goodwin and Green (1985), consists in flat posture, forepaw treading and hindlimb abduction. A control animal does not show this behavioural pattern. The test consists of administering the compound and registering the presence of the above mentioned

symptoms within 50 min giving them a score. The results are expressed as the sum of said scores for each rat (Tab. 2).

TABLE 2 - INDUCTION OF 5-HT_{1A} RELATED SYNDROME

Compound	Dose mg/kg/ip	Total score
VEHICLE	-	0
1	8	7.3 ± 0.9
3	16	5.3 ± 1.2
4	8	0.8 ± 0.5
	16	3.3 ± 1.0
8	8	3.0 ± 2.7
9	8	2.0 ± 0.6
10	8	3.0 ± 0.4
17	8	1.5 ± 0.7
	32	10.3 ± 2.9
18	16	3.8 ± 2.2
	32	12.0 ± 3.2
22	32	3.8 ± 1.3
24	8	4.3 ± 1.1
25	8	2.0 ± 1.2

Values represent mean ± s.e. from 4 rats

Antagonism of Quipazine-induced head twitches

Head twitches depend on the stimulation of 5-HT₂ receptors (Goodwin and Green (1985)). The test consists of administering

the compound in quipazine-treated animals and scoring the number of head twitches within 20 minutes (Table 3).

TABLE 3 - DOSE OF COMPOUND (ID₅₀) WHICH ANTAGONIZES THE SYNDROME INDUCED BY QUIPAZINE

Compound	ID50 μg/kg/ip
1	720
3	498
4	250
8	385
9	1720
10	3300
17	178
18	170
22	8200
24	102
25	1420

According to a further feature of the present invention there are provided pharmaceutical compositions comprising as active ingredient at least one compound of formula (I), as hereinbefore defined, or a physiologically acceptable acid addition salt thereof in association with one or more pharmaceutical carriers, diluents or excipients. For pharmaceutical administration the compounds of general formula (I) and their physiologically acceptable acid addition salts may be incorporated into the conventional pharmaceutical preparations in solid, liquid or spray form. The compositions may, for example, be presented in a form suitable for oral, rectal, parenteral administration or for nasal inhalation. Preferred forms include, for example, capsules, tablets, coated tablets, ampoules, suppositories and nasal spray.

The active ingredient may be incorporated in excipients or carriers conventionally used in pharmaceutical compositions such as, for example, talc, arabic gum, lactose, gelatine, magnesium stearate, corn starch, aqueous or non-aqueous vehicles, polyvinylpyrrolidone, semisynthetic glycerides of fatty acids, benzalcon chloride, sodium phosphate, EDTA, polysorbate 80.

In order to increase the solubility of the compounds of general formula (I) or their physiological acceptable salts, surfactants, non-ionic surfactants such as PEG 400, cyclodextrins, metastable polymorphs, inert absorbents such as bentonite may be incorporated. Furthermore some techniques

may be employed by preparing for example eutectic mixtures and/or solid dispersions by using mannitol, sorbitol, saccharose, succinic acid, or physical modified forms by using hydrosoluble polymers, PVP, PEG 4000-20000.

The compositions are advantageously formulated in dosage units, each dosage unit being adapted to supply a single dose of the active ingredient. Each dosage unit may conveniently contain from 0,01 mg to 100 mg and preferably from 0,1 mg to 50 mg.

The following examples illustrate the preparation of some new compounds according to the present invention and will enable other skilled in the art to understand it more completely. It should be understood, however, that the invention is not limited solely to the particular examples given below.

Description 1

1-(6-Chlorohexyl)-2,3-dihydro-1H-benzimidazol-2-one

The above mentioned compound was prepared analogously to the procedure described in J. Het. Chem. 18, 85 (1981) from 1-(β -phenylvinyl)-2,3-dihydro-1H-benzimidazol-2-one and 1,6-dichloro hexane. The protecting group was then removed by acid hydrolysis with hydrochloric acid following the above procedure. M.p. 80-84°C.

Analogously were prepared:

1-(2-Chloroethyl)-2,3-dihydro-1H-benzimidazol-2-one

M.p. 146-148°C

1-(3-Chloropropyl)-2,3-dihydro-1H-benzimidazol-2-one

M.p. 113-115°C

1-(4-Chlorobutyl)-2,3-dihydro-1H-benzimidazol-2-one

M.p. 80-81°C

Description 2

1-(α -phenylvinyl)-3-n-hexyl-2,3-dihydro-1H-benzimidazol-2-one

The above mentioned compound was prepared similarly to the procedure described in J. Het. Chem. 18, 85 (1981) for the preparation of the analogues bearing in position 1 a methyl, ethyl, allyl and isopropyl residue. The compound may be obtained from 1-(α -phenylvinyl)-2,3-dihydro-1H-benzimidazol-2-one and 1-bromohexane. The compound was used as such without further purification.

Description 3

1-n-hexyl-2,3-dihydro-1H-benzimidazol-2-one

The compound was prepared similarly to the procedure described in J. Het. Chem. 18, 85 (1981) for the preparation of the analogues bearing in position 3 a methyl, ethyl, allyl and isopropyl residue. The compound may be obtained from 1-(α -phenylvinyl)-3-n-hexyl-2,3-dihydro-1H-benzimidazol-2-one by acid hydrolysis with hydrochloric acid. The compound was used as such without further purification.

Description 4

1-(4-Chlorobutyl)-3-n-hexyl-2,3-dihydro-1H-benzimidazol-2-one

The compound may be prepared according to the procedure described in J. Het. Chem. 18, 85 (1981) from 3-n-hexyl-2,3-dihydro-1H-benzimidazol-2-one and 1,4-dichlorobutane.

Analogously may be prepared:

1-(4-Chlorobutyl)-3-methyl-2,3-dihydro-1H-benzimidazol-2-one

1-(4-Chlorobutyl)-3-isopropyl-2,3-dihydro-1H-benzimidazol-2-one

1-(4-Chlorobutyl)-3-allyl-2,3-dihydro-1H-benzimidazol-2-one

1-(6-Chlorohexyl)-3-ethyl-2,3-dihydro-1H-benzimidazol-2-one

All the above mentioned compounds were used as such without further purification.

Description 5

N-(2-Amino-4-methoxyphenyl)-ethyl carbamate

A solution of N-(2-nitro-4-methoxyphenyl)-ethyl carbamate (4 g) (prepared by allowing 4-methoxy-2-nitroaniline to react with ethylchloroformate in pyridine at reflux for 4 hrs, m.p. 56-59°C) in absolute ethanol (150 ml) was hydrogenated at room pressure and temperature in the presence of 10% palladium on charcoal (0,2 g). After absorption of the calculated amount of hydrogen was over, the catalyst was

filtered on celite and the alcoholic solution was evaporated. The desired compound (3,8 g) was obtained as a solid. M.p. 74-76°C.

Description 6

5-Methoxy-2,3-dihydro-2-oxo-1H-benzimidazol-1-ethyl carboxylate

A solution of N-(2-amino-4-methoxyphenyl)ethyl carbamate (0,5 g) and triethylamine (0,4 ml) in CH_2Cl_2 (10 ml), was dropped into a solution of trichloromethylchloroformate (0,32 ml) in CH_2Cl_2 (5 ml) under stirring at 5°C. When the addition was over, the reaction was allowed to reach room temperature and stirring was continued for 1 hour. Water was then added and the product was extracted with CH_2Cl_2 . After evaporation of the solvent the solid residue was purified by washing with diethyl ether. 0,2 g of the desired product was obtained. M.p. 176-178°C.

Description 7

5-Methoxy-3-(2-bromoethyl)-2,3-dihydro-2-oxo-1H-benzimidazol-1-ethyl carboxylate

To a suspension of 80% sodium hydride (0,38 g) in anhydrous dimethylformamide (45 ml) 5-methoxy-2,3-dihydro-2-oxo-1H-benzimidazol-1-ethyl carboxylate (3 g) was added portionwise. After the mixture was stirred for 1 hour at room temperature, a solution of 1,2-dibromoethane (1,1 ml) in 6 ml of dimethylformamide was added. Stirring was kept on

12 hours at room temperature. The water was then added and the product which separated was collected by filtration. The raw solid was purified by chromatography on Silicagel; eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}/32\% \text{NH}_4\text{OH}$ 98:2:02. 1,0 g of the desired product were obtained. M.p. 118-120°C.

Similarly, starting from 1,4-dibromobutane, 5-methoxy-3-(4-bromobutyl)-2,3-dihydro-2-oxo-1H-benzimidazole-1-ethyl-carboxylate may be prepared.

Description 8

1-(3-Trifluoromethylphenyl)-hexahydro-1,4-diazepine

i) Tri-(3-trifluoromethylphenyl)bismuth

3-Trifluoromethylbromobenzene (3.1 ml) dissolved in dry diethyl ether (100 ml) was cautiously added in 60 min to a suspension of magnesium (0.6 g) in the same solvent (5 ml), to which a small crystal of iodine was previously added. The formation of the Grignard reagent was induced by local heating and, once started, the reaction was continued at reflux for further 60 min.

After cooling to 0°C, previously dried bismuth trichloride (3,5 g) was added portionwise and then reflux and stirring was continued for 3 hrs. Water was then added and the intermediate tri-(3-trifluoromethylphenyl)bismuth was extracted into ethyl acetate. After drying, evaporation of the solvent under vacuum, left 3,9 g of the desired

compound as an oil. The compound was further purified by column chromatography on Silicagel using hexane/ethylacetate 95:5 as the eluent. Yield 2,46 g. Yellow oil. MS (C.I.): $[M+H]^+$ 645, 625, 498, 353 m/z.

ii) 1-(3-Trifluoromethylphenyl) piperidin-4-one

Piperidin-4-one monohydrochloride, monohydrate (3.0 g) was dissolved in water (15 ml) and 10% sodium hydroxide (8 ml) was added. The free base was extracted four times into methylene dichloride (150 ml); after separation of the layers, the organic phase was dessicated over $MgSO_4$.

In another flask, tri-(3-trifluoromethylphenyl) bismuth (6 g) was dissolved in dry methylenedichloride (70 ml), and copper acetate (1.55 g) was added. The previously prepared solution of piperidin-4-one was then added dropwise under stirring at room temperature and under a nitrogen stream. The reaction mixture became light blue and then turned green. Stirring was kept on for 2 days, then water was added. The insoluble material that separated was filtered off, the organic layer was separated, dessicated over $MgSO_4$ concentrated to dryness under vacuum. After chromatographic purification on Silicagel with eluent hexane/ethylacetate 80:20, 1.26 g of the desired compound were obtained.

MS (C.I.): $[M+H]^+$ 244, 224 m/z.

iii) 1-(3-Trifluoromethylphenyl)hexahydro-1,4-diazepin-5-one

1-(3-trifluoromethylphenyl)piperidin-4-one (0.4 g) was dissolved into a mixture of glacial acetic (2.5 ml) and concentrated sulphuric acid (1.5 ml). The reaction mixture was cooled to 0°C and sodium azide (118 mg) was added portionwise in eight hours. Stirring was continued overnight and then solid sodium hydroxide in pellets was added under external cooling until pH 8-10 was reached. A small amount of water was also added. The desired compound was extracted into chloroform, the organic phase was dessicated over magnesium sulphate and evaporation of the solvent left 0.36 g of a white solid. M.p. 114-115°C. MS (C.I.): $[M+H]^+$ 259, 239 m/z.

iiii) 1-(3-Trifluoromethylphenyl)hexahydro-1,4-diazepine

A solution of 1-(3-trifluoromethylphenyl)-hexahydro-1,4-diazepin-5-one (0.36 g) in anhydrous tetrahydrofuran was added dropwise to a suspension of $LiAlH_4$ (0.11 g) in the same solvent (10 ml) at room temperature under stirring. Stirring was continued for 3 hrs at room temperature and 4 hrs at reflux. After cooling, water was cautiously added and then the reaction mixture was filtered; the solvent was removed under vacuum and the compound was purified by column chromatography. Yield 0,11 g. Oil.
MS (C.I.): $[M+H]^+$ 245, 225 m/z.

Description 9

7-Methoxy-1-naphtylpiperazine was prepared according to J. Med. Chem. 32, 1921-26 (1989).

Example 1

1-[4-(4-(2-methoxy-phenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 1)

A mixture of 1-(4-chlorobutyl)-2,3-dihydro-1H-benzimidazol-2-one (2 g) and of 1-(2-methoxyphenyl)piperazine hydrochloride (2,03 g) with sodium carbonate (1,88 g) and potassium iodide (0,01 g) in absolute ethanol (100 ml) was refluxed for 18 hours. After filtering the inorganic salts and evaporating the solvent the residue was dissolved in diluted HCl and washed with ethyl acetate. The aqueous phase was made strongly alkaline with 30% NaOH and the product, which separated, was extracted into ethyl acetate. After dehydration, the solvent was removed under vacuum, and a white solid was obtained; it was treated with diethylether, filtered and recrystallized from isopropanol. 2,1 g of the desired compound were obtained. M.p. 160-161°C.

Analysis

$C_{22}H_{28}N_4O_2$

Found % C 69,00 H 7,44 N 14,15

Calc. % C 69,45 H 7,42 N 14,73

1H NMR ($CDCl_3$) 9.82 (s, 1H), 7.1-6.7 (8H), 3.93 (t, 2H), 3.84 (s, 3H), 3.1-2.9 (4H), 2.8-2.5 (4H), 2.45 (t, 2H), 1.9-1.4 (4H)

Following the above described process and using the appropriate benzimidazol-2-one derivatives and arylpiperazine the following may be prepared:

1-[4-(3-(3-chlorophenyl)piperazin-1-yl)propyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 2)

Dihydrochloride salt (isopropanol).

M.p. 165-170°C.

Analysis

$C_{20}H_{23}ClN_4O \cdot 2HCl$ Found % C 53,48 H 5,71 N 12,39

Calc. % C 54,13 H 5,68 N 12,62

1H NMR (DMSO- d_6 /CDCl $_3$ 5:2) 11.09 (b, 1H), 10.81 (s, 1H), 7.3-6.7 (8H), 4.71 (s, 1H + H $_2$ O), 3.94 (t, 2H), 4.1-3.0 (10H), 2.26 (m, 2H)

1-[2-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)ethyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 3)

Hydrochloride salt (isopropanol) M.p. 230-231°C

Analysis

$C_{20}H_{21}F_3N_4O \cdot HCl$ Found % C 56,37 H 5,20 N 13,12

Calc. % C 56,27 H 5,20 N 13,13

1H NMR (DMSO- d_6 /CDCl $_3$ 5:2) 11.09 (b, 1H), 11.04 (s, 1H), 7.5-6.9 (8H), 4.36 (t, 2H), 4.1-3.1 (10H)

1-[4-(4-(3-chlorophenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 4)

Hydrochloride salt (isopropanol) M.p. 217-220°C

Analysis

$C_{21}H_{25}ClN_4O \cdot 2HCl$ Found % C 54,87 H 5,87 N 12,34
Calc. % C 55,09 H 5,94 N 12,24

1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 11.01 (b, 1H), 10.92 (s, 1H),
7.3-6.8 (8H), 4.42 (b, 1H), 4.0-3.8 (4H), 3.50 (d, 2H), 3.3-
3.0 (6H), 1.9-1.6 (4H)

1-[4-(4-(pyrimidin-2-yl)piperazin-1-yl)butyl]-2,3-dihydro-
1H-benzimidazol-2-one

(Compound 5)

Hydrochloride salt (isopropanol) M.p. 237-242°C

Analysis

$C_{19}H_{24}N_6O \cdot 2HCl$ Found % C 53,47 H 6,34 N 19,51
Calc. % C 53,65 H 6,16 N 19,76

1H NMR (CDCL $_3$) 9.75 (s, 1H), 8.29 (d, 2H), 7.2-6.9 (4H),
6.47 (m, 1H), 4.1-3.7 (6H), 2.7-2.4 (6H), 2.1-1.5 (4H). The
spectrum was recorded on the compound in the form of free
base.

1-[2-(4-(3-chlorophenyl)piperazin-1-yl)ethyl]-2,3-dihydro-
1H-benzimidazol-2-one

(Compound 6)

Hydrochloride salt (isopropanol) M.p. 230-231°C

Analysis

$C_{19}H_{21}ClN_4O \cdot 2HCl$ Found % C 53,39 H 5,64 N 13,06
Calc. % C 53,10 H 5,39 N 13,04
 1H NMR (DMSO- d_6) 11.10 (b, 2H), 7.35 (m, 1H), 7.26 (m, 1H), 7.1-7.0 (4H), 6.97 (d, 1H), 6.87 (d, 1H), 4.32 (t, 2H), 4.1-3.0 (10H)

1-[2-(4-(2-methoxyphenyl)piperazin-1-yl)ethyl]-2,3-dihydro-1H-benzimidazol-2-one
(Compound 7)

Hydrochloride salt (isopropanol) M.p. 241-242°C

Analysis

$C_{20}H_{24}N_4O_2 \cdot 2HCl$ Found % C 56,00 H 6,24 N 12,79
Calc. % C 56,47 H 6,16 N 13,17
 1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 11.0 (b, 1H), 10.97 (s, 1H), 7.4-6.7 (9H), 4.35 (t, 2H), 3.82 (s, 3H), 4.0-2.9 (10H)

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one
(Compound 8)

M.p. 114-115°C

Analysis

$C_{22}H_{25}F_3N_4O$ Found % C 62,99 H 6,18 N 13,41
Calc. % C 63,15 H 6,02 N 13,39
 1H NMR (CDCL $_3$) 10.16 (s, 1H), 7.34 (m, 1H), 7.2-7.0 (7H), 3.94 (t, 2H), 3.30 (m, 4H), 2.74 (m, 4H), 2.61 (t, 2H), 2.0-1.6 (4H)

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-3-methyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 9)

Hydrochloride salt (isopropanol) M.p. 215-216°C

Analysis

$C_{23}H_{27}F_3N_4O \cdot HCl$ Found % C 59,19 H 6,20 N 12,14
Calc. % C 58,91 H 6,02 N 11,95
 1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 10.81 (b, 1H), 7.5-6.9 (8H),
3.90 (t, 2H), 3.36 (s, 3H), 4.1-3.0 (10H), 2.0-1.6 (4H)

1-[3-(4-(2-methoxyphenyl)piperazin-1-yl)propyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 10)

Hydrochloride salt (isopropanol) M.p. 200-204°C

Analysis

$C_{21}H_{26}N_4O_2 \cdot 2HCl$ Found % C 56,93 H 6,50 N 12,57
Calc. % C 57,41 H 6,42 N 12,75
 1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 11.12 (b, 1H), 10.87 (s, 1H),
7.3-6.9 (8H), 6.75 (s, 1H + HDO), 3.92 (t, 2H), 3.81 (s,
3H), 3.8-3.1 (10H), 2.19 (m, 2H)

1-[4-(4-phenyl-piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 11)

Hydrochloride salt (isopropanol) M.p. 255-259°C

Analysis

$C_{21}H_{26}N_4O \cdot 2HCl$ Found % C 59,32 H 6,69 N 12,99
Calc. % C 59,57 H 6,67 N 13,23
 1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 11.11 (b, 1H), 10.82 (s, 1H),
9.10 (s, 1H + HDO), 7.4-6.7 (9H), 4.0-3.1 (12H), 1.9-1.7
(4H)

1-[6-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)hexyl-2,3-
dihydro-1H-benzimidazol-2-one

(Compound 12)

Hydrochloride salt (isopropanol) M.p. 118-120°C

Analysis

$C_{24}H_{29}F_3N_4O \cdot HCl$ Found % C 57,21 H 6,28 N 11,00
 $\cdot H_2O$ Calc. % C 55,54 H 6,44 N 11,18
 1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 10.89 (b, 1H), 10.76 (s, 1H),
7.6-6.9 (8H), 4.1-3.0 (12H), 2.0-1.2 (8H)

1-[6-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)hexyl]-3-
ethyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 13)

Hydrochloride salt (isopropanol) M.p. 138-139°C

Analysis

$C_{26}H_{33}F_3N_4O \cdot HCl$ Found % C 61,21 H 6,64 N 10,67
Calc. % C 61,11 H 6,71 N 10,96
 1H NMR (CDCL $_3$) 12.91 (b, 1H), 7.4-6.8 (8H), 4.1-2.8 (14H),
1.33 (t, 3H), 2.2-1.2 (8H)

1-[4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl]-3-allyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 14)

Hydrochloride salt (isopropanol) M.p. 201-204°C

Analysis

$C_{25}H_{32}N_4O_2 \cdot 2HCl$ Found % C 61,35 H 7,09 N 11,18
Calc. % C 60,85 H 6,94 N 11,35
 1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 11.07 (b, 1H), 8.04 (s, 1H + HDO), 7.2-6.8 (8H), 5.90 (m, 1H), 5.2-5.0 (2H), 4.48 (d, 2H), 3.92 (t, 2H), 3.82 (s, 3H), 3.7-3.0 (10H), 2.0-1.7 (4H)

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-3-isopropyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 15)

Hydrochloride salt (isopropanol) M.p. 181-184°C

Analysis

$C_{25}H_{31}F_3N_4O \cdot HCl$ Found % C 60,01 H 6,51 N 11,24
Calc. % C 60,42 H 6,49 N 11,27
 1H NMR (CDCL $_3$) 12.85 (b, 1H), 7.4-6.9 (8H), 4.70 (m, 1H), 3.92 (t, 2H), 4.0-2.8 (10H), 2.2-1.8 (4H), 1.53 (d, 6H)

1-[4-(4-(3-chlorophenyl)piperazin-1-yl)butyl]-3-n-hexyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 16)

Hydrochloride salt (isopropanol) M.p. 107-111°C

Analysis

$C_{27}H_{37}ClN_4O \cdot HCl$ Found % C 64,57 H 7,53 N 11,10
Calc. % C 64,15 H 7,58 N 11,08
 1H NMR ($CDCl_3$) 12.80 (b, 1H), 7.2-6.5 (8H), 3.9-2.7 (14H),
2.3-1.2 (12H), 0.87 (m, 3H)

1-[4-(4-(3-chlorophenyl)piperazin-1-yl)butyl]-3-methyl-2,3-
dihydro-1H-benzimidazol-2-one

(Compound 17)

Hydrochloride salt (isopropanol) M.p. 214-216°C

Analysis

$C_{22}H_{27}ClN_4O \cdot HCl$ Found % C 60,88 H 6,58 N 12,86
Calc. % C 60,69 H 6,48 N 12,87
 1H NMR ($CDCl_3$) 12.84 (b, 1H), 7.3-6.7 (8H), 3.93 (t, 2H),
3.42 (s, 3H), 4.0-2.9 (10H), 2.1-1.8 (4H)

1-[3-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)propyl]-
2,3-dihydro-1H-benzimidazol-2-one

(Compound 18)

Hydrochloride salt (isopropanol) M.p. 160-162°C

Analysis

$C_{21}H_{23}F_3N_4O \cdot 2HCl$ Found % C 52,84 H 5,29 N 11,77
Calc. % C 52,84 H 5,28 N 11,74
 1H NMR ($DMSO-d_6/CDCl_3$ 5:2) 11.14 (b, 1H), 10.87 (s, 1H),
7.6-6.9 (8H), 6.64 (s, 1H + HDO), 3.93 (t, 2H), 4.1-3.0
(10H), 2.21 (m, 2H)

1-[4-(4-(2-chlorophenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 19)

Hydrochloride salt (ethanol) M.p. 247-250°C

Analysis

$C_{21}H_{25}ClN_4O \cdot HCl$	Found %	C 59,36	H 6,34	N 12,96
	Calc. %	C 59,86	H 6,22	N 13,30

1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 10.81 (b, 1H), 10.80 (s, 1H), 7.4-6.9 (8H), 3.86 (t, 2H), 3.7-3.1 (10H), 2.0-1.7 (4H)

1-[4-(4-(3-methoxyphenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 20)

Hydrochloride salt (ethanol) M.p. 190-192°C

Analysis

$C_{22}H_{28}N_4O_2 \cdot 2HCl$	Found %	C 58,36	H 6,69	N 12,38
	Calc. %	C 58,28	H 6,67	N 12,36

1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 11.05 (b, 1H), 10.80 (s, 1H), 7.3-7.0 (5H), 6.6-6.4 (3H), 5.40 (s, 1H + HDO), 3.74 (s, 3H), 4.0-3.0 (12H), 2.0-1.6 (4H)

1-[4-(2-(7-methoxynaphth-1-yl)piperazin-1-yl)ethyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 22)

Hydrochloride salt (ethanol) M.p. 240-242°C

Analysis

$C_{24}H_{26}N_4O_2 \cdot HCl$ Found % C 65,49 H 6,31 N 12,58

Calc. % C 65,67 H 6,20 N 12,76

1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 10.97 (s, 1H), 10.63 (b, 1H),
7.9-7.0 (10H), 4.37 (t, 2H), 3.94 (s, 3H), 4.1-3.2 (10H)

1-[4-(4-(5-benzodioxan-piperazin-1-yl)butyl)-2,3-dihydro-1H-
benzimidazol-2-one

(Compound 29)

Hydrochloride salt (isopropanol) M.p. 186-188°C

Analysis

$C_{23}H_{28}N_4O_3 \cdot 2HCl$ Found % C 56,96 H 6,57 N 11,83

Calc. % C 57,38 H 6,28 N 11,64

1H NMR (DMSO- d_6) 10.93 (s, 1H), 10.9 (b, 1H), 7.15 (m,
1H), 7.1-6.9 (3H), 6.76 (m, 1H), 6.6-6.5 (2H), 4.53 (s, 1H +
HDO), 4.24 (s, 4H), 3.83 (t, 2H), 3.6-3.4 (4H), 3.3-3.0
(6H), 1.9-1.6 (4H)

1-[4-(4-(1-naphthyl)piperazin-1-yl)butyl)-2,3-dihydro-1H-
benzimidazol-2-one

(Compound 30)

Hydrochloride salt (isopropanol) M.p. 264-267°C

Analysis

$C_{25}H_{28}N_4O \cdot 2HCl$ Found % C 63,66 H 6,59 N 11,99

Calc. % C 63,42 H 6,39 N 11,83

1H NMR (DMSO- d_6) 10.86 (s, 1H), 10.51 (b, 1H), 8.12 (m,
1H), 7.89 (m, 1H), 7.64 (d, 1H), 7.6-7.5 (2H), 7.42 (d, 1H),
7.2-7.0 (5H), 3.89 (t, 2H), 3.7-3.2 (10H), 2.0-1.7 (4H)

1-[2-(4-(3-trifluoromethyl-phenyl)hexahydro-1H-1,4-diazepin-1-yl)ethyl]-2,3-dihydro-1H-benzimidazol-2-one
(Compound 31)

Hydrochloride salt (Ethylacetate - diethyl ether)

M.p. 128-130°C

Analysis

$C_{21}H_{23}F_3N_4O \cdot HCl$	Found %	C 55,81	H 5,56	N 12,19
	Calc. %	C 57,21	H 5,49	N 12,71

1H NMR ($CDCl_3$) 12.77 (b, 1H), 9.86 (b, 1H), 7.5-6.8 (8H), 4.44 (b, 2H), 4.2-2.0 (10H), 2.40 (m, 2H)

1-[2-(4-phenyl-piperazin-1-yl)ethyl]-2,3-dihydro-1H-benzimidazol-2-one
(Compound 35)

Hydrochloride salt (isopropanol - ethanol)

M.p. 232-234°C

Analysis

$C_{19}H_{22}N_4O \cdot HCl$	Found %	C 62,84	H 6,46	N 15,43
	Calc. %	C 63,59	H 6,46	N 15,61

1H NMR ($DMSO-d_6/CDCl_3$ 5:2) 11.0 (b, 1H), 10.97 (s, 1H), 7.4-6.7 (9H), 4.34 (t, 2H), 4.0-3.1 (10H)

1-[3-(3-phenylimidazolidin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one
(Compound 36)

Example 2

6-methoxy-1-[2-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)
ethyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 23)

5-Methoxy-3-(2-bromoethyl)-2,3-dihydro-2-oxo-1H-benzimidazol-1-ethyl carboxylate (0.6 g) was suspended into a mixture of ethanol (60 ml) and dry dimethylformamide (20 ml), in the presence of sodium carbonate (0.23 g). 3-Trifluoromethyl-phenylpiperazine (0.33 ml) was added dropwise to the suspension at room temperature under stirring, then the reaction mixture was heated to reflux for 14 hrs. The solvents were removed under vacuum and the raw material was purified by column chromatography on Silicagel; eluent methylenedichloride/methanol/32% ammonium hydroxide 98:2:0.2. The title compound was further purified by crystallization from 50% aqueous ethanol. Yield 0.1 g. The hydrochloride was obtained by adding the stoichiometric amount of aqueous hydrochloric acid and freeze-drying.

Hydrochloride salt (water) M.p. 208-210°C

Analysis

$C_{21}H_{23}F_3N_4O_2 \cdot HCl$	Found %	C 55,28	H 5,22	N 11,85
	Calc. %	C 55,21	H 5,29	N 12,26

1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 11.09 (b, 1H), 10.75 (s, 1H), 7.5-6.8 (6H), 6.57 (m, 1H), 4.31 (t, 2H), 3.79 (s, 3H), 4.1-3.0 (10H)

Similarly were prepared:

6-methoxy-1-[4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 24)

Hydrochloride salt (ethanol) M.p. 163-165°C

Analysis

$C_{23}H_{30}N_4O_3 \cdot HCl$ Found % C 61,55 H 6,81 N 12,70
Calc. % C 61,80 H 6,99 N 12,53

1H NMR (DMSO- d_6 /CDCl $_3$ 5:2) 10.96 (b, 1H), 10.61 (s, 1H), 7.2-6.6 (6H), 6.53 (m, 1H), 3.81 (s, 3H), 3.77 (s, 3H), 3.9-3.0 (12H), 1.9-1.6 (4H)

6-methoxy-1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 25)

Hydrochloride salt (ethanol) M.p. 122-124°C

Analysis

$C_{23}H_{27}F_3N_4O_2 \cdot HCl$ Found % C 56,79 H 5,74 N 11,38
Calc. % C 56,97 H 5,82 N 11,55

1H NMR (DMSO- d_6 /CDCl $_3$ 5:2) 11.04 (b, 1H), 10.59 (s, 1H), 7.5-6.7 (6H), 6.53 (m, 1H), 3.77 (s, 3H), 4.1-3.0 (12H), 2.0-1.6 (4H)

Description 10

4-(2-methoxyphenyl)-N-(2-nitro-5-chlorophenyl)-1-piperazin-butanamine

The compound was prepared according to the method described in Farmaco Ed. Sci. 36, 359 (1981) from 2,4-dichloronitrobenzene and 4-(2-methoxyphenyl)-1-piperazinbutanamine. M.p.

227-229°C as hydrochloride salt.

Analogously may be prepared:

4-(2-methoxyphenyl)-N-(2-nitro-4,5-dichlorophenyl)-1-piperazinbutanamine. Oil

4-(2-methoxyphenyl)-N-(2-nitro-5-methylphenyl)-1-piperazinbutanamine hydrochloride salt. M.p. 227°C

4-(2-methoxyphenyl)-N-(2-nitrophenyl)-1-piperazinbutanamine

Description 11

4-(2-methoxyphenyl)-N-(2-amino-5-chlorophenyl)-1-piperazinbutanamine

The compound was prepared according to the method described in Farmaco Ed. Sci. 36, 359 (1981) reducing 4-(2-methoxyphenyl)-N-(2-nitro-5-chlorophenyl)-1-piperazinbutanamine by catalytic hydrogenation. Oil.

Analogously may be prepared:

4-(2-methoxyphenyl)-N-(2-amino-5-methylphenyl)-1-piperazinbutanamine, hydrochloride salt. M.p. 233-235°C

4-(2-methoxyphenyl)-N-(2-amino-4,5-dichlorophenyl)-1-piperazinbutanamine. Oil

4-(2-methoxyphenyl)-N-(2-aminophenyl)-1-piperazinbutanamine

Description 12

N-(2-Nitrophenyl)-4-(3-trifluoromethylphenyl)-1-piperazin-
propionamide

The compound was prepared from 3-bromo-N-(2-nitrophenyl)-propionamide and 1-(3-trifluoromethylphenyl)piperazine according to the method described in J. Med. Chem. 33, 2970 (1990). Monohydrochloride salt. M.p. 185-188°C.

Description 13

N-(2-Aminophenyl)-4-(3-trifluoromethylphenyl)-1-piperazin-
propionamide

The compound was prepared according to the method described in J. Med. Chem. 30, 13 (1987) by catalytic reduction of N-(2-nitrophenyl)-4-(3-trifluoromethylphenyl)-1-piperazin propionamide in the presence of 10% Pd/C. Monohydrochloride salt. M.p. 194-196°C.

Example 3

N-[3-(4-(3-trifluoromethylphenyl)piperazin-1-yl)propionyl]-
2,3-dihydro-1H-benzimidazol-2-one

(Compound 21)

A solution of N-(2-aminophenyl)-4-(3-trifluoromethylphenyl)-1-piperazinpropionamide (4 g) and triethylamine (2 ml) in anhydrous tetrahydrofuran (50 ml) was dropped to a solution of trichloromethylchloroformate (1 ml) in tetrahydrofuran (20 ml) under stirring at 5°C. When the addition was finished, the reaction was allowed to reach the room temperature and the stirring was continued for 1 hour. Then

water was added and the product was extracted with ethyl acetate. After evaporating the solvent, the residue was purified by column chromatography over silica gel and a mixture of CH_2Cl_2 , MeOH, NH_3 90/10/1 as eluent. 1.8 g of the desired product were obtained. The hydrochloride was obtained from ethanol-diethyl ether.

Hydrochloride salt (ethanol-diethyl ether)

M.p. 227-230°C.

Analysis

$\text{C}_{21}\text{H}_{21}\text{F}_3\text{N}_4\text{O}_2 \cdot \text{HCl}$ Found % C 54,65 H 4,76 N 12,10

Calc. % C 55,45 H 4,88 N 12,32

^1H NMR ($\text{DMSO}-d_6/\text{CDCl}_3$ 5:2) 11.43 (s, 1H), 10.84 (b, 1H), 8.02 (m, 1H), 7.6-7.0 (7H), 4.1-3.2 (12H)

Similarly and using the suitable intermediates the following compounds were obtained:

6-chloro-1-[4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 26)

Hydrochloride salt (diethyl ether) M.p. 206-209°C

Analysis

$\text{C}_{22}\text{H}_{27}\text{ClN}_4\text{O}_2 \cdot 2\text{HCl}$ Found % C 54,11 H 6,08 N 10,82

Calc. % C 54,16 H 5,99 N 11,48

^1H NMR ($\text{DMSO}-d_6/\text{CDCl}_3$ 5:2) 10.98 (s, 1H), 10.54 (b, 1H), 7.3-6.8 (8H), 3.81 (s, 3H), 4.0-3.0 (12H), 2.0-1.6 (4H)

5,6-dichloro-1-[4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl]-

2,3-dihydro-1H-benzimidazol-2-one

(Compound 27)

Hydrochloride salt (diethyl ether) M.p. 157-160°C

Analysis

$C_{22}H_{26}Cl_2N_4O_2 \cdot 2HCl$ Found % C 50,19 H 5,60 N 10,52

Calc. % C 50,59 H 5,40 N 10,73

1H NMR (DMSO- d_6 /CDCl $_3$ 5:2) 11.16 (s, 1H), 10.75 (b, 1H),
7.42 (s, 1H), 7.13 (s, 1H), 7.2-6.7 (5H), 3.82 (s, 3H), 4.0-
3.0 (12H), 2.1-1.6 (4H)

6-methyl-1-[4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 28)

Hydrochloride salt (isopropanol) M.p. 210-213°C

Analysis

$C_{23}H_{30}N_4O_2 \cdot 2HCl$ Found % C 59,53 H 7,19 N 11,68

Calc. % C 59,10 H 6,90 N 11,99

1H NMR (DMSO- d_6 /CDCl $_3$ 5:2) 10.98 (b, 1H), 10.66 (s, 1H),
7.08 (b, 1H), 7.1-6.7 (7H), 3.82 (s, 3H), 3.9-3.0 (12H),
2.35 (s, 3H), 2.1-1.7 (4H)

1-[4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

(Compound 1)

M.p. 159-161°C

Analysis

$C_{22}H_{28}N_4O_2$

Found % C 69,30 H 7,50 N 14,50

Calc. % C 69,45 H 7,42 N 14,73

1H NMR ($CDCl_3$) 9.82 (s, 1H), 7.1-6.7 (8H), 3.93 (t, 2H), 3.84 (s, 3H), 3.1-2.9 (4H), 2.8-2.5 (4H), 2.45 (t, 2H), 1.9-1.4 (4H)

Description 14

1-(4-Chlorobutyl)-4-(3-trifluoromethylphenyl)-piperazine

The product was prepared according to the methods described in J. Org. Chem. 24, 764 (1958) from 1-(3-trifluoromethylphenyl)piperazine and 1-chloro-4-bromobutane. The compound was used as such without further purification.

Analogously may be prepared:

1-(6-Chlorohexyl)-4-(3-trifluoromethylphenyl)piperazine

1-(4-Chlorobutyl)-4-(2-methoxyphenyl)piperazine

1-(4-Chlorobutyl)-4-(3-chlorophenyl)piperazine

Example 4

1-[4-(4-(3-chlorophenyl)piperazin-1-yl)butyl]-3-methyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 17)

To a suspension of 80% sodium hydride (1 g) in anhydrous dimethylformamide (50 ml). 1-methyl-benzimidazol-2-one (5 g) was added portionwise. After stirring the reaction mixture for 1 hour at room temperature, a solution of 4-(3-chlorophenyl)-1-chlorobutyl-piperazine (6 g) in dimethylformamide

(15 ml) was added. The reaction mixture was allowed to react for 10 hours at 60°C, then after cooling, water was added and the product was extracted with ethyl acetate. The raw product was purified by column chromatography over silica gel and a mixture CH₂Cl₂/MeOH 95/5 as eluent. 5 g of the desired product were obtained.

Hydrochloride salt (isopropanol) M.p. 213-216°C

Analysis

C₂₂H₂₇ClN₄O . HCl Found % C 60,65 H 6,53 N 12,34

Calc. % C 60,69 H 6,48 N 12,87

¹H NMR (CDCl₃) 12.84 (b, 1H), 7.3-6.7 (8H), 3.93 (t, 2H),

3.42 (s, 3H), 4.0-2.9 (10H), 2.1-1.8 (4H)

Analogously were obtained:

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-3-methyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 9)

Hydrochloride salt (isopropanol) M.p. 215-216°C

Analysis

C₂₃H₂₇F₃N₄O . HCl Found % C 58,74 H 6,08 N 12,03

Calc. % C 58,91 H 6,02 N 11,95

¹H NMR (DMSO-d₆/CDCl₃ 5:2) 10.81 (b, 1H), 7.5-6.9 (8H),

3.90 (t, 2H), 3.36 (s, 3H), 4.1-3.0 (10H), 2.0-1.6 (4H)

1-[6-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)hexyl]-3-ethyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 13)

Hydrochloride salt (isopropanol) M.p. 136-139°C

Analysis

$C_{26}H_{33}F_3N_4O \cdot HCl$ Found % C 60,89 H 6,51 N 11,03

Calc. % C 61,11 H 6,71 N 10,96

1H NMR ($CDCl_3$) 12.91 (b, 1H), 7.4-6.8 (8H), 4.1-2.8 (14H),
1.33 (t, 3H), 2.2-1.2 (8H)

1-[4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl]-3-allyl-2,3-
dihydro-1H-benzimidazol-2-one

(Compound 14)

Hydrochloride salt (isopropanol) M.p. 201-204°C

Analysis

$C_{25}H_{32}N_4O_2 \cdot 2HCl$ Found % C 60,50 H 6,59 N 11,53

Calc. % C 60,85 H 6,94 N 11,35

1H NMR ($DMSO-d_6/CDCl_3$ 5:2) 11.07 (b, 1H), 8.04 (s, 1H +
HDO), 7.2-6.8 (8H), 5.90 (m, 1H), 5.2-5.0 (2H), 4.48 (d,
2H), 3.92 (t, 2H), 3.82 (s, 3H), 3.7-3.0 (10H), 2.0-1.7 (4H)

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-3-
isopropyl-2,3-dihydro-1H-benzimidazol-2-one

(Compound 15)

Hydrochloride salt (isopropanol) M.p. 181-184°C

Analysis

$C_{25}H_{31}F_3N_4O \cdot HCl$ Found % C 60,00 H 6,69 N 11,01

Calc. % C 60,42 H 6,49 N 11,27

1H NMR ($CDCl_3$) 12.85 (b, 1H), 7.4-6.9 (8H), 4.70 (m, 1H),
3.92 (t, 2H), 4.0-2.8 (10H), 2.2-1.8 (4H), 1.53 (d, 6H)

1-[4-(4-(3-chlorophenyl)piperazin-1-yl)butyl]-3-n-hexyl-2,3-

dihydro-1H-benzimidazol-2-one

(Compound 16)

Hydrochloride salt (isopropanol) M.p. 107-111°C

Analysis

$C_{27}H_{37}ClN_4O \cdot HCl$ Found % C 63,87 H 7,37 N 11,27

Calc. % C 64,15 H 7,58 N 11,08

1H NMR ($CDCl_3$) 12.80 (b, 1H), 7.2-6.5 (8H), 3.9-2.7 (14H),

2.3-1.2 (12H), 0.87 (m, 3H)

Description 15

1-(4-aminobutyl)-4-(3-trifluoromethyl-phenyl)-piperazine

Lithium aluminium hydride (1 g) was suspended in 30 ml of anhydrous tetrahydrofuran. 4-[4-(3-trifluoromethylphenyl) piperazin-1-yl]butirronitrile (5,4 g) dissolved in 30 ml of the same solvent was dropped into the suspension of the reducing agent under stirring by cooling at 5°C. Finally, the temperature was allowed to reach the room temperature and the suspension was stirred for all over night. An amount of water, necessary to decompose the reaction complexes, was added. The insoluble material was eliminated by filtration and the residual organic solution was concentrated to dryness under vacuum. The product was purified by column chromatography on 60 Merck silica gel with eluent: methylene chloride/methanol/ 32% ammonium hydroxide 80:20:2. 2,3 g of a colourless oil were obtained.

Analogously may be prepared:

1-(2-aminoethyl)-4-(2-pyrimidinyl)-piperazine. Oil

1-(2-aminoethyl)-4-(3-trifluoromethyl)phenyl-piperazine. Oil

Example 5

N-[2-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)ethyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-carboxamide

(Compound 32)

A mixture of 1-chlorocarbonyl-benzimidazol-2-one (1 g) prepared as described in EP 309423 and 4-(3-trifluoromethyl-phenyl)-1-(2-aminoethyl)piperazine (3,2 g) with Na_2CO_3 (1,57 g) in anhydrous dimethylformamide (50 ml) was heated with stirring at 100°C for 4 hours. The reaction mixture was cooled, diluted with water and extracted with ethyl acetate. After removal of the solvent, the residual solid was converted into the corresponding hydrochloride salt by adding anhydrous hydrogen chloride to a solution of the base in isopropanol. 1,3 g of the desired compound were obtained. Hydrochloride salt (isopropanol) M.p. 230-233°C

Analysis

$\text{C}_{21}\text{H}_{22}\text{F}_3\text{N}_5\text{O}_2 \cdot \text{HCl}$	Found %	C	53,51	H	4,92	N	14,99
	Calc. %	C	53,68	H	4,93	N	14,90

^1H NMR ($\text{DMSO}-d_6/\text{CDCl}_3$ 5:2) 11.60 (s, 1H), 10.80 (b, 1H), 9.01 (t, 1H), 8.00 (m, 1H), 7.5-6.6 (7H), 4.0-2.8 (12H)

Analogously may be obtained:

N-[2-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-carboxamide

(Compound 33)

¹H NMR (DMSO-d₆/CDCl₃ 5:2) 11.57 (s, 1H), 11.09 (b, 1H), 8.83 (t, 1H), 7.98 (m, 1H), 7.6-7.0 (7H), 4.1-3.0 (12H), 2.1-1.6 (4H)

Similarly, starting from 3-methyl-1-chlorocarbonyl-benzimidazol-2-one, prepared as described in EP 309,423, it was obtained

N-[2-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)ethyl]-2-oxo-3-methyl-2,3-dihydro-1H-benzimidazol-1-carboxamide
(Compound 34)

Hydrochloride salt (isopropanol) M.p. 214-215°C

Analysis

C ₂₂ H ₂₄ F ₃ N ₅ O ₂ . HCl	Found %	C 54,40	H 5,25	N 14,39
	Calc. %	C 54,60	H 5,21	N 14,47

¹H NMR (DMSO-d₆) 10.72 (b, 1H), 8.96 (t, 1H), 8.04 (d, 1H), 7.47 (m, 1H), 7.4-7.1 (6H), 3.41 (s, 3H), 4.1-3.1 (12H)

Example 6

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-3-isopropyl-2,3-dihydro-1H-benzimidazol-2-one
(Compound 15)

1-[4-(4-(3-Trifluoromethyl-phenyl)-piperazin-1-yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one (1 g) was added portionwise to a suspension of 85% powdered potassium hydroxide (0,24 g) in dimethylformamide, under stirring at room temperature.

The addition was completed in 10 minutes and the reaction mixture so obtained was stirred at the same temperature for 1 hour. Then, isopropylbromide (0,27 ml) was added and heated for 5 hours at 40°C. The reaction mixture was poured in water and the product was extracted with ethyl acetate. For concentration to dryness the desired product as residual solid was obtained. It was purified by preparing the hydrochloride salt from ethyl acetate.

Hydrochloride salt M.p. 181-184°C

Analysis

$C_{25}H_{31}F_3N_4O \cdot HCl$	Found %	C 60,31	H 6,48	N 11,20
	Calc. %	C 60,42	H 6,49	N 11,27

1H NMR ($CDCl_3$) 12.85 (b, 1H), 7.4-6.9 (8H), 4.70 (m, 1H), 3.92 (t, 2H), 4.0-2.8 (10H), 2.2-1.8 (4H), 1.53 (d, 6H)

Analogously were prepared:

1-[6-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)hexyl]-3-ethyl-2,3-dihydro-1H-benzimidazol-2-one
(Compound 13)

Hydrochloride salt (isopropanol) M.p. 138-139°C

Analysis

$C_{26}H_{33}F_3N_4O \cdot HCl$	Found %	C 61,32	H 6,70	N 10,91
	Calc. %	C 61,11	H 6,71	N 10,96

1H NMR ($CDCl_3$) 12.91 (b, 1H), 7.4-6.8 (8H), 4.1-2.8 (14H), 1.33 (t, 3H), 2.2-1.2 (8H)

1-[4-(4-(2-methoxyphenyl)piperazin-1-yl)butyl]-3-allyl-2,3-

dihydro-1H-benzimidazol-2-one

(Compound 14)

Hydrochloride salt (isopropanol) M.p. 201-204°C

Analysis

$C_{25}H_{32}N_4O_2 \cdot 2HCl$ Found % C 60,93 H 7,01 N 11,24
Calc. % C 60,85 H 6,94 N 11,35
 1H NMR (DMSO- d_6 /CDCL $_3$ 5:2) 11.07 (b, 1H), 8.04 (s, 1H +
HDO), 7.2-6.8 (8H), 5.90 (m, 1H), 5.2-5.0 (2H), 4.48 (d,
2H), 3.92 (t, 2H), 3.82 (s, 3H), 3.7-3.0 (10H), 2.0-1.7 (4H)

1-[4-(4-(3-chlorophenyl)piperazin-1-yl)butyl]-3-n-hexyl-2,3-
dihydro-1H-benzimidazol-2-one

(Compound 16)

Hydrochloride salt (isopropanol) M.p. 108-111°C

Analysis

$C_{27}H_{37}ClN_4O \cdot HCl$ Found % C 64,31 H 7,56 N 11,12
Calc. % C 64,15 H 7,58 N 11,08
 1H NMR (CDCL $_3$) 12.80 (b, 1H), 7.2-6.5 (8H), 3.9-2.7 (14H),
2.3-1.2 (12H), 0.87 (m, 3H)

1-[4-(4-(3-chlorophenyl)piperazin-1-yl)butyl]-3-methyl-2,3-
dihydro-1H-benzimidazol-2-one

(Compound 17)

Hydrochloride salt (isopropanol) M.p. 214-216°C

Analysis

$C_{22}H_{27}ClN_4O \cdot HCl$ Found % C 60,51 H 6,53 N 12,81

Calc. % C 60,69 H 6,48 N 12,87

¹H NMR (CDCL₃) 12.84 (b, 1H), 7.3-6.7 (8H), 3.93 (t, 2H),
3.42 (s, 3H), 4.0-2.9 (10H), 2.1-1.8 (4H)

The following not limitative examples of pharmaceutical compositions according to the invention are given:

Example 7

Tablets

- active ingredient	10 mg
- lactose	187 mg
- corn starch	50 mg
- magnesium stearate	3 mg

Method of preparation: the active ingredient, lactose and corn starch were mixed and homogeneously moistened with water. After screening of the moist mass and drying in a tray drier, the mixture was again passed through a screen and magnesium stearate was added. Then the mixture was pressed into tablets weighing 250 mg each. Each tablet contains 10 mg of active ingredient.

Example 8

Capsules

- active ingredient	10 mg
- lactose	188 mg
- magnesium stearate	2 mg

Method of preparation: the active ingredient was mixed with the auxiliary products, and the mixture was passed through a screen and mixed homogeneously in a suitable device. The resulting mixture was filled into hard gelatine capsules (200 ml per capsule); each capsule contains 10 mg of active ingredient.

Example 9

Ampoules

- | | |
|---------------------|------|
| - active ingredient | 2 mg |
| - sodium chloride | 9 mg |

Method of preparation: the active ingredient and sodium chloride were dissolved in an appropriate amount of water for injection. The resulting solution was filtered and filled into vials under sterile conditions.

Example 10

Suppositories

- | | |
|---|---------|
| - active ingredient | 25 mg |
| - semisynthetic glycerides of fatty acids | 1175 mg |

Method of preparation: the semisynthetic glycerides of fatty acids were melted and the active ingredient was added while stirring homogeneously. After cooling at a proper temperature the mass was poured into preformed moulds for suppositories weighing 1200 mg each. Each suppository contains 25 mg of active ingredient.

Example 11

Nasal spray


- active ingredient	80	mg
- benzalconium chloride	0,1	mg
- sodium chloride	8	mg
- EDTA	1	mg
-sodium phosphate (buffer pH 6,5)	10	mg
polysorbate 80	10	mg
- bidistilled water	q.s. to 2 ml	

Method of preparation: the single components were added in the suitable volume of bidistilled water by stirring until a complete dissolution before an further addition. After taking to volume, the solution was filtered upon sterilising filter, introduced in suitable bottles and blocked up by the opportune dosage system.



CLAIMS

1. Compounds of general formula I



(I)

wherein

R_1 and R_2 may be at the same time or not a hydrogen atom, halogen, trifluoromethyl, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkylthio, C_{1-6} acyl, carboxyl, C_{1-6} alkoxycarbonyl, hydroxy, nitro, amino optionally C_{1-4} alkyl N-mono or di-substituted, C_{1-6} acylamino, C_{1-6} alkoxycarbonylamino, carbamoyl optionally C_{1-4} alkyl N-mono or di-substituted, cyano, C_{1-6} alkylsulphonyl, C_{1-6} alkylsulphonyl, amino sulphonyl optionally C_{1-4} alkyl N-mono or di-substituted, C_{1-4} alkyl N-mono or di-substituted aminosulphonylamino,





aminosulphonylamino;

R₃ is hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl or C₂₋₆ alkynyl;

A is -CO- or -CONH- or it is absent;

B is a straight or branched, saturated or unsaturated C₂₋₆ alkyl;

m and n are both independently an integer from 1 to 3;

R₄ is an aryl, aralkyl, a heteroaryl or heteroaralkyl group, each group being optionally substituted by one or more substituents selected from halogen, trifluoromethyl, cyano, C₁₋₃ alkoxy, C₁₋₄ alkyl and acid addition salts thereof.

2) Compounds of general formula (I) according to claim 1 characterized in that A is absent, B is a straight, saturated C₂₋₄ alkyl, m and n are the integer 2, R₄ is a substituted phenyl ring wherein the substituents are selected from methoxy, chloro or trifluoromethyl, and acid addition salts thereof.

3. Compound of formula (I) selected from

1-[2-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)ethyl]-
2,3-dihydro-1H-benzimidazol-2-one

1-[4-(4-(3-chloro-phenyl)piperazin-1-yl)butyl]-2,3-

This Sheet Has Been Replaced





dihydro-1H-benzimidazol-2-one

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-
2,3-dihydro-1H-benzimidazol-2-one

1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)butyl]-
3-methyl-2,3-dihydro-1H-benzimidazol-2-one

1-[4-(4-(3-trifluoromethylphenyl)piperazin-1-yl)butyl]-3-
isopropyl-2,3-dihydro-1H-benzimidazol-2-one

1-[3-(4-(3-trifluoromethyl-phenyl)piperazin-1-yl)propyl]-
2,3-dihydro-1H-benzimidazol-2-one

6-methoxy-1-[4-(4-(3-trifluoromethyl-phenyl)piperazin-1-
yl)butyl]-2,3-dihydro-1H-benzimidazol-2-one

1-[4-(4-(1-naphthyl)piperazin-1-yl)butyl]-2,3-dihydro-1H-
benzimidazol-2-one


4. Physiologically acceptable acid addition salts of compounds of general formula (I) according to claim 1-3.
5. Salts according to claim 4, characterized in that the physiologically acceptable acids are hydrochloric, maleic or fumaric acid.
6. Process for the preparation of compounds of general

This Sheet Has Been Replaced






formula (I) according to claim 1, characterized in that a compound of general formula (II)



(II)

wherein G is R₃, or a protecting group; A is absent and R₁, R₂, R₃ and B are as defined in claim 1 and X is a leaving group, is reacted with a compound of general formula (III)



(III)

wherein m, n and R₄ are as defined in claim 1, in an organic solvent at a temperature ranging from 0° to 150°C and when G is a protecting group, it is removed during the process or by the treatment with acids or alkyls to give the compound with R₃ is H.


7. Process according to claim 6, characterized in that the





protecting group is selected from ethoxycarbonyl, α -phenylvinyl or α -methylvinyl group.

8. Process according to claim 6, characterized in that the leaving group is selected from halogen, methansulfonate or 4-methylbenzensulfonate.
9. Process for the preparation of compounds of general formula (I) according to claim 1, characterized in that a compound of general formula (V)



wherein R_1 , R_2 , R_4 , A, B, m and n are as defined in claim 1 is reacted with a carbonyl derivative of formula (VI)




in which Y and Y' are leaving groups, identical or






different from each other in an aprotic solvent at a temperature ranging from 0° to 100°C.

10. Process according to claim 9, characterized in that the leaving group is selected from chlorine, trichloromethoxy, methoxy, ethoxy or imidazolyl.
11. Process for the preparation of compounds of general formula (I) according to claim 1, in which A is absent or it is a carbonyl group CO, characterized in that a compound of general formula (XIV)







wherein R_1 , R_2 and R_3 are as defined in claim 1 and M is a metal atom, is reacted with a compound of formula (XV)




wherein R_4 , A, B, m and n are as defined in claim 1 and Hal represents a halogen atom in a polar solvent at a temperature ranging 0° to 100°C.

12. Process according to claim 11, in which the metal atom is selected from sodium, potassium or lithium.
13. Process for the preparation of compounds of general formula (I) according to claim 1, in which A is a -CONH- group, characterized in that a compound of formula (XVI)





in which R_1 , R_2 and R_3 are as defined in claim 1 and L is a leaving group, is reacted with a compound of formula (IX)



in which B, m, n and R_4 are as defined in claim 1, in an aprotic solvent in the presence of an organic or acid inorganic/acceptor at a temperature ranging from -10° to the boiling point of the choosed solvent.

14. Process according to claim 13, characterized in that the leaving group is selected from halogen or alkoxy.
15. Process for the preparation of compound of general formula (I) according to claim 1, characterized in that a compound of formula I, in which R_3 is H obtained by a process according to any claims 6-14, is transformed by alkylation in another compound of formula I, in which R_3 is C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, with a suitable alkyl halide in the presence of a strong base and of an aprotic solvent.





16. Pharmaceutical compositions comprising as active ingredient an effective amount of a compound of general formula (I), as defined in claim 1, or physiologically acceptable acid addition salts thereof, in association with pharmaceutically acceptable carriers, diluents or excipients.
17. Pharmaceutical compositions according to claim 16 for the use in the treatment of patients suffering from central nervous system disorders, in particular in affective disorders (e.g. depression, bipolar disorders), anxiety, sleep and sexual disorders, psychosis, schizophrenia, personality disorders, mental organic disorders, mental disorders in childhood, aggressiveness and age associated memory impairment.
18. Pharmaceutical compositions according to claim 16 for the use in the treatment of patients suffering from cardiovascular disorders, e.g. hypertension and thrombosis.

