


CLAIMS

What is claimed is:

1. A compound of formula (I)



(I)

or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein:

R¹ is hydrogen, C₁-C₈ alkyl, or C₃-C₇ cycloalkyl;

R² and R³ are each independently hydrogen, C₁-C₈ alkyl, C₃-C₇ cycloalkyl, or C₃-C₇ cycloalkylmethyl;

R⁴ is hydrogen, halogen, C₁-C₈ alkyl, or C₃-C₇ cycloalkyl; and

R⁵ is hydrogen or halogen.

2. The compound of claim 0, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R¹ is C₁-C₆ alkyl.

3. The compound of claim 0, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R¹ is methyl, ethyl, propyl, or isopropyl.


4. The compound of claim 2, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R² is C₁-C₆ alkyl, C₃-C₆ cycloalkyl, or C₃-C₆ cycloalkylmethyl.

5. The compound of claim 3, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R² is methyl, ethyl, propyl, isopropyl, cyclopropyl, cyclopentyl, cyclopropylmethyl, or cyclopentylmethyl.

6. The compound of claim 4, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R³ is C₁-C₆ alkyl or C₃-C₆ cycloalkyl.

7. The compound of claim 5, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R³ is methyl, ethyl, propyl, isopropyl, or cyclopropyl.

8. The compound of claim 0, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R⁴ is located at the 7-position of the indazole ring, characterized by formula Ia:




(Ia)

9. The compound of claim 8, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R⁴ is hydrogen or halogen.


10. The compound of claim 0, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R⁵ is hydrogen or fluoro.

11. The compound of claim 8, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, wherein R¹ is methyl or ethyl; R² is isopropyl, cyclopropyl, cyclopropylmethyl, or cyclopentyl; R³ is methyl or ethyl; R⁴ is hydrogen or fluoro; and R⁵ is hydrogen or fluoro.


12. The compound of claim 6, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, selected from the group consisting of:



(Ic)









(Id)









(Ie)

13. The compound of claim 0, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, selected from the list in the table below:



Compound No.	Structure	Name
1		<i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoro-4-(3-isopropyl-2-methyl-2H-indazol-5-yl)pyrimidin-2-amine
2		<i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoro-4-(7-fluoro-3-isopropyl-2-methyl-2H-indazol-5-yl)pyrimidin-2-amine
3		<i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-4-(7-fluoro-3-isopropyl-2-methyl-2H-indazol-5-yl)pyrimidin-2-amine
4		4-(3-cyclopentyl-2-methyl-2H-indazol-5-yl)- <i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
5		4-(3-cyclopentyl-7-fluoro-2-methyl-2H-indazol-5-yl)- <i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine

6		4-(3-cyclopentyl-7-fluoro-2-methyl-2H-indazol-5-yl)-N-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)pyrimidin-2-amine
7		4-(3-cyclopropyl-2-methyl-2H-indazol-5-yl)-N-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
8		4-(3-cyclopropyl-7-fluoro-2-methyl-2H-indazol-5-yl)-N-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
9		4-(3-cyclohexyl-2-methyl-2H-indazol-5-yl)-N-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
10		4-(3-cyclohexyl-7-fluoro-2-methyl-2H-indazol-5-yl)-N-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
11		5-fluoro-4-(3-isopropyl-2-methyl-2H-indazol-5-yl)-N-(5-((4-isopropylpiperazin-1-yl)methyl)pyridin-2-yl)pyrimidin-2-amine

12		5-fluoro-4-(7-fluoro-3-isopropyl-2-methyl-2H-indazol-5-yl)-N-(5-((4-isopropylpiperazin-1-yl)methyl)pyridin-2-yl)pyrimidin-2-amine
13		4-(3-cyclopentyl-2-methyl-2H-indazol-5-yl)-5-fluoro-N-(5-((4-isopropylpiperazin-1-yl)methyl)pyridin-2-yl)pyrimidin-2-amine
14		4-(3-cyclopentyl-7-fluoro-2-methyl-2H-indazol-5-yl)-5-fluoro-N-(5-((4-isopropylpiperazin-1-yl)methyl)pyridin-2-yl)pyrimidin-2-amine
15		5-fluoro-4-(3-isopropyl-2-methyl-2H-indazol-5-yl)-N-(5-((4-propylpiperazin-1-yl)methyl)pyridin-2-yl)pyrimidin-2-amine
16		5-fluoro-4-(7-fluoro-3-isopropyl-2-methyl-2H-indazol-5-yl)-N-(5-((4-propylpiperazin-1-yl)methyl)pyridin-2-yl)pyrimidin-2-amine
17		4-(3-cyclopentyl-2-methyl-2H-indazol-5-yl)-5-fluoro-N-(5-((4-propylpiperazin-1-yl)methyl)pyridin-2-yl)pyrimidin-2-amine

18		4-(3-cyclopentyl-7-fluoro-2-methyl-2H-indazol-5-yl)-5-fluoro- <i>N</i> -((4-propylpiperazin-1-yl)methyl)pyridin-2-amine
19		4-(3-ethyl-2-methyl-2H-indazol-5-yl)- <i>N</i> -((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
20		4-(3-ethyl-7-fluoro-2-methyl-2H-indazol-5-yl)- <i>N</i> -((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
21		4-(3-(sec-butyl)-2-methyl-2H-indazol-5-yl)- <i>N</i> -((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
22		4-(3-(sec-butyl)-7-fluoro-2-methyl-2H-indazol-5-yl)- <i>N</i> -((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
23		4-(2-ethyl-3-isopropyl-2H-indazol-5-yl)- <i>N</i> -((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine

24		4-(2-ethyl-7-fluoro-3-isopropyl-2H-indazol-5-yl)- <i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
25		4-(3-cyclopropyl-2-ethyl-2H-indazol-5-yl)- <i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
26		4-(3-cyclopropyl-2-ethyl-7-fluoro-2H-indazol-5-yl)- <i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
27		4-(3-(cyclopropylmethyl)-2-methyl-2H-indazol-5-yl)- <i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
28		4-(3-(cyclopropylmethyl)-7-fluoro-2-methyl-2H-indazol-5-yl)- <i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
29		4-(3-cyclopropyl-2-ethyl-7-fluoro-2H-indazol-5-yl)- <i>N</i> -(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine

30		4-(3-(sec-butyl)-2-methyl-2H-indazol-5-yl)-N-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine
31		4-(3-(sec-butyl)-7-fluoro-2-methyl-2H-indazol-5-yl)-N-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoropyrimidin-2-amine


14. A pharmaceutical composition comprising a compound of claim 0, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, and one or more pharmaceutically acceptable adjuvants, diluents, and/or carriers.

15. An in vitro method of inhibiting cell proliferation comprising treating proliferative cells with an effective amount of a compound according to claim 0, or a salt, solvate, or prodrug thereof.

16. An in vitro method of inhibiting a cyclin-dependent kinase (CDK) comprising treating said kinase with an effective amount of a compound of claim 0, or a salt, solvate, prodrug, or composition thereof.


17. The in vitro method of claim 16, wherein said cyclin-dependent kinase is CDK4, CDK6, or a combination thereof.

18. A method of preparing a compound of formula (I), comprising a step of coupling intermediate **E** with intermediate **G**:




wherein R^1 through R^5 are defined as in claim 0, and X^3 is Cl, Br, or I.

19. The method of claim 18, further comprising the steps of converting intermediate **C** to intermediate **D** and coupling the intermediate **D** with a pyrimidine compound **H** to form the intermediate **E**:




wherein R^x and R^y are independent alkyl, aryl, cycloalkyl, or alternatively together form an alkylene group, each optionally substituted by one or more substituents independently selected from C_1 - C_4 alkyl, halogen and phenyl; and wherein X^1 , X^2 , and X^3 are each independently Cl, Br, or I, on condition that the intermediate **D** couples with the compound **H** selectively at the X^2 site over the X^3 site.

20. The method of claim 18, further comprising the steps of converting starting material **S1** to intermediate **A** and converting the intermediate **A** to the intermediate **C**:




wherein X is Cl, Br, I, or MeSO_3^- ; and R^2 and R^3 are defined as in claim 18.

21. The method of claim 20, wherein said converting the intermediate **A** to the intermediate **C**, alternatively, comprises converting the intermediate **A** to an alcohol intermediate **B** followed by reduction of the alcohol intermediate **B** to form the intermediate **C**.



wherein R^{2a} and R^{2b} are each independently hydrogen, alkyl, cycloalkyl, or together form an alkylene group so that the group $R^{2a}R^{2b}$ formed in the intermediate C is R^2 as defined in claim 20.

22. The method of claim 18, further comprising a step of forming the intermediate **G** through coupling the pyridine aldehyde compound **S2** and the piperazine compound **S3** to form an intermediate F, followed by converting the intermediate F to the intermediate G:



wherein X^4 is selected from the group consisting of Cl, Br, I, and $-\text{NO}_2$; and

wherein said converting the intermediate F to the intermediate G comprises replacing $-\text{X}^4$ with $-\text{NH}_2$ when X^4 is Cl, Br, or I; or reducing the nitro group ($-\text{NO}_2$) to amino group ($-\text{NH}_2$) when X^4 is NO_2 .