

## NOVEL ANTI-HUMAN GPVI ANTIBODIES AND USES THEREOF

### FIELD OF INVENTION

- 5 The present invention relates to novel anti-human glycoprotein VI antibodies, and to the uses thereof for treating cardiovascular diseases.

### BACKGROUND OF INVENTION

- 10 Acute coronary and cerebrovascular accidents are currently a major cause of death in the world. In addition, the global incidence of recurrence and death in the 6-month post-treatment period after an acute coronary syndrome is still 8-15%.

- In the case of acute coronary syndrome with ST segment elevation, mechanical treatment with coronary angioplasty and introduction of a stent is highly efficient to urgently restore coronary artery flow, but does not prevent morbidity/mortality for about 15% of patients  
15 in the next 6 months. Thrombolytic treatments, which are based on long term fibrinolytic, anticoagulant and anti-aggregating drugs associations, give even less encouraging results. Indeed, despite improvements in medical treatment of thrombosis, morbidity/mortality at 6 months is similar to that observed for acute coronary syndrome without ST segment elevation.

- 20 Regarding cerebrovascular ischemic accidents, treatments are still very limited due to the generally late caring of most patients and to the hemorrhagic risk of currently available anti-thrombotic treatments.

- There is thus still a clinical need for improving treatments for cardiovascular diseases, and especially for new molecules with improved features compared to available  
25 molecules. The challenge to face is to obtain molecules with excellent efficiency on the pathological thrombosis but devoid of risk of bleeding.

To answer to this requirement, the target must have a greater role in thrombus formation occurring in a diseased vessel than in physiological hemostasis required to limit the bleeding in a healthy vessel. This is the case of platelet Glycoprotein VI that has been demonstrated in animals to play a role in experimental thrombosis including stroke,  
5   vascular remodeling and to be critical in atherothrombosis.

Contrary to  $\alpha\text{IIb}\beta 3$  integrin antagonists, which are currently used in thrombosis treatment and inhibit platelet final activation phase, i.e. platelet aggregation, and to antagonists of platelet recruitment (inhibitors of the P2Y<sub>12</sub> ADP receptor and aspirin), GPVI is implicated into several steps of the platelet plug formation: initiation via the interaction  
10   with the injured vascular wall, amplification via initial platelet activation leading to the secretion of secondary agonists, the activation of integrins and of platelet procoagulant activity, growth and stabilization via the interaction with fibrin (Mammadova *et al.*, Blood 2015). Thus, GPVI antagonists may prevent not only platelet aggregation, but also secondary agonists liberation as well a growth factors and cytokines secretion resulting  
15   into vascular lesions development. Finally, GPVI expression is limited to platelets and megakaryocytes, and thus represents a perfectly specific target for anti-thrombosis treatment.

GPVI antagonists were thus developed for treating cardiovascular diseases.

WO 2001/000810 and WO 2003/008454 both describe a soluble GPVI recombinant  
20   protein which is a fusion protein between the GPVI extracellular domain and a human Ig Fc domain. This soluble recombinant GPVI protein competes with platelet GPVI for binding collagen. Encouraging results were first obtained with this soluble GPVI protein in a thrombosis murine model, but these results were not confirmed. In addition, this approach involves structural, functional and pharmacological disadvantages. First, this  
25   compound is a high molecular weight chimeric protein (~ 160 kDa). GPVI-Fc targets the collagen exposed at the site of the vascular injury, the amount and accessibility of which are poorly predictable and, thus, the amount of product to be injected constitutes a potential limitation to the use of GPVI-Fc. Another limitation could be the risk of immunization against neoepitopes potentially exposed on the fusion protein.

Neutralizing monoclonal antibodies directed against human GPVI were also described in the art.

For example, EP 1224942 and EP 1228768 disclose the rat monoclonal anti-GPVI antibody JAQ1, which specifically binds to mouse GPVI, for the treatment of thrombotic disease. JAQ1 antibody induces irreversible internalization of the GPVI receptor on mouse platelets.

EP 1538165 describes another rat monoclonal anti-GPVI antibody (hGP 5C4), which Fab fragment was found to have marked inhibitory effects on the main physiological functions of platelets induced by collagen stimulation: stimulation of collagen-mediated physiological activation markers PAC-I and CD62P-Selectin was completely prevented by hGP 5C4 Fab, and hGP 5C4 Fab potently inhibited human platelet aggregation *ex vivo* without any intrinsic activity. However, 5C4 is a rat antibody, and therefore only presents a very limited therapeutic potential.

WO 2005/111083 describes 4 mouse monoclonal anti-GPVI antibodies OM1, OM2, OM3 and OM4, that were found to inhibit GPVI binding to collagen, collagen-induced secretion and thromboxane A2 (TXA2) formation *in vitro*, as well as *ex vivo* collagen-induced platelet aggregation after intravenous injection to Cynomolgus monkeys. OM4 also appears to inhibit thrombus formation in a rat thrombosis model.

WO 2001/000810 also describes various murine monoclonal anti-GPVI antibodies named 8M14.3, 3F8.1, 9E18.3, 3J24.2, 6E12.3, 1P10.2, 4L7.3, 7H4.6, 9O12.2, 7H14.1, and 9E18.2, as well as several human phage antibodies scFv fragments named A9, A10, C9, A4, C10, B4, C3 and D11. Some of these antibodies and scFv fragments were found to inhibit GPVI binding to collagen, including antibodies 8M14.3, 3F8.1, 9E18.3, 3J24.2, 6E12.3, 1P10.2, 4L7.3, 7H4.6, and 9O12.2, and scFv fragments A10, A4, C10, B4, C3 and D11. In addition, 9O12.2 Fab fragments were found to completely block collagen-induced platelet aggregation and secretion, to block fibrin-induced platelet aggregation, to inhibit the procoagulant activity of collagen-stimulated or fibrin-stimulated platelets and platelet adhesion to collagen or fibrin in static conditions, to impair platelet adhesion and to prevent thrombi formation under arterial flow conditions.

WO 2008/049928 describes a scFv fragment derived from 9O12.2, constituted of the VH and VL domains of 9O12.2 monoclonal antibody linked via a (Gly<sub>4</sub>Ser)<sub>3</sub> peptide.

However, none of the currently known anti-GPVI antibodies was shown to be efficient *in vivo* for preventing and/or treating cardiovascular diseases. In particular, the majority of  
5 anti-GPVI antibodies that have been reported appeared not fitted for the development of an antithrombotic for medical use in human, especially due to their animal origin.

In particular, some human phage antibodies scFvs directed to human GPVI have been reported to be inhibitory but their affinity appears to be low. Moreover, cross-linking of GPVI at the platelet surface by a divalent or multivalent ligand such as 9O12.2 whole IgG  
10 results in platelet activation via GPVI dimerization and via cross-linking of GPVI to the low affinity Fc receptor (FcγRIIA). In contrast, monovalent 9O12.2 Fab and scFv fragments are inhibitory.

However, these fragments could not be used in therapeutic due to their size and to their animal origin which makes them immunogenic in human patients. Moreover, scFv  
15 fragments present a short half-life, which limits their therapeutic potentials.

There is thus still a need for neutralizing GPVI antagonists without immunogenicity in human.

US 2006/0088531 describes a human scFv fragment 10B12, presenting a K<sub>D</sub> for binding to human GPVI of about 7.9.10<sup>-7</sup> M. Smethurst *et al.*, 2004, further describes the epitope  
20 bound by 10B12 on the Ig-like C2-type domain 1 (D1) of human GPVI. This epitope comprises residues R58, K61, R66, K79 and R80 of human GPVI (numbering based on UniProtKB accession number Q9HCN6).

O'Connor *et al.*, 2006, also describes a human scFv fragment 1C3, of low affinity for GPVI (5.4 10<sup>-7</sup> M), which neither blocked collagen-induced platelet aggregation nor  
25 GPVI binding to collagen but which potentiated the inhibitory effect of the 10B12 antibody. The 1C3 epitope in GPVI comprises amino acid I168, and it is further postulated that this epitope might encompass a region between residues S164 and S182, a region which is highly conserved from mouse to human.

There is thus a need for a humanized antibody (i.e. an antibody without immunogenicity in human) with improved affinity, efficacy and half-life as compared to the antagonists of the prior art, as a means for efficiently and contentedly preventing and/or treating cardiovascular diseases in human. Moreover, these antagonists should preferably be easily purified.

In the art, anti-GPVI antibodies inducing a GPVI depletion phenotype were described (WO 2006/118350, WO 2011/073954, EP 2363416). In particular, WO 2006/118350 discloses an anti-GPVI antibody directed against all mammals. The epitope of GPVI bound by this antibody is described and corresponds to the loops 9 and 11 of the Ig-like C2-type domain 2 of GPVI, corresponding to amino acid residues 136-142 and 158-162, respectively.

However, GPVI depletion is undesirable since it cannot be controlled, and is irreversible (i.e. it lasts the lifetime of platelets, or even longer due to GPVI depletion on megakaryocytes).

In therapy, a rapid and safe antiplatelet effect is required. Therefore, antibodies not inducing a GPVI depletion phenotype, and preferably not inducing a decrease in platelet count *in vivo* (i.e. with a reversible effect) should be developed.

The Applicant developed an antibody anti-GPVI binding to a novel, undescribed, conformational epitope. Said antibody anti-GPVI has a strong affinity for human GPVI, and blocks GPVI interaction with its ligands (including collagen and fibrin), without decreasing the platelet count nor depleting GPVI *in vivo*. The present invention thus relates to an improved humanized neutralizing antibody specific to human GPVI.

## SUMMARY

The present disclosure relates to an isolated humanized protein binding to human Glycoprotein VI (hGPVI), wherein said protein binds to a conformational epitope comprising:

- at least one amino acid residue from amino acid residues 114 to 142 of hGPVI (SEQ ID NO: 13) or from a sequence sharing at least 60% of identity over amino acid residues 114 to 142 of hGPVI (SEQ ID NO: 13); and
- at least one amino acid residue from amino acid residues 165 to 187 of hGPVI (SEQ ID NO: 13) or from a sequence sharing at least 60% of identity over amino acid residues 165 to 187 of hGPVI (SEQ ID NO: 13).

In one aspect of the present disclosure, there is provided an isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or a fragment thereof,

wherein the variable region of the heavy chain comprises the following CDRs:

**VH-CDR1:** GYTFTSYNMH (SEQ ID NO: 1);

**VH-CDR2:** GIYPGNGDTSYNQKFQG (SEQ ID NO: 2); and

**VH-CDR3:** GTVVGDWYFDV (SEQ ID NO: 3); and

wherein the variable region of the light chain comprises the following CDRs:

**VL-CDR1:** RSSQSLENSNGNTYLN (SEQ ID NO: 4);

**VL-CDR2:** RVSNRFS (SEQ ID NO: 5); and

**VL-CDR3:** LQLTHVPWT (SEQ ID NO: 6).

In one embodiment, the conformational epitope bound by the isolated humanized protein of the present disclosure comprises at least one amino acid residue from amino acid residues 121 to 135 of hGPVI (SEQ ID NO: 13) or from a sequence sharing at least 60% of identity over amino acid residues 121 to 135 of hGPVI (SEQ ID NO: 13); and at  
5 least one amino acid residue from amino acid residues 169 to 183 of hGPVI (SEQ ID NO: 13) or from a sequence sharing at least 60% of identity over amino acid residues 169 to 183 of hGPVI (SEQ ID NO: 13).

The present disclosure also relates to an isolated humanized protein binding to human Glycoprotein VI (hGPVI), wherein said protein has a  $K_D$  for binding to hGPVI less than  
10 15 nM, wherein said  $K_D$  is measured by surface plasmon resonance using 960 to 1071 RU of soluble human GPVI and using PBS pH 7.4 as running buffer. In one embodiment, the isolated humanized protein of the present disclosure does not induce a GPVI depletion phenotype *in vivo*.

In one embodiment, the protein of the present disclosure is an antibody molecule  
15 selected from the group consisting of a whole antibody, a humanized antibody, a single chain antibody, a Fv, a Fab; or an antibody fragment selected from the group consisting of a unibody, a domain antibody, and a nanobody; or a monomeric antibody mimetic selected from the group consisting of an affibody, an affilin, an affitin, an adnectin, an atrimer, an evasin, a DARPin, an anticalin, an avimer, a fynomer, and a versabody.

20 In one embodiment, the antibody of the present disclosure is monovalent.

In one embodiment, the variable region of the heavy chain comprises at least one of the following CDRs:

**VH-CDR1:** GYTFTSYNMH (SEQ ID NO: 1);

**VH-CDR2:** GIYPGNGDTSYNQKFQG (SEQ ID NO: 2); and

**VH-CDR3:** GTVVGDWYFDV (SEQ ID NO: 3),

or any CDR having an amino acid sequence that shares at least 60% of identity with SEQ ID NO: 1-3,

- 5 In one embodiment, the variable region of the light chain comprises at least one of the following CDRs:

**VL-CDR1:** RSSQSLENSNGNTYLN (SEQ ID NO: 4);

**VL-CDR2:** RVSNRFS (SEQ ID NO: 5); and

**VL-CDR3:** LQLTHVPWT (SEQ ID NO: 6),

- 10 or any CDR having an amino acid sequence that shares at least 60% of identity with SEQ ID NO: 4-6.

In one embodiment, the variable region of the heavy chain comprises at least one of the CDRs as defined hereinabove and the variable region of the light chain comprises at least one of the CDRs as defined hereinabove.

- 15 In one embodiment, the variable region of the heavy chain comprises the following CDRs: GYTFTSYNMH (SEQ ID NO: 1), GIYPGNGDTSYNQKFQG (SEQ ID NO: 2) and GTVVGDWYFDV (SEQ ID NO: 3) and the variable region of the light chain comprises the following CDRs: RSSQSLENSNGNTYLN (SEQ ID NO: 4), RVSNRFS (SEQ ID NO: 5) and LQLTHVPWT (SEQ ID NO: 6) or any CDR having an amino acid  
20 sequence that shares at least 60% of identity with said SEQ ID NO: 1-6.

In one embodiment, the amino acid sequence encoding the heavy chain variable region is SEQ ID NO: 7 and the amino acid sequence encoding the light variable region is SEQ ID NO: 8, or any sequence having an amino acid sequence that shares at least 60% of identity with said SEQ ID NO: 7 or 8.

- 25 In one embodiment, the amino acid sequence encoding the heavy chain variable region is SEQ ID NO: 7 and the amino acid sequence encoding the light variable region is SEQ ID NO: 9, or any sequence having an amino acid sequence that shares at least 60% of identity with said SEQ ID NO: 7 or 9.

The present disclosure further relates to a composition comprising the protein binding to human GPVI as defined hereinabove.

In another aspect of the present disclosure, there is provided a composition comprising the isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or  
5 fragment thereof disclosed herein.

The present disclosure further relates to a pharmaceutical composition comprising the protein binding to human GPVI as defined hereinabove and at least one pharmaceutically acceptable excipient.

In a further aspect of the present disclosure, there is provided a pharmaceutical  
10 composition comprising the isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof disclosed herein and at least one pharmaceutically acceptable excipient.

In one embodiment, the pharmaceutical composition of the present disclosure is for treating, or for use in treating, a GPVI-related condition.

15 In one embodiment, said GPVI-related condition is a cardiovascular disease selected from arterial and venous thrombosis, restenosis, acute coronary syndrome and ischemic cerebrovascular accidents.

In a further aspect of the present disclosure, there is provided use of the isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof as  
20 defined hereinabove for the manufacture of a medicament for treating a GPVI-related condition.

The present disclosure further relates to the use of the antibody against human GPVI of the present disclosure for detecting GPVI in a biological sample.

Another object of the present disclosure is an expression vector comprising at least one of SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NO: 12 or any sequence having a nucleic acid sequence that shares at least 60% of identity with said SEQ ID NO: 10-12.

5 In another aspect of the present disclosure, there is provided an expression vector comprising at least one nucleic acid sequence encoding a heavy chain or a light chain of an antibody disclosed herein, wherein said expression vector comprises at least one of SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NO: 12 or any sequence having a nucleic acid sequence that shares at least 95% of identity with said SEQ ID NO: 10-12.

## 10 DETAILED DESCRIPTION

“**Glycoprotein VI (GPVI)**” is a platelet membrane glycoprotein that is involved in platelet-collagen interactions. GPVI is a transmembrane collagen receptor expressed on the surface of platelets. In one embodiment, the amino acid sequence of human GPVI is SEQ ID NO: 13 (accession number: BAA89353.1) or any amino acid sequence  
15 presenting at least about 90% identity with SEQ ID NO: 13, preferably at least about 91, 92, 93, 94, 95, 96, 97, 98, 99% identity or more with SEQ ID NO: 13.

(SEQ ID NO: 13)

MSPSPTALFCLGLCLGRVPA (Signal peptide)

QSGPLPKPSLQALPSSLVPLEKPVTLRCQGPPGVDLYRLEKLSSSRYQDQAVLFI  
 PAMKRSLAGRYRCSYQNGSLWSLPSDQLELVATGVFAKPSLSAQPGPAVSSGG  
 DVTLCQTRYGFDQFALYKEGDPAPYKNPERWYRASFPITVTAHSGTYRCY  
 SFSSRDPLYWSAPSDPLELVVTGTSVTPSRLPTEPPSSVAEFSEATAELTVSFTNK  
 5 VFTTETSRSITTSPKESDSPAGPARQYYTKGN (Extracellular domain)  
 LVRICLGAVILIILAGFLAEDWHSRRKRLRHRGRAVQRPLPPLPQTRKSHGG  
 QDGGRQDVHSRGLCS (Transmembrane and cytoplasmic domains)

The extracellular domain of GPVI is composed of two Ig-like C2-type domains, namely D1 and D2, linked by a hinge interdomain. In one embodiment, D1 comprises amino acid  
 10 residues 21 to 109 of SEQ ID NO: 13. In one embodiment, the hinge interdomain between D1 and D2 comprises amino acid residues 110 to 113 of SEQ ID NO: 13. In one embodiment, D2 comprises amino acid residues 114 to 207 of SEQ ID NO: 13.

“**About**” preceding a figure means plus or less 10% of the value of said figure.

“**Antibody**” or “**Immunoglobulin**” - As used herein, the term "immunoglobulin"  
 15 includes a protein having a combination of two heavy and two light chains whether or not it possesses any relevant specific immunoreactivity. "Antibodies" refers to such assemblies which have significant known specific immunoreactive activity to an antigen of interest (e.g. human GPVI). The term "anti-GPVI antibodies" is used herein to refer to antibodies which exhibit immunological specificity for human GPVI protein. As  
 20 explained elsewhere herein, “specificity” for human GPVI does not exclude cross-reaction with species homologues of GPVI. Antibodies and immunoglobulins comprise light and heavy chains, with or without an interchain covalent linkage between them. Basic immunoglobulin structures in vertebrate systems are relatively well understood. The generic term “immunoglobulin” comprises five distinct classes of antibody that can  
 25 be distinguished biochemically. All five classes of antibodies are within the scope of the present invention; the following discussion will generally be directed to the IgG class of immunoglobulin molecules. With regard to IgG, immunoglobulins comprise two identical light polypeptide chains of molecular weight of about 23,000 Daltons, and two identical heavy chains of molecular weight of about 53,000-70,000 Daltons. The four  
 30 chains are joined by disulfide bonds in a “Y” configuration wherein the light chains

bracket the heavy chains starting at the mouth of the “Y” and continuing through the variable region. The light chains of an antibody are classified as either kappa or lambda ( $[\kappa]$ ,  $[\lambda]$ ). Each heavy chain class may be bonded with either a kappa or lambda light chain. In general, the light and heavy chains are covalently bonded to each other, and the “tail” regions of the two heavy chains are bonded to each other by covalent disulfide linkages or non-covalent linkages when the immunoglobulins are generated either by hybridomas, B cells or genetically engineered host cells. In the heavy chain, the amino acid sequences run from an N-terminus at the forked ends of the Y configuration to the C-terminus at the bottom of each chain. Those skilled in the art will appreciate that heavy chains are classified as gamma, mu, alpha, delta, or epsilon ( $\gamma$ ,  $\mu$ ,  $\alpha$ ,  $\delta$ ,  $\epsilon$ ) with some subclasses among them (e.g.,  $\gamma 1 - \gamma 4$ ). It is the nature of this chain that determines the “class” of the antibody as IgG, IgM, IgA, IgD, or IgE, respectively. The immunoglobulin subclasses (isotypes) e.g., IgG1, IgG2, IgG3, IgG4, IgA1, etc. are well characterized and are known to confer functional specialization. Modified versions of each of these classes and isotypes are readily discernable to the skilled artisan in view of the instant disclosure and, accordingly, are within the scope of the instant invention. As indicated above, the variable region of an antibody allows the antibody to selectively recognize and specifically bind epitopes on antigens. That is, the light chain variable domain (VL domain) and heavy chain variable domain (VH domain) of an antibody combine to form the variable region that defines a three-dimensional antigen binding site. This quaternary antibody structure forms the antigen binding site presents at the end of each arm of the “Y”. More specifically, the antigen binding site is defined by three complementarity determining regions (CDRs) on each of the VH and VL chains.

**“An isolated antibody”** - As used herein, an “isolated antibody” is one that has been separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses of the antibody, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous components. In preferred embodiments, the antibody is purified: (1) to greater than 80, 85, 90, 91, 92, 93, 94, 95% or more by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal

amino acid sequence by use of a spinning cup sequenator; or (3) to homogeneity as shown by SDS-PAGE under reducing or non-reducing conditions and using Coomassie blue or, preferably, silver staining. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

**"Affinity variants"** - As used herein, the term "affinity variant" refers to a variant antibody which exhibits one or more changes in amino acid sequence compared to a reference anti-GPVI antibody, wherein the affinity variant exhibits an altered affinity for the human GPVI protein in comparison to the reference antibody. Typically, affinity variants will exhibit an improved affinity for human GPVI, as compared to the reference anti-GPVI antibody. The improvement may be either a lower  $K_D$  for human GPVI, a higher  $K_A$  for human GPVI, a faster on-rate for human GPVI or a slower off-rate for human GPVI or an alteration in the pattern of cross-reactivity with non-human GPVI homologues. Affinity variants typically exhibit one or more changes in amino acid sequence in the CDRs, as compared to the reference anti-GPVI antibody. Such substitutions may result in replacement of the original amino acid present at a given position in the CDRs with a different amino acid residue, which may be a naturally occurring amino acid residue or a non-naturally occurring amino acid residue. The amino acid substitutions may be conservative or non-conservative.

**"Binding Site"** - As used herein, the term "binding site" comprises a region of a protein which is responsible for selectively binding to a target antigen of interest (e.g. human GPVI). Binding domains or binding regions comprise at least one binding site. Exemplary binding domains include an antibody variable domain. The protein of the invention may comprise a single antigen binding site or multiple (e.g., two, three or four) antigen binding sites. Preferably, however, the protein of the invention comprises a single antigen binding site.

**"Conservative amino acid substitution"** - As used herein, a "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art, including basic side chains (e.g., lysine, arginine, histidine),

acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a nonessential amino acid residue in an immunoglobulin polypeptide may be replaced with another amino acid residue from the same side chain family. In another embodiment, a string of amino acids can be replaced with a structurally similar string that differs in order and/or composition of side chain family members.

10 **“Chimeric”** - As used herein, a “chimeric” protein comprises a first amino acid sequence linked to a second amino acid sequence with which it is not naturally linked in nature. The amino acid sequences may normally exist in separate proteins that are brought together in the fusion protein or they may normally exist in the same protein but are placed in a new arrangement in the fusion protein. A chimeric protein may be created, for example, by chemical synthesis, or by creating and translating a polynucleotide in which the peptide regions are encoded in the desired relationship.

**“CDR”** - As used herein, the term “CDR” or “complementarity determining region” means the non-contiguous antigen combining sites found within the variable region of both heavy and light chain polypeptides. CDRs were identified according to the following rules as deduced from Kabat *et al.* (1991) and Chotia and Lesk (1987):

- CDR-L1:
  - Start - Approx residue 24
  - Residue before is always a Cys
  - Residue after is always a Trp. Typically TRP-TYR-GLN, but also, TRP-LEU-GLN, TRP-PHE-GLN, TRP-TYR-LEU
  - Length 10 to 17 residues
- CDR-L2:
  - Start - always 16 residues after the end of L1
  - Residues before generally ILE-TYR, but also, VAL-TYR, ILE-LYS, ILE-PHE

Length always 7 residues.

- CDR-L3:

Start - always 33 residues after end of L2

Residue before is always Cys

5 Residues after always PHE-GLY-XXX-GLY (SEQ ID NO: 21)

Length 7 to 11 residues

- CDR-H1:

Start - Approx residue 26 (always 4 after a CYS) [Chothia / AbM definition] Kabat definition starts 5 residues later

10 Residues before always CYS-XXX-XXX-XXX (SEQ ID NO: 22)

Residues after always a TRP. Typically TRP-VAL, but also, TRP-ILE, TRP-ALA

Length 10 to 12 residues (AbM definition) Chothia definition excludes the last 4 residues

- CDR-H2:

15 Start - always 15 residues after the end of Kabat / AbM definition) of CDR-H1  
Residues before typically LEU-GLU-TRP-ILE-GLY (SEQ ID NO: 23), but a number of variations

Residues after LYS/ARG-LEU/ILE/VAL/PHE/THR/ALA-THR/SER/ILE/ALA  
Length Kabat definition 16 to 19 residues (AbM definition ends 7 residues earlier)

20 - CDR-H3:

Start - always 33 residues after end of CDR-H2 (always 2 after a CYS)

Residues before always CYS-XXX-XXX (typically CYS-ALA-ARG)

Residues after always TRP-GLY-XXX-GLY (SEQ ID NO: 24)

Length 3 to 25 residues

25 **“CH2 domain”** - As used herein, the term “CH2 domain” includes the region of a heavy chain molecule that usually extends from about amino acid 231 to about amino acid 340.

The CH2 domain is unique in that it is not closely paired with another domain. Rather, two N-linked branched carbohydrate chains are interposed between the two CH2 domains of an intact native IgG molecule. It has been speculated that the carbohydrate may provide a substitute for the domain-domain pairing and help stabilize the CH2 domain (Burton, Molec. Immunol. 22 (1985) 161-206).

**“Derived from”** - As used herein, the term “derived from” a designated protein (e.g. an anti-GPVI antibody or antigen-binding fragment thereof) refers to the origin of the protein. In an embodiment, the protein or amino acid sequence which is derived from a particular starting protein is a CDR sequence or sequence related thereto. In an embodiment, the amino acid sequence which is derived from a particular starting protein is not contiguous. For example, in an embodiment, one, two, three, four, five, or six CDRs are derived from a starting antibody. In an embodiment, the protein or amino acid sequence which is derived from a particular starting protein or amino acid sequence has an amino acid sequence that is essentially identical to that of the starting sequence, or a region thereof wherein the region consists of at least 3-5 amino acids, at least 5-10 amino acids, at least 10-20 amino acids, at least 20-30 amino acids, or at least 30-50 amino acids, or which is otherwise identifiable to one of ordinary skill in the art as having its origin in the starting sequence. In an embodiment, the one or more CDR sequences derived from the starting antibody are altered to produce variant CDR sequences, e.g. affinity variants, wherein the variant CDR sequences maintain GPVI binding activity.

**“Diabodies”** - As used herein, the term “diabodies” refers to small antibody fragments prepared by constructing sFv fragments (see sFv paragraph) with short linkers (about 5-10 residues) between the VH and VL domains such that inter-chain but not intra-chain pairing of the V domains is achieved, resulting in a bivalent fragment, i.e., fragment having two antigen-binding sites. Bispecific diabodies are heterodimers of two “crossover” sFv fragments in which the VH and VL domains of the two antibodies are present on different polypeptide chains. Diabodies are described more fully in, for example, EP 0404097; WO 1993/011161; and Holliger *et al.*, Proc. Natl. Acad. Sci., 90:6444-6448 (1993).

**“Engineered”** - As used herein, the term “engineered” includes manipulation of nucleic acid or polypeptide molecules by synthetic means (e.g. by recombinant techniques, *in*

*vitro* peptide synthesis, by enzymatic or chemical coupling of peptides or some combination of these techniques). Preferably, the antibodies of the invention are engineered, including for example, humanized and/or chimeric antibodies, and antibodies which have been engineered to improve one or more properties, such as antigen binding, stability/half-life or effector function.

**“Epitope”** - As used herein, the term “epitope” refers to a specific arrangement of amino acids located on a protein or proteins to which an antibody binds. Epitopes often consist of a chemically active surface grouping of molecules such as amino acids or sugar side chains, and have specific three dimensional structural characteristics as well as specific charge characteristics. Epitopes can be linear or conformational, i.e., involving two or more sequences of amino acids in various regions of the antigen that may not necessarily be contiguous.

**“Framework region”** – As used herein, the term “framework region” or “FR region” includes the amino acid residues that are part of the variable region, but are not part of the CDRs (e.g., using the Kabat/Chothia definition of CDRs). Therefore, a variable region framework is between about 100-120 amino acids in length but includes only those amino acids outside of the CDRs. For the specific example of a heavy chain variable region and for the CDRs as defined by Kabat/Chothia, framework region 1 may correspond to the domain of the variable region encompassing amino acids 1-25; framework region 2 may correspond to the domain of the variable region encompassing amino acids 36-49; framework region 3 may correspond to the domain of the variable region encompassing amino acids 67-98, and framework region 4 may correspond to the domain of the variable region from amino acids 110 to the end of the variable region. The framework regions for the light chain are similarly separated by each of the light chain variable region CDRs. In naturally occurring antibodies, the six CDRs present on each monomeric antibody are short, non-contiguous sequences of amino acids that are specifically positioned to form the antigen binding site as the antibody assumes its three dimensional configuration in an aqueous environment. The remainders of the heavy and light variable domains show less inter-molecular variability in amino acid sequence and are termed the framework regions. The framework regions largely adopt a [beta]-sheet conformation and the CDRs form loops which connect, and in some cases form part of, the [beta]-sheet structure. Thus,

these framework regions act to form a scaffold that provides for positioning the six CDRs in correct orientation by inter-chain, non-covalent interactions. The antigen binding site formed by the positioned CDRs defines a surface complementary to the epitope on the immunoreactive antigen. This complementary surface promotes the non-covalent binding  
5 of the antibody to the immunoreactive antigen epitope. The position of CDRs can be readily identified by one of ordinary skill in the art.

**“Fragment”** - As used herein, the term “fragment” refers to a part or region of an antibody or antibody chain comprising fewer amino acid residues than an intact or complete antibody or antibody chain. The term "antigen-binding fragment" refers to a  
10 protein fragment of an immunoglobulin or antibody that binds antigen or competes with intact antibody (i.e., with the intact antibody from which they were derived) for antigen binding (i.e., specific binding to human GPVI). As used herein, the term "fragment" of an antibody molecule includes antigen-binding fragments of antibodies, for example, an antibody light chain variable domain (VL), an antibody heavy chain variable domain  
15 (VH), a single chain antibody (scFv), a F(ab')<sub>2</sub> fragment, a Fab fragment, an Fd fragment, an Fv fragment, a single domain antibody fragment (DAb), a one-armed (monovalent) antibody, diabodies or any antigen-binding molecule formed by combination, assembly or conjugation of such antigen binding fragments. Fragments can be obtained, e.g., via chemical or enzymatic treatment of an intact or complete antibody or antibody chain or  
20 by recombinant means.

**“Fv”** - As used herein, the term “Fv” is the minimum antibody fragment that contains a complete antigen-recognition and -binding site. This fragment consists of a dimer of one heavy- and one light-chain variable region domain in tight, non-covalent association. From the folding of these two domains emanate six hypervariable loops (three loops each  
25 from the H and L chain) that contribute to antigen binding and confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

**“Heavy chain region”** - As used herein, the term “heavy chain region” includes amino  
30 acid sequences derived from the constant domains of an immunoglobulin heavy chain. A protein comprising a heavy chain region comprises at least one of: a CH1 domain, a hinge

(e.g., upper, middle, and/or lower hinge region) domain, a CH2 domain, a CH3 domain, or a variant or fragment thereof. In an embodiment, a binding molecule of the invention may comprise the Fc region of an immunoglobulin heavy chain (e.g., a hinge portion, a CH2 domain, and a CH3 domain). In another embodiment, a binding molecule of the invention lacks at least a region of a constant domain (e.g., all or part of a CH2 domain). In certain embodiments, at least one, and preferably all, of the constant domains are derived from a human immunoglobulin heavy chain. For example, in one preferred embodiment, the heavy chain region comprises a fully human hinge domain. In other preferred embodiments, the heavy chain region comprising a fully human Fc region (e.g., hinge, CH2 and CH3 domain sequences from a human immunoglobulin). In certain embodiments, the constituent constant domains of the heavy chain region are from different immunoglobulin molecules. For example, a heavy chain region of a protein may comprise a CH2 domain derived from an IgG1 molecule and a hinge region derived from an IgG3 or IgG4 molecule. In other embodiments, the constant domains are chimeric domains comprising regions of different immunoglobulin molecules. For example, a hinge may comprise a first region from an IgG1 molecule and a second region from an IgG3 or IgG4 molecule. As set forth above, it will be understood by one of ordinary skill in the art that the constant domains of the heavy chain region may be modified such that they vary in amino acid sequence from the naturally occurring (wild-type) immunoglobulin molecule. That is, the proteins of the invention disclosed herein may comprise alterations or modifications to one or more of the heavy chain constant domains (CH1, hinge, CH2 or CH3) and/or to the light chain constant domain (CL). Exemplary modifications include additions, deletions or substitutions of one or more amino acids in one or more domains.

25 **“Hinge region”** - As used herein, the term “hinge region” includes the region of a heavy chain molecule that joins the CH1 domain to the CH2 domain. This hinge region comprises approximately 25 residues and is flexible, thus allowing the two N-terminal antigen binding regions to move independently. Hinge regions can be subdivided into three distinct domains: upper, middle, and lower hinge domains (Roux *et al.*, J. Immunol. 1998 161 :4083).

30

The terms “**hypervariable loop**” and “**complementarity determining region**” are not strictly synonymous, since the hypervariable loops (HVs) are defined on the basis of structure, whereas complementarity determining regions (CDRs) are defined based on sequence variability (Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD., 1983) and the limits of the HVs and the CDRs may be different in some VH and VL domains. The CDRs of the VL and VH domains can typically be defined by the Kabat/Chothia definition (see above). In one embodiment, the CDRs of the VL and VH domains may comprise the following amino acids: residues 24-39 (CDRL1), 55-61 (CDRL2) and 94-102 (CDRL3) in the light chain variable domain, and residues 26-35 (CDRH1), 50-66 (CDRH2) and 99-109 (CDRH3) in the heavy chain variable domain. Thus, the HVs may be comprised within the corresponding CDRs and references herein to the “hypervariable loops” of VH and VL domains should be interpreted as also encompassing the corresponding CDRs, and vice versa, unless otherwise indicated. The more highly conserved regions of variable domains are called the framework region (FR), as defined below. The variable domains of native heavy and light chains each comprise four FRs (FR1, FR2, FR3 and FR4, respectively), largely adopting a [beta]-sheet configuration, connected by the three hypervariable loops. The hypervariable loops in each chain are held together in close proximity by the FRs and, with the hypervariable loops from the other chain, contribute to the formation of the antigen-binding site of antibodies. Structural analysis of antibodies revealed the relationship between the sequence and the shape of the binding site formed by the complementarity determining regions (Chothia *et al.*, J. Mol. Biol. 227: 799-817 (1992)); Tramontano *et al.*, J. Mol. Biol. 215: 175-182 (1990)). Despite their high sequence variability, five of the six loops adopt just a small repertoire of main-chain conformations, called “canonical structures”. These conformations are first of all determined by the length of the loops and secondly by the presence of key residues at certain positions in the loops and in the framework regions that determine the conformation through their packing, hydrogen bonding or the ability to assume unusual main-chain conformations.

30 “**Humanized**” - As used herein, the term “humanized” refers to chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', or

other antigen-binding subsequences of antibodies) which contain minimal sequence derived from a murine immunoglobulin. For example, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary-determining region (CDR) of the recipient are replaced by residues from a CDR of the original antibody (donor antibody) while maintaining the desired specificity, affinity, and capacity of the original antibody.

**“Humanizing substitutions”** - As used herein, the term “humanizing substitutions” refers to amino acid substitutions in which the amino acid residue present at a particular position in the VH or VL domain of a non-human anti-GPVI antibody (for example a murine anti-GPVI antibody) is replaced with an amino acid residue which occurs at an equivalent position in a reference human VH or VL domain. The reference human VH or VL domain may be a VH or VL domain encoded by the human germline, in which case the substituted residues may be referred to as “germlining substitutions”. Humanizing/germlining substitutions may be made in the framework regions and/or the CDRs of an anti-GPVI antibody, defined herein.

**“High human homology”** - An antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL) will be considered as having high human homology if the VH domains and the VL domains, taken together, exhibit at least 70, 75, 80, 85, 90, 95% or more amino acid sequence identity to the closest matching human germline VH and VL sequences. Antibodies having high human homology may include antibodies comprising VH and VL domains of native non-human antibodies which exhibit sufficiently high % sequence identity human germline sequences, as well as engineered, especially humanized, variants of such antibodies and also “fully human” antibodies. In an embodiment the VH domain of the antibody with high human homology may exhibit an amino acid sequence identity or sequence homology of 75%, 80% or greater with one or more human VH domains across the framework regions FR1, FR2, FR3 and FR4. In other embodiments the amino acid sequence identity or sequence homology between the VH domain of the protein of the invention and the closest matching human germline VH domain sequence may be 85% or greater, 90% or greater, 95% or greater, 97% or greater, or up to 99% or even 100%. In an embodiment the VH domain of the antibody with high human homology may contain one or more (e.g. 1 to

20) amino acid sequence mismatches across the framework regions FR1, FR2, FR3 and FR4, in comparison to the closest matched human VH sequence. In another embodiment the VL domain of the antibody with high human homology may exhibit a sequence identity or sequence homology of 80% or greater with one or more human VL domains  
5 across the framework regions FR1, FR2, FR3 and FR4. In other embodiments the amino acid sequence identity or sequence homology between the VL domain of the protein of the invention and the closest matching human germline VL domain sequence may be 85% or greater 90% or greater, 95% or greater, 97% or greater, or up to 99% or even 100%.

In an embodiment the VL domain of the antibody with high human homology may  
10 contain one or more (e.g. 1 to 20, preferably 1 to 10 and more preferably 1 to 5) amino acid sequence mismatches across the framework regions FR1, FR2, FR3 and FR4, in comparison to the closest matched human VL sequence. Before analyzing the percentage sequence identity between the antibody with high human homology and human germline VH and VL, the canonical folds may be determined, which allow the identification of the  
15 family of human germline segments with the identical combination of canonical folds for H1 and H2 or L1 and L2 (and L3). Subsequently the human germline family member that has the highest degree of sequence homology with the variable region of the antibody of interest is chosen for scoring the sequence homology. The determination of Chothia canonical classes of hypervariable loops L1, L2, L3, H1 and H2 can be performed with  
20 the bioinformatics tools publicly available on webpage [www.bioinf.org.uk/abs/chothia.html](http://www.bioinf.org.uk/abs/chothia.html). The output of the program shows the key residue requirements in a data file. In these data files, the key residue positions are shown with the allowed amino acids at each position. The sequence of the variable region of the antibody of interest is given as input and is first aligned with a consensus antibody  
25 sequence to assign the Kabat/Chothia numbering scheme. The analysis of the canonical folds uses a set of key residue templates derived by an automated method developed by Martin and Thornton (Martin *et al.*, J. Mol. Biol. 263:800-815 (1996)). With the particular human germline V segment known, which uses the same combination of canonical folds for H1 and H2 or L1 and L2 (and L3), the best matching family member in terms of  
30 sequence homology can be determined. With bioinformatics tools the percentage sequence identity between the VH and VL domain framework amino acid sequences of

the antibody of interest and corresponding sequences encoded by the human germline can be determined, but actually manual alignment of the sequences can be applied as well. Human immunoglobulin sequences can be identified from several protein data bases, such as VBase (<http://vbase.mrc-cpe.cam.ac.uk/>) or the Pluckthun/Honegger database (<http://www.bioc.unizh.ch/antibody/Sequences/Germlines>). To compare the human sequences to the V regions of VH or VL domains in an antibody of interest a sequence alignment algorithm such as available via websites like [www.expasy.ch/tools/#align](http://www.expasy.ch/tools/#align) can be used, but also manual alignment with the limited set of sequences can be performed. Human germline light and heavy chain sequences of the families with the same combinations of canonical folds and with the highest degree of homology with the framework regions 1, 2, and 3 of each chain are selected and compared with the variable region of interest; also the FR4 is checked against the human germline JH and JK or JL regions. Note that in the calculation of overall percent sequence homology the residues of FR1, FR2 and FR3 are evaluated using the closest match sequence from the human germline family with the identical combination of canonical folds. Only residues different from the closest match or other members of the same family with the same combination of canonical folds are scored (NB - excluding any primer-encoded differences). However, for the purposes of humanization, residues in framework regions identical to members of other human germline families, which do not have the same combination of canonical folds, can be considered "human", despite the fact that these are scored "negative" according to the stringent conditions described above. This assumption is based on the "mix and match" approach for humanization, in which each of FR1, FR2, FR3 and FR4 is separately compared to its closest matching human germline sequence and the humanized molecule therefore contains a combination of different FRs as was done by Qu and colleagues (Qu *et al.*, Clin. Cancer Res. 5:3095-3100 (1999)) and Ono and colleagues (Ono *et al.*, Mol. Immunol. 36:387-395 (1999)). The boundaries of the individual framework regions may be assigned using the IMGT numbering scheme, which is an adaptation of the numbering scheme of Chothia (Lefranc *et al.*, Nucleic acid res 27: 209-212 (1999); <http://im.gt.cines.fr>). Antibodies with high human homology may comprise hypervariable loops or CDRs having human or human-like canonical folds, as discussed in detail below. In an embodiment at least one hypervariable loop or CDR in either the VH domain or the VL domain of the antibody with high human homology may

be obtained or derived from a VH or VL domain of a non-human antibody, yet exhibit a predicted or actual canonical fold structure which is substantially identical to a canonical fold structure which occurs in human antibodies. It is well established in the art that although the primary amino acid sequences of hypervariable loops present in both VH domains and VL domains encoded by the human germline are, by definition, highly variable, all hypervariable loops, except CDR H3 of the VH domain, adopt only a few distinct structural conformations, termed canonical folds (Chothia *et al.*, J. Mol. Biol. 196:901-917 (1987); Tramontano *et al.*, Proteins 6:382- 94 (1989)), which depend on both the length of the hypervariable loop and presence of the so-called canonical amino acid residues (Chothia *et al.*, J. Mol. Biol. 196:901-917 (1987)). Actual canonical structures of the hypervariable loops in intact VH or VL domains can be determined by structural analysis (e.g. X-ray crystallography), but it is also possible to predict canonical structure on the basis of key amino acid residues which are characteristic of a particular structure (discussed further below). In essence, the specific pattern of residues that determines each canonical structure forms a "signature" which enables the canonical structure to be recognized in hypervariable loops of a VH or VL domain of unknown structure; canonical structures can therefore be predicted on the basis of primary amino acid sequence alone. The predicted canonical fold structures for the hypervariable loops of any given VH or VL sequence in an antibody with high human homology can be analyzed using algorithms which are publicly available from [www.bioinf.org.uk/abs/chothia.html](http://www.bioinf.org.uk/abs/chothia.html), [www.biochem.ucl.ac.uk/~martin/antibodies.html](http://www.biochem.ucl.ac.uk/~martin/antibodies.html) and [www.bioc.unizh.ch/antibody/Sequences/Germlines/Vbase\\_hVk.html](http://www.bioc.unizh.ch/antibody/Sequences/Germlines/Vbase_hVk.html). These tools permit query VH or VL sequences to be aligned against human VH or VL domain sequences of known canonical structure, and a prediction of canonical structure made for the hypervariable loops of the query sequence. In the case of the VH domain, H1 and H2 loops may be scored as having a canonical fold structure "substantially identical" to a canonical fold structure known to occur in human antibodies if at least the first, and preferable both, of the following criteria are fulfilled:

1. An identical length, determined by the number of residues, to the closest matching human canonical structural class.

2. At least 33% identity, preferably at least 50% identity with the key amino acid residues described for the corresponding human H1 and H2 canonical structural classes (note for the purposes of the foregoing analysis the H1 and H2 loops are treated separately and each compared against its closest matching human canonical structural class). The foregoing analysis relies on prediction of the canonical structure of the H1 and H2 loops of the antibody of interest. If the actual structures of the H1 and H2 loops in the antibody of interest are known, for example based on X-ray crystallography, then the H1 and H2 loops in the antibody of interest may also be scored as having a canonical fold structure "substantially identical" to a canonical fold structure known to occur in human antibodies if the length of the loop differs from that of the closest matching human canonical structural class (typically by +1 or +2 amino acids) but the actual structure of the H1 and H2 loops in the antibody of interest matches the structure of a human canonical fold. Key amino acid residues found in the human canonical structural classes for the first and second hypervariable loops of human VH domains (H1 and H2) are described by Chothia *et al.*, J. Mol. Biol. 227:799-817 (1992), the contents of which are incorporated herein in their entirety by reference. In particular, Table 3 on page 802 of Chothia *et al.*, which is specifically incorporated herein by reference, lists preferred amino acid residues at key sites for H1 canonical structures found in the human germline, whereas Table 4 on page 803, also specifically incorporated by reference, lists preferred amino acid residues at key sites for CDR H2 canonical structures found in the human germline. In an embodiment, both H1 and H2 in the VH domain of the antibody with high human homology exhibit a predicted or actual canonical fold structure which is substantially identical to a canonical fold structure which occurs in human antibodies. Antibodies with high human homology may comprise a VH domain in which the hypervariable loops H1 and H2 form a combination of canonical fold structures which is identical to a combination of canonical structures known to occur in at least one human germline VH domain. It has been observed that only certain combinations of canonical fold structures at H1 and H2 actually occur in VH domains encoded by the human germline. In an embodiment H1 and H2 in the VH domain of the antibody with high human homology may be obtained from a VH domain of a non-human species, yet form a combination of predicted or actual canonical fold structures which is identical to a combination of canonical fold structures known to occur in a human germline or somatically mutated VH domain. In non-limiting

embodiments H1 and H2 in the VH domain of the antibody with high human homology may be obtained from a VH domain of a non-human species, and form one of the following canonical fold combinations: 1-1, 1-2, 1-3, 1-6, 1-4, 2- 1, 3-1 and 3-5. An antibody with high human homology may contain a VH domain which exhibits both high  
5 sequence identity/sequence homology with human VH, and which contains hypervariable loops exhibiting structural homology with human VH. It may be advantageous for the canonical folds present at H1 and H2 in the VH domain of the antibody with high human homology, and the combination thereof, to be "correct" for the human VH germline sequence which represents the closest match with the VH domain of the antibody with  
10 high human homology in terms of overall primary amino acid sequence identity. By way of example, if the closest sequence match is with a human germline VH3 domain, then it may be advantageous for H1 and H2 to form a combination of canonical folds which also occurs naturally in a human VH3 domain. This may be particularly important in the case of antibodies with high human homology which are derived from non-human species, e.g.  
15 antibodies containing VH and VL domains which are derived from camelid conventional antibodies, especially antibodies containing humanized camelid VH and VL domains. Thus, in an embodiment the VH domain of the anti-GPVI antibody with high human homology may exhibit a sequence identity or sequence homology of 70% or greater, 80% or greater, 85% or greater, 90% or greater, 95% or greater, 97% or greater, or up to 99%  
20 or even 100% with a human VH domain across the framework regions FR1, FR2 , FR3 and FR4, and in addition H1 and H2 in the same antibody are obtained from a non-human VH domain, but form a combination of predicted or actual canonical fold structures which is the same as a canonical fold combination known to occur naturally in the same human VH domain. In other embodiments, L1 and L2 in the VL domain of the antibody with  
25 high human homology are each obtained from a VL domain of a non-human species, and each exhibits a predicted or actual canonical fold structure which is substantially identical to a canonical fold structure which occurs in human antibodies. As with the VH domains, the hypervariable loops of VL domains of both VLambda and VKappa types can adopt a limited number of conformations or canonical structures, determined in part by length  
30 and also by the presence of key amino acid residues at certain canonical positions. Within an antibody of interest having high human homology, L1, L2 and L3 loops obtained from a VL domain of a non-human species, e.g. a Camelidae species, may be scored as having

a canonical fold structure "substantially identical" to a canonical fold structure known to occur in human antibodies if at least the first, and preferable both, of the following criteria are fulfilled:

1. An identical length, determined by the number of residues, to the closest matching human structural class.
2. At least 33% identity, preferably at least 50% identity with the key amino acid residues described for the corresponding human L1 or L2 canonical structural classes, from either the VLambda or the VKappa repertoire (note for the purposes of the foregoing analysis the L1 and L2 loops are treated separately and each compared against its closest matching human canonical structural class). The foregoing analysis relies on prediction of the canonical structure of the L1, L2 and L3 loops in the VL domain of the antibody of interest. If the actual structure of the L1, L2 and L3 loops is known, for example based on X-ray crystallography, then L1, L2 or L3 loops derived from the antibody of interest may also be scored as having a canonical fold structure "substantially identical" to a canonical fold structure known to occur in human antibodies if the length of the loop differs from that of the closest matching human canonical structural class (typically by +1 or +2 amino acids) but the actual structure of the loops in the antibody of interest matches a human canonical fold. Key amino acid residues found in the human canonical structural classes for the CDRs of human VLambda and VKappa domains are described by Morea *et al.*, Methods, 20: 267-279 (2000) and Martin *et al.*, J. Mol. Biol., 263:800-815 (1996). The structural repertoire of the human VKappa domain is also described by Tomlinson *et al.*, EMBO J. 14:4628-4638 (1995), and that of the VLambda domain by Williams *et al.*, J. Mol. Biol., 264:220-232 (1996). The contents of all these documents are to be incorporated herein by reference. L1 and L2 in the VL domain of an antibody with high human homology may form a combination of predicted or actual canonical fold structures which is identical to a combination of canonical fold structures known to occur in a human germline VL domain. In non-limiting embodiments L1 and L2 in the VLambda domain of an antibody with high human homology may form one of the following canonical fold combinations: 11-7, 13-7(A,B,C), 14-7(A,B), 12-11, 14-11 and 12-12 (as defined in Williams *et al.*, J. Mol. Biol. 264:220 -32 (1996) and as shown on [http://www.bioc.uzh.ch/antibody/Sequences/Germlines/VBase\\_hVL.html](http://www.bioc.uzh.ch/antibody/Sequences/Germlines/VBase_hVL.html)). In non-

limiting embodiments L1 and L2 in the VKappa domain may form one of the following canonical fold combinations: 2- 1, 3-1, 4- 1 and 6- 1 (as defined in Tomlinson *et al.*, EMBO J. 14:4628-38 (1995) and as shown on [http://www.bioc.uzh.ch/antibody/Sequences/Germlines/VBase\\_hVK.html](http://www.bioc.uzh.ch/antibody/Sequences/Germlines/VBase_hVK.html)).

- 5 In a further embodiment, all three of L1, L2 and L3 in the VL domain of an antibody with high human homology may exhibit a substantially human structure. It is preferred that the VL domain of the antibody with high human homology exhibit both high sequence identity/sequence homology with human VL, and also that the hypervariable loops in the VL domain exhibit structural homology with human VL.
- 10 In an embodiment, the VL domain of the anti-GPVI antibody with high human homology may exhibit a sequence identity of 70% or greater, 80% or greater, 85% or greater, 90% or greater, 95% or greater, 97% or greater, or up to 99% or even 100% with a human VL domain across the framework regions FR1, FR2 , FR3 and FR4, and in addition hypervariable loop L1 and hypervariable loop L2 may form a combination of predicted
- 15 or actual canonical fold structures which is the same as a canonical fold combination known to occur naturally in the same human VL domain. It is, of course, envisaged that VH domains exhibiting high sequence identity/sequence homology with human VH, and also structural homology with hypervariable loops of human VH will be combined with VL domains exhibiting high sequence identity/sequence homology with human VL, and
- 20 also structural homology with hypervariable loops of human VL to provide antibodies with high human homology containing VH/VL pairings with maximal sequence and structural homology to human-encoded VH/VL pairings.

**“Immunospecific”, “specific for” or to “specifically bind”** - As used herein, an antibody is said to be “immunospecific”, “specific for” or to “specifically bind” an

- 25 antigen if it reacts at a detectable level with the antigen, preferably with an affinity constant,  $K_A$ , of greater than or equal to about  $10^6 \text{ M}^{-1}$ , greater than or equal to about  $10^7 \text{ M}^{-1}$ , or greater than or equal to  $10^8 \text{ M}^{-1}$ , or greater than or equal to  $1.5 \cdot 10^8 \text{ M}^{-1}$ , or greater than or equal to  $10^9 \text{ M}^{-1}$  or greater than or equal to  $5 \cdot 10^9 \text{ M}^{-1}$ . Affinity of an antibody for its cognate antigen is also commonly expressed as a dissociation constant  $K_D$ , and in
- 30 certain embodiments, an antibody specifically binds to antigen if it binds with a  $K_D$  of less than or equal to  $10^{-6} \text{ M}$ , less than or equal to  $10^{-7} \text{ M}$ , or less than or equal to  $1.5 \cdot 10^{-8}$

M, or less than or equal to  $10^{-8}$  M, or less than or equal to  $5 \cdot 10^{-9}$  M or less than or equal to  $10^{-9}$  M. Affinities of antibodies can be readily determined using conventional techniques, for example, those described by Scatchard G *et al.* (The attractions of proteins for small molecules and ions. *Ann NY Acad Sci* 1949;51: 660-672). Binding properties of an antibody to antigens, cells or tissues thereof may generally be determined and assessed using immunodetection methods including, for example, ELISA, immunofluorescence-based assays, such as immuno-histochemistry (IHC) and/or fluorescence-activated cell sorting (FACS) or by surface plasmon resonance (SPR, BIAcore).

**“Isolated nucleic acid”** - As used herein, an “isolated nucleic acid” is a nucleic acid that is substantially separated from other genome DNA sequences as well as proteins or complexes such as ribosomes and polymerases, which naturally accompany a native sequence. The term embraces a nucleic acid sequence that has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogues or analogues biologically synthesized by heterologous systems. A substantially pure nucleic acid includes isolated forms of the nucleic acid. Of course, this refers to the nucleic acid as originally isolated and does not exclude genes or sequences later added to the isolated nucleic acid by the hand of man.

The term **“polypeptide”** is used in its conventional meaning, i.e., as a sequence of less than 100 amino acids. A polypeptide usually refers to a monomeric entity. The term **“protein”** refers to a sequence of more than 100 amino acids and/or to a multimeric entity. The proteins of the invention are not limited to a specific length of the product. This term does not refer to or exclude post-expression modifications of the protein, for example, glycosylation, acetylation, phosphorylation and the like, as well as other modifications known in the art, both naturally occurring and non-naturally occurring. A protein may be an entire protein, or a subsequence thereof. Particular proteins of interest in the context of this invention are amino acid subsequences comprising CDRs and being capable of binding an antigen. An “isolated protein” is one that has been identified and separated and/or recovered from a component of its natural environment. In preferred embodiments, the isolated protein will be purified (1) to greater than 80, 85, 90, 95% by weight of protein as determined by the Lowry method, and most preferably more than 96, 97, 98, or 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal

amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or non-reducing conditions using Coomassie blue or, preferably, silver staining. Isolated protein includes the protein in situ within recombinant cells since at least one component of the protein's natural environment will not be present.

5 Ordinarily, however, isolated protein will be prepared by at least one purification step.

**“Identity” or “identical”** - As used herein, the term “identity” or “identical”, when used in a relationship between the sequences of two or more amino acid sequences, refers to the degree of sequence relatedness between amino acid sequences, as determined by the number of matches between strings of two or more amino acid residues. “Identity”  
10 measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., “algorithms”). Identity of related amino acid sequences can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University  
15 Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York,  
20 1991; and Carillo *et al.*, SIAM J. Applied Math. 48, 1073 (1988). Preferred methods for determining identity are designed to give the largest match between the sequences tested. Methods of determining identity are described in publicly available computer programs. Preferred computer program methods for determining identity between two sequences include the GCG program package, including GAP (Devereux *et al.*, Nucl. Acid. Res. \2,  
25 387 (1984); Genetics Computer Group, University of Wisconsin, Madison, Wis.), BLASTP, BLASTN, and FASTA (Altschul *et al.*, J. Mol. Biol. 215, 403-410 (1990)). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST Manual, Altschul *et al.*, NCB/NLM/NIH Bethesda, Md. 20894; Altschul *et al.*, supra). The well-known Smith Waterman algorithm  
30 may also be used to determine identity.

**“Modified antibody”** - As used herein, the term “modified antibody” includes synthetic forms of antibodies which are altered such that they are not naturally occurring, e.g., antibodies that comprise at least two heavy chain regions but not two complete heavy chains (such as, domain deleted antibodies or minibodies); multispecific forms of antibodies (e.g., bispecific, trispecific, etc.) altered to bind to two or more different antigens or to different epitopes on a single antigen; heavy chain molecules joined to scFv molecules and the like. ScFv molecules are known in the art and are described, e.g., in US patent 5,892,019. In addition, the term "modified antibody" includes multivalent forms of antibodies (e.g., trivalent, tetravalent, etc., antibodies that bind to three or more copies of the same antigen). In another embodiment, a modified antibody of the invention is a fusion protein comprising at least one heavy chain region lacking a CH2 domain and comprising a binding domain of a protein comprising the binding region of one member of a receptor ligand pair.

**“Mammal”** - As used herein, the term “mammal” refers to any mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is a primate, more preferably a human.

**“Monoclonal antibody”** - As used herein, the term “monoclonal antibody” refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprised in the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibody preparations that include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they may be synthesized uncontaminated by other antibodies. The modifier “monoclonal” is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies useful in the present invention may be prepared by the hybridoma methodology first described by Kohler *et al.*, Nature, 256:495 (1975), or may be made using recombinant DNA methods in bacterial, eukaryotic animal or plant cells (see, e.g., U.S. Pat. No 4,816,567). The

“monoclonal antibodies” may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, Nature, 352:624-628 (1991) and Marks *et al.*, J. Mol. Biol., 222:581-597 (1991), for example.

**“Native sequence”** - As used herein, the term “native sequence” nucleotide refers to a polynucleotide that has the same nucleotide sequence as a polynucleotide derived from nature. Accordingly, a “native sequence” protein is one that has the same amino acid sequence as a protein (e.g., antibody) derived from nature (e.g., from any species). Such native sequence polynucleotides and proteins can be isolated from nature or can be produced by recombinant or synthetic means. A polynucleotide “variant”, as the term is used herein, is a polynucleotide that typically differs from a polynucleotide specifically disclosed herein in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the polynucleotide sequences of the invention and evaluating one or more biological activities of the encoded proteins as described herein and/or using any of a number of techniques well known in the art. A protein “variant”, as the term is used herein, is a protein that typically differs from a protein specifically disclosed herein in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the above protein sequences of the invention and evaluating one or more biological activities of the protein as described herein and/or using any of a number of techniques well known in the art. Modifications may be made in the structure of the polynucleotides and proteins of the present invention and still obtain a functional molecule that encodes a variant or derivative protein with desirable characteristics. When it is desired to alter the amino acid sequence of a protein to create an equivalent, or even an improved, variant or region of a protein of the invention, one skilled in the art will typically change one or more of the codons of the encoding DNA sequence. For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of its ability to bind other proteins (e.g., antigens) or cells. Since it is the binding capacity and nature of a protein that defines that protein’s biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and of course, its underlying DNA coding sequence, and nevertheless obtain a protein with

similar properties. It is thus contemplated that various changes may be made in the amino acid sequences of the disclosed compositions, or corresponding DNA sequences that encode said proteins without appreciable loss of their biological utility or activity. In many instances, a protein variant will contain one or more conservative substitutions. A  
5 “conservative substitution” is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the protein to be substantially unchanged. As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their  
10 hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take several of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine. Amino acid substitutions may further be made on the basis of similarity in polarity, charge, solubility,  
15 hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine,  
20 threonine, phenylalanine and tyrosine. Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain non-conservative changes. In a preferred embodiment, variant proteins differ from a native sequence by substitution, deletion or addition of five  
25 amino acids or fewer. Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydropathic nature of the protein.

**“Pharmaceutically acceptable excipient”** - As used herein, the term “pharmaceutically acceptable excipient” includes any and all solvents, dispersion media, coatings,  
30 antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. Said excipient does not produce an adverse, allergic or other untoward reaction when

administered to an animal, preferably a human. For human administration, preparations should meet sterility, pyrogenicity, and general safety and purity standards as required by regulatory offices, such as, for example, FDA Office or EMA.

**“Specificity”** - As used herein, the term "specificity" refers to the ability to specifically bind (e.g., immunoreact with) a given target, e.g., GPVI. A protein may be monospecific and contain one or more binding sites which specifically bind a target, or a protein may be multispecific and contain two or more binding sites which specifically bind the same or different targets. In an embodiment, an antibody of the invention is specific for more than one target. For example, in an embodiment, a multispecific binding molecule of the invention binds to GPVI and a second molecule.

**“Single-chain Fv”** also abbreviated as **“sFv”** or **“scFv”** - As used herein, the terms “Single-chain Fv”, “sFv” or “scFv” are antibody fragments that comprise the VH and VL antibody domains connected into a single amino acid chain. Preferably, the sFv amino acid sequence further comprises a peptidic linker between the VH and VL domains that enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994); Borrebaeck 1995, *infra*.

**“Subject”** - As used herein, the term “subject” refers to a mammal, preferably a human. In one embodiment, a subject may be a "patient", i.e. a warm-blooded animal, more preferably a human, who/which is awaiting the receipt of, or is receiving medical care or was/is/will be the object of a medical procedure, or is monitored for the development of a disease.

**“Synthetic”** - As used herein, the term "synthetic" with respect to proteins includes proteins which comprise an amino acid sequence that is not naturally occurring. For example, non-naturally occurring proteins are modified forms of naturally occurring proteins (e.g., comprising a mutation such as an addition, substitution or deletion) or proteins which comprise a first amino acid sequence (which may or may not be naturally occurring) that is linked in a linear sequence of amino acids to a second amino acid sequence (which may or may not be naturally occurring) to which it is not naturally linked in nature.

“**Therapeutically effective amount**” means level or amount of agent that is aimed at, without causing significant negative or adverse side effects to the target, (1) delaying or preventing the onset of GPVI-related disease; (2) slowing down or stopping the progression, aggravation, or deterioration of one or more symptoms of the GPVI-related disease; (3) bringing about ameliorations of the symptoms of the GPVI-related disease; (4) reducing the severity or incidence of the GPVI-related disease; or (5) curing the GPVI-related disease. A therapeutically effective amount may be administered prior to the onset of the GPVI-related disease, for a prophylactic or preventive action. Alternatively or additionally, the therapeutically effective amount may be administered after initiation of the GPVI-related disease, for a therapeutic action.

“**Variable region**” or “**variable domain**” - As used herein, the term “variable” refers to the fact that certain regions of the variable domains VH and VL differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its target antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called "hypervariable loops" in each of the VL domain and the VH domain which form part of the antigen binding site. The first, second and third hypervariable loops of the VLambda light chain domain are referred to herein as L1 ( $\lambda$ ), L2 ( $\lambda$ ) and L3 ( $\lambda$ ) and may be defined as comprising residues 24-33 (L1( $\lambda$ ), consisting of 9, 10 or 11 amino acid residues), 49-53 L2 ( $\lambda$ ), consisting of 3 residues) and 90-96 (L3( $\lambda$ ), consisting of 6 residues) in the VL domain (Morea *et al.*, Methods 20:267-279 (2000)). The first, second and third hypervariable loops of the VKappa light chain domain are referred to herein as L1( $\kappa$ ), L2( $\kappa$ ) and L3( $\kappa$ ) and may be defined as comprising residues 25-33 (L1( $\kappa$ ), consisting of 6, 7, 8, 11, 12 or 13 residues), 49-53 (L2( $\kappa$ ), consisting of 3 residues) and 90-97 (L3( $\kappa$ ), consisting of 6 residues) in the VL domain (Morea *et al.*, Methods 20:267-279 (2000)). The first, second and third hypervariable loops of the VH domain are referred to herein as H1, H2 and H3 and may be defined as comprising residues 25-33 (H1, consisting of 7, 8 or 9 residues), 52-56 (H2, consisting of 3 or 4 residues) and 91-105 (H3, highly variable in length) in the VH domain (Morea *et al.*, Methods 20:267-279 (2000)). Unless otherwise indicated, the terms L1, L2 and L3 respectively refer to the first, second and third hypervariable loops of a VL domain, and encompass hypervariable loops obtained from

both VKappa and VLambda isotypes. The terms H1, H2 and H3 respectively refer to the first, second and third hypervariable loops of the VH domain, and encompass hypervariable loops obtained from any of the known heavy chain isotypes, including [gamma], [epsilon], [delta], [alpha] or [mu]. The hypervariable loops L1, L2, L3, H1, H2  
5 and H3 may each comprise part of a “complementarity determining region” or “CDR”, as defined hereinabove.

**“Valency”** -As used herein, the term “valency” refers to the number of potential target binding sites in a protein. Each target binding site specifically binds one target molecule or specific site on a target molecule. When a protein comprises more than one target  
10 binding site, each target binding site may specifically bind the same or different molecules (e.g., may bind to different ligands or different antigens, or different epitopes on the same antigen). The subject binding molecules preferably have at least one binding site specific for a human GPVI molecule. Preferably, the proteins provided herein are monovalent.

**“Treating” or “treatment” or “alleviation”** - As used herein, the terms “treating” or  
15 “treatment” or “alleviation” refers to both therapeutic treatment and prophylactic or preventative measures; wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented. A subject or mammal is successfully “treated” for the targeted pathologic  
20 condition or disorder if, after receiving a therapeutic amount of a protein according to the present invention, the patient shows observable and/or measurable reduction in or absence of one or more of the following: reduction in the number of pathogenic cells; reduction in the percent of total cells that are pathogenic; and/or relief to some extent, of one or more of the symptoms associated with the specific disease or condition; reduced  
25 morbidity and mortality, and improvement in quality of life issues. The above parameters for assessing successful treatment and improvement in the disease are readily measurable by routine procedures familiar to a physician.

The present invention relates to isolated humanized proteins binding to human GPVI, preferably to isolated humanized antibodies against human GPVI. In one embodiment, the isolated protein of the invention is purified.

In one embodiment, the protein of the invention binds to the extracellular domain of  
5 GPVI.

In one embodiment, the protein of the invention binds to the Ig-like C2-type domain 2 (D2) of human GPVI.

Thus, in one embodiment, the protein of the invention binds to an epitope comprising at least one amino acid residue from amino acid residues 114 to 207 of human GPVI (SEQ  
10 ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 207 of human GPVI (SEQ ID NO: 13).

In one embodiment, said epitope comprises at least one amino acid residue from amino acid residues 114 to 187, preferably from 115 to 187, more preferably from 116 to 187,  
15 more preferably from 117 to 187, more preferably from 118 to 186, more preferably from 119 to 185, more preferably from 120 to 184, even more preferably from 121 to 183 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 187, preferably from 115 to 187, more preferably from 116 to 187, more preferably from 117  
20 to 187, more preferably from 118 to 186, more preferably from 119 to 185, more preferably from 120 to 184, even more preferably from 121 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said epitope comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37,  
25 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 or more amino acid residues from amino acid residues 114 to 187, preferably from 115 to 187, more preferably from 116 to 187, more preferably from 117 to 187, more preferably from 118 to 186, more preferably from 119 to 185, more preferably from 120 to 184, even more preferably from 121 to 183 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%,

80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 187, preferably from 115 to 187, more preferably from 116 to 187, more preferably from 117 to 187, more preferably from 118 to 186, more preferably from 119 to 185, more preferably from 120 to 184, even more preferably from 121 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said epitope comprises at least one amino acid residue from amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13).

In one embodiment, said epitope comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 of amino acid residues from 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13).

In one embodiment, said epitope comprises at least one amino acid residue from amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%,

80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13).

In one embodiment, said epitope comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21 of amino acid residues from 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13).

In one embodiment, said epitope comprises at least one amino acid residue from amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said epitope comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22 of amino acid residues from 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, the isolated protein of the invention binds to a conformational epitope.

In one embodiment, said conformational epitope comprises at least one amino acid residue of human GPVI.

- 5 In one embodiment, said conformational epitope comprises at least one amino acid residue of the Ig-like C2-type domain 2 (D2) of human GPVI.

In one embodiment, said conformational epitope comprises at least one amino acid residue from 114 to 207 of human GPVI (SEQ ID NO: 13).

- 10 In one embodiment, said conformational epitope comprises at least one amino acid residue from amino acid residues 114 to 187, preferably from 115 to 187, more preferably from 116 to 187, more preferably from 117 to 187, more preferably from 118 to 186, more preferably from 119 to 185, more preferably from 120 to 184, even more preferably from 121 to 183 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid  
15 residues 114 to 187, preferably from 115 to 187, more preferably from 116 to 187, more preferably from 117 to 187, more preferably from 118 to 186, more preferably from 119 to 185, more preferably from 120 to 184, even more preferably from 121 to 183 of human GPVI (SEQ ID NO: 13).

- 20 In one embodiment, said conformational epitope comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 or more amino acid residues from amino acid residues 114 to 187, preferably from 115 to 187, more preferably from 116 to 187, more preferably from 117 to 187, more preferably from 118 to 186, more preferably from 119 to 185, more preferably from 120 to 184, even more  
25 preferably from 121 to 183 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 187, preferably from 115 to 187, more preferably from 116 to 187, more preferably from 117 to 187, more preferably from 118 to 186, more preferably from

119 to 185, more preferably from 120 to 184, even more preferably from 121 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises at least one amino acid residue from amino acid residues 114 to 142, preferably from 115 to 141, more preferably  
5 from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more  
10 preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 amino acid  
15 residues from amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino  
20 acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises at least one amino acid  
25 residue from amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid  
30 residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more

preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21 amino acid residues from amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13), or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises at least one amino acid residue from amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22 amino acid residues from amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises:

- at least one amino acid residue from amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13); and
- at least one amino acid residue from amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises:

- at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 amino acid residues from amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 142, preferably from 115 to 141, more preferably from 116 to 140, more preferably from 117 to 139, more preferably from 118 to 138, more preferably from 119 to 137, more preferably from 120 to 136, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13); and

- 5       - at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22 amino acid residues from amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises:

- 10       - at least one amino acid residue from amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%,  
15       95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13); and  
20       - at least one amino acid residue from amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%,  
25       98%, 99% of identity over amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises:

- at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21 amino acid residues from amino acid residues 114 to 135, preferably from 115 to 135,

- more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 114 to 135, preferably from 115 to 135, more preferably from 116 to 135, more preferably from 117 to 135, more preferably from 118 to 135, more preferably from 119 to 135, more preferably from 120 to 135, even more preferably from 121 to 135 of human GPVI (SEQ ID NO: 13); and
- 10 - at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22 amino acid residues from amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 165 to 187, preferably from 166 to 186, more preferably from 167 to 185, more preferably from 168 to 184, even more preferably from 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope comprises at least one amino acid residue from amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13); and at least one amino acid residue from amino acid residues 169 to 183 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 169 to 183 of human GPVI (SEQ ID NO: 13).

Thus, in one embodiment, the protein of the invention binds to a conformational epitope comprising at least one amino acid residue from amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13); and at least one amino acid residue from amino acid residues

169 to 183 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 169 to 183 of human GPVI (SEQ ID NO: 13).

5 In one embodiment, said conformational epitope comprises at least one amino acid residue from amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13); and at least one amino acid residue from amino acid residues 169 to 180 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%,  
10 97%, 98%, 99% of identity over amino acid residues 169 to 180 of human GPVI (SEQ ID NO: 13).

Thus, in one embodiment, the protein of the invention binds to a conformational epitope comprising at least one amino acid residue from amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%,  
15 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13); and at least one amino acid residue from amino acid residues 169 to 180 of human GPVI (SEQ ID NO: 13) or from a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 169 to 180 of human GPVI (SEQ ID NO: 13).

20 In one embodiment, said conformational epitope consists of:

- amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13) or of a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13); and
- 25 - amino acid residues 169 to 183 of human GPVI (SEQ ID NO: 13) or of a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 169 to 183 of human GPVI (SEQ ID NO: 13).

Thus, in one embodiment, the protein of the invention binds to a conformational epitope consisting of:

- amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13) or of a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13); and
- 5     - amino acid residues 169 to 183 of human GPVI (SEQ ID NO: 13) or of a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 169 to 183 of human GPVI (SEQ ID NO: 13).

In one embodiment, said conformational epitope consists of:

- 10     - amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13) or of a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13); and
- 15     - amino acid residues 169 to 180 of human GPVI (SEQ ID NO: 13) or of a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 169 to 180 of human GPVI (SEQ ID NO: 13).

Thus, in one embodiment, the protein of the invention binds to a conformational epitope consisting of:

- 20     - amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13) or of a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 121 to 135 of human GPVI (SEQ ID NO: 13); and
- amino acid residues 169 to 180 of human GPVI (SEQ ID NO: 13) or of a sequence sharing at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity over amino acid residues 169 to 180 of human GPVI (SEQ ID NO: 13).

25     Another object of the invention is an isolated protein binding to human GPVI, wherein said protein has a  $K_D$  for binding to human GPVI, preferably soluble human GPVI, less than or equal to 15 nM, preferably less than or equal to 10 nM and more preferably less than or equal to 5 nM.

In one embodiment, the isolated protein of the invention has a  $k_{\text{off}}$  for binding to human GPVI of less than or equal to about  $8 \cdot 10^{-4} \text{ sec}^{-1}$ , preferably less than or equal to about  $6 \cdot 10^{-4} \text{ sec}^{-1}$ , and more preferably less than or equal to about  $4 \cdot 10^{-4} \text{ sec}^{-1}$ .

In one embodiment, the isolated protein of the invention has a  $k_{\text{on}}$  for binding to human  
5 GPVI of at least about  $5 \cdot 10^4 \text{ M}^{-1}\text{sec}^{-1}$ , preferably at least about  $5.5 \cdot 10^4 \text{ M}^{-1}\text{sec}^{-1}$ , and more preferably at least about  $5.9 \cdot 10^4 \text{ M}^{-1}\text{sec}^{-1}$  and more preferably at least about  $6 \cdot 10^4 \text{ M}^{-1}\text{sec}^{-1}$ .

In one embodiment, the  $K_D$  may be determined by Surface plasmon resonance (SPR, BIAcore), using immobilized soluble GPVI at a dose ranging from about 700 to about 1600 resonance units (RU) (corresponding to about 8.5 to about  $11.3 \text{ fmol/mm}^2$ ),  
10 preferably from about 850 to about 1200 RU, more preferably from about 950 to about 1075 RU and/or using PBS pH 7.4 as running buffer, and/or using BIAevaluation version 3.0 software for analyzing data. In one embodiment, soluble GPVI corresponds to the extracellular domain of GPVI fused at its C-terminus via a linker (such as, for example, via a Gly-Gly-Arg linker) to a hFc sequence. This soluble GPVI may be referred to as  
15 soluble GPVI-Fc.

A method for determining the affinity of the protein of the invention for soluble GPVI is shown below:

Binding of the protein of the invention to soluble human GPVI is analyzed with surface plasmon resonance using a BIAcore 2000 system (Uppsala, Sweden).

20 Soluble GPVI-Fc is immobilized at a dose ranging from about 700 to about 1600 RU (corresponding to about 8.5 to about  $11.3 \text{ fmol/mm}^2$ ), preferably from about 850 to about 1200 RU, more preferably from about 950 to about 1075 RU, and even more preferably from about 960 to about 1071 RU onto a Carboxy-Methyl Dextran CM5 sensor chip using the amine coupling method (Wizard procedure). The protein is then passed over the  
25 immobilized GPVI-Fc in PBS pH 7.4 (10 mM phosphate, 138 mM NaCl, 2.7 mM KCl, pH 7.42 at  $25.4^\circ\text{C}$ ) at a flow rate of  $20 \mu\text{L/min}$  at  $25^\circ\text{C}$ . Kinetic constants ( $k_{\text{on}}$ ,  $k_{\text{off}}$ ) and affinity are determined using BIAevaluation version 3.0 software, by fitting data to binding model. PBS pH 7.4 is the running buffer.

In an embodiment, said protein is an antibody molecule selected from the group consisting of a whole antibody, a humanized antibody, a single chain antibody, a dimeric single chain antibody, a Fv, a Fab, a F(ab)'<sub>2</sub>, a defucosylated antibody, a bi-specific antibody, a diabody, a triabody and a tetrabody.

- 5 In another embodiment, said protein is an antibody fragment selected from the group consisting of a unibody, a domain antibody, and a nanobody.

In another embodiment, said protein is an antibody mimetic selected from the group consisting of an affibody, an affilin, an affitin, an adnectin, an atrimer, an evasin, a DARPin, an anticalin, an avimer, a fynomer, a versabody and a duocalin.

- 10 A domain antibody is well known in the art and refers to the smallest functional binding units of antibodies, corresponding to the variable regions of either the heavy or light chains of antibodies.

- A nanobody is well known in the art and refers to an antibody-derived therapeutic protein that contains the unique structural and functional properties of naturally-occurring heavy chain antibodies. These heavy chain antibodies may contain a single variable domain (VHH) and two constant domains (CH2 and CH3).
- 15

- A unibody is well known in the art and refers to an antibody fragment lacking the hinge region of IgG4 antibodies. The deletion of the hinge region results in a molecule that is essentially half the size of traditional IgG4 antibodies and has a univalent binding region rather than the bivalent binding region of IgG4 antibodies.
- 20

An affibody is well known in the art and refers to affinity proteins based on a 58 amino acid residue protein domain, derived from one of the IgG binding domain of staphylococcal protein A.

- DARPin (Designed Ankyrin Repeat Proteins) are well known in the art and refer to an antibody mimetic DRP (designed repeat protein) technology developed to exploit the binding abilities of non-antibody proteins.
- 25

Anticalins are well known in the art and refer to another antibody mimetic technology, wherein the binding specificity is derived from lipocalins. Anticalins may also be formatted as dual targeting protein, called Duocalins.

Avimers are well known in the art and refer to another antibody mimetic technology.

- 5   Versabodies are well known in the art and refer to another antibody mimetic technology. They are small proteins of 3-5 kDa with >15% cysteines, which form a high disulfide density scaffold, replacing the hydrophobic core the typical proteins have.

10   In one embodiment, the protein of the invention is monovalent, and is preferably selected from a whole monovalent antibody, a humanized monovalent antibody, a single chain antibody, a Fv, a Fab, or an antibody fragment selected from the group consisting of a unibody, a domain antibody, and a nanobody; or a monomeric antibody mimetic selected from the group consisting of an affibody, an affilin, an affitin, an adnectin, an atrimer, an evasin, a DARPIn, an anticalin, an avimer, a fynomer, and a versabody.

15   In another embodiment, said protein is an immunoconjugate comprising an antibody or fragment thereof conjugated to a therapeutic agent.

In another embodiment, said protein is a conjugate comprising the protein of the invention conjugated to an imaging agent. Said protein could be used for example for imaging applications.

In an embodiment, said protein is a monoclonal antibody.

- 20   In another embodiment, said protein is a polyclonal antibody.

Another object of the invention is an anti-hGPVI antibody or antigen binding fragment thereof wherein the variable region of the heavy chain comprises at least one of the followings CDRs:

**VH-CDR1:** GYTFTSYNMH (SEQ ID NO: 1);

- 25   **VH-CDR2:** GIYPGNGDTSYNQKFQG (SEQ ID NO: 2); and

**VH-CDR3:** GTVVGDWYFDV (SEQ ID NO: 3).

CDR numbering and definition are according to the Kabat/Chothia definition.

Another object of the invention is an anti-hGPVI antibody or antigen binding fragment thereof wherein the variable region of the light chain comprises at least one of the followings CDRs:

5 **VL-CDR1:** RSSQSLENSNGNTYLN (SEQ ID NO: 4);

**VL-CDR2:** RVSNRFS (SEQ ID NO: 5); and

**VL-CDR3:** LQLTHVPWT (SEQ ID NO: 6).

CDR numbering and definition are according to the Kabat/Chothia definition.

10 In one embodiment, the anti-hGPVI antibody or antigen binding fragment thereof of the invention comprises:

- in the heavy chain, at least one of the following CDR: GYTFTSYNMH (SEQ ID NO: 1), GIYPGNGDTSYNQKFQG (SEQ ID NO: 2) and GTVVGDWYFDV (SEQ ID NO: 3), and/or
  - in the light chain, at least one of the following CDR: RSSQSLENSNGNTYLN (SEQ ID NO: 4), RVSNRFS (SEQ ID NO: 5), and LQLTHVPWT (SEQ ID NO: 6).
- 15

In another embodiment of the invention, the anti-hGPVI antibody or antigen binding fragment thereof comprises in its heavy chain the 3 CDRs SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3.

20 In another embodiment of the invention, the anti-hGPVI antibody or antigen binding fragment thereof comprises in its light chain the 3 CDRs SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6.

In another embodiment of the invention, the anti-hGPVI antibody or antigen binding fragment thereof comprises in its heavy chain the 3 CDRs SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, and in its light chain the 3 CDRs SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6.

25

In another embodiment of the invention, the anti-hGPVI antibody or antigen binding fragment thereof comprises in its heavy chain the 3 CDRs SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, and in its light chain the 3 CDRs SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6, optionally wherein one, two, three or more of the amino acids in any of  
 5 said sequences may be substituted by a different amino acid.

According to the invention, any of the CDRs 1, 2 and 3 of the heavy and light chains may be characterized as having an amino acid sequence that shares at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity with the particular CDR or sets of CDRs listed in the corresponding SEQ ID NO.

10 In another embodiment of the invention, the anti-hGPVI antibody or antigen binding fragment thereof is selected from the group consisting of an antibody having:

- (i) the heavy chain CDR 1, 2 and 3 (VH-CDR1, VH-CDR2, VH-CDR3) amino acid sequences as shown in SEQ ID NO: 1, 2 and 3 respectively; and
- (ii) the light chain CDR 1, 2 and 3 (VL-CDR1, VL-CDR2, VL-CDR3) amino acid  
 15 sequences as shown in SEQ ID NO: 4, 5 and 6 respectively;

optionally wherein one, two, three or more of the amino acids in any of said sequences may be substituted by a different amino acid.

In one embodiment, the anti-GPVI antibody or antigen binding fragment thereof of the invention comprises a heavy chain variable region comprising or consisting of the  
 20 sequence SEQ ID NO: 7.

(SEQ ID NO: 7)

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYNMHWVRQAPGQGLEWMGGI  
 YPGNGDTSYNQKFQGRVTMTRDTSTSTVYMELSSLRSEDVAVYYCARGTVVG  
 DWYFDVWGQGTLVTVSS

In one embodiment, the anti-GPVI antibody or antigen binding fragment thereof of the invention comprises a light chain variable region comprising or consisting of the sequence SEQ ID NO: 8.

(SEQ ID NO: 8)

5 DIQMTQSPSSLSASVGDRVITICRSSQSLENSNGNTYLNWYQQKPGKAPKLLIY  
RVSNRFSGVPSRFSGSGSGTDFTFTISSLQPEDATYYCLQLTHVPWTFGQGTKV  
EITR

In one embodiment, the anti-GPVI antibody or antigen binding fragment thereof of the invention comprises a light chain variable region comprising or consisting of the sequence  
10 SEQ ID NO: 9.

(SEQ ID NO: 9)

DIQMTQSPSSLSASVGDRVITICSSQSLENSNGNTYLNWYQQKPGKAPKLLIY  
RVSNRFSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCLQLTHVPWTFGQGTKV  
EIKR

15 In another embodiment of the invention, the anti-GPVI antibody (ACT017 antibody) or antigen binding fragment thereof comprises a heavy chain variable region comprising or consisting of the sequence SEQ ID NO: 7 and the light chain variable region comprising or consisting of the sequence SEQ ID NO: 8.

In another embodiment of the invention, the anti-GPVI antibody (ACT006 antibody) or  
20 antigen binding fragment thereof comprises the heavy chain variable region comprising or consisting of the sequence SEQ ID NO: 7 and the light chain variable region comprising or consisting of the sequence SEQ ID NO: 9.

According to the invention, one, two, three or more of the amino acids of the heavy chain or light chain variable regions as described hereinabove may be substituted by a different  
25 amino acid.

In another embodiment, an antibody of the invention comprises heavy and light chain variable regions comprising amino acid sequences that are homologous to the amino acid sequences of the ACT017 antibody described herein, and wherein the antibodies retain the desired functional properties of the protein of the invention.

- 5 In another embodiment, an antibody of the invention comprises heavy and light chain variable regions comprising amino acid sequences that are homologous to the amino acid sequences of the ACT006 antibody described herein, and wherein the antibodies retain the desired functional properties of the protein of the invention.

According to the invention, the heavy chain variable region encompasses sequences that  
10 have 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% or more of identity with SEQ ID NO: 7.

According to the invention, the light chain variable region encompasses sequences that have 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% or more of identity with SEQ ID NO: 8 or SEQ ID NO: 9.

- 15 In any of the antibodies of the invention, e.g. ACT017 or ACT006, the specified variable region and CDR sequences may comprise conservative sequence modifications. Conservative sequence modifications refer to amino acid modifications that do not significantly affect or alter the binding characteristics of the antibody containing the amino acid sequence. Such conservative modifications include amino acid substitutions,  
20 additions and deletions. Modifications can be introduced into an antibody of the invention by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Conservative amino acid substitutions are typically those in which an amino acid residue is replaced with an amino acid residue having a side chain with similar physicochemical properties. Specified variable region and CDR sequences may  
25 comprise one, two, three, four or more amino acid insertions, deletions or substitutions. Where substitutions are made, preferred substitutions will be conservative modifications. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side

chains (e.g. glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g. threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, one or more amino acid residues within the CDR regions of an antibody of the invention can be replaced with other amino acid residues from the same side chain family and the altered antibody can be tested for retained function (i.e., the properties set forth herein) using the assays described herein.

In one embodiment, the invention also provides an antibody that binds essentially the same epitope as ACT017 or ACT006 antibodies. In the present invention, an antibody that binds essentially the same epitope as ACT017 or ACT006 antibodies will be referred as an ACT017-like or ACT006-like antibody, respectively.

In some embodiments of this invention, anti-hGPVI antibodies comprising VH and VL domains, or CDRs thereof may comprise CH1 domains and/or CL domains, the amino acid sequence of which is fully or substantially human. Where the antigen binding protein of the invention is an antibody intended for human therapeutic use, it is typical for the entire constant region of the antibody, or at least a part thereof, to have a fully or substantially human amino acid sequence. Therefore, one or more or any combination of the CH1 domain, hinge region, CH2 domain, CH3 domain and CL domain (and CH4 domain if present) may be fully or substantially human with respect to its amino acid sequence. Advantageously, the CH1 domain, hinge region, CH2 domain, CH3 domain and CL domain (and CH4 domain if present) may all have a fully or substantially human amino acid sequence. In the context of the constant region of a humanized or chimeric antibody, or an antibody fragment, the term “substantially human” refers to an amino acid sequence identity of at least 70%, or at least 80%, or at least 90%, or at least 95%, or at least 97%, or at least 99% with a human constant region. The term “human amino acid sequence” in this context refers to an amino acid sequence which is encoded by a human immunoglobulin gene, which includes germline, rearranged and somatically mutated genes. The invention also contemplates proteins comprising constant domains of “human” sequence which have been altered, by one or more amino acid additions,

deletions or substitutions with respect to the human sequence, excepting those embodiments where the presence of a “fully human” hinge region is expressly required. The presence of a “fully human” hinge region in the anti-hGPVI antibodies of the invention may be beneficial both to minimize immunogenicity and to optimize stability of the antibody. It is considered that one or more amino acid substitutions, insertions or deletions may be made within the constant region of the heavy and/or the light chain, particularly within the Fc region. Amino acid substitutions may result in replacement of the substituted amino acid with a different naturally occurring amino acid, or with a non-natural or modified amino acid. Other structural modifications are also permitted, such as for example changes in glycosylation pattern (e.g. by addition or deletion of N- or O-linked glycosylation sites). Depending on the intended use of the antibody, it may be desirable to modify the antibody of the invention with respect to its binding properties to Fc receptors, for example to modulate effector function. For example cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved effector function. See Caron *et al.*, J. Exp. Med. 176: 1191 - 1195 (1992) and Shopes, B. J. Immunol. 148:2918-2922 (1992).

Another object of the invention is an isolated nucleic sequence encoding the heavy chain variable region of sequence SEQ ID NO: 7. Preferably, said nucleic sequence is SEQ ID NO: 10:

(SEQ ID NO: 10)

CAGGTTTCAGCTGGTTCAGTCAGGGGCTGAGGTGAAGAAGCCTGGAGCCTCA  
GTGAAGGTGTCCTGCAAGGCTTCTGGCTACACATTTACCAGTTACAATATGC  
ACTGGGTAAGACAGGCTCCTGGACAGGGCCTGGAATGGATGGGAGGTATTT  
ATCCAGGAAATGGTGATACTTCTTCTACAATCAGAAGTTCCAGGGCCGAGTTA  
CTATGACTCGGGACACTTCCACCTCTACAGTGTACATGGAGCTCAGCAGCCT  
GAGATCTGAGGACACCGCGGTCTATTACTGTGCAAGAGGCACCGTGGTTCGG  
CGACTGGTACTTCGATGTGTGGGGCCAAGGCACCCTGGTCACCGTGAGCAG  
T

Another object of the invention is an isolated nucleic sequence encoding the light chain variable region of sequence SEQ ID NO: 8. Preferably, said nucleic sequence is SEQ ID NO: 11:

(SEQ ID NO: 11)

5 GACATCCAGATGACCCAGAGCCCAAGCAGCCTGAGCGCCAGCGTGGGTGAC  
AGAGTGACCATCACCTGTAGAAGTAGTCAGAGCCTTGAGAACAGCAACGGA  
AACACCTACCTGAATTGGTACCAGCAGAAGCCAGGTAAGGCTCCAAAGCTG  
CTGATCTACAGAGTTTCCAACCGATTCTCTGGTGTGCCAAGCAGATTCAGCG  
GTAGCGGTAGCGGTACCGACTTCACCTTCACCATCAGCAGCCTCCAGCCAG  
10 AGGACATCGCCACCTACTACTGCCTCCAGCTGACTCATGTCCCATGGACCTT  
CGGTCAGGGCACCAAGGTGGAGATCACCCGG

Another object of the invention is an isolated nucleic sequence encoding the light chain variable region of sequence SEQ ID NO: 9. Preferably, said nucleic sequence is SEQ ID NO: 12

15 (SEQ ID NO: 12)

GACATCCAGATGACCCAGAGCCCAAGCAGCCTGAGCGCCAGCGTGGGTGAC  
AGAGTGACCATCACCTGTAGTGCCAGTCAGAGCCTTGAGAACAGCAACGGA  
AACACCTACCTGAATTGGTACCAGCAGAAGCCAGGTAAGGCTCCAAAGCTG  
CTGATCTACAGAGTTTCCAACCGATTCTCTGGTGTGCCAAGCAGATTCAGCG  
20 GTAGCGGTAGCGGTACCGACTTCACCCTCACCATCAGCAGCCTCCAGCCAG  
AGGACTTCGCCACCTACTACTGCCTCCAGCTGACTCATGTCCCATGGACCTT  
CGGTCAGGGCACCAAGGTGGAGATCAAACGC

Another object of the invention is an expression vector comprising the nucleic sequences encoding the anti-GPVI antibody of the invention. In one embodiment, the expression  
25 vector of the invention comprises at least one of SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NO: 12 or any sequence having a nucleic acid sequence that shares at least 60%, 70%, 75%, 80%, 90%, 95%, 96%, 97%, 98%, 99% of identity with said SEQ ID NO: 10-12.

In one embodiment, the vector of the invention comprises the sequence SEQ ID NO: 10 and a sequence encoding a constant region of a heavy chain. A non-limiting example of a sequence encoding a constant region of a heavy chain is SEQ ID NO: 14.

(SEQ ID NO: 14)

5 GCCTCCACCAAGGGTCCCTCAGTCTTCCCCTGGCACCCCTCCTCCAAGAGCA  
CCTCTGGTGGCACAGCTGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCAGA  
ACCAGTGACTGTGTCATGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACAC  
CTTCCCTGCTGTCTTGCAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTG  
ACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAAT  
10 CACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTCGAGCCTAAGTCATG  
CGACAAGACTCAC

In one embodiment, the vector of the invention comprises the sequence SEQ ID NO: 11 or SEQ ID NO: 12 and a sequence encoding a constant region of a light chain. A non-limiting example of a sequence encoding a constant region of a light chain is SEQ ID NO:  
15 15.

(SEQ ID NO: 15)

ACTGTGGCTGCACCAAGTGTGTTTCATCTTCCCACCTAGCGATGAGCAGTTGA  
AATCTGGAAGTGCCTCTGTCGTGTGCCTCCTGAACAACCTTCTACCCACGGGA  
GGCCAAGGTACAGTGGAAGGTGGATAACGCCCTCCAATCCGGTAACTCCCA  
20 GGAGAGTGTACAGAGCAAGATAGCAAGGACAGCACCTACAGCCTCAGCA  
GCACCCTGACTCTGAGCAAAGCAGACTACGAGAAGCACAAGGTCTACGCCT  
GCGAAGTCACCCATCAGGGCCTGAGTTCCCCTGTCACAAAGAGCTTCAACC  
GGGGAGAGTGT

In one embodiment, the vector of the invention comprises a sequence encoding a signal  
25 peptide. Non-limiting examples of signal peptides sequences include, but are not limited to, SEQ ID NO: 16 and SEQ ID NO: 17.

(SEQ ID NO: 16)

ATGGATATGCGTGTACCAGCTCAACTACTTGGACTTCTATTGCTTTGGCTTC  
GTGGTGCTAGATGT

(SEQ ID NO: 17)

5 ATGGACTGGACTTGGAGAATCCTATTCTTGGTTGCTGCAGCTACAGGTGCTC  
ATTCA

In one embodiment, the vector of the invention comprises SEQ ID NO: 10, and a sequence encoding a constant region of a heavy chain (such as, for example, SEQ ID NO: 14), and a signal peptide sequence. An example of such a vector is a vector comprising SEQ ID  
10 NO: 18. SEQ ID NO: 18 further comprises cloning sites.

(SEQ ID NO: 18)

GCGGCCGCCACCATGGACTGGACTTGGAGAATCCTATTCTTGGTTGCTGCAG  
CTACAGGTGCTCATTACAGGTTTCAGCTGGTTTCAGTCAGGGGCTGAGGTGA  
AGAAGCCTGGAGCCTCAGTGAAGGTGTCCTGCAAGGCTTCTGGCTACACAT  
15 TTACCAGTTACAATATGCACTGGGTAAGACAGGCTCCTGGACAGGGCCTGG  
AATGGATGGGAGGTATTTATCCAGGAAATGGTGATACTTCCTACAATCAGA  
AGTTCCAGGGCCGAGTTACTATGACTCGGGACACTTCACCTCTACAGTGTA  
CATGGAGCTCAGCAGCCTGAGATCTGAGGACACCGCGGTCTATTACTGTGC  
AAGAGGCACCGTGGTCGGCGACTGGTACTTCGATGTGTGGGGCCAAGGCAC  
20 CCTGGTCACCGTGAGCAGTGCCTCCACCAAGGGTCCCTCAGTCTTCCCACTG  
GCACCCTCCTCCAAGAGCACCTCTGGTGGCACAGCTGCCCTGGGCTGCCTGG  
TCAAGGACTACTTCCCAGAACCAGTGACTGTGTCATGGAACCTCAGGCGCCC  
TGACCAGCGGCGTGACACCTTCCCTGCTGTCTTGCAGTCCTCAGGACTCTA  
CTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGAC  
25 CTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAA  
AGTCGAGCCTAAGTCATGCGACAAGACTCACTGATGAGGATCC

In one embodiment, the vector of the invention comprises SEQ ID NO: 8, and a sequence encoding a constant region of a light chain (such as, for example, SEQ ID NO: 15), and a signal peptide sequence. An example of such a vector is a vector comprising SEQ ID NO: 19. SEQ ID NO: 19 further comprises cloning sites.

5 (SEQ ID NO: 19)

GACGTCACCATGGATATGCGTGTACCAGCTCAACTACTTGGACTTCTATTGC  
 TTTGGCTTCGTGGTGCTAGATGTGACATCCAGATGACCCAGAGCCCAAGCA  
 GCCTGAGCGCCAGCGTGGGTGACAGAGTGACCATCACCTGTAGAAGTAGTC  
 AGAGCCTTGAGAACAGCAACGGAAACACCTACCTGAATTGGTACCAGCAGA  
 10 AGCCAGGTAAGGCTCCAAAGCTGCTGATCTACAGAGTTTCCAACCGATTCTC  
 TGGTGTGCCAAGCAGATTCAGCGGTAGCGGTACCGACTTCACCTT  
 CACCATCAGCAGCCTCCAGCCAGAGGACATCGCCACCTACTACTGCCTCCA  
 GCTGACTCATGTCCCATGGACCTTCGGTCAGGGCACCAAGGTGGAGATCAC  
 CCGGACTGTGGCTGCACCAAGTGTGTTTCATCTTCCCACCTAGCGATGAGCAG  
 15 TTGAAATCTGGAAGTGCCTCTGTCGTGTGCCTCCTGAACAACCTTCTACCCAC  
 GGGAGGCCAAGGTACAGTGGAAGGTGGATAACGCCCTCCAATCCGGTAACT  
 CCCAGGAGAGTGTACAGAGCAAGATAGCAAGGACAGCACCTACAGCCTC  
 AGCAGCACCTGACTCTGAGCAAAGCAGACTACGAGAAGCACAAGGTCTAC  
 GCCTGCGAAGTCACCCATCAGGGCCTGAGTTCCCCTGTCACAAAGAGCTTC  
 20 AACCGGGGAGAGTGTTGATGATATC

In one embodiment, the vector of the invention comprises SEQ ID NO: 9, and a sequence encoding a constant region of a light chain (such as, for example, SEQ ID NO: 15), and a signal peptide sequence. An example of such a vector is a vector comprising SEQ ID  
 25 NO: 20. SEQ ID NO: 20 further comprises cloning sites.

(SEQ ID NO: 20)

GACGTCACCATGGATATGCGTGTACCAGCTCAACTACTTGGACTTCTATTGC  
 TTTGGCTTCGTGGTGCTAGATGTGACATCCAGATGACCCAGAGCCCAAGCA

GCCTGAGCGCCAGCGTGGGTGACAGAGTGACCATCACCTGTAGTGCCAGTC  
 AGAGCCTTGAGAACAGCAACGGAAACACCTACCTGAATTGGTACCAGCAGA  
 AGCCAGGTAAGGCTCCAAAGCTGCTGATCTACAGAGTTTCCAACCGATTCTC  
 TGGTGTGCCAAGCAGATTCAGCGGTAGCGGTAGCGGTACCGACTTCACCCT  
 5 CACCATCAGCAGCCTCCAGCCAGAGGACTTCGCCACCTACTACTGCCTCCAG  
 CTGACTCATGTCCCATGGACCTTCGGTCAGGGCACCAAGGTGGAGATCAAA  
 CGCACTGTGGCTGCACCAAGTGTGTTTCATCTTCCCACCTAGCGATGAGCAGT  
 TGAAATCTGGAAGTGCCTCTGTCGTGTGCCTCCTGAACAACTTCTACCCACG  
 GGAGGCCAAGGTACAGTGGAAGGTGGATAACGCCCTCCAATCCGGTAACTC  
 10 CCAGGAGAGTGTACAGAGCAAGATAGCAAGGACAGCACCTACAGCCTCA  
 GCAGCACCTGACTCTGAGCAAAGCAGACTACGAGAAGCACAAGGTCTACG  
 CCTGCGAAGTCACCCATCAGGGCCTGAGTTCCCCTGTCACAAAGAGCTTCA  
 ACCGGGGAGAGTGTTGATGATATC

Another object of the invention is an isolated host cell comprising said vector. Said host  
 15 cell may be used for the recombinant production of the antibodies of the invention. In an  
 embodiment, host cells may be prokaryote, yeast, or eukaryote cells preferably  
 mammalian cells, such as, for example: monkey kidney CV1 line transformed by SV40  
 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned  
 for growth in suspension culture, Graham *et al.*, J. Gen. Virol. 36:59 (1977)); baby  
 20 hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO,  
 Urlaub *et al.*, Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse Sertoli cells (TM4,  
 Mather, Biol. Reprod. 23:243-251 (1980)); mouse myeloma cells SP2/0-AG14 (ATCC  
 CRL 1581 ; ATCC CRL 8287) or NSO (HPA culture collections no. 85110503); monkey  
 kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC  
 25 CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells  
 (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human  
 lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse  
 mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather *et al.*, Annals N.Y.  
 Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep  
 30 G2), as well as DSM's PERC-6 cell line. Expression vectors suitable for use in each of

these host cells are also generally known in the art. It should be noted that the term "host cell" generally refers to a cultured cell line. Whole human beings into which an expression vector encoding an antigen binding protein according to the invention has been introduced are explicitly excluded from the definition of a "host cell".

- 5 Another object of the invention is a method of producing an anti-hGPVI antibody or antigen binding fragment thereof which comprises culturing host cells containing the isolated polynucleotide sequence encoding the anti-hGPVI antibody under conditions suitable for expression of the anti-hGPVI antibody, and recovering the expressed anti-hGPVI antibody. This recombinant process can be used for large scale production of anti-hGPVI antibodies according to the invention, including monoclonal antibodies intended  
10 for *in vitro*, *ex vivo*, *in vivo* therapeutic, diagnostic uses. These processes are available in the art and will be known by the skilled person.

In one embodiment, the protein of the invention may be purified by chromatography, preferably by affinity chromatography, more preferably by affinity chromatography on  
15 protein L agarose.

Therefore, in one embodiment, the protein of the invention comprises a domain for binding protein L (PpL). Methods for transferring PpL-binding activity onto proteins of the invention are described in Muzard *et al.*, Analytical Biochemistry 388, 331-338, 2009 and in Lakhri *et al.*, MAbs. 2016;8(2):379-88, which are incorporated herein by  
20 reference.

Fragments and derivatives of antibodies of this invention (which are encompassed by the term "antibody" or "antibodies" as used in this application, unless otherwise stated or clearly contradicted by context), preferably a ACT017-like or ACT006-like antibody, can  
25 be produced by techniques that are known in the art. "Fragments" comprise a portion of the intact antibody, generally the antigen binding site or variable region. Examples of antibody fragments include Fab, Fab', Fab'-SH, F (ab')<sub>2</sub>, and Fv fragments; diabodies; any antibody fragment that is a protein having a primary structure consisting of one

uninterrupted sequence of contiguous amino acid residues (referred to herein as a “single-chain antibody fragment” or “single chain protein”), including without limitation (1) single-chain Fv molecules (2) single chain proteins containing only one light chain variable domain, or a fragment thereof that contains the three CDRs of the light chain variable domain, without an associated heavy chain moiety and (3) single chain proteins containing only one heavy chain variable region, or a fragment thereof containing the three CDRs of the heavy chain variable region, without an associated light chain moiety; and multispecific antibodies formed from antibody fragments. Fragments of the present antibodies can be obtained using standard methods. For instance, Fab or F(ab')<sub>2</sub> fragments may be produced by protease digestion of the isolated antibodies, according to conventional techniques. It will be appreciated that immunoreactive fragments can be modified using known methods, for example to slow clearance *in vivo* and obtain a more desirable pharmacokinetic profile the fragment may be modified with polyethylene glycol (PEG). Methods for coupling and site-specifically conjugating PEG to a Fab' fragment are described in, for example, Leong *et al.*, Cytokines 16 (3): 106-119 (2001) and Delgado *et al.*, Br. J. Cancer 73 (2): 175- 182 (1996), the disclosures of which are incorporated herein by reference.

Alternatively, the DNA encoding an antibody of the invention, preferably an ACT017-like or ACT006-like antibody, may be modified so as to encode a fragment of the invention. The modified DNA is then inserted into an expression vector and used to transform or transfect an appropriate cell, which then expresses the desired fragment.

An object of the invention is a composition comprising at least one of the protein of the invention as described here above.

Another object of the invention is a pharmaceutical composition comprising at least one of the protein of the invention as described here above and at least one pharmaceutically acceptable excipient.

Pharmaceutically acceptable excipients that may be used in these compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine,

sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances (for example  
5 sodium carboxymethylcellulose), polyethylene glycol, polyacrylates, waxes, polyethylene- polyoxypropylene- block polymers, polyethylene glycol and wool fat.

Another object of the invention is a medicament comprising at least one of the protein of the invention as described here above.

In an embodiment, said protein is an anti-hGPVI antibody or antigen binding fragment  
10 thereof that inhibits GPVI signaling and/or signaling of a ligand of GPVI. As used herein, the term “inhibit” means that the protein is capable of blocking, reducing, preventing or neutralizing GPVI interaction with its ligands, GPVI signaling and/or the activation of molecules from the GPVI signaling pathway.

Examples of ligands of GPVI include, but are not limited to, collagen, fibrin, fibronectin,  
15 vitronectin and laminins.

In an embodiment, said protein is a neutralizing anti-hGPVI antibody.

In an embodiment, said protein inhibits the binding of GPVI to a ligand thereof (such as, for example, collagen, fibrin or any other GPVI ligand capable of inducing downstream signaling and platelet activation).

20 In an embodiment, said protein inhibits and/or prevents platelet aggregation in response to a ligand of GPVI, such as, for example, collagen. In an embodiment, said protein inhibits and/or prevents platelet adhesion to a ligand of GPVI, such as, for example, collagen.

In an embodiment, said protein inhibits and/or prevents GPVI-dependent thrombin  
25 production in response to a ligand of GPVI, such as, for example, fibrin. In an embodiment, said protein inhibits and/or prevent platelet-catalyzed thrombin production in response to collagen and/or to tissue factor.

In an embodiment, said protein inhibits and/or prevents platelet recruitment by a ligand of GPVI (such as, for example, fibrin) via GPVI.

In one embodiment, the protein of the invention induces saturation of platelets in whole blood or in platelet rich plasma when present at a concentration ranging from about 0.1  
5 to about 10  $\mu\text{g/mL}$ , preferably from about 0.5 to about 5  $\mu\text{g/mL}$ , and more preferably from about 1 to about 2  $\mu\text{g/mL}$  (corresponding to about 2 to about 200 nM, preferably from about 10 to about 100 nM, and more preferably from about 20 to about 40 nM).

In one embodiment, the protein of the invention inhibits collagen-induced platelet aggregation when used at a concentration of at least about 15  $\mu\text{g/mL}$ , preferably of at  
10 least about 10  $\mu\text{g/mL}$ , and more preferably of at least about 5  $\mu\text{g/mL}$ . Preferably, the protein of the invention fully inhibits collagen-induced platelet aggregation when used at such concentrations.

In one embodiment, the IC<sub>50</sub> of the protein of the invention for inhibiting collagen-induced platelet aggregation ranges from about 0.5 to about 10  $\mu\text{g/mL}$ , preferably from  
15 about 1 to about 6  $\mu\text{g/mL}$ , more preferably from about 2 to about 3.2  $\mu\text{g/mL}$ .

In one embodiment, the concentration of the protein of the invention reducing by 50% the velocity of collagen-induced platelet aggregation ranges from about 0.5 to about 5  $\mu\text{g/mL}$ , preferably from about 1 to about 3  $\mu\text{g/mL}$ , more preferably of about 2  $\mu\text{g/mL}$ .

In one embodiment, the concentration of the protein of the invention reducing the  
20 intensity of collagen-induced platelet aggregation ranges from about 0.5 to about 10  $\mu\text{g/mL}$ , preferably from about 1 to about 6  $\mu\text{g/mL}$ , more preferably of about 3.2  $\mu\text{g/mL}$ .

In one embodiment, the protein of the invention does not induce depletion of GPVI when administered *in vivo*, such as, for example, when administered at a dose ranging from 0.01 to 500 mg.

25 In one embodiment, the protein of the invention does not induce a decrease in platelet count, i.e., thrombocytopenia, when administered *in vivo*, such as, for example, when administered at a dose ranging from 0.01 to 500 mg.

The present invention also relates to a protein, a composition, pharmaceutical composition or medicament as described hereinabove for treating or for use in treating a disease, disorder or condition related to GPVI.

5 In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to modulate leukocyte-platelet and platelet-endothelium interactions in inflammation and/or thrombosis. Therefore, according to an embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat inflammation and/or thrombosis.

10 In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to modulate, preferably to prevent, platelet aggregation and degranulation.

15 In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat disorders associated with abnormal or aberrant megakaryocyte and/or platelet proliferation, differentiation, morphology, migration, aggregation, degranulation and/or function. Examples of these disorders include, but are not limited to, bleeding disorders (such as, for example, bleeding tendency and/or prolonged bleeding time) such as thrombocytopenia such as, for example, idiopathic thrombocytopenic purpura (ITP) or immune thrombocytopenia.

20 In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat thrombotic disorders (such as, for example, thrombotic occlusion of coronary arteries), hemorrhagic disorders, diseases exhibiting quantitative or qualitative platelet dysfunction and diseases displaying endothelial dysfunction. These diseases include, but are not limited to, coronary artery and cerebral artery diseases.

25 In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat cerebral vascular diseases, including stroke and ischemia, venous thromboembolism diseases (such as, for example, diseases involving leg swelling, pain and ulceration, pulmonary embolism, abdominal venous thrombosis), thrombotic microangiopathies, vascular purpura.

In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat symptoms associated with platelet disorders and/or diseases (such as, for example, bleeding disorders). In particular, the protein, composition, pharmaceutical composition or medicament as described  
5 hereinabove can be used to modulate symptoms associated with ITP such as purpura and severe bleeding problems.

In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat coronary diseases (such as, for example, cardiovascular diseases including unstable angina pectoris, myocardial  
10 infarction, acute myocardial infarction, coronary artery disease, coronary revascularization, coronary restenosis, ventricular thromboembolism, atherosclerosis, coronary artery disease (e. g., arterial occlusive disorders), plaque formation, cardiac ischemia, including complications related to coronary procedures, such as percutaneous coronary artery angioplasty (balloon angioplasty) procedures). With respect to coronary  
15 procedures, such treatment can be achieved via administration of a protein of the invention prior to, during, or subsequent to the procedure. In a preferred embodiment, such administration can be utilized to prevent acute cardiac ischemia following angioplasty.

In another embodiment, the protein, composition, pharmaceutical composition or  
20 medicament as described hereinabove is used to treat disorders resulting from any blood vessel insult that can result in platelet aggregation. Such blood vessel insults include, but are not limited to, vessel wall injury, such as vessel injuries that result in a highly thrombogenic surface exposed within an otherwise intact blood vessel e. g., vessel wall injuries that result in release of ADP, thrombin and/or epinephrine, fluid shear stress that  
25 occurs at the site of vessel narrowing, ruptures and/or tears at the sites of atherosclerotic plaques, and injury resulting from balloon angioplasty or atherectomy.

Further, in certain embodiments, it is preferred that the protein of the invention does not affect other platelet attributes or functions, such as agonist-induced platelet shape change (e.g., GPIb-vWF-mediated platelet activation), release of internal platelet granule

components, activation of signal transduction pathways or induction of calcium mobilization upon platelet activation by agonists that do not interact with GPVI.

In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat disorders associated with aberrant  
5 signal transduction in response to ligands of GPVI (including, without limitation, collagen, fibrin, fibronectin, vitronectin and laminins) or to other extracellular matrix proteins.

In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat disorders associated with aberrant  
10 levels of GPVI expression and/or activity either in cells that normally express GPVI or in cells that do not express GPVI. For example, the protein of the invention can be used to modulate disorders associated with aberrant expression of GPVI in cancerous (e. g., tumor) cells that do not normally express GPVI. Such disorders can include, for example, ones associated with tumor cell migration and progression to metastasis.

15 In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to modulate immunoregulatory functions of platelets.

In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat disorders of liver, bone marrow and  
20 peripheral blood.

In another embodiment, the protein, composition, pharmaceutical composition or medicament as described hereinabove is used to treat disorders in which platelets contribute by modulating inflammatory responses including, without limitation, sustained or prolonged inflammation associated with infection, arthritis, fibrosis or disorders in  
25 which platelets modulate cell functions including, without limitation, cancer cells proliferation and/or dissemination.

Further examples of diseases, disorders or conditions related to GPVI include, but are not limited to, cardiovascular diseases and/or cardiovascular events, such as, for example,

arterial thrombosis including atherothrombosis, ischemic events, acute coronary artery syndrome, myocardial infarction (heart attack), acute cerebrovascular ischemia (stroke), percutaneous coronary intervention, stenting thrombosis, ischemic, restenosis, ischemia, (acute and chronic), diseases of the aorta and its branches (such as aortic aneurysm, thrombosis), peripheral artery disease, venous thrombosis, acute phlebitis and pulmonary embolism, cancer-associated thrombosis (Trousseau syndrome), inflammatory thrombosis and thrombosis associated to infection.

In one embodiment, the pharmaceutical composition or medicament of the invention is for treating or for use in treating arterial or venous thrombosis, restenosis, acute coronary syndrome or cerebrovascular accidents due to atherosclerosis, preferably thrombosis.

Another object of the invention is a method of treating a disease, disorder or condition related to GPVI, wherein said method comprises administering to a subject in need thereof a protein, a composition, a pharmaceutical composition or a medicament of the present invention.

Preferably, a therapeutically effective amount of the protein of the invention is administered to the subject in need thereof.

It will be understood that the total daily usage of the protein of the invention, composition, pharmaceutical composition or medicament of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disease being treated and the severity of the disease; activity of the specific protein employed; the specific composition employed, the age, body weight, general health, sex and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific protein employed; the duration of the treatment; drugs used in combination or coincidental with the specific protein employed; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the compound at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved. However, the daily dosage of the protein may be varied over a wide range from

about 0.01 to 500 mg per adult per day. Preferably, the compositions contain about 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250, 300, 350, 400, 450 or 500 mg of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. A medication typically contains from about 0.01 mg to about 1000 mg of the active ingredient, preferably from 1 mg to about 500 mg of the active ingredient.

In one embodiment, the subject is affected by, preferably is diagnosed with a disease, disorder or condition related to GPVI, preferably a cardiovascular disease and/or event.

In another embodiment, the subject is at risk of developing a disease, disorder or condition related to GPVI, preferably a cardiovascular disease and/or event. Examples of risk include, but are not limited to, family history (such as, for example, genetic predisposition), ethnicity, age, tobacco exposure, high blood pressure (hypertension), high cholesterol, obesity, physical inactivity, diabetes (in particular type 2 diabetes), unhealthy diets, and harmful use of alcohol.

For use in administration to a subject, the composition will be formulated for administration to the subject. The compositions of the present invention may be administered parenterally, by inhalation spray, rectally, nasally, or via an implanted reservoir. The term administration used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques.

Examples of forms adapted for injection include, but are not limited to, solutions, such as, for example, sterile aqueous solutions, gels, dispersions, emulsions, suspensions, solid forms suitable for using to prepare solutions or suspensions upon the addition of a liquid prior to use, such as, for example, powder, liposomal forms and the like.

Sterile injectable forms of the compositions of this invention may be aqueous or an oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent. Among the acceptable vehicles and

solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

Schedules and dosages for administration of the protein in the pharmaceutical compositions of the present invention can be determined in accordance with known methods for these products, for example using the manufacturers' instructions. For example, a protein present in a pharmaceutical composition of this invention can be supplied at a concentration ranging from about 1 to about 100 mg/mL, such as, for example, at a concentration of 1 mg/mL, 5 mg/mL, 10 mg/mL, 50 mg/mL or 100 mg/mL. In one embodiment, the protein is supplied at a concentration of about 10 mg/mL in either 100 mg (10 mL) or 500 mg (50 mL) single-use vials. In one embodiment, the pharmaceutical composition of the invention may comprise a protein of the invention in PBS pH 7.2-7.7. It will be appreciated that these schedules are exemplary and that an optimal schedule and regimen can be adapted taking into account the affinity and tolerability of the particular antibody in the pharmaceutical composition that must be determined in clinical trials.

In an embodiment, the protein of the invention may be used *in vitro* or *in vivo* to identify samples, tissues, organs or cells that express GPVI.

Examples of assays in which the protein of the invention may be used, include, but are not limited to, ELISA, sandwich ELISA, RIA, FACS, tissue immunohistochemistry, Western-blot, and immunoprecipitation.

In an embodiment of the invention, the sample is a biological sample. Examples of biological samples include, but are not limited to, bodily fluids, preferably blood, more preferably blood serum, plasma, synovial fluid, bronchoalveolar lavage fluid, sputum, lymph, ascitic fluids, urine, amniotic fluid, peritoneal fluid, cerebrospinal fluid, pleural fluid, pericardial fluid, and alveolar macrophages, tissue lysates, biopsies and extracts prepared from diseased tissues.

10 In an embodiment of the invention, the term “sample” is intended to mean a sample taken from an individual prior to any analysis.

In another embodiment, the protein of the invention may be labeled for diagnostic or detection purposes. By labeled herein is meant that a compound has at least one element, isotope or chemical compound attached to enable the detection of the compound.

15 Examples of labels include, but are not limited to, isotopic labels such as radioactive or heavy isotopes; magnetic, electric or thermal labels and colored or luminescent dyes. Examples of luminescent dyes include, but are not limited to, lanthanide complexes, quantum dots, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methyl-coumarins, pyrene, malachite green, stilbene, Lucifer yellow, cascade

20 blue, texas red, alexa dyes, and cy dyes.

Another object of the invention is a kit comprising at least one protein of the invention.

By “kit” is intended any manufacture (e.g., a package or a container) comprising at least one reagent, i.e. for example an antibody, for specifically detecting the expression of GPVI. The kit may be promoted, distributed, or sold as a unit for performing the methods

25 of the present invention. Furthermore, any or all of the kit reagents may be provided within containers that protect them from the external environment, such as in sealed containers. The kits may also contain a package insert describing the kit and methods for its use.

The present invention further relates to a method for inhibiting GPVI receptor function and downstream signalling, thereby treating a GPVI related condition, wherein said method comprises administering a protein of the invention to a subject.

In one embodiment, the method for inhibiting GPVI receptor function and downstream  
5 signalling does not impact platelet count, expression of GPVI at the platelet surface nor bleeding time.

In one embodiment, the method for inhibiting GPVI receptor function and downstream signalling is efficient and reversible.

The present invention further relates to a method for inhibiting the binding of GPVI to its  
10 ligands (preferably, but not exclusively, collagen), thereby treating a GPVI related condition, wherein said method comprises administering a protein of the invention to a subject.

In one embodiment, the method for inhibiting the binding of GPVI to its ligands does not impact platelet count, expression of GPVI at the platelet surface nor bleeding time.

15 In one embodiment, the method for inhibiting the binding of GPVI to its ligands is efficient and reversible.

The present invention further relates to a method for inhibiting and/or preventing platelet adhesion to collagen, thereby treating a GPVI related condition, wherein said method comprises administering a protein of the invention to a subject.

20 The present invention further relates to a method for inhibiting and/or preventing collagen-induced platelet aggregation, thereby treating a GPVI related condition, wherein said method comprises administering a protein of the invention to a subject.

The present invention further relates to a method for inhibiting and/or preventing platelet activation, in particular platelet aggregation, in response to collagen, thereby treating a  
25 GPVI related condition, wherein said method comprises administering a protein of the invention to a subject.

The present invention further relates to a method for inhibiting and/or preventing thrombin production in response to collagen and/or to tissue factor, thereby treating a

GPVI related condition, wherein said method comprises administering a protein of the invention to a subject.

The present invention further relates to a method for inhibiting the binding of GPVI to fibrin, thereby treating a GPVI related condition, wherein said method comprises  
5 administering a protein of the invention to a subject.

The present invention further relates to a method for inhibiting and/or preventing platelet recruitment by fibrin via GPVI, thereby treating a GPVI related condition, wherein said method comprises administering a protein of the invention to a subject.

The present invention further relates to a method for inhibiting and/or preventing GPVI-  
10 dependent thrombin production in response to fibrin, thereby treating a GPVI related condition, wherein said method comprises administering a protein of the invention to a subject.

In one embodiment, administering a protein of the invention to a subject does not induce depletion of GPVI *in vivo*.

15 In one embodiment, administering a protein of the invention to a subject does not induce a decrease in platelet count. Thus, in one embodiment, administering a protein of the invention to a subject does not induce thrombocytopenia.

In one embodiment, administering a protein of the invention to a subject does not induce an increase in bleeding time.

20 In one embodiment, the method of the invention comprises administering a therapeutically effective amount of the protein of the invention to the subject.

## BRIEF DESCRIPTION OF THE DRAWINGS

**Figure 1** is a photograph of a gel stained with Coomassie Blue, showing the protein of  
25 the invention. MM: molecular weight marker, FT: flow through fraction, PX: protein purified (eluted from the Protein-L agarose column). CM cell culture conditioned media

**Figure 2** is a graph showing the isotherm binding curves of two humanized Fab of the invention. Purified GPVI-Fc was coated onto microtitration plates and the purified proteins added at increasing concentrations. After washing, bound proteins were revealed using an alkaline phosphatase coupled-specific antibody to human Fab fragment and hydrolysis of the alkaline phosphatase substrate was measured at 485 nm.

**Figure 3** is a curve obtained by flow cytometry with Alexa488-coupled ACT017. ACT017 was coupled to A488 according to the manufacturer's instructions. A488-ACT0017 was added at increasing concentrations to human whole anticoagulated blood or platelet rich plasma from a healthy volunteer. After 20 min at room temperature, samples were diluted and immediately analyzed by flow cytometry. The mean fluorescence of platelets is represented as function of A488-coupled ACT017 concentration. Data from one representative experiment out of 3.

**Figure 4** is a curve showing aggregation of platelets as measured in human platelet-rich plasma (PRP). PRP was pre-incubated with increasing concentrations of purified ACT017 for 10 min at 37°C without stirring before platelet aggregation was triggered by collagen (1  $\mu\text{g.mL}^{-1}$ ). Incubation was prolonged at 37°C with stirring and continuous registration of changes in light transmission. Data from one representative experiment out of 3.

**Figure 5** is a graph showing the capacity of ACT017 to inhibit collagen-induced platelet aggregation quantified on the velocity and intensity of the response. The intensity and velocity of platelet aggregation were measured at each concentration of ACT017. The residual responses were calculated as the ratio of the response in the presence of ACT017 to the response in the absence of ACT017 x100 and are plotted as a function of ACT017 concentration using the non-linear regression competition curve (log (inhibitor) vs response) (three parameters) from Graph Pad Prism (5.0) software. Curves are from one representative experiment out of 3.

**Figure 6** is a combination of graphs showing collagen induced platelet aggregation. The maximal extent of platelet aggregation triggered by the addition of collagen (1  $\mu\text{g/mL}$ ) to

the platelet rich plasma (PRP) of mice having received a vehicle or increasing doses of ACT017 is reported.

**Figure 7** is a combination of graphs showing the expression of GPVI on platelets. Blood obtained before injection and 30 min after injection of vehicle or of increasing doses of ACT017 was incubated with a FITC-coupled anti-GPVI monoclonal antibody 3J24. Left panel: mean fluorescence of platelets of mice treated by vehicle (ACT17 0 mg/kg) or ACT017 at different doses. Middle and right panels: Evolution of GPVI expression between the pre-injection time and 30 min post injection is reported for individual animals treated with the vehicle or with ACT017 (4 mg/kg).

**Figure 8** is a graph showing the number of platelets in blood of mice after administration of ACT017 (4 mg/kg) or of the vehicle (ACT17 0 mg/kg).

**Figure 9** is a combination of graphs showing the bleeding time (**A**) and the blood loss (**B**) of mice 30 min after administration of vehicle (Ctrl), ACT017 (4 mg/kg) or clopidogrel, an antagonist of the P2Y<sub>12</sub> ADP receptor on platelets (10 mg/kg).

**Figure 10** is a combination of graphs showing the mean  $\pm$  SEM intensity (**A**) and velocity (**B**) of collagen-induced platelet aggregation measured 30 minutes and 2 hours after the end of a 15 minutes' infusion of increasing doses of ACT017 (1, 2, 4, 8 mg/kg) to cynomolgus monkeys (n=4).

**Figure 11** is showing the blood platelet count of cynomolgus measured before any treatment (T<sub>0</sub>) or 24 hours after the infusion of vehicle or ACT017 at increasing doses (1, 2, 4, 8 mg/kg).

**Figure 12** is a combination of graphs showing the bleeding time measured in 4 subjects, before treatment (Time = 0) and 30 min and 2 hours after the infusion of ACT017 (1, 2, 4, 8 mg/kg) or vehicle to cynomolgus monkeys.

**Figure 13** is a combination of graphs showing the mean  $\pm$  SD intensity (**A**) and velocity (**B**) of collagen-induced platelet aggregation measured at different time after the beginning of a one-hour infusion of two doses of ACT017 (2 and 8 mg/kg) to cynomolgus monkeys (n=8).

**Figure 14** is a graph showing the level of GPVI expression measured by flow cytometry on the platelets of cynomolgus monkeys ( $n = 8$ ) at different times after the beginning of a one-hour infusion of ACT017 at 2 mg/kg (MFI: Mean Fluorescence Intensity).

## 5 EXAMPLES

The present invention is further illustrated by the following examples.

### Example 1: Production of ACT017 and ACT006

Antibodies ACT017 and ACT006 (Fab fragments) were produced in CHO-S cells  
10 (Invitrogen) by transient transfection, using standard conditions according to the manufacturer instructions.

In brief, immediately before transfection, CHO-S cells were splitted and seeded in 10 mL of serum-free growth medium at a density of  $0.5-1.0 \times 10^6$  cells/mL. Cells were then transfected with a vector comprising a sequence encoding a light chain and a heavy chain  
15 of an antibody of the invention. Culture supernatants were recovered 4 to 5 days after transfection, when cell viability was lower than 80%.

For ACT017, cells were transfected using a vector comprising SEQ ID NO: 19 and SEQ ID NO: 18, wherein SEQ ID NO: 19 comprises sequences encoding the constant and variable regions of the light chain of ACT017 fused to a signal peptide sequence, and  
20 cloning sites, and wherein SEQ ID NO: 18 comprises sequences encoding the constant and variable regions of the heavy chain of ACT017 fused to a signal peptide sequence, and cloning sites.

For ACT006, cells were transfected using a vector comprising SEQ ID NO: 20 and SEQ ID NO: 18, wherein SEQ ID NO: 20 comprises sequences encoding the constant and variable regions of the light chain of ACT006 fused to a signal peptide sequence, and  
25 cloning sites, and wherein SEQ ID NO: 18 comprises sequences encoding the constant

and variable regions of the heavy chain of ACT006 fused to a signal peptide sequence, and cloning sites.

For purification, the conditioned media of cells transiently transfected with the cDNA of the humanized Fab was applied on a Protein L agarose column (PIERCE protein L  
5 Agarose cat n° 20510017) according the manufacturer instructions. The Fab was eluted using Glycine 0.1 M pH 2.5 and fractions collected on Tris 1 M pH 11 to neutralize the pH. After dialysis against PBS, the concentration of the Fab was determined by measuring absorbance at 280 nm and 320 nm for light scattering correction and using an absorbance value for 0.1% solutions ( $A_{280\text{nm}}^{0.1\%}$ ) of 1.5 as determined from the sequence. The protein  
10 concentration was confirmed by a BCA assay. The purity of the Fab was assessed by SDS-PAGE and Coomassie Blue staining.

As shown in Figure 1, in non-reducing conditions, the purified humanized Fab migrated as a major band of 42 kDa in agreement with its amino-acid sequence. After disulfide  
15 bridge reduction, a doublet at 23-24 kDa was observed corresponding to the heavy and light chains.

### Example 2: Binding to GPVI-Fc

Soluble GPVI-Fc was produced as follows: the open-reading frame of the predicted extracellular domain of GPVI was PCR-amplified from the Kozak sequence before the  
20 first methionine to asparagine 269, immediately prior to the predicted transmembrane sequence. The PCR fragment was ligated into a pCDM8 host vector containing the genomic sequence of the human IgG1 Fc domain, such that the extracellular part of the hGPVI cDNA was fused at its C-terminus via a 3 alanine linker to the hFc sequence. The sequenced DNA construct was transfected into HEK 293T cells. GPVI-Fc was purified  
25 from the conditioned media of the cells by affinity chromatography on Protein A agarose according to manufacturer recommendations.

Microtitration plates were coated with GPVI-Fc in PBS (2 µg/mL, 100 µL per well) overnight at 4°C. Nonspecific binding sites were saturated with 100 µL of 1% BSA in PBS for 120 min. The plates were then incubated with increasing concentrations of the

- antibody preparations (100  $\mu$ L in PBS containing 0.1% BSA and 0.1% Tween 20) for 120 min. After 4 washing rounds plates were incubated with an alkaline phosphatase -coupled mouse antibody against human Fab (100  $\mu$ L, in PBS containing 0.1% BSA and 0.1% Tween 20) for 120 min. Finally, 100  $\mu$ L of the substrate solution (paranitrophenylphosphate) were added to each well for 5 min, and the colorimetric reaction stopped by 25  $\mu$ L NaOH 3 M before reading the absorbance at 405 nm. Alternatively, GPVI bound antibodies were detected using peroxidase-coupled protein L and O-Phenylenediamine Dihydrochloride (OPD) as substrate, reading being then performed at 485 nm.
- Figure 2 shows isotherm binding curves of two humanized Fab of the invention on immobilized GPVI-Fc. Analysis of the curves allowed calculating a half-saturation constant of 1 nM for ACT006 and 0.5 nM for ACT017.

### Example 3: Surface plasmon resonance (BIAcore)

- Binding of ACT017, ACT006 and Fab 9O12 to soluble human GPVI were analyzed with surface plasmon resonance using a BIAcore 2000 system (Uppsala, Sweden).

Soluble GPVI-Fc (produced as described in Example 2) was immobilized in acetate buffer 10 mM pH 6 at a dose ranging from 960 to 1071 RU (corresponding about 6.4 to 7.15 fmol/mm<sup>2</sup>) onto a Carboxy-Methyl Dextran CM5 sensor chip using the amine coupling method (Wizard procedure). The protein is then passed over the immobilized GPVI-Fc in PBS pH 7.4 (10 mM phosphate, 138 mM NaCl, 2.7 mM KCl, pH 7.42 at 25.4°C) at a flow rate of 20  $\mu$ L/min at 25°C. Kinetic constants ( $K_A$ ,  $K_D$ ,  $k_{on}$ ,  $k_{off}$ ) and affinity are determined using BIAevaluation version 3.0 software, by fitting data to binding model. PBS pH 7.4 is the running buffer.

- Results are shown in the Table 1 below:

|                                                | <b>ACT017</b>         | <b>ACT006</b>         | <b>mFab 9O12</b>      |
|------------------------------------------------|-----------------------|-----------------------|-----------------------|
| $k_{on}$ (M <sup>-1</sup> .sec <sup>-1</sup> ) | 5.97.10 <sup>4</sup>  | 6.65.10 <sup>4</sup>  | 4.96.10 <sup>4</sup>  |
| $k_{off}$ (sec <sup>-1</sup> )                 | 2.32.10 <sup>-4</sup> | 4.76.10 <sup>-4</sup> | 8.58.10 <sup>-4</sup> |
| $K_A$ (M <sup>-1</sup> )                       | 2.57.10 <sup>7</sup>  | 1.4.10 <sup>7</sup>   | 5.8.10 <sup>7</sup>   |

|                     |             |             |               |
|---------------------|-------------|-------------|---------------|
| K <sub>D</sub> (nM) | 4.1 (±0.28) | 7.3 (±0.24) | 17.03 (±0.73) |
|---------------------|-------------|-------------|---------------|

Table 1

Antibodies of the invention (ACT017 and ACT006) thus present an enhanced affinity for soluble GPVI as compared to monoclonal Fab 9O12 described in the prior art and studied in the same experimental condition.

5

#### **Example 4: Biologic data**

##### *Materials and Methods*

Human platelets: Blood was collected from healthy volunteers who had not taken medication since 10 days by venipuncture onto citrate de sodium 3,2% (Vacutainer  
10 Beckton Dickinson, Le Pont-de-Clais, France). All blood donors were volunteers who gave their free and informed written consent to this research study, which conforms to the ethical standards of the Declaration of Helsinki.

Flow cytometry: The humanized Fab ACT017 was coupled to Alexa488 (Molecular Probes) according to the manufacturer recommendation. Separation was performed using  
15 an antibody conjugate purification kit (Molecular Probes) and concentration determined at 280 and 494 nm. The A488-coupled humanized Fab at different concentrations was incubated with human platelets in whole blood or platelet rich plasma for 20 min at room temperature in the dark. After dilution in PBS, the cells were analyzed by a fluorescence-activated cell sorter (FACS LSR II BD) flow cytometer.

20 Platelet aggregation: Human platelet rich plasma (PRP) was obtained after centrifugation (120 x g; 15 min; 20°C) and immediately used. Platelet aggregation was measured using the APACT® aggregometer and initiated by the addition of 1 µg.mL<sup>-1</sup> type I collagen (Horm collagen, Nycomed, DE) to the PRP containing various concentration of the humanized Fab and continuously recorded. The velocity and intensity of the aggregation  
25 were measured.

GPVI-humanized mice: The *gp6* knock-in mutant mouse line was established as

previously reported by introducing the sequence of the human *gp6* gene into exon 1 at ATG of mouse *gp6* gene (Mangin P *et al.*, A Humanized Glycoprotein VI (GPVI) Mouse Model to Assess the Antithrombotic Efficacies of Anti-GPVI Agents. J Pharmacol Exp Ther. 2012;341(1):156-63. doi: 10.1124/jpet.111.189050. Epub 2012 Jan 11). The mice  
5 were viable and fertile and did not present any hematological defects. Approximately 3700 copies of human GPVI were detected at the platelet surface.

Platelet count: Whole blood was collected into EDTA (6 mM) after severing the mouse tail. The platelet count was determined in a scil Vet abc automatic cell counter (Scil Animal Care Company, Holtzheim, France) set to murine parameters.

10 *Ex vivo* platelet aggregation: Blood collected onto trisodium citrate before injection and 30 minutes post injection of increasing doses of ACT017 or an equivalent volume of vehicle and PRP obtained by centrifugation. Platelet count was adjusted at  $300 \times 10^9 \cdot L^{-1}$  and platelet aggregation triggered by  $1 \mu g \cdot mL^{-1}$  type I collagen was continuously recorded in four-channel CARAT TX4 aggregometer.

15 GPVI expression: Blood samples were collected onto EDTA before injection and 30 minutes post injection of increasing doses of ACT017 or an equivalent volume of vehicle and were incubated with the FITC-coupled anti-GPVI monoclonal antibody 3J24 ( $10 \mu g \cdot mL^{-1}$ ) that recognizes human GPVI at an epitope different from the epitope of ACT017, and fluorescence was measured on a Beckman Coulter Gallios Flow cytometer.

20 Tail bleeding time and blood lost: The extremity of the tail of anesthetized mice was cut transversally with a scalpel (3 mm) and immediately immersed in tube containing 13 mL of 0.9% isotonic saline at 37°C. The bleeding was observed and measured during 30 min, and when necessary bleeding was stopped manually at the 10-minute time point to prevent death. The volume of lost blood was measured based on the hemoglobin content.

25 Cynomolgus monkeys: 8 cynomolgus monkeys free of any previous treatment were used in the study. Platelet aggregation measurements were performed mostly as described for human platelets and using a collagen concentration of  $2.5 mg \cdot mL^{-1}$ . Platelet count was measured in EDTA anticoagulated blood. The bleeding time was measured on vigil monkeys at the surface of the forearm, according to standard clinical procedure and using

0.5 cm Surgicutt™ bleeding time device. GPVI expression was measured mostly as described above using a commercial FITC coupled anti-human GPVI monoclonal antibody (clone 1G5, Biocytex) that cross react with cynomolgus GPVI.

### Results

- 5    *In vitro* binding of ACT017 to human platelets - A typical curve of Alexa488-coupled ACT017 binding to human platelets as determined by flow cytometry is shown on Figure 3. Saturation of platelets is obtained in whole blood and PRP for an antibody concentration of 1 to 2 µg/mL (20 to 40 nM).

- 10    Inhibition, *in vitro*, of collagen-induced aggregation of human platelets - Typical aggregation curves obtained after preincubation of human PRP with increasing concentrations of ACT017 are shown on Figure 4. In these conditions ACT017 fully inhibited collagen-induced platelet aggregation for concentration  $\geq 5$  µg/mL.

- 15    The capacity of ACT017 to inhibit collagen-induced platelet aggregation was quantified on the velocity and intensity of the response as shown on Figure 5. On three different donors, the mean IC<sub>50</sub> was of  $3.2 \pm 2.5$  and  $2 \pm 0.7$  µg.mL<sup>-1</sup> for the intensity and velocity respectively, while a total inhibition was observed for a concentration of  $6.7 \pm 2.9$  µg.mL<sup>-1</sup> in both cases that is in good agreement with the results of binding to platelets.

These results thus demonstrate that the proteins of the invention are potent inhibitors of collagen-induced platelet aggregation.

20

- Willing to analyze *ex vivo* the effect of ACT017 on collagen induced platelet aggregation, platelet count and GPVI expression, GPVI-humanized mice received an intravenous injection of ACT017 (1, 2 or 4 mg/kg) or of vehicle. 30 minutes after injection, blood was collected for analysis of collagen induced platelet aggregation, platelet count and GPVI expression.
- 25

As shown in Figure 6, a complete inhibition of collagen-induced platelet aggregation is observed when mice received ACT017 at a dose  $\geq 0.5$  mg/kg.

Regarding the expression of GPVI, there was no statistical difference in the mean fluorescence of platelets in samples recovered pre and post treatment as well as between the samples from mice treated by vehicle (ACT017 0 mg/kg) or ACT017 at different doses (Figure 7, left panel). Middle and right panels of Figure 7 show the evolution of GPVI expression between the preinjection time and 30 min post injection in individual animals treated with the vehicle or ACT017 4mg/kg.

These results demonstrate that ACT017 does not induce GPVI depletion *in vivo*.

Moreover, the platelet counts of mice having received vehicle (ACT017 0 mg/kg) or ACT017 (4mg/kg) were measured. As shown in Figure 8, no statistical difference between the two groups was evidenced (mean  $\pm$  SD: 847.7 $\pm$ 110.5 vs 842 $\pm$ 115.5).

The bleeding time and blood loss measured by transection of the tail of mice 30 min after the administration of ACT017 (4 mg/kg) were unmodified as compared to values obtained in control mice (vehicle injection) (Figure 9). For comparison, the bleeding time of clopidogrel treated mice (10 mg/kg the day before and two hours before experiment) was significantly increased as well as the blood loss.

This result demonstrates that ACT017 does not induce a decrease in platelet count, i.e., thrombocytopenia *in vivo*.

The effect of ACT017 administration was further characterized in non-human primates. Eight cynomolgus monkeys were enrolled in the study. First, increasing doses of ACT017 (1, 2, 4, 8 mg/kg) were intravenously administrated over 15 minutes. At time 30 minutes and 2 hours after administration, blood was collected. Platelet aggregation was reversibly inhibited in a dose dependent manner (Figure 10). Increasing the dose from 1 to 2 and from 2 to 4 mg/kg increased the effect whereas 8 mg/kg had no additional effect as compared to 4 mg/kg. The platelet count measured 24 hours after the injection was not modified as compared to the values obtained before injection (Figure 11). No significant increase in the bleeding time was observed after treatment with vehicle or 2, 4 or 8 mg/kg ACT017 (Figure 12). Next, after a washing out period, the cynomolgus received ACT017 (2 or 8 mg/kg) administered as a one-hour perfusion. Platelet aggregation and GPVI expression were measured at different time after the beginning of the treatment: (1, 1.5,

2, 4 and 7 hours) (Figures 13 and 14). Collagen-induced platelet aggregation was reversibly inhibited and the duration of the effect was prolonged in cynomolgus treated with 8 mg/kg compared to 2 mg/kg (Figure 13). GPVI expression on platelets remained stable at the different time after the beginning of the treatment as compared to the pretreatment values (Figure 14).

Together, these results confirm in non-human primates that administration of ACT017 efficiently and reversibly inhibits GPVI function without impact on the platelet count, on expression of GPVI at the platelet surface nor on bleeding time.

## Example 5: Epitope mapping

### *Materials and Methods*

To reconstruct epitopes of the target molecule, i.e., human GPVI, a library of peptides was synthesized. An amino functionalized polypropylene support was obtained by grafting with a proprietary hydrophilic polymer formulation, followed by reaction with t-butyloxycarbonyl-hexamethylenediamine (BocHMDA) using dicyclohexylcarbodiimide (DCC) with N-hydroxybenzotriazole (HOBt) and subsequent cleavage of the Boc-groups using trifluoroacetic acid (TFA). Standard Fmoc-peptide synthesis was used to synthesize peptides on the amino-functionalized solid support by custom modified JANUS liquid handling stations (Perkin Elmer).

Synthesis of structural mimics was done using Pepscan's proprietary Chemically Linked Peptides on Scaffolds (CLIPS) technology. CLIPS technology allows to structure peptides into single loops, double-loops, triple loops, sheet-like folds, helix-like folds and combinations thereof. CLIPS templates are coupled to cysteine residues. The side-chains of multiple cysteines in the peptides are coupled to one or two CLIPS templates. For example, a 0.5 mM solution of the P2 CLIPS (2,6-bis(bromomethyl)pyridine) is dissolved in ammonium bicarbonate (20 mM, pH 7.8)/acetonitrile (1:3 (v/v)). This solution is added onto the peptide arrays. The CLIPS template bind to side-chains of two cysteines as present in the solid-phase bound peptides of the peptide-arrays (455 wells plate with 3  $\mu$ L

wells). The peptide arrays are gently shaken in the solution for 30 to 60 minutes while completely covered in solution. Finally, the peptide arrays are washed extensively with excess of H<sub>2</sub>O and sonicated in disrupt-buffer containing 1% SDS/0.1% beta-mercaptoethanol in PBS (pH 7.2) at 70°C for 30 minutes, followed by sonication in H<sub>2</sub>O  
5 for another 45 minutes. The T3 CLIPS carrying peptides were made in a similar way but with three cysteines.

The binding of ACT017 to each of the synthesized peptides was tested in a PEPSCAN-based ELISA. The peptide arrays were incubated with primary antibody solution (overnight at 4°C). After washing, the peptide arrays were incubated with a 1/1000  
10 dilution of an appropriate antibody peroxidase conjugate (SBA; goat anti-human HRP conjugate, Southern Biotech, 2010-05) for one hour at 25°C. After washing, the peroxidase substrate 2,2'-azino-di-3-ethylbenzthiazoline sulfonate (ABTS) and 20 µl/mL of 3 percent H<sub>2</sub>O<sub>2</sub> were added. After one hour, the color development was measured. The color development was quantified with a charge coupled device (CCD)-camera and an  
15 image processing system.

The values obtained from the CCD-camera range from 0 to 3000 mAU, similar to a standard 96-well plate ELISA-reader. The results are quantified and stored into the Peplab database.

To verify the quality of the synthesized peptides, a separate set of positive and negative  
20 control peptides was synthesized in parallel. These were screened with antibody 57.9 (Posthumus *et al.*, J. Virology, 1990, 64:3304-3309).

Intensity profiles for each set of peptides (linear, helical, looped, discontinuous) were analyzed and systematic binding was evaluated.

### Results

25 After several rounds of condition optimization, it was shown that ACT017 systematically recognizes a discontinuous epitope composed of peptide stretches GPAVSSGGDVTLCQ and TVTAAHSGTYRCYSF, where GPAVSSGGDVTLCQ is the dominant part of the recognition site, since CLIPS constrained peptides containing

this sequence can also be bound by ACT017; however, there is no detectable binding with linear peptides containing this sequence.

Without willing to be bound by any theory, it is hypothesized that peptide stretch TVTAAHSGTYRCYSF probably provides additional structural context for the binding.

- 5 The peptide stretch GPAVSSGGDVTLCQ is significantly modified in the mouse GPVI, with several non-synonymous mutations; in contrast, it is highly conserved in non-human primates. Without willing to be bound by any theory, it is suggested that this explains why ACT017 binds to human GPVI and non-human primates GPVI, but not to murine GPVI.
- 10 Without willing to be bound by any theory, it is suggested that the binding of ACT17 to the identified epitope should impair the binding of collagen to the GPVI Ig-like C2-type domain 1 (D1) and/or GPVI dimerization and subsequent high affinity binding to collagen.

## CLAIMS

1. An isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or a fragment thereof,

5 wherein the variable region of the heavy chain comprises the following CDRs:  
**VH-CDR1:** GYTFTSYNMH (SEQ ID NO: 1);  
**VH-CDR2:** GIYPGNGDTSYNQKFQG (SEQ ID NO: 2); and  
**VH-CDR3:** GTVVGDWYFDV (SEQ ID NO: 3); and  
 wherein the variable region of the light chain comprises the following CDRs:  

10 **VL-CDR1:** RSSQSLENSNGNTYLN (SEQ ID NO: 4);  
**VL-CDR2:** RVSNRFS (SEQ ID NO: 5); and  
**VL-CDR3:** LQLTHVPWT (SEQ ID NO: 6).
2. The isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to claim 1, wherein said antibody or fragment thereof

15 binds to an epitope comprising at least one amino acid residue from amino acid residues 114 to 142 of hGPVI (SEQ ID NO: 13) or from a sequence sharing at least 60% of identity over amino acid residues 114 to 142 of hGPVI (SEQ ID NO: 13); and at least one amino acid residue from amino acid residues 165 to 187 of hGPVI (SEQ ID NO: 13) or from a sequence sharing at least 60% of identity over

20 amino acid residues 165 to 187 of hGPVI (SEQ ID NO: 13).
3. The isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to claim 1 or claim 2, wherein said antibody or fragment thereof has a KD for binding to hGPVI less than 15 nM, wherein said KD is measured by surface plasmon resonance using 960 to 1071 RU of soluble

25 human GPVI and using PBS pH 7.4 as running buffer and wherein said isolated humanized antibody does not induce a GPVI depletion phenotype in vivo.
4. The isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to any one of claims 1 to 3, being an antibody molecule selected from the group consisting of a whole antibody, a humanized

antibody, a single chain antibody, a Fv, a Fab; or an antibody fragment selected from the group consisting of a unibody, a domain antibody, and a nanobody.

- 5       **5.**   The isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to any one of claims **1** to **4**, being a monovalent antibody.
- 10       **6.**   The isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to any one of claims **1** to **5**, wherein the amino acid sequence encoding the heavy chain variable region is SEQ ID NO: **7** and the amino acid sequence encoding the light variable region is SEQ ID NO: **8**, or any  
15       sequence having an amino acid sequence that shares at least 60% of identity with said SEQ ID NO: **7** or **8**.
- 15       **7.**   The isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to any one of claims **1** to **5**, wherein the amino acid sequence encoding the heavy chain variable region is SEQ ID NO: **7** and the  
20       amino acid sequence encoding the light variable region is SEQ ID NO: **9**, or any sequence having an amino acid sequence that shares at least 60% of identity with said SEQ ID NO: **7** or **9**.
- 20       **8.**   A composition comprising the isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to any one of claims **1** to **7**.
- 9.**   A pharmaceutical composition comprising the isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to any one of claims **1** to **7** and at least one pharmaceutically acceptable excipient.
- 25       **10.** Use of the isolated humanized antibody binding to human Glycoprotein VI (hGPVI) or fragment thereof according to any one of claims **1** to **7** for the manufacture of a medicament for treating a GPVI-related condition.
- 11.** The use according to claim **10**, wherein said GPVI-related condition is a cardiovascular disease selected from arterial and venous thrombosis, restenosis, acute coronary syndrome, ischemic cerebrovascular accidents, cerebral vascular

diseases, venous thromboembolism diseases, thrombotic microangiopathies, and vascular purpura.

- 5       **12.** The use according to claim **10**, wherein said GPVI-related condition is a cardiovascular disease selected from coronary artery diseases and cerebral artery diseases.
- 10       **13.** The use according to claim **10**, wherein said GPVI-related condition is a cardiovascular disease selected from atherothrombosis, ischemic events, acute coronary artery syndrome, myocardial infarction, stroke, percutaneous coronary intervention, stenting thrombosis, ischemic restenosis, acute ischemia, chronic ischemia, diseases of the aorta and its branches, peripheral artery disease, venous thrombosis, acute phlebitis, pulmonary embolism, cancer-associated thrombosis, inflammatory thrombosis and thrombosis associated to infection.
- 15       **14.** An expression vector comprising at least one nucleic acid sequence encoding a heavy chain or a light chain of an antibody according to any one of claims **1** to **7**, wherein said expression vector comprises at least one of SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NO: 12 or any sequence having a nucleic acid sequence that shares at least 95% of identity with said SEQ ID NO: 10-12.