

Y742

- 1 -

## DESCRIPTION

COMPOSITION FOR REGENERATING HAIR FOLLICLES COMPRISING  
CD36-EXPRESSING DERMAL SHEATH CELLS

5

TECHNICAL FIELD**[0001]**

The present invention relates to a composition comprising dermal sheath (DS) cells expressing CD36 antigen (thrombospondin receptor) (to be referred to as "CD36-expressing DSc") and arbitrarily, dermal papilla (DP) cells (to be referred to as "DPc"), a method for regenerating hair follicles using such a composition, and an animal or three-dimensional skin model having hair follicles regenerated by such a method.

15

BACKGROUND ART**[0002]**

Hair is viewed to be extremely important in terms of aesthetic appearance. Thus, hair loss caused by congenital or acquired factors is a serious problem for many people. In today's society having a large elderly population and containing numerous sources of stress in particular, there are a growing number of opportunities for being at risk to the loss of scalp hair due to various acquired factors. In order to deal with this situation, various attempts have been made to provide superior hair tonics that more effectively demonstrate hair growth effects including the promotion of hair growth and thicker hair.

20

25

**[0003]**

Hair follicles are exceptional organs that continue to repeatedly self-regenerate throughout nearly the entire life of a mature living organism. Elucidation of the mechanisms of this self-regeneration are expected to lead to clinical applications for which there are considerable demand, such as hair loss treatment using tissue or cell transplants, and the construction of natural and functionally superior skin sheets containing

30

35

- 2 -

hair follicles and sebaceous glands. In recent years, research has progressed rapidly on follicular epithelial stem cells (epithelial cells) accompanying the growing interest in stem cell research, and the properties of dermal papilla cells, which are follicle-specific mesenchymal cells, have also been gradually determined. Dermal papilla cells fulfill the role of a so-called control tower that induces activation signals to follicular epithelial stem cells for self-regeneration of hair follicles, and have been determined, along with follicular epithelial stem cells, to be essential cells in follicle reconstruction evaluation systems (Kishimoto, et al., Proc. Natl. Acad. Sci. USA (1999), Vol. 96, pp. 7336-7341: Non-Patent Document 1).

**[0004]**

Dermal papilla (DP) and dermal sheath (DS) surrounding the periphery of hair follicles both differ from epithelial cells composing the majority of hair follicles in that they are composed of mesenchymal-derived cell populations. With respect to DS, numerous findings have been reported in recent years suggesting its importance with respect to hair follicle formation. DS has been reported to be regenerated from DP in an experiment involving transplantation of hair bulbs-removed hair follicles of rat whiskers (1), and follicle regeneration has been reported to be induced in mice by transplanting DS of hair follicles from which the lower half has been severed (2). In addition, Jahoda, et al. (Development, 1992, Apr:114(4), pp. 887-897: Non-Patent Document 2) reported that follicle regeneration can be induced by transplanting DS to humans (Horne, K.A. and Jahoda, C.A., Development, 1992, Nov:116(3), pp. 563-571: Non-Patent Document 3). Moreover, the Tobin, Paus et al. group reported that cells migration occur between DS and DP, and that proliferation of dermal sheath cells (DSc) begins prior to DPc that begin to proliferate during the hair growth phase (Tobin, D.J. et al., J. Invest.

- 3 -

Dermatol., 120, pp. 895-904, 2003: Non-Patent Document 4).

**[0005]**

5 In this manner, although DS has a high possibility of fulfilling an important role in the formation of hair follicles, the mechanisms of action have not been well known, and even the properties of DSc are not understood. Therefore, we investigated the gene expression profile that characterizes DSc, and analyzed the properties for  
10 the purpose of clarifying the mechanism of action in the follicle formation.

Prior Art Documents

Non-Patent Documents

**[0006]**

15 Non-Patent Document 1: Kishimoto, et al., Proc. Natl. Acad. Sci. USA (1999), Vol. 96, pp. 7336-7341

Non-Patent Document 2: Jahoda, et al., Development, 1992, Apr:114(4), pp. 887-897

20 Non-Patent Document 3: Horne, K.A. and Jahoda, C.A., Development, 1992, Nov:116(3), pp. 563-571

Non-Patent Document 4: Tobin, D.J. et al., J. Invest. Dermatol., 120, pp. 895-904, 2003

25 Non-Patent Document 5: Linder, J. et al., Federation of American Societies for Experimental Biology, 14(2), 319 (2000)

SUMMARY OF THE INVENTION

Problems to be Solved by the Invention

**[0007]**

30 An object of the present invention is to provide a novel follicle regeneration system.

Means for Solving the Problems

**[0008]**

35 As a result of investigating the gene expression profile of DSc using a microarray, 304 genes were identified as genes with two or more higher expression rate in DSc in comparison with that in DPc and fibroblasts (FBc). As a result of categorizing these

- 4 -

genes by function using GeneSpring's GeneOntology, many of the genes belonged to the category of vascular-related factors, thereby suggesting involvement between DS and blood vessels. Among these, DSc were found to highly  
5 express CD36, and this expression of CD36 by DSc was determined to be correlated with expression of HGF (hepatocyte growth factor) that demonstrates hair growth promotional effects (J. Linder, et al.; Federation of American Societies for Experimental Biology, 14(2), 319  
10 (2000): Non-Patent Document 5).

**[0009]**

Thus, the present application includes the inventions indicated below.

[1] A composition for regenerating hair follicles, comprising CD36-expressing dermal sheath (DS) cells.  
15

[2] The composition for regenerating hair follicles of [1], further comprising dermal papilla (DP) cells.

[3] The composition of [2], wherein the ratio of the number of CD36-expressing DSc to the number of DPc is about 10:1 to 1:10.  
20

[4] The composition of [2] or [3], wherein the CD36-expressing DSc and the DPc are both derived from mice or both derived from rats.

[5] The composition of [2] or [3], wherein the CD36-expressing DSc and the DPc are heterologous cells, and each of them are derived from mice, rats or humans.  
25

[6] A method for regenerating hair follicles, comprising transplanting the composition of any of [1] to [5] to a human.

[7] A method for regenerating hair follicles by transplanting the composition of any of [1] to [5] to a recipient animal.  
30

[8] The method of [7], wherein the recipient animal is an immunosuppressed animal.

[9] The method of [7] or [8], wherein the recipient animal is an immunosuppressed animal selected from the group consisting of a nude mouse, SCID mouse and nude  
35

- 5 -

rat.

[10] A method for regenerating hair follicles, comprising producing a three-dimensional skin model containing the composition of any of [2] to [5].

5 [11] A chimeric animal imparted with regenerated hair follicles by transplanting the composition of any of [1] to [5] to a recipient animal.

[12] The chimeric animal of [11], wherein the recipient animal is an immunosuppressed animal.

10 [13] The chimeric animal of [11] or [12], wherein the recipient animal is an immunosuppressed animal selected from the group consisting of a nude mouse, SCID mouse and nude rat.

[14] A three-dimensional skin model imparted with regenerated hair follicles by producing a three-dimensional skin model containing the composition of any of [2] to [5].

Effects of the Invention

[0010]

20 The hair follicle regeneration composition of the present invention can be used in transplant surgery for regenerating hair follicles and in research and development on hair follicle reconstruction.

BRIEF DESCRIPTION OF THE DRAWINGS

25 [0011]

FIG. 1 is a schematic diagram showing the structure of hair follicle tissue.

FIG. 2 indicates expression level of various types of vascular-related factors in various cells.

30 FIG. 3 is an image obtained by CD36 and CD31 immunohistochemical staining.

FIG. 4 is an image of whole mount stained hair follicles obtained by CD36 and CD31 immunostaining.

35 FIG. 5 indicates the results of co-culturing CD36-positive DSc and vascular endothelial cells.

FIG. 6 indicates the results of HGF expression level in CD36-positive DSc.

- 6 -

EMBODIMENTS TO CARRY OUT THE INVENTION**[0012]**

The present invention provides a composition for regenerating hair follicles comprising DSc and  
5 arbitrarily comprising DPc, a method for regenerating hair follicles using such a composition, and an animal or three-dimensional skin model having hair follicles regenerated by such a method.

**[0013]**

10 CD36 antigen is also referred to as thrombospondin receptor. CD36 is an integral membrane protein found on the surfaces of numerous cell types of vertebrate animals, and is also known as FAT, SCARB3, GP88, glycoprotein IV (gpIV) and glycoprotein IIb (gpIIb).  
15 CD36 is a member of the class B scavenger receptor family of cell surface proteins. In addition to thrombospondin, CD36 also binds with numerous other ligands such as collagen, erythrocytes parasitized by falciparum malaria parasites, oxidized low density lipoproteins, naturally-  
20 occurring lipoproteins, oxidized phospholipids and long-chain fatty acids. In recent research using genetically modified rodents, CD36 was able to be confirmed to fulfill a definite role in fatty acid and sugar metabolism, heart disease, sense of taste and the  
25 transport of vegetable fats and oils in the intestinal tract. CD36 can also be involved in impaired glucose tolerance, atherosclerosis, arterial hypertension, diabetes, cardiomyopathy and Alzheimer's disease.

Furthermore, the relationship between CD36 antigen  
30 and hair growth is completely unknown.

**[0014]**

DSc are cells that compose the sheath portion that surrounds the periphery of DP in hair follicles, and are mesenchymal cells in the same manner as DPc. DP are  
35 considered to be derived from DS, and since DS proliferates prior to the proliferation of DP during the hair growth phase, DS is thought to supply DPc (Tobin,

- 7 -

D.J. et al., J. Invest. Dermatol., 120, pp. 895-904, 2003: Non-Patent Document 4).

**[0015]**

Although there are no particular limitations  
5 thereon, DSc that express CD36 can be sorted from DPc and other cells by, for example, commonly used cell sorting techniques using antibodies, and preferably monoclonal antibodies, to CD36.

**[0016]**

10 The DSc of the present invention can be derived from the skin of all mammals, such as humans, chimpanzees, other primates, domestic animals such as dogs, cats, rabbits, horses, sheep, goats, cows or pigs, and experimental animals such as rats, mice or guinea pigs,  
15 and preferably nude mice, SCID mice or nude rats. In addition, the epidermal site may be a site where there is hair growth such as the scalp, or a site where there is no hair growth such as the foreskin.

**[0017]**

20 DPc (dermal papilla cells) refer to cells that are mesenchymal cells located in the lowest portion of hair follicles, and fulfill the role of a so-called control tower by transducing activation signals to follicular epithelial stem cells for self-regeneration of hair  
25 follicles. Dermal papilla cell preparations containing only activated dermal papilla cells can be prepared according to, for example, the method described in Kishimoto, et al., Proc. Natl. Acad. Sci. USA (1999), Vol. 96, pp. 7336-7341 using transgenic mice. However,  
30 in consideration of yield and the like, these preparations are preferably prepared by, for example, preparing a cell suspension by treating a dermal tissue fraction, obtained by removing epidermal tissue from skin tissue, with collagenase, followed by destroying the  
35 follicular epithelial cells by subjecting the cell suspension to cryopreservation.

**[0018]**

- 8 -

The above-mentioned cryopreservation method can be specifically carried out, for example, in the manner indicated below.

1. Mammalian skin is acquired.

5           2. The skin is then allowed to stand undisturbed for a suitable amount of time, such as overnight, in a protease solution such as a trypsin solution as necessary, the epidermal portion is subsequently removed with a forceps and the like, and the remaining dermis is  
10       treated with a collagenase to prepare a cell suspension.

3. The cell suspension is then filtered with a cell strainer as necessary, and the sediment is removed by allowing to stand undisturbed.

15           4. After measuring the number of cells, the cells are re-suspended in a cryopreservation solution at a suitable cell density such as  $1 \times 10^5/\text{ml}$  to  $1 \times 10^8/\text{ml}$ , and the suspension is dispensed into small aliquots as necessary and then placed in cryopreservation in accordance with ordinary cell storage methods.

20           5. The cells are stored for a suitable amount of time and used after thawing.

[0019]

Although there are no particular limitations thereon, the freezing method consists of storing at a  
25       temperature of  $-20^\circ\text{C}$  or lower, preferably  $-50^\circ\text{C}$  or lower and more preferably  $-80^\circ\text{C}$  or lower in an ultra-deep freezer, or in liquid nitrogen. Although there are no particular limitations on the duration of cryopreservation, it is a period of, for example, 1 day  
30       or more, preferably 3 days or more and more preferably 1 week or more to ensure that the epithelial cells are killed. Furthermore, dermal papilla cells have been confirmed to remain viable even after storing for 4 months in liquid nitrogen. An ordinary cryopreservation  
35       solution used to store cells, such as CellBanker 2 cryopreservation solution (Catalog No. BLC-2, Nippon

- 9 -

Zenyaku Kogyo Co., Ltd.), can be used for the cryopreservation solution.

**[0020]**

5 Measurement of the number of cells can be carried out by a method commonly known among persons with ordinary skill in the art. For example, the number of cells can be measured by placing a cell suspension obtained by diluting the cells with an equal volume of 0.4% Trypan blue stain (No. 15250-061, Invitrogen Corp.)  
10 on a hemocytometer (Eosinophil Counter, SLGC Co., Ltd.), and calculating the number of cells in accordance with the method described in the instruction manual provided with the hemocytometer.

**[0021]**

15 Similar to DSc, the DPc of the present invention can be derived from the skin of all mammals, such as humans, chimpanzees, other primates, domestic animals such as dogs, cats, rabbits, horses, sheep, goats, cows or pigs, and experimental animals such as rats, mice or guinea  
20 pigs, and preferably nude mice, SCID mice or nude rats.

**[0022]**

Preferably, the composition for regenerating hair follicles of the present invention further comprises epithelial cells. Epithelial cells are cells that  
25 compose the majority of the dermis or epidermis of skin, and arise from a single layer of basal cells in contact with the dermis. In using the example of mice, although epithelial cells derived from newborn mice (or fetuses) can be preferably used as epithelial cells, they may also  
30 be cells derived from mature skin, such as the epithelium of resting phase hair or epithelium of growth stage hair, or cultured cells in the form of keratinocytes. These cells can be prepared from the skin of a desired donor animal according to methods commonly known among persons  
35 with ordinary skill in the art.

**[0023]**

In a preferable aspect thereof, the epithelial cells

- 10 -

can be prepared in the manner described below.

1. Mammalian skin is acquired.

2. The epidermis of this skin is treated with trypsin by allowing to stand undisturbed overnight at 4°C in PBS containing 0.25% trypsin as necessary.

3. After peeling off only the epidermal portion with a forceps and the like and slicing into thin sections, the epidermal tissue is treated by suspending for about 1 hour at 4°C in a suitable culture broth (such as keratinocyte culture broth).

4. The suspension is passed through a cell strainer having a suitable pore size, followed by subjecting to centrifugal separation and recovering the epithelial cells.

5. The cell preparation is suspended in KGM or SFM medium at a desired cell density, and then allowed to stand undisturbed on ice until just before the time of use.

**[0024]**

Similar to DSc and DPc, the epithelial cells of the present invention can be derived from the skin of all mammals, such as humans, chimpanzees, other primates, domestic animals such as dogs, cats, rabbits, horses, sheep, goats, cows or pigs, and experimental animals such as rats, mice or guinea pigs, and preferably nude mice, SCID mice or nude rats. In addition, the epidermal site may be a site where there is hair growth such as the scalp, or a site where there is no hair growth such as the foreskin.

**[0025]**

Although there are no particular limitations on the ratio of DSc to DPc used, in the composition of the present invention, DSc and DPc are contained at a ratio of preferably 1:10 to 10:1, and more preferably 1:3 to 3:1. Moreover, epithelial cells are contained at a ratio to the total number of DSc and DPc of 1:10 to 10:1, preferably 1:1 to 10:1, more preferably 1:1 to 3:1 and

- 11 -

most preferably 1:1.

**[0026]**

The combination of DSc, DPc and arbitrarily  
epithelial cells may be from the same species or  
5 different species. Thus, the composition for  
regenerating hair follicles of the present invention may  
be, for example, a combination in which all of the DSc,  
DPc and epithelial cells are derived from a human, a  
combination in which all of the DSc, DPc and epithelial  
10 cells are derived from the same species of mammal other  
than human (the above combinations are combinations of  
the same species), a combination in which the DSc and DPc  
are derived from a human while the epithelial cells are  
derived a mammal other than human, a combination in which  
15 one of either the DSc or DPc are derived from a human,  
and the other and the epithelial cells are derived from  
the same species or different species of mammal other  
than a human, and a combination in which one of either  
the DSc or DPc are derived a mammal other than a human  
20 and the other and the epithelial cells are derived from a  
human (the above combinations are combinations of  
different species).

**[0027]**

The method used to transplant the composition for  
25 regenerating hair follicles of the present invention into  
a recipient animal can be in accordance with a known  
transplant method. Reference can be made to, for  
example, Weinberg, et al., J. Invest. Dermatol., Vol. 100  
(1993), pp. 229-236. For example, in the case of  
30 transplanting into nude mice, the cells that have been  
acquired are mixed just before or within 1 hour before  
transplant, the culture broth is removed by  
centrifugation (9000×g, 10 min.), and after forming a  
cell aggregate of about 50 μL to 100 μL, the cell  
35 aggregate is promptly poured into a silicone dome-shaped  
chamber embedded in the skin on the backs of the nude  
mice. Two weeks later, the chamber is carefully removed

- 12 -

and starting an additional 3 weeks later, the presence of hair formation at the transplant site can be observed macroscopically. Transplantation for the purpose of growing hair in animals including humans can be carried out by a similar method, and an appropriate method may be suitably determined by a physician or veterinarian.

**[0028]**

In the case of transplanting the aforementioned composition into a recipient animal, that transplantation may be a homotransplantation, namely an autotransplantation, isotransplantation or allotransplantation, or may be a heterotransplantation. In the case of a homotransplantation, the dermal papilla cell preparation and the epithelial cells are from the same species as the recipient. In the case of a heterotransplantation, either the dermal papilla cell preparation or the epithelial cells are from a different species than the recipient while the other may be from the same species as the recipient, or both may be from a different species than the recipient. Examples of recipient animals include all mammals, such as humans, chimpanzees, other primates, domestic animals such as dogs, cats, rabbits, horses, sheep, goats, cows or pigs, and experimental animals such as rats, mice or guinea pigs, and preferably nude mice, SCID mice or nude rats.

**[0029]**

In addition, a chimeric animal having regenerated hair follicles can be provided by transplanting the aforementioned composition according to the present invention into a suitable recipient animal. The resulting chimeric animal can serve as a useful animal model for, for example, researching and elucidating the mechanism of hair follicle regeneration, or screening drugs or herbs effective for regenerating hair follicles, growing hair or preventing hair loss. The recipient animal is preferably an immunosuppressed animal regardless of the source of each of the cells contained

in system transplanted into the recipient animal. In addition, an animal species capable of being used as an experimental animal can be used for the animal species, and although any animal species may be used provided it coincides with the object of the present invention, preferable examples thereof include mice and rats. Among these animals, examples of immunosuppressed animals when using the example of mice include those in having the trait of a missing thymus in the manner of nude mice. Furthermore, in consideration of the object of the present invention, particularly preferable examples of recipient animals include commercially available nude mice (such as Balb-c nu/nu strain), SCID mice (such as Balb/c-SCID), and nude rats (such as F344/N Jcl-run).

**[0030]**

Moreover, by incorporating the composition according to the present invention in a three-dimensional skin model, a three-dimensional skin model can be provided that has regenerated hair follicles. In this case, however, dermal papilla cells serving as a control tower for hair growth are essential. A three-dimensional skin model can be produced in the manner described below, for example, according to a method commonly known among persons with ordinary skill in the art (Amano, S. et al., Exp. Cell Res. (2001), Vol. 271, pp. 249-362). The three-dimensional skin model respectively contains DSc and DPc at  $1 \times 10^6$  cells/cm<sup>2</sup> to  $1 \times 10^8$  cells/cm<sup>2</sup>, preferably at  $1.0 \times 10^7$  cells/cm<sup>2</sup> to  $1.5 \times 10^7$  cells/m<sup>2</sup>, and more preferably at about  $1.27 \times 10^7$  cells/cm<sup>2</sup>.

**[0031]**

**Method for Producing Three-Dimensional Skin Model**

A suitable number of human fibroblasts are dispersed in DMEM containing 0.1% collagen and 10% FBS followed by dispensing into a Petri dish and immediately allowing to stand undisturbed in a CO<sub>2</sub> incubator at 37°C. After the medium has gelled, the gel is scraped from the side walls

- 14 -

and bottom of the Petri dish so as to suspend the gel in the Petri dish. The cells are then cultured while shaking the collagen gel to contract the gel to about one-fifth its original size and obtain a dermal model.

5 The dermal model is then placed on a stainless steel grid, a glass ring is placed thereon, and 0.4 ml of cultured human epithelial cells ( $1.0 \times 10^6$  cells/ml) dispersed in KGM (keratinocyte growth medium) are injected into the ring and cultured. At this time, DSc  
10 and DPc are simultaneously mixed and injected. Mouse neonatal epithelial cells can also be used instead of cultured human epithelial cells.

Medium consisting of DMEM, KGM, 5% FBS and  $\text{Ca}^{2+}$  is placed in the Petri dish to a degree that the upper  
15 portion of the dermal model is exposed to air followed by culturing, and after about one week, the skin model is observed and assessed for the presence or absence of rudimentary hair follicle formation and regeneration.

**[0032]**

20 Similar to the aforementioned chimeric animal having regenerated hair follicles, a three-dimensional skin model having regenerated hair follicles can be used for researching and elucidating the mechanism of hair follicle regeneration, or screening drugs or herbs  
25 effective for growing hair or preventing hair loss.

**[0033]**

The following provides a more detailed explanation of the present invention by indicating examples thereof.

Example 1

30 **[0034]**

(Method)

Cell Isolation and Culturing

Removal of the dermal portion of human scalp tissue was carried out with a scalpel in DMEM containing 10%  
35 fetal bovine serum (Gibco/ Invitrogen Corp.), and hair follicles were extracted from the severed surface. Hair shafts containing outer root sheath cells (ORS,

- 15 -

follicular epithelial cells) were removed from the hair  
follicles using micro-forceps so as to extract DP and DS.  
The isolated DP were static-cultured in a 35 mm collagen-  
coated tissue dish containing Medium-1 (Nissui low-serum  
5 fibroblast medium containing 10% fetal bovine serum, 10  
ng/ml EGF, 20 ng/ml bFGF, 0.00075%  $\beta$ -mercaptoethanol, 100  
units/ml (titer) penicillin, 0.1 mg/ml (titer)  
streptomycin and 0.25  $\mu$ g/ml (titer) amphotericin B),  
while the isolated DS were treated with collagenase for  
10 40 minutes at 37°C followed by similarly static- culturing  
in a 35 mm collagen-coated tissue dish. After confirming  
the growth of both DP and DS one week later, the  
resulting DPc and DSc were used as experimental samples.  
Commercially available cells (Toyobo Co., Ltd.) were used  
15 for the fibroblasts (FBc). The DSc, DPc and FBc were  
static-cultured for 7 to 10 days using Medium-1.  
Subsequently, the cells were subcultured using trypsin.  
Culturing conditions consisted of culturing in a  
collagen-coated flask (T-75, Iwaki Glass Co., Ltd.) at  
20 37°C and 5% CO<sub>2</sub>. In addition, each of the cells used in  
the experiment had been subcultured one to three times.

Normal human adult dermal microvascular endothelial  
cells (HMVEC, Kurabo Industries Ltd.) were used as  
vascular endothelial cells, and the cells were cultured  
25 in low-serum growth medium (Humedia-MvG, Kurabo  
Industries Ltd.) and then subcultured five times prior to  
use in the experiment.

#### [0035]

Comparison of Gene Expression Profiles of DSc, DPc  
30 and FBc Using the Microarray Method

Total RNA containing mRNA was collected from the  
DSc, DPc and FBc using the RNeasy Micro Kit (Qiagen  
Corp.). The collected RNA was subjected to double-  
stranded cDNA synthesis using Agilent's protocol followed  
35 by synthesis of cRNA labeled with cyanine 3.5. The  
labeled cRNA was hybridized for 17 hours at 65°C on a

- 16 -

microarray chip slide (Agilent, Whole Human Genome (4x44K), G4110) using a two-color assay. Comparisons of the gene expression levels of two types of cells each consisting of DSc and DPc, DPc and FBc and FBc and DSc were carried out on each chip slide using two types each of RNA derived from the DS of two individuals (total of 4 types), two types each of RNA derived from the DP of two individuals (total of 4 types), and two types each of RNA derived from the FB of one individual. After washing the slides, images of fluorescent signals (cyanine 3.5) of the cDNA on the chip were obtained with a dual-laser microarray scanner (Agilent Technologies Inc.). The image data was quantified using Feature Extraction Software Ver. 9.1, and at that time, taggings were made (tagged) to indicate abnormal values and low values at about the same level as background noise followed by analyzing the data. Each expression level was compared by comparing two sets of quantitative values of the acquired signals.

**[0036]**

(Analysis of Microarray Data)

Gene Spring GX 7.3.1 software (Agilent Technologies Inc.) was used to analyze each gene expression level in greater detail using bioinformatics techniques. Abnormal values and low values at about the same level as background noise were tagged in a procedure using Feature Extraction Software Ver. 9.1, and analyses were conducted using those genes that were not tagged. The genes for which there was a difference in expression level between two types of cells were extracted and subjected to functional categorization by using GeneOntology (<http://www.geneontology.org/>). At that time, the degree of statistical significance was tested using Fisher's exact test.

**[0037]**

Cell Staining

Cell staining using CD36 antibody consisted of

- 17 -

seeding DSc in a four well-chamber slide (Nalgene, Nalge  
Nunc International Corp.) subjected to collagen surface  
treatment using an acidic collagen solution (Koken Co.,  
Ltd.) and using the cells after culturing for 1 to 2  
5 days. After washing with PBS, the cells were fixed for  
30 minutes with 4% PFA, washed with PBS and treated for  
10 minutes with PBS solution containing 0.1% Triton X-  
100. After blocking for 30 minutes with PBS containing  
3% BSA, the cells were allowed to react for 1 hour with a  
10 primary antibody solution obtained by diluting CD36  
antibody (ab17044, Abcam Inc.) 50-fold with PBS  
containing 1% BSA. After washing four times with PBS,  
the cells were allowed to react for 1 hour with a  
secondary antibody solution obtained by diluting Alexa  
15 594-labeled anti-mouse IgG antibody (Invitrogen Corp.)  
200-fold with PBS containing 1% BSA. After reacting with  
DAPI solution in order to carry out nuclear staining, the  
cells were washed four times with PBS and sealed with an  
anti-fade reagent (Prolong Gold Antifade Reagent,  
20 Invitrogen Corp.) and a cover glass. The cells were  
observed using a fluorescence microscope (Olympus Corp.).

**[0038]**

## Tissue Staining

Human scalp tissue was embedded in a frozen tissue  
25 embedding agent (OTC Compound, Sakura Finetek Japan Co.,  
Ltd.), and frozen section slides were prepared with a  
frozen section production system (Cryostat, Leica Camera  
AG). After fixing for 15 minutes with 4% PFA, the tissue  
was washed with PBS and allowed to react for 1 hour using  
30 a blocking solution obtained by adding 5% skim milk, 1%  
donkey serum and 0.1% Triton X-100 to PBS. Next, the  
tissue was allowed to react for 1 hour at room  
temperature or overnight at 4°C using a primary antibody  
solution obtained by diluting CD36 antibody solution  
35 (ab17044, Abcam Inc.) or CD31 antibody solution (AF806,  
R&D Co., Ltd.) 50-fold and 100-fold, respectively, with  
the blocking solution. Furthermore, CD31 antibody was

- 18 -

used to label CD31 used as a vascular endothelial cell marker. After washing 3 times with PBS, the tissue was allowed to react for 1 hour at room temperature using a secondary antibody solution obtained by diluting Alexa 594-labeled anti-mouse IgG antibody (Invitrogen Corp.) or Alexa 488-labeled anti-rabbit IgG antibody (Invitrogen Corp.) 200-fold each with blocking solution. After reacting with DAPI solution, the tissue was washed 3 times with PBS and sealed with an anti-fade reagent (Prolong Gold Antifade Reagent) and a cover glass. The tissue was observed using a fluorescence microscope (Olympus Corp.).

**[0039]****Hair Follicle Whole-Mount Staining**

Hair follicles isolated from human tissue were fixed while shake-culturing for 2 hours at 4°C with 4% PFA. The follicles were sequentially subjected to dehydration treatment consisting of treating for 5 minutes each using 0.1% Tween PBS containing 25%, 50% or 75% ethanol (to be referred to as PBST) and treating for 5 minutes each using 100% ethanol. The treated samples were stored in ethanol at -20°C. At the time of use, after rehydrating with the same ethanol series PBST, the follicles were treated for 10 minutes with PBS containing 5% Triton X-100. Subsequently, the follicles were sequentially reacted with the blocking solution used for tissue staining, a primary antibody solution containing CD36 antibody (ab17044, Abcam Inc.) and CD31 antibody (AF806, R&D Co., Ltd.), a secondary antibody solution containing Alexa 594-labeled anti-mouse IgG antibody and Alexa 488-labeled anti-rabbit IgG antibody, and DAPI solution. Furthermore, the follicles were washed 8 times each using PBS containing 0.1% Triton X-100 both between the antibody reaction procedures and after staining. Reaction conditions consisted of reacting overnight at 4°C in the case of the primary antibody solution, and

- 19 -

reacting for 2 to 3 hours at 4°C in the case of the  
secondary antibody solution. The follicles were observed  
with a fluorescence microscope (Olympus Corp.) after  
sealing with an anti-fade reagent (Prolong Gold Antifade  
5 Reagent) and a cover glass.

**[0040]**

RT-PCR

RNA was collected from the cells using TRIzol  
(Invitrogen Corp.) in accordance with the protocol  
10 provided. The concentration of the collected RNA was  
measured with a nucleic acid quantification system  
(Nanodrop, Thermo Scientific Inc.). After making the  
concentrations of RNA targeted for comparison to a same  
level, cDNA was synthesized using oligo(dT) primers from  
15 the RNA using reverse transcriptase (Superscript III,  
Invitrogen Corp.) in accordance with the Invitrogen  
protocol. Quantitative RT-PCR was then carried out using  
the synthesized cDNA as template and using LightCycler®  
FastStart DNA MasterPLUS SYBR Green (Roche Diagnostics  
20 GmbH) for the reaction reagent and LightCycler (Roche  
Diagnostics GmbH) for the reaction device. Composition  
conditions were in accordance with the Roche protocol.  
PCR conditions consisted of initial denaturation for 10  
minutes at 95°C, denaturation for 10 seconds at 95°C,  
25 annealing for 10 seconds at 60°C, and elongation for 10  
seconds at 72°C. The primer data used is as indicated  
below.

G3PDH:

Forward primer: 5'-GCACCGTCAAGGCTGAGAAC-3' (SEQ ID  
30 NO: 1)

Reverse primer: 5'-ATGGTGGTGAAGACGCCAGT-3' (SEQ ID  
NO: 2)

CD36:

Forward primer: 5'-GAGGAACTATATTGTGCCTATTCTTTGGC-3'  
35 (SEQ ID NO: 3)

Reverse primer: 5'-CATAAAAGCAACAAACATCACCACACCAAC-3'

- 20 -

(SEQ ID NO: 4)

CD31:

Forward primer: 5'-ATGCCGTGGAAAGCAGATACTCTAG-3' (SEQ  
ID NO: 5)

5 Reverse primer: 5'-AATTGCTGTGTTCTGTGGGAGCAG-3' (SEQ  
ID NO: 6)

HGF:

Forward primer: 5'-GAGGGAAGGTGACTCTGAATGAG-3' (SEQ  
ID NO: 7)

10 Reverse primer: 5'-AATACCAGGACGATTTGGAATGGCAC-3'  
(SEQ ID NO: 8)

The expression levels of each gene were measured  
using the software provided. Furthermore, G3PDH was used  
as an internal standard, and this was used to correct the  
amount of cDNA of a control group when quantifying each  
gene.

**[0041]**

## Cell Sorting

The cells were sorted using the Cell Separation  
20 Magnet (BD Biosciences Inc.). Operating conditions were  
in accordance with the protocol provided by BD  
Biosciences Inc. After separating the cells using  
trypsin solution, the cell suspension was passed through  
a cell strainer having a pore size of 70  $\mu$ m (Falcon Inc.)  
25 followed by counting the number of cells. 5 million to  
10 million of cells were suspended in 500 ml of PBS  
solution containing 3% fetal bovine serum, followed by  
the addition of CD36 antibody (ab17044, Abcam Inc.) so as  
to be diluted 50-fold and allowing to react for about 15  
30 minutes on ice. After recovering the cells by washing  
and centrifuging using 1xImag Buffer (BD Biosciences  
Inc.), the cells were re-suspended in 30  $\mu$ l of anti-mouse  
IgG1 magnetic particles (BD Biosciences Inc.) and allowed  
to stand undisturbed for 30 minutes on ice. 500  $\mu$ l of  
35 1xImag Buffer (BD Biosciences Inc.) were added followed  
by placing in the Cell Separation Magnet (BD Biosciences

- 21 -

Inc.) and allowing to stand undisturbed for 8 minutes. The supernatant was recovered while being careful so as not to dislodge cells adhered to the lateral surfaces by the magnet, and the resulting supernatant was used as  
5 CD36-negative DSc. Continuing, after again adding 500  $\mu$ l of 1xImag Buffer (BD Biosciences Inc.) and suspending cells adhered to the lateral surfaces, the suspension was placed in the Cell Separation Magnet and allowed to stand undisturbed for 4 minutes followed by removal of the  
10 supernatant. This procedure was further repeated once and the cells adhered to the lateral surfaces were used as CD36-positive DSc. The recovered CD36-positive and CD36-negative DSc were suspended in Medium-1 followed by culturing for 2 to 4 days at 37°C and 5% CO<sub>2</sub> using a  
15 collagen-coated flask (T-25, Iwaki Glass Co., Ltd.) for the culture vessel, followed by use in the experiment.

**[0042]**

## Co-Culturing Experiment

An experiment was conducted using the CD36-positive  
20 and CD-36 negative DSc derived from each specimen for N=3 and 4 times, respectively. 300,000 of the sorted CD36-positive and CD36-negative DSc were respectively seeded to a collagen-coated flask (T-75). Subsequently, after culturing for 2 days in Medium-1, 400,000 HMVEC were  
25 added and co-cultured for 1 day in Humedica-MvG (Kurabo Industries Ltd.). Subsequently, the medium was replaced with medium obtained by adding 100 units/ml (titer) of penicillin, 0.1 mg/ml (titer) of streptomycin, 0.25  $\mu$ g/ml (titer) of amphotericin B and 0.1% BSA (Sigma Corp.) to  
30 vascular endothelial cell basal medium (Humedia-EB2, Kurabo Industries Ltd.). After further co-culturing for 1 day, the cells were separated with trypsin solution followed by proceeding with analysis using FACS. The cells were then passed through a 70  $\mu$ m cell strainer  
35 (Falcon Inc.), suspended in PBS solution containing 3% fetal bovine serum, and allowed to react for 20 minutes

- 22 -

on ice using a primary antibody solution, i.e., CD31 antibody solution (AF806, R&D Co., Ltd.). After washing the cells with PBS solution containing 3% fetal bovine serum, the cells were allowed to react for 20 minutes on ice with secondary antibody solution, i.e., Alexa 488-labeled anti-rabbit IgG antibody (Invitrogen Corp.) and re-suspended in 0.5 ml of PBS solution followed by proceeding with analysis using Cell Lab Quanta SC (Beckman Coulter Inc.). Preparations, including laser accuracy management, were made using the protocol and working reagent designated by Beckman Coulter. The number of CD31-positive cells was measured using the FL1 channel (525 nm). Furthermore, correction was made to eliminate autofluorescence by using endothelial cells that did not react with CD31 antibody. Following measurement, the number of CD31-positive cells was calculated based on total number of cells obtained and the ratio of CD31-positive cells.

**[0043]**

## (Results)

Table 1 shows the expression level of some vascular-related factors. Although vascular-related factors were determined to be highly expressed in DSs, CD36 and HGF were determined to be specifically highly expressed in DSs. FIG. 2 indicates the expression level of various types of vascular-related factors in DSc, DPc, ORS cells and VEC (vascular endothelial cells). CD36 and HGF were determined to be expressed extremely specifically in DS. On the basis of cell staining results as well, CD36-positive cells were observed to be present only in isolated DS cultured cells, while CD36-positive cells were determined to be absent in DPc or FBc (data not shown).

- 23 -

[Table 1]

## Vascular-Related Factors Highly Expressed in DSc

|                                                 | DS/DP | DS/FB | FB/DP |
|-------------------------------------------------|-------|-------|-------|
| CD36                                            | 40.24 | 10.97 | 3.93  |
| HGF                                             | 20.19 | 3.28  | 10.34 |
|                                                 | 14.52 | 4.33  | 4.45  |
|                                                 | 17.85 | 4.14  | 4.47  |
| TBX2 (T-box2 (transcription factor))            | 20.67 | 4.68  | 6.07  |
| VEGFA (vascular endothelial cell growth factor) | 3.41  | 5.12  | 0.50  |
|                                                 | 4.34  | 7.40  | 0.48  |
|                                                 | 2.22  | 4.16  | 0.52  |
|                                                 | 2.68  | 4.55  | 0.60  |
| PDGFA (platelet-derived growth factor)          | 5.82  | 3.67  | 1.80  |
| COX-1 (prostaglandin synthase)                  | 11.73 | 2.24  | 4.88  |
|                                                 | 9.64  | 1.60  | 5.83  |

## [0044]

5 Specific staining of CD36 was also observed in the dermal sheaths of hair follicles, namely DS as a result of immunohistochemical staining of CD36 and CD31 on the sections of hair follicle (FIG. 3). Moreover, the results of hair follicle whole-mount staining indicated

10 dense areas of blood vessels in a portion of the DS, and CD36-positive DSc were determined to be localized in those dense areas. Thus, CD36-positive DSc was suggested to be intimately involved with blood vessels. In addition, although CD36-positive DSc cells are nearly

15 always co-localized with blood vessels, CD36-positive DSc are absent in the vicinity of some vessels (FIG. 4).

Thus, CD36-positive DSc cells were suggested to promote vascularization by, for example, promoting the proliferation of vascular endothelial cells.

## 20 [0045]

In an experiment in which CD36-positive DSc isolated by cell sorting were co-cultured with vascular endothelial cells, the CD36-positive DSc were indicated to significantly promote the proliferation of vascular

25 endothelial cells in comparison with CD36-negative DSc (FIG. 5). Moreover, isolated CD36-positive DSc were also indicated to highly express HGF in comparison with CD36-negative DSc cells (FIG. 6). As mentioned above, HGF is

- 24 -

commonly known as a factor that promotes the growth of  
new hair and hair growth (Non-Patent Document 5). Thus,  
transplantation of CD36-positive DSc into hair follicles  
is clearly effective for the growth of new hair and hair  
5 growth.

- 25 -

CLAIMS

1. A composition for regenerating hair follicles, comprising CD36-expressing dermal sheath cells (DSc).

5 2. The composition for regenerating hair follicles according to claim 1, further comprising dermal papilla cells (DPc).

3. The composition according to claim 2, wherein the ratio of the number of CD36-expressing DSc to the number of DPc is about 10:1 to 1:10.

10 4. The composition according to claim 2 or 3, wherein the CD36-expressing DSc and the DPc are both derived from mice or both derived from rats.

5 5. The composition according to claim 2 or 3, wherein the CD36-expressing DSc and the DPc are heterologous cells, and each of them are derived from mice, rats or humans.

6. A method for regenerating hair follicles, comprising transplanting the composition according to any of claims 1 to 5 to a human.

20 7. A method for regenerating hair follicles by transplanting the composition according to any of claims 1 to 5 to a recipient animal.

8. The method according to claim 7, wherein the recipient animal is an immunosuppressed animal.

25 9. The method according to claim 7 or 8, wherein the recipient animal is an immunosuppressed animal selected from the group consisting of a nude mouse, SCID mouse and nude rat.

30 10. A method for regenerating hair follicles, comprising producing a three-dimensional skin model containing the composition according to any of claims 2 to 5.

35 11. A chimeric animal imparted with regenerated hair follicles by transplanting the composition according to any of claims 1 to 5 to a recipient animal.

12. The chimeric animal according to claim 11, wherein the recipient animal is an immunosuppressed

- 26 -

animal.

13. The chimeric animal according to claim 11 or  
12, wherein the recipient animal is an immunosuppressed  
animal selected from the group consisting of a nude  
5 mouse, SCID mouse and nude rat.

14. A three-dimensional skin model imparted with  
regenerated hair follicles by producing a three-  
dimensional skin model containing the composition  
according to any of claims 2 to 5.