

Description

Background

5 The present invention pertains in general to hematopoietic growth factors and to polynucleotides encoding such factors. The present application pertains in particular to mammalian pluripotent colony stimulating factors, specifically human pluripotent granulocyte colony-stimulating factor (hpG-CSF), to fragments and polypeptide analogs thereof and to polynucleotides encoding the same.

10 The human blood-forming (hematopoietic) system replaces a variety of white blood cells (including neutrophils, macrophages, and basophils/mast cells), red blood cells (erythrocytes) and clot-forming cells (megakaryocytes/platelets). The hematopoietic system of the average human male has been estimated to produce on the order of 4.5×10^{11} granulocytes and erythrocytes every year, which is equivalent to an annual replacement of total body weight. Dexter et al., BioEssays, 2, 154-158 (1985).

15 It is believed that small amounts of certain hematopoietic growth factors account for the differentiation of a small number of progenitor "stem cells" into the variety of blood cell lines, for the tremendous proliferation of those lines, and for the ultimate differentiation of mature blood cells from those lines. Because the hematopoietic growth factors are present in extremely small amounts, the detection and identification of these factors has relied upon an array of assays which as yet only distinguish among the different factors on the basis of stimulative effects on cultured cells under artificial conditions. As a result, a large number of names have been coined
20 to denote a much smaller number of factors. As an example of the resultant confusion the terms, IL-3, BPA, multi-CSF, HCGF, MCGF and PSF are all acronyms which are now believed to apply to a single murine hematopoietic growth factor. Metcalf, Science, 229, 16-22 (1985). See also, Burgess, et al. J.Biol.Chem., 252, 1988 (1977), Das, et al. Blood, 58, 600 (1980), Ihle, et al., J.Immunol., 129, 2431 (1982), Nicola, et al., J.Biol.Chem., 258, 9017 (1983), Metcalf, et al., Int.J.Cancer, 30, 773 (1982), and Burgess, et al. Int.J.Cancer, 26, 647 (1980), relating to various murine growth regulatory glycoproteins.

25 The application of recombinant genetic techniques has brought some order out of this chaos. For example, the amino acid and DNA sequences for human erythropoietin, which stimulates the production of erythrocytes, have been obtained. (See, Lin, PCT Published Application No. 85/02610, published June 20, 1985.) Recombinant methods have also been applied to the isolation of cDNA for a human granulocyte-macrophage colony-stimulating factor. See, Lee, et al., Proc. Natl. Acad. Sci. (USA), 82, 4360-4364 (1985) and Wong, et al., Science, 228, 810-814 (1985). See also Yokota, et al. Proc. Natl. Acad. Sci. (USA), 81, 1070 (1984), Fung, et al., Nature, 307, 233 (1984), and Gough, et al., Nature, 309, 763 (1984) relating to cloning of murine genes, as well as Kawasaki, et al., Science, 230, 291 (1985) relating to human M-CSF.

35 A human hematopoietic growth factor, called human pluripotent colony-stimulating factor (hpCSF) or pluri-poitin, has been shown to be present in the culture medium of a human bladder carcinoma cell line denominated 5637 and deposited under restrictive conditions with the American Type Culture Collection, Rockville, Maryland as A.T.C.C. Deposit No. HTB-9. The hpCSF purified from this cell line has been reported to stimulate proliferation and differentiation of pluripotent progenitor cells leading to the production of all major blood cell types in assays using human bone marrow progenitor cells. Welte et al., Proc. Natl. Acad. Sci. (USA), 82, 1526-1530 (1985). Purification of hpCSF employed: $(\text{NH}_4)_2\text{SO}_4$ precipitation; anion exchange chromatography (DEAE cellulose, DE52); gel filtration (AcA54 column); and C18 reverse phase high performance liquid chromatography. A protein identified as hpCSF, which is eluted in the second of two peaks of activity in C18 reverse phase HPLC fractions, was reported to have a molecular weight (MW) of 18,000 as determined by sodium dodecyl sulphate (SDS)-polyacrylamide gel electrophoresis (PAGE) employing silver staining. HpCSF
40 was earlier reported to have an isoelectric point of 5.5 [Welte, et al., J. Cell. Biochem., Supp 9A, 116 (1985)] and a high differentiation activity for the mouse myelomonocytic leukemic cell line WEHI-3B D⁺ [Welte, et al., UCLA Symposia on Molecular and Cellular Biology, Gale, et al., eds., New Series, 28 (1985)]. Preliminary studies indicate that the factor identified as hpCSF has predominately granulocyte colony-stimulating activity during the first seven days in a human CFU-GM assay.

50 Another factor, designated human CSF- β , has also been isolated from human bladder carcinoma cell line 5637 and has been described as a competitor of murine ¹²⁵I-labelled granulocyte colony-stimulating factor (G-CSF) for binding to WEHI-3B D⁺ cells in a dose-response relationship identical to that of unlabelled murine G-CSF [Nicola, et al., Nature, 314, 625-628 (1985)]. This dose-response relationship had previously been reported to be unique to unlabelled murine G-CSF and not possessed by such factors as M-CSF, GM-CSF, or multi-CSF [Nicola, et al., Proc. Natl. Acad. Sci. (USA), 81, 3765-3769 (1984)]. CSF- β and G-CSF are also unique among
55 CSF's in that they share a high degree of ability to induce differentiation of WEHI-3B D⁺ cells. Nicola, et al., Immunology Today, 5, 76-80 (1984). At high concentrations, G-CSF stimulates mixed granulocyte/macrophage colony-forming cells [Nicola, et al., (1984) supra], which is consistent with preliminary results indicating the

appearance of granulocytic, monocytic, mixed granulocytic/monocytic and eosinophilic colonies (CFU-GEMM) after 14 days incubation of human bone marrow cultures with hpCSF. CSF- β has also been described as stimulating formation of neutrophilic granulocytic colonies in assays which employed mouse bone marrow cells, a property which has been a criterion for identification of a factor as a G-CSF. On the basis of these similarities, human CSF- β has been identified with G-CSF (granulocytic colony stimulating factor). Nicola et al., *Nature*, 314, 625-628 (1985).

Based upon their common properties, it appears that human CST- β of Nicola, et al., *supra*, and the hpCSF of Welte, et al., *supra*, are the same factor which could properly be referred to as a human pluripotent granulocyte colony-stimulating factor (hpG-CSF). Characterization and recombinant production of hpG-CSF would be particularly desirable in view of the reported ability of murine G-CSF to completely suppress an *in vitro* WEHI-3B D⁺ leukemic cell population at "quite normal concentrations", and the reported ability of crude, injected preparations of murine G-CSF to suppress established transplanted myeloid leukemias in mice. Metcalf, *Science*, 229, 16-22 (1985). See also, Sachs, *Scientific American*, 284(1), 40-47 (1986).

To the extent that hpG-CSF may prove to be therapeutically significant and hence need to be available in commercial scale quantities, isolation from cell cultures is unlikely to provide an adequate source of material. It is noteworthy, for example, that restrictions appear to exist against commercial use of Human Tumor Bank cells such as the human bladder carcinoma cell line 5637 (A.T.C.C. HTB9) which have been reported as sources of natural hpCSF isolates in Welte, et al. (1985, *supra*).

The European patent application EP-A-0169566 discloses the isolation of a human CSF from the tumor cell line CHU-1 and its physics chemical characterisation.

The subsequent European patent application EP-A-0215126 which is based on seven different priority documents of which only the first one precedes the first priority date in the present case discloses the further characterisation of said CSF and its production by recombinant DNA technology.

Summary of the Invention

According to the present invention, DNA sequences coding for all or part of hpG-CSF are provided. Such sequences may include : the incorporation of codons "preferred" for expression by selected non-mammalian hosts ; the provision of sites for cleavage by restriction endonuclease enzymes ; and the provision of additional initial, terminal or intermediate DNA sequences which facilitate construction of readily expressed vectors. The present invention also provides DNA sequences coding for microbial expression of polypeptide analogs or derivatives of hpG-CSF which differ from naturally-occurring forms in terms of the identity or location of one or more amino acid residues (i.e., deletion analogs containing less than all of the residues specified for hpG-CSF ; substitution analogs, such as [Ser¹⁷]hpG-CSF, wherein one or more residues specified are replaced by other residues ; and addition analogs wherein one or more amino acid residues is added to a terminal or medial portion of the polypeptide) and which share some or all the properties of naturally-occurring forms.

Novel DNA sequences of the invention include sequences useful in securing expression in procaryotic or eucaryotic host cells of polypeptide products having at least a part of the primary structure and one or more of the biological properties of naturally occurring pluripotent granulocyte colony-stimulating factor. DNA sequences of the invention are specifically seen to comprise : (a) the DNA sequence set forth in Table VII or the complementary strands ; (b) a DNA sequence which hybridizes (under hybridization conditions such as illustrated herein or more stringent conditions) to the DNA sequences in Table VII or to fragments thereof ; and (c) a DNA sequence which, but for the degeneracy of the genetic code, would hybridize to the DNA sequences defined in part (a) or (b) and which sequences code for a polypeptide having the same amino acid sequence. Specifically comprehended in part (b) are genomic DNA sequences encoding allelic variant forms of hpG-CSF and/or encoding other mammalian species of pluripotent granulocyte colony-stimulating factor. Specifically comprehended by part (c) are manufactured DNA sequences encoding hpG-CSF, fragments of hpG-CSF and analogs of hpG-CSF which DNA sequences may incorporate codons facilitating translation of messenger RNA in microbial hosts. Such manufactured sequences may readily be constructed according to the methods of Alton, et al., PCT published application WO 83/04053.

Also comprehended by the present invention is that class of polypeptides coded for by portions of the DNA complement to the top strand human cDNA or genomic DNA sequences of Tables VII or VIII herein, i.e., "complementary inverted proteins" as described by Tramontano, et al., *Nucleic Acids Res.*, 12, 5049-5059 (1984).

The present invention provides purified and isolated polypeptide products having part or all of the primary structure (i.e., continuous sequence of amino acid residues) and one or more of the biological properties (e.g., immunological properties and *in vitro* biological activity) and physical properties (e.g., molecular weight) of naturally-occurring hpG-CSF including allelic variants thereof. These polypeptides are also characterized by being the product of procaryotic or eucaryotic host expression (e.g., by bacterial, yeast, higher plant, insect and mam-

malian cells in culture) of exogenous DNA sequences obtained by genomic or cDNA cloning or by gene synthesis. The products of typical yeast (e.g., *Saccharomyces cerevisiae*) or procaryote [e.g., *Escherichia coli* (*E. coli*)] host cells are free of association with any mammalian proteins. The products of microbial expression in vertebrate (e.g., non-human mammalian and avian) cells are free of association with any human proteins. Depending upon the host employed, polypeptides of the invention may be glycosylated with mammalian or other eucaryotic carbohydrates or may be non-glycosylated. Polypeptides of the invention may also include an initial methionine amino acid residue (at position -1).

Also comprehended by the invention are pharmaceutical compositions comprising effective amounts of polypeptide products of the invention together with suitable diluents, adjuvants and/or carriers useful in hpG-CSF therapy.

Polypeptide products of the invention may be "labelled" by association with a detectable marker substance (e.g., radiolabelled with ^{125}I) to provide reagents useful in detection and quantification of human hpG-CSF in solid tissue and fluid samples such as blood or urine. DNA products of the invention may also be labelled with detectable markers (such as radiolabels and non-isotopic labels such as biotin) and employed in DNA hybridization processes to locate the human hpG-CSF gene position and/or the position of any related gene family in a chromosomal map. They may also be used for identifying human hpG-CSF gene disorders at the DNA level and used as gene markers for identifying neighboring genes and their disorders.

Polypeptide products of the present invention may be useful, alone or in combination with other hematopoietic factors or drugs in the treatment of hematopoietic disorders, such as aplastic anemia. They may also be useful in the treatment of hematopoietic deficits arising from chemotherapy or from radiation therapy. The success of bone marrow transplantation, for example, may be enhanced by application of hpG-CSF. Wound healing burn treatment and the treatment of bacterial inflammation may also benefit from the application of hpG-CSF. In addition, hpG-CSF may also be useful in the treatment of leukemia based upon a reported ability to differentiate leukemic cells. Welte, et al., *Proc. Natl. Acad. Sci. (USA)*, 82, 1526-1530 (1985) and Sachs, *supra*.

Numerous aspects and advantages of the invention will be apparent to those skilled in the art upon consideration of the following detailed description which provides illustrations of the practice of the invention in its presently preferred embodiments.

Brief Description of the Drawings

The Figure is a partial restriction endonuclease map of the hpG-CSF gene accompanied by arrows depicting the sequencing strategy used to obtain the genomic sequence.

Detailed Description

According to the present invention, DNA sequences encoding part or all of the polypeptide sequence of hpG-CSF have been isolated and characterized.

The following examples are presented by way of illustration of the invention and are specifically directed to procedures carried out prior to identification of hpG-CSF cDNA and genomic clones, to procedures resulting in such identification, and to the sequencing, development of expression systems based on cDNA, genomic and manufactured genes and verification of expression hpG-CSF and analog products in such systems.

More particularly, Example 1 is directed to amino acid sequencing of hpG-CSF. Example 2 is directed to the preparation of a cDNA library for colony hybridization screening. Example 3 relates to construction of hybridization probes. Example 4 relates to hybridization screening, identification of positive clones, DNA sequencing of a positive cDNA clone and the generation of polypeptide primary structure (amino acid sequence) information. Example 5 is directed to the identification and sequencing of a genomic clone encoding hpG-CSF. Example 6 is directed to the construction of a manufactured gene encoding hpG-CSF wherein *E. coli* preference codons are employed.

Example 7 is directed to procedures for construction of an *E. coli* transformation vector incorporating hpG-CSF-encoding DNA, the use of the vector in procaryotic expression of hpG-CSF, and to analysis of properties of recombinant products of the invention. Example 8 is directed to procedures for generating analogs of hpG-CSF wherein cysteine residues are replaced by another suitable amino acid residue by means of mutagenesis performed on DNA encoding hpG-CSF. Example 9 is directed to procedures for the construction of a vector incorporating hpG-CSF analog-encoding DNA derived from a positive cDNA clone, the use of the vector for transfection of COS-1 cells, and the cultured growth of the transfected cells. Example 10 relates to physical and biological properties or recombinant polypeptide products of the invention.

Example 1**(A) Sequencing of Material Provided By Literature Methods**

5 A sample (3-4 µg, 85-90% pure of SDS, silver stain-PAGE) of hpG-CSF was obtained from Sloan Kettering Institute, New York, New York, as isolated and purified according to Welte, et al., Proc. Natl. Acad. Sci. (USA), 82, 1526-1530 (1985).

10 The N-terminal amino acid sequence of this sample of hpG-CSF was determined in a Run #1 by micro-sequence analysis using an AB407A gas phase sequencer (Applied Biosystems, Foster City, California) to provide the sequence information set out in Table I below. In Tables I-IV single letter codes are employed, "X" designates a residue which was not unambiguously determined and residues in parentheses were only alternatively or tentatively assigned.

TABLE I

15

1	5	10	15
K-P-L-G-P-A-S-K-L-K-Q-(G,V,S)-G-L-X-X-X			

20 A high background was present in every cycle of the run for which results are reported in Table I, indicating that the sample had many contaminating components, probably in the form of chemical residues from purification. The sequence was retained only for reference use.

25 In Run #2, a second sample (5-6 µg, ~95% pure) was obtained from Sloan Kettering as for Run #1 and a sequencing procedure was performed as for Run #1. This sample was from the same lot of material employed to generate Fig. 4 of Welte, et al., Proc. Natl. Acad. Sci.(USA), 82, 1526-1530 (1985). The results are given in Table II.

TABLE II

30

1	5	10	15	20
35 T-P-L-G-P-A-S-(S)-L-P-Q-($\frac{S}{X}$)-M-($\frac{L}{M}$)-X-K-(R)-X-X-(R)-(L)-X-				

40 Although more residues were identified, Run #2 did not provide a sufficiently long, unambiguous sequence from which a reasonable number of probes could be constructed to search for hpG-CSF DNA. It was calculated that at least 1536 probes would have been required to attempt isolation of cDNA based on the sequence of Table II. Again, contamination of the sample was believed to be the problem.

45 Accordingly, a third sample (3-5 µg, ~40% pure) was obtained from Sloan Kettering as above. This preparation was electroblotted after separation by SDS-PAGE in an attempt at further purification. Sequence analysis of this sample yielded no data.

(B) Sequencing of Materials Provided by Revised Methods

50 In order to obtain a sufficient amount of pure material to perform suitably definitive amino acid sequence analysis, cells of a bladder carcinoma cell line 5637 (subclone 1A6) as produced at Sloan-Kettering were obtained from Dr. E. Platzer. Cells were initially cultured in Iscove's medium (GIBCO, Grand Island, New York) in flasks to confluence. When confluent, the cultures were trypsinized and seeded into roller bottles (1-1/2 flasks/bottle) each containing 25 ml of preconditioned Iscove's medium under 5% CO₂. The cells were grown overnight at 37°C. at 0.3 rpm.

55 Cytodex-1® beads (Pharmacia, Uppsala, Sweden) were washed and sterilized using the following procedures. Eight grams of beads were introduced into a bottle and 400 ml of PBS was added. Beads were suspended by swirling gently for 3 hours. After allowing the beads to settle, the PBS was drawn off, the beads were rinsed in PBS and fresh PBS was added. The beads were autoclaved for 15 minutes. Prior to use, the beads were washed in Iscove's medium plus 10% fetal calf serum (FCS) before adding fresh medium plus 10% FCS to

obtain treated beads.

After removing all but 30 ml of the medium from each roller bottle, 30 ml of fresh medium plus 10% FCS and 40 ml of treated beads were added to the bottles. The bottles were gassed with 5% CO₂ and all bubbles were removed by suction. The bottles were placed in roller racks at 3 rpm for 1/2 hour before reducing the speed to 0.3 rpm. After 3 hours, an additional flask was trypsinized and added to each roller bottle containing beads.

At 40% to 50% of confluence the roller bottle cultures were washed with 50 ml PBS and rolled for 10 min. before removing the PBS. The cells were cultured for 48 hours in medium A [Iscoe's medium containing 0.2% FCS, 10⁻⁸M hydrocortisone, 2mM glutamine, 100 units/ml penicillin, and 100 µg/ml streptomycin]. Next, the culture supernatant was harvested by centrifugation at 3000 rpm for 15 min., and stored at -70°C. The cultures were refed with medium A containing 10% FCS and were cultured for 48 hours. After discarding the medium, the cells were washed with PBS as above and cultured for 48 hours in medium A. The supernatant was again harvested and treated as previously described.

Approximately 30 liters of medium conditioned by 1A6 cells were concentrated to about 2 liters on a Millipore Pellicon unit equipped with 2 cassettes having 10,000 M.W. cutoffs at a filtrate rate of about 200 ml/min. and at a retentate rate of about 1000 ml/min. The concentrate was diafiltered with about 10 liters of 50 mM Tris (pH 7.8) using the same apparatus and same flow rates. The diafiltered concentrate was loaded at 40 ml/min. onto a 1 liter DE cellulose column equilibrated in 50 mM Tris (pH 7.8). After loading, the column was washed at the same rate with 1 liter of 50 mM Tris (pH 7.8) and then with 2 liters of 50 mM Tris (pH 7.8) with 50 mM NaCl. The column was then sequentially eluted with six 1 liter solutions of 50 mM Tris (pH 7.5) containing the following concentrations of NaCl: 75 mM; 100 mM; 125 mM; 150 mM; 200 mM; and 300 mM. Fractions (50 ml) were collected, and active fractions were pooled and concentrated to 65 ml on an Amicon ultrafiltration stirred cell unit equipped with a YM5 membrane. This concentrate was loaded onto a 2 liter AcA54 gel filtration column equilibrated in PBS. The column was run at 80 ml/hr. and 10 ml fractions were collected. Active fractions were pooled and loaded directly onto a C4 high performance liquid chromatography (HPLC) column.

Samples, ranging in volume from 125 ml to 850 ml and containing 1-8 mg of protein, about 10% of which was hpG-CSF, were loaded onto the column at a flow rate ranging from 1 ml to 4 ml per minute. After loading and an initial washing with 0.1 M ammonium acetate (pH 6.0-7.0) in 80% 2-propanol at a flow rate of 1 ml/min. One milliliter fractions were collected and monitored for proteins at 220 nm, 260 nm and 280 nm.

As a result of purification, fractions containing hpG-CSF were clearly separated (as fractions 72 and 73 of 80) from other protein-containing fractions. HpG-CSF was isolated (150-300 µg) at a purity of about 85 ± 5% and at a yield of about 50%. From this purified material 9 µg was used in Run #4, an amino acid sequence analysis wherein the protein sample was applied to a TFA-activated glass fiber disc without polybrene. Sequence analysis was carried out with an AB 470A sequencer according to the methods of Hewick, et al., *J. Biol. Chem.*, 256, 7990-7997 (1981) and Lai, *Anal. Chim. Acta*, 163, 243-248 (1984). The results of Run #4 appear in Table III.

TABLE III

1	5	10
Thr - Pro - Leu - Gly - Pro - Ala - Ser - Ser - Leu - Pro -		
	15	20
Gln - Ser - Phe - Leu - Leu - Lys - (Lys) - Leu - (Glu) - Glu -		
	25	30
Val - Arg - Lys - Ile - (Gln) - Gly - Val - Gly - Ala - Ala -		
Leu - X - X -		

In Run #4, beyond 31 cycles (corresponding to residue 31 in Table III) no further significant sequence information was obtained. In order to obtain a longer unambiguous sequence, in a Run #5, 14 µg of hpG-CSF purified from conditioned medium were reduced with 10 µl of β-mercaptoethanol for one hour at 45°C, then thoroughly

dried under a vacuum. The protein residue was then redissolved in 5% formic acid before being applied to a polybrenized glass fiber disc. Sequence analysis was carried out as for Run #4 above. The results of Run #5 are given in Table IV.

TABLE IV

5

1	5	10
Thr - Pro - Leu - Gly - Pro - Ala - Ser - Ser - Leu - Pro -		
15	20	
Gln - Ser - Phe - Leu - Leu - Lys - Cys - Leu - Glu - Gln -		
25	30	
Val - Arg - Lys - Ile - Gln - Gly - Asp - Gly - Ala - Ala -		
35	40	
Leu - Gln - Phe - Lys - Leu - Gly - Ala - Thr - Tyr - Lys -		
45		
Val - Phe - Ser - Thr - (Arg) - (Phe) - (Met) -X-		

25 The amino acid sequence give in Table IV was sufficiently long (44 residues) and unambiguous to construct probes for obtaining hpG-CSF cDNA as described infra.

Example 2

30 Among standard procedures for isolating cDNA sequences of interest is the preparation of plasmid-borne cDNA "libraries" derived from reverse transcription of mRNA abundant in donor cells selected on the basis of their expression of a target gene. Where substantial portions of the amino acid sequence of a polypeptide are known, labelled, single-stranded DNA probe sequences duplicating a sequence putatively present in the "target" cDNA may be employed in DNA/DNA hybridization procedures carried out on cloned copies of the cDNA which have been denatured to single stranded form. Weissman, et al., U.S. Patent No. 4,394,443 ; Wallace, et al., Nucleic Acids Res., 6, 3543-3557 (1979), and Reyes, et al., Proc. Natl. Acad. Sci. (USA), 79, 3270-3274 (1982), and Jaye, et al., Nucleic Acids Res., 11, 2325-2335 (1983). See also, U.S. Patent No. 4,358,535 to Falkow, et al., relating to DNA/DNA hybridization procedures in effecting diagnosis ; and Davis, et al., "A Manual for Genetic Engineering, Advanced Bacterial Genetics", Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1980) at pp. 55-58 and 174-176, relating to colony and plaque hybridization techniques.

40 Total RNA was extracted from approximately 1 gram of cells from a bladder carcinoma cell line 5637 (1A6) using a guanidinium thiocyanate procedure for quantitative isolation of intact RNA. [Chirgwin, et al., Biochemistry, 18, 5294-5299 (1979)].

45 The sterile aqueous RNA solution contained total RNA from the 1A6 cells. To obtain only the messenger RNA from the total RNA solution, the solution was passed through a column containing oligodeoxythymidylate [oligo (dT)] (Collaborative Research, Inc., Waltham, Massachusetts. Poly-Adenylated (poly-A⁺) tails characteristic of messenger RNA adhere to the column while ribosomal RNA is eluted. As a result of this procedure, approximately 90 µg of poly-adenylated messenger RNA (poly-A⁺ mRNA) were isolated. The isolated poly-A⁺ messenger RNA was pre-treated with methylmercury hydroxide (Alpha Ventron, Danvers, Massachusetts) at a final concentration of 4 mM for 5 minutes at room temperature prior to use in a cDNA reaction. The methylmercury hydroxide treatment denatured interactions of messenger RNA, both with itself and with contaminating molecules that inhibit translation. Payvar, et al., J Biol.Chem., 258, 7636-7642 (1979).

50 According to the Okayama procedure [Okayama, et al., Molecular & Cellular Biology, 2, 161-170 (1982)], a cDNA bank was prepared using mRNA obtained from 1A6 cells. The cDNAs were then transformed by incubation into a host microorganism E.coli K-12 strain HB101 for amplification.

55

Example 3

Hybridization probes designed on the basis of the hpG-CSF amino terminal sequence of Table IV consisted of a set of 24 oligonucleotides each being 23 bases in length and containing three inosine residues. The probe oligonucleotides were manufactured according to the procedure of Caruthers, et al., Genetic Engineering, 4, 1-18 (1982) and labeled with γ -³²P ATP by kinasing with polynucleotide kinase. The probe oligonucleotides, corresponding to the messenger RNA for residues 23-30 of the sequence of Table IV, are illustrated in Table V.

TABLE V

hpG-CSF Probes

5' GC IGC ICC $\begin{smallmatrix} A \\ G \end{smallmatrix}$ TC ICC $\begin{smallmatrix} T \\ C \end{smallmatrix}$ TG $\begin{smallmatrix} G \\ A \\ T \end{smallmatrix}$ AT $\begin{smallmatrix} T \\ C \end{smallmatrix}$ TT^{3'}

The assignment of neutrality to I's was based on the published work of Takahashi, et al., Proc. Natl. Acad. Sci. (USA), 82, 1931-1935 (1985) and Ohtsuka, et al., J. Biol. Chem., 260, 2605-2608 (1985). However, inosine may have a destabilizing effect if base paired with a G or T. In Takahashi, et al., inosines may appear to have a neutral effect because they average out as a group to near neutrality (e.g., three having paired favorably with C and two not favorable to pairing with T).

To test the effect of having I's base pair with G's, control experiments were designed using an N-myc gene sequence and clone. The sequences picked from the N-myc gene had the same overall G and C content at the first two positions of each codon as was prescribed by the hpG-CSF probes. Thus, the N-myc test probes were of the same length, contained I's in the same relative positions and had potentially the same average Tm (62-66°C., not accounting for the 3 or 4 inosine residues included) as the hpG-CSF probes.

Two sets of N-myc test probes were constructed according to the procedure of Caruthers, et al., supra. Set I, as illustrated in Table VI included: 1, a 23 mer with perfect match; 2, in which three third position C's were replaced with I's generating the worst possible case for adding I's; and 3, in which four third position C's were replaced with I's. The second set of test probes was designed to represent a more random distribution of inosine base pairs, that might give an overall neutral base pairing effect. Set II, as illustrated in Table VI, included: 4, containing two I's that will base pair with C's and one with a G; and 5, identical to 4 with the addition of one more I: G base pair.

TABLE VI

N-myc Test Probes

1. 5' CAC AAC TAT GCC GCC CCC TCC CC^{3'}

2. 5' CAC AAC TAT GCI GCC CCI TCI CC^{3'}

3. 5' CAI AAC TAT GCI GCC CCI TCI CC^{3'}

4. 5' AAC GAG CTG TGI GGC AGI CCI GC^{3'}

5. 5' AAI GAG CTG TGI GGC AGI CCI GC^{3'}

Five replica filters containing N-myc DNA sequences and chicken growth hormone DNA sequences (as a

negative control) were baked in a vacuum oven for 2 hours at 80°C. prior to hybridization. All filters were hybridized as described in Example 4 for the hpG-CSF probes except the period of hybridization was only 6 hours. Filters were washed three times at room temperature then once at 45°C., 10 minutes each. The filters were monitored with a Geiger counter.

The filter representing N-myc probe 3 gave a very weak signal relative to the other four probed filters and was not washed any further. After a 10 minute 50°C. wash, the Geiger counter gave the following percent signal with probe one being normalized to 100% : Probe 2, 20% ; Probe 3 (45°C.), 2% ; Probe 4, 92% ; and Probe 5, 75%. After a 55°C. wash, the percentages were : Probe 2, 16% ; Probe 4, 100% ; and Probe 5, 80%. A final wash at 60°C. yielded the following percentages : Probe 2, 1.6% ; Probe 4, 90% ; and Probe 5, 70%.

Thus, in the presence of three I's, as in probes 2 and 4, up to a 60-fold difference in signal is observed as the theoretical T_m (I's not included in the calculation) is approached [based upon a worst case I base pairing (Probe 2) and a relatively neutral I base pairing case (Probe 4)].

The standardization information gained by the N-myc test hybridizations was utilized in washing and monitoring of the hpG-CSF hybridization as indicated below, to gauge the degree of confidence with which the results of less than stringent washing might be accepted.

Example 4

According to the procedure of Hanahan, et al., *J. Mol. Biol.*, **166**, 557-580 (1983), bacteria containing recombinants with cDNA inserts as prepared in Example 2 were spread on 24 nitrocellulose filters (Millipore, Bedford, Massachusetts) laid on agar plates. The plates were then incubated to establish approximately 150,000 colonies which were replica plated to 24 other nitrocellulose filters. The replicas were incubated until distinct colonies appeared. The bacteria on the filters were lysed on sheets of Whatman 3 MM paper barely saturated with sodium hydroxide (0.5 M) for 10 minutes, then blotted with Tris (1 M) for 2 minutes, followed by blotting with Tris (0.5 M) containing NaCl (1.5 M) for 10 minutes. When the filters were nearly dry, they were baked for 2 hours at 80°C. in a vacuum oven prior to nucleic acid hybridization. [Wahl, et al., *Proc. Natl. Acad. Sci. (USA)*, **76**, 3683-3687 (1979)] ; and Maniatis, et al., *Cell*, **81**, 163-182 (1976).

The filters were prehybridized for 2 hours at 65°C. in 750 ml of 10X Denhardt's, 0.2% SDS and 6X SSC. The filters were rinsed in 6X SSC, then placed four in a bag and hybridized for 14 hours in 6X SSC and 10X Denhardt's. There was approximately 15 ml of solution per bag containing 50×10^6 cpm of ³²p-labeled probe (oligonucleotides).

After hybridization, the filters were washed three times in 6X SSC (1 liter/wash) at room temperature for 10 minutes each. The filters were then washed two times at 45°C. for 15 minutes each, once at 50° for 15 minutes and once at 55°C. for 15 minutes using 1 liter volumes of 6X SSC. The filters were autoradiographed for 2 hours at -70°C. using an intensifying screen and Kodak XAR-2 film. On this autoradiograph, there were 40-50 positive signals detected including 5 very intense signals.

The areas containing the strongest five signals and an additional five positives were scraped from the master plates and replated for a secondary screening using the same probe mixture under the same conditions. The wash procedure differed in that the high temperature washes consisted of two at 55°C. for 15 minutes each and then one at 60°C. for 15 minutes. Based on the N-myc probe study of Example 3, the final wash temperature in the second screening was raised because the aggregate melting temperature for the 24 23-mers was 60-68°C., similar to that of the N-myc probes. Just after the second 55°C. wash, the filters were left damp and an autoradiograph was made. Comparison of this autoradiograph with a second autoradiograph taken for a similar period of time after a final wash at 60°C. showed that only two of the 10 clones being tested did not suffer a substantial loss in signal in rising from 55-60°C. These two clones were later shown to be of nearly identical lengths and restriction endoclease patterns. One clone designated Ppo2, was selected for sequencing.

Sequencing of the recombinant hpG-CSF cDNA clone, Ppo2, obtained by the above procedure was accomplished by the dideoxy method of Sanger, et al., *Proc. Natl. Acad. Sci. (USA)* **74**, 5463-5467 (1977). The single-stranded DNA phage M-13 was used as a cloning vector for supplying single-stranded DNA templates from the double-stranded cDNA clones. The Sanger, et al., method revealed the sequence as set forth in Table VII accompanied by its amino acid translation and a complementary strand in the polypeptide coding region.

Hindu

5' - AGCTTGACTCAGGGGGGGGGGGGGGGGGGGGGGGG[ⁿNNNNN]

-12	Leu	Trp	His	Ser	Ala	Leu	Trp	Thr	Val	Gln	Glu	Ala	Thr	Pro	Leu	Gly	Pro
	CTG	TGG	CAC	AGT	GCA	CTC	TGG	ACA	GTG	CAG	GAA	GCC	ACC	CCC	CTG	GGC	CCT
	GAC	ACC	GTG	TCA	CGT	GAG	ACC	TGT	CAC	GTC	CTT	CGG	TGG	GGG	GAC	CCG	GGA
					<u>HgiAI</u>										<u>ApaI</u>		
		10										20					
	Ala	Ser	Leu	Ser	Phe	Leu	Leu	Lys	Cys	Leu	Glu	Gln	Val	Arg	Lys	Ile	Gln
	GCC	AGC	TCC	AGC	TTC	CTG	CTC	AAG	TGC	TTA	GAG	CAA	GTG	AGG	AAG	ATC	CAG
	CGG	TCC	AGG	TCC	AAG	GAC	GAG	TTC	ACG	AAT	CTC	GTT	CAC	TCC	TTC	TAG	GTC
		30										40					
	Gly	Asp	Gly	Gln	Glu	Lys	Leu	Cys	Ala	Thr	Tyr	Lys	Leu	Cys	His	Pro	Glu
	GGC	GAT	GGC	CAG	GAG	AAG	CTG	TGT	GCC	ACC	TAC	AAG	CTG	TGC	CAC	CCC	GAG
	CCG	CTA	CCG	GAG	CTC	TTC	GAC	ACA	CGG	TGG	ATG	TTC	GAC	ACG	GTG	GGG	CTC
		50										60					
	Glu	Leu	Val	Gly	His	Leu	Gly	Ile	Pro	Trp	Ala	Pro	Leu	Ser	Ser	Cys	Pro
	GAG	CTG	GTG	GGA	CAC	CTG	GGC	ATC	CCC	TGG	GCT	CCC	CTG	AGC	AGC	TGC	CCC
	CTC	GAC	CAC	GAG	GTG	GAC	CCG	TAG	GGG	ACC	CGA	GGG	GAC	TCG	TCG	ACG	GGG
		70										80					
	Ser	Gln	Ala	Leu	Ala	Gly	Cys	Leu	Ser	Gln	Leu	His	Gly	Leu	Phe	Leu	Tyr
	AGC	CAG	GCC	CTG	GCA	GGC	TGC	TTG	AGC	CAA	CTC	CAT	AGC	CTT	TTC	CTC	TAC
	TCG	GTC	CGG	GAC	CGT	CCG	ACG	AAC	TCG	GTT	GAG	GTA	TCG	GAA	AAG	GAG	ATG
		90										100					
	Gln	Gly	Leu	Ala	Leu	Glu	Gly	Ile	Ser	Pro	Glu	Leu	Gly	Pro	Thr	Leu	Thr
	CAG	GGG	CTC	GCC	CTG	GAA	GGG	ATC	TCC	CCC	GAG	TTG	GGT	CCC	ACC	TTG	ACC
	GTC	CCC	GAG	GTC	GAC	CTT	CCC	TAG	AGG	GGG	CTC	AAC	CCA	GGG	TGG	AAC	TGT

5
10
15
20
25
30
35
40
45
50
55

110	Leu	Gln	Leu	Asp	Val	Ala	Asp	Phe	Ala	Thr	Thr	Ile	Trp	Gln	Gln	Met	Glu	Glu	Leu	Gly
	CTG	CAG	CTG	GAC	GTC	GCC	GAC	TTT	GCC	ACC	ACC	ATC	TGG	CAG	CAG	ATG	GAA	GAA	CTG	GGA
	GAC	GTC	GAC	CTG	CAG	CGG	CTG	AAA	CGG	TGG	TGG	TAG	ACC	GTC	GTG	TAC	CTT	CTT	GAC	CCT
130	Met	Ala	Pro	Ala	Leu	Gln	Pro	Thr	Gln	Gly	Ala	Met	Pro	Ala	Phe	Ala	Ser	Ala	Phe	Gln
	ATG	GCC	CCT	GCC	CTG	CAG	CAG	CCC	ACC	ACC	GGT	GCC	ATG	CCG	TTC	GCC	TCT	GCT	TTC	CAG
	TAC	CGG	GGA	CGG	GAC	GTC	GTC	GGG	TGG	CTC	CCA	CGG	TAC	GGC	AAG	CGG	AGA	CGA	AAG	GTC
150	Arg	Arg	Ala	Gly	Gly	Val	Leu	Val	Ala	Ser	His	Leu	Gln	Ser	Phe	Leu	Glu	Val	Ser	Tyr
	CGC	CGG	GCA	GGA	GGG	GTC	CTG	GTT	GCC	TCC	CAT	CTG	CAG	AGC	TTC	CTG	GAG	GTG	TCG	TAC
	GCG	GCC	CGT	CCT	CCC	CAG	GAC	CAA	CGG	AGG	GTA	GAC	GTC	TCG	AAG	GAC	CTC	CAC	AGC	ATG
170	Arg	Val	Leu	Arg	His	Leu	Ala	Gln	Pro	OP										
	CGC	GTT	CTA	CGC	CAC	CTT	GCC	CAG	CCC	TGA	GCC	AAG	CCC	TCC	CCA	TCC	CAT	GTA	TTT	ATC
	CGC	CAA	GAT	GCG	GTG	GAA	CGG	GTC	GGG	ACT										
	TCT	ATT	TAA	TAT	TTA	TGT	CTA	TTT	AAG	CCT	CAT	ATT	TAA	AGA	CAG	GGA	AGA	GCA	GAA	CGG
	AGC	CCC	AGG	CCT	CTG	TGT	CCT	TCC	CTG	CAT	TTC	TGA	GTT	TCA	TTC	TCC	TGC	CTG	TAG	CAG
	TGA	GAA	AAA	GCT	CCT	GTC	CTC	CCA	TCC	CCT	GGA	CTG	GGA	GGT	AGA	TAG	GTA	AAT	ACC	AAG
	TAT	TTA	TTA	CTA	TGA	CTG	CTC	CCC	AGC	CCT	GGC	TCT	GCA	ATG	GGC	ACT	GGG	ATG	AGC	CGC
	TGT	GAG	CCC	CTG	GTC	CTG	AGG	GTC	CCC	ACC	TGG	GAC	CCT	TGA	GAG	TAT	CAG	GTC	TCC	CAC

TABLE VII (cont'd.)

GTG GGA GAC AAG AAA TCC CTG TTT AAT ATT TAA ACA GCA GTG TTC CCC ATC TGG GTC CTT	5
GCA CCC CTC ACT CTG GCC TCA GCC GAC TGC ACA GCG GCC CCT GCA TCC CCT TGG CTG TGA	10
GGC CCC TGG ACA AGC AGA GGT GGC CAG AGC TGG GAG GCA TGG CCC TGG GGT CCC ACG AAT	15
TTG CTG GGG AAT CTC GTT TTT CTT CTT AAG ACT TTT GGG ACA TGG TTT GAC TCC CGA ACA	20
TCA CCG ACG TGT CTC CTG TTT TTC TGG GTG GCC TCG GGA CAC CTG CCC TGC CCC CAC GAG	25
GGT CAG GAC TGT GAC TCT TTT TAG GGC CAG GCA GGT GCC TGG ACA TTT GCC TTG CTG GAC	30
GGG GAC TGG GGA TGT GGG AGG GAG CAG ACA GGA GGA ATC ATG TCA GGC CTG TGT GTG AAA	35
GGA AGC TCC ACT GTC ACC CTC CAC CTC TTC ACC CCC CAC TCA CCA GTG TCC CCT CCA CTG	40
TCA CAT TGT AAC TGA ACT TCA GGA TAA <u>AGT GTT TGC CTC CA</u>	45

StuI

[f150-200 base poly A plus 25-30 bases plasmid DNA preceding a PvuII restriction site]-3'

The following characteristics of the sequence of Table VII are of note. At the 5' end of the sequence there are shown bases corresponding to those of the poly G cDNA linker. There then occur about five bases (designated as "N") whose sequence could not readily be determined unambiguously by the Sanger, et al. method due to the preceding multiple G's. The sequence thereafter reveals a series of 12 codons encoding a portion of a putative leader sequence for the polypeptide. Based on correspondence to the amino terminal sequence of natural isolates of hpCSF described in Example 1, the initial threonine residue of the putative "mature" form of hpG-CSF is indicated by +1. Mature hpG-CSF is thereafter revealed to include 174 residues as indicated. Following the "stop" codon (the OP codon, TGA) are approximately 856 bases of an untranslated 3' sequence and multiple A's of the poly A "tail". Unique HgiAI, and Apal restriction endonuclease recognition sites, as well as two StuI sites (discussed infra with respect to construction of procaryotic and eucaryotic expression systems) are also designated in Table VII. Owing to the lack of asparagine residues in the polypeptide, there are no apparent sites for N-glycosylation. The underscored 6 bases near the end of the 3' untranslated sequence represent a potential polyadenylation site.

It is noteworthy that each of two additional cDNA clones identified by the hybridization procedures described above from among a total of 450,000 clones failed to include DNA encoding the entire leader sequence from the transcription initiation site onward. Indeed, all three hpG-CSF clones terminated in the 5' region at exactly the same site, indicating that secondary structure of the mRNA transcribed severely hinders cDNA formation beyond this site. As a practical matter, therefore, cDNA expression screening such as described in Okayama, et al., Mol. and Cell. Biol., 3, 280-289 (1983) and as actually employed to isolate GM-CSF in Wong, et al., Science, 228, 810-814 (1985) could not have readily applied to isolation of hpCSF DNA because such isolation systems ordinarily rely upon the presence of a full length cDNA transcript in the clones assayed.

The above sequence is not readily susceptible for securing direct expression of hpG-CSF in a microbial host. To secure such expression, the hpG-CSF coding region should be provided with an initial ATG codon and the sequence should be inserted in a transformation vector at a site under control of a suitable promoter/regulator DNA sequence.

Example 5

In this example, cDNA encoding hpG-CSF as isolated in the previous example was used to screen a genomic clone. A phage lambda human fetal liver genomic library [prepared according to the procedure of Lawn, et al. Cell, 15, 1157-1174 (1978) and obtained from T. Maniatis] was screened using a nick translated probe consisting of two hpG-CSF cDNA fragments isolated by digestion with HgiAI and StuI (HgiAI to StuI, 649 b.p. ; StuI to StuI, 639 b.p.). A total of approximately 500,000 phage were plated on 12 (15 cm) petri dishes and plaque lifted and hybridized to probe using the Benton/Davison procedure [Benton, et al., Science, 196, 180 (1977)]. A total of 12 positive clones were observed. Three clones (1-3) yielding the strongest signals upon autoradiography in a secondary screening were grown in 1 liter cultures and mapped by restriction enzyme digestion and Southern blotting using a radiolabeled 24-mer oligonucleotide (kinased with γ -³²P ATP) 5' CTGCACTGTCCAGAGTGCACTGTG3'. The mapping results showed that isolates 1 and 3 were identical and 2 contained 2000 additional bases 5' to the hpG-CSF gene. Therefore, clone 2 was used for further characterization. DNA from clone 2 was digested with R1 to release an 8500 bp hpG-CSF containing fragment which was subsequently subcloned into pBR322 and further mapped by restriction endonuclease digests, Southern Blotting, M13 subcloning and sequencing. The sequence obtained is as set out in Table VIII.

TABLE VIII

GGGGACAGGCTTGAGATCCCAAGGAGAGGGGCAAGGACACTGCCCCCGCAAGTCTGCCAGAGCAGAGAGGGAGACCCCGACTCAGCTGCCACTTCCC	100
CACAGGCTCGTGCCGCTTCCAGGCGTCTATCAGGGGCTCAGCCCTTTGTTTCAGCTGTTCTGTTCAAAACACTCTGGGGCCATTTCAGGCCTGGGTGGGGCAGC	200
GGGAGGAAGGGAGTTTGAGGGGGGCAAGGCGACGTCAAGGAGGATCAGAGATTCCACAATTTACAAAACCTTTTCGCAACAGCTTTTGTGTTCCAAACCCC	300
CCTGCATTGTCTTGGACACCAAAATTTGTCATAAATCCTGGGAAGTTATTACTAAGCCTTAGTCGTGGCCCCCAGGTAATTTCTCCAGGCCCTCCATGGGGT	400
-30 MetAlaGlyProAlaThrGlnSerProM	
TATGTATAAGGGCCCCCTAGAGCTGGGGCCCCCAAAACAGCCCCGAGCCTGTCAGCCCCAGCCCCCAGACCCCATGGCTGGACCTGCCACCCAGAGAGCCCCA	500
-20 -18 etLysLeuMetA	
TGAAGCTGATGGGTGAGTGTCTTGGCCCCCAGGATGGGAGAGCCCCCTGCCCTGGCATGGAGGGAGGCTGGTGTGACAGAGGGGCTGGGGATCCCCGTTCT	600
-16 laLeuGlnLeuL	
GGGAATGGGCATTAAAGGCACCCAGTGTCCCCGAGAGGGCCCTCAGGTGGTAGGGAACAGCATGTCTCTGAGCCCGCTGTCTCCCCAGCCCTGCAGCTGC	700

TABLE VIII (cont'd.)

	-10	-1 +1	10	20	
					5
					10
					15
					20
					25
					30
					35
					40
					45
					50
					55
					800
					900
					1000
					1100
					1200
					1300

-10

-1 +1

10

20

euLeuTrpHisSerAlaLeuTrpThrValGlnGluAlaThrProLeuGlyProAlaSerSerLeuProGlnSerPheLeuLeuLysCysLeuGluGlnVa
TGCTGTGGCACAGTGCACACTCTGGACAGTGCAGGAAGCCACCCCCCTGGGCCCTGCCAGCTCCCTGCCAGAGCTTCTGCTCAAGTGTCTTAGAGCAAGT

800

IArgLysIleGlnGlyAspGlyAlaAlaLeuGlnGluLysLeu
GAGGANGATCCAGGGCGATGGCGCAGCGCTCCAGGAGAAAGCTGGTGAGTGAGGTGGTGAGAGGGCTGTGGAGGGAAGCCCGGTGGGAGAGCTAAGGGG

-6

900

GATGGAACTGCAGGGCCCAACATCCTCTGTGAAGGGACATGGGAGAATATTAGGAGCAGTGGAGCTGGGGAAGGCTTGGGGAGGAGGACCT

1000

TGGTGGGACAGTGCTCGGGAGGGCTGGCTGGGATGGGAGTGGAGGCATCACATTTCAGGAGAAAGGGCAAGGGCCCCCTGTGAGATCAGAGAGTGGGGTG

1100

CAGGGCAGAGAGGAACCTGAACAGCCTGGCAGGACATGGAGGGAGGGGAAGACCAGAGAGTCTGGGAGGACCCCGGAAGGAGCGCGGACCCCGCCACGGC

1200

36 40

50

CysAlaThrTyrLysLeuCysHisProGluGluLeuValLeuLeuGlyHisSerLeuGlyIleProTrpA
GAGTCTCACTCAGCATCCTTCCATCCCCAGTGTGCCACCTACAAAGCTGTGCCACCCCGAGGAGCTGGTGCTGCTCGGACACTCTCTGGGCATCCCTCTGGG

1300

TABLE VIII (cont'd.)

60	70	71	
1aProLeuSerSerCysProSerGlnAlaLeuGlnLeu			1400
CTCCCCGTAGCAGCTGCCCCAGCCAGGCCCTGCAGCTGGTGAGTGTCAAGGAAGGATAAGGCTAATGAGGAGGGGAAGGAGGAGGAAACACCCCATGGG			
	72		
	AlaGlyCysLeuSerGln		1500
CTCCCCCATGTCTCCAGGTTCCAAAGCTGGGGGGCTGACGTATCTCAGGCAGCACCCCCCTAACTCTTCCGCTCTGTCTCACAGGCAGGCTGCTTGAGCCAA			
80	90	100	110
LeuHisSerGlyLeuPheLeuTyrglnGlyLeuLeuGlnAlaLeuGluGlyIleSerProGluLeuGlyProThrLeuAspThrLeuGlnLeuAspVala			1600
CTCCATAGCGGCCCTTTTCCCTCTACCAGGGGCTCCCTGCAGGCCCTTGAAGGGATCTCCCCCGAGTTGGGTCCCACCTTGGACACACTGCAGCTGGACCGTCG			
	120		1700
1aAspPheAlaThrThrIleTrpGlnGln			
CGGACTTGGCCACCACCATCTGGCAGCAGGTGAGCCTTGTTGGCAGGGTGGCCAAAGGTCGTGCTGGCATTTCTGGGCACCACAGCCGGGGCCTGTGTATGG			
	121		1800
GGCCCTGTCCATGCTGCAGCCCCCAGCATTTTCTCATTTTGTATAAACGCCCCACTCAGAAGGGGCCCAACCACTGATCACAGCTTTCCCCCACAGATGGGAAG			
	MetGluG		

TABLE VIII (cont'd.)

130	140	150	
1uLeuGlyMetAlaProAlaLeuGlnProThrGlnGlyAlaMetProAlaPheAlaSerAlaPheGlnArgAlaGlyGlyValLeuValAlaSerHi			1900
AACTGGGAATGGCCCTGCTGCAGCCACCAGGGTGCCATGCCGGCCCTTCGCTGCTTCCAGCGCGGGCAGGAGGGTCTGCTGGTGCCTCCCA			
160	170	174	
sLeuGlnSerPheLeuGluValSerTyrArgValLeuArgHisLeuAlaGlnProOP			2000
TCTGCAGAGCTTCCTGGAGGTGTCGTACCGCGTCTACGCCACCTTGCCAGCCCTTGAGCCCAAGCCCTCCCCATGCCCATGTATTATCTCTATTAAATAT			
TTATGTCATTTAAGCCTCATATTTAAAGACAGGGAAGCAGAACGAGGCCCTCTGTGTCTTCCCTGCATTCTGAGTTTCATTCTCTCCTGCC			2100
TGTAGCAGTGAGAAAGCTCCTGTCTCCTCCATCCCTGGACTGGGAGGTAGATAGGTAAATACCAAGTATTATTAATACTATGACTGCTCCCCAGCCCTGG			2200
CTCTGCAATGGGCACTGGGATGAGCCGCTGTGAGCCCTGTCTGAGGGTCCCCACCTGGGACCCCTTGAGAGTATCAGGTCTCCCACGTGGGAGACACAG			2300
AAATCCCTGTTTAATATTTAAACAGCAGTGTTCCCCCATCTGGGTCTTGCACCCCTCACTCTGGCCTCAGCCGACTGCACAGGGCCCTTGCATCCCCCTT			2400
GGCTGTGAGGCCCCCTGGACAAGCAGAGGTGGCCAGAGCTGGGAGGCATGGCCCTGGGGTCCACAGAAATTGTGCTGGGAAATCTCGTTTTCCTTAAGAC			2500
TTTTGGACATGGTTTGACTCCCGAACATCACCGAGGTGTCTCTGTTTTTCTGGGTGGCCTCGGGACACCTGCCCTGCCCCACGAGGGTCAGGACTGT			2600

TABLE VIII (cont'd.)

5
10
15
20
25
30
35
40
45
50
55

2700	GACTCTTTTAGGGCCAGGCAGGTGCCTGGACATTGGCTTGCTGGATGGGGACTGGGGATGTGGGAGGGAGCAGACAGGAGGAATCATGTTCAGGCCCTGT
2800	GTGTGAAGGAAGCTCCACTGTCAACCTCCACCTCTTACCCCCCACTCACAGTGTCCCCCTCCA CTGTCA CATTTGTA ACTGAACTTCAGGATAATAAAG
2900	TGTTTGCCCTCCAGTCACGTCCCTTCCTCTCTCTGAGTCCAGCTGGTGCCTTGCC CAGGGGCTGGGGAGGTGGCTGAAGGCTGGGAGAGGCCAGAGGGAGGT
3000	CGGGAGGAGGTCTGGGGAGGAGGTCCAGGGAGGAGGAGGAAAGTTCTCAAGTTCCCTCTGACATTCATTCGGTTAGCACATATTTATCTGAGCACCTACT
3070	CTGTGCAGACGCTGGGCTAAGTGTCTGGGGACACAGCAGGGGAACAAGGCAGACATGGAATCTGCACCTCGAG

A restriction endonuclease map (approximately 3.4 Kb) of genomic DNA containing the hpG-CSF gene is detailed in Figure 1. The restriction endonucleases shown in Figure 1 are : NcoI, N ; PstI, P ; BamHI, B ; Apal, A ; XhoI, X ; and KpnI, K. The arrows below the map depict the sequencing strategy used to obtain the genomic sequence. The boxed regions are those found in the cDNA clone with the dashed open ended box representing sequence not present in the cDNA clone, but identified by probing mRNA blots. The identification of coding sequences proposed for exon one was carried out by Northern blot analysis. A 24 mer oligonucleotide probe, 5' CAGCAGCTGCAGGGCCATCAGCTT³, spanning the predicted splice junctures for exons 1 and 2 was hybridized to hpG-CSF mRNA in a Northern blot format. The resulting blot shows an mRNA the same size (~1650 bp) as that seen with an exon 2 oligonucleotide probe. This data combined with the ability to direct expression of hpG-CSF from the pSVGM-PpoI vector (Example 9) using the Met initiation codon depicted in Table VIII, defines the coding sequences contained in exon 1. Exons 2-5 are defined by the coding sequences obtained in the cDNA clone (Ppo2) of the hpG-CSF gene (Table VII).

Example 6

This example relates to preparation of a manufactured gene encoding hpG-CSF and including E. coli preference codons.

Briefly stated, the protocol employed was generally as set out in the disclosure of co-owned Alton, et al., PCT Publication No. W083/04053, which is incorporated by reference herein. The genes were designed for initial assembly of component oligonucleotides into multiple duplexes which, in turn, were assembled into three discrete sections. These sections were designed for ready amplification and, upon removal from the amplification system, could be assembled sequentially or through a multiple fragment ligation in a suitable expression vector.

The construction of Sections I, II and III is illustrated in Table IX though XIV. In the construction of Section I, as illustrated in Tables IX and X, oligonucleotides 1-14 were assembled into 7 duplexes (1 and 8) ; 2 and 9 ; 3 and 10 ; 4 and 11 ; 5 and 12 ; 6 and 13 ; and 7 and 14). The 7 duplexes were then ligated to form Section I as shown in Table X. It may also be noted in Table X that Section I includes an upstream XbaI sticky end and a downstream BamHI sticky end useful for ligation to amplification and expression vectors and for ligation to Section II.

TABLE IX

EChpG-CSFDNA SECTION I

CTAGAAAAACCAAGGAGGTAATAAA	1
TAATGACTCCATTAGGTCCTGCTTCTTCT	2
CTGCCGCAAAGCTTTCTGCTGAAATGTCTGG	3
AACAGGTTCGTAAAATCCAGGGTGACGGT	4
GCTGCACTGCAAGAAAACTGTGCGCTA	5
CTTACAACTGTGCCATCCGGAAGAGC	6
TGGTACTGCTGGGTCAATCTCTTGG	7
CATTATTTATTACCTCCTTGGTTTTTT	8
GCAGAGAAGAAGCAGGACCTAATGGAGT	9
TGTTCCAGACATTTACGAGAAAGCTTTGCG	10
CAGCACCGTCACCCTGGATTTTACGAACC	11
TAAGTAGCGCACAGTTTTTCTTGCAGTG	12
ACCAGCTCTTCCGGATGGCACAGTTTG	13
GATCCCAAGAGAATGACCCAGCAGT	14

TABLE X

EChpG-CSFDNA SECTION I

10 1 20 30 40 2 50 60
 CTAGAAAA ACCAAGGAGG TAATAAATAA TGACTCCATT AGGTCCTGGT TCTTCTCTGC
 TTTT TGGTTCCTCC ATTATTATT ACTGAGGTAA TCCAGGACGA AGAAGAGACG

8XbaI

70 3 80 90 100 4 110 120
 CGCAAGCTT TCTGCTGAAA TGTCTGGAAC AGGTCGTAA AATCCAGGT GACGGTGTG
 GCGTTTCGAA AGACGACTTT ACAGACCTTG TCCAAGCATT TTAGGTCCCA CTGCCACGAC

1011

130 5 140 150 160 6 170 180
 CACTGCAAGA AAAACTGTGC GCTACTTACA AACTGTGCCA TCCGGAAGAG CTGGTACTGC
 GTGACGTTCT TTTTGACACG CGATGAATGT TTGACACGGT AGGCTTCTG GACCATGACG

1213

7 190 100
 TGGGTCATTC TCTTGG
 ACCCAGTAAG AGAACCCCTAG

14BamHI

As illustrated in Tables XI and XII, in the construction of Section II, oligonucleotides 15-30 were assembled into 8 duplexes (15 and 23 ; 16 and 24 ; 17 and 25 ; 18 and 26 ; 19 and 27 ; 20 and 28 ; 21 and 29 ; and 22 and 30). These 8 duplexes were then ligated to form Section II, as shown in Table XII. As further shown in Table XII, Section II has an upstream BamHI sticky end and a downstream EcoRI sticky end useful for ligation to an amplification vector and for ligation to Section I. Near its downstream end, Section II also includes a downstream SstI site useful in the eventual ligation Sections II and III.

TABLE XI

EChpG-CSFDNA SECTION II

GATCCCGTGGGCTCCGCTGTCTTCT	15
TGTCCATCTCAAGCTCTTCAGCTGGC	16
TGGTTGTCTGTCTCAACTGCATTCTGGT	17
CTGTTCCCTGTATCAGGGTCTTCTG	18
CAAGCTCTGGAAGGTATCTCTCCGGA	19
ACTGGGTCCGACTCTGGACACTCTGCA	20
GCTAGATGTAGCTGACTTTGCTACTACT	21
ATTTGGCAACAGATGGAAGAGCTCAAAG	22
GACAAGAAGACAGCGGAGCCACGG	23
ACCAGCCAGCTGAAGAGCTTGAGATG	24
ACAGACCAGAATGCAGTTGAGACAGACA	25
CTTGCAGAAGACCCTGATACAGGA	26
CAGTTCCGGAGAGATACCTTCCAGAG	27
TAGCTGCAGAGTGTCCAGAGTCGGACC	28
AAATAGTAGTAGCAAAGTCAGCTACATC	29
AATTCTTTGAGCTCTTCCATCTGTTGCC	30

TABLE XII

EChpG-CSFDNA SECTION II

10 15 20 30 40 50 60
 GATCCCGTG GGCTCCGCTG TCTTCTTGTC CATCTCAAGC TCTTCAGCTG GCTGGTTGTC
 GGCAC CCGAGGCGAC AGAAGAACAG GTAGAGTTCC AGAGTTCGAC CGACCAACAG

BamHI

70 17 80 90 110 120
 TGTCTCAACT GCATTCTGGT CTGTTCCCTGT ATCAGGGTCT TCTGCAAGCT CTGGAAGGTA
 ACAGAGTTGA CGTAAGACCA GACAAGGACA TAGTCCCAGA AGAGTTCCGA GACCTTCCAT

130 140 150 160 170 180
 TCTCTCCGGA ACTGGGTCGG ACTCTGGACA CTCTGCAGCT AGATGTAGCT GACTTTTGCTA
 AGAGAGGCCT TGACCCAGGC TGAGACCTGT GAGACGTCGA TCTACATCGA CTGAAACGAT

190 200 210
 CTACTATTG GCAACAGATG GAAGAGCTCA AAG
 GATGATAAAC CGTTGTCCTAC CTCTCCGAGT TTCTTAA

SstI EcoRI

Finally, Section III was constructed as shown in Tables XIII and XIV. For this construction, oligonucleotides 31-42 were assembled into 6 duplexes (31 and 37 ; 32 and 38 ; 33 and 39 ; 34 and 40 ; 35 and 41 ; and 36 and 42). The 6 duplexes were then ligated to form Section III as depicted in Table XIV. As also shown in Table XIV, Section III includes an upstream BamHI sticky end and a downstream EcoRI sticky end useful for ligating into an amplification vector, and at least in the case of the EcoRI end, into an expression vector. In addition, Section II has an upstream SstI site useful in the eventual ligation of Sections II and III.

TABLE XIII

EChpG-CSFDNA SECTION III

GATCCAAAGAGCTCGGTATGGCACCAG	31
CTCTGCAACCGACTCAAGGTGCTATGCCG	32
GCATTGCTTCTGCATTCCAGCGTCGTGC	33
AGGAGGTGTACTGGTTGCTTCTCATCTG	34
CAATCTTTCCTGGAAGTATCTTACCGTGT	35
TCTGCGTCATCTGGCTCAGCCGTAATAG	36
AGAGCTGGTGCCATACCGAGCTCTTTG	37
ATGCCGGCATAGCACCTTGAGTCGGTTGC	38
TCCTGCACGACGCTGGAATGCAGAAGCGA	39
ATTGCAGATGAGAAGCAACCAGTACACC	40
CAGAACACGGTAAGATACTTCCAGGAAAG	41
AATTCTATTACGGCTGAGCCAGATGACG	42

TABLE XIV

EChpG-CSFDNA SECTION III

10	31	20	30	40	32	50	60
GATCCAAAG	AGCTCGGTAT	GGCACCAGCT	CTGCAACCGA	CTCAAGGTGC	TATGCCGGCA		
GTTC	TCGAGCCATA	CCGTGGTCCA	GACGTGGCT	GAGTTCCACG	ATACGGCCGT		
	<u>37</u>				<u>38</u>		
<u>BamHI</u>	<u>SstI</u>						
70	33	80	90	100	34	110	120
TTCGCTTCG	CATTCCAGCG	TCGTGCAGGA	GGTGACTGG	TTGCTTCTCA	TCTGCAATCT		
AAGCGAAGAC	GTAAGGTCCG	AGCACGTCCT	CCACATGACC	AACGAAGAGT	AGACGTTAGA		
	<u>39</u>				<u>40</u>		
35	130	140	150	36	160	170	
TTCCCTGGGAG	TATCTTACCG	TGTTCTGCGT	CATCTGGCTC	AGCCGTATA G			
AAGGACCTTC	ATAGAATGGC	ACAAGACCGA	GTAGACCGAG	TCGGCATTAT	CTTAA		
	<u>41</u>			<u>42</u>			
							<u>ECORI</u>

The XbaI to BamHI fragment formed by Section I is ligated into an M 13 mp 11 phage vector opened with XbaI and BamHI. The vector is then reopened by digestion with BamHI and EcoRI, followed by ligation with the BamHI to EcoRI fragment formed by Section II. At this stage, Sections I and II have been joined in proper orientation. Next, another M 13 mp 11 vector is opened by BamHI to EcoRI digestion and then ligated with the

5

BamHI to EcoRI fragment formed by Section III.

The vector containing Sections I and II is digested with XbaI and SstI. Likewise, the vector containing Section III is digested with SstI and EcoRI. Both of the smaller of the two fragments resulting from each digestion are ligated into a plasmid pCFM1156 which is previously opened with XbaI and EcoRI. The product of this reaction is an expression plasmid containing a continuous DNA sequence, as shown in Table XV, encoding the entire hpG-CSF polypeptide with an amino terminal methionine codon (ATG) for E. coli translation initiation.

10

15

20

25

30

35

40

45

50

55

[illegible]

5

10

15

20

25

30

35

40

45

50

55

TABLE XV (cont'd.)

150	Val	Leu	Val	Ala	Ser	His	Leu	Gln	Ser	Phe	Leu	Glu	Val	Ser	Tyr	Arg	Val	Leu	Arg
Gly	GTA	CTG	GTT	GCT	TCT	CAT	CTG	CAA	TCT	TTC	CTG	GAA	GTA	TCT	TAC	CGT	GTT	CTG	CGT
160																			
170	His	Leu	Ala	Gln	Pro														
CAT	CTG	GCT	CAG	CCG	TAA	TAG	AAT	T											
174																			

Although any suitable vector may be employed to express this DNA, the expression plasmid pCFM1156 may readily be constructed from a plasmid pCFM836, the construction of which is described in published European Patent Application No. 136,490. pCFM836 is first cut with NdeI and then blunt-ended with Poll such that both existing NdeI sites are destroyed. Next, the vector is digested with ClaI and SacII to remove an existing polylinker before ligation to a substitute polylinker as illustrated in Table XVI. This substitute polylinker may be constructed according to the procedure of Alton, et al., supra. Control of expression in the expression pCFM1156 plasmid is by means of a lambda P_L promoter, which itself may be under the control of a C₁₈₅₇ repressor gene (such as is provided in E. coli strain K12ΔHtrp).

10

15

20

25

30

35

40

45

50

55

TABLE XVI

1 ATCGATTGATTCTAGAGGAGGGAATAACATATGGTTAACCGGTGGGAATTCGGTACCAT
TAGCTAAACTAAGATCTTCCTCCTTATTGTATACCAATTGCGCAACCTTAAGCCCATGGTA
1 ClaI, 12 XbaI, 29 NdeI, 35 HincII, HpaI, 39 MluI, 47 EcoRII,
53 HgiCI KpnI, 57 NcoI StyI,
61 GGAAGCTTACTCGAGGATCCGGGGATAAATAAGTAACGATCC
CCTTCGAATGAGCTCCTAGCGGCTATTATTATTCATTGCTAGG
63 HindIII, 70 Aval XhoI, 75 BamHI Xho2, 79 Sac2,

Example 7

This example relates to *E. coli* expression of an hpG-CSF polypeptide by means of a DNA sequence encoding (Met¹) hpCSF. The sequence employed was partially synthetic and partially cDNA-derived. The synthetic sequence employed *E. coli* preference codons.

Plasmid Ppo2, containing the hpG-CSF gene shown in Table VII, was digested with HgiAI and StuI providing an approximately 645 base pair fragment including the gene for mature hpCSF (as shown in Table VII) with seven of the leader sequence residue codons at the 5' end and about 100 base pairs of the 3' non-coding region. HgiAI digestion leaves a 5', 4-base sticky end identical to that of PstI, and StuI leaves a blunt end. This allows for ready insertion of the fragment into M13 mp8 (Rf) cut with PstI and with the blunt-end-forming restriction enzyme, HincII. Upon amplification in M13, the hpG-CSF DNA was excised by digestion with Apal and BamHI which cut, respectively, at the Apal site spanning the codons for residues + 3 to + 5 of hpCSF and at a BamHI site "downstream" of the HincII site in the M13 mp8 restriction polylinker. In order to allow for *E. coli* expression of the hpG-CSF polypeptide, a synthetic fragment was prepared as set out in Table XVII below.

TABLE XVII

```

5' - C TAG AAA AAA CCA AGG AGG TAA TAA ATA
      3' - TTT TTT GGT TCC TCC ATT ATT TAT
      XbaI

-1  +1
Met Thr Pro Leu
ATG ACA CCT CTG GGC C - 5'
TAC TGT GGA GAC -3'
                        Apal

```

As may be determined from analysis of Table XVII, the linker includes an Apal sticky end, codons specifying the initial three residues of the amino terminal of hpG-CSF ("restoring" the Thr¹, Pro², Leu³-specifying codons deleted upon Apal digestion of the M13 DNA described above and employing codons preferentially expressed in *E. coli*), a translation initiating ATG, a sequence of 24 base pairs providing a ribosome binding site, and an XbaI sticky end.

The expression vector employed for *E. coli* expression was that described as pCFM536 in European Patent Application No. 136,490, by Morris, published April 10, 1985. (See also, A.T.C.C. 39934, *E. coli* JM103 harboring pCFM536). Briefly, plasmid pCFM536 was digested with XbaI and BamHI. The hpG-CSF fragment (Apal/BamHI) and linker (XbaI/Apal) described above were then ligated therein to form a plasmid designated p536Ppo2.

Plasmid p536Ppo2 was transformed into a phage resistant variant of the *E. coli* AM7 strain which has previously been transformed with plasmid pMWI (A.T.C.C. No. 39933) harboring a *Cl⁸⁵⁷* gene. Transformation was verified on the basis of the antibiotic (amp) resistance marker gene carried on the pCFM536 progenitor plasmid. Cultures of cells in LB broth (ampicillin 50 µg 1ml) were maintained at 28°C. and upon growth of cells in culture to A600 = 0.5, hpCSF expression was induced by raising the culture temperature to 42°C. for 3 hours. The final O.D. of the culture was A600 = 1.2.

The level of expression of hpG-CSF by the transformed cells was estimated on a SDS-poly acrylamide gel stained with coomassie blue dye to be 3-5% of total cellular protein.

Cells were harvested by centrifugation at 3500 g for 10 minutes in a JS-4.2 rotor. Cells at 25% (w/v) in water were broken by passing 3 times through a French Pressure Cell at 10,000 p.s.i. The broken cell suspension was centrifuged at 10,000 g for 15 minutes in a JA-20 rotor. The pellet was resuspended in water and solubilized at about 5 mg/ml total protein in 1% lauric acid, 50 mM Tris, pH 8.7. The solubilized pellet material was centrifuged at 15,000 g for 10 minutes and to the supernatant CuSO₄ was added to 20 mM. After 1 hour, this sample was loaded onto a C4 HPLC column for purification according to the procedures of example 1 (B) with adjustments made for volume and concentration.

A second purification procedure was developed to yield larger quantities of hpG-CSF formulated in a nonorganic-containing buffer. This material is suitable for *in vivo* studies. One hundred and fifty grams of cell paste was resuspended in about 600 ml of 1 mM DTT and passed 4 times through a Manton Gualin Homogenizer at about 7000 PSI. The broken cell suspension was centrifuged at 10,000 g for 30 minutes and the pellet was

resuspended in 400 ml of 1% deoxycholate (DOC), 5 mM EDTA, 5 mM DTT, and 50 mM Tris, pH 9. This suspension was mixed at room temperature for 30 minutes and centrifuged at 10,000 g for 30 minutes. The pellet was resuspended in about 400 ml of water and centrifuged at 10,000 g for 30 minutes. The pellet was solubilized in 100 ml of 2% Sarkosyl and 50 mM at pH 8. CuSO₄ was added to 20 µM and the mixture was stirred 16 hours at room temperature, and then centrifuged at 20,000 g for 30 minutes. To the supernatant was added 300 ml acetone. This mixture was put on ice for 20 minutes and then centrifuged at 5000 g for 30 minutes. The pellet was dissolved in 250 ml of 6 M guanidine and 40 mM sodium acetate at pH 4, and put over a 1,200 ml G-25 column equilibrated and run in 20 mM sodium acetate at pH 5.4. The hpG-CSF peak (about 400 ml) was pooled and put on a 15 ml CM-cellulose column equilibrated in 20 mM sodium acetate at pH 5.4. After loading, the column was washed with 60 ml of 20 mM sodium acetate at pH 5.4 and with 25 mM sodium chloride, and then the column was eluted with 200 ml of 20 mM sodium acetate at pH 5.4 and with 37 mM sodium chloride. 150 ml of this eluent was concentrated to 10 ml and applied to a 300 ml G-75 column equilibrated and run in 20 mM sodium acetate and 100 mM sodium chloride at pH 5.4. The peak fractions comprising 35 ml were pooled and filter sterilized. The final concentration of hpG-CSF was 1.5 mg/ml, is greater than 95% pure as determined by analysis on a gel, and contained less than 0.5 ng of pyrogen per 0.5 mg of hpG-CSF. The pyrogen level was determined using a Limulus Amebocyte Lysate (LAL) test kit (M. A. Bioproducts, Walkersville, Maryland).

Example 8

This example relates to the use of recombinant methods to generate analogs of hpG-CSF wherein cysteine residues present at positions 17, 36, 42, 64 and 74 were individually replaced by a suitable amino acid residue. Site directed mutagenesis procedures according to Souza, et al., published PCT Application No. WO85/00817, published February 28, 1985, were carried out on [Met⁻¹] encoding DNA of plasmid p536Ppo2, described *infra*, using synthetic oligonucleotides ranging in size from 20 to 23 bases as set out in Table XVIII below. Oligonucleotide No. 1 allowed for formation of a gene encoding [Ser¹⁷] hpG-CSF; oligonucleotide No. 2 allowed for formation of [Ser³⁶] hpG-CSF, and so on.

TABLE XVIII

<u>Oligonucleotide</u>	<u>Sequence</u>
1.	5'-CTG CTC AAG TCC TTA GAG CAA GT-3'
2.	5'-GAG AAG CTG TCT GCC ACC TACA-3'
3.	5'-TAC AAG CTG TCC CAC CCC GAG-3'
4.	5'-TGA GCA GCT CCC CCA GCC AG-3'
5.	5'-CTG GCA GGC TCC TTG AGC CAA-3'

The Cys to Ser site directed mutagenesis restrictions were carried out using M13 mp10 containing an XbaI-BamHI hpG-CSF fragment isolated from p536Ppo2 as a template. DNA from each M13 mp10 clone containing a Cys-Ser substitution was treated with XbaI and BamHI. The resulting fragment was cloned into expression vector pCFM746 and expression products were isolated as in Example 7.

The plasmid pCFM746 may be constructed by cleaving a plasmid pCFM736 (the construction of which from deposited and publically available materials is described in Morris, published PCT Application No. WO85/00829, published February 28, 1985) with ClaI and BamHI to remove an existing polylinker and by substituting the following polylinker.

TABLE XIXClaI

5' CGATTGATTCTAGAATTCGTTAACGGTACCATGGAA

3' TAAACTAAGATCTTAAGCAATTGCCATGGTACCTT

GCTTACTCGAGGATCCGCGGATAAATAAGTAAC^{3'}CGAATGAGCTCCTAGGCGCCTATTTATTCATTGCTAG^{5'}Sau3a

In a purification procedure for Cys to Ser analogs according to the present invention, about 10-15 g of cell paste was resuspended in 40 ml of 1 mM DTT and passed 3 times through a French Pressure Cell at 10,000 psi. The broken cell suspension was centrifuged at 1,000 g for 30 minutes. The pellet was resuspended in 1% DOC, 5 mM EDTA, 5 mM DTT, 50 mM Tris, pH 9 and allowed to mix 30 minutes at room temperature. The mixture was centrifuged at 10,000 g for 30 minutes, resuspended in 40 ml H₂O, and recentrifuged as 10,000 g for 30 minutes. The pellet was dissolved in 10 ml of 2% Sarkosyl, 50 mM DTT, 50 mM Tris, pH 8. After mixing for 1 hour, the mixture was clarified by centrifugation at 20,000 g for 30 minutes, and then applied to a 300 ml G-75 column equilibrated and run in 1% Sarkosyl, 50 mM Tris, pH 8. Fractions containing the analog were pooled and allowed to air oxidize by standing with exposure to air for at least one day. Final concentrations ranged from 0.5-5 mg/ml.

Example 9

In this example, a mammalian cell expression system was devised to ascertain whether an active polypeptide product of hpG-CSF DNA could be expressed in and secreted by mammalian cells (COS-1, A.T.C.C. CRL-1650). This system was designed to provide for secretion of a polypeptide analog of hpGCSF via expression and secretory processing of a partially synthetic, partially cDNA-derived construction encoding [Ala¹] hpG-CSF preceded by a leader polypeptide having the sequence of residues attributed to human GM-CSF in Wong, et al., Science 228, 810-815 (1985) and Lee, et al., Proc. Natl. Acad. Sci. (USA), 82, 4360-4364 (1985).

The expression vector employed for preliminary studies of expression of polypeptide products of the invention was a "shuttle" vector incorporating both pBR322 and SV40 DNA which had been designed to allow for autonomous replication in both *E. coli* and mammalian cells, with mammalian cell expression of inserted exogenous DNA under control of a viral promoter/regulator DNA sequence. This vector, designated pSVDM-19, harbored in *E. coli* HB101, was deposited August 23, 1985, with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland, and received the accession No. A.T.C.C. 53241.

The specific manipulations involved in the expression vector construction were as follows. A leader-encoding DNA sequence was synthesized as set out in Table XX below.

Finally, the genomic sequence whose isolation is described in Example 5 was employed to form an expression vector for mammalian cell expression of hpG-CSF. More specifically, pSVDM-19 was digested with KpnI and HindIII and the large fragment used in a fourway ligation with a synthetic linker with HindIII and NcoI sticky ends, as shown in Table XXI. An NcoI-BamHI fragment containing exon 1 isolated from pBR322 (8500 hpG-CSF), a genomic subclone, and a BamHI-KpnI fragment containing exons 2-5 isolated from the plasmid

pBR322 (8500 hpG-CSF genomic subclone). The resulting mammalian expression vector, pSV/ghG-CSF produced 1 to 2.5 µg/ml of hpG-CSF from transformed COS cells.

TABLE XXI

HindIII
 5' AGCTTCCAACAC
 AGGTTGTGGTAC 5'
NcoI

Example 10

This example relates to physical and biological properties of recombinant polypeptide products of the invention.

1. Molecular Weight

Recombinant hpG-CSF products of *E. coli* expression as in Example 7 had an apparent molecular weight of 18.8 kD when determined in reducing SDS-PAGE (as would be predicted from the deduced amino acid analysis of Table VII), whereas natural isolates purified as described in Example 1 had an apparent molecular weight of 19.6 kD. The presence of N-glycans associated with the natural isolates could effectively be ruled out on the basis of the lack of asparagine residues in the primary sequence of hpG-CSF in Table VII and therefore a procedure was devised to determine if O-glycans were responsible for molecular weight differences between natural isolates and the non-glycosylated recombinant products. Approximately 5 µg of the natural isolate material was treated with neuraminidase (Calbiochem, LaJolla, California), a 0.5 µg sample was removed, and the remaining material was incubated with 4 mU O-Glycanase (endo-x-n-acetylgalactoseaminidase, Genzyme, Boston, Massachusetts) at 37°C. Aliquots were removed after 1/2, 2 and 4 hours of incubation. These samples were subjected to SDS-PAGE side by side with the *E. coli* derived recombinant material. After neuraminidase treatment, the apparent molecular weight of the isolate shifted from 19.6 kD to 19.2 kD, suggestive of removal of a sialic acid residue. After 2 hours of treatment with O-glycanase, the molecular weight shifted to 18.8 kD-identical to the apparent molecular weight of the *E. coli* derived material. The sensitivity of the carbohydrate structure to neuraminidase and O-glycanase suggests the following structure for the carbohydrate component: N-acetylneuraminic acid-α(2-6)(galactose β (1-3) N-acetylgalactoseamine-R, wherein R is serine or threonine.

2. ³H-Thymidine Uptake

Proliferation induction of human bone marrow cells was assayed on the basis of increased incorporation of ³H-thymidine. Human bone marrow from healthy donors was subjected to a density cut with Ficoll-Hypaque (1.077 g/ml, Pharmacia) and low density cells were suspended in Iscove's medium (GIBCO) containing 10% fetal bovine serum and glutamine pen-strep. Subsequently, 2 × 10⁴ human bone marrow cells were incubated with either control medium or the recombinant *E. coli* material of Example 7 in 96 flat bottom well plates at 37°C in 5% CO₂ in air for 2 days. The samples were assayed in duplicate and the concentration varied over a 10,000 fold range. Cultures were then pulsed for 4 hours with 0.5 µCi/well of ³H-Thymidine (New England Nuclear, Boston, Massachusetts). ³H-Thymidine uptake was measured as described in Ventua, et al., *Blood*, 61, 781 (1983). In this assay human hpG-CSF isolates can induce ³H-Thymidine incorporation into human bone marrow cells at levels approximately 4-10 times higher than control supernatants. The *E. coli*-derived hpG-CSF material of Example 6 had similar properties.

A second human bone marrow cell proliferation study was carried out using culture medium of transfected COS-1 cells as prepared in Example 9 and yielded similar results, indicating that encoded polypeptide products were indeed secreted into culture medium as active materials.

3. WEHI-3B D⁺ Differentiation Induction

Capacity of recombinant, *E. coli*-derived materials to induce differentiation of the murine myelomonocytic leukemic cell line WEHI-3B D⁺ was assayed in semi-solid agar medium as described in Metcalf, *Int. J. Cancer*,

25, 225 (1980). The recombinant hpG-CSF product and media controls were incubated with ~60 WEHI-3B D⁺ cells/well at 37°C. in 5% CO₂ in air for 7 days. The samples were incubated in 24 flat bottom well plates and the concentration varied over a 2000-fold range. Colonies were classified as undifferentiated, partially differentiated or wholly differentiated and colony cell counts were counted microscopically. The E. coli recombinant material was found to induce differentiation.

4. CFU-GM, BFU-E and CFU-GEMM Assays

Natural isolates of pluripotent human G-CSF (hpG-CSF) and the recombinant pluripotent human G-CSF (rhpG-CSF) were found to cause human bone marrow cells to proliferate and differentiate. These activities were measured in CFU-GM [Broxmeyer, et al., Exp. Hematol., 5, 87, (1971)] BFU-E and CFU-GEMM assays [Lu, et al., Blood, 61, 250 (1983)] using low density, non-adherent bone marrow cells from healthy human volunteers. A comparison of CFU-GM, BFU-E and CFU-GEMM biological activities using either 500 units of hpG-CSF or rhpG-CSF are shown in Table XXII below.

All the colony assays were performed with low density non-adherent bone marrow cells. Human bone marrow cells were subject to a density cut with Ficoll-Hypaque (density, 1.077 g/cm³ ; Pharmacia). The low density cells were then resuspended in Iscove's modified Dulbecco's medium containing fetal calf serum and placed for adherence on Falcon tissue culture dishes (No. 3003, Becton Dickenson, Cockeysville, MD.) for 1-1/2 hours at 37°C.

TABLE XXII

	<u>CFU-GM</u>	<u>BFU-E</u>	<u>CFU-GEMM</u>
Medium	0±0	26±1	0±0
natural			
hpG-CSF	83±5.4	83±6.7	4±0
rhpG-CSF	87±5	81±0.1	6±2

Medium control consisted of Iscove's modified Dulbecco medium plus 10% FCS, 0.2 mM hemin and 1 unit of recombinant erythropoietin.

For the CFU-GM assay target cells were plated at 1×10^5 in 1 ml of 0.3% agar culture medium that included supplemented McCoy's 5A medium and 10% heat inactivated fetal calf serum. Cultures were scored for colonies (greater than 40 cells per aggregate) and morphology assessed on day 7 of culture. The number of colonies is shown as the mean $\pm$ SEM as determined from quadruplicate plates.

For the BFU-E and CFU-GEMM assays, cells (1×10^5) were added to a 1 ml mixture of Iscove's modified Dulbecco medium (Gibco), 0.8% methylcellulose, 30% fetal calf serum 0.05 nM 2-mercaptoethanol, 0.2 mM hemin and 1 unit of recombinant erythropoietin. Dishes were incubated in a humidified atmosphere of 5% CO₂ and 5% O₂. Low oxygen tension was obtained using an oxyreducer from Reming Bioinstruments (Syracuse, N.Y.). Colonies were scored after 14 days of incubation. The number of colonies is shown as the mean $\pm$ SEM, as determined from duplicate plates.

Colonies formed in the CFU-GM assay were all found to be chloracetate esterase positive and non-specific esterase (alpha-naphthyl acetate esterase) negative, consistent with the colonies being granulocyte in type. Both natural hpG-CSF and rhpG-CSF were found to have a specific activity of approximately 1×10^8 U/mg pure protein, when assayed by serial dilution in a CFU-GM assay. The BFU-E and CFU-GEMM data in Table XXII are representative of three separate experiments and similar to the data reported previously for natural hpG-CSF. It is important to note that the rhpG-CSF is extremely pure and free of other potential mammalian growth factors by virtue of its production in E. coli. Thus rhpG-CSF is capable of supporting mixed colony formation (CFU-GEMM) and BFU-E when added in the presence of recombinant erythropoietin.

5. Cell Binding Assays

It was previously reported that WEHI-3B(D⁺) cells and human leukemic cells from newly diagnosed

leukemias will bind ^{125}I -labeled murine G-CSF and that this binding can be complete for by addition of unlabeled G-CSF or human CSF- β . The ability of natural hpG-CSF and rhpG-CSF to compete for binding of ^{125}I -hpG-CSF to human and murine leukemic cells was tested. Highly purified natural hpG-CSF (>95% pure ; 1 μg) was iodinated [Tejedor, et al., *Anal.Biochem.*, 127, 143 (1982)] was separated from reactants by gel filtration and ion exchange chromatography. The specific activity of the natural ^{125}I -hpG-CSF was approximately $\mu\text{Ci}/\mu\text{g}$ protein. Murine WEHI-3B(D*) and two human peripheral blood myeloid leukemic cell preparations (ANLL, one classified as M4, the other as M5B) were tested for their ability to bind ^{125}I -hpG-CSF.

The murine and freshly obtained human peripheral blood myeloid leukemic cells were washed three times with PBS/1% BSA. WEHI-3B(D*) cells (5×10^6) or fresh leukemic cells (3×10^6) were incubated in duplicate in PBS/1% BSA (100 μl) in the absence or presence of various concentrations (volume : 10 μl) of unlabeled hpG-CSF, rhpG-CSF or GM-CSF and in the presence of ^{125}I -hpG-CSF (approx. 100,000 cpm or 1 ng) at 0°C. for 90 min. (total volume : 120 μl). Cells were then resuspended and layered over 200 μl ice cold FCS in a 350 μl plastic centrifuge tube and centrifuged (1000 g ; 1 min.). The pellet was collected by cutting off the end of the tube and pellet and supernatant counted separately in a gamma counter (Packard).

Specific binding (cpm) was determined as total binding in the absence of a competitor (mean of duplicates) minus binding (cpm) in the presence of 100-fold excess of unlabeled hpG-CSF (non-specific binding). The non-specific binding was maximally 2503 cpm for WEHI-3B(D*) cells, 1072 cpm for ANLL (M4) cells and 1125 cpm for ANLL (M5B) cells. Experiments one and two were run on separate days using the same preparation of ^{125}I -hpG-CSF and display internal consistency in the percent inhibition noted for 2000 units of hpG-CSF. Data obtained are reported in Table XXIII below.

TABLE XXIII

Competitor	(U/ml)	WEHI-3B(D ⁺) cpm	<u>% Inhib.</u>	ANLL (M4) cpm	<u>% Inhib.</u>	ANLL (M5B) cpm	<u>% Inhib.</u>
<u>Exp. 1</u>							
none	0	6,608	-	1,218	-	122	-
natural hpG-CSF:							
	10,000	685	90				
	2,000	1,692	74	34	97	-376	0
	200	2,031	69				
rhG-CSF:							
	10,000	0	100				
	2,000	1,185	82	202	83	0	0
	200	2,330	65				
<u>Exp. 2</u>							
none	0	2,910	0				
natural hpG-CSF:							
	2,000	628	78				
GM-CSF:							
	2,000	3,311	0				

As shown in Table XXIII, ^{125}I -hpG-CSF demonstrated binding to the WEHI-3B(D⁺) leukemic cells. The binding was inhibited in a dose dependent manner by unlabeled natural hpG-CSF or rhpG-CSF, but not by GM-CSF. In addition, binding of natural hpG-CSF to human myelomonocytic leukemic cells (ANLL, M4) was observed. The binding to these cells is paralleled in response to natural hpG-CSF in liquid cultures by differentiation into mature macrophages as judged by morphology. The absence of binding of natural ^{125}I -hpG-CSF to monocytic leukemic cells from another patient (ANLL, M5B) suggests that certain leukemias may differentially express or lack receptors for hpG-CSF. The ability of rhpG-CSF to compete for the binding of natural ^{125}I -hpG-CSF, similar to natural hpG-CSF, suggests that the receptors recognize both forms equally well.

These studies demonstrating the binding of natural ^{125}I -labeled hpG-CSF to leukemic cells are paralleled in culture by the ability of natural hpG-CSF to induce granulocytic and monocytic differentiation of light density bone marrow cells obtained from one patient with an acute promyelocytic leukemia (M3) and a second patient with an acute myeloblastic leukemia (M2). Cells from each patient were cultured for four days in medium alone or in the presence of 1×10^5 units of rhpG-CSF. Cells from the M3 control cultures incubated in medium alone were still promyelocyte in type; while cells cultured in the presence of rhpG-CSF showed mature cells of the myeloid type including a metamyelocyte, giant band form and segmented neutrophils and monocyte. The actual differentials for this patient, on 100 cells evaluated for the control, 100% promyelocytes, and for the rhpG-CSF treated cells, 22% blasts plus promyelocytes, 7% myelocytes, 35% metamyelocytes, 20% band forms plus segmented neutrophils, 14% monocytes and 2% macrophages. Of note is the fact that one of the polymorphonuclear granulocytes still contained a prominent auer rod, suggesting that at least this cell represented a differentiated cell belonging to the leukemic clone. Cells from the second patient with a myeloblastic leukemia (M2) were also cultured for four days in the presence of absence of rhpG-CSF. Visual analysis of M2 cells cultured in medium alone revealed large "blast-like" cells, some of which had nucleoli. Some of the M2 cells, when treated with rhpG-CSF, differentiated to mature segmented neutrophils displaying residual auer rods in the center neutrophil suggesting differentiation occurring in a cell belonging to the leukemic clone. The actual differentiation of 100 cells evaluated morphologically revealed that control cells consisted of 100% blasts. The rhpG-CSF treated cells consisted of 43% blasts, 1% myelocytes, 15% metamyelocytes, 28% band forms plus segmented neutrophils, 2% promonocytes and 11% monocytes. The leukemic cells were also examined for differentiation at four other concentrations of rhpG-CSF (5×10^3 , 1×10^4 , 2.5×10^4 and 5×10^4 U/ml, data not shown). Even at the lowest concentration of rhpG-CSF tested (5×10^3 U/ml), there was significant differentiation (cells differentiated beyond myelocytes) of the M3 (50%) and M2 (37%) leukemic cells.

6. Immunoassay

To prepare polyclonal antibodies for immunoassay use the antigen employed was pluripotent G-CSF purified from the human bladder carcinoma cell line 5637 (1A6) as prepared in Example 1 (B). This material was judged to be 85% pure based on silver nitrate staining of polyacrylamide gels. Six week-old Balb/C mice were immunized with multiple-site subcutaneous injections of antigen. The antigen was resuspended in PBS and emulsified with equal volumes of Freund's complete adjuvant. The dose was 5 to 7 μg of antigen per mouse per injection. A booster immunization was administered 18 days later with the same amount of antigen emulsified with an equal volume of Freund's incomplete adjuvant. 4 days later mouse serum was taken to test for the antibody specific to human pluripotent G-CSF.

Dynatech Immulon II Removawell strips in holders (Dynatech Lab., Inc., Alexandria, Virginia) were coated with hpG-CSF 5 $\mu\text{g}/\text{ml}$ in 50 mM carbonate-bicarbonate buffer, pH 9.2. Wells were coated with 0.25 μg in a volume of 50 μl . Antigen coated plates were incubated 2 hours at room temperature and overnight at 4°C. The solution was decanted and the plates were incubated 30 minutes with PBS containing 5% BSA to block the reactive surface. This solution was decanted and the diluted preimmune or test sera were added to the wells and incubated for 2 hours at room temperature. Sera were diluted with PBS, pH 7.0 containing 1% BSA. The serum solution was decanted and plates were washed three times with Wash Solution (KPL, Gaithersburg, Maryland). Approximately 200,000 cpm of iodinated rabbit anti-mouse IgG (NEN, Boston, Massachusetts) in 50 μl PBS, pH 7.0 containing 1% BSA was added to each well. After incubating 1-1/2 hours at room temperature, the solution was decanted and plates were washed 5 times with Wash Solution. Wells were removed from holder and counted in a Beckman 5500 gamma counter. High-titered mouse sera showed greater than 12-fold higher reactivity than the corresponding preimmune sera at a dilution of 1 : 100.

The immunological properties of *E. coli*-derived hpG-CSF were determined by reactivity to high-titered mouse serum specific to mammalian-cell derived hpG-CSF. 0.25 μg of 90% pure *E. coli*-derived protein was coated to Immulon II Removawells in a volume of 50 μl and mouse serum was assayed as described above.

High-titered mouse sera showed a 24-fold higher reactivity to the *E. coli*-derived material than did the corresponding preimmune sera at a dilution of 1 : 100.

7. Serine Analog Bioassays

[Ser¹⁷]hpG-CSF, [Ser³⁶]hpG-CSF, [Ser⁴²]hpG-CSF, [Ser⁶⁴]hpG-CSF, and [Ser⁷⁴]hpG-CSF products prepared according to Example 9 were assayed for hpG-CSF activity in the ³H-thymidine uptake, CFU-GM, and WEHI3B D⁺ assays. In each assay, the [Ser¹⁷] analog had activity comparable to that of recombinant molecules having the native structure. The remaining analogs had on the order of 100-fold lesser activity in the ³H-thymidine uptake assay, 250-fold lesser activity in the CFU-GM assay, and 500-fold lesser activity in the WEHI-3B D⁺ assay. This data is supportive of the proposition that cysteines at positions 36, 42, 64 and 74 may be needed for full biological activity.

8. In vivo Bioassay

Alzet® osmotic pumps (Alzet Corp., Palo Alto, CA ; Model 2001) were connected to indwelling right jugular vein catheters and implanted subcutaneously in seven male Syrian golden hamster. Four of the pumps contained a buffer (20 mM sodium acetate (pH 5.4) and 37 mM sodium chloride) and 1.5 mg/ml *E. coli*-derived hpG-CSF while 3 contained buffer alone. The claimed pumping rate for the osmotic pumps was 1 microliter/hr. for up to seven days. At the third day after implantation of the pumps, the mean granulocyte count of the four treated hamsters was six-fold higher than that of the three (buffer) controls and the increased granulocyte count was reflected in a four-fold increase in total lymphocytes. Erythrocyte count was unchanged by treatment. These results indicate that the recombinant material produces a specific enhancement of production and/or release of granulocytes in a mammal.

The present invention embraces hpG-CSF products such as polypeptide analogs of hpG-CSF and fragments of hpG-CSF. Following the procedures of the above-noted published application by Alton, et al. (WO/83/04053) one may readily design and manufacture genes coding for microbial expression of polypeptides having primary conformations which differ from that herein specified for in terms of the identity or location of one or more residues (e.g., substitutions, terminal and intermediate additions and deletions). Alternately, modifications of cDNA and genomic genes may be readily accomplished by well-known site-directed mutagenesis techniques and employed to generate analogs and derivatives of. Such products would share at least one of the biological properties of hpG-CSF but may differ in others. As examples, projected products of the invention include those which are foreshortened by e.g., deletions ; or those which are more stable to hydrolysis (and, therefore, may have more pronounced or longer lasting effects than naturally-occurring) ; or which have been altered to delete one or more a potential sites for o-glycosylation (which may result in higher activities for yeast-produced products) ; or which have one or more cysteine residues deleted or replaced by, e.g., alanine or serine residues and are potentially more easily isolated in active form from microbial systems ; or which have one or more tyrosine residues replaced by phenylalanine and may bind more or less readily to hpG-CSF receptors on target cells. Also comprehended are polypeptide fragments duplicating only a part of the continuous amino acid sequence or secondary conformations within hpG-CSF, which fragments may possess one activity of (e.g., receptor binding) and not others (e.g., colony growth stimulating activity). It is noteworthy that activity is not necessary for any one or more of the products of the invention to have therapeutic utility (see, Weiland, et al., *Blut*, 44, 173-175 (1982)) or utility in other contexts, such as in assays of hpG-CSF antagonism. Competitive antagonists may be quite useful in, for example, cases of overproduction of hpG-CSF.

According to another aspect of the present invention, the DNA sequence described herein which encodes hpG-CSF polypeptides is valuable for the information which it provides concerning the amino acid sequence of the mammalian protein which has heretofore been unavailable despite analytical processing of isolates of naturally-occurring products. The DNA sequences are also conspicuously valuable as products useful in effecting the large scale microbial synthesis of hpG-CSF by a variety of recombinant techniques. Put another way, DNA sequences provided by the invention are useful in generating new and useful viral and circular plasmid DNA vectors, new and useful transformed and transfected microbial procaryotic and eucaryotic host cells (including bacterial and yeast cells and mammalian cells grown in culture), and new and useful methods for cultured growth of such microbial host cells capable of expression of hpG-CSF and its related products. DNA sequences of the invention are also conspicuously suitable materials for use as labelled probes in isolating hpG-CSF and related protein encoding human genomic DNA as well as cDNA and genomic DNA sequences of other mammalian species. DNA sequences may also be useful in various alternative methods of protein synthesis (e.g., in insect cells) or in genetic therapy in humans and other mammals. DNA sequences of the invention are expected to be useful in developing transgenic mammalian species which may serve as eucaryotic "hosts" for production of hpG-CSF and hpG-CSF products in quantity. See, generally, Palmiter, et al., *Science*, 222(4625), 809-814 (1983).

Of applicability to hpG-CSF fragments and polypeptide analogs of the invention are reports of the

immunological activity of synthetic peptides which substantially duplicate the amino acid sequence extant in naturally-occurring proteins, glycoproteins and nucleoproteins. More specifically, relatively low molecular weight polypeptides have been shown to participate in immune reactions which are similar in duration and extent to the immune reactions of physiologically significant proteins such as viral antigens, polypeptide hormones, and the like. Included among the immune reactions of such polypeptides is the provocation of the formation of specific antibodies in immunologically active animals. See, e.g., Lerner, et al., *Cell*, 23, 309-310 (1981); Ross, et al., *Nature*, 294, 654-656 (1981); Walter, et al., *Proc. Natl. Acad. Sci. (USA)*, 77, 5197-5200 (1980); Lerner, et al., *Proc. Natl. Acad. Sci. (USA)*, 78, 3403-3407 (1981); Walter, et al., *Proc. Natl. Acad. Sci. (USA)*, 78, 4882-4886 (1981); Wong, et al., *Proc. Natl. Acad. Sci. (USA)*, 78, 7412-7416 (1981); Green, et al., *Cell*, 28, 477-487 (1982); Nigg, et al., *Proc. Natl. Acad. Sci. (USA)*, 79, 5322-5326 (1982); Baron, et al., *Cell*, 28, 395-404 (1982); Dreesman, et al., *Nature*, 295, 185-160 (1982); and Lerner, *Scientific American*, 248, No. 2, 66-74 (1983). See, also, Kaiser, et al., *Science*, 223, 249-255 (1984) relating to biological and immunological activities of synthetic peptides which approximately share secondary structures of peptide hormones but may not share their primary structural conformation.

While the present invention has been described in terms of preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations which come within the scope of the invention as claimed.

20 Claims

Claims for the Contracting States BE. CH. DE. FR. GB. IT. LI. LU. NL. SE

1. A polypeptide which :
 - a) has part or all of the primary structure and one or more of the biological properties typical of naturally-occurring human pluripotent granulocyte colony-stimulating factor (hpG-CSF) of the sequence set forth in Table VII,
 - b) is a non-naturally occurring polypeptide ; and
 - c) is the product of procaryotic or eucaryotic expression of an exogenous DNA sequence.
2. A polypeptide according to Claim 1 wherein the exogenous DNA sequence is a cDNA sequence.
3. A polypeptide according to Claim 1 wherein the exogenous DNA sequence is a genomic DNA sequence.
4. A polypeptide according to Claim 1 wherein the exogenous DNA sequence is carried on an autonomously replicating DNA plasmid or viral vector.
5. A polypeptide according to Claim 1 further characterized by being covalently associated with a detectable label substance.
6. A polypeptide according to Claim 5 wherein said detectable label is a radiolabel.
7. A DNA sequence which codes upon expression in a procaryotic or eucaryotic host cell for a polypeptide product having at least a part of the primary structure and one or more of the biological properties of naturally-occurring pluripotent granulocyte colony-stimulating factor, said DNA sequence being selected from among:
 - (a) the DNA sequence set out in Table VII or the complementary stands thereof ;
 - (b) DNA sequences which hybridize to the DNA sequences defined in (a) or fragments thereof ; and
 - (c) DNA sequences which, but for the degeneracy of the genetic code, would hybridize to the DNA sequences defined in (a) or (b) and which sequences code for a polypeptide having the same amino acid sequence.
8. A procaryotic or eucaryotic host cell transformed or transfected with a DNA sequence according to Claim 7 in a manner allowing the host cell to express said polypeptide product.
9. A host cell according to Claim 8 wherein the host is E. coli.
10. A host cell according to Claim 8 wherein the host cell is a mammalian cell.
11. A non-naturally occurring polypeptide product of the expression of a DNA sequence of Claim 7 in a procaryotic or eucaryotic host.
12. A cDNA sequence according to Claim 7.
13. A genomic DNA sequence according to Claim 7.
14. A DNA sequence according to Claim 7 and including one or more codons preferred for expression in E. coli cells.
15. A DNA sequence according to Claim 7 and coding for expression of human pluripotent granulocyte colony-stimulating factor.
16. A DNA sequence according to Claim 15 and including one or more codons preferred for expression in yeast cells.



immunological activity of synthetic peptides which substantially duplicate the amino acid sequence extant in naturally-occurring proteins, glycoproteins and nucleoproteins. More specifically, relatively low molecular weight polypeptides have been shown to participate in immune reactions which are similar in duration and extent to the immune reactions of physiologically significant proteins such as viral antigens, polypeptide hormones, and the like. Included among the immune reactions of such polypeptides is the provocation of the formation of specific antibodies in immunologically active animals. See, e.g., Lerner, et al., *Cell*, 23, 309-310 (1981) ; Ross, et al., *Nature*, 294, 654-656 (1981) ; Walter, et al., *Proc. Natl. Acad. Sci. (USA)*, 77, 5197-5200 (1980) ; Lerner, et al., *Proc. Natl. Acad. Sci. (USA)*, 78, 3403-3407 (1981) ; Walter, et al., *Proc. Natl. Acad. Sci. (USA)*, 78, 4882-4886 (1981) ; Wong, et al., *Proc. Natl. Acad. Sci. (USA)*, 78, 7412-7416 (1981) ; Green, et al., *Cell*, 28, 477-487 (1982) ; Nigg, et al., *Proc. Natl. Acad. Sci. (USA)*, 79, 5322-5326 (1982) ; Baron, et al., *Cell*, 28, 395-404 (1982) ; Dreesman, et al., *Nature*, 295, 185-186 (1982) ; and Lerner, *Scientific American*, 248, No. 2, 66-74 (1983). See, also, Kaiser, et al., *Science*, 223, 249-255 (1984) relating to biological and immunological activities of synthetic peptides which approximately share secondary structures of peptide hormones but may not share their primary structural conformation.

While the present invention has been described in terms of preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations which come within the scope of the invention as claimed.

Claims

Claims for the Contracting States BE. CH. DE. FR. GB. IT. LI. LU. NL. SE

1. A polypeptide which :

a) has part or all of the primary structure and one or more of the biological properties typical of naturally-occurring human pluripotent granulocyte colony-stimulating factor (hpG-CSF) of the sequence set forth in Table VII,

b) is a non-naturally occurring polypeptide ; and

c) is the product of procaryotic or eucaryotic expression of an exogenous DNA sequence.

2. A polypeptide according to Claim 1 wherein the exogenous DNA sequence is a cDNA sequence.

3. A polypeptide according to Claim 1 wherein the exogenous DNA sequence is a genomic DNA sequence.

4. A polypeptide according to Claim 1 wherein the exogenous DNA sequence is carried on an autonomously replicating DNA plasmid or viral vector.

5. A polypeptide according to Claim 1 further characterized by being covalently associated with a detectable label substance.

6. A polypeptide according to Claim 5 wherein said detectable label is a radiolabel.

7. A DNA sequence which codes upon expression in a procaryotic or eucaryotic host cell for a polypeptide product having at least a part of the primary structure and one or more of the biological properties of naturally-occurring pluripotent granulocyte colony-stimulating factor, said DNA sequence being selected from among:

(a) the DNA sequence set out in Table VII or the complementary stands thereof ;

(b) DNA sequences which hybridize to the DNA sequences defined in (a) or fragments thereof ; and

(c) DNA sequences which, but for the degeneracy of the genetic code, would hybridize to the DNA sequences defined in (a) or (b) and which sequences code for a polypeptide having the same amino acid sequence.

8. A procaryotic or eucaryotic host cell transformed or transfected with a DNA sequence according to Claim 7 in a manner allowing the host cell to express said polypeptide product.

9. A host cell according to Claim 8 wherein the host is *E. coli*.

10. A host cell according to Claim 8 wherein the host cell is a mammalian cell.

11. A non-naturally occurring polypeptide product of the expression of a DNA sequence of Claim 7 in a procaryotic or eucaryotic host.

12. A cDNA sequence according to Claim 7.

13. A genomic DNA sequence according to Claim 7.

14. A DNA sequence according to Claim 7 and including one or more codons preferred for expression in *E. coli* cells.

15. A DNA sequence according to Claim 7 and coding for expression of human pluripotent granulocyte colony-stimulating factor.

16. A DNA sequence according to Claim 15 and including one or more codons preferred for expression in yeast cells.



17. A DNA sequence according to Claim 12 or 13 coding for expression of human pluripotent granulocyte colony-stimulating factor.
18. A DNA sequence according to Claim 7 covalently associated with a detectable label substance.
19. A DNA sequence according to Claim 18 wherein the detectable label is a radiolabel.
20. A single-stranded DNA sequence according to Claim 18.
21. A DNA sequence coding for (Ala¹) hpG-CSF.
22. A biologically functional plasmid or viral DNA vector including a DNA sequence according to Claim 7.
23. A procaryotic or eucaryotic host cell stably transformed or transfected with a DNA vector according to Claim 22.
24. A process for the production of a polypeptide having part or all of the primary structure and one or more of the biological properties of naturally occurring pluripotent granulocyte colony-stimulating factor, which process is characterized by culturing under suitable nutrient conditions, procaryotic or eucaryotic host cells transformed or transfected with a DNA sequence according to Claim 7 in a manner allowing the host cell to express said polypeptide, and isolating desired polypeptide products of the expression of DNA sequence.
25. A process for the production of a polypeptide having the primary structure of human pluripotent granulocyte colony-stimulating factor, which process is characterized by culturing under suitable nutrient conditions, procaryotic or eucaryotic host cells transformed or transfected with a DNA sequence set forth in Table VII in a manner allowing the host cells to express said polypeptide, and isolating desired polypeptide products of the expression of the DNA sequence.
26. A pharmaceutical composition comprising an effective amount of a polypeptide according to Claim 1 and/or produced by the process of Claim 24 or 25 and a pharmaceutically acceptable diluent, adjuvant or carrier.
27. A pharmaceutical composition according to Claim 26 further characterized by being free of association with any human protein.
28. Use of a polypeptide according to Claim 1 for the manufacture of a medicament for providing hematopoietic therapy to a mammal.
29. Use of a polypeptide according to Claim 1 for the manufacture of a medicament for arresting proliferation of leukemic cells.
30. A DNA sequence coding for a polypeptide analog of hpG-CSF having one or more cysteine residues deleted or replaced by alanine or serine residues.
31. A non-naturally occurring polypeptide product of the expression in a procaryotic or eucaryotic host cell of a DNA sequence according to Claim 30.
32. A biologically functional plasmid or viral DNA vector including a DNA sequence according to Claim 30.
33. A procaryotic or eucaryotic host cell stably transformed or transfected with a DNA vector according to Claim 32.
34. A polypeptide according to Claim 1 preceded by a methionine residue.
35. A polypeptide having the hematopoietic biological properties of naturally occurring pluripotent granulocyte colony-stimulating factor, said polypeptide having an amino acid sequence selected from the polypeptide sequence set forth in Table VII, or any allelic variants, derivatives, deletion analogs, substitution analogs, or addition analogs thereof, and characterized by being non-naturally occurring and by being the product of procaryotic or eucaryotic expression of an exogenous DNA sequence.
36. A non-naturally occurring polypeptide consisting only of the amino acid sequence 1-174 set forth in Table VII.
37. A pharmaceutical composition comprising an effective amount of the polypeptide according to Claim 34 or 36 and a pharmaceutically acceptable diluent, adjuvant or carrier.

Claims for the Contracting State AT

1. A process for the production of a polypeptide having part or all of the primary structure and one or more of the biological properties of naturally occurring pluripotent granulocyte colony-stimulating factor, which process is characterized by culturing, under suitable nutrient conditions, procaryotic or eucaryotic host cells transformed or transfected with an exogenous DNA sequence in a manner allowing the host cell to express the polypeptide, and isolating desired polypeptide having the sequence set forth in Table VII and characterized by being a non-naturally occurring polypeptide.
2. A process according to Claim 1 wherein the exogenous DNA sequence is a cDNA sequence.
3. A process according to Claim 1 wherein the exogenous DNA sequence is a genomic DNA sequence.
4. A process according to Claim 1 wherein the exogenous DNA sequence is carried on an autonomously replicating DNA plasmid or viral vector.
5. A process according to Claim 1 wherein the polypeptide is covalently associated with a detectable label



substance.

6. A process according to Claim 5 wherein said detectable label is a radiolabel.

7. A process according to Claim 1 wherein the exogenous DNA sequence codes upon expression in a pro-
caryotic or eucaryotic host cell for a polypeptide product having at least a part of the primary structure and one
or more of the biological properties of naturally-occurring pluripotent granulocyte colony-stimulating factor, said
DNA sequence being selected from among :

(a) the DNA sequence set out in Table VII or the complementary stands thereof ;

(b) DNA sequences which hybridize to the DNA sequences defined in (a) or fragments thereof ; and

(c) DNA sequences which, but for the degeneracy of the genetic code, would hybridize to the DNA sequ-
ences defined in (a) or (b) and which sequences code for a polypeptide having the same amino acid sequ-
ence.

8. A process according to Claim 1, using a procaryotic or eucaryotic host cell transformed or transfected
with a DNA sequence as set out in Claim 7 in a manner allowing the host cell to express said polypeptide pro-
duct.

9. A process according to Claim 8 wherein the host is E. coli.

10. A process according to Claim 8 wherein the host cell is a mammalian cell.

11. A cDNA sequence as set out in Claim 7.

12. A genomic DNA sequence as set out in Claim 7.

13. A DNA sequence as set out in Claim 7 and including one or more codons preferred for expression in
E. coli cells.

14. A DNA sequence as set out in Claim 7 and coding for expression of human pluripotent granulocyte
colony-stimulating factor.

15. A DNA sequence according to Claim 14 and including one or more codons preferred for expression in
yeast cells.

16. A DNA sequence according to Claim 11 or 12 and coding for expression of a human pluripotent granulo-
cyte colony-stimulating factor-coding.

17. A DNA sequence as set out in Claim 7 covalently associated with a detectable label substance.

18. A DNA sequence according to Claim 17 wherein the detectable label is a radiolabel.

19. A single-stranded DNA sequence according to Claim 17.

20. A DNA sequence coding for (Ala¹) hpG-CSF.

21. A biologically functional plasmid or viral DNA vector including a DNA sequence as set out in Claim 7.

22. A procaryotic or eucaryotic host cell stably transformed or transfected with a DNA vector according to
Claim 21.

23. A process for the production of a polypeptide having the primary structure of human pluripotent granulo-
cyte colony-stimulating factor, which process is characterized by culturing under suitable nutrient conditions,
procaryotic or eucaryotic host cells transformed or transfected with a DNA sequence set forth in Table VII in a
manner allowing the host cells to express said polypeptide, and isolating desired polypeptide products of the
expression of the DNA sequence.

24. A process of producing a pharmaceutical composition which process incorporating an effective amount
of a polypeptide produced by the process according to Claim 1 and/or Claim 23 in a pharmaceutically accept-
able diluent, adjuvant or carrier.

25. A process according to Claim 24 wherein the polypeptide is free of association with any human protein.

26. Use of a polypeptide having part or all of the primary structure and one or more of the biological proper-
ties of naturally occurring pluripotent granulocyte colony-stimulating factor, which process is characterized by
culturing, under suitable nutrient conditions, procaryotic or eucaryotic host cells transformed or transfected with
an exogenous DNA sequence in a manner allowing the host cell to express the polypeptide, and isolating des-
ired polypeptide having the sequence set forth in Table VII and characterized by being a non-naturally occurring
polypeptide, for the manufacture of a medicament for providing hematopoietic therapy to a mammal.

27. Use of a polypeptide having part or all of the primary structure and one or more of the biological proper-
ties of naturally occurring pluripotent granulocyte colony-stimulating factor, which process is characterized by
culturing, under suitable nutrient conditions, procaryotic or eucaryotic host cells transformed or transfected with
an exogenous DNA sequence in a manner allowing the host cell to express the polypeptide, and isolating des-
ired polypeptide having the sequence set forth in Table VII and characterized by being a non-naturally occurring
polypeptide, for the manufacture of a medicament for arresting proliferation of leukemic cells.

28. A DNA sequence coding for a polypeptide analog of hpG-CSF having one or more cysteine residues
deleted or replaced by alanine or serine residues.

29. A non-naturally occurring polypeptide product of the expression in a procaryotic or eucaryotic host cell
of a DNA sequence according to Claim 28.



30. A biologically functional plasmid or viral DNA vector including a DNA sequence according to Claim 30.
31. A procaryotic or eucaryotic host cell stably transformed or transfected with a DNA vector according to Claim 30.
32. A polypeptide produced according to the process of claim 1 which is preceded by a methomine residue.
33. A polypeptide having the hematopoietic biological properties of naturally occurring pluripotent granulocyte colony stimulating factor, said polypeptide having an amino acid sequence selected from the polypeptide sequence set forth in Table VII, or any allelic variants, derivatives, deletion analogs, substitution analogs, or addition analogs thereof, and characterized by being non-naturally occurring and by being the product of procaryotic or eucaryotic expression of an exogenous DNA sequence.
34. A non-naturally occurring polypeptide consisting only of the amino acid sequence 1-174 set forth in Table VII.
35. A pharmaceutical composition comprising an effective amount of the polypeptide having only the sequence 1-174 set forth in Table VII, and a pharmaceutically acceptable diluent, adjuvant or carrier.
36. A process of producing a pharmaceutical composition which process comprises incorporating an effective amount of the polypeptide having only the sequence 1-174 set forth in Table VII in a pharmaceutically acceptable diluent, adjuvant or carrier.

Ansprüche

Für die Vertragsstaaten BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Polypeptid, welches :
 - a) einen Teil oder die Gesamtheit der Primärstruktur und eine oder mehrere der typischen biologischen Eigenschaften von natürlich auftretendem menschlichem pluripotente Granulozytkolonien erregendem Faktor (hpG-CSF) der in Tabelle VII dargestellten Sequenz besitzt,
 - b) ein nicht-natürlich auftretendes Polypeptid ist ; und
 - c) das Produkt prokaryontischer oder eukaryontischer Expression einer exogenen DNA-Sequenz ist.
2. Polypeptid nach Anspruch 1, dadurch gekennzeichnet, daß die exogene DNA-Sequenz eine cDNA-Sequenz ist.
3. Polypeptid nach Anspruch 1, dadurch gekennzeichnet, daß die exogene DNA-Sequenz eine genomische DNA-Sequenz ist.
4. Polypeptid nach Anspruch 1, dadurch gekennzeichnet, daß die exogene DNA-Sequenz auf einem autonom replizierenden DNA-Plasmid- oder viralen Vektor vorliegt.
5. Polypeptid nach Anspruch 1, dadurch gekennzeichnet, daß es kovalent mit einer nachweisbaren Markierungssubstanz verknüpft ist.
6. Polypeptid nach Anspruch 5, dadurch gekennzeichnet, daß besagte nachweisbare Markierung eine radioaktive Markierung ist
7. DNA-Sequenz, die bei Expression in einer prokaryontischen oder eukaryontischen Wirtszelle für ein Polypeptidprodukt kodiert, das wenigstens einen Teil der Primärstruktur und eine oder mehrere der biologischen Eigenschaften von natürlich auftretendem pluripotente Granulozytkolonien erregendem Faktor besitzt, wobei besagte DNA-Sequenz ausgewählt ist aus :
 - a) der DNA-Sequenz, die in Tabelle VII dargestellt ist, oder den komplementären Strängen derselben ;
 - b) DNA-Sequenzen, die zu den in (a) definierten DNA-Sequenzen oder Fragmenten derselben hybridisieren ; und
 - c) DNA-Sequenzen, die, ohne die Degeneration des genetischen Codes, zu den in (a) oder (b) definierten DNA-Sequenzen hybridisieren würden und die für ein Polypeptid mit derselben Aminosäuresequenz kodieren.
8. Prokaryontische oder eukaryontische Wirtszelle, die mit einer DNA-Sequenz nach Anspruch 7 in einer Weise transformiert oder transfiziert ist, die es der Wirtszelle erlaubt, besagtes Polypeptidprodukt zu exprimieren.
9. Wirtszelle nach Anspruch 8, dadurch gekennzeichnet, daß der Wirt E. coli ist.
10. Wirtszelle nach Anspruch 8, dadurch gekennzeichnet, daß die Wirtszelle eine Säugetierzelle ist.
11. Nicht-natürlich auftretendes Polypeptidprodukt der Expression einer DNA-Sequenz nach Anspruch 7 in einem prokaryontischen oder eukaryontischen Wirt.
12. cDNA-Sequenz nach Anspruch 7.
13. Genomische DNA-Sequenz nach Anspruch 7.
14. DNA-Sequenz nach Anspruch 7, die ein oder mehrere für die Expression in E. coli-Zellen bevorzugte



Codons einschließt.

15. DNA-Sequenz nach Anspruch 7, die für die Expression von menschlichem pluripotente Granulozytkolonien erregendem Faktor kodiert.

16. DNA-Sequenz nach Anspruch 15, die ein oder mehrere für die Expression in Hefezellen bevorzugte Codons einschließt.

17. DNA-Sequenz nach Anspruch 12 oder 13, die für die Expression von menschlichem pluripotente Granulozytkolonien erregendem Faktor kodiert.

18. DNA-Sequenz nach Anspruch 7, kovalent verknüpft mit einer nachweisbaren Markierungssubstanz.

19. DNA-Sequenz nach Anspruch 18, dadurch gekennzeichnet, daß die nachweisbaren Markierung eine radioaktive Markierung ist.

20. Einzelsträngige DNA-Sequenz nach Anspruch 18.

21. DNA-Sequenz, die für (Ala¹) hpG-CSF kodiert.

22. Biologisch funktionaler Plasmid- oder viraler DNA-Vektor, der eine DNA-Sequenz nach Anspruch 7 einschließt.

23. Prokaryontische oder eukaryontische Wirtszelle, stabil transformiert oder transfiziert mit einem DNA-Vektor nach Anspruch 22.

24. Verfahren zur Herstellung eines Polypeptids, das einen Teil oder die Gesamtheit der Primärstruktur und eine oder mehrere der biologischen Eigenschaften von natürlich auftretendem pluripotente Granulozytkolonien erregendem Faktor besitzt, wobei dieses Verfahren gekennzeichnet ist durch Kultivieren von prokaryontischen oder eukaryontischen Wirtszellen, die mit einer DNA-Sequenz nach Anspruch 7 in einer Weise transformiert oder transfiziert sind, die es der Wirtszelle erlaubt, besagtes Polypeptid zu exprimieren, unter geeigneten Nährstoffbedingungen und Isolieren besagter Polypeptidprodukte der Expression der DNA-Sequenz.

25. Verfahren zur Herstellung eines Polypeptid, das die Primärstruktur des menschlichen pluripotente Granulozytkolonien erregenden Faktors besitzt, wobei dieses Verfahren gekennzeichnet ist durch Kultivieren von prokaryontischen oder eukaryontischen Wirtszellen, die mit einer DNA-Sequenz, wie sie in Tabelle VII dargestellt ist, in einer Weise transformiert oder transfiziert sind, die es den Wirtszellen erlaubt, besagtes Polypeptid zu exprimieren unter geeigneten Nährstoffbedingungen und Isolieren der gewünschten Polypeptidprodukte der Expression der DNA-Sequenz.

26. Pharmazeutische Zusammensetzung, die eine wirksame Menge eines Polypeptids nach Anspruch 1 und/oder hergestellt nach dem Verfahren von Anspruch 24 oder 25 und ein pharmazeutisch akzeptables Streckmittel, Adjuvans oder Trägermittel umfaßt.

27. Pharmazeutische Zusammensetzung nach Anspruch 26, dadurch gekennzeichnet, daß sie von Verknüpfung mit irgendeinem menschlichen Protein frei ist.

28. Verwendung eines Polypeptids nach Anspruch 1 für die Herstellung eines Arzneimittels für die hämatopoetische Therapie eines Säugetier.

29. Verwendung eines Polypeptids nach Anspruch 1 für die Herstellung eines Arzneimittels, um die Proliferation von Leukämiezellen zum Stillstand zu bringen.

30. DNA-Sequenz, die für ein Polypeptid-Analogon von hpG-CSF kodiert, wobei ein oder mehrere Cystein-Reste weggelassen oder durch Alanin- oder Serin-Reste ersetzt sind.

31. Nicht-natürlich auftretendes Polypeptidprodukt der Expression einer DNA-Sequenz nach Anspruch 30 in einer prokaryontischen oder eukaryontischen Wirtszelle.

32. Biologisch funktionaler Plasmid- oder viraler DNA-Vektor, der eine DNA-Sequenz nach Anspruch 30 einschließt.

33. Prokaryontische oder eukaryontische Wirtszelle, stabil transformiert oder transfiziert mit einem DNA-Vektor nach Anspruch 32.

34. Polypeptid nach Anspruch 1, dem ein Methionin-Rest vorangeht.

35. Polypeptid mit den hämatopoetischen biologischen Eigenschaften von natürlich auftretendem pluripotente Granulozytkolonien erregendem Faktor, wobei besagtes Polypeptid eine Aminosäuresequenz besitzt, die ausgewählt ist aus der Polypeptidsequenz, die in Tabelle VII dargestellt ist oder irgendwelchen allelischen Varianten, Derivaten, Deletionsanaloga, Substitutionsanaloga oder Additionsanaloga derselben und dadurch gekennzeichnet, daß sie nicht-natürlich auftritt und das Produkt prokaryontischer oder eukaryontischer Expression einer exogenen DNA-Sequenz ist.

36. Nicht-natürlich auftretendes Polypeptid, das nur aus der Aminosäuresequenz 1-174 besteht, die in Tabelle VII dargestellt ist.

37. Pharmazeutische Zusammensetzung, die eine wirksame Menge des Polypeptids nach Anspruch 34 oder 36 und ein pharmazeutisch akzeptables Verdünnungsmittel, Adjuvans oder Trägermittel umfaßt.



Für den Vertragsstaat AT

1. Verfahren zur Herstellung eines Polypeptids, das einen Teil oder die Gesamtheit der Primärstruktur und eine oder mehrere der biologischen Eigenschaften von natürlich auftretendem pluripotente Granulozytkolonien erregendem Faktor besitzt, wobei dieses Verfahren gekennzeichnet ist durch Kultivieren von prokaryontischen oder eukaryontischen Wirtszellen, die mit einer exogenen DNA-Sequenz in einer Weise transformiert oder transfiziert sind, die es der Wirtszelle erlaubt, das Polypeptid zu exprimieren, unter geeigneten Nährstoffbedingungen und Isolieren des gewünschten Polypeptids, das die in Tabelle VII dargestellte Sequenz besitzt und dadurch gekennzeichnet ist, daß es ein nicht-natürlich auftretendes Polypeptid ist.
2. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß die exogene DNA-Sequenz eine cDNA-Sequenz ist.
3. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß die exogene DNA-Sequenz eine genomische DNA-Sequenz ist.
4. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß die exogene DNA-Sequenz auf einem autonom replizierenden DNA-Plasmid- oder viralen Vektor vorliegt.
5. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß das Polypeptid kovalent mit einer nachweisbaren Markierungssubstanz verknüpft ist.
6. Verfahren nach Anspruch 5, dadurch gekennzeichnet, daß besagte nachweisbare Markierung eine radioaktive Markierung ist.
7. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß die exogene DNA-Sequenz bei Expression in einer prokaryontischen oder eukaryontischen Wirtszelle für ein Polypeptidprodukt kodiert, das wenigstens einen Teil der Primärstruktur und eine oder mehrere der biologischen Eigenschaften von natürlich auftretendem pluripotente Granulozytkolonien erregendem Faktor besitzt, wobei besagte DNA-Sequenz ausgewählt ist aus:
 - a) der DNA-Sequenz, die in Tabelle VII dargestellt ist, oder den komplementären Strängen derselben ;
 - b) DNA-Sequenzen, die zu den in (a) definierten DNA-Sequenzen oder Fragmenten derselben hybridisieren ; und
 - c) DNA-Sequenzen, die, ohne die Degeneration des genetischen Codes, zu den in (a) oder (b) definierten DNA-Sequenzen hybridisieren würden und die für ein Polypeptid mit derselben Aminosäuresequenz kodieren.
8. Verfahren nach Anspruch 1, das eine prokaryontische oder eukaryontische Wirtszelle verwendet, die mit einer DNA-Sequenz, wie in Anspruch 7 dargelegt, in einer Weise transformiert oder transfiziert ist, die es der Wirtszelle erlaubt, besagtes Polypeptidprodukt zu exprimieren.
9. Verfahren nach Anspruch 8, dadurch gekennzeichnet, daß der Wirt E. coli ist.
10. Verfahren nach Anspruch 8, dadurch gekennzeichnet, daß die Wirtszelle eine Säugetierzelle ist.
11. cDNA-Sequenz, wie in Anspruch 7 dargelegt.
12. Genomische DNA-Sequenz, wie in Anspruch 7 dargelegt.
13. DNA-Sequenz, wie in Anspruch 7 dargelegt, die ein oder mehrere für die Expression in E. coli-Zellen bevorzugte Codons einschließt.
14. DNA-Sequenz, wie in Anspruch 7 dargelegt, die für die Expression von menschlichem pluripotente Granulozytkolonien erregenden Faktor kodiert.
15. DNA-Sequenz nach Anspruch 14, die ein oder mehrere für die Expression in Hefezellen bevorzugte Codons einschließt.
16. DNA-Sequenz nach Anspruch 11 oder 12, die für die Expression einer Kodierung für menschlichen pluripotente Granulozytkolonien erregenden Faktor kodiert.
17. DNA-Sequenz wie in Anspruch 7 dargelegt, kovalent verknüpft mit einer nachweisbaren Markierungssubstanz.
18. DNA-Sequenz nach Anspruch 17, dadurch gekennzeichnet, daß die nachweisbare Markierung eine radioaktive Markierung ist.
19. Einzelsträngige DNA-Sequenz nach Anspruch 17.
20. DNA-Sequenz, die für (Ala¹) hpG-CSF kodiert.
21. Biologisch funktionaler Plasmid- oder viraler DNA-Vektor, der eine DNA-Sequenz, wie in Anspruch dargelegt, einschließt.
22. Prokaryontische oder eukaryontische Wirtszelle, stabil transformiert oder transfiziert mit einem DNA-Vektor nach Anspruch 21.
23. Verfahren zur Herstellung eines Polypeptids, das die Primärstruktur von menschlichem pluripotente Granulozytkolonien erregendem Faktor besitzt, wobei dieses Verfahren gekennzeichnet ist durch Kultivieren von prokaryontischen oder eukaryontischen Wirtszellen, die mit einer DNA-Sequenz, wie sie in Tabelle VII dargestellt ist, in einer Weise transformiert oder transfiziert sind, die es den Wirtszellen erlaubt, besagtes Poly-



peptid zur exprimieren unter geeigneten Nährstoffbedingungen und Isolieren der gewünschten Polypeptidprodukte der Expression der DNA-Sequenz.

24. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, wobei dieses Verfahren eine wirksame Menge eines Polypeptids, hergestellt nach dem Verfahren nach Anspruch 1 und/oder Anspruch 23, in ein pharmazeutisch akzeptables Verdünnungsmittel, Adjuvans oder Trägermittel einarbeitet.

25. Verfahren nach Anspruch 24, dadurch gekennzeichnet, daß das Polypeptid frei von Verknüpfung mit irgendeinem menschlichen Protein ist.

26. Verwendung eines Polypeptids, das einen Teil oder die Gesamtheit der Primärstruktur und eine oder mehrere der biologischen Eigenschaften von natürlich auftretendem pluripotente Granulozytcolonien erregendem Faktor besitzt, wobei dieses Verfahren gekennzeichnet ist durch Kultivieren von prokaryontischen oder eukaryontischen Wirtszellen, die mit einer exogenen DNA-Sequenz in einer Weise transformiert und transfiziert sind, die es der Wirtszelle erlaubt, das Polypeptid zu exprimieren, unter geeigneten Nährstoffbedingungen und Isolieren des gewünschten Polypeptids, das die in Tabelle VII dargestellte Sequenz besitzt und dadurch gekennzeichnet ist, daß es ein nicht-natürlich auftretendes Polypeptid ist, für die Herstellung eines Arzneimittels zum für die hematopoyetische Therapie eines Säugetiers.

27. Verwendung eines Polypeptids, das einen Teil oder die Gesamtheit der Primärstruktur und eine oder mehrere der biologischen Eigenschaften von natürlich auftretendem pluripotente Granulozytcolonien erregenden Faktor besitzt, wobei dieses Verfahren gekennzeichnet ist durch Kultivieren von prokaryontischen oder eukaryontischen Wirtszellen, die mit einer exogenen DNA-Sequenz in einer Weise transformiert oder transfiziert sind, die es der Wirtszelle erlaubt, das Polypeptid zu exprimieren, unter geeigneten Nährstoffbedingungen und Isolieren des gewünschten Polypeptids, das die in Tabelle VII dargestellte Sequenz besitzt und dadurch gekennzeichnet ist, daß es sich um eine nichtnatürlich auftretendes Peptid handelt, für die Herstellung eines Arzneimittels, um die Proliferation von Leukämiezellen zum Stillstand zu bringen.

28. DNA-Sequenz, die für ein Polypeptid-Analogon von hpG-CSF kodiert, wobei ein oder mehrere Cystein-Reste weggelassen oder durch Alanin- oder Serin-Reste ersetzt sind.

29. Nicht-natürlich auftretendes Polypeptidprodukt der Expression einer DNA-Sequenz nach Anspruch 28 in einer prokaryontischen oder eukaryontischen Wirtszelle.

30. Biologisch funktionaler Plasmid- oder viraler DNA-Vektor, der eine DNA-Sequenz nach Anspruch 30 einschließt.

31. Prokaryontische oder eukaryontische Wirtszelle, stabil transformiert oder transfiziert mit einem DNA-Vektor nach Anspruch 30.

32. Polypeptid, hergestellt nach dem Verfahren von Anspruch 1, dem ein Methionin-Rest vorangeht.

33. Polypeptid mit den hematopoetischen biologischen Eigenschaften von natürlich auftretendem pluripotente Granulozytcolonien erregendem Faktor, wobei besagtes Polypeptid eine Aminosäuresequenz besitzt, die ausgewählt ist aus der in Tabelle VII dargestellten Polypeptidsequenz oder irgendwelchen allelischen Varianten, Derivaten, Deletionsanaloga, Substitutionsanaloga oder Additionsanaloga derselben und dadurch gekennzeichnet ist, daß es nicht-natürlich auftritt und das Produkt von prokaryontischer oder eukaryontischer Expression einer exogenen DNA-Sequenz ist.

34. Nicht-natürlich auftretendes Polypeptid, das nur aus der Aminosäuresequenz 1-174, die in Tabelle VII dargestellt ist, besteht.

35. Pharmazeutische Zusammensetzung, die eine wirksame Menge des Polypeptids, die nur die Sequenz 1-74, die in Tabelle VII dargestellt ist, besitzt, und ein pharmazeutisch akzeptables Verdünnungsmittel, Adjuvans oder Trägermittel umfaßt.

36. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, wobei dieses Verfahren das Einarbeiten einer wirksamen Menge des Polypeptids, das nur die Sequenz 1-174, die in Tabelle VII dargestellt ist, besitzt, in ein pharmazeutisch akzeptables Verdünnungsmittel, Adjuvans oder Trägermittel umfaßt.

Revendications

Revendications pour les Etats Contractants : BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Polypeptide qui,

a) présente une partie ou toute la structure principale et une ou plusieurs des propriétés biologiques typiques du facteur de stimulation de colonies de granulocytes pluriactifs humains existant naturellement (hpG-CSF) de la séquence indiquée dans le tableau VII,

b) est un polypeptide existant non-naturellement ; et

c) est le produit de l'expression eucaryote ou procaryote d'une séquence d'ADN exogène.



2. Polypeptide selon la revendication 1, dans lequel la séquence d'ADN exogène est une séquence d'ADNc.

3. Polypeptide selon la revendication 1, dans lequel la séquence d'ADN exogène est une séquence d'ADN génomique.

5 4. Polypeptide selon la revendication 1, dans lequel la séquence d'ADN exogène est portée par un vecteur viral ou plasmide d'ADN à répllication autonome.

5. Polypeptide selon la revendication 1, caractérisé de plus en ce qu'il est associé, de façon covalente, à un traceur détectable.

6. Polypeptide selon la revendication 5, dans lequel ledit traceur détectable est un radiotraceur.

10 7. Séquence d'ADN qui code, lors de l'expression dans une cellule hôte eucaryote ou procaryote, pour un polypeptide ayant au moins une partie de la structure principale et une ou plusieurs des propriétés biologiques du facteur de stimulation de colonies de granulocytes pluriactifs existant naturellement, ladite séquence d'ADN étant choisie parmi :

(a) la séquence d'ADN indiquée dans le tableau VII ou les brins complémentaires de celle-ci ;

15 (b) des séquences d'ADN qui s'hybrident en séquences d'ADN définies dans a) ou des fragments de celles-ci ; et

(c) des séquences d'ADN qui, n'étant la dégénérescence du code génétique, s'hybrideraient en séquences d'ADN définies dans a) ou b) et lesquelles séquences codent pour un polypeptide ayant la même séquence d'acides aminés.

20 8. Cellule hôte eucaryote ou procaryote, transformée ou ayant subi une transfection avec une séquence d'ADN selon la revendication 7 d'une façon permettant à la cellule hôte d'exprimer ledit polypeptide.

9. Cellule hôte selon la revendication 8, dans laquelle l'hôte est *E. coli*.

10. Cellule hôte selon la revendication 8, dans laquelle la cellule hôte est une cellule de mammifère.

25 11. Polypeptide existant non-naturellement produit par l'expression d'une séquence d'ADN de la revendication 7 dans un hôte eucaryote ou procaryote.

12. Séquence d'ADNc selon la revendication 7.

13. Séquence d'ADN génomique selon la revendication 7.

14. Séquence d'ADN selon la revendication 7 et comportant un ou plusieurs codons préférés pour l'expression dans des cellules *E. Coli*.

30 15. Séquence d'ADN selon la revendication 7 et codant pour l'expression du facteur de stimulation de colonies de granulocytes pluriactifs humains.

16. Séquence d'ADN selon la revendication 15 et incluant un ou plusieurs codons préférés pour l'expression dans des cellules de levure.

35 17. Séquence d'ADN selon la revendication 12 ou 13 codant pour l'expression du facteur de stimulation de colonies de granulocytes pluriactifs humains.

18. Séquence d'ADN selon la revendication 7 associée, de façon covalente, à un traceur détectable.

19. Séquence d'ADN selon la revendication 18, dans laquelle le traceur détectable est un radiotraceur.

20. Séquence d'ADN à brin unique selon la revendication 18.

21. Séquence d'ADN codant pour (Ala¹) hpG-CSF.

40 22. Vecteur d'ADN viral ou plasmide biologiquement fonctionnel incluant une séquence d'ADN selon la revendication 7.

23. Cellule hôte eucaryote ou procaryote, transformée ou ayant subi une transfection, de façon stable, avec un vecteur d'ADN selon la revendication 22.

45 24. Procédé pour la production d'un polypeptide ayant une partie ou toute la structure principale et une ou plusieurs des propriétés biologiques du facteur de stimulation de colonies de granulocytes pluriactifs existant naturellement, lequel procédé est caractérisé par la mise en culture, dans des conditions nutritives appropriées, de cellules hôtes eucaryotes or procaryotes, transformées ou ayant subi une transfection avec une séquence d'ADN selon la revendication 7 d'une façon permettant à la cellule hôte d'exprimer ledit polypeptide, et l'isolation des produits polypeptidiques souhaités de l'expression de la séquence d'ADN.

50 25. Procédé pour la production d'un polypeptide ayant la structure principale du facteur de stimulation de colonies de granulocytes pluriactifs humains, lequel procédé est caractérisé par la mise en culture, dans des conditions nutritives appropriées, de cellules hôtes eucaryotes ou procaryotes, transformées ou ayant subi une transfection avec une séquence d'ADN indiquée dans le tableau VII d'une façon permettant aux cellules hôtes d'exprimer ledit polypeptide, et l'isolation des produits polypeptidiques souhaités de l'expression de la séquence d'ADN.

55 26. Composition pharmaceutique comprenant une quantité efficace d'un polypeptide selon la revendication 1 et/ou produit par le procédé de la revendication 24 ou 25 et un support, adjuvant ou diluant pharmaceutiquement acceptable.



27. Composition pharmaceutique selon la revendication 26, caractérisée de plus en ce qu'elle est dépourvue de toute association avec une protéine humaine quelconque.

28. Utilisation d'un polypeptide selon la revendication 1 pour la fabrication d'un médicament fournissant une thérapie hématopoïétique à un mammifère.

5 29. Utilisation d'un polypeptide selon la revendication 1 pour la fabrication d'un médicament pour stopper la prolifération de cellules leucémiques.

30. Séquence d'ADN codant pour un polypeptide analogue à hpG-CSF, ayant un ou plusieurs résidus de cystéine, supprimés ou remplacés par des résidus d'alanine ou de sérine.

10 31. Produit polypeptidique existant non-naturellement de l'expression dans une cellule hôte eucaryote ou procaryote d'une séquence d'ADN selon la revendication 30.

32. Vecteur d'ADN viral ou plasmide biologiquement fonctionnel incluant une séquence d'ADN selon la revendication 30.

33. Cellule hôte eucaryote ou procaryote, transformée ou ayant subi une transfection, de façon stable, avec un vecteur d'ADN selon la revendication 32.

15 34. Polypeptide selon la revendication 1 précédé par un résidu de méthionine.

35. Polypeptide ayant les propriétés biologiques hématopoïétiques du facteur de stimulation de colonies de granulocytes pluriactifs existant naturellement, ledit polypeptide ayant une séquence d'acides aminés, choisie à partir de la séquence de polypeptide indiquée dans le tableau VII, ou tout analogue d'addition, analogue de substitution, analogue d'élimination, dérivé, variant alléomorphe de celle-ci, et caractérisé en ce qu'il existe non-naturellement et en ce qu'il est le produit de l'expression eucaryote ou procaryote d'une séquence d'ADN exogène.

36. Polypeptide existant non-naturellement, constitué seulement de la séquence d'acides aminés 1-174, indiquée dans le tableau VII.

25 37. Composition pharmaceutique comprenant une quantité efficace du polypeptide selon la revendication 34 ou 36 et un support, adjuvant ou diluant pharmaceutiquement acceptable.

Revendications pour l'Etat Contractant AT

30 1. Procédé pour la production d'un polypeptide ayant une partie ou toute la structure principale et une ou plusieurs des propriétés biologiques du facteur de stimulation de colonies de granulocytes pluriactifs existant naturellement, lequel procédé est caractérisé par la mise en culture, dans des conditions nutritives appropriées, de cellules hôtes eucaryotes ou procaryotes, transformées ou ayant subi une transfection avec une séquence d'ADN exogène d'une façon permettant à la cellule hôte d'exprimer le polypeptide, et l'isolation du polypeptide souhaité ayant la séquence indiquée dans le tableau VII et caractérisé en ce qu'il est un polypeptide existant non-naturellement.

2. Procédé selon la revendication 1, dans lequel la séquence d'ADN exogène est une séquence d'ADNc.

3. Procédé selon la revendication 1, dans lequel la séquence d'ADN exogène est une séquence d'ADN génomique.

40 4. Procédé selon la revendication 1, dans lequel la séquence d'ADN exogène est portée par un vecteur viral ou plasmide d'ADN à répllication autonome.

5. Procédé selon la revendication 1, dans lequel le polypeptide est associé, de façon covalente, à un traceur détectable.

6. Procédé selon la revendication 5, dans lequel ledit traceur détectable est un radiotraceur.

45 7. Procédé selon la revendication 1, dans lequel la séquence d'ADN exogène code, lors de l'expression dans une cellule hôte eucaryote ou procaryote, pour un polypeptide ayant au moins une partie de la structure principale et une ou plusieurs des propriétés biologiques du facteur de stimulation de colonies de granulocytes pluriactifs existant naturellement, ladite séquence d'ADN étant choisie parmi :

a) la séquence d'ADN indiquée dans le tableau VII ou les brins complémentaires de celle-ci ;

50 b) des séquences d'ADN qui s'hybrident en séquences d'ADN définies dans (a) ou des fragments de celles-ci ; et

c) des séquences d'ADN qui, n'étant la dégénérescence du code génétique, s'hybrideraient en séquences d'ADN définies dans (a) ou (b) et lesquelles séquences codent pour un polypeptide ayant la même séquence d'acides aminés.

8. Procédé selon la revendication 1, utilisant une cellule hôte eucaryote ou procaryote, transformée ou ayant subi une transfection avec une séquence d'ADN comme indiqué dans la revendication 7 d'une façon permettant à la cellule hôte d'exprimer ledit polypeptide.

9. Procédé selon la revendication 8, dans lequel l'hôte est E. coli.

10. Procédé selon la revendication 8, dans lequel la cellule hôte est une cellule de mammifère.



11. Séquence d'ADNc comme indiqué dans la revendication 7.
12. Séquence d'ADN génomique comme indiqué dans la revendication 7.
13. Séquence d'ADN comme indiqué dans la revendication 7 et incluant un ou plusieurs codons préférés pour l'expression dans des cellules *E. coli*.
- 5 14. Séquence d'ADN comme indiqué dans la revendication 7 et codant pour l'expression du facteur de stimulation de colonies de granulocytes pluriactifs humains.
15. Séquence d'ADN selon la revendication 14 et incluant un ou plusieurs codons préférés pour l'expression dans des cellules de levure.
16. Séquence d'ADN selon la revendication 11 ou 12 et codant pour l'expression d'un facteur de stimulation de colonies de granulocytes pluriactifs humains.
- 10 17. Séquence d'ADN comme indiqué dans la revendication 7 associée, de façon covalente, à un traceur détectable.
18. Séquence d'ADN selon la revendication 17, dans laquelle le traceur détectable est un radiotraceur.
19. Séquence d'ADN à brin unique selon la revendication 17.
- 15 20. Séquence d'ADN codant pour (Ala¹) hpG-CSF.
21. Vecteur d'ADN viral ou plasmide biologiquement fonctionnel incluant une séquence d'ADN comme indiqué dans la revendication 7.
22. Cellule hôte eucaryote ou procaryote, transformée ou ayant subi une transfection, de façon stable, avec un vecteur d'ADN selon la revendication 21.
- 20 23. Procédé pour la production d'un polypeptide ayant la structure principale du facteur de stimulation de colonies de granulocytes pluriactifs humains, lequel procédé est caractérisé par la mise en culture, dans des conditions nutritives appropriées, de cellules hôtes eucaryotes ou procaryotes, transformées ou ayant subi une transfection avec une séquence d'ADN indiquée dans le tableau VII d'une façon permettant aux cellules hôtes d'exprimer ledit polypeptide, et l'isolation des produits polypeptidiques souhaités de l'expression de la
- 25 séquence d'ADN.
24. Procédé de production d'une composition pharmaceutique, lequel procédé comprend l'incorporation d'une quantité efficace d'un polypeptide produit par le procédé selon la revendication 1 et/ou la revendication 23 dans un support, adjuvant ou diluant pharmaceutiquement acceptable.
25. Procédé selon la revendication 24, dans lequel le polypeptide n'est pas associé à une protéine humaine
- 30 quelconque.
26. Utilisation d'un polypeptide ayant une partie ou toute la structure principale et une ou plusieurs des propriétés biologiques du facteur de stimulation de colonies de granulocytes pluriactifs existant naturellement, lequel procédé est caractérisé par la mise en culture, dans des conditions nutritives appropriées, de cellules hôtes eucaryotes ou procaryotes, transformées ou ayant subi une transfection avec une séquence d'ADN exogène d'une façon permettant à la cellule hôte d'exprimer le polypeptide, et l'isolation du polypeptide souhaité
- 35 ayant la séquence indiquée dans le tableau VII et caractérisé en ce qu'il est un polypeptide existant non-naturellement, pour la fabrication d'un médicament pour fournir une thérapie hématopoïétique à un mammifère.
27. Utilisation d'un polypeptide ayant une partie ou toute la structure principale et une ou plusieurs des propriétés biologiques du facteur de stimulation de colonies de granulocytes pluriactifs existant naturellement, lequel procédé est caractérisé par la mise en culture, dans des conditions nutritives appropriées, de cellules hôtes eucaryotes ou procaryotes, transformées ou ayant subi une transfection avec une séquence d'ADN exogène d'une façon permettant à la cellule hôte d'exprimer le polypeptide, et l'isolation du polypeptide souhaité
- 40 ayant la séquence indiquée dans le tableau VII et caractérisé en ce qu'il est un polypeptide existant non-naturellement, pour la fabrication d'un médicament pour stopper la prolifération de cellules leucémiques.
28. Séquence d'ADN codant pour un polypeptide analogue à hpG-CSF, ayant un ou plusieurs résidus de cystéine supprimés ou remplacés par des résidus d'alanine ou de sérine.
29. Produit polypeptidique existant non-naturellement de l'expression dans une cellule hôte eucaryote ou procaryote d'une séquence d'ADN selon la revendication 28.
30. Vecteur d'ADN viral ou plasmide biologiquement fonctionnel incluant une séquence d'ADN selon la
- 50 revendication 28.
31. Cellule hôte eucaryote ou procaryote, transformée ou ayant subi une transfection, de façon stable, avec un vecteur d'ADN selon la revendication 30.
32. Polypeptide produit selon le procédé de la revendication 1 qui est précédé par un résidu de méthionine.
33. Polypeptide ayant les propriétés biologiques hématopoïétiques du facteur de stimulation de colonies de granulocytes pluriactifs existant naturellement, ledit polypeptide ayant une séquence d'acides aminés choisie à partir de la séquence de polypeptide indiquée dans le tableau VII, ou tout analogue d'addition, analogue de substitution, analogue d'élimination, dérivé ou variant allélomorphe de celle-ci, et caractérisé en ce qu'il existe non-naturellement et en ce qu'il est le produit de l'expression eucaryote ou procaryote d'une séquence



d'ADN exogène.

34. Polypeptide existant non-naturellement constitué seulement de la séquence d'acides aminés 1-174 indiquée dans le tableau VII.

5 35. Composition pharmaceutique comprenant une quantité efficace du polypeptide ayant seulement la séquence 1-174 indiquée dans le tableau VII, et un support, adjuvant ou diluant pharmaceutiquement acceptable.

36. Procédé de production d'une composition pharmaceutique, lequel procédé comprend l'incorporation d'une quantité efficace du polypeptide ayant seulement la séquence 1-174 indiquée dans le tableau VII dans un support, adjuvant ou diluant pharmaceutiquement acceptable.

10

15

20

25

30

35

40

45

50

55