

A CHECKPOINT INHIBITOR AND A WHOLE CELL MYCOBACTERIUM FOR
USE IN CANCER THERAPY

FIELD OF THE INVENTION

- 5 The present invention relates to the field of cancer therapy. In particular, the present invention relates to a method of preventing, treating or inhibiting the development of tumours and/or metastases in a subject.

BACKGROUND OF THE INVENTION

10

In humans with advance cancer, anti-tumour immunity is often ineffective due to the tightly regulated interplay of pro- and anti-inflammatory, immune-stimulatory and immunosuppressive signals. For example, loss of the anti-inflammatory signals leads to chronic inflammation and prolonged proliferative signalling.

- 15 Interestingly, cytokines that both promote and suppress proliferation of the tumour cells are produced at the tumour site. It is the imbalance between the effects of these various processes that results in tumour promotion.

- To date, a major barrier to attempts to develop effective immunotherapy for cancer has been an inability to break immunosuppression at the cancer site and restore normal networks of immune reactivity. The physiological approach of immunotherapy is to normalize the immune reactivity so that, for example, the endogenous tumour antigens would be recognized and effective cytolytic responses would be developed against tumour cells. Although it was once
20 unclear if tumour immunosurveillance existed, it is now believed that the immune system constantly monitors and eliminates newly transformed cells. Accordingly, cancer cells may alter their phenotype in response to immune pressure in order to escape attack (immunoediting) and upregulate expression of inhibitory signals. Through immunoediting and other subversive processes, primary
25 tumour and metastasis maintain their own survival.
30

One of the major mechanisms of anti-tumour immunity subversion is known as 'T-cell exhaustion', which results from chronic exposure to antigens and is characterized by the up-regulation of inhibitory receptors. These inhibitory

receptors serve as immune checkpoints in order to prevent uncontrolled immune reactions.

PD-1 and co-inhibitory receptors such as cytotoxic T-lymphocyte antigen 4
5 (CTLA-4, B and T Lymphocyte Attenuator (BTLA; CD272), T cell
Immunoglobulin and Mucin domain-3 (Tim-3), Lymphocyte Activation Gene-3
(Lag-3; CD223), and others are often referred to as a checkpoint regulators.
They act as molecular “tollbooths,” which allow extracellular information to
dictate whether cell cycle progression and other intracellular signalling
10 processes should proceed.

In addition to specific antigen recognition through the TCR, T-cell activation is
regulated through a balance of positive and negative signals provided by co-
stimulatory receptors. These surface proteins are typically members of either the
15 TNF receptor or B7 superfamilies. Agonistic antibodies directed against
activating co-stimulatory molecules and blocking antibodies against negative co-
stimulatory molecules may enhance T-cell stimulation to promote tumour
destruction.

20 Programmed Cell Death Protein 1, (PD-1 or CD279), a 55-kD type 1
transmembrane protein, is a member of the CD28 family of T cell co-stimulatory
receptors that include immunoglobulin superfamily member CD28, CTLA-4,
inducible co-stimulator (ICOS), and BTLA. PD-1 is highly expressed on activated
T cells and B cells. PD-1 expression can also be detected on memory T-cell
25 subsets with variable levels of expression. Two ligands specific for PD-1 have
been identified: programmed death- ligand 1 (PD-L1, also known as B7-H1 or
CD274) and PD-L2 (also known as B7-DC or CD273). PD-L1 and PD-L2 have
been shown to down-regulate T cell activation upon binding to PD-1 in both
mouse and human systems (Okazaki *et al.*, Int Immunol., 2007; 19: 813-824).
30 The interaction of PD-1 with its ligands, PD-L1 and PD-L2, which are expressed
on antigen-presenting cells (APCs) and dendritic cells (DCs), transmits negative
regulatory stimuli to down-modulate the activated T cell immune response.
Blockade of PD-1 suppresses this negative signal and amplifies T cell
responses.

Numerous studies indicate that the cancer microenvironment manipulates the PD-L1/PD-1 signalling pathway and that induction of PD-L1 expression is associated with inhibition of immune responses against cancer, thus permitting cancer progression and metastasis. The PD-L1/PD-1 signalling pathway is a primary mechanism of cancer immune evasion for several reasons. First, and most importantly, this pathway is involved in negative regulation of immune responses of activated T effector cells, found in the periphery. Second, PD-L1 is up-regulated in cancer microenvironments, while PD-1 is also up-regulated on activated tumour infiltrating T cells, thus possibly potentiating a vicious cycle of inhibition. Third, this pathway is intricately involved in both innate and adaptive immune regulation through bi-directional signalling. These factors make the PD-1/PD-L1 complex a central point through which cancer can manipulate immune responses and promote its own progression.

The first immune-checkpoint inhibitor to be tested in a clinical trial was ipilimumab (Yervoy, Bristol-Myers Squibb), an CTLA-4 mAb. CTLA-4 belongs to the immunoglobulin superfamily of receptors, which also includes PD-1, BTLA, TIM-3, and V-domain immunoglobulin suppressor of T cell activation (VISTA). Anti-CTLA-4 mAb is a powerful checkpoint inhibitor which removes “the break” from both naïve and antigen-experienced cells. Therapy enhances the antitumor function of CD8⁺ T cells, increases the ratio of CD8⁺ T cells to Foxp3⁺ T regulatory cells, and inhibits the suppressive function of T regulatory cells. The major drawback to anti-CTLA-4 mAb therapy is the generation of autoimmune toxicities due to on-target effects of an over-exuberant immune system which has lost the ability to turn itself down. It has been reported that up to 25% of patients treated with ipilimumab developed serious grade 3–4 adverse events/autoimmune-type side effects including dermatitis, enterocolitis, hepatitis, endocrinopathies (including hypophysitis, thyroiditis, and adrenalitis), arthritis, uveitis, nephritis, and aseptic meningitis. In contrast to the anti-CTLA-4 experience, anti-PD-1 therapy appears to be better-tolerated and induces a relatively lower rate of autoimmune-type side effects.

TIM-3 has been identified as another important inhibitory receptor expressed by exhausted CD8⁺ T cells. In mouse models of cancer, it has been shown that the

most dysfunctional tumour-infiltrating CD8+ T cells actually co-express PD-1 and TIM-3.

LAG-3 is another recently identified inhibitory receptor that acts to limit effector
5 T-cell function and augment the suppressive activity of T regulatory cells. It has recently been revealed that PD-1 and LAG-3 are extensively co-expressed by tumour-infiltrating T cells in mice, and that combined blockade of PD-1 and LAG-3 provokes potent synergistic antitumor immune responses in mouse models of cancer.

10

PD-1 pathway blockade can be combined with vaccines or other immunomodulatory antibodies for improved therapeutic efficacy (Hirano, F. et al, Cancer Res., 65(3): 1089-1096 (2005); Li, B. et al, Clin. Cancer Res., 15: 1507-1509 (2009); and Curran, M.A. et al, Proc. Natl. Acad. Set, 107(9):4275-4280
15 (2010)).

Currently, antagonist mAbs against both PD-1 and their ligand PD-L1 are in various stages of development for the treatment of cancer, and recent human trials have shown promising results in cancer patients with advanced, treatment-
20 refractory disease.

The first of the agents blocking the B7-H1/PD-1 pathway to enter phase I clinical trials was Nivolumab (MDX-1106/BMS-936558/ONO-4538), a fully human IgG4 anti-PD-1 mAb developed by Bristol-Myers Squibb. Another PD-1 mAb
25 undergoing clinical evaluation is CT-011, a humanized IgG1 mAb specific for PD-1 developed by CureTech Ltd. Other agents include Lambrolizumab (MK-3475 - Merck), a humanized monoclonal IgG4 PD-1 antibody; BMS-936559, a fully human IgG4 PD-L1 antibody and Roche's MPDL3280A, a human monoclonal antibody that targets the PD-L1 pathway.

30

Accordingly, an aim of the present invention is a combination therapy for treating cancer comprising an immunomodulator and blockade of checkpoint inhibitors with the potential to elicit potent and durable immune responses.

SUMMARY OF THE INVENTION

The present invention provides an effective method for treating and/or preventing cancer and/or the establishment of metastases by administering a
5 checkpoint inhibitor which acts synergistically with a whole cell *Mycobacterium*.

In a first aspect of the invention, there is an immunomodulator for use in the treatment, reduction, inhibition or control of a neoplastic disease in a patient intended to undergo checkpoint inhibition therapy simultaneously, separately or
10 sequentially with administration of the immunomodulator.

In a second aspect of the invention, there is a method of treating, reducing, inhibiting or controlling a neoplasia, tumour or cancer in a subject, wherein said method comprises simultaneously, separately or sequentially administering to
15 the subject, (i) a checkpoint inhibitor, and (ii) an immunomodulator, wherein said method results in enhanced therapeutic efficacy relative to administration of the checkpoint inhibitor or immunomodulator alone.

In a third aspect of the invention, there is a method of treating, reducing,
20 inhibiting or controlling a neoplasia, tumour or cancer in a subject, wherein said method comprises simultaneously, separately or sequentially administering to the subject, (i) a sub-therapeutic amount and/or duration of checkpoint inhibitor, and (ii) an immunomodulator, wherein said method results in enhanced therapeutic efficacy relative to administration of the checkpoint inhibitor or
25 immunomodulator alone.

The present invention therefore provides a combination therapy of checkpoint inhibitor therapy together with a specific type of immunotherapy comprising administration of an immunomodulator. The inventors have found that the
30 combination of both therapies is synergistic beyond simple additive effects of each therapy used individually.

Description of the Drawings

The invention is described with reference to the following drawings, in which:

- 5 Figure 1 shows the effect of a preparation of heat-killed *Mycobacterium obuense* (NCTC 13365) with or without co-administration of a checkpoint inhibitor (anti-PD-L1 mAb), in a xenograft model of pancreatic cancer (KPC cells injected subcutaneously).

10 Detailed Description of the Invention

The invention provides a method for treating, reducing, inhibiting or controlling a neoplasia, tumour or cancer in a subject involving administering an immunomodulator and a checkpoint inhibitor. It is based upon the discovery that
15 administration of an immunomodulator (whole cell heat-killed *Mycobacterium*) in combination with an anti- PD-L1 antibody (a checkpoint inhibitor) results in synergistic anti-tumour activity and/or antitumor activity that is more potent than administration of immunomodulator or anti-PD-L1 antibody alone.

20 In order that the present invention may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.

A “checkpoint inhibitor” is an agent which acts on surface proteins which are
25 members of either the TNF receptor or B7 superfamilies, including agents which bind to negative co-stimulatory molecules selected from CTLA-4, PD-1, TIM-3, BTLA, VISTA, LAG-3, and/or their respective ligands, including PD-L1. (Mellman et al., *supra*).

30 An immunomodulator, as defined according to the present invention, is a component which stimulates innate and type-1 immunity, including Th1 and macrophage activation and cytotoxic cell activity, as well as independently down-regulating inappropriate anti-Th2 responses via immunoregulatory mechanisms.

The terms "tumour," "cancer" and "neoplasia" are used interchangeably and refer to a cell or population of cells whose growth, proliferation or survival is greater than growth, proliferation or survival of a normal counterpart cell, e.g. a cell proliferative or differentiative disorder. Typically, the growth is uncontrolled.

- 5 The term "malignancy" refers to invasion of nearby tissue. The term "metastasis" refers to spread or dissemination of a tumour, cancer or neoplasia to other sites, locations or regions within the subject, in which the sites, locations or regions are distinct from the primary tumour or cancer.

- 10 The terms "Programmed Death 1," "Programmed Cell Death 1," "Protein PD-1," "PD-1," and "PD1," are used interchangeably, and include variants, isoforms, species homologs of human PD-1, and analogs having at least one common epitope with PD-1. The complete PD-1 sequence can be found under GenBank Accession No. U64863.

15

The terms "cytotoxic T lymphocyte-associated antigen-4," "CTLA-4," "CTLA4," and "CTLA-4 antigen" (see, e.g., Murata, Am. J. Pathol. (1999) 155:453-460) are used interchangeably, and include variants, isoforms, species homologs of human CTLA-4, and analogs having at least one common epitope with CTLA-4
20 (see, e.g., Balzano (1992) Int. J. Cancer Suppl. 7:28-32). The complete CTLA-4 nucleic acid sequence can be found under GenBank Accession No. L15006.

- As used herein, "sub-therapeutic dose" means a dose of a therapeutic compound (e.g., an antibody) or duration of therapy which is lower than the
25 usual or typical dose of the therapeutic compound or therapy of shorter duration, when administered alone for the treatment of cancer. For example, a sub-therapeutic dose of CTLA-4 antibody is a single dose of the antibody at less than about 3 mg/kg, i.e., the known dose of anti-CTLA-4 antibody.

- 30 The term "therapeutically effective amount" is defined as an amount of a checkpoint inhibitor, in combination with an immunomodulator, that preferably results in a decrease in severity of disease symptoms, an increase in frequency and duration of disease symptom-free periods, or a prevention of impairment or disability due to the disease affliction. The terms "effective amount" or

"pharmaceutically effective amount" refer to a sufficient amount of an agent to provide the desired biological or therapeutic result. That result can be reduction, amelioration, palliation, lessening, delaying, and/or alleviation of one or more of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. In reference to cancer, an effective amount may comprise an amount sufficient to cause a tumour to shrink and/or to decrease the growth rate of the tumour (such as to suppress tumour growth) or to prevent or delay other unwanted cell proliferation. In some embodiments, an effective amount is an amount sufficient to delay development, or prolong survival or induce stabilisation of the cancer or tumour.

In some embodiments, a therapeutically effective amount is an amount sufficient to prevent or delay recurrence. A therapeutically effective amount can be administered in one or more administrations. The therapeutically effective amount of the drug or combination may result in one or more of the following: (i) reduce the number of cancer cells; (ii) reduce tumour size; (iii) inhibit, retard, slow to some extent and preferably stop cancer cell infiltration into peripheral organs; (iv) inhibit (i.e., slow to some extent and preferably stop) tumour metastasis; (v) inhibit tumour growth; (vi) prevent or delay occurrence and/or recurrence of tumour; and/or (vii) relieve to some extent one or more of the symptoms associated with the cancer.

For example, for the treatment of tumours, a "therapeutically effective dosage" may induce tumour shrinkage by at least about 5 % relative to baseline measurement, such as at least about 10 %, or about 20 %, or about 60 % or more. The baseline measurement may be derived from untreated subjects.

A therapeutically effective amount of a therapeutic compound can decrease tumour size, or otherwise ameliorate symptoms in a subject. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

The term "immune response" refers to the action of, for example, lymphocytes, antigen presenting cells, phagocytic cells, granulocytes, and soluble macromolecules produced by the above cells or the liver (including antibodies, cytokines, and complement) that results in selective damage to, destruction of, or elimination from the human body of cancerous cells.

The term "antibody" as referred to herein includes whole antibodies and any antigen-binding fragment (i.e., "antigen-binding portion") or single chains thereof.

The term "antigen-binding portion" of an antibody (or simply "antibody portion"), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to a receptor and its ligand (e.g., PD-1). including: (i) a Fab fragment, (ii) a F(ab')₂ fragment,; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment, (v) a dAb fragment (Ward et al, Nature, 341 :544-546 (1989)), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are screened for utility in the same manner as are intact antibodies.

The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of single molecular composition. A monoclonal antibody composition displays a single binding specificity and affinity for a particular epitope.

The term "human antibody", as used herein, is intended to include antibodies having variable regions in which both the framework and CDR regions are derived from human germline immunoglobulin sequences.

30

The term "humanized antibody" is intended to refer to antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences. Additional

framework region modifications may be made within the human framework sequences.

5 In addition to antibodies, other biological molecules may act as checkpoint inhibitors, including peptides having binding affinity to the appropriate target.

10 The term "treatment" or "therapy" refers to administering an active agent with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect a condition (e.g., a disease), the symptoms of the condition, or to prevent or delay the onset of the symptoms, complications, biochemical indicia of a disease, or otherwise arrest or inhibit further development of the disease, condition, or disorder in a statistically significant manner.

15 As used herein, the term "subject" is intended to include human and non-human animals. Preferred subjects include human patients in need of enhancement of an immune response. The methods are particularly suitable for treating human patients having a disorder that can be treated by augmenting the T-cell mediated immune response. In a particular embodiment, the methods are particularly suitable for treatment of cancer cells in vivo.

20 As used herein, the terms "concurrent administration" or "concurrently" or "simultaneous" mean that administration occurs on the same day. The terms "sequential administration" or "sequentially" or "separate" mean that administration occurs on different days.

25 "Simultaneous" administration, as defined herein, includes the administration of the immunomodulator and agent or procedure comprising checkpoint inhibitor therapy within about 2 hours or about 1 hour or less of each other, even more preferably at the same time.

30 "Separate" administration, as defined herein, includes the administration of the immunomodulator and agent or procedure comprising checkpoint inhibitor therapy, more than about 12 hours, or about 8 hours, or about 6 hours or about 4 hours or about 2 hours apart.

“Sequential” administration, as defined herein, includes the administration of the immunomodulator and chemotherapeutic agent each in multiple aliquots and/or doses and/or on separate occasions. The immunomodulator may be administered to the patient after before and/or after administration of the checkpoint inhibitor. Alternatively, the immunomodulator is continued to be applied to the patient after treatment with a checkpoint inhibitor.

The use of the alternative (e.g., “or”) should be understood to mean either one, both, or any combination thereof of the alternatives. As used herein, the indefinite articles “a” or “an” should be understood to refer to “one or more” of any recited or enumerated component.

As used herein, “about” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. For example, “about” can mean within 1 or more than 1 standard deviation per the practice in the art. Alternatively, “about” can mean a range of up to 20%. When particular values are provided in the application and claims, unless otherwise stated, the meaning of “about” should be assumed to be within an acceptable error range for that particular value.

In one aspect of the present invention the immunomodulator comprises heat-killed *Mycobacterium*, preferably a whole cell *Mycobacterium*. Examples of mycobacterial species for use in the present invention include *M. vaccae*, *M. thermoresistibile*, *M. flavescens*, *M. duvalii*, *M. phlei*, *M. obuense*, *M. parafortuitum*, *M. sphagni*, *M. aichiense*, *M. rhodesiae*, *M. neoaurum*, *M. chubuense*, *M. tokaiense*, *M. komossense*, *M. aurum*, *M. w*, *M. tuberculosis*, *M. microti*; *M. africanum*; *M. kansasii*, *M. marinum*; *M. simiae*; *M. gastri*; *M. nonchromogenicum*; *M. terrae*; *M. triviale*; *M. gordonae*; *M. scrofulaceum*; *M. paraffinicum*; *M. intracellulare*; *M. avium*; *M. xenopi*; *M. ulcerans*; *M. diernhoferi*, *M. smegmatis*; *M. thamnopheos*; *M. flavescens*; *M. fortuitum*; *M. peregrinum*; *M. chelonae*; *M. paratuberculosis*; *M. leprae*; *M. lepraemurium* and combinations thereof.

Preferably, the heat-killed *Mycobacterium* is non-pathogenic. The non-pathogenic heat-killed *Mycobacterium* is preferably selected from *M. vaccae*, *M. obuense*, *M. parafortuitum*, *M. aurum*, *M. indicus pranii*, *M. phlei* and combinations thereof. More preferably the non-pathogenic heat-killed *Mycobacterium* is a rough variant. The amount of *Mycobacterium* administered to the patient is sufficient to elicit a protective immune response in the patient such that the patient's immune system is able to mount an effective immune response to the cancer or tumour. In certain embodiments of the invention, there is provided a containment means comprising the effective amount of heat-killed *Mycobacterium* for use in the present invention, which typically may be from 10^3 to 10^{11} organisms, preferably from 10^4 to 10^{10} organisms, more preferably from 10^6 to 10^{10} organisms, and even more preferably from 10^6 to 10^9 organisms. The effective amount of heat-killed *Mycobacterium* for use in the present invention may be from 10^3 to 10^{11} organisms, preferably from 10^4 to 10^{10} organisms, more preferably from 10^6 to 10^{10} organisms, and even more preferably from 10^6 to 10^9 organisms. Most preferably the amount of heat-killed *Mycobacterium* for use in the present invention is from 10^7 to 10^9 cells or organisms. Typically, the composition according to the present invention may be administered at a dose of from 10^8 to 10^9 cells for human and animal use. Alternatively the dose is from 0.01 mg to 1 mg or 0.1mg to 1mg organisms presented as either a suspension or dry preparation.

M. vaccae and *M. obuense* are particularly preferred.

M. vaccae and *M. obuense* induce a complex immune response in the host. Treatment with these preparations will stimulate innate and type-1 immunity, including Th1 and macrophage activation and cytotoxic cell activity. They also independently down-regulate inappropriate Th2 responses via immunoregulatory mechanisms. This restores the healthy balance of the immune system.

The present invention may be used to treat a neoplastic disease, such as solid or non-solid cancers. As used herein, "treatment" encompasses the prevention, reduction, control and/or inhibition of a neoplastic disease. Such diseases

include a sarcoma, carcinoma, adenocarcinoma, melanoma, myeloma, blastoma, glioma, lymphoma or leukemia. Exemplary cancers include, for example, carcinoma, sarcoma, adenocarcinoma, melanoma, neural (blastoma, glioma), mesothelioma and reticuloendothelial, lymphatic or haematopoietic

5 neoplastic disorders (e.g., myeloma, lymphoma or leukemia). In particular aspects, a neoplasm, tumour or cancer includes a lung adenocarcinoma, lung carcinoma, diffuse or interstitial gastric carcinoma, colon adenocarcinoma, prostate adenocarcinoma, esophagus carcinoma, breast carcinoma, pancreas adenocarcinoma, ovarian adenocarcinoma, adenocarcinoma of the adrenal

10 gland, adenocarcinoma of the endometrium or uterine adenocarcinoma.

Neoplasia, tumours and cancers include benign, malignant, metastatic and non-metastatic types, and include any stage (I, II, III, IV or V) or grade (G1, G2, G3, etc.) of neoplasia, tumour, or cancer, or a neoplasia, tumour, cancer or

15 metastasis that is progressing, worsening, stabilized or in remission. Cancers that may be treated according to the invention include but are not limited to cells or neoplasms of the bladder, blood, bone, bone marrow, brain, breast, colon, esophagus, gastrointestinal, gum, head, kidney, liver, lung, nasopharynx, neck, ovary, prostate, skin, stomach, testis, tongue, or uterus. In addition, the cancer

20 may specifically be of the following histological type, though it is not limited to the following: neoplasm, malignant; carcinoma; carcinoma, undifferentiated; giant and spindle cell carcinoma; small cell carcinoma; papillary carcinoma; squamous cell carcinoma; lymphoepithelial carcinoma; basal cell carcinoma; pilomatrix carcinoma; transitional cell carcinoma; papillary transitional cell carcinoma;

25 adenocarcinoma; gastrinoma, malignant; cholangiocarcinoma; hepatocellular carcinoma; combined hepatocellular carcinoma and cholangiocarcinoma; trabecular adenocarcinoma; adenoid cystic carcinoma; adenocarcinoma in adenomatous polyp; adenocarcinoma, familial polyposis coli; solid carcinoma; carcinoid tumour, malignant; bronchiolo-alveolar adenocarcinoma; papillary

30 adenocarcinoma; chromophobe carcinoma; acidophil carcinoma; oxyphilic adenocarcinoma; basophil carcinoma; clear cell adenocarcinoma; granular cell carcinoma; follicular adenocarcinoma; papillary and follicular adenocarcinoma; nonencapsulating sclerosing carcinoma; adrenal cortical carcinoma; endometroid carcinoma; skin appendage carcinoma; apocrine adenocarcinoma;

sebaceous adenocarcinoma; ceruminous adenocarcinoma; mucoepidermoid carcinoma; cystadenocarcinoma; papillary cystadenocarcinoma; papillary serous cystadenocarcinoma; mucinous cystadenocarcinoma; mucinous adenocarcinoma; signet ring cell carcinoma; infiltrating duct carcinoma; 5 medullary carcinoma; lobular carcinoma; inflammatory carcinoma; Paget's disease, mammary; acinar cell carcinoma; adenosquamous carcinoma; adenocarcinoma with squamous metaplasia; thymoma, malignant; ovarian stromal tumour, malignant; thecoma, malignant; granulosa cell tumour, malignant; androblastoma, malignant; Sertoli cell carcinoma; Leydig cell tumour, 10 malignant; lipid cell tumour, malignant; paraganglioma, malignant; extra-mammary paraganglioma, malignant; pheochromocytoma; glomangiosarcoma; malignant melanoma; amelanotic melanoma; superficial spreading melanoma; malignant melanoma in giant pigmented nevus; epithelioid cell melanoma; blue nevus, malignant; sarcoma; fibrosarcoma; fibrous histiocytoma, malignant; 15 myxosarcoma; liposarcoma; leiomyosarcoma; rhabdomyosarcoma; embryonal rhabdomyosarcoma; alveolar rhabdomyosarcoma; stromal sarcoma; mixed tumour; Mullerian mixed tumour; nephroblastoma; hepatoblastoma; carcinosarcoma; mesenchymoma, malignant; Brenner tumour, malignant; phyllodes tumour, malignant; synovial sarcoma; mesothelioma, malignant; 20 dysgerminoma; embryonal carcinoma; teratoma, malignant; struma ovarii, malignant; choriocarcinoma; mesonephroma, malignant; hemangiosarcoma; hemangioendothelioma, malignant; Kaposi's sarcoma; hemangiopericytoma, malignant; lymphangiosarcoma; osteosarcoma; juxtacortical osteosarcoma; chondrosarcoma; chondroblastoma, malignant; mesenchymal chondrosarcoma; 25 giant cell tumour of bone; Ewing's sarcoma; odontogenic tumour, malignant; ameloblastic odontosarcoma; ameloblastoma, malignant; ameloblastic fibrosarcoma; pinealoma, malignant; chordoma; glioma, malignant; ependymoma; astrocytoma; protoplasmic astrocytoma; fibrillary astrocytoma; astroblastoma; glioblastoma; oligodendroglioma; oligodendroblastoma; primitive 30 neuroectodermal; cerebellar sarcoma; ganglioneuroblastoma; neuroblastoma; retinoblastoma; olfactory neurogenic tumour; meningioma, malignant; neurofibrosarcoma; neurilemmoma, malignant; granular cell tumour, malignant; malignant lymphoma; Hodgkin's disease; Hodgkin's; paragranuloma; malignant lymphoma, small lymphocytic; malignant lymphoma, large cell, diffuse; malignant

lymphoma, follicular; mycosis fungoides; other specified non-Hodgkin's lymphomas; malignant histiocytosis; multiple myeloma; mast cell sarcoma; immunoproliferative small intestinal disease; leukemia; lymphoid leukemia; plasma cell leukemia; erythroleukemia; lymphosarcoma cell leukemia; myeloid
5 leukemia; basophilic leukemia; eosinophilic leukemia; monocytic leukemia; mast cell leukemia; megakaryoblastic leukemia; myeloid sarcoma; and hairy cell leukemia. Preferably, the neoplastic disease may be tumours associated with a cancer selected from prostate cancer, liver cancer, renal cancer, lung cancer, breast cancer, colorectal cancer, pancreatic cancer, brain cancer, hepatocellular
10 cancer, lymphoma, leukaemia, gastric cancer, cervical cancer, ovarian cancer, thyroid cancer, melanoma, head and neck cancer, skin cancer and soft tissue sarcoma and/or other forms of carcinoma. The tumour may be metastatic or a malignant tumour.

15 More preferably, the neoplastic disease to be treated is pancreatic cancer, breast cancer, lung cancer, prostate cancer and skin cancer. Most preferably, the neoplastic disease to be treated is pancreatic cancer.

In an embodiment of the invention, the checkpoint inhibitor therapy, in
20 combination therapy with an immunomodulator, most preferably a whole cell, heat-killed *Mycobacterium*, is used to reduce or inhibit metastasis of a primary tumour or cancer to other sites, or the formation or establishment of metastatic tumours or cancers at other sites distal from the primary tumour or cancer thereby inhibiting or reducing tumour or cancer relapse or tumour or cancer
25 progression.

In a further embodiment of the invention, there is provided a combination therapy for treating cancer, comprising an immunomodulator and blockade of checkpoint inhibitors with the potential to elicit potent and durable immune responses with
30 enhanced therapeutic benefit and more manageable toxicity.

In a further embodiment of the invention, there is provided a combination therapy for treating cancer, comprising an immunomodulator which; (i) stimulates innate and type-1 immunity, including Th1 and macrophage activation and cytotoxic cell

activity, and, (ii) independently down-regulates inappropriate Th2 responses via immunoregulatory mechanisms; and, blockade of a checkpoint inhibitor, optionally wherein the immunomodulator is a whole cell *Mycobacterium* selected from *M. vaccae* or *M. obuense*.

5

In an embodiment of the invention is provided a method for treating cancer and/or preventing the establishment of metastases by employing a checkpoint inhibitor which act synergistically with a whole cell *Mycobacterium*.

- 10 In further embodiments, methods of the invention include, one or more of the following: 1) reducing or inhibiting growth, proliferation, mobility or invasiveness of tumour or cancer cells that potentially or do develop metastases, 2) reducing or inhibiting formation or establishment of metastases arising from a primary tumour or cancer to one or more other sites, locations or regions distinct from
- 15 the primary tumour or cancer; 3) reducing or inhibiting growth or proliferation of a metastasis at one or more other sites, locations or regions distinct from the primary tumour or cancer after a metastasis has formed or has been established, 4) reducing or inhibiting formation or establishment of additional metastasis after the metastasis has been formed or established, 5) prolonged
- 20 overall survival, 6) prolonged progression free survival, or 7) disease stabilisation.

In an embodiment of the invention, administration of the checkpoint inhibitor therapy, in combination therapy with an immunomodulator, preferably a whole

25 cell, heat-killed *Mycobacterium*, provides a detectable or measurable improvement in a condition of a given subject, such as alleviating or ameliorating one or more adverse (physical) symptoms or consequences associated with the presence of a cell proliferative or cellular hyperproliferative disorder, neoplasia, tumour or cancer, or metastasis, i e., a therapeutic benefit or a beneficial effect.

30

A therapeutic benefit or beneficial effect is any objective or subjective, transient, temporary, or long-term improvement in the condition or pathology, or a reduction in onset, severity, duration or frequency of an adverse symptom associated with or caused by cell proliferation or a cellular hyperproliferative

disorder such as a neoplasia, tumour or cancer, or metastasis. It may lead to improved survival. A satisfactory clinical endpoint of a treatment method in accordance with the invention is achieved, for example, when there is an incremental or a partial reduction in severity, duration or frequency of one or more associated pathologies, adverse symptoms or complications, or inhibition or reversal of one or more of the physiological, biochemical or cellular manifestations or characteristics of cell proliferation or a cellular hyperproliferative disorder such as a neoplasia, tumour or cancer, or metastasis. A therapeutic benefit or improvement therefore may be, but is not limited to destruction of target proliferating cells (e.g., neoplasia, tumour or cancer, or metastasis) or ablation of one or more, most or all pathologies, adverse symptoms or complications associated with or caused by cell proliferation or the cellular hyperproliferative disorder such as a neoplasia, tumour or cancer, or metastasis. However, a therapeutic benefit or improvement need not be a cure or complete destruction of all target proliferating cells (e.g., neoplasia, tumour or cancer, or metastasis) or ablation of all pathologies, adverse symptoms or complications associated with or caused by cell proliferation or the cellular hyperproliferative disorder such as a neoplasia, tumour or cancer, or metastasis. For example, partial destruction of a tumour or cancer cell mass, or a stabilization of the tumour or cancer mass, size or cell numbers by inhibiting progression or worsening of the tumour or cancer, can reduce mortality and prolong lifespan even if only for a few days, weeks or months, even though a portion or the bulk of the tumour or cancer mass, size or cells remain.

Specific non-limiting examples of therapeutic benefit include a reduction in neoplasia, tumour or cancer, or metastasis volume (size or cell mass) or numbers of cells, inhibiting or preventing an increase in neoplasia, tumour or cancer volume (e.g., stabilizing), slowing or inhibiting neoplasia, tumour or cancer progression, worsening or metastasis, or inhibiting neoplasia, tumour or cancer proliferation, growth or metastasis.

In an embodiment of the invention, administration of the checkpoint inhibitor, in combination therapy with an immunomodulator, preferably a whole cell, heat-killed *Mycobacterium*, provides a detectable or measurable improvement or

overall response according to the irRC (as derived from time-point response assessments and based on tumour burden), including one of more of the following: (i) irCR - complete disappearance of all lesions, whether measurable or not, and no new lesions (confirmation by a repeat, consecutive assessment
5 no less than 4 weeks from the date first documented), (ii) irPR - decrease in tumour burden $\geq 50\%$ relative to baseline (confirmed by a consecutive assessment at least 4 weeks after first documentation).

10 An invention method may not take effect immediately. For example, treatment may be followed by an increase in the neoplasia, tumour or cancer cell numbers or mass, but over time eventual stabilization or reduction in tumour cell mass, size or numbers of cells in a given subject may subsequently occur.

15 Additional adverse symptoms and complications associated with neoplasia, tumour, cancer and metastasis that can be inhibited, reduced, decreased, delayed or prevented include, for example, nausea, lack of appetite, lethargy, pain and discomfort. Thus, a partial or complete decrease or reduction in the severity, duration or frequency of an adverse symptom or complication associated with or caused by a cellular hyperproliferative disorder, an
20 improvement in the subjects quality of life and/or well-being, such as increased energy, appetite, psychological well-being, are all particular non-limiting examples of therapeutic benefit.

25 A therapeutic benefit or improvement therefore can also include a subjective improvement in the quality of life of a treated subject. In an additional embodiment, a method prolongs or extends lifespan (survival) of the subject. In a further embodiment, a method improves the quality of life of the subject.

30 In a most preferred embodiment, administration of the checkpoint inhibitor, in combination therapy with an immunomodulator, preferably a whole cell, heat-killed *Mycobacterium* results in a clinically relevant improvement in one or more markers of disease status and progression selected from one or more of the following: (i): overall survival, (ii): progression-free survival, (iii): overall response rate, (iv): reduction in metastatic disease, (v): circulating levels of tumour

antigens such as carbohydrate antigen 19.9 (CA19.9) and carcinoembryonic antigen (CEA) or others depending on tumour, (vii) nutritional status (weight, appetite, serum albumin), (viii): pain control or analgesic use, (ix): CRP/albumin ratio.

5

Treatment with heat-killed whole cell *M. vaccae* and *M. obuense* gives rise to more complex immunity including not only the development of innate immunity and type-1 immunity, but also immunoregulation which more efficiently restores appropriate immune functions.

10

The immunomodulator according to the invention is administered in combination with a checkpoint inhibitor.

In a preferred embodiment, the checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against CTLA-4, PD-1, PD-L1, TIM-3, BTLA, VISTA, LAG-3 and combinations thereof.

In another preferred embodiment, the checkpoint inhibition therapy comprises administration of a sub-therapeutic amount and/or duration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against CTLA-4, PD-1, PD-L1, TIM-3, BTLA, VISTA, LAG-3 and combinations thereof.

In a further preferred embodiment, the checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against PD-1 and/or, PD-L1.

In a further preferred embodiment, the checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against PD-1 and/or, PD-L1, simultaneously, separately or sequentially with administration of a blocking antibody or antigen binding fragment thereof, directed against CTLA-4.

The term "combination" as used throughout the specification, is meant to encompass the administration of the checkpoint inhibitor simultaneously, separately or sequentially with administration of the *Mycobacterium*.
5 Accordingly, the checkpoint inhibitor and the *Mycobacterium* may be present in the same or separate pharmaceutical formulations, and administered at the same time or at different times.

Thus, a *Mycobacterium* and the checkpoint inhibitor may be provided as
10 separate medicaments for administration at the same time or at different times.

Preferably, a *Mycobacterium* and checkpoint inhibitor are provided as separate medicaments for administration at different times. When administered separately and at different times, either the *Mycobacterium* or checkpoint inhibitor may be
15 administered first; however, it is suitable to administer checkpoint inhibitor followed by the *Mycobacterium*. In addition, both can be administered on the same day or at different days, and they can be administered using the same schedule or at different schedules during the treatment cycle.

20 In an embodiment of the invention, a treatment cycle consists of the administration of a *Mycobacterium* daily, weekly fortnightly or monthly, simultaneously with checkpoint inhibitor weekly. Alternatively, the *Mycobacterium* is administered before and/or after the administration of the checkpoint inhibitor.

25 In another embodiment of the invention, the *Mycobacterium* is administered to the patient before and after administration of a checkpoint inhibitor. That is, in one embodiment, the immunomodulator is administered to the patient before and after checkpoint inhibitor.

30 Dose delays and/ or dose reductions and schedule adjustments are performed as needed depending on individual patient tolerance to treatments.

Alternatively, the administration of checkpoint inhibitor may be performed simultaneously with the administration of the effective amounts of the *Mycobacterium*.

- 5 The immunomodulator, preferably a whole cell heat-killed *Mycobacterium*, may be administered to the patient via the parenteral, oral, sublingual, nasal or pulmonary route. In a preferred embodiment, the immunomodulator is administered via a parenteral route selected from subcutaneous, intradermal, subdermal, intraperitoneal, intravenous and intravesicular injection. More
10 preferably, administration by the parenteral route does not comprise intratumoural injection of mycobacterial cell wall extract.

The subject whom is to undergo checkpoint inhibition therapy according to the present invention may do so simultaneously, separately or sequentially with
15 administration of the immunomodulator, preferably a whole cell heat-killed *Mycobacterium*.

In an aspect of the invention, the effective amount of the immunomodulator may be administered as a single dose. Alternatively, the effective amount of the
20 immunomodulator may be administered in multiple (repeat) doses, for example two or more, three or more, four or more, five or more, ten or more, or twenty or more repeat doses. The immunomodulator may be administered between about 4 weeks and about 1 day prior to checkpoint inhibition therapy, such as between about 4 weeks and 1 week, or about between 3 weeks and 1 week, or about
25 between 3 weeks and 2 weeks. Administration may be presented in single or multiple doses.

In a preferred embodiment of the invention there is a *Mycobacterium* for use in the treatment of neoplastic disease in combination with a checkpoint inhibitor
30 wherein said *Mycobacterium* is administered to the subject before, concurrently with and/or after the checkpoint inhibitor is administered.

In another preferred embodiment of the invention is a method of treating, reducing, inhibiting or controlling a neoplasia, tumour or cancer in a subject,

wherein said method comprises simultaneously, separately or sequentially administering to the subject, (i) a checkpoint inhibitor, and (ii) an immunomodulator, preferably a whole cell heat-killed *Mycobacterium*, wherein said method results in enhanced therapeutic efficacy relative to administration of
5 the checkpoint inhibitor or immunomodulator alone.

In another preferred embodiment of the invention is a method of treating, reducing, inhibiting or controlling a neoplasia, tumour or cancer in a subject, wherein said method comprises simultaneously, separately or sequentially
10 administering to the subject, (i) a checkpoint inhibitor, and (ii) an immunomodulator, preferably a whole cell heat-killed *Mycobacterium*, wherein said checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against CTLA-4, PD-1, PD-L1, TIM-3, BTLA, VISTA, LAG-3 and
15 combinations thereof.

In another preferred embodiment of the invention is a method of treating, reducing, inhibiting or controlling a neoplasia, tumour or cancer in a subject, wherein said method comprises simultaneously, separately or sequentially
20 administering to the subject, (i) a checkpoint inhibitor, and (ii) an immunomodulator, preferably a whole cell heat-killed *Mycobacterium*, wherein said checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against PD-1 and/or PD-L1.

25

In yet another preferred embodiment of the invention is a method of treating, reducing, inhibiting or controlling a neoplasia, tumour or cancer in a subject, wherein said method comprises simultaneously, separately or sequentially administering to the subject, (i) a checkpoint inhibitor, and (ii) an
30 immunomodulator, preferably a whole cell heat-killed *Mycobacterium*, wherein said checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against CTLA-4, PD-1, PD-L1, TIM-3, BTLA, VISTA, LAG-3 and combinations thereof, wherein said checkpoint inhibition therapy comprises

administration of a sub-therapeutic amount and/or duration of said blocking antibody or antigen binding fragment thereof.

In yet another embodiment of the invention is a method of treating, reducing, inhibiting or controlling a neoplasia, tumour or cancer in a subject, wherein said method comprises simultaneously, separately or sequentially administering to the subject, (i) two or more checkpoint inhibitors, and (ii) an immunomodulator, preferably a whole cell heat-killed *Mycobacterium*, wherein said checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against CTLA-4, PD-1, PD-L1, TIM-3, BTLA, VISTA, LAG-3 and combinations thereof, wherein said checkpoint inhibition therapy optionally comprises administration of a sub-therapeutic amount and/or duration of said blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof.

In one embodiment of the present invention, the *Mycobacterium* may be in the form of a medicament administered to the patient in a dosage form.

A container according to the invention in certain instances, may be a vial, an ampoule, a syringe, capsule, tablet or a tube. In some cases, the mycobacteria may be lyophilized and formulated for resuspension prior to administration. However, in other cases, the mycobacteria are suspended in a volume of a pharmaceutically acceptable liquid. In some of the most preferred embodiments there is provided a container comprising a single unit dose of mycobacteria suspended in pharmaceutically acceptable carrier wherein the unit dose comprises about 1×10^6 to about 1×10^{10} organisms. In some very specific embodiments the liquid comprising suspended mycobacteria is provided in a volume of between about 0.1 ml and 10 ml, or between about 0.3 ml and 2ml or between about 0.5 ml and 2 ml. The foregoing compositions provide ideal units for immunotherapeutic applications described herein.

Embodiments discussed in the context of a methods and/or composition of the invention may be employed with respect to any other method or composition

described herein. Thus, an embodiment pertaining to one method or composition may be applied to other methods and compositions of the invention as well.

5 In some cases non-pathogenic heat-killed mycobacteria are administered to specific sites on or in a subject. For example, the mycobacterial compositions according to the invention, such as those comprising *M. obuense* in particular, may be administered adjacent to tumours or adjacent to lymph nodes, such as those that drain tissue surrounding a tumour. Thus, in certain instances sites administration of mycobacterial composition may be near the posterior cervical,
10 tonsillar, axillary, inguinal, anterior cervical, sub-mandibular, sub mental or superclavicular lymph nodes.

The immunomodulator may be administered for the length of time the cancer or tumour(s) is present in a patient or until such time the cancer has regressed or
15 stabilized. The immunomodulator may also be continued to be administered to the patients once the cancer or tumour has regressed or stabilised.

Mycobacterial compositions according to the invention will comprise an effective amount of mycobacteria typically dispersed in a pharmaceutically acceptable
20 carrier. The phrases "pharmaceutically or pharmacologically acceptable" refers to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, such as, for example, a human, as appropriate. The preparation of an pharmaceutical composition that contains mycobacteria will be known to those of skill in the art
25 in light of the present disclosure, as exemplified by Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, Moreover, for animal (e.g., human) administration, it will be understood that preparations should meet sterility, pyrogenicity, general safety and purity standards. A specific example of a pharmacologically acceptable carrier as described herein is borate buffer or
30 sterile saline solution (0.9% NaCl).

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, surfactants, antioxidants, preservatives {e.g., antibacterial agents, antifungal agents}, isotonic agents, absorption

delaying agents, salts, preservatives, drugs, drug stabilizers, gels, binders, excipients, disintegration agents, lubricants, sweetening agents, flavouring agents, dyes, such like materials and combinations thereof, as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, pp. 1289-1329).

In a preferred embodiment, the immunomodulator is administered via a parenteral route selected from subcutaneous, intradermal, subdermal, intraperitoneal, intravenous and intravesicular injection. Intradermal injection enables delivery of an entire proportion of the mycobacterial composition to a layer of the dermis that is accessible to immune surveillance and thus capable of electing an anti-cancer immune response and promoting immune cell proliferation at local lymph nodes.

Though in highly preferred embodiments of the invention mycobacterial compositions are administered by direct intradermal injection, it is also contemplated that other methods of administration may be used in some case. Thus in certain instances heat-killed mycobacteria of the present invention can be administered by injection, infusion, continuous infusion, intravenously, intradermally, intraarterially, intraperitoneally, intralesionally, intravitreally, intravaginally, intrarectally, topically, intratumourally, intramuscularly, intraperitoneally, subcutaneously, subconjunctival, intravesicularly, mucosally, intrapericardially, intraumbilically, intraocularly, orally, intracranially, intraarticularly, intraprostatically, intrapleurally, intratracheally, intranasally, topically, locally, inhalation (e.g. aerosol inhalation), via a catheter, via a lavage, or by other method or any combination of the foregoing as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990). More preferably, administration by the parenteral route does not comprise intratumoural injection of mycobacterial cell wall extract.

All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described methods and system of the present invention will be apparent to those skilled in the art without

departing from the scope and spirit of the present invention. Although the present invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in biochemistry and immunology or related fields are intended to be within the scope of the following claims.

The invention is further described with reference to the following non-limiting Example.

Example 1

Adult C57BL/6 mice were injected subcutaneously on the flank with 10^5 cells from a pancreatic cancer cell line obtained from KPC mice (Hingorani *et al.* Cancer Cell, 2005, 7:469-48). These murine pancreatic cancer cells bear mutations in Kras, p53 and Pdx-Cre (Hingorani *et al.* Cancer Cell, 2005, 7:469-48).

When the injected tumour cells had grown to become a palpable tumour (day 0), mice were left untreated or received treatment with:

- 1) 0.1mg *M. obuense* NTCT 13365/mouse, subcutaneously alternating injections in the scruff of the neck with those at the base of the tail on alternating days over 5 day period with a 2 day break for the length of the study;
- 2) 10mg/kg anti-PDL-1 mAb intraperitoneally once weekly;
- 3) the combination of anti-PDL-1 and *M. obuense* NTCT 13365 at a dose and schedule described above for the two compounds used singly.

Tumour growth was monitored over the course of the study to determine whether the treatment administered had an effect on reducing tumour size and improving prospects of survival.

Data presented in Figure 1 show that mice which received the treatment combination of anti-PDL-1 and *M. obuense* NTCT 13365 demonstrated a

continued reduction in tumour size and appeared to control the tumour. This reduction in tumour size was more pronounced compared to mice receiving either treatment alone. Mice left untreated had uncontrolled tumour growth and soon succumbed to the disease.

CLAIMS

1. An immunomodulator for use in the treatment, reduction, inhibition or control of a neoplastic disease in a patient intended to undergo checkpoint inhibition therapy simultaneously, separately or sequentially with administration of the immunomodulator.
2. The immunomodulator according to claim 1, wherein said checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against CTLA-4, PD-1, PD-L1, TIM-3, BTLA, VISTA, LAG-3 and combinations thereof.
3. The immunomodulator according to claim 1 or claim 2, wherein said checkpoint inhibition therapy comprises administration of a sub-therapeutic amount and/or duration of said blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof.
4. The immunomodulator according to any of claims 1 to 3, wherein said neoplastic disease is associated with a cancer selected from prostate cancer, liver cancer, renal cancer, lung cancer, breast cancer, colorectal cancer, breast cancer, pancreatic cancer, brain cancer, hepatocellular cancer, lymphoma, leukaemia, gastric cancer, cervical cancer, ovarian cancer, thyroid cancer, melanoma, carcinoma, head and neck cancer, skin cancer and soft tissue sarcoma.
5. The immunomodulator according to claim 4, wherein said neoplastic disease is associated with pancreatic cancer.
6. The immunomodulator according to any of claims 1 to 5, wherein said neoplastic disease is metastatic.
7. The immunomodulator according to any of claims 1 to 6, wherein the immunomodulator comprises a whole cell *Mycobacterium*.

8. The immunomodulator according to claim 7, wherein the *Mycobacterium* is a non-pathogenic heat-killed *Mycobacterium*.
- 5 9. The immunomodulator according to claim 8, wherein the non-pathogenic heat-killed *Mycobacterium* is selected from *M. vaccae*, *M. obuense*, *M. parafortuitum*, *M. aurum*, *M. indicus pranii* and combinations thereof.
- 10 10. The immunomodulator according to any of claims 8 or 9, wherein the nonpathogenic heat-killed *Mycobacterium* is preferentially the rough variant.
11. The immunomodulator according to any of claims 8 to 10, wherein the nonpathogenic heat-killed *Mycobacterium* is for administration via the parenteral, oral, sublingual, nasal or pulmonary route.
- 15 12. The immunomodulator according to claim 11, wherein the parenteral route is selected from subcutaneous, intradermal, subdermal, intraperitoneal, or intravenous.
- 20 13. The immunomodulator according to claim 12, wherein the parenteral route does not comprise intratumoural injection.
14. The immunomodulator according to any preceding claim, wherein said checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against PD-L1 or PD-1.
- 25 15. The immunomodulator according to any preceding claim, wherein administration of said immunomodulator is prior to and/or after the checkpoint inhibition therapy.
- 30 16. A method of treating, reducing, inhibiting or controlling a neoplasia, tumour or cancer in a subject, wherein said method comprises simultaneously, separately or sequentially administering to the subject, (i) a checkpoint inhibitor,

and (ii) an immunomodulator, wherein said method results in enhanced therapeutic efficacy relative to administration of the checkpoint inhibitor or immunomodulator alone.

5 17. A method according to claim 16, wherein said checkpoint inhibition therapy comprises administration of a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against CTLA-4, PD-1, PD-L1, TIM-3, BTLA, VISTA, LAG-3 and combinations thereof.

10 18. A method according to claim 16 or 17, wherein said checkpoint inhibitor is a blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof, directed against PD-1 or PD-L1.

15 19. A method according to any of claims 16 to 18, wherein said checkpoint inhibition therapy comprises administration of a sub-therapeutic amount of said blocking agent, selected from a cell, protein, peptide, antibody or antigen binding fragment thereof.

20 20. A method according to any of claims 16 to 19, wherein said neoplasia, tumour or cancer is associated with a cancer selected from prostate cancer, liver cancer, renal cancer, lung cancer, breast cancer, colorectal cancer, breast cancer, pancreatic cancer, brain cancer, hepatocellular cancer, lymphoma, leukaemia, gastric cancer, cervical cancer, ovarian cancer, thyroid cancer, melanoma, carcinoma, head and neck cancer, skin cancer and soft tissue
25 sarcoma.

21. A method according to 20, wherein said neoplasia, tumour or cancer is associated with pancreatic cancer.

30 22. A method according to any of claims 16 to 21, wherein the neoplasia, tumour or cancer is metastatic.

23. A method according to any of claims 16 to 22, wherein the immunomodulator comprises a whole cell *Mycobacterium*.

24. A method according to claim 23, wherein the *Mycobacterium* is a non-pathogenic heat-killed *Mycobacterium*.

5 25. A method according to claim 24, wherein the non-pathogenic heat-killed *Mycobacterium* is selected from *M. vaccae*, *M. obuense*, *M. parafortuitum*, *M. aurum*, *M. indicus pranii* and combinations thereof.

26. The method according to claim 24 or 25, wherein the non-pathogenic
10 heat-killed *Mycobacterium* is the rough variant.

27. A method according to claim 25, wherein the non-pathogenic heat-killed *Mycobacterium* is administered via the parenteral, oral, sublingual, nasal or pulmonary route.

15

28. A method according to claim 27, wherein the parenteral route is selected from subcutaneous, intradermal, subdermal, intraperitoneal, or intravenous.

29. A method according to claim 27 or 28, wherein the parenteral route does
20 not comprise intratumoural injection.

30. A method according to any of claims 24 to 29, wherein the amount of non-pathogenic heat-killed *Mycobacterium* administered is from 10^7 to 10^9 cells.

25 31. A method according to claim 16, wherein enhanced therapeutic efficacy is measured by increased overall survival time.

32. A method according to claim 16, wherein enhanced therapeutic efficacy is measured by increased progression-free survival.

30

33. A method according to claim 16, wherein enhanced therapeutic efficacy is measured by a decrease or stabilisation of tumour size.

34. A method according to claim 16, wherein enhanced therapeutic efficacy is measured by increased quality of life.