



CLAIMS

What is claimed is:


1. A compound having the formula:



Ring A is a ring system selected from



Ring B is a ring system selected from



wherein Ring B is optionally substituted with up to 4 fluorine atoms, up to two OH or up to two C_{1-4} alkyl which is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC_{1-4} alkyl groups;

Ring C is a cyclohexane or a cyclobutane ring;

X is $-NH-$, $-O-$, or $-OC_{1-4}$ alkyl-;

each of R^1 and R^2 is, independently, hydrogen, $-C(O)NHR^4$, $-C(O)OR^4$, $-NHC(O)R^4$, $-NHC(O)OR^4$, $NHC(O)NHR^4$, $-NHS(O)_2R^4$, $-C_{0-4}$ alkyl- NHR^4 , or $-OR^4$, wherein R^1 and R^2 cannot simultaneously be hydrogen, and wherein R^1 and R^2 and the intervening carbon atom can form a dioxane or dioxolane ring;


R^3 is hydrogen, $-C_{1-4}$ alkyl, fluoro, chloro, $-OC_{1-2}$ alkyl, $-C(O)H$, $-C(O)OH$, $-C(O)OC_{1-2}$ alkyl, $-CN$, $-C(O)NHC_{1-2}$ alkyl, or $-C(O)NH_2$, wherein each of said R^3 alkyl is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC_{1-2} alkyl groups;

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole,


pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, or quinoline, or a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, or tetrahydropyridopyrimidine, wherein each of said R⁴ groups is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl, up to three CN, NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂, C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and

each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, a spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


2. The compound of claim 1, wherein Ring C is cyclobutane.
3. The compound of claim 2, having the following formula:




4. The compound of claim 3, having the following formula:




5. The compound of claim 4, having the following formula:




6. The compound of claim 4 or 5, wherein R^2 is $-C_{0-4} \text{ alkyl-NHR}^4$.
7. The compound of claim 3, having the following formula:




8. The compound of claim 7, having the following formula:



9. The compound of claim 7 or 8, wherein R^1 is $-C_{0-4}$ alkyl-NHR⁴.
10. The compound of any one of claims 2-9, wherein X is $-O-$ or $-OC_{1-4}$ alkyl-.
11. The compound of any one of claims 2-10 wherein



12. The compound of claim 11 wherein



13. The compound of any one of claims 2-12, wherein R^3 is hydrogen.

14. The compound of any one of claims 2-13, wherein

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-C(O)NH₂, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}alkyl-C_{3-5}cycloalkyl$; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

15. The compound of any one of claims 2-13, wherein

R^4 is a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, or quinoline, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl,

C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}alkyl-C_{3-5}cycloalkyl$; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


16. The compound of any one of claims 2-13, wherein R^4 is a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, or tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , $C_{2-4}alkenyl$, $C_{2-4}alkynyl$, $C_{3-6}cycloalkyl$, $C_{0-4}alkyl-C_{3-5}cycloalkyl$, $C_{0-4}alkyl-O-C_{1-4}alkyl$, $C_{0-4}alkyl-O-C_{0-4}alkyl-C_{3-5}cycloalkyl$, $C(O)OC_{1-4}alkyl$, $C(O)OC_{0-4}alkyl-C_{3-5}cycloalkyl$, $C_{0-4}alkyl-C(O)NH_2$, $C(O)NHC_{1-4}alkyl$, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine,

pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{1-4}$ alkyl- C_{3-5} cycloalkyl; and

each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


17. The compound of claim 1, wherein Ring C is cyclohexane.

18. The compound of claim 17, having the following formula:




19. The compound of claim 17 or 18, wherein X is $-NH-$.

20. The compound of any one of claims 17-19, having the following formula:



21. The compound of claim 20, having the following formula:




22. The compound of claim 20 or 21, wherein R^2 is $-C_{0-4}$ alkyl-NHR⁴ or $-OR^4$.


23. The compound of claim 22, wherein R^2 is $-NHR^4$.

24. The compound of claim 22, wherein R^2 is $-OR^4$.

25. The compound of claim 17-19, having the following formula:



26. The compound of claim 25 having the following formula:




27. The compound of claim 25 or 26, wherein R^1 is $-C_{0-4}$ alkyl-NHR⁴ or $-OR^4$.


28. The compound of claim 27, wherein R^1 is $-NHR^4$.

29. The compound of claim 27, wherein R^1 is $-OR^4$.

30. The compound of any one of claims 17-29 wherein




31. The compound of claim 30 wherein




32. The compound of claim 17 or 18, wherein X is $-O-$.


33. The compound of claim 17, 18 or 32, having the following formula:




34. The compound of claim 33, having the following formula:



35. The compound of claim 33 or 34, wherein R^2 is $-C_{0-4}$ alkyl-NHR⁴ or -OR⁴.
36. The compound of claim 35, wherein R^2 is -NHR⁴.
37. The compound of claim 35, wherein R^2 is -OR⁴.
38. The compound of claim 17, 18 or 32, having the following formula:




39. The compound according to claim 38 having the following formula:




40. The compound of claim 38 or 39, wherein R^1 is $-C_{0-4}$ alkyl-NHR⁴ or -OR⁴.
41. The compound of claim 40, wherein R^1 is -NHR⁴.
42. The compound of claim 40, wherein R^1 is -OR⁴.

43. The compound of claim 38 or 39 having the following formula:




Y is -O- or -NH-.

44. The compound of any one of claims 32-43 wherein



45. The compound of claim 44 wherein



46. The compound of any one of claims 17-45, wherein R^3 is hydrogen, C_{1-4} alkyl, OC_{1-2} alkyl, or $C(O)NH_2$, or $C(O)H$, wherein each of said R^3 alkyl is optionally substituted with OH.

47. The compound of claim 46, wherein R^3 is hydrogen.

48. The compound of any one of claims 17-47, wherein

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl,

$C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$, $C(O)NH(C_{0-4}$ alkyl- C_{3-5} cycloalkyl), CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$,
 C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl- $NHC(O)R^5$, C_{0-4} alkyl- $N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and
 each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

49. The compound of any one of claims 17-47, wherein

R^4 is a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, or quinoline, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$, $C(O)NH(C_{0-4}$ alkyl- C_{3-5} cycloalkyl), CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$, C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl- $NHC(O)R^5$, C_{0-4} alkyl- $N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from

furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two

SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and

each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, a spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

50. The compound of claim 49, wherein

R⁴ is pyridine or pyrimidine, which is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl, up to three CN, NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂,

C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two

SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and


each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl

selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

51. The compound of any one of claims 17-47, wherein

R^4 is a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuroypyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, or tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}alkyl-C_{3-5}cycloalkyl$; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


52. The compound of claim 32 having the formula:



(IV), wherein

R^3 is hydrogen, $-C_{1-4}$ alkyl, fluoro, chloro, $-OC_{1-2}$ alkyl, or $-C(O)NH_2$, $-C(O)H$, or $-CN$, wherein each of said R^3 alkyl is optionally substituted with OH or up to 3 fluorine atoms;

R^4 is



X^1 is N, CH, CF, CCl, or CC_{1-2} alkyl optionally substituted with up to 3 fluorine atoms;
 X^2 is N or CR^{4c} , wherein X^1 and X^2 cannot simultaneously be N;
 each of R^{4a} , R^{4b} , and R^{4c} is, independently, hydrogen, F, Cl, Br, CN, NO_2 , C_{1-4} alkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $O-C_{1-4}$ alkyl, C_{0-4} alkyl- $O-C_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, or piperazine, or a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, or tetrazole, or R^{4c} , R^{4a} , and the intervening atoms form a dihydrofuran, a dihydropyran, or a tetrahydropiperidine heterocyclic ring system;
 wherein each of said R^{4a} , R^{4b} , or R^{4c} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl, up to two OH groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and
 wherein each of said R^{4a} , R^{4b} , or R^{4c} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.


53. The compound of claim 52, wherein R^3 is hydrogen, $-C_{1-4}$ alkyl, $-OC_{1-2}$ alkyl, or $-C(O)NH_2$, or $-C(O)H$, wherein each of said R^3 alkyl is optionally substituted with OH.
54. The compound of claim 52 or 53, wherein R^3 is hydrogen.
55. The compound of any one of claims 52-54, wherein each of X^1 and X^2 is, independently, CH or N, wherein X^1 and X^2 cannot simultaneously be N.
56. The compound of any one of claims 52-55, wherein each of R^{4a} and R^{4b} is, independently, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, or piperazine, wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl, up to two OH groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.
57. The compound of any one of claims 52-55, wherein each of R^{4a} and R^{4b} is, independently, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, or tetrazole, wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl, up to two OH groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.
58. The compound of any one of claims 52-55, wherein each of R^{4a} and R^{4b} is, independently, hydrogen, F, Cl, Br, CN, NO_2 , C_{1-4} alkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4} \text{ alkyl})_2$, or $C(O)NH(C_{0-4} \text{ alkyl-}C_{3-5} \text{ cycloalkyl})$,

wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.


59. A compound selected from the list of compounds in Table 1.
60. A compound selected from the list of compounds in Table 2.
61. A pharmaceutical composition comprising a compound of any one of claims 1-60 and a pharmaceutically acceptable excipient.
62. A method of sensitizing a cell to a therapeutic agent or a disease state that induces a DNA lesion comprising the step of contacting the cell with the compound of any one of claims 1-60, or a pharmaceutical composition comprising said compound.
63. A method of potentiating a therapeutic regimen for the treatment of cancer in a patient comprising the step of administering to said patient an effective amount of the compound of any one of claims 1-60, or a pharmaceutical composition comprising said compound.
64. A method of treating cancer or inhibiting cancer cell growth in a patient comprising administering to said patient an effective amount of the compound of any one of claims 1-60, or a pharmaceutical composition comprising said compound, either alone or in combination with one or more additional therapeutic agent.

Claims


1. A compound having the formula:



Ring A is a ring system selected from



Ring B is a ring system selected from



Ring B is optionally substituted with up to 4 fluorine atoms, up to two OH or up to two C₁₋₄alkyl which is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC₁₋₂alkyl groups;

Ring C is a cyclohexane or a cyclobutane ring;

X is -NH-, -O-, or -OC₁₋₄ alkyl-;


each of R¹ and R² is, independently, hydrogen, -C(O)NHR⁴, -C(O)OR⁴, -NHC(O)R⁴, -NHC(O)OR⁴, -NHC(O)NHR⁴, -NHS(O)₂R⁴, -C₀₋₄ alkyl-NHR⁴, or -OR⁴, wherein R¹ and R² cannot simultaneously be hydrogen, and wherein R¹ and R² and the intervening carbon atom can form a dioxane or dioxolane ring;

R³ is hydrogen, -C₁₋₄alkyl, fluoro, chloro, -OC₁₋₂alkyl, -C(O)H, -C(O)OH, -C(O)OC₁₋₂alkyl, -CN, -C(O)NHC₁₋₂alkyl, or -C(O)NH₂, wherein each of said R³ alkyl is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC₁₋₂alkyl groups;


R⁴ is hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₅cycloalkyl, phenyl, a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole,

thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, or quinoline, or a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, or tetrahydropyridopyrimidine, wherein each of said R⁴ groups is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl, up to three CN, NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂, C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, a spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


2. The compound of claim 1, wherein Ring C is cyclobutane.
3. The compound of claim 2, having the following formula:




4. The compound of claim 3, having the following formula:




5. The compound of claim 4, having the following formula:




6. The compound of claim 4 or 5, wherein R^2 is $-C_{0-4}$ alkyl-NHR⁴.
7. The compound of claim 3, having the following formula:




8. The compound of claim 7, having the following formula:



9. The compound of claim 7 or 8, wherein R^1 is $-C_{0-4}$ alkyl-NHR⁴.
10. The compound of any one of claims 2-9, wherein X is $-O-$ or $-OC_{1-4}$ alkyl-.
11. The compound of any one of claims 2-10 wherein



12. The compound of claim 11 wherein



13. The compound of any one of claims 2-12, wherein R^3 is hydrogen.

14. The compound of any one of claims 2-13, wherein

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$, $C(O)NH(C_{0-4}$ alkyl- C_{3-5} cycloalkyl), CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$, C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl- $NHC(O)R^5$, C_{0-4} alkyl- $N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

15. The compound of any one of claims 2-13, wherein

R^4 is a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, or quinoline, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl,

C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$, $C(O)NH(C_{0-4}$ alkyl- C_{3-5} cycloalkyl), CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$, C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl- $NHC(O)R^5$, C_{0-4} alkyl- $N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

16. The compound of any one of claims 2-13, wherein


R^4 is a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, or tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$, $C(O)NH(C_{0-4}$ alkyl- C_{3-5} cycloalkyl), CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$, C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl- $NHC(O)R^5$, C_{0-4} alkyl- $N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents

is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and

each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


17. The compound of claim 1, wherein Ring C is cyclohexane.

18. The compound of claim 17, having the following formula:




19. The compound of claim 17 or 18, wherein X is $-NH-$.

20. The compound of any one of claims 17-19, having the following formula:



21. The compound of claim 20, having the following formula:




22. The compound of claim 20 or 21, wherein R^2 is $-C_{0-4}$ alkyl-NHR⁴ or $-OR^4$.


23. The compound of claim 22, wherein R^2 is $-NHR^4$.

24. The compound of claim 22, wherein R^2 is $-OR^4$.

25. The compound of claim 17-19, having the following formula:



26. The compound of claim 25 having the following formula:




27. The compound of claim 25 or 26, wherein R^1 is $-C_{0-4}$ alkyl-NHR⁴ or $-OR^4$.


28. The compound of claim 27, wherein R^1 is $-NHR^4$.

29. The compound of claim 27, wherein R^1 is $-OR^4$.

30. The compound of any one of claims 17-29 wherein




31. The compound of claim 30 wherein




32. The compound of claim 17 or 18, wherein X is $-O-$.

33. The compound of claim 17, 18 or 32, having the following formula:



34. The compound of claim 33, having the following formula:




35. The compound of claim 33 or 34, wherein is R^2 is $-C_{0-4}$ alkyl-NHR⁴ or -OR⁴.


36. The compound of claim 35, wherein is R^2 is -NHR⁴.

37. The compound of claim 35, wherein is R^2 is -OR⁴.

38. The compound of claim 17, 18 or 32, having the following formula:



39. The compound according to claim 38 having the following formula:




40. The compound of claim 38 or 39, wherein is R^1 is $-C_{0-4}$ alkyl-NHR⁴ or -OR⁴.

41. The compound of claim 40, wherein is R^1 is -NHR⁴.


42. The compound of claim 40, wherein is R^1 is -OR⁴.

43. The compound of claim 38 or 39 having the following formula:




Y is -O- or -NH-.

44. The compound of any one of claims 32-43 wherein



45. The compound of claim 44 wherein



46. The compound of any one of claims 17-45, wherein R^3 is hydrogen, C_{1-4} alkyl, OC_{1-2} alkyl, or $C(O)NH_2$, or $C(O)H$, wherein each of said R^3 alkyl is optionally substituted with OH.

47. The compound of claim 46, wherein R^3 is hydrogen.

48. The compound of any one of claims 17-47, wherein

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl,

$C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$,
 $C(O)NH(C_{0-4}$ alkyl- C_{3-5} cycloalkyl), CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$,
 C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl- $NHC(O)R^5$, C_{0-4} alkyl- $N(R^5)_2$, a heterocyclic ring system selected
from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine,
pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole,
pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said
optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4}
alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two
 SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and
each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from
imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from
oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with
chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two
 OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the
intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine
ring, or piperazine ring.

49. The compound of any one of claims 17-47, wherein

R^4 is a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole,
imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone,
pyrazine, pyridazine, or quinoline, wherein each of said R^4 groups is optionally substituted with
up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 ,
 C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl,
 C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl,
 C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$,
 $C(O)NH(C_{0-4}$ alkyl- C_{3-5} cycloalkyl), CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$,
 C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl- $NHC(O)R^5$, C_{0-4} alkyl- $N(R^5)_2$, a heterocyclic ring system selected
from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine,
pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole,
pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said

optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, a spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


50. The compound of claim 49, wherein

R⁴ is pyridine or pyrimidine, which is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl, up to three CN, NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂, C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, a spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

51. The compound of any one of claims 17-47, wherein


R^4 is a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, or tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

52. The compound of claim 32 having the formula:



R^3 is hydrogen, $-C_{1-4}$ alkyl, fluoro, chloro, $-OC_{1-2}$ alkyl, or $-C(O)NH_2$, $-C(O)H$, or $-CN$, wherein each of said R^3 alkyl is optionally substituted with OH or up to 3 fluorine atoms;

R^4 is



X^1 is N, CH, CF, CCl, or CC_{1-2} alkyl optionally substituted with up to 3 fluorine atoms;

X^2 is N or CR^{4c} , wherein X^1 and X^2 cannot simultaneously be N;

each of R^{4a} , R^{4b} , and R^{4c} is, independently, hydrogen, F, Cl, Br, CN, NO_2 , C_{1-4} alkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, or piperazine, or a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, or tetrazole, or R^{4c} , R^{4a} , and the intervening atoms form a dihydrofuran, a dihydropyran, or a tetrahydropiperidine heterocyclic ring system;

wherein each of said R^{4a} , R^{4b} , or R^{4c} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl, up to two OH groups, a

$C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and

wherein each of said R^{4a} , R^{4b} , or R^{4c} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

53. The compound of claim 52, wherein R^3 is hydrogen, $-C_{1-4}$ alkyl,

-OC₁₋₂alkyl, or -C(O)NH₂, or -C(O)H, wherein each of said R³ alkyl is optionally substituted with OH.

54. The compound of claim 52 or 53, wherein R³ is hydrogen.

55. The compound of any one of claims 52-54, wherein each of X¹ and X² is, independently, CH or N, wherein X¹ and X² cannot simultaneously be N.

56. The compound of any one of claims 52-55, wherein each of R^{4a} and R^{4b} is, independently, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, or piperazine, wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

57. The compound of any one of claims 52-55, wherein each of R^{4a} and R^{4b} is, independently, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, or tetrazole, wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

58. The compound of any one of claims 52-55, wherein each of R^{4a} and R^{4b} is, independently, hydrogen, F, Cl, Br, CN, NO₂, C₁₋₄ alkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, or C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl),

wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

- 59. A compound selected from the list of compounds in Table 1.
- 60. A compound selected from the list of compounds in Table 2.
- 61. A pharmaceutical composition comprising a compound of any one of claims 1-60 and a pharmaceutically acceptable excipient.
- 62. A compound of any one of claims 1-61, or a pharmaceutical composition comprising said compound for use in the sensitization of a cell to a therapeutic agent or a disease state that induces a DNA lesion.
- 63. A compound of any one of claims 1-60, or a pharmaceutical composition comprising said compound for use in the potentiation of a therapeutic regimen for the treatment of cancer in a patient.
- 64. A compound of any one of claims 1-60, or a pharmaceutical composition either alone or in combination with one or more additional therapeutic agent for use in the treatment of cancer or inhibiting cancer cell growth in a patient.