
MEDICAMENTS FOR THE TREATMENT OR PREVENTION OF FIBROTIC DISEASES

The present invention relates to a new use of indolinones of general formula



(I),

substituted in the 6 position, the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof, particularly the physiologically acceptable salts thereof.

BACKGROUND

Compounds of the above general formula I, the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof, particularly the physiologically acceptable salts thereof, have been described in WO 01/27081 and WO 04/13099 as having valuable pharmacological properties, in particular an inhibiting effect on various kinases, especially receptor tyrosine kinases such as VEGFR2, PDGFRα, PDGFRβ, FGFR1, FGFR3, EGFR, HER2, IGF1R and HGFR, as well as complexes of CDK's (Cyclin Dependent Kinases) such as CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK8 and CDK9 with their specific cyclins (A, B1, B2, C, D1, D2, D3, E, F, G1, G2, H, I and K) and to viral cyclin (cf. L. Mengtao in J. Virology 71(3), 1984-1991 (1997)), and on the proliferation of cultivated human cells, in particular endothelial cells, e.g. in angiogenesis, but also on the proliferation of other cells, in particular tumour cells.

However, none of these compounds have been described for their use in the treatment or prevention of the fibrotic diseases referred to in the present invention.

Remodeling is a normal response to tissue injury and inflammation that is observed in many tissues throughout the body. After resolution of the inflammation and repair of tissue damage, the tissue is generally returned to its original condition. Excessive uncontrolled tissue repair or the failure to stop remodeling when it is no longer required leads to condition known as fibrosis. Fibrosis is characterized by excessive deposition of extracellular matrix components and overgrowth of fibroblasts. Fibrosis can occur in all tissues but is especially prevalent in

organs with frequent exposure to chemical and biological insults including the lung, skin, digestive tract, kidney, and liver (Eddy, 1996, J Am Soc Nephrol, 7(12):2495-503; Dacic et al., 2003, Am J Respir Cell Mol Biol, 29S: S5-9; Wynn, 2004, Nat Rev Immunol, 4(8):583-94). Fibrosis often severely compromises the normal function(s) of the organ and many fibrotic diseases are, in fact, life-threatening or severely disfiguring, such as idiopathic pulmonary fibrosis (IPF), liver cirrhosis, scleroderma, or renal fibrosis. Treatment options for these diseases are often limited to organ transplantation, a risky and expensive procedure.

A large body of literature implicates the platelet-derived growth factor (PDGF), fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), epidermal growth factor (EGF), and transforming growth factor beta (TGFb) growth factor families in the induction or persistence of fibrosis (Levitzki, Cytokine Growth Factor Rev, 2004, 15(4):229-35; Strutz et al., Kidney Intl, 2000, 57:1521-38; Strutz et al., 2003, Springer Semin Immunopathol, 24:459-76; Rice et al., 1999, Amer J Pathol, 155(1):213-221; Broekelmann et al., 1991, Proc Nat Acad Sci, 88:6642-6; Wynn, 2004, Nat Rev Immunol, 4(8):583-94).

PDGF, EGF and FGF family members are potent mitogens for mesenchymal cells such as smooth muscle cells, myofibroblasts and fibroblasts (Benito et al., 1993, Growth Regul 3(3):172-9; Simm et al, 1998, Basic Res Cardiol, 93(S3):40-3; Klagsburn, Prog Growth Factor Res, 1989, 1(4):207-35; Kirkland et al., 1998, J Am Soc Nephrol, 9(8):1464-73), the very cells which supplant normal tissue

in fibrosis and are believed to play a role in tissue remodeling (Abboud, 1995, Annu Rev Physiol., 57:297-309; Jinnin et al., 2004, J Cell Physiol, online; Martinet et al., 1996, Arch Toxicol 18:127-39; Desmouliere, Cell Biology International, 1995, 19:471-6; Jelaska et al., Springer Semin Immunopathol, 2000, 21:385-95).

Inhibition of PDGF attenuates both liver fibrosis and lung fibrosis in experimental models, suggesting fibrosis in different organs may have a common origin (Borkham-Kamphorst et al., 2004, Biochem Biophys Res Commun; Rice et al., 1999, Amer J Pathol, 155(1):213-221). An EGF receptor kinase inhibitor was also active in this lung fibrosis model. Three-fold overexpression of an EGF family member, HB-EGF, in mouse pancreas islets was sufficient to cause development of fibrosis in both the exocrine and endocrine compartments (Means et al., 2003, Gastroenterology, 124(4):1020-36).

Similarly, FGF1/FGF2-deficient mice show dramatically decreased liver fibrosis after chronic carbon tetrachloride (CCl4) exposure (Yu et al., 2003, Am J Pathol, 163(4):1653-62). FGF expression is increased in human renal interstitial fibrosis where it strongly correlates with interstitial scarring (Strutz et al., 2000, Kidney Intl, 57:1521-38) as well as in a model of experimental lung fibrosis (Barrios et al., 1997, Am J Physiol, 273 (2 Pt 1):L451-8), again lending credence to the idea that fibrosis in various tissues has a common basis.

In addition, elevated levels of VEGF have been observed in several studies in persons with asthma (Hoshino et al.,

2001, J Allergy Clin Immunol 107:1034-39; Hoshino et al. , 2001, J Allergy Clin Immunol 107:295-301; Kanazawa et al. , 2002, Thorax 57:885-8; Asai et al., J Allergy Clin Immunol 110:571-5, 2002; Kanazawa et al., 2004, Am J Respir Crit Care Med, 169:1125-30). Inducible expression of VEGF in a transgenic mouse model induces an asthma-like phenotype, edema, angiogenesis and smooth muscle hyperplasia (Lee et al., 2004, Nature Med 10:1095-1103).

Finally, TGFb stimulates production of extracellular matrix proteins including fibronectin and collagens and is believed to play an important role in fibrosis in many tissues (Leask et al., 2004, FASEB J 18(7):816-27; Bartram et al., 2004, Chest 125(2):754-65; Strutz et al., 2003, Springer Semin Immunopathol, 24:459-76; Wynn, 2004, Nat Rev Immunol, 4(8):583-94). Inhibitors of TGFb production and signaling pathways are active in a number of fibrosis animal models (Wang et al., 2002, Exp Lung Res, 28:405-17; Laping, 2003, Curr Opin Pharmacol, 3(2):204-8).

As summarized above, several growth factors are upregulated in fibrosis and the inhibition of a single factor seems to reduce the severity of fibrosis in the fibrosis models.

SUMMARY OF THE INVENTION

Surprisingly, we found that the compounds of above general formula I are effective in the treatment or prevention of specific fibrotic diseases.


The present invention thus relates to the use of the compounds of above general formula I for the preparation of a medicament for the treatment or prevention of specific fibrotic diseases.

The present invention also relates to a method for the treatment or prevention of specific fibrotic diseases, by administration to a patient in need thereof of a pharmaceutical composition comprising a compound of above general formula I, together with a pharmaceutically suitable carrier. The expression "patient" is meant to comprise the mammalian animal body, preferably the human body.

The present invention further relates to a pharmaceutical composition for the treatment or prevention of specific fibrotic diseases which comprises a compound of above general formula I alone or in combination with one or more further therapeutic agents.

DETAILLED DESCRIPTION OF THE INVENTION

In accordance with the present invention, the compounds of above general formula I are the compounds



(I),

in which

X denotes an oxygen or sulphur atom,

R_1 denotes a hydrogen atom or a prodrug group such as a C_{1-4}-alkoxycarbonyl or C_{2-4}-alkanoyl group,

R_2 denotes a carboxy group, a straight-chain or branched C_{1-6}-alkoxy-carbonyl group, a C_{4-7}-cycloalkoxy-carbonyl or an aryloxycarbonyl group,

a straight-chain or branched C_{1-6}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, heteroaryl, carboxy, C_{1-3}-alkoxy-carbonyl, aminocarbonyl, C_{1-3}-alkylamino-carbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

a straight-chain or branched C_{2-6}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a chlorine atom or a hydroxy, C_{1-3}-alkoxy, amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group,

an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C_{1-3}-alkoxy group or a di-(C_{1-2}-alkyl)-aminocarbonyl group,

R_3 denotes a hydrogen atom, a C_{1-6}-alkyl, C_{3-7}-cycloalkyl, trifluoromethyl or heteroaryl group,

a phenyl or naphthyl group, a phenyl or naphthyl group mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a trifluoromethyl, C_{1-3}-alkyl or C_{1-3}-alkoxy group, whilst in the event of disubstitution the substituents may be identical or different and wherein the abovementioned unsubstituted as well as the mono- and disubstituted phenyl and naphthyl groups may additionally be substituted

by a hydroxy, hydroxy-C_{1-3}-alkyl or C_{1-3}-alkoxy-C_{1-3}-alkyl group,

by a cyano, carboxy, carboxy-C_{1-3}-alkyl, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylamino-carbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

by a nitro group,

by an amino, C_{1-3}-alkylamino, di-(C_{1-3}-alkyl)-amino or amino-C_{1-3}-alkyl group,

by a C_{1-3}-alkylcarbonylamino, N-(C_{1-3}-alkyl)-C_{1-3}-alkyl-carbonylamino, C_{1-3}-alkylcarbonylamino-C_{1-3}-alkyl, N-(C_{1-3}-alkyl)-C_{1-3}-alkylcarbonylamino-C_{1-3}-alkyl, C_{1-3}-alkyl-

sulphonylamino, C_{1-3}-alkylsulphonylamino-C_{1-3}-alkyl, N-(C_{1-3}-alkyl)-C_{1-3}-alkylsulphonylamino-C_{1-3}-alkyl or aryl-C_{1-3}-alkylsulphonylamino group,

by a cycloalkylamino, cycloalkyleneimino, cycloalkyleneiminocarbonyl, cycloalkyleneimino-C_{1-3}-alkyl, cycloalkyleneiminocarbonyl-C_{1-3}-alkyl or cycloalkyleneiminosulphonyl-C_{1-3}-alkyl group having 4 to 7 ring members in each case, whilst in each case the methylene group in position 4 of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH or -N(C_{1-3}-alkyl) group,

or by a heteroaryl or heteroaryl-C_{1-3}-alkyl group,

R_4 denotes a C_{3-7}-cycloalkyl group,

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group or replaced by an -NH or -N(C_{1-3}-alkyl) group,


or a phenyl group substituted by the group R_6, which may additionally be mono- or disubstituted by fluorine, chlorine, bromine or iodine atoms, by C_{1-5}-alkyl, trifluoromethyl, hydroxy, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, amino, acetylamino, C_{1-3}-alkyl-sulphonylamino, aminocarbonyl, C_{1-3}-alkyl-aminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, aminosulphonyl, C_{1-3}-alkyl-aminosulphonyl, di-(C_{1-3}-alkyl)-aminosulphonyl, nitro or cyano groups,

wherein the substituents may be identical or different and wherein

R_6 denotes a hydrogen, fluorine, chlorine, bromine or iodine atom,

a cyano, nitro, amino, C_{1-5}-alkyl, C_{3-7}-cycloalkyl, trifluoromethyl, phenyl, tetrazolyl or heteroaryl group,

the group of formula



,

wherein the hydrogen atoms bound to a nitrogen atom may in each case be replaced independently of one another by a C_{1-3}-alkyl group,

a C_{1-3}-alkoxy group, a C_{1-3}-alkoxy-C_{1-3}-alkoxy, phenyl-C_{1-3}-alkoxy, amino-C_{2-3}-alkoxy, C_{1-3}-alkylamino-C_{2-3}-alkoxy, di-(C_{1-3}-alkyl)-amino-C_{2-3}-alkoxy, phenyl-C_{1-3}-alkylamino-C_{2-3}-alkoxy, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino-C_{2-3}-alkoxy, C_{5-7}-cycloalkyleneimino-C_{2-3}-alkoxy or C_{1-3}-alkylmercapto group,

a carboxy, C_{1-4}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkyl-amino-carbonyl, N-(C_{1-5}-alkyl)-C_{1-3}-alkylaminocarbonyl, phenyl-C_{1-3}-alkylamino-carbonyl, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino-carbonyl, piperazinocarbonyl or N-(C_{1-3}-alkyl)-piperazinocarbonyl group,

a C_{1-3}-alkylaminocarbonyl or N-(C_{1-5}-alkyl)-C_{1-3}-alkylaminocarbonyl group wherein an alkyl moiety is substituted by a carboxy or C_{1-3}-alkoxycarbonyl group or in the 2 or 3 position by a di-(C_{1-3}-alkyl)-amino, piperazino, N-(C_{1-3}-alkyl)-piperazino or a 4- to 7-membered cycloalkyleneimino group,

a C_{3-7}-cycloalkyl-carbonyl group,

wherein the methylene group in the 4 position of the 6- or 7-membered cycloalkyl moiety may be substituted by an amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group or replaced by an -NH or -N(C_{1-3}-alkyl) group,

a 4- to 7-membered cycloalkyleneimino group wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

the cycloalkylene moiety may be fused to a phenyl ring or

one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl group and/or

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, –NH, –N(C_{1-3}–alkyl), –N(phenyl), –N(C_{1-3}–alkyl–carbonyl) or –N(benzoyl) group,

a C_{1-4}–alkyl group substituted by the group R_7, wherein

R_7 denotes a C_{3-7}–cycloalkyl group,

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C_{1-3}–alkylamino or di–(C_{1-3}–alkyl)–amino group or replaced by an –NH or –N(C_{1-3}–alkyl) group or

in a 5- to 7-membered cycloalkyl group a $-(CH_2)_2$ group may be replaced by a –CO–NH group, a $-(CH_2)_3$ group may be replaced by a –NH–CO–NH or –CO–NH–CO group or a $-(CH_2)_4$ group may be replaced by a –NH–CO–NH–CO group, whilst in each case a hydrogen atom bound to a nitrogen atom may be replaced by a C_{1-3}–alkyl group,

an aryl or heteroaryl group,

a hydroxy or C_{1-3}–alkoxy group,

an amino, C_{1-7}–alkylamino, di–(C_{1-7}–alkyl)–amino, phenylamino, N–phenyl–C_{1-3}–alkyl–amino, phenyl–C_{1-3}–alkyl–amino, N–(C_{1-3}–alkyl)–phenyl–C_{1-3}–alkylamino or di–(phenyl–C_{1-3}–alkyl)–amino group,

an ω–hydroxy–C_{2-3}–alkyl–amino, N–(C_{1-3}–alkyl)–ω–hydroxy–C_{2-3}–alkyl–amino, di–(ω–hydroxy–C_{2-3}–alkyl)–amino,

di-(ω-(C_{1-3}-alkoxy)-C_{2-3}-alkyl)-amino or N-(dioxolan-2-yl)-C_{1-3}-alkyl-amino group,

a C_{1-3}-alkylcarbonylamino-C_{2-3}-alkyl-amino or C_{1-3}-alkylcarbonylamino-C_{2-3}-alkyl-N-(C_{1-3}-alkyl)-amino group,

a C_{1-3}-alkylsulphonylamino, N-(C_{1-3}-alkyl)-C_{1-3}-alkyl-sulphonylamino, C_{1-3}-alkylsulphonylamino-C_{2-3}-alkyl-amino or C_{1-3}-alkylsulphonylamino-C_{2-3}-alkyl-N-(C_{1-3}-alkyl)-amino group,

a hydroxycarbonyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-hydroxycarbonyl-C_{1-3}-alkyl-amino group,

a guanidino group wherein one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl group,

a group of formula

$-N(R_8)-CO-(CH_2)_n-R_9$ (II),

wherein

R_8 denotes a hydrogen atom or a C_{1-3}-alkyl group,

n denotes one of the numbers 0, 1, 2 or 3 and

R_9 denotes an amino, C_{1-4}-alkylamino, di-(C_{1-4}-alkyl)-amino, phenylamino, N-(C_{1-4}-alkyl)-phenylamino, benzylamino, N-(C_{1-4}-alkyl)-benzylamino or C_{1-4}-alkoxy group, a 4- to 7-membered cycloalkyleneimino group, whilst in each case the methylene group in the 4 position of a 6- or 7-membered

cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, –NH, –N(C_{1-3}–alkyl), –N(phenyl), –N(C_{1-3}–alkyl–carbonyl) or –N(benzoyl) group, or, if n denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

$$-N(R_{10})-(CH_2)_m-(CO)_o-R_{11} \quad (III),$$

wherein

R_{10} denotes a hydrogen atom, a C_{1-3}–alkyl group, a C_{1-3}–alkylcarbonyl, arylcarbonyl, phenyl–C_{1-3}–alkyl–carbonyl, C_{1-3}–alkylsulphonyl, arylsulphonyl or phenyl–C_{1-3}–alkylsulphonyl group,

m denotes one of the numbers 1, 2, 3 or 4,

o denotes the number 1 or, if m denotes one of the numbers 2, 3 or 4, o may also denote the number 0 and

R_{11} denotes an amino, C_{1-4}–alkylamino, di–(C_{1-4}–alkyl)–amino, phenylamino, N–(C_{1-4}–alkyl)–phenylamino, benzylamino, N–(C_{1-4}–alkyl)–benzylamino, C_{1-4}–alkoxy or C_{1-3}–alkoxy–C_{1-3}–alkoxy group, a di–(C_{1-4}–alkyl)–amino–C_{1-3}–alkylamino group optionally substituted in the 1 position by a C_{1-3}–alkyl group or a 4– to 7–membered cycloalkyleneimino group, wherein the cycloalkylene moiety may be fused to a phenyl ring or in each case the methylene group in the 4 position of a 6– or 7–membered cycloalkyleneimino group may be replaced by an oxygen or

sulphur atom, by a sulphinyl, sulphonyl, –NH, –N(C_{1-3}–alkyl), –N(phenyl), –N(C_{1-3}–alkyl–carbonyl) or –N(benzoyl) group,

a C_{4-7}–cycloalkylamino, C_{4-7}–cycloalkyl–C_{1-3}–alkylamino or C_{4-7}–cycloalkenylamino group wherein position 1 of the ring is not involved in the double bond and wherein the abovementioned groups may each additionally be substituted at the amino–nitrogen atom by a C_{5-7}–cycloalkyl, C_{2-4}–alkenyl or C_{1-4}–alkyl group,

a 4– to 7–membered cycloalkyleneimino group, wherein

the cycloalkylene moiety may be fused to a phenyl group or to an oxazolo, imidazolo, thiazolo, pyridino, pyrazino or pyrimidino group optionally substituted by a fluorine, chlorine, bromine or iodine atom, by a nitro, C_{1-3}–alkyl, C_{1-3}–alkoxy or amino group, and/or

one or two hydrogen atoms may each be replaced by a C_{1-3}–alkyl, C_{5-7}–cycloalkyl or phenyl group and/or

the methylene group in the 3 position of a 5–membered cycloalkyleneimino group may be substituted by a hydroxy, hydroxy–C_{1-3}–alkyl, C_{1-3}–alkoxy or C_{1-3}–alkoxy–C_{1-3}–alkyl group,

the methylene group in the 3 or 4 position of a 6– or 7–membered cycloalkyleneimino group may in each case be substituted by a hydroxy, hydroxy–C_{1-3}–alkyl, C_{1-3}–alkoxy, C_{1-3}–alkoxy–C_{1-3}–alkyl, carboxy, C_{1-4}–alkoxycarbonyl, aminocarbonyl, C_{1-3}–alkylaminocarbonyl, di–(C_{1-3}–alkyl)–

aminocarbonyl, phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkyl-amino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl-), -N(phenyl), -N(phenyl-C_{1-3}-alkyl-), -N(C_{1-3}-alkyl-carbonyl-), -N(C_{1-4}-alkyl-hydroxy-carbonyl-), -N(C_{1-4}-alkoxy-carbonyl-), -N(benzoyl-) or -N(phenyl-C_{1-3}-alkyl-carbonyl-) group,

wherein a methylene group linked to an imino-nitrogen atom of the cycloalkyleneimino group may be replaced by a carbonyl or sulphonyl group or in a 5- to 7-membered monocyclic cycloalkyleneimino group or a cycloalkyleneimino group fused to a phenyl group the two methylene groups linked to the imino-nitrogen atom may each be replaced by a carbonyl group,

or R_6 denotes a C_{1-4}-alkyl group which is substituted by a carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group or by a 4- to 7-membered cycloalkyleneiminocarbonyl group,

an N-(C_{1-3}-alkyl)-C_{2-4}-alkanoylamino group which is additionally substituted in the alkyl moiety by a carboxy or C_{1-3}-alkoxycarbonyl group,

a group of formula

$-N(R_{12})-CO-(CH_2)_p-R_{13}$ (IV),

wherein

R_{12} denotes a hydrogen atom, a C_{1-6}-alkyl or C_{3-7}-cycloalkyl group or a C_{1-3}-alkyl group terminally substituted by a phenyl, heteroaryl, trifluoromethyl, hydroxy, C_{1-3}-alkoxy, aminocarbonyl, C_{1-4}-alkylamino-carbonyl, di-(C_{1-4}-alkyl)-amino-carbonyl, C_{1-3}-alkyl-carbonyl, C_{1-3}-alkyl-sulphonyl-amino, N-(C_{1-3}-alkyl)-C_{1-3}-alkyl-sulphonylamino, C_{1-3}-alkyl-aminosulphonyl or di-(C_{1-3}-alkyl)-aminosulphonyl group and

p denotes one of the numbers 0, 1, 2 or 3 and

R_{13} assumes the meanings of the abovementioned group R_7, or, if p denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

$$-N(R_{14})-(CH_2)_q-(CO)_r-R_{15} \qquad (V),$$

wherein

R_{14} denotes a hydrogen atom, a C_{1-4}-alkyl group, a C_{1-3}-alkylcarbonyl, arylcarbonyl, phenyl-C_{1-3}-alkylcarbonyl, heteroarylcarbonyl, heteroaryl-C_{1-3}-alkylcarbonyl, C_{1-4}-alkylsulphonyl, arylsulphonyl, phenyl-C_{1-3}-alkylsulphonyl, heteroarylsulphonyl or heteroaryl-C_{1-3}-alkyl-sulphonyl group,

q denotes one of the numbers 1, 2, 3 or 4,

r denotes the number 1 or, if q is one of the numbers 2, 3 or 4, it may also denote the number 0 and

R_{15} assumes the meanings of the abovementioned group R_7,

a group of formula

$-N(R_{16})-SO_2-R_{17}$ (VI),

wherein

R_{16} denotes a hydrogen atom or a C_{1-4}-alkyl group optionally terminally substituted by a cyano, trifluoromethyl-carbonylamino or N-(C_{1-3}-alkyl)-trifluoromethyl-carbonyl-amino group and

R_{17} denotes a C_{1-3}-alkyl group,

an amino group substituted by a di-(C_{1-3}-alkyl)-amino-C_{1-3}-alkyl-carbonyl or di-(C_{1-3}-alkyl)-amino-C_{1-3}-alkyl-sulphonyl group and a di-(C_{1-3}-alkyl)-aminocarbonyl-C_{1-3}-alkyl group,

or an N-(C_{1-3}-alkyl)-C_{1-5}-alkylsulphonylamino or N-(C_{1-3}-alkyl)-phenylsulphonylamino group wherein the alkyl moiety is additionally substituted by a cyano or carboxy group,

wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R_6 may be mono- or disubstituted by fluorine, chlorine, bromine or iodine

atoms, by C_{1-5}-alkyl, trifluoromethyl, hydroxy, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-4}-alkylamino-carbonyl, di-(C_{1-4}-alkyl)-amino-carbonyl, aminosulphonyl, C_{1-3}-alkyl-aminosulphonyl, di-(C_{1-3}-alkyl)-aminosulphonyl, C_{1-3}-alkyl-sulphonylamino, nitro or cyano groups, wherein the substituents may be identical or different, or two adjacent hydrogen atoms of the phenyl groups may be replaced by a methylenedioxy group,

and

R_5 denotes a hydrogen atom or a C_{1-3}-alkyl group,

wherein by an aryl group is meant a phenyl or naphthyl group optionally mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a cyano, trifluoromethyl, nitro, carboxy, aminocarbonyl, C_{1-3}-alkyl or C_{1-3}-alkoxy group and

by a heteroaryl group is meant a monocyclic 5- or 6-membered heteroaryl group optionally substituted by a C_{1-3}-alkyl group in the carbon skeleton, wherein

the 6-membered heteroaryl group contains one, two or three nitrogen atoms and

the 5-membered heteroaryl group contains an imino group optionally substituted by a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group, an oxygen or sulphur atom or

an imino group optionally substituted by a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group or an oxygen or sulphur atom and additionally a nitrogen atom or

an imino group optionally substituted by a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group and two nitrogen atoms,

and moreover a phenyl ring may be fused to the abovementioned monocyclic heterocyclic groups via two adjacent carbon atoms and the bonding takes place via a nitrogen atom or via a carbon atom of the heterocyclic moiety or a fused phenyl ring,

some or all of the hydrogen atoms in the abovementioned alkyl and alkoxy groups or in the alkyl moieties contained in the above-defined groups of formula I optionally being replaced by fluorine atoms,

the saturated alkyl and alkoxy moieties with more than 2 carbon atoms which are present in the groups defined hereinbefore also include the branched isomers thereof, such as for example the isopropyl, tert.butyl, isobutyl group, unless otherwise stated, and

additionally the hydrogen atom of any carboxy group present or a hydrogen atom bound to a nitrogen atom, e.g. a hydrogen atom of an amino, alkylamino or imino group or a saturated N-heterocycle such as the piperidinyl group, may each be replaced by a group which can be cleaved *in vivo*.

By a group which can be cleaved *in vivo* from an imino or amino group is meant, for example, a hydroxy group, an acyl

group such as the benzoyl or pyridinoyl group or a C_{1-16}-alkanoyl group such as the formyl, acetyl, propionyl, butanoyl, pentanoyl or hexanoyl group, an allyloxycarbonyl group, a C_{1-16}-alkoxycarbonyl group such as the methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, tert.butoxycarbonyl, pentoxycarbonyl, hexyloxycarbonyl, octyloxycarbonyl, nonyloxycarbonyl, decyloxycarbonyl, undecyloxycarbonyl, dodecyloxycarbonyl or hexadecyloxycarbonyl group, a phenyl-C_{1-6}-alkoxycarbonyl group such as the benzyloxycarbonyl, phenylethoxycarbonyl or phenylpropoxycarbonyl group, a C_{1-3}-alkylsulphonyl-C_{2-4}-alkoxycarbonyl, C_{1-3}-alkoxy-C_{2-4}-alkoxy-C_{2-4}-alkoxycarbonyl or $R_eCO-O-(R_fCR_g)-O-CO$ group wherein

R_e denotes a C_{1-8}-alkyl, C_{5-7}-cycloalkyl, phenyl or phenyl-C_{1-3}-alkyl group,

R_f denotes a hydrogen atom, a C_{1-3}-alkyl, C_{5-7}-cycloalkyl or phenyl group and

R_g denotes a hydrogen atom, a C_{1-3}-alkyl or $R_eCO-O-(R_fCR_g)-O$ group wherein R_e to R_g are as hereinbefore defined,

wherein additionally the amino group may be a phthalimido group, whilst the abovementioned ester groups may also be used as a group which can be converted *in vivo* into a carboxy group.

One sub-group of compounds of general formula I which deserves special mention comprises those wherein

X, R_1 and R_3 to R_5 are as hereinbefore defined and

R_2 denotes a straight-chain or branched C_{1-6}-alkoxy-carbonyl group, a C_{4-7}-cycloalkoxycarbonyl or a aryloxycarbonyl group,

a straight-chain or branched C_{1-6}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, heteroaryl, carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

a straight-chain or branched C_{2-6}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a chlorine atom or a hydroxy, C_{1-3}-alkoxy, amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

A second sub-group of compounds of general formula I which deserves special mention comprises those wherein

X, R_1 and R_3 to R_5 are as hereinbefore defined and

R_2 denotes an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C_{1-3}-alkoxy group or a di-(C_{1-2}-alkyl)-aminocarbonyl group,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

A third sub-group of compounds of general formula I which deserves special mention comprises those wherein

X, R_1 to R_3 and R_5 are as hereinbefore defined and

R_4 denotes an R_7-(C_{1-4}-alkyl)-phenyl group, wherein

R_7 denotes an amino, C_{1-7}-alkylamino, di-(C_{1-7}-alkyl)-amino, phenylamino, N-phenyl-C_{1-3}-alkyl-amino, phenyl-C_{1-3}-alkyl-amino, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino or di-(phenyl-C_{1-3}-alkyl)-amino group,

or a phenyl group substituted by the group of formula

$$-N(R_{12})-CO-(CH_2)_p-R_{13} \quad (IV),$$

wherein R_{12}, p and R_{13} are as hereinbefore defined,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

Preferred compounds of general formula I are those wherein

R_1 and R_3 are as hereinbefore defined and

X denotes an oxygen atom,

R_2 denotes a carboxy group, a straight-chain or branched C_{1-6}-alkoxy-carbonyl group, a C_{5-7}-cycloalkoxycarbonyl or a phenoxycarbonyl group,

a straight-chain or branched C_{1-3}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, heteroaryl, carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

a straight-chain or branched C_{2-3}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a chlorine atom, by a hydroxy, C_{1-3}-alkoxy, amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group,

an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C_{1-3}-alkoxy group or a di-(C_{1-2}-alkyl)-aminocarbonyl group,

R_4 denotes a C_{3-7}-cycloalkyl group,

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group or replaced by an -NH or -N(C_{1-3}-alkyl) group,


or a phenyl group substituted by the group R_6, which may additionally be mono- or disubstituted by fluorine, chlorine or bromine atoms, by C_{1-3}-alkyl, trifluoromethyl, hydroxy, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, amino, acetylamino, aminocarbonyl, C_{1-3}-alkyl-aminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, nitro or cyano groups, wherein the substituents may be identical or different and

wherein

R_6 denotes a hydrogen, fluorine, chlorine, bromine or iodine atom,

a cyano, nitro, amino, C_{1-5}-alkyl, C_{3-7}-cycloalkyl, trifluoromethyl, phenyl, tetrazolyl or heteroaryl group,

the group of formula



,

wherein a hydrogen atom bound to the nitrogen atom may be replaced by a C_{1-3}-alkyl group,

a C_{1-3}-alkoxy group, an amino-C_{2-3}-alkoxy, C_{1-3}-alkylamino-C_{2-3}-alkoxy, di-(C_{1-3}-alkyl)-amino-C_{2-3}-alkoxy, phenyl-C_{1-3}-alkylamino-C_{2-3}-alkoxy, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino-C_{2-3}-alkoxy, pyrrolidino-C_{2-3}-alkoxy, piperidino-C_{2-3}-alkoxy or C_{1-3}-alkylmercapto group,

a carboxy, C_{1-4}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkyl-amino-carbonyl, phenyl-C_{1-3}-alkylamino-carbonyl or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino-carbonyl group,

a C_{3-7}-cycloalkyl-carbonyl group,

wherein the methylene group in the 4 position of the 6- or 7-membered cycloalkyl moiety may be replaced by an -NH or -N(C_{1-3}-alkyl) group,

a 4- to 7-membered cycloalkyleneimino group, wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl group and/or

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH or -N(C_{1-3}-alkyl) group,

a C_{1-4}-alkyl group terminally substituted by the group R_7, wherein

R_7 denotes a C_{5-7}-cycloalkyl group,

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be replaced by an -NH or -N(C_{1-3}-alkyl) group or

in a 5- to 7-membered cycloalkyl group a $-(CH_2)_2$ group may be replaced by a -CO-NH group, a $-(CH_2)_3$ group may be replaced by a -NH-CO-NH- or a $-(CH_2)_4$ group may be replaced by a -NH-CO-NH-CO group, whilst in each case a hydrogen atom bound to a nitrogen atom may be replaced by a C_{1-3}-alkyl group,

a phenyl or heteroaryl group,

a hydroxy or C_{1-3}-alkoxy group,

an amino, C_{1-6}-alkylamino, di-(C_{1-6}-alkyl)-amino, phenylamino, N-phenyl-C_{1-3}-alkyl-amino, phenyl-C_{1-3}-alkylamino, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino or di-(phenyl-C_{1-3}-alkyl)-amino group,

a ω-hydroxy-C_{2-3}-alkyl-amino, N-(C_{1-3}-alkyl)-ω-hydroxy--C_{2-3}-alkyl-amino, di-(ω-hydroxy-C_{2-3}-alkyl)-amino, di-(ω-(C_{1-3}-alkoxy)-C_{2-3}-alkyl)-amino or N-(dioxolan-2-yl)-C_{1-3}-alkyl-amino group,

a C_{1-3}-alkylcarbonylamino-C_{2-3}-alkyl-amino or C_{1-3}-alkylcarbonylamino-C_{2-3}-alkyl-N-(C_{1-3}-alkyl)-amino group,

a C_{1-3}-alkylsulphonylamino, N-(C_{1-3}-alkyl)-C_{1-3}-alkylsulphonylamino, C_{1-3}-alkylsulphonylamino--C_{2-3}-alkyl-amino or C_{1-3}-alkylsulphonylamino-C_{2-3}-alkyl--N-(C_{1-3}-alkyl)-amino group,

a hydroxycarbonyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-hydroxycarbonyl-C_{1-3}-alkyl-amino group

a guanidino group wherein a hydrogen atom may be replaced by a C_{1-3}-alkyl group,

a group of formula

$$-N(R_8)-CO-(CH_2)_n-R_9 \qquad (II),$$

wherein

R_8 denotes a hydrogen atom or a C_{1-3}-alkyl group,

n denotes one of the numbers 0, 1, 2 or 3 and

R_9 denotes an amino, C_{1-3}-alkylamino, di-(C_{1-3}-alkyl)-amino, phenylamino, benzylamino or C_{1-4}-alkoxy group, a 5- to 7-membered cycloalkyleneimino group, wherein the methylene group in position 4 of the piperidino group may be replaced by an oxygen or sulphur atom, by an -NH, -N(C_{1-3}-alkyl), -N(phenyl), -N(C_{1-3}-alkyl-carbonyl) or -N(benzoyl) group, or, if n denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

$$-N(R_{10})-(CH_2)_m-(CO)_o-R_{11} \quad (III),$$

wherein

R_{10} denotes a hydrogen atom, a C_{1-3}-alkyl group, a C_{1-3}-alkylcarbonyl or C_{1-3}-alkylsulphonyl group,

m denotes one of the numbers 1, 2 or 3,

o denotes the number 1 or, if m is one of the numbers 2 or 3, o may also denote the number 0 and

R_{11} denotes an amino, C_{1-3}-alkylamino, di-(C_{1-3}-alkyl)-amino, C_{1-4}-alkoxy or C_{1-3}-alkoxy-C_{1-3}-alkoxy group or a 5- to 7-

membered cycloalkyleneimino group, wherein the methylene group in position 4 of the piperidino group may be replaced by an oxygen or sulphur atom, by an –NH, –N(C_{1-3}-alkyl), –N(phenyl), –N(C_{1-3}-alkyl-carbonyl) or –N(benzoyl) group,

a C_{4-7}-cycloalkylamino or C_{4-7}-cycloalkenylamino group wherein position 1 of the ring is not involved in the double bond,

a 4- to 7-membered cycloalkyleneimino group, wherein

the cycloalkylene moiety may be fused to a phenyl group or

one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl group and/or

the methylene group in position 3 of the pyrrolidino group may be substituted by a hydroxy or C_{1-3}-alkoxy group,

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a hydroxy, hydroxy-C_{1-3}-alkyl, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, –NH, –N(C_{1-3}-alkyl), –N(phenyl), –N(phenyl-C_{1-3}-alkyl), –N(C_{1-3}-alkyl-carbonyl), –N(C_{1-4}-alkoxy-carbonyl), –N(benzoyl) or –N(phenyl-C_{1-3}-alkyl-carbonyl) group,

wherein a methylene group linked to an imino-nitrogen atom of the cycloalkyleneimino group may be replaced by a carbonyl or sulphonyl group or in a 5- to 6-membered monocyclic cycloalkyleneimino group or a cycloalkyleneimino group fused to a phenyl group the two methylene groups linked to the imino-nitrogen atom may each be replaced by a carbonyl group,

or R_6 denotes a C_{1-4}-alkyl group which is terminally substituted by a carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group or by a 4- to 7-membered cycloalkyleneiminocarbonyl group,

a group of formula

$-N(R_{12})-CO-(CH_2)_p-R_{13}$ (IV),

wherein

R_{12} denotes a hydrogen atom, a C_{1-3}-alkyl, C_{5-7}-cycloalkyl, phenyl-C_{1-3}-alkyl or heteroaryl-C_{1-3}-alkyl group and

p denotes one of the numbers 0, 1, 2 or 3 and

R_{13} assumes the meanings of the abovementioned group R_7, or, if p denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

$-N(R_{14})-(CH_2)_q-(CO)_r-R_{15}$ (V),

wherein

R_{14} denotes a hydrogen atom, a C_{1-4}-alkyl group, a C_{1-3}-alkylcarbonyl, phenylcarbonyl, phenyl-C_{1-3}-alkyl-carbonyl, heteroarylcarbonyl, heteroaryl-C_{1-3}-alkylcarbonyl, C_{1-4}-alkylsulphonyl, phenylsulphonyl, phenyl-C_{1-3}-alkyl-sulphonyl- heteroarylsulphonyl or heteroaryl-C_{1-3}-alkyl-sulphonyl group,

q denotes one of the numbers 1, 2, 3 or 4,

r denotes the number 1 or, if q is one of the numbers 2, 3 or 4, it may also denote the number 0 and

R_{15} assumes the meanings of the abovementioned group R_7,

a group of formula

$-N(R_{16})-SO_2-R_{17}$ (VI),

wherein

R_{16} denotes a hydrogen atom or a C_{1-4}-alkyl group optionally terminally substituted by a cyano, trifluoromethyl-carbonylamino or N-(C_{1-3}-alkyl)-trifluoromethyl--carbonyl-amino group and

R_{17} denotes a C_{1-3}-alkyl group,

an amino group substituted by a di-(C_{1-3}-alkyl)-amino-C_{1-3}-alkyl-carbonyl or di-(C_{1-3}-alkyl)-amino-C_{1-3}-alkyl-sulphonyl group and a di-(C_{1-3}-alkyl)-aminocarbonyl-C_{1-3}-alkyl group,

wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R_6 may be mono- or disubstituted by fluorine, chlorine or bromine atoms, by C_{1-3}-alkyl, trifluoromethyl, hydroxy, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkyl-aminocarbonyl, aminosulphonyl, C_{1-3}-alkyl-aminosulphonyl, nitro or cyano groups, wherein the substituents may be identical or different, or two adjacent hydrogen atoms of the phenyl groups may be replaced by a methylenedioxy group, and

R_5 denotes a hydrogen atom or a C_{1-3}-alkyl group,

whilst by a heteroaryl group as mentioned above is meant a pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, furyl, thienyl, oxazolyl, thiazolyl, pyrazolyl, imidazolyl or triazolyl group optionally substituted in the carbon skeleton by a C_{1-3}-alkyl group wherein a hydrogen atom bound to a nitrogen atom may be replaced by a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group and wherein the 5-membered heteroaryl groups containing at least one imino group are bound via a carbon or nitrogen atom,

a hydrogen atom bound to a nitrogen atom in the abovementioned groups may be replaced by a group which can be cleaved *in vivo*, particularly by an acetyl or tert.butoxycarbonyl group,

the carboxy groups contained in the abovementioned groups may each be substituted by a group which can be cleaved *in vivo* and may occur, for example, in the form of the tert.butoxycarbonyl group,

some or all of the hydrogen atoms in the abovementioned alkyl and alkoxy groups or in the alkyl moieties contained in the above-defined groups of formula I optionally being replaced by fluorine atoms and

the saturated alkyl and alkoxy moieties contained in the abovementioned groups, which contain more than 2 carbon atoms, may be straight-chain or branched, unless otherwise stated,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

One subgroup of preferred compounds of general formula I deserving special mention comprises those wherein

X, R_1 and R_3 to R_5 are as hereinbefore defined and

R_2 denotes a straight-chain or branched C_{1-6}-alkoxy-carbonyl group, a C_{5-7}-cycloalkoxycarbonyl or a phenoxycarbonyl group,

a straight-chain or branched C_{1-3}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl- carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

a straight-chain or branched C_{2-3}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a hydroxy, C_{1-3}-alkoxy, amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

A second sub-group of preferred compounds of general formula I deserving special mention comprises those wherein

X, R_1 and R_3 to R_5 are as hereinbefore defined and

R_2 denotes an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C_{1-3}-alkoxy group or a di-(C_{1-2}-alkyl)-aminocarbonyl group,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

A third sub-group of preferred compounds of general formula I deserving special mention comprises those wherein

X, R_1 to R_3 and R_5 are as hereinbefore defined and

R_4 denotes an R_7-(n-C_{1-4}-alkyl)-phenyl group, wherein

R_7 denotes an amino, C_{1-6}-alkylamino, di-(C_{1-6}-alkyl)-amino, phenylamino, N-phenyl-C_{1-3}-alkyl-amino, phenyl-C_{1-3}-alkyl-

amino, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino or di-(phenyl-C_{1-3}-alkyl)-amino group,

or a phenyl group substituted by the group of formula

-N(R_{12})-CO-(CH_2)$_p$-R_{13} (IV),

wherein R_{12}, p and R_{13} are as hereinbefore defined,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

Particularly preferred compounds of general formula I are those wherein

X denotes an oxygen atom,

R_1 denotes a hydrogen atom,

R_2 denotes a carboxy group, a straight-chain or branched C_{1-4}-alkoxycarbonyl group or a phenoxycarbonyl group,

a straight-chain or branched C_{1-3}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

a straight-chain or branched C_{2-3}-alkoxy-carbonyl group which is terminally substituted in the alkyl moiety by a

hydroxy, C_{1-3}-alkoxy, amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group,

an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C_{1-3}-alkoxy group or a di-(C_{1-2}-alkyl)-aminocarbonyl group,

R_3 denotes a C_{1-4}-alkyl group or a phenyl group which may be substituted by a fluorine, chlorine or bromine atom, by a trifluoromethyl, C_{1-3}-alkyl, hydroxy or C_{1-3}-alkoxy group,

R_4 denotes a C_{5-6}-cycloalkyl group,

wherein the methylene group in position 4 of the cyclohexyl group may be substituted by an amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group or replaced by an -NH or -N(C_{1-3}-alkyl) group,


a phenyl group, a phenyl group disubstituted by C_{1-3}-alkyl, C_{1-3}-alkoxy or nitro groups, wherein the substituents may be identical or different, or

a phenyl group substituted by the group R_6, which may additionally be substituted by a fluorine, chlorine or bromine atom or by an amino or nitro group, wherein R_6 denotes a fluorine, chlorine or bromine atom,

a C_{1-3}-alkyl, C_{1-3}-alkoxy, nitro, amino or C_{5-6}-cycloalkyl group,

a pyrrolyl, pyrazolyl, imidazolyl, triazolyl or tetrazolyl group bound via a carbon atom, wherein the abovementioned heteroaromatic groups in the carbon skeleton may be substituted by a C_{1-3}-alkyl group or a hydrogen atom bound to a nitrogen atom may be replaced by a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group,

the group of formula



,

a carboxy, C_{1-4}-alkoxycarbonyl, phenyl-C_{1-3}-alkyl-amino-carbonyl or C_{5-7}-cycloalkyl-carbonyl group,

a 5 or 6-membered cycloalkyleneimino group, wherein

the methylene group in position 4 of the piperidino group may be replaced by an oxygen or sulphur atom, by an -NH or -N(C_{1-3}-alkyl) group,

an unbranched C_{1-3}-alkyl group terminally substituted by the group R_7, wherein

R_7 denotes a C_{5-7}-cycloalkyl group,

wherein in a 5 or 6-membered cycloalkyl group a $-(CH_2)_2$ group may be replaced by a -CO-NH group, a $-(CH_2)_3$ group may be replaced by an -NH-CO-NH- or a $-(CH_2)_4$ group may be replaced by an -NH-CO-NH-CO group, whilst in each case a

hydrogen atom bound to a nitrogen atom may be replaced by a C_{1-3}-alkyl group,

a phenyl or pyridinyl group or a pyrrolyl, pyrazolyl, imidazolyl or triazolyl group bound via a carbon or nitrogen atom, wherein the abovementioned heteroaromatic groups in the carbon skeleton may be substituted by a C_{1-3}-alkyl group or a hydrogen atom bound to a nitrogen atom may be replaced by a C_{1-3}-alkyl group,

a hydroxy or C_{1-3}-alkoxy group,

an amino, C_{1-6}-alkylamino, di-(C_{1-6}-alkyl)-amino, phenylamino, N-phenyl-C_{1-3}-alkylamino, phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino group,

a ω-hydroxy-C_{2-3}-alkyl-amino,
N-(C_{1-3}-alkyl)-ω-hydroxy-C_{2-3}-alkylamino,
di-(ω-hydroxy-C_{2-3}-alkyl)-amino or
di-(ω-(C_{1-3}-alkoxy)-C_{2-3}-alkyl)-amino group,

a C_{1-3}-alkylcarbonylamino-C_{2-3}-alkyl-amino or C_{1-3}-alkylcarbonylamino-C_{2-3}-alkyl-N-(C_{1-3}-alkyl)-amino group,

a C_{1-3}-alkylsulphonylamino, N-(C_{1-3}-alkyl)-C_{1-3}-alkylsulphonylamino, C_{1-3}-alkylsulphonylamino-C_{2-3}-alkylamino or C_{1-3}-alkylsulphonylamino-C_{2-3}-alkyl-N-(C_{1-3}-alkyl)-amino group,

a hydroxycarbonyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-hydroxycarbonyl-C_{1-3}-alkyl-amino group,

a guanidino group wherein a hydrogen atom may be replaced by a C_{1-3}-alkyl group,

a group of formula

$-N(R_8)-CO-(CH_2)_n-R_9$ (II),

wherein

R_8 denotes a hydrogen atom or a C_{1-3}-alkyl group,

n denotes one of the numbers 0, 1, 2 or 3 and

R_9 denotes an amino, C_{1-3}-alkylamino, di-(C_{1-3}-alkyl)-amino or C_{1-4}-alkoxy group, a 5- or 6-membered cycloalkyleneimino group, wherein the methylene group in position 4 of the piperidino group may be replaced by an -NH, -N(C_{1-3}-alkyl) or -N(C_{1-3}-alkyl-carbonyl) group, or, if n denotes one of the numbers 1, 2 or 3, R_9 may also denote a hydrogen atom,

a group of formula

$-N(R_{10})-(CH_2)_m-(CO)_o-R_{11}$ (III),

wherein

R_{10} denotes a hydrogen atom or a C_{1-3}-alkyl group,

m denotes one of the numbers 1, 2 or 3,

o denotes the number 1 or, if m is one of the numbers 2 or 3, o may also denote the number 0 and

R_{11} denotes an amino, C_{1-3}-alkylamino, di-(C_{1-3}-alkyl)-amino, C_{1-4}-alkoxy or methoxy-C_{1-3}-alkoxy group or a 5- or 6-membered cycloalkyleneimino group, wherein the methylene group in position 4 of the piperidino group may be replaced by an -NH, -N(C_{1-3}-alkyl) or -N(C_{1-3}-alkyl-carbonyl) group,

an azetidino, pyrrolidino, piperidino, 2,6-dimethyl-piperidino, 3,5-dimethyl-piperidino or azepino group, wherein

the methylene group in position 3 of the pyrrolidino group may be substituted by a hydroxy group,

the methylene group in position 4 of the piperidino group may be substituted by a hydroxy, hydroxy-C_{1-3}-alkyl or C_{1-3}-alkoxy group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl), -N(C_{1-3}-alkyl-carbonyl), -N(benzoyl) or -N(phenyl-C_{1-3}-alkyl-carbonyl) group,

wherein a methylene group linked to an imino-nitrogen atom of the pyrrolidino, piperidino or piperazino group may be replaced by a carbonyl group,

or R_6 denotes a straight-chain C_{1-3}-alkyl group which is terminally substituted by a carboxy or C_{1-3}-alkoxy-carbonyl group,

a group of formula

$-N(R_{12})-CO-(CH_2)_p-R_{13}$ (IV),

wherein

R_{12} denotes a hydrogen atom, a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group,

p denotes one of the numbers 0, 1 or 2 and

R_{13} denotes an amino, C_{1-4}-alkylamino, di-(C_{1-4}-alkyl)-amino, benzylamino, N-(C_{1-3}-alkyl)-benzylamino, C_{1-3}-alkoxy-C_{1-3}-alkylamino, N-(C_{1-3}-alkyl)-C_{1-3}-alkoxy-C_{1-3}-alkylamino, di-(2-methoxy-ethyl)-amino, di-(ω-hydroxy-C_{2-3}-alkyl)-amino or aminocarbonyl-methyl-N-(methyl)-amino group,

a pyrrolyl, pyrazolyl or imidazolyl group bound via a nitrogen atom and optionally substituted by a C_{1-3}-alkyl group,

a pyrrolidino, piperidino, morpholino, thiomorpholino or a piperazino group optionally substituted in the 4 position by a C_{1-3}-alkyl, phenyl-C_{1-3}-alkyl, C_{1-3}-alkylcarbonyl or C_{1-4}-alkoxycarbonyl group or, if n denotes the number 1 or 2, it may also denote a hydrogen atom,

a group of formula

$-N(R_{14})-(CH_2)_q-(CO)_r-R_{15}$ (V),

wherein

R_{14} denotes a hydrogen atom, a C_{1-4}-alkyl, C_{1-3}-alkyl-carbonyl, phenylcarbonyl, phenyl-C_{1-3}-alkylcarbonyl, furyl-carbonyl, pyridinyl-carbonyl, furyl-C_{1-3}-alkylcarbonyl, pyridinyl-C_{1-3}-alkylcarbonyl, C_{1-4}-alkylsulphonyl, phenylsulphonyl or phenyl-C_{1-3}-alkylsulphonyl group,

q denotes one of the numbers 1, 2 or 3,

r denotes the number 1 or, if q is one of the numbers 2 or 3, it may also denote the number 0 and

R_{15} denotes an amino, C_{1-4}-alkylamino, di-(C_{1-4}-alkyl)-amino, phenylamino, N-(C_{1-4}-alkyl)-phenylamino, benzylamino or N-(C_{1-4}-alkyl)-benzylamino group,

or a group of formula

$-N(R_{16})-SO_2-R_{17}$ (VI),

wherein

R_{16} denotes a hydrogen atom or a C_{1-3}-alkyl group optionally terminally substituted by a cyano, trifluoromethyl-carbonylamino or N-(C_{1-3}-alkyl)-trifluoromethyl-carbonyl-amino group and

R_{17} denotes a C_{1-3}-alkyl group,

wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R_6 may be

substituted by a fluorine, chlorine or bromine atom, by a methyl, trifluoromethyl, methoxy, nitro or cyano group and

R_5 denotes a hydrogen atom,

wherein a hydrogen atom bound to a nitrogen atom in the abovementioned groups may be replaced by an acetyl or tert.butoxycarbonyl group,

the carboxy groups contained in the abovementioned groups may also be present in the form of the tert.butoxycarbonyl precursor group and

the saturated alkyl and alkoxy moieties contained in the abovementioned groups, which contain more than 2 carbon atoms, may be straight-chain or branched, unless otherwise stated,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

One subgroup of particularly preferred compounds of general formula I deserving special mention comprises those wherein

X, R_1, R_3 and R_5 are as hereinbefore defined,

R_2 denotes a straight-chain or branched C_{1-4}-alkoxycarbonyl group or a phenoxycarbonyl group,

a straight-chain or branched C_{1-3}-alkoxycarbonyl group, which is terminally substituted in the alkyl moiety by a phenyl- carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl,

C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group, or

a straight-chain or branched C_{2-3}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a hydroxy, C_{1-3}-alkoxy, amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group, and

R_4 denotes an R_7-(n-C_{1-3}-alkyl)-phenyl group, wherein

R_7 denotes an amino, C_{1-6}-alkylamino, di-(C_{1-4}-alkyl)-amino, ω-hydroxy-C_{2-3}-alkyl-amino, N-(C_{1-3}-alkyl)-ω-hydroxy-C_{2-3}-alkyl-amino, di-(ω-hydroxy-C_{2-3}-alkyl)-amino or di-(ω-(C_{1-3}-alkoxy)-C_{2-3}-alkyl)-amino group,

or a phenyl group substituted by the group of formula

$-N(R_{12})-CO-(CH_2)_p-R_{13}$ (IV),

wherein R_{12}, p and R_{13} are as hereinbefore defined,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

A second subgroup of particularly preferred compounds of general formula I deserving special mention comprises those wherein

X, R_1, R_3 and R_5 are as hereinbefore defined,

R_2 denotes an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C_{1-3}-alkoxy group or a di-(C_{1-2}-alkyl)-aminocarbonyl group and

R_4 denotes a R_7-(n-C_{1-3}-alkyl)-phenyl group, wherein

R_7 denotes an amino, C_{1-6}-alkylamino, di-(C_{1-4}-alkyl)-amino, ω-hydroxy-C_{2-3}-alkyl-amino, N-(C_{1-3}-alkyl)-ω-hydroxy--C_{2-3}-alkyl-amino, di-(ω-hydroxy-C_{2-3}-alkyl)-amino or di-(ω-(C_{1-3}-alkoxy)-C_{2-3}-alkyl)-amino group,

or a phenyl group substituted by the group of formula

$-N(R_{12})-CO-(CH_2)_p-R_{13}$ (IV),

wherein R_{12}, p and R_{13} are as hereinbefore defined,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

Most particularly preferred compounds of general formula I are those wherein

X denotes an oxygen atom,

R_1 and R_5 each denote a hydrogen atom,

R_2 denotes a methoxycarbonyl, ethoxycarbonyl or aminocarbonyl group,

R_3 denotes a phenyl group and

R_4 denotes a phenyl group monosubstituted by the group R_6, wherein

R_6 denotes an N-methyl-imidazol-2-yl group,

an unbranched C_{1-3}-alkyl group which is terminally substituted by a C_{1-4}-alkylamino, di-(C_{1-4}-alkyl)-amino, piperidino or 2,6-dimethyl-piperidino group,

a group of formula

$-N(R_{12})-CO-(CH_2)_p-R_{13}$ (IV),

wherein

R_{12} denotes a C_{1-3}-alkyl group,

p denotes one of the numbers 1 or 2 and

R_{13} denotes a di-(C_{1-3}-alkyl)-amino group,

or a group of formula

$-N(R_{14})-(CH_2)_q-(CO)_r-R_{15}$ (V),

wherein

R_{14} denotes a C_{1-3}-alkyl-carbonyl or C_{1-3}-alkylsulphonyl group,

q denotes one of the numbers 1, 2 or 3,

r denotes the number 1 or, if q is one of the numbers 2 or 3, r may also denote the number 0 and

R_{15} denotes a di-(C_{1-3}-alkyl)-amino group,

wherein the saturated alkyl moieties contained in the abovementioned groups which contain more than 2 carbon atoms may be straight-chain or branched, unless otherwise stated,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

A subgroup of most particularly preferred compounds of general formula I deserving special mention comprises those wherein

X, R_1, R_3 and R_5 are as hereinbefore defined,

R_2 denotes a methoxycarbonyl or ethoxycarbonyl group and

R_4 denotes a di-(C_{1-3}-alkyl)-amino-C_{1-3}-alkylphenyl group or a phenyl group substituted by the group of formula

$-N(R_{12})-CO-(CH_2)_p-R_{13}$ (IV),

wherein R_{12}, p and R_{13} are as hereinbefore defined,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

The following are mentioned as examples of particularly preferred compounds:

(a) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(b) 3-Z-[(1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone,

(c) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(d) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(e) 3-Z-[1-(4-((2,6-dimethyl-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(f) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(g) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(h) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(i) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(j) 3-Z-[1-(4-(N-acetyl-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(k) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(l) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(m) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(n) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(o) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(p) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(q) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(r) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone and

(s) 3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

their tautomers, their stereoisomers or the physiologically acceptable salts thereof.

Another subgroup of compounds of general formula I comprises those wherein
X denotes an oxygen or sulphur atom,

R_1 denotes a hydrogen atom or a prodrug group such as a C_{1-4}-alkoxycarbonyl or C_{2-4}-alkanoyl group,

R_2 denotes a carboxy group, a straight-chain or branched C_{1-6}-alkoxycarbonyl group, a C_{5-7}-cycloalkoxycarbonyl or phenyl-C_{1-3}-alkoxycarbonyl group, an aminocarbonyl or C_{1-2}-alkylaminocarbonyl group or, if R_4 does not denote an aminosulphonyl-phenyl or N-(C_{1-5}-alkyl)-C_{1-3}-alkylaminocarbonyl-phenyl group, a di-(C_{1-2}-alkyl)-aminocarbonyl group,

R_3 denotes a hydrogen atom, a C_{1-6}-alkyl, C_{3-7}-cycloalkyl, trifluoromethyl or heteroaryl group,

a phenyl or naphthyl group, a phenyl or naphthyl group mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a trifluoromethyl, C_{1-3}-alkyl or C_{1-3}-alkoxy

group, whilst in the event of disubstitution the substituents may be identical or different and wherein the abovementioned unsubstituted as well as the mono- and disubstituted phenyl and naphthyl groups may additionally be substituted

by a hydroxy, hydroxy-C_{1-3}-alkyl or C_{1-3}-alkoxy-C_{1-3}-alkyl group,

by a cyano, carboxy, carboxy-C_{1-3}-alkyl, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylamino-carbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

by a nitro group,

by an amino, C_{1-3}-alkylamino, di-(C_{1-3}-alkyl)-amino or amino-C_{1-3}-alkyl group,

by a C_{1-3}-alkylcarbonylamino, N-(C_{1-3}-alkyl)-C_{1-3}-alkyl-carbonylamino, C_{1-3}-alkylcarbonylamino-C_{1-3}-alkyl, N-(C_{1-3}-alkyl)-C_{1-3}-alkylcarbonylamino-C_{1-3}-alkyl, C_{1-3}-alkyl-sulphonylamino, C_{1-3}-alkylsulphonylamino-C_{1-3}-alkyl, N-(C_{1-3}-alkyl)-C_{1-3}-alkylsulphonylamino-C_{1-3}-alkyl or aryl-C_{1-3}-alkylsulphonylamino group,

by a cycloalkylamino, cycloalkyleneimino, cyclo-alkyleneiminocarbonyl, cycloalkyleneimino-C_{1-3}-alkyl, cycloalkyleneiminocarbonyl-C_{1-3}-alkyl or cycloalkyleneiminosulphonyl-C_{1-3}-alkyl group having 4 to 7 ring members in each case, whilst in each case the methylene group in position 4 of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or

sulphur atom, by a sulphinyl, sulphonyl, -NH or -N(C_{1-3}-alkyl) group,

or by a heteroaryl or heteroaryl-C_{1-3}-alkyl group,

R_4 denotes a C_{3-7}-cycloalkyl group,

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group or replaced by an -NH or -N(C_{1-3}-alkyl) group,

or a phenyl group substituted by the group R_6, which may additionally be substituted by a fluorine, chlorine, bromine or iodine atom, by a C_{1-5}-alkyl, trifluoromethyl, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, aminosulphonyl, nitro or cyano group, wherein

R_6 denotes a hydrogen, fluorine, chlorine, bromine or iodine atom,

a cyano, nitro, C_{1-5}-alkyl, C_{3-7}-cycloalkyl, trifluoromethyl, phenyl, tetrazolyl or heteroaryl group,

a C_{1-3}-alkoxy group optionally sunstituted by 1 to 3 fluorine atoms, a C_{1-3}-alkoxy-C_{1-3}-alkoxy, phenyl-C_{1-3}-alkoxy, amino-C_{2-3}-alkoxy, C_{1-3}-alkylamino-C_{2-3}-alkoxy, di-(C_{1-3}-alkyl)-amino-C_{2-3}-alkoxy, phenyl-C_{1-3}-alkylamino-C_{2-3}-alkoxy, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino-C_{2-3}-alkoxy, C_{5-7}-cycloalkyleneimino-C_{2-3}-alkoxy or C_{1-3}-alkylmercapto group,

a carboxy, C_{1-4}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkyl-amino-carbonyl, N-(C_{1-5}-alkyl)-C_{1-3}-alkylaminocarbonyl, phenyl-C_{1-3}-alkylamino-carbonyl, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino-carbonyl, piperazinocarbonyl or N-(C_{1-3}-alkyl)-piperazinocarbonyl group,

a C_{1-3}-alkylaminocarbonyl or N-(C_{1-5}-alkyl)-C_{1-3}-alkylaminocarbonyl group wherein an alkyl moiety is substituted by a carboxy or C_{1-3}-alkoxycarbonyl group or is substituted in the 2 or 3 position by a di-(C_{1-3}-alkyl)-amino, piperazino, N-(C_{1-3}-alkyl)-piperazino or a 4- to 7-membered cycloalkyleneimino group,

a 4- to 7-membered cycloalkyleneimino group, wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

the cycloalkylene moiety may be fused to a phenyl ring or

one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl group and/or

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, –NH, –N(C_{1-3}–alkyl), –N(phenyl), –N(C_{1-3}–alkyl–carbonyl) or –N(benzoyl) group,

a C_{1-4}–alkyl group which may be substituted

by a hydroxy or C_{1-3}–alkoxy group,

by an amino, C_{1-7}–alkylamino, di–(C_{1-7}–alkyl)–amino, di–N–(C_{1-3}–alkyl)–amino–C_{2-3}–alkylamino, tri–N,N,N'–(C_{1-3}–alkyl)–amino–C_{2-3}–alkylamino, phenylamino, N–phenyl–C_{1-3}–alkyl–amino, phenyl–C_{1-3}–alkylamino, N–(C_{1-3}–alkyl)–phenyl–C_{1-3}–alkylamino or di–(phenyl–C_{1-3}–alkyl)–amino group,

by a C_{1-3}–alkylcarbonylamino, N–(C_{1-3}–alkyl)–C_{1-3}–alkylcarbonylamino, C_{1-3}–alkoxycarbonyl–C_{1-3}–alkylamino or N–(C_{1-3}–alkyl)–C_{1-3}–alkoxycarbonyl–C_{1-3}–alkylamino group,

by a C_{4-7}–cycloalkylamino, C_{4-7}–cycloalkyl–C_{1-3}–alkylamino or C_{4-7}–cycloalkenylamino group wherein position 1 of the ring is not involved in the double bond and wherein the abovementioned groups may each additionally be substituted at the amino–nitrogen atom by a C_{1-3}–alkyl group wherein some or all of the hydrogen atoms are replaced by fluorine atoms, by a C_{5-7}–cycloalkyl, C_{2-4}–alkenyl or C_{1-4}–alkyl group,

by a 4– to 7–membered cycloalkyleneimino group, wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

the cycloalkylene moiety may be fused to a phenyl group or to an oxazolo, imidazolo, thiazolo, pyridino, pyrazino or pyrimidino group optionally substituted by a fluorine, chlorine, bromine or iodine atom, by a nitro, C_{1-3}-alkyl, C_{1-3}-alkoxy or amino group or

one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl, C_{5-7}-cycloalkyl or phenyl group and/or

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a hydroxy, carboxy, C_{1-4}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl), -N(phenyl), -N(C_{1-3}-alkyl-carbonyl) or -N(benzoyl) group,

by a carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group or

by a 4- to 7-membered cycloalkyleneiminocarbonyl group,

an amino, pyrrolidino, piperidino, morpholino, benzoylamino or N-(C_{1-3}-alkyl)-benzoylamino group,

an N-(C_{1-3}-alkyl)-C_{2-4}-alkanoylamino group which is additionally substituted in the alkyl moiety by a carboxy or C_{1-3}-alkoxycarbonyl group,

a group of formula

$-N(R_8)-CO-(CH_2)_n-R_9$ (II),

wherein

R_8 denotes a hydrogen atom or a C_{1-3}-alkyl group,

n denotes one of the numbers 0, 1, 2 or 3 and

R_9 denotes an amino, C_{1-4}-alkylamino, phenylamino, N-(C_{1-4}-alkyl)-phenylamino, benzylamino, N-(C_{1-4}-alkyl)-benzylamino or di-(C_{1-4}-alkyl)-amino group, a 4- to 7-membered cycloalkyleneimino group, whilst in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl), -N(phenyl), -N(C_{1-3}-alkyl-carbonyl) or -N(benzoyl) group, or, if n denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

$-N(R_{10})-(CH_2)_m-(CO)_o-R_{11}$ (III),

wherein

R_{10} denotes a hydrogen atom, a C_{1-3}-alkyl group, a C_{1-3}-alkylcarbonyl, arylcarbonyl, phenyl-C_{1-3}-alkylcarbonyl, C_{1-3}-alkylsulphonyl, arylsulphonyl or phenyl-C_{1-3}-alkylsulphonyl group,

m denotes one of the numbers 1, 2, 3 or 4,

o denotes one of the numbers 0 or 1 and

R_{11} denotes an amino, C_{1-4}-alkylamino, phenylamino, N-(C_{1-4}-alkyl)-phenylamino, benzylamino, N-(C_{1-4}-alkyl)-benzylamino or di-(C_{1-4}-alkyl)-amino group, a 4- to 7-membered cycloalkyleneimino group, wherein the cycloalkylene moiety may be fused to a phenyl ring or in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl), -N(phenyl), -N(C_{1-3}-alkyl-carbonyl) or -N(benzoyl) group, a C_{1-3}-alkoxy group or a di-(C_{1-4}-alkyl)-amino-C_{1-3}-alkylamino group optionally substituted in the 1 position by a C_{1-3}-alkyl group,

or an N-(C_{1-3}-alkyl)-C_{1-5}-alkylsulphonylamino or N-(C_{1-3}-alkyl)-phenylsulphonylamino group wherein the alkyl moiety is additionally substituted by a cyano or carboxy group,

wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R_6 may be mono- or disubstituted by fluorine, chlorine, bromine or iodine atoms, by C_{1-5}-alkyl, trifluoromethyl, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, aminosulphonyl, nitro or cyano groups,

wherein the substituents may be identical or different, or two adjacent hydrogen atoms of the phenyl groups may be replaced by a methylenedioxy group,

and

R_5 denotes a hydrogen atom or a C_{1-3}-alkyl group,

wherein by an aryl group is meant a phenyl or naphthyl group optionally mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a trifluoromethyl, C_{1-3}-alkyl or C_{1-3}-alkoxy group and

by a heteroaryl group is meant a monocyclic 5- or 6-membered heteroaryl group optionally substituted by a C_{1-3}-alkyl group, wherein the 6-membered heteroaryl group contains one, two or three nitrogen atoms and the 5-membered heteroaryl group contains an imino group optionally substituted by a C_{1-3}-alkyl group, an oxygen or sulphur atom or an imino group optionally substituted by a C_{1-3}-alkyl group and an oxygen or sulphur atom or one or two nitrogen atoms, and moreover a phenyl ring may be fused to the abovementioned monocyclic heterocyclic groups via two adjacent carbon atoms,

the saturated alkyl and alkoxy moieties present in the groups defined above which contain more than 2 carbon atoms also include the branched isomers thereof such as, for example, the isopropyl, tert.butyl or isobutyl group, unless otherwise stated, and

additionally any carboxy, amino or imino group present may be substituted by a group which can be cleaved *in vivo*,

the isomers and the salts thereof.

A further subgroup of compounds of general formula I which deserves special mention is the subgroup wherein the substituent in the 6 position of the substituted indolinone of general formula I comprises a substituted amido group.

The above exemplified compounds, their tautomers, their stereoisomers or the physiologically acceptable salts thereof, as well as their manufacturing process, have been described in WO 01/27081, the content of which is incorporated herein by reference.

Further compounds in accordance with the above general formula I which are preferred within the meaning of the present invention are the following compounds:

(t) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(u) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(v) 3-Z-[1-(3-cyano-4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(w) 3-Z-[1-(3-methoxy-4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(x) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(y) 3-Z-[1-(4-(N-(N-(2-dimethylamino-ethyl)-N-methyl-aminomethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(z) 3-Z-[1-(4-(N-(di-(2-hydroxy-ethyl)-amino-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(aa) 3-Z-[1-(4-(N-(imidazol-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ab) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ac) 3-Z-[1-(4-(N-((4-methyl-[1,4]diazepan-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ad) 3-Z-[1-(4-(N-((1-methyl-piperidin-4-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ae) 3-Z-[1-(2,3-dimethyl-4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(af) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ag) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ah) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ai) 3-Z-[1-(4-(N-((3-dimethylamino-propyl)-aminocarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(aj) 3-Z-[1-anilino-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ak) 3-Z-[1-(4-(N-methyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(al) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(am) 3-Z-[1-(4-(4-methylpiperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(an) 3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(ao) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(ap) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(aq) 3-Z-[1-(4-(di-(2-hydroxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(ar) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(as) 3-Z-[1-(4-(N-(morpholin-4-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(at) 3-Z-[1-(4-(N-(N-(2-dimethylamino-ethyl)-N-methyl-aminomethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(au) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(av) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-carboxy-2-indolinone

(aw) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(ax) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(ay) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(az) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-(4-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(ba) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bb) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bc) 3-Z-[1-((1-methyl-piperidin-4-yl)-amino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bd) 3-Z-[1-(trans-4-dimethylamino-cyclohexylamino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(be) 3-Z-[1-(4-(2-diethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bf) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-propionyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bg) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bh) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bi) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bj) 3-Z-[1-(3-diethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bk) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bl) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bm) 3-Z-[1-anilino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bn) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bo) 3-Z-[1-(4-((2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bp) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bq) 3-Z-[1-(4-(N-methyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(br) 3-Z-[1-(4-methoxycarbonyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bs) 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bt) 3-Z-[1-(4-(N-(dimethylamino-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bu) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(bv) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(bw) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(bx) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(by) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(bz) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(ca) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cb) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cc) 3-Z-[1-(4-carbamoyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(cd) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(ce) 3-Z-[1-(4-((4-methyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(cf) 3-Z-[1-(4-((4-methyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cg)3-Z-[1-(4-((4-ethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(ch) 3-Z-[1-(4-(N-ethyl-N-(2-dimethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(ci) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-diethylcarbamoyl-2-indolinone

(cj) 3-Z-[1-(4-((cis-3,5-dimethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(ck) 3-Z-[1-(4-((4-ethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cl) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(cm) 3-Z-[1-(4-((cis-3,5-dimethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cn) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(co) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(cp) 3-Z-[1-(4-dimethylaminomethyl-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(cq) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(cr) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

(cs) 3-Z-[1-(4-dimethylaminomethyl-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

(ct) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

(cu) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone,

their tautomers, their stereoisomers or the physiologically acceptable salts thereof.

These compounds may be prepared analogously to the compounds of WO 01/27081 and using the methods described hereafter.

Abbreviations used:

HOBt = 1-hydroxy-1H-benzotriazole


TBTU = O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium-tetrafluoroborate

DEPC = diethyl pyrocarbonate

n.d. = not determined









Examples (t) to (al):


The following compounds of general formula II are prepared analogously to the compounds described in WO 01/27081:



(II)

Example	A	B	R^6	Formula	Mass spectrum	Melting point [°C]	R_f-value*
(t)	-H	-H	$-O(CH_2)_2-NMe_2$	$C_{27}H_{27}N_3O_4$	456 $[m-H]^-$	n. b.	0.30 (A)

(u)	-H	-H		$C_{31}H_{33}N_5O_4$	540 $[m+H]^+$	250-252	0.60 (B)
(v)	-CN	-H	-N(Me)-(CO)-CH_2-NMe_2	$C_{29}H_{27}N_5O_4$	510 $[m+H]^+$	163-165	0.35 (A)
(w)	-OMe	-H	-N(Me)-(CO)-CH_2-NMe_2	$C_{29}H_{30}N_4O_5$	515 $[m+H]^+$	160-163	0.40 (A)
(x)	-H	-H	-N(Me)-(CO)-CH_2-NH_2	$C_{26}H_{24}N_4O_4$	457 $[m+H]^+$	221	0.45 (C)
(y)	-H	-H		$C_{24}H_{21}N_3O_3$	542 $[m+H]^+$	265	n.d.
(Z)	-H	-H		$C_{30}H_{32}N_4O_6$	545 $[m+H]^+$	199-202	0.40 (A)
(aa)	-H	-H		$C_{29}H_{25}N_5O_4$	508 $[m+H]^+$	271	0.45 (A)
(ab)	-H	-H	-NH-(CO)-CH_2-NMe_2	$C_{27}H_{26}N_4O_4$	471 $[m+H]^+$	250-255	0.50 (A)
(ac)	-H	-H		$C_{32}H_{35}N_5O_4$	554 $[m+H]^+$	180-185	0.50 (D)
(ad)	-H	-H		$C_{32}H_{34}N_4O_4$	539 $[m+H]^+$	190-193	0.40 (D)
(ae)	$-CH_3$	$-CH_3$		$C_{32}H_{35}N_5O_4$	554 $[m+H]^+$	254-257	0.50 (C)
(af)	-H	-H		$C_{30}H_{31}N_5O_4$	526 $[m+H]^+$	170-175	0.40 (A)

(ag)	-H	-H		$C_{30}H_{31}N_5O_4$	526 $[m+H]^+$	205-208	0.40 (A)
(ah)	-H	-H	-N(Me)-(CO)-$(CH_2)_3$-NMe_2	$C_{30}H_{32}N_4O_4$	511 $[m-H]^-$	166-170	0.40 (C)
(ai)	-H	-H	-N(Me)-(CO)-NH-$(CH_2)_3$-NMe_2	$C_{30}H_{33}N_5O_4$	528 $[m+H]^+$	166-170	0.30 (E)
(aj)	-H	-H	-H	$C_{23}H_{18}N_2O_3$	371 $[m+H]^+$	275-280	0.80 (C)
(ak)	-H	-H	-N(SO_2Me)-CH_3	$C_{25}H_{23}N_3O_5S$	478 $[m+H]^+$	278-282	0.70 (C)

*Solvents:

(A): silica gel, methylene chloride/methanol 9:1

(B): aluminum oxide, methylene chloride/methanol 20:1

(C): silica gel, methylene chloride/methanol/ammonia 9:1:0.1

(D): silica gel, methylene chloride/methanol/ammonia 5:1:0.01

(E): silica gel, methylene chloride/methanol/ammonia 9:1:0.01

The following compound is prepared analogously:

(al) 3-Z-(1-cyclohexylamino-1-phenyl-methylene)-6-methoxycarbonyl-2-indolinone

R_f value: 0.60 (silica gel, methylene chloride/methanol = 9:1)


Melting point: 236-243 °C

$C_{23}H_{24}N_2O_3$



Mass spectrum: m/z = 377 $[m+H]^+$





Examples (am) to (av)

The following compounds of general formula III are prepared analogously to the compounds described in WO 01/27081:



(III)

Example	R^6	Formula	Mass spectrum	Melting point [°C]	R_f-value*
(am)		$C_{28}H_{28}N_4O_3$	467 $[m-H]^-$	275	0.50 (A)
(an)	$-CH_2-NHMe$	$C_{24}H_{21}N_3O_3$	398 $[m-H]^-$	287	0.70 (A)
(ao)		$C_{27}H_{25}N_3O_4$	454 $[m-H]^-$	335	0.70 (A)
(ap)	$-N(SO_2Me)-(CH_2)_2-NMe_2$	$C_{27}H_{28}N_4O_5S$	519 $[m-H]^-$	280	0.70 (A)

(aq)		$C_{27}H_{27}N_3O_5$	496 $[m+Na]^+$	256-257	0.75 (A)
(ar)		$C_{30}H_{31}N_5O_4$	526 $[m+H]^+$	346	0.60 (A)
(as)		$C_{29}H_{28}N_4O_5$	513 $[m+H]^+$	237-238	0.70 (A)
(at)		$C_{30}H_{33}N_5O_4$	528 $[m+H]^+$	238-240	0.50 (A)
(au)	$-O(CH_2)_2-NMe_2$	$C_{26}H_{25}N_3O_4$	444 $[m+H]^+$	n. b.	0.35 (B)

*Solvents:

(A): reversed phase RP8, methanol/brine(5%) = 4:1

(B): silica gel, methylene chloride/methanol 4:1

The following compound is prepared analogously:

(av) 3-Z-(1-cyclohexylamin-1-phenyl-methylene)-6-carboxy-2-indolinone

Rf value: 0.50 (silica gel, methylene chloride/methanol = 9:1)

Melting point: 347-350 °C

$C_{22}H_{22}N_2O_3$

Mass spectrum: m/z = 363 $[m+H]^+$

Examples (aw) to (az)

(aw) 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(ax) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinon

(ay) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(az) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-(4-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

Preparation of the starting compounds:

(I.1) 1-acetyl-3-(1-hydroxy-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone

6.00 g 1-acetyl-6-methoxycarbonyl-2-indolinone, 6.30 g 3-(2-ethoxycarbonyl-ethyl)-benzoic acid (preparation analogously to Tetrahedron **1997,** *53*, 7335-7340) and 9.10 g TBTU are dissolved in 80 ml of dimethylformamide, 13.5 ml diisopropylmethylamine and 4.34 g HOBt are added and the mixture is stirred for 12 hrs at ambient temperature. After this time the solvent is removed, diluted hydrochloric acid is added and the residue is recrystallized from methylene chloride/methanol.

Yield: 10.6 g (94 % of theory)

R_f value: 0.50 (silica gel, methylene chloride/methanol = 19:1)

Melting point: 80-84 °C

$C_{24}H_{23}NO_7$

Mass spectrum: m/z = 438 $[m+H]^+$

The following compounds are prepared analogously:

(I.2) 1-acetyl-3-(1-hydroxy-1-(4-(2-methoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone
prepared from 1-acetyl-6-methoxycarbonyl-2-indolinone and 4-(2-methoxycarbonyl-ethyl)-benzoic acid (preparation analogously to Tetrahedron **1997,** *53,* 7335-7340)
R_f value: 0.60 (silica gel, methylene chloride/methanol = 19:1)
Melting point: 188-192 °C
$C_{23}H_{21}NO_7$
Mass spectrum: m/z = 422 $[m-H]^-$

(II.1) 1-acetyl-3-(1-methoxy-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone
7.17 g trimethyloxoniumtetrafluoroborate are slowly added to a solution of 10.6 g 1-acetyl-3-(1-hydroxy-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone (starting material I.1) and 12.5 ml ethyl-diisopropylamine in 100 ml methylene chloride. After stirring for 4 hrs at ambient temperature another 3.50 g trimethyloxoniumtetrafluoroborate are added and the mixture is stirred for 12 hrs at ambient temperature. After that time the mixture is washed twice with water, the organic phase is dried over magnesium sulphate and the solvent is removed. The residue is purified over a silica gel column with methylene chloride/methanol (97:3) as eluant.
Yield: 4.56 g (42 % of theory)
R_f value: 0.90 (silica gel, methylene chloride/methanol = 20:1)
$C_{25}H_{25}NO_7$

Mass spectrum: m/z = 452 $[m+H]^+$

The following compounds are prepared analogously:

(II.2)1-acetyl-3-(1-methoxy-1-(4-(2-methoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone
prepared from 1-acetyl-3-(1-hydroxy-1-(4-(2-methoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone (starting material I.2)
R_f value: 0.80 (silica gel, methylene chloride/methanol = 19:1)
Melting point: 112-117 °C
$C_{24}H_{23}NO_7$
Mass spectrum: m/z = 438 $[m+H]^+$

(III.1) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone
1.2 g 1-acetyl-3-(1-methoxy-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone (starting material II.1) and 0.32 g 4-(dimethylaminomethyl)-aniline are dissolved in 10 ml of dimethylformamide and stirred for 3 days at 110°C. After cooling the solvent is evaporated, the residue is taken up in 5 ml of methanol and 200 mg 20 percent sodiumethylat-solution in ethanol are added. The mixture is stirred for 1.5 hrs at ambient temperature, the solvent is removed and the residue is taken up in water. The aqueous phase is three times extracted with ethyl acetate and the combined organic phases are dried over sodium sulphate. After evaporation of the solvent the

residue is purified over a silica gel column with methylene chloride/methanol (9:1) as eluant.

Yield: 0.33 g (35% of theory),

R_f value: 0.35 (silica gel, methylene chloride/methanol = 9:1)

Melting point: 129-134 °C

$C_{31}H_{33}N_3O_5$

Mass spectrum: m/z = 528 $[m+H]^+$

The following compounds are prepared analogously:

(III.2) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

prepared from 1-acetyl-3-(1-methoxy-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone (starting material II.1)

R_f value: 0.30 (silica gel, methylene chloride/methanol = 9:1)

Melting point: 174-177 °C

$C_{32}H_{35}N_3O_5$

Mass spectrum: m/z = 542 $[m+H]^+$

(III.3)3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

prepared from 1-acetyl-3-(1-methoxy-1-(3-(2-ethoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone (starting material II.1)

R_f value: 0.45 (silica gel, methylene chloride/methanol = 9:1)

Melting point: 102 °C

$C_{32}H_{30}N_4O_5$

Mass spectrum: m/z = 551 $[m+H]^+$

(III.4) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-(4-(2-methoxycarbonyl-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

prepared from 1-acetyl-3-(1-methoxy-1-(4-(2-methoxycarbonyl-ethyl)-phenyl)-methylene)-6-methoxycarbonyl-2-indolinone (starting material II.2)

R_f value: 0.50 (silica gel, methylene chloride/methanol = 9:1)


Melting point: 226-229 °C

$C_{30}H_{31}N_3O_5$






Mass spectrum: m/z = 512 $[m-H]^-$

Preparation of the final compounds:

The following compounds of general formula IV are prepared analogously to the compounds described in WO 01/27081, starting from the above mentioned starting materials:



(IV)

Example	R^3	R^6	Starting compound	Formula	Mass spectrum	Melting point [°C]	R_f value*
(aw)		$-CH_2-NMe_2$	III.1	$C_{29}H_{29}N_3O_5$	500 $[m+H]^+$	163-167	0.40 (A)
(ax)		$-(CH_2)_2-NMe_2$	III.2	$C_{30}H_{31}N_3O_5$	514 $[m+H]^+$	248-255	0.35 (A)
(ay)			III.3	$C_{30}H_{26}N_4O_5$	523 $[m+H]^+$	184-190	0.35 (A)
(az)		$-CH_2-NMe_2$	III.4	$C_{29}H_{29}N_3O_5$	498 $[m-H]^-$	190-195	0.20 (B)

*Solvents:

(A): Reversed Phase RP8, methanol/brine(5%) = 4:1

(B): silica gel, methylene chloride/methanol 9:1

Examples (ba) to (cn)

Preparation of the starting compounds:

(IV) 3-(1-hydroxy-1-phenyl-methylene)-6-carboxy-2-indolinone

11.0 g 1-acetyl-3-(1-methoxy-1-phenyl-methylene)-6-methoxycarbonyl-2-indolinone (preparation described in WO

01/27081) are disolved in 500 ml of methanol and 160 ml of 1N sodium hydroxide solution are added. The mixture is stirred for 1 hr at ambient temperature and for 6 hrs at reflux. After that time another 20 ml of 1N sodium hydroxide solution are added and the mixture is stirred for another 3 hrs at reflux. 160 ml of 1N hydrochloric acid are added, the resulting residue is filtered off and dried at 100°C. The residue is used without further purification.
Yield: 7.60 g (86 % of theory)

(V.1) 3-(1-hydroxy-1-phenyl-methylene)-6-(N-ethyl-methylcarbamoyl)-2-indolinone
5.50 g 3-(1-hydroxy-1-phenyl-methylene)-6-carboxy-2-indolinone (starting material IV), 7.54 g TBTU, 3.60 g HOBt and 17.1 ml ethyldiisopropylamine are dissolved in 200 ml of dimethylformamide. 2.70 ml of a 94-percent solution of N-methyl-ethylamine are added and the mixture is stirred for 12 hrs at ambient temperature. After that time the solvent is evaporated and the residue is purified over a silica gel column with methylene chloride/methanol/ammonia (9:1:0.1) as eluant.
Yield: 6.10 g (97 % of theory)
R_f value: 0.35 (silica gel, methylene chloride/methanol/ammonia = 9:1:0.1)
$C_{19}H_{18}N_2O_3$
Mass spectrum: m/z = 323 $[m+H]^+$

The following compound is prepared analogously:

(V.2) 3-(1-hydroxy-1-phenyl-methylene)-6-ethylcarbamoyl-2-indolinone

prepared from 3-(1-hydroxy-1-phenyl-methylene)-6-carboxy-2-indolinone (starting material IV) und ethylamine

$C_{18}H_{16}N_2O_3$

Mass spectrum: m/z = 309 $[m+H]^+$

Preparation of the final compounds:

(ba) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-(N-ethyl-methylcarbamoyl)-2-indolinone

250 mg 3-(1-hydroxy-1-phenyl-methylene)-6-(N-ethyl-methylcarbamoyl)-2-indolinone (starting material V.1) and 382 mg 4-(2-dimethylamino-ethyl)-aniline are dissolved in 3 ml of tetrahydrofuran, 569 ml trimethylsilylimidazole are added and the mixture is stirred at 170°C in a microwave oven. After cooling the solvent is evaporated and the residue is taken up in water. The residue is filtered off and vacuum-dried at 90°C.

Yield: 0.18 g (50% of theory),


R_f value: 0.30 (silica gel, methylene chloride/methanol/ammonia = 9:1:0.1)

Melting point: 195-200 °C






$C_{29}H_{32}N_4O_2$








Mass spectrum: m/z = 469 $[m+H]^+$









The following compounds of general formula V are prepared analogously to the above compound (ba), following the procedures described in WO 01/27081:




















(V)

Example	X	R^4	Formula	Mass spectrum	Melting point [°C]	R_f value*
(bb)	$-CH_3$		$C_{28}H_{30}N_4O_2$	455 $[m+H]^+$	239-243	0.35 (A)
(bc)	$-CH_3$		$C_{25}H_{30}N_4O_2$	419 $[m+H]^+$	267-271	0.35 (B)
(bd)	$-CH_3$		$C_{27}H_{34}N_4O_2$	447 $[m+H]^+$	133-138	0.30 (B)
(be)	$-CH_3$		$C_{31}H_{36}N_4O_3$	513 $[m+H]^+$	191-196	0.45 (B)
(bf)	$-CH_3$		$C_{33}H_{39}N_5O_3$	554 $[m+H]^+$	258-262	0.40 (B)

(bg)	$-CH_3$		$C_{33}H_{38}N_6O_3$	567 $[m+H]^+$	214-218	0.20 (B)
(bh)	$-CH_3$		$C_{25}H_{29}N_3O_2$	404 $[m+H]^+$	239-242	0.70 (A)
(bi)	$-CH_3$		$C_{31}H_{35}N_5O_3$	526 $[m+H]^+$	237-240	0.30 (B)
(bj)	$-CH_3$		$C_{30}H_{34}N_4O_2$	483 $[m+H]^+$	105-108	0.40 (B)
(bk)	$-CH_3$		$C_{30}H_{33}N_5O_3$	512 $[m+H]^+$	208-211	0.40 (B)
(bl)	$-CH_3$		$C_{30}H_{35}N_5O_4S$	562 $[m+H]^+$	197-201	0.40 (B)
(bm)	$-CH_3$		$C_{25}H_{23}N_3O_2$	398 $[m+H]^+$	296-301	0.40 (B)
(bn)	$-CH_3$		$C_{28}H_{30}N_4O_2$	455 $[m+H]^+$	243-247	0.30 (A)
(bo)	$-H$		$C_{28}H_{30}N_4O_2$	455 $[m+H]^+$	328-332	0.30 (A)

(bp)	$-CH_3$		$C_{32}H_{37}N_5O_3$	540 $[m+H]^+$	224-228	0.25 (A)
(bq)	$-CH_3$		$C_{27}H_{28}N_4O_4S$	505 $[m+H]^+$	265-269	0.40 (B)
(br)	$-CH_3$		$C_{27}H_{25}N_3O_4$	456 $[m+H]^+$	254-257	0.60 (B)
(bs)	$-CH_3$		$C_{26}H_{23}N_3O_4$	442 $[m+H]^+$	316-321	0.10 (B)
(bt)	$-CH_3$		$C_{30}H_{33}N_5O_5S$	576 $[m+H]^+$	258-262	0.35 (B)
(bu)	$-H$		$C_{28}H_{30}N_4O_3$	471 $[m+H]^+$	308-311	0.35 (B)
(bv)	$-H$		$C_{32}H_{36}N_6O_3$	553 $[m+H]^+$	279-283	0.60 (C)
(bw)	$-H$		$C_{29}H_{33}N_5O_4S$	548 $[m+H]^+$	213-217	0.35 (B)

(bx)	$-CH_3$		$C_{29}H_{32}N_4O_3$	485 $[m+H]^+$	218-222	0.40 (A)
(by)	-H		$C_{29}H_{31}N_5O_3$	498 $[m+H]^+$	130-134	0.35 (D)
(bz)	-H		$C_{27}H_{28}N_4O_2$	441 $[m+H]^+$	341-344	0.45 (D)
(ca)	-H		$C_{30}H_{33}N_5O_3$	512 $[m+H]^+$	266-270	0.40 (D)
(cb)	-H		$C_{31}H_{35}N_5O_3$	526 $[m+H]^+$	198-202	0.40 (D)
(cc)	$-CH_3$		$C_{26}H_{24}N_4O_3$	441 $[m+H]^+$	290-295	0.25 (B)
(cd)	$-CH_3$		$C_{32}H_{37}N_5O_3$	540 $[m+H]^+$	120-126	0.40 (B)
(ce)	$-CH_3$		$C_{31}H_{33}N_5O_3$	524 $[m+H]^+$	100-105	0.50 (B)
(cf)	-H		$C_{30}H_{31}N_5O_3$	510 $[m+H]^+$	288-292	0.40 (A)

(cg)	$-CH_3$		$C_{32}H_{35}N_5O_3$	538 $[m+H]^+$	157-163	0.30 (B)
(ch)	$-CH_3$		$C_{32}H_{37}N_5O_3$	540 $[m+H]^+$	162-169	0.20 (B)
(ci)	$-CH_2CH_3$		$C_{34}H_{40}N_6O_3$	581 $[m+H]^+$	195-198	0.50 (E)
(cj)	$-CH_3$		$C_{32}H_{35}N_5O_3$	538 $[m+H]^+$	238-242	0.35 (B)
(ck)	-H		$C_{31}H_{33}N_5O_3$	524 $[m+H]^+$	127-130	0.50 (D)
(cl)	-H		$C_{31}H_{35}N_5O_3$	526 $[m+H]^+$	250-253	0.40 (D)
(cm)	-H		$C_{32}H_{35}N_5O_3$	524 $[m+H]^+$	217-220	0.40 (D)
(cn)	-H		$C_{29}H_{33}N_5O_4S$	560 $[m-H]^-$	171-175	0.45 (D)

*Solvents:
(A): silica gel, methylene chloride/methanol 9:1
(B): silica gel, methylene chloride/methanol/ammonia 9:1:0.1
(C): aluminum oxide, methylene chloride/methanol 9:1
(D): aluminum oxide, methylene chloride/methanol 19:1
(E): Reversed Phase RP8, acetonitrile/water/trifluoroacetic acid = 1:1:0.01


Examples (co) to (cq)

Preparation of the starting compounds:


(VI) 1-acetyl-3-(1-ethoxy-methylene)-6-methoxycarbonyl-2-indolinone
8.00 g 1-acetyl-6-methoxycarbonyl-2-indolinone and 17.2 ml triethyl orthoformate are dissolved in 70 ml of acetic anhydride and stirred for 5.5 hrs at 110°C. After cooling the residue is filtered off, washed with ether and vacuum-dried at 100°C.
Yield: 8.80 g (89 % of theory)
R_f value: 0.35 (silica gel, petrol ether/methylene chloride/ethylacetate = 5:4:1)
Melting point: 187-189 °C
$C_{15}H_{15}NO_5$
Mass spectrum: m/z = 290 $[m+H]^+$

Preparation of the final compounds:

The following compounds of general formula VI are prepared analogously to the compounds described in WO 01/27081:



(VI)

Example	R^6	Formula	Mass spectrum	Melting point [°C]	R_f value*
(co)	$-NMe-(CO)-CH_2-NMe_2$	$C_{22}H_{24}N_4O_4$	409 $[m+H]^+$	250-255	0.40 (A)
(cp)	$-CH_2-NMe_2$	$C_{20}H_{21}N_3O_3$	352 $[m+H]^+$	234-238	0.35 (A)
(cq)		$C_{25}H_{29}N_5O_4$	464 $[m+H]^+$	203-207	0.45 (A)

*Solvents:

(A): silica gel, methylene chloride/methanol/ammonia 9:1:0.1

Examples (cr) to (cu)

Preparation of the starting compounds:

(VII) 3-(1-hydroxy-methylene)-6-carboxy-2-indolinone

5.00 g 1-acetyl-3-(1-ethoxy-methylene)-6-methoxycarbonyl-2-indolinone (starting material VI) are dissoled in 150 ml of methanol and 86.4 ml of 1N sodium hydroxide solution are added. The mixture is refluxed for 8.5 hrs. After that time 86.4 ml of 1N hydrochloric acid are added. The residue is filtered off and dried at 90°C.

Yield: 2.50 g (71 % of theory)

$C_{10}H_7NO_4$

Mass spectrum: m/z = 206 $[m+H]^+$

(VIII) 3-(1-hydroxy-methylene)-6-ethylcarbamoyl-2-indolinone

400 mg 3-(1-hydroxy-methylene)-6-carboxy-2-indolinone (starting material VII), 689 mg TBTU, 291 mg HOBt and 1.35 ml triethylamine are dissolved in 20 ml of dimethylformamide. At 0°C 1.95 ml of a 2M ethylamine-solution in THF are added and the mixture is stirred for additional 12 hrs at ambient temperature. After that time the solvent is evaporated and the residue is purified over a silica gel cloumn with methylene chloride/ethanol/acetic acid (5:1:0.01) as eluant.

Yield: 160 mg (35 % of theory)

R_f value: 0.20 (silica gel, methylene chloride/ethanol/acetic acid = 5:1:0.01)

Melting point: 146-150 °C

$C_{12}H_{12}N_2O_3$

Mass spectrum: m/z = 233 $[m+H]^+$

Preparation of the final compounds:

(cr) 3-Z-[1-(4-(N-(4-methyl-piperazin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

160 mg 3-(1-hydroxy-methylene)-6-ethylcarbamoyl-2-indolinone (starting material VIII) and 543 mg N-[(4-methyl-piperazin-1-yl)-methylcarbonyl]-N-methyl-p-phenylendiamine are dissolved in 3 ml of tetrahydrofuran, 506 ml trimethylsilylimidazole are added and the mixture is stirred for 25 minutes at 170°C in a microwave oven. After cooling the solvent is evaporated and the residue is purified over an aluminum oxide column (activity 2-3) with methylene chloride/ethanol (19:1) as eluant. The residue is recrystallized from ether and vacuum-dried at 80°C.

Yield: 0.17 g (52% of theory),


R_f value: 0.60 (aluminum oxide, methylene chloride/methanol = 9:1)

Melting point: 255-260 °C




$C_{26}H_{32}N_6O_3$

Mass spectrum: m/z = 477 $[m+H]^+$

The following compounds of general formula VII are prepared analogously to the above compound (ct), following the procedures described in WO 01/27081:



(VII)

Example	X	R^4	Formula	Mass spectrum	Melting point [°C]	R_f value*
(cs)	-H		$C_{23}H_{27}N_5O_3$	422 $[m+H]^+$	280-283	0.70 (A)
(ct)	-H		$C_{24}H_{28}N_4O_2$	405 $[m+H]^+$	245-248	0.80 (A)
(cu)	-H		$C_{25}H_{31}N_5O_3$	450 $[m+H]^+$	130	0.40 (B)

*Solvents:

(A): aluminum oxide, methylene chloride/methanol 9:1

(B): silica gel, methylene chloride/ethanol/ammonia 5:2:0.01

Tautomers, stereoisomers or physiologically acceptable salts of these compounds are also contemplated within the scope of the present invention, and may be obtained using the methods described in WO 01/27081, the content of which is herein incorporated by reference.

A particularly preferred compound is the monoethanesulphonate salt of 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone, disclosed

for example in WO 04/13099, the content of which is incorporated herein by reference.

The metabolites of the compound 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone-monoethanesulphonate and the prodrugs of this compound or of these metabolites obtained via, for example, chemical or non-chemical derivatization of the entire molecule or of one or more chemical groups on the molecule, are also contemplated compounds within the meaning of the present invention. In this matter, reference is made to WO 04/13099, which describes metabolites and prodrugs of the compound 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone-monoethanesulphonate.

The following list of specific compounds is illustrative of the present invention, without constituting any limitation of its scope:

(1) 3-Z-(1-anilino-1-phenyl-methylene)-6-ethoxycarbonyl-2-indolinone

(2) 3-Z-[1-(4-nitro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(3) 3-Z-[1-(4-fluoro-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone

(4) 3-Z-[1-(4-chloro-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone

(5) 3-Z-[1-(4-iodo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(6) 3-Z-[1-(4-cyano-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone

(7) 3-Z-[1-(4-methoxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(8) 3-Z-[1-(4-ethoxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(9) 3-Z-[1-(4-trifluoromethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(10) 3-Z-[1-(4-methyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(11) 3-Z-[1-(4-methylmercapto-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(12) 3-Z-[1-(4-aminomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(13) 3-Z-[1-(4-(isopropylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(14) 3-Z-[1-(4-(anilinomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(15) 3-Z-[1-(4-(propylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(16) 3-Z-[1-(4-(butylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(17) 3-Z-[1-(4-(isobutylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(18) 3-Z-[1-(4-(cyclohexylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(19) 3-Z-[1-(4-(benzylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(20) 3-Z-[1-(4-((N-ethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(21) 3-Z-[1-(4-((N-methyl-N-propyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(22) 3-Z-[1-(4-((N-isopropyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(23) 3-Z-[1-(4-((N-ethyl-N-propyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(24) 3-Z-[1-(4-((N-ethyl-N-isopropyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(25) 3-Z-[1-(4-(dipropylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(26) 3-Z-[1-(4-(diisopropylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(27) 3-Z-[1-(4-((N-benzyl-N-ethyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(28) 3-Z-[1-(4-(dibenzylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(29) 3-Z-[1-(4-(3,6-dihydro-2H-pyridin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(30) 3-Z-[1-(4-(3,5-dimethyl-piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(31) 3-Z-[1-(4-(azepan-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(32) 3-Z-[1-(4-(piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(33) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(34) 3-Z-[1-(4-(thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(35) 3-Z-[1-(4-(1-oxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(36) 3-Z-[1-(4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(37) 3-Z-[1-(4-(acetylamino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(38) 3-Z-[1-(4-(2-amino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(39) 3-Z-[1-(4-(2-methylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(40) 3-Z-[1-(4-(2-ethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(41) 3-Z-[1-(4-(2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(42) 3-Z-[1-(4-(2-piperidin-1-yl-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(43) 3-Z-[1-(4-(2-acetylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(44) 3-Z-[1-(4-(3-amino-propyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(45) 3-Z-[1-(4-(3-dimethylamino-propyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(46) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(47) 3-Z-[1-(4-(N-methylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(48) 3-Z-[1-(4-(N-ethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(49) 3-Z-[1-(4-(N-diethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(50) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(51) 3-Z-[1-(4-(N-(morpholin-4-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(52) 3-Z-[1-(4-(N-(piperazin-1-yl-methylcarbonyl)-N-methyl-amino]-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(53) 3-Z-[1-(4-(N-(2-amino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(54) 3-Z-[1-(4-(N-(2-methylamino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(55) 3-Z-[1-(4-(N-(2-diethylamino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(56) 3-Z-[1-(4-(N-acetyl-N-(2-aminoethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(57) 3-Z-[1-(4-(N-acetyl-N-(2-methylamino-ethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(58) 3-Z-[1-(4-(N-acetyl-N-(2-methylamino-propyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(59) 3-Z-[1-(4-(N-acetyl-N-(2-piperidin-1-yl-ethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(60) 3-Z-[1-(4-(N-acetyl-N-(aminocarbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(61) 3-Z-[1-(4-(N-acetyl-N-(dimethylaminocarbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(62) 3-Z-[1-(4-(N-acetyl-N-(piperidin-1-yl-carbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(63) 3-Z-[1-(4-(N-methyl-N-(aminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(64) 3-Z-[1-(4-(N-methyl-N-(methylaminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(65) 3-Z-[1-(4-(N-methyl-N-(dimethylaminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(66) 3-Z-[1-(4-(N-methyl-N-(piperidin-1-yl-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(67) 3-Z-[1-(4-(N-(2-aminoethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(68) 3-Z-[1-(4-(N-(2-methylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(69) 3-Z-[1-(4-(N-(2-ethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(70) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(71) 3-Z-[1-(4-(N-(2-pyrrolidin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(72) 3-Z-[1-(4-(N-(2-piperidin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(73) 3-Z-[1-(4-(N-(2-piperazin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(74) 3-Z-[1-(4-(N-(2-(morpholin-4-yl)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(75) 3-Z-[1-(4-(N-(aminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(76) 3-Z-[1-(4-(N-(methylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(77) 3-Z-[1-(4-(N-(ethylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(78) 3-Z-[1-(4-(N-(N-(2-dimethylamino-ethyl)-N-methyl-amino)-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(79) 3-Z-[1-(4-(N-(diethylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(80) 3-Z-[1-(4-(N-(pyrrolidin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(81) 3-Z-[1-(4-(N-(piperidin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(82) 3-Z-[1-(4-(N-(piperazin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(83) 3-Z-[1-(4-(N-((morpholin-4-yl)-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(84) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(85) 3-Z-[1-(4-(3-dimethylamino-propoxy)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(86) 3-Z-[1-(4-(aminocarbonylmethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(87) 3-Z-[1-(4-(2-aminocarbonyl-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(88) 3-Z-[1-(4-(pyridin-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(89) 3-Z-[1-(4-(pyridine-3-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(90) 3-Z-[1-(4-(pyridin-4-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(91) 3-Z-[1-(4-(N-acetyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(92) 3-Z-[1-(4-(N-ethylcarbonyl-N-(dimethylaminocarbonyl-methyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(93) 3-Z-[1-(carbamoylmethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(94) 3-Z-[1-(4-dimethylcarbamoylmethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(95) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(96) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(97) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(98) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(99) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-ethylidene]-6-ethoxycarbonyl-2-indolinone

(100) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(101) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(102) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(103) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(104) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(105) 3-Z-[1-(4-tetrazol-5-yl-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(106) 3-Z-[1-(4-tetrazol-5-yl-anilino)-ethylidene]-6-ethoxycarbonyl-2-indolinone

(107) 3-Z-[1-(4-tetrazol-5-yl-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(108) 3-Z-[1-(4-tetrazol-5-yl-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(109) 3-Z-[1-(4-carboxy-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(110) 3-Z-[1-(4-carboxy-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(111) 3-Z-[1-(4-carboxy-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(112) 3-Z-[1-(4-(N-(3-dimethylamino-propionyl)-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(113) 3-Z-[1-(4-(N-(4-dimethylamino-butyryl)-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(114) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-(2-dimethylamino-ethylsulphonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(115) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-(3-dimethylamino-propylsulphonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(116) 3-Z-[1-(4-((2-hydroxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(117) 3-Z-[1-(4-((2-methoxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(118) 3-Z-[1-(4-((2-dimethylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(119) 3-Z-[1-(4-((3-dimethylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(120) 3-Z-[1-(4-((N-tert.butoxycarbonyl-2-amino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(121) 3-Z-[1-(4-((N-tert.butoxycarbonyl-3-amino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(122) 3-Z-[1-(4-((2-amino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(123) 3-Z-[1-(4-((3-amino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(124) 3-Z-[1-(4-((2-acetylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(125) 3-Z-[1-(4-((3-acetylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(126) 3-Z-[1-(4-((2-methylsulphonylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(127) 3-Z-[1-(4-((3-methylsulphonylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(128) 3-Z-[1-(4-(N-(N-tert.butoxycarbonyl-2-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(129) 3-Z-[1-(4-(N-(2-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(130) 3-Z-[1-(4-(N-(2-acetylamino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(131) 3-Z-[1-(4-(N-(2-methylsulphonylamino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(132) 3-Z-[1-(4-(carboxymethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(133) 3-Z-[1-(4-(ethoxycarbonylmethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(134) 3-Z-[1-(4-(carbamoylmethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(135) 3-Z-[1-(4-(dimethylcarbamoyl-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(136) 3-Z-[1-(4-(methylcarbamoyl-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(137) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(138) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-nitro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(139) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-acetylamino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(140) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methylsulphonylamino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(141) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-cyano-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(142) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-hydroxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(143) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methoxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(144) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-ethoxycarbonyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(145) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-carboxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(146) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-carbamoyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(147) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-chloro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(148) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-fluoro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(149) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-bromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(150) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(151) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-trifluoromethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(152) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3,5-dibromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(153) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3,5-dichloro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(154) 3-Z-[1-(4-(dimethylaminomethyl)-3-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(155) 3-Z-[1-(4-(dimethylaminomethyl)-3-nitro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(156) 3-Z-[1-(4-(dimethylaminomethyl)-3-acetylamino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(157) 3-Z-[1-(4-(dimethylaminomethyl)-3-(methylsulphonylamino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(158) 3-Z-[1-(4-(dimethylaminomethyl)-3-cyano-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(159) 3-Z-[1-(4-(dimethylaminomethyl)-3-hydroxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(160) 3-Z-[1-(4-(dimethylaminomethyl)-3-methoxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(161) 3-Z-[1-(4-(dimethylaminomethyl)-3-(ethoxycarbonyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(162) 3-Z-[1-(4-(dimethylaminomethyl)-3-carboxy-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(163) 3-Z-[1-(4-(dimethylaminomethyl)-3-carbamoyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(164) 3-Z-[1-(4-(dimethylaminomethyl)-3-chloro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(165) 3-Z-[1-(4-(dimethylaminomethyl)-3-fluoro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(166) 3-Z-[1-(4-(dimethylaminomethyl)-3-bromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(167) 3-Z-[1-(4-(dimethylaminomethyl)-3-methyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(168) 3-Z-[1-(4-(dimethylaminomethyl)-3-trifluoromethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(169) 3-Z-[1-(4-(dimethylaminomethyl)-3,5-dibromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(170) 3-Z-[1-(4-(dimethylaminomethyl)-3,5-dichloro-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(171) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(172) 3-Z-[1-(4-(N-(imidazo-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(173) 3-Z-[1-(4-(N-(phthalimido-2-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(174) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(175) 3-Z-[1-(4-(N-acetylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(176) 3-Z-[1-(4-(N-methylsulphonylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(177) 3-Z-[1-(4-(N-((N-(2-methoxyethyl)-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(178) 3-Z-[1-(4-(N-((N-(2-dimethylaminoethyl)-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(179) 3-Z-[1-(4-(N-((di-(2-hydroxyethyl)-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(180) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(181) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-ethylidene]-6-ethoxycarbonyl-2-indolinone

(182) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(183) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(184) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-methylene]-6-ethoxycarbonyl-2-indolinone

(185) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-ethylidene]-6-ethoxycarbonyl-2-indolinone

(186) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-propylidene]-6-ethoxycarbonyl-2-indolinone

(187) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-butylidene]-6-ethoxycarbonyl-2-indolinone

(188) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(189) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(190) 3-Z-[1-(4-((imidazolidin-2,4-dion-5-ylidene)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(191) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(192) 3-Z-[1-(4-(N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(193) 3-Z-[1-(4-(2-oxo-pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(194) 3-Z-[1-(4-(N-aminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(195) 3-Z-[1-(4-(N-cyanomethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(196) 3-Z-[1-(4-(2-(imidazol-4-yl)-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(197) 3-Z-[1-(4-((2-(N-benzyl-N-methyl-amino)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(198) 3-Z-[1-(4-cyclohexylamino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(199) 3-Z-[1-(4-(imidazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(200) 3-Z-[1-(4-(imidazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(201) 3-Z-[1-(N-methyl-piperidine-4-yl-amino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(202) 3-Z-[1-(4-(imidazol-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(203) 3-Z-[1-(4-((4-hydroxy-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(204) 3-Z-[1-(4-((4-methoxy-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(205) 3-Z-[1-(4-benzyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(206) 3-Z-[1-(4-(N-(3-trifluoroacetylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(207) 3-Z-[1-(4-(4-tert.butoxycarbonyl-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(208) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(209) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(210) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(211) 3-Z-[1-(4-((3-(N-benzyl-N-methyl-amino)-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(212) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(213) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-butyryl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(214) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-isobutyryl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(215) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-benzoyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(216) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(217) 3-Z-[1-(4-(4-hydroxymethyl-piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(218) 3-Z-[1-(4-(2-(4-hydroxy-piperidin-1-yl)-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(219) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-propylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(220) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-butylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(221) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-phenylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(222) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-benzylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(223) 3-Z-[1-(4-((imidazolidin-2,4-dion-5-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(224) 3-Z-[1-(4-((3-hydroxy-pyrrolidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(225) 3-Z-[1-(4-(cyclohexylyl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(226) 3-Z-[1-(4-(cyclohexyl-carbonyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(227) 3-Z-[1-(4-diethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(228) 3-Z-[1-(4-(N-(n-hexyl)-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(229) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(furan-2-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(230) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(2-methoxy-benzoyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(231) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(pyridine-3-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(232) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(phenyl-acetyl)-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(233) 3-Z-[1-(4-(imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(234) 3-Z-[1-(4-(1-ethyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(235) 3-Z-[1-(4-(1-benzyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(236) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-isopropylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(237) 3-Z-[1-(4-(N-((4-benzyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(238) 3-Z-[1-(4-(N-(pyrrolidin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(239) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-bromo-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(240) 3-Z-[1-(4-(5-methyl-imidazol-4-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(241) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(242) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(243) 3-Z-[1-(4-(N-butyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(244) 3-Z-[1-(4-(N-((N-aminocarbonylmethyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(245) 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(246) 3-Z-[1-(4-(N-(di-(2-methoxyethyl)-amino-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(247) 3-Z-[1-(4-(N-((2-(4-tert.butoxycarbonyl-piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(248) 3-Z-[1-(4-(N-((2-(piperidin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(249) 3-Z-[1-(4-(N-((2-(N-benzyl-N-methyl-amino)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(250) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(251) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(252) 3-Z-[1-(4-(N-((4-tert.butoxycarbonyl-piperazin-1-yl)-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(253) 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(254) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(255) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(256) 3-Z-[1-(4-(1,2,4-triazol-2-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(257) 3-Z-[1-(4-(1,2,3-triazol-2-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(258) 3-Z-[1-(4-(1,2,3-triazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(259) 3-Z-[1-(4-((N-aminocarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(260) 3-Z-[1-(4-((di-(2-methoxy-ethyl)-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(261) 3-Z-[1-(4-((di-(2-hydroxy-ethyl)-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(262) 3-Z-[1-(4-((N-ethoxycarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(263) 3-Z-[1-(4-(azetidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(264) 3-Z-[1-(4-(N-propyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(265) 3-Z-[1-(4-((N-(2-(2-methoxy-ethoxy)-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(266) 3-Z-[1-(4-((N-(tert.butoxycarbonyl-3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(267) 3-Z-[1-(4-((N-(methylcarbamoyl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(268) 3-Z-[1-(4-((N-(dimethylcarbamoyl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(269) 3-Z-[1-(4-((N-propyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(270) 3-Z-[1-(4-((N-(2-dimethylamino-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(271) 3-Z-[1-(4-((N-(3-dimethylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(272) 3-Z-[1-(4-((N-(2-methoxy-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(273) 3-Z-[1-(4-((N-(2-hydroxy-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(274) 3-Z-[1-(4-((N-(dioxolan-2-yl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(275) 3-Z-[1-(4-(3-oxo-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(276) 3-Z-[1-(4-(N-(piperazin-1-yl-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(277) 3-Z-[1-(4-(N-((2-(piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(278) 3-Z-[1-(4-((N-(3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(279) 3-Z-[1-(4-(N-(3-methylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(280) 3-Z-[1-(4-Ureidomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(281) 3-Z-[1-(4-guanidinomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(282) 3-Z-[1-(4-(N-methylsulphonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(283) 3-Z-[1-(4-(4-benzoyl-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(284) 3-Z-[1-(4-((N-(3-acetylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(285) 3-Z-[1-(4-((N-(3-methylsulphonylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(286) 3-Z-[1-(4-((N-carboxymethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(287)3-Z-(1-anilino-1-phenyl-methylene)-6-methoxycarbonyl-2-indolinone

(288) 3-Z-[1-(4-nitro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(289) 3-Z-[1-(4-fluoro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(290) 3-Z-[1-(4-chloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(291) 3-Z-[1-(4-bromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(292) 3-Z-[1-(4-iodo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(293) 3-Z-[1-(4-cyano-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(294) 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(295) 3-Z-[1-(4-methoxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(296) 3-Z-[1-(4-ethoxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(297) 3-Z-[1-(4-trifluoromethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(298) 3-Z-[1-(4-methylmercapto-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(299) 3-Z-[1-(4-(isopropylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(300) 3-Z-[1-(4-(anilinomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(301) 3-Z-[1-(4-(isobutylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(302) 3-Z-[1-(4-(cyclohexylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(303) 3-Z-[1-(4-(benzylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(304) 3-Z-[1-(4-((N-methyl-N-propyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(305) 3-Z-[1-(4-((N-isopropyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(306) 3-Z-[1-(4-((N-ethyl-N-propyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(307) 3-Z-[1-(4-((N-ethyl-N-isopropyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(308) 3-Z-[1-(4-(dipropylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(309) 3-Z-[1-(4-(diisopropylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(310) 3-Z-[1-(4-((N-benzyl-N-ethyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(311) 3-Z-[1-(4-(dibenzylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(312) 3-Z-[1-(4-(3,6-dihydro-2H-pyridin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(313) 3-Z-[1-(4-(3,5-dimethyl-piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(314) 3-Z-[1-(4-(azepan-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(315) 3-Z-[1-(4-(2-amino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(316) 3-Z-[1-(4-(2-methylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(317) 3-Z-[1-(4-(2-ethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(318) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(319) 3-Z-[1-(4-(2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(320) 3-Z-[1-(4-(2-piperidin-1-yl-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(321) 3-Z-[1-(4-(2-acetylamino-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(322) 3-Z-[1-(4-(3-amino-propyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(323) 3-Z-[1-(4-(3-dimethylamino-propyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(324) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(325) 3-Z-[1-(4-(N-ethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(326) 3-Z-[1-(4-(N-diethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(327) 3-Z-[1-(4-(N-dipropylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(328) 3-Z-[1-(4-(N-((N-ethyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(329) 3-Z-[1-(4-(N-((N-ethyl-N-propyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(330) 3-Z-[1-(4-(N-((N-methyl-N-propyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(331) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-ethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(332) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-propyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(333) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-butyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(334) 3-Z-[1-(4-(N-(2-amino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(335) 3-Z-[1-(4-(N-(2-diethylamino-ethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(336) 3-Z-[1-(4-(N-acetyl-N-(2-aminoethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(337) 3-Z-[1-(4-(N-acetyl-N-(2-methylamino-ethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(338) 3-Z-[1-(4-(N-acetyl-N-(3-methylamino-propyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(339) 3-Z-[1-(4-(N-acetyl-N-(2-piperidin-1-yl-ethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(340) 3-Z-[1-(4-(N-acetyl-N-(aminocarbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(341) 3-Z-[1-(4-(N-acetyl-N-(piperidin-1-yl-carbonylmethyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(342) 3-Z-[1-(4-(N-methyl-N-(aminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(343) 3-Z-[1-(4-(N-methyl-N-(methylaminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(344) 3-Z-[1-(4-(N-methyl-N-(dimethylaminocarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(345) 3-Z-[1-(4-(N-methyl-N-(piperidin-1-yl-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(346) 3-Z-[1-(4-(N-(2-ethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(347) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(348) 3-Z-[1-(4-(N-(2-pyrrolidin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(349) 3-Z-[1-(4-(N-(2-piperidin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(350) 3-Z-[1-(4-(N-(2-piperazin-1-yl-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(351) 3-Z-[1-(4-(N-(2-(4-morpholin-1-yl)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(352) 3-Z-[1-(4-(N-(ethylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(353) 3-Z-[1-(4-(N-(diethylaminocarbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(354) 3-Z-[1-(4-(N-(pyrrolidin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(355) 3-Z-[1-(4-(N-(piperidin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(356) 3-Z-[1-(4-(N-(piperazin-1-yl-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(357) 3-Z-[1-(4-(N-((morpholin-4-yl)-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(358) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(359) 3-Z-[1-(4-(3-dimethylamino-propoxy)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(360) 3-Z-[1-(4-(aminocarbonylmethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(361) 3-Z-[1-(4-(2-aminocarbonyl-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(362) 3-Z-[1-(4-(pyridin-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(363) 3-Z-[1-(4-(pyridine-3-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(364) 3-Z-[1-(4((N-phenethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(365) 3-Z-[1-(4-(N-acetyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(366) 3-Z-[1-(4-(N-ethylcarbonyl-N-(dimethylaminocarbonyl-methyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(367) 3-Z-[1-(4-(N-methyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(368) 3-Z-[1-(4-carboxymethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(369) 3-Z-[1-(4-carbamoylmethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(370) 3-Z-[1-(4-dimethylcarbamoylmethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(371) 3-Z-[1-(4-tetrazol-5-yl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(372) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(373) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(374) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(375) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(376) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(377) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(378) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(379) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(380) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(381) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(382) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(383) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(384) 3-Z-[1-(4-tetrazol-5-yl-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(385) 3-Z-[1-(4-tetrazol-5-yl-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(386) 3-Z-[1-(4-tetrazol-5-yl-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(387) 3-Z-[1-(4-tetrazol-5-yl-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(388) 3-Z-[1-(4-carboxy-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(389) 3-Z-[1-(4-carboxy-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(390) 3-Z-[1-(4-carboxy-anilino)-propylidene]-6-methoxy-carbonyl-2-indolinone

(391) 3-Z-[1-(4-carboxy-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(392) 3-Z-[1-(4-(N-benzyl-N-methyl-aminomethyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(393) 3-Z-[1-(4-(2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(394) 3-Z-[1-(4-((benzo(1,3)dioxol-5-yl-methyl)-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(395) 3-Z-[1-(4-(N-phenethyl-N-methyl-aminomethyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(396) 3-Z-[1-(4-(N-(3,4-dimethoxy-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(397) 3-Z-[1-(4-(N-(4-chloro-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(398) 3-Z-[1-(4-(N-(4-methylbenzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(399) 3-Z-[1-(4-(N-(4-fluoro-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(400) 3-Z-[1-(4-(N-(4-bromo-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-methoxycarbonyl-2-indolinone

(401) 3-Z-[1-(4-(N-(3-dimethylamino-propionyl)-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(402) 3-Z-[1-(4-(N-(4-dimethylamino-butyryl)-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(403) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-(2-dimethylamino-ethylsulphonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(404) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-(3-dimethylamino-propylsulphonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(405) 3-Z-[1-(4-((2-hydroxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(406) 3-Z-[1-(4-((2-methoxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(407) 3-Z-[1-(4-((2-dimethylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(408) 3-Z-[1-(4-((3-dimethylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(409) 3-Z-[1-(4-((N-tert.butoxycarbonyl-2-amino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(410) 3-Z-[1-(4-((N-tert.butoxycarbonyl-3-amino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(411) 3-Z-[1-(4-((2-amino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(412) 3-Z-[1-(4-((3-amino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(413) 3-Z-[1-(4-((2-acetylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(414) 3-Z-[1-(4-((3-acetylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(415) 3-Z-[1-(4-((2-methylsulphonylamino-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(416) 3-Z-[1-(4-((3-methylsulphonylamino-propyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(417) 3-Z-[1-(4-(N-(N-tert.butoxycarbonyl-2-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(418) 3-Z-[1-(4-(N-(2-amino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(419) 3-Z-[1-(4-(N-(2-acetylamino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(420) 3-Z-[1-(4-(N-(2-methylsulphonylamino-ethyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(421) 3-Z-[1-(4-(carboxymethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(422) 3-Z-[1-(4-(ethoxycarbonylmethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(423) 3-Z-[1-(4-(carbamoylmethyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(424) 3-Z-[1-(4-(dimethylcarbamoyl-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(425) 3-Z-[1-(4-(methylcarbamoyl-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(426) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(427) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-nitro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(428) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-acetylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(429) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methylsulphonylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(430) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-cyano-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(431) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-hydroxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(432) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methoxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(433) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-ethoxycarbonyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(434) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-carboxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(435) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-carbamoyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(436) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-chloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(437) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-fluoro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(438) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-bromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(439) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-methyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(440) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3-trifluoromethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(441) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3,5-dibromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(442) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-3,5-dichloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(443) 3-Z-[1-(4-(dimethylaminomethyl)-3-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(444) 3-Z-[1-(4-(dimethylaminomethyl)-3-nitro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(445) 3-Z-[1-(4-(dimethylaminomethyl)-3-acetylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(446) 3-Z-[1-(4-(dimethylaminomethyl)-3-methylsulphonylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(447) 3-Z-[1-(4-(dimethylaminomethyl)-3-cyano-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(448) 3-Z-[1-(4-(dimethylaminomethyl)-3-hydroxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(449) 3-Z-[1-(4-(dimethylaminomethyl)-3-methoxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(450) 3-Z-[1-(4-(dimethylaminomethyl)-3-ethoxycarbonyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(451) 3-Z-[1-(4-(dimethylaminomethyl)-3-carboxy-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(452) 3-Z-[1-(4-(dimethylaminomethyl)-3-carbamoyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(453) 3-Z-[1-(4-(dimethylaminomethyl)-3-chloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(454) 3-Z-[1-(4-(dimethylaminomethyl)-3-fluoro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(455) 3-Z-[1-(4-(dimethylaminomethyl)-3-bromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(456) 3-Z-[1-(4-(dimethylaminomethyl)-3-methyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(457) 3-Z-[1-(4-(dimethylaminomethyl)-3-trifluoromethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(458) 3-Z-[1-(4-dimethylaminomethyl-3,5-dibromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(459) 3-Z-[1-(4-(dimethylaminomethyl)-3,5-dichloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(460) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(2-hydroxy-ethoxy)-carbonyl]-2-indolinone

(461) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(ethoxycarbonyl-methoxy)-carbonyl]-2-indolinone

(462) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(carboxy-methoxy)-carbonyl]-2-indolinone

(463) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(carbamoyl-methoxy)-carbonyl]-2-indolinone

(464) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-hydroxy-ethoxy)-carbonyl]-2-indolinone

(465) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(ethoxycarbonyl-methoxy)-carbonyl]-2-indolinone

(466) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(carboxy-methoxy)-carbonyl]-2-indolinone

(467) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(carbamoyl-methoxy)-carbonyl]-2-indolinone

(468) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-methoxy-ethoxy)-carbonyl]-2-indolinone

(469) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-dimethylamino-ethoxy)-carbonyl]-2-indolinone

(470) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-(N-tert.butoxycarbonyl-amino)-ethoxy)-carbonyl]-2-indolinone

(471) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2-amino-ethoxy)-carbonyl]-2-indolinone

(472) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-[(2,2,2-trifluoroethoxy)-carbonyl]-2-indolinone

(473) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(474) 3-Z-[1-(4-(N-(imidazo-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(475) 3-Z-[1-(4-(N-(phthalimido-2-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(476) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(477) 3-Z-[1-(4-(N-acetylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(478) 3-Z-[1-(4-(N-methylsulphonylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(479) 3-Z-[1-(4-(N-((N-(2-methoxyethyl)-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(480) 3-Z-[1-(4-(N-((N-(2-dimethylaminoethyl)-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(481) 3-Z-[1-(4-(N-((di-(2-hydroxyethyl)-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(482) 3-Z-[1-(4-tert.butoxycarbonylmethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(483) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(484) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(485) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(486) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(487) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(488) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-ethylidene]-6-methoxycarbonyl-2-indolinone

(489) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-propylidene]-6-methoxycarbonyl-2-indolinone

(490) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-butylidene]-6-methoxycarbonyl-2-indolinone

(491) 3-Z-[1-(4-tert.butyloxycarbonyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(492) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(493) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(494) 3-Z-[1-(4-(N-methyl-acetylamino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(495) 3-Z-[1-(4-(imidazol-4-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(496) 3-Z-[1-(4-((N-(dioxolan-2-yl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(497) 3-Z-[1-(4-(N-benzyl-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(498) 3-Z-[1-(4-(2,3,4,5-tetrahydro-benzo(d)azepin-3-yl-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(499) 3-Z-[1-(4-((benzo(1,3)dioxol-5-yl-methyl)-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(500) 3-Z-[1-(4-(N-phenethyl-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(501) 3-Z-[1-(4-(N-(3,4-dimethoxy-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(502) 3-Z-[1-(4-(N-(4-chloro-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(503) 3-Z-[1-(4-(N-(4-methyl-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(504) 3-Z-[1-(4-(N-(4-fluoro-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(505) 3-Z-[1-(4-(N-(4-bromo-benzyl)-N-methyl-amino-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(506) 3-Z-[1-(4-((N-(2-methoxy-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(507) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-[(2-amino-ethoxy)-carbonyl]-2-indolinone

(508) 3-Z-[1-(4-((N-(3-methylsulfonylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(509) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(510) 3-Z-(1-anilino-1-phenyl-methylene)-6-carbamoyl-2-indolinone

(511) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(512) 3-Z-[1-(4-(2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(513) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(514) 3-Z-[1-(4-(1-oxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(515) 3-Z-[1-(4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(516) 3-Z-[1-(4-(benzylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(517) 3-Z-[1-(4-(amino-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(518) 3-Z-[1-(4-(2,6-dimethylpiperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(519) 3-Z-[1-(4-(pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(520) 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(521) 3-Z-[1-(3-(N-methyl-N-ethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(522) 3-Z-[1-(3-(methylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(523) 3-Z-[1-(3-hydroxymethyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(524) 3-Z-[1-(4-(methoxycarbonylmethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(525) 3-Z-[1-(4-(N-methylsulphonyl-N-(dimethylaminocarbonyl-methyl)-amino)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(526) 3-Z-[1-(4-(N-acetyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(527) 3-Z-[1-(3,4-dimethoxy-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(528) 3-Z-[1-(4-(morpholin-4-yl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(529) 3-Z-[1-(4-acetylamino-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(530) 3-Z-[1-(4-amino-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(531) 3-Z-[1-(4-N-methyl-N-acetyl-amino-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(532) 3-Z-[1-(4-ethoxycarbonyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(533) 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(534) 3-Z-[1-(4-benzylcarbamoyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(535) 3-Z-[1-(cyclohexyl-amino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(536) 3-Z-[1-(4-amino-cyclohexyl-amino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(537) 3-Z-[1-(N-methyl-piperidine-4-yl-amino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(538) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(539) 3-Z-[1-(3-dimethylaminomethyl-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(540) 3-Z-[1-(4-(N-methyl-N-benzyl-aminomethyl)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(541) 3-Z-[1-(4-(N-methylsulphonyl-N-(2-dimethylamino-ethyl)-amino)-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(542) 3-Z-[1-(4-chloro-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(543) 3-Z-[1-(3-chloro-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(544) 3-Z-[1-(4-methoxycarbonyl-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(545) 3-Z-[1-(4-carboxy-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(546) 3-Z-[1-(4-methyl-3-nitro-anilino)-1-methyl-methylene]-6-carbamoyl-2-indolinone

(547) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(548) 3-Z-[1-(3-dimethylaminomethyl-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(549) 3-Z-[1-(4-(N-methyl-N-benzyl-aminomethyl)-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(550) 3-Z-[1-(4-(N-methylsulphonyl-N-(2-dimethylamino-ethyl)-amino)-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(551) 3-Z-[1-(4-chloro-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

(552) 3-Z-[1-(3-chloro-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

(553) 3-Z-[1-(4-methoxycarbonyl-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

(554) 3-Z-[1-(4-carboxy-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

(555) 3-Z-[1-(4-methyl-3-nitro-anilino)-1-propyl-methylene]-6-carbamoyl-2-indolinone

(556) 3-Z-[1-(3-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(557) 3-Z-[1-(3-(diethylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(558) 3-Z-[1-(3-(benzylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(559) 3-Z-[1-(3-(N-methyl-N-benzyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(560) 3-Z-[1-(3-(butylaminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(561) 3-Z-[1-(3-(aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(562) 3-Z-[1-(3-(N-(3-dimethylaminopropyl)-N-methyl-amino-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(563) 3-Z-[1-(3-(N-(2-dimethylaminoethyl)-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone-trifluoroacetate

(564) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(565) 3-Z-[1-(4-bromo-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone

(566) 3-Z-[1-(3-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(567) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(568) 3-Z-[1-(4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(569) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(570) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(571) 3-Z-[1-(4-tert.butyloxycarbonyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(572) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(573) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(574) 3-Z-[1-(4-(4-methyl-piperazin-1-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(575) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(576) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(577) 3-Z-[1-(4-(N-methyl-acetylamino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(578) 3-Z-[1-(4-(N-methyl-methylsulphonylamino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(579) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(580) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(581) 3-Z-[1-(4-(imidazol-4-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(582) 3-Z-[1-(4-(tetrazol-5-yl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(583) 3-Z-[1-(4-(N-benzyl-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(584) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-propionyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(585) 3-Z-[1-(4-(pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(586) 3-Z-[1-(4-(N-methyl-N-phenethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(587) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(588) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-ethylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(589) 3-Z-[1-(4-(N-tert.butoxycarbonyl-N-ethyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(590) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

(591) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

(592) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(593) 3-Z-[1-(4-[(2,6-dimethyl-piperidin-1-yl)-methyl]-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(594) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(595) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(596) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(597) 3-Z-[1-(4-(N-acetyl-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(598) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(599) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(600) 3-Z-[1-(4-(N-methylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(601) 3-Z-[1-(4-((imidazolidin-2,4-dion-5-ylidene)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(602) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(603) 3-Z-[1-(4-(N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(604) 3-Z-[1-(4-(2-oxo-pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(605) 3-Z-[1-(4-(N-aminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(606) 3-Z-[1-(4-(thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(607) 3-Z-[1-(4-(1,1-dioxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(608) 3-Z-[1-(4-(N-cyanomethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(609) 3-Z-[1-(4-(N-tert.butoxycarbonyl-ethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(610) 3-Z-[1-(4-(N-benzyl-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(611) 3-Z-[1-(4-(1-oxo-thiomorpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(612) 3-Z-[1-(4-(2-(imidazol-4-yl)-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(613) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(614) 3-Z-[1-(4-((4-methyl-piperazin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(615) 3-Z-[1-(4-((2-(N-benzyl-N-methyl-amino)-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(616) 3-Z-[1-(4-cyclohexylamino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(617) 3-Z-[1-(4-(pyridin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(618) 3-Z-[1-(4-(imidazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(619) 3-Z-[1-(4-(imidazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(620) 3-Z-[1-(N-methyl-piperidine-4-yl-amino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(621) 3-Z-[1-(4-(imidazol-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(622) 3-Z-[1-(4-((4-hydroxy-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(623) 3-Z-[1-(4-((4-methoxy-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(624) 3-Z-[1-(4-benzyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(625) 3-Z-[1-(4-(N-(3-trifluoroacetylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(626) 3-Z-[1-(4-tert.butoxycarbonylmethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(627) 3-Z-[1-(4-tert.butoxycarbonyl-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

(628) 3-Z-[1-(4-(4-tert.butoxycarbonyl-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(629) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(630) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-nitro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(631) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(632) 3-Z-[1-(4-((3-(N-benzyl-N-methyl-amino)-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(633) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-3-chloro-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(634) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(635) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(636) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-propionyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(637) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-butyryl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(638) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-isobutyryl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(639) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-benzoyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(640) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-amino-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(641) 3-Z-[1-(4-(4-hydroxymethyl-piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(642) 3-Z-[1-(4-(2-(4-hydroxy-piperidin-1-yl)-ethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(643) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-propylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(644) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-butylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(645) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-phenylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(646) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-benzylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(647) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-ethylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(648) 3-Z-[1-(4-((imidazolidin-2,4-dion-5-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(649) 3-Z-[1-(4-((3-hydroxy-pyrrolidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(650) 3-Z-[1-(4-(cyclohexylyl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(651) 3-Z-[1-(4-(cyclohexyl-carbonyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(652) 3-Z-[1-(4-diethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(653) 3-Z-[1-(4-(N-(n-hexyl)-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(654) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(furan-2-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(655) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(2-methoxy-benzoyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxy-carbonyl-2-indolinone

(656) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(pyridine-3-carbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(657) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-(phenyl-acetyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(658) 3-Z-[1-(4-(N-ethyl-N-methyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(659) 3-Z-[1-(4-(imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(660) 3-Z-[1-(4-(1-ethyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(661) 3-Z-[1-(4-(1-benzyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(662) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-isopropylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(663) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(664) 3-Z-[1-(4-(N-(morpholin-4-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(665) 3-Z-[1-(4-(N-((4-benzyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(666) 3-Z-[1-(4-(N-(pyrrolidin-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(667) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-3-bromo-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(668) 3-Z-[1-(4-(5-methyl-imidazol-4-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(669) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(670) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(671) 3-Z-[1-(4-(N-butyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(672) 3-Z-[1-(4-(N-((N-aminocarbonylmethyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(673) 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(674) 3-Z-[1-(4-(N-(di-(2-methoxyethyl)-amino-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(675) 3-Z-[1-(4-(N-((2-(4-tert.butoxycarbonyl-piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(676) 3-Z-[1-(4-(N-((2-(piperidin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(677) 3-Z-[1-(4-(N-((2-(N-benzyl-N-methyl-amino)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(678) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(679) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(680) 3-Z-[1-(4-(N-((4-tert.butoxycarbonyl-piperazin-1-yl)-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(681) 3-Z-[1-(4-(N-((N-benzyl-N-methyl-amino)-methylcarbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(682) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(683) 3-Z-[1-(4-(N-(piperidin-1-yl-methylcarbonyl)-N-benzyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(684) 3-Z-[1-(4-(1,2,4-triazol-2-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(685) 3-Z-[1-(4-(1,2,3-triazol-2-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(686) 3-Z-[1-(4-(1,2,3-triazol-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(687) 3-Z-[1-(4-((N-aminocarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(688) 3-Z-[1-(4-((di-(2-methoxy-ethyl)-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(689) 3-Z-[1-(4-(pyrrolidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(690) 3-Z-[1-(4-((di-(2-hydroxy-ethyl)-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(691) 3-Z-[1-(4-((N-ethoxycarbonylmethyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(692) 3-Z-[1-(4-(azetidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(693) 3-Z-[1-(4-(N-propyl-N-tert.butoxycarbonyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(694) 3-Z-[1-(4((N-(2-(2-methoxy-ethoxy)-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(695) 3-Z-[1-(4-((N-(tert.butoxycarbonyl-3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(696) 3-Z-[1-(4-((N-(methylcarbamoyl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(697) 3-Z-[1-(4-((N-(dimethylcarbamoyl-methyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(698) 3-Z-[1-(4-methyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(699) 3-Z-[1-(4-((N-propyl-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(700) 3-Z-[1-(4-((N-(2-hydroxy-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(701) 3-Z-[1-(4-((N-(2-dimethylamino-ethyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(702) 3-Z-[1-(4-((N-(3-dimethylamino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(703) 3-Z-[1-(4-(3-oxo-piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(704) 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-ethoxy-carbonyl-2-indolinone

(705) 3-Z-[1-(4-aminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(706) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(707) 3-Z-[1-(4-carboxymethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(708) 3-Z-[1-(4-carboxy-anilino)-1-ethyl-methylene]-6-ethoxycarbonyl-2-indolinone

(709) 3-Z-[1-(4-(piperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(710) 3-Z-[1-(4-butylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(711) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone

(712) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone

(713) 3-Z-[1-(4-(N-(piperazin-1-yl-methylcarbonyl)-N-isopropyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(714) 3-Z-[1-(4-(N-((2-(piperazin-1-yl)-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(715) 3-Z-[1-(4-(N-propyl-aminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(716) 3-Z-[1-(4-((N-(3-amino-propyl)-N-methyl-amino)-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(717) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(718) 3-Z-[1-(3-cyano-4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(719) 3-Z-[1-(3-methoxy-4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(720) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(721) 3-Z-[1-(4-(N-(N-(2-dimethylamino-ethyl)-N-methyl-aminomethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(722) 3-Z-[1-(4-(N-(di-(2-hydroxy-ethyl)-amino-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(723) 3-Z-[1-(4-(N-(imidazol-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(724) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(725) 3-Z-[1-(4-(N-((4-methyl-[1,4]diazepan-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(726) 3-Z-[1-(4-(N-((1-methyl-piperidin-4-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(727) 3-Z-[1-(2,3-dimethyl-4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(728) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(729) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(730) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(731) 3-Z-[1-(4-(N-((3-dimethylamino-propyl)-aminocarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(732) 3-Z-[1-anilino-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(733) 3-Z-[1-(4-(N-methyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(734) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(735) 3-Z-[1-(4-(4-methylpiperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(736) 3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(737) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(738) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(739) 3-Z-[1-(4-(di-(2-hydroxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(740) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(741) 3-Z-[1-(4-(N-(morpholin-4-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(742) 3-Z-[1-(4-(N-(N-(2-dimethylamino-ethyl)-N-methyl-aminomethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(743) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(744) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-carboxy-2-indolinone

(745) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(746) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(747) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(748) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-(4-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(749) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(750) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(751) 3-Z-[1-((1-methyl-piperidin-4-yl)-amino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(752) 3-Z-[1-(trans-4-dimethylamino-cyclohexylamino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(753) 3-Z-[1-(4-(2-diethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(754) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-propionyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(755) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(756) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(757) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(758) 3-Z-[1-(3-diethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(759) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(760) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(761) 3-Z-[1-anilino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(762) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(763) 3-Z-[1-(4-((2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(764) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(765) 3-Z-[1-(4-(N-methyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(766) 3-Z-[1-(4-methoxycarbonyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(767) 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(768) 3-Z-[1-(4-(N-(dimethylamino-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(769) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(770) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(771) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(772) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(773) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(774) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(775) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(776) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(777) 3-Z-[1-(4-carbamoyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(778) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(779) 3-Z-[1-(4-((4-methyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(780) 3-Z-[1-(4-((4-methyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(781) 3-Z-[1-(4-((4-ethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(782) 3-Z-[1-(4-(N-ethyl-N-(2-dimethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(783) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-diethylcarbamoyl-2-indolinone

(784) 3-Z-[1-(4-((cis-3,5-dimethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(785) 3-Z-[1-(4-((4-ethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(786) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(787) 3-Z-[1-(4-((cis-3,5-dimethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(788) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(789) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(790) 3-Z-[1-(4-dimethylaminomethyl-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(791) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(792) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

(793) 3-Z-[1-(4-dimethylaminomethyl-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

(794) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

(795) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

as well as their tautomers, their stereoisomers or the physiologically acceptable salts thereof.

The compounds of general formula I, their tautomers, their stereoisomers or the physiologically acceptable salts thereof are thus suitable for the prevention or treatment of a specific fibrotic disease selected from the group consisting of:

Fibrosis and remodeling of lung tissue in chronic obstructive pulmonary disease (COPD), chronic bronchitis, and emphysema;

Lung fibrosis and pulmonary diseases with a fibrotic component including but not limited to idiopathic pulmonary fibrosis (IPF), giant cell interstitial pneumonia (GIP), sarcodosis, cystic fibrosis, respiratory distress syndrome

(ARDS), granulomatosis, silicosis, drug-induced lung fibrosis (for example, induced by drugs such as bleomycin, bis-chloronitrosourea, cyclophosphamide, amiodarone, procainamide, penicillamine, gold or nitrofurantoin), silicosis, asbestosis, systemic scleroderma;

Fibrosis and remodeling in asthma;

Fibrosis in rheumatoid arthritis;

Virally induced hepatic cirrhosis, for example hepatitis C;

Radiation-induced fibrosis;

Restenosis, post angioplasty;

Renal disorders including chronic glomerulonephritis, renal fibrosis in patients receiving cyclosporine and renal fibrosis due to high blood pressure;

Diseases of the skin with a fibrotic component including but not limited to, scleroderma, sarcodosis, systemic lupus erythematosus;

Excessive scarring.

In a preferred embodiment in accordance with the present invention, the compounds of general formula I, their tautomers, their stereoisomers or the physiologically acceptable salts thereof are especially suitable for the prevention or treatment of idiopathic pulmonary fibrosis.

BIOLOGICAL ACTIVITY

The following experimental results illustrate the present invention without representing a limitation of its scope.

Example B1:

In the following experiments of Example B1, Example A denotes the compound 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone, which is compound (m) of the list of the preferred compounds.

(A) Effect of a representative compound on lung morphology following bleomycin-induced pulmonary fibrosis.

Materials and Methods

Bleomycin sulfate (Bleomycin HEXAL™) was purchased from a local pharmacy.

Bleomycin administration and treatment protocols

All experiments were performed in accordance with German guidelines for animal welfare, performed by persons certified to work with animals and approved by the responsible authorities. Male Wistar rats were intratracheally injected with Bleomycin sulfate (10U/kg body weight in 300µl saline) or saline alone (saline control) using a catheter (0,5mm internal diameter, 1.0mm external diameter) through the nasal passage, following exposure to the anaesthetic Isofluorane for 5 minutes. The

following day, the rats were orally treated with Example A (compound (m)) or saline suspended in 1ml 0.1% Natrosol. Control rats were administered 1ml 0.1% Natrosol (vehicle control).

A total of 25 rats were investigated and were grouped and treated as shown in Table 1.

Table 1

Intratracheal instillation	No. of animals	Compound	Treatment Schedule
Bleomycin 10U /kg	10	Example A (Compound (m))	Days 1-21
Bleomycin 10U /kg	10	Vehicle only	Days 1-21
Saline (300µl)	5	Vehicle only	Days 1-21

21 days following bleomycin instillation, the rats were killed with a lethal intraperitoneal injection of Narcoren™ (Pentobarbital Sodium, Rhone Merieux). The lungs were then removed, blotted dry and half was snap frozen in liquid nitrogen and stored at -80°C. The other half was fixed in 4% formalin for subsequent paraffin embedding and histology.

Histology

The lung tissues fixed in 4% formalin were embedded into paraffin and 5μm sections were cut using a microtome (Leica SM200R) and placed on poly-L-lysine coated slides. The sections were then dried onto the slides (60°C 2 hours) and then left to cool at room temperature. Collagen deposition was assessed using Masson's Trichrome staining.

Results

Figure 1A shows the result obtained with the control group, which received saline and the vehicle instead of bleomycin intratracheally.

Rats treated intratracheally with bleomycin and the vehicle developed severe lung fibrosis, as seen in Figure 1B. The alveoli have been largely replaced by fibroblasts and extracellular matrix and the normal lung structure is nearly obliterated.

Daily treatment of bleomycin-treated rats with 50 mg/kg of Example A (compound (m)) showed a consistent, nearly complete reversal of lung fibrosis in this model. A typical example is shown in Figure 1C. Alveoli are intact and little or no fibroblast infiltration or extracellular matrix deposition has occurred. Normal lung structure has been maintained, which is evidenced by a comparison of Figure 1C with Figure 1A.

(B) Effect of a representative compound on expression of fibrotic marker genes following bleomycin-induced pulmonary fibrosis.

mRNA extractions and synthesis of cDNA

One part of the frozen lung tissue dedicated to investigation of gene expression was cut into small pieces using a sterile scalpel blade. Approximately 100mg of tissue was then placed into a 2ml Eppendorf tube and 1.5ml of Trizol (Invitrogen) was added. A sterile tungsten carbide bead (Qiagen) was then added to the tube and the tube was placed in a Retsch MM300 Tissue disruptor (Qiagen) at a frequency of 30.0Hz for 8 minutes. After this time, the bead was removed and the sample centrifuged at 12000rpm for 10 minutes to remove tissue debris. The RNA was extracted using a modified version of the manufacturer's protocol supplied with Trizol. Briefly, 0.3ml chloroform was added to the tube and the tube shaken vigorously and then left to incubate at room temperature for 5 minutes, after which the tube was centrifuged for 15 minutes at 12000 rpm at 4°C. The upper colorless aqueous phase was then collected and added to 750µl isopropanol. This was then shaken vigorously and stored at -80°C overnight. The samples were then incubated at room temperature for 15 minutes, after which they were centrifuged for 40 minutes at 12000 rpm at 4°C. The supernatant was then removed and 500µl of 70% ethanol was added to wash the pellet then the sample was centrifuged for 10 minutes at 12000 rpm an 4°C, this wash step was repeated twice, after which the pellet was left to dry for 10 - 15 minutes. Finally the pellet was resuspended in 20µl RNase free water and stored at -80°C. The concentration of each sample was then measured using a spectrophotometer.

Using the Superscript™ III (Invitrogen, Paisley, UK) RT-first strand synthesis kit, 2µg of each mRNA sample was reversed transcribed using a modified version of the manufacturer's protocol. Briefly, a mixture of 2µg RNA, 1µl random hexamer primers (50ng/µl), 1µl dNTP mix (10mM) was made up to 10µl with DEPC-treated water and incubated at 65°C for 5 minutes, after which it was placed on ice for 5 minutes. Following this, to each reaction, 2µl RT buffer (10X), 4µl $MgCl_2$ (25mM), 2µl DTT (0.1M), 1µl RNaseOUT™ (40U/µl) and 1µl SuperScript™ III enzyme (200U/µl) was added and the mixture placed in a thermal cycler (Applied Biosystems) under the following conditions: 25° C for 10 minutes, 50°C for 50 minutes and 85°C for 5 minutes, after which 1µl of RNase H was added and incubated at 37°C for 20 minutes. The synthesized cDNA was diluted to 5ng/µl using the assumption that the RT reaction fully transcribed all of the mRNA to cDNA and was a concentration of 100ng/µl.

Investigation of gene expression using real time PCR

Gene expression was investigated in each of the samples using the Applied Biosystems 7700 sequence detection system. Primers for the 18S endogenous control and TGFb1 were purchased as pre-developed assay reagent kits, whereas primers and probes (see Table 2 below) for pro-collagen I and fibronectin were designed using PrimerExpress™ (Applied Biosystems), ensuring that at least one of the primers or probes in each set overlapped an intron / exon junction, thus eliminating the possibility of amplifying any contaminating genomic DNA in the cDNA sample. The purchased PDARs also amplified only cDNA.

Table 2

Target		Sequence
Fibronectin	Forward	5′-GAT GCC GAT CAG AAG TTT GGA-3′
	Reverse	5′-TCG TTG GTC GTG CAG ATC TC-3′
	Probe	5′-FAM-CTG CCC AAT GGC TGC CCA TGA-TAMRA-3′
Pro-Collagen I	Forward	5′-CAG ACT GGC AAC CTG AAG AAG TC-3′
	Reverse	5′-TCG CCC CTG AGC TCG AT-3′
	Probe	5′-FAM-CTG CTC CTC CAG GGC TCC AAC GA-TAMRA3′

Real Time PCR was carried out in 25µl reactions, using 25ng (5µl) of cDNA per reaction. A quantitative PCR core kit was purchased (Eurogentec) and a master-mix was made up as follows for 100 reactions: 500µl 10X reaction buffer, 500µl $MgCl_2$ (50mM), 200µl dNTP mix solution (5mM), 25µl Hot Goldstar enzyme, 75µl 18S PDAR, 22.5µl forward primer, 22.5µl reverse primer, 15µl probe and 640µl DEPC treated water. 20µl of this master-mix was then added to 25ng (5µl) target cDNA. Each analysis was carried out in triplicate.

In order to quantify the gene expression, a standard curve was constructed for each primer set and was included on each plate. The standards were made up of a mix of all the

cDNA's under investigation; this mix of cDNA's was serially diluted 10, 20, 50, 100, 100 times. A standard curve was constructed of the obtained C_T (**C**ycle at which amplification reaches a set **T**hreshold) against the LOG_{10} of the dilution factor. Curves were drawn for the target gene and the 18S rRNA endogenous control. The C_T value for both targets for each of the samples was then converted to a fold dilution using the standard curve and the target gene value was normalized to the 18S gene value.

Statistics

All statistical analyses were carried out using GraphPad Prism V 4.02 software. Comparisons were made using a non-parametric T-test (Mann-Whitney U test) and a significant value was considered to be $p = 0.05$.

Results

The results are shown in Figures 2 (procollagen I) and 3 (fibronectin). Each data point represents RNA isolated from the lung of a single rat.

Intratracheal administration of bleomycin and subsequent treatment with vehicle only showed large increases in procollagen I and fibronectin gene expression in the lung, as seen in Figures 2 and 3, consistent with the histologically apparent lung fibrosis seen in Figure 1B.

Daily treatment of Bleomycin-treated rats with 50 mg/kg of Example A (compound (m)) showed a significant ($p \leq 0.0001$)

inhibition of expression of fibrotic marker genes in this model, as seen in Figures 2 and 3.

This experiment thus demonstrates that expression of fibrotic markers, and therefore deposition of extracellular matrix, may be dramatically reduced by treatment with Example A (compound (m)).

Example B2:

In the following experiments of Example B2, the compound 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone, which is compound (u) of the list of the preferred compounds, was used.

All the methods employed are the same as the methods described in the experiments of Example B1, however using compound (u) instead of compound (m).

(A) Effect of a representative compound on lung morphology following bleomycin-induced pulmonary fibrosis.

Samples were prepared from rats treated as outlined in above Table 1 of experimental Example B1 (A).

Results

Figure 4A shows the result obtained with the control group, which received saline and the vehicle instead of bleomycin intratracheally.

Rats treated intratracheally with bleomycin and the vehicle developed severe lung fibrosis, as seen in Figure 4B. The alveoli have been largely replaced by fibroblasts and extracellular matrix and the normal lung structure is nearly obliterated.

Daily treatment of bleomycin-treated rats with 50 mg/kg of compound (u) showed a consistent, nearly complete reversal of lung fibrosis in this model. A typical example is shown in Figure 4C. Alveoli are intact and little or no fibroblast infiltration or extracellular matrix deposition has occurred. Normal lung structure has been maintained, which is evidenced by a comparison of Figure 4C with Figure 4A.

(B) Effect of a representative compound on expression of fibrotic marker genes following Bleomycin-induced pulmonary fibrosis.

The experiment was carried out using the methods as outlined above for Example B1 (B).

The results are shown in Figure 5 (procollagen I) and Figure 6 (TGFb). Each data point represents RNA isolated from the lung of a single rat.

Intratracheal administration of bleomycin and subsequent treatment with vehicle only showed large increases in procollagen I and TGFb gene expression in the lung, as seen in Figures 5 and 6, consistent with the histologically apparent lung fibrosis seen in Figure 1B.

Daily treatment of bleomycin-treated rats with 50 mg/kg of (compound (u) showed a significant ($p \leq 0.0001$) inhibition of expression of fibrotic marker genes in this model, as seen in Figures 5 and 6.

This experiment also demonstrates that expression of fibrotic markers, and therefore deposition of extracellular matrix, may be dramatically reduced by treatment with another compound representative of this invention, namely compound (u).

By reason of their biological properties the compounds according to the invention may be used in monotherapy or in conjunction with other pharmacologically active compounds. Such pharmacologically active compounds may be compounds which are, for example, also pharmacologically active in the treatment of fibrosis. Such pharmacologically active compounds may also be substances with a secretolytic, broncholytic and/or anti-inflammatory activity.

In a preferred embodiment in accordance with the present invention, such pharmacologically active compounds are preferably selected from the group consisting of anticholinergic agents, beta-2 mimetics, steroids, PDE-IV inhibitors, p38 MAP kinase inhibitors, NK_1 antagonists, LTD4 antagonists, EGFR inhibitors and endothelin-antagonists.

Anticholinergic agents may preferably be selected from the group consisting of the tiotropium salts, oxitropium salts,

flutropium salts, ipratropium salts, glycopyrronium salts and trospium salts.

Beta-2 mimetics may preferably be selected from the beta-2 mimetics disclosed, for example, in US 4,460,581, which is incorporated herein by reference.

PDE-IV inhibitors may preferably be selected from the group consisting of enprofyllin, theophyllin, roflumilast, ariflo (cilomilast), CP-325,366, BY343, D-4396 (Sch-351591), AWD-12-281 (GW-842470), N-(3,5-dichloro-1-oxo-pyridin-4-yl)-4-difluoromethoxy-3-cyclopropylmethoxybenzamide, NCS-613, pumafentine, (-)p-[(4*a*R*,10*b*S*)-9-ethoxy-1,2,3,4,4a,10b-hexahydro-8-methoxy-2-methylbenzo[s][1,6]naphthyridin-6-yl]-N,N-diisopropylbenzamide, (R)-(+)-1-(4-bromobenzyl)-4-[(3-cyclopentyloxy)-4-methoxyphenyl]-2-pyrrolidone, 3-(cyclopentyloxy-4-methoxyphenyl)-1-(4-N'-[N-2-cyano-S-methyl-isothioureido]benzyl)-2-pyrrolidone, cis[4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carbonic acid], 2-carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-one, cis[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol], (R)-(+)-ethyl[4-(3-cyclopentyloxy-4-methoxyphenyl)pyrrolidin-2-yliden]acetate, (S)-(-)-ethyl[4-(3-cyclopentyloxy-4-methoxyphenyl)pyrrolidin-2-yliden]acetate, CDP840, Bay-198004, D-4418, PD-168787, T-440, T-2585, arofyllin, atizoram, V-11294A, Cl-1018, CDC-801, CDC-3052, D-22888, YM-58997, Z-15370, 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(2-thienyl)-9*H*-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine and 9-cyclopentyl-5,6-dihydro-7-ethyl-3-(*tert*-butyl)-9*H*-pyrazolo[3,4-c]-1,2,4-triazolo[4,3-a]pyridine. These compounds may be used, as available, in


the form of their racemates, enantiomers or diastereoisomers, or in the form of pharamacologically acceptable acid addition salts thereof, or in the form of their solvates and/or hydrates.

Steroids may preferably be selected from the group consisting of prednisolone, prednisone, butixocortpropionate, RPR-106541, flunisolid, beclomethasone, triamcinolone, budesonid, fluticasone, mometasone, ciclesonid, rofleponid, ST-126, dexamethasone, 6α,9α-difluoro-17α-[(2-furanylcarbonyl)oxy]-11β-hydroxy-16α-methyl-3-oxo-androsta-1,4-dien-17β-carbothionic acid (S)-fluoromethylester, and 6α,9α-difluoro-11β-hydroxy-16α-methyl-3-oxo-17α-propionyloxy-androsta-1,4-diene-17β-carbothionic acid (S)-(2-oxo-tetrahydro-furan-3S-yl)ester. These compounds may be used, as available, in the form of their racemates, enantiomers or diastereoisomers, or in the form of pharamacologically acceptable acid addition salts thereof, or in the form of their solvates and/or hydrates.

p38 MAP kinase inhibitors may preferably be selected from the group consisting of the p38 Kinase inhibitors that are disclosed for instance in US Patents 5,716,972, US 5,686,455, US 5,656,644, US 5,593,992, US 5,593,991, US 5,663,334, US 5,670,527, US 5,559,137, 5,658,903, US 5,739,143, US 5,756,499, US 6,277,989, US 6,340,685, and US 5,716,955 and PCT applications WO 92/12154, WO 94/19350, WO 95/09853, WO 95/09851, WO 95/09847, WO 95/09852, WO 97/25048, WO 97/25047, WO 97/33883, WO 97/35856, WO 97/35855, WO 97/36587, WO 97/47618, WO 97/16442, WO 97/16441, WO 97/12876, WO 98/25619, WO 98/06715, WO

98/07425, WO 98/28292, WO 98/56377, WO 98/07966, WO 98/56377, WO 98/22109, WO 98/24782, WO 98/24780, WO 98/22457, WO 98/52558, WO 98/52559, WO 98/52941, WO 98/52937, WO 98/52940, WO 98/56788, WO 98/27098, WO 98/47892, WO 98/47899, WO 98/50356, WO 98/32733, WO 99/58523, WO 99/01452, WO 99/01131, WO 99/01130, WO 99/01136, WO 99/17776, WO 99/32121, WO 99/58502, WO 99/58523, WO 99/57101, WO 99/61426, WO 99/59960, WO 99/59959, WO 99/00357, WO 99/03837, WO 99/01441, WO 99/01449, WO 99/03484, WO 99/15164, WO 99/32110, WO 99/32111, WO 99/32463, WO 99/64400, WO 99/43680, WO 99/17204, WO 99/25717, WO 99/50238, WO 99/61437, WO 99/61440, WO 00/26209, WO 00/18738, WO 00/17175, WO 00/20402, WO 00/01688, WO 00/07980, WO 00/07991, WO 00/06563, WO 00/12074, WO 00/12497, WO 00/31072, WO 00/31063, WO 00/23072, WO 00/31065, WO 00/35911, WO 00/39116, WO 00/43384, WO 00/41698, WO 00/69848, WO 00/26209, WO 00/63204, WO 00/07985, WO 00/59904, WO 00/71535, WO 00/10563, WO 00/25791, WO 00/55152, WO 00/55139, WO 00/17204, WO 00/36096, WO 00/55120, WO 00/55153, WO 00/56738, WO 01/21591, WO 01/29041, WO 01/29042, WO 01/62731, WO 01/05744, WO 01/05745, WO 01/05746, WO 01/05749, WO 01/05751, WO 01/27315, WO 01/42189, WO 01/00208, WO 01/42241, WO 01/34605, WO 01/47897, WO 01/64676, WO 01/37837, WO 01/38312, WO 01/38313, WO 01/36403, WO 01/38314, WO 01/47921, WO 01/27089, DE 19842833, and JP 2000 86657 whose disclosures are all incorporated herein by reference in their entirety. Of particular interest for the combinations according to the invention are those p38 inhibitors disclosed in US 6,277,989, US 6,340,685, WO 00/12074, WO 00/12497, WO 00/59904, WO 00/71535, WO 01/64676, WO 99/61426, WO

00/10563, WO 00/25791, WO 01/37837, WO 01/38312, WO 01/38313, WO 01/38314, WO 01/47921, WO 99/61437, WO 99/61440, WO 00/17175, WO 00/17204, WO 00/36096, WO 98/27098, WO 99/00357, WO 99/58502, WO 99/64400, WO 99/01131, WO 00/43384, WO 00/55152, WO 00/55139, and WO 01/36403. In a preferred embodiment the p38 kinase inhibitor is selected from the compounds of following formula (**I**) as disclosed in WO 99/01131



(**I**),

wherein

R_1 is 4-pyridyl, pyrimidinyl, 4-pyridazinyl, 1,2,4-triazin-5-yl, quinolyl, isoquinolinyl, or quinazolin-4-yl ring, which ring is substituted with $Y-R_a$ and optionally with an additional independent substituent selected from C_{1-4} alkyl, halogen, hydroxyl, C_{1-4} alkoxy, C_{1-4} akylthio, C_{1-4} aklylsulfinyl, CH_2OR_{12}, amino, mono and di- C_{1-6} alkyl substituted amino, an N-heterocyclyl ring which ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR_{15}, $N(R_{10})C(O)R_b$ or NHR_a;

Y is oxygen or sulfur;

R_4 is phenyl, naphth-1-yl or naphthyl, or a heteroaryl, which is optionally substituted by one or two substituents, each of which is independently selected, and which, for a 4-phenyl, 4naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl substituent, is halogen, cyano, nitro, $C(Z)NR_7R_{17}$, $C(Z)OR_{16}$, $(CR_{10}R_{20})_vCOR_{12}$, SR_5, SOR_5, OR_{12}, halo-substituted-C_{1-4} alkyl,

C_{1-4} alkyl, $ZC(Z)R_{12}$, $NR_{10}C(Z)R_{16}$, or $(CR_{10}R_{20})_vNR_{10}R_{20}$ and which, for other positions of substitution, is halogen, cyano, $C(Z)NR_{13}R_{14}$, $C(Z)OR_3$, $(CR_{10}R_{20})_{m''}COR_3$, $S(O)_mR_3$, OR_3, halo-substituted-C_{1-4} alkyl, C_{1-4} alkyl, $(CR_{10}R_{20})_{m''}R_{10}C(Z)R_3$, $NR_{10}S(O)_{m'}R_8$, $NR_{10}S(O)_{m'}NR_7R_{17}$, $ZC(Z)R_3$ or $(CR_{10}R_{20})_{m''}NR_{13}R_{14}$;

Z is oxygen or sulfur;

n is an integer having a value of 1 to 10;

m is 0, or integer 1 or 2;

m′ is an integer having a value of 1 or 2;

m″ is 0, or an integer having a value of 1 to 5;

v is 0, or an integer having a value of 1 to 2;

R_2 is $-C(H)(A)(R_{22})$;

A is optionally substituted aryl, heterocyclyl, or heteroaryl ring, or A is substituted C_{1-10} alkyl;

R_{22} is an optionally substituted C_{1-10} alkyl;

R_a is aryl, arylC_{1-6} alkyl, heterocyclic, heterocyclylC_{1-6} alkyl, heteroaryl, heteroarylC_{1-6}alkyl, wherein each of these moieties may be optionally substituted;

R_b is hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aryl, aryl C_{1-4} alkyl, heteroaryl, heteroarylC_{1-4} alkyl, heterocyclyl, or heterocyclylC_{1-4} alkyl, wherein each of these moieties may be optionally substituted;

R_3 is heterocyclyl, heterocyclyl C_{1-10} alkyl or R_8;

R_5 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl or NR_7R_{17}, excluding the moieties SR_5 being $SNR_7R_{17}$and SOR_5 being SOH;

R_6 is hydrogen, a pharmaceutically acceptable cation, C_{1-10} alkyl, C_{3-7} cycloalkyl, aryl, aryl C_{1-4} alkyl, heteroaryl, heteroaryl C_{1-4} alkyl, heterocyclyl, aryl, or C_{1-10} alkanoyl;

R_7 and R_{17} is each independently selected from hydrogen or C_{1-4} alkyl or R_7 and R_{17} together with the nitrogen to which they are attached form a heterocyclic ring of 5 to 7

members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR_{15};

R_8 is C_{1-10} alkyl, halo-substituted C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-7} cycloalkyl, C_{5-7} cycloalkenyl, aryl, aryl C_{1-10} alkyl, heteroaryl, heteroaryl C_{1-10} alkyl, $(CR_{10}R_{20})_nOR_{11}$, $(CR_{10}R_{20})_nS(O)_mR_{18}$, $(CR_{10}R_{20})_nNHS(O)_2R_{18}$, $(CR_{10}R_{20})_nNR_{13}R_{14}$; wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl may be optionally substituted;

R_9 is hydrogen, C(Z) R_{11} or optionally substituted C_{1-10} alkyl, $S(O)_2R_{18}$, optionally substituted aryl or optionally substituted aryl C_{1-4} alkyl;

R_{10} and R_{20} is each independently selected from hydrogen or C_{1-4} alkyl;

R_{11} is hydrogen, C_{1-10} alkyl, C_{3-7} cycloalkyl, heterocyclyl, heterocyclyl C_{1-10} alkyl, aryl, arylC_{1-10} alkyl, heteroaryl or heteroaryl C_{1-10} alkyl, wherein these moieties may be optionally substituted;

R_{12} is hydrogen or R_{16};

R_{13} an R_{14} is each independently selected from hydrogen or optionally substituted

C_{1-4} alkyl, optionally substituted aryl or optionally substituted arylC_{1-4} alkyl, or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR_9;


R_{15} is R_{10} or C(Z)-C_{1-4} alkyl;

R_{16} is C_{1-4} alkyl, halo-substituted-C_{1-4} alkyl, or C_{3-7} cycloalkyl;

R_{18} is C_{1-10} alkyl, C_{3-7} cycloalkyl, heterocyclyl, aryl, aryl$_{1-10}$ alkyl, heterocyclyl, heterocyclyl- C_{1-10}alkyl, heteroaryl or heteroaryl$_{1-10}$ alkyl;

or a pharmaceutically acceptable salt thereof.


NK_1 antagonists may preferably be selected from the group consisting of N-[2-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-2-{4-cyclopropylmethyl-piperazin-1-yl}-N-methyl-2-phenyl-acetamide (BIIF 1149), CP-122721, FK-888, NKP 608C, NKP 608A, CGP 60829, SR 48968 (Saredutant), SR 140333 (Nolpitantium besilate/chloride), LY 303 870 (Lanepitant), MEN-11420 (Nepadutant), SB 223412, MDL-105172A, MDL-103896, MEN-11149, MEN-11467, DNK 333A, SR-144190, YM-49244, YM-44778, ZM-274773, MEN-10930, S-19752, Neuronorm, YM-35375, DA-5018, Aprepitant (MK-869), L-754030, CJ-11974, L-758298, DNK-33A, 6b-I, CJ-11974, TAK-637, GR 205171 and the arylglycine amide derivates of general formula (**VIII**)




(**VIII**)

wherein

R^1 and R^2 together with the N-atom they are bound to form a ring of formula



or



,

wherein r and s independently denote the number 2 or 3;


R^6 denotes H, $-C_1-C_5$-alkyl, C_3-C_5-alkenyl, propinyl, hydroxy(C_2-C_4)alkyl, methoxy(C_2-C_4)alkyl, di(C_1-C_3)alkylamino(C_2-C_4)alkyl, amino(C_2-C_4)alkyl, amino, di(C_1-C_3)alkylamino, monofluoro- up to perfluoro(C_1-C_2)alkyl, N-methylpiperidinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl,

R^7 denotes any of the groups defined under (a) to (d):

(a) hydroxy

(b) 4-piperidinopiperidyl,

(c)



wherein R^{16} and R^{17} independently denote H, (C_1-C_4)alkyl, (C_3-C_6)cycloalkyl, hydroxy(C_2-C_4)alkyl, dihydroxy(C_2-C_4)alkyl, (C_1-C_3)alkoxy(C_2-C_4)alkyl, phenyl(C_1-C_4)alkyl or di(C_1-C_3)alkylamino(C_2-C_4)alkyl, and

R^8 denotes H,

optionally in the form of enantiomers, mixtures of enantiomers or the racemates.

The compounds of formula (**VIII**) mentioned hereinbefore are described in WO 96/32386, WO 97/32865 and WO 02/32865. The

disclosure of these international patent applications is incorporated herein by reference in its entirety.

LTD4 antagonists may preferably be selected from the group consisting of montelukast, 1-(((R)-(3-(2-(6,7-difluoro-2-quinolinyl)ethenyl)phenyl)-3-(2-(2- hydroxy-2-propyl)phenyl)thio)methylcyclopropane-acetate, 1-(((1(R)-3(3-(2-(2,3-dichlorothieno[3,2-b]pyridin-5-yl)-(E)-ethenyl)phenyl)-3-(2-(1-hydroxy-1-methylethyl)phenyl)propyl)thio)methyl)cyclopropane-acetate, pranlukast, zafirlukast, [2-[[2-(4-tert-butyl-2-thiazolyl)-5-benzofuranyl]oxymethyl]phenyl]acetate, MCC-847 (ZD-3523), MN-001, MEN-91507 (LM-1507), VUF-5078, VUF-K-8707 and L-733321. These compounds may be used, as available, in the form of their racemates, enantiomers or diastereoisomers, or in the form of pharamacologically acceptable acid addition salts thereof, or in the form of their solvates and/or hydrates.

EGFR inhibitors may preferably be selected from the group consisting of 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-7-cyclopropylmethoxy-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N,N-diethylamino)-1-oxo-2-buten-1-yl]amino}-7-cyclopropylmethoxy-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino}-7-cyclopropylmethoxy-chinazoline, 4-[(*R*)-(1-phenyl-ethyl)amino]-6-{[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-7-cyclopentyloxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{[4-((*R*)-6-methyl-2-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-7-cyclopropylmethoxy-chinazoline, 4-[(3-chlor-4-fluor-

phenyl)amino]-6-{[4-((*R*)-6-methyl-2-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-7-[(*S*)-(tetrahydrofuran-3-yl)oxy]-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{[4-((*R*)-2-methoxymethyl-6-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-7-cyclopropylmethoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[2-((*S*)-6-methyl-2-oxo-morpholin-4-yl)-ethoxy]-7-methoxy-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-({4-[N-(2-methoxy-ethyl)-N-methyl-amino]-1-oxo-2-buten-1-yl}amino)-7-cyclopropylmethoxy-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino}-7-cyclopentyloxy-chinazoline, 4-[(*R*)-(1-phenyl-ethyl)amino]-6-{[4-(N,N-bis-(2-methoxy-ethyl)-amino)-1-oxo-2-buten-1-yl]amino}-7-cyclopropylmethoxy-chinazoline, 4-[(*R*)-(1-phenyl-ethyl)amino]-6-({4-[N-(2-methoxy-ethyl)-N-ethyl-amino]-1-oxo-2-buten-1-yl}amino)-7-cyclopropylmethoxy-chinazoline, 4-[(*R*)-(1-phenyl-ethyl)amino]-6-({4-[N-(2-methoxy-ethyl)-N-methyl-amino]-1-oxo-2-buten-1-yl}amino)-7-cyclopropylmethoxy-chinazoline, 4-[(*R*)-(1-phenyl-ethyl)amino]-6-({4-[N-(tetrahydropyran-4-yl)-N-methyl-amino]-1-oxo-2-buten-1-yl}amino)-7-cyclopropylmethoxy-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino}-7-((*R*)-tetrahydrofuran-3-yloxy)-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino}-7-((*S*)-tetrahydrofuran-3-yloxy)-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-({4-[N-(2-methoxy-ethyl)-N-methyl-amino]-1-oxo-2-buten-1-yl}amino)-7-cyclopentyloxy-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N-cyclopropyl-N-methyl-amino)-1-oxo-2-buten-1-yl]amino}-7-cyclopentyloxy-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N,N-dimethylamino)-1-oxo-2-buten-

1-yl]amino}-7-[(*R*)-(tetrahydrofuran-2-yl)methoxy]-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino}-7-[(*S*)-(tetrahydrofuran-2-yl)methoxy]-chinazoline, 4-[(3-ethinyl-phenyl)amino]-6,7-bis-(2-methoxy-ethoxy)-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-7-[3-(morpholin-4-yl)-propyloxy]-6-[(vinylcarbonyl)amino]-chinazoline, 4-[(*R*)-(1-phenyl-ethyl)amino]-6-(4-hydroxy-phenyl)-7H-pyrrolo[2,3-d]pyrimidine, 3-cyano-4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino}-7-ethoxy-chinoline, 4-{[3-chlor-4-(3-fluor-benzyloxy)-phenyl]amino}-6-(5-{[(2-methansulfonyl-ethyl)amino]methyl}-furan-2-yl)chinazoline, 4-[(R)-(1-phenyl-ethyl)amino]-6-{[4-((R)-6-methyl-2-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-{[4-(morpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-7-[(tetrahydrofuran-2-yl)methoxy]-chinazoline, 4-[(3-chlor-4-fluorphenyl)amino]-6-({4-[N,N-bis-(2-methoxy-ethyl)-amino]-1-oxo-2-buten-1-yl}amino)-7-[(tetrahydrofuran-2-yl)methoxy]-chinazoline, 4-[(3-ethinyl-phenyl)amino]-6-{[4-(5,5-dimethyl-2-oxo-morpholin-4-yl)-1-oxo-2-buten-1-yl]amino}-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholin-4-yl)-ethoxy]-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholin-4-yl)-ethoxy]-7-[(R)-(tetrahydrofuran-2-yl)methoxy]-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-7-[2-(2,2-dimethyl-6-oxo-morpholin-4-yl)-ethoxy]-6-[(S)-(tetrahydrofuran-2-yl)methoxy]-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{2-[4-(2-oxo-morpholin-4-yl)-piperidin-1-yl]-ethoxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[1-(tert.-butyloxycarbonyl)-piperidin-4-yloxy]-7-methoxy-chinazoline,

4-[(3-chlor-4-fluor-phenyl)amino]-6-(trans-4-amino-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(trans-4-methansulfonylamino-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(tetrahydropyran-3-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-methyl-piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(methoxymethyl)carbonyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(piperidin-3-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[1-(2-acetylamino-ethyl)-piperidin-4-yloxy]-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-ethoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-((S)-tetrahydrofuran-3-yloxy)-7-hydroxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methoxy-ethoxy)-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{trans-4-[(dimethylamino)sulfonylamino]-cyclohexan-1-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{trans-4-[(morpholin-4-yl)carbonylamino]-cyclohexan-1-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{trans-4-[(morpholin-4-yl)sulfonylamino]-cyclohexan-1-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-acetylamino-ethoxy)-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(tetrahydropyran-4-yloxy)-7-(2-methansulfonylamino-ethoxy)-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(piperidin-1-yl)carbonyl]-piperidin-4-yloxy}-7-methoxy-

chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-aminocarbonylmethyl-piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(cis-4-{N-[(tetrahydropyran-4-yl)carbonyl]-N-methyl-amino}-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(cis-4-{N-[(morpholin-4-yl)carbonyl]-N-methyl-amino}-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(cis-4-{N-[(morpholin-4-yl)sulfonyl]-N-methyl-amino}-cyclohexan-1-yloxy)-7-methoxy- chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(trans-4-ethansulfonylamino-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-methansulfonyl-piperidin-4-yloxy)-7-ethoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-methansulfonyl-piperidin-4-yloxy)-7-(2-methoxy-ethoxy)-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[1-(2-methoxy-acetyl)-piperidin-4-yloxy]-7-(2-methoxy-ethoxy)-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(cis-4-acetylamino-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-Ethinyl-phenyl)amino]-6-[1-(tert.-butyloxycarbonyl)-piperidin-4-yloxy]-7-methoxy-chinazoline, 4-[(3-Ethinyl-phenyl)amino]-6-(tetrahydropyran-4-yloxy]-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(cis-4-{N-[(piperidin-1-yl)carbonyl]-N-methyl-amino}-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(cis-4-{N-[(4-methyl-piperazin-1-yl)carbonyl]-N-methyl-amino}-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{cis-4-[(morpholin-4-yl)carbonylamino]-cyclohexan-1-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[2-(2-oxopyrrolidin-1-yl)ethyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-

[(morpholin-4-yl)carbonyl]-piperidin-4-yloxy}-7-(2-methoxy-ethoxy)-chinazoline, 4-[(3-Ethinyl-phenyl)amino]-6-(1-acetyl-piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-Ethinyl-phenyl)amino]-6-(1-methyl-piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-Ethinyl-phenyl)amino]-6-(1-methansulfonyl-piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-methyl-piperidin-4-yloxy)-7(2-methoxy-ethoxy)-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-isopropyloxycarbonyl-piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(cis-4-methylamino-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{cis-4-[N-(2-methoxy-acetyl)-N-methyl-amino]-cyclohexan-1-yloxy}-7-methoxy-chinazoline, 4-[(3-Ethinyl-phenyl)amino]-6-(piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-Ethinyl-phenyl)amino]-6-[1-(2-methoxy-acetyl)-piperidin-4-yloxy]-7-methoxy-chinazoline, 4-[(3-Ethinyl-phenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(cis-2,6-dimethyl-morpholin-4-yl)carbonyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(2-methyl-morpholin-4-yl)carbonyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(S,S)-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)carbonyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(N-methyl-N-2-methoxyethyl-amino)carbonyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-ethyl-piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(2-methoxyethyl)carbonyl]-piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-{1-[(3-methoxypropyl-amino)-carbonyl]-

piperidin-4-yloxy}-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[cis-4-(N-methansulfonyl-N-methyl-amino)-cyclohexan-1-yloxy]-7-methoxy-chinazoline , 4-[(3-chlor-4-fluor-phenyl)amino]-6-[cis-4-(N-acetyl-N-methyl-amino)-cyclohexan-1-yloxy]-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(trans-4-methylamino-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[trans-4-(N-methansulfonyl-N-methyl-amino)-cyclohexan-1-yloxy]-7-methoxy-chinazoline , 4-[(3-chlor-4-fluor-phenyl)amino]-6-(trans-4-dimethylamino-cyclohexan-1-yloxy)-7-methoxy-chinazoline , 4-[(3-chlor-4-fluor-phenyl)amino]-6-(trans-4-{N-[(morpholin-4-yl)carbonyl]-N-methyl-amino}-cyclohexan-1-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-[2-(2,2-dimethyl-6-oxo-morpholin-4-yl)-ethoxy]-7-[(S)-(tetrahydrofuran-2-yl)methoxy]-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-methansulfonyl-piperidin-4-yloxy)-7-methoxy-chinazoline, 4-[(3-chlor-4-fluor-phenyl)amino]-6-(1-cyano-piperidin-4-yloxy)-7-methoxy-chinazoline, Cetuximab, Trastuzumab, ABX-EGF and Mab ICR-62. These compounds may be used, as available, in the form of their racemates, enantiomers or diastereoisomers, or in the form of pharamacologically acceptable acid addition salts thereof, or in the form of their solvates and/or hydrates. These compounds are disclosed in the prior art, e.g. in WO 96/30347, WO 97/02266, WO 99/35146, WO 00/31048, WO 00/78735, WO 01/34574, WO 01/61816, WO 01/77104, WO02/18351, WO 02/18372, WO 02/18373, WO 02/18376, WO 02/50043, WO 03/082290, Cancer Research 2004, **64**:11 (3958-3965), Am J Health-Syst Pharm 2000, 57(15), 2063-2076, Clinical Therapeutics 1999, 21(2), 309-318, WO 98/50433, and WO 95/20045.

Endothelin-antagonists may preferably be selected from the group consisting of tezosentan, bosentan, enrasentan, sixtasentan, T-0201, BMS-193884, K-8794, PD-156123, PD-156707, PD-160874, PD-180988, S-0139 and ZD-1611. Any reference to endothelin-antagonists within the scope of the present invention includes a reference to the salts, preferably pharmacologically acceptable acid addition salts, or derivatives which may be formed from the endothelin-antagonists.

These combinations may be administered either simultaneously or sequentially.

For pharmaceutical use the compounds according to the invention are preferably used for warm-blooded vertebrates, particularly humans, in doses of 0.0001-100 mg/kg of body weight.

These compounds may be administered either on their own or in conjunction with other active substances by intravenous, subcutaneous, intramuscular, intraperitoneal or intranasal route, by inhalation, or transdermally, or orally, whilst aerosol formulations are particularly suitable for inhalation.

For administration they are formulated with one or more conventional inert solid, semisolid or liquid carriers e.g. with starch, different types of cellulose, lactose, mannitol, sorbitol, glucose, calcium phosphate, hard fat, fatty alcohols, glycerol, medium chained triglycerides and related esters, polyethylene glycol, refined specialty

oils, water, water/ethanol, water/glycerol, water/sorbitol, water/polyethylene glycol, propylene glycol,and/or functional excipients, e.g. with polyvinylpyrrolidone, hydroxypropylmethylcellulose, sodium carboxymethyl-cellulose, sodium starch glycolate, silicon dioxide, polysorbates, poloxamers, gelucires, magnesium stearate, citric acid, tartaric acid, or suitable mixtures thereof in conventional galenic preparations such as plain or coated tablets, capsules, powders, injectable solutions, ampoules, suspensions, solutions, sprays or suppositories.

The following examples of formulations illustrate the present invention without representing a limitation of its scope.

Example F1: Coated tablet containing 75 mg of active substance

Composition

1 tablet core contains:

active substance	75.0 mg
calcium phosphate	131.0 mg
polyvinylpyrrolidone	10.0 mg
carboxymethylcellulose sodium	10.0 mg
silicon dioxide	2.5 mg
magnesium stearate	1.5 mg
	230.0 mg

Preparation (direct compression)

The active substance is mixed with all components, sieved and compressed in a tablet-making machine to form tablets of the desired shape.

Weight of core:	230 mg
Appearance of core:	9 mm, biconvex

The tablet cores thus produced are coated with a film consisting essentially of hydroxypropylmethylcellulose.

Weight of coated tablet: 240 mg.

Example F2: Tablet containing 100 mg of active substance

Composition

1 tablet contains:

active substance	100.0 mg
lactose	80.0 mg
corn starch	34.0 mg
hydroxypropylmethylcellulose	4.0 mg
magnesium stearate	2.0 mg
	220.0 mg

Preparation (wet granulation)

The active substance, lactose and starch are mixed together and uniformly moistened with an aqueous solution of the hydroxypropylmethylcellulose. After the moist composition has been screened (2.0 mm mesh size) and dried in a rack-type drier at 50°C it is screened again (1.5 mm mesh size)

and the lubricant is added. The finished mixture is compressed to form tablets.

Weight of tablet:	220 mg
Appearance of tablet:	10 mm, flat faced

with bevelled edges and breaking notch on one side.

Example F3: Tablet containing 150 mg of active substance

Composition

1 tablet contains:

active substance	150.0 mg
lactose	85.0 mg
microcrystalline cellulose	40.0 mg
polyvinylpyrrolidone	10.0 mg
silicon dioxide	10.0 mg
magnesium stearate	5.0 mg
	300.0 mg

Preparation (dry granulation)

The active substance mixed with lactose, polyvinyl-pyrrolidone,and parts of the microcrystalline cellulose, magnesium stearate is compacted e.g. on a roller compactor. The ribbons are broken up in fine granules through a screen with a mesh size of 0.8 mm. After subsequent sieving through a screen with a mesh size of 0.5 mm and blending with the remaining components, tablets are pressed from the mixture.

Weight of tablet:	300 mg

Appearance of tablet: 10 mm, flat

Example F4: Hard gelatine capsule containing 150 mg of active substance

Composition

1 capsule contains:

active substance	150.0 mg
lactose	85.0 mg
microcrystalline cellulose	40.0 mg
polyvinylpyrrolidone	10.0 mg
silicon dioxide	10.0 mg
magnesium stearate	5.0 mg
	300.0 mg

Preparation

The active substance mixed with lactose, polyvinyl-pyrrolidone,and parts of the microcrystalline cellulose, magnesium stearate is compacted e.g. on a roller compactor. The ribbons are broken up in fine granules through a screen with a mesh size of 0.8 mm. After subsequent sieving through a screen with a mesh size of 0.5 mm and blending with the remaining components, the finished mixture is packed into size 1 hard gelatine capsules.

Capsule filling: approx. 300 mg
Capsule shell: size 1 hard gelatine capsule.

Example F5: Suppository containing 150 mg of active substance

1 suppository contains:

active substance	150.0 mg
polyethyleneglycol 1500	800.0 mg
polyethyleneglycol 6000	850.0 mg
polyoxyl 40 hydrogenated castor oil	200.0 mg
	2,000.0 mg

Preparation

After the suppository mass has been melted the active substance is homogeneously distributed therein and the melt is poured into chilled moulds.

Example F6: Suspension containing 50 mg of active substance

100 ml of suspension contains

active substance		1.00 g
carboxymethylcellulose sodium		0.10 g
methyl p-hydroxybenzoate		0.05 g
propyl p-hydroxybenzoate		0.01 g
glucose		10.00 g
glycerol		5.00 g
70% sorbitol solution		20.00 g
flavouring		0.30 g
dist. water	ad	100 ml

Preparation

The distilled water is heated to 70°C. The methyl and propyl p-hydroxybenzoates together with the glycerol and sodium salt of carboxymethylcellulose are dissolved therein

with stirring. The solution is cooled to ambient temperature and the active substance is added and homogeneously dispersed therein with stirring. After the sugar, the sorbitol solution and the flavouring have been added and dissolved, the suspension is evacuated with stirring to eliminate air.

Thus, 5 ml of suspension contains 50 mg of active substance.

Example F7: Ampoule containing 10 mg active substance

Composition

active substance	10.0 mg
0.01 N hydrochloric acid	q.s.
double-distilled water	ad 2.0 ml

Preparation

The active substance is dissolved in the necessary amount of 0.01 N HCl, made isotonic with sodium chloride, filtered sterile and transferred into a 2 ml ampoule.

Example F8: Ampoule containing 50 mg of active substance

Composition

active substance	50.0 mg
0.01 N hydrochloric acid	q.s.
double-distilled water	ad 10.0 ml

Preparation

The active substance is dissolved in the necessary amount of 0.01 N HCl, made isotonic with sodium chloride, filtered sterile and transferred into a 10 ml ampoule.

Example F9: Capsule for powder inhalation containing 5 mg of active substance

1 capsule contains

active substance	5.0 mg
lactose for inhalation	15.0 mg
	20.0 mg

Preparation

The active substance is mixed with lactose for inhalation. The mixture is packed into capsules in a capsule-making machine (weight of the empty capsule approx. 50 mg).

weight of capsule: 70.0 mg
size of capsule = size 3

Example F10: Solution for inhalation for a hand-held nebuliser containing 2.5 mg active substance

1 spray contains

active substance	2.500 mg
benzalkonium chloride	0.001 mg

1N hydrochloric acid	q.s.
ethanol/water (50/50)	ad 15.000 mg


Preparation

The active substance and benzalkonium chloride are dissolved in ethanol/water (50/50). The pH of the solution is adjusted with 1N hydrochloric acid. The resulting solution is filtered and transferred into suitable containers for use in hand-held nebulisers (cartridges).

Contents of the container: 4.5 g

CLAIMS

1. Use of the indolinones of general formula



(I),

substituted in the 6 position, wherein

X denotes an oxygen or sulphur atom,

R_1 denotes a hydrogen atom or a prodrug group,

R_2 denotes a carboxy group, a straight-chain or branched C_{1-6}-alkoxy-carbonyl group, a C_{4-7}-cycloalkoxy-carbonyl or an aryloxycarbonyl group,

a straight-chain or branched C_{1-6}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a phenyl, heteroaryl, carboxy, C_{1-3}-alkoxy-carbonyl, aminocarbonyl, C_{1-3}-alkylamino-carbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

a straight-chain or branched C_{2-6}-alkoxy-carbonyl group, which is terminally substituted in the alkyl moiety by a

chlorine atom or a hydroxy, C_{1-3}-alkoxy, amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group,

an aminocarbonyl or methylaminocarbonyl group, an ethylaminocarbonyl group optionally substituted in the 2 position of the ethyl group by a hydroxy or C_{1-3}-alkoxy group or a di-(C_{1-2}-alkyl)-aminocarbonyl group,

R_3 denotes a hydrogen atom, a C_{1-6}-alkyl, C_{3-7}-cycloalkyl, trifluoromethyl or heteroaryl group,

a phenyl or naphthyl group, a phenyl or naphthyl group mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a trifluoromethyl, C_{1-3}-alkyl or C_{1-3}-alkoxy group, whilst in the event of disubstitution the substituents may be identical or different and wherein the abovementioned unsubstituted as well as the mono- and disubstituted phenyl and naphthyl groups may additionally be substituted

by a hydroxy, hydroxy-C_{1-3}-alkyl or C_{1-3}-alkoxy-C_{1-3}-alkyl group,

by a cyano, carboxy, carboxy-C_{1-3}-alkyl, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylamino-carbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group,

by a nitro group,

by an amino, C_{1-3}-alkylamino, di-(C_{1-3}-alkyl)-amino or amino-C_{1-3}-alkyl group,

by a C_{1-3}-alkylcarbonylamino, N-(C_{1-3}-alkyl)-C_{1-3}-alkyl-carbonylamino, C_{1-3}-alkylcarbonylamino-C_{1-3}-alkyl, N-(C_{1-3}-alkyl)-C_{1-3}-alkylcarbonylamino-C_{1-3}-alkyl, C_{1-3}-alkyl-sulphonylamino, C_{1-3}-alkylsulphonylamino-C_{1-3}-alkyl, N-(C_{1-3}-alkyl)-C_{1-3}-alkylsulphonylamino-C_{1-3}-alkyl or aryl-C_{1-3}-alkylsulphonylamino group,

by a cycloalkylamino, cycloalkyleneimino, cyclo-alkyleneiminocarbonyl, cycloalkyleneimino-C_{1-3}-alkyl, cycloalkyleneiminocarbonyl-C_{1-3}-alkyl or cycloalkyleneiminosulphonyl-C_{1-3}-alkyl group having 4 to 7 ring members in each case, whilst in each case the methylene group in position 4 of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH or -N(C_{1-3}-alkyl) group,

or by a heteroaryl or heteroaryl-C_{1-3}-alkyl group,

R_4 denotes a C_{3-7}-cycloalkyl group,

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group or replaced by an -NH or -N(C_{1-3}-alkyl) group,


or a phenyl group substituted by the group R_6, which may additionally be mono- or disubstituted by fluorine, chlorine, bromine or iodine atoms, by C_{1-5}-alkyl, trifluoromethyl, hydroxy, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, amino, acetylamino, C_{1-3}-alkyl-sulphonylamino, aminocarbonyl,

C_{1-3}-alkyl-aminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, aminosulphonyl, C_{1-3}-alkyl-aminosulphonyl, di-(C_{1-3}-alkyl)-aminosulphonyl, nitro or cyano groups, wherein the substituents may be identical or different and wherein

R_6 denotes a hydrogen, fluorine, chlorine, bromine or iodine atom,

a cyano, nitro, amino, C_{1-5}-alkyl, C_{3-7}-cycloalkyl, trifluoromethyl, phenyl, tetrazolyl or heteroaryl group,

the group of formula



,

wherein the hydrogen atoms bound to a nitrogen atom may in each case be replaced independently of one another by a C_{1-3}-alkyl group,

a C_{1-3}-alkoxy group, a C_{1-3}-alkoxy-C_{1-3}-alkoxy, phenyl-C_{1-3}-alkoxy, amino-C_{2-3}-alkoxy, C_{1-3}-alkylamino-C_{2-3}-alkoxy, di-(C_{1-3}-alkyl)-amino-C_{2-3}-alkoxy, phenyl-C_{1-3}-alkylamino-C_{2-3}-alkoxy, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino-C_{2-3}-alkoxy, C_{5-7}-cycloalkyleneimino-C_{2-3}-alkoxy or C_{1-3}-alkylmercapto group,

a carboxy, C_{1-4}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkyl-amino-carbonyl, N-(C_{1-5}-alkyl)-C_{1-3}-alkylaminocarbonyl, phenyl-C_{1-3}-alkylamino-carbonyl, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino-carbonyl,

piperazinocarbonyl or N-(C_{1-3}-alkyl)-piperazinocarbonyl group,

a C_{1-3}-alkylaminocarbonyl or N-(C_{1-5}-alkyl)-C_{1-3}-alkylaminocarbonyl group wherein an alkyl moiety is substituted by a carboxy or C_{1-3}-alkoxycarbonyl group or in the 2 or 3 position by a di-(C_{1-3}-alkyl)-amino, piperazino, N-(C_{1-3}-alkyl)-piperazino or a 4- to 7-membered cycloalkyleneimino group,

a C_{3-7}-cycloalkyl-carbonyl group,

wherein the methylene group in the 4 position of the 6- or 7-membered cycloalkyl moiety may be substituted by an amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group or replaced by an -NH or -N(C_{1-3}-alkyl) group,

a 4- to 7-membered cycloalkyleneimino group wherein

a methylene group linked to the imino group may be replaced by a carbonyl or sulphonyl group or

the cycloalkylene moiety may be fused to a phenyl ring or

one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl group and/or

in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be substituted by a carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl,

phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl), -N(phenyl), -N(C_{1-3}-alkyl-carbonyl) or -N(benzoyl) group,

a C_{1-4}-alkyl group substituted by the group R_7, wherein

R_7 denotes a C_{3-7}-cycloalkyl group,

whilst the methylene group in the 4 position of a 6- or 7-membered cycloalkyl group may be substituted by an amino, C_{1-3}-alkylamino or di-(C_{1-3}-alkyl)-amino group or replaced by an -NH or -N(C_{1-3}-alkyl) group or

in a 5- to 7-membered cycloalkyl group a $-(CH_2)_2$ group may be replaced by a -CO-NH group, a $-(CH_2)_3$ group may be replaced by a -NH-CO-NH or -CO-NH-CO group or a $-(CH_2)_4$ group may be replaced by a -NH-CO-NH-CO group, whilst in each case a hydrogen atom bound to a nitrogen atom may be replaced by a C1-3-alkyl group,

an aryl or heteroaryl group,

a hydroxy or C_{1-3}-alkoxy group,

an amino, C_{1-7}-alkylamino, di-(C_{1-7}-alkyl)-amino, phenylamino, N-phenyl-C_{1-3}-alkyl-amino, phenyl-C_{1-3}-alkyl-amino, N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkylamino or di-(phenyl-C_{1-3}-alkyl)-amino group,

an ω-hydroxy-C_{2-3}-alkyl-amino, N-(C_{1-3}-alkyl)-ω-hydroxy-C_{2-3}-alkyl-amino, di-(ω-hydroxy-C_{2-3}-alkyl)-amino, di-(ω-(C_{1-3}-alkoxy)-C_{2-3}-alkyl)-amino or N-(dioxolan-2-yl)-C_{1-3}-alkyl-amino group,

a C_{1-3}-alkylcarbonylamino-C_{2-3}-alkyl-amino or C_{1-3}-alkylcarbonylamino-C_{2-3}-alkyl-N-(C_{1-3}-alkyl)-amino group,

a C_{1-3}-alkylsulphonylamino, N-(C_{1-3}-alkyl)-C_{1-3}-alkyl-sulphonylamino, C_{1-3}-alkylsulphonylamino-C_{2-3}-alkyl-amino or C_{1-3}-alkylsulphonylamino-C_{2-3}-alkyl-N-(C_{1-3}-alkyl)-amino group,

a hydroxycarbonyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-hydroxycarbonyl-C_{1-3}-alkyl-amino group,

a guanidino group wherein one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl group,

a group of formula

$$-N(R_8)-CO-(CH_2)_n-R_9 \quad (II),$$

wherein

R_8 denotes a hydrogen atom or a C_{1-3}-alkyl group,

n denotes one of the numbers 0, 1, 2 or 3 and

R_9 denotes an amino, C_{1-4}-alkylamino, di-(C_{1-4}-alkyl)-amino, phenylamino, N-(C_{1-4}-alkyl)-phenylamino, benzylamino, N-(C_{1-4}-alkyl)-benzylamino or C_{1-4}-alkoxy group, a 4- to 7-membered cycloalkyleneimino group, whilst in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl), -N(phenyl), -N(C_{1-3}-alkyl-carbonyl) or -N(benzoyl) group, or, if n denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

$$-N(R_{10})-(CH_2)_m-(CO)_o-R_{11} \quad (III),$$

wherein

R_{10} denotes a hydrogen atom, a C_{1-3}-alkyl group, a C_{1-3}-alkylcarbonyl, arylcarbonyl, phenyl-C_{1-3}-alkyl-carbonyl, C_{1-3}-alkylsulphonyl, arylsulphonyl or phenyl-C_{1-3}-alkylsulphonyl group,

m denotes one of the numbers 1, 2, 3 or 4,

o denotes the number 1 or, if m denotes one of the numbers 2, 3 or 4, o may also denote the number 0 and

R_{11} denotes an amino, C_{1-4}-alkylamino, di-(C_{1-4}-alkyl)-amino, phenylamino, N-(C_{1-4}-alkyl)-phenylamino, benzylamino, N-(C_{1-4}-alkyl)-benzylamino, C_{1-4}-alkoxy or C_{1-3}-alkoxy-C_{1-3}-alkoxy group, a di-(C_{1-4}-alkyl)-amino-C_{1-3}-alkylamino group optionally substituted in the 1 position

by a C_{1-3}-alkyl group or a 4- to 7-membered cycloalkyleneimino group, wherein the cycloalkylene moiety may be fused to a phenyl ring or in each case the methylene group in the 4 position of a 6- or 7-membered cycloalkyleneimino group may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl), -N(phenyl), -N(C_{1-3}-alkyl-carbonyl) or -N(benzoyl) group,

a C_{4-7}-cycloalkylamino, C_{4-7}-cycloalkyl-C_{1-3}-alkylamino or C_{4-7}-cycloalkenylamino group wherein position 1 of the ring is not involved in the double bond and wherein the abovementioned groups may each additionally be substituted at the amino-nitrogen atom by a C_{5-7}-cycloalkyl, C_{2-4}-alkenyl or C_{1-4}-alkyl group,

a 4- to 7-membered cycloalkyleneimino group, wherein

the cycloalkylene moiety may be fused to a phenyl group or to an oxazolo, imidazolo, thiazolo, pyridino, pyrazino or pyrimidino group optionally substituted by a fluorine, chlorine, bromine or iodine atom, by a nitro, C_{1-3}-alkyl, C_{1-3}-alkoxy or amino group, and/or

one or two hydrogen atoms may each be replaced by a C_{1-3}-alkyl, C_{5-7}-cycloalkyl or phenyl group and/or

the methylene group in the 3 position of a 5-membered cycloalkyleneimino group may be substituted by a hydroxy, hydroxy-C_{1-3}-alkyl, C_{1-3}-alkoxy or C_{1-3}-alkoxy-C_{1-3}-alkyl group,

the methylene group in the 3 or 4 position of a 6- or 7-membered cycloalkyleneimino group may in each case be substituted by a hydroxy, hydroxy-C_{1-3}-alkyl, C_{1-3}-alkoxy, C_{1-3}-alkoxy-C_{1-3}-alkyl, carboxy, C_{1-4}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl, di-(C_{1-3}-alkyl)-aminocarbonyl, phenyl-C_{1-3}-alkylamino or N-(C_{1-3}-alkyl)-phenyl-C_{1-3}-alkyl-amino group or

may be replaced by an oxygen or sulphur atom, by a sulphinyl, sulphonyl, -NH, -N(C_{1-3}-alkyl-), -N(phenyl), -N(phenyl-C_{1-3}-alkyl-), -N(C_{1-3}-alkyl-carbonyl-), -N(C_{1-4}-alkyl-hydroxy-carbonyl-), -N(C_{1-4}-alkoxy-carbonyl-), -N(benzoyl-) or -N(phenyl-C_{1-3}-alkyl-carbonyl-) group,

wherein a methylene group linked to an imino-nitrogen atom of the cycloalkyleneimino group may be replaced by a carbonyl or sulphonyl group or in a 5- to 7-membered monocyclic cycloalkyleneimino group or a cycloalkyleneimino group fused to a phenyl group the two methylene groups linked to the imino-nitrogen atom may each be replaced by a carbonyl group,

or R_6 denotes a C_{1-4}-alkyl group which is substituted by a carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-3}-alkylaminocarbonyl or di-(C_{1-3}-alkyl)-aminocarbonyl group or by a 4- to 7-membered cycloalkyleneiminocarbonyl group,

an N-(C_{1-3}-alkyl)-C_{2-4}-alkanoylamino group which is additionally substituted in the alkyl moiety by a carboxy or C_{1-3}-alkoxycarbonyl group,

a group of formula

$-N(R_{12})-CO-(CH_2)_p-R_{13}$ (IV),

wherein

R_{12} denotes a hydrogen atom, a C_{1-6}-alkyl or C_{3-7}-cycloalkyl group or a C_{1-3}-alkyl group terminally substituted by a phenyl, heteroaryl, trifluoromethyl, hydroxy, C_{1-3}-alkoxy, aminocarbonyl, C_{1-4}-alkylamino-carbonyl, di-(C_{1-4}-alkyl)-amino-carbonyl, C_{1-3}-alkyl-carbonyl, C_{1-3}-alkyl-sulphonyl-amino, N-(C_{1-3}-alkyl)-C_{1-3}-alkyl-sulphonylamino, C_{1-3}-alkyl-aminosulphonyl or di-(C_{1-3}-alkyl)-aminosulphonyl group and

p denotes one of the numbers 0, 1, 2 or 3 and

R_{13} assumes the meanings of the abovementioned group R_7, or, if p denotes one of the numbers 1, 2 or 3, it may also denote a hydrogen atom,

a group of formula

$-N(R_{14})-(CH_2)_q-(CO)_r-R_{15}$ (V),

wherein

R_{14} denotes a hydrogen atom, a C_{1-4}-alkyl group, a C_{1-3}-alkylcarbonyl, arylcarbonyl, phenyl-C_{1-3}-alkylcarbonyl, heteroarylcarbonyl, heteroaryl-C_{1-3}-alkylcarbonyl, C_{1-4}-alkylsulphonyl, arylsulphonyl, phenyl-C_{1-3}-alkylsulphonyl, heteroarylsulphonyl or heteroaryl-C_{1-3}-alkyl-sulphonyl group,

q denotes one of the numbers 1, 2, 3 or 4,

r denotes the number 1 or, if q is one of the numbers 2, 3 or 4, it may also denote the number 0 and

R_{15} assumes the meanings of the abovementioned group R_7,

a group of formula

$-N(R_{16})-SO_2-R_{17}$ (VI),

wherein

R_{16} denotes a hydrogen atom or a C_{1-4}-alkyl group optionally terminally substituted by a cyano, trifluoromethyl-carbonylamino or N-(C_{1-3}-alkyl)-trifluoromethyl-carbonyl-amino group and

R_{17} denotes a C_{1-3}-alkyl group,

an amino group substituted by a di-(C_{1-3}-alkyl)-amino-C_{1-3}-alkyl-carbonyl or di-(C_{1-3}-alkyl)-amino-C_{1-3}-alkyl-sulphonyl group and a di-(C_{1-3}-alkyl)-aminocarbonyl-C_{1-3}-alkyl group,

or an N-(C_{1-3}-alkyl)-C_{1-5}-alkylsulphonylamino or N-(C_{1-3}-alkyl)-phenylsulphonylamino group wherein the alkyl moiety is additionally substituted by a cyano or carboxy group,

wherein all the single-bonded or fused phenyl groups contained in the groups mentioned under R_6 may be mono- or disubstituted by fluorine, chlorine, bromine or iodine atoms, by C_{1-5}-alkyl, trifluoromethyl, hydroxy, C_{1-3}-alkoxy, carboxy, C_{1-3}-alkoxycarbonyl, aminocarbonyl, C_{1-4}-alkylamino-carbonyl, di-(C_{1-4}-alkyl)-amino-carbonyl, aminosulphonyl, C_{1-3}-alkyl-aminosulphonyl, di-(C_{1-3}-alkyl)-aminosulphonyl, C_{1-3}-alkyl-sulphonylamino, nitro or cyano groups, wherein the substituents may be identical or different, or two adjacent hydrogen atoms of the phenyl groups may be replaced by a methylenedioxy group,

and

R_5 denotes a hydrogen atom or a C_{1-3}-alkyl group,

wherein by an aryl group is meant a phenyl or naphthyl group optionally mono- or disubstituted by a fluorine, chlorine, bromine or iodine atom, by a cyano, trifluoromethyl, nitro, carboxy, aminocarbonyl, C_{1-3}-alkyl or C_{1-3}-alkoxy group and

by a heteroaryl group is meant a monocyclic 5- or 6-membered heteroaryl group optionally substituted by a C_{1-3}-alkyl group in the carbon skeleton, wherein

the 6-membered heteroaryl group contains one, two or three nitrogen atoms and

the 5-membered heteroaryl group contains an imino group optionally substituted by a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group, an oxygen or sulphur atom or

an imino group optionally substituted by a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group or an oxygen or sulphur atom and additionally a nitrogen atom or

an imino group optionally substituted by a C_{1-3}-alkyl or phenyl-C_{1-3}-alkyl group and two nitrogen atoms,

and moreover a phenyl ring may be fused to the abovementioned monocyclic heterocyclic groups via two adjacent carbon atoms and the bonding takes place via a nitrogen atom or via a carbon atom of the heterocyclic moiety or a fused phenyl ring,

some or all of the hydrogen atoms in the abovementioned alkyl and alkoxy groups or in the alkyl moieties contained in the above-defined groups of formula I may be replaced by fluorine atoms,

and the hydrogen atom of any carboxy group present or a hydrogen atom bound to a nitrogen atom may each be replaced by a group which can be cleaved *in vivo*,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof,

for the preparation of a medicament for the prevention or treatment of a fibrotic disease selected from the group consisting of fibrosis and remodeling of lung tissue in chronic obstructive pulmonary disease, fibrosis and remodeling of lung tissue in chronic bronchitis, fibrosis and remodeling of lung tissue in emphysema, lung fibrosis and pulmonary diseases with a fibrotic component, fibrosis and remodeling in asthma, fibrosis in rheumatoid arthritis, virally induced hepatic cirrhosis, radiation-induced fibrosis, post angioplasty restenosis, chronic glomerulonephritis, renal fibrosis in patients receiving cyclosporine and renal fibrosis due to high blood pressure, diseases of the skin with a fibrotic component, and excessive scarring.

2. Use in accordance with claim 1 wherein the substituent in the 6 position of the substituted indolinone of general formula I comprises a substituted amido group.

3. Use in accordance with claim 1 wherein the substituted indolinone of general formula I is selected from the group consisting of:

(a) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(b) 3-Z-[(1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carbamoyl-2-indolinone,

(c) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(d) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(e) 3-Z-[1-(4-((2,6-dimethyl-piperidin-1-yl)-methyl)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(f) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(g) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(h) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethoxycarbonyl-2-indolinone,

(i) 3-Z-[1-(4-(dimethylaminomethyl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(j) 3-Z-[1-(4-(N-acetyl-N-dimethylaminocarbonylmethyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(k) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(l) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(m) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(n) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(o) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(p) 3-Z-[1-(4-(N-dimethylaminocarbonylmethyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(q) 3-Z-[1-(4-(N-((2-dimethylamino-ethyl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(r) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone and

(s) 3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

4. Use in accordance with claim 1 wherein the substituted indolinone of general formula I is selected from the group consisting of:

(t) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(u) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone,

(v) 3-Z-[1-(3-cyano-4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(w)3-Z-[1-(3-methoxy-4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(x) 3-Z-[1-(4-(N-aminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(y) 3-Z-[1-(4-(N-(N-(2-dimethylamino-ethyl)-N-methyl-aminomethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(z) 3-Z-[1-(4-(N-(di-(2-hydroxy-ethyl)-amino-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(aa) 3-Z-[1-(4-(N-(imidazol-1-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ab) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ac) 3-Z-[1-(4-(N-((4-methyl-[1,4]diazepan-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ad) 3-Z-[1-(4-(N-((1-methyl-piperidin-4-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ae) 3-Z-[1-(2,3-dimethyl-4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(af) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ag) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-carbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ah) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ai) 3-Z-[1-(4-(N-((3-dimethylamino-propyl)-aminocarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(aj) 3-Z-[1-anilino-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(ak) 3-Z-[1-(4-(N-methyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(al) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone

(am) 3-Z-[1-(4-(4-methylpiperazin-1-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(an) 3-Z-[1-(4-methylaminomethyl-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(ao) 3-Z-[1-(4-(morpholin-4-yl-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(ap) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(aq) 3-Z-[1-(4-(di-(2-hydroxy-ethyl)-amino-methyl)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(ar) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(as) 3-Z-[1-(4-(N-(morpholin-4-yl-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(at) 3-Z-[1-(4-(N-(N-(2-dimethylamino-ethyl)-N-methyl-aminomethylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(au) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-carboxy-2-indolinone

(av) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-carboxy-2-indolinone

(aw) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(ax) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(ay) 3-Z-[1-(4-(1-methyl-imidazol-2-yl)-anilino)-1-(3-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(az) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-(4-(2-carboxy-ethyl)-phenyl)-methylene]-6-methoxycarbonyl-2-indolinone

(ba) 3-Z-[1-(4-(2-dimethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bb) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bc) 3-Z-[1-((1-methyl-piperidin-4-yl)-amino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bd) 3-Z-[1-(trans-4-dimethylamino-cyclohexylamino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(be) 3-Z-[1-(4-(2-diethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bf) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-propionyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bg) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bh) 3-Z-[1-cyclohexylamino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bi) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bj) 3-Z-[1-(3-diethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bk) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bl) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bm) 3-Z-[1-anilino-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bn) 3-Z-[1-(4-ethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bo) 3-Z-[1-(4-((2-diethylamino-ethyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bp) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bq) 3-Z-[1-(4-(N-methyl-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(br) 3-Z-[1-(4-methoxycarbonyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bs) 3-Z-[1-(4-carboxy-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bt) 3-Z-[1-(4-(N-(dimethylamino-carbonylmethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(bu) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(bv) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(bw) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(bx) 3-Z-[1-(4-(2-dimethylamino-ethoxy)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(by) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(bz) 3-Z-[1-(4-dimethylaminomethyl-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(ca) 3-Z-[1-(4-(N-(2-dimethylamino-ethyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cb) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cc) 3-Z-[1-(4-carbamoyl-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(cd) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(ce) 3-Z-[1-(4-((4-methyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(cf) 3-Z-[1-(4-((4-methyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cg) 3-Z-[1-(4-((4-ethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(ch) 3-Z-[1-(4-(N-ethyl-N-(2-dimethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(ci) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-methylene]-6-diethylcarbamoyl-2-indolinone

(cj) 3-Z-[1-(4-((cis-3,5-dimethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(ck) 3-Z-[1-(4-((4-ethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cl) 3-Z-[1-(4-(N-(2-diethylamino-ethyl)-carbamoyl)-anilino)-1-phenyl-methylene]-6-ethylmethylcarbamoyl-2-indolinone

(cm) 3-Z-[1-(4-((cis-3,5-dimethyl-piperazin-1-yl)-carbonyl)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(cn) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-methylsulphonyl-amino)-anilino)-1-phenyl-methylene]-6-ethylcarbamoyl-2-indolinone

(co) 3-Z-[1-(4-(N-dimethylaminomethylcarbonyl-N-methyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(cp) 3-Z-[1-(4-dimethylaminomethyl-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(cq)3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-methylene]-6-methoxycarbonyl-2-indolinone

(cr) 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

(cs) 3-Z-[1-(4-dimethylaminomethyl-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone

(ct) 3-Z-[1-(4-(piperidin-1-yl-methyl)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone, and

(cu) 3-Z-[1-(4-(N-(3-dimethylamino-propyl)-N-acetyl-amino)-anilino)-methylene]-6-ethylcarbamoyl-2-indolinone,

the tautomers, the diastereomers, the enantiomers, the mixtures thereof and the salts thereof.

5. Use in accordance with claim 1 wherein the substituted indolinone of general formula I is selected from 3-Z-[1-(4-(N-((4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino)-1-phenyl-

methylene]-6-methoxycarbonyl-2-indolinone or the monoethanesulfonate salt of this compound.

6. Use in accordance with any one of claims 1 to 5 wherein the disease is selected from the group consisting of the lung fibrosis and pulmonary diseases with a fibrotic component selected from idiopathic pulmonary fibrosis, giant cell interstitial pneumonia, sarcodosis, cystic fibrosis, respiratory distress syndrome, drug-induced lung fibrosis, granulomatosis, silicosis, asbestosis, systemic scleroderma, the virally induced hepatic cirrhosis selected from hepatitis C induced hepatic cirrhosis, and the diseases of the skin with a fibrotic component selected from scleroderma, sarcodosis and systemic lupus erythematosus.

7. Use in accordance with any one of claims 1 to 6 wherein the disease is idiopathic pulmonary fibrosis.

8. Use in accordance with any one of claims 1 to 7 wherein the treatment is a combined treatment with a further pharmacologically active substance selected from the group consisting of anticholinergic agents, beta-2 mimetics, steroids, PDE-IV inhibitors, p38 MAP kinase inhibitors, NK_1 antagonists, LTD4 antagonists, EGFR inhibitors and endothelin-antagonists.

9. A method for the prevention or treatment of a fibrotic disease selected from the group consisting of fibrosis and remodeling of lung tissue in chronic obstructive pulmonary disease, fibrosis and remodeling of lung tissue in chronic bronchitis, fibrosis and remodeling of lung tissue in emphysema, lung fibrosis and pulmonary diseases with a fibrotic component, fibrosis and remodeling in asthma, fibrosis in rheumatoid arthritis, virally induced hepatic cirrhosis, radiation-induced fibrosis, post angioplasty restenosis, chronic glomerulonephritis, renal fibrosis in patients receiving cyclosporine and renal fibrosis due to high blood pressure, diseases of the skin with a fibrotic component, and excessive scarring, wherein a substituted indolinone of formula I as defined in any one of claims 1 to 5 is administered to a patient in need thereof, optionally together with one or more pharmaceutically acceptable carriers or excipients.

10. The method in accordance with claim 9, wherein the disease is selected from the group consisting of the lung fibrosis and pulmonary diseases with a fibrotic component selected from idiopathic pulmonary fibrosis, giant cell interstitial pneumonia, sarcodosis, cystic fibrosis, respiratory distress syndrome, drug-induced lung fibrosis, granulomatosis, silicosis, asbestosis, systemic scleroderma, the virally induced hepatic cirrhosis selected from hepatitis C induced hepatic cirrhosis, and the diseases of the skin with a fibrotic component selected from scleroderma, sarcodosis and systemic lupus erythematosus.

11. The method in accordance with claim 10 wherein the disease is idiopathic pulmonary fibrosis.

12. The method in accordance with any one of claims 9 to 11, wherein the treatment is a combined treatment with a further pharmacologically active substance selected from the group consisting of anticholinergic agents, beta-2 mimetics, steroids, PDE-IV inhibitors, p38 MAP kinase inhibitors, NK_1 antagonists, LTD4 antagonists, EGFR inhibitors and endothelin-antagonists.said further pharmacologically active substance being administered either simultaneously or sequentially.

13. Pharmaceutical composition containing a substituted indolinone of formula I as defined in any one of claims 1 to 5 in combination with a further pharmacologically active substance selected from the group consisting of anticholinergic agents, beta-2 mimetics, PDE-IV inhibitors, p38 MAP kinase inhibitors, NK_1 antagonists, LTD4 antagonists and endothelin-antagonists, optionally together with one or more pharmaceutically acceptable carriers or excipients.