



1/11

*Fig. 1A**Fig. 1B*

2/11



Fig. 1C




Fig. 1D



Fig. 1E

3/11

siGFP si β gluc

Fig. 1F



Fig. 1G






Fig. 1H

4/11

*Fig. 1I**Fig. 1J*

5/11






Fig. 2A

siGFP/dsRED

siβgluc/dsRED

6/11

*Fig. 2B**Fig. 2C*

7/11




Fig. 3A





Fig. 3B

Fig. 3C

8/11

*Fig. 3D*

9/11




Fig. 4A




Fig. 4B




Fig. 4C

10/11




Fig. 4D






Fig. 4E

11/11

*Fig. 4F**Fig. 4G*

SEQUENCE LISTING

<110> University of Iowa Research Foundation

5 Davidson, B.

Xia, H.

Mao, Q.

<120> siRNA-Mediated Gene Silencing with Viral Vectors

10

<130> 875.078WO1

<150> US 10/430,351

<151> 2003-05-05

15

<150> US 10/322,086

<151> 2003-12-17

<150> US 10/212,322

20<151> 2002-08-05

<160> 6

<170> FastSEQ for Windows Version 4.0

25

<210> 1

<211> 40

<212> DNA

<213> Artificial Sequence

30

<220>

<223> A primer

<400> 1

35aagggtaccag atcttagtta ttaatagtaa tcaattacgg

40

<210> 2

<211> 43

<212> DNA

40<213> Artificial Sequence

<220>

<223> A primer

<400> 2

gaatcgatgc atgcctcgag acggttcact aaaccagctc tgc

43

5<210> 3

<211> 69

<212> DNA

<213> Artificial Sequence

10<220>

<223> An oligonucleotide, used with SEQ ID NO:4, to form a minimal polyA

<400> 3

ctagaactag taataaagga tcctttatctt tcattggatc cgtgtgttgg ttttttgtgt

60

15gcggccgcg

69

<210> 4

<211> 69

<212> DNA

20<213> Artificial Sequence

<220>

<223> An oligonucleotide, used with SEQ ID NO:3, to form a minimal polyA

25<400> 4

tcgacgcggc cgcacacaaa aaaccaacac acggatccaa tgaaaataaa ggatccttta

60

ttactagtt

69

<210> 5

30<211> 21

<212> DNA

<213> Artificial Sequence

<220>

35<223> A P32 labeled sense oligonucleotide used to probe a blot

<400> 5

cacaagctgg agtacaacta c

21

40<210> 6

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> A P32 labeled antisense oligonucleotide used to probe a blot

5

<400> 6

gtacttgtag tccagctttg tg

22