


3-substituted pyridine derivatives as H3 antagonists

The present disclosure is concerned with novel 3-substituted 6-piperazinyl-pyridine derivatives, their manufacture, pharmaceutical compositions containing them and their use as medicaments. The active compounds of the present invention are useful in treating obesity and other disorders.

In particular, the present disclosure relates to compounds of the general formula



I

wherein

R¹ is selected from the group consisting of hydrogen, lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl, C₃-C₇-cycloalkyl, and lower C₃-C₇-cycloalkylalkyl;

X is C(O) or SO₂;

m is 0 or 1;

R² is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl, unsubstituted C₃-C₇-cycloalkyl or C₃-C₇-cycloalkyl substituted by phenyl, lower C₃-C₇-cycloalkylalkyl, lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, unsubstituted pyridyl or pyridyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and -NR³R⁴, or, in case X is C(O), R² can also be lower alkoxy or lower alkoxyalkoxy,

- 2 -

or, in case m is 1, R² can also be unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl,

R³ is hydrogen or lower alkyl;

R⁴ is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl,
5 lower alkoxyalkyl,
C₃-C₇-cycloalkyl,
C₃-C₇-cycloalkyl substituted by phenyl,
lower C₃-C₇-cycloalkylalkyl,
unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower
10 alkoxy, halogen or lower halogenalkyl, and
lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by
lower alkyl, lower alkoxy, halogen or lower halogenalkyl; or

R³ and R⁴ together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or
7-membered heterocyclic ring optionally containing a further heteroatom selected
15 from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or
substituted by one, two or three groups independently selected from lower alkyl,
lower alkoxy, lower alkoxycarbonyl, oxo, halogen and halogenalkyl, or being
condensed with a C₅-C₆-cycloalkyl ring or a phenyl ring, said cycloalkyl ring or
phenyl ring being unsubstituted or substituted by one, two or three groups
20 independently selected from lower alkyl, lower alkoxy, halogen and halogenalkyl;

and pharmaceutically acceptable salts thereof;

with the exception of 2,2-dimethyl-N-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-propionamide.

2,2-Dimethyl-N-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-propionamide has
25 been described in EP 1 035 115 A1 (F. Hoffmann-La Roche AG) as intermediate for the
synthesis of 4-phenylpyridine derivatives useful as NK-1 receptor antagonists.

It has been found that the compounds of formula I are antagonists and/or inverse agonists at the histamine 3 receptor (H3 receptor).

Histamine (2-(4-imidazolyl) ethylamine) is one of the aminergic neurotransmitters
30 which is widely distributed throughout the body, e. g. the gastrointestinal tract (Burks
1994 in Johnson L.R. ed., Physiology of the Gastrointestinal Tract, Raven Press, NY, pp.
211 – 242). Histamine regulates a variety of digestive pathophysiological events like
gastric acid secretion, intestinal motility (Leurs et al., Br J. Pharmacol. 1991, 102, pp 179-

185), vasomotor responses, intestinal inflammatory responses and allergic reactions (Raithel et al., *Int. Arch. Allergy Immunol.* 1995, 108, 127-133). In the mammalian brain, histamine is synthesized in histaminergic cell bodies which are found centrally in the tuberomammillary nucleus of the posterior basal hypothalamus. From there, the cell
5 bodies project to various brain regions (Panula et al., *Proc. Natl. Acad. Sci. USA* 1984, 81, 2572-2576; Inagaki et al., *J. Comp. Neurol* 1988, 273, 283 – 300).

According to current knowledge, histamine mediates all its actions in both the CNS and in the periphery through four distinct histamine receptors, the histamine H1, H2 H3 and H4 receptors.

10 H3 receptors are predominantly localized in the CNS. As an autoreceptor, H3 receptors constitutively inhibit the synthesis and secretion of histamine from histaminergic neurons (Arrang et al., *Nature* 1983, 302, 832-837; Arrang et al., *Neuroscience* 1987, 23, 149-157). As heteroreceptors, H3 receptors also modulate the release of other neurotransmitters such as acetylcholine, dopamine, serotonin and
15 norepinephrine among others in both the central nervous system and in peripheral organs, such as lungs, cardiovascular system and gastrointestinal tract (Clapham & Kilpatrick, *Br. J. Pharmacol.* 1982, 107, 919- 923; Blandina et al. in *The Histamine H3 Receptor* (Leurs RL and Timmermann H eds, 1998, pp 27-40, Elsevier, Amsterdam, The Netherlands). H3 receptors are constitutively active, meaning that even without
20 exogenous histamine, the receptor is tonically activated. In the case of an inhibitory receptor such as the H3 receptor, this inherent activity causes tonic inhibition of neurotransmitter release. Therefore it may be important that a H3R antagonist would also have inverse agonist activity to both block exogenous histamine effects and to shift the receptor from its constitutively active (inhibitory) form to a neutral state.


25 The wide distribution of H3 receptors in the mammalian CNS indicates the physiological role of this receptor. Therefore the therapeutic potential as a novel drug development target in various indications has been proposed.

The administration of H3R ligands - as antagonists, inverse agonists, agonists or partial agonists - may influence the histamine levels or the secretion of neurotransmitters
30 in the brain and the periphery and thus may be useful in the treatment of several disorders. Such disorders include obesity, (Masaki et al; *Endocrinol.* 2003, 144, 2741-2748; Hancock et al., *European J. of Pharmacol.* 2004, 487, 183-197), cardiovascular disorders such as acute myocardial infarction, dementia and cognitive disorders such as attention deficit hyperactivity disorder (ADHD) and Alzheimer's disease, neurological
35 disorders such as schizophrenia, depression, epilepsy, Parkinson's disease, and seizures or convulsions, sleep disorders, narcolepsy, pain, gastrointestinal disorders, vestibular

dysfunction such as Morbus Meniere, drug abuse and motion sickness (Timmermann, J. Med. Chem. 1990, 33, 4-11).

It is therefore an aim of the present disclosure to provide selective, directly acting H3 receptor antagonists respectively inverse agonists. Such antagonists / inverse agonists are useful as therapeutically active substances, particularly in the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.

In a first aspect, there is provided compounds of the general formula



I

wherein

R¹ is lower alkyl or C₃-C₇-cycloalkyl;

X is C(O) or SO₂;

m is 0 or 1;

R² is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl, unsubstituted C₃-C₇-cycloalkyl or C₃-C₇-cycloalkyl substituted by phenyl, lower C₃-C₇-cycloalkylalkyl, lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, unsubstituted pyridyl or pyridyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and -NR³R⁴,

or, in case X is C(O), R² can also be lower alkoxy or lower alkoxyalkoxy,

or, in case m is 1, R² can also be unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl,

R³ is hydrogen or lower alkyl;

R⁴ is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, lower alkoxyalkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl substituted by phenyl, lower C₃-C₇-cycloalkylalkyl,

unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl; or


R³ and R⁴ together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, lower alkoxycarbonyl, oxo, halogen and halogenalkyl, or being condensed with a C₅-C₆-cycloalkyl ring or a phenyl ring, said cycloalkyl ring or phenyl ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, halogen and halogenalkyl;

and pharmaceutically acceptable salts thereof;

with the exception of 2,2-dimethyl-N-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-propionamide.

In a second aspect, there is provided a process for the manufacture of compounds as defined above, which process comprises


a) reacting a compound of the formula II



II


wherein R^1 and m are as defined above,

with a sulfonylchloride or sulfamoylchloride of the formula III



wherein R^2 is as defined above,


to obtain a compound of the formula I-A



I-A

wherein R^1 , R^2 and m are as defined above, or

b) reacting a compound of the formula II



II


wherein R^1 and m are as defined above,

with a chloride of the formula IV



wherein R^2 is as defined above,


to obtain a compound of the formula I-B



I-B


wherein R^1 , R^2 and m are as defined above, or

c) reacting a compound of the formula II




II

wherein R^1 and m are as defined above,
with an isocyanate of the formula V



wherein R^4 is as defined above,
to obtain a compound of the formula I-C



I-C

wherein R^3 is hydrogen and R^1 , R^4 and m are as defined above,

and if desired, converting the compound of formula I-A, I-B or I-C into a
pharmaceutically acceptable salt.

In a third aspect, there is provided compounds as defined above when manufactured
by a process as defined above.

In a fourth aspect, there is provided pharmaceutical compositions comprising a
compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

In a fifth aspect, there is provided compounds as defined above for use as
therapeutically active substances.

In a sixth aspect, there is provided compounds as defined above for use as therapeutically active substances for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.

In a seventh aspect, there is provided the use of compounds as defined for the preparation of medicaments for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.

In the present description the term "alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of one to twenty carbon atoms, preferably one to sixteen carbon atoms, more preferably one to ten carbon atoms.

The term "lower alkyl" or "C₁-C₈-alkyl", alone or in combination, signifies a straight-chain or branched-chain alkyl group with 1 to 8 carbon atoms, preferably a straight or branched-chain alkyl group with 1 to 6 carbon atoms and particularly preferred a straight or branched-chain alkyl group with 1 to 4 carbon atoms. Examples of straight-chain and branched C₁-C₈ alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert.-butyl, the isomeric pentyls, the isomeric hexyls, the isomeric heptyls and the isomeric octyls, preferably methyl, ethyl and isopropyl, and most preferred methyl and ethyl.

The term "lower alkenyl" or "C₃-C₈-alkenyl", alone or in combination, signifies a straight-chain or branched alkyl group comprising an olefinic bond and up to 8, preferably up to 6, particularly preferred up to 4 carbon atoms. Examples of alkenyl groups are 1-propenyl, 2-propenyl, isopropenyl, 1-butenyl, 2-butenyl, 3-butenyl and isobutenyl. A preferred example is 2-propenyl.

The term "lower alkynyl" or "C₃-C₈-alkynyl", alone or in combination, signifies a straight-chain or branched alkyl group comprising a triple bond and up to 8, preferably up to 6, particularly preferred up to 4 carbon atoms. Examples of alkynyl groups include 2-propynyl (propargyl), 1-methyl-2-propynyl, 2-butylnyl, 3-butylnyl, 2-pentylnyl and 1-pentyln-3-yl.

The term "alkoxy" refers to the group R'-O-, wherein R' is alkyl. The term "lower alkoxy" refers to the group R'-O-, wherein R' is lower alkyl and the term "lower alkyl" has the previously given significance ("C₁-C₈-alkoxy"). Examples of lower alkoxy groups are e.g. methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec. butoxy and tert.butoxy, preferably methoxy and ethoxy and most preferred methoxy.

The term "lower alkoxyalkyl" or "C₁-C₈-alkoxy-C₁-C₈-alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl group is replaced by an alkoxy group as defined above. Among the preferred lower alkoxyalkyl groups are methoxymethyl, methoxyethyl and ethoxymethyl, with
5 methoxymethyl being especially preferred.

The term "lower alkoxyalkoxy" or "C₁-C₈-alkoxy-C₁-C₈-alkoxy" refers to lower alkoxy groups as defined above wherein at least one of the hydrogen atoms of the lower alkoxy group is replaced by an alkoxy group as defined above. Among the preferred lower alkoxyalkoxy groups are methoxyethoxy, methoxypropyloxy and ethoxyethoxy,
10 with methoxyethoxy being especially preferred.

The term "halogen" refers to fluorine, chlorine, bromine and iodine, with fluorine, chlorine and bromine being preferred.

The term "lower halogenalkyl" or "halogen-C₁-C₈-alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl
15 group is replaced by a halogen atom, preferably fluoro or chloro, most preferably fluoro. Among the preferred halogenated lower alkyl groups are trifluoromethyl, difluoromethyl, fluoromethyl and chloromethyl, with trifluoromethyl being especially preferred.

The term "lower hydroxyalkyl" or "hydroxy-C₁-C₈-alkyl" refers to lower alkyl
20 groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl group is replaced by a hydroxy group. Examples of lower hydroxyalkyl groups are hydroxymethyl or hydroxyethyl.

The term "cycloalkyl" or "C₃-C₇-cycloalkyl" means a cycloalkyl ring containing 3 to 7 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.
25 The cycloalkyl ring may be substituted as defined herein. Especially preferred is cyclopropyl or cyclopentyl.

The term "lower cycloalkylalkyl" or "C₃-C₇-cycloalkyl-C₁-C₈-alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl group is replaced by a cycloalkyl group as defined above. Examples of preferred
30 lower cycloalkylalkyl groups are cyclopropylmethyl or cyclopentylmethyl.

The term "lower phenylalkyl" or "phenyl-C₁-C₈-alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl group is replaced by a phenyl group. The phenyl ring may be substituted as defined herein. Examples of preferred lower phenylalkyl groups are benzyl, 4-methylbenzyl, 4-

fluorobenzyl, 3-methoxybenzyl and 3,4-dimethoxybenzyl.

The term "form a 4-, 5-, 6- or 7-membered saturated heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur" refers to a saturated N-heterocyclic ring, which may optionally contain a further nitrogen, oxygen or sulfur atom, such as azetidiny, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, or azepanyl. The heterocyclic ring may be unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, oxo, halogen and halogenalkyl. The heterocyclic ring may also be condensed with a C₅-C₆-cycloalkyl ring or a phenyl ring, said cycloalkyl ring or phenyl ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, halogen and halogenalkyl. Examples for such a condensed heterocyclic rings are 3,4-dihydro-1H-isoquinoline, 1,3-dihydro-isoindole and octahydroquinoline.


The term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. The salts are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, preferably hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, salicylic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, N-acetylcystein and the like. In addition these salts may be prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium salts and the like. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, lysine, arginine, N-ethylpiperidine, piperidine, polymine resins and the like. The compound of formula I can also be present in the form of zwitterions. Particularly preferred pharmaceutically acceptable salts of compounds of formula I are the hydrochloride salts.

The compounds of formula I can also be solvated, e.g. hydrated. The solvation can be effected in the course of the manufacturing process or can take place e.g. as a consequence of hygroscopic properties of an initially anhydrous compound of formula I

(hydration). The term pharmaceutically acceptable salts also includes physiologically acceptable solvates.

“Isomers” are compounds that have identical molecular formulae but that differ in the nature or the sequence of bonding of their atoms or in the arrangement of their
 5 atoms in space. Isomers that differ in the arrangement of their atoms in space are termed “stereoisomers”. Stereoisomers that are not mirror images of one another are termed “diastereoisomers”, and stereoisomers that are non-superimposable mirror images are termed “enantiomers”, or sometimes optical isomers. A carbon atom bonded to four nonidentical substituents is termed a “chiral center”.

10 In detail, the present invention relates to compounds of the general formula



I

wherein

R^1 is selected from the group consisting of hydrogen, lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl, C₃-C₇-cycloalkyl, and lower C₃-C₇-cycloalkylalkyl;
 15

X is C(O) or SO₂;

m is 0 or 1;

R^2 is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl, unsubstituted C₃-C₇-cycloalkyl or C₃-C₇-cycloalkyl substituted by phenyl, lower C₃-C₇-cycloalkylalkyl, lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, unsubstituted pyridyl or pyridyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and
 20 -NR³R⁴,
 25 or, in case X is C(O), R^2 can also be lower alkoxy or lower alkoxyalkoxy, or, in case m is 1, R^2 can also be unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl,

- R³ is hydrogen or lower alkyl;
- R⁴ is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, lower alkoxyalkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl substituted by phenyl, lower C₃-C₇-cycloalkylalkyl, unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl; or
- R³ and R⁴ together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, lower alkoxycarbonyl, oxo, halogen and halogenalkyl, or being condensed with a C₅-C₆-cycloalkyl ring or a phenyl ring, said cycloalkyl ring or phenyl ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, halogen and halogenalkyl; and pharmaceutically acceptable salts thereof;
- with the exception of 2,2-dimethyl-N-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-propionamide.

Thus, the substituent R¹ is selected from the group consisting of hydrogen, lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl, C₃-C₇-cycloalkyl and lower C₃-C₇-cycloalkylalkyl. Preferred compounds of formula I according to the present invention are those, wherein R¹ is lower alkyl or C₃-C₇-cycloalkyl, with those compounds, wherein R¹ is C₃-C₇-cycloalkyl being more preferred, and those compounds, wherein R¹ is cyclopentyl, being most preferred. Compounds of formula I, wherein R¹ is ethyl or isopropyl, are also very preferred.

m is an integer of 0 or 1. Especially preferred are those compounds of formula I, wherein m is 1. However, compounds of formula I, wherein m is 0, are also a preferred embodiment of the invention.

Further preferred are compounds of formula I according to the present invention, wherein R² is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl,

lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl,
 unsubstituted C₃-C₇-cycloalkyl or C₃-C₇-cycloalkyl substituted by phenyl,
 lower C₃-C₇-cycloalkylalkyl,
 lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower
 5 alkyl, lower alkoxy, halogen or lower halogenalkyl,
 unsubstituted pyridyl or pyridyl mono- or disubstituted by lower alkyl, lower alkoxy,
 halogen or lower halogenalkyl, and
 -NR³R⁴,
 or, in case X is C(O), R² can also be lower alkoxy or lower alkoxyalkoxy.

10 Within this group, compounds of formula I, wherein R² is selected from the group
 consisting of lower alkyl, C₃-C₇-cycloalkyl and C₃-C₇-cycloalkyl substituted by phenyl,
 are preferred.

Furthermore, compounds of formula I are preferred, wherein R² is lower
 phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl,
 15 lower alkoxy, halogen or lower halogenalkyl.

Also preferred are compounds of formula I of the present invention, wherein R² is
 the group -NR³R⁴.

More preferred are those compounds of formula I, wherein R² is the group -NR³R⁴
 and R³ is hydrogen or lower alkyl;

20 R⁴ is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl,
 C₃-C₇-cycloalkyl,
 C₃-C₇-cycloalkyl substituted by phenyl,
 lower C₃-C₇-cycloalkylalkyl,
 unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower
 25 alkoxy, halogen or lower halogenalkyl, and
 lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by
 lower alkyl, lower alkoxy, halogen or lower halogenalkyl; or

R³ and R⁴ together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or
 7-membered heterocyclic ring optionally containing a further heteroatom selected
 30 from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or
 substituted by one, two or three groups independently selected from lower alkyl,
 lower alkoxy, oxo, halogen and halogenalkyl, or being condensed with a phenyl
 ring, said phenyl ring being unsubstituted or substituted by one, two or three
 groups independently selected from lower alkyl, lower alkoxy, halogen and
 35 halogenalkyl.

Especially preferred are those compounds of formula I, wherein R^3 and R^4 are lower alkyl.


Another group of preferred compounds of formula I are those, wherein R^2 is the group $-NR^3R^4$, R^3 is hydrogen and R^4 is selected from the group consisting of

- 5 lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl,
C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl substituted by phenyl,
lower C₃-C₇-cycloalkylalkyl,
unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy,
halogen or lower halogenalkyl, and
- 10 lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl.

Those compounds, wherein R^3 is hydrogen and R^4 is lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, are especially preferred.

- 15 Furthermore, compounds of formula I of the present invention are preferred, wherein m is 1 and R^2 is unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl.

A preferred group of compounds of formula I are those, wherein X is SO₂. These are the compounds having the formula




I-A

20

wherein R^1 , R^2 and m are as defined hereinbefore.

Furthermore, compounds of formula I are preferred, wherein X is C(O). These are compounds having the formula


- 11 -



I-B

wherein R^1 , R^2 and m are as defined hereinbefore.

Within this group, those compounds are especially preferred wherein X is $C(O)$ and R^2 is $-NR^3R^4$, thus meaning compounds of the formula



I-C

5

wherein R^1 , R^3 , R^4 and m are as defined hereinbefore.

Examples of preferred compounds of formula I are the following:

- 1-[6-(4-ethylpiperazin-1-yl)-pyridin-3-yl]-3-propyl-urea,
- 1-cyclohexyl-3-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 10 1-benzyl-3-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-p-tolyl-urea,
- 1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea,
- 1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea,
- 1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-fluoro-phenyl)-urea,
- 15 1-cyclohexyl-3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-benzyl-3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea,

- 1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea,
- 1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-fluoro-phenyl)-urea,
- 1-cyclohexyl-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 5 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-phenyl-urea,
- 1-benzyl-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-o-tolyl-urea,
- 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-m-tolyl-urea,
- 1-(2-chloro-phenyl)-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 10 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea,
- 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea,
- 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea,
- 1-(4-fluoro-phenyl)-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-propyl-urea,
- 15 1-cyclohexyl-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-benzyl-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-m-tolyl-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-p-tolyl-urea,
- 1-(2-chloro-phenyl)-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 20 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea,
- 1-(4-fluoro-phenyl)-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-dimethylaminosulfonamide,

- N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-dimethylaminosulfonamide,
N-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-dimethylaminosulfonamide,
cyclohexanecarboxylic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-amide,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-phenyl-acetamide,
5 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-(4-fluoro-phenyl)-acetamide,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-(3-methoxy-phenyl)-acetamide,
2-(3,4-dimethoxy-phenyl)-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-acetamide,
[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid 2-methoxy-ethyl ester,
[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester,
10 cyclopropanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-butyramide,
cyclobutanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
cyclopentanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-ethyl-butyramide,
15 cyclohexanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
2-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-nicotinamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-phenyl-acetamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(4-fluoro-phenyl)-acetamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(3-methoxy-phenyl)-acetamide,
20 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(3,4-dimethoxy-phenyl)-
acetamide,
[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid ethyl ester,
[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid 2-methoxy-ethyl ester,
[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester,

- cyclohexanecarboxylic acid [6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-amide,
2-(3,4-dimethoxy-phenyl)-N-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-acetamide,
[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid ethyl ester,
[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester,
- 5 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-butyramide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-methoxy-benzamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-phenyl-acetamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-(4-fluoro-phenyl)-
acetamide,
- 10 ethanesulfonic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide,
propane-1-sulfonic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-dimethylaminosulfonamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-C-phenyl-
- 15 methanesulfonamide,
C-(4-chloro-phenyl)-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-
methanesulfonamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-fluoro-benzenesulfonamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-fluoro-benzenesulfonamide,
- 20 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-4-fluoro-benzenesulfonamide,
2-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
4-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-(4-fluoro-benzyl)-urea,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-methoxy-benzamide,
- 25 N-[6-(4-Ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-phenyl-acetamide,

- N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-(4-fluoro-phenyl)-acetamide,
ethanesulfonic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide,
propane-1-sulfonic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-dimethylaminosulfonamide,
5 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-C-phenyl-methanesulfonamide,
C-(4-chloro-phenyl)-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-
methanesulfonamide,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-fluoro-benzenesulfonamide,
10 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-fluoro-benzenesulfonamide,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-4-fluoro-benzenesulfonamide,
2-chloro-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
4-chloro-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
4-methyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-
15 amide,
2,6-dimethyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-
yl]-amide,
4-trifluoromethyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-
pyridin-3-yl]-amide,
20 1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylcarbamoyl]-piperidine-4-carboxylic
acid ethyl ester,
octahydro-quinoline-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-
amide,
octahydro-isoquinoline-2-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-
25 yl]-amide,
2-trifluoromethyl-pyrrolidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-
pyridin-3-yl]-amide,

- 2-isopropyl-pyrrolidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- 1,3-dihydro-isoindole-2-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- 5 3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-isopropyl-1-(2-methoxy-ethyl)-urea,
- azepane-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- 3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-ethyl-1-phenyl-urea,
- 3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-(4-methoxy-phenyl)-1-methyl-urea,
- 10 3,4-dihydro-2H-quinoline-1-carboxylic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- 3,4-dihydro-2H-quinoline-1-carboxylic acid [6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- and pharmaceutically acceptable salts thereof.
- 15 Particularly preferred compounds of formula I of the present invention are the following:
- 1-benzyl-3-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-benzyl-3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea,
- 20 1-benzyl-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-dimethylaminosulfonamide,
- N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(3,4-dimethoxy-phenyl)-acetamide,
- N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-methoxy-benzamide,
- 25 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-phenyl-acetamide,
- N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-dimethylaminosulfonamide,

N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-C-phenyl-methanesulfonamide,

C-(4-chloro-phenyl)-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-methanesulfonamide,

- 5 2,6-dimethyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,

and pharmaceutically acceptable salts thereof.

Furthermore, the pharmaceutically acceptable salts of the compounds of formula I and the pharmaceutically acceptable esters of the compounds of formula I individually
10 constitute preferred embodiments of the present invention.

Compounds of formula I may form acid addition salts with acids, such as conventional pharmaceutically acceptable acids, for example hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, salicylate, sulphate, pyruvate, citrate, lactate, mandelate, tartrate, and methanesulphonate. Preferred are the
15 hydrochloride salts. Also solvates and hydrates of compounds of formula I and their salts form part of the present invention.

Furthermore, the N-atom of the pyridine ring can be present as an N-oxide group. Such N-oxides of compounds of formula I also form part of the present invention.


Compounds of formula I can have one or more asymmetric carbon atoms and can
20 exist in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates. The optically active forms can be obtained for example by resolution of the racemates, by asymmetric synthesis or asymmetric chromatography (chromatography with a chiral adsorbent or
25 eluant). The invention embraces all of these forms.

It will be appreciated, that the compounds of general formula I in this invention may be derivatised at functional groups to provide derivatives which are capable of conversion back to the parent compound in vivo. Physiologically acceptable and metabolically labile derivatives, which are capable of producing the parent compounds of
30 general formula I in vivo are also within the scope of this invention.

- 18 -

A further aspect of the present invention is the process for the manufacture of compounds of formula I as defined above, which process comprises

a) reacting a compound of the formula II




II

5 wherein R^1 and m are as defined herein before,
with a sulfonylchloride or sulfamoylchloride of the formula III



wherein R^2 is as defined herein before,
to obtain a compound of the formula I-B




I - A

10

wherein R^1 , R^2 and m are as defined herein before, or


b) reacting a compound of the formula II



II


15

wherein R^1 and m are as defined herein before,
with a chloride of the formula IV



wherein R^2 is as defined herein before,
to obtain a compound of the formula I-A


- 19 -



I - B


wherein R^1 , R^2 and m are as defined herein before, or

c) reacting a compound of the formula II




II

5 wherein R^1 and m are as defined herein before,
with an isocyanate of the formula V



wherein R^4 is as defined herein before,
to obtain a compound of the formula I-C



I - C

10

wherein R^3 is hydrogen and R^1 , R^4 and m are as defined herein before,
and if desired, converting the compound of formula I-A, I-B or I-C into a
pharmaceutically acceptable salt.

In more detail, the compounds of formula I can be manufactured by the methods
15 given below, by the methods given in the examples or by analogous methods.
Appropriate reaction conditions for the individual reaction steps are known to a person
skilled in the art. Starting materials are either commercially available or can be prepared

by methods analogous to the methods given below, by methods described in references cited in the text or in the examples, or by methods known in the art.


The preparation of compounds of formula I of the present invention may be carried out in sequential or convergent synthetic routes. Syntheses of the invention are shown in the following scheme. The skills required for carrying out the reaction and purification of the resulting products are known to those in the art. The substituents and indices used in the following description of the processes have the significance given herein before unless indicated to the contrary.

Compounds of general formula I can be prepared according to scheme 1 as follows:

- a) The coupling of chloro substituted pyridine derivatives with piperazines is widely described in literature and the procedures are known to those in the art (For reaction conditions described in literature affecting such reactions see for example: Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2nd Edition, Richard C. Larock. John Wiley & Sons, New York, NY. 1999). 2-Chloro-5-cyanopyridine (VI) or 2-chloro-5-nitropyridine (IX) can conveniently be transformed to the respective pyridine derivative VIII or X through reaction with a piperazine derivative VII (either commercially available or accessible by methods described in references or by methods known in the art; as appropriate). The reaction can be carried out in the presence or the absence of a solvent and in the presence or the absence of a base. We find it convenient to carry out the reaction in a solvent like water and/or dimethylformamide (DMF) and in the presence of a base like diisopropyl-ethylamine (DIPEA). There is no particular restriction on the nature of the solvent to be employed, provided that it has no adverse effect on the reaction or the reagents involved and that it can dissolve the reagents, at least to some extent. Examples for suitable solvents include DMF, dichloromethane (DCM), dioxane, tetrahydrofuran (THF), and the like. There is no particular restriction on the nature of the base used in this stage, and any base commonly used in this type of reaction may equally be employed here. Examples of such bases include triethylamine and diisopropylethylamine, and the like. The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention. We find it convenient to carry out the reaction with heating from ambient temperature to reflux. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents. A period of from 0.5 h to several days will usually suffice to yield pyridine derivatives VIII or X.

- 21 -

Scheme 1



However, in the cases where the desired piperazine derivative is not easily accessible, an alternative route can be pursued leading to the pyridine derivatives VIII or X.

5 Piperazine (VII; R¹ = H) is reacted with 2-chloro-5-cyanopyridine (VI) or 2-chloro-5-nitropyridine (IX) to access the respective pyridine derivative VIII or X (R¹ = H). The reaction can be carried out in the presence or the absence of a solvent and in the

- presence or the absence of a base. We find it convenient to carry out the reaction in a solvent like water and/or dimethylformamide (DMF) and in the presence of a base like triethylamine. There is no particular restriction on the nature of the solvent to be employed, provided that it has no adverse effect on the reaction or the reagents involved and that it can dissolve the reagents, at least to some extent. Examples for suitable solvents include: DMF, dichloromethane (DCM), dioxane, THF, and the like. There is no particular restriction on the nature of the base used in this stage, and any base commonly used in this type of reaction may equally be employed here. Examples of such bases include triethylamine and diisopropylethylamine, and the like. The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention. We find it convenient to carry out the reaction with heating from ambient temperature to reflux. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents. However, a period of from 0.5 h to several days will usually suffice to yield pyridine derivatives VIII or X. Subsequently, to access the pyridine derivatives VIII or X ($R^1 \neq H$) the intermediate is subjected to reductive amination reaction conditions with suitable aldehydes or alkylating reaction conditions with suitable alkylating reagents. The reaction conditions for either reaction are widely described in literature and the procedures are known to those in the art (For reaction conditions described in literature affecting such reactions see for example: Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2nd Edition, Richard C. Larock. John Wiley & Sons, New York, NY. 1999).
- b) The reduction of the cyano or nitro functionality in VIII or X respectively can be achieved under various reducing reaction conditions to access the aminomethyl pyridine derivatives II-A (compounds of formula II wherein m is 1) or the aminopyridine derivatives II-B (compounds of formula II wherein m is 0). The reaction conditions for either reaction are widely described in literature and the procedures are known to those in the art (For reaction conditions described in literature affecting such reactions see for example: Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2nd Edition, Richard C. Larock. John Wiley & Sons, New York, NY. 1999). We find it convenient to hydrogenate VIII or X over Raney Nickel or palladium/charcoal (Pd/C) in a solvent and in the presence or absence of an acid. We find it convenient to carry out the reaction in a solvent like methanol or ethyl acetate. There is no particular restriction on the nature of the solvent to be employed, provided that it has no adverse effect on the reaction or the reagents involved and that it can dissolve the reagents, at least to some extent. Examples for suitable solvents include: methanol, ethanol, ethyl acetate,

and the like. There is no particular restriction on the nature of the acid used in this stage, and any base commonly used in this type of reaction may equally be employed here. Examples of such acids include acetic acid or HCl, and the like. The reduction can be achieved through hydrogen, however any other reducing agents employed in such reaction might equally be employed here. Neither the precise hydrogen pressure nor the precise reaction temperature are critical to the invention. The reaction can take place over a wide range of temperatures, and a wide range of hydrogen pressure. We find it convenient to carry out the reaction with heating from ambient temperature to reflux. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents. However, a period of from 0.5 h to several days will usually suffice to yield pyridine derivatives II-A (m is 1).

c) Sulfonamides, amides, carbamates and ureas can be prepared from suitable starting materials according to methods known in the art. The conversion of the amino-moiety in II-A or II-B to access sulfonamides, amides, carbamates and ureas can be affected by methods described in literature. For example the conversion of the amine derivatives II to access compounds of the general formula I is affected by reaction of II with suitable with sulfonyl chlorides or sulfamoyl chlorides (compounds of formula III) or acid chlorides, chloroformates, or carbonate esters (compounds of formula IV as defined herein before) or with isocyanates (compounds of formula V), respectively, in a solvent like dichloromethane and in the presence or the absence of a base. The compounds of formula III, IV or V are known or can be prepared by known methods. There is no particular restriction on the nature of the solvent to be employed, provided that it has no adverse effect on the reaction or the reagents involved and that it can dissolve the reagents, at least to some extent. Examples for suitable solvents include: chloroform, or dioxane, THF, and the like. There is no particular restriction on the nature of the base used in this stage, and any base commonly used in this type of reaction may equally be employed here. Examples of such bases include triethylamine and diisopropylethyl-amine, and the like. The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention. We find it convenient to carry out the reaction with heating from ambient temperature to reflux. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents. However, a period of from 0.5 h to several days will usually suffice to yield pyridine derivatives I. For reaction conditions described in literature affecting such reactions see for example: Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2nd Edition, Richard C. Larock. John Wiley & Sons, New York, NY. 1999.

As described above, the compounds of formula I of the present invention can be used as medicaments for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders. The use as medicament for the treatment and/or prevention of obesity is preferred.

The invention therefore also relates to pharmaceutical compositions comprising a compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

Further, the invention relates to compounds as defined above for use as therapeutically active substances, particularly as therapeutic active substances for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders.

In another embodiment, the invention relates to a method for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders. A method for the treatment and/or prevention of obesity is preferred.

The invention further relates to the use of compounds of formula I as defined above for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders. The use of compounds of formula I as defined above for the treatment and/or prevention of obesity is preferred.

In addition, the invention relates to the use of compounds of formula I as defined above for the preparation of medicaments for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders. The use of compounds of formula I as defined above for the preparation of medicaments for the treatment and/or prevention of obesity is preferred.

The compounds of formula I and their pharmaceutically acceptable salts possess valuable pharmacological properties. Specifically, it has been found that the compounds of the present invention are good histamine 3 receptor (H3R) antagonists and/or inverse agonists.

The following test was carried out in order to determine the activity of the compounds of formula (I).

30 Binding assay with ^3H -(R) α -methylhistamine

Saturation binding experiments were performed using HR3-CHO membranes prepared as described in Takahashi, K, Tokita, S., Kotani, H. (2003) J. Pharmacol. Exp. Therapeutics 307, 213-218.

An appropriate amount of membrane (60 to 80 μg protein/well) was incubated with increasing concentrations of ^3H -(R) α -Methylhistamine di-hydrochloride (0.10 to 10

nM). Non specific binding was determined using a 200 fold excess of cold (R) α -Methylhistamine dihydrobromide (500 nM final concentration). The incubation was carried out at room temperature (in deep-well plates shaking for three hours). The final volume in each well was 250 μ l. The incubation was followed by rapid filtration on GF/B
5 filters (pre-soaked with 100 μ l of 0.5% PEI in Tris 50 mM shaking at 200 rpm for two hours). The filtration was made using a cell-harvester and the filter plates were then washed five times with ice cold washing buffer containing 0.5 M NaCl. After harvesting, the plates were dried at 55 °C for 60min, then we added scintillation fluid (Microscint 40, 40 microl in each well) and the amount of radioactivity on the filter was determined in
10 Packard top-counter after shaking the plates for two hours at 200 rpm at room temperature.

Binding Buffer: 50 mM Tris-HCl pH 7.4 and 5 mM MgCl₂·6H₂O pH 7.4. Washing Buffer: 50 mM Tris-HCl pH 7.4 and 5 mM MgCl₂·6H₂O and 0.5 M NaCl pH 7.4.

Indirect measurement of affinity of H₃R inverse agonists: twelve increasing
15 concentrations (ranging from 10 μ M to 0.3 nM) of the selected compounds were always tested in competition binding experiments using membrane of the human HR3-CHO cell line. An appropriate amount of protein, e.g. approximately 500cpm binding of RAMH at K_d, were incubated for 1 hour at room temperature in 250 μ l final volume in 96-well plates in presence of ³H(R) α -Methylhistamine (1 nM final concentration = K_d).
20 Non-specific binding was determined using a 200 fold excess of cold (R) α -Methylhistamine dihydrobromide.

All compounds were tested at a single concentration in duplicates. Compounds that showed an inhibition of [³H]-RAMH by more than 50% were tested again to determine IC₅₀ in a serial dilution experiment. K_i's were calculated from IC₅₀ based on Cheng-
25 Prusoff equation (Cheng, Y, Prusoff, WH (1973) Biochem Pharmacol 22, 3099-3108).

The compounds of the present invention exhibit K_i values within the range of about 1 nM to about 1000 nM, preferably of about 1 nM to about 100 nM, and more preferably of about 1 nM to about 30 nM. The following table shows measured values for some selected compounds of the present invention.

30 The following table shows measured values for some selected compounds of the present invention.

- 27 -

	K _i (nM)
Example 3	39
Example 34	81
Example 53	40

The compounds of formula (I) and their pharmaceutically acceptable salts and esters can be used as medicaments, e.g. in the form of pharmaceutical preparations for enteral, parenteral or topical administration. They can be administered, for example, 5 perorally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions, rectally, e.g. in the form of suppositories, parenterally, e.g. in the form of injection solutions or infusion solutions, or topically, e.g. in the form of ointments, creams or oils.

The production of the pharmaceutical preparations can be effected in a manner 10 which will be familiar to any person skilled in the art by bringing the described compounds of formula (I) and their pharmaceutically acceptable, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

Suitable carrier materials are not only inorganic carrier materials, but also organic 15 carrier materials. Thus, for example, lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used as carrier materials for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carrier materials for soft gelatine capsules are, for example, vegetable oils, waxes, fats and semi-solid and liquid polyols (depending on the nature of the active ingredient no carriers are, however, required in the case of soft 20 gelatine capsules). Suitable carrier materials for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar and the like. Suitable carrier materials for injection solutions are, for example, water, alcohols, polyols, glycerol and vegetable oils. Suitable carrier materials for suppositories are, for example, natural or hardened oils, waxes, fats and semi-liquid or liquid polyols. Suitable carrier materials for 25 topical preparations are glycerides, semi-synthetic and synthetic glycerides, hydrogenated oils, liquid waxes, liquid paraffins, liquid fatty alcohols, sterols, polyethylene glycols and cellulose derivatives.

Usual stabilizers, preservatives, wetting and emulsifying agents, consistency-improving agents, flavour-improving agents, salts for varying the osmotic pressure, buffer substances, solubilizers, colorants and masking agents and antioxidants come into consideration as pharmaceutical adjuvants.

- 5 The dosage of the compounds of formula (I) can vary within wide limits depending on the disease to be controlled, the age and the individual condition of the patient and the mode of administration, and will, of course, be fitted to the individual requirements in each particular case. For adult patients a daily dosage of about 1 mg to about 1000 mg, especially about 1 mg to about 100 mg, comes into consideration.
- 10 Depending on the dosage it is convenient to administer the daily dosage in several dosage units.

The pharmaceutical preparations conveniently contain about 0.1-500 mg, preferably 0.5-100 mg, of a compound of formula (I).

The following examples serve to illustrate the present invention in more detail.

- 15 They are, however, not intended to limit its scope in any manner.

Examples

Intermediate 1

1-Ethyl-4-(5-nitro-pyridin-2-yl)-piperazine

- 20 A mixture of 2 g (13 mmol) 2-chloro-5-nitro-pyridine and 0.76 g (6 mmol) N,N-diisopropylethylamine in 21 ml water and 4 ml DMF was heated to 80 °C. During 2 min 1.73 g (15 mmol) N-ethylpiperazine was added and the mixture was kept for an additional hour at 80 °C. The yellow precipitate was filtered off and washed three times with 4 ml water and dried for 16 h under vacuum to yield 2.48 g (83 %) of the title compound as yellow crystals. (m/e): 237.1 (MH⁺; 100%).

25 Intermediate 2

6-(4-Ethyl-piperazin-1-yl)-pyridin-3-ylamine

- 30 A mixture of 2.47 g (10 mmol) 1-ethyl-4-(5-nitro-pyridin-2-yl)-piperazine and 0.247 g Pd/C (10 %) in 25 ml methanol was treated with 1 bar hydrogen at room temperature for 2 h. After filtration the mixture was evaporated to dryness to yield 2.12 g (98 %) of the title compound as colourless solid. (m/e): 207.3 (MH⁺; 100%).

Intermediate 3

1-Isopropyl-4-(5-nitro-pyridin-2-yl)-piperazine

According to the procedure described for the synthesis of intermediate 1 1-isopropyl-4-(5-nitro-pyridin-2-yl)-piperazine was synthesised from 2-chloro-5-nitro-pyridine and N-isopropylpiperazine to yield 97 % of the title compound as yellow
5 crystals. (m/e): 251.1 (MH^+ ; 100%).

Intermediate 4

6-(4-Isopropyl-piperazin-1-yl)-pyridin-3-ylamine

According to the procedure described for the synthesis of intermediate 2 6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine was synthesised from 1-isopropyl-4-(5-nitro-pyridin-2-yl)-piperazine through hydrogenation to yield the title compound which
10 was used without further purification in the following step. (m/e): 221.1 (MH^+ ; 100%).

Intermediate 5

1-Cyclopentyl-4-(5-nitro-pyridin-2-yl)-piperazine

According to the procedure described for the synthesis of intermediate 1 1-cyclopentyl-4-(5-nitro-pyridin-2-yl)-piperazine was synthesised from 2-chloro-5-nitro-pyridine and N-cyclopentylpiperazine to yield 87 % of the title compound as yellow
15 crystals. (m/e): 277.1 (MH^+ ; 100%).

Intermediate 6

20 6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine

According to the procedure described for the synthesis of intermediate 2 -(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine was synthesised from 1-cyclopentyl-4-(5-nitro-pyridin-2-yl)-piperazine through hydrogenation to yield 98% of the title
compound as colourless crystals. (m/e): 247.1 (MH^+ ; 100%).

25 Intermediate 7

1-Isobutyl-4-(5-nitro-pyridin-2-yl)-piperazine

A mixture of 7 g (34 mmol) 1-(5-nitro-pyridin-2-yl)-piperazine (commercially available), 3.15 g (44 mmol) isobutyraldehyde, 10.7 g (50 mmol) sodium trisacetoxyborohydride in 140 ml THF and 3 ml acetic acid was stirred at room

temperature for 16 h. After addition of 50 ml water, THF was removed under vacuum. The residue was taken up in 300 ml water and 400 ml ethyl acetate and made alkaline by addition of 2M aq. Na_2CO_3 solution. The mixture was afterwards extracted two times with 300 ml ethyl acetate each. The combined organic phases were washed two times
5 with 200 ml water each, dried with MgSO_4 and evaporated to dryness. The residue was used without further purification in the synthesis of intermediate 9. (m/e): 265.0 (MH^+ ; 100%).

Intermediate 8

6-(4-Isobutyl-piperazin-1-yl)-pyridin-3-ylamine

10 A mixture of 2.48 g 1-isobutyl-4-(5-nitro-pyridin-2-yl)-piperazine (intermediate 7) and 0.8 g Pd/C (10 %) in 30 ml methanol was treated with 1 bar hydrogen at room temperature for 2 h. After filtration the mixture was evaporated to dryness to yield 2.09 g (95 %) of the title compound as colourless solid. (m/e): 235.0 (MH^+ ; 100%).

Example 1

15 1-[6-(4-Ethyl-piperazin-1-yl)-pyridin-3-yl]-3-propyl-urea

A mixture of 20 mg (0.1 mmol) 6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine, 15 mg (0.15 mmol) triethylamine and 9.4 mg (0.11 mmol) 1-isocyanato-propane in 1 ml DCM was stirred at room temperature for 16 h. After evaporation the residue was taken up in 1 ml methanol/acetonitrile 1/1 and subjected to preparative HPLC purification on
20 reversed phase eluting with a gradient of acetonitrile / water (0.05 % triethylamine). The combined product fractions were evaporated to dryness to yield 15.5 mg (53 %) of the title compound. (m/e): 292.3 (MH^+ ; 100%).

According to the procedure described for the synthesis of Example 1 further pyridine derivatives have been synthesized from the intermediates 2, 4, 6 and 8,
25 respectively, and commercially available reagents listed in table 1. The examples are compiled in table 1 and comprise Examples No 2 to 60.

Table 1

No	MW	Name	Starting materials	MH+ found
2	331.46	1-cyclohexyl-3-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and 1-isocyanato-cyclohexane	332
3	339.44	1-benzyl-3-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and isocyanatomethyl-benzene	340.3
4	339.44	1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-p-tolyl-urea	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and 1-isocyanato-4-methyl-benzene	340.3
5	365.48	1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and ((1R,2S)-2-isocyanato-cyclopropyl)-benzene	366.3
6	355.44	1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and 1-isocyanato-3-methoxy-benzene	355.8
7	343.41	1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-fluoro-phenyl)-urea	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and 1-isocyanato-4-fluoro-benzene	344.2
8	371.53	1-cyclohexyl-3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and isocyanato-cyclohexane	372.3
9	379.51	1-benzyl-3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and isocyanato-methylbenzene	380.4

No	MW	Name	Starting materials	MH+ found
10	393.53	1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and 1-isocyanatomethyl-4-methylbenzene	394.2
11	405.55	1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and ((1R,2S)-2-isocyanato-cyclopropyl)-benzene	406.5
12	383.47	1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-fluoro-phenyl)-urea	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and 1-isocyanato-4-fluorobenzene	383.9
13	345.49	1-cyclohexyl-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and isocyanato-cyclohexane	345.3
14	339.44	1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-phenyl-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and isocyanato-benzene	340.2
15	353.47	1-benzyl-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and isocyanatomethyl-benzene	354.3
16	353.47	1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-o-tolyl-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and 1-isocyanato-2-methyl-benzene	353.9

No	MW	Name	Starting materials	MH+ found
17	353.47	1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-m-tolyl-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and 1-isocyanato-3-methyl-benzene	354.3
18	373.89	1-(2-chloro-phenyl)-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and 1-isocyanato-2-chloro-benzene	374.2
19	367.5	1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and isocyanatomethyl-4-methyl-benzene	368.2
20	379.51	1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and ((1R,2S)-2-isocyanato-cyclopropyl)-benzene	379.9
21	369.47	1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and 1-isocyanato-3-methoxy-benzene	370.3
22	357.43	1-(4-fluoro-phenyl)-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and 1-isocyanato-4-fluoro-benzene	358.2
23	319.45	1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-propyl-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and 1-isocyanato-propane	327.3

No	MW	Name	Starting materials	MH+ found
24	359.52	1-cyclohexyl-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and isocyanato-cyclohexane	359.9
25	367.5	1-benzyl-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and isocyanatomethyl-benzene	368.2
26	367.5	1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-m-tolyl-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and 1-isocyanato-3-methyl-benzene	368.2
27	367.5	1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-p-tolyl-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and 1-isocyanato-4-methyl-benzene	368.2
28	387.91	1-(2-chloro-phenyl)-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and 1-isocyanato-2-chloro-benzene	387.9
29	381.52	1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and isocyanatomethyl-4-methyl-benzene	382
30	393.53	1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and ((1R,2S)-2-isocyanato-cyclopropyl)-benzene	393.9

No	MW	Name	Starting materials	MH+ found
31	383.5	1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and 1-isocyanato-3-methoxy-benzene	384.3
32	371.46	1-(4-fluoro-phenyl)-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea	6-(4-isobutyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 8) and 1-isocyanato-4-fluoro-benzene	371.4
33	313.42	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-diethylaminosulfonamide	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and dimethylaminosulfamoylchloride	
34	353.49	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-diethylaminosulfonamide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and dimethylaminosulfamoylchloride	
35	327.45	N-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-diethylaminosulfonamide	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and dimethylaminosulfamoylchloride	
36	316.45	cyclohexanecarboxylic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-amide	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and cyclohexanecarbonyl chloride	317.3
37	324.43	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-phenyl-acetamide	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and phenyl-acetyl chloride	325.4
38	342.42	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-(4-fluoro-phenyl)-acetamide	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and (4-fluoro-phenyl)-acetyl chloride	343.2

No	MW	Name	Starting materials	MH+ found
39	354.45	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-(3-methoxy-phenyl)-acetamide	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and (3-methoxy-phenyl)-acetyl chloride	355.4
40	384.48	2-(3,4-dimethoxy-phenyl)-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-acetamide	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and (3,4-dimethoxy-phenyl)-acetyl chloride	385.3
41	308.38	[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid 2-methoxy-ethyl ester	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and 2-methoxy-ethanol carbonyl chloride	309.4
42	306.41	[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester	6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 2) and carbonic acid chloride monoisobutyl ester	307.4
43	314.43	cyclopropanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and cyclopropanecarbonyl chloride	315.2
44	316.45	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-butyramide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and butyryl chloride	317.3
45	328.46	cyclobutanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and cyclobutanecarbonyl chloride	329.4
46	342.49	cyclopentanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and cyclopentanecarbonyl chloride	343.4

No	MW	Name	Starting materials	MH+ found
47	344.5	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-ethyl-butyramide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and 2-ethyl-butyryl chloride	345.3
48	356.51	cyclohexanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and cyclohexanecarbonyl chloride	357.3
49	385.9	2-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-nicotinamide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and 2-chloro-nicotinoyl chloride	386.3
50	364.49	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-phenyl-acetamide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and phenyl-acetyl chloride	365.4
51	382.48	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(4-fluoro-phenyl)-acetamide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and (4-fluoro-phenyl)-acetyl chloride	383.3
52	394.52	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(3-methoxy-phenyl)-acetamide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and (3-methoxy-phenyl)-acetyl chloride	395.3
53	424.54	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(3,4-dimethoxy-phenyl)-acetamide	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and (3,4-dimethoxy-phenyl)-acetyl chloride	425.3

No	MW	Name	Starting materials	MH+ found
54	318.42	[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid ethyl ester	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and carbonic acid monoethyl ester	319.3
55	348.45	[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid 2-methoxy-ethyl ester	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and carbonic acid mono (2-methoxyethyl) ester	349.4
56	346.47	[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester	6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 6) and carbonic acid monoisobutyl ester	347.3
57	330.48	cyclohexanecarboxylic acid [6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-amide	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and cyclohexanecarbonyl chloride	331.3
58	398.51	2-(3,4-dimethoxy-phenyl)-N-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-acetamide	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and (3,4-dimethoxy-phenyl)-acetyl chloride	399.2
59	292.38	[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid ethyl ester	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and carbonic acid monoethyl ester	293.3
60	320.44	[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester	6-(4-isopropyl-piperazin-1-yl)-pyridin-3-ylamine (intermediate 4) and carbonic acid monoisobutyl ester	321.3

Intermediate 9

6-(4-Ethyl-piperazin-1-yl)-nicotinonitrile

A mixture of 2 g (14 mmol) 6-chloronicotinonitrile (commercially available) and 0.88 g (7 mmol) N,N-diisopropylethylamine in 20 ml water and 4 ml DMF was heated to 80 °C. During 2 min 1.98 g (17 mmol) N-ethylpiperazine was added and stirred at 80 °C for 1 h. 100 ml 1M aq. Na₂CO₃ solution was added and the mixture was extracted three times with 100 ml ethyl acetate each. The combined organic phases were washed twice with 100 ml brine each and dried with MgSO₄. After evaporation the residue was purified with flash chromatography on Alox eluting with a gradient of ethyl acetate / heptan to yield 1.4 g (45 %) of the title compound as slightly yellow crystals. (m/e): 217.02 (MH⁺; 100%).

Intermediate 10

C-[6-(4-Ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine

A mixture of 2 g (8 mmol) 6-(4-ethyl-piperazin-1-yl)-nicotinonitrile, 1.3 g Raney Nickel (B113Z, Degussa) in 25 ml methanol, 15 ml ethyl acetate and 5 ml ammonia aq. (ca. 25 %) was treated with 1 bar hydrogen at 30-35 °C for 4 h. The mixture was filtered and the residue washed three times with 20 ml ethyl acetate each and the combined organic phases evaporated to dryness to yield 1.83 g (97 %) of the title compound as white crystals. (m/e): 261.1 (MH⁺; 100%).

Intermediate 11

6-(4-Cyclopentyl-piperazin-1-yl)-nicotinonitrile

A mixture of 2 g (14 mmol) 6-chloronicotinonitrile (commercially available) 2.45 g (16 mmol) N-cyclopentylpiperazine (commercially available) and 1.86 g (14 mmol) N,N-diisopropylethylamine in 5 ml water and 15 ml DMF was heated to 90 °C for 24 h. After addition of 250 ml 1M NaHCO₃ aq. solution the mixture was extracted three times with 250 ml ethyl acetate each. The combined organic phases were washed twice with 150 ml brine each, dried and evaporated to dryness. Recrystallisation from ethyl acetate yielded a first batch of 2.83 g of white crystals. An additional batch was yielded from the filtrate and in total 3.14 g (85 %) of the title compound was obtained as white crystals. (m/e): 257.1 (MH⁺; 100%).

Intermediate 12

C-[6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine

According to the procedure described for the synthesis of intermediate 10 C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine was synthesized from 6-(4-cyclopentyl-piperazin-1-yl)-nicotinonitrile through hydrogenation over Raney Nickel.
 5 1.99 g (98 %) of the title compounds was obtained as white crystals. (m/e): 261.1 (MH⁺; 100%).

Example 61

N-[6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-butyramide

10 A mixture of 26 mg (0.1 mmol) C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine, 20 mg (0.2 mmol) triethylamine and 12 mg (0.11 mmol) butyryl chloride in DCM was stirred at room temperature. After evaporation acetonitrile / DMF was added and the mixture was subjected to preparative HPLC purification on reversed phase eluting with a gradient of acetonitrile / water (0.05 % triethylamine). The
 15 combined product fractions were evaporated to dryness to yield 20.3 mg (61 %) of the title compound. (m/e): 331.3

According to the procedure described for the synthesis of example 73 further pyridine derivatives have been synthesized from C-[6-(4-Ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine and C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine
 20 and the respective commercially available reagent mentioned in Table 2. The Examples are shown in table 2 and comprise Examples No 62 to 90.

Table 2

No	MW	Name	Starting materials	MH ⁺ found
62	394.52	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-methoxy-benzamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and 3-methoxy-benzoyl chloride	395.3
63	378.52	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-phenyl-acetamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and phenyl-acetyl chloride	379.4

No	MW	Name	Starting materials	MH+ found
64	396.51	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-(4-fluoro-phenyl)-acetamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and (4-fluoro-phenyl)-acetyl chloride	397.4
65	352.5	ethanesulfonic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and ethanesulfonyl chloride	353.3
66	366.53	propane-1-sulfonic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and propanesulfonyl chloride	367.2
67	367.52	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and dimethylaminosulfamoylchloride	368.2
68	400.54	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and benzenesulfonyl chloride	401.5
69	414.57	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-C-phenyl-methanesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and phenyl-methanesulfonyl chloride	415.4
70	449.02	C-(4-chloro-phenyl)-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-methanesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and 4-chloro-phenyl-methanesulfonyl chloride	449.3

No	MW	Name	Starting materials	MH+ found
71	418.53	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-fluoro-benzenesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and 2-fluoro-benzenesulfonyl chloride	419.3
72	418.53	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-fluoro-benzenesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and 3-fluoro-benzenesulfonyl chloride	419.3
73	418.53	N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-4-fluoro-benzenesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and 4-fluoro-benzenesulfonyl chloride	419.3
74	434.99	2-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and 2-chloro-benzenesulfonyl chloride	435.3
75	434.99	4-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide	C-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 12) and 4-chloro-benzenesulfonyl chloride	435.3
76	371.46	1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-(4-fluoro-benzyl)-urea	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 1-fluoro-4-isocyanatomethyl-benzene	372.3
77	354.45	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-methoxy-benzamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 3-methoxybenzoylchloride	355.3

No	MW	Name	Starting materials	MH+ found
78	338.45	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-phenyl-acetamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and phenyl-acetyl chloride	339.2
79	356.44	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-(4-fluoro-phenyl)-acetamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 4-fluoro-phenyl acetyl chloride	357.3
80	312.44	ethanesulfonic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and ethanesulfonylchloride	313
81	326.46	propane-1-sulfonic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and propanesulfonylchloride	327.3
82	327.45	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-dimethylaminosulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and dimethylaminosulfamoylchloride	328.2
83	360.48	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and benzenesulfonyl chloride	361.3
84	374.51	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-C-phenyl-methanesulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and phenyl-methanesulfonyl chloride	375.4

No	MW	Name	Starting materials	MH+ found
85	408.95	C-(4-Chloro-phenyl)-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-methanesulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 4-chloro-phenyl-methanesulfonyl chloride	409.3
86	378.47	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-fluoro-benzenesulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 2-fluoro-benzenesulfonyl chloride	379.3
87	378.47	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-fluoro-benzenesulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 3-fluoro-benzenesulfonyl chloride	379.3
88	378.47	N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-4-fluoro-benzenesulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 4-fluoro-benzenesulfonyl chloride	379.3
89	394.92	2-chloro-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 2-chloro-benzenesulfonyl chloride	395.3
90	394.92	4-chloro-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide	C-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-methylamine (intermediate 10) and 4-chloro-benzenesulfonyl chloride	395.3

- 45 -

Example 91

4-Methyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

A mixture of 29 mg (0.08 mmol) [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid phenyl ester (intermediately built from intermediate 6 and phenylchloroformate) and 12 mg (0.12 mmol) 4-methyl-piperidine (commercially available) in 1ml DCM and 0.1 ml DMF was stirred at room temperature for 16 h. After evaporation the residue was taken up in methanol/DMF and subjected to preparative HPLC purification on reversed phase eluting with a gradient of acetonitrile / water (0.05 % triethylamine). The combined product fractions were evaporated to dryness to yield 13.6 mg (46 %) of the title compound. (m/e): 372.3 (MH⁺; 100%).

Example 92

2,6-Dimethyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

According to the procedure described for the synthesis of Example 91 2,6-dimethyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide was synthesised from intermediate 6 and 2,6-dimethyl-piperidine (commercially available). (m/e): 372.3 (MH⁺; 100%).

Example 93

4-Trifluoromethyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

According to the procedure described for the synthesis of Example 91 4-trifluoromethyl-piperidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide was synthesised from intermediate 6 and 4-trifluoromethyl-piperidine (commercially available). (m/e): 426.3 (MH⁺; 100%).

Example 94

1-[6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-ylcarbamoyl]-piperidine-4-carboxylic acid ethyl ester

According to the procedure described for the synthesis of Example 91 1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylcarbamoyl]-piperidine-4-carboxylic

- 46 -

acid ethyl ester was synthesised from intermediate 6 and piperidine-4-carboxylic acid ethyl ester (commercially available). (m/e): 430.4 (MH⁺; 100%).

Example 95

5 Octahydro-quinoline-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

According to the procedure described for the synthesis of Example 91 octahydro-quinoline-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide was synthesised from intermediate 6 and 2,6-dimethyl-piperidine (commercially available). (m/e): 412.5 (MH⁺; 100%).

10 Example 96

Octahydro-isoquinoline-2-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

15 According to the procedure described for the synthesis of Example 91 octahydro-isoquinoline-2-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide was synthesised from intermediate 6 and octahydro-isoquinoline (commercially available). (m/e): 412.5 (MH⁺; 100%).

Example 97

2-trifluoromethyl-pyrrolidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

20 According to the procedure described for the synthesis of Example 91 2-trifluoromethyl-pyrrolidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide was synthesised from intermediate 6 and 2-trifluoromethyl-pyrrolidine (commercially available). (m/e): 412.4 (MH⁺; 100%).

Example 98

25 2-Isopropyl-pyrrolidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

According to the procedure described for the synthesis of Example 91 2-isopropyl-pyrrolidine-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide was synthesised from intermediate 6 and 2-isopropyl-pyrrolidine
30 (commercially available). (m/e): 386.3 (MH⁺; 100%).

Example 99

1,3-Dihydro-isoindole-2-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

According to the procedure described for the synthesis of Example 91 1,3-dihydro-isoindole-2-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide was synthesised from intermediate 6 and 1,3-dihydro-isoindole (commercially available). (m/e): 392.2 (MH⁺; 100%).

Example 100

3-[6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-isopropyl-1-(2-methoxy-ethyl)-urea

According to the procedure described for the synthesis of Example 91 3-[6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-isopropyl-1-(2-methoxy-ethyl)-urea was synthesised from intermediate 6 and isopropyl-2-methoxyethyl-amine (commercially available). (m/e): 390.4 (MH⁺; 100%).

Example 101

Azepane-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide

According to the procedure described for the synthesis of Example 91 azepane-1-carboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide was synthesised from intermediate 6 and azepine (commercially available). (m/e): 372.3 (MH⁺; 100%).

Example 102

3-[6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-ethyl-1-phenyl-urea

According to the procedure described for the synthesis of Example 91 3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-ethyl-1-phenyl-urea was synthesised from intermediate 6 and ethylphenylamine (commercially available). (m/e): 394.4 (MH⁺; 100%).

Example 103

3-[6-(4-Cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-(4-methoxy-phenyl)-1-methyl-urea

According to the procedure described for the synthesis of Example 91 3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-1-(4-methoxy-phenyl)-1-methyl-urea was

- 48 -

synthesised from intermediate 6 and (4-methoxy-phenyl)-methyl-amine (commercially available). (m/e): 410.4 (MH^+ ; 100%).

Example 104

3,4-Dihydro-2H-quinoline-1-carboxylic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-
5 amide

According to the procedure described for the synthesis of Example 91
3,4-dihydro-2H-quinoline-1-carboxylic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-
amide was synthesised from intermediate 2 and 3,4-dihydro-2H-quinoline
(commercially available). (m/e): 366.3 (MH^+ ; 100%).

10

Example 105

3,4-Dihydro-2H-quinoline-1-carboxylic acid [6-(4-isopropyl-piperazin-1-yl)-pyridin-3-
yl]-amide

According to the procedure described for the synthesis of Example 91 3,4-dihydro-
2H-quinoline-1-carboxylic acid [6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-amide
15 was synthesised from intermediate 4 and 3,4-dihydro-2H-quinoline (commercially
available). (m/e): 380.3 (MH^+ ; 100%).

Example A

Film coated tablets containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per tablet</u>	
Kernel:		
Compound of formula (I)	10.0 mg	200.0 mg
Microcrystalline cellulose	23.5 mg	43.5 mg
Lactose hydrous	60.0 mg	70.0 mg
Povidone K30	12.5 mg	15.0 mg
Sodium starch glycolate	12.5 mg	17.0 mg
Magnesium stearate	1.5 mg	4.5 mg
(Kernel Weight)	120.0 mg	350.0 mg
Film Coat:		
Hydroxypropyl methyl cellulose	3.5 mg	7.0 mg
Polyethylene glycol 6000	0.8 mg	1.6 mg
Talc	1.3 mg	2.6 mg
Iron oxyde (yellow)	0.8 mg	1.6 mg
Titanium dioxide	0.8 mg	1.6 mg

- 5 The active ingredient is sieved and mixed with microcrystalline cellulose and the mixture is granulated with a solution of polyvinylpyrrolidone in water. The granulate is mixed with sodium starch glycolate and magnesiumstearate and compressed to yield kernels of 120 or 350 mg respectively. The kernels are lacquered with an aqueous solution / suspension of the above mentioned film coat.

Example B

Capsules containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per capsule</u>
Compound of formula (I)	25.0 mg
Lactose	150.0 mg
Maize starch	20.0 mg
Talc	5.0 mg

- 5 The components are sieved and mixed and filled into capsules of size 2.

Example C

Injection solutions can have the following composition:

Compound of formula (I)	3.0 mg
Gelatine	150.0 mg
Phenol	4.7 mg
Sodium carbonate	to obtain a final pH of 7
Water for injection solutions	ad 1.0 ml

- 51 -

Example D

Soft gelatin capsules containing the following ingredients can be manufactured in a conventional manner:

Capsule contents

Compound of formula (I)	5.0 mg
Yellow wax	8.0 mg
Hydrogenated Soya bean oil	8.0 mg
Partially hydrogenated plant oils	34.0 mg
Soya bean oil	110.0 mg
Weight of capsule contents	165.0 mg

Gelatin capsule

Gelatin	75.0 mg
Glycerol 85 %	32.0 mg
Karion 83	8.0 mg (dry matter)
Titanium dioxide	0.4 mg
Iron oxide yellow	1.1 mg

- 5 The active ingredient is dissolved in a warm melting of the other ingredients and the mixture is filled into soft gelatin capsules of appropriate size. The filled soft gelatin capsules are treated according to the usual procedures.

Example E


Sachets containing the following ingredients can be manufactured in a conventional manner:

Compound of formula (I)	50.0 mg
Lactose, fine powder	1015.0 mg
Microcrystalline cellulose (AVICEL PH 102)	1400.0 mg
Sodium carboxymethyl cellulose	14.0 mg
Polyvinylpyrrolidone K 30	10.0 mg
Magnesiumstearate	10.0 mg
Flavoring additives	1.0 mg

- 5 The active ingredient is mixed with lactose, microcrystalline cellulose and sodium carboxymethyl cellulose and granulated with a mixture of polyvinylpyrrolidone in water. The granulate is mixed with magnesiumstearate and the flavouring additives and filled into sachets.

Claims

1. Compounds of the general formula



I

5 wherein

R^1 is lower alkyl or C_3 - C_7 -cycloalkyl;

X is $\text{C}(\text{O})$ or SO_2 ;

m is 0 or 1;

10 R^2 is selected from the group consisting of lower alkyl, C_3 - C_7 -alkenyl, C_3 - C_7 -alkinyl, lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl, unsubstituted C_3 - C_7 -cycloalkyl or C_3 - C_7 -cycloalkyl substituted by phenyl, lower C_3 - C_7 -cycloalkylalkyl, lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, 15 unsubstituted pyridyl or pyridyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and $-\text{NR}^3\text{R}^4$,

or, in case X is $\text{C}(\text{O})$, R^2 can also be lower alkoxy or lower alkoxyalkoxy,

20 or, in case m is 1, R^2 can also be unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl,

R^3 is hydrogen or lower alkyl;

R^4 is selected from the group consisting of lower alkyl, C_3 - C_7 -alkenyl, C_3 - C_7 -alkinyl, lower alkoxyalkyl, C_3 - C_7 -cycloalkyl,

25 C_3 - C_7 -cycloalkyl substituted by phenyl, lower C_3 - C_7 -cycloalkylalkyl,

unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl; or

- 5 R^3 and R^4 together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, lower alkoxycarbonyl, oxo, halogen and halogenalkyl, or being
10 condensed with a C_5 - C_6 -cycloalkyl ring or a phenyl ring, said cycloalkyl ring or phenyl ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, halogen and halogenalkyl;

and pharmaceutically acceptable salts thereof;

- with the exception of 2,2-dimethyl-N-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-
15 propionamide.

2. Compounds of formula I according to claim 1, wherein R^1 is C_3 - C_7 -cycloalkyl.

3. Compounds of formula I according to claims 1 or 2, wherein R^1 is ethyl or isopropyl.

4. Compounds of formula I according to any of claims 1 to 3, wherein m is 1.

- 20 5. Compounds of formula I according to any of claims 1 to 3, wherein m is 0.

6. Compounds of formula I according to any of claims 1 to 5, wherein R^2 is selected from the group consisting of
lower alkyl, C_3 - C_7 -alkenyl, C_3 - C_7 -alkinyl,
lower halogenalkyl, lower hydroxyalkyl, lower alkoxyalkyl,
25 unsubstituted C_3 - C_7 -cycloalkyl or C_3 - C_7 -cycloalkyl substituted by phenyl,
lower C_3 - C_7 -cycloalkylalkyl,
lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl,
unsubstituted pyridyl or pyridyl mono- or disubstituted by lower alkyl, lower alkoxy,
30 halogen or lower halogenalkyl, and

-NR³R⁴,

or, in case X is C(O), R² can also be lower alkoxy or lower alkoxyalkoxy.

7. Compounds of formula I according to any of claims 1 to 6, wherein R² is selected from the group consisting of lower alkyl, C₃-C₇-cycloalkyl and C₃-C₇-cycloalkyl substituted by phenyl.

8. Compounds of formula I according to any of claims 1 to 6, wherein R² is lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl.

9. Compounds of formula I according to any of claims 1 to 6, wherein R² is -NR³R⁴ and R³ and R⁴ are as defined in claim 1.

10. Compounds of formula I according to any of claims 1 to 6 or 9, wherein

R³ is hydrogen or lower alkyl;

R⁴ is selected from the group consisting of lower alkyl, C₃-C₇-alkenyl, C₃-C₇-alkinyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkyl substituted by phenyl, lower C₃-C₇-cycloalkylalkyl, unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl; or

R³ and R⁴ together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, oxo, halogen and halogenalkyl, or being condensed with a phenyl ring, said phenyl ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy, halogen and halogenalkyl.

11. Compounds of formula I according to claims 9 or 10, wherein R³ and R⁴ are lower alkyl.

12. Compounds of formula I according to claims 9 or 10, wherein R^3 is hydrogen and R^4 is selected from the group consisting of lower alkyl, C_3 - C_7 -alkenyl, C_3 - C_7 -alkinyl, C_3 - C_7 -cycloalkyl, C_3 - C_7 -cycloalkyl substituted by phenyl,
- 5 lower C_3 - C_7 -cycloalkylalkyl, unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl, and lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl.
- 10 13. Compounds of formula I according to claims 9 or 10, wherein R^3 is hydrogen and R^4 is lower phenylalkyl wherein phenyl is unsubstituted or mono- or disubstituted by lower alkyl, lower alkoxy, halogen or lower halogenalkyl.
14. Compounds of formula I according to any of claims 1 to 4, wherein m is 1 and R^2 is unsubstituted phenyl or phenyl mono- or disubstituted by lower alkyl, lower
- 15 alkoxy, halogen or lower halogenalkyl.
15. Compounds of formula I according to any of claims 1 to 14, wherein X is SO_2 .
16. Compounds of formula I according to any of claims 1 to 14, wherein X is $C(O)$.
17. Compounds of formula I according to any of claims 1 to 14, wherein X is $C(O)$ and R^2 is $-NR^3R^4$.
- 20 18. Compounds of formula I according to claim 1, selected from the group consisting of
- 1-[6-(4-ethylpiperazin-1-yl)-pyridin-3-yl]-3-propyl-urea,
1-cyclohexyl-3-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-urea,
1-benzyl-3-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-urea,
25 1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-p-tolyl-urea,
1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea,
1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea,
1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-fluoro-phenyl)-urea,
1-cyclohexyl-3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-urea,
30 1-benzyl-3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-urea,
1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea,

- 1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea,
- 1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-fluoro-phenyl)-urea,
- 1-cyclohexyl-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 5 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-phenyl-urea,
- 1-benzyl-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-o-tolyl-urea,
- 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-m-tolyl-urea,
- 1-(2-chloro-phenyl)-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 10 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea,
- 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea,
- 1-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea,
- 1-(4-fluoro-phenyl)-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-propyl-urea,
- 15 1-cyclohexyl-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-benzyl-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-m-tolyl-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-p-tolyl-urea,
- 1-(2-chloro-phenyl)-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- 20 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-((1R,2S)-2-phenyl-cyclopropyl)-urea,
- 1-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-3-(3-methoxy-phenyl)-urea,
- 1-(4-fluoro-phenyl)-3-[6-(4-isobutyl-piperazin-1-yl)-pyridin-3-yl]-urea,
- N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-dimethylaminosulfonamide,
- 25 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-dimethylaminosulfonamide,
- N-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-dimethylaminosulfonamide,
- cyclohexanecarboxylic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-phenyl-acetamide,
- N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-(4-fluoro-phenyl)-acetamide,
- 30 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-2-(3-methoxy-phenyl)-acetamide,
- 2-(3,4-dimethoxy-phenyl)-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-acetamide,
- [6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid 2-methoxy-ethyl ester,
- [6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester,
- cyclopropanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- 35 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-butyramide,
- cyclobutanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- cyclopentanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,
- N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-ethyl-butyramide,
- cyclohexanecarboxylic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-amide,

- 2-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-nicotinamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-phenyl-acetamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(4-fluoro-phenyl)-acetamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(3-methoxy-phenyl)-acetamide,
 5 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(3,4-dimethoxy-phenyl)-
 acetamide,
 [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid ethyl ester,
 [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid 2-methoxy-ethyl ester,
 [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester,
 10 cyclohexanecarboxylic acid [6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-amide,
 2-(3,4-dimethoxy-phenyl)-N-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-acetamide,
 [6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid ethyl ester,
 [6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-carbamic acid isobutyl ester,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-butyramide,
 15 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-methoxy-benzamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-phenyl-acetamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-(4-fluoro-phenyl)-
 acetamide,
 ethanesulfonic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide,
 20 propane-1-sulfonic acid [6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-dimethylaminosulfonamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-C-phenyl-
 methanesulfonamide,
 25 C-(4-chloro-phenyl)-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-
 methanesulfonamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-fluoro-benzenesulfonamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-fluoro-benzenesulfonamide,
 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-4-fluoro-benzenesulfonamide,
 30 2-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
 4-chloro-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
 1-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-(4-fluoro-benzyl)-urea,
 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-methoxy-benzamide,
 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-phenyl-acetamide,
 35 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-(4-fluoro-phenyl)-acetamide,
 ethanesulfonic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide,
 propane-1-sulfonic acid [6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-amide,
 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-dimethylaminosulfonamide,
 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,

N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-C-phenyl-methanesulfonamide,
C-(4-chloro-phenyl)-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-
methanesulfonamide,


- 5 N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-fluoro-benzenesulfonamide,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-fluoro-benzenesulfonamide,
N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-4-fluoro-benzenesulfonamide,
2-chloro-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
4-chloro-N-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-ylmethyl]-benzenesulfonamide,
and pharmaceutically acceptable salts thereof.

- 10 19. Compounds of formula I according to claim 1, selected from the group
consisting of

- 1-benzyl-3-[6-(4-ethyl-piperazin-1-yl)-pyridin-3-yl]-urea,
1-benzyl-3-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-urea,
1-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-3-(4-methyl-benzyl)-urea,
15 1-benzyl-3-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-urea,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-dimethylaminosulfonamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-yl]-2-(3,4-dimethoxy-phenyl)-
acetamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-3-methoxy-benzamide,
20 N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-2-phenyl-acetamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-dimethylaminosulfonamide,
N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-C-phenyl-
methanesulfonamide,
C-(4-chloro-phenyl)-N-[6-(4-cyclopentyl-piperazin-1-yl)-pyridin-3-ylmethyl]-
25 methanesulfonamide,
and pharmaceutically acceptable salts thereof.


20. A process for the manufacture of compounds according to any one of claims 1
to 19, which process comprises

- a) reacting a compound of the formula II




II

wherein R^1 and m are as defined in claim 1,
with a sulfonylchloride or sulfamoylchloride of the formula III




wherein R^2 is as defined in claim 1,
to obtain a compound of the formula I-A



I-A

wherein R^1 , R^2 and m are as defined in claim 1, or

b) reacting a compound of the formula II




II

wherein R^1 and m are as defined in claim 1,
with a chloride of the formula IV




wherein R^2 is as defined in claim 1,
to obtain a compound of the formula I-B



I-B

wherein R^1 , R^2 and m are as defined in claim 1, or

c) reacting a compound of the formula II




II

wherein R^1 and m are as defined in claim 1,
with an isocyanate of the formula V

5



wherein R^4 is as defined in claim 1,
to obtain a compound of the formula I-C



I-C

wherein R^3 is hydrogen and R^1 , R^4 and m are as defined in claim 1,

10 and if desired, converting the compound of formula I-A, I-B or I-C into a
pharmaceutically acceptable salt.

21. Compounds according to any one of claims 1 to 19 when manufactured by a
process according to claim 20.

22. Pharmaceutical compositions comprising a compound according to any one of
15 claims 1 to 19 and a pharmaceutically acceptable carrier and/or adjuvant.

23. Pharmaceutical compositions according to claim 22 for the treatment and/or
prevention of diseases which are associated with the modulation of H3 receptors.

24. Compounds according to any one of claims 1 to 19 for use as therapeutically
active substances.

25. Compounds according to any one of claims 1 to 19 for use as therapeutically active substances for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.
26. The use of compounds according to any one of claims 1 to 19 for the preparation of medicaments for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.
27. The use according to claim 26 for the treatment and/or prevention of obesity.