



## CLAIM

1. A method of preparing Compound (N), comprising:




(q) reacting Compound (K) or a pharmaceutically acceptable salt thereof with acetaldehyde in the presence of a palladium catalyst to form Compound (M)



(M) or a pharmaceutically acceptable salt thereof, wherein X<sup>1</sup> is independently -F or -Cl, and X<sup>3</sup> is -Br or -I; and

(p) tosylating Compound (M) or a pharmaceutically acceptable salt thereof to form




Compound (N) .

2. The method of claim 1, wherein the palladium catalyst of step (q) comprises a mixture of bis(dibenzylideneacetone) palladium and a tertiary phosphine ligand, PR<sub>3</sub>, wherein R is C<sub>1-6</sub> alkyl or C<sub>5-6</sub> cycloalkyl.


3. The method of claim 2, wherein the tertiary phosphine ligand, PR<sub>3</sub> is P(<sup>t</sup>Bu)<sub>3</sub>, PCy<sub>3</sub>, P(*i*-Pr)<sub>3</sub>, P(Bu)<sub>3</sub>, PEt<sub>3</sub>, PMe<sub>3</sub>, or any combination thereof.

4. The method of either of claims 2 or 3, wherein the tertiary phosphine ligand comprises P(<sup>t</sup>Bu)<sub>3</sub>.

5. A method of preparing compound (N) comprising:




(1) reacting Compound (J) **(J)** or a pharmaceutically acceptable salt




thereof with an iodinating agent or a brominating agent to form Compound (K) **(K)** or a pharmaceutically acceptable salt thereof, wherein X<sup>1</sup> is -F or -Cl, and X<sup>3</sup> is Br or I;

(m) reacting Compound (K) or a pharmaceutically acceptable salt thereof with




trimethylsilyl acetylene to form Compound (L) **(L)** or a pharmaceutically acceptable salt thereof, wherein TMS is trimethylsilyl;

(n) reacting Compound (L) or a pharmaceutically acceptable salt thereof with a C<sub>1-6</sub>




alkoxide base to form Compound (P) **(P)** or a pharmaceutically acceptable salt thereof;

(o) reacting Compound (P) with potassium *tert*-butoxide, potassium *tert*-amylate, or



any combination thereof to form Compound (M) **(M)** or a pharmaceutically acceptable salt thereof; and

(p) tosylating Compound (M) or a pharmaceutically acceptable salt thereof to form



Compound (N) **(N)** .

6. The method of claim 5, wherein the C<sub>1-6</sub> alkoxide base of step (n) comprises potassium *tert*-amylate, potassium *tert*-butoxide, potassium methoxide, sodium *tert*-amylate, sodium *tert*-butoxide, sodium methoxide, or any combinations thereof.
7. The method of either of claims 5 or 6, wherein the reaction of Compound (K) or a pharmaceutically acceptable salt thereof with trimethylsilyl acetylene in step (m) is performed in the presence of a palladium catalyst comprising Pd(Ph<sub>3</sub>P)<sub>4</sub>, Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>, Pd(dppf)<sub>2</sub>Cl<sub>2</sub>, or any combination thereof, a copper (I) halide catalyst, or any combination thereof.
8. The method of any one of claims 5-7, wherein the reaction of Compound (K) or a pharmaceutically acceptable salt thereof with trimethylsilyl acetylene in step (m) is performed in the presence of CuI, Pd(Ph<sub>3</sub>P)<sub>4</sub>, Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>, Pd(dppf)<sub>2</sub>Cl<sub>2</sub>, or any combination thereof.
9. The method of any one of claims 5-8, wherein the tosylation of step (p) is performed by reacting Compound (M) or a pharmaceutically acceptable salt thereof with TsCl.
10. The method of any one of claims 5-9, wherein Compound (J) or a pharmaceutically acceptable salt thereof is reacted with an iodinating agent comprising I<sub>2</sub>, ICl, *N*-iodosuccinimide, and wherein X<sup>3</sup> is I.
11. The method of claim 10, wherein the iodinating agent is I<sub>2</sub>.
12. The method of any one claims 5-9, wherein Compound (J) or a pharmaceutically acceptable salt thereof is reacted with a brominating agent comprising Br<sub>2</sub>, *N*-bromosuccinimide, 1,3-dibromo-5,5-dimethylhydantoin, or any combination thereof, and wherein X<sup>3</sup> is Br.
13. The method of claim 12, wherein the brominating agent is Br<sub>2</sub>.
14. The method of claim 1, further comprising treating the reaction mixture of step (q) with a carbonate base prior to the step (p) reaction.
15. The method of claim 14, wherein the carbonate base is Na<sub>2</sub>CO<sub>3</sub>.