

Field of Invention

10 The present invention relates to antibody-based polypeptides with binding specificity for CD137, which have utility in the treatment of diseases such as cancer. The invention also relates to pharmaceutical compositions, uses, methods and kits comprising such antibodies.

15 Background

CD137 (4-1BB, TNFRSF9) is a TNF receptor (TNFR) superfamily member and is expressed on activated CD4⁺ and CD8⁺ T cells, Treg, DC, monocytes, mast cells and eosinophils. CD137 activation plays an important role in CD8⁺ T cell activation and survival
20 (Lee *et al.*, 2002; Pulle *et al.*, 2006). It sustains and augments, rather than initiates, effector functions and preferentially supports Th1 cytokine production (Shuford *et al.*, 1997). In CD4⁺ T cells, CD137 stimulation initially results in activation and later in activation-induced cell death, explaining why CD137 agonistic antibodies have shown therapeutic effect in tumour immunity as well as in autoimmunity (Zhang, JCI, 2007, Sun, Trends Mol Med, 2003).
25 CD137 also suppresses Treg function (So, Cytokine Growth Factor Rev, 2008). Activation of CD137 is dependent on receptor oligomerization (Rabu *et al.*, 2005; Wyzgol *et al.*, 2009).

CD137 agonistic antibody has been shown to activate endothelial cells in the tumour
30 environment, leading to upregulation of ICAM-1 and VCAM-1 and improved T cell recruitment (Palazon, Cancer Res, 2011).

CD137 is upregulated on NK cells activated by cytokines or CD16, in mice or humans, respectively (see Melero, CCR 19 (5)1044-53, 2013 and references cited therein). CD137
35 has been shown to activate NK cells in mice as well as humans, potentiating ADCC (Kohrt *et al.*, 2014), though there are reports suggesting opposite effects on NK cells in mice and

humans, leading to NK cell activation in mice and inhibition in humans (Baessler, Blood, 2010).

Several studies have demonstrated induction of tumour immunity by treatment with agonistic CD137 antibody (Dubrot *et al.*, 2010; Gauttier *et al.*, 2014; Kim *et al.*, 2001; 5 McMillin *et al.*, 2006; Melero *et al.*, 1997; Miller *et al.*, 2002; Sallin *et al.*, 2014; Taraban *et al.*, 2002; Uno *et al.*, 2006; Vinay and Kwon, 2012; Wilcox *et al.*, 2002). In addition, it synergizes with several immunomodulators, including CpG, TRAIL, CD40, OX-40, DR5, PD-1/PD-L1, CTLA-4 Tim-3, IL-2, IL-12 (Curran *et al.*, 2011; Gray *et al.*, 2008; Guo *et al.*, 10 2013; Kwong *et al.*, 2013; Lee *et al.*, 2004; Morales-Kastresana *et al.*, 2013; Pan *et al.*, 2002; St Rose *et al.*, 2013; Uno *et al.*, 2006; Wei *et al.*, 2013; Westwood *et al.*, 2010; Westwood *et al.*, 2014a; Westwood *et al.*, 2014b) in pre-clinical models.

Two CD137 antibodies are in clinical development. Urelumab (BMS-66513) is a fully 15 human IgG4 antibody developed by Bristol-Myers Squibb. Several phase I and II studies in various indications are currently ongoing. The other CD137 antibody in development is PF-05082566, a fully human IgG2 antibody developed by Pfizer. It is currently in phase I development in lymphoma and various solid cancers.

20 The agonistic effect of CD137 antibodies is affected by the isotype of the Fc region. The antibodies tested in the clinic are either IgG2 or IgG4. Like most TNFR family members, CD137 depends on cross linking for activation (Wilson 2011, Cancer Cell). The CD137L expressed on the membrane of an APC may induce significant multiple cross linking of the receptor. An antibody can by itself only cross link two CD137 receptors, and to induce a 25 strong signal, further cross linking via FcγRs expressed on other cells (in trans) may be necessary for induction of a strong CD137 mediated signal. An exception to this may be IgG2 antibodies, which induce a cross linking independent signaling by an unknown mechanism (White et al, 2015 Cancer Cell). T cells do not express FcγRs, and the FcγR mediated cross linking *in vivo* is thought to be mediated by monocytes, macrophages, DCs 30 and potentially B cells and other cell types. It has been suggested that interaction with the inhibitory FcγR FcγRIIB plays a major role for this effect in mouse models for CD40 agonists (Li 2011, Science), whereas for OX40 antibodies, interactions with activating receptors may be of greater importance (Bulliard 2014, Imm and Cell Biol). For CD137 antibodies, FcγRII is not critical (Sanmamed 2015, Semin Onc). The translational 35 relevance of this is uncertain, since the human FcγR distribution as well as the affinity of different IgG isotypes to different FcγR differs from mice. Further, human IgG1 binds to

mFcγRIIb with relatively low affinity, similar to mIgGIIa and considerably lower than mIgG1, the latter having the most potent effect in vivo (Li Science 2011, Overdijk 2012 JI, Horton et al 2008, White et al 2011 and 2014).

5 Another factor to take into account is that engagement of FcγR receptors may also induce ADCC, antibody-dependent cellular phagocytosis (ADCP) and complement-dependent cytotoxicity (CDC) on cells coated with antibodies (for simplicity ADCC below includes ADCP and CDC). Typically, human IgG1 is a strong inducer of NK/Macrophage dependent ADCC, depending on the nature of the target, the cell type and the receptor density. IgG4
10 antibodies may also induce ADCC but to a lower extent than IgG1 (Wang 2015, Front Imm; Vidarson 2014 Front Imm).

The effect of a CD137 agonistic antibody with different isotypes may thus be affected by the balance between 1) inducing cross linking, which results in a stronger immune
15 activation, and 2) inducing ADCC, which may lead to killing of both effector T cells (predominantly CD8 T cells) and Tregs. The net effect of 1) and 2) will likely depend on the distribution of CD137 expressing cells, the possibility of the target cells to engage with FcγR expressing immune cells, the receptor density and affinity and the sensitivity of Teff vs Treg to ADCC. The CD137 expression is high both on CD8 and Tregs in melanoma
20 tumours (Quezada, presentation SITC 2015). The IgG4 format would allow for FcγRI mediated cross linking by macrophages and monocytes, yet minimizing NK mediated ADCC of effector CD8 T cells.

However, as outlined above, it is difficult to translate comparison of different human Fc in
25 mouse models due to differences in expression and affinity between murine and human FcRs. Further, the functional consequence in vivo of antibodies blocking the binding of the CD137L to CD137 is currently debated.

Several studies have demonstrated induction of tumour immunity by treatment with
30 agonistic CD137 mAb (Dubrot et al., 2010; Gauttier et al., 2014; Kim et al., 2001; McMillin et al., 2006; Melero et al., 1997; Miller et al., 2002; Sallin et al., 2014; Taraban et al., 2002; Uno et al., 2006; Vinay and Kwon, 2012; Wilcox et al., 2002). Two different antibodies are commonly used for in vivo studies in mice, Lob12.3 and 3H3 (Shuford 1997 J Exp Med).

35 The toxicity seen in mouse models has been detected following repeated dosing in a time dependent but not dose dependent manner (Ascierto 2010 Semin Onc, Dubrot 2010 Can

Imm, Niu 2007 JI). The toxicity includes skin toxicity and liver toxicity: aspartate amino transferase/alanine amino transferase ratio (ASAT/ALAT) and cytokine release. This suggests that either the toxicity requires CD137 mediated pre-activation of immune cell populations (likely T cells) or it depends on secondary effects caused by antidrug-antibodies (ADA) response, potentially forming aggregations of CD137 antibodies that may lead to enhanced cross-linking. The toxicities seen in mice are reversible and seems to depend on TNFa/CD8 cell dependent manner (Ascierto 2010 Sem Onc). Toxicology studies in monkeys showed that both single and repeated dosing of up to 100mg/kg once weekly for four weeks was tolerable with no skin or liver toxicity detected (Ascierto 2010, Semin Onc).

Prolonged and continuous activation through TNF receptor family members may lead to immune exhaustion. Therefore, it may be of advantage to administer such antibodies in a manner allowing resting periods for the cells expressing the receptors. One approach to increase the resting period in a specific dosing protocol is to reduce the half-life of an antibody by for example decreasing the binding to the neonatal Fc receptor (FcRn). This could, depending on the administration route, also reduce the toxicity associated with the treatment.

There remains a need for improved anti-tumour therapies, particularly anti-CD137 antibodies suitable for clinical use and with improved properties, such as reduced toxicity.

Summary of Invention

A first aspect of the invention provides an antibody or an antigen-binding fragment thereof ('antibody polypeptides') with binding specificity for domain 2 of CD137, wherein the antibody or antigen-binding fragment is a CD137 agonist and is capable of inhibiting the binding of reference antibody '1630/1631' to human CD137.

A second aspect of the invention, provides an antibody or an antigen-binding fragment thereof ('antibody polypeptides') with binding specificity for domain 2 of CD137, wherein the antibody or antigen-binding fragment is a CD137 agonist and is capable of inhibiting the binding of reference antibody '2674/2675' to human CD137.

In one embodiment of the above aspects of the invention, the antibody or antigen binding fragment is capable of inhibiting the binding of reference antibody '1630/1631' and/or '2674/2675' to human CD137.

- 5 According to the first aspect of the invention, antibody polypeptides are provided which are capable of inhibiting the binding of one or more reference antibodies to human CD137.

For the avoidance of doubt, the following disclosures are applicable to both the first and second aspect of the invention.

10

By "CD137" we specifically include the human CD137 protein, for example as described in GenBank Accession No. AAH06196.1 (the sequence of which is set out in SEQ ID NO: 11, below). CD137 is also known in the scientific literature as 4-1BB and TNFRSF9.

- 15 Human CD137, amino acid sequence: >gi|571321|gb|AAA53133.1| 4-1BB [Homo sapiens]

MGNSCYNIVATLLLVLNFERTRSLQDPCSNCPAGTFCDNNRNQICSPCPPNSESS
AGGQRTCDIC**RQCKGVFTRTRKECSSTSNAECDCTPGFHCLGAGCSMCEQDCKQ**GGQ
ELTKKGCKDCCFGTFNDQKRIGICRPWTNCSLDGKSVLVNGTKERDVVCGPSPADL
20 SPGASSVTTPAPAREPGHSPQIIISFFLALTSTALLELLFFLTLRFSVVKRGRKKL
LYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCEL

[SEQ ID NO: 11]

- 25 By "domain 2" corresponds to amino acids 66 to 107 of human CD137 (see bold, underlined region in SEQ ID NO:11 above).

Thus, the antibody polypeptides of the invention have specificity for CD137. By "specificity" we mean that the antibody polypeptide is capable of binding to CD137 *in vivo*, *i.e.* under the physiological conditions in which CD137 exists within the human
30 body. Preferably, the antibody polypeptide does not bind to any other protein *in vivo*. Such binding specificity may be determined by methods well known in the art, such as ELISA, immunohistochemistry, immunoprecipitation, Western blots and flow cytometry using transfected cells expressing CD137.

- 35 The antibody preferably binds to human CD137 with a K_d value which is less than 10x10⁻⁹M or less than 7x10⁻⁹M, more preferably less than 4, or 2x10⁻⁹M, most preferably less than

1.2x10⁻⁹M. Advantageously, the antibody polypeptide is capable of binding selectively to CD137, *i.e.* it binds at least 10-fold more strongly to CD137 than to any other proteins. The anti-CD137 antibody preferably specifically binds to CD137, *i.e.* it binds to CD137 but does not bind, or binds at a lower affinity, to other molecules. Therefore, typically, the K_d for the antibody with respect to human CD137 will be 2-fold, preferably 5-fold, more preferably 10-fold less than K_d with respect to the other, non-target molecule, such as murine CD137, other TNFR superfamily members, or any other unrelated material or accompanying material in the environment. More preferably, the K_d will be 50-fold less, even more preferably 100-fold less, and yet more preferably 200-fold less.

Methods for measuring the overall affinity (K_D) and on-rate (k_a) and off-rate (k_d) of an interaction (such as an interaction between an antibody and a ligand) are well known in the art. Exemplary *in vitro* methods are described in the accompanying Examples. It is also conceivable to use flow cytometry based methods (Sklar *et al.*, Annu Rev Biophys Biomol Struct, (31), 97-119, 2002).

The term CD137 as used herein typically refers to human CD137. The antibody may have some binding affinity for CD137 from other mammals, such as CD137 from a non-human primate, for example *Macaca fascicularis* (cynomolgus monkey). The antibody preferably does not bind to murine CD137 and/or does not bind to other human TNFR superfamily members, for example human OX40 or CD40.

Typically, the invention provides an antibody or antigen-binding fragment with affinity for CD137 in its native state, and in particular for CD137 localised on the surface of a cell.

By "localised on the surface of a cell" it is meant that CD137 is associated with the cell such that one or more region of CD137 is present on the outer face of the cell surface. For example, CD137 may be inserted into the cell plasma membrane (*i.e.* orientated as a transmembrane protein) with one or more regions presented on the extracellular surface. This may occur in the course of expression of CD137 by the cell. Thus, in one embodiment, "localised on the surface of a cell" may mean "expressed on the surface of a cell." Alternatively, CD137 may be outside the cell with covalent and/or ionic interactions localising it to a specific region or regions of the cell surface.

The antibodies and antigen-binding fragments thereof as defined herein are CD137 agonists. For example, they may be capable of inducing the release of interferon-gamma

from CD8+ T cells. Agonistic activity of anti-CD137 antibodies may be evaluated in a T cell assay based on primary CD8+ T cells (see Examples).

Thus, the antibody may modulate the activity of a cell expressing CD137, wherein said modulation is an increase or decrease in the activity of said cell. The cell is typically a T cell. The antibody may increase the activity of a CD4+ or CD8+ effector cell, or may decrease the activity of, or deplete, a regulatory T cell (T reg). In either case, the net effect of the antibody will be an increase in the activity of effector T cells, particularly CD4+, CD8+ or NK effector T cells. Methods for determining a change in the activity of effector T cells are well known and are as described earlier.

The antibody preferably causes an increase in activity in a CD8+ T cell *in vitro*, optionally wherein said increase in activity is an increase in proliferation, IFN- γ production and/or IL-2 production by the T cell. The increase is preferably at least 2-fold, more preferably at least 10-fold and even more preferably at least 25-fold higher than the change in activity caused by an isotype control antibody measured in the same assay.

As outlined above, antibody polypeptides which are capable of inhibiting the binding of one or more reference antibodies to human CD137 are provided. The reference antibodies described herein are reference antibody 1630/1631 and reference antibody 2674/2675.

By reference antibody "1630/1631" we mean an intact IgG antibody comprising heavy and light chains having the amino acid sequences of SEQ ID NOS: 17 and 18, respectively.

1630/1631- Full sequence Heavy chain

EVQLLES GGGLVQPGGSLRLSCAASGFTFGYSYMSWVRQAPGKGLEWVSSIGSGSSY
TYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARVYSSPGIDYWGGQGLVT
VSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA
VLQSSGLYSLSSVVTVPSSSLGTKTYTCNV DHKPSNTKVDKRVESKYGP PCPPCPAPE
FLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVSQEDPEVQFNWYVDGVEVHNAKTKP
REEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYT
LPQSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS
RLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLSLGK

[SEQ ID NO: 17]

1630/1631 - Full sequence Light chain

DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVP
SRFSGSGSGTDFTLTISSLQPEDFATYYCQQYYTWVPFTFGQGTKLEIKRTVAAPSVFIF
5 PPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSTSL
SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

[SEQ ID NO: 18]

By reference antibody “2674/2675” we mean an intact IgG antibody comprising heavy and
10 light chains having the amino acid sequences of SEQ ID NOS: 29 and 30, respectively.

2674/2675 – Full sequence heavy chain

EVQLLESGGGLVQPGGSLRLSCAASGPNFGYSYMSWVRQAPGKGLEWVSSIGSTSSH
15 TYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARVYSSPGIDYWGQGTLLV
VSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA
VLQSSGLYSLSSVTVPSSSLGKTYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPE
FLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKP
REEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVY
20 LPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS
RLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLGLK

[SEQ ID NO: 29]

2674/2675 – Full sequence light chain

25 DIQMTQSPSSLSASVGDRVTITCRASQSIGSTLNWYQQKPGKAPKLLIYGASSLQSGVP
SRFSGSGSGTDFTLTISSLQPEDFATYYCQQYYTWVPFTFGQGTKLEIKRTVAAPSVFIF
PPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSTSL
SSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

30 [SEQ ID NO: 30]

As discussed below, the reference antibody ‘1630/1631’ binds to domain 2 of CD137.
Reference antibody 2674/2675 also binds to domain 2 of CD137. Thus, it will be
appreciated that the antibody or an antigen-binding fragment of the invention also binds to
35 domain 2 of CD137.

By “capable of inhibiting the binding of reference antibody ‘1630/1631’ to human CD137” we mean that the presence of the antibody polypeptides of the invention inhibits, in whole or in part, the binding of ‘1630/1631’ to human CD137. Similarly, by “capable of inhibiting the binding of reference antibody ‘2674/2675’ to human CD137” we mean that the presence of the antibody polypeptides of the invention inhibits, in whole or in part, the binding of ‘2674/2675’ to human CD137. Such competitive binding inhibition can be determined using assays and methods well known in the art, for example using BIAcore chips with immobilised CD137 and incubating with the reference antibody ‘1630/1631’ or ‘2674/2675’ with and without an antibody polypeptide to be tested. Alternatively, a pairwise mapping approach can be used, in which the reference antibody ‘1630/1631’ or ‘2674/2675’ is immobilised to the surface of the BIAcore chip, CD137 antigen is bound to the immobilised antibody, and then a second antibody is tested for simultaneous CD137-binding ability (see ‘BIAcore Assay Handbook’, GE Healthcare Life Sciences, 29-0194-00 AA 05/2012; the disclosures of which are incorporated herein by reference).

15

In a further alternative, competitive binding inhibition can be determined using flow cytometry. For example, to test whether a test antibody is able to inhibit the binding of the 1630/1631 or 2674/2675 reference antibody to a cell surface antigen, cells expressing the antigen can be pre-incubated with the test antibody for 20 min before cells are washed and incubated with the reference 1630/1631 or 2674/2675 antibody conjugated to a fluorophore, which can be detected by flow cytometry. If the pre-incubation with the test antibody reduces the detection of the reference 1630/1631 or 2674/2675 antibody in flow cytometry, the test antibody inhibits the binding of the reference antibody to the cell surface antigen. If the antibody to be tested exhibits high affinity for CD137, then a reduced pre-incubation period may be used (or even no pre-incubation at all).

25

In a further alternative, competitive binding inhibition can be determined using an ELISA (e.g. as described in Example 8).

By “an antibody or an antigen-binding fragment thereof” we include substantially intact antibody molecules, as well as chimaeric antibodies, humanised antibodies, isolated human antibodies, single chain antibodies, bispecific antibodies, antibody heavy chains, antibody light chains, homodimers and heterodimers of antibody heavy and/or light chains, and antigen-binding fragments and derivatives of the same. Suitable antigen-binding fragments and derivatives include, but are not necessarily limited to, Fv fragments (e.g. single chain Fv and disulphide-bonded Fv), Fab-like fragments (e.g. Fab fragments,

35

Fab' fragments and F(ab)₂ fragments), single variable domains (e.g. V_H and V_L domains) and domain antibodies (dAbs, including single and dual formats [*i.e.* dAb-linker-dAb]). The potential advantages of using antibody fragments, rather than whole antibodies, are several-fold. The smaller size of the fragments may lead to improved pharmacological properties, such as better penetration of solid tissue. Moreover, antigen-binding fragments such as Fab, Fv, ScFv and dAb antibody fragments can be expressed in and secreted from *E. coli*, thus allowing the facile production of large amounts of the said fragments.

For example, the antigen-binding fragment may comprise an scFv molecule, *i.e.* wherein the V_H and V_L partner domains are linked via a flexible oligopeptide.

The phrase "an antibody or an antigen-binding fragment thereof" is also intended to encompass antibody mimics (for example, non-antibody scaffold structures that have a high degree of stability yet allow variability to be introduced at certain positions). Those skilled in the art of biochemistry will be familiar with many such molecules, as discussed in Gebauer & Skerra, 2009, *Curr Opin Chem Biol* **13**(3): 245–255 (the disclosures of which are incorporated herein by reference). Exemplary antibody mimics include: affibodies (also called Trinetins; Nygren, 2008, *FEBS J*, **275**, 2668-2676); CTLDs (also called Tetranectins; *Innovations Pharmac. Technol.* (2006), 27-30); adnectins (also called monobodies; *Meth. Mol. Biol.*, **352** (2007), 95-109); anticalins (*Drug Discovery Today* (2005), **10**, 23-33); DARPins (ankyrins; *Nat. Biotechnol.* (2004), **22**, 575-582); avimers (*Nat. Biotechnol.* (2005), **23**, 1556-1561); microbodies (*FEBS J*, (2007), **274**, 86-95); peptide aptamers (*Expert. Opin. Biol. Ther.* (2005), **5**, 783-797); Kunitz domains (*J. Pharmacol. Exp. Ther.* (2006) **318**, 803-809); affilins (*Trends. Biotechnol.* (2005), **23**, 514-522); affimers (Avacta Life Sciences, Wetherby, UK).

Persons skilled in the art will further appreciate that the invention also encompasses modified versions of antibodies and antigen-binding fragments thereof, whether existing now or in the future, e.g. modified by the covalent attachment of polyethylene glycol or another suitable polymer (see below).

Methods of generating antibodies and antibody fragments are well known in the art. For example, antibodies may be generated via any one of several methods which employ induction of *in vivo* production of antibody molecules, screening of immunoglobulin libraries (Orlandi. *et al*, 1989. *Proc. Natl. Acad. Sci. U.S.A.* **86**:3833-3837; Winter *et al.*, 1991, *Nature* **349**:293-299, the disclosures of which are incorporated herein by reference) or generation of monoclonal antibody molecules by cell lines in culture. These include, but

are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the Epstein-Barr virus (EBV)-hybridoma technique (Kohler *et al.*, 1975. *Nature* **256**:4950497; Kozbor *et al.*, 1985. *J. Immunol. Methods* **81**:31-42; Cote *et al.*, 1983. *Proc. Natl. Acad. Sci. USA* **80**:2026-2030; Cole *et al.*, 1984. *Mol. Cell. Biol.* **62**:109-120, the disclosures of which are incorporated herein by reference).

Suitable methods for the production of monoclonal antibodies are also disclosed in “*Monoclonal Antibodies: A manual of techniques*”, H Zola (CRC Press, 1988, the disclosures of which are incorporated herein by reference) and in “*Monoclonal Hybridoma Antibodies: Techniques and Applications*”, J G R Hurrell (CRC Press, 1982, the disclosures of which are incorporated herein by reference).

Likewise, antibody fragments can be obtained using methods well known in the art (see, for example, Harlow & Lane, 1988, “*Antibodies: A Laboratory Manual*”, Cold Spring Harbor Laboratory, New York, the disclosures of which are incorporated herein by reference). For example, antibody fragments according to the present invention can be prepared by proteolytic hydrolysis of the antibody or by expression in *E. coli* or mammalian cells (e.g. Chinese hamster ovary cell culture or other protein expression systems) of DNA encoding the fragment. Alternatively, antibody fragments can be obtained by pepsin or papain digestion of whole antibodies by conventional methods.

The antibodies of the invention are defined by reference to the variable regions of reference antibodies 1630/1631 and 2674/2675.

The reference antibody designated ‘1630/1631’ comprises:

(a) a heavy chain variable region having the amino acid sequence of SEQ ID NO: 1:

EVQLLES G GGLVQP G GSLRLSCAASGFTFGYSYMSWVRQAPGKGLEWVSSIGSGS
SYTTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARVYSSPGIDYWGQ
GTLVTVSS

[SEQ ID NO:1]

and

(b) a light chain variable region having the amino acid sequence of SEQ ID NO: 2:

DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQ
SGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTWVPFTFGQGTKLEIK

5 [SEQ ID NO:2]

The reference antibody designated '2674/2675' comprises:

(a) a heavy chain variable region having the amino acid sequence of SEQ ID NO: 19:

10

EVQLLES GGGLVQPGGSLRLSCAASG FNF GYSYMSWVRQAPGKGLEWVSSIGSTS
SHTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARVYSSPGIDYWGQ
GTLVTVSS

[SEQ ID NO:19]

15 and

(b) a light chain variable region having the amino acid sequence of SEQ ID NO: 20:

20

DIQMTQSPSSLSASVGDRVTITCRASQSIGSTLNWYQQKPGKAPKLLIYGASSLQ
SGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTWVPFTFGQGTKLEIK

[SEQ ID NO:20]

25 The term "amino acid" as used herein includes the standard twenty genetically-encoded amino acids and their corresponding stereoisomers in the 'D' form (as compared to the natural 'L' form), omega-amino acids and other naturally-occurring amino acids, unconventional amino acids (e.g. α,α -disubstituted amino acids, N-alkyl amino acids, etc.) and chemically derivatised amino acids (see below).

30 When an amino acid is being specifically enumerated, such as "alanine" or "Ala" or "A", the term refers to both L-alanine and D-alanine unless explicitly stated otherwise. Other unconventional amino acids may also be suitable components for polypeptides of the present invention, as long as the desired functional property is retained by the polypeptide. For the peptides shown, each encoded amino acid residue, where appropriate, is
35 represented by a single letter designation, corresponding to the trivial name of the conventional amino acid.

In one embodiment, the antibody polypeptides as defined herein comprise or consist of L-amino acids.

5 It will be appreciated by persons skilled in the art that the binding specificity of an antibody or antigen-binding fragment thereof is conferred by the presence of complementarity determining regions (CDRs) within the variable regions of the constituent heavy and light chains, such as those CDRs described herein.

10 It will be further appreciated by persons skilled in the art that any intact IgG antibody comprising the above variable regions may be used as the reference antibody to identify antibody polypeptides of the invention that competitively inhibit 1630/1631 or 2674/2675 binding to CD137. Preferably however, reference antibody 1630/1631 consists of heavy and light chains as defined in SEQ ID NOs:17 and 18, respectively, and reference antibody
15 2674/2675 consists of heavy and light chains as defined in SEQ ID NOs:29 and 30, respectively

Competitive binding typically arises because the test antibody binds at, or at least very close to, the epitope on the antigen to which binds the reference antibody (in this case,
20 1630/1631 or 2674/2675). However, it will be appreciated by persons skilled in the art that competitive binding may also arise by virtue of steric interference; thus, the test antibody may bind at an epitope different from that to which the reference antibody binds but may still be of sufficient size or configuration to hinder the binding of the reference antibody to the antigen.

25

The antibodies and antigen-binding fragments of the present invention were identified after screening of anti-CD137 antibodies, on the basis of exhibiting properties that make them particularly suitable as diagnostic and therapeutic agents for cancer.

30 Thus, in one embodiment, the antibody or antigen-binding fragment exhibits one or more of the following properties:

- a) the ability to stimulate CD137 and activate T cells and other immune cells via a cross-linking dependent mechanism (e.g. to induce release of interferon-gamma from CD8+ T cells; see Examples); and/or
- 35 b) cross-reactivity with cynomolgus CD137 (see Examples).

For example, the antibody or antigen-binding fragment may exhibit both of the above properties.

5 As described above, the antibodies of the invention may have a cross linking dependent mechanism. By “cross linking dependent mechanism”, we include an Fc cross linking dependent mechanism wherein the antibody has to bind both CD137 and an Fc receptor in order to stimulate CD137. As such, the antibody has to be capable of binding both CD137 and an Fc receptor.

10

In a preferred embodiment, the Fc receptor that is targeted is an FcγR. Examples of FcγRs include, FcγRI, FcγRIIA and FcγRIIB. Thus, in one embodiment, the FcγR may be FcγRIIA. By FcγRIIA, we include both the R131 and H131 allotypes of FcγRIIA. Thus, in one embodiment, the FcγR to be targeted is the R131 allotype of FcγRIIA.

15

In an alternative embodiment, the antibody could be Fc crosslinking independent, such that it can stimulate CD137 in the absence of binding to an Fc receptor.

Thus, exemplary antibodies 2674/2675 and 1630/1631 are FcγR-crosslinking dependent
20 agonistic antibodies targeting the co-stimulatory CD137 receptor. They are therefore only active in tissues or tumours containing cells expressing CD137 and FcγR. By “tumours containing cells expressing CD137 and FcγR” we include tumours or tumour draining lymph nodes comprising tumour cells and/or tumour infiltrating immune cells (such as monocytes, macrophages, dendritic cells, NK cells, T cells, B cells and granulocytes)
25 expressing CD137 and FcγR. It will be appreciated that CD137 and FcγR may be expressed on separate cells within the tumour and/or co-expressed in the same cells. Reference antibodies 2674/2675 and 1630/1631 will thus provide a tumour directed immune activation in indications associated with cells that express both CD137 and FcγR in the tumour micro environment; this contrasts with FcγR *independent* CD137 agonists
30 (e.g. Urelumab), which capable of inducing systemic immune activation. The tumour localizing effect of antibodies 2674/2675 and 1630/1631 will primarily depend on the number of tumour infiltrating macrophages/myeloid cells expressing different FcγRs.

It is known that IgG4 binds with high affinity to FcγRI and with moderate/low affinity to
35 FcγRIIa and FcγRIIb. FcγRI and FcγRIIa are expressed on monocytes and FcγRIIb is expressed with a high density on B cells. Crosslinking of antibodies 2674/2675 and 1630/1631 will preferentially occur intratumorally as well as in adjacent draining lymph

nodes. Systemically in the blood, where serum IgG levels are high, the availability of free non-blocked FcγRs are believed to be too low for an effective crosslinking to occur. Therefore, the risk for a systemic immune activation of is believed to be low which improves the risk-benefit profile compared to other CD137 mAbs.

5

Patient selection and a biomarker rationale for treatment with antibodies of the invention, such as 2674/2675 and 1630/1631, may be guided by tumour types that have infiltrating cells expressing CD137 and FcγRs. Thus, the antibodies of the invention may be for use in patients selected on the basis of having a tumour containing cells expressing CD137
10 and FcγRs (*i.e.* as a companion diagnostic test).

By “infiltrating cells” we include tumour infiltrating immune cells such as monocytes, macrophages, dendritic cells, NK cells, T cells, B cells and granulocytes

15 Advantageously, the antibody or antigen-binding fragment is capable of inducing tumour immunity. Tumour immunity can be demonstrated using methods well known in the art, for example by re-challenging mice that have been cured from a given tumour by CD137 antibody treatment with the same tumour. If tumour immunity has been induced by the antibody therapy, then the tumour is rejected upon re-challenge.

20

In one embodiment, the antibody or antigen binding fragment substantially incapable of inducing the following upon binding to cells expressing CD137:

- a) antibody-dependent cellular cytotoxicity (ADCC);
- 25 b) antibody-dependent cellular phagocytosis (ADCP); and/or
- c) complement-dependent cytotoxicity (CDC).

In one embodiment, the antibody or antigen-binding fragment is capable of binding to an epitope on the extracellular domain of CD137 which overlaps, at least in part, with the
30 epitope on CD137 to which reference antibody 1630/1631 and/or 2674/2675 is capable of binding. Thus, the antibody or antigen-binding fragment may be capable of binding to an epitope located at/within domain 2 of CD137.

In one embodiment, the antibody polypeptide of the invention comprises or consists of an
35 intact antibody (such as an IgG1 or IgG4 antibody). In a preferred embodiment, the antibody is an IgG4 antibody.

In an alternative embodiment, the antibody polypeptide of the invention comprises or consists of an antigen-binding fragment selected from the group consisting of Fv fragments (e.g. single chain Fv and disulphide-bonded Fv), Fab-like fragments (e.g. Fab fragments, Fab' fragments and F(ab)₂ fragments) and domain antibodies (e.g. single V_H variable domains or V_L variable domains). In particular, the antibody polypeptide may be a scFv.

In a further embodiment, as discussed above, the polypeptide of the invention comprises or consists of an antibody mimic selected from the group comprising or consisting of affibodies, tetranectins (CTLDS), adnectins (monobodies), anticalins, DARPin (ankyrins), avimers, iMabs, microbodies, peptide aptamers, Kunitz domains and affilins.

In one embodiment, the antibody or antigen binding fragment thereof according to the first or second aspect of the invention comprises:

- a) a heavy chain CDR1 sequence with the consensus sequence G, F, T/N, F, G, Y, S, Y;
- b) a heavy chain CDR2 sequence with the consensus sequence I, G, S, G/T, S, S, Y/H, T; and
- c) a heavy chain CDR3 sequence with the sequence ARVYSSPGIDY.

In one embodiment, the antibody or antigen binding fragment thereof comprises:

- a) a light chain CDR1 sequence with the consensus sequence Q, S, I, S/G, S, Y/T;
- b) a light chain CDR2 sequence with the consensus sequence A/G, A, S; and
- c) a light chain CDR3 sequence with the sequence QQYYTWVPFT.

In a preferred embodiment, the antibody or antigen-binding fragment thereof according to the first aspect of the invention comprises a heavy chain variable region comprising the following CDRs:

- a) GFTFGYSY [SEQ ID NO: 3] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 3, for example 1, 2 or 3 mutations;

- b) IGSGSSYT [SEQ ID NO: 4] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 4, for example 1, 2 or 3 mutations; and
- c) ARVYSSPGIDY [SEQ ID NO: 5] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 5, for example 1, 2 or 3 mutations.

Thus, the antibody or antigen-binding fragment thereof may comprise a heavy chain variable region comprising the CDRs of SEQ ID NOs 3, 4 and 5.

For example, the antibody or antigen-binding fragment thereof may comprise a heavy chain variable region having the amino acid sequence of the corresponding region of the 1630/1631 reference antibody, *i.e.* SEQ ID NO:1.

In an alternative preferred embodiment, the antibody or antigen-binding fragment thereof according to the first or second aspect of the invention comprises a heavy chain variable region comprising the following CDRs:

- a) GFNFGYSY [SEQ ID NO: 21] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 21, for example 1, 2 or 3 mutations;
- b) IGSTSSHT [SEQ ID NO: 22] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 22, for example 1, 2 or 3 mutations; and
- c) ARVYSSPGIDY [SEQ ID NO: 23] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 23, for example 1, 2 or 3 mutations.

Thus, the antibody or antigen-binding fragment thereof may comprise a heavy chain variable region comprising the CDRs of SEQ ID NOs 21, 22 and 23.

For example, the antibody or antigen-binding fragment thereof may comprise a heavy chain variable region having the amino acid sequence of the corresponding region of the 2674/2675 reference antibody, *i.e.* SEQ ID NO:19.

However, it will be appreciated (in relation to either embodiment, 1630/1631 or 2674/2675) that a low level of mutation (typically, just one, two or three amino acids) within a CDR sequence may be tolerated without loss of the specificity of the antibody or antigen-binding fragment for CD137.

For example, in an alternative embodiment, the antibody or antigen-binding fragment thereof may comprise a heavy chain variable region comprising the CDRs as defined above, wherein the H1 and H2 CDRs are mutated versions of SEQ ID NO: 3 and 4, respectively, and wherein the H3 CDR is SEQ ID NO: 5.

In a further alternative embodiment, the antibody or antigen-binding fragment thereof may comprise a heavy chain variable region comprising the CDRs as defined above, wherein the H1 and H2 CDRs are mutated versions of SEQ ID NO: 21 and 22, respectively, and wherein the H3 CDR is SEQ ID NO: 23.

Percent identity can be determined by, for example, the LALIGN program (Huang and Miller, *Adv. Appl. Math.* (1991) **12**:337-357, the disclosures of which are incorporated herein by reference) at the Expasy facility site (http://www.ch.embnet.org/software/LALIGN_form.html) using as parameters the global alignment option, scoring matrix BLOSUM62, opening gap penalty -14, extending gap penalty -4. Alternatively, the percent sequence identity between two polypeptides may be determined using suitable computer programs, for example the GAP program of the University of Wisconsin Genetic Computing Group and it will be appreciated that percent identity is calculated in relation to polypeptides whose sequence has been aligned optimally.

The alignment may alternatively be carried out using the Clustal W program (as described in Thompson *et al.*, 1994, *Nucl. Acid Res.* **22**:4673-4680, which is incorporated herein by reference). The parameters used may be as follows:

- Fast pair-wise alignment parameters: K-tuple(word) size; 1, window size; 5, gap penalty; 3, number of top diagonals; 5. Scoring method: x percent.
- Multiple alignment parameters: gap open penalty; 10, gap extension penalty; 0.05.
- Scoring matrix: BLOSUM.

Alternatively, the BESTFIT program may be used to determine local sequence alignments.

In a further preferred embodiment, the antibody or antigen-binding fragment thereof according to the first aspect of the invention comprises a light chain variable region comprising the following CDRs:

- 5 a) QSISSY [SEQ ID NO: 6] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 6, for example 1, 2 or 3 mutations;
- b) AAS [SEQ ID NO: 7] or an amino acid sequence containing up to 2 amino acid mutations compared to SEQ ID NO: 7; for example 1 or 2 mutations and
- 10 c) QQYYTWVPFT [SEQ ID NO: 8] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 8, for example 1, 2 or 3 mutations.

Thus, the antibody polypeptide may comprise a light chain variable region comprising the CDRs of SEQ ID NOs 6, 7 and 8.

15

For example, the antibody or antigen-binding fragment thereof may comprise a light chain variable region having the amino acid sequence of the corresponding region of the 1630/1631 reference antibody, *i.e.* SEQ ID NO: 2.

- 20 In an alternative embodiment, the antibody or antigen-binding fragment thereof may comprise a light chain variable region comprising the CDRs as defined above, wherein the L1 and L2 CDRs are mutated versions of SEQ ID NO: 6 and 7, respectively, and wherein the L3 CDR is SEQ ID NO:8.

- 25 In a further preferred embodiment, the antibody or antigen-binding fragment thereof according to the first or second aspect of the invention comprises a light chain variable region comprising the following CDRs:

- 30 a) QSIGST [SEQ ID NO: 24] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 24, for example 1, 2 or 3 mutations;
- b) GAS [SEQ ID NO: 25] or an amino acid sequence containing up to 2 amino acid mutations compared to SEQ ID NO: 25; for example 1 or 2 mutations and
- 35 c) QQYYTWVPFT [SEQ ID NO: 26] or an amino acid sequence containing up to 3 amino acid mutations compared to SEQ ID NO: 26, for example 1, 2 or 3 mutations.

Thus, the antibody polypeptide may comprise a light chain variable region comprising the CDRs of SEQ ID NOs 24, 25 and 26.

For example, the antibody or antigen-binding fragment thereof may comprise a light chain variable region having the amino acid sequence of the corresponding region of the 1630/1631 reference antibody, *i.e.* SEQ ID NO: 20.

In an alternative embodiment, the antibody or antigen-binding fragment thereof may comprise a light chain variable region comprising the CDRs as defined above, wherein the L1 and L2 CDRs are mutated versions of SEQ ID NO: 24 and 25, respectively, and wherein the L3 CDR is SEQ ID NO: 26.

It will be appreciated by persons skilled in the art that for human therapy, human or humanised antibodies are preferably used. Humanised forms of non-human (*e.g.* murine) antibodies are genetically engineered chimaeric antibodies or antibody fragments having preferably minimal-portions derived from non-human antibodies. Humanised antibodies include antibodies in which complementary determining regions of a human antibody (recipient antibody) are replaced by residues from a complementary determining region of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired functionality. In some instances, Fv framework residues of the human antibody are replaced by corresponding non-human residues. Humanised antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported complementarity determining region or framework sequences. In general, the humanised antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the complementarity determining regions correspond to those of a non-human antibody and all, or substantially all, of the framework regions correspond to those of a relevant human consensus sequence. Humanised antibodies optimally also include at least a portion of an antibody constant region, such as an Fc region, typically derived from a human antibody (see, for example, Jones *et al.*, 1986, *Nature* **321**:522-525; Riechmann *et al.*, 1988, *Nature* **332**:323-329; Presta, 1992, *Curr. Op. Struct. Biol.* **2**:593-596, the disclosures of which are incorporated herein by reference).

Methods for humanising non-human antibodies are well known in the art. Generally, the humanised antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues, often referred to as imported residues, are typically taken from an imported variable domain. Humanisation can be essentially performed as described (see, for example, Jones *et al.*, 1986, *Nature* **321**:522-

525; Reichmann *et al.*, 1988. *Nature* **332**:323-327; Verhoeyen *et al.*, 1988, *Science* **239**:1534-1536; US 4,816,567, the disclosures of which are incorporated herein by reference) by substituting human complementarity determining regions with corresponding rodent complementarity determining regions. Accordingly, such humanised antibodies are
5 chimeric antibodies, wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanised antibodies may be typically human antibodies in which some complementarity determining region residues and possibly some framework residues are substituted by residues from analogous sites in rodent antibodies. Chimeric antibodies are discussed by
10 Neuberger *et al* (1998, *8th International Biotechnology Symposium* Part 2, 792-799).

Human antibodies can also be identified using various techniques known in the art, including phage display libraries (see, for example, Hoogenboom & Winter, 1991, *J. Mol. Biol.* **227**:381; Marks *et al.*, 1991, *J. Mol. Biol.* **222**:581; Cole *et al.*, 1985, In: *Monoclonal antibodies and Cancer Therapy*, Alan R. Liss, pp. 77; Boerner *et al.*, 1991. *J. Immunol.* **147**:86-95, the disclosures of which are incorporated herein by reference).

It will be appreciated by persons skilled in the art that humanised antibodies or antigen-binding fragments of the invention may further comprise a heavy chain constant region, or
20 part thereof (see below).

In one embodiment, the antibody polypeptide comprises a CH1, CH2 and/or CH3 region of an IgG heavy chain (such as an IgG1, IgG2, IgG3 or IgG4 heavy chain). Thus, the antibody polypeptide may comprise part or all of the constant regions from an IgG4 heavy
25 chain. For example, the antibody polypeptide may be a Fab fragment comprising CH1 and CL constant regions, combined with any of the above-defined heavy and light variable regions respectively.

Likewise, the above-defined antibodies or antigen-binding fragments of the invention may
30 further comprise a light chain constant region, or part thereof (see below). For example, the antibody polypeptide may comprise a CL region from a kappa or lambda light chain.

In one embodiment, the antibodies or antigen-binding fragments of the invention comprise an antibody Fc-region. It will be appreciated by a skilled person that the Fc portion may
35 be from an IgG antibody, or from a different class of antibody (such as IgM, IgA, IgD or

IgE). In one embodiment, the Fc region is from an IgG1, IgG2, IgG3 or IgG4 antibody. Advantageously, however, the Fc region is from an IgG4 antibody.

5 The Fc region may be naturally-occurring (e.g. part of an endogenously produced antibody) or may be artificial (e.g. comprising one or more point mutations relative to a naturally-occurring Fc region). A variant of an Fc region typically binds to Fc receptors, such as FcγR and/or neonatal Fc receptor (FcRn) with altered affinity providing for improved function and/or half-life of the polypeptide. The biological function and/ or the half-life may be either increased or a decreased relative to the half-life of a polypeptide
10 comprising a native Fc region. Examples of such biological functions which may be modulated by the presence of a variant Fc region include antibody dependent cell cytotoxicity (ADCC), antibody-dependent cellular phagocytosis (ADCP), complement-dependent cytotoxicity (CDC), and/or apoptosis.

15 Thus, the Fc region may be naturally-occurring (e.g. part of an endogenously produced human antibody) or may be artificial (e.g. comprising one or more point mutations relative to a naturally-occurring human Fc region).

As is well documented in the art, the Fc region of an antibody mediates its serum half-life
20 and effector functions, such as complement-dependent cytotoxicity (CDC), antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cell phagocytosis (ADCP).

Engineering the Fc region of a therapeutic monoclonal antibody or Fc fusion protein allows
25 the generation of molecules that are better suited to the pharmacology activity required of them (Strohl, 2009, *Curr Opin Biotechnol* **20**(6):685-91, the disclosures of which are incorporated herein by reference).

(a) Engineered Fc regions for increased half-life

30 One approach to improve the efficacy of a therapeutic antibody is to increase its serum persistence, thereby allowing higher circulating levels, less frequent administration and reduced doses.

35 The half-life of an IgG depends on its pH-dependent binding to the neonatal receptor FcRn. FcRn, which is expressed on the surface of endothelial cells, binds the IgG in a pH-dependent manner and protects it from degradation.

Some antibodies that selectively bind the FcRn at pH 6.0, but not pH 7.4, exhibit a higher half-life in a variety of animal models.

5 Several mutations located at the interface between the CH2 and CH3 domains, such as T250Q/M428L (Hinton *et al.*, 2004, *J Biol Chem.* **279**(8):6213-6, the disclosures of which are incorporated herein by reference) and M252Y/S254T/T256E + H433K/N434F (Vaccaro *et al.*, 2005, *Nat. Biotechnol.* **23**(10):1283-8, the disclosures of which are incorporated herein by reference), have been shown to increase the binding affinity to
10 FcRn and the half-life of IgG1 *in vivo*.

(b) Engineered Fc regions for altered effector function

Depending on the therapeutic antibody or Fc fusion protein application, it may be desired
15 to either reduce or increase the effector function (such as ADCC).

For antibodies that target cell-surface molecules, especially those on immune cells, abrogating effector functions may be required for certain clinical indications.

20 The four human IgG isotypes bind the activating Fcγ receptors (FcγRI, FcγRIIa, FcγRIIIa), the inhibitory FcγRIIb receptor, and the first component of complement (C1q) with different affinities, yielding very different effector functions (Bruhns *et al.*, 2009, *Blood.* **113**(16):3716-25, the disclosures of which are incorporated herein by reference).

25 FcγRI binding affinity of IgG4 vs IgG2

Bruhns *et al* performed a series of experiments that evaluated the specificity and affinity of the known human FcγRs, and their polymorphic variants, for the different human IgG subclasses (Bruhns *et al.*, 2009, *Blood.* **113**(16):3716-25, the disclosures of which are
30 incorporated herein by reference). In this study, it was clearly demonstrated that while IgG2 had no detectable affinity for FcγRI, IgG1, IgG3 and IgG4 all displayed a binding affinity for FcγRI in the nanomolar range (Bruhns *et al.*, 2009, *Blood.* **113**(16):3716-25, Lu *et al.*, 2015, *Proc Natl Acad Sci U S A.* **112**(3):833-8, the disclosures of which are incorporated herein by reference). A summary of the relative binding affinities between
35 the major human FcγRs and their variants and IgG isotypes is summarized in Table A. (Stewart *et al.* 2014, *J Immunother.* **2**(29), the disclosures of which are incorporated herein by reference)

Table A. Binding affinity between human FcγRs and IgG isotypes.

FcγR	IgG1	IgG2	IgG4
FcγRI	++++	-	++++
FcγRIIA H131	+++	++	++
FcγRIIA R131	+++	+	++
FcγRIIB	++	-	++
FcγRIIA V158	+++	+	++
FcγRIIA F158	++	-	++

5 However, cellular activation influences the affinity of FcγRI for IgG immune complexes and the data generated by surface plasmon resonance in the Bruhns paper may not correctly reproduce what occurs at an inflammatory site. A review paper by Hogarth et al (Hogarth *et al.* 2012, *Nat Rev Drug Discov* 11(4):311-31, the disclosures of which are incorporated herein by reference) summarizes this as well as other studies focusing on FcγR binding
10 for IgG.

FcγRI expression on myeloid cell subsets

Human FcγRs are primarily expressed by cells of the myeloid lineage, which has been
15 demonstrated in numerous studies for circulating myeloid cell subsets. Classical monocytes, generally identified as CD14⁺ CD16⁻ display high levels of FcγRII (CD32), intermediate levels of FcγRI and low levels of FcγRIII (CD16) (Almeida *et al.* 2001, **100**(3):325-38, Cheeseman *et al.* 2016, PLoS One **11**(5):e0154656, the disclosures of which are incorporated herein by reference). CD14⁻ CD16⁺ non-classical monocytes,
20 however, display high levels of FcγRIII, intermediate levels of FcγRII and low levels of FcγRI (Almeida *et al.* 2001). A summary and compilation of several published microarray data sets showing the expression of human FcγR genes on different myeloid cell subsets confirms these observations (Guilliams *et al.* 2014, *Nat Rev Immunol.* **14**(2):94-108, the disclosures of which are incorporated herein by reference).

25 Once within tissues, monocytes differentiate towards macrophages and, depending on environmental cues, these macrophages obtain specific phenotypes. In a study by Roussel et al (Roussel *et al.* 2017, *J Leukoc Biol.* **102**(2):437-447, the disclosures of which are incorporated herein by reference), peripheral blood monocytes were polarized towards
30 different macrophage lineages by using various inflammatory stimuli and the expression

profile of these cells evaluated. Here, IFN- γ stimulated monocytes resulted in a highly elevated expression specifically of CD64. A similar observation was made in SLE patients where increased CD64 expression was detected on circulating CD14⁺ monocytes, which correlated with expression of interferon-stimulated genes (Li *et al.* 2010, Arthritis Res Ther 12(3): R90, the disclosures of which are incorporated herein by reference).

Myeloid cell infiltration within various human tumors

Various myeloid cell subsets such as inflammatory monocytes, monocytic myeloid-derived suppressor cells (MDSC) and macrophages have, in numerous studies, been shown to accumulate in cancer patients (Solito *et al.* 2014, Ann N Y Acad Sci 1319:47-65., Hu *et al.* 2016, Clin Transl Oncol.18(3):251-8, the disclosures of which are incorporated herein by reference). Although recent attempts have aimed at proposing strategies to standardize the characterization of these cells (Bronte *et al.* 2016, Nat Commun. 7:12150, the disclosures of which are incorporated herein by reference), many phenotypic definitions of these cell populations can still be found throughout the literature (Elliott *et al.* 2017, Front Immunol. 8:86, the disclosures of which are incorporated herein by reference). Most commonly, these cells are defined by the expression of the markers CD11b, CD14, CD33 and the low expression of HLA-DR (monocytic MDSC) (Bronte *et al.* 2016). Additionally, tumor-associated macrophages (TAM) are commonly identified by the expression of CD64 and CD68 (M1-polarized, anti-tumorigenic), or CD163 and CD206 (M2-polarized, pro-tumorigenic) (Elliott *et al.* 2017).

A recent review by Elliott *et al* (referenced above) summarizes the numerous phenotypes used to identify myeloid cell subsets in cancer patients. Most of these studies have focused their analyses on circulating cells and increased frequencies of myeloid CD11b⁺ cells have been observed in the blood of patients with e.g. bladder, breast, colorectal, hepatocellular, pancreatic, prostate and renal cell carcinoma (Solito *et al.* 2014, Elliott *et al.* 2017). Other studies have also attempted to characterize the level of infiltration of these cells into tumor tissue. In colorectal tumors, a high frequency of CD14⁺ CD169⁺ cells was observed. These cells also expressed CD163 and CD206 and were thus suggested to be M2-polarized TAM (Li *et al.* 2015, PLoS One 10(10):e0141817, the disclosures of which are incorporated herein by reference). Another study in colorectal cancer patients also detected increased numbers of CD11b⁺ CD33⁺ HLA-DR⁻ cells, compared to healthy individuals (Zhang *et al.* 2013, PLoS One 8(2):e57114, the disclosures of which are incorporated herein by reference).

Similarly, CD11b⁺ myeloid cells were also identified in bladder tumors, where they accounted for 10-20% of all nucleated cells (Eruslanov *et al.* 2012, *Int J Cancer* **130**(5):1109-19, the disclosures of which are incorporated herein by reference). An even higher frequency of CD11b⁺ cells was observed in pancreatic cancer where over 60% of the CD45⁺ cells were CD11b⁺ CD15⁺ CD33⁺ (Porembka *et al.* 2012, *Cancer Immunol Immunother* **61**(9):1373-85, the disclosures of which are incorporated herein by reference). Also, one study concluded that the major myeloid cell population within non-small cell lung carcinoma is a CD11b⁺ CD15⁺ CD66b⁺ neutrophil-like population. Interestingly, once these cells migrate from blood to the tumor tissue, these cells display an altered expression profile, including upregulated FcγRI (Eruslanov *et al.* 2014, *J Clin Invest.* **124**(12):5466-80, the disclosures of which are incorporated herein by reference).

FcγRI expression on tumor-infiltrating cells

Although numerous studies have identified a high infiltration of myeloid cells within human tumors, no study has thoroughly explored the expression of FcγRs on these cells in detail. Several publications have, however, demonstrated the presence of FcγRI-expressing cells within tumor tissue.

A study by Morimura *et al.* (Morimura *et al.* 1990, *Acta Neuropathol.* **80**(3):287-94, the disclosures of which are incorporated herein by reference) evaluated gliomas from 12 human samples by immunocytochemistry and compared these to peritumoral control tissue. This study demonstrated a high presence of macrophages (using the marker CD163, RM3/1) in gliomas, compared to peritumoral tissue, as well as an increase in FcγRI and FcγRII (CD32). A more recent study by Griesinger *et al.* (Griesinger *et al.* 2013, *J Immunol.* **191**(9):4880-8, the disclosures of which are incorporated herein by reference) confirmed these observations by performing flow cytometric analyses of various pediatric brain tumor types. Here, a high frequency of CD45⁺ CD11b⁺ myeloid cells was observed for tissues from pilocytic astrocytoma and ependymoma patients. These cells also expressed high levels of FcγRI.

In addition to brain tumors, FcγRI expression has also been shown for other types of tumors. Grugan *et al.* (Grugan *et al.* 2012, *J Immunol.* **189**(11):5457-66, the disclosures of which are incorporated herein by reference) demonstrated the presence of CD11b⁺ CD14⁺ cells within human breast tumor tissue. These cells were shown to express high levels of FcγRI and FcγRIIa, as well as FcγRIIb and FcγRIII. Also, CD45⁺ CD11b⁺ CD14⁺ CD68⁺ TAM were identified in gastrointestinal stromal tumors displaying expression of FcγRI

(Cavnar *et al.* 2013, *J Exp Med.* 210(13):2873-86, the disclosures of which are incorporated herein by reference). CD45⁺ CD11b⁺ FcγRI⁺ cells were also identified in colorectal cancer patients and these cells displayed a higher expression of FcγRI in tumor tissue, compared to healthy control tissue (Norton *et al.* 2016, *Clin Transl Immunology.* 5(4):e76, the disclosures of which are incorporated herein by reference). FcγRI expression has also been demonstrated for melanoma metastases (Hansen *et al.* 2006, *Acta Oncol* 45(4):400-5, the disclosures of which are incorporated herein by reference).

Binding of IgG to the FcγRs or C1q depends on residues located in the hinge region and the CH2 domain. Two regions of the CH2 domain are critical for FcγRs and C1q binding, and have unique sequences in IgG2 and IgG4. Substitutions into human IgG1 of IgG2 residues at positions 233-236 and IgG4 residues at positions 327, 330 and 331 were shown to greatly reduce ADCC and CDC (Armour *et al.*, 1999, *Eur J Immunol.* 29(8):2613-24; Shields *et al.*, 2001, *J Biol Chem.* 276(9):6591-604, the disclosures of which are incorporated herein by reference). Furthermore, Idusogie *et al.* demonstrated that alanine substitution at different positions, including K322, significantly reduced complement activation (Idusogie *et al.*, 2000, *J Immunol.* 164(8):4178-84, the disclosures of which are incorporated herein by reference). Similarly, mutations in the CH2 domain of murine IgG2A were shown to reduce the binding to FcγRI, and C1q (Steurer. *et al.*, 1995. *J Immunol.* 155(3):1165- 74, the disclosures of which are incorporated herein by reference).

Numerous mutations have been made in the CH2 domain of human IgG1 and their effect on ADCC and CDC tested in vitro (see references cited above). Notably, alanine substitution at position 333 was reported to increase both ADCC and CDC (Shields *et al.*, 2001, *supra*; Steurer *et al.*, 1995, *supra*). Lazar *et al.* described a triple mutant (S239D/I332E/A330L) with a higher affinity for FcγRIIIa and a lower affinity for FcγRIIb resulting in enhanced ADCC (Lazar *et al.*, 2006, *PNAS* 103(11):4005–4010, the disclosures of which are incorporated herein by reference). The same mutations were used to generate an antibody with increased ADCC (Ryan *et al.*, 2007, *Mol. Cancer Ther.* 6:3009-3018, the disclosures of which are incorporated herein by reference). Richards *et al.* studied a slightly different triple mutant (S239D/I332E/G236A) with improved FcγRIIIa affinity and FcγRIIa/FcγRIIb ratio that mediates enhanced phagocytosis of target cells by macrophages (Richards *et al.*, 2008. *Mol Cancer Ther.* 7(8):2517-27, the disclosures of which are incorporated herein by reference).

Due to their lack of effector functions, IgG4 antibodies represent a preferred IgG subclass for receptor modulation without cell depletion. IgG4 molecules can exchange half-molecules in a dynamic process termed Fab-arm exchange. This phenomenon can also occur *in vivo* between therapeutic antibodies and endogenous IgG4.

5

The S228P mutation has been shown to prevent this recombination process allowing the design of less unpredictable therapeutic IgG4 antibodies (Labrijn *et al.*, 2009, *Nat Biotechnol.* **27**(8):767-71, the disclosures of which are incorporated herein by reference).

10 In a further embodiment, the effector function of the Fc region may be altered through modification of the carbohydrate moieties within the CH2 domain therein, for example by modifying the relative levels of fucose, galactose, bisecting N-acetylglucosamine and/or sialic acid during production (see Jefferis, 2009, *Nat Rev Drug Discov.* **8**(3):226-34 and Raju, 2008, *Curr Opin Immunol.*, **20**(4):471-8; the disclosures of which are incorporated
15 herein by reference)

Thus, it is known that therapeutic antibodies lacking or low in fucose residues in the Fc region may exhibit enhanced ADCC activity in humans (for example, see Peipp *et al.*, 2008, *Blood* **112**(6):2390-9, Yamane-Ohnuki & Satoh, 2009, *MAbs* **1**(3):230-26, Iida *et al.*,
20 2009, *BMC Cancer* **9**:58 (the disclosures of which are incorporated herein by reference). Low fucose antibody polypeptides may be produced by expression in cells cultured in a medium containing an inhibitor of mannosidase, such as kifunensine (see Example I below).

25 Other methods to modify glycosylation of an antibody into a low fucose format include the use of the bacterial enzyme GDP-6-deoxy-D-lyxo-4-hexulose reductase in cells not able to metabolise rhamnose (e.g. using the *GlymaxX*® technology of ProBioGen AG, Berlin, Germany).

30 Another method to create low fucose antibodies is by inhibition or depletion of alpha-(1,6)-fucosyltransferase in the antibody-producing cells (e.g. using the *Potelligent*® CHOK1SV technology of Lonza Ltd, Basel, Switzerland).

An exemplary heavy chain constant region amino acid sequence which may be combined
35 with any VH region sequence disclosed herein (to form a complete heavy chain) is the IgG1 heavy chain constant region sequence reproduced here:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA
VLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP
PCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV
EVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTIS
5 KAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
[SEQ ID NO: 12]

Other heavy chain constant region sequences are known in the art and could also be
10 combined with any VH region disclosed herein. For example, as indicated above, a
preferred constant region is a modified IgG4 constant region such as that reproduced here:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA
15 VLQSSGLYSLSSVVTVPSSSLGTQTYTCNVDHKPSNTKVDKRVESKYGPPCPPCP
APEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVSQEDPEVQFNWYVDGVEVH
NAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAK
GQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTP
PVLDSGDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLSLGK
20 [SEQ ID NO: 13]

This modified IgG4 sequence results in stabilization of the core hinge of IgG4 making the
IgG4 more stable, preventing Fab arm exchange.

25 Another preferred constant region is a modified IgG4 constant region such as that
reproduced here:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA
VLQSSGLYSLSSVVTVPSSSLGTQTYTCNVDHKPSNTKVDKRVESKYGPPCPPCP
30 APEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVSQEDPEVQFNWYVDGVEVH
NAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAK
GQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTP
PVLDSGDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNRYTQKSLSLSLGK
[SEQ ID NO: 14]

35

This modified IgG4 sequence exhibits reduced FcRn binding and hence results in a reduced serum half-life relative to wild type IgG4. In addition, it exhibits stabilization of the core hinge of IgG4 making the IgG4 more stable, preventing Fab arm exchange.

- 5 Also suitable for use in the polypeptides of the invention is a wild type IgG4 constant region such as that reproduced here:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA
VLQSSGLYSLSSVVTVPSSSLGTQKTYTCNVDPKPSNTKVDKRVERESKYGPPCPSCP
10 APEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVSQEDPEVQFNWYVDGVEVH
NAKTKPREEQFNSTYRVVSVLTVQLHQLDNLNGKEYKCKVSNKGLPSSIEKTISKAK
GQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP
PVLDSGDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQKSLSLGLGK

[SEQ ID NO: 15]

15

An exemplary light chain constant region amino acid sequence which may be combined with any VL region sequence disclosed herein (to form a complete light chain) is the kappa chain constant region sequence reproduced here:

20 RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES
VTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

[SEQ ID NO: 16]

25 Other light chain constant region sequences are known in the art and could also be combined with any VL region disclosed herein.

In an exemplary embodiment of the invention, the antibody polypeptide may comprise the IgG4 constant regions of SEQ ID NOs: 13 and 16, respectively.

30 Thus, exemplary antibody polypeptides of the invention comprise:

(a) a heavy chain comprising a variable region of SEQ ID NO: 1 together with a constant region of SEQ ID NO: 13; and

35 (b) a light chain comprising a variable region of SEQ ID NO: 2 together with a constant region of SEQ ID NO: 16.

For example, the antibody polypeptides may be an intact IgG4 molecule comprising or consisting of two heavy chains having an amino acid sequence of SEQ ID NO: 17 and two light chains having an amino acid sequence of SEQ ID NO: 18.

5

Alternative exemplary polypeptides of the invention comprise:

(a) a heavy chain comprising a variable region of SEQ ID NO: 19 together with a constant region of SEQ ID NO: 13; and

10

(b) a light chain comprising a variable region of SEQ ID NO: 20 together with a constant region of SEQ ID NO: 16.

15

For example, the antibody polypeptides may be an intact IgG4 molecule comprising or consisting of two heavy chains having an amino acid sequence of SEQ ID NO: 29 and two light chains having an amino acid sequence of SEQ ID NO: 30.

In one embodiment of the first or second aspect of the invention, the antibody polypeptide of the invention is or comprises a "fusion" polypeptide.

20

In addition to being fused to a moiety in order to improve pharmacokinetic properties, it will be appreciated that the polypeptide of the invention may also be fused to a polypeptide such as glutathione-S-transferase (GST) or protein A in order to facilitate purification of said polypeptide. Examples of such fusions are well known to those skilled in the art.

25

Similarly, the said polypeptide may be fused to an oligo-histidine tag, such as His6, or to an epitope recognised by an antibody such as the well-known Myc tag epitope. Fusions to any variant or derivative of said polypeptide are also included in the scope of the invention. It will be appreciated that fusions (or variants, derivatives or fusions thereof) which retain or improve desirable properties, such as IL-1R binding properties or *in vivo* half-life are preferred.

30

Thus, the fusion may comprise an amino acid sequence as detailed above together with a further portion which confers a desirable feature on the said polypeptide of the invention; for example, the portion may be useful in detecting or isolating the polypeptide, or promoting cellular uptake of the polypeptide. The portion may be, for example, a biotin moiety, a radioactive moiety, a fluorescent moiety, for example a small fluorophore or a green fluorescent protein (GFP) fluorophore, as well known to those skilled in the art. The moiety

35

may be an immunogenic tag, for example a Myc tag, as known to those skilled in the art or may be a lipophilic molecule or polypeptide domain that is capable of promoting cellular uptake of the polypeptide, as known to those skilled in the art.

- 5 It will be appreciated by persons skilled in the art that the antibody polypeptides of the invention may comprise or consist of one or more amino acids which have been modified or derivatised.

Chemical derivatives of one or more amino acids may be achieved by reaction with a
10 functional side group. Such derivatised molecules include, for example, those molecules in which free amino groups have been derivatised to form amine hydrochlorides, *p*-toluene sulphonyl groups, carboxybenzoxy groups, *t*-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatised to form salts, methyl and ethyl esters or other types of esters and hydrazides. Free hydroxyl groups may be derivatised
15 to form O-acyl or O-alkyl derivatives. Also included as chemical derivatives are those peptides which contain naturally occurring amino acid derivatives of the twenty standard amino acids. For example: 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine and ornithine for lysine. Derivatives
20 also include peptides containing one or more additions or deletions as long as the requisite activity is maintained. Other included modifications are amidation, amino terminal acylation (e.g. acetylation or thioglycolic acid amidation), terminal carboxylamidation (e.g. with ammonia or methylamine), and the like terminal modifications.

- 25 It will be further appreciated by persons skilled in the art that peptidomimetic compounds may also be useful. The term 'peptidomimetic' refers to a compound that mimics the conformation and desirable features of a particular peptide as a therapeutic agent.

For example, the said polypeptide includes not only molecules in which amino acid
30 residues are joined by peptide (-CO-NH-) linkages but also molecules in which the peptide bond is reversed. Such retro-inverso peptidomimetics may be made using methods known in the art, for example such as those described in Meziere *et al.* (1997) *J. Immunol.* 159, 3230-3237, which is incorporated herein by reference. This approach involves making pseudo-peptides containing changes involving the backbone, and not the orientation of
35 side chains. Retro-inverse peptides, which contain NH-CO bonds instead of CO-NH peptide bonds, are much more resistant to proteolysis. Alternatively, the said polypeptide

may be a peptidomimetic compound wherein one or more of the amino acid residues are linked by a $-y(\text{CH}_2\text{NH})-$ bond in place of the conventional amide linkage.

5 In a further alternative, the peptide bond may be dispensed with altogether provided that an appropriate linker moiety which retains the spacing between the carbon atoms of the amino acid residues is used; it may be advantageous for the linker moiety to have substantially the same charge distribution and substantially the same planarity as a peptide bond.

10 It will also be appreciated that the said polypeptide may conveniently be blocked at its N- or C-terminus so as to help reduce susceptibility to exo-proteolytic digestion.

A variety of un-coded or modified amino acids such as D-amino acids and N-methyl amino acids have also been used to modify mammalian peptides. In addition, a presumed
15 bioactive conformation may be stabilised by a covalent modification, such as cyclisation or by incorporation of lactam or other types of bridges, for example see Veber *et al.*, 1978, *Proc. Natl. Acad. Sci. USA* 75:2636 and Thursell *et al.*, 1983, *Biochem. Biophys. Res. Comm.* 111:166, which are incorporated herein by reference.

20 Typically, the antibody polypeptide of the invention will be a 'naked' antibody polypeptide, *i.e.* without any additional functional moieties such as cytotoxic or detectable moieties. For example, where the therapeutic effect is mediated by a direct effect of the antibody of the invention on immune cells, *e.g.* to reduce inflammation, it may be advantageous for the antibody to lack any cytotoxic activity.

25 However, in alternative embodiment, the antibody polypeptides of the invention may be augmented with a functional moiety to facilitate their intended use, for example as a diagnostic (*e.g. in vivo* imaging) agent or therapeutic agent. Thus, in one embodiment, the antibody polypeptide is linked, directly or indirectly, to a therapeutic moiety. A suitable
30 therapeutic moiety is one that is capable of reducing or inhibiting the growth, or in particular killing, a cancer cell (or associated stem cells or progenitor cells). For example, the therapeutic agent may be a cytotoxic moiety, such as a radioisotope (*e.g.* ^{90}Y , ^{177}Lu , $^{99}\text{Tc}^{\text{m}}$, *etc*) or cytotoxic drug (*e.g.* antimetabolites, toxins, cytostatic drugs, *etc*).

35 Alternatively, the cytotoxic moiety may comprise or consist of one or more moieties suitable for use in activation therapy, such as photon activation therapy, neutron activation

therapy, neutron-induced Auger electron therapy, synchrotron irradiation therapy or low energy X-ray photon activation therapy.

Optionally, the antibody polypeptide of the invention may further comprise a detectable moiety. For example, a detectable moiety may comprise or consist of a radioisotope, such as a radioisotope selected from the group consisting of ^{99m}Tc , ^{111}In , ^{67}Ga , ^{68}Ga , ^{72}As , ^{89}Zr , ^{123}I and ^{201}Tl . Optionally, the agent may comprise a pair of detectable and cytotoxic radionuclides, such as $^{86}\text{Y}/^{90}\text{Y}$ or $^{124}\text{I}/^{211}\text{At}$. Alternatively, the antibody polypeptide may comprise a radioisotope that is capable of simultaneously acting in a multi-modal manner as a detectable moiety and also as a cytotoxic moiety to provide so-called "Multimodality theragnostics". The binding moieties may thus be coupled to nanoparticles that have the capability of multi-imaging (for example, SPECT, PET, MRI, Optical, or Ultrasound) together with therapeutic capability using cytotoxic drugs, such as radionuclides or chemotherapy agents.

15

Therapeutic and/or detectable moieties (such as a radioisotope, cytotoxic moiety or the like) may be linked directly, or indirectly, to the antibody or fragment thereof. Suitable linkers are known in the art and include, for example, prosthetic groups, non-phenolic linkers (derivatives of N-succinimidyl- benzoates; dodecaborate), chelating moieties of both macrocyclics and acyclic chelators, such as derivatives of 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA), deferoxamine (DFO), derivatives of diethylenetriaminepentaacetic acid (DTPA), derivatives of S-2-(4-Isothiocyanatobenzyl)-1,4,7-triazacyclononane-1,4,7-triacetic acid (NOTA) and derivatives of 1,4,8,11-tetraazacyclododecan-1,4,8,11-tetraacetic acid (TETA), derivatives of 3,6,9,15-Tetraazabicyclo[9.3.1]-pentadeca-1(15),11,13-triene-4-(S)-(4-isothiocyanatobenzyl)-3,6,9-triacetic acid (PCTA), derivatives of 5-S-(4-Aminobenzyl)-1-oxa-4,7,10-triazacyclododecane-4,7,10-tris(acetic acid) (DO3A) and other chelating moieties.

25

One preferred linker is DTPA, for example as used in ^{177}Lu -DTPA-[antibody polypeptide of the invention]. A further preferred linker is deferoxamine, DFO, for example as used in ^{89}Zr -DFO-[antibody polypeptide of the invention].

30

However, it will be appreciated by persons skilled in the art that many medical uses of the antibody polypeptides of the invention will not require the presence of a cytotoxic or diagnostic moiety.

35

As discussed above, methods for the production of antibody polypeptides of the invention are well known in the art.

Conveniently, the antibody polypeptide is or comprises a recombinant polypeptide. Suitable methods for the production of such recombinant polypeptides are well known in the art, such as expression in prokaryotic or eukaryotic hosts cells (for example, see Green & Sambrook, 2012, *Molecular Cloning, A Laboratory Manual*, Fourth Edition, Cold Spring Harbor, New York, the relevant disclosures in which document are hereby incorporated by reference).

Although the antibody may be a polyclonal antibody, it is preferred if it is a monoclonal antibody, or that the antigen-binding fragment, variant, fusion or derivative thereof, is derived from a monoclonal antibody.

Suitable monoclonal antibodies may be prepared by known techniques, for example those disclosed in *"Monoclonal Antibodies; A manual of techniques"*, H Zola (CRC Press, 1988) and in *"Monoclonal Hybridoma Antibodies: Techniques and Application"*, SGR Hurrell (CRC Press, 1982). Polyclonal antibodies may be produced which are poly-specific or mono-specific. It is preferred that they are mono-specific.

Antibody polypeptides of the invention can also be produced using a commercially available *in vitro* translation system, such as rabbit reticulocyte lysate or wheatgerm lysate (available from Promega). Preferably, the translation system is rabbit reticulocyte lysate. Conveniently, the translation system may be coupled to a transcription system, such as the TNT transcription-translation system (Promega). This system has the advantage of producing suitable mRNA transcript from an encoding DNA polynucleotide in the same reaction as the translation.

It will be appreciated by persons skilled in the art that antibody polypeptides of the invention may alternatively be synthesised artificially, for example using well known liquid-phase or solid phase synthesis techniques (such as *t*-Boc or Fmoc solid-phase peptide synthesis).

A third aspect of the invention provides an isolated nucleic acid molecule encoding an antibody or antigen-binding fragment of the first or second aspect of the invention, or a component polypeptide chain thereof. By "nucleic acid molecule" we include DNA (e.g. genomic DNA or complementary DNA) and mRNA molecules, which may be single- or

double-stranded. By “isolated” we mean that the nucleic acid molecule is not located or otherwise provided within a cell.

In one embodiment, the nucleic acid molecule is a cDNA molecule.

5

Preferably, the nucleic acid molecule comprises one or more nucleotide sequence selected from either SEQ ID NO: 9 and SEQ ID NO: 10, reproduced below.

Nucleotide sequence encoding VH region of “1630”

10

GAGGTGCAGCTGTTGGAGAGCGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGC
GCCTCTCCTGTGCAGCCAGCGGATTCACCTTTGGTTACTCTTACATGTCTTGGGT
CCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTCTCATCTATTGGTTCTGGTTCT
TCTTACACATACTATGCAGACTCCGTGAAGGGCCGGTTCACCATCTCCCGTGACA
ATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGC
TGTATATTATTGTGCGCGCGTTTACTCTTCTCCGGGTATTGACTATTGGGGCCAG
GGAACCCTGGTCACCGTCTCCTCA [SEQ ID NO:9]

15

Nucleotide sequence encoding VL region of “1631”

20

GACATCCAGATGACCCAGTCTCCATCCTCCCTGAGCGCATCTGTAGGAGACCGCG
TCACCATCACTTGCCGGGCAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCA
GCAGAAACCAGGGAAAGCCCCTAAGCTCCTGATCTATGCTGCATCCAGTTTGCAA
AGTGGGGTCCCATCACGTTTCAGTGGCAGTGGAAGCGGGACAGATTTCACTCTCA
CCATCAGCAGTCTGCAACCTGAAGATTTTGCAACTTATTACTGTCAACAGTACTA
CACTTGGGTTCCGTTCACTTTTGGCCAGGGGACCAAGCTGGAGATCAAA

25

[SEQ ID NO:10]

In an alternative preferred embodiment, the nucleic acid molecule comprises one or more nucleotide sequence selected from either SEQ ID NO: 27 and SEQ ID NO: 28, reproduced below.

30

Nucleotide sequence encoding VH region of "2674"

gaggtgcagttgttggaaatctggcggaggattggtgcagcctggcggatctctgagactgtcttgt
gccgcctctggcttcaacttcggctactcctacatgtcctgggtccgacaggctcctggcaaagga
5 ctggaatgggtgtcctccatcggctccaccagctctcacacctaactacgccgattccgtgaagggc
agattcaccatcagccgggacaactccaagaacaccctgtacctgcagatgaactccctgagagcc
gaggacaccgccgtgtactactgtgccagagtgtactcctctcctggcatcgattattggggccag
ggcacactggtcaccgtgtcctctgtcttctaccaagggaccctctgtgttccctctggctccttgc
tccagatccacctctgagtctaccgtgtctctgggtgcctgggtcaaggattactttcctgagcct
10 gtgaccgtgtcttggaaactccgggtgtctgtacatccggcgtgcacacatttccagctgtgtctgag
tcctccggcctgtactctctgtcctctgtcgtgaccgtgccttctagctctctgggcaccaagacc
tacacctgtaacgtggaccacaagccttccaacaccaaggtggacaagcgcgtggaatctaagtac
ggcctccatgtccaccatgtcctgtctcagaattcctcggcggaccaagcgtgttccctgtttcct
ccaaagcctaaggacacctgatgatctctcggacccctgaagtgacctgcgtgggtggatgtg
15 tctcaagaggaccagaagtgcagttcaattggtagctggacggcgtggaagtgcacaacgccaag
accaagcctagagaggaacagttcaactccacctacagagtgggtgtccgtgctgaccgtgtgtcac
caggattgggtgaacggcaaagagtacaagtgaaggtgtccaacaagggcctgccttccagcatc
gaaaagaccatctccaaggctaagggccagcctcgggaacctcaggtttacacctgcctccaagc
caagaggaaatgaccaagaaccaggtgtccttgacctgcctcgtgaagggattctaccttccgat
20 atcgccgtggaatgggagtctaacggccagccagagaacaactacaagacaacccctcctgtgtgtg
gactccgacggctctttcttctctgtattctcgctgacctggacaagtctcggtggcaagagggc
aacgtgttctcctgctctgtgatgcacgaggccctgcacaaccactacacacagaagtccctgtct
ctgtccctgggcaag

[SEQ ID NO: 27]

25

Nucleotide sequence encoding VL region of "2675"

gacatccagatgaccagctctccatcctctctgtctgcctctgtgggcgacagagtgaccatcacc
tgtcgggcttctcagtcctcggcagcaccctgaactgggtatcagcagaagcctggcaaggccct
30 aagctgtgtgatctatggcgctagctctctgcagctctggcgtgcctctagattttccggctctggc
tctggcacccgacttcacctgacaatcagttccctgcagcctgaggacttcgccacctactactgc
cagcagtactacacctgggtgccctttacctttggccagggcaccaagctggaaatcaagagaacc
gtggcgcgtccttccgtgttcatcttccaccatctgacgagcagctgaagtccggcacagcttct
gtcgtgtgcctgctgaacaacttctacctcgggaagccaaggtgcagtggaaggtggacaatgcc
35 ctgcagtccggcaactccaagagtctgtgaccgagcaggactccaaggactctacctacagcctg
tcctccacactgacctgtctaaggccgactacgagaagcacaaggtgtacgcctgcgaagtgacc
catcagggactgtctagccccgtgaccaagtccttcaacagagggcaggtgt

[SEQ ID NO: 28]

It will be appreciated by persons skilled in the art that the nucleic acid molecule may be codon-optimised for expression of the antibody polypeptide in a particular host cell, e.g. for expression in human cells (for example, see Angov, 2011, *Biotechnol. J.* **6**(6):650-659, the disclosures of which are incorporated herein by reference).

5

Also included within the scope of the invention are the following:

10

(a) a fourth aspect of the invention provides a vector (such as an expression vector) comprising a nucleic acid molecule according to the third aspect of the invention;

(b) a fifth aspect of the invention provides a host cell (such as a mammalian cell, e.g. human cell, or Chinese hamster ovary cell, e.g. CHOK1SV cells) comprising a nucleic acid molecule according to the third aspect of the invention or a vector according to the fourth aspect of the invention; and

15

(c) a sixth aspect of the invention provides a method of making an antibody polypeptide according to the first or second aspect of the invention comprising culturing a population of host cells according to the fifth aspect of the invention under conditions in which said polypeptide is expressed, and isolating the polypeptide therefrom.

20

A seventh aspect of the invention provides a pharmaceutical composition comprising a pharmaceutically effective amount of an antibody or antigen-binding fragment according to the first or second aspect of the invention and a pharmaceutically-acceptable diluent, carrier, adjuvant or excipient.

25

It will be appreciated by persons skilled in the art that additional compounds may also be included in the pharmaceutical compositions, including, chelating agents such as EDTA, citrate, EGTA or glutathione.

30

The pharmaceutical compositions may be prepared in a manner known in the art that is sufficiently storage stable and suitable for administration to humans and animals. For example, the pharmaceutical compositions may be lyophilised, e.g. through freeze drying, spray drying, spray cooling, or through use of particle formation from supercritical particle formation.

35

By "pharmaceutically acceptable" we mean a non-toxic material that does not decrease the effectiveness of the CD137-binding activity of the antibody polypeptide of the invention.

Such pharmaceutically acceptable buffers, carriers or excipients are well-known in the art (see Remington's Pharmaceutical Sciences, 18th edition, A.R Gennaro, Ed., Mack Publishing Company (1990) and handbook of Pharmaceutical Excipients, 3rd edition, A. Kibbe, Ed., Pharmaceutical Press (2000), the disclosures of which are incorporated herein
5 by reference).

The term "buffer" is intended to mean an aqueous solution containing an acid-base mixture with the purpose of stabilising pH. Examples of buffers are Trizma, Bicine, Tricine, MOPS, MOPSO, MOBS, Tris, Hepes, HEPBS, MES, phosphate, carbonate, acetate, citrate,
10 glycolate, lactate, borate, ACES, ADA, tartrate, AMP, AMPD, AMPSO, BES, CABS, cacodylate, CHES, DIPSO, EPPS, ethanolamine, glycine, HEPPSO, imidazole, imidazolelactic acid, PIPES, SSC, SSPE, POPSO, TAPS, TABS, TAPSO and TES.

The term "diluent" is intended to mean an aqueous or non-aqueous solution with the
15 purpose of diluting the antibody polypeptide in the pharmaceutical preparation. The diluent may be one or more of saline, water, polyethylene glycol, propylene glycol, ethanol or oils (such as safflower oil, corn oil, peanut oil, cottonseed oil or sesame oil).

The term "adjuvant" is intended to mean any compound added to the formulation to
20 increase the biological effect of the antibody polypeptide of the invention. The adjuvant may be one or more of zinc, copper or silver salts with different anions, for example, but not limited to fluoride, chloride, bromide, iodide, thiocyanate, sulfite, hydroxide, phosphate, carbonate, lactate, glycolate, citrate, borate, tartrate, and acetates of different acyl composition. The adjuvant may also be cationic polymers such as cationic cellulose ethers,
25 cationic cellulose esters, deacetylated hyaluronic acid, chitosan, cationic dendrimers, cationic synthetic polymers such as poly(vinyl imidazole), and cationic polypeptides such as polyhistidine, polylysine, polyarginine, and peptides containing these amino acids.

The excipient may be one or more of carbohydrates, polymers, lipids and minerals.
30 Examples of carbohydrates include lactose, glucose, sucrose, mannitol, and cyclodextrines, which are added to the composition, e.g. for facilitating lyophilisation. Examples of polymers are starch, cellulose ethers, cellulose carboxymethylcellulose, hydroxypropylmethyl cellulose, hydroxyethyl cellulose, ethylhydroxyethyl cellulose, alginates, carageenans, hyaluronic acid and derivatives thereof, polyacrylic acid,
35 polysulphonate, polyethyleneglycol/polyethylene oxide, polyethyleneoxide/polypropylene oxide copolymers, polyvinylalcohol/polyvinylacetate of different degree of hydrolysis, and polyvinylpyrrolidone, all of different molecular weight, which are added to the composition,

e.g., for viscosity control, for achieving bioadhesion, or for protecting the lipid from chemical and proteolytic degradation. Examples of lipids are fatty acids, phospholipids, mono-, di-, and triglycerides, ceramides, sphingolipids and glycolipids, all of different acyl chain length and saturation, egg lecithin, soy lecithin, hydrogenated egg and soy lecithin, which are added to the composition for reasons similar to those for polymers. Examples of minerals are talc, magnesium oxide, zinc oxide and titanium oxide, which are added to the composition to obtain benefits such as reduction of liquid accumulation or advantageous pigment properties.

The antibody polypeptides of the invention may be formulated into any type of pharmaceutical composition known in the art to be suitable for the delivery thereof.

In one embodiment, the pharmaceutical compositions of the invention may be in the form of a liposome, in which the antibody polypeptide is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids, which exist in aggregated forms as micelles, insoluble monolayers and liquid crystals. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Suitable lipids also include the lipids above modified by poly(ethylene glycol) in the polar headgroup for prolonging bloodstream circulation time. Preparation of such liposomal formulations is can be found in for example US 4,235,871, the disclosures of which are incorporated herein by reference.

The pharmaceutical compositions of the invention may also be in the form of biodegradable microspheres. Aliphatic polyesters, such as poly(lactic acid) (PLA), poly(glycolic acid) (PGA), copolymers of PLA and PGA (PLGA) or poly(caprolactone) (PCL), and polyanhydrides have been widely used as biodegradable polymers in the production of microspheres. Preparations of such microspheres can be found in US 5,851,451 and in EP 0 213 303, the disclosures of which are incorporated herein by reference.

In a further embodiment, the pharmaceutical compositions of the invention are provided in the form of polymer gels, where polymers such as starch, cellulose ethers, cellulose carboxymethylcellulose, hydroxypropylmethyl cellulose, hydroxyethyl cellulose, ethylhydroxyethyl cellulose, alginates, carageenans, hyaluronic acid and derivatives thereof, polyacrylic acid, polyvinyl imidazole, polysulphonate, polyethylene-glycol/polyethylene oxide, polyethyleneoxide/polypropylene oxide copolymers,

polyvinylalcohol/polyvinylacetate of different degree of hydrolysis, and polyvinylpyrrolidone are used for thickening of the solution containing the agent. The polymers may also comprise gelatin or collagen.

- 5 Alternatively, the antibody polypeptide may simply be dissolved in saline, water, polyethylene glycol, propylene glycol, ethanol or oils (such as safflower oil, corn oil, peanut oil, cottonseed oil or sesame oil), tragacanth gum, and/or various buffers.

10 It will be appreciated that the pharmaceutical compositions of the invention may include ions and a defined pH for potentiation of action of the active antibody polypeptide. Additionally, the compositions may be subjected to conventional pharmaceutical operations such as sterilisation and/or may contain conventional adjuvants such as preservatives, stabilisers, wetting agents, emulsifiers, buffers, fillers, *etc.*

15 The pharmaceutical compositions according to the invention may be administered via any suitable route known to those skilled in the art. Thus, possible routes of administration include parenteral (intravenous, subcutaneous, and intramuscular), topical, ocular, nasal, pulmonar, buccal, oral, parenteral, vaginal and rectal. Also, administration from implants is possible.

20 In one preferred embodiment, the pharmaceutical compositions are administered parenterally, for example, intravenously, intracerebroventricularly, intraarticularly, intra-arterially, intraperitoneally, intrathecally, intraventricularly, intrasternally, intracranially, intramuscularly or subcutaneously, or they may be administered by infusion techniques.

25 They are conveniently used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The aqueous solutions should be suitably buffered (preferably to a pH of from 3 to 9), if necessary. The preparation of suitable parenteral formulations under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to
30 those skilled in the art.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and
35 aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried

(lyophilised) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

5

Thus, the pharmaceutical compositions of the invention are particularly suitable for parenteral, e.g. intravenous, administration.

Alternatively, the pharmaceutical compositions may be administered intranasally or by
10 inhalation (for example, in the form of an aerosol spray presentation from a pressurised container, pump, spray or nebuliser with the use of a suitable propellant, such as dichlorodifluoromethane, trichlorofluoro-methane, dichlorotetrafluoro-ethane, a hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane (HFA 134A3 or 1,1,1,2,3,3,3-heptafluoropropane (HFA 227EA3), carbon dioxide or other suitable gas). In the case of
15 a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurised container, pump, spray or nebuliser may contain a solution or suspension of the active polypeptide, e.g. using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g. sorbitan trioleate. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or
20 insufflator may be formulated to contain a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

Aerosol or dry powder formulations are preferably arranged so that each metered dose or 'puff' contains at least 1 mg of a compound of the invention for delivery to the patient. It
25 will be appreciated that the overall daily dose with an aerosol will vary from patient to patient, and may be administered in a single dose or, more usually, in divided doses throughout the day.

Alternatively, the antibody polypeptides of the invention can be administered in the form of
30 a suppository or pessary, or they may be applied topically in the form of a lotion, solution, cream, ointment or dusting powder. The compounds of the invention may also be transdermally administered, for example, by the use of a skin patch. They may also be administered by the ocular route.

35 For ophthalmic use, the antibody polypeptides of the invention can be formulated as micronised suspensions in isotonic, pH adjusted, sterile saline, or, preferably, as solutions in isotonic, pH adjusted, sterile saline, optionally in combination with a preservative such

as a benzylalkonium chloride. Alternatively, they may be formulated in an ointment such as petrolatum.

For application topically to the skin, the antibody polypeptide of the invention can be formulated as a suitable ointment containing the active compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, they can be formulated as a suitable lotion or cream, suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, a polyethylene glycol, liquid paraffin, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The pharmaceutical compositions will be administered to a patient in a pharmaceutically effective dose. A 'therapeutically effective amount', or 'effective amount', or 'therapeutically effective', as used herein, refers to that amount which provides a therapeutic effect for a given condition and administration regimen. This is a predetermined quantity of active material calculated to produce a desired therapeutic effect in association with the required additive and diluent, *i.e.* a carrier or administration vehicle. Further, it is intended to mean an amount sufficient to reduce and most preferably prevent, a clinically significant deficit in the activity, function and response of the host. Alternatively, a therapeutically effective amount is sufficient to cause an improvement in a clinically significant condition in a host. As is appreciated by those skilled in the art, the amount of a compound may vary depending on its specific activity. Suitable dosage amounts may contain a predetermined quantity of active composition calculated to produce the desired therapeutic effect in association with the required diluent. In the methods and use for manufacture of compositions of the invention, a therapeutically effective amount of the active component is provided. A therapeutically effective amount can be determined by the ordinary skilled medical or veterinary worker based on patient characteristics, such as age, weight, sex, condition, complications, other diseases, *etc.*, as is well known in the art. The administration of the pharmaceutically effective dose can be carried out both by single administration in the form of an individual dose unit or else several smaller dose units and also by multiple administrations of subdivided doses at specific intervals. Alternatively, the dose may be provided as a continuous infusion over a prolonged period.

In the context of diagnostic use of the antibody polypeptides of the invention, a 'pharmaceutically effective amount', or 'effective amount', or 'diagnostically effective', as

used herein, refers to that amount which provides a detectable signal for diagnosis, *e.g.* for *in vivo* imaging purposes.

5 The antibody polypeptides can be formulated at various concentrations, depending on the efficacy/toxicity of the polypeptide being used. For example, the formulation may comprise the active antibody polypeptide at a concentration of between 0.1 μM and 1 mM, more preferably between 1 μM and 500 μM , between 500 μM and 1 mM, between 300 μM and 700 μM , between 1 μM and 100 μM , between 100 μM and 200 μM , between 200 μM and 300 μM , between 300 μM and 400 μM , between 400 μM and 500 μM , between 500 μM and 600 μM , between 600 μM and 700 μM , between 800 μM and 900 μM or between 900 μM and 1 mM. Typically, the formulation comprises the active antibody polypeptide at a concentration of between 300 μM and 700 μM .

15 Typically, the therapeutic dose of the antibody polypeptide (with or without a therapeutic moiety) in a human patient will be in the range of 100 μg to 1 g per administration (based on a body weight of 70kg, *e.g.* between 300 μg to 700 mg per administration). For example, the maximum therapeutic dose may be in the range of 0.1 to 10 mg/kg per administration, *e.g.* between 0.1 and 5 mg/kg or between 1 and 5 mg/kg or between 0.1 and 2 mg/kg. It will be appreciated that such a dose may be administered at different intervals, as determined by the oncologist/physician; for example, a dose may be administered daily, twice-weekly, weekly, bi-weekly or monthly.

25 It will be further appreciated by persons skilled in the art that the polypeptides and pharmaceutical formulations of the present invention have utility in both the medical and veterinary fields. Thus, the methods of the invention may be used in the treatment of both human and non-human animals (such as horses, dogs and cats). Preferably, however, the patient is human.

30 For veterinary use, the agents, medicaments and pharmaceutical compositions of the invention are administered as a suitably acceptable formulation in accordance with normal veterinary practice and the veterinary surgeon will determine the dosing regimen and route of administration which will be most appropriate for a particular animal.

35 An eighth aspect of the invention provides an antibody or antigen-binding fragment thereof according to the first or second aspect of the invention for use in medicine.

In one embodiment, the antibody polypeptides and formulations of the invention may be used to treat patients or subjects who suffer from or are at risk of suffering from a cancer.

5 By 'treatment' we include both therapeutic and prophylactic treatment of the patient. The term 'prophylactic' is used to encompass the use of an agent, or formulation thereof, as described herein which either prevents or reduces the likelihood of a cancer, or the spread, dissemination, or metastasis of cancer cells in a patient or subject. The term 'prophylactic' also encompasses the use of an agent, or formulation thereof, as described herein to prevent recurrence of a cancer in a patient who has previously been treated for the
10 neoplastic disorder.

The cancer may be associated with formation of solid tumours or may be a haematologic cancer. Cancer types that may be treated include carcinomas, sarcomas, lymphomas, leukemias, blastomas and germ cell tumours.

15 For example, the antibody or antigen-binding fragment thereof may be for use in the treatment of a cancer selected from the group consisting of prostate cancer; breast cancer; colorectal cancer; kidney cancer; pancreatic cancer; ovarian cancer; lung cancer; cervical cancer; rhabdomyosarcoma; neuroblastoma; bone cancer; multiple myeloma; leukemia
20 (such as acute lymphoblastic leukemia [ALL] and acute myeloid leukemia [AML]), skin cancer (e.g. melanoma), bladder cancer and glioblastoma.

In one embodiment, the cancer may be selected from the list of cancers in Table 16 or Table 17.

25 Typically, the therapeutic agents of the invention will be administered in parenteral form, for example by injection into the bloodstream or at/near the site of a tumour.

In one embodiment, the agent for treating a patient who has been pre-screened and
30 identified as having a tumour with cells expressing CD137 and FcγR, such as FcγRI, FcγRIIA, FcγRIIB or combinations thereof.

Related aspects of the invention provide the following:

- (i) use of an antibody or antigen-binding fragment thereof according to the first or second aspect of the invention in the preparation of a medicament for treating cancer; and
- (ii) a method for treating an individual with cancer, the method comprising the step of administering to an individual in need thereof an effective amount of an antibody or antigen-binding fragment thereof according to the first or second aspect of the invention.

It will be further appreciated that the antibody-based agents of the invention may be used as a sole treatment for cancer in a patient or as part of a combination treatment (which further treatment may be a pharmaceutical agent, radiotherapy and/or surgery).

Thus, the patient may also receive one or more further treatments for cancer, for example pharmaceutical agents (such as chemotherapeutic agents), radiotherapy and/or surgery.

For example, the pharmaceutical compositions of the invention may be administered in combination with other therapeutic agents used in the treatment of cancers, such as antimetabolites, alkylating agents, anthracyclines and other cytotoxic antibiotics, vinca alkyloids, etoposide, platinum compounds, taxanes, topoisomerase I inhibitors, antiproliferative immunosuppressants, corticosteroids, sex hormones and hormone antagonists, and other therapeutic antibodies (such as trastuzumab).

In one embodiment, the one or more further treatments are selected from the group consisting of conventional chemotherapeutic agents (such as alkylating agents, antimetabolites, plant alkaloids and terpenoids, topoisomerase inhibitors and antineoplastics), radiotherapeutic agents, antibody-based therapeutic agents (such as gemtuzumab, alemtuzumab, rituximab, trastuzumab, nimotuzumab, cetuximab, bevacizumab), and steroids.

A ninth aspect of the invention provides a method of identifying a patient susceptible to treatment with an antibody according to the first or second aspect of the invention, comprising screening a patient to identify if they have a tumour with cells expressing CD137 and FcγR.

Optionally, the FcγR that is screened for is FcγRIIA. In one embodiment, the FcγRIIA is the R131 allotype.

5 Suitable biomarker screening methods are well known in the art. For example, a tumour biopsy sample may be taken from the patient and analysed to determine the level of expression (at an RNA level and/or protein level) of CD137 and/or FcγRs therein; for example using immunohistochemistry, flow cytometry or proteomic approaches.

10 Preferences and options for a given aspect, feature or parameter of the invention should, unless the context indicates otherwise, be regarded as having been disclosed in combination with any and all preferences and options for all other aspects, features and parameters of the invention.

15 The listing or discussion of an apparently prior-published document in this specification should not necessarily be taken as an acknowledgement that the document is part of the state of the art or is common general knowledge.

20 The use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.”

25 These, and other, embodiments of the invention will be better appreciated and understood when considered in conjunction with the above description and the accompanying drawings. It should be understood, however, that the above description, while indicating various embodiments of the invention and numerous specific details thereof, is given by way of illustration and not of limitation. Many substitutions, modifications, additions and/or rearrangements may be made within the scope of the invention without departing from the spirit thereof, and the invention includes all such substitutions, modifications, additions and/or rearrangements.

30 The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

35

Preferred, non-limiting examples which embody certain aspects of the invention will now be described, with reference to the following figures:

5 Figure 1 shows binding to CD137 human and cynomolgus CD137. Data from separate two experiments included.

Figure 2 shows the CD137 variants in Example 6.

10 Figure 3 shows a summary of two experiments of CD137 mAb competition with CD137L binding to CHO-huCD137 cells when titrated (from left to right 25 µg/ml).

Figure 4 shows the stimulation index of clones normalized to reference REF1.

15 Figure 5 shows the induction of NF-κB mediated signaling by the antibodies with and without cross-linking.

Figure 6 shows the effect of treatment with the 1630/1631 antibody on tumour volume in a mouse tumour model.

20 Figure 7 shows determined patch size and Aggscore for 2674/2675 as well as the parental clone 1630/1631 clone via Schrödinger analysis.

Figure 8 shows binding of 2674/2675 to human and cynomolgus CD137.

25 Figure 9 shows crosslinking of 2674/2675 and parental clone 1630/1631 with FcγRI transfected CHO cells in the CD137 reporter assay.

Figure 10 shows crosslinking of 2674/2675 and parental clone 1630/1631 with FcγRIIIa R131 and FcγRIIb in the CD137 reporter assay.

30 Figure 11 shows CHO cells transfected with empty vector (pcDNA3.1), used to determine crosslinking independent activation in the reporter cell line.

35 Figure 12 shows IFN-γ production of CD8⁺ T cells after stimulation with CD137 mAbs when crosslinked with FcγRI transfected CHO cells. Summary of IFN-γ response normalized against 2674/2675 agonist response in a CD8⁺ T cell agonist assay (n=5).

Figure 13 shows Dot plots showing the correlation between the mean expression values of Fcγ receptor (X axes) and TNFRSF9 (CD137, Y axes) for various human cancers. Cancers with an above average expression (mean expression level ≥ 10) of both Fcγ receptor and CD137 have been highlighted as clear symbols.

5

EXAMPLES

Example 1 - Selection of CD137 antibodies from Alligator GOLD

5 Phage display selections were performed using a human antibody (scFv) library, Alligator GOLD. Selections towards recombinant CD137 in soluble form, coated onto the surface of beads or tubes, or expressed on the surface of CD137-transfected cells were performed. CTLA4-Fc and an irrelevant His-tagged protein were used as non-targets included in excess in the selections. Prior to each selection round, the phage stocks were pre-selected
10 towards non-target proteins, beads or CD137 negative cells to remove unspecific binders.

To identify specific binders from the phage selection, approximately 4500 individual clones were screened in phage format using ELISA coated with either recombinant target (CD137-Fc) or non-target protein, followed by confirmation as soluble scFv for some
15 clones. Clones exhibiting specific binding to CD137 were sequenced and unique clones were produced as IgG for further characterization.

Example 2 - Binding to human CD137 measured by ELISA

20 Aim

The aim was to determine binding potency of the CD137 antibody.

Material and methods

Binding of CD137 antibodies to recombinant human CD137 was determined by sandwich
25 ELISA. Briefly, ELISA plates (Greiner # 655074) coated with recombinant human CD137-Fc (R&D # 838-4B) were incubated with serial dilutions of the various CD137 antibodies to be investigated. CD137 antibodies were detected using HRP-conjugated goat-anti-human kappa light chain (AbD Serotec # STAR127P) and developed with SuperSignal ELISA Pico Chemiluminescent substrate (Pierce # 37069). EC50 values of the various
30 antibodies were determined in 2-6 separate experiments.

Two different reference antibodies with specificity for CD137, synthesized from published amino acid sequence information, were used in this study (designated "REF1" and "REF2").
35

The other reference antibodies used, namely REF3, REF4 and REF8, are human CD137-specific monospecific IgG antibodies obtained from the *Alligator GOLD* library. They are

agonistic and stimulate T cells upon binding to CD137. The binding epitopes of the reference antibodies has been established as outlined in Examples 6-8 (see below).

5 The reference antibodies were selected because they have previously undergone at least some clinical testing and so represent the benchmark against which new anti-CD137 antibodies can be judged for improved properties and/or function.

Results and conclusion

10 Exemplary antibody 1630/1631 exhibits EC50 values in a similar range as those of the reference antibodies, *i.e.* sub nM. Data is summarized in **Table 1**.

Table 1 - EC50 values (nM) of CD137 antibodies determined by ELISA for human CD137.

Antibody	Mean	SD	n
REF1	0.75	0.137	8
REF2	0.33	0.069	5
REF3	0.39	0.037	3
REF4	0.41	0.050	4
REF8	0.38	0.137	2
1630/1631	0.27	0.078	4

n = number of data points.

15

Example 3 - Binding to human and cynomolgus CD137 measured by flow cytometry

Aim

20 The aim of this study was to determine the binding to human and cynomolgus (Macaca fascicularis) CD137.

Material and methods

25 Binding and EC50 was determined using flow cytometric of CHO cells transfected with human CD137, cynomolgus CD137 or empty vector. The extracellular part of human or cynomolgus CD137 was fused to the transmembrane and intracellular part of human CD40 and cloned into pcDNA3.1. The vector was subsequently stably transfected into CHO cells. Expression of CD137 was confirmed by flow cytometry using CD137 antibody (human CD137-PE, BD Biosciences # 555956) for 30 min at 4°C. CD137-transfected and empty vector-transfected cells were incubated with CD137 antibodies for at least 1h at 4°C to

saturate the binding. In order to minimize antibody internalization, 0.05% sodium azide was used in the incubation buffer and all work was performed on ice. The CD137 antibodies were detected using PE-conjugated anti-hlgG antibody (109-115-098, Jackson Immunoresearch laboratories), incubated for 30 min at 4°C. Directly after staining the cells were fixed with a paraformaldehyde solution (10x concentrate BD CellFIX, BD biosciences # 340181). Cells were analyzed by flow cytometry using FACSVerse (BD Biosciences). The median fluorescence intensity (MFI) for each sample was determined and the dose response data was analysed using Graph Pad Prism.

MFI data was normalized for each antibody, where 0% is defined as the lowest value and 100% is the highest value in the dose titration for each antibody. EC50 and 95% confidence interval were calculated with Graph Pad Prism based on data from the two experiments (non-linear regression (curve fit), constraints set to 0 and 100).

Results and conclusion

Binding to CHO-huCD137, CHO-cyCD137 and CHO-pcDNA was confirmed in two separate experiments (Figure 1). 1630/1631 binds to human CD137 with EC50 comparable with the two reference antibodies REF1 and REF2. 1630/1631 binds well to cynomolgus CD137. Reference antibody REF1 and REF8 (Figure 1) binds very weakly or not at all to cynomolgus CD137. REF8 exhibits weak binding and does not reach a complete saturation.

The EC50 determination is presented as 95% confidence intervals for each CD137 antibody tested in order to include the inter and intra assay variations (Table 2).

Table 2 - 95% confidence intervals for the EC50 of each CD137 antibody determined as an average from two experiments of normalized data. Nd: Not detectable

Antibody	Binding to human CD137, EC50 (µg/mL)	Binding to cyno CD137, EC50 (µg/mL)	Ratio, cyno:human
REF1	1.00 - 1.99	Nd	Nd
REF2	0.21 - 0.31	0.13 - 0.24	0.69
REF3	0.20 - 0.36	Nd	Nd
REF4	0.16 - 0.27	0.11 - 0.17	0.67
REF8	0.20 - 0.42	> 3	>14
1630/1631	0.17 - 0.26	0.12 - 0.16	0.63

5 **Example 4 - Affinity measured by Biacore**

Aim

The aim was to estimate the affinity, on rate and off rate of the different CD137 antibodies.

10 Material and methods

Human CD137 (R&D systems) was immobilized to the Biacore™ sensor chip, CM5, using conventional amine coupling. The tested antibody and control (serially diluted 1/2 10-0.63 nM) were analyzed for binding in HBS-P (GE, #BR-1003-68) at a flow rate of 30 µl/ml. The association was followed for 5 minutes and the dissociation for 15 minutes. Regeneration was performed twice using 10 mM Glycine pH 1.7 for 30 seconds. The kinetic parameters and the affinity constants were calculated using 1:1 Langmuir model.

Results and conclusion

The affinities of the antibodies were in the nanomolar to sub-nanomolar range (Table 3) measured using bivalent antibodies flowed over CD137 coated on the chip surface.

Table 3 - Kinetic parameters measured by surface plasmon resonance

Antibody	ka (1/Ms)	kd (1/s)	KD (M)
REF4	6.76E+05	6.60E-04	9.76E-10
REF8	3.92E+05	5.19E-04	1.32E-09
1630/1631	1.85E+06	1.18E-03	6.41E-10
REF2	1.05E+06	4.45E-04	4.24E-10

Example 5 - Target specificity of the CD137 antibodies

5

Aim

The aim with this study was to evaluate the risk that any of the CD137 antibody binds targets other than CD137.

10 Material and methods

Binding to TNFR superfamily members for which ELISA methods had already been established (CD40 and OX40) was evaluated to detect potential propensity to cross react to non-target proteins. In addition, a BLAST search was performed identifying TNFRSF21 as the most similar sequence (34% sequence identity). Since this sequence similarity is rather low, determination of non-target binding to OX40 and CD40 was considered sufficient.

ELISA plates (Greiner # 655074) were coated with 50µl/well of recombinant human OX40 (R&D # 1493-CD), CD40-Fc (Ansell # 504-820) or CD137 (R&D # 838-4B) diluted to a final concentration of 0.5µg/ml in PBS for 1h at 37°C or overnight at 4°C. Plates were washed with PBS+0.05% TWEEN20 (PBST), followed by block with PBST+1% bovine serum albumin (BSA). Antibody samples were prepared as serial 1/10 dilutions from 10 – 0.01 µg/ml in PBST+1% BSA and incubated for 1h in room temperature, followed by detection using a horse radish peroxidase-conjugated anti-human kappa light chain antibody (AbD Serotec # STAR127P) and developed using SuperSignal ELISA Pico Chemiluminescent substrate (Pierce ThermoScientific #37069).

Results and conclusion

Table 4 - Summary of CD137 antibody unspecific binding to OX40 and CD40

Antibody	Binding to OX40 and CD40	EC50 CD137
REF3	No	
REF4	Weak; EC50> 6 µg/ml (40 nM)	0.4 nM
REF8	No	
1630/1631	No	0.4 nM
2674/2675	No	0.3 nM

- 5 The results from the two experiments were similar. One antibody (REF4) exhibited weak binding to OX40 and CD40, whereas none of the remaining antibody showed any detectable binding to either OX40 or CD40. An overview of antibodies analyzed, and results from the two experiments is shown in Table 4.
- 10 Further, binding to primary PBL from multiple blood donors was tested. The binding of 1630/1631 and 2674/2675 to PBL was similar to Reference antibodies. No relevant unspecific binding to non-target proteins was detected.

Example 6 - Domain mapping of antibodies binding to CD137

15

Aim

The aim was to define distinct classes of epitope specificity, and compare to the properties of reference antibody.

Material and methods

20 The ability of each antibody to bind to a panel of human/mouse CD137 chimeras expressed on the surface of transfected cells was analyzed by flow cytometry.

The chimeras were designed by exchanging domains or modules of the human CD137
25 with the corresponding mouse domain. Genes of CD137 human/mouse chimeras were synthesized (GenScript) and constructs cloned into pcDNA3.1 vector (Invitrogen) and transiently transfected into FreeStyle 293-F cells (Invitrogen). The transfected cells were incubated with CD137 antibodies and control antibodies, followed by incubation with anti-human IgG-PE (Jackson Immunoresearch) for detection and analyzed with FACS Verse
30 (BD Biosciences). Binding to the different chimeric constructs was calculated as relative

MFI compared to the binding of the isotype control, followed by normalization to the full-length human CD137 construct to minimize the effect of affinity differences between individual antibodies.

5 Results and conclusion

Three binding patterns was observed as described below Figure 2. Data is summarized in Table 5.

Pattern A

10 Antibody REF1 depends on domain 1 for binding to human CD137.

Pattern B

Antibodies REF3, REF4, 2674/2675 and 1630/1631 are mainly dependent on domain 2 for binding to human CD137.

15

Pattern C

Antibodies REF2 (Reference antibody) and REF8 appear to be mainly dependent on domains 3B-4A for binding to human CD137.

20

Table 5. Median fluorescence intensity (MFI) for antibody sample/isotype control, normalized to full-length human CD137.

Domain		1	2				3B-4A	
Clone	Description	REF1	1630/ 1631	2674/ 2675	REF3	REF4	REF2	REF8
1550	Human CD137 with mouse domains 1, 2A and 2B (aa 24-86)	0.12	0.11	0.13	0.05	0.05	0.22	0.17
1551	Human CD137 with mouse domains 2A, 2B and 3A (aa 47-96)	0.41	0.10	0.15	0.04	0.05	0.37	0.33
1552	Human CD137 with mouse domains 2B, 3A and 3B (aa 64-118)	0.76	0.25	0.13	0.05	0.06	0.19	0.18
1553	Human CD137 with mouse domains 3A, 3B and 4A (aa 87-133)	1.07	0.91	1.08	0.65	0.65	0.17	0.17
1554	Human CD137 with mouse domains 3B, 4A and 4B (aa 97-159)	0.82	0.85	0.88	0.84	0.51	0.16	0.17
1555	Human CD137 with mouse domains 1 and 4B and region of unknown function (aa 24-46 and aa 139-186)	0.11	0.35	0.38	0.24	0.26	0.26	0.32
1030*	Human full length CD137	1	1	1	1	1	1	1

Example 7 - CD137 ligand blocking

Aim and background

The aim was to determine if the CD137 antibodies block the CD137 ligand binding.

5

If the CD137 antibodies bind to epitopes close to the ligand binding region, binding to the antigen can lead to partly or total block of ligand binding. Binding close to the CD137 ligand binding epitope may also affect the ligand binding due to steric hindrance or conformational changes of the CD137 ligand binding epitope. All CD137 antibodies were titrated against a fixed concentration of CD137L for evaluation of ligand blocking properties.

10

Material and method

CHO-cells transfected with human CD137 were used for the ligand competition. The extracellular part of human CD137 was fused to the transmembrane and intracellular part of hCD40 and cloned into pcDNA3.1. The vector was subsequently stably transfected into CHO cells. The expression of CD137 was confirmed by staining with commercial antibody targeting CD137.

15

The CHO-huCD137 were pre-incubated with CD137 monoclonal antibodies, titrating down from a predetermined saturating concentration (0.25 µg/ml), for 1h at +4°C before the addition of CD137 ligand at a concentration at EC50. After co-incubation for another 30 min at +4°C, the cells were washed and bound CD137 ligand was detected with anti-FLAG-APC (Cell signaling technology). Before analysis the cells were fixed with paraformaldehyde (10x concentrate BD CellFIX, BD biosciences). Analysis was performed with FACSverse and the MFI (Median Fluorescence Intensity) was calculated with FlowJo software.

20

25

Results and conclusion

It can be concluded that all CD137 mAbs tested were not blocking the CD137 ligand binding (Table 6, Figure 3). CD137 mAbs belonging to group B and C, binding to domain 2B-4A, block the CD137L (including 2674/2675 and 1630/1631). REF1 belonging to group A which bound to domain 1, did not block CD137 ligand. REF1 increased the binding of the CD137L.

30

Table 6 - Maximal CD137 ligand competition of the CD137 antibodies.

Group (domain mapping)	CD137 mAb	CD137L, max inhib.
A	REF1	-167%
B	1630	69%
B	2674/2675	66%
C	REF2	-26%

Example 8 - Competition ELISA

5

Aim and background

By competing each CD137 antibody with each another, it is possible to determine antibodies binding to similar epitopes based on their blocking pattern. The competition ELISA is performed by co-incubating biotinylated CD137 antibodies with non-biotinylated CD137 antibodies when binding to coated CD137-Fc. Competition is defined as loss of signal from the biotinylated CD137 antibody. Low competition values could either be due to no competition between the antibodies or binding kinetics of the antibodies. Binding of one antibody could also lead to steric hindrance or conformational changes when binding the antigen which affects the binding of the other CD137 antibody.

15

Material and method

CD137 antibodies were biotinylated (EZ-link NHS-LC-Biotin, ThermoFisher) and intact binding properties to CD137-Fc was verified with ELISA by comparing EC50 between biotinylated and non-biotinylated anti-CD137 mAbs. Non-biotinylated anti-CD137 (anti-CD137-bio) was pre-incubated to CD137-Fc at concentrations 30 times higher than the determined EC50 for 0.5 h. Without washing, anti-CD137-bio was added and co-incubated for another 1h. The binding of anti-CD137-bio was detected with Streptavidin-HRP (Pierce). Competition was calculated as the relative number by dividing the binding measured to other antibodies relative to its maximum competition (competing with itself).

The relative values obtained were normalized against the maximum blocking capacity (Table 7).

Table 7 - Summary of CD137 antibody competition ELISA from two experiments.

	REF1	REF4	1630/1631	REF2	REF8
REF1	100	7	5	5	4
REF2	15	41	70	94	61
REF4	18	58	91	63	50
REF8	4	49	91	100	82
1630/1631	14	31	56	23	16

5

Result and conclusion

The competition ELISA was repeated two times. In both experiments, several of the CD137 mAbs did not fully compete with itself (Table 7). The antibody REF1 that belongs to domain mapping group A, displayed a unique pattern in the competition ELISA. The other CD137 antibodies that were analyzed displayed similar blocking patterns. Differences in binding kinetics between those antibodies may explain some of the minor variations in the binding patterns among these antibodies, although it cannot be excluded that the small variations within groups reflects actual differences in the binding epitope.

15 ***Example 9 - In vitro efficacy of CD137 antibodies***

Aim

The aim was to identify CD137 antibodies with agonistic activity.

20 Material and methods

Agonistic activity of CD137 antibodies was evaluated in a T cell assay based on primary human CD8⁺ T cells. Briefly, CD8⁺ T cells were separated from human peripheral blood mononuclear cells by MACS separation (Miltenyi # 130-096-495) according to the manufacturer's protocol. Cells were incubated in 96-well microtiter plates (NuncThermo Scientific #268200), pre-coated with anti-CD3 antibody (clone OKT3, Affymetrix eBioscience # 16-0037) and titrated concentrations of the CD137 antibody to be tested. Following 72 or 96-hour incubation, culture medium was harvested and IFN- γ levels were determined by ELISA (BD #555142).

Each clone was analyzed in at least 6 donors and compared to the reference CD137 antibody REF1 and the negative control antibody.

Due to large intra-donor variations the stimulation index (SI, fold induction by antibody compared to negative control) was determined for each sample and normalized to the stimulation index for the reference antibody REF1.

Results and conclusion

Several clones with efficacy comparable to the reference REF1 were identified. Data is summarized in Figure 4.

Table 8 Table 8 indicates the absolute IFN- γ levels induced by CD137 stimulation. However, all antibodies were not analyzed head-to-head in all donors, and the normalized SI is more relevant for comparison of the efficacy.

Table 8 – IFN- γ production levels induced by the various antibody.

Clone name	Mean IFN- γ (pg/ml)	Min IFN- γ (pg/ml)	Max IFN- γ (pg/ml)	n
Ctrl IgG	2502	337	8526	13
REF1	42268	2256	136802	12
REF4	26749	11952	51832	8
REF8	52448	7727	123127	8
1630/1631	51236	3361	145055	8

Example 10 - In vitro NF κ B reporter assay

293T cells (30 million) were transfected with plasmids which encode for human CD137, firefly luciferase under NF- κ B promoter and renilla. After 5 hours of transfection, the antibodies were added at three different concentrations. 18 hours later, cells were harvested and luciferase reporter assay (Promega) was performed. The cells were cultured with soluble antibodies without crosslinking, as well as with cross-linking, at 5 μ g/ml with crosslinking using anti-IgG antibody.

Results

1630/1631 stimulates CD137 inducing NF- κ B mediated signaling when cross linked but not in the absence of a cross linking agent. In contrast, REF1 induce CD137 signaling also in the absence of a cross linking agent.

5

Example 11- In vivo anti-tumour effect in HT-29 colon cancer model

Summary

The anti-tumour effect of 1630/1631 was investigated using hPBMC humanized
10 immunodeficient mice and subcutaneous tumour models of HT-29 colon carcinoma.

1630/1631 demonstrated statistically significant tumour volume inhibition.

Material and methods

15 Leukocyte concentrates were obtained from Lund University Hospital.

Female SCID-Beige mice (7-8w) from Taconic's Denmark were used in the experiments. All experiments were done by approval of Malmö/Lund ethical committee.

20 HT-29 colon cancer were obtained from ATCC and cultivated according to ATCC recommendations. The HT-29 cell line growing in log phase was injected subcutaneously (4×10^6 cells in 200 μ L at day 0 (D0)). Human PBMC (7×10^6 in 100 μ L) isolated from leukocyte concentrates was injected intraperitoneally at the same day. Intraperitoneal treatments (100 μ g) were done at days 6, 13, and 20.

25

Tumour was measured with a calliper in width, length and height of which the tumour volume calculated ($w/2 \times l/2 \times h/2 \times \pi \times (4/3)$). The animals were terminated before the tumour volume reached 2cm³, at wounding, or affected health of the mice.

30 The data were analyzed by Mann-Whitney test using the GraphPad Prism program.

Results

Pooled data from mice engrafted with 4 different donors demonstrated statistically significant anti-tumour efficacy at days 12-16 in the form of inhibition of tumour growth
35 when treated with the 1630/1631 antibody ($p = 0.0675$ to $p = 0.0132$, Mann-Whitney non parametric, 2-tail) in comparison to the vehicle group. The percentage of tumour volume

inhibition ranged from 29-42% with 1630/1631 between days 10 and 21 (see Figure 6 and Table 9).

In conclusion, the anti-tumour effect of 1630/1631 was investigated using hPBMC humanized immunodeficient mice and subcutaneous tumour models of HT-29 colon carcinoma. 1630/1631 demonstrated statistically significant tumour volume inhibition.

Table 9. Statistical analysis and percent tumour inhibition

Day after tumour inoculation	Tumour growth inhibition (tumour volume) compared to vehicle (%)	p-value (Mann-Whitney 2-tail)
D12	42.1	0.0132
D14	32.6	0.0675
D16	38.7	0.0304
D19	32.7	0.1918
D21	29.5	0.0911

Example 12 - Optimization of CD137 parental antibody clone 1630/1631

The aim of the optimization was to generate improved variants of the 1630/1631 antibody with regard to affinity and biophysical properties. Phage selections towards recombinant CD137 coated onto the surface of beads were performed and prior to each selection round, the phage stocks were pre-selected towards non-target proteins as well as beads. Prior to the fourth round of selection a thermal incubation step at 65°C were performed. Overall the selection strategy was designed to promote the isolation of clones with a slow off-rate as well as a fast on-rate by prolonging the washing steps and decreasing the incubation time between the phage pool and CD137.

After phage selections, screening was performed in a soluble scFv format to identify target binding clones as well as to evaluate the diversity. An extended primary screening was performed to identify clones with improved temperature stability, cynomolgus reactivity as well as affinity or off-rate. A total of 50 clones were re-cloned into the final IgG4 format having the S228P stabilizing mutation. Further evaluation of optimized variants was performed in the final format and was focused on binding in an ELISA set-up, cell-binding as determined by FACS, affinity, temperature stability as determined by both an ELISA set-up as well as DSF, SE-HPLC, Schrödinger modelling and specificity.

Example 13 - Improved stability of clone 2674/2675

The aim with the DSF analysis was to determine the T_m of the clone 2674/2675 compared to the parental clone 1630/1631 to evaluate the improvement in temperature stability after optimization.

Material and Method

All antibodies were analyzed with differential scanning fluorometry (DSF) at SARomics Biostructure. Samples were diluted to 0.1 mg/ml in sterile filtered PBS and a volume of 150 µl were delivered to SARomics.

The samples for the DSF measurements (0.1 mg/ml in PBS buffer) were made up of 63 µl sample + 7 µl PBS buffer, 1:100 fold diluted SYPRO Orange). In total, the SYPRO Orange was diluted 1:1000-fold. Duplicate measurements were made for each construct using a Stratagene MX3000P, qPCR machine. Measurements were performed in the temperature range 25 °C – 95 °C. The average melting temperature T_m was calculated for all samples.

Results and conclusions

Melting curves for all samples were obtained and the determined T_{m1} as well as T_{m2} for 2674/2675 and parental clone 1630/1631 can be seen in Table 10 below. 2674/2675 had an improved T_{m2} by 1-2 °C as compared to the parental clone 1630/1631.

Table 10 - Determined T_{m1} as well as T_{m2} values and the average T_{m2} difference of 2674/2675 compared to 1630/1631 parental clone as measured by DSF

	T _{m1} (°C)	T _{m2} (°C)	Improved T _{m2} (°C)
2674	66.3	73.4	2
1630	66.3	71.4	-

Example 14 – Reduced aggregation propensity of clone 2674/2675 analyzed by Antibody Aggregation Prediction at Schrödinger.

The aim with the Schrödinger analysis of the optimized variants was to evaluate the size of the hydrophobic patch as well as the aggregation propensity of 2674/2675 in comparison to the parental clone 1630.

Material and methods

Sequences for the different variants were sent to Schrödinger and 3D structures were generated. The 3D models were analyzed with Protein Surface Analyzer and ranked with Aggscore. REF9-24 are clones obtained during optimization based on binding capacity for human CD137.

Results and conclusions

The defined patch size and aggscore for the clones 2674/2675 and REF9-24 (clones obtained during selection) as well as parental clone 1630/1631 can be seen in Figure 7. It can be concluded that the introduced mutations clearly disrupted the hydrophobic patch and reduced the aggregation propensity according to the modelling analysis.

Example 15 – Binding of 2674/2675 to Human and Cynomolgus CD137 measured by ELISA

Aim and background

The aim of the evaluation was to determine binding of 2674/2675 compared with parental clone 1630/1631 in ELISA to both human and cynomolgus CD137.

Material and Methods

Binding of CD137 antibodies to recombinant human CD137 was determined by sandwich ELISA. Briefly, ELISA plates (Greiner # 655074) coated with recombinant human CD137-Fc (R&D # 838-4B) were incubated with serial dilutions of the various CD137 antibodies to be investigated. CD137 antibodies were detected using HRP-conjugated goat-anti-human kappa light chain (AbD Serotec # STAR127P) and developed with SuperSignal ELISA Pico Chemiluminescent substrate (Pierce # 37069). EC50 values of the various antibodies were determined in 2-6 separate experiments.

Results and conclusions

2674/2675 exhibits EC50 values in a similar range as parental clone 1630/1631, i.e. sub nM. Data is summarized in Table 11 below.

Table 11. Determined EC50 values as measured by ELISA for 2674/2675 and parental clone 1630/1631.

	EC50 (nM) human CD137	EC50 (nM) cyno CD137
2674/2675	0.34	0.57
1630/1631	0.41	0.85

5 **Example 16 - Binding of 2674/2675 to human and cynomolgus CD137, Octet**

Aim

The aim was to compare relative binding affinities for 2674/2675 and 1630/1631 to human and cynomolgus CD137 using the Octet platform.

10

Materials and method

CD137 affinity was determined using the Octet Red 96 platform (ForteBio). 2674/2675, 1630/1631, REF1, REF 2 and 1188 isotype control were coupled at 10µg/ml to ARG2 biosensors (ForteBio #18-5092) by amine coupling with EDC and NHS. 7 2-fold serial dilutions of CD137 (Acro Biosystems # 41B-H5227 and # 41B-C52H4) from 100nM was prepared in 1x kinetic buffer (ForteBio #18-1092). Association was measured for 180 s followed by dissociation for 180s in 1x kinetic buffer. 10mM Glycine pH 2.2 was used for regeneration.

15

20 Data generated was referenced by reference well subtraction (1188), the baseline was aligned with the y-axis, inter-step correlation by alignment against association was performed and the data was smoothed by a Savitzky–Golay filtering in the data analysis software (v.9.0.0.14). The processed data was fitted using a 1:1 Langmuir binding model with X^2 as a measurement of fitting accuracy.

25

Results and conclusion

The binding affinities of 2674/2675, 1630/1631 and REF antibodies to human and cynomolgus CD137 are presented in Table 12. The affinity for 2674/2675 to human CD137 was improved by a factor 2 compared to 1630/1631. The affinity for 2674/2675 to cynomolgus CD137 was in the same range as 1630/1631.

30

Table 12 - Affinity for human and cynomolgus CD137 to immobilized 2674/2675, 1630/1631 and REF antibodies

Immobilized antibodies	Human CD137			Cynomolgus CD137		
	KD (M)	kon (1/Ms)	koff (1/s)	KD (M)	kon (1/Ms)	koff (1/s)
2674/2675	6.9E-09	3.5E+05	2.4E-03	2.5E-08	1.6E+05	4.0E-03
1630/1631	1.4E-08	1.6E+05	2.3E-03	1.8E-08	1.4E+05	2.5E-03
REF1	3.0E-09	4.6E+05	1.4E-03	no binding		
REF 2	6.1E-09	6.8E+05	4.1E-03	1.0E-08	5.2E+05	5.3E-03

5 **Example 17 – Binding affinity of 2674/2675 to human FcγRs**

Aim

The aim was to determine relative binding affinities for 2674/2675 to human FcγRs using the Octet platform.

Materials and methods

FcγR affinity was determined using the Octet RED96 platform equipped with Anti-Human Fab-CH1 (FAB2G) sensor tips (ForteBio). Antibodies were diluted to 200nM in 1X Kinetics Buffer (ForteBio) and loaded to a set of 8 parallel sensors for 300 seconds to reach an immobilization response of >1.5nm. The immobilized antibodies were then assayed against 7 2-fold dilutions of FcγRs, starting at 100nM. One immobilized sensor was assayed against 1X Kinetics Buffer for referencing and the entire assay was repeated without immobilization of antibodies to allow for double referencing. FcγRs included were obtained from R&D Systems (human FcγRI, #1257-FC-050; human FcγRIIa, #1330-CD-050; human FcγRIIb, #1460-CD-050; human FcγRIIIa (V158), #4325-FC-050; human FcγRIIIa (F158), #8894-FC-050). Binding to FcγRs was carried out for 60 seconds, followed by dissociation for 60 seconds in 1X Kinetics Buffer and regeneration of sensor tips using 10 mM glycine, pH 1.7. Data generated was referenced by standard double referencing, the baseline was aligned with the y-axis, inter-step correlation by alignment against dissociation was performed and the data was smoothed by a Savitzky–Golay filtering in the data analysis software (v.9.0.0.14). The processed data was fitted using a 1:1 Langmuir binding model with X^2 as a measurement of fitting accuracy. To improve curve fitting quality of dissociation curves generated against FcγRs with very fast dissociation rates, only the initial 10 seconds of the dissociation curves were included in the curve fitting.

Results and conclusions

The binding affinities human FcγRs of 2674/2675 and REF antibodies are presented in Table 13. 2674/2675 has a stronger binding to human FcγRI than all other assayed Fc receptors, as expected of an IgG4 antibody binding to the high affinity receptor FcγRI.

5 2674/2675 has a comparable binding to human FcγRs as REF1 antibody.

Table 13 - Determined binding affinities, KD (M), for 2674/2675 and REF antibodies to human FcγRs

KD (M)	FcγRI	FcγRIIa	FcγRIIb	FcγRIIIa 176V	FcγRIIIa 176F
2674/2675	2.11E-09	1.00E-06	7.79E-07	<det. limit	<det. limit
REF1	1.88E-09	6.74E-07	6.23E-07	<det. limit	<det. limit
REF2	<det. limit	5.87E-07	2.70E-06	<det. limit	<det. limit

10

Example 18 – Binding of 2674/2675 to Human and Cynomolgus CD137 measured by FACS

The aim of this study was to determine the binding to human and cynomolgus CD137.

15

Material and methods

Binding and EC50 was determined using flow cytometric of CHO cells transfected with human CD137, cynomolgus CD137 or empty vector. The extracellular part of human or cynomolgus CD137 was fused to the transmembrane and intracellular part of human CD40 and cloned into pcDNA3.1. The vector was subsequently stably transfected into CHO cells. Expression of CD137 was confirmed by flow cytometry using CD137 antibody (human CD137-PE, BD Biosciences # 555956) for 30 min at 4°C. CD137-transfected and empty vector-transfected cells were incubated with CD137 antibodies for at least 1h at 4°C to saturate the binding. In order to minimize antibody internalization, 0.05% sodium azide was used in the incubation buffer and all work was performed on ice. The CD137 antibodies were detected using PE-conjugated anti-hIgG antibody (109-115-098, Jackson ImmunoResearch laboratories), incubated for 30 min at 4°C. Directly after staining the cells were fixed with a paraformaldehyde solution (10x concentrate BD CellFIX, BD biosciences # 340181). Cells were analyzed by flow cytometry using FACSVerse (BD Biosciences). The median fluorescence intensity (MFI) for each sample was determined and the dose response data was analysed using Graph Pad Prism.

30

MFI data was normalized for each antibody, where 0% is defined as the lowest value and 100% is the highest value in the dose titration for each antibody. EC50 and 95% confidence interval were calculated with Graph Pad Prism based on data from the two experiments (non-linear regression (curve fit), constraints set to 0 and 100).

5

Results and conclusion

2674/2675 have a comparable binding to human CD137 as the parental clone 1630 and the REF1 and REF2 CD137 mAbs Figure 8 and Table 14. 2674/2675 and 1630/1631 have comparable binding to cynomolgus CD137 while REF1 does not bind to cynomolgus CD137 at all.

10

Table 14 - 95% confidence intervals for the EC50 of each CD137 antibody determined as an average from 3 experiments of normalized data. Nd: Not detectable

15

	Hu CD137 EC50 (nM) 95% conf interval	Cy CD137 EC50 (nM) 95% conf interval
2674/2675	0.26 - 0.37	0.46 – 0.77
1630/1631	0.23 - 0.34	0.55 – 0.87
REF1	0.33 - 0.73	n.d.
REF2	0.16 - 0.27	0.41 - 0.56

Example 19 – CD137 reporter assay with FcγR expressing cells for crosslink of CD137 mAb

20

Aim and background

Functional evaluation of 2674/2675 with the parental clone 1630/1631 in the CD137 reporter assay when crosslinking CD137 mAbs with FcγR transfected CHO cells.

25

Materials and methods

CHO-cells transfected with human FcγRI, FcγRIIa R131, FcγRIIb or empty vector (pcDNA3.1) were used for crosslinking. FcγR genes were cloned into pcDNA3.1. The vector was subsequently stably transfected into CHO cells. The expression of FcγRs was confirmed by staining with commercial antibody targeting CD32 or CD64.

30

Agonistic function of the CD137 mAbs was evaluated using a CD137 reporter assay (Promega, CD137 Bioassay Kit CS196005). The assay was performed according to the

manufacturer's protocol. In brief, FcγR transfected CHO cells and titrating concentrations of CD137 mAbs were diluted in RPMI containing 10% FCS and added to the assay plates before the addition of CD137 (Jurkat/CD137 cells) reporter cells. The assay plate was incubated for 6 h at 37°C until addition of Bio-Glo™ Luciferase Assay Detection and plate read in the BMG reader.

Results and conclusions

Crosslinking of the CD137 mAbs in the CD137 reporter assay with FcγRI transfected CHO cells demonstrates that 2674/2675 as well as the parental clone 1630/1631 induces a CD137 dependent activation of NF-κB in the reporter cell line (Figure 9). It can be concluded that if the CD137 antibody is crosslinking dependent, binding affinities to FcγRI, FcγRIIa R131 and FcγRIIb (shown in previous example) correlate well with the agonistic activity induced in the CD137 reporter assay after FcγR crosslinking (Figure 10). FcγR cross-linking independent activation of REF1 but not of 2674/2675, 1630/1631 or REF2 was determined using CHO cells transfected with empty vector (Figure 11).

Example 20 – CD8⁺ T cell agonist assay with FcγRI expressing CHO cells for crosslinking of CD137 antibodies

Aim and background

Functional evaluation of 2674/2675 with the parental clone 1630/1631 in a CD8⁺ T cell agonist assay when crosslinking the CD137 mAbs with FcγRI expressing cells.

Materials and methods

CHO-cells transfected with human FcγRI were used for crosslinking. The FcγRI gene were cloned into pcDNA3.1. The vector was subsequently stably transfected into CHO cells. The expression of FcγRI was confirmed by staining with commercial antibody targeting CD64.

Agonistic activity of CD137 antibodies was evaluated in a T cell assay based on primary human CD8⁺ T cells. Briefly, CD8⁺ T cells were separated from human peripheral blood mononuclear cells by MACS separation (Miltenyi # 130-096-495) according to the manufacturer's protocol. Cells were incubated in 96-well microtiter plates (NuncThermo Scientific #268200) pre-incubated with CHO cells transfected with FcγRI and incubated with tosyl beads coated with anti-CD3 antibody (clone OKT3, Affymetrix eBioscience # 16-0037) and titrated concentrations of the CD137 antibody to be tested. Following 72-hour

incubation, culture medium was harvested and IFN- γ levels were determined by ELISA (BD #555142).

Each clone was analyzed in at least 5 donors and compared to the reference CD137 antibody REF2. Due to large intra-donor variations, IFN- γ levels were normalized with 2674/2675 within each donor for comparison.

Results and conclusions

Crosslinking with Fc γ RI expressing CHO cells in the CD8 T cell agonist assay of the CD137 mAbs 2674/2675 and the parental clone 1630/1631, but not REF2, induces T cell activation, measured as an increase in IFN- γ production after 72 h (Figure 12). IgG binding affinities to Fc γ Rs of 2674/2675, 1630/1631 and REF2 have been determined and was shown in a previous example. It can be concluded that IgG binding affinity to Fc γ R correlate with the agonistic activity induced in the CD8⁺ T cells.

15

Example 21 - In vivo anti-tumour effect in HT-29 colon cancer model

Aim

The anti-tumour effect of 2674/2675 was investigated using hPBMC humanized immunodeficient mice and subcutaneous tumour models of HT-29 colon carcinoma.

20

Material and methods

Leukocyte concentrates were obtained from Lund University Hospital. Female SCID-Beige mice (7-8w) from Taconic's Denmark were used in the experiments. All experiments were done by approval of Malmö/Lund ethical committee.

25

HT-29 colon cancer was obtained from ATCC and cultivated according to ATCC recommendations. The HT-29 cell line growing in log phase was injected subcutaneously (4×10^6 cells in 200 μ L at day 0 (D0)). Human PBMC (10×10^6 in 100 μ L) isolated from leukocyte concentrates was injected intraperitoneally at the same day. Intraperitoneal treatments (100 μ g) were done twice weekly for three weeks starting at day 7.

30

Tumour was measured with a calliper in width, length and height of which the tumour volume calculated ($w/2 \times l/2 \times h/2 \times \pi \times (4/3)$). The animals were terminated before the tumour volume reached 2 cm³, at wounding, or affected health of the mice.

35

Results and conclusion

2674/2675 demonstrated anti-tumor efficacy in humanized mouse models in comparison to the vehicle group. The percentage of tumour volume inhibition ranged from 0-35% with 2674/2675 between days 19 and 28 (Table 15).

5

In conclusion, the anti-tumour effect of 2674/2675 was investigated using hPBMC humanized immunodeficient mice and subcutaneous tumour models of HT-29 colon carcinoma. 2674/2675 demonstrated tumour volume inhibition.

10 **Table 15.** Percent tumour inhibition

Day after tumour inoculation	Tumour growth inhibition (tumour volume) compared to vehicle (%)
D19	9.4
D21	24.0
D24	27.4
D26	24.7
D28	35.1

Example 22 - Gene expression analyses of FcγR and CD137 co expression in human tumor tissue

15

Aim

Assessing the gene expression of various Fcγ receptors, as well as CD137, in a wide range of human cancers using a curated and quality-controlled database of microarray and RNA-seq datasets.

20

Methods

Mean expression values for FcγRI, FcγRIIa, FcγRIIb, FcγRIIc, FcγRIIIa, FcγRIIIb and CD137 were obtained by performing gene expression profiling using Genevestigator, a curated and quality-controlled database of RNA microarray studies for human tissues (Hruz *et al.* 2008, *Adv Bioinformatics* **2008**:420747, the disclosures of which are incorporated herein by reference). Correlation plots were obtained by plotting the mean expression values of the various Fcγ receptor genes versus CD137 (Figure 13). Cancers displaying an above average expression (mean expression level ≥ 10) of both Fcγ receptor

25

and CD137 were identified and the top 10 solid tumor types and hematological malignancies are presented in Table 16 and Table 17, respectively.

Results and conclusions

5 Several human tumors with a high expression of Fcγ receptors and an above average expression of CD137 were identified using this method. The tables below provide an example of indications that could be highly sensitive to the antibody defined in this invention. This approach could be used to identify patient cohorts or individual patients that may benefit from treatment with agonistic CD137 antibodies. In fact, this type of
10 approach could be used to identify patients on an individual level that may benefit from the treatment. One could also envisage other methods being used to molecularly characterize the tumor, such as next generation sequencing or methods based on protein analysis such as immunohistochemistry, flow cytometry or proteomic approaches.

15 **Table 16:** Mean expression values of solid human tumors with an above average expression (mean expression level ≥ 10) of both Fcγ receptor and CD137 (TNFRSF9), as identified in Figure 13. The ten tumors with the highest expression of the six Fcγ receptors are shown.

Cancers with cells expressing FcγRIA and TNFRSF9	FcγRIA (Mean expression level)	TNFRSF9 (Mean expression level)
fibrous histiocyte, malignant, metastatic	13.43	10.22
adenosquamous carcinoma	12.78	10.80
undifferentiated sarcoma	12.67	10.16
clear cell adenocarcinoma, NOS, metastatic	12.66	10.62
acinar cell carcinoma	12.47	10.82
dedifferentiated liposarcoma	12.45	10.67
renal cell carcinoma, unstated behavior	12.44	10.38
intraductal micropapillary carcinoma	12.36	10.68
fibrous histiocyte, malignant	12.35	11.05
large cell neuroendocrine carcinoma	12.35	10.29
Cancers with cells expressing FcγRIIA and TNFRSF9	FcγRIIA (Mean expression level)	TNFRSF9 (Mean expression level)
Langerhans-cell histiocytosis, unifocal	14.58	13.85
acinar cell carcinoma	13.55	10.82

fibrous histiocytoma, malignant, metastatic	13.54	10.22
undifferentiated sarcoma	13.48	10.16
adenocarcinoma with mixed subtypes	13.43	10.77
dedifferentiated liposarcoma	13.32	10.67
undifferentiated sarcoma	13.22	10.52
fibrous histiocytoma, malignant	13.10	11.05
adenosquamous carcinoma	13.01	10.80
dedifferentiated liposarcoma	12.95	10.36
Cancers with cells expressing FcγRIIB and TNFRSF9	FcγRIIB (Mean expression level)	TNFRSF9 (Mean expression level)
Langerhans-cell histiocytosis, unifocal	13.23	13.85
fibrous histiocytoma, malignant, metastatic	12.64	10.22
adenocarcinoma with mixed subtypes	12.42	10.77
adenosquamous carcinoma	12.22	10.80
acinar cell carcinoma	12.19	10.82
Hodgkin's disease, NOS	12.01	10.88
renal cell carcinoma, unstated behavior	12.00	10.38
dedifferentiated liposarcoma	11.98	10.67
papillary adenocarcinoma, NOS	11.81	10.02
undifferentiated sarcoma	11.76	10.16
Cancers with cells expressing FcγRIIC and TNFRSF9	FcγRIIC (Mean expression level)	TNFRSF9 (Mean expression level)
Langerhans-cell histiocytosis, unifocal	12.72	13.85
Cancers with cells expressing FcγRIIIA and TNFRSF9	FcγRIIIA (Mean expression level)	TNFRSF9 (Mean expression level)
fibrous histiocytoma, malignant, metastatic	14.14	10.22
acinar cell carcinoma	13.14	10.82
large cell neuroendocrine carcinoma	12.97	10.29
adenocarcinoma with mixed subtypes	12.91	10.77
renal cell carcinoma, unstated behavior	12.85	10.38
intraductal micropapillary carcinoma	12.83	10.68
carcinoma, NOS	12.70	10.19
carcinoma, NOS	12.68	10.06
undifferentiated sarcoma	12.52	10.16
carcinoma, NOS	12.42	10.10

Cancers with cells expressing FcγRIIIB and TNFRSF9	FcγRIIIB (Mean expression level)	TNFRSF9 (Mean expression level)
carcinoma, NOS	13.09	10.19
carcinoma, NOS	12.66	10.06
carcinoma, NOS	12.61	10.10
tubular adenocarcinoma	12.06	10.18
carcinoma, NOS, micro-dissected	12.00	10.92
adenocarcinoma, intestinal type	11.97	10.26
fibrous histiocytoma, malignant, metastatic	11.89	10.22
neoplasm, malignant	11.81	10.05
adenocarcinoma, NOS	11.74	10.06
renal cell carcinoma, unstated behavior	11.61	10.38

Table 17: Mean expression values of hematological malignancies with an above average expression (mean expression level ≥ 10) of both Fcγ receptor and CD137, as identified in Figure 13. The ten malignancies with the highest expression of the six Fcγ receptors are shown.

Cancers with cells expressing FcγRIA and TNFRSF9	FcγRIA (Mean expression level)	TNFRSF9 (Mean expression level)
(extranodal) NK/T-cell lymphoma, nasal and nasal-type	13.97	11.56
Hodgkin's disease, NOS	13.65	10.88
malignant lymphoma, large B-cell, diffuse	12.15	12.23
primary mediastinal B-cell lymphoma	12.13	12.43
anaplastic large cell lymphoma, T-cell and Null cell type (ALCL), unstated behavior	12.11	11.38
angioimmunoblastic T-cell lymphoma	11.93	13.85
mature T-cell lymphoma, NOS	11.86	12.53
mature T-cell lymphoma, NOS, unstated behavior	11.69	11.72
anaplastic large cell lymphoma, T-cell and Null cell type (ALCL)	11.67	11.43
angioimmunoblastic T-cell lymphoma, unstated behavior	11.63	12.98

Cancers with cells expressing FcγRIIA and TNFRSF9	FcγRIIA (Mean expression level)	TNFRSF9 (Mean expression level)
mycosis fungoides	13.27	11.34
Hodgkin's disease, NOS	12.33	10.88
(extranodal) NK/T-cell lymphoma, nasal and nasal-type	12.26	11.56
mature T-cell lymphoma, NOS	11.76	12.53
angioimmunoblastic T-cell lymphoma	11.69	13.85
malignant lymphoma, large B-cell, diffuse	11.37	12.23
angioimmunoblastic T-cell lymphoma, unstated behavior	11.34	12.98
mature T-cell lymphoma, NOS, unstated behavior	11.33	11.72
adult T-cell leukemia/lymphoma (HTLV-1 positive), unstated behavior	11.28	11.13
anaplastic large cell lymphoma, T-cell and Null cell type (ALCL), unstated behavior	11.23	11.38
Cancers with cells expressing FcγRIIB and TNFRSF9	FcγRIIB (Mean expression level)	TNFRSF9 (Mean expression level)
chronic lymphocytic B-cell leukemia, unstated behavior, micro-dissected	14.69	10.46
mantle cell lymphoma	14.06	10.11
(extranodal) marginal zone B-cell lymphoma, NOS	12.73	11.85
malignant lymphoma, nodular, NOS, unstated behavior	12.69	12.97
malignant lymphoma, nodular, NOS	12.16	12.02
Hodgkin's disease, NOS	12.01	10.88
malignant lymphoma, large B-cell, diffuse	11.97	12.23
angioimmunoblastic T-cell lymphoma	11.89	13.85
mycosis fungoides	11.76	11.34
Hodgkin's disease, NOS, micro-dissected	11.38	11.32
Cancers with cells expressing FcγRIIC and TNFRSF9	FcγRIIC (Mean expression level)	TNFRSF9 (Mean expression level)
juvenile myelomonocytic leukemia	10.00	10.00

Cancers with cells expressing FcγRIIIA and TNFRSF9	FcγRIIIA (Mean expression level)	TNFRSF9 (Mean expression level)
juvenile myelomonocytic leukemia	13.19	10.00
(extranodal) NK/T-cell lymphoma, nasal and nasal-type	12.09	11.56
mycosis fungoides	11.71	11.34
mature T-cell lymphoma, NOS	11.16	12.53
anaplastic large cell lymphoma, T-cell and Null cell type (ALCL)	10.80	11.43
angioimmunoblastic T-cell lymphoma	10.65	13.85
Hodgkin's disease, NOS	10.51	10.88
anaplastic large cell lymphoma, T-cell and Null cell type (ALCL), unstated behavior	10.38	11.38
malignant lymphoma, large B-cell, diffuse	10.31	12.23
primary mediastinal B-cell lymphoma	10.26	12.43
Cancers with cells expressing FcγRIIIB and TNFRSF9	FcγRIIIB (Mean expression level)	TNFRSF9 (Mean expression level)
juvenile myelomonocytic leukemia	15.03	10.00
(extranodal) NK/T-cell lymphoma, nasal and nasal-type	11.95	11.56
anaplastic large cell lymphoma, T-cell and Null cell type (ALCL), unstated behavior	11.85	11.38
mature T-cell lymphoma, NOS	11.57	12.53
mature T-cell lymphoma, NOS, unstated behavior	11.56	11.72
angioimmunoblastic T-cell lymphoma, unstated behavior	11.54	12.98
primary mediastinal B-cell lymphoma	11.44	12.43
angioimmunoblastic T-cell lymphoma	11.30	13.85
adult T-cell leukemia/lymphoma (HTLV-1 positive), unstated behavior	11.08	11.13
(extranodal) marginal zone B-cell lymphoma, NOS	10.95	11.85

REFERENCES

- Almeida J, Bueno C, Algueró MC et al. Comparative analysis of the morphological, cytochemical, immunophenotypical, and functional characteristics of normal human peripheral blood lineage(-)/CD16(+)/HLA-DR(+)/CD14(-/lo) cells, CD14(+) monocytes, and CD16(-) dendritic cells. Clin Immunol. 2001 Sep;100(3):325-38.
- Akhmetzyanova, I., Zelinskyy, G., Littwitz-Salomon, E., Malyshkina, A., Dietze, K. K., Streeck, H., Brandau, S., and Dittmer, U. (2016) CD137 Agonist Therapy Can Reprogram Regulatory T Cells into Cytotoxic CD4+ T Cells with Antitumour Activity. J. Immunol. 196, 484-492.
- Ascierto, P. A., Simeone, E., Sznol, M., Fu, Y. X., and Melero, I. (2010) Clinical experiences with anti-CD137 and anti-PD1 therapeutic antibodies. Semin. Oncol. 37, 508-516.
- Baessler T, Charton JE, Schmiedel BJ, Grünebach F, Krusch M, Wacker A, Rammensee HG, Salih HR. CD137 ligand mediates opposite effects in human and mouse NK cells and impairs NK-cell reactivity against human acute myeloid leukemia cells. Blood. 2010 Apr 15;115(15):3058-69. doi: 10.1182/blood-2009-06-227934. Erratum in: Blood. 2010 Dec 23;116(26):6152. PubMed PMID: 20008791.
- Bartkowiak, T. and Curran, M. A. (2015) 4-1BB Agonists: Multi-Potent Potentiators of Tumour Immunity. Front Oncol. 5, 117.
- Bronte V, Brandau S, Chen SH et al. Recommendations for myeloid-derived suppressor cell nomenclature and characterization standards. Nat Commun. 2016 Jul 6;7:12150.
- Bruhns P, Iannascoli B, England P et al. Specificity and affinity of human Fcγ receptors and their polymorphic variants for human IgG subclasses. Blood. 2009 Apr 16;113(16):3716-25.
- Bulliard Y, Jolicoeur R, Zhang J, Dranoff G, Wilson NS, Brogdon JL. OX40 engagement depletes intratumoural Tregs via activating FcγRs, leading to antitumour efficacy. Immunol Cell Biol. 2014 Jul;92(6):475-80. doi: 10.1038/icb.2014.26. PubMed PMID: 24732076.
- Cavnar MJ, Zeng S, Kim TS et al. KIT oncogene inhibition drives intratumoral macrophage M2 polarization. J Exp Med. 2013 Dec 16;210(13):2873-86.
- Cheeseman HM, Carias AM, Evans AB et al. Expression Profile of Human Fc Receptors in Mucosal Tissue: Implications for Antibody-Dependent Cellular Effector Functions Targeting HIV-1 Transmission. PLoS One. 2016 May 10;11(5):e0154656.
- Curran, M. A., Kim, M., Montalvo, W., Al-Shamkhani, A., and Allison, J. P. (2011) Combination CTLA-4 blockade and 4-1BB activation enhances tumour rejection by increasing T-cell infiltration, proliferation, and cytokine production. PLoS. ONE. 6, e19499.
- Dubrot, J., Milheiro, F., Alfaro, C., Palazon, A., Martinez-Forero, I., Perez-Gracia, J. L., Morales-Kastresana, A., Romero-Trevejo, J. L., Ochoa, M. C., Hervas-Stubbs, S., Prieto, J., Jure-Kunkel, M., Chen, L., and Melero, I. (2010) Treatment with anti-CD137 mAbs causes intense accumulations of liver T cells without selective antitumour immunotherapeutic effects in this organ. Cancer Immunol. Immunother. 59, 1223-1233.

- Gauttier, V., Judor, J. P., Le, G., V, Cany, J., Ferry, N., and Conchon, S. (2014) Agonistic anti-CD137 antibody treatment leads to antitumour response in mice with liver cancer. *Int. J. Cancer* 135, 2857-2867.
- 5 Gray, J. C., French, R. R., James, S., Al-Shamkhani, A., Johnson, P. W., and Glennie, M. J. (2008) Optimising anti-tumour CD8 T-cell responses using combinations of immunomodulatory antibodies. *Eur. J. Immunol.* 38, 2499-2511.
- Guilliams M, Bruhns P, Saeys Y et al. The function of Fcγ receptors in dendritic cells and
10 macrophages. *Nat Rev Immunol.* 2014 Feb;14(2):94-108.
- Guo, Z., Cheng, D., Xia, Z., Luan, M., Wu, L., Wang, G., and Zhang, S. (2013) Combined TIM-3 blockade and CD137 activation affords the long-term protection in a murine model of ovarian cancer. *J. Transl. Med.* 11, 215.
- 15 Elliott LA, Doherty GA, Sheahan K et al. Human Tumor-Infiltrating Myeloid Cells: Phenotypic and Functional Diversity. *Front Immunol.* 2017 Feb 6;8:86
- Eruslanov EB, Bhojnagarwala PS, Quatromoni JG et al. Tumor-associated neutrophils stimulate T cell responses in early-stage human lung cancer. *J Clin Invest.* 2014 Dec;124(12):5466-80.
- 20 Eruslanov E, Neuberger M, Daurkin I et al. Circulating and tumor-infiltrating myeloid cell subsets in patients with bladder cancer. *Int J Cancer.* 2012 Mar 1;130(5):1109-19.
- 25 Griesinger AM, Birks DK, Donson AM et al. Characterization of distinct immunophenotypes across pediatric brain tumor types. *J Immunol.* 2013 Nov 1;191(9):4880-8.
- 30 Grugan KD, McCabe FL, Kinder M et al. Tumor-associated macrophages promote invasion while retaining Fc-dependent anti-tumor function. *J Immunol.* 2012 Dec 1;189(11):5457-66.
- Hansen BD, Schmidt H, von der Maase H et al. Tumour-associated macrophages are
35 related to progression in patients with metastatic melanoma following interleukin-2 based immunotherapy. *Acta Oncol.* 2006;45(4):400-5.
- Hogarth PM, Pietersz GA. Fc receptor-targeted therapies for the treatment
40 of inflammation, cancer and beyond. *Nat Rev Drug Discov.* 2012 Mar 30;11(4):311-31.
- Holbrook E, Kohrt, A, Dimitrios Colevas, Roch Houot,1,2,3 Kipp Weiskopf, Matthew J. Goldstein, Peder Lund, Antonia Mueller, Idit Sagiv-Barfi, Aurelien Marabelle, Ruth Lira, Emily Troutner, Lori Richards,1 Amanda Rajapaska, Jonathan Hebb, Cariad Chester, Erin
45 Waller,Anton Ostashko, Wen-Kai Weng,Lieping Chen, Debra Czerwinski, Yang-Xin Fu, John Sunwoo, and Ronald Levy. Targeting CD137 enhances the efficacy of cetuximab. *The Journal of Clinical Investigation* Volume 124 Number 6 June 2014
- Horton HM, Bennett MJ, Peipp M, Pong E, Karki S, Chu SY, Richards JO, Chen H, Repp
50 R, Desjarlais JR, Zhukovsky EA. Fc-engineered anti-CD40 antibody enhances multiple effector functions and exhibits potent in vitro and in vivo antitumour activity against hematologic malignancies. *Blood.* 2010 Oct 21;116(16):3004-12.
doi: 10.1182/blood-2010-01-265280. PubMed PMID: 20616215.

- Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W, Zimmermann P: Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. *Adv Bioinformatics* 2008, **2008**:420747.
- 5 Hu W, Li X, Zhang C et al. Tumor-associated macrophages in cancers. *Clin Transl Oncol.* 2016 Mar;18(3):251-8.
- Kim, J. A., Averbook, B. J., Chambers, K., Rothchild, K., Kjaergaard, J., Papay, R., and Shu, S. (2001) Divergent effects of 4-1BB antibodies on antitumour immunity and on tumour-reactive T-cell generation. *Cancer Res* 61, 2031-2037.
- 10 Kwong, B., Gai, S. A., Elkhader, J., Wittrup, K. D., and Irvine, D. J. (2013) Localized immunotherapy via liposome-anchored Anti-CD137 + IL-2 prevents lethal toxicity and elicits local and systemic antitumour immunity. *Cancer Res.* 73, 1547-1558.
- 15 Lee, H. W., Park, S. J., Choi, B. K., Kim, H. H., Nam, K. O., and Kwon, B. S. (2002) 4-1BB promotes the survival of CD8+ T lymphocytes by increasing expression of Bcl-xL and Bfl-1. *J Immunol* 169, 4882-4888.
- 20 Lee, S. J., Myers, L., Muralimohan, G., Dai, J., Qiao, Y., Li, Z., Mittler, R. S., and Vella, A. T. (2004) 4-1BB and OX40 dual costimulation synergistically stimulate primary specific CD8 T cells for robust effector function. *J. Immunol.* 173, 3002-3012.
- 25 Li Y, Lee PY, Kellner ES et al. Monocyte surface expression of Fcγ receptor RI (CD64), a biomarker reflecting type-I interferon levels in systemic lupus erythematosus. *Arthritis Res Ther.* 2010;12(3):R90
- Li C, Luo X, Lin Y et al. A Higher Frequency of CD14+ CD169+ Monocytes/Macrophages in Patients with Colorectal Cancer. *PLoS One.* 2015 Oct 28;10(10):e0141817.
- 30 Li, F. and Ravetch, J. V. (2011) Inhibitory Fcγ receptor engagement drives adjuvant and anti-tumour activities of agonistic CD40 antibodies. *Science* 333, 1030-1034.
- 35 Lu J, Chu J, Zou Z et al. Structure of FcγRI in complex with Fc reveals the importance of glycan recognition for high-affinity IgG binding. *Proc Natl Acad Sci U S A.* 2015 Jan 20;112(3):833-8
- McMillin, D. W., Hewes, B., Gangadharan, B., Archer, D. R., Mittler, R. S., and Spencer, H. T. (2006) Complete regression of large solid tumours using engineered drug-resistant hematopoietic cells and anti-CD137 immunotherapy. *Hum. Gene Ther* 17, 798-806.
- 40 Melero, I., Shuford, W. W., Newby, S. A., Aruffo, A., Ledbetter, J. A., Hellstrom, K. E., Mittler, R. S., and Chen, L. (1997) Monoclonal antibodies against the 4-1BB T-cell activation molecule eradicate established tumours. *Nat Med* 3, 682-685.
- 45 Melero I, Daniel Hirschhorn-Cymerman, Aizea Morales-Kastresana, et al. Agonist Antibodies to TNFR Molecules That Costimulate T and NK cells *Clin Cancer Res* 2013;19:1044-1053. Published online March 3, 2013.
- 50 Melero I, Antonio M. Grimaldi, Jose L. Perez-Gracia, et al. Clinical Development of Immunostimulatory Monoclonal Antibodies and Opportunities for Combination. *Clin Cancer Res* 2013;19:997-1008.
- 55 Miller, R. E., Jones, J., Le, T., Whitmore, J., Boiani, N., Gliniak, B., and Lynch, D. H. (2002) 4-1BB-specific monoclonal antibody promotes the generation of tumour-specific immune

- responses by direct activation of CD8 T cells in a CD40-dependent manner. *J Immunol* 169, 1792-1800.
- 5 Morales-Kastresana, A., Sanmamed, M. F., Rodriguez, I., Palazon, A., Martinez-Forero, I., Labiano, S., Hervas-Stubbs, S., Sangro, B., Ochoa, C., Rouzaut, A., Azpilikueta, A., Bolanos, E., Jure-Kunkel, M., Gutgemann, I., and Melero, I. (2013) Combined immunostimulatory monoclonal antibodies extend survival in an aggressive transgenic hepatocellular carcinoma mouse model. *Clin. Cancer Res.* 19, 6151-6162.
- 10 Morimura T, Neuchrist C, Kitz K et al. Monocyte subpopulations in human gliomas: expression of Fc and complement receptors and correlation with tumor proliferation. *Acta Neuropathol.* 1990;80(3):287-94.
- 15 Norton SE, Dunn ET, McCall JL et al. Gut macrophage phenotype is dependent on the tumor microenvironment in colorectal cancer. *Clin Transl Immunology.* 2016 Apr 29;5(4):e76.
- 20 Niu, L., Strahotin, S., Hewes, B., Zhang, B., Zhang, Y., Archer, D., Spencer, T., Dillehay, D., Kwon, B., Chen, L., Vella, A. T., and Mittler, R. S. (2007) Cytokine-mediated disruption of lymphocyte trafficking, hemopoiesis, and induction of lymphopenia, anemia, and thrombocytopenia in anti-CD137-treated mice. *J. Immunol.* 178, 4194-4213.
- 25 Overdijk MB, Verploegen S, Ortiz Buijsse A, Vink T, Leusen JH, Bleeker WK, Parren PW. Crosstalk between human IgG isotypes and murine effector cells. *J Immunol.* 2012 Oct 1;189(7):3430-8. PubMed PMID: 22956577.
- 30 Palazón A, Iván Martínez-Forero, Alvaro Teijeira, et al. The HIF-1a Hypoxia Response in Tumour-Infiltrating T Lymphocytes Induces Functional CD137 (4-1BB) for Immunotherapy *Cancer Discovery* 2012;2:608-623. Published OnlineFirst June 19, 2012.
- 35 Palazón A, Teijeira A, Martínez-Forero I, Hervás-Stubbs S, Roncal C, Peñuelas I, Dubrot J, Morales-Kastresana A, Pérez-Gracia JL, Ochoa MC, Ochoa-Callejero L, Martínez A, Luque A, Dinchuk J, Rouzaut A, Jure-Kunkel M, Melero I. Agonist anti-CD137 mAb act on tumour endothelial cells to enhance recruitment of activated T lymphocytes. *Cancer Res.* 2011 Feb 1;71(3):801-11. doi:10.1158/0008-5472.CAN-10-1733. PubMed PMID: 21266358.
- 40 Pan, P. Y., Zang, Y., Weber, K., Meseck, M. L., and Chen, S. H. (2002) OX40 ligation enhances primary and memory cytotoxic T lymphocyte responses in an immunotherapy for hepatic colon metastases. *Mol Ther* 6, 528-536.
- 45 Porembka MR, Mitchem JB, Belt BA et al. Pancreatic adenocarcinoma induces bone marrow mobilization of myeloid-derived suppressor cells which promote primary tumor growth. *Cancer Immunol Immunother.* 2012 Sep;61(9):1373-85.
- 50 Pulle, G., Vidric, M., and Watts, T. H. (2006) IL-15-dependent induction of 4-1BB promotes antigen-independent CD8 memory T cell survival. *J Immunol* 176, 2739-2748.
- 55 Rabu, C., Quemener, A., Jacques, Y., Echasserieau, K., Vusio, P., and Lang, F. (2005) Production of recombinant human trimeric CD137L (4-1BBL). Cross-linking is essential to its T cell co-stimulation activity. *J Biol Chem* 280, 41472-41481.
- Roussel M, Ferrell PB Jr, Greenplate AR et al. Mass cytometry deep phenotyping of human mononuclear phagocytes and myeloid-derived suppressor cells from human blood and bone marrow. *J Leukoc Biol.* 2017 Aug;102(2):437-447

- Sallin, M. A., Zhang, X., So, E. C., Burch, E., Cai, L., Lin, W., Chapoval, A. I., and Strome, S. E. (2014) The anti-lymphoma activities of anti-CD137 monoclonal antibodies are enhanced in FcγRIII(-/-) mice. *Cancer Immunol. Immunother.* 63, 947-958.
- 5 Sanmamed, M. F., Pastor, F., Rodriguez, A., Perez-Gracia, J. L., Rodriguez-Ruiz, M. E., Jure-Kunkel, M., and Melero, I. (2015) Agonists of Co-stimulation in Cancer Immunotherapy Directed Against CD137, OX40, GITR, CD27, CD28, and ICOS. *Semin. Oncol.* 42, 640-655.
- 10 Shuford, W. W., Klussman, K., Tritchler, D. D., Loo, D. T., Chalupny, J., Siadak, A. W., Brown, T. J., Emswiler, J., Raecho, H., Larsen, C. P., Pearson, T. C., Ledbetter, J. A., Aruffo, A., and Mittler, R. S. (1997) 4-1BB costimulatory signals preferentially induce CD8+ T cell proliferation and lead to the amplification in vivo of cytotoxic T cell responses. *J Exp. Med* 186, 47-55.
- 15 So, T., Lee, S. W., and Croft, M. (2008) Immune regulation and control of regulatory T cells by OX40 and 4-1BB. *Cytokine Growth Factor Rev.* 19, 253-262.
- 20 Solito S, Marigo I, Pinton L et al. Myeloid-derived suppressor cell heterogeneity in human cancers. *Ann N Y Acad Sci.* 2014 Jun;1319:47-65.
- Stewart R, Hammond S, Oberst M, Wilkinson R. (2014) The role of Fc gamma receptors in the activity of immunomodulatory antibodies for cancer. *J Immunother.* 2:29
- 25 St Rose, M. C., Taylor, R. A., Bandyopadhyay, S., Qui, H. Z., Hagymasi, A. T., Vella, A. T., and Adler, A. J. (2013) CD134/CD137 dual costimulation-elicited IFN-γ maximizes effector T-cell function but limits Treg expansion. *Immunol. Cell Biol.* 91, 173-183.
- 30 Sun Y, Subudhi SK, Fu YX. Co-stimulation agonists as a new immunotherapy for autoimmune diseases. *Trends Mol Med.* 2003 Nov;9(11):483-9. Review. PubMed PMID: 14604826.
- 35 Taraban, V. Y., Rowley, T. F., O'Brien, L., Chan, H. T., Haswell, L. E., Green, M. H., Tutt, A. L., Glennie, M. J., and Al-Shamkhani, A. (2002) Expression and costimulatory effects of the TNF receptor superfamily members CD134 (OX40) and CD137 (4-1BB), and their role in the generation of anti-tumour immune responses. *Eur J Immunol* 32, 3617-3627.
- 40 Uno, T., Takeda, K., Kojima, Y., Yoshizawa, H., Akiba, H., Mittler, R. S., Gejyo, F., Okumura, K., Yagita, H., and Smyth, M. J. (2006) Eradication of established tumours in mice by a combination antibody-based therapy. *Nat. Med.* 12, 693-698.
- Vidarsson G, Dekkers G, Rispens T. IgG subclasses and allotypes: from structure to effector functions. *Front Immunol.* 2014 Oct 20;5:520. doi:10.3389/fimmu.2014.00520. Review. PubMed PMID: 25368619; PubMed Central PMCID: PMC4202688.
- 45 Vinay, D. S. and Kwon, B. S. (2012) Immunotherapy of cancer with 4-1BB. *Mol. Cancer Ther.* 11, 1062-1070.
- 50 Wang W, Erbe AK, Hank JA, Morris ZS, Sondel PM. NK Cell-Mediated Antibody-Dependent Cellular Cytotoxicity in Cancer Immunotherapy. *Front Immunol.* 2015 Jul 27;6:368. doi: 10.3389/fimmu.2015.00368. Review. PubMed PMID: 26284063; PubMed Central PMCID: PMC4515552.
- 55

Wei, H., Zhao, L., Li, W., Fan, K., Qian, W., Hou, S., Wang, H., Dai, M., Hellstrom, I., Hellstrom, K. E., and Guo, Y. (2013) Combinatorial PD-1 blockade and CD137 activation has therapeutic efficacy in murine cancer models and synergizes with cisplatin. *PLoS ONE*. 8, e84927.

Westwood, J. A., Darcy, P. K., Guru, P. M., Sharkey, J., Pegram, H. J., Amos, S. M., Smyth, M. J., and Kershaw, M. H. (2010) Three agonist antibodies in combination with high-dose IL-2 eradicate orthotopic kidney cancer in mice. *J. Transl. Med.* 8, 42.

Westwood, J. A., Matthews, G. M., Shortt, J., Faulkner, D., Pegram, H. J., Duong, C. P., Chesi, M., Bergsagel, P. L., Sharp, L. L., Huhn, R. D., Darcy, P. K., Johnstone, R. W., and Kershaw, M. H. (2014a) Combination anti-CD137 and anti-CD40 antibody therapy in murine myc-driven hematological cancers. *Leuk. Res.* 38, 948-954.

White AL, Chan HT, French RR, Willoughby J, Mockridge CI, Roghanian A, Penfold CA, Booth SG, Dodhy A, Polak ME, Potter EA, Ardern-Jones MR, Verbeek JS, Johnson PW, Al-Shamkhani A, Cragg MS, Beers SA, Glennie MJ. Conformation of the human immunoglobulin G2 hinge imparts superagonistic properties to immunostimulatory anticancer antibodies. *Cancer Cell*. 2015 Jan 12;27(1):138-48. doi:10.1016/j.ccell.2014.11.001. PubMed PMID: 25500122; PubMed Central PMCID:PMC4297290.

Westwood, J. A., Potdevin Hunnam, T. C., Pegram, H. J., Hicks, R. J., Darcy, P. K., and Kershaw, M. H. (2014b) Routes of delivery for CpG and anti-CD137 for the treatment of orthotopic kidney tumours in mice. *PLoS ONE*. 9, e95847.

Wilcox, R. A., Flies, D. B., Zhu, G., Johnson, A. J., Tamada, K., Chapoval, A. I., Strome, S. E., Pease, L. R., and Chen, L. (2002) Provision of antigen and CD137 signaling breaks immunological ignorance, promoting regression of poorly immunogenic tumours. *J Clin Invest* 109, 651-659.

Wilson, N. S., Yang, B., Yang, A., Loeser, S., Marsters, S., Lawrence, D., Li, Y., Pitti, R., Totpal, K., Yee, S., Ross, S., Vernes, J. M., Lu, Y., Adams, C., Offringa, R., Kelley, B., Hymowitz, S., Daniel, D., Meng, G., and Ashkenazi, A. (2011b) An Fcgamma receptor-dependent mechanism drives antibody-mediated target-receptor signaling in cancer cells. *Cancer Cell* 19, 101-113.

Wyzgol, A., Muller, N., Fick, A., Munkel, S., Grigoleit, G. U., Pfizenmaier, K., and Wajant, H. (2009) Trimer stabilization, oligomerization, and antibody-mediated cell surface immobilization improve the activity of soluble trimers of CD27L, CD40L, 41BBL, and glucocorticoid-induced TNF receptor ligand. *J Immunol* 183, 1851-1861.

Zhang, N., Sadun, R. E., Arias, R. S., Flanagan, M. L., Sachsman, S. M., Nien, Y. C., Khawli, L. A., Hu, P., and Epstein, A. L. (2007) Targeted and untargeted CD137L fusion proteins for the immunotherapy of experimental solid tumours. *Clin. Cancer Res.* 13, 2758-2767.

Zhang B, Wang Z, Wu L et al. Circulating and tumor-infiltrating myeloid-derived suppressor cells in patients with colorectal carcinoma. *PLoS One*. 2013;8(2):e57114

CLAIMS

1. An antibody or an antigen-binding fragment thereof with binding specificity for domain
2 of human CD137 in which the antibody or antigen-binding fragment is a CD137 agonist and
is capable of inhibiting the binding of reference antibody '1630/1631' to human CD137; in
5 which the antibody or an antigen-binding fragment thereof comprises a heavy chain variable
region comprising the following CDRs:

(a) a heavy chain CDR1 sequence with the consensus sequence G, F, T/N, F, G, Y, S,
Y;

10 (b) a heavy chain CDR2 sequence with the consensus sequence I, G, S, G/T, S, S,
Y/H, T; and

(c) a heavy chain CDR3 sequence with the sequence ARVYSSPGIDY;

and comprises a light chain variable region comprising the following CDRs:

(a) a light chain CDR1 sequence with the consensus sequence Q, S, I, S/G, S, Y/T;

(b) a light chain CDR2 sequence with the consensus sequence A/G, A, S; and

15 (c) a light chain CDR3 sequence with the sequence QQYYTWVPFT.

2. An antibody or an antigen-binding fragment thereof according to Claim 1, in which the
antibody or antigen-binding fragment is capable of inhibiting the binding of reference
antibody '2674/2675' to human CD137.

20 3. An antibody or antigen-binding fragment thereof according to Claim 1 or 2, in which the
antibody or antigen-binding fragment exhibits one or more of the following properties:

(a) the ability to stimulate CD137 and activate T cells and other immune cells via a
cross-linking dependent mechanism; and/or

(b) cross-reactivity with cyno-CD137 antibodies; and/or

25 (c) is capable of binding an Fc receptor; optionally, in which the ability of the
antibody to activate T cells is dependent upon binding to both CD137 and Fc receptors;
and/or

(d) is capable of inducing tumour immunity.

4. An antibody or antigen-binding fragment thereof according to any of Claims 1, 2 or 3, in which the antibody or antigen-binding fragment is incapable of inducing the following upon binding to cells expressing CD137:
 - (a) antibody-dependent cellular cytotoxicity (ADCC);
 - 5 (b) antibody-dependent cellular phagocytosis (ADCP); and/or
 - (c) complement-dependent cytotoxicity (CDC).
5. An antibody or antigen-binding fragment thereof according to any preceding claim, in which the antibody or antigen-binding fragment is capable of binding to an epitope on the extracellular domain of CD137 which overlaps, at least in part, with the epitope on CD137 to
10 which reference antibody 1630/1631 is capable of binding; optionally, in which the antibody or antigen-binding fragment is capable of binding to an epitope on the extracellular domain of CD137 which overlaps, at least in part, with the epitope on CD137 to which reference antibody 2674/2675 is capable of binding; further optionally, in which the epitope is located at or within amino acids 66 to 107 of human CD137.
- 15 6. An antibody or antigen-binding fragment thereof according to any preceding claim, comprising or consisting of:
 - (a) an intact antibody, for example an IgG1, IgG2, IgG3 or IgG4 antibody; or
 - (b) an antigen-binding fragment selected from the group consisting of Fv fragments (*e.g.* single chain Fv and disulphide-bonded Fv), Fab-like fragments (*e.g.* Fab fragments, Fab' fragments and F(ab)₂ fragments) and domain antibodies (*e.g.* single V_H variable domains or V_L variable domains).
 - 20 (c) an antigen-binding fragment selected from the group consisting of Fv fragments (*e.g.* single chain Fv and disulphide-bonded Fv), Fab-like fragments (*e.g.* Fab fragments, Fab' fragments and F(ab)₂ fragments) and domain antibodies (*e.g.* single V_H variable domains or V_L variable domains).
7. An antibody or antigen-binding fragment according to any preceding claim, in which:
 - (a) the antibody or antigen-binding fragment thereof is a recombinant polypeptide; and/or
 - 25 (b) the antibody or antigen-binding fragment thereof is monoclonal; and/or
 - (c) the antibody or antigen-binding fragment thereof is human or humanised.
8. An antibody or antigen-binding fragment thereof according to any preceding claim, comprising:

5 (a) a heavy chain variable region comprising the CDRs of SEQ ID NOs 3, 4 and 5; optionally, in which the antibody or antigen-binding fragment thereof comprises a heavy chain variable region having the amino acid sequence of SEQ ID NO: 1 or an amino acid sequence having at least 60% sequence identity therewith, for example at least 70%, 80%, or 90% sequence identity; and

10 (b) a light chain variable region comprising the CDRs of SEQ ID NOs: 6, 7 and 8; optionally, in which the antibody or antigen-binding fragment thereof comprises a light chain variable region having the amino acid sequence of SEQ ID NO: 2 or an amino acid sequence having at least 60% sequence identity therewith, for example at least 70%, 80%, or 90% sequence identity.

9. An antibody or antigen-binding fragment thereof according to any of Claims 1 to 7, comprising:

15 (a) a heavy chain variable region comprising the CDRs of SEQ ID NOs 21, 22 and 23; optionally, in which the antibody or antigen-binding fragment thereof comprises a heavy chain variable region having the amino acid sequence of SEQ ID NO 19: or an amino acid sequence having at least 60% sequence identity therewith, for example at least 70%, 80%, or 90% sequence identity; and

20 (b) a light chain variable region comprising the CDRs of SEQ ID NOs: 24, 25 and 26; optionally, in which the antibody or antigen-binding fragment thereof comprises a light chain variable region having the amino acid sequence of SEQ ID NO: 20 or an amino acid sequence having at least 60% sequence identity therewith, for example at least 70%, 80%, or 90% sequence identity.

10. An antibody or antigen-binding fragment thereof according to any preceding claim, comprising:

25 (a) a heavy chain constant region; optionally, in which the heavy chain constant region is of an immunoglobulin subtype selected from the group consisting of IgG1, IgG2, IgG3 and IgG4, further optionally in which the heavy chain constant region comprises or consists of an amino acid sequence selected from the group consisting of SEQ ID NO: 12, 13, 14 and 15; and/or

(b) a light chain constant region; optionally, in which the light chain constant region is of a kappa or lambda light chain; further optionally, in which the light chain constant region comprises or consists of an amino acid sequence of SEQ ID NO: 16.

11. An antibody or antigen-binding fragment thereof according to any preceding claim,
5 comprising an Fc region; optionally, in which

(a) the Fc region is naturally occurring; or

(b) the Fc region is non-naturally occurring; further optionally, in which the Fc region comprises mutations to shorten the half-life of the antibody or antigen binding fragment.

12. An antibody or antigen-binding fragment thereof according to any of Claims 1 to 8,
10 comprising:

(a) a heavy chain comprising a variable region of SEQ ID NO: 1 together with a constant region of SEQ ID NO: 13; and

(b) a light chain comprising a variable region of SEQ ID NO: 2 together with a constant region of SEQ ID NO: 16.

13. An antibody or antigen-binding fragment thereof according to any of Claims 1 to 7 or 9,
15 comprising:

(a) a heavy chain comprising a variable region of SEQ ID NO: 19 together with a constant region of SEQ ID NO: 13; and

(b) a light chain comprising a variable region of SEQ ID NO: 20 together with a
20 constant region of SEQ ID NO: 16.

14. An antibody or antigen-binding fragment thereof according to Claim 12, in which the antibody is an intact IgG4 molecule comprising or consisting of two heavy chains having an amino acid sequence of SEQ ID NO: 17 and two light chains having an amino acid sequence of SEQ ID NO: 18.

25 15. An antibody or antigen-binding fragment thereof according to Claim 13, in which the antibody is an intact IgG4 molecule comprising or consisting of two heavy chains having an amino acid sequence of SEQ ID NO: 29 and two light chains having an amino acid sequence of SEQ ID NO: 30.

16. An antibody or antigen-binding fragment thereof according to any preceding claim, further comprising:

(i) a cytotoxic moiety;

optionally, in which the cytotoxic moiety comprises or consists of:

5 (a) a radioisotope; or

(b) a cytotoxic drug; and/or

(ii) a detectable moiety;

optionally, in which the detectable moiety comprises or consists of a radioisotope, and in which optionally the cytotoxic moiety and/or detectable moiety is joined to the

10 antibody or antigen-binding fragment thereof indirectly, via a linking moiety; optionally in which the linking moiety is a chelator; further optionally, in which the chelator is selected from the group consisting of derivatives of 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA), deferoxamine (DFO), derivatives of diethylenetriaminepentaacetic acid (DTPA), derivatives of S-2-(4-
15 Isothiocyanatobenzyl)-1,4,7-triazacyclononane-1,4,7-triacetic acid (NOTA) and derivatives of 1,4,8,11-tetraazacyclododecane-1,4,8,11-tetraacetic acid (TETA).

17. An isolated nucleic acid molecule encoding an antibody or antigen-binding fragment thereof according to any preceding claim.

18. A nucleic acid molecule according to Claim 17, encoding an antibody heavy chain or
20 variable region thereof comprising or consisting of a nucleotide sequence of SEQ ID NO: 9 or a nucleotide sequence of SEQ ID NO: 27.

19. A nucleic acid molecule according to Claim 17, encoding an antibody light chain or variable region thereof comprising or consisting of a nucleotide sequence of SEQ ID NO: 10 or a nucleotide sequence of SEQ ID NO: 28.

20. A vector comprising a nucleic acid molecule according to any of Claims 17 to 19,
25 optionally in which the vector is an expression vector.

21. A recombinant host cell comprising a nucleic acid molecule according to any of Claims 17 to 19 or a vector according to Claim 20, optionally in which the host cell is a bacterial cell or a mammalian cell; optionally, in which the mammalian cell is a human cell.

22. A method for producing an antibody or antigen-binding fragment according to any of Claims 1 to 16, the method comprising culturing a host cell as defined in Claim 21 under conditions which permit expression of the encoded antibody or antigen-binding fragment thereof.
- 5 23. A pharmaceutical composition comprising an effective amount of an antibody or antigen-binding fragment thereof according to any of Claims 1 to 16 and a pharmaceutically-acceptable diluent, carrier or excipient.
- 10 24. An antibody or antigen-binding fragment thereof according to any of Claims 1 to 16 for use in the treatment of cancer, in which the cancer is preferably selected from the group consisting of: prostate cancer; breast cancer; colorectal cancer; kidney cancer; pancreatic cancer; ovarian cancer; lung cancer; cervical cancer; rhabdomyosarcoma; neuroblastoma; bone cancer; multiple myeloma; leukemia (such as acute lymphoblastic leukemia [ALL] and acute myeloid leukemia [AML]), skin cancer (e.g. melanoma), bladder cancer and glioblastoma.
- 15 25. Use of an antibody or antigen-binding fragment according to any of Claims 1 to 16 in the manufacture of a medicament for the treatment of cancer.
- 20 26. Use according to Claim 25 in which the cancer is selected from the group consisting of: prostate cancer; breast cancer; colorectal cancer; kidney cancer; pancreatic cancer; ovarian cancer; lung cancer; cervical cancer; rhabdomyosarcoma; neuroblastoma; bone cancer; multiple myeloma; leukemia (such as acute lymphoblastic leukemia [ALL] and acute myeloid leukemia [AML]), skin cancer (e.g. melanoma), bladder cancer and glioblastoma; optionally in which the medicament is for treating a patient who has been pre-screened and identified as having a tumour with cells expressing CD137 and FcγR.
- 25 27. A combination therapy comprising an antibody or antigen-binding fragment thereof according to any of Claims 1 to 16 and one or more further therapeutic agents; optionally in which the one or more further therapeutic agents are cancer therapies.