

DESCRIPTION

GLYCINE TRANSPORTER-INHIBITING SUBSTANCES

TECHNICAL FIELD

[0001] The present invention relates to compounds having a glycine transporter-inhibiting action.

BACKGROUND ART

[0002] The NMDA receptor, which is one of glutamate receptors, is located on the nerve cell membranes in the brain and involved in various neurophysiologic events such as neuronal plasticity, cognition, attention, and memory. The NMDA receptor has a plurality of allosteric binding sites, one of which is the glycine binding site (glycine binding site on NMDA receptor complex). It has been reported that the glycine binding site on NMDA receptor complex is involved in the activation of NMDA receptors (Non-Patent Document 1).

[0003] Action potential arriving at the presynaptic terminals of glycinergic nerves triggers the release of glycine into synaptic clefts. The released glycine binds to the postsynaptic receptors or the like and is then removed from the synaptic clefts by transporters. Based on this fact, glycine transporters are believed to regulate the functions of NMDA receptors through regulation of the amount of glycine in the extracellular fluid.

[0004] Glycine transporters (GlyTs) are proteins involved in the reuptake of extracellular glycine into cells, and two subtypes, GlyT1 and GlyT2, have so far been identified. GlyT1, which is expressed primarily in the cerebral cortex, hippocampus, thalamus and the like, has been reported to be associated with diseases such as schizophrenia, Alzheimer's disease, cognitive impairment, dementia, anxiety disorders (e.g., generalized anxiety disorder, panic disorder, obsessive-compulsive disorder, social anxiety disorder, post-traumatic stress disorder, specific phobias, acute stress disorder), depression, drug dependence, spasm, tremor, pain, Parkinson's disease, attention deficit hyperactivity disorder, bipolar disorder, eating disorder, and sleep disorders (Non-Patent Documents 2-4).

[0005] Compounds having a GlyT1-inhibiting action and having an imidazolidin-2-one

structure have been reported in the documents shown below (Patent Documents 1 and 2).

These compounds described in Patent Documents 1 and 2 are characterized in that a phenyl group is attached via amide or carbonyl to one of the endocyclic nitrogen atoms of the imidazolidine, while another phenyl group is attached to the other endocyclic nitrogen atom of the imidazolidine, and that an endocyclic carbon atom of the imidazolidinone is a spiro carbon atom.

CITATION LIST

PATENT DOCUMENTS

[0006] Patent Document 1: WO2008092878

Patent Document 2: WO2009034062

NON-PATENT DOCUMENTS

[0007] Non-Patent Document 1: Molecular Psychiatry (2004) 9, 984-997

Non-Patent Document 2: Current Medicinal Chemistry, 2006, 13, 1017-1044

Non-Patent Document 3: Neuropsychopharmacology (2005), 30, 1963-1985

Non-Patent Document 4: Expert Opinion on Therapeutic Patents (2004) 14 (2) 201-

214

SUMMARY OF INVENTION

TECHNICAL PROBLEM

[0008] The present invention aims to provide novel compounds or pharmaceutically acceptable salts thereof which are useful in the prevention or treatment of diseases such as schizophrenia, Alzheimer's disease, cognitive impairment, dementia, anxiety disorders (e.g., generalized anxiety disorder, panic disorder, obsessive-compulsive disorder, social anxiety disorder, post-traumatic stress disorder, specific phobias, acute stress disorder), depression, drug dependence, spasm, tremor, pain, Parkinson's disease, attention deficit hyperactivity disorder, bipolar disorder, eating disorder, or sleep disorders, which is based on the glycine uptake-inhibiting action.

SOLUTION TO PROBLEM


[0009] As a result of extensive and intensive studies on structurally novel compounds with

an inhibitory action against GlyT1, the present inventors found that the compounds represented by the following formula, in which a nitrogen-containing aromatic ring group is attached to one of the endocyclic nitrogen atoms of the imidazolidine and the endocyclic carbon atoms of the imidazolidinone are not spiro carbon atoms, are superior GlyT1-inhibiting substances. This finding has led to the completion of the present invention.

[0010] The present invention will be described below in detail. Embodiments of the present invention (hereinafter each referred to as “the inventive compound”) are as shown below.

(1) A compound of formula [I] or a pharmaceutically acceptable salt thereof:

[0011] [Formula 1]



[0012] wherein

R^1 and $R^{1'}$ are the same or different, and each represent a hydrogen atom, a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a cyano group, a heteroaryl group (which may be substituted by a C_{1-6} alkyl group), a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a C_{1-6} alkylamino group, or the formula $CONR^7R^8$ (R^7 and R^8 are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group),

R^2 represents a hydrogen atom or a C_{1-6} alkyl group,

R^3 represents a phenyl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, cyano groups, C_{1-6} alkyl groups, C_{1-6} alkoxy groups, C_{1-6} alkylamino groups, C_{1-6} alkylsulfonyl groups, halo C_{1-6} alkyl groups, halo C_{1-6} alkoxy groups, halo C_{1-6} alkylsulfanyl groups, phenyl groups, phenoxy groups, heteroaryl groups (which may be substituted by a C_{1-6} alkyl group), and the formula $-SO_2NR^9R^{10}$ (R^9 and R^{10} are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group)) or a heteroaryl group or a bicyclic heteroaryl group (the each heteroaryl group may be substituted by 1 to 3 substituents selected from halogen atoms, C_{1-6} alkyl groups, C_{3-6} cycloalkyl groups, C_{1-6}

alkoxy groups, cyano groups, C₁₋₆ alkanoyl groups, and haloC₁₋₆ alkyl groups),

R⁴ represents a C₁₋₆ alkyl group (which may be substituted by 1 to 3 halogen atoms, a C₁₋₆ alkoxy group, a C₃₋₆ cycloalkyl group, or a phenyl group), a C₃₋₆ cycloalkyl group, or a phenyl group,

R⁵ and R⁶ are the same or different, and each represent a hydrogen atom or a C₁₋₆ alkyl group, and

A¹, A², A³, and A⁴ are the same or different, and each represent the formula CH or a nitrogen atom, provided that one or two of A¹, A², A³, and A⁴ represent a nitrogen atom.

(2) The compound or pharmaceutically acceptable salt thereof according to (1), wherein

R¹ is a hydrogen atom, a halogen atom, a C₁₋₆ alkoxy group, a haloC₁₋₆ alkyl group, a cyano group, a heteroaryl group (which may be substituted by a C₁₋₆ alkyl group), a C₁₋₆ alkyl group, a C₁₋₆ alkylamino group, or the formula CONR⁷R⁸ (R⁷ and R⁸ are the same or different, and each represent a hydrogen atom or a C₁₋₆ alkyl group),

R^{1'} is a hydrogen atom,

R³ is a phenyl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, cyano groups, C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, C₁₋₆ alkylamino groups, C₁₋₆ alkylsulfonyl groups, haloC₁₋₆ alkyl groups, haloC₁₋₆ alkoxy groups, haloC₁₋₆ alkylsulfanyl groups, phenyl groups, phenoxy groups, heteroaryl groups (which may be substituted by a C₁₋₆ alkyl group), and the formula -SO₂NR⁹R¹⁰ (R⁹ and R¹⁰ are the same or different, and each represent a hydrogen atom or a C₁₋₆ alkyl group)) or a heteroaryl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₁₋₆ alkoxy groups, cyano groups, and haloC₁₋₆ alkyl groups), and

R⁴ is a C₁₋₆ alkyl group (which may be substituted by a C₃₋₆ cycloalkyl group or a phenyl group) or a phenyl group.

(3) The compound or pharmaceutically acceptable salt thereof according to (1), wherein R⁴ is a C₁₋₆ alkyl group which may be substituted by 1 to 3 halogen atoms.

(4) The compound or pharmaceutically acceptable salt thereof according to (1) or (2), wherein R⁴ is a C₁₋₆ alkyl group.

(5) The compound or pharmaceutically acceptable salt thereof according to any one of (1) to (4), wherein

R^2 is a hydrogen atom, and

R^5 and R^6 are both a hydrogen atom.

(6) The compound or pharmaceutically acceptable salt thereof according to (5), wherein

R^1 is a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a cyano group, a heteroaryl group (which may be substituted by a C_{1-6} alkyl group), a C_{1-6} alkyl group, a C_{1-6} alkylamino group, or the formula $CONR^7R^8$ (R^7 and R^8 are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group), and

$R^{1'}$ is a hydrogen atom.

(7) The compound or pharmaceutically acceptable salt thereof according to any one of (1) and (3) to (5), wherein

R^1 is a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a C_{1-6} alkyl group, or a C_{3-6} cycloalkyl group, and

$R^{1'}$ is a hydrogen atom, a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a C_{1-6} alkyl group, or a C_{3-6} cycloalkyl group.

(8) The compound or pharmaceutically acceptable salt thereof according to any one of (1) to (7), wherein R^1 is attached in the para position.

(9) The compound or pharmaceutically acceptable salt thereof according to any one of (1) to (8), wherein any one of A^1 , A^2 , A^3 and A^4 is a nitrogen atom or A^1 and A^3 are both a nitrogen atom.

(10) The compound or pharmaceutically acceptable salt thereof according to any one of (1) to (8), wherein

A^1 is a nitrogen atom,

A^2 and A^4 are both the formula CH, and

A^3 is the formula CH or a nitrogen atom.

(11) The compound or pharmaceutically acceptable salt thereof according to any one of (1) and (3) to (10), wherein R^3 is a heteroaryl group (which may be substituted by 1 to 3

substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₃₋₆ cycloalkyl groups, C₁₋₆ alkoxy groups, cyano groups, C₁₋₆ alkanoyl groups, and haloC₁₋₆ alkyl groups).

(12) The compound or pharmaceutically acceptable salt thereof according to any one of (1) and (3) to (10), wherein R³ is a pyridyl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₃₋₆ cycloalkyl groups, C₁₋₆ alkoxy groups, cyano groups, C₁₋₆ alkanoyl groups, and haloC₁₋₆ alkyl groups).

(13) The compound or pharmaceutically acceptable salt thereof according to any one of (1) and (3) to (10), wherein R³ is a pyridyl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₃₋₆ cycloalkyl groups, and haloC₁₋₆ alkyl groups).

(14) The compound or pharmaceutically acceptable salt thereof according to (1), wherein the compound is selected from the group consisting of:

2-[(5S)-3-(5-methoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-3-(5-ethylpyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]acetamide,

2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]-N-(4-ethylpyridin-2-yl)acetamide,

2-[(5S)-3-(5-fluoropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-3-(5-fluoropyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-cyclopropylpyridin-2-yl)-2-[(5S)-2-oxo-5-(propan-2-yl)-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-2-oxo-5-(propan-2-yl)-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]acetamide,

2-[(5S)-3-(5-cyclopropylpyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-3-(5-cyclopropylpyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-cyclopropylpyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-chloropyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-ethylpyridin-2-yl)acetamide,

N-(4-chloropyridin-2-yl)-2-[(5R)-5-[(1S)-1-fluoropropyl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5R)-3-(5-chloropyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl]acetamide,

N-(4-cyclopropylpyridin-2-yl)-2-[(5R)-5-[(1S)-1-fluoropropyl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]acetamide,

2-[(5R)-5-[(1S)-1-fluoropropyl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5R)-3-(5-chloropyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5R)-5-[(1S)-1-fluoropropyl]-3-(5-fluoropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-{(5R)-3-(5-cyclopropylpyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl}-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-{(5R)-3-(5-ethoxypyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl}-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-cyclopropylpyridin-2-yl)-2-{(5R)-3-(5-ethoxypyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl}acetamide,

N-(4-chloropyridin-2-yl)-2-{(5R)-3-(5-ethoxypyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl}acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-fluoropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-{(5S)-5-[(2S)-butan-2-yl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}-N-(4-cyclopropylpyridin-2-yl)acetamide,

2-{(5S)-5-[(2S)-butan-2-yl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}-N-(4-chloropyridin-2-yl)acetamide,

2-[(5S)-3-(5-ethoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-3-(5-ethoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide,

N-(4-cyclopropylpyridin-2-yl)-2-[(5S)-3-(5-ethoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-ethoxypyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-cyclopropylpyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(5-chloropyridin-2-yl)-2-{(5S)-2-oxo-5-propyl-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}acetamide,

N-(5-chloro-6-methylpyridin-2-yl)-2-{(5S)-2-oxo-5-propyl-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}acetamide,

N-(5-chloro-6-methylpyridin-2-yl)-2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]acetamide,

2-[(5S)-2-oxo-5-(propan-2-yl)-3-[5-(propan-2-yloxy)pyrimidin-2-yl]imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-ethoxypyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-chloropyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-cyclopropylpyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-chloropyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-ethoxypyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-cyclopropylpyridin-2-yl)acetamide, and

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-cyclopropylpyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-cyclopropylpyridin-2-yl)acetamide.

(15) A pharmaceutical composition comprising, as an active ingredient, the compound or pharmaceutically acceptable salt thereof according to any one of (1) to (14).

(16) An agent for preventing or treating diseases of schizophrenia, Alzheimer's disease, cognitive impairment, dementia, anxiety disorders, depression, drug dependence, spasm, tremor, pain, Parkinson's disease, attention deficit hyperactivity disorder, bipolar disorder, eating disorder, or sleep disorders, which comprises, as an active ingredient, the compound or pharmaceutically acceptable salt thereof according to any one of (1) to (14).

ADVANTAGEOUS EFFECTS OF INVENTION

[0013] The inventive compounds have glycine transporter (GlyT1)-inhibiting activity.

DESCRIPTION OF EMBODIMENTS

[0014] The term "C_{x-y} (x and y each denote a natural number)" as used herein means that the number of carbon atoms is x to y.

[0015] The term "C₁₋₆ alkyl group" as used herein refers to a straight-chain or branched-chain alkyl group having 1 to 6 carbon atoms, and includes, for example, a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, an isobutyl group, a tert-butyl group, a pentyl group, an isopentyl group, and a hexyl group.

[0016] The term “C₃₋₆ cycloalkyl group” as used herein refers to a cycloalkyl group having 3 to 6 carbon atoms, which is a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, or a cyclohexyl group.

[0017] The term “C₁₋₆ alkoxy group” as used herein refers to a straight-chain or branched-chain alkoxy group having 1 to 6 carbon atoms, and includes, for example, a methoxy group, an ethoxy group, a propoxy group, an isopropoxy group, a butoxy group, an isobutoxy group, a pentyloxy group, an isopentyloxy group, and a hexyloxy group.

[0018] The term “C₁₋₆ alkanoyl group” as used herein refers to a straight-chain or branched-chain alkanoyl group having 1 to 6 carbon atoms, and includes, for example, a formyl group, an acetyl group, a propanoyl group, a butanoyl group, and a pivaloyl group.

[0019] The term “halogen (halo)” as used herein refers to fluorine, chlorine, bromine, or iodine.

[0020] The term “haloC₁₋₆ alkyl group” as used herein refers to a straight-chain or branched-chain alkyl group which has 1 to 6 carbon atoms and which has been substituted by a halogen atom or halogen atoms. The preferred number of the substituting halogen atom(s) is 1 to 3. Examples of the haloC₁₋₆ alkyl group include a fluoromethyl group, a difluoromethyl group, a trifluoromethyl group, and a trichloromethyl group. Among these groups, a trifluoromethyl group is preferred.

[0021] The term “C₁₋₆ alkylamino group” as used herein refers to a group characterized in that 1 or 2 straight-chain or branched-chain alkyl groups each having 1 to 6 carbon atoms are attached to an amino group. Examples of the C₁₋₆ alkylamino group include a methylamino group, a dimethylamino group, a diethylamino group, an N-ethyl-N-methylamino group, and the like.

[0022] The term “C₁₋₆ alkylamine” as used herein refers to an amine which has one or two straight-chain or branched-chain alkyl groups having 1 to 6 carbon atoms. Examples of the C₁₋₆ alkylamine include methylamine, dimethylamine, diethylamine, N-ethyl-N-methylamine, and the like.

[0023] The term “C₁₋₆ alkylsulfonyl group” as used herein refers to a straight-chain or

branched-chain alkylsulfonyl group having 1 to 6 carbon atoms, and includes, for example, a methylsulfonyl group, an ethylsulfonyl group, a propylsulfonyl group, an isopropylsulfonyl group, a butylsulfonyl group, an isobutylsulfonyl group, a tert-butylsulfonyl group, a pentylsulfonyl group, an isopentylsulfonyl group, and a hexylsulfonyl group.

[0024] The term “haloC₁₋₆ alkylsulfanyl group” as used herein refers to a straight-chain or branched-chain alkylsulfanyl group which has 1 to 6 carbon atoms and which has been substituted by a halogen atom or halogen atoms. The preferred number of the substituting halogen atom(s) is 1 to 3. Examples of the haloC₁₋₆ alkylsulfanyl group include a fluoromethylsulfanyl group, a difluoromethylsulfanyl group, a trifluoromethylsulfanyl group, and a trichloromethylsulfanyl group.

[0025] The term “haloC₁₋₆ alkoxy group” as used herein refers to a straight-chain or branched-chain alkoxy group which has 1 to 6 carbon atoms and which has been substituted by a halogen atom or halogen atoms. The preferred number of the substituting halogen atom(s) is 1 to 3. Examples of the haloC₁₋₆ alkoxy group include a fluoromethoxy group, a difluoromethoxy group, and a trifluoromethoxy group.

[0026] The term “heteroaryl group” as used herein refers to a monocyclic heteroaryl group having in the ring at least one atom selected from the group consisting of a nitrogen atom, an oxygen atom, and a sulfur atom. When the monocyclic heteroaryl group has a nitrogen atom or nitrogen atoms in the ring, the each nitrogen atom may be an N-oxide.

[0027] The heteroaryl group is preferably a 5- or 6-membered heteroaryl group, and includes, for example, a pyridyl group, a pyridazyl group, a pyrimidyl group, a pyrazyl group, a pyrazolyl group, a thiazolyl group, an imidazolyl group, an oxazolyl group, an isoxazolyl group, a thienyl group, a triazolyl group, an oxadiazolyl group, and a thiadiazolyl group.

[0028] The term “bicyclic heteroaryl group” as used herein refers to a bicyclic heteroaryl group having in the ring at least one atom selected from the group consisting of a nitrogen atom, an oxygen atom, and a sulfur atom. When the bicyclic heteroaryl group has a nitrogen atom or nitrogen atoms in the ring, the each nitrogen atom may be an N-oxide.

[0029] The bicyclic heteroaryl group is preferably a 9- or 10-membered heteroaryl group, and includes, for example, a quinolyl group, an isoquinolyl group, and an indolyl group.

[0030] The term "pharmaceutically acceptable salt" as used herein refers to an acid addition salt that may be accepted in pharmaceutical terms. Examples of the acid that may be used include inorganic acids such as sulfuric acid, hydrochloric acid, hydrobromic acid, nitric acid and phosphoric acid, and organic acids such as acetic acid, oxalic acid, lactic acid, citric acid, malic acid, gluconic acid, tartaric acid, fumaric acid, maleic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid and p-toluenesulfonic acid. The free forms may be converted to these salts in a conventional manner.

[0031] In connection with the inventive compounds, preferred embodiments will be shown below.

Preferred compounds are those wherein R^1 is a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a cyano group, a heteroaryl group (which may be substituted by a C_{1-6} alkyl group), a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a C_{1-6} alkylamino group, or the formula $CONR^7R^8$ (R^7 and R^8 are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group). More preferred compounds are those wherein R^1 is a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a C_{1-6} alkyl group, or a C_{3-6} cycloalkyl group. The halo C_{1-6} alkyl group is more preferably a trifluoromethyl group, and the halogen atom is more preferably a chlorine atom. Compounds wherein R^1 is attached in the para position are preferred.

Preferred compounds are those wherein $R^{1'}$ is a hydrogen atom, a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a C_{1-6} alkyl group, or a C_{3-6} cycloalkyl group. More preferred compounds are those wherein $R^{1'}$ is a hydrogen atom or a halogen atom. When $R^{1'}$ is an atom or group other than a hydrogen atom, $R^{1'}$ is preferably attached in the ortho position.


Preferred compounds are those wherein R^2 is a hydrogen atom.

Preferred compounds are those wherein R^3 is a phenyl group (which has been substituted by 1 to 3 substituents selected from halogen atoms, cyano groups, C_{1-6} alkyl

groups, C₁₋₆ alkoxy groups, C₁₋₆ alkylamino groups, C₁₋₆ alkylsulfonyl groups, haloC₁₋₆ alkyl groups, haloC₁₋₆ alkoxy groups, haloC₁₋₆ alkylsulfonyl groups, phenyl groups, phenoxy groups, heteroaryl groups (which may be substituted by a C₁₋₆ alkyl group), and the formula -SO₂NR⁹R¹⁰ (R⁹ and R¹⁰ are the same or different, and each represent a hydrogen atom or a C₁₋₆ alkyl group)) or a heteroaryl group (which has been substituted by 1 to 3 substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₃₋₆ cycloalkyl groups, C₁₋₆ alkoxy groups, cyano groups, C₁₋₆ alkanoyl groups, and haloC₁₋₆ alkyl groups). More preferred compounds are those wherein R³ is a pyridyl group (which has been substituted by 1 to 3 substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₃₋₆ cycloalkyl groups, C₁₋₆ alkoxy groups, cyano groups, C₁₋₆ alkanoyl groups, and haloC₁₋₆ alkyl groups). Still more preferred compounds are those wherein R³ is a pyridyl group (which has been substituted by 1 to 3 substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₃₋₆ cycloalkyl groups, and haloC₁₋₆ alkyl groups). In the embodiments of R³, the haloC₁₋₆ alkyl group mentioned as a substituent by which a phenyl group or a heteroaryl group (more preferably, a pyridyl group) is substituted is more preferably a trifluoromethyl group, and the halogen atom also mentioned as a substituent is more preferably a chlorine atom. The pyridyl group is preferably a pyridin-2-yl group having a substituent in the 4-position.

Preferred compounds are those wherein R⁴ is a C₁₋₆ alkyl group which may be substituted by 1 to 3 halogen atoms, and more preferred compounds are those wherein R⁴ is a branched-chain C₁₋₆ alkyl group or a straight-chain C₁₋₆ alkyl group substituted by 1 to 3 halogen atoms. The configuration of the carbon atom to which R⁴ is attached is preferably as shown below.

[0032] [Formula 2]



[0033] Preferred compounds are those wherein R^5 and R^6 are both a hydrogen atom.

Preferred compounds are those wherein any one of A^1 , A^2 , A^3 and A^4 is a nitrogen atom or A^1 and A^3 are both a nitrogen atom. More preferred compounds are those wherein A^1 is a nitrogen atom, A^2 and A^4 are both the formula CH, and A^3 is the formula CH or a nitrogen atom.

[0034] The inventive compounds may contain a plurality of asymmetric centers. Thus, the aforementioned compounds may exist not only in optically active forms but also in their racemates. Further, a plurality of diastereomers may also exist. All of these forms are included in the scope of the present invention. Individual isomers may be obtained by known methods, for example, by use of optically active starting materials or intermediates, by an optically selective reaction or a diastereoselective reaction in the preparation of intermediates or final products, or by chromatographic separation or the like in the preparation of intermediates or final products. If the inventive compounds form hydrates or solvates, these hydrates or solvates are also included in the scope of the present invention. Likewise, pharmaceutically acceptable salts of hydrates or solvates of the inventive compounds are also included in the scope of the present invention.

[0035] The compounds according to the present invention may be administered orally or parenterally. The dosage forms are tablets, capsules, granules, dispersions, powders, lozenges, ointments, creams, emulsions, suspensions, suppositories, injections and the like, all of which may be produced by conventional formulation techniques (for example, the methods set forth in the 15th revised Japanese Pharmacopoeia). These dosage forms may be selected as appropriate, according to the symptoms and age of patients and the purpose of treatment.

[0036] To produce these preparations, a composition containing the compound of the present invention may be blended with one or more pharmacologically acceptable carriers, namely, excipients (e.g., crystalline cellulose, starch, lactose, mannitol), binders (e.g., hydroxypropylcellulose, polyvinylpyrrolidone), lubricants (e.g., magnesium stearate, talc), disintegrants (e.g., carboxymethylcellulose calcium), and/or various other pharmacologically

acceptable additives.

[0037] The compounds of the present invention may be used in combination with one or more other therapeutic agents, namely, various antipsychotics, antidepressants, for example, 5HT₃ antagonists, 5HT₂ antagonists, serotonin agonists, NK-1 antagonists, selective serotonin reuptake inhibitors (SSRIs), serotonin noradrenaline reuptake inhibitors (SNRIs), tricyclic antidepressants, dopaminergic antidepressants, H₃ antagonists, 5HT_{1A} antagonists, 5HT_{1B} antagonists, 5HT_{1D} antagonists, D₁ agonists, M₁ agonists, anticonvulsants, cognitive enhancement drugs, and other psychoactive drugs.

[0038] Examples of other therapeutic agents that may be used in combination with the compounds of the present invention include ondansetron, granisetron, metoclopramide, sumatriptan, rauwolscine, yohimbine, fluoxetine, citalopram, escitalopram, femoxetine, fluvoxamine, paroxetine, indalpine, sertraline (registered trademark), zimeldine, venlafaxine, reboxetine, Milnacipran, duloxetine, imipramine, amitriptyline, chlomipramine, nortriptyline, bupropion, amineptine, divalproex, carbamazepine, diazepam, risperidone, olanzapine, ziprasidone, aripiprazole, quetiapine, perospirone, clozapine, haloperidol, pimozide, droperidol, chlorpromazine, thioridazine, mesoridazine, trifluoperazine, perphenazine, fluphenazine, thiflupromazine, prochlorperazine, acetophenazine, thiothixene, chlorprothixene, lamotrigine, loxapine, molindone, and the like. Such combinations may be administered simultaneously (in the same pharmaceutical formulation or in different pharmaceutical formulations), separately, or sequentially.

[0039] Particular advantages associated with the use of, and methods for treatment with, combinations of the compounds of the present invention include comparable or improved effects achieved by using individual ingredients at lower doses than their usual doses. Such use and treatment methods are also expected to further enhance the therapeutic effects on positive and/or negative symptoms of psychiatric disorders and/or cognitive impairment. The use of and methods for treatment with combinations of the compounds of the present invention also may provide benefits in the treatment of patients who do not sufficiently respond to, or who are resistant to, treatment with some type of neuroleptic.

[0040] The compounds according to the present invention may be administered in doses which, in the case of treating adults, range from 1 to 2000 mg per day, either once daily or in divided portions. The dose may be increased or decreased as appropriate, depending on the age, body weight and symptom of a patient.

[0041] The compounds of formula [I] may be produced by various methods of synthesis. The methods described below are only illustrative of the process for producing the inventive compounds and should not be taken as limiting.

[0042] In the general production processes, the term “inert solvent” refers to, for example, an alcohol such as methanol, ethanol, isopropanol, n-butanol, or ethylene glycol; an ether such as diethyl ether, t-butyl methyl ether, diisopropyl ether, tetrahydrofuran, 1,4-dioxane, or 1,2-dimethoxyethane; a hydrocarbon such as pentane, hexane, heptane, toluene, benzene, or xylene; an ester such as ethyl acetate or ethyl formate; a ketone such as acetone or methyl ethyl ketone; a halogenated carbon-based solvent such as chloroform or dichloromethane; an amide such as dimethylformamide or N-methylpyrrolidone; acetonitrile; dimethyl sulfoxide; water; or a mixed solvent thereof.

[0043] The term “base” refers to, for example, an alkali metal or alkaline earth metal hydride such as lithium hydride, sodium hydride, potassium hydride, or calcium hydride; an alkali metal or alkaline earth metal amide such as lithium amide, sodium amide, lithium diisopropylamide, lithium dicyclohexylamide, lithium hexamethyldisilazide, sodium hexamethyldisilazide, or potassium hexamethyldisilazide; an alkali metal or alkaline earth metal lower alkoxide such as sodium methoxide, sodium ethoxide, or potassium tert-butoxide; an alkyl lithium such as butyl lithium, sec-butyl lithium, tert-butyl lithium, or methyl lithium; an alkali metal or alkaline earth metal hydroxide such as sodium hydroxide, potassium hydroxide, lithium hydroxide, or barium hydroxide; an alkali metal or alkaline earth metal carbonate such as sodium carbonate, potassium carbonate, or cesium carbonate; an alkali metal or alkaline earth metal hydrogencarbonate such as sodium hydrogen carbonate or potassium hydrogen carbonate; an amine such as triethylamine, N-methylmorpholine, N,N-diisopropylethylamine, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), 1,5-

diazabicyclo[4.3.0]non-5-ene (DBN), or N,N-dimethylaniline; or a basic heterocyclic compound such as pyridine, imidazole, or 2,6-lutidine. These bases are selected as appropriate, according to various reaction conditions known to skilled artisans.


[0044] The term “acid” refers to, for example, an inorganic acid such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, or phosphoric acid; or an organic acid such as p-toluenesulfonic acid, methanesulfonic acid, trifluoroacetic acid, formic acid, acetic acid, citric acid, or oxalic acid. These acids are selected as appropriate, according to various reaction conditions known to skilled artisans.

[0045] In the general production processes, X^1 represents a halogen atom or a hydroxyl group; X^2 represents a chlorine atom, a bromine atom, an iodine atom, or a trifluoromethanesulfonyloxy group; P^1 represents an ester-protective group such as a methyl group or a benzyl group (refer to Theodora W. Green, Peter G. M. Wuts, “Green’s Protective Groups in Organic Synthesis, Fourth Edition”, Wiley Interscience); P^2 represents a nitrogen atom-protective group such as a tert-butoxycarbonyl group or a benzyloxycarbonyl group (refer to the same reference as mentioned above); R^{1a} represents a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a halo C_{1-6} alkyl group, or a heteroaryl group; R^{1b} represents a C_{1-6} alkylamino group or a heteroaryl group; and the other symbols are as defined above.

General production process 1

[0046]

[Formula 3]



[0047] Step 1: Compound (1) and compound (2) where X^1 is a halogen atom may be reacted in an inert solvent in the presence or absence of a base, to obtain compound (3). Alternatively, a Mitsunobu reaction between compound (1) and compound (2) where X^1 is a hydroxyl group may be performed using either an organophosphorus compound and an azo compound or a phosphorus ylide reagent, in an inert solvent in the presence or absence of a base, to obtain compound (3). Examples of the organophosphorus compound include triphenylphosphine, tributylphosphine, and the like. Examples of the azo compound include azodicarboxylic acid diethyl, azodicarboxylic acid diisopropyl, azodicarboxylic acid di-tert-butyl, and the like. Examples of the phosphorus ylide reagent include cyanomethylene tributylphosphorane and the like.


Step 2: A deprotection reaction described in Theodora W. Green, Peter G. M. Wuts, "Green's Protective Groups in Organic Synthesis, Fourth Edition" may be performed to obtain compound (4).

Step 3: An amidation reaction of compound (4) with compound (5) may be performed in an inert solvent in the presence or absence of a base, to obtain the inventive compound [I]. Such an amidation reaction may be performed in accordance with many standard procedures known to skilled artisans, and includes, for example, amidation via a

mixed acid anhydride using ethyl chlorocarbonate, isobutyl chlorocarbonate, pivaloyl chloride or the like; amidation via an acid chloride using oxalyl chloride, thionyl chloride or the like; and amidation using a condensing agent such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC•HCl), 1,3-dicyclohexylcarbodiimide (DCC), diphenylphosphoryl azide (DPPA), diethyl phosphorocyanidate, carbonyldiimidazole (CDI), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), or benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP reagent). To perform the amidation reaction using a condensing agent, an additive such as 1-hydroxybenzotriazole (HOBt) or hydroxysuccinimide (HOSu) may be used as needed.

General production process 2

[0048] [Formula 4]




[0049] Step 4: Compound (1) and compound (6) may be reacted in the same manner as shown in Step 1 of General production process 1, to obtain the inventive compound [I].

Compound (1) mentioned above may be produced in accordance with the process described below.

General production process 3

[0050]

[Formula 5]



[0051] Step 5: A general oxidation reaction of an alcohol into an aldehyde may be performed using an oxidizing agent in an inert solvent to obtain compound (8). Examples of the oxidation reaction include processes using an oxidizing agent such as IBX, TEMPO, PCC, or PDC; Swern oxidation; and the like.

Step 6: Compound (8) and compound (9) may be subjected to a reductive amination reaction using a reducing agent in an inert solvent in the presence or absence of an acid, to obtain compound (10). Examples of the reducing agent include sodium triacetoxyborohydride, sodium cyanoborohydride, sodium borohydride, and the like.

Step 7: A deprotection reaction described in Theodora W. Green, Peter G. M. Wuts, "Green's Protective Groups in Organic Synthesis, Fourth Edition" may be performed to obtain compound (11).


Step 8: Compound (11) may be cyclized using a reagent such as triphosgene, phosgene, or carbonyldiimidazole, in an inert solvent in the presence or absence of a base, to obtain compound (1).

Compound (1) mentioned above may also be produced in accordance with the process described below.

General production process 4

[0052]

[Formula 6]




[0053] Step 9: A urea formation reaction may be performed by reacting compound (12) with, for example, compound (13) or isocyanate in an inert solvent in the presence or absence of a base, to obtain compound (14).

Step 10: Compound (14) may be subjected to an intramolecular cyclization reaction to obtain compound (1). Examples of the intramolecular cyclization reaction that occurs in this case include Mitsunobu reactions using either an organophosphorus compound and an azo compound or a phosphorus ylide reagent. The intramolecular cyclization reaction may also be performed after conversion of the hydroxyl group of compound (14) to a leaving group by mesylation, tosylation, halogenation or the like in the presence of a base.

Compound (1) mentioned above may also be produced in accordance with the process described below.

General production process 5

[0054] [Formula 7]



[0055] Step 11: Compound (15) may be cyclized using a reagent such as triphosgene, phosgene, carbonyldiimidazole, or 4-nitrophenyl chloroformate, in an inert solvent in the

presence or absence of a base, to obtain compound (16).

Step 12: Compound (16) may be reacted with a reducing agent in an inert solvent to obtain compound (17). Examples of the reducing agent include lithium aluminum hydride, sodium bis(2-methoxyethoxy)aluminum hydride, and the like. Further, heating and stirring or use of aluminum trichloride, as needed, is preferred.


Step 13: The same process as shown in Step 12 may be performed to obtain compound (18) from compound (15).

Step 14: The same process as shown in Step 11 may be performed to obtain compound (17) from compound (18).

Step 15: Compound (17) and compound (19) may be reacted using a palladium catalyst or a copper catalyst and, if needed, a ligand of such a metal catalyst, in an inert solvent in the presence or absence of a base, to obtain compound (1). Examples of the palladium catalyst include $\text{Pd}(\text{OAc})_2$, $\text{Pd}_2(\text{dba})_3$, $\text{Pd}(\text{PPh}_3)_4$ and the like, and examples of the copper catalyst include CuI , CuBr and the like. Examples of the ligand of the palladium catalyst include triphenylphosphine, Xantphos, BINAP and the like, and examples of the ligand of the copper catalyst include N,N' -dimethylethylenediamine, 1,2-cyclohexanediamine, phenanthroline, proline and the like.

General production process 6

[0056] [Formula 8]




[0057] Step 16: Compound (20) may be reacted with compound (21) using a metal catalyst such as palladium, copper, iron or nickel and, if needed, a ligand, in an inert solvent in the presence or absence of a base, to obtain the inventive compound [I2]. Compound (21) represents an organometallic reagent and includes, for example, Grignard reagents (e.g., $R^{1a}MgCl$), zinc reagents (e.g., $R^{1a}ZnCl$), boron reagents (e.g., those where R^{1a} is attached to boric acid or a boric acid ester), and tin reagents (e.g., $R^{1a}SnBu_3$). Examples of the iron reagent include tris(2,4-pentanedionato)iron(III), and examples of the nickel reagent include 1,2-bis(diphenylphosphino)ethane nickel(II) chloride and the like.

Step 17: Compound (20) may be reacted with, for example, a C_{1-6} alkylamine or a heteroaryl group having an NH group in the ring, using a metal catalyst such as palladium or copper and, if needed, a ligand, in an inert solvent in the presence or absence of a base, to obtain the inventive compound [I3].

General production process 7

[0058] [Formula 9]



[0059] Step 18: The same process as shown in Step 15 may be performed to obtain the inventive compound [I] from compound (22).

EXAMPLES

[0060] Next, the present invention will be described in more detail by means of Production Examples, Working Examples and Test Example, but these Examples are in no way intended to limit the scope of the present invention.

[0061] The microwave reaction apparatus used in the Production Examples and Working Examples described below was Initiator from Biotage.

[0062] In the Production Examples and Working Examples below, the “NH silica gel

cartridge” and “silica gel cartridge” used for purification by column chromatography were Biotage SNAP Cartridge KP-NH and Biotage SNAP Cartridge KP-Sil or HP-Sil, respectively.

[0063] In the Production Examples and Working Examples below, the “NH silica gel” and “silica gel” used for purification by preparative thin-layer chromatography (PTLC) were NH₂ Silica Gel 60F254 Plate Wako (20 cm × 20 cm) from Wako Pure Chemical Industries, Ltd. and Silica Gel 60F254 (20 cm × 20 cm) from Merck, respectively.

[0064] In the Production Examples and Working Examples below, the purification by preparative high-performance liquid chromatography (HPLC) was performed under the conditions shown below. It should be noted that when trifluoroacetic acid was used in the main procedure for producing compounds having a basic functional group, neutralization operation or the like was conducted as appropriate for obtaining the compounds in free form.

Apparatus: Preparative HPLC System from Gilson, Inc.

Column: Capcelpak C18 MGII 5 μm 20 × 150 mm from Shiseido, Co., Ltd., or Waters SunFire Prep C18 OBD 5 μm 30 × 50 mm

Solvent: A-liquid (0.1 % trifluoroacetic acid-containing water), B-liquid (0.1% trifluoroacetic acid-containing acetonitrile)

Gradient Condition 1: 0 min (A-liquid/B-liquid = 90/10), 22 min (A-liquid/B-liquid = 20/80), 25 min (A-liquid/B-liquid = 10/90); flow rate, 20 mL/min

Gradient Condition 2: 0 min (A-liquid/B-liquid = 80/20), 20 min (A-liquid/B-liquid = 5/95), 25 min (A-liquid/B-liquid = 1/99); flow rate, 20 mL/min

Gradient Condition 3: 0 min (A-liquid/B-liquid = 90/10), 11 min (A-liquid/B-liquid = 20/80), 12 min (A-liquid/B-liquid = 5/95); flow rate, 40 mL/min

Gradient Condition 4: 0 min (A-liquid/B-liquid = 80/20), 10 min (A-liquid/B-liquid = 5/95), 11 min (A-liquid/B-liquid = 1/99); flow rate, 40 mL/min

Detection method: UV 254 nm

In the Production Examples and Working Examples below, mass spectra (MS) were measured under the following conditions:

MS spectra: Shimadzu LCMS-2010EV, Micromass Platform LC, or Shimadzu LCMS-IT-TOF

In the Production Examples and Working Examples below, nuclear magnetic resonance spectra (NMR) were used for structure confirmation. The nuclear magnetic resonance spectra (NMR) were measured under the following conditions:

NMR spectra: [¹H-NMR] 600 MHz, JNM-ECA600 (JEOL Ltd.); 500 MHz, JNM-ECA500 (JEOL Ltd.); 300 MHz, UNITYNOVA300 (Varian Inc.); 200 MHz, GEMINI2000/200 (Varian Inc.)

The RT (retention time) shown in Tables 1-9 to 1-26 are values measured with a high-performance liquid chromatography mass spectrometer (LCMS) under any one of the conditions shown below.

Condition A:

Instrument: Agilent 1290 Infinity and Agilent 6150

Column: Waters Acquity CSH C18, 1.7 μm, φ2.1 × 50 mm

Solvent: A-liquid (0.1 % formic acid-containing water), B-liquid (0.1 % formic acid-containing acetonitrile)

Gradient: 0 min (A-liquid/B-liquid = 80/20), 1.2-1.4 min (A-liquid/B-liquid = 1/99)

Flow rate: 0.8 mL/min, Detection method: 254 nm

Condition B:

Instrument: Shimadzu LCMS-2010EV

Column: Shimpack XR-ODS, 2.2 μm, φ2.0 × 30 mm

Solvent: A-liquid (0.1 % formic acid-containing water), B-liquid (0.1 % formic acid-containing acetonitrile)

Gradient: 0 min (A-liquid/B-liquid = 90/10), 3 min (A-liquid/B-liquid = 0/100)

Flow rate: 0.6 mL/min, Detection method: 254 nm

Condition C:

Instrument: Agilent 1100 and Micromass Platform LC

Column: Waters SunFire C18, 2.5 μm, φ4.6 × 50 mm

Solvent: A-liquid (0.1 % trifluoroacetic acid-containing water), B-liquid (0.1 % trifluoroacetic acid-containing acetonitrile)

Gradient: 0 min (A-liquid/B-liquid = 90/10), 0.5 min (A-liquid/B-liquid = 90/10), 5.5 min (A-liquid/B-liquid = 20/80), 6.0 min (A-liquid/B-liquid = 1/99), 6.3 min (A-liquid/B-liquid = 1/99)


Flow rate: 1 mL/min, Detection method: 254 nm

In the Production Examples and Working Examples below, compounds were named in accordance with ACD/Name (ACD/Labs 12.01, Advanced Chemistry Development Inc.)

Production Example 1

4-(2-Methylpropyl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

[0065] [Formula 10]



[0066] (1) Di-tert-butyl dicarbonate (4.1 g) was added to a solution of 2-amino-4-methylpentan-1-ol (2.0 g) in THF (40 mL), and the mixture was stirred at room temperature overnight. The reaction mixture was concentrated under reduced pressure, and the residue was washed with a mixed solvent of hexane/ethyl acetate (= 5:1) to afford tert-butyl (1-hydroxy-4-methylpentan-2-yl)carbamate (3.7 g).

(ESI pos.) m/z : 240([M+Na]⁺)

(2) To a solution of tert-butyl (1-hydroxy-4-methylpentan-2-yl)carbamate (2.7 g) in DMSO (60 mL) was added 2-iodoxybenzoic acid (3.5 g), and the mixture was stirred at room temperature for 3 hr. Water was added thereto, the resulting insoluble matter was filtered off, and the filtrate was extracted with ethyl acetate. The organic layer was washed with

water and brine, and dried over anhydrous magnesium sulfate. After the dessicant was filtered off, the filtrate was concentrated under reduced pressure to afford a crude product of tert-butyl (4-methyl-1-oxopentan-2-yl)carbamate.

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.96 - 1.00 (m, 6 H), 1.61 - 1.84 (m, 3 H), 3.42 - 4.98 (m, 2 H), 9.60 (s, 1 H)

(3) The crude product of tert-butyl(4-methyl-1-oxopentan-2-yl)carbamate was dissolved in chloroform, 2-trifluoromethyl-5-aminopyridine (1.35 g) and sodium triacetoxyborohydride (3.52 g) were added thereto, and the mixture was stirred at room temperature overnight. Saturated aqueous sodium hydrogen carbonate solution was added to the mixture, followed by extraction with chloroform and drying over anhydrous magnesium sulfate. The dissicant was filtered off, and then the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate) to afford tert-butyl (4-methyl-1-{{[6-(trifluoromethyl)pyridin-3-yl]amino}pentan-2-yl)carbamate (2.58 g).

(ESI pos.) m/z : 362([M+H]⁺)

(4) A solution of 4M hydrochloric acid in 1,4-dioxane was added to a solution of tert-butyl 4-methyl-1-{{[6-(trifluoromethyl)pyridin-3-yl]amino}pentan-2-yl)carbamate (2.3 g) in ethanol (30 mL), and the mixture was stirred at room temperature overnight. After the reaction mixture was concentrated under reduced pressure, saturated aqueous sodium hydrogen carbonate solution was added thereto, followed by extraction with chloroform. After drying over anhydrous magnesium sulfate, the desiccant was filtered off and the filtrate was concentrated under reduced pressure. The residue was dissolved in THF (20 mL) and cooled in ice, triethylamine (2.0 mL) and triphosgene (0.41 g) were then added thereto, and the resulting mixture was stirred for 30 min. Saturated aqueous sodium hydrogen carbonate solution was added thereto, followed by extraction with ethyl acetate and drying over anhydrous magnesium sulfate. After the desiccant was filtered off, the filtrate was concentrated under reduced pressure, and the residue was washed with hexane/ethyl acetate (= 2:1) to affore the title compound (0.58 g).

(ESI pos.) m/z : 288([M+H]⁺)

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.97 - 1.03 (m, 6 H), 1.48 - 1.55 (m, 1 H), 1.61 - 1.66 (m, 1 H), 1.69 - 1.79 (m, 1 H), 3.56 (dd, $J=8.7, 6.4$ Hz, 1 H), 3.94 - 4.01 (m, 1 H), 4.06 - 4.11 (m, 1 H), 5.28 (br. s., 1 H), 7.64 (d, $J=8.7$ Hz, 1 H), 8.37 (dd, $J=8.7, 2.3$ Hz, 1 H), 8.67 (d, $J=2.8$ Hz, 1 H)

The following compounds were synthesized according to the similar procedure.

4-(Propan-2-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 274([M+H]⁺)


1-(6-Methoxypyridin-3-yl)-4-propylimidazolidin-2-one

(ESI pos.) m/z : 236([M+H]⁺)

Production Example 2

(4S)-4-(Propan-2-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

[0067] [Formula 11]



[0068] (1) Pyridine (2.9 mL) was added to a solution of 2-trifluoromethyl-5-aminopyridine (1.95 g) in chloroform (15 mL), the mixture was cooled in ice, and phenyl chloroformate (1.8 mL) was added thereto. The resulting mixture was stirred at room temperature overnight, and the reaction mixture was concentrated under reduced pressure. The residue was washed with isopropyl ether to afford phenyl [6-(trifluoromethyl)pyridin-3-yl]carbamate (2.16 g).

(ESI pos.) m/z : 283([M+H]⁺)

(2) Triethylamine (0.54 mL) and phenyl [6-(trifluoromethyl)pyridin-3-yl]carbamate (600 mg)

were added to a solution of L-valinol (200 mg) in chloroform, and the mixture was stirred at 80°C for 1 hr. The reaction mixture was concentrated under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate) to afford 1-[(2S)-1-hydroxy-3-methylbutan-2-yl]-3-[6-(trifluoromethyl)pyridin-3-yl]urea (640 mg).

(ESI pos.) m/z : 292([M+H]⁺)

(3) Triphenylphosphine (770 mg) and diethyl azodicarboxylate (2.2 M solution in toluene, 1.3 mL) were added to a solution of 1-[(2S)-1-hydroxy-3-methylbutan-2-yl]-3-[6-(trifluoromethyl)pyridin-3-yl]urea (640 mg) in THF (10 mL), and the mixture was stirred at room temperature for 1 hr. The reaction mixture was concentrated under reduced pressure, the residue was purified by column chromatography (silica gel cartridge and NH silica gel cartridge, hexane/ethyl acetate), and the resulting solid was washed with isopropyl ether to afford the title compound (500 mg). This product contained by-products originating in reaction reagents.

(ESI pos.) m/z : 274([M+H]⁺)

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.96 - 1.05 (m, 6 H), 1.76 - 1.86 (m, 1 H), 3.59 - 3.66 (m, 2 H), 3.99 - 4.08 (m, 1 H), 7.64 (d, J=8.7 Hz, 1 H), 8.41 (dd, J=8.7, 2.5 Hz, 1 H), 8.67 (d, J=2.5 Hz, 1 H)

The following compounds were synthesized according to the similar procedure.

(4R)-4-(Propan-2-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(4S)-4-(2-Methylpropyl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 288([M+H]⁺)

(4S)-4-(Cyclohexylmethyl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 328([M+H]⁺)

(4S)-4-(Butan-2-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 288([M+H]⁺)

(4S)-4-Phenyl-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 308([M+H]⁺)

(4S)-4-tert-Butyl-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 288([M+H]⁺)

(4S)-4-Ethyl-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 260([M+H]⁺)

(4S)-1-(5-Fluoropyridin-2-yl)-4-(propan-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 224([M+H]⁺)

(4S)-1-(6-Bromopyridin-3-yl)-4-(propan-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 284([M+H]⁺)

(4S)-4-Benzyl-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 322([M+H]⁺)

(4S)-4-Propyl-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 274([M+H]⁺)

(4S)-1-(6-Bromopyridin-3-yl)-4-tert-butylimidazolidin-2-one

(ESI pos.) m/z : 298([M+H]⁺)

(4S)-4-(Cyclopropylmethyl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 286([M+H]⁺)

(4S)-4-Propyl-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

¹H NMR (200 MHz, DMSO-d₆) δ ppm 0.82 - 0.98 (m, 3 H), 1.23 - 1.64 (m, 4 H), 3.48 - 3.62 (m, 1 H), 3.66 - 3.84 (m, 1 H), 3.98 - 4.14 (m, 1 H), 7.68 (s, 1 H), 7.81 (d, J=8.79 Hz, 1 H), 8.17 - 8.28 (m, 1 H), 8.86 - 8.93 (m, 1 H)

(4R)-4-(2-Fluoropropan-2-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 292([M+H]⁺)

(4R)-4-(2-Methoxypropan-2-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 304([M+H]⁺)

(4R)-4-[(1S)-1-Fluoropropyl]-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.06 - 1.16 (m, 3 H), 1.64 - 1.79 (m, 2 H), 3.91 (dd, J=9.29, 5.16 Hz, 1 H), 3.96 - 4.04 (m, 1 H), 4.06 - 4.16 (m, 1 H), 4.34 - 4.50 (m, 1 H), 5.45 (br. s., 1 H), 7.66 (d, J=8.67 Hz, 1 H), 8.30 - 8.38 (m, 1 H), 8.70 - 8.76 (m, 1 H)

(4S)-4-Cyclopropyl-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

(ESI pos.) m/z : 272([M+H]⁺)

(4S)-4-(2-Fluoro-2-methylpropyl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one


(ESI pos.) m/z : 306([M+H]⁺)

(4S)-4-[2-(Benzyloxy)ethyl]-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

Production Example 3

{(5S)-2-Oxo-5-(propan-2-yl)-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl} acetic acid

[0069] [Formula 12]



[0070] (1) Sodium hydride (0.53 g) was added to a solution of (4S)-4-(propan-2-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one (5.6 g) in DMF (20 mL), and the mixture was stirred at room temperature for 10 min. Thereafter, tert-butyl bromoacetate (2.0 mL) was added portionwise thereto, and the mixture was stirred at room temperature for 2 hr.

Water and saturated aqueous sodium hydrogen carbonate solution were added thereto, and the mixture was extracted with chloroform. The organic layer was separated out by phase-separation cartridge (Biotage, Isolute Phase Separator) and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate) to afford tert-butyl {(5S)-2-oxo-5-(propan-2-yl)-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl} acetate (802 mg).

(ESI pos.) m/z : 388([M+H]⁺)

(2) Trifluoroacetic acid (30 mL) was added to a solution of tert-butyl {(5S)-2-oxo-5-(propan-2-yl)-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl} acetate (4.44 g) in chloroform (20 mL), and the mixture was stirred at room temperature for 64 hr. Additional trifluoroacetic acid (9 mL) was added thereto, the resulting mixture was further stirred for

1 hr, and the reaction mixture was then concentrated under reduced pressure. A portion (6.5 g) of the residue was dissolved in diethyl ether, followed by extraction with 6M aqueous sodium hydroxide solution and water. The water layer was washed with diethyl ether and then made acidic using 1M hydrochloric acid. After extraction with chloroform, the organic layer was separated out by phase-separation cartridge, and the solvent was distilled off under reduced pressure to afford the title compound (2.3 g).

(ESI pos.) m/z : 332([M+H]⁺)

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=6.6 Hz, 3 H), 1.01 (d, J=7.0 Hz, 3 H), 2.06 - 2.16 (m, 1 H), 3.59 (dd, J=8.9, 6.4 Hz, 1 H), 3.78 (d, J=18.2 Hz, 1 H), 3.86 - 3.92 (m, 1 H), 3.92 - 3.99 (m, 1 H), 4.48 (d, J=18.2 Hz, 1 H), 7.64 (d, J=9.1 Hz, 1 H), 8.30 - 8.37 (m, 1 H), 8.75 (d, J=2.5 Hz, 1 H)

The following compounds were synthesized according to the similar procedure.

[3-(6-Methoxypyridin-3-yl)-2-oxo-5-propylimidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 294([M+H]⁺)

{(5S)-2-Oxo-5-(propan-2-yl)-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl} acetic acid

(ESI neg.) m/z : 331([M-H]⁻)

[(5S)-3-(5-Chloropyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 299([M+H]⁺)

2-{(5S)-2-Oxo-5-(propan-2-yl)-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}propanoic acid (mixture of diastereomers)

(ESI pos.) m/z : 346([M+H]⁺)

[(5S)-3-(5-Fluoropyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 283([M+H]⁺)

[(5S)-3-(6-Bromopyridin-3-yl)-5-tert-butyl-2-oxoimidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 356([M+H]⁺)

{(5S)-5-(Cyclopropylmethyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 344([M+H]⁺)

{(5S)-2-Oxo-5-propyl-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 332([M+H]⁺)

[(5S)-3-[3-Fluoro-5-(trifluoromethyl)pyridin-2-yl]-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 350([M+H]⁺)

{(5S)-2-Oxo-5-(propan-2-yl)-3-[2-(trifluoromethyl)pyrimidin-5-yl]imidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 333([M+H]⁺)

[(5S)-3-(5-Methoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 295([M+H]⁺)

[(5S)-3-(5-Ethylpyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 293([M+H]⁺)

{(5S)-5-(2-Methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 346([M+H]⁺)

{(5S)-5-tert-Butyl-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetic acid

[(5S)-3-[3-Fluoro-5-(trifluoromethyl)pyridin-2-yl]-5-(2-methylpropyl)-2-oxoimidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 364([M+H]⁺)

{(5S)-5-Cyclopropyl-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 330([M+H]⁺)

{(5S)-3-[3-Fluoro-5-(trifluoromethyl)pyridin-2-yl]-2-oxo-5-propylimidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 372([M+Na]⁺)

{(5R)-5-[(1S)-1-Fluoropropyl]-3-[3-fluoro-5-(trifluoromethyl)pyridin-2-yl]-2-oxoimidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 390([M+Na]⁺)

[(5S)-3-(5-Chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 321([M+Na]⁺)

{(5S)-5-[(2S)-Butan-2-yl]-3-[3-fluoro-5-(trifluoromethyl)pyridin-2-yl]-2-oxoimidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 364([M+H]⁺)

[(5S)-5-[(2S)-Butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 313([M+H]⁺)

[(5S)-5-[(2S)-Butan-2-yl]-3-(5-fluoropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 297([M+H]⁺)

{(5S)-5-[(2S)-Butan-2-yl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 247([M+H]⁺)

[(5S)-5-[(2S)-Butan-2-yl]-3-(5-chloro-3-fluoropyridin-2-yl)-2-oxoimidazolidin-1-yl]acetic acid

(ESI pos.) m/z : 330([M+H]⁺)


{(5S)-5-(2-Fluoro-2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetic acid

(ESI pos.) m/z : 364([M+H]⁺)

Production Example 4

(4S)-4-(Propan-2-yl)-1-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-2-one

[0071] [Formula 13]



[0072] (1) To a suspension of L-valinamide hydrochloride (1.0 g) and sodium hydrogen carbonate (3.03 g) in acetonitrile (100 mL) was added p-nitrophenyl chloroformate (1.32 g), and the mixture was stirred at room temperature for 3 hr. Water was added thereto, the resulting mixture was stirred at room temperature overnight, and then acetonitrile was distilled off under reduced pressure. After extraction with ethyl acetate, the organic layer was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel cartridge, chloroform/methanol) to afford (5S)-5-(propan-2-yl)imidazolidine-2,4-dione (540 mg).

(ESI pos.) m/z : 143([M+H]⁺)

(2) To a suspension of lithium aluminum hydride (290 mg) in diethyl ether (20 mL) was added (5S)-5-(propan-2-yl)imidazolidine-2,4-dione (540 mg) while cooling in ice, and the mixture was stirred at room temperature overnight. After cooling in ice, water (1 mL), 4M aqueous sodium hydroxide solution (1 mL), THF (15 mL) and ethanol (2 mL) were added thereto, and the resulting mixture was stirred at room temperature for 10 min. After filtration through Celite (registered trademark), the filtrate was concentrated under reduced pressure to afford (4S)-4-(propan-2-yl)imidazolidin-2-one (160 mg).

(ESI pos.) m/z : 129([M+H]⁺)

(3) A solution of (4S)-4-(propan-2-yl)imidazolidin-2-one (100 mg), 2-chloro-5-trifluoromethylpyrimidine (142 mg), Pd₂(dba)₃ (40 mg), Xantphos (45 mg) and sodium tert-butoxide (70 mg) in toluene (2 mL) was stirred at 100°C for 1 hr. The reaction mixture was purified by column chromatography (NH silica gel cartridge and silica gel cartridge, hexane/ethyl acetate) to afford the title compound (47 mg).

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.95 - 1.02 (m, 6 H), 1.75 - 1.85 (m, 1 H), 3.50 - 3.61 (m, 1 H), 3.87 (dd, J=11.1, 6.6 Hz, 1 H), 4.18 - 4.26 (m, 1 H), 5.17 (br. s., 1 H), 8.84 (s, 2 H)

The following compounds were synthesized according to the similar procedure.

(4S)-1-(5-Chloropyrimidin-2-yl)-4-(propan-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 241([M+H]⁺)

6-[(4S)-2-Oxo-4-(propan-2-yl)imidazolidin-1-yl]pyridine-3-carbonitrile

(ESI pos.) m/z : 231([M+H]⁺)

(4S)-4-(Propan-2-yl)-1-[5-(trifluoromethyl)pyrazin-2-yl]imidazolidin-2-one

(ESI pos.) m/z : 275([M+H]⁺)

(4S)-1-(5-Fluoropyrimidin-2-yl)-4-(propan-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 225([M+H]⁺)

(4S)-1-[3-Fluoro-5-(trifluoromethyl)pyridin-2-yl]-4-(propan-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 292([M+H]⁺)

(4S)-4-(Propan-2-yl)-1-[2-(trifluoromethyl)pyrimidin-5-yl]imidazolidin-2-one

(ESI pos.) m/z : 275([M+H]⁺)

(4S)-1-(5-Chloropyridin-2-yl)-4-(propan-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 240([M+H]⁺)

(4S)-1-(5-Methoxypyrimidin-2-yl)-4-(propan-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 237([M+H]⁺)

(4S)-1-(5-Ethylpyrimidin-2-yl)-4-(propan-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 235([M+H]⁺)

(4S)-1-[3-Fluoro-5-(trifluoromethyl)pyridin-2-yl]-4-(2-methylpropyl)imidazolidin-2-one

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.95 - 1.01 (m, 6 H), 1.47 - 1.56 (m, 1 H), 1.61 - 1.77 (m, 2 H), 3.76 - 3.85 (m, 1 H), 3.93 - 4.03 (m, 1 H), 4.07 - 4.19 (m, 1 H), 5.12 (br. s., 1 H), 7.63 - 7.71 (m, 1 H), 8.45 (s, 1 H)

(4S)-1-[3-Fluoro-5-(trifluoromethyl)pyridin-2-yl]-4-propylimidazolidin-2-one

(ESI pos.) m/z : 314([M+Na]⁺)

(4S)-1-(5-Chloro-3-fluoropyridin-2-yl)-4-propylimidazolidin-2-one

(ESI pos.) m/z : 258([M+H]⁺)


(4S)-4-(Cyclopropylmethyl)-1-[3-fluoro-5-(trifluoromethyl)pyridin-2-yl]imidazolidin-2-one

(ESI pos.) m/z : 304([M+H]⁺)

Production Example 5

2-[(5S)-2-Oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide

[0073] [Formula 14]



[0074] (1) A solution of (2S)-2-aminopentan-1-ol (500 mg) in THF (6 mL) was cooled in a dry ice-acetone bath under nitrogen atmosphere, and a solution of 4-methoxyphenyl isocyanate (657 mg) in THF (4 mL) was added dropwise thereto. After the mixture was stirred for 17 hr while gradually warming to room temperature, methanol was added thereto, and the solvent was distilled off under reduced pressure to afford 1-[(2S)-1-hydroxypentan-2-yl]-3-(4-methoxyphenyl)urea (1.17 g).

(ESI pos.) m/z : 253([M+H]⁺)

(2) A solution of 1-[(2S)-1-hydroxypentan-2-yl]-3-(4-methoxyphenyl)urea (550 mg) and potassium tert-butoxide (593 mg) in THF (25 mL) was cooled in ice under nitrogen atmosphere, and a solution of p-toluenesulfonylchloride (672 mg) in THF (10 mL) was added dropwise thereto. After the mixture was stirred for 1 hr while cooling in ice, water was added thereto, and the resulting mixture was extracted with chloroform. The organic layer was separated out by phase-separation cartridge, and the solvent was distilled off under reduced pressure. The residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 88:12 - 0:100), and further recrystallized from chloroform/hexane. The solid was collected by filtration to afford (4S)-1-(4-methoxyphenyl)-4-propylimidazolidin-2-one (34 mg). In addition, the filtrate was concentrated under reduced pressure to afford (4S)-1-(4-methoxyphenyl)-4-propylimidazolidin-2-one (254 mg).

(ESI pos.) m/z : 235([M+H]⁺)

(3) Sodium hydride (60%, 27 mg) was added to a solution of (4S)-1-(4-methoxyphenyl)-4-propylimidazolidin-2-one (32 mg) in DMF (1.5 mL), and the mixture was stirred at room temperature for 15 min. To the mixture was added 2-chloro-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide (39 mg), and the resulting mixture was stirred for 15 hr. Thereafter, the reaction mixture was subjected to filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by preparative HPLC, and further the resulting solid was washed with isopropyl ether to afford 2-[(5S)-3-(4-methoxyphenyl)-2-oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide (19 mg).

(ESI pos.) m/z : 437([M+H]⁺)

(4) A suspension of 2-[(5S)-3-(4-methoxyphenyl)-2-oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide (137 mg) in acetonitrile (3 mL) was cooled in ice, and a solution of ammonium cerium nitrate (340 mg) in water (3 mL) was added dropwise thereto. After stirring the mixture at room temperature for 2 hr, an additional solution of ammonium cerium nitrate (100 mg) in water (0.5 mL) was added thereto, and the mixture was stirred at room temperature for 10 min. Saturated aqueous sodium hydrogen carbonate solution was added thereto, the resulting mixture was extracted with ethyl acetate, and then the organic layer was washed with brine, followed by drying over anhydrous magnesium sulfate. After the dissicant was filtered off, the solvent was distilled off under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, chloroform/methanol = 98:2 - 80:20) to afford the title compound (76 mg).

(ESI pos.) m/z : 331([M+H]⁺)

The following compounds were obtained according to the similar procedure.

N-(4-Chloropyridin-2-yl)-2-[(5R)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl]acetamide

(ESI pos.) m/z : 315([M+H]⁺)

N-(4-Cyclopropylpyridin-2-yl)-2-[(5R)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl]acetamide

(ESI pos.) m/z : 321([M+H]⁺)

2-{(5R)-5-[(1S)-1-Fluoropropyl]-2-oxoimidazolidin-1-yl}-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide

(ESI pos.) m/z : 371([M+Na]⁺)

N-(5-Chloropyridin-2-yl)-2-[(5S)-2-oxo-5-propylimidazolidin-1-yl]acetamide

(ESI pos.) m/z : 297([M+H]⁺)

N-(5-Chloro-6-methylpyridin-2-yl)-2-[(5S)-2-oxo-5-propylimidazolidin-1-yl]acetamide

(ESI pos.) m/z : 311([M+H]⁺)

2-[(5S)-2-Oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide

(ESI pos.) m/z : 331([M+H]⁺)

N-(4-Cyclopropylpyridin-2-yl)-2-[(5S)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide

(ESI pos.) m/z : 303([M+H]⁺)

N-(4-Chloropyridin-2-yl)-2-[(5S)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide

(ESI pos.) m/z : 297([M+H]⁺)

2-{(5S)-5-[(2S)-Butan-2-yl]-2-oxoimidazolidin-1-yl}-N-(4-chloropyridin-2-yl)acetamide

(ESI pos.) m/z : 333([M+Na]⁺)

2-{(5S)-5-[(2S)-Butan-2-yl]-2-oxoimidazolidin-1-yl}-N-(4-cyclopropylpyridin-2-yl)acetamide

(ESI pos.) m/z : 317([M+H]⁺)

2-{(5S)-5-[(2S)-Butan-2-yl]-2-oxoimidazolidin-1-yl}-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide


(ESI pos.) m/z : 345([M+H]⁺)

Production Example 6

2-[(5S)-2-Oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[6-(trifluoromethyl)pyridin-3-yl]acetamide

[0075]

[Formula 15]



[0076] (1) To a solution of L-valine methyl ester hydrochloride (2.6 g) and triethylamine (1.6 g) in chloroform (30 mL) was added 2,4-dimethoxybenzyl isocyanate (3.0 g), and the mixture was stirred at room temperature overnight. After the solvent was distilled off under reduced pressure, the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate) to afford methyl N-[(2,4-dimethoxybenzyl)carbamoyl]-L-valinate (4.2 g).

(ESI pos.) m/z : 325([M+H]⁺)

(2) A solution of methyl N-[(2,4-dimethoxybenzyl)carbamoyl]-L-valinate (4.2 g) and triethylamine (1.3 g) in methanol was stirred while heating to reflux for 5 hr. After the solvent was distilled off under reduced pressure, the residue was purified by column chromatography (NH silica gel cartridge, hexane/ethyl acetate) to afford (5S)-3-(2,4-dimethoxybenzyl)-5-(propan-2-yl)imidazolidine-2,4-dione (4.3 g).

(ESI pos.) m/z : 293([M+H]⁺)

(3) Bis(2-methoxyethoxy)aluminum sodium hydride (65%, solution in toluene) was added to a solution of (5S)-3-(2,4-dimethoxybenzyl)-5-(propan-2-yl)imidazolidine-2,4-dione (4.3 g) in THF (300 mL), and the mixture was stirred while heating to reflux for 3 hr. After cooling in ice, sodium sulfate decahydrate was added thereto, followed by stirring. After filtration, the solvent was distilled off under reduced pressure, and the residue was purified by column

chromatography (silica gel cartridge, hexane/ethyl acetate) to afford (4S)-1-(2,4-dimethoxybenzyl)-4-(propan-2-yl)imidazolidin-2-one (2.4 g).

(4) Sodium hydride (60%, 380 mg) was added to a solution of (4S)-1-(2,4-dimethoxybenzyl)-4-(propan-2-yl)imidazolidin-2-one (2.4 g) in DMF (35 mL), and the mixture was stirred for 20 min. Ethyl bromoacetate (1.14 mL) was added thereto, and the mixture was stirred at 90°C for 1 hr. Additional ethyl bromoacetate (1.14 g) and sodium hydride (60%, 380 mg) were added thereto, and the resulting mixture was further stirred at 90°C for 1 hr. After the solvent was distilled off under reduced pressure, the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate) to afford ethyl [(5S)-3-(2,4-dimethoxybenzyl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetate (1.5 g).

(ESI pos.) m/z : 365([M+H]⁺)

(5) An aqueous solution of 2M sodium hydroxide (2.6 mL) was added to a solution of ethyl [(5S)-3-(2,4-dimethoxybenzyl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetate (850 mg) in methanol (10 mL), and the mixture was stirred at room temperature overnight. To the reaction mixture was added 2M hydrochloric acid (2.6 mL), and the reaction mixture was concentrated under reduced pressure. DMF (10 mL), 5-amino-2-trifluoromethylpyridine (380 mg), HATU (890 mg), and diisopropylethylamine (300 mg) were added thereto, and the resulting mixture was stirred at 80°C for 3 hr. The solvent was distilled off under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate) to afford 2-[(5S)-3-(2,4-dimethoxybenzyl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[6-(trifluoromethyl)pyridin-3-yl]acetamide (900 mg).

(ESI pos.) m/z : 481([M+H]⁺)

(6) Trifluoroacetic acid (20 mL) was added to 2-[(5S)-3-(2,4-dimethoxybenzyl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[6-(trifluoromethyl)pyridin-3-yl]acetamide (900 mg), and the mixture was stirred at room temperature for 3 hr. The solvent was distilled off under reduced pressure, and saturated sodium hydrogen carbonate was added thereto, followed by extraction with chloroform. The solvent was distilled off under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate) to


afford the title compound (310 mg).

(ESI pos.) m/z : 331([M+H]⁺)

Production Example 7

(2R,3S)-2-Amino-3-fluoropentan-1-ol hydrochloride

[0077] [Formula 16]



[0078] (1) A solution of tert-butyl (4R)-4-formyl-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (14.0 g) in THF (250 mL) was cooled in a dry ice/acetone bath under nitrogen atmosphere, and ethyl magnesium bromide (24 mL, 3M solution in diethyl ether) was added dropwise thereto. The resulting mixture was stirred for 3 hr while elevating the temperature gradually to 0°C, and saturated aqueous ammonium chloride solution was added thereto. The mixture was separated into phases, the aqueous layer was then extracted with ethyl acetate, and the organic layer was washed with water and brine. After drying over anhydrous magnesium sulfate, the desiccant was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 92:8 - 37:63) to afford tert-butyl (4R)-4-(1-hydroxypropyl)-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (11.14 g).

(ESI pos.) m/z : 282([M+Na]⁺)

(2) A solution of tert-butyl (4R)-4-(1-hydroxypropyl)-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (10.0 g) in chloroform (200 mL) was cooled in ice under nitrogen atmosphere, diethylaminosulfur trifluoride (6.1 mL) was added dropwise thereto, and the mixture was stirred under cooling in ice for 1 hr. Saturated aqueous sodium hydrogen carbonate solution was added thereto, and the resulting mixture was extracted with chloroform, followed by drying over anhydrous magnesium sulfate. The desiccant was filtered off, the filtrate was concentrated under reduced pressure, and then the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 95:5 - 60:40) to afford tert-butyl

((4R)-4-[(1S)-1-fluoropropyl]-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (4.74 g).

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.91 (t, J=7.22 Hz, 3 H), 1.28 - 1.62 (m, 17 H), 3.72 - 4.01 (m, 3 H), 4.30 - 4.63 (m, 1 H)


(3) A 5-10% HCl/methanol solution (70 mL) was added to tert-butyl ((4R)-4-[(1S)-1-fluoropropyl]-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (4.9 g), and the mixture was stirred at room temperature overnight. The solvent was distilled off under reduced pressure to afford the title compound (2.96 g).

(ESI pos.) m/z : 122([M+H]⁺)

Production Example 8

(4R)-4-[(1S)-1-Fluoropropyl]-1-[3-fluoro-5-(trifluoromethyl)pyridin-2-yl]imidazolidin-2-one

[0079] [Formula 17]



[0080] (1) A solution of (2R,3S)-2-amino-3-fluoropentan-1-ol hydrochloride (2.0 g) in THF (60 mL) was cooled in ice, a solution of potassium carbonate (4.4 g) in water (6 mL) and a solution of benzyl chloroformate (2.4 g) in THF (6 mL) were added thereto, and the mixture was stirred at room temperature overnight. Saturated aqueous sodium hydrogen carbonate solution was added thereto, followed by extraction with ethyl acetate and drying over anhydrous magnesium sulfate. After the desiccant was filtered off, the solvent was distilled off under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 88:12 - 0:100) to afford benzyl [(2R,3S)-3-fluoro-1-hydroxypentan-2-yl]carbamate (2.5 g).

¹H NMR (600 MHz, DMSO-d₆) δ ppm 0.88 (t, J=7.43 Hz, 3 H), 1.45 - 1.65 (m, 2 H), 3.36 - 3.50 (m, 2 H), 3.59 - 3.70 (m, 1 H), 4.30 - 4.44 (m, 1 H), 4.65 - 4.71 (m, 1 H), 4.95 - 5.04 (m,

2 H), 7.12 - 7.36 (m, 5 H)

(2) A solution of benzyl [(2R,3S)-3-fluoro-1-hydroxypentan-2-yl]carbamate (2.4 g) in chloroform (40 mL) was cooled in ice, triethylamine (2 mL) and methanesulfonyl chloride (0.8 mL) were added thereto, and the mixture was stirred for 30 min. Saturated sodium hydrogen carbonate was added thereto, followed by extraction with chloroform and drying over anhydrous magnesium sulfate. After the desiccant was filtered off, the solvent was distilled off under reduced pressure, and the residue was dissolved in DMF (40 mL). Sodium azide (3.06 g) was added thereto, and the mixture was stirred at 60°C for 2 hr. Water was added thereto, the resulting mixture was extracted with ethyl acetate, and the organic layer was washed with water and brine. After drying over anhydrous magnesium sulfate, the desiccant was filtered off, and the solvent was distilled off under reduced pressure. The residue was dissolved in THF (50 mL), triphenylphosphine (3.59 g) and water (10 mL) were added thereto, and the resulting mixture was stirred at room temperature overnight. The solvent was distilled off under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, chloroform/methanol = 98:2 - 80:20) to afford benzyl [(2R,3S)-1-amino-3-fluoropentan-2-yl]carbamate (2.34 g).

(ESI pos.) m/z : 255([M+H]⁺)

(3) Potassium carbonate was added to a solution of benzyl [(2R,3S)-1-amino-3-fluoropentan-2-yl]carbamate (2.2 g) and 2,3-difluoro-5-trifluoromethylpyridine (1.74 g) in acetonitrile (45 mL), and the mixture was stirred at 80°C for 5 hr. The reaction mixture was diluted with ethyl acetate and washed with water and brine, and then the solvent was distilled off under reduced pressure. The residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 95:5 - 20:80) to afford benzyl [(2R,3S)-3-fluoro-1-{[3-fluoro-5-(trifluoromethyl)pyridin-2-yl]amino}pentan-2-yl]carbamate (2.95 g).

(ESI pos.) m/z : 418([M+H]⁺)

(4) Sodium hydride (about 60%, 546 mg) was added to a solution of benzyl [(2R,3S)-3-fluoro-1-{[3-fluoro-5-(trifluoromethyl)pyridin-2-yl]amino}pentan-2-yl]carbamate (2.85 g) in THF (35 mL), and the mixture was stirred at room temperature for 6 hr. Water was added

to the reaction mixture, the resulting mixture was extracted with ethyl acetate, and the organic layer was washed with brine. The solvent was distilled off under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 80:20 - 40:60) to afford the title compound (2.07 g).

(ESI pos.) m/z : 310([M+H]⁺)

The following compounds were obtained according to the similar procedure.

(4S)-4-[(2S)-Butan-2-yl]-1-[3-fluoro-5-(trifluoromethyl)pyridin-2-yl]imidazolidin-2-one

(ESI pos.) m/z : 306([M+H]⁺)

(4S)-4-[(2S)-Butan-2-yl]-1-(5-chloropyrimidin-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 255([M+H]⁺)

(4S)-4-[(2S)-Butan-2-yl]-1-(5-fluoropyrimidin-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 239([M+H]⁺)

(4S)-4-[(2S)-Butan-2-yl]-1-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-2-one

(ESI pos.) m/z : 289([M+H]⁺)


(4S)-4-[(2S)-Butan-2-yl]-1-(5-chloro-3-fluoropyridin-2-yl)imidazolidin-2-one

(ESI pos.) m/z : 277([M+H]⁺)

Production Example 9

(4S)-4-(2-Fluoroethyl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one

[0081] [Formula 18]



[0082] (1) Palladium hydroxide (120 mg) was added to a solution of (4S)-4-[2-(benzyloxy)ethyl]-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one (1.12 g) in methanol (25 mL), the system was purged with hydrogen gas, and the mixture was stirred at room temperature for 3 hr. After filtration of the reaction mixture, the filtrate was concentrated under reduced pressure, and the resulting solid was washed with ethyl acetate to afford (4S)-4-(2-hydroxyethyl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one (235 mg).


(2) A suspension of (4S)-4-(2-hydroxyethyl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one (0.10 g) in chloroform (10 mL) was cooled in ice, diethylaminosulfur trifluoride (0.144 mL) was added thereto, and the mixture was stirred at room temperature for 3 hr. Saturated aqueous sodium hydrogen carbonate solution was added thereto, and the organic layer was washed with water and brine, followed by drying over anhydrous sodium sulfate. After the desiccant was filtered off, the filtrate was concentrated under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 4:1 - 0:100) to afford the title compound (0.05 g).

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.86 - 2.01 (m, 2 H), 3.65 (dd, J=7.9, 5.2 Hz, 1 H), 3.89 - 3.941 (m, 1 H), 4.10 (d, J=7.9 Hz, 1 H), 4.60 (dd, J=39.2, 4.5 Hz, 2 H), 7.73 (s, 1 H), 7.82 (d, J=7.2 Hz, 1 H), 8.21 (d, J=7.2 Hz, 1 H), 8.90 (s, 1 H).

Production Example 10

(2S)-2-Amino-5-fluoropentan-1-ol hydrochloride

[0083] [Formula 19]



[0084] (1) A solution of tert-butyl [(2S)-1,5-dihydroxypentan-2-yl]carbamate (9.45 g), 2,2-dimethoxypropane (48 mL) and p-toluenesulfonic acid monohydrate (0.41 g) in chloroform (100 mL) was stirred at room temperature for 3 hr. Saturated aqueous sodium hydrogen carbonate solution was added thereto, and the solvent was distilled off under reduced pressure. Ethyl acetate was added thereto, and the resulting mixture was washed with water and brine, followed by drying over anhydrous sodium sulfate. After the desiccant was filtered off, the filtrate was concentrated under reduced pressure and purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 10:1 - 0:100) to afford tert-butyl (4S)-4-(3-hydroxypropyl)-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (3.05 g).

(ESI pos.) m/z : 282([M+Na]⁺)

(2) A solution of tert-butyl (4S)-4-(3-hydroxypropyl)-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (3.05 g) in chloroform (40 mL) was cooled in a dry ice-acetone bath, bis(2-


methoxyethyl)aminosulfur trifluoride (2.77 mL) was added thereto, and the mixture was stirred at room temperature for 3 hr. Saturated aqueous sodium hydrogen carbonate solution was added thereto, and the organic layer was washed with water and brine, followed by drying with anhydrous sodium sulfate. After the desiccant was filtered off, the filtrate was concentrated under reduced pressure, and the residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate = 10:1 - 2:1) to afford tert-butyl (4S)-4-(3-fluoropropyl)-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (900 mg).

(3) According to the similar procedure as in Production Example 7 (3), the title compound (580 mg) was obtained from tert-butyl (4S)-4-(3-fluoropropyl)-2,2-dimethyl-1,3-oxazolidine-3-carboxylate (900 mg) as a crude product.

Working Example 1

N-(3,5-Difluorophenyl)-2-[3-(6-methoxypyridin-3-yl)-2-oxo-5-propylimidazolidin-1-yl]acetamide

[0085] [Formula 20]




[0086] N,N-diisopropylethylamine (0.06 mL), HATU (142 mg), and 3,5-difluoroaniline (48 mg) were added to a solution of [3-(6-methoxypyridin-3-yl)-2-oxo-5-propylimidazolidin-1-yl]acetic acid (100 mg) in chloroform (2 mL), and the mixture was stirred at room temperature overnight. Saturated aqueous sodium hydrogen carbonate solution was added thereto, followed by extraction with chloroform and drying over anhydrous magnesium sulfate. After the desiccant was filtered off, the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel cartridge, hexane/ethyl acetate) to afford the title compound (114 mg).

Working Example 2

2-{2-Oxo-5-(propan-2-yl)-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}-N-[3-

(trifluoromethyl)phenyl]acetamide

[0087] [Formula 21]




[0088] Sodium hydride (60%, 22 mg) was added to a solution of 4-(propan-2-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-2-one (50 mg) in DMF (1.5 mL), and the mixture was stirred at room temperature for 15 min. After 2-chloro-N-[3-(trifluoromethyl)phenyl]acetamide (52 mg) was added thereto, the resulting mixture was stirred at room temperature for 21 hr. Water and saturated aqueous sodium hydrogen carbonate solution were added to the reaction mixture, followed by extraction with chloroform. The organic layer was separated out by phase separation cartridge and concentrated under reduced pressure. The residue was purified by column chromatography (NH cartridge, and silica gel cartridge, hexane/ethyl acetate) and preparative HPLC to afford the title compound (79 mg).

Working Example 3

2-((5S)-2-Oxo-5-(propan-2-yl)-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl)-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide

[0089] [Formula 22]




[0090] Oxalyl chloride (57 μ L) and DMF (0.05 mL) were added to a solution of ((5S)-2-oxo-5-(propan-2-yl)-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl)acetic acid (200 mg) in chloroform (2.5 mL), and the mixture was stirred at room temperature for 1 hr. The reaction mixture was concentrated under reduced pressure to afford the residue (236 mg). Half of this residue was dissolved in chloroform (0.5 mL), and triethylamine

(158 μ L) was added thereto. A solution of 2-amino-4-trifluoromethylpyridine (40 mg) in chloroform (1 mL) was added thereto while cooling in ice, and the resulting mixture was stirred at room temperature for 3 hr. After the reaction mixture was concentrated under reduced pressure, the residue was purified by preparative HPLC and PTLC (hexane/ethyl acetate = 1:1) to afford the title compound (20 mg).

Working Example 4

2-[(5S)-3-(6-Cyanopyridin-3-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[3-(trifluoromethyl)phenyl]acetamide

[0091] [Formula 23]




[0092] A solution of 2-[(5S)-3-(6-bromopyridin-3-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[3-(trifluoromethyl)phenyl]acetamide (150 mg), zinc cyanide (91 mg), $\text{Pd}_2(\text{dba})_3$ (8.5 mg), Xantphos (11 mg), and N,N,N',N' -tetramethylethylenediamine (14 μ L) in DMF (2 mL) was stirred under irradiation with microwave at 180°C for 30 min. Additional zinc cyanide (91 mg) was added thereto, and the resulting mixture was stirred under the same condition for 30 min. NH silica gel was added thereto, the mixture was stirred, followed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by preparative HPLC to afford the title compound (47 mg).

Working Example 5

2-{(5S)-2-Oxo-5-(propan-2-yl)-3-[6-(1H-pyrazol-1-yl)pyridin-3-yl]imidazolidin-1-yl}-N-[3-(trifluoromethyl)phenyl]acetamide

[0093]

[Formula 24]




[0094] A solution of 2-[(5S)-3-(6-bromopyridin-3-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[3-(trifluoromethyl)phenyl]acetamide (150 mg), pyrazole (105 mg), trans-1,2-bis(methylamino)cyclohexane (18 mg), copper(I) iodide (12 mg), and cesium carbonate (302 mg) in 1,4-dioxane (2 mL) was stirred at 120°C for 2 hr. After filtration through Celite, the filtrate was concentrated under reduced pressure. The residue was purified by preparative HPLC and PTLC (hexane/ethyl acetate = 1:1) to afford the title compound (74 mg).

Working Example 6

2-[(5S)-3-[6-(Dimethylamino)pyridin-3-yl]-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[3-(trifluoromethyl)phenyl]acetamide

[0095] [Formula 25]




[0096] A solution of 2-[(5S)-3-(6-bromopyridin-3-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[3-(trifluoromethyl)phenyl]acetamide (250 mg) and dimethyl amine (2M solution in THF, 1.3 mL), Pd₂(dba)₃ (24 mg), BINAP (32 mg), and potassium tert-butoxide (87 mg) in toluene (5 mL) was stirred at 120°C for 2 hr. Water was added thereto, the mixture was extracted with chloroform, and the organic layer was concentrated under reduced pressure. The residue was purified by preparative HPLC and PTLC (hexane/ethyl acetate = 1:1) to afford the title compound (34 mg).

Working Example 7

2-[(5S)-2-Oxo-5-(propan-2-yl)-3-(6-propylpyridin-3-yl)imidazolidin-1-yl]-N-[3-(trifluoromethyl)phenyl]acetamide

[0097] [Formula 26]




[0098] To a suspension of 2-[(5S)-3-(6-bromopyridin-3-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[3-(trifluoromethyl)phenyl]acetamide (50 mg) and tris(2,4-pentanedionato)iron(III) (55 mg) in THF (1 mL) and NMP (0.1 mL) was added n-propylmagnesium bromide (2M solution in THF, 0.31 mL), and the mixture was stirred at room temperature for 1 hr. Saturated aqueous ammonium chloride solution was added thereto, the resulting mixture was extracted with acetic acid, and the organic layer was washed with water and brine. After drying over anhydrous sodium sulfate, the desiccant was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography (hexane/ethyl acetate) to afford the title compound (24 mg).

Working Example 8

N,N-Dimethyl-5-[(4S)-2-oxo-3-(2-oxo-2-{[3-(trifluoromethyl)phenyl]amino}ethyl)-4-(propan-2-yl)imidazolidin-1-yl]pyridine-2-carboxamide

[0099]

[Formula 27]



[0100] (1) A suspension of 2-[(4S)-3-(6-bromopyridin-3-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[3-(trifluoromethyl)phenyl]acetamide (100 mg), potassium carbonate (43 mg), and Pd(PPh₃)₄ (24 mg) in DMF/ethanol (2:1, 2.1 mL) was stirred under atmosphere of carbon monoxide gas at 80°C for 1.5 hr. Saturated aqueous sodium hydrogen carbonate solution was added thereto, the mixture was washed with ethyl acetate, and the organic layer was washed with water and brine. After concentration under reduced pressure, the residue was purified by preparative HPLC to afford methyl 5-[(4S)-2-oxo-3-(2-oxo-2-{[3-(trifluoromethyl)phenyl]amino}ethyl)-4-(propan-2-yl)imidazolidin-1-yl]pyridine-2-carboxylate (50 mg).

(ESI pos.) m/z : 465([M+H]⁺)

(2) To a solution of methyl 5-[(4S)-2-oxo-3-(2-oxo-2-{[3-(trifluoromethyl)phenyl]amino}ethyl)-4-(propan-2-yl)imidazolidin-1-yl]pyridine-2-carboxylate (50 mg) in methanol/water (2:1, 1.5 mL) was added 6M aqueous sodium hydroxide solution (27 µL), and the mixture was stirred at room temperature for 1 hr, followed by at 60°C for 1 hr. After 2M hydrochloric acid was added thereto to make the mixture acidic, the resulting mixture was extracted with chloroform. The organic layer was concentrated under reduced pressure to afford 5-[(4S)-2-oxo-3-(2-oxo-2-{[3-(trifluoromethyl)phenyl]amino}ethyl)-4-(propan-2-yl)imidazolidin-1-yl]pyridine-2-

carboxylic acid (43 mg).


(ESI pos.) m/z : 451([M+H]⁺)

(3) According to the similar procedure as in Working Example 1, the title compound (33 mg) was obtained from 5-[(4S)-2-oxo-3-(2-oxo-2-{[3-(trifluoromethyl)phenyl]amino}ethyl)-4-(propan-2-yl)imidazolidin-1-yl]pyridine-2-carboxylic acid (43 mg) and dimethylamine (solution in THF).

Working Example 9

2-[(5S)-3-(5-Fluoropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide

[0101] [Formula 28]




[0102] A solution of 2-[(5S)-2-oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide (70 mg), 2-chloro-5-fluoropyrimidine (33 mg), Pd₂(dba)₃ (22 mg), Xantphos (24 mg), and sodium tert-butoxide (30 mg) in toluene was stirred at 80°C for 2 hr. Saturated aqueous sodium hydrogen carbonate solution was added thereto, followed by extraction with ethyl acetate and drying over anhydrous magnesium sulfate. The desiccant was filtered off, the solvent was distilled off under reduced pressure, and the residue was purified by preparative HPLC to afford the title compound (27 mg).

Working Example 10

N-[6-(Difluoromethyl)pyridin-3-yl]-2-[(5S)-5-(2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl]acetamide

[0103]

[Formula 29]



[0104] (1) Lithium borohydride (27 mg) was added to a solution of methyl 5-[(5S)-5-(2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl]acetamido]pyridine-2-carboxylate (160 mg) in THF (2 mL), and the mixture was stirred at room temperature for 18 hr. Saturated aqueous ammonium chloride solution was added thereto, followed by extraction with chloroform. The solvent was distilled off under reduced pressure, and the residue was purified by preparative HPLC to afford N-[6-(hydroxymethyl)pyridin-3-yl]-2-[(5S)-5-(2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl]acetamide (43 mg).


(ESI pos.) m/z : 452([M+H]⁺)

(2) To a solution of N-[6-(hydroxymethyl)pyridin-3-yl]-2-[(5S)-5-(2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl]acetamide (35 mg) in DMSO (1 mL) was added 2-iodoxybenzoic acid (26 mg), and the mixture was stirred at room temperature for 2 hr. Water and ethyl acetate were added thereto, the resulting mixture was subjected to filtration, and then the filtrate was extracted with ethyl acetate. The organic layer was washed with water and brine and then dried over anhydrous magnesium sulfate. After the desiccant was filtered off, the solvent was distilled off under reduced pressure. The residue was dissolved in chloroform (1 mL) and cooled in ice, and then diethylaminosulfur trifluoride (0.08 mL) was added thereto. After the resulting mixture was stirred at room temperature for 3 days, water was added thereto, followed by extraction with chloroform. After the solvent was distilled off under reduced pressure, the residue was purified by PTLC (silica gel, hexane/ethyl acetate = 1:1, and NH silica gel, hexane/ethyl acetate = 1:1), and the solid was washed with isopropyl ether to afford the title compound (3.8 mg).

Working Example 11

N-(6-Acetylpyridin-3-yl)-2-{{(5S)-5-(2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetamide

[0105] [Formula 30]




[0106] A suspension of N-(6-bromopyridin-3-yl)-2-{{(5S)-5-(2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetamide (78 mg), tributyl(1-ethoxyvinyl)tin (145 mg), bis(triphenylphosphine)palladium(II) dichloride (22 mg), and copper(I) iodide (6.3 mg) in acetonitrile (1.5 mL) was stirred under irradiation with microwave at 150°C for 2 hr. Saturated aqueous sodium hydrogen carbonate solution was added thereto, followed by extraction with chloroform. The organic layer was separated by phase separation cartridge, and the solvent was distilled off under reduced pressure. The residue was purified by preparative HPLC, 1M HCl (1.5 mL) was added to the resulting product, and the mixture was stirred at room temperature for 10 min. The mixture was made basic with saturated sodium hydrogen carbonate, followed by extraction with chloroform, and the solvent was distilled off under reduced pressure to afford the title compound (60 mg).

Working Example 12

N-(6-Cyclopropylpyridin-3-yl)-2-{{(5S)-5-(2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetamide

[0107] [Formula 31]












[0108] A solution of N-(6-bromopyridin-3-yl)-2-((5S)-5-(2-methylpropyl)-2-oxo-3-[6-(trifluoromethyl)pyridin-3-yl]imidazolidin-1-yl}acetamide (50 mg), cyclopropyl boronic acid (17 mg), palladium acetate (1.1 mg), triphenylphosphine (2.6 mg), and potassium carbonate (41 mg) in toluene/water (= 20:1, 1.5 mL) was stirred at 110°C for 9 hr. Additional cyclopropyl boronic acid (35 mg) was added thereto, and the mixture was further stirred at 110°C for 6 hr, followed by at 80°C for 2 hr. The solvent was distilled off under reduced pressure, and the residue was purified by preparative HPLC. After further purification by PTLC (NH silica gel, hexane/ethyl acetate = 1:1), the resulting solid was washed with isopropyl ether to afford the title compound (2.5 mg).

[0109] Tables 1-1 to 1-26 show the structural formulae and instrumental data of the compounds shown in Working Examples 1 to 12 and compounds synthesized in manners similar to those of the exemplified compounds. Each number shown in the column of "Example" in the tables indicates which one of the above Working Examples 1 to 12 the synthesis of each intended compound was based on. The column of "Configuration" shows the configuration of the carbon atom to which R⁴ is attached in the inventive compound as represented by formula [I], and the term "racemate" refers to a racemic mixture.

[0110]

[Table 1-1]

Compound	Example	Structure	Configuration	(ESI pos.) m/z (ESI neg.) m/z	¹ H-NMR
1	1		racemate	405([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.98 (t, J=7.3 Hz, 3 H), 1.33 – 1.42 (m, 2 H), 1.51 – 1.57 (m, 1 H), 1.81 – 1.89 (m, 1 H), 3.52 (dd, J=8.7, 7.3 Hz, 1 H), 3.77 – 3.86 (m, 2 H), 3.92 (s, 3 H), 3.96 (t, J=8.9 Hz, 1 H), 4.13 (d, J=15.6 Hz, 1 H), 6.53 (tt, J=8.7, 2.3 Hz, 1 H), 6.77 (d, J=9.2 Hz, 1 H), 7.09 – 7.15 (m, 2 H), 8.03 (dd, J=9.2, 2.8 Hz, 1 H), 8.10 (d, J=2.8 Hz, 1 H), 8.94 (br. s., 1 H)
2	2		racemate	489([M+H] ⁺)	
3	2		racemate	475([M+H] ⁺)	
4	2		racemate	443([M+H] ⁺)	
5	2		racemate	457([M+H] ⁺)	
6	2		R	475([M+H] ⁺)	
7	2		S	475([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 1.21 – 1.35 (m, 2 H), 2.22 – 2.32 (m, 1 H), 3.60 – 3.71 (m, 1 H), 3.90 – 3.97 (m, 2 H), 3.99 – 4.06 (m, 1 H), 4.08 – 4.15 (m, 1 H), 7.34 – 7.39 (m, 1 H), 7.40 – 7.47 (m, 1 H), 7.63 – 7.70 (m, 2 H), 7.85 (s, 1 H), 8.28 – 8.37 (m, 1 H), 8.71 (br. s., 1 H), 8.80 (d, J=2.9 Hz, 1 H)
8	2		S	529([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.93 – 1.89 (m, 13 H), 3.56 – 3.63 (m, 1 H), 3.94 – 4.04 (m, 2 H), 4.07 – 4.16 (m, 2 H), 7.33 – 7.38 (m, 1 H), 7.38 – 7.44 (m, 1 H), 7.63 – 7.69 (m, 2 H), 7.83 (s, 1 H), 8.27 (dd, J=8.7, 2.5 Hz, 1 H), 8.57 (s, 1 H), 8.81 (d, J=2.5 Hz, 1 H)
9	2		S	489([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.86 (d, J=6.6 Hz, 3 H), 1.03 (t, J=7.4 Hz, 3 H), 1.23 – 1.35 (m, 1 H), 1.37 – 1.47 (m, 1 H), 1.97 – 2.06 (m, 1 H), 3.65 (dd, J=9.5, 6.6 Hz, 1 H), 3.91 (t, J=9.5 Hz, 1 H), 4.01 – 4.06 (m, 2 H), 4.07 – 4.12 (m, 1 H), 7.35 – 7.40 (m, 1 H), 7.41 – 7.47 (m, 1 H), 7.64 – 7.72 (m, 2 H), 7.86 (s, 1 H), 8.29 – 8.37 (m, 1 H), 8.70 (br. s., 1 H), 8.79 (d, J=2.5 Hz, 1 H)










[0111] [Table 1-2]

Compound	Example	Structure	Configuration	(ESI pos.) m/z (ESI neg.) m/z	¹ H-NMR
10	2		S	489([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.01 (d, J=6.2 Hz, 3 H), 1.03 (d, J=6.2 Hz, 3 H), 1.46 – 1.88 (m, 3 H), 3.61 (dd, J=8.9, 7.6 Hz, 1 H), 3.93 – 4.02 (m, 2 H), 4.08 – 4.13 (m, 1 H), 4.16 (d, J=15.7 Hz, 1 H), 7.35 – 7.40 (m, 1 H), 7.41 – 7.47 (m, 1 H), 7.64 – 7.71 (m, 2 H), 7.84 (s, 1 H), 8.26 – 8.34 (m, 1 H), 8.53 (br. s., 1 H), 8.78 (d, J=2.9 Hz, 1 H)
11	2		S	447([M+Na] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.87 (d, J=7.0 Hz, 3 H), 1.01 (d, J=7.0 Hz, 3 H), 2.14 – 2.25 (m, 1 H), 3.79 – 3.92 (m, 2 H), 3.99 – 4.14 (m, 3 H), 7.28 – 7.49 (m, 3 H), 7.58 – 7.69 (m, 1 H), 7.83 (br. s., 1 H), 8.15 (d, J=2.9 Hz, 1 H), 8.22 – 8.35 (m, 1 H), 9.05 (br. s., 1 H)
12	2		S	509([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 3.53 (d, J=16.1 Hz, 1 H), 3.77 (dd, J=9.3, 7.6 Hz, 1 H), 4.16 (d, J=16.5 Hz, 1 H), 4.26 (t, J=9.3 Hz, 1 H), 4.98 (dd, J=9.3, 7.6 Hz, 1 H), 7.22 – 7.40 (m, 7 H), 7.46 – 7.52 (m, 1 H), 7.59 (d, J=9.1 Hz, 1 H), 7.70 (s, 1 H), 7.99 (br. s., 1 H), 8.27 (dd, J=8.7, 2.5 Hz, 1 H), 8.68 (d, J=2.5 Hz, 1 H)
13	2		S	489([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.05 (s, 9 H), 3.63 (dd, J=9.5, 5.4 Hz, 1 H), 3.72 (dd, J=9.5, 5.0 Hz, 1 H), 4.00 (d, J=16.1 Hz, 1 H), 4.06 (t, J=9.5 Hz, 1 H), 4.34 (d, J=16.1 Hz, 1 H), 7.35 – 7.40 (m, 1 H), 7.41 – 7.46 (m, 1 H), 7.66 – 7.71 (m, 2 H), 7.85 (s, 1 H), 8.33 (dd, J=8.7, 2.5 Hz, 1 H), 8.76 – 8.82 (m, 2 H)
14	2		S	461([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.00 (t, J=7.4 Hz, 3 H), 1.64 – 1.73 (m, 1 H), 1.92 – 2.00 (m, 1 H), 3.59 – 3.64 (m, 1 H), 3.90 – 3.96 (m, 1 H), 4.01 (d, J=15.7 Hz, 1 H), 4.06 – 4.16 (m, 2 H), 7.33 – 7.38 (m, 1 H), 7.40 – 7.46 (m, 1 H), 7.63 – 7.70 (m, 2 H), 7.85 (s, 1 H), 8.32 (dd, J=8.7, 2.5 Hz, 1 H), 8.56 (br. s., 1 H), 8.79 (d, J=2.5 Hz, 1 H)
15	2		S	485([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=6.6 Hz, 3 H), 1.01 (d, J=6.6 Hz, 3 H), 2.19 – 2.30 (m, 1 H), 3.58 (dd, J=8.9, 6.4 Hz, 1 H), 3.83 – 3.89 (m, 1 H), 3.90 – 3.95 (m, 1 H), 4.00 – 4.11 (m, 2 H), 7.31 – 7.48 (m, 3 H), 7.64 (d, J=8.3 Hz, 1 H), 7.85 (s, 1 H), 8.06 – 8.15 (m, 1 H), 8.40 (d, J=2.9 Hz, 1 H), 8.84 (br. s., 1 H)
16	2		S	443([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.03 (d, J=6.6 Hz, 3 H), 2.20 – 2.31 (m, 1 H), 3.66 (dd, J=7.8, 5.0 Hz, 1 H), 3.86 – 3.99 (m, 3 H), 4.13 (d, J=15.7 Hz, 1 H), 6.52 – 6.60 (m, 1 H), 7.10 – 7.18 (m, 2 H), 7.69 (d, J=8.7 Hz, 1 H), 8.33 (dd, J=8.7, 2.5 Hz, 1 H), 8.75 – 8.83 (m, 2 H)
17	1		S	475([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.20 – 2.29 (m, 1 H), 3.66 (dd, J=8.5, 5.6 Hz, 1 H), 3.87 – 4.05 (m, 3 H), 4.28 (d, J=16.1 Hz, 1 H), 7.26 – 7.30 (m, 1 H), 7.55 – 7.59 (m, 1 H), 7.61 – 7.64 (m, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 8.12 – 8.21 (m, 1 H), 8.36 – 8.45 (m, 2 H), 8.76 (d, J=2.5 Hz, 1 H)
18	3		S	476([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.92 (d, J=6.6 Hz, 3 H), 1.04 (d, J=6.6 Hz, 3 H), 2.20 (t, J=1.0 Hz, 1 H), 3.61 – 3.70 (m, 1 H), 3.91 – 4.03 (m, 3 H), 4.32 (d, J=16.5 Hz, 1 H), 7.44 (d, J=7.8 Hz, 1 H), 7.67 (d, J=9.1 Hz, 1 H), 7.88 (t, J=8.1 Hz, 1 H), 8.37 (d, J=8.7 Hz, 1 H), 8.40 – 8.46 (m, 1 H), 8.55 (br. s., 1 H), 8.74 (d, J=2.5 Hz, 1 H)










[0112] [Table 1-3]

Compound	Example	Structure	Configuration	(ESI pos.) m/z (ESI neg.) m/z	¹ H-NMR
19	3		S	476([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.91 (d, J=7.0 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.19 – 2.25 (m, 1 H), 3.61 – 3.68 (m, 1 H), 3.92 – 3.98 (m, 2 H), 4.06 (d, J=16.5 Hz, 1 H), 4.23 (d, J=16.5 Hz, 1 H), 7.26 – 7.30 (m, 1 H), 7.66 (d, J=8.7 Hz, 1 H), 8.40 – 8.43 (m, 1 H), 8.44 – 8.48 (m, 2 H), 8.74 (d, J=2.9 Hz, 1 H), 8.85 (br. s., 1 H)
20	4		S	432([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=7.0 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.23 – 2.31 (m, 1 H), 3.60 – 3.69 (m, 1 H), 3.89 – 4.01 (m, 2 H), 4.04 – 4.14 (m, 2 H), 7.34 – 7.41 (m, 1 H), 7.41 – 7.47 (m, 1 H), 7.62 – 7.73 (m, 2 H), 7.85 (s, 1 H), 8.30 (dd, J=8.7, 2.5 Hz, 1 H), 8.59 (br. s., 1 H), 8.81 (d, J=2.5 Hz, 1 H)
21	5		S	473([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.91 (d, J=6.6 Hz, 3 H), 1.03 (d, J=6.6 Hz, 3 H), 2.26 (td, J=6.9, 3.1 Hz, 1 H), 3.66 (dd, J=7.6, 5.2 Hz, 1 H), 3.87 – 3.94 (m, 2 H), 3.97 (d, J=15.7 Hz, 1 H), 4.16 (d, J=15.3 Hz, 1 H), 6.45 – 6.49 (m, 1 H), 7.33 – 7.47 (m, 1 H), 7.65 – 7.70 (m, 1 H), 7.71 – 7.75 (m, 1 H), 7.88 (s, 1 H), 7.99 (d, J=9.1 Hz, 1 H), 8.19 – 8.25 (m, 1 H), 8.52 (d, J=2.5 Hz, 1 H), 8.55 (d, J=2.5 Hz, 1 H), 8.91 (br. s., 1 H)
22	6		S	450([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 0.97 (d, J=7.0 Hz, 3 H), 2.14 – 2.25 (m, 1 H), 3.07 (s, 6 H), 3.55 (dd, J=8.5, 6.0 Hz, 1 H), 3.72 – 3.88 (m, 3 H), 4.15 (d, J=15.3 Hz, 1 H), 6.55 (d, J=9.5 Hz, 1 H), 7.30 – 7.45 (m, 2 H), 7.64 – 7.72 (m, 1 H), 7.80 – 7.88 (m, 2 H), 8.12 (d, J=2.5 Hz, 1 H), 9.19 (br. s., 1 H)
23	7		S	449([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=6.6 Hz, 3 H), 0.96 (t, J=7.2 Hz, 3 H), 1.00 (d, J=7.0 Hz, 3 H), 1.65 – 1.81 (m, 2 H), 2.15 – 2.28 (m, 1 H), 2.69 – 2.80 (m, 2 H), 3.55 – 3.65 (m, 1 H), 3.83 – 3.93 (m, 2 H), 3.96 – 4.14 (m, 2 H), 7.05 – 7.18 (m, 1 H), 7.28 – 7.42 (m, 2 H), 7.57 – 7.67 (m, 1 H), 7.85 (s, 1 H), 8.01 – 8.09 (m, 1 H), 8.57 (d, J=2.5 Hz, 1 H), 9.10 (br. s., 1 H)
24	3		S	432([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.22 – 2.30 (m, 1 H), 3.66 (dd, J=7.4, 4.5 Hz, 1 H), 3.89 – 4.00 (m, 3 H), 4.16 (d, J=15.7 Hz, 1 H), 7.38 – 7.45 (m, 2 H), 7.62 – 7.66 (m, 1 H), 7.69 (d, J=8.7 Hz, 1 H), 7.99 (s, 1 H), 8.32 (dd, J=8.7, 2.5 Hz, 1 H), 8.81 (d, J=2.5 Hz, 1 H), 8.87 (br. s., 1 H)
25	3		S	491([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.22 – 2.31 (m, 1 H), 3.62 – 3.68 (m, 1 H), 3.90 – 3.97 (m, 2 H), 3.99 – 4.04 (m, 1 H), 4.08 – 4.14 (m, 1 H), 6.96 – 7.01 (m, 1 H), 7.31 – 7.36 (m, 2 H), 7.59 (s, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 8.33 (dd, J=8.7, 2.5 Hz, 1 H), 8.63 (br. s., 1 H), 8.80 (d, J=2.5 Hz, 1 H)
26	3		S	507([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.23 – 2.31 (m, 1 H), 3.63 – 3.68 (m, 1 H), 3.91 – 3.97 (m, 2 H), 4.00 – 4.05 (m, 1 H), 4.10 – 4.14 (m, 1 H), 7.36 – 7.43 (m, 2 H), 7.62 – 7.66 (m, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 7.87 (s, 1 H), 8.34 (dd, J=8.7, 2.5 Hz, 1 H), 8.62 (br. s., 1 H), 8.80 (d, J=2.5 Hz, 1 H)
27	3		S	441([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=6.6 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.21 – 2.31 (m, 1 H), 3.60 – 3.69 (m, 1 H), 3.89 – 3.96 (m, 2 H), 3.98 – 4.05 (m, 1 H), 4.07 – 4.14 (m, 1 H), 7.06 – 7.14 (m, 1 H), 7.24 (t, J=8.3 Hz, 1 H), 7.31 – 7.36 (m, 1 H), 7.61 – 7.73 (m, 2 H), 8.33 (dd, J=8.7, 2.5 Hz, 1 H), 8.57 (br. s., 1 H), 8.80 (d, J=2.5 Hz, 1 H)










[0113] [Table 1-4]

Compound	Example	Structure	Configuration	(ESI pos.) m/z (ESI neg.) m/z	¹ H-NMR
28	3		S	435([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=6.6 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.23 (s, 3 H), 2.27 – 2.31 (m, 1 H), 2.31 (s, 3 H), 3.63 (dd, J=6.8, 4.3 Hz, 1 H), 3.86 – 3.97 (m, 2 H), 4.01 – 4.14 (m, 2 H), 6.88 (d, J=7.8 Hz, 1 H), 7.05 (d, J=7.4 Hz, 1 H), 7.66 (d, J=8.7 Hz, 1 H), 7.76 (s, 1 H), 8.24 (br. s., 1 H), 8.27 – 8.34 (m, 1 H), 8.82 (d, J=2.5 Hz, 1 H)
29	3		S	467([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=6.6 Hz, 3 H), 1.02 (d, J=7.0 Hz, 3 H), 2.19 – 2.27 (m, 1 H), 3.63 (dd, J=9.1, 6.2 Hz, 1 H), 3.78 (s, 3 H), 3.82 (s, 3 H), 3.87 – 4.03 (m, 3 H), 4.31 (d, J=16.1 Hz, 1 H), 6.59 (dd, J=8.9, 3.1 Hz, 1 H), 6.79 (d, J=9.1 Hz, 1 H), 7.66 (d, J=9.1 Hz, 1 H), 8.04 (d, J=3.3 Hz, 1 H), 8.36 (dd, J=8.7, 2.5 Hz, 1 H), 8.45 (br. s., 1 H), 8.81 (d, J=2.5 Hz, 1 H)
30	3		S	475([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.19 – 2.31 (m, 1 H), 3.60 – 3.71 (m, 1 H), 3.89 – 3.98 (m, 2 H), 3.99 – 4.06 (m, 1 H), 4.10 – 4.18 (m, 1 H), 7.53 – 7.60 (m, 2 H), 7.63 (d, J=8.3 Hz, 2 H), 7.68 (d, J=8.7 Hz, 1 H), 8.30 (dd, J=8.7, 2.5 Hz, 1 H), 8.77 – 8.85 (m, 2 H)
31	3		S	450([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 1.02 (d, J=7.0 Hz, 3 H), 2.23 – 2.32 (m, 1 H), 2.95 (s, 6 H), 3.63 (dd, J=8.7, 5.8 Hz, 1 H), 3.88 – 3.98 (m, 2 H), 3.98 – 4.05 (m, 1 H), 4.07 – 4.13 (m, 1 H), 6.46 – 6.52 (m, 1 H), 6.73 – 6.78 (m, 1 H), 7.03 (s, 1 H), 7.16 (t, J=8.3 Hz, 1 H), 7.66 (d, J=8.7 Hz, 1 H), 8.16 (br. s., 1 H), 8.34 (dd, J=8.7, 2.5 Hz, 1 H), 8.80 (d, J=2.5 Hz, 1 H)
32	3		S	485([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.22 – 2.31 (m, 1 H), 3.06 (s, 3 H), 3.63 – 3.71 (m, 1 H), 3.89 – 3.98 (m, 2 H), 4.00 – 4.07 (m, 1 H), 4.10 – 4.18 (m, 1 H), 7.52 (t, J=8.1 Hz, 1 H), 7.65 – 7.73 (m, 2 H), 7.83 – 7.90 (m, 1 H), 8.05 – 8.10 (m, 1 H), 8.31 – 8.37 (m, 1 H), 8.78 – 8.82 (m, 1 H), 8.84 (s, 1 H)
33	2		S	523([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 2.84 (dd, J=13.6, 8.7 Hz, 1 H), 3.29 (dd, J=13.6, 5.0 Hz, 1 H), 3.65 (dd, J=9.3, 6.4 Hz, 1 H), 3.93 (t, J=9.1 Hz, 1 H), 4.04 – 4.09 (m, 1 H), 4.15 – 4.19 (m, 1 H), 4.23 – 4.31 (m, 1 H), 7.18 – 7.24 (m, 2 H), 7.27 – 7.39 (m, 4 H), 7.41 – 7.48 (m, 1 H), 7.59 – 7.70 (m, 2 H), 7.84 (s, 1 H), 8.15 – 8.23 (m, 1 H), 8.36 (br. s., 1 H), 8.74 (d, J=2.5 Hz, 1 H)
34	2		S	475([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.02 (t, J=7.2 Hz, 3 H), 1.36 – 1.48 (m, 2 H), 1.58 – 1.68 (m, 1 H), 1.86 – 1.96 (m, 1 H), 3.62 (dd, J=9.1, 7.0 Hz, 1 H), 3.92 – 3.98 (m, 1 H), 4.01 (d, J=16.1 Hz, 1 H), 4.06 – 4.18 (m, 2 H), 7.34 – 7.40 (m, 1 H), 7.40 – 7.47 (m, 1 H), 7.63 – 7.71 (m, 2 H), 7.85 (s, 1 H), 8.31 (dd, J=8.7, 2.5 Hz, 1 H), 8.57 (br. s., 1 H), 8.78 (d, J=2.5 Hz, 1 H)
35	5		S	507([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.91 (d, J=6.6 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.24 – 2.30 (m, 1 H), 3.62 – 3.69 (m, 1 H), 3.89 – 3.96 (m, 2 H), 4.01 (d, J=15.7 Hz, 1 H), 4.14 (d, J=15.7 Hz, 1 H), 7.21 (s, 1 H), 7.34 – 7.40 (m, 2 H), 7.42 – 7.47 (m, 1 H), 7.60 – 7.63 (m, 1 H), 7.68 – 7.73 (m, 1 H), 7.86 (s, 1 H), 8.29 (s, 1 H), 8.35 – 8.40 (m, 1 H), 8.49 (d, J=2.5 Hz, 1 H), 8.86 (br. s., 1 H)
36	3		S	437([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.84 (d, J=6.6 Hz, 3 H), 0.98 (d, J=7.0 Hz, 3 H), 2.17 – 2.27 (m, 1 H), 3.57 – 3.63 (m, 1 H), 3.75 (s, 3 H), 3.83 – 3.92 (m, 2 H), 3.96 – 4.05 (m, 2 H), 6.61 – 6.65 (m, 1 H), 6.91 – 6.96 (m, 1 H), 7.17 (t, J=8.3 Hz, 1 H), 7.20 – 7.23 (m, 1 H), 7.62 (d, J=8.7 Hz, 1 H), 8.25 – 8.32 (m, 2 H), 8.74 (d, J=2.5 Hz, 1 H)

[0114] [Table 1-5]

Compound	Example	Structure	Configuration	(ESI pos.) m/z (ESI neg.) m/z	¹ H-NMR
37	3		S	499([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.87 (d, J=6.6 Hz, 3 H), 1.01 (d, J=7.0 Hz, 3 H), 2.17 – 2.30 (m, 1 H), 3.59 – 3.67 (m, 1 H), 3.86 – 3.95 (m, 2 H), 3.99 – 4.11 (m, 2 H), 6.71 – 6.78 (m, 1 H), 7.00 (d, J=7.8 Hz, 2 H), 7.07 – 7.14 (m, 1 H), 7.17 – 7.37 (m, 5 H), 7.66 (d, J=8.7 Hz, 1 H), 8.33 (dd, J=8.7, 2.5 Hz, 1 H), 8.39 (br. s., 1 H), 8.72 – 8.80 (m, 1 H)
38	3		S	499([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.23 – 2.34 (m, 1 H), 2.79 (s, 3 H), 3.62 – 3.67 (m, 1 H), 3.90 – 3.99 (m, 2 H), 4.10 (s, 2 H), 7.43 – 7.53 (m, 2 H), 7.64 – 7.73 (m, 2 H), 7.81 (d, J=7.4 Hz, 1 H), 8.19 – 8.25 (m, 1 H), 8.34 (dd, J=8.7, 2.5 Hz, 1 H), 8.53 (s, 1 H), 8.67 (d, J=5.0 Hz, 1 H), 8.82 (d, J=2.5 Hz, 1 H)
39	3		S	483([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.24 – 2.34 (m, 1 H), 3.65 (dd, J=8.1, 5.2 Hz, 1 H), 3.89 – 3.99 (m, 2 H), 4.04 – 4.14 (m, 2 H), 7.32 – 7.47 (m, 5 H), 7.52 (s, 1 H), 7.56 – 7.60 (m, 2 H), 7.67 (d, J=8.7 Hz, 1 H), 7.75 (s, 1 H), 8.35 (dd, J=8.7, 2.5 Hz, 1 H), 8.46 (br. s., 1 H), 8.77 – 8.84 (m, 1 H)
40	3		S	426([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.20 – 2.29 (m, 1 H), 3.67 (dd, J=8.7, 5.8 Hz, 1 H), 3.87 – 4.00 (m, 3 H), 4.17 (d, J=15.7 Hz, 1 H), 7.16 (d, J=5.4 Hz, 1 H), 7.29 – 7.34 (m, 1 H), 7.69 (d, J=9.1 Hz, 1 H), 8.10 (d, J=5.8 Hz, 1 H), 8.26 – 8.32 (m, 1 H), 8.82 (d, J=2.9 Hz, 1 H), 9.17 (br. s., 1 H)
41	3		S	426([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=7.0 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.22 – 2.31 (m, 1 H), 3.67 (dd, J=7.6, 4.3 Hz, 1 H), 3.91 – 3.97 (m, 2 H), 4.00 (d, J=15.7 Hz, 1 H), 4.16 (d, J=15.7 Hz, 1 H), 6.92 (dd, J=8.7, 3.3 Hz, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 8.09 – 8.16 (m, 1 H), 8.23 – 8.28 (m, 1 H), 8.30 – 8.35 (m, 1 H), 8.74 (br. s., 1 H), 8.80 (d, J=2.5 Hz, 1 H)
42	3		S	477([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.94 (d, J=7.0 Hz, 3 H), 1.02 (d, J=7.0 Hz, 3 H), 2.06 – 2.22 (m, 1 H), 3.62 (dd, J=8.9, 6.4 Hz, 1 H), 3.91 – 3.98 (m, 1 H), 4.00 – 4.06 (m, 1 H), 4.34 (d, J=17.8 Hz, 1 H), 4.86 (d, J=17.8 Hz, 1 H), 7.36 (d, J=5.0 Hz, 1 H), 7.64 (d, J=9.1 Hz, 1 H), 8.33 – 8.41 (m, 1 H), 8.51 (br. s., 1 H), 8.76 (d, J=2.9 Hz, 1 H), 8.86 (d, J=5.0 Hz, 1 H)
43	3		S	482([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=6.6 Hz, 3 H), 1.03 (d, J=6.6 Hz, 3 H), 2.13 – 2.26 (m, 1 H), 3.66 (dd, J=9.1, 5.8 Hz, 1 H), 3.84 – 4.01 (m, 2 H), 4.11 – 4.24 (m, 2 H), 7.37 – 7.46 (m, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 8.37 – 8.47 (m, 1 H), 8.71 (d, J=2.5 Hz, 1 H), 9.95 (br. s., 1 H)
44	5		S	487([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.91 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.23 – 2.28 (m, 1 H), 2.31 (s, 3 H), 3.59 – 3.70 (m, 1 H), 3.85 – 3.95 (m, 2 H), 4.02 (d, J=15.7 Hz, 1 H), 4.09 – 4.18 (m, 1 H), 7.29 – 7.33 (m, 2 H), 7.34 – 7.40 (m, 1 H), 7.40 – 7.46 (m, 1 H), 7.66 – 7.75 (m, 1 H), 7.86 (s, 1 H), 8.20 (s, 1 H), 8.34 (dd, J=9.1, 2.9 Hz, 1 H), 8.46 (d, J=2.9 Hz, 1 H), 8.92 (br. s., 1 H)
45	5		S	487([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.91 (d, J=7.0 Hz, 3 H), 1.04 (d, J=6.6 Hz, 3 H), 2.23 – 2.31 (m, 1 H), 2.36 (s, 3 H), 3.62 – 3.71 (m, 1 H), 3.90 – 3.97 (m, 2 H), 4.01 (d, J=15.7 Hz, 1 H), 4.14 (d, J=15.7 Hz, 1 H), 6.90 (s, 1 H), 7.31 – 7.40 (m, 2 H), 7.41 – 7.47 (m, 1 H), 7.66 – 7.73 (m, 1 H), 7.81 – 7.89 (m, 2 H), 8.39 (dd, J=8.7, 2.9 Hz, 1 H), 8.59 (d, J=2.9 Hz, 1 H), 8.81 (br. s., 1 H)

[0115] [Table 1-6]

Compound	Example	Structure	Configuration	(ESI pos.) m/z (ESI neg.) m/z	¹ H-NMR
46	7		S	435 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=7.0 Hz, 3 H), 1.01 (d, J=7.0 Hz, 3 H), 1.30 (t, J=7.6 Hz, 3 H), 2.19 – 2.30 (m, 1 H), 2.82 (q, J=7.4 Hz, 2 H), 3.62 (dd, J=8.5, 6.0 Hz, 1 H), 3.81 – 3.96 (m, 3 H), 4.16 (d, J=15.3 Hz, 1 H), 7.18 (d, J=8.3 Hz, 1 H), 7.32 – 7.38 (m, 1 H), 7.40 – 7.46 (m, 1 H), 7.65 – 7.71 (m, 1 H), 7.86 (s, 1 H), 8.04 – 8.12 (m, 1 H), 8.56 (d, J=2.5 Hz, 1 H), 8.99 (br. s., 1 H)
47	7		S	421 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.01 (d, J=7.0 Hz, 3 H), 2.18 – 2.27 (m, 1 H), 2.54 (s, 3 H), 3.61 (dd, J=8.3, 5.8 Hz, 1 H), 3.80 – 3.96 (m, 3 H), 4.15 (d, J=15.7 Hz, 1 H), 7.16 (d, J=8.7 Hz, 1 H), 7.33 – 7.38 (m, 1 H), 7.39 – 7.46 (m, 1 H), 7.64 – 7.71 (m, 1 H), 7.85 (s, 1 H), 8.02 – 8.09 (m, 1 H), 8.52 (d, J=2.5 Hz, 1 H), 8.96 (br. s., 1 H)
48	8		S	478 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=7.0 Hz, 3 H), 1.02 (d, J=7.0 Hz, 3 H), 2.21 – 2.29 (m, 1 H), 3.14 (s, 3 H), 3.15 (s, 3 H), 3.61 – 3.67 (m, 1 H), 3.86 – 3.96 (m, 2 H), 3.98 (d, J=15.7 Hz, 1 H), 4.14 (d, J=15.7 Hz, 1 H), 7.34 – 7.39 (m, 1 H), 7.40 – 7.46 (m, 1 H), 7.64 – 7.69 (m, 1 H), 7.73 (d, J=8.7 Hz, 1 H), 7.86 (s, 1 H), 8.09 – 8.17 (m, 1 H), 8.75 (d, J=2.5 Hz, 1 H), 8.83 (br. s., 1 H)
49	1		S	477 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=6.6 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.17 – 2.28 (m, 1 H), 3.85 – 3.94 (m, 2 H), 4.03 – 4.17 (m, 2 H), 4.27 (d, J=16.1 Hz, 1 H), 8.40 – 8.47 (m, 2 H), 8.82 – 8.87 (m, 3 H)
50	3		S	425 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.02 (d, J=6.6 Hz, 3 H), 2.20 – 2.32 (m, 1 H), 3.56 – 3.67 (m, 1 H), 3.87 – 3.95 (m, 2 H), 3.97 – 4.04 (m, 1 H), 4.05 – 4.12 (m, 1 H), 6.96 – 7.06 (m, 2 H), 7.42 – 7.54 (m, 2 H), 7.66 (d, J=8.7 Hz, 1 H), 8.31 (dd, J=8.7, 2.5 Hz, 1 H), 8.44 (br. s., 1 H), 8.80 (d, J=2.5 Hz, 1 H)
51	3		S	441 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.19 – 2.33 (m, 1 H), 3.60 – 3.69 (m, 1 H), 3.89 – 3.96 (m, 2 H), 3.97 – 4.04 (m, 1 H), 4.06 – 4.14 (m, 1 H), 7.28 (d, J=9.1 Hz, 2 H), 7.47 (d, J=9.1 Hz, 2 H), 7.67 (d, J=8.7 Hz, 1 H), 8.32 (dd, J=8.7, 2.5 Hz, 1 H), 8.53 (br. s., 1 H), 8.81 (d, J=2.5 Hz, 1 H)
52	3		S	432 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.21 – 2.31 (m, 1 H), 3.66 (dd, J=7.8, 5.0 Hz, 1 H), 3.86 – 4.02 (m, 3 H), 4.16 (d, J=15.7 Hz, 1 H), 7.56 – 7.72 (m, 5 H), 8.26 – 8.33 (m, 1 H), 8.83 (d, J=2.5 Hz, 1 H), 8.99 (br. s., 1 H)
53	3		S	491 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.21 – 2.34 (m, 1 H), 3.61 – 3.69 (m, 1 H), 3.89 – 3.97 (m, 2 H), 4.02 (s, 1 H), 4.08 – 4.15 (m, 1 H), 7.18 (d, J=8.7 Hz, 2 H), 7.55 (d, J=9.1 Hz, 2 H), 7.68 (d, J=8.7 Hz, 1 H), 8.31 (dd, J=8.7, 2.5 Hz, 1 H), 8.62 (br. s., 1 H), 8.81 (d, J=2.5 Hz, 1 H)
54	3		S	483 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.91 (d, J=6.6 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.12 – 2.24 (m, 1 H), 3.69 (dd, J=8.9, 6.0 Hz, 1 H), 3.88 – 4.03 (m, 2 H), 4.27 – 4.39 (m, 2 H), 7.68 (d, J=9.1 Hz, 1 H), 8.37 – 8.45 (m, 1 H), 8.73 (d, J=2.5 Hz, 1 H)

[0116] [Table 1-7]

Compound	Example	Structure	Configuration	(ESI pos.) m/z (ESI neg.) m/z	¹ H-NMR
55	3		S	438([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.87 (d, J=6.6 Hz, 3 H), 1.02 (d, J=7.0 Hz, 3 H), 2.20 – 2.29 (m, 1 H), 3.65 (dd, J=8.1, 5.2 Hz, 1 H), 3.87 – 4.01 (m, 6 H), 4.12 (d, J=15.7 Hz, 1 H), 6.94 – 6.99 (m, 1 H), 7.00 – 7.05 (m, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 8.06 (d, J=5.8 Hz, 1 H), 8.30 – 8.37 (m, 1 H), 8.71 (br. s., 1 H), 8.79 (d, J=2.5 Hz, 1 H)
56	3		S	433([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.14 – 2.28 (m, 1 H), 3.67 (dd, J=8.3, 5.8 Hz, 1 H), 3.87 – 4.00 (m, 2 H), 4.07 – 4.21 (m, 2 H), 7.67 (d, J=9.1 Hz, 1 H), 7.95 (dd, J=8.7, 2.5 Hz, 1 H), 8.30 (d, J=8.7 Hz, 1 H), 8.37 – 8.45 (m, 1 H), 8.55 – 8.61 (m, 1 H), 8.75 (d, J=2.9 Hz, 1 H), 9.03 (s, 1 H)
57	3		S	438([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.02 (d, J=6.6 Hz, 3 H), 2.18 – 2.27 (m, 1 H), 3.63 (dd, J=8.5, 5.6 Hz, 1 H), 3.85 (s, 3 H), 3.89 – 4.03 (m, 3 H), 4.20 – 4.31 (m, 1 H), 6.50 (d, J=8.3 Hz, 1 H), 7.54 – 7.60 (m, 1 H), 7.62 – 7.73 (m, 2 H), 8.24 (br. s., 1 H), 8.37 (dd, J=8.7, 2.5 Hz, 1 H), 8.78 (d, J=2.5 Hz, 1 H)
58	3		S	438([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.22 – 2.32 (m, 1 H), 3.60 – 3.69 (m, 1 H), 3.88 – 3.97 (m, 5 H), 3.99 – 4.06 (m, 1 H), 4.08 – 4.15 (m, 1 H), 6.73 (d, J=8.7 Hz, 1 H), 7.67 (d, J=9.1 Hz, 1 H), 7.84 (dd, J=8.9, 2.7 Hz, 1 H), 8.22 (d, J=2.5 Hz, 1 H), 8.31 – 8.39 (m, 2 H), 8.79 (d, J=2.5 Hz, 1 H)
59	3		S	476([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.21 – 2.33 (m, 1 H), 3.68 (dd, J=7.8, 5.0 Hz, 1 H), 3.88 – 4.05 (m, 3 H), 4.20 (d, J=15.7 Hz, 1 H), 7.64 (d, J=8.7 Hz, 1 H), 7.69 (d, J=8.7 Hz, 1 H), 8.25 – 8.35 (m, 2 H), 8.70 (d, J=2.5 Hz, 1 H), 8.80 (d, J=2.5 Hz, 1 H), 9.10 (s, 1 H)
60	3		S	438([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.22 – 2.31 (m, 1 H), 3.61 – 3.69 (m, 1 H), 3.86 (s, 3 H), 3.90 – 3.98 (m, 2 H), 4.00 – 4.06 (m, 1 H), 4.10 – 4.16 (m, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 7.77 – 7.84 (m, 1 H), 8.08 (d, J=2.5 Hz, 1 H), 8.15 (d, J=2.1 Hz, 1 H), 8.32 – 8.39 (m, 1 H), 8.63 (s, 1 H), 8.78 (d, J=2.9 Hz, 1 H)
61	3		S	442([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.14 – 2.24 (m, 1 H), 3.60 – 3.69 (m, 1 H), 3.89 – 3.99 (m, 2 H), 4.04 (d, J=16.5 Hz, 1 H), 4.21 (d, J=16.5 Hz, 1 H), 7.04 – 7.10 (m, 1 H), 7.66 (d, J=8.7 Hz, 1 H), 8.18 (d, J=5.4 Hz, 1 H), 8.24 (s, 1 H), 8.39 – 8.45 (m, 1 H), 8.67 (br. s., 1 H), 8.73 (d, J=2.5 Hz, 1 H)
62	3		S	443([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.02 (d, J=7.0 Hz, 3 H), 2.13 – 2.26 (m, 1 H), 3.80 – 3.91 (m, 2 H), 4.00 – 4.14 (m, 2 H), 4.18 – 4.28 (m, 1 H), 7.24 – 7.26 (m, 1 H), 8.40 – 8.46 (m, 2 H), 8.58 (s, 2 H), 8.97 (br. s., 1 H)
63	3		S	422([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.02 (d, J=7.0 Hz, 3 H), 2.15 – 2.26 (m, 1 H), 2.36 (s, 3 H), 3.59 – 3.67 (m, 1 H), 3.89 – 4.03 (m, 3 H), 4.25 – 4.32 (m, 1 H), 6.86 – 6.91 (m, 1 H), 7.65 (d, J=8.7 Hz, 1 H), 8.00 (br. s., 1 H), 8.13 (d, J=5.0 Hz, 1 H), 8.40 – 8.49 (m, 2 H), 8.72 (d, J=2.9 Hz, 1 H)











[0117] [Table 1-8]

Compound	Example	Structure	Configuration	(ESI pos.) m/z (ESI neg.) m/z	¹ H-NMR
64	3		S	426 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=6.6 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.17 – 2.27 (m, 1 H), 3.59 – 3.68 (m, 1 H), 3.89 – 3.98 (m, 2 H), 4.03 (d, J=16.5 Hz, 1 H), 4.22 (d, J=16.1 Hz, 1 H), 7.37 – 7.49 (m, 1 H), 7.66 (d, J=9.1 Hz, 1 H), 8.11 – 8.22 (m, 2 H), 8.43 (dd, J=8.7, 2.5 Hz, 1 H), 8.63 (br. s., 1 H), 8.74 (d, J=2.5 Hz, 1 H)
65	3		S	442 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.91 (d, J=6.6 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.14 – 2.26 (m, 1 H), 3.61 – 3.69 (m, 1 H), 3.89 – 3.96 (m, 2 H), 4.04 (d, J=16.5 Hz, 1 H), 4.22 (d, J=16.5 Hz, 1 H), 7.02 – 7.10 (m, 1 H), 7.66 (d, J=8.7 Hz, 1 H), 8.18 (d, J=5.4 Hz, 1 H), 8.25 (s, 1 H), 8.40 – 8.45 (m, 1 H), 8.69 (br. s., 1 H), 8.74 (d, J=2.5 Hz, 1 H)
66	3		S	489 ([M+H] ⁺)	
67	3		S	425 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.88 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.21 – 2.32 (m, 1 H), 3.61 – 3.68 (m, 1 H), 3.89 – 3.96 (m, 2 H), 3.98 – 4.04 (m, 1 H), 4.07 – 4.14 (m, 1 H), 6.77 – 6.85 (m, 1 H), 7.10 – 7.16 (m, 1 H), 7.21 – 7.30 (m, 1 H), 7.46 – 7.53 (m, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 8.27 – 8.35 (m, 1 H), 8.60 (br. s., 1 H), 8.80 (d, J=2.5 Hz, 1 H)
68	3		S	421 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=6.6 Hz, 3 H), 1.02 (d, J=7.0 Hz, 3 H), 2.22 – 2.30 (m, 1 H), 2.34 (s, 3 H), 3.64 (dd, J=7.6, 4.7 Hz, 1 H), 3.87 – 3.98 (m, 2 H), 4.06 (s, 2 H), 6.91 – 6.97 (m, 1 H), 7.20 (t, J=7.8 Hz, 1 H), 7.27 – 7.32 (m, 1 H), 7.35 (s, 1 H), 7.67 (d, J=8.7 Hz, 1 H), 8.25 – 8.38 (m, 2 H), 8.80 (d, J=2.5 Hz, 1 H)
69	3		S	486 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=7.0 Hz, 3 H), 1.03 (d, J=6.6 Hz, 3 H), 2.19 – 2.29 (m, 1 H), 3.64 (dd, J=8.7, 5.8 Hz, 1 H), 3.94 – 4.08 (m, 3 H), 4.18 (d, J=16.1 Hz, 1 H), 5.04 (s, 2 H), 7.34 (t, J=7.8 Hz, 1 H), 7.54 – 7.60 (m, 1 H), 7.66 (d, J=8.7 Hz, 2 H), 7.93 (s, 1 H), 8.15 – 8.21 (m, 1 H), 8.86 – 8.91 (m, 1 H), 8.98 (s, 1 H)
70	3		S	433 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.04 (d, J=7.0 Hz, 3 H), 2.17 – 2.26 (m, 1 H), 3.61 – 3.70 (m, 1 H), 3.88 – 3.99 (m, 2 H), 4.07 – 4.13 (m, 1 H), 4.14 – 4.21 (m, 1 H), 7.26 – 7.28 (m, 1 H), 7.66 (d, J=8.7 Hz, 1 H), 8.39 (dd, J=8.9, 2.7 Hz, 1 H), 8.43 – 8.49 (m, 2 H), 8.76 (d, J=2.5 Hz, 1 H), 9.00 (br. s., 1 H)
71	3		S	476 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.15 – 2.26 (m, 1 H), 3.58 – 3.69 (m, 1 H), 3.86 – 3.99 (m, 2 H), 4.08 (d, J=16.5 Hz, 1 H), 4.17 – 4.25 (m, 1 H), 7.66 (d, J=9.1 Hz, 1 H), 7.92 (dd, J=8.7, 2.1 Hz, 1 H), 8.25 – 8.33 (m, 1 H), 8.41 (dd, J=8.7, 2.5 Hz, 1 H), 8.51 – 8.57 (m, 1 H), 8.74 (d, J=2.5 Hz, 1 H), 8.90 (br. s., 1 H)
72	3		S	442 ([M+H] ⁺)	¹ H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=6.6 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 2.14 – 2.27 (m, 1 H), 3.58 – 3.68 (m, 1 H), 3.88 – 3.99 (m, 2 H), 4.04 (d, J=16.1 Hz, 1 H), 4.21 (d, J=16.5 Hz, 1 H), 7.64 – 7.71 (m, 2 H), 8.10 – 8.19 (m, 1 H), 8.24 (d, J=2.5 Hz, 1 H), 8.43 (dd, J=8.7, 2.5 Hz, 1 H), 8.64 (br. s., 1 H), 8.74 (d, J=2.9 Hz, 1 H)

[0118] [Table 1-9]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
73	3		S	494 ([M+H] ⁺)	1.164 condition A
74	3		S	477 ([M+H] ⁺)	1.079 condition A
75	2		S	499 ([M+H] ⁺)	1.155 condition A
76	2		S	500 ([M+H] ⁺)	1.126 condition A
77	3		S	477 ([M+H] ⁺)	1.066 condition A
78	2		S	432 ([M+H] ⁺)	2.63 condition C
79	2		S	476 ([M+H] ⁺)	2.80 condition C
80	4		S	446 ([M+H] ⁺)	1.081 condition A
81	7		S	449 ([M+H] ⁺)	0.750 condition A
82	3		S	426 ([M+H] ⁺)	1.001 condition A

[0119] [Table 1-10]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
83	3		S	438 ([M+H] ⁺)	0.740 condition A
84	2		S	441 ([M+H] ⁺)	1.221 condition A
85	2		S	426 ([M+H] ⁺)	0.985 condition A
86	3		S	446 ([M+H] ⁺)	1.052 condition A
87	3		S	490 ([M+H] ⁺)	1.146 condition A
88	3		S	452 ([M+H] ⁺)	0.933 condition A
89	3		S	490 ([M+H] ⁺)	1.096 condition A
90	4		S	447 ([M+H] ⁺)	1.044 condition A
91	3		S	476 ([M+H] ⁺)	1.037 condition A
92	3		S	466 ([M+H] ⁺)	1.098 condition A

[0120] [Table 1-11]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
93	1		S	439 ([M+H] ⁺)	0.922 condition A
94	1		S	437 ([M+H] ⁺)	0.977 condition A
95	7		S	450 ([M+H] ⁺)	0.702 condition A
96	3		S	436 ([M+H] ⁺)	0.994 condition A
97	3		S	458 ([M+H] ⁺)	0.947 condition A
98	3		S	454 ([M+H] ⁺)	1.109 condition A
99	3		S	456 ([M+H] ⁺)	1.135 condition A
100	3		S	456 ([M+H] ⁺)	1.061 condition A
101	3		S	490 ([M+H] ⁺)	1.061 condition A
102	1		S	427 ([M+H] ⁺)	1.87 condition C

[0121] [Table 1-12]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
103	3		S	454 ([M+H] ⁺)	1.098 condition A
104	1		S	437 ([M+H] ⁺)	0.918 condition A
105	3		S	500 ([M+H] ⁺)	1.076 condition A
106	3		S	456 ([M+H] ⁺)	1.90 condition B
107	7		S	450 ([M+H] ⁺)	1.48 condition B
108	11		S	464 ([M+H] ⁺)	1.84 condition B
109	3		S	470 ([M+H] ⁺)	1.96 condition B
110	3		S	470 ([M+H] ⁺)	2.06 condition B
111	10		S	486 ([M+H] ⁺)	1.95 condition B
112	10		S	472 ([M+H] ⁺)	1.88 condition B











[0122] [Table 1-13]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
113	2		S	479 ([M+H] ⁺)	1.92 condition B
114	12		S	462 ([M+H] ⁺)	0.813 condition A
115	3		S	470 ([M+H] ⁺)	1.081 condition A
116	3		S	456 ([M+H] ⁺)	1.037 condition A
117	3		S	456 ([M+H] ⁺)	1.121 condition A
118	3		S	474 ([M+H] ⁺)	1.063 condition A
119	3		S	508 ([M+H] ⁺)	1.113 condition A
120	3		S	486 ([M+H] ⁺)	1.195 condition A
121	3		S	466 ([M+H] ⁺)	1.078 condition A
122	1		S	508 ([M+H] ⁺)	1.92 condition B

[0123] [Table 1-14]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
123	9		S	443 ([M+H] ⁺)	1.74 condition B
124	1		S	474 ([M+H] ⁺)	1.92 condition B
125	3		S	474 ([M+H] ⁺)	1.024 condition A
126	3		S	440 ([M+H] ⁺)	1.034 condition A
127	9		S	477 ([M+H] ⁺)	1.87 condition B
128	9		S	494 ([M+H] ⁺)	1.97 condition B
129	9		S	477 ([M+H] ⁺)	1.82 condition B
130	3		S	488 ([M+H] ⁺)	1.073 condition A
131	3		S	454 ([M+H] ⁺)	1.088 condition A
132	3		S	476 ([M+H] ⁺)	1.068 condition A

[0124] [Table 1-15]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
133	3		S	442 ([M+H] ⁺)	1.084 condition A
134	3		S	442 ([M+H] ⁺)	1.003 condition A
135	3		S	508 ([M+H] ⁺)	1.162 condition A
136	3		S	474 ([M+H] ⁺)	1.188 condition A
137	2		R	494 ([M+H] ⁺)	1.016 condition A
138	3		S	490 ([M+H] ⁺)	1.230 condition A
139	2		R	506 ([M+H] ⁺)	1.021 condition A
140	3		S	470 ([M+H] ⁺)	1.193 condition A
141	2		R	494 ([M+H] ⁺)	1.026 condition A
142	3		S	463 ([M+H] ⁺)	1.010 condition A

[0125] [Table 1-16]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
143	3		S	488 ([M+H] ⁺)	1.114 condition A
144	2		S	460 ([M+H] ⁺)	1.131 condition A
145	2		S	508 ([M+H] ⁺)	1.100 condition A
146	1		S	454 ([M+H] ⁺)	1.009 condition A
147	1		S	466 ([M+H] ⁺)	0.965 condition A
148	2		R	506 ([M+H] ⁺)	1.076 condition A
149	2		S	494 ([M+H] ⁺)	12.16 condition A
150	2		R	494 ([M+H] ⁺)	1.068 condition A
151	2		S	506 ([M+H] ⁺)	1.168 condition A
152	2		S	472 ([M+H] ⁺)	1.133 condition A











[0126] [Table 1-17]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
153	2		R	512 ([M+H] ⁺)	1.123 condition A
154	2		S	460 ([M+H] ⁺)	2.27 condition B
155	2		R	494 ([M+H] ⁺)	1.075 condition A
156	1		R	472 ([M+H] ⁺)	0.969 condition A
157	1		R	484 ([M+H] ⁺)	0.924 condition A
158	3		S	443 ([M+H] ⁺)	1.015 condition A
159	3		S	409 ([M+H] ⁺)	0.958 condition A
160	3		S	403 ([M+H] ⁺)	0.788 condition A
161	9		S	427 ([M+H] ⁺)	0.943 condition A
162	3		S	460 ([M+H] ⁺)	1.114 condition A

[0127] [Table 1-18]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
163	3		S	443 ([M+H] ⁺)	1.029 condition A
164	3		S	454 ([M+H] ⁺)	0.986 condition A
165	3		S	437 ([M+H] ⁺)	0.906 condition A
166	3		S	466 ([M+H] ⁺)	0.943 condition A
167	3		S	449 ([M+H] ⁺)	0.866 condition A
168	3		S	448 ([M+H] ⁺)	0.890 condition A
169	9		S	426 ([M+H] ⁺)	1.109 condition A
170	9		S	442 ([M+H] ⁺)	1.194 condition A
171	9		S	476 ([M+H] ⁺)	1.218 condition A
172	9		S	438 ([M+H] ⁺)	1.038 condition A

[0128] [Table 1-19]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
173	9		S	452 ([M+H] ⁺)	1.107 condition A
174	9		S	460 ([M+H] ⁺)	1.109 condition A
175	9		S	427 ([M+H] ⁺)	0.927 condition A
176	9		S	424 ([M+H] ⁺)	0.861 condition A
177	9		S	432 ([M+H] ⁺)	0.861 condition A
178	9		S	399 ([M+H] ⁺)	0.643 condition A
179	9		S	415 ([M+H] ⁺)	0.727 condition A
180	9		S	449 ([M+H] ⁺)	0.812 condition A
181	9		S	411 ([M+H] ⁺)	0.637 condition A
182	9		S	409 ([M+H] ⁺)	0.707 condition A











[0129] [Table 1-20]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
183	2		S	508 ([M+H] ⁺)	2.39 condition B
184	2		S	457 ([M+H] ⁺)	2.18 condition B
185	9		S	408 ([M+H] ⁺)	1.162 condition A
186	9		S	418 ([M+H] ⁺)	1.059 condition A
187	9		S	426 ([M+H] ⁺)	1.063 condition A
188	9		S	393 ([M+H] ⁺)	0.864 condition A
189	9		S	409 ([M+H] ⁺)	0.937 condition A
190	9		S	443 ([M+H] ⁺)	0.995 condition A
191	9		S	405 ([M+H] ⁺)	0.852 condition A
192	9		S	403 ([M+H] ⁺)	0.907 condition A

[0130] [Table 1-21]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
193	9		S	449 ([M+H] ⁺)	0.991 condition A
194	9		S	421 ([M+H] ⁺)	0.737 condition A
195	9		S	415 ([M+H] ⁺)	0.934 condition A
196	3		S	480 ([M+H] ⁺)	2.10 condition B
197	3		S	468 ([M+H] ⁺)	2.18 condition B
198	3		S	474 ([M+H] ⁺)	2.35 condition B
199	3		S	429 ([M+H] ⁺)	1.79 condition B
200	3		S	423 ([M+H] ⁺)	2.10 condition B
201	3		S	417 ([M+H] ⁺)	1.86 condition B
202	9		R	461 ([M+H] ⁺)	0.968 condition A











[0131] [Table 1-22]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
203	9		R	427 ([M+H] ⁺)	0.908 condition A
204	9		R	433 ([M+H] ⁺)	0.914 condition A
205	9		R	467 ([M+H] ⁺)	0.802 condition A
206	9		R	433 ([M+H] ⁺)	0.716 condition A
207	9		R	439 ([M+H] ⁺)	0.729 condition A
208	9		R	478 ([M+H] ⁺)	1.082 condition A
209	9		R	470 ([M+H] ⁺)	1.087 condition A
210	9		R	436 ([M+H] ⁺)	1.039 condition A
211	9		R	442 ([M+H] ⁺)	0.860 condition A
212	9		S	448 ([M+H] ⁺)	1.085 condition A











[0132] [Table 1-23]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
213	9		S	414 ([M+H] ⁺)	1.028 condition A
214	9		S	420 ([M+H] ⁺)	0.857 condition A
215	9		R	495 ([M+H] ⁺)	1.024 condition A
216	9		R	461 ([M+H] ⁺)	0.968 condition A
217	9		R	457 ([M+H] ⁺)	0.893 condition A
218	9		R	445 ([M+H] ⁺)	0.903 condition A
219	9		R	467 ([M+H] ⁺)	0.971 condition A
220	9		R	471 ([M+H] ⁺)	0.955 condition A
221	9		R	443 ([M+H] ⁺)	0.704 condition A
222	9		R	437 ([M+H] ⁺)	0.897 condition A










[0133] [Table 1-24]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
223	9		R	466 ([M+H] ⁺)	1.095 condition A
224	9		R	438 ([M+H] ⁺)	0.882 condition A
225	9		R	432 ([M+H] ⁺)	1.044 condition A
226	3		S	441 ([M+H] ⁺)	2.08 condition B
227	3		S	413 ([M+H] ⁺)	1.66 condition B
228	3		S	407 ([M+H] ⁺)	1.99 condition B
229	3		S	401 ([M+H] ⁺)	1.72 condition B
230	9		R	417 ([M+H] ⁺)	0.636 condition A
231	9		R	444 ([M+H] ⁺)	1.030 condition A
232	3		S	491 ([M+H] ⁺)	2.24 condition B

[0134] [Table 1-25]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
233	3		S	463 ([M+H] ⁺)	1.92 condition B
234	3		S	457 ([M+H] ⁺)	2.18 condition B
235	3		S	474 ([M+H] ⁺)	2.34 condition B
236	9		S	453 ([M+H] ⁺)	0.982 condition A
237	9		S	419 ([M+H] ⁺)	0.923 condition A
238	9		S	425 ([M+H] ⁺)	0.716 condition A
239	3		S	446 ([M+H] ⁺)	2.02 condition B
240	9		S	467 ([M+H] ⁺)	1.048 condition A
241	9		S	463 ([M+H] ⁺)	1.069 condition A
242	9		S	443 ([M+H] ⁺)	1.008 condition A

[0135] [Table 1-26]

Compound	Example	Structure	Configuration	(ESI pos.) m/z	RT (min) condition
243	9		S	409 ([M+H] ⁺)	0.951 condition A
244	9		S	457 ([M+H] ⁺)	1.073 condition A
245	9		S	423 ([M+H] ⁺)	1.025 condition A
246	9		S	462 ([M+H] ⁺)	1.159 condition A
247	9		S	467 ([M+H] ⁺)	1.039 condition A
248	9		S	433 ([M+H] ⁺)	0.995 condition A
249	9		S	429 ([M+H] ⁺)	1.008 condition A
250	9		S	439 ([M+H] ⁺)	0.800 condition A
251	9		S	435 ([M+H] ⁺)	0.822 condition A

[0136] The ¹H-NMR data of the following compounds listed below among those shown in Tables 1-9 to 1-26 are shown below:

Compound 73

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.96 (d, J=7.02 Hz, 3 H), 1.02 (d, J=6.61 Hz, 3 H), 2.12 - 2.30 (m, 1 H), 3.84 - 3.97 (m, 2 H), 4.02 - 4.15 (m, 2 H), 4.17 - 4.28 (m, 1 H), 7.20 - 7.27 (m, 1 H), 7.64 - 7.74 (m, 1 H), 8.41 - 8.50 (m, 3 H), 8.97 (br. s., 1 H)

Compound 74

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.92 (d, J=6.61 Hz, 3 H), 1.05 (d, J=7.02 Hz, 3 H), 2.19 - 2.28 (m, 1 H), 3.63 (dd, J=9.08, 6.19 Hz, 1 H), 3.90 - 4.07 (m, 3 H), 4.32 (d, J=16.51 Hz, 1 H), 7.28 - 7.31 (m, 1 H), 8.41 - 8.48 (m, 2 H), 8.68 (br. s., 1 H), 9.18 (s, 2 H)

Compound 93

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=7.02 Hz, 3 H), 1.01 (d, J=7.02 Hz, 3 H), 2.12 - 2.23 (m, 1 H), 3.78 - 3.83 (m, 1 H), 3.84 - 3.88 (m, 1 H), 3.90 (s, 3 H), 4.03 - 4.08 (m, 1 H), 4.10 - 4.18 (m, 2 H), 7.24 - 7.26 (m, 1 H), 8.34 (s, 2 H), 8.41 - 8.47 (m, 2 H), 9.09 (br. s., 1 H)

Compound 94

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.90 (d, J=6.61 Hz, 3 H), 1.01 (d, J=7.02 Hz, 3 H), 1.25 (t, J=7.64 Hz, 3 H), 2.15 - 2.24 (m, 1 H), 2.61 (q, J=7.84 Hz, 2 H), 3.77 - 3.84 (m, 1 H), 3.87 (dd, J=10.94, 6.40 Hz, 1 H), 4.07 (dd, J=10.73, 9.50 Hz, 1 H), 4.10 - 4.17 (m, 2 H), 7.23 - 7.25 (m, 1 H), 8.41 - 8.45 (m, 2 H), 8.48 (s, 2 H), 9.05 (br. s., 1 H)

Compound 103

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.89 (d, J=7.0 Hz, 3 H), 1.03 (d, J=7.0 Hz, 3 H), 1.32 (s, 9 H), 2.13 - 2.26 (m, 1 H), 3.66 (dd, J=8.9, 6.0 Hz, 1 H), 3.83 - 4.01 (m, 2 H), 4.05 - 4.17 (m, 2 H), 6.27 (s, 1 H), 7.68 (d, J=8.7 Hz, 1 H), 8.40 (dd, J=8.7, 2.5 Hz, 1 H), 8.72 (d, J=2.5 Hz, 1 H), 9.32 (br. s., 1 H)

Compound 107

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.98 - 1.04 (m, 2 H), 1.29 (t, J=7.6 Hz, 1 H), 1.72 - 1.85 (m, 1 H), 2.76 - 2.86 (m, 1 H), 3.57 - 3.65 (m, 1 H), 3.91 - 4.04 (m, 1 H), 4.11 (t, J=8.9 Hz, 1 H), 4.18 (d, J=16.1 Hz, 1 H), 7.15 - 7.29 (m, 2 H), 7.64 - 7.71 (m, 1 H),

8.32 - 8.38 (m, 1 H), 8.52 - 8.59 (m, 1 H), 8.73 - 8.77 (m, 1 H)

Compound 108

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.02 (s, 6 H), 1.57 - 1.67 (m, 2 H), 1.75 - 1.84 (m, 1 H), 2.69 (s, 3 H), 3.58 - 3.68 (m, 1 H), 3.91 - 4.00 (m, 2 H), 4.08 - 4.16 (m, 1 H), 4.21 (d, J=16.1 Hz, 1 H), 7.69 (d, J=9.1 Hz, 1 H), 8.05 (d, J=8.7 Hz, 1 H), 8.15 - 8.22 (m, 1 H), 8.29 - 8.35 (m, 1 H), 8.68 - 8.72 (m, 1 H), 8.76 - 8.80 (m, 1 H), 8.83 - 8.87 (m, 1 H)

Compound 112

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.85 - 0.98 (m, 6 H), 1.41 - 1.48 (m, 2 H), 1.62 - 1.76 (m, 1 H), 3.49 - 3.58 (m, 1 H), 3.80 - 3.91 (m, 2 H), 3.98 - 4.05 (m, 1 H), 4.11 (d, J=15.7 Hz, 1 H), 6.36 - 6.64 (m, 1 H), 7.52 (d, J=8.3 Hz, 1 H), 7.59 (d, J=9.1 Hz, 1 H), 8.09 - 8.24 (m, 2 H), 8.53 - 8.58 (m, 1 H), 8.61 (s, 1 H), 8.65 - 8.69 (m, 1 H)

Compound 114

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.94 - 0.97 (m, 4 H), 0.97 - 1.03 (m, 6 H), 1.46 - 1.53 (m, 1 H), 1.69 - 1.83 (m, 2 H), 1.96 - 2.03 (m, 1 H), 3.54 - 3.61 (m, 1 H), 3.90 - 4.00 (m, 2 H), 4.04 - 4.18 (m, 2 H), 7.08 (d, J=8.3 Hz, 1 H), 7.66 (d, J=8.7 Hz, 1 H), 7.86 - 7.93 (m, 1 H), 8.21 (br. s., 1 H), 8.29 - 8.36 (m, 1 H), 8.39 - 8.43 (m, 1 H), 8.71 - 8.76 (m, 1 H)

Compound 148

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.22 (s, 3 H), 1.24 (s, 3 H), 3.24 (s, 3 H), 3.60 (dd, J=9.1, 6.2 Hz, 1 H), 3.96 (dd, J=9.9, 6.2 Hz, 1 H), 4.02 - 4.10 (m, 1 H), 4.26 (d, J=16.5 Hz, 1 H), 4.45 (d, J=16.5 Hz, 1 H), 7.66 (d, J=9.1 Hz, 1 H), 8.41 - 8.47 (m, 2 H), 8.50 (s, 1 H), 8.69 (br. s., 1 H), 8.70 - 8.73 (m, 1 H)

Compound 152

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.11 - 0.24 (m, 2 H), 0.49 - 0.61 (m, 2 H), 0.65 - 0.76 (m, 1 H), 1.63 - 1.75 (m, 2 H), 3.94 - 4.04 (m, 2 H), 4.04 - 4.10 (m, 1 H), 4.14 - 4.20 (m, 1 H), 4.21 - 4.29 (m, 1 H), 7.06 (dd, J=5.4, 1.7 Hz, 1 H), 7.71 (dd, J=9.5, 1.7 Hz, 1 H), 8.17 (d, J=5.4 Hz, 1 H), 8.27 (s, 1 H), 8.48 (s, 1 H), 8.74 (br. s., 1 H)

Compound 155

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 1.12 (t, J=7.4 Hz, 3 H), 1.59 - 1.72 (m, 1 H), 1.74 - 1.88 (m, 1 H), 3.90 (dd, J=8.5, 6.8 Hz, 1 H), 4.01 - 4.17 (m, 2 H), 4.18 - 4.24 (m, 1 H), 4.25 - 4.32 (m, 1 H), 4.62 - 4.78 (m, 1 H), 7.28 (d, J=5.8 Hz, 1 H), 7.67 (d, J=8.7 Hz, 1 H), 8.36 - 8.42 (m, 1 H), 8.44 - 8.50 (m, 2 H), 8.72 - 8.80 (m, 2 H)

Compound 161

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.98 (t, J=7.4 Hz, 3 H), 1.32 - 1.48 (m, 2 H), 1.54 - 1.63 (m, 1 H), 1.80 - 1.89 (m, 1 H), 3.73 - 3.84 (m, 2 H), 4.04 (d, J=16.5 Hz, 1 H), 4.20 - 4.28 (m, 2 H), 7.23 - 7.25 (m, 1 H), 8.40 - 8.44 (m, 1 H), 8.46 (s, 1 H), 8.51 (s, 2 H), 8.86 (br. s., 1 H)

Compound 193

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.68 - 0.72 (m, 2 H), 0.88 (d, J=6.6 Hz, 3 H), 1.00 (d, J=7.0 Hz, 3 H), 1.01 - 1.05 (m, 2 H), 1.79 - 1.86 (m, 1 H), 2.14 - 2.22 (m, 1 H), 3.61 - 3.67 (m, 1 H), 3.77 - 3.88 (m, 2 H), 4.05 (dd, J=10.7, 9.5 Hz, 1 H), 4.13 (s, 2 H), 7.22 - 7.25 (m, 1 H), 8.38 (s, 2 H), 8.40 - 8.45 (m, 2 H), 9.10 (br. s., 1 H)

Compound 195

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.66 - 0.74 (m, 2 H), 0.88 (d, J=7.0 Hz, 3 H), 0.95 - 1.08 (m, 5 H), 1.76 - 1.88 (m, 1 H), 2.10 - 2.23 (m, 1 H), 3.74 - 3.92 (m, 2 H), 3.99 - 4.19 (m, 3 H), 6.98 - 7.06 (m, 1 H), 8.15 (d, J=5.4 Hz, 1 H), 8.21 - 8.28 (m, 1 H), 8.38 (s, 2 H), 8.94 (br. s., 1 H)

Compound 248

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.87 (d, J=6.6 Hz, 3 H), 0.99 (t, J=7.4 Hz, 3 H), 1.25 - 1.32 (m, 1 H), 1.36 - 1.42 (m, 1 H), 1.45 (t, J=7.0 Hz, 3 H), 1.90 - 1.96 (m, 1 H), 3.81 - 3.86 (m, 1 H), 3.87 - 3.92 (m, 1 H), 4.00 - 4.17 (m, 5 H), 7.05 (dd, J=5.4, 2.1 Hz, 1 H), 8.16 (d, J=5.4 Hz, 1 H), 8.25 - 8.28 (m, 1 H), 8.32 (s, 2 H), 8.95 - 8.99 (m, 1 H)

Compound 249

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.68 - 0.73 (m, 2 H), 0.86 (d, J=6.6 Hz, 3 H), 0.99 (t, J=7.4 Hz, 3 H), 1.01 - 1.05 (m, 2 H), 1.22 - 1.33 (m, 1 H), 1.35 - 1.45 (m, 1 H), 1.76 - 1.87 (m, 1 H), 1.89 - 1.98 (m, 1 H), 3.80 - 3.87 (m, 1 H), 3.88 - 3.93 (m, 1 H), 3.99 -

4.18 (m, 3 H), 7.01 - 7.06 (m, 1 H), 8.16 (d, J=5.4 Hz, 1 H), 8.23 - 8.29 (m, 1 H), 8.38 (s, 2 H), 8.95 (br. s., 1 H)

Compound 250

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.84 - 0.91 (m, 5 H), 0.98 (t, J=7.4 Hz, 3 H), 1.09 - 1.14 (m, 2 H), 1.25 - 1.33 (m, 1 H), 1.36 - 1.42 (m, 1 H), 1.44 (t, J=7.0 Hz, 3 H), 1.88 - 1.94 (m, 2 H), 3.83 (dd, J=10.5, 6.4 Hz, 1 H), 3.91 - 3.96 (m, 1 H), 3.96 - 4.00 (m, 1 H), 4.03 - 4.07 (m, 1 H), 4.10 (q, J=6.7 Hz, 2 H), 4.23 - 4.30 (m, 1 H), 6.71 - 6.75 (m, 1 H), 7.93 - 7.98 (m, 1 H), 8.03 - 8.06 (m, 1 H), 8.31 (s, 2 H)

Compound 251

¹H NMR (600 MHz, CHLOROFORM-d) δ ppm 0.57 - 0.63 (m, 2 H), 0.69 - 0.74 (m, 2 H), 0.76 (d, J=6.6 Hz, 3 H), 0.87 (t, J=7.4 Hz, 3 H), 0.90 - 0.95 (m, 2 H), 0.95 - 1.00 (m, 2 H), 1.14 - 1.23 (m, 1 H), 1.25 - 1.34 (m, 1 H), 1.69 - 1.76 (m, 1 H), 1.76 - 1.85 (m, 2 H), 3.72 (dd, J=10.7, 6.6 Hz, 1 H), 3.79 - 3.85 (m, 1 H), 3.87 (d, J=16.5 Hz, 1 H), 3.90 - 3.96 (m, 1 H), 4.14 (d, J=16.5 Hz, 1 H), 6.60 (dd, J=5.4, 1.2 Hz, 1 H), 7.74 - 7.80 (m, 1 H), 7.97 (d, J=5.4 Hz, 1 H), 8.28 (s, 2 H), 8.53 (br. s., 1 H)

Test Example 1 [Glycine Uptake Inhibition Experiment]

A glycine uptake experiment was conducted in accordance with the method published in Neuron, 8, 927-935, 1992. In the experiment, T98G cells (glioma cells) expressing human type 1 glycine transporter (GlyT1) were used. The T98G cells were seeded in a 96-well plate at 2.0×10^4 cells/well and cultured overnight in a CO₂ incubator. The test substance was dissolved in a 100% DMSO solution and then dissolved in a 10 mM HEPES buffer solution (pH 7.4) containing 150 mM sodium chloride, 1 mM calcium chloride, 5 mM potassium chloride, 1 mM magnesium chloride, 10 mM glucose and 0.2% bovine serum albumin. After removing the cell culture medium, the test substance was subjected to a 10-min pretreatment. Subsequently, the test substance and [³H] glycine (final concentration: 250 nM) were added to the cells and reaction was performed at room temperature for 15 minutes. After the end of the reaction, the extracellular fluid was aspirated with a manifold to remove excess labeled glycine present outside the cells, and then

the cells were lysed with a 0.5 M aqueous sodium hydroxide solution. The glycine content in the cells was determined by measuring the radioactivity in the cell lysate with a liquid scintillation counter. Glycine uptake in the presence of 10 μ M ALX5407 was defined as non-specific uptake, and the value calculated by subtracting the amount of the non-specific uptake from the total uptake in the absence of 10 μ M ALX5407 was defined as specific uptake. In addition, glycine uptake inhibitory activity (IC_{50} value) was calculated from an inhibition curve at the concentrations of each test substance ranging from 10^{-9} to 10^{-5} M.

[0137] It should be noted that ALX5407 is an HCl salt of N-[(3R)-3-([1,1'-biphenyl]-4-yloxy)-3-(4-fluorophenyl)propyl]-N-methylglycine.

[0138] All the compounds of the Working Examples in the present invention were found to have IC_{50} values of less than 10 μ M. Specific examples of their IC_{50} values are as follows: Compound 8, 0.66 μ M; Compound 11, 0.089 μ M; Compound 12, 0.071 μ M; Compound 13, 0.074 μ M; Compound 54, 0.80 μ M; Compound 61, 0.053 μ M; Compound 62, 0.033 μ M; Compound 93, 0.054 μ M; Compound 98, 0.26 μ M; Compound 103, 0.23 μ M; Compound 104, 0.24 μ M; Compound 148, 0.075 μ M; Compound 152, 0.043 μ M; Compound 165, 0.15 μ M; Compound 186, 0.045 μ M; Compound 188, 0.60 μ M; Compound 193, 0.022 μ M; Compound 220, 0.017 μ M; Compound 248, 0.016 μ M; Compound 249, 0.018 μ M; and Compound 250, 0.024 μ M.


INDUSTRIAL APPLICABILITY

[0139] The inventive compounds have glycine transporter (GlyT1)-inhibiting activity, and thus, are effective in the prevention or treatment of diseases associated with the glycine transporter which are, specifically, schizophrenia, Alzheimer's disease, cognitive impairment, dementia, anxiety disorders (e.g., generalized anxiety disorder, panic disorder, obsessive-compulsive disorder, social anxiety disorder, post-traumatic stress disorder, specific phobias, acute stress disorder), depression, drug dependence, spasm, tremor, pain, Parkinson's disease, attention deficit hyperactivity disorder, bipolar disorder, eating disorder, sleep disorders or the like.

CLAIMS

1. A compound of formula [I] or a pharmaceutically acceptable salt thereof:

[Formula 1]



wherein

R^1 and $R^{1'}$ are the same or different, and each represent a hydrogen atom, a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a cyano group, a heteroaryl group (which may be substituted by a C_{1-6} alkyl group), a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a C_{1-6} alkylamino group, or the formula $CONR^7R^8$ (R^7 and R^8 are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group),

R^2 represents a hydrogen atom or a C_{1-6} alkyl group,

R^3 represents a phenyl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, cyano groups, C_{1-6} alkyl groups, C_{1-6} alkoxy groups, C_{1-6} alkylamino groups, C_{1-6} alkylsulfonyl groups, halo C_{1-6} alkyl groups, halo C_{1-6} alkoxy groups, halo C_{1-6} alkylsulfanyl groups, phenyl groups, phenoxy groups, heteroaryl groups (which may be substituted by a C_{1-6} alkyl group), and the formula $-SO_2NR^9R^{10}$ (R^9 and R^{10} are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group)) or a heteroaryl group or a bicyclic heteroaryl group (the each heteroaryl group may be substituted by 1 to 3 substituents selected from halogen atoms, C_{1-6} alkyl groups, C_{3-6} cycloalkyl groups, C_{1-6} alkoxy groups, cyano groups, C_{1-6} alkanoyl groups, and halo C_{1-6} alkyl groups),

R^4 represents a C_{1-6} alkyl group (which may be substituted by 1 to 3 halogen atoms, a C_{1-6} alkoxy group, a C_{3-6} cycloalkyl group, or a phenyl group), a C_{3-6} cycloalkyl group, or a phenyl group,

R^5 and R^6 are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group, and

A^1 , A^2 , A^3 , and A^4 are the same or different, and each represent the formula CH or a nitrogen atom, provided that one or two of A^1 , A^2 , A^3 , and A^4 represent a nitrogen atom.

2. The compound or pharmaceutically acceptable salt thereof according to claim 1, wherein

R^1 is a hydrogen atom, a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a cyano group, a heteroaryl group (which may be substituted by a C_{1-6} alkyl group), a C_{1-6} alkyl group, a C_{1-6} alkylamino group, or the formula $CONR^7R^8$ (R^7 and R^8 are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group),

$R^{1'}$ is a hydrogen atom,

R^3 is a phenyl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, cyano groups, C_{1-6} alkyl groups, C_{1-6} alkoxy groups, C_{1-6} alkylamino groups, C_{1-6} alkylsulfonyl groups, halo C_{1-6} alkyl groups, halo C_{1-6} alkoxy groups, halo C_{1-6} alkylsulfanyl groups, phenyl groups, phenoxy groups, heteroaryl groups (which may be substituted by a C_{1-6} alkyl group), and the formula $-SO_2NR^9R^{10}$ (R^9 and R^{10} are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group)) or a heteroaryl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, C_{1-6} alkyl groups, C_{1-6} alkoxy groups, cyano groups, and halo C_{1-6} alkyl groups), and

R^4 is a C_{1-6} alkyl group (which may be substituted by a C_{3-6} cycloalkyl group or a phenyl group) or a phenyl group.

3. The compound or pharmaceutically acceptable salt thereof according to claim 1, wherein R^4 is a C_{1-6} alkyl group which may be substituted by 1 to 3 halogen atoms.

4. The compound or pharmaceutically acceptable salt thereof according to claim 1 or 2, wherein R^4 is a C_{1-6} alkyl group.

5. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 4, wherein

R^2 is a hydrogen atom, and

R^5 and R^6 are both a hydrogen atom.

6. The compound or pharmaceutically acceptable salt thereof according to claim 5,

wherein

R^1 is a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a cyano group, a heteroaryl group (which may be substituted by a C_{1-6} alkyl group), a C_{1-6} alkyl group, a C_{1-6} alkylamino group, or the formula $CONR^7R^8$ (R^7 and R^8 are the same or different, and each represent a hydrogen atom or a C_{1-6} alkyl group), and

$R^{1'}$ is a hydrogen atom.

7. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 and 3 to 5, wherein

R^1 is a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a C_{1-6} alkyl group, or a C_{3-6} cycloalkyl group, and

$R^{1'}$ is a hydrogen atom, a halogen atom, a C_{1-6} alkoxy group, a halo C_{1-6} alkyl group, a C_{1-6} alkyl group, or a C_{3-6} cycloalkyl group.

8. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 7, wherein R^1 is attached in the para position.

9. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 8, wherein any one of A^1 , A^2 , A^3 and A^4 is a nitrogen atom or A^1 and A^3 are both a nitrogen atom.

10. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 8, wherein

A^1 is a nitrogen atom,

A^2 and A^4 are both the formula CH, and

A^3 is the formula CH or a nitrogen atom.

11. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 and 3 to 10, wherein R^3 is a heteroaryl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, C_{1-6} alkyl groups, C_{3-6} cycloalkyl groups, C_{1-6} alkoxy groups, cyano groups, C_{1-6} alkanoyl groups, and halo C_{1-6} alkyl groups).

12. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 and 3 to 10, wherein R^3 is a pyridyl group (which may be substituted by 1 to 3

substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₃₋₆ cycloalkyl groups, C₁₋₆ alkoxy groups, cyano groups, C₁₋₆ alkanoyl groups, and haloC₁₋₆ alkyl groups).

13. The compound or pharmaceutically acceptable salt thereof according to any one of claims 1 and 3 to 10, wherein R³ is a pyridyl group (which may be substituted by 1 to 3 substituents selected from halogen atoms, C₁₋₆ alkyl groups, C₃₋₆ cycloalkyl groups, and haloC₁₋₆ alkyl groups).

14. The compound or pharmaceutically acceptable salt thereof according to claim 1, wherein the compound is selected from the group consisting of:

2-[(5S)-3-(5-methoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-3-(5-ethylpyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]acetamide,

2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]-N-(4-ethylpyridin-2-yl)acetamide,

2-[(5S)-3-(5-fluoropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-3-(5-fluoropyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-cyclopropylpyridin-2-yl)-2-[(5S)-2-oxo-5-(propan-2-yl)-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-{(5S)-2-oxo-5-(propan-2-yl)-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}acetamide,

2-[(5S)-3-(5-cyclopropylpyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-3-(5-cyclopropylpyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-cyclopropylpyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-chloropyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-chloropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-ethylpyridin-2-yl)acetamide,

N-(4-chloropyridin-2-yl)-2-{(5R)-5-[(1S)-1-fluoropropyl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}acetamide,

N-(4-chloropyridin-2-yl)-2-{(5R)-3-(5-chloropyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl}acetamide,

N-(4-cyclopropylpyridin-2-yl)-2-{(5R)-5-[(1S)-1-fluoropropyl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}acetamide,

2-{(5R)-5-[(1S)-1-fluoropropyl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl}-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-{(5R)-3-(5-chloropyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl}-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5R)-5-[(1S)-1-fluoropropyl]-3-(5-fluoropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-{(5R)-3-(5-cyclopropylpyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl}-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-{(5R)-3-(5-ethoxypyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl}-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-cyclopropylpyridin-2-yl)-2-[(5R)-3-(5-ethoxypyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5R)-3-(5-ethoxypyrimidin-2-yl)-5-[(1S)-1-fluoropropyl]-2-oxoimidazolidin-1-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-fluoropyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]-N-(4-cyclopropylpyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-2-oxo-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]-N-(4-chloropyridin-2-yl)acetamide,

2-[(5S)-3-(5-ethoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(4-chloropyridin-2-yl)-2-[(5S)-3-(5-ethoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide,

N-(4-cyclopropylpyridin-2-yl)-2-[(5S)-3-(5-ethoxypyrimidin-2-yl)-2-oxo-5-(propan-2-yl)imidazolidin-1-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-ethoxypyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-cyclopropylpyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

N-(5-chloropyridin-2-yl)-2-[(5S)-2-oxo-5-propyl-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]acetamide,

N-(5-chloro-6-methylpyridin-2-yl)-2-[(5S)-2-oxo-5-propyl-3-[5-(trifluoromethyl)pyrimidin-2-yl]imidazolidin-1-yl]acetamide,

N-(5-chloro-6-methylpyridin-2-yl)-2-[(5S)-3-(5-chloropyrimidin-2-yl)-2-oxo-5-propylimidazolidin-1-yl]acetamide,

2-[(5S)-2-oxo-5-(propan-2-yl)-3-[5-(propan-2-yloxy)pyrimidin-2-yl]imidazolidin-1-yl]-N-[4-(trifluoromethyl)pyridin-2-yl]acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-ethoxypyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-chloropyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-cyclopropylpyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-chloropyridin-2-yl)acetamide,

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-ethoxypyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-cyclopropylpyridin-2-yl)acetamide, and

2-[(5S)-5-[(2S)-butan-2-yl]-3-(5-cyclopropylpyrimidin-2-yl)-2-oxoimidazolidin-1-yl]-N-(4-cyclopropylpyridin-2-yl)acetamide.

15. A pharmaceutical composition comprising, as an active ingredient, the compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 14.

16. An agent for preventing or treating diseases of schizophrenia, Alzheimer's disease, cognitive impairment, dementia, anxiety disorders, depression, drug dependence, spasm, tremor, pain, Parkinson's disease, attention deficit hyperactivity disorder, bipolar disorder, eating disorder, or sleep disorders, which comprises, as an active ingredient, the compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 14.