

NOVEL SUBSTITUTED SULFAMIDE DERIVATIVES

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U. S. Provisional Application
5 61/082,654, filed on July 22, 2008, which is incorporated by reference herein in
its entirety.

FIELD OF THE INVENTION

The present invention is directed to novel substituted sulfamide
10 derivatives, pharmaceutical compositions containing said derivatives and the
use of said derivatives form in the treatment of anxiety and related disorders;
bipolar depression and mania; depression; epilepsy and related disorders;
epileptogenesis; glucose related disorders; lipid related disorders; migraine;
obesity; pain; substance abuse and as neuroprotective agents. The present
15 invention is further directed to a process for the preparation of the novel
substituted sulfamide derivatives.


BACKGROUND OF THE INVENTION

US Patent Publication US2006-0041008 A1, published February 23,
20 2006 discloses benzo-fused sulfamide derivatives useful for the treatment of
epilepsy and related disorders; US Patent Publication 2007-0293441 A1,
published November 18, 2008 discloses co-therapy for the treatment of
epilepsy and related disorder comprising administration of benzo-fused
ssulfamide derivatives and on or more anticonvulsants and / or anti-epileptic
25 agents; US Patent Publication US2007-0155826 A1, published July 5, 2007
discloses the use of benzo-fused sulfamide derivatives for the treatment of
bipolar disorder and mania; US Patent Publication US2007-0155827 A1,
published July 5, 2007 discloses the use of benzo-fused sulfamide derivatives
for the treatment of depression; US Patent Publication US2007-0155824 A1,
30 published July 5, 2007 discloses the use of benzo-fused sulfamide derivatives
for the treatment of epileptogenesis; US Patent Publication US2007-0155821
A1, published July 5, 2007 discloses the use of benzo-fused sulfamide
derivatives for the treatment of glucose related disorders and for the treatment

of lipid related disorders; US Patent Publication US2007-0191474 A1,
published August 16, 2007 discloses the use of benzo-fused sulfamide
derivatives for the treatment of migraine; US Patent Publication US2007-
015823 A1, published July 5, 2007 discloses the use of benzo-fused sulfamide
5 derivatives for neuroprotection; US Patent Publication US2008-0027131 A1,
published January 31, 2008 discloses the use of benzo-fused sulfamide
derivatives for the treatment of obesity; US Patent Publication US2007-
0155822 A1, published July 5, 2007 discloses the use of benzo-fused
sulfamide derivatives for the treatment of pain; US Patent Publication US2007-
10 0155825 A1, published July 5, 2007 discloses the use of benzo-fused
sulfamide derivatives for the treatment of substance abuse and / or addiction;
which are herein incorporated by reference in their entirety.

SUMMARY OF THE INVENTION

15 The present invention is directed to compounds of formula (I)




wherein

R^1 is selected from the group consisting of C_{1-4} alkyl and -O-phenyl;
wherein the phenyl is optionally substituted with one to three substituents,
20 independently selected from the group consisting of halogen, C_{1-4} alkyl,
fluorinated C_{1-4} alkyl, C_{1-4} alkoxy, fluorinated C_{1-4} alkoxy, cyano, nitro, $-NR^A R^B$
and $-C(O)-NR^C R^D$; wherein R^A and R^B are each independently selected from
the group consisting of C_{1-4} alkyl; and wherein R^C and R^D are each
independently selected from the group consisting of hydrogen and C_{1-4} alkyl;
25 and pharmaceutically acceptable salts thereof;

provided that when R^1 is methyl, then the compound of formula (I) is
present as an isolated or substantially pure form.

The present invention is further directed to an acetyl-substituted
sulfamide derivative, a compound of formula (A)



and pharmaceutically acceptable salts thereof; also known as (2S)-(-)-N'-acetyl-N-(6-chloro-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl)-sulfamide; wherein the compound of formula (A) is present as an isolated or substantially pure form; preferably the compound of formula (A) is present as an isolated and substantially pure form.

The present invention is further directed to a process for the preparation of compounds of formula (I), as described in more detail herein. In an embodiment, the present invention is directed to a process for the preparation of a compound of formula (A), as described in more detail herein. The present invention is further directed to a product prepared according to any of the processes described herein.

Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of formula (I), preferably a compound of formula (A). An illustration of the invention is a pharmaceutical composition made by mixing a compound of formula (I), preferably a compound of formula (A) and a pharmaceutically acceptable carrier. Illustrating the invention is a process for making a pharmaceutical composition comprising mixing a compound of formula (I), preferably a compound of formula (A) and a pharmaceutically acceptable carrier.

The present invention is further directed to methods for the treatment of anxiety and related disorders; bipolar depression and mania; depression; epilepsy and related disorders; epileptogenesis; glucose related disorders; lipid related disorders; migraine; obesity; pain; substance abuse or neuroprotection comprising administering to a subject in need thereof a therapeutically effective amount of the compound of formula (I), preferably a compound of formula (A).


Another example of the invention is the use of the compound of formula (I), preferably a compound of formula (A) in the preparation of a medicament for treating: (a) anxiety and related disorders; (b) bipolar depression (c) mania; (d) depression; (e) epilepsy and related disorders; (f) epileptogenesis; (g)

glucose related disorders; (h) lipid related disorders; (i) migraine; (j) obesity; (k) pain; (l) substance abuse and (m) for neuroprotection, in a subject in need thereof.

5


DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to compounds of formula (I)




wherein R¹ is as herein defined; and pharmaceutically acceptable salts thereof; useful as prodrugs of the compound of formula (S)

10




also known as (2S)-(-)-N-(6-chloro-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl)-sulfamide (disclosed in US Patent Publication US2006-0041008 A1, published February 23, 2006). In an embodiment, the present invention is directed to an acetyl-substituted sulfamide derivative, a compound of formula

15 (A)



and pharmaceutically acceptable salts thereof. The compound of formula (A) is a human metabolite of the compound of formula (S)



The compound of formula (A) was initially isolated from human plasma samples from clinical trials dosing with the compound of formula (S).

Additionally, the compound of formula (A) is a pro-drug of the compound of formula (S). In mice studies dosing with the compound of formula (A), plasma
5 and brain analyses showed the presence of both the compound of formula (A) and the compound of formula (S), with the levels of the compound of formula (S) higher than those of the compound of formula (A).

The present invention is further directed to the use of compounds of formula (I), preferably the compound of formula (A), for the treatment of anxiety
10 and related disorders; bipolar depression and mania; depression; epilepsy and related disorders; epileptogenesis; glucose related disorders; lipid related disorders; migraine; obesity; pain; substance abuse and for neuroprotection.

The present invention is further directed to pharmaceutical compositions comprising a compound of formula (I), preferably a compound of formula (A).

15 In an embodiment of the present invention, R^1 is selected from the group consisting of C_{1-4} alkyl. In another embodiment of the present invention, R^1 is selected from the group consisting of C_{2-4} alkyl and $-O$ -phenyl. In another embodiment of the present invention, R^1 is selected from the group consisting
20 of C_{2-4} alkyl. In another embodiment of the present invention, R^1 is selected from the group consisting of methyl, ethyl and t-butyl. In yet another embodiment of the present invention, R^1 is methyl.

In an embodiment of the present invention, R^1 is selected from the group consisting of $-O$ -phenyl; wherein the phenyl is optionally substituted with one to
25 three substituents (preferably one to two substituents, more preferably one substituent), independently selected from the group consisting of halogen, C_{1-4} alkyl, fluorinated C_{1-4} alkyl, C_{1-4} alkoxy, fluorinated C_{1-4} alkoxy, cyano, nitro, $-NR^A R^B$ and $-C(O)-NR^C R^D$; wherein R^A and R^B are each independently selected from the group consisting of C_{1-4} alkyl; and wherein R^C and R^D are each
30 independently selected from the group consisting of hydrogen and C_{1-4} alkyl. In another embodiment of the present invention, R^1 is selected from the group consisting of $-O$ -phenyl; wherein the phenyl is optionally substituted with a

substituent selected from the group consisting of halogen, C₁₋₄alkyl, -CF₃, C₁₋₄alkoxy, -OCF₃, cyano, nitro, -NR^AR^B and -C(O)-NR^CR^D.

In an embodiment of the present invention, R^A and R^B are each independently selected from the group consisting of methyl, ethyl and t-butyl.

- 5 In another embodiment of the present invention, R^C and R^D are each independently selected from the group consisting of hydrogen, methyl, ethyl and t-butyl.

- 10 As used herein, "**halogen**" shall mean chlorine, bromine, fluorine and iodine. Preferably, the halogen is chlorine or fluorine.

As used herein, the term "**C₁₋₄alkyl**" whether used alone or as part of a substituent group, shall include straight and branched chains containing one to four carbon atoms. For example, alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and t-butyl.

- 15 As used herein, unless otherwise noted, the term "**fluorinated C₁₋₄alkyl**" shall mean any C₁₋₄alkyl group as defined above substituted with at least one fluoro atom. Suitable examples include but are not limited to -CH₂F, -CF₃, -CH₂-CF₃, -CF₂-CF₂-CF₂-CF₃, and the like.

- 20 As used herein, unless otherwise noted, "**C₁₋₄alkoxy**" shall denote an oxygen ether radical of any the above described straight or branched chain alkyl groups containing one to four carbon atoms. For example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec-butoxy and, t-butoxy.

- 25 As used herein, unless otherwise noted, the term "**fluorinated C₁₋₄alkoxy**" shall mean any C₁₋₄alkoxy group as defined above substituted with at least one fluoro atom. Suitable examples include but are not limited to -OCH₂F, -OCF₃, -OCH₂-CF₃, -OCF₂-CF₂-CF₂-CF₃, and the like.

- 30 Where the compound according to this invention has at least one **chiral center**, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Preferably, wherein the compound is present as an enantiomer, the enantiomer is present

at an enantiomeric excess of greater than or equal to about 80%, more preferably, at an enantiomeric excess of greater than or equal to about 90%, more preferably still, at an enantiomeric excess of greater than or equal to about 95%, more preferably still, at an enantiomeric excess of greater than or equal to about 98%, most preferably, at an enantiomeric excess of greater than or equal to about 99%.

As used herein, unless otherwise noted, the term “**isolated form**” shall mean that the compound is present in a form which is separate from any biological environment (e.g. plasma, blood, gastric fluids, urine, cerebrospinal fluid, and the like). In an embodiment of the present invention, the compound of formula (I), preferably, the compound of formula (A) is present as an isolated form.

As used herein, unless otherwise noted, the term “**substantially pure form**” shall mean that the mole percent of impurities in the isolated compound is less than about 5 mole percent, preferably less than about 2 mole percent, more preferably, less than about 0.5 mole percent, most preferably, less than about 0.1 mole percent. In an embodiment of the present invention, the compound of formula (I), preferably, the compound of formula (A) is a substantially pure form.

As used herein, unless otherwise noted, the term “**substantially free of corresponding salt form(s)**” when used to described the compound of formula (I), preferably a compound of formula (A) shall mean that mole percent of salt forms of the compound of formula (I), preferably a compound of formula (A) is less than about 5 mole percent, preferably less than about 2 mole percent, more preferably, less than about 0.5 mole percent, most preferably less than about 0.1 mole percent. In an embodiment of the present invention, the compound of formula (I), preferably, the compound of formula (A) is substantially free of corresponding salt form(s).

As used herein, unless otherwise noted, the terms “**treating**”, “**treatment**” and the like, shall include the management and care of a subject or patient (preferably mammal, more preferably human) for the purpose of combating a disease, condition, or disorder and includes the administration of a

compound of the present invention to prevent the onset of the symptoms or complications, alleviate the symptoms or complications, or eliminate the disease, condition, or disorder.

As used herein, unless otherwise noted, the term “**prevention**” shall
5 include (a) reduction in the frequency of one or more symptoms; (b) reduction in the severity of one or more symptoms; (c) the delay or avoidance of the development of additional symptoms; and / or (d) delay or avoidance of the development of the disorder or condition.

One skilled in the art will recognize that wherein the present invention is
10 directed to methods of prevention, a subject in need of thereof (i.e. a subject in need of prevention) shall include any subject or patient (preferably a mammal, more preferably a human) who has experienced or exhibited at least one symptom of the disorder, disease or condition to be prevented. Further, a subject in need thereof may additionally be a subject (preferably a mammal,
15 more preferably a human) who has not exhibited any symptoms of the disorder, disease or condition to be prevented, but who has been deemed by a physician, clinician or other medical profession to be at risk of developing said disorder, disease or condition. For example, the subject may be deemed at risk of developing a disorder, disease or condition (and therefore in need of
20 prevention or preventive treatment) as a consequence of the subject's medical history, including, but not limited to, family history, pre-disposition, co-existing (comorbid) disorders or conditions, genetic testing, and the like.

As used herein, the terms “**anxiety and related disorders**” and “**anxiety or a related disorder**” shall be defined to include anxiety and related
25 disorders including generalized anxiety disorder, acute stress disorder, post traumatic stress disorder, obsessive-compulsive disorder, social phobia (also known as social anxiety disorder), specific phobia, panic disorder with or without agoraphobia, agoraphobia without a history of panic disorder, anxiety disorder due to general medical condition, substance abuse induced anxiety
30 disorder and anxiety disorder not otherwise specified (as these conditions are described by their diagnostic criteria, as listed in the *Diagnostic and Statistical Manual of Mental Disorders*, 4th Edition, Text Revision, American Psychiatric Association, **2000**, incorporated herein by reference). Preferably, the anxiety or

related disorder is selected from the group consisting of generalized anxiety disorder, acute stress disorder, post traumatic stress disorder and obsessive-compulsive disorder. More preferably, the anxiety and related disorder is generalized anxiety disorder.

5 Bipolar disorder is psychiatric disorder characterized by unpredictable swings in mood from mania (or hypomania) to depression. As used herein, the term “**bipolar disorder**” shall include bipolar disorder I, bipolar disorder II, cyclothymic disorder and bipolar disorder not otherwise specified. Preferably, the bipolar disorder is characterized by depressive and manic (or hypomanic)
10 phases, wherein the phases cycle. Preferably, the bipolar disorder is bipolar disorder I or bipolar disorder II.

 As used herein, the term “**bipolar depression**” is intended to mean the depression associated with, characteristic of or symptomatic of a bipolar disorder. Thus, methods of treating bipolar depression of the present invention
15 are directed to methods which treat the depression and / or depressed phase of bipolar disorders.

 As used herein, unless otherwise noted the terms “**cycling**” or “**bipolar cycling**” shall refer to the alternation of mood between depressive and manic phases characteristic of bipolar disorders. Thus, the present invention includes
20 methods for the stabilization of said cycling, including, but not limited to, decreasing the frequency of the cycling and / or decreasing the magnitude of the manic and / or depressive phases.

 As used herein, the term “**mania**” shall include mania or a manic mood phase, regardless of underlying cause. As used herein, the term “**bipolar**
25 **mania**” is intended to mean the mania associated with, characteristic of or symptomatic of a bipolar disorder. Thus, methods of treating bipolar mania of the present invention are directed to methods which treat the mania and / or manic phase of bipolar disorders.

 As used herein, the term “**depression**” shall be defined to include major
30 depressive disorder (including single episode and recurrent), unipolar depression, treatment-refractory depression, resistant depression, anxious depression and dysthymia (also referred to as dysthymic disorder). Further, the term “depression” shall encompass any major depressive disorder,

dysthymic disorder and depressive disorder not otherwise specific as defined by their diagnostic criteria, as listed in the *Diagnostic and Statistical Manual of Mental Disorders*, 4th Edition, Text Revision, American Psychiatric Association, **2000**. Preferably, the depression is major depressive disorder, unipolar

5 depression, treatment-refractory depression, resistant depression or anxious depression. More preferably, the depression is major depressive disorder.

As used herein, unless otherwise noted, the terms “**epilepsy and related disorders**” or “**epilepsy or related disorder**” shall mean any disorder in which a subject (preferably a human adult, child or infant) experiences one or more

10 seizures and / or tremors. Suitable examples include, but are not limited to, epilepsy (including, but not limited to, localization-related epilepsies, generalized epilepsies, epilepsies with both generalized and local seizures, and the like), seizures as a complication of a disease or condition (such as seizures associated with encephalopathy, phenylketonuria, juvenile Gaucher’s disease, Lundborg’s

15 progressive myoclonic epilepsy, stroke, head trauma, stress, hormonal changes, drug use or withdrawal, alcohol use or withdrawal, sleep deprivation, and the like), essential tremor, restless limb syndrome, and the like. Preferably, the disorder is selected from epilepsy (regardless of type, underlying cause or origin), essential tremor or restless limb syndrome, more preferably, the disorder is epilepsy

20 (regardless of type, underlying cause or origin) or essential tremor.

As used herein, the term “**epileptogenesis**” shall mean the biochemical, genetic, histological or other structural or functional processes or changes that make nervous tissue, including the central nervous system (CNS) susceptible to recurrent, spontaneous seizures. In addition, the term “epileptogenesis” is

25 also used herein in a broader sense to refer to the changes and / or processes that contribute to the clinical progression observed in patients with epilepsy or other seizure disorder or an analogous seizure-related disorder including but not limited to; the worsening or progression of the disorder and it’s symptoms or the development of “pharmacoresistance,” in which the disorder becomes

30 more difficult to treat as a result of neurobiological changes which result in reduced drug sensitivity or the recruitment by the process of epileptogenesis of non seizure prone nervous tissue. Furthermore the term “epileptogenesis” is used herein in the broadest possible sense to refer to the similar phenomena of

progressive worsening over time of the signs and symptoms of apparently non-epileptic disorders, including psychiatric disorders the etiology of which appear to be seizure related.

Epileptogenesis is a Two Phase Process: "Phase 1 epileptogenesis" is the initiation of the epileptogenic process prior to the first epileptic seizure or symptom of an analogous seizure-related disorder, and is often the result of some kind of injury or trauma to the brain, i.e., stroke, disease (e.g., infection such as meningitis), or trauma, such as an accidental blow to the head or a surgical procedure performed on the brain. "Phase 2 epileptogenesis" refers to the process during which brain tissue that is already susceptible to epileptic seizures or seizure related phenomena of an analogous seizure-related disorder, becomes still more susceptible to seizures of increasing frequency and/or severity and/or becomes less responsive to treatment.

As used herein, the term "**glucose related disorder**" shall be defined as any disorder which is characterized by elevated glucose levels. Glucose related disorders include elevated glucose level, pre-diabetes, impaired oral glucose tolerance, poor glycemic control, Type II Diabetes Mellitus, Syndrome X (also known as metabolic syndrome), gestational diabetes, insulin resistance, hyperglycemia and loss of muscle mass as a results of hyperglycemia (cachexia).

Treatment of glucose related disorders may comprise lowering glucose levels, improving glycemic control, decreasing insulin resistance and / or preventing the development of a glucose related disorder (for example preventing a patient suffering from impaired oral glucose tolerance or elevated glucose levels from developing Type II diabetes mellitus).

As used herein, the term "**lipid related disorder**" shall be defined as any disorder which is characterized by non-normal lipid levels. Lipid related disorders include elevated triglyceride levels, low HDL cholesterol and dyslipidemia, preferably elevated triglyceride levels or low HDL cholesterol levels Treatment of lipid related disorder may comprise lowering triglycerides, elevating HDL cholesterol and / or improving the triglyceride/HDL ratio.

As used herein, the term "**migraine**" shall mean a chronic, episodic and debilitating clinical condition that is diagnosed by the presence of moderate to

severe pulsating unilateral headaches lasting between 4 and 72 h, which includes migraine without aura and migraine with aura.

As used herein, “**migraine without aura**” shall mean at least five attacks fulfilling the following criteria: (a) the headache attack lasts 4-72 hours with the
5 headache having at least two of the following features: unilateral location, pulsating quality, moderate or severe intensity with direct influence on activities of daily living, and aggravation by walking up stairs or similar routines; and (b) during the headache at least one of the following occurs: nausea and/or vomiting, and photophobia and phonophobia.

10 As used herein, “**migraine with aura**” shall mean at least two attacks accompanied by at least 3 of the 4 following features: (a) one or more fully reversible aura symptoms; (b) at least one aura symptom which develops gradually over more than four minutes or two or more symptoms which occur in succession; (c) no aura symptom which lasts more than 60 minutes; (d) a
15 headache occurs prior to, simultaneously with or following the aura, with a free interval between aura and headache of less than about 60 minutes.

As used herein, the term “**prevention**” shall include the prevention of migraine attacks (headaches), a decrease in the frequency of migraine attacks (headaches), a decrease in the severity of migraine attacks (headaches) and/or
20 a decrease in the duration of migraine attacks (headaches).

As used herein, the term “**obesity**” shall be defined as a body mass index (BMI) of greater than or equal to about 25, preferably a BMI of greater than or equal to about 30. Thus as used herein, the term “**obesity**” shall include both overweight and clinically obese subjects / patients.

25 As used herein, the term “**pain**” shall be defined to include acute, chronic, inflammatory and neuropathic pain (preferably diabetic neuropathy). Further, the pain may be centrally mediated, peripherally mediated, caused by structural tissue injury, caused by soft tissue injury or caused by progressive disease. Any centrally mediated, peripherally mediated, structural tissue injury,
30 soft tissue injury or progressive disease related pain may be acute or chronic.

As used herein, unless otherwise noted, pain shall include inflammatory pain, centrally mediated pain, peripherally mediated pain, visceral pain, structural related pain, cancer pain, soft tissue injury related pain, progressive

disease related pain, neuropathic pain, acute pain from acute injury, acute pain from trauma, acute pain from surgery, headache, dental pain, back pain (preferably lower back pain), chronic pain from neuropathic conditions and chronic pain from post-stroke conditions.

- 5 In an embodiment of the present invention, is a method for the treatment of pain, wherein the pain is acute pain. In another embodiment of the present invention, is a method for the treatment of pain, wherein the pain is chronic pain. In another embodiment of the present invention, is a method for the treatment of pain, wherein the pain is neuropathic pain, more preferably
- 10 diabetic neuropathy. In yet another embodiment of the present invention, is a method for the treatment of pain, wherein the pain is inflammatory pain.

- In an embodiment, the pain is selected from the group consisting of osteoarthritis, rheumatoid arthritis, fibromyalgia, headache, toothache, burn, sunburn, animal bite (such as dog bite, cat bite, snake bite, spider bite, insect
- 15 sting, and the like), neurogenic bladder, benign prostatic hypertrophy, interstitial cystitis, rhinitis, contact dermatitis/hypersensitivity, itch, eczema, pharyngitis, mucositis, enteritis, cellulites, causalgia, sciatic neuritis, mandibular joint neuralgia, peripheral neuritis, polyneuritis, stump pain, phantom limb pain, post-operative ileus, cholecystitis, postmastectomy pain syndrome, oral neuropathic
- 20 pain, Charcot's pain, reflex sympathetic dystrophy, Guillain-Barre syndrome, meralgia paresthetica, burning-mouth syndrome, post-herpetic neuralgia, trigeminal neuralgia, peripheral neuropathy, bilateral peripheral neuropathy, diabetic neuropathy, postherpetic neuralgia, trigeminal neuralgia, optic neuritis, postfebrile neuritis, migrating neuritis, segmental neuritis, Gombault's neuritis,
- 25 neuronitis, cervicobrachial neuralgia, cranial neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, migrainous neuralgia, idiopathic neuralgia, intercostals neuralgia, mammary neuralgia, Morton's neuralgia, nasociliary neuralgia, occipital neuralgia, red neuralgia, Sluder's neuralgia, splenopalatine neuralgia, supraorbital neuralgia, vidian neuralgia, inflammatory bowel disease,
- 30 irritable bowel syndrome, labor, childbirth, menstrual cramps, cancer, back pain, lower back pain and carpal tunnel syndrome pain.

Acute pain includes pain caused by acute injury, trauma, illness or surgery (for example, open-chest surgery (including open-heart or bypass

surgery)). Acute pain also includes, and is not limited to, headache, post-operative pain, kidney stone pain, gallbladder pain, gallstone pain, obstetric pain, rheumatological pain, dental pain or pain caused by sports-medicine injuries, carpal tunnel syndrome, burns, musculoskeletal sprains and strains, musculotendinous strain, cervicobrachial pain syndromes, dyspepsia, gastric ulcer, duodenal ulcer, dysmenorrhea or endometriosis.

Chronic pain includes pain caused by an inflammatory condition, osteoarthritis, rheumatoid arthritis or as sequela to disease, acute injury or trauma. Chronic pain also includes, and is not limited to, headache, upper back pain or lower back pain (selected from back pain resulting from systematic, regional or primary spine disease (selected from radiculopathy)), bone pain (selected from bone pain due to osteoarthritis, osteoporosis, bone metastases or unknown reasons), pelvic pain, spinal cord injury-associated pain, cardiac chest pain, non-cardiac chest pain, central post-stroke pain, myofascial pain, cancer pain, AIDS pain, sickle cell pain, geriatric pain or pain caused by headache, migraine, trigeminal neuralgia, temporomandibular joint syndrome, fibromyalgia syndrome, osteoarthritis, rheumatoid arthritis, gout, fibrositis or thoracic outlet syndromes.

Neuropathic pain includes pain resulting from chronic or debilitating conditions or disorders. The chronic or debilitating conditions or disorders which can lead to neuropathic pain include, but are not limited to, painful diabetic peripheral neuropathy, post-herpetic neuralgia, trigeminal neuralgia, post-stroke pain, multiple sclerosis-associated pain, neuropathies-associated pain such as in idiopathic or post-traumatic neuropathy and mononeuritis, HIV-associated neuropathic pain, cancer-associated neuropathic pain, carpal tunnel-associated neuropathic pain, spinal cord injury-associated pain, complex regional pain syndrome, fibromyalgia-associated neuropathic pain, lumbar and cervical pain, reflex sympathetic dystrophy, phantom limb syndrome and other chronic and debilitating condition-associated pain syndromes.

As used herein, the term “**treatment of substance abuse**” shall include treatment of substance abuse or addiction, including but not limited to the treatment of cravings, withdrawal, and other symptoms of addiction or abuse. As used herein, unless otherwise noted the term “**substance**” when referring to

substances of abuse and / or addiction shall include any legal or illegal substance to which a subject or patient may develop an addiction. Suitable examples include, but are not limited to alcohol, cocaine, heroine, methamphetamine, ketamine, Ecstasy, nicotine, oxycontin / oxycodone, codeine, morphine, and the like.

As used herein, the term “**neuroprotection**” shall mean the protecting neurons in the brain, central nervous system or peripheral nervous system (preferably in the brain or spinal cord) from death and / or damage. Preferably, the neurons are protected from death or damage caused by oxidative stress, for example oxygen radicals.

“**Acute neurodegenerative disorders**” included within the methods of the present invention include, but are not limited, to various types of acute neurodegenerative disorders associated with neuron death or damage including cerebrovascular insufficiency, focal brain trauma, diffuse brain damage, and spinal cord injury, that is, cerebral ischemia or infarction including embolic occlusion and thrombotic occlusion, reperfusion following acute ischemia, perinatal hypoxic-ischemic injury, cardiac arrest, as well as intracranial hemorrhage of any type (including, but not limited to, epidural, subdural, subarachnoid and intracerebral), and intracranial and intravertebral lesions (including, but not limited to, contusion, penetration, shear, compression and laceration), and whiplash shaken infant syndrome. Preferably, the acute neurodegenerative disorder is a result of stroke, acute ischemic injury, head injury or spinal injury.

“**Chronic neurodegenerative disorders**” included within the methods of the present invention included, but are not limited to, Alzheimer's disease, Pick's disease, diffuse Lewy body disease, progressive supranuclear palsy (Steel-Richardson syndrome), multisystem degeneration (Shy-Drager syndrome), chronic epileptic conditions associated with neurodegeneration, motor neuron diseases including amyotrophic lateral sclerosis, degenerative ataxias, cortical basal degeneration, ALS–Parkinson's–Dementia complex of Guam, subacute sclerosing panencephalitis, Huntington's disease, Parkinson's disease, synucleinopathies (including multiple system atrophy), primary progressive aphasia, striatonigral degeneration, Machado-Joseph disease /

- spinocerebellar ataxia type 3 and olivopontocerebellar degenerations, Gilles De La Tourette's disease, bulbar and pseudobulbar palsy, spinal and spinobulbar muscular atrophy (Kennedy's disease), multiple sclerosis, primary lateral sclerosis, familial spastic paraplegia, Werdnig-Hoffmann disease, Kugelberg-
5 Welander disease, Tay-Sach's disease, Sandhoff disease, familial spastic disease, Wohlfart-Kugelberg-Welander disease, spastic paraparesis, progressive multifocal leukoencephalopathy, familial dysautonomia (Riley-Day syndrome), and prion diseases (including, but not limited to Creutzfeldt-Jakob, Gerstmann-Sträussler-Scheinker disease, Kuru and fatal familial insomnia).
- 10 Preferably, the chronic neurodegenerative disorder is selected from Alzheimer's disease, Parkinson's disease, multiple sclerosis or cerebral palsy, Other disorders which manifest neuron death or damage and as such are intended to be included within the methods of the present invention include dementias, regardless of underlying etiology, including age-related dementia
15 and other dementias and conditions with memory loss including dementia associated with Alzheimer's disease, vascular dementia, diffuse white matter disease (Binswanger's disease), dementia of endocrine or metabolic origin, dementia of head trauma and diffuse brain damage, dementia pugilistica and frontal lobe dementia.
- 20 Also included within the present invention are methods of **neuroprotection** (i.e. methods for the prevention of neuron death and / or damage) following injury to the brain, central nervous system or peripheral nervous system, wherein the injury resulting from chemical, toxic, infectious, radiation and / or traumatic injury. Preferably, the methods of the present
25 invention are directed to preventing neuron death or damage following brain, head and / or spinal cord trauma or injury, regardless of cause.
- The term "**subject**" as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment. Preferably, the subject has experienced and / or
30 exhibited at least one symptom of the disease or disorder to be treated and / or prevented.

The term "**therapeutically effective amount**" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or

medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

As used herein, the term “**composition**” is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

One skilled in the art will recognize that, where not otherwise specified, the reaction step(s) is performed under suitable conditions, according to known methods, to provide the desired product. One skilled in the art will further recognize that wherein a reaction step of the present invention may be carried out in a variety of solvents or solvent systems, said reaction step may also be carried out in a mixture of the suitable solvents or solvent systems.

One skilled in the art will recognize that, in the specification and claims as presented herein, wherein a reagent or reagent class/type (e.g. base, solvent, etc.) is recited in more than one step of a process, the individual reagents are independently selected for each reaction step and may be the same or different from each other. For example wherein two steps of a process recite an organic or inorganic base as a reagent, the organic or inorganic base selected for the first step may be the same or different than the organic or inorganic base of the second step.


To provide a more concise description, some of the quantitative expressions given herein are not qualified with the term “**about**”. It is understood that whether the term “about” is used explicitly or not, every quantity given herein is meant to refer to the actual given value, and it is also meant to refer to the approximation to such given value that would reasonably be inferred based on the ordinary skill in the art, including approximations due to the experimental and/or measurement conditions for such given value.

To provide a more concise description, some of the quantitative expressions herein are recited as a range from about amount X to about amount Y. It is understood that wherein a range is recited, the range is not

limited to the recited upper and lower bounds, but rather includes the full range from about amount X through about amount Y, or any range therein.

As used herein, unless otherwise noted, the term “**leaving group**” shall mean a charged or uncharged atom or group which departs during a substitution or displacement reaction. Suitable examples include, but are not limited to, Br, Cl, I, acetoxy, phenoxy, and the like.

Compounds of formula (I) may be prepared according to the process outlined in Scheme 1.




Scheme 1

Accordingly, a compound of formula (S) is reacted with a suitably substituted compound of formula (X), a known compound or compound prepared by known methods, wherein X is a suitably selected leaving group such as Br, Cl, I, acetoxy, phenyloxy, and the like, preferably X is Cl; a known compound or compound prepared by known methods;

in an organic solvent such as THF, 1,4-dioxane, chloroform and the like; preferably at a temperature in the range of from about 60°C to about 100°C, for example, at about solvent reflux temperature; to yield the corresponding compound of formula (I).

In an embodiment, the compound of formula (A) may be prepared as outlined in Scheme 2, below.



Scheme 2

Accordingly, a compound of formula (S) is reacted with a suitably substituted compound of formula (X-A), a known compound or compound
 5 prepared by known methods, wherein X is a suitably selected leaving group such as Br, Cl, I, acetoxy, phenyloxy, and the like, preferably X is Cl; a known compound or compound prepared by known methods;

in an organic solvent such as THF, 1,4-dioxane, chloroform and the like; preferably at a temperature in the range of from about 60°C to about 100°C, for
 10 example, at about solvent reflux temperature; to yield the corresponding compound of formula (A).

For use in medicine, the salts of the compounds of this invention refer to non-toxic “**pharmaceutically acceptable salts.**” Other salts may, however, be
 15 useful in the preparation of compounds according to this invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds include acid addition salts which may, for example, be formed by mixing a solution of the compound with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulfuric acid,
 20 fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g., sodium or

potassium salts; alkaline earth metal salts, e.g., calcium or magnesium salts; and salts formed with suitable organic ligands, e.g., quaternary ammonium salts. Thus, representative pharmaceutically acceptable salts include, but are not limited to, the following: acetate, benzenesulfonate, benzoate, bicarbonate, 5 bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, 10 maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, oleate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide and valerate.

15 Representative acids which may be used in the preparation of pharmaceutically acceptable salts include, but are not limited to, the following: acids including acetic acid, 2,2-dichloroacetic acid, acylated amino acids, adipic acid, alginic acid, ascorbic acid, L-aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, (+)-camphoric acid, camphorsulfonic acid, (+)- 20 (1S)-camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxy-ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, D-gluconic acid, D-glucuronic acid, L-glutamic acid, α -oxo-glutaric acid, glycolic acid, hipuric acid, 25 hydrobromic acid, hydrochloric acid, (+)-L-lactic acid, ($\pm$)-DL-lactic acid, lactobionic acid, maleic acid, (-)-L-malic acid, malonic acid, ($\pm$)-DL-mandelic acid, methanesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, nitric acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, phosphoric acid, L- 30 pyroglutamic acid, salicylic acid, 4-amino-salicylic acid, sebaic acid, stearic acid, succinic acid, sulfuric acid, tannic acid, (+)-L-tartaric acid, thiocyanic acid, p-toluenesulfonic acid and undecylenic acid.

Representative bases which may be used in the preparation of pharmaceutically acceptable salts include, but are not limited to, the following: bases including ammonia, L-arginine, benethamine, benzathine, calcium hydroxide, choline, deanol, diethanolamine, diethylamine, 2-(diethylamino)-ethanol, ethanolamine, ethylenediamine, N-methyl-glucamine, hydrabamine, 1H-imidazole, L-lysine, magnesium hydroxide, 4-(2-hydroxyethyl)-morpholine, piperazine, potassium hydroxide, 1-(2-hydroxyethyl)-pyrrolidine, secondary amine, sodium hydroxide, triethanolamine, tromethamine and zinc hydroxide.

10 The present invention further comprises pharmaceutical compositions containing a compound of formula (I) (for example the compound of formula (A)), with a pharmaceutically acceptable carrier. Pharmaceutical compositions containing one or more of the compounds of the invention described herein as the active ingredient can be prepared by intimately mixing the compound or
15 compounds with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending upon the desired route of administration (e.g., oral, parenteral). Thus for liquid oral preparations such as suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, stabilizers, coloring agents and the like; for
20 solid oral preparations, such as powders, capsules and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Solid oral preparations may also be coated with substances such as sugars or be enteric-coated so as to modulate
25 major site of absorption. For parenteral administration, the carrier will usually consist of sterile water and other ingredients may be added to increase solubility or preservation. Injectable suspensions or solutions may also be prepared utilizing aqueous carriers along with appropriate additives.

30 To prepare the pharmaceutical compositions of this invention, one or more compounds of the present invention as the active ingredient is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration,

e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, 5 flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the 10 most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, through other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable 15 suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The 20 pharmaceutical compositions herein will contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, of from about 0.1-1000 mg or any range therein, and may be given at a dosage of from about 0.1-1000 mg/kg/day, or any range therein, preferably from about 0.5-500 mg/kg/day, or any range therein. The dosages, however, may be varied 25 depending upon the requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms from such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or 30 suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-

monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g.

5 conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to
10 these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing
15 from 0.1 to about 1000 mg, or any amount or range therein, of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of
20 an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as
25 shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or
30 peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

The methods of treatment described in the present invention may also be carried out using a pharmaceutical composition comprising any of the compounds as defined herein and a pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.1 mg and 1000 mg of the compound, or any range therein; preferably about 10 to 500 mg of the compound, and may be constituted into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixers, emulsions, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable
5 preservatives are employed when intravenous administration is desired.

To prepare a pharmaceutical composition of the present invention, a compound of formula (I) as the active ingredient is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the
10 form of preparation desired for administration (e.g. oral or parenteral). Suitable pharmaceutically acceptable carriers are well known in the art. Descriptions of some of these pharmaceutically acceptable carriers may be found in The Handbook of Pharmaceutical Excipients, published by the American Pharmaceutical Association and the Pharmaceutical Society of Great Britain.

15 Methods of formulating pharmaceutical compositions have been described in numerous publications such as Pharmaceutical Dosage Forms: Tablets, Second Edition, Revised and Expanded, Volumes 1-3, edited by Lieberman et al; Pharmaceutical Dosage Forms: Parenteral Medications, Volumes 1-2, edited by Avis et al; and Pharmaceutical Dosage Forms: Disperse Systems, Volumes 1-2, edited by Lieberman et al; published by
20 Marcel Dekker, Inc.

Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the
25 art whenever treatment of anxiety and related disorders; bipolar depression and mania; depression; epilepsy and related disorders; epileptogenesis; glucose related disorders; lipid related disorders; migraine; obesity; pain; substance abuse or neuroprotection is required.

The daily dosage of the products may be varied over a wide range from
30 1.0 to 10,000 mg per adult human per day, or any range therein. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250, 500 and 1,000 milligrams of the active ingredient for the symptomatic

adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.1 mg/kg to about 1000 mg/kg of body weight per day, or any range therein. Preferably, the range is from about 0.5 to about 500 mg/kg of body weight per day, or any range therein.

5 More preferably, from about 1.0 to about 250 mg/kg of body weight per day, or any range therein. More preferably, from about 0.1 to about 100 mg/kg of body weight per day, or any range therein. In an example, the range may be from about 0.1 to about 50.0 mg/kg of body weight per day, or any amount or range therein. In another example, the range may be from about 0.1 to about 15.0
10 mg/kg of body weight per day, or any range therein. In yet another example, the range may be from about 0.5 to about 7.5 mg/kg of body weight per day, or any amount or range therein. The compounds may be administered on a regimen of 1 to 4 times per day.


Optimal dosages to be administered may be readily determined by those
15 skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.

20 One skilled in the art will recognize that, both *in vivo* and *in vitro* trials using suitable, known and generally accepted cell and / or animal models are predictive of the ability of a test compound to treat or prevent a given disorder.

One skilled in the art will further recognize that human clinical trials including first-in-human, dose ranging and efficacy trials, in healthy patients
25 and / or those suffering from a given disorder, may be completed according to methods well known in the clinical and medical arts.

The following Examples are set forth to aid in the understanding of the invention, and are not intended and should not be construed to limit in any way
30 the invention set forth in the claims which follow thereafter.

Example 1: Preparation of the Compound of Formula (A)



Acetyl chloride (0.71 g, 9.0 mmol) was added to (2S)-(-)-N-(6-chloro-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl)-sulfamide (2.1 g, 7.53 mmol) in 1,4-dioxane (30 mL) and the resulting mixture was brought to reflux under argon.

- 5 After about 2 h, the resulting mixture was cooled to room temperature and evaporated in vacuo to yield a white solid (2.63 g). The white solid was purified using reverse phase HPLC (20 – 90% ACN) to yield the title compound as a white solid (1.03 g). The white solid was then recrystallized from ethyl acetate and hexane to yield the title compound as a white powdery solid.

10 mp: 184 – 185°C

Optical rotation: $[\alpha]_D = -47.3^\circ$ (c = 1.20, CH₃OH)

Elemental Analysis:

Calculated: C, 41.19; H, 4.09; N, 8.73; Cl, 11.05; S, 10.00.

Measured: C, 41.15; H, 4.00; N, 8.62; Cl, 11.02; S, 9.87.

- 15 ¹H NMR (DMSO d₆): δ 11.45 (s, 1H), 8.02 (dd, J = 6.0, 6.0 Hz, 1H), 6.98 (dd, 1.4, 1.4 Hz, 1H), 6.89 (m, 2H), 4.28 (m, 2H), 4.0 (dd, J = 6.4, 11.3 Hz, 1H), 3.19 (dd, J = 6.0, 6.0 Hz, 2H), 1.94 (s, 3H).

Example 2: Oral Formulation – Prophetic Example

- As a specific embodiment of an oral composition, 100 mg of the
20 compound prepared as in Example 1 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

Example 3: In Vivo Assay: Maximal Electroshock Test (MES)

- Anticonvulsant activity was determined using the MES test, run
25 according to the procedure described in detail below. Swinyard EA, Woodhead JH, White HS, Franklin MR. Experimental selection, quantification, and evaluation of anticonvulsants. In Levy RH, et al., eds. *Antiepileptic Drugs*. 3rd ed. New York: Raven Press, 1989:85-102.

- CF-1 male albino mice (25-35g) were fasted for 16 hours before testing.
30 Mice were randomly selected into control and test groups, with the animals

dosed with vehicle or test compound, at varying concentrations, respectively. On the study date, at 30 minutes prior to shock, the mice were orally dosed with vehicle (0.5% methylcellulose) or test compound (200 mg/kg). Seizures were induced by trans-corneal electric shock using a 60-Hz alternating current, 50 mA, delivered for 0.2 sec. The mice in the test groups were subjected to electrical stimulus at time intervals between 15 minutes and 4 hours following administration of test compound. The shock resulted in an immediate full body tonic extension. The test was complete when the entire course of the convulsion has been observed (typically, less than 1 minute after electrical stimulation), and the mice were then immediately euthanized by carbon dioxide inhalation.

Abolition of the full body tonic extensor component of the seizure was taken as the endpoint of the test. Absence of this component indicated that the test compound had the ability to prevent the spread of seizure discharge through neural tissue. The ED₅₀ value of the test compound (calculated when appropriate) was the calculated dose required to block the hind limb tonic-extensor component of the MES-induced seizure in 50% of the rodents tested. A probit analysis was used to calculate the ED₅₀ and 95% fiducial limits (FL).

The compound of formula (A) was tested according to the procedure described above, with results as listed in Table 1 below. Results are listed as (number of mice with full body tonic extension prevented) / (total number of mice tested) (@ a given time).


Table 1: MES Activity

	0.50 Hrs	1.0 Hrs	2 Hrs	3 Hrs	6 Hrs
Vehicle	0 / 5				
Compound (A) 200 mg/kg po	0 / 5	1 / 5	4 / 5	5 / 5	2 / 5
Compound (S) 150 mg/kg po				5 / 5	

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

We Claim:

1. A compound of formula (I)




wherein

- 5 R^1 is selected from the group consisting of C_{1-4} alkyl and -O-phenyl;
 wherein the phenyl is optionally substituted with one to three substituents,
 independently selected from the group consisting of halogen, C_{1-4} alkyl,
 fluorinated C_{1-4} alkyl, C_{1-4} alkoxy, fluorinated C_{1-4} alkoxy, cyano, nitro, $-NR^A R^B$
 and $-C(O)-NR^C R^D$;
 10 wherein R^A and R^B are each independently selected from the group
 consisting of C_{1-4} alkyl;
 and wherein R^C and R^D are each independently selected from the group
 consisting of hydrogen and C_{1-4} alkyl;
 or a pharmaceutically acceptable salt thereof;
 15 provided that when R^1 is methyl, then the compound of formula (I) is
 present as an isolated or a substantially pure form.

2. A compound as in Claim 1, wherein R^1 is selected from the group
 consisting of C_{2-4} alkyl and -O-phenyl.

20

3. A compound of formula (A)




or a pharmaceutically acceptable salt thereof;

- 25 wherein the compound of formula (A) is present as an isolated or a
 substantially pure form.



4. A compound as in Claim 3, wherein the compound of formula (A) is present as an isolated and a substantially pure form.
5. A compound as in Claim 3, wherein the compound of formula (A) is present as an isolated, substantially pure form, having an (-) optical rotation.
6. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Claim 1.
- 10 7. A pharmaceutical composition made by mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.
8. A process for making a pharmaceutical composition comprising mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.
- 15 9. A method of treating a disorder selected from the group consisting of anxiety and related disorders; bipolar depression and mania; depression; epilepsy and related disorders; epileptogenesis; glucose related disorders; lipid related disorders; migraine; obesity; pain and substance abuse; or a method of neuroprotection; comprising administering to a subject in need thereof a therapeutically effective amount of a compound as in Claim 1.
- 20 10. A method as in Claim 9, wherein the disorder is selected from the group consisting of epilepsy and related disorders; depression; glucose related disorders; and obesity.
- 25 11. A method of treating of treating a disorder selected from the group consisting of anxiety and related disorders; bipolar depression and mania; depression; epilepsy and related disorders; epileptogenesis; glucose related disorders; lipid related disorders; migraine; obesity; pain and substance abuse; or a method of neuroprotection; comprising administering to a subject in need thereof a therapeutically effective amount of the composition of Claim 6.
- 30

12. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Claim 3.
13. A pharmaceutical composition made by mixing a compound of Claim 3
5 and a pharmaceutically acceptable carrier.
14. A process for making a pharmaceutical composition comprising mixing a compound of Claim 3 and a pharmaceutically acceptable carrier.
- 10 15. A method of treating a disorder selected from the group consisting of anxiety and related disorders; bipolar depression and mania; depression; epilepsy and related disorders; epileptogenesis; glucose related disorders; lipid related disorders; migraine; obesity; pain and substance abuse; or a method of neuroprotection; comprising administering to a subject in need thereof a
15 therapeutically effective amount of a compound as in Claim 3.
16. A method as in Claim 15, wherein the disorder is selected from the group consisting of epilepsy and related disorders; depression; glucose related disorders; and obesity.
- 20 17. A method of treating of treating a disorder selected from the group consisting of anxiety and related disorders; bipolar depression and mania; depression; epilepsy and related disorders; epileptogenesis; glucose related disorders; lipid related disorders; migraine; obesity; pain and substance abuse;
25 or a method of neuroprotection; comprising administering to a subject in need thereof a therapeutically effective amount of the composition of Claim 12.
18. A process for the preparation of a compound of formula (I)




30 wherein

- 5 R^1 is selected from the group consisting of C_{2-4} alkyl and -O-phenyl; wherein the phenyl is optionally substituted with one to three substituents, independently selected from the group consisting of halogen, C_{1-4} alkyl, fluorinated C_{1-4} alkyl, C_{1-4} alkoxy, fluorinated C_{1-4} alkoxy, cyano, nitro, $-NR^A R^B$ and $-C(O)-NR^C R^D$; wherein R^A and R^B are each independently selected from the group consisting of C_{1-4} alkyl; and wherein R^C and R^D are each independently selected from the group consisting of hydrogen and C_{1-4} alkyl; or a pharmaceutically acceptable salt thereof; comprising




- 10 reacting a compound of formula (S) with a compound of formula (X), wherein X is a leaving group; in an organic solvent; to yield the corresponding compound of formula (I).

19. A process for the preparation of a compound of formula (A)



- 15 or a pharmaceutically acceptable salt thereof; comprising



reacting a compound of formula (S) with a compound of formula (X-A), wherein X is a leaving group; in an organic solvent; to yield the corresponding compound of formula (A).

5

20. A process as in Claim 19, wherein X is chloro; wherein the organic solvent is 1,4-dioxane; and wherein the compound of formula (S) is reacted with the compound of formula (X-A) at about solvent reflux temperature.

10 21. The use of a compound as in Claim 1, for the preparation of a medicament for treating: (a) anxiety and related disorders; (b) bipolar depression (c) mania; (d) depression; (e) epilepsy and related disorders; (f) epileptogenesis; (g) glucose related disorders; (h) lipid related disorders; (i) migraine; (j) obesity; (k) pain; (l) substance abuse and (m) for neuroprotection,
15 in a subject in need thereof.

22. The use of a compound as in Claim 3, for the preparation of a medicament for treating: (a) anxiety and related disorders; (b) bipolar depression (c) mania; (d) depression; (e) epilepsy and related disorders; (f)
20 epileptogenesis; (g) glucose related disorders; (h) lipid related disorders; (i) migraine; (j) obesity; (k) pain; (l) substance abuse and (m) for neuroprotection, in a subject in need thereof.