

Description

Title of Invention: NOVEL BENZAMIDE DERIVATIVES

Technical Field

- [1] The present invention relates to a novel benzamide derivative of formula 1 which will be illustrated hereinafter or a pharmaceutically acceptable salt thereof, a method for preparing the same, and a 5-HT₄ receptor agonist containing the same as an active ingredient.

Background Art

- [2] Serotonin (5-HT) is a neurotransmitter widely distributed throughout the body. Seven subtypes of serotonin are currently known. In particular, great attention has been focused on the elucidation of a 5-HT₄ receptor and the confirmation of pharmaceutical action thereof.
- [3] In general, 5-HT₄ receptor agonists are found to be therapeutically effective for the treatment of various disease conditions such as gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive impairment, emesis, migraine, neurological disease, pain, cardiovascular disorder, cardiac failure, cardiac arrhythmia, diabetes and apnea syndrome (See Tips, 1992, 13, 141; Ford A.P.D.W. et al., *Med. Res. Rev.*, 1993, 13, 633; Gullikson G.W. et al., *Drug Dev. Res.*, 1992, 26, 405; Richard M. Eglen et al., *Tips*, 1995, 16, 391; Bockaert J. et al., *CNS Drugs*, 1, 6; Romanelli M.N. et al., *Arzheim Forsch./Drug Res.*, 1993, 43, 913; Kaumann A. et al., *Naunyn-Schmiedeberg's*, 1991, 344, 150; and Romanelli M.N. et al., *Arzheim Forsch./Drug Res.*, 1993, 43, 913).
- [4] Despite extensive uses of 5-HT₄ receptor agonists, there are little 5-HT₄ receptor agonist compounds that are currently clinically used. To this end, there is a need for a 5-HT₄ receptor agonist which is capable of exhibiting excellent medicinal effects while having minimum adverse side effects.
- [5] Benzamide derivatives have several prominent pharmacological actions. These excellent pharmacological activities of the benzamide derivatives are due to their effects on the nervous system which is controlled by serotonin that is a neurotransmitter. The role of serotonin, that is, the pharmacological action of benzamide derivatives has been broadly involved in a variety of diseases and conditions for many years. Further, a great deal of study and research has focused on production and storage sites of serotonin as well as the location of serotonin receptors in order to determine the relationship between the location of serotonin receptors and various

disease states or conditions in humans.

- [6] Cisapride, which is a typical 5-HT₄ receptor agonist, is one of benzamide derivatives. U.S. Patent Nos. 4,962,115, 5,057,525 and 5,137,896 disclose N-(3-hydroxy-4-piperidiny)benzamides including cisapride. These compounds are known to stimulate gastrointestinal motility. Further, U.S. Patent No. 5,864,039 discloses benzamide derivatives.

- [7] To this end, the inventors of the present invention succeeded in the synthesis of novel benzamide derivatives which exhibit an agonistic activity via strong binding with a 5-HT₄ receptor and good gastrointestinal absorption and which are capable of minimizing adverse side effects. The present invention has been completed based on this finding.

Disclosure of Invention

Technical Problem


- [8] The present invention is intended to provide a novel benzamide derivative or a pharmaceutically acceptable salt thereof and a method for preparing the same.
- [9] Further, the present invention is intended to provide a 5-HT₄ receptor agonist containing a novel benzamide derivative or a pharmaceutically acceptable salt thereof or a hydrate thereof as an active ingredient.

Solution to Problem

- [10] The present invention provides a novel benzamide derivative represented by formula 1(a compound represented by formula 1):

[11] [Formula 1]

[12]



- [13] wherein m represents an integer of 1 to 10; and Q represents a heteroaromatic ring or phenyl, wherein the heteroaromatic ring or phenyl is independently substituted by 0, 1, 2, or 3 substituents selected from alkyl, alkoxy, hydroxy, cyano, nitro and halogen; or a pharmaceutically acceptable salt thereof.
- [14]
- [15] Unless otherwise indicated, the following terms will have the following meanings throughout this specification.
- [16] As used herein, the term "alkyl" refers to a linear or branched, monovalent saturated

C₁-C₂₀ hydrocarbon radical containing only carbon atoms and hydrogen atoms.

Examples of the alkyl radical include methyl, ethyl, propyl, isopropyl, 2,2-dimethylpropyl, butyl, isobutyl, sec-butyl, tert-butyl, 3-methylbutyl, pentyl, 3-methylpentyl, 4-methylpentyl, n-hexyl, 2-ethylhexyl, octyl, and dodecyl.

[17] As used herein, the term "alkoxy" refers to a radical -OR wherein R represents alkyl as defined above. Examples of the alkoxy radical include methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentoxy, 3-methylpentoxy, 4-methylpentoxy, n-hexoxy, and 2-ethylhexoxy.

[18] As used herein, the term "heteroaromatic ring" refers to an aromatic ring or bicyclic aromatic ring contains 1 to 4 hetero atoms selected from O, N or S. Examples of the heteroaromatic ring include pyrrole, imidazole, triazole, tetrazole, pyridine, pyrimidine, oxazole, oxadiazole, isoxazole, indole, quinoline and benzofuran.

[19] Further, the present invention provides a novel benzamide derivative represented by formula 1 wherein m represents an integer of 1 to 5; and Q represents a heteroaromatic ring or phenyl wherein the heteroaromatic ring or phenyl is independently substituted by 0, 1, 2 or 3 substituents selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy and halogen, wherein the heteroaromatic ring being a C₁-C₁₂ aromatic ring or C₁-C₁₂ bicyclic aromatic ring independently contains 1 to 4 hetero atoms selected from N, O or S; or a pharmaceutically acceptable salt thereof.

[20] In the present invention, the pharmaceutically acceptable salt may be an acid addition salt with an acceptable free acid. The free acid may be an inorganic or organic acid. Examples of the inorganic acid include hydrochloric acid, hydrobromic acid, sulfuric acid, and phosphoric acid. Examples of the organic acid include citric acid, acetic acid, lactic acid, maleic acid, fumaric acid, gluconic acid, methanesulfonic acid, glycolic acid, succinic acid, 4-toluenesulfonic acid, trifluoroacetic acid, galacturonic acid, embonic acid, glutamic acid, and aspartic acid.

[21] Further, the compound of formula 1 or a pharmaceutically acceptable salt thereof may exhibit polymorphism and may also be present in the form of a solvate (e.g., hydrate, etc).

[22] Further, the present invention relates to a novel benzamide derivative selected from the group consisting of the following compounds:

[23] (1) N-((1-(3-(1,2,4-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

[24] (2) N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5

[25] -chloro-2-methoxybenzamide,

[26] (3) N-((1-(3-(indol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5

[27] -chloro-2-methoxybenzamide,

- [28] (4) N-((1-(3-(2-methylimidazol-1-yl)propyl)piperidin-4-yl)methyl)-4
[29] -amino-5-chloro-2-methoxybenzamide,
[30] (5) N-((1-(5-(indol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5
[31] -chloro-2-methoxybenzamide,
[32] (6) N-
((1-(5-(1,2,3-triazol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
[33] (7) N-
((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
[34] (8) N-
((1-(3-(1,2,3-triazol-2-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
[35] (9) N-((1-(pyridin-3-ylmethyl)piperidin-4-yl)methyl)-4-amino-5-chloro
[36] -2-methoxybenzamide,
[37] (10) N-((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino
[38] -5-chloro-2-methoxybenzamide,
[39] (11) N-((1-(imidazol-2-ylmethyl)piperidin-4-yl)methyl)-4-amino-5
[40] -chloro-2-methoxybenzamide,
[41] (12) N-
((1-((1-methylpyrrol-2-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
[42] (13) N-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2
[43] -methoxybenzamide,
[44] (14) N-((1-(4-hydroxybenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2
[45] -methoxybenzamide,
[46] (15) N-((1-(2-(indol-3-yl)ethyl)piperidin-4-yl)methyl)-4-amino-5
[47] -chloro-2-methoxybenzamide,
[48] (16) N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5
[49] -chloro-2-methoxybenzamide hydrochloride,
[50] (17) N-((1-(5-(1,2,3-triazol-1-yl)pentyl)piperidin-4-yl)methyl)-4
[51] -amino-5-chloro-2-methoxybenzamide hydrochloride,
[52] (18) N-((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4
[53] -amino-5-chloro-2-methoxybenzamide hydrochloride,
[54] (19) N-((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino
[55] -5-chloro-2-methoxybenzamide hydrochloride,
[56] (20) N-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2
[57] -methoxybenzamide hydrochloride,


[58] (21) N-((1-(4-hydroxybenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride and a pharmaceutically acceptable salt thereof.

[60] Further, the present invention provides a method for preparing a benzamide derivative of formula 1 or a pharmaceutically acceptable salt thereof.

[61] The present invention provides a method for preparing a compound of formula 1 or a pharmaceutically acceptable salt thereof, which includes reacting a compound of formula 2 or a pharmaceutically acceptable salt thereof with a compound of formula 3 in the presence of a base to introduce the compound of formula 3 at an amine of the 1-position of the piperidine ring of the compound of formula 2 or the pharmaceutically acceptable salt thereof, thereby preparing the compound of formula 1 (hereinafter, referred to as "Preparation Method 1").


[62] [Formula 1]

[63]



[64] [Formula 2]

[65]



[66] [Formula 3]

[67]



[68] In the above formulae, m and Q are as defined in formula 1, and Y represents a halogen atom or C₁-C₄ alkylsulfonate.


[69] In Preparation Method 1 of the present invention, the base is preferably selected from potassium carbonate, potassium iodide, triethylamine, diisopropylethylamine and their mixture, the solvent may be dimethylformamide, dimethylacetamide, acetone,

1,4-dioxane or the like, and the reaction may be carried out at a temperature of 50°C to 140°C.

[70] The present invention provides a method for preparing a compound of formula 1 or a pharmaceutically acceptable salt thereof, which includes reacting a compound of formula 2 or a pharmaceutically acceptable salt thereof with a compound of formula 11 in the presence of a reducing agent to prepare the compound of formula 1 (hereinafter, referred to as "Preparation Method 2").


[71] [Formula 1]

[72]




[73] [Formula 2]

[74]



[75] [Formula 11]

[76]



[77] In the above formulae, Q is as defined in formula 1, and m represents an integer of 1.

[78] In Preparation Method 2 of the present invention, the reducing agent is preferably sodium cyanoborohydride and acetic acid, or sodium borohydride, the solvent may be a C₁-C₆ lower alcohol, preferably ethanol or methanol, and the reaction may be carried out at a temperature of 50°C to 100°C.


[79] The compound of formula 2 or the pharmaceutically acceptable salt thereof in

Preparation Method 1 or Preparation Method 2 of the present invention may be prepared by the following steps of:

- [80] (1) reacting a compound of formula 4 with an amine protecting group-introducing reagent to introduce an amine protecting group at an amine of the 1-position of the piperidine ring of the compound of formula 4, thereby obtaining a compound of formula 5;
- [81] (2) reacting hydroxy of the compound of formula 5 with N-bromosuccinimide and carbon tetrabromide, or with C₁-C₄ alkyl sulfonyl halide in the presence of a base to obtain a compound of formula 6;
- [82] (3) reacting a substituent Y of the compound of formula 6 with sodium azide to obtain a compound of formula 7;
- [83] (4) reducing the azido substituent of the compound of formula 7 into an amine in the presence of a reducing agent to obtain a compound of formula 8;
- [84] (5) reacting the compound of formula 8 with a compound of formula 9 in the presence of an amide bond formation-inducing reagent to obtain a compound of formula 10; and
- [85] (6) deprotecting the amine protecting group of the piperidine ring of the compound of formula 10 in the presence of a base or acid.


[86] [Formula 2]

[87]




[88] [Formula 4]

[89]




[90] [Formula 5]

[91]




[92] [Formula 6]

[93]




[94] [Formula 7]

[95]




[96] [Formula 8]

[97]




[98] [Formula 9]

[99]



[100] [Formula 10]

[101]



[102] In the above formulae, Y represents a halogen atom or C₁-C₄ alkylsulfonate, and Z represents C₁-C₄ alkyl.

[103] In the method for preparing a compound of formula 2 in accordance with the present invention, the amine protecting group-introducing reagent of Step (1) refers to a reagent used conventionally for the protection of an amine in order to prevent an amine group from participating in the reaction. For example, such a reagent is preferably selected from di-*t*-butyl dicarbonate or ethyl chloroformate in the presence of a tertiary amine such as triethylamine. The solvent may be a C₁-C₆ lower alcohol. The reaction may be carried out with gradual elevation of a temperature from 0°C to room temperature.

[104] In the method for preparing a compound of formula 2 in accordance with the present invention, the C₁-C₄ alkyl sulfonyl halide of Step (2) is preferably methane sulfonyl chloride, methane sulfonyl bromide or methane sulfonyl fluoride, the tertiary amine may be triethylamine, diisopropylethylamine or the like, and the solvent may be dichloromethane, chloroform, or the like. The reaction may be carried out with gradual elevation of a temperature from 0°C to room temperature.

[105] In the method for preparing a compound of formula 2 in accordance with the present invention, the solvent used in Step (3) may be dimethylformamide, dimethylacetamide or the like, and the reaction temperature may be in the range of 80 to 140°C.

[106] In the method for preparing a compound of formula 2 in accordance with the present invention, the reducing agent used in Step (4) is preferably triphenylphosphine or lithium aluminum hydride, and the solvent used may be tetrahydrofuran. The reaction may be carried out with gradual elevation of a temperature from 0°C to room temperature or may be carried out in the range of 60 to 80°C.


[107] In the method for preparing a compound of formula 2 in accordance with the present invention, the amide bond formation-inducing reagent used in Step (5) refers to a common reagent which is used by a person of ordinary skill and which is used to remove water generated after the reaction, in order to promote the amide bond between carboxylic acid and an amine, or is used to activate an amine or carboxylic acid. Examples of the amide bond formation-inducing reagent include N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride and 1-hydroxybenzotriazole in the presence of a base, ethyl chloroformate in the presence

of a base, and carbodiimidazole in the absence of a base. Here, examples of the base used in combination with N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride and 1-hydroxybenzotriazole, or ethyl chloroformate include triethylamine and diisopropylethylamine. Here, the reaction solvent used may be dimethyl formamide, dimethyl acetamide, dichloromethane, or the like. The reaction may be carried out with gradual elevation of a temperature from 0°C to room temperature.

[108] In the method for preparing a compound of formula 2 in accordance with the present invention, the base or acid of Step (6) refers to a base or acid which is conventionally used to deprotect the carbamate of an amine, and examples thereof include hydrochloric acid, trifluoroacetic acid, and potassium hydroxide. The reaction solvent used may be 1,4-dioxane, dichloromethane, C₁-C₆ lower alcohol, or the like. The reaction may be carried out with gradual elevation of a temperature from 0°C to room temperature.

[109] The compound of formula 3 in the present invention may be prepared according to the method as in Reaction Scheme 1 or Reaction Scheme 2 provided below.


[110] [Reaction Scheme 1]



[112] In Reaction Scheme 1, the substituents Q and m are as defined in formula 1, and both of Y and Y' represent halogen atoms which are preferably different from each other, for example, one of Y and Y' represents chloro (Cl) and the other one of Y and Y' represents bromo (Br).

[113] The reaction of Reaction Scheme 1 may be carried out in the presence of a strong base such as lithium hydride, sodium hydride or potassium hydride, in an organic solvent such as dimethylformamide, dimethylacetamide or tetrahydrofuran, at a reaction temperature of 0 to 40°C for 1 to 24 hours.


[114] [Reaction Scheme 2]



[116] In Reaction Scheme 2, the substituents Q and m are as defined in formula 1, Y represents a C₁-C₄ alkyl sulfonyl halide or a halogen atom, Y'' represents a hydroxyl group, and Y''' represents a halogen atom selected from Cl, Br or I.


[117] The first step reaction of Reaction Scheme 2 may be carried out in the presence of a base such as potassium carbonate and potassium iodide, in a solvent such as 1,4-dioxane or acetone, at a temperature of 60 to 120°C for 1 to 12 hours.

[118] When Y of formula 3 represents a C₁-C₄ alkyl sulfonyl halide, the second step reaction of Reaction Scheme 2 may be carried out by reacting



methane sulfonyl chloride, methane sulfonyl bromide, or methane sulfonyl fluoride) in the presence of a base such as triethylamine or diisopropylethylamine, in an organic solvent such as dichloromethane or chloroform, at a reaction temperature of 0 to 40°C for 1 to 24 hours.

- [119] When the substituent Y of formula 3 represents a halogen atom, the second step reaction of Reaction Scheme 2 may be carried out under the conventionally known reaction conditions for substitution of the hydroxyl group of



formula 3 represents bromo (Br), the reaction may be carried out in the presence of one species selected from N-bromosuccinimide or carbon tetrabromide and triphenylphosphine, in an organic solvent such as dichloromethane, at a reaction temperature of 0 to 40°C for 1 to 24 hours.

- [120] The acid addition salt of a free base of a compound represented by formula 1 may be prepared using a conventional method known in the art, for example by mixing a free base of the compound of formula 1 with an appropriate acid in a suitable solvent, which is then followed by evaporation to form a salt or the addition of a non-solvent to precipitate a salt. For example, mention may be made of a method which involves treating a solution or suspension of a free base with a desired acid in a reaction-inert solvent, followed by concentration under reduced pressure or crystallization or any standard chemical manipulation to form a desired salt. In one embodiment, a hydrochloride of the compound of formula 1 may be prepared by dissolving a free base of the compound of formula 1 in a C₁-C₄ alcohol solvent such as ethanol or methanol, adding hydrochloric acid thereto, and then stirring the mixture at room temperature.

- [121] Further, the present invention provides a 5-HT₄ receptor agonist containing a compound of formula 1 or a pharmaceutically acceptable salt thereof or a hydrate thereof as an active ingredient.

- [122] In the present invention, the 5-HT₄ receptor agonist may be a composition for the prevention or treatment of a disease selected from gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive impairment, emesis, migraine, neurological disease, pain, cardiovascular disorder, cardiac failure, cardiac arrhythmia, diabetes or apnea syndrome.

- [123] In the present invention, the 5-HT₄ receptor agonist of the present invention may further contain one or more active ingredients exhibiting an identical or similar function, in addition to the compound of formula 1 in accordance with the present invention or a pharmaceutically acceptable salt thereof.

- [124] For the purpose of desired administration, the agonist or composition of the present invention may be formulated into a variety of dosage forms by further inclusion of one or more pharmaceutically acceptable carriers in combination with the above-mentioned active ingredient. Examples of the pharmaceutically acceptable carrier include saline, sterile water, Ringer's solution, buffered saline, a dextrose solution, a maltodextrin solution, glycerol, and ethanol. These materials may be used alone or in any combination thereof. If necessary, other conventional additives may be added such as an antioxidant, a buffer and a bacteriostatic agent. Further, a diluent, a dispersant, a surfactant, a binder and a lubricant may be additionally added to prepare an injectable formulation such as aqueous solution, suspension or emulsion, or an oral formulation such as pill, capsule, granule or tablet. Furthermore, the desired dosage form may be preferably formulated depending on diseases to be treated and ingredients, using any appropriate method known in the art, as disclosed in "Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, PA.
- [125] Further, when the 5-HT₄ receptor agonist of the present invention is intended for oral administration, the content of the compound of formula 1 in accordance with the present invention or a pharmaceutically acceptable salt thereof in the formulation may be in the range of 1 to 95% by weight and preferably 1 to 70% by weight.
- [126] The agonist or composition of the present invention may be administered parenterally (for example, intravenously, subcutaneously, intraperitoneally or locally) or orally, depending on desired applications. The dose of the active ingredient may vary depending on various factors such as weight, age, sex, health status and dietary habits of patients, administration times and routes, excretion rates, and severity of diseases. The benzamide derivative of the present invention may be administered at a dose of 1 to 1000 µg/kg, preferably about 10 to 500 µg/kg and more preferably about 83 to 167 µg/kg, once or several times a day.
- [127] Further, the present invention provides a method for the prevention, treatment or alleviation of a disease due to attenuated efficacy of a 5-HT₄ receptor, including administering a 5-HT₄ receptor agonist containing a compound of formula 1 in accordance with the present invention or a pharmaceutically acceptable salt thereof as an active ingredient to a mammal including a human in need of 5-HT₄ receptor agonistic effects. Here, the 5-HT₄ receptor agonist may be a composition for the prevention or treatment of a disease selected from gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive impairment, emesis, migraine, neurological disease, pain, cardiovascular disorder, cardiac failure, cardiac arrhythmia, diabetes or apnea syndrome.

- [128] For the prevention and treatment of a disease due to attenuated efficacy of a 5-HT₄ receptor, the 5-HT₄ receptor agonist of the present invention may be used alone or in combination with methods employing surgical operation, hormone therapy, medication therapy and biological response modifiers.

Advantageous Effects of Invention

- [129] Benzamide derivatives of the present invention have a superior affinity for 5-HT₄ receptors, a capability to reduce a gastric emptying time and a low toxicity, and consequently are therapeutically effective for the treatment of a variety of diseases associated with 5-HT₄ receptors.

Mode for the Invention

- [130] Hereinafter, the present invention will be described in more detail with reference to the following Examples. However, the following Examples are provided only for illustrating the present invention and should not be construed as limiting the scope and spirit of the present invention.
- [131] Unless otherwise indicated, reagents used hereinafter were purchased from Aldrich Korea, Acros, Lancaster, TCI, etc., and ¹H NMR experiments were performed using a Varian 400MHz spectrometer.
- [132]
- [133] **Example 1: Preparation of N-((1-(3-(1,2,4-triazol-1-yl)propyl)**
- [134] **piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**
- [135]
- [136] **1-1. Preparation of 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)**
- [137] **benzamide (compound of formula 2) hydrochloride**
- [138] **Step 1: Preparation of tertiary butyl 4-(hydroxymethyl)piperidine-1-carboxylate**
(compound of formula 5)

- [139] 4-piperidine methanol (compound of formula 4) (20g, 174mmol) was dissolved in methanol (30mL) and the solution was cooled to 0°C. Then, triethylamine (48.8mL, 347mmol) and di-tertiary-butyl dicarbonate (56.8g, 260mmol) were added thereto. The reaction mixture was warmed to room temperature, stirred for 2 hours, and concentrated under reduced pressure to remove the solvent. After extraction with dichloromethane and water, the organic layer was washed with a citric acid solution, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound (35.25g, 94%).
- [140] ¹H NMR (CDCl₃, 400MHz): δ 4.18-4.04 (m, 2H), 3.47 (t, *J*=5.6Hz, 2H), 2.70-2.64 (m, 2H), 1.70-1.56 (m, 3H), 1.42 (s, 9H), 1.16-1.09 (m, 2H).

[141]

- [142] **Step 2: Preparation of tertiary butyl 4-((methylsulfonyloxy)methyl)**

[143] **piperidine-1-carboxylate (compound of formula 6)**

[144] Tertiary butyl 4-(hydroxymethyl)piperidine-1-carboxylate (35.25g, 164mmol) was dissolved in dichloromethane (300mL) and the solution was cooled to 0°C. Then, triethylamine (46.01mL, 327mmol) and methane sulfonyl chloride (19.14mL, 246mmol) were added thereto. The reaction mixture was warmed to room temperature, stirred for 2 hours, and extracted with dichloromethane and water. The organic layer was washed with a citric acid solution, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound (48g, 100%).

[145] ¹H NMR (CDCl₃, 400MHz): δ 4.18-4.06 (m, 2H), 4.04 (d, *J* = 6.4Hz, 2H), 2.99 (s, 3H), 2.78-2.62 (m, 2H), 1.92-1.85 (m, 1H), 1.73-1.70 (m, 2H), 1.43 (s, 9H), 1.24-1.13 (m, 2H).

[146]

[147] **Step 3: Preparation of tertiary butyl 4-(azidomethyl)piperidine-1-carboxylate (compound of formula 7)**

[148] Tertiary butyl 4-((methylsulfonyloxy)methyl)piperidine-1-carboxylate (48g, 163.6mmol) was dissolved in dimethylformamide (500mL) and sodium azide (19.15g, 295mmol) was added thereto, followed by stirring at 120°C for 6 hours. After the reaction was completed, the reactants were cooled to room temperature, extracted with ethyl acetate and water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to afford the title compound (35g, 89%).

[149] ¹H NMR (CDCl₃, 400MHz): δ 4.18-4.00 (m, 2H), 3.16 (d, *J* = 6.0Hz, 2H), 2.72-2.58 (m, 2H), 1.70-1.64 (m, 3H), 1.43 (s, 9H), 1.20-1.08 (m, 2H).

[150]

[151] **Step 4: Preparation of tertiary butyl 4-(aminomethyl)piperidine-1-carboxylate (compound of formula 8)**

[152] Tertiary butyl 4-(azidomethyl)piperidine-1-carboxylate (30g, 124.8mmol) was dissolved in tetrahydrofuran (300mL), and triphenylphosphine (39.3g, 149.8mmol) was added thereto, followed by stirring under reflux for 2 hours. Water (120mL) was added thereto, followed by stirring under reflux for another 3 hours and concentration under reduced pressure. The resulting residue was extracted with ethyl acetate and a 1N hydrochloride solution, and the aqueous layer was neutralized with a 2N sodium hydroxide solution, followed by re-extraction with dichloromethane. The resulting organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound (21.16g, 79%).

[153] ¹H NMR (CDCl₃, 400MHz): δ 4.16-3.98 (m, 2H), 2.69-2.63 (m, 2H), 2.56 (d, *J* = 6.8Hz, 2H), 1.69-1.66 (m, 2H), 1.46-1.38 (m, 10H), 1.10-1.04 (m, 2H).

[154]

[155] **Step 5: Preparation of tertiary butyl**

4-((4-amino-5-chloro-2-methoxybenzamido)methyl)piperidine-1-carboxylate (compound of formula 10)

[156] 4-amino-5-chloro-2-methoxybenzoic acid (compound of formula 9) (16.6g, 82.28mmol) was dissolved in dimethylformamide (166mL) and the solution was cooled to 0°C. Then, tertiary butyl 4-(aminomethyl)piperidine-1-carboxylate (21.16g, 98.74mmol), triethylamine (11.56mL, 246.84mmol), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (EDC, 20.51g, 106.96mmol), and 1-hydroxybenzotriazole (HOBT, 16.68g, 123.42mmol) were added thereto. The reaction mixture was warmed to room temperature, stirred for 4 hours, and extracted with ethyl acetate and water. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound (31.4g, 96%).

[157] ¹H NMR (CDCl₃, 400MHz): δ 8.08 (s, 1H), 7.76-7.70 (m, 1H), 6.27 (s, 1H), 4.37 (s, 2H), 4.16-4.04 (m, 2H), 3.88 (s, 3H), 3.34-3.26 (m, 2H), 2.74-2.60 (m, 2H), 1.80-1.63 (m, 3H), 1.42 (s, 9H), 1.20-1.10 (m, 2H).

[158]

[159] **Step 6: Preparation of 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide (compound of formula 2) hydrochloride**

[160] Tertiary butyl 4-((4-amino-5-chloro-2-methoxybenzamido)methyl)piperidine-1-carboxylate (30.7g, 77.16mmol) was dissolved in 1,4-dioxane (300mL), and a 4M hydrochloride solution (220mL) in 1,4-dioxane was added thereto. The mixture was stirred at room temperature for 2 hours and concentrated under reduced pressure to afford the title compound (24g, 93%).

[161] ¹H NMR (CD₃OD, 400MHz): δ 7.84 (s, 1H), 6.84 (s, 1H), 3.95 (s, 3H), 3.42-3.29 (m, 6H), 3.01-2.95 (m, 2H), 1.98-1.93 (m, 3H), 1.54-1.42 (m, 2H).

[162]

[163] **1-2. Preparation of 1-(3-chloropropyl)-1,2,4-triazole (compound of formula 3)**

[164] A 1,2,4-triazole sodium salt (5g, 54.91mmol) was dissolved in dimethylformamide (50mL) and the solution was cooled to 0°C. Then, sodium hydride (60%, 2.86g, 71.38mmol) was added thereto, followed by stirring for 30 minutes. 1-bromo-3-chloropropane (6.5mL, 65.89mmol) was added thereto, followed by stirring at room temperature for 12 hours, and the reaction was terminated with the addition of an ammonium chloride saturated solution. After extraction with ethyl acetate and water, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by column chromatography to afford the title compound (2.77g, 35%).

[165] ¹H NMR (CDCl₃, 400MHz): δ 8.08 (s, 1H), 7.94 (s, 1H), 4.36 (t, *J* = 6.4Hz, 2H), 3.45 (t, *J* = 6Hz, 2H), 2.35-2.30 (m, 2H).

[166]

[167] **1-3: Preparation of N-((1-(3-(1,2,4-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide (compound of formula 1)**

[168] 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide (compound of formula 2) hydrochloride (500mg, 1.5mmol) was dissolved in dimethylformamide (10mL), and 1-(3-chloropropyl)-1,2,4-triazole (compound of formula 3) (306mg, 2.1mmol), potassium carbonate (497mg, 3.6mmol), and potassium iodide (50mg, 0.3mmol) were added thereto. The reactants were stirred at 100°C for 8 hours, and the reaction was terminated with the addition of water. After extraction with ethyl acetate and water, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by column chromatography to afford the title compound (55mg, 9%).

[169] ¹H NMR (CDCl₃, 400MHz): δ 8.09 (s, 1H), 8.04 (s, 1H), 7.91 (s, 1H), 7.78-7.70 (m, 1H), 6.27 (s, 1H), 4.35 (s, 2H), 4.22 (t, *J* = 6.4Hz, 2H), 3.88 (s, 3H), 3.30 (t, *J* = 6.4Hz, 2H), 2.85-2.81 (m, 2H), 2.24 (t, *J* = 6.4Hz, 2H), 2.06-2.00 (m, 2H), 1.97-1.90 (m, 2H), 1.78-1.50 (m, 3H), 1.39-1.28 (m, 2H).

[170]

[171] **Example 2: Preparation of N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**

[172] 1-(3-chloropropyl)-tetrazole was prepared using 1H-tetrazole and 1-bromo-3-chloropropane as starting materials of <Example 1-2>, and then the title compound (186mg) was obtained using 1-(3-chloropropyl)-tetrazole and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials of <Example 1-3>.

[173] ¹H NMR (CDCl₃, 400MHz): δ 8.60 (s, 1H), 8.07 (s, 1H), 7.75-7.68 (m, 1H), 6.28 (s, 1H), 4.49 (t, *J* = 6.8Hz, 2H), 4.39 (s, 2H), 3.88 (s, 3H), 3.30 (t, *J* = 6.4Hz, 2H), 2.79-2.76 (m, 2H), 2.23 (t, *J* = 6.8Hz, 2H), 2.10-2.05 (m, 2H), 1.92-1.86 (m, 2H), 1.75-1.69 (m, 2H), 1.62-1.50 (m, 1H), 1.35-1.23 (m, 2H).

[174]

[175] **Example 3: Preparation of N-((1-(3-(indol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**

[176] 1-(3-chloropropyl)-indole was prepared using indole and 1-bromo-3-chloropropane as starting materials of <Example 1-2>, and then the title compound (26mg) was

obtained using 1-(3-chloropropyl)-indole and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials of <Example 1-3>.

- [177] ¹H NMR (CDCl₃, 400MHz): δ 8.09 (s, 1H), 7.85-7.76 (m, 1H), 7.60 (d, *J* = 7.6Hz, 1H), 7.34 (d, *J* = 8.0Hz, 1H), 7.15 (t, *J* = 7.2Hz, 1H), 7.10-7.07 (m, 2H), 6.45 (d, *J* = 3.2Hz, 1H), 6.27 (s, 1H), 4.34 (s, 2H), 4.18 (t, *J* = 7.2Hz, 2H), 3.88 (s, 3H), 3.32 (t, *J* = 6.4Hz, 2H), 2.88-2.85 (m, 2H), 2.26 (t, *J* = 7.2Hz, 2H), 2.03-1.97 (m, 2H), 1.90-1.80 (m, 2H), 1.73-1.60 (m, 3H), 1.40-1.32 (m, 2H).

[178]

[179] **Example 4: Preparation of N-((1-(3-(2-methylimidazol-1-yl)propyl)**
[180] **piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**

- [181] 1-(3-chloropropyl)-2-methylimidazole was prepared using 2-methylimidazole and 1-bromo-3-chloropropane as starting materials of <Example 1-2>, and then the title compound (226mg) was obtained using 1-(3-chloropropyl)-2-methylimidazole and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials of <Example 1-3>.

- [182] ¹H NMR (CDCl₃, 400MHz): δ 8.06 (s, 1H), 7.78-7.69 (m, 1H), 6.85 (s, 1H), 6.78 (s, 1H), 6.27 (s, 1H), 4.46 (s, 2H), 3.87-3.83 (m, 5H), 3.29 (t, *J* = 6.0Hz, 2H), 2.85-2.79 (m, 2H), 2.34 (s, 3H), 2.22 (t, *J* = 6.8Hz, 2H), 1.90-1.83 (m, 4H), 1.72-1.67 (m, 2H), 1.63-1.50 (m, 1H), 1.32-1.26 (m, 2H).

[183]

[184] **Example 5: Preparation of N-((1-(5-(indol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**

- [185] 1-(5-chloropentyl)-indole was prepared using indole and 1-bromo-5-chloropentane as starting materials of <Example 1-2>, and then the title compound (250mg) was obtained using 1-(5-chloropentyl)-indole and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials of <Example 1-3>.

- [186] ¹H NMR (CDCl₃, 400MHz): δ 8.09 (s, 1H), 7.75-7.69 (m, 1H), 7.60 (d, *J* = 8.0Hz, 1H), 7.31 (d, *J* = 7.6Hz, 1H), 7.19-7.16 (m, 1H), 7.09-7.05 (m, 2H), 6.47-6.44 (m, 1H), 6.26 (s, 1H), 4.34 (s, 2H), 4.11-4.08 (m, 2H), 3.87 (s, 3H), 3.30 (t, *J* = 6.0Hz, 2H), 2.92-2.86 (m, 2H), 2.28-2.20 (m, 2H), 1.91-1.80 (m, 4H), 1.72-1.48 (m, 5H), 1.39-1.26 (m, 4H).

[187]

[188] **Example 6: Preparation of N-((1-(5-(1,2,3-triazol-1-yl)pentyl)**
[189] **piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**

- [190] 1-(5-chloropentyl)-1,2,3-triazole was prepared using 1,2,3-triazole and

1-bromo-5-chloropentane as starting materials of <Example 1-2>, and then the title compound (225mg) was obtained using 1-(5-chloropentyl)-1,2,3-triazole and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials of <Example 1-3>.

[191] ^1H NMR (CDCl_3 , 400MHz): δ 8.06 (s, 1H), 7.79-7.70 (m, 1H), 7.67 (s, 1H), 7.52 (s, 1H), 6.27 (s, 1H), 4.41 (s, 2H), 4.35 (t, $J=6.8\text{Hz}$, 2H), 3.86 (s, 3H), 3.28 (t, $J=6.0\text{Hz}$, 2H), 2.92-2.88 (m, 2H), 2.28 (t, $J=7.6\text{Hz}$, 2H), 1.94-1.84 (m, 4H), 1.72-1.68 (m, 2H), 1.66-1.46 (m, 3H), 1.34-1.21 (m, 4H).

[192]

[193] **Example 7: Preparation of N-((1-(3-(1,2,3-triazol-1-yl)propyl) piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**

[194] **Step 1: Preparation of 3-(1,2,3-triazol-1-yl)propanol**

[195] 1,2,3-triazole (2g, 28.96mmol) was dissolved in 1,4-dioxane (40mL), and potassium carbonate (8g, 57.92mmol), potassium iodide (962mg, 5.79mmol) and 3-bromopropanol (3.3mL, 43.43mmol) were added thereto, followed by stirring at 100°C for 3 hours. After the reaction was completed, the reactants were cooled to room temperature and filtered. The filtrate was concentrated under reduced pressure. The resulting residue was purified by column chromatography to afford the title compound (2.5g, 68%).

[196] ^1H NMR (CDCl_3 , 400MHz): δ 7.68 (s, 1H), 7.59 (s, 1H), 4.54 (t, $J=6.8\text{Hz}$, 2H), 3.62 (q, $J=5.6\text{Hz}$, 2H), 2.44 (t, $J=5.2\text{Hz}$, 1H), 2.15-2.10 (m, 2H).

[197] **Step 2: Preparation of 3-(1,2,3-triazol-1-yl)propyl methanesulfonate**

[198] 3-(1,2,3-triazol-1-yl)propanol (2.5g, 19.66mmol) was dissolved in dichloromethane (50mL), and the solution was cooled to 0°C. Then, triethylamine (5.53mL, 39.32mmol) and methane sulfonyl chloride (2.3mL, 29.5mmol) were added thereto. The reaction mixture was warmed to room temperature, stirred for 2 hours, extracted with dichloromethane and water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting residue was purified by column chromatography to afford the title compound (3.01g, 75%).

[199] ^1H NMR (CDCl_3 , 400MHz): δ 7.72 (s, 1H), 7.63 (s, 1H), 4.55 (t, $J=6.8\text{Hz}$, 2H), 4.22 (t, $J=6\text{Hz}$, 2H), 3.03 (s, 3H), 2.40-2.36 (m, 2H).

[200]

[201] **Step 3: Preparation of N-**

[202] **((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**

[203] 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide (compound of formula 2) hydrochloride (400mg, 1.2mmol) was dissolved in dimethylformamide (10mL), and 3-(1,2,3-triazol-1-yl)propyl methanesulfonate (compound of formula 3)

(344mg, 1.68mmol), triethylamine (0.5mL, 3.591mmol), potassium carbonate (232mg, 1.68mmol), and potassium iodide (40mg, 0.24mmol) were added thereto, followed by stirring at 120°C for 4 hours. After the reaction was completed, the reactants were cooled to room temperature, extracted with ethyl acetate and water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting residue was purified by column chromatography to afford the title compound (185mg, 38%).

[204] ¹H NMR (CDCl₃, 400MHz): δ 8.09 (s, 1H), 7.74-7.70 (m, 1H), 7.67 (s, 1H), 7.54 (s, 1H), 6.27 (s, 1H), 4.43 (t, *J*=6.8Hz, 2H), 4.35 (s, 2H), 3.88 (s, 3H), 3.31 (t, *J*=6.4Hz, 2H), 2.85-2.82 (m, 2H), 2.27 (t, *J*=6.8Hz, 2H), 2.07-2.04 (m, 2H), 1.92-1.86 (m, 2H), 1.73-1.69 (m, 2H), 1.62-1.50 (m, 1H), 1.34-1.22 (m, 2H).

[205]

[206] **Example 8: Preparation of N-((1-(3-(1,2,3-triazol-2-yl)propyl)**
[207] **piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide**

[208] Using 3-(1,2,3-triazol-2-yl)propanol obtained as a by-product in Step 1 of Example 7 and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials, the title compound (250mg) was obtained in the same manner as in <Example 7>.

[209] ¹H NMR (CDCl₃, 400MHz): δ 8.03 (s, 1H), 7.83-7.75 (m, 1H), 7.53 (s, 2H), 6.25 (s, 1H), 4.50-4.42 (m, 4H), 3.82 (s, 3H), 3.26 (t, *J*=6.0Hz, 2H), 2.86-2.83 (m, 2H), 2.31-2.27 (m, 2H), 2.12-2.05 (m, 2H), 1.89-1.83 (m, 2H), 1.67-1.64 (m, 2H), 1.60-1.50 (m, 1H), 1.31-1.20 (m, 2H).

[210]

[211] **Example 9: Preparation of N-**
((1-(pyridin-3-ylmethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide

[212] 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride (1g, 2.99mmol) was dissolved in methanol (40mL), and 3-pyridine carboxaldehyde (0.42mL, 4.49mmol), acetic acid (1mL) and sodium cyanoborohydride (470mg, 7.48mmol) were added thereto, followed by stirring at 60°C for 5 hours. After the reaction was completed, the reaction mixture was concentrated under reduced pressure to remove the solvent, extracted with dichloromethane and water, and then washed with a sodium bicarbonate solution. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by column chromatography to afford the title compound (355mg, 30%).

[213] ¹H NMR (CDCl₃, 400MHz): δ 8.07 (s, 1H), 7.74-7.69 (m, 1H), 7.65 (d, *J*=7.6Hz, 1H), 7.37 (s, 1H), 7.30 (d, *J*=8.4Hz, 1H), 7.25-7.22 (m, 1H), 7.19-7.15 (m, 1H), 6.26 (s, 1H), 4.42 (s, 2H), 3.83 (s, 3H), 3.77 (s, 3H), 3.28 (t, *J*=5.6Hz, 2H), 2.95-2.82 (m,

4H), 2.19-2.14 (m, 4H), 1.91-1.85 (m, 2H), 1.80-1.72 (m, 2H), 1.69-1.63 (m, 2H), 1.61-1.53 (m, 1H), 1.32-1.15 (m, 5H).

[214]

[215] **Example 10: Preparation of N-**

((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide

[216] Using 1-methylindole-3-carboxaldehyde and

4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials, the title compound (60mg) was obtained in the same manner as in <Example 9>.

[217] ¹H NMR (CDCl₃, 400MHz): δ 8.02 (s, 1H), 7.82-7.75 (m, 1H), 7.56 (d, *J*=8.4Hz, 1H), 7.33-7.22 (m, 3H), 7.12 (t, *J*=7.2Hz, 1H), 6.27 (s, 1H), 4.38 (s, 2H), 4.01 (s, 2H), 3.87 (s, 3H), 3.79 (s, 3H), 3.30-3.23 (m, 2H), 2.46-2.37 (m, 2H), 1.88-1.75 (m, 3H), 1.59-1.50 (m, 2H).

[218]

[219] **Example 11: Preparation of N-**

((1-(imidazol-2-ylmethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide

[220] Using 2-imidazole carboxaldehyde and

4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials, the title compound (123mg) was obtained in the same manner as in <Example 9>.

[221] ¹H NMR (CDCl₃, 400MHz): δ 8.07 (s, 1H), 7.75-7.66 (m, 1H), 6.96 (s, 2H), 6.27 (s, 1H), 4.38 (s, 2H), 3.87 (s, 3H), 3.61 (s, 2H), 3.31 (t, *J*=6.8Hz, 2H), 2.85-2.80 (m, 2H), 2.11-2.04 (m, 2H), 1.74-1.69 (m, 2H), 1.63-1.53 (m, 1H), 1.34-1.23 (m, 2H).

[222]

[223] **Example 12: Preparation of N-((1-((1-methylpyrrol-2-yl)methyl)**

piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide

[225] Using 1-methyl-2-pyrrole carboxaldehyde and

4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials, the title compound (50mg) was obtained in the same manner as in <Example 9>.

[226] ¹H NMR (CDCl₃, 400MHz): δ 8.08 (s, 1H), 7.75-7.64 (m, 1H), 6.57-6.53 (m, 1H), 6.26 (s, 1H), 6.00-5.96 (m, 1H), 5.95-5.93 (m, 1H), 4.38 (s, 2H), 3.85 (s, 3H), 3.60 (s, 2H), 3.28 (t, *J*=6.8Hz, 2H), 2.86-2.82 (m, 2H), 1.92-1.84 (m, 2H), 1.68-1.63 (m, 2H), 1.61-1.53 (m, 1H), 1.30-1.19 (m, 2H).

[227]

[228] **Example 13: Preparation of N-((1-(4-fluorobenzyl)piperidin-4-yl)**

[229] **methyl)-4-amino-5-chloro-2-methoxybenzamide**

[230] Using 4-fluorobenzaldehyde and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials, the title compound (215mg) was obtained in the same manner as in <Example 9>.

[231] ¹H NMR (CDCl₃, 400MHz): δ 8.07 (s, 1H), 7.75-7.68 (m, 1H), 7.27-7.21 (m, 2H), 7.00-6.94 (m, 2H), 6.26 (s, 1H), 4.36 (s, 2H), 3.86 (s, 3H), 3.46-3.43 (m, 2H), 3.28 (t, *J* = 6.4Hz, 2H), 2.87-2.84 (m, 2H), 2.01-1.91 (m, 2H), 1.70-1.56 (m, 3H), 1.36-1.20 (m, 2H).

[232]

[233] **Example 14: Preparation of N-((1-(4-hydroxybenzyl)piperidin-4-yl)**

[234] **methyl)-4-amino-5-chloro-2-methoxybenzamide**

[235] Using 4-hydroxybenzaldehyde and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials, the title compound (75mg) was obtained in the same manner as in <Example 9>.

[236] ¹H NMR (CDCl₃, 400MHz): δ 7.81 (s, 1H), 7.19 (d, *J* = 8.8Hz, 2H), 6.75 (s, *J* = 8.8Hz, 2H), 6.20 (s, 1H), 4.40 (s, 2H), 3.76 (s, 2H), 3.32-3.24 (m, 5H), 3.12-3.05 (m, 2H), 2.68-2.59 (m, 2H), 1.80-1.72 (m, 3H), 1.27-1.12 (m, 2H).

[237]

[238] **Example 15: Preparation of N-**

((1-(2-(indol-3-yl)ethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide

[239] **Step 1: Preparation of 2-(indol-3-yl)ethanol**

[240] Indole-3-acetic acid (5g, 28.54mmol) was dissolved in diethyl ether (100mL) and the solution was cooled to 0°C. Lithium aluminum hydride (1.19g, 31.39mmol) was added thereto, followed by stirring for 4 hours. The reaction was terminated with the addition of water and a 10% sodium hydroxide solution. The reaction mixture was filtered through celite and concentrated under reduced pressure. The resulting residue was purified by column chromatography to afford the title compound (1.24g, 27%).

[241] ¹H NMR (CDCl₃, 400MHz): δ 8.10-7.96 (bs, 1H), 7.61 (d, *J* = 7.6Hz, 1H), 7.36 (dd, *J* = 8Hz, 0.8Hz, 1H), 7.24-7.17 (m, 1H), 7.15-7.07 (m, 2H), 3.92-3.87 (m, 2H), 3.05-3.00 (m, 2H).

[242]

[243] **Step 2: Preparation of 3-(2-bromoethyl)-indole (compound of formula 3)**

[244] 2-(indol-3-yl)ethanol (623mg, 3.86mmol) was dissolved in dichloromethane (20mL) and the solution was cooled to 0°C. Triphenylphosphine (1.12g, 4.25mmol) and tetra-bromomethane (1.41g, 4.25mmol) were then added thereto, followed by stirring for 1

hour. After the reaction was completed, the reaction mixture was concentrated under reduced pressure to remove the solvent. The resulting residue was purified by column chromatography to afford the title compound (765mg, 88%).

[245] ¹H NMR (CDCl₃, 400MHz): δ 8.10-7.88 (bs, 1H), 7.58 (d, *J*=8Hz, 1H), 7.38-7.34 (m, 1H), 7.24-7.20 (m, 1H), 7.19-7.08 (m, 2H), 3.65-3.60 (m, 2H), 3.35-3.30 (m, 2H).

[246]

[247] **Step 3: Preparation of N-((1-(2-(indol-3-yl)ethyl)piperidin-4-yl)**

[248] **methyl)-4-amino-5-chloro-2-methoxybenzamide**

[249] Using 3-(2-bromoethyl)-indole and 4-amino-5-chloro-2-methoxy-(piperidin-4-ylmethyl)benzamide hydrochloride as starting materials, the title compound (190mg) was obtained in the same manner as in <Example 1-3>.

[250] ¹H NMR (CDCl₃, 400MHz): δ 8.09 (s, 1H), 8.05 (s, 1H), 7.81-7.75 (m, 1H), 7.59 (d, *J*=7.6Hz, 1H), 7.33 (d, *J*=8.4Hz, 1H), 7.16 (t, *J*=7.2Hz, 1H), 7.09 (t, *J*=7.6Hz, 1H), 7.00 (d, *J*=1.6Hz, 1H), 6.27 (s, 1H), 4.36 (s, 2H), 3.88 (s, 3H), 3.32 (t, *J*=6.4Hz, 2H), 3.12-3.07 (m, 2H), 3.01-2.96 (m, 2H), 2.73-2.69 (m, 2H), 2.11-2.04 (m, 2H), 1.79-1.75 (m, 2H), 1.74-1.63 (m, 1H), 1.49-1.38 (m, 2H).

[251]

[252] **Example 16: Preparation of N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride**

[253] N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide (Example 2) (517mg, 1.27mmol) was dissolved in ethanol (10mL) and a 12N hydrochloride solution (0.16mL, 1.90mmol) was added thereto. The reaction mixture was stirred at room temperature for 12 hours and then filtered to afford the title compound (417mg, 74%).

[254] ¹H NMR (DMSO-*d*₆, 400MHz): δ 10.52 (s, 1H), 9.47 (s, 1H), 8.02-7.99 (m, 1H), 7.64 (s, 1H), 6.47 (s, 1H), 5.95 (s, 2H), 4.59-4.56 (m, 2H), 3.81 (s, 3H), 3.45-3.42 (m, 2H), 3.18-3.08 (m, 2H), 3.04-2.98 (m, 2H), 2.87-2.79 (m, 2H), 2.35-2.31 (m, 2H), 1.90-1.89 (m, 1H), 1.78-1.75 (m, 2H), 1.58-1.52 (m, 2H).

[255]

[256] **Example 17: Preparation of N-((1-(5-(1,2,3-triazol-1-yl)pentyl)**

[257] **piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride**

[258] Using N-((1-(5-(1,2,3-triazol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide (Example 6) as a starting material, the title compound (112mg, 35%) was obtained in the same manner as in <Example 16>.

[259] ¹H NMR (DMSO-*d*₆, 400MHz): δ 10.06 (s, 1H), 8.14 (s, 1H), 8.02-7.99 (m, 1H),

7.71 (s, 1H), 7.64 (s, 1H), 6.47 (s, 1H), 5.95 (s, 2H), 4.40-4.37 (m, 2H), 3.81 (s, 3H), 3.40-3.32 (m, 2H), 3.18-3.12 (m, 2H), 2.96-2.90 (m, 2H), 2.84-2.75 (m, 2H), 1.87-1.70 (m, 7H), 1.59-1.48 (m, 2H), 1.25-1.17 (m, 2H).

[260]

[261] **Example 18: Preparation of N-((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride**
[262]

[263] Using N-((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide (Example 7) as a starting material, the title compound (291mg, 93%) was obtained in the same manner as in <Example 16>.

[264] ¹H NMR (DMSO-*d*₆, 400MHz): δ 10.67 (s, 1H), 8.19 (s, 1H), 8.02-7.99 (m, 1H), 7.74 (s, 1H), 7.64 (s, 1H), 6.48 (s, 1H), 5.93 (bs, 2H), 4.51-4.47 (m, 2H), 3.81 (s, 3H), 3.44-3.41 (m, 2H), 3.34-3.31 (m, 2H), 2.98-2.96 (m, 2H), 2.87-2.79 (m, 2H), 2.34-2.29 (m, 2H), 1.96-1.86 (m, 1H), 1.77-1.74 (m, 2H), 1.60-1.54 (m, 2H).

[265]

[266] **Example 19: Preparation of N-((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride**

[267] Using N-((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide (Example 10) as a starting material, the title compound (237mg, 44%) was obtained in the same manner as in <Example 16>.

[268] ¹H NMR (DMSO-*d*₆, 400MHz): δ 10.34 (s, 1H), 7.96-7.93 (m, 1H), 7.79 (d, *J* = 8Hz, 1H), 7.64-7.59 (m, 2H), 7.47 (d, *J* = 8Hz, 1H), 7.22-7.19 (m, 1H), 7.14-7.10 (m, 1H), 6.47 (s, 1H), 4.36 (d, *J* = 4.4Hz, 2H), 3.81 (s, 3H), 3.79 (s, 3H), 3.40-3.37 (m, 2H), 3.14-3.11 (m, 2H), 2.90-2.82 (m, 2H), 1.77-1.71 (m, 3H), 1.53-1.48 (m, 2H).

[269]

[270] **Example 20: Preparation of N-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride**
[271]

[272] Using N-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide (Example 13) as a starting material, the title compound (569mg, 87%) was obtained in the same manner as in <Example 16>.

[273] ¹H NMR (DMSO-*d*₆, 400MHz): δ 10.78 (s, 1H), 7.98-7.96 (m, 1H), 7.73-7.63 (m, 3H), 7.30-7.25 (m, 2H), 6.47 (s, 1H), 5.94 (s, 2H), 4.22 (d, *J* = 4.8Hz, 2H), 3.80 (s, 3H), 3.29-3.26 (m, 2H), 3.14-3.07 (m, 2H), 2.86-2.78 (m, 2H), 1.89-1.85 (m, 1H), 1.77-1.74 (m, 2H), 1.59-1.53 (m, 2H).

[274]

[275] **Example 21: Preparation of N-((1-(4-hydroxybenzyl)piperidin-4-yl)**
[276] **methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride**

[277] Using N-((1-(4-hydroxybenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide (Example 14) as a starting material, the title compound (158mg, 97%) was obtained in the same manner as in <Example 16>.

[278] ¹H NMR (DMSO-*d*₆, 400MHz): δ 10.25 (s, 1H), 7.98-7.96 (m, 1H), 7.63 (s, 1H), 7.34 (d, *J* = 6.4Hz, 2H), 6.80 (d, *J* = 6.4Hz, 2H), 6.47 (s, 1H), 4.08 (d, *J* = 3.6Hz, 2H), 3.80 (s, 3H), 3.28-3.25 (m, 2H), 3.15-3.12 (m, 2H), 2.82-2.74 (m, 2H), 1.77-1.62 (m, 3H), 1.54-1.48 (m, 2H).

[279]

[280] **Experimental Example 1: Binding affinity of compounds for 5-HT₄ receptor**

[281] The binding affinity of the compounds for a human 5-HT₄ receptor was assayed according to the method as disclosed in the literature [Wyngaert et al., Journal of Neurochemistry, (1997) 69, 1810-1819]. For this purpose, COS-7 cells expressing the human 5-HT₄ receptor were constructed and homogenized to obtain membrane homogenates which were then used in binding assay experiments. For the binding assay, the membrane homogenates were respectively mixed and incubated with different concentrations of test materials and [H3]-GR113808 (Amersham Biosciences). The concentrations of the individual test materials were set to 4μM, 1μM, 0.25μM, and 0.0625μM, respectively, and the concentration of [H3]-GR113808 was set to 0.595nM. After the incubation was completed, the reaction products were collected in GF/B glass fiber filters using a Packard cell harvester, and the bound radioactivity was then determined using a liquid cell scintillation counter (Packard TopCount NXT™, Perkin Elmer). Specific binding of the radioligand to the 5-HT₄ receptor was calculated by subtracting the non-specific binding of the radioligand from the total radioligand binding. IC₅₀ was calculated from % inhibition of specific binding of the radioligand to the 5-HT₄ receptor, with respect to varying concentrations of the test materials. The results are given in Table 1 below.

[282] [Table 1]

[283]

Compounds	IC ₅₀ (nM)
Cisapride	0.362
Example 1	0.226
Example 2	0.164
Example 3	0.039
Example 4	0.229
Example 5	0.005
Example 6	0.103
Example 7	0.067
Example 8	0.182
Example 9	0.267
Example 10	0.076
Example 11	0.454
Example 12	0.421
Example 13	0.134
Example 14	0.375
Example 15	0.002

[284] As can be seen from Table 1, the compounds of the present invention inhibited specific binding of the radioligand to the 5-HT₄ receptor at a concentration similar to or lower than that of cisapride as a control, thus representing that the inventive compounds have a strong binding affinity for 5-HT₄ receptor.

[285]

[286]

[287] **Experimental Example 2: Evaluation of gastric emptying**

[288] Evaluation of gastric emptying was carried out based on the method described in Iwanaga Y, et al., *Jpn J Pharmacol.* 1991, 56(3), 261-269. As experimental animals, SD rats (240 to 250g) were used and divided into groups (n= 5 to 6). On the day prior to the experiment, animals were fasted for 18 hours. Test materials were formulated as suspensions thereof in 0.5%(w/v) methylcellulose and were orally administered to the animals at a dose of 5mg/kg. 1 hour after the oral administration of the test materials, 2mL of a semi-solid diet was orally administered to the animals. The semi-solid diet was in the form of porridge formulated by pulverizing a standard solid diet together with distilled water in a mixer. 50 minutes after the administration of the semi-solid diet, the animals were sacrificed by cervical dislocation. After carrying out an abdominal incision, the stomach was extracted from the animals. The weight of diet

remaining in the stomach was measured to thereby determine the gastric emptying rate (%). The gastric emptying rate was calculated according to the following equation. The results are given in Table 2 below.

[289]

[290] **Gastric emptying rate (%) = (1-X/Y)*100**

[291] X: Weight of diet remaining in the stomach extracted 50 minutes after administration of meal

[292] Y: Weight of diet remaining in the stomach extracted immediately after administration of meal

[293]

[294] [Table 2]

[295]

Compounds	Gastric emptying rate (%)
Control (0.5% methylcellulose)	32.2
Example 1	43.0
Example 2	52.8
Example 3	45.2
Example 4	46.4
Example 5	52.0
Example 6	46.0
Example 7	56.1
Example 8	49.1
Example 9	51.2
Example 10	59.1
Example 11	43.1
Example 12	62.0
Example 13	52.3
Example 14	42.1
Example 15	41.3

[296] As can be seen from Table 2, the administration of the compounds of the present invention resulted in a remarkable improvement of gastric emptying, as compared to the methylcellulose-treated group as a control, thus representing that the inventive compounds promote gastrointestinal motility.

[297] **Experimental Example 3: Acute oral toxicity of compounds in mice**

[298] In order to examine acute toxicity of the compounds in accordance with the present invention, the following experiment was carried out.

[299] 200mg of each of Example compounds was triturated in 1% hydroxypropyl methyl-cellulose, and the resulting triturations were then orally administered to 5-week old male ICR mice (20g±2g, n = 5) at a dose of 1g/10mL/kg. The minimum lethal dose (MLD, mg/kg) of the individual compounds was investigated by the observation of the mortality, body weight, clinical symptoms and the like of animals over the entire experimental period of 2 weeks. The results are given in Table 3 below.

[300] [Table 3]

[301]

Compounds	Minimum lethal dose (MLD, mg/kg)
Example 1	>1000
Example 2	>1000
Example 3	>1000
Example 4	>1000
Example 5	>1000
Example 6	>1000
Example 7	>1000
Example 8	>1000
Example 9	>1000
Example 10	>1000
Example 11	>1000
Example 12	>1000
Example 13	>1000
Example 14	>1000
Example 15	>1000

[302] As can be seen from the acute toxicity test results of Table 3, all of the compounds used in the test exhibited an MLD of more than 1000mg/kg, thus representing that the inventive compounds are safe for use.

[303]

[304]

[305] **Experimental Example 4: Binding affinity of drug compounds for hERG receptor**

[306] The binding affinity of the compounds for the human ether-a-go-go-related gene (hERG) potassium (K⁺) channel which is associated with cardiac QT prolongation was assayed in MDS Pharma Service (Catalog No. 265900). Membrane homogenates were obtained from mammalian HEK-293 cells expressing the hERG potassium channel and then used in the binding assay experiment. For the binding assay experiment, the membrane homogenates were respectively mixed and incubated with 10 μ M of test materials and 1.5nM of [H₃]-Astemizole (Perkin Elmer). After the incubation was completed, the radioactivity bound to the hERG K⁺ channel was counted. The affinity of each test material for the hERG K⁺ channel was calculated from % inhibition of specific binding of the radioligand to the hERG K⁺ channel, resulting from the action of the test material. The results are given in Table 4 below.

[307]

[308] [Table 4]

[309]

Compounds	Inhibition (% at 10 μ M)
Example 1	1
Example 2	4
Example 4	14
Example 6	8
Example 7	7
Example 8	15
Example 9	6
Example 11	21
Example 12	32
Example 13	52
Example 14	40

[310] Incidence of cardiac arrhythmia which is a fatal adverse effect is due to the cardiac QT prolongation that results from an excessively high affinity of the drug for the hERG receptor. The compounds of the present invention exhibited an inhibition rate of less than 50% even at a dose of 10 μ M and therefore have a low binding affinity for the


hERG receptor, thus suggesting that the inventive compounds exhibit a significantly low risk of causing arrhythmia.

Claims

[Claim 1]

A compound represented by formula 1:

[Formula 1]



wherein m represents an integer of 1 to 10; and Q represents a heteroaromatic ring or phenyl, wherein the heteroaromatic ring or phenyl is independently substituted by 0, 1, 2, or 3 substituents selected from alkyl, alkoxy, hydroxy, cyano, nitro and halogen; or a pharmaceutically acceptable salt thereof.

[Claim 2]

The compound of formula 1 or a pharmaceutically acceptable salt thereof according to claim 1, wherein m represents an integer of 1 to 5; and Q represents a heteroaromatic ring or phenyl wherein the heteroaromatic ring or phenyl is independently substituted by 0, 1, 2 or 3 substituents selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy and halogen, wherein the heteroaromatic ring being a C₁-C₁₂ aromatic ring or C₁-C₁₂ bicyclic aromatic ring independently contains 1 to 4 hetero atoms selected from N, O or S;

[Claim 3]

The compound of formula 1 or a pharmaceutically acceptable salt thereof according to claim 1, wherein the compound of formula 1 or a pharmaceutically acceptable salt thereof is a compound selected from the group consisting of the following compounds:

- (1) N-((1-(3-(1,2,4-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
- (2) N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
- (3) N-((1-(3-(indol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
- (4) N-((1-(3-(2-methylimidazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
- (5) N-((1-(5-(indol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5


- chloro-2-methoxybenzamide,
(6) N-((1-(5-(1,2,3-triazol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(7) N-((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(8) N-((1-(3-(1,2,3-triazol-2-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(9) N-((1-(pyridin-3-ylmethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(10) N-((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(11) N-((1-(imidazol-2-ylmethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(12) N-((1-((1-methylpyrrol-2-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(13) N-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(14) N-((1-(4-hydroxybenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(15) N-((1-(2-(indol-3-yl)ethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,
(16) N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,
(17) N-((1-(5-(1,2,3-triazol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,
(18) N-((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,
(19) N-((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,
(20) N-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,
(21) N-((1-(4-hydroxybenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride, and a pharmaceutically acceptable

salt thereof.


[Claim 4]

A method for preparing a compound of formula 1 or a pharmaceutically acceptable salt thereof, comprising reacting a compound of formula 2 or a pharmaceutically acceptable salt thereof with a compound of formula 3 in the presence of a base to introduce the compound of formula 3 at an amine of the 1-position of the piperidine ring of the compound of formula 2 or the pharmaceutically acceptable salt thereof, thereby preparing the compound of formula 1.

[Formula 1]



[Formula 2]



[Formula 3]



wherein m and Q are as defined in claim 1, and Y represents a halogen atom or C₁-C₄ alkylsulfonate.

[Claim 5]


The method according to claim 4, wherein the base is selected from potassium carbonate, potassium iodide, triethylamine, diisopropylethylamine and their mixture.

[Claim 6]


A method for preparing a compound of formula 1 or a pharmaceutically acceptable salt thereof, comprising reacting a compound of formula 2 or a pharmaceutically acceptable salt thereof with a compound of formula 11 in the presence of a reducing agent to prepare the

compound of formula 1.


[Formula 1]



[Formula 2]



[Formula 11]



wherein Q is as defined in claim 1, and m represents an integer of 1.

[Claim 7]

The method according to claim 6, wherein the reducing agent is sodium cyanoborohydride and acetic acid, or sodium borohydride.

[Claim 8]

The method according to claim 4 or 6, wherein the compound of formula 2 or the pharmaceutically acceptable salt thereof is prepared by:

- (1) reacting a compound of formula 4 with an amine protecting group-introducing reagent to introduce an amine protecting group at an amine of the 1-position of the piperidine ring of the compound of formula 4, thereby obtaining a compound of formula 5;
- (2) reacting hydroxy of the compound of formula 5 with N-bromosuccinimide and carbon tetrabromide, or with C₁-C₄ alkyl

sulfonyl halide in the presence of a base to obtain a compound of formula 6;


(3) reacting a substituent Y of the compound of formula 6 with sodium azide to obtain a compound of formula 7;

(4) reducing the azido substituent of the compound of formula 7 into an amine in the presence of a reducing agent to obtain a compound of formula 8;


(5) reacting the compound of formula 8 with a compound of formula 9 in the presence of an amide bond formation-inducing reagent to obtain a compound of formula 10; and

(6) deprotecting the amine protecting group of the piperidine ring of the compound of formula 10 in the presence of a base or acid.


[Formula 2]




[Formula 4]




[Formula 5]




[Formula 6]




[Formula 7]




[Formula 8]



[Formula 9]




[Formula 10]



wherein Y represents a halogen atom or C₁-C₄ alkylsulfonate, and Z represents C₁-C₄ alkyl.

- [Claim 9] The method according to claim 8, wherein the amine protecting group-introducing reagent of Step (1) is di-t-butyl dicarbonate or ethyl chloroformate in the presence of a tertiary amine base.
- [Claim 10] The method according to claim 8, wherein the C₁-C₄ alkyl sulfonyl halide of Step (2) is methane sulfonyl chloride, methane sulfonyl bromide or methane sulfonyl fluoride.
- [Claim 11] The method according to claim 8, wherein the reducing agent of Step (4) is triphenylphosphine or lithium aluminum hydride.
- [Claim 12] The method according to claim 8, wherein the amide bond formation-inducing reagent of Step (5) is selected from N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride and 1-hydroxybenzotriazole in the presence of a base, ethyl chloroformate in the presence of a base, or carbodiimidazole in the absence of a base.
- [Claim 13] The method according to claim 8, wherein the base or acid of Step (6) is selected from hydrochloric acid, trifluoroacetic acid or potassium hydroxide.
- [Claim 14] A 5-HT₄ receptor agonist comprising a compound of formula 1:
[Formula 1]



wherein m represents an integer of 1 to 10; and Q represents a heteroaromatic ring or phenyl, wherein the heteroaromatic ring or phenyl is independently substituted by 0, 1, 2, or 3 substituents selected from alkyl, alkoxy, hydroxy, cyano, nitro and halogen; or a pharmaceutically acceptable salt thereof, as an active ingredient.

- [Claim 15] The agonist according to claim 14, wherein m represents an integer of 1 to 5; and Q represents a heteroaromatic ring or phenyl wherein the heteroaromatic ring or phenyl is independently substituted by 0, 1, 2 or 3 substituents selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy and halogen, wherein the heteroaromatic ring being a C₁-C₁₂ aromatic ring

or C₁-C₁₂ bicyclic aromatic ring independently contains 1 to 4 hetero atoms selected from N, O or S.

[Claim 16]

The agonist according to claim 14, wherein the compound of formula 1 or a pharmaceutically acceptable salt thereof is a compound selected from the group consisting of the following compounds:

(1) N-

((1-(3-(1,2,4-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(2) N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(3) N-((1-(3-(indol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(4) N-((1-(3-(2-methylimidazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(5) N-((1-(5-(indol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(6) N-

((1-(5-(1,2,3-triazol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(7) N-

((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(8) N-

((1-(3-(1,2,3-triazol-2-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(9) N-((1-(pyridin-3-ylmethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(10) N-

((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(11) N-((1-(imidazol-2-ylmethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(12) N-

((1-((1-methylpyrrol-2-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(13) N-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(14) N-((1-(4-hydroxybenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-

2-methoxybenzamide,

(15) N-((1-(2-(indol-3-yl)ethyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide,

(16) N-((1-(3-(tetrazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,

(17) N-((1-(5-(1,2,3-triazol-1-yl)pentyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,

(18) N-((1-(3-(1,2,3-triazol-1-yl)propyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,

(19) N-((1-((1-methylindol-3-yl)methyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,

(20) N-((1-(4-fluorobenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride,

(21) N-((1-(4-hydroxybenzyl)piperidin-4-yl)methyl)-4-amino-5-chloro-2-methoxybenzamide hydrochloride and a pharmaceutically acceptable salt thereof

[Claim 17] The agonist according to any one of claims 14 to 16, wherein the 5-HT₄ receptor agonist is a composition for the prevention or treatment of a disease selected from gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive impairment, emesis, migraine, neurological disease, pain, cardiovascular disorder, cardiac failure, cardiac arrhythmia, diabetes or apnea syndrome.

[Claim 18] A method for the prevention, treatment or alleviation of a disease due to attenuated efficacy of a 5-HT₄ receptor, comprising administering a 5-HT₄ receptor agonist containing a compound of formula 1 of any one of claims 1 to 3 or a pharmaceutically acceptable salt thereof as an active ingredient to a mammal including a human in need of 5-HT₄ receptor agonistic effects.

[Claim 19] The method according to claim 18, wherein the 5-HT₄ receptor agonist is a composition for the prevention or treatment of a disease selected from gastroesophageal reflux disease, gastrointestinal disease, gastric motility disorder, non-ulcer dyspepsia, functional dyspepsia, irritable bowel syndrome (IBS), constipation, dyspepsia, esophagitis, gastroesophageal disease, nausea, central nervous system disease, Alzheimer's disease, cognitive impairment, emesis, migraine, neurological disease,

pain, cardiovascular disorder, cardiac failure, cardiac arrhythmia, diabetes or apnea syndrome.