


DLX2, or *MEIS1*. The contacting may comprise contacting a cell that expresses menin. In some embodiments, the method comprises administering a second therapeutic agent. In some embodiments, the contacting takes place *in vivo*. In some embodiments, the contacting takes place *in vitro*.

[0029] In certain aspects, the present disclosure provides a method of treating a disease or condition associated with MLL fusion proteins, comprising administering to a subject in need thereof an effective amount of a compound or salt of Formula (I) or (II). In certain aspects, the present disclosure provides a method of treating a disease or condition in a subject, comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition of a compound or salt of Formula (I) or (II). In some embodiments, the disease or condition comprises a leukemia, hematologic malignancy, solid tumor cancer, prostate cancer, breast cancer, liver cancer, brain tumor, or diabetes. In some embodiments, the leukemia comprises AML, ALL, Mixed Lineage Leukemia or a leukemia with Partial Tandem Duplications of MLL.

[0030] In certain aspects, the present disclosure provides a method of treating a disorder mediated by chromosomal rearrangement on chromosome 11q23 in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound or salt of Formula (I) or (II). In certain aspects, the present disclosure provides a method of treating a disorder mediated by an interaction between menin and another protein, comprising administering to a subject in need thereof a therapeutically effective amount of a compound or salt of Formula (I) or (II). In some embodiments, the subject is a human.

[0031] In certain aspects, the present disclosure provides a kit comprising a pharmaceutical composition described herein and instructions for using the composition to treat a subject suffering from a disease or condition mediated by an interaction between menin and another protein.

[0031a] In certain aspects, the present disclosure provides, a compound of Formula (II-A):



or a pharmaceutically acceptable salt thereof, wherein:

C is selected from C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle;

L² is selected from bond, -O-, -S-, -N(R⁵¹)-, -N(R⁵¹)CH₂-, -C(O)-, -C(O)O-, -OC(O)-, -OC(O)O-, -C(O)N(R⁵¹)-, -C(O)N(R⁵¹)C(O)-, -C(O)N(R⁵¹)C(O)N(R⁵¹)-, -N(R⁵¹)C(O)-, -N(R⁵¹)C(O)N(R⁵¹)-, -N(R⁵¹)C(O)O-, -OC(O)N(R⁵¹)-, -C(NR⁵¹)-, -N(R⁵¹)C(NR⁵¹)-, -C(NR⁵¹)N(R⁵¹)-, -N(R⁵¹)C(NR⁵¹)N(R⁵¹)-, -S(O)₂-, -OS(O)-, -S(O)O-, -S(O)-, -OS(O)₂-, -S(O)₂O-, -N(R⁵¹)S(O)₂-, -S(O)₂N(R⁵¹)-, -N(R⁵¹)S(O)-, -S(O)N(R⁵¹)-, -N(R⁵¹)S(O)₂N(R⁵¹)-, -N(R⁵¹)S(O)N(R⁵¹)-; alkylene, alkenylene, alkynylene, heteroalkylene, heteroalkenylene, and heteroalkynylene, each of which is optionally substituted with one or more R⁵⁰;

L³ is selected from alkylene, alkenylene, and alkynylene, each of which is substituted with one or more R⁵⁶ and optionally further substituted with one or more R⁵⁰;

R^1 , R^2 and R^3 are each independently selected at each occurrence from hydrogen and R^{50} ;

R^A , R^B and R^C are each independently selected at each occurrence from R^{50} , or two R^A groups, two R^B groups or two R^C groups attached to the same atom or different atoms can together optionally form a bridge or ring;

m, n and p are each independently an integer from 0 to 6;

R^{50} is independently selected at each occurrence from:

halogen, $-\text{NO}_2$, $-\text{CN}$, $-\text{OR}^{52}$, $-\text{SR}^{52}$, $-\text{N}(\text{R}^{52})_2$, $-\text{NR}^{53}\text{R}^{54}$, $-\text{S}(=\text{O})\text{R}^{52}$, $-\text{S}(=\text{O})_2\text{R}^{52}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^{52})_2$, $-\text{S}(=\text{O})_2\text{NR}^{53}\text{R}^{54}$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{R}^{52}$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{N}(\text{R}^{52})_2$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{NR}^{53}\text{R}^{54}$, $-\text{C}(\text{O})\text{R}^{52}$, $-\text{C}(\text{O})\text{OR}^{52}$, $-\text{OC}(\text{O})\text{R}^{52}$, $-\text{OC}(\text{O})\text{OR}^{52}$, $-\text{OC}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{OC}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{NR}^{52}\text{C}(\text{O})\text{R}^{52}$, $-\text{NR}^{52}\text{C}(\text{O})\text{OR}^{52}$, $-\text{NR}^{52}\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{NR}^{52}\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{P}(\text{O})(\text{OR}^{52})_2$, $-\text{P}(\text{O})(\text{R}^{52})_2$, $=\text{O}$, $=\text{S}$, $=\text{N}(\text{R}^{52})$;

C_{1-10} alkyl, C_{2-10} alkenyl, and C_{2-10} alkynyl, each of which is independently optionally substituted at each occurrence with one or more substituents selected from halogen, $-\text{NO}_2$, $-\text{CN}$, $-\text{OR}^{52}$, $-\text{SR}^{52}$, $-\text{N}(\text{R}^{52})_2$, $-\text{NR}^{53}\text{R}^{54}$, $-\text{S}(=\text{O})\text{R}^{52}$, $-\text{S}(=\text{O})_2\text{R}^{52}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^{52})_2$, $-\text{S}(=\text{O})_2\text{NR}^{53}\text{R}^{54}$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{R}^{52}$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{N}(\text{R}^{52})_2$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{NR}^{53}\text{R}^{54}$, $-\text{C}(\text{O})\text{R}^{52}$, $-\text{C}(\text{O})\text{OR}^{52}$, $-\text{OC}(\text{O})\text{R}^{52}$, $-\text{OC}(\text{O})\text{OR}^{52}$, $-\text{OC}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{OC}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{NR}^{52}\text{C}(\text{O})\text{R}^{52}$, $-\text{NR}^{52}\text{C}(\text{O})\text{OR}^{52}$, $-\text{NR}^{52}\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{NR}^{52}\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{P}(\text{O})(\text{OR}^{52})_2$, $-\text{P}(\text{O})(\text{R}^{52})_2$, $=\text{O}$, $=\text{S}$, $=\text{N}(\text{R}^{52})$, C_{3-12} carbocycle, and 3- to 12-membered heterocycle; and

C_{3-12} carbocycle and 3- to 12-membered heterocycle,

wherein each C_{3-12} carbocycle and 3- to 12-membered heterocycle in R^{50} is independently optionally substituted with one or more substituents selected from halogen, $-\text{NO}_2$, $-\text{CN}$, $-\text{OR}^{52}$, $-\text{SR}^{52}$, $-\text{N}(\text{R}^{52})_2$, $-\text{NR}^{53}\text{R}^{54}$, $-\text{S}(=\text{O})\text{R}^{52}$, $-\text{S}(=\text{O})_2\text{R}^{52}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^{52})_2$, $-\text{S}(=\text{O})_2\text{NR}^{53}\text{R}^{54}$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{R}^{52}$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{N}(\text{R}^{52})_2$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{NR}^{53}\text{R}^{54}$, $-\text{C}(\text{O})\text{R}^{52}$, $-\text{C}(\text{O})\text{OR}^{52}$, $-\text{OC}(\text{O})\text{R}^{52}$, $-\text{OC}(\text{O})\text{OR}^{52}$, $-\text{OC}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{OC}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{NR}^{52}\text{C}(\text{O})\text{R}^{52}$, $-\text{NR}^{52}\text{C}(\text{O})\text{OR}^{52}$, $-\text{NR}^{52}\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{NR}^{52}\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{P}(\text{O})(\text{OR}^{52})_2$, $-\text{P}(\text{O})(\text{R}^{52})_2$, $=\text{O}$, $=\text{S}$, $=\text{N}(\text{R}^{52})$, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

R^{51} is independently selected at each occurrence from:

hydrogen, $-\text{C}(\text{O})\text{R}^{52}$, $-\text{C}(\text{O})\text{OR}^{52}$, $-\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$,

C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl, each of which is independently optionally substituted at each occurrence with one or more substituents selected from halogen, $-\text{NO}_2$, $-\text{CN}$, $-\text{OR}^{52}$, $-\text{SR}^{52}$, $-\text{N}(\text{R}^{52})_2$, $-\text{NR}^{53}\text{R}^{54}$, $-\text{S}(=\text{O})\text{R}^{52}$, $-\text{S}(=\text{O})_2\text{R}^{52}$, $-\text{S}(=\text{O})_2\text{N}(\text{R}^{52})_2$, $-\text{S}(=\text{O})_2\text{NR}^{53}\text{R}^{54}$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{R}^{52}$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{N}(\text{R}^{52})_2$, $-\text{NR}^{52}\text{S}(=\text{O})_2\text{NR}^{53}\text{R}^{54}$, $-\text{C}(\text{O})\text{R}^{52}$, $-\text{C}(\text{O})\text{OR}^{52}$, $-\text{OC}(\text{O})\text{R}^{52}$, $-\text{OC}(\text{O})\text{OR}^{52}$, $-\text{OC}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{OC}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{NR}^{52}\text{C}(\text{O})\text{R}^{52}$, $-\text{NR}^{52}\text{C}(\text{O})\text{OR}^{52}$, $-\text{NR}^{52}\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{NR}^{52}\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{C}(\text{O})\text{N}(\text{R}^{52})_2$, $-\text{C}(\text{O})\text{NR}^{53}\text{R}^{54}$, $-\text{P}(\text{O})(\text{OR}^{52})_2$, $-\text{P}(\text{O})(\text{R}^{52})_2$, $=\text{O}$, $=\text{S}$, $=\text{N}(\text{R}^{52})$, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle; and

C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle,

wherein each C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle in R⁵¹ is independently optionally substituted with one or more substituents selected from halogen, -NO₂, -CN, -OR⁵², -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

R⁵² is independently selected at each occurrence from hydrogen; and C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, 1- to 6-membered heteroalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted by halogen, -CN, -NO₂, -NH₂, -NHCH₃, -NHCH₂CH₃, =O, -OH, -OCH₃, -OCH₂CH₃, C₃₋₁₂ carbocycle, or 3- to 6-membered heterocycle;

R⁵³ and R⁵⁴ are taken together with the nitrogen atom to which they are attached to form a heterocycle, optionally substituted with one or more R⁵⁰;

R⁵⁶ is independently selected at each occurrence from:

-NO₂, -OR⁵⁹, -SR⁵², -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle,

wherein each C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, and C₂₋₁₀ alkynyl in R⁵⁶ is independently optionally substituted at each occurrence with one or more substituents selected from halogen, -NO₂, -CN, -OR⁵⁹, -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

wherein each C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle in R⁵⁶ is independently optionally substituted with one or more substituents selected from halogen, -NO₂, -CN, -OR⁵², -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

$P(O)(OR^{52})_2$, $-P(O)(R^{52})_2$, $=O$, $=S$, $=N(R^{52})$, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and

further wherein R^{56} optionally forms a bond to ring C; and

R^{59} is independently selected at each occurrence from C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl, 1- to 6-membered heteroalkyl, C_{3-12} carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted by halogen, $-CN$, $-NO_2$, $-NH_2$, $-NHCH_3$, $-NHCH_2CH_3$, $=O$, $-OH$, $-OCH_3$, $-OCH_2CH_3$, C_{3-12} carbocycle, or 3- to 6-membered heterocycle, wherein for a compound or salt of Formula (II-A), when R^{56} is $-CH_3$, L^3 is not further substituted with $-OH$, $-NH_2$, or $-CN$.

[0031b] In certain aspects, the present disclosure provides a pharmaceutical composition comprising a compound as described herein and a pharmaceutically acceptable carrier.

[0031c] In certain aspects, the present disclosure provides a method of inhibiting an interaction of menin with one or more of MLL1, MLL2, a MLL fusion protein, and a MLL Partial Tandem Duplication, comprising contacting menin with an effective amount of a compound as described herein.

[0031d] In certain aspects, the present disclosure provides a use of a compound or salt as described herein in the manufacture of a medicament for the treatment of leukemia, a hematologic malignancy, a solid tumor cancer, prostate cancer, breast cancer, liver cancer, a brain tumor, or diabetes.

INCORPORATION BY REFERENCE

[0032] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0033] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0034] **FIG. 1** is an amino acid sequence of human menin, isoform 1 (SEQ ID NO: 1).

[0035] **FIG. 2** is an amino acid sequence of human menin, isoform 2 (SEQ ID NO: 2).

[0036] **FIG. 3** is an amino acid sequence of human menin, isoform 3 (SEQ ID NO: 3).

[0037] **FIG. 4** depicts the change in volume of MV4;11 tumors in vehicle and compound treated mice.

[0038] **FIG. 5** depicts the luminescence of MV4;11-luc tumors in vehicle and compound treated xenotransplantation mouse models of MLL leukemia after 6 days of treatment.

What is claimed is:

-
- (II-A).

-168-

carbocycle, and 3- to 12-membered heterocycle; and

C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle,

wherein each C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle in R⁵⁰ is independently optionally substituted with one or more substituents selected from halogen, -NO₂, -CN, -OR⁵², -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

R⁵¹ is independently selected at each occurrence from:

hydrogen, -C(O)R⁵², -C(O)OR⁵², -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴;

C₁₋₆ alkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl, each of which is independently optionally substituted at each occurrence with one or more substituents selected from halogen, -NO₂, -CN, -OR⁵², -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle; and

C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle,

wherein each C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle in R⁵¹ is independently optionally substituted with one or more substituents selected from halogen, -NO₂, -CN, -OR⁵², -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

R⁵² is independently selected at each occurrence from hydrogen; and C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, 1- to 6-membered heteroalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted by halogen, -CN, -NO₂, -NH₂, -NHCH₃, -NHCH₂CH₃, =O, -OH, -OCH₃, -OCH₂CH₃, C₃₋₁₂ carbocycle, or 3- to 6-membered heterocycle;

R⁵³ and R⁵⁴ are taken together with the nitrogen atom to which they are attached to form a heterocycle, optionally substituted with one or more R⁵⁰;

R⁵⁶ is independently selected at each occurrence from:

-NO₂, -OR⁵⁹, -SR⁵², -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -

C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle,

wherein each C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, and C₂₋₁₀ alkynyl in R⁵⁶ is independently optionally substituted at each occurrence with one or more substituents selected from halogen, -NO₂, -CN, -OR⁵⁹, -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle;

wherein each C₃₋₁₂ carbocycle and 3- to 12-membered heterocycle in R⁵⁶ is independently optionally substituted with one or more substituents selected from halogen, -NO₂, -CN, -OR⁵², -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl; and

further wherein R⁵⁶ optionally forms a bond to ring C; and


R⁵⁹ is independently selected at each occurrence from C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, 1- to 6-membered heteroalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted by halogen, -CN, -NO₂, -NH₂, -NHCH₃, -NHCH₂CH₃, =O, -OH, -OCH₃, -OCH₂CH₃, C₃₋₁₂ carbocycle, or 3- to 6-membered heterocycle,

wherein for a compound or salt of Formula (II-A), when R⁵⁶ is -CH₃, L³ is not further substituted with -OH, -NH₂, or -CN.

2. The compound of claim 1, wherein R^C is selected from -C(O)R⁵², -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², =O, C₁₋₃ alkyl, and C₁₋₃ haloalkyl, or two R^C groups attached to different atoms can together form a C₁₋₃ bridge.

3. The compound of claim 1 or 2, wherein C is 5- to 12-membered saturated heterocycle, wherein the heterocycle comprises at least one nitrogen atom.

4. The compound of claim 1, wherein C is selected from:



wherein R⁵⁷ is selected from -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵²; and C₁₋₁₀ alkyl substituted with one or more substituents selected from -S(=O)R⁵², -

$S(=O)_2R^{52}$, $-S(=O)_2N(R^{52})_2$, $-S(=O)_2NR^{53}R^{54}$, and $-NR^{52}S(=O)_2R^{52}$.

5. The compound of any one of the preceding claims, wherein:

R^2 is selected from hydrogen, halogen, $-OH$, $-OR^{52}$, $-NH_2$, $-N(R^{52})_2$, $-CN$, C_{1-3} alkyl, C_{1-3} alkyl- $N(R^{52})_2$, C_{1-3} haloalkyl, C_{2-3} alkenyl, and C_{2-3} alkynyl; and

6. R^3 is selected from hydrogen, halogen, $-OH$, $-N(R^{52})_2$, $-CN$, $-C(O)OR^{52}$, C_{1-3} alkyl, and C_{1-3} haloalkyl. The compound of any one of the preceding claims, wherein R^1 is C_{1-3} haloalkyl.


7. The compound of any one of the preceding claims, wherein m is 0.

8. The compound of any one of the preceding claims, wherein n is an integer from 1 to 3.

9. The compound of any one of the preceding claims, wherein L^2 is C_{1-4} alkylene, optionally substituted with one or more R^{50} .

10. The compound of any one of the preceding claims, wherein L^3 is substituted with $=O$, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-3} alkyl(cyclopropyl), C_{1-3} alkyl($NR^{52}C(O)R^{52}$) or $-O(C_{1-6}$ alkyl).

11. The compound of any one of claims 1 to 9, wherein L^3 is selected from



12. The compound of claim 11, wherein R^{56} is methyl.

13. The compound of claim 1, wherein:

L^2 is selected from $-O-$, $-S-$, $-NH-$, and $-CH_2-$;

L^3 is selected from C_{1-6} alkylene, C_{2-6} alkenylene, and C_{2-6} alkynylene, each of which is substituted with one or more R^{56} and optionally further substituted with one or more R^{50} ;

R^A , R^B and R^C are each independently selected at each occurrence from R^{50} , or two R^A groups, two R^B groups or two R^C groups attached to the same atom or different atoms can together optionally form a bridge or ring;

m is an integer from 0 to 3;

n is an integer from 1 to 3;

p is an integer from 0 to 6;

R^{56} is independently selected at each occurrence from:

$-OR^{59}$, $=O$, C_{1-10} alkyl, C_{2-10} alkenyl, and C_{2-10} alkynyl,

wherein each C_{1-10} alkyl, C_{2-10} alkenyl, and C_{2-10} alkynyl in R^{56} is independently optionally substituted at each occurrence with one or more substituents selected from halogen, $-NO_2$, $-CN$, $-OR^{59}$, $-SR^{52}$, $-N(R^{52})_2$, $-NR^{53}R^{54}$, $-S(=O)R^{52}$, $-S(=O)_2R^{52}$, $-S(=O)_2N(R^{52})_2$, $-S(=O)_2NR^{53}R^{54}$, $-NR^{52}S(=O)_2R^{52}$, $-NR^{52}S(=O)_2N(R^{52})_2$, $-NR^{52}S(=O)_2NR^{53}R^{54}$, $-C(O)R^{52}$, $-C(O)OR^{52}$, $-OC(O)R^{52}$, $-OC(O)OR^{52}$, $-OC(O)N(R^{52})_2$, $-OC(O)NR^{53}R^{54}$, $-NR^{52}C(O)R^{52}$, $-NR^{52}C(O)OR^{52}$, $-NR^{52}C(O)N(R^{52})_2$, $-NR^{52}C(O)NR^{53}R^{54}$, $-C(O)N(R^{52})_2$, $-C(O)NR^{53}R^{54}$, $-P(O)(OR^{52})_2$, $-P(O)(R^{52})_2$, $=O$, $=S$, $=N(R^{52})$, C_{3-12} carbocycle, and 3- to 12-membered heterocycle;

wherein each C_{3-12} carbocycle and 3- to 12-membered heterocycle in R^{56} is independently optionally substituted with one or more substituents selected from halogen,

-NO₂, -CN, -OR⁵², -SR⁵², -N(R⁵²)₂, -NR⁵³R⁵⁴, -S(=O)R⁵², -S(=O)₂R⁵², -S(=O)₂N(R⁵²)₂, -S(=O)₂NR⁵³R⁵⁴, -NR⁵²S(=O)₂R⁵², -NR⁵²S(=O)₂N(R⁵²)₂, -NR⁵²S(=O)₂NR⁵³R⁵⁴, -C(O)R⁵², -C(O)OR⁵², -OC(O)R⁵², -OC(O)OR⁵², -OC(O)N(R⁵²)₂, -OC(O)NR⁵³R⁵⁴, -NR⁵²C(O)R⁵², -NR⁵²C(O)OR⁵², -NR⁵²C(O)N(R⁵²)₂, -NR⁵²C(O)NR⁵³R⁵⁴, -C(O)N(R⁵²)₂, -C(O)NR⁵³R⁵⁴, -P(O)(OR⁵²)₂, -P(O)(R⁵²)₂, =O, =S, =N(R⁵²), C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl; and

further wherein R⁵⁶ optionally forms a bond to ring C; and

R⁵⁹ is independently selected at each occurrence from C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, 1- to 6-membered heteroalkyl, C₃₋₁₂ carbocycle, and 3- to 12-membered heterocycle, each of which is optionally substituted by halogen, -CN, -NO₂, -NH₂, -NHCH₃, -NHCH₂CH₃, =O, -OH, -OCH₃, -OCH₂CH₃, C₃₋₁₂ carbocycle, or 3- to 6-membered heterocycle.

14. The compound of claim 13, wherein C is substituted with -S(=O)₂R⁵⁸, -S(=O)₂N(R⁵²)₂, or -S(=O)₂NR⁵³R⁵⁴.


15. The compound of claim 13, wherein the compound is a stereoisomer provided in at least 90% enantiomeric excess.

16. A pharmaceutical composition comprising a compound of any one of the preceding claims and a pharmaceutically acceptable carrier.


17. A method of inhibiting an interaction of menin with one or more of MLL1, MLL2, a MLL fusion protein, and a MLL Partial Tandem Duplication, comprising contacting menin with an effective amount of a compound of any one of claims 1 to 15.

18. Use of a compound or salt of any one of claims 1 to 15 in the manufacture of a medicament for the treatment of leukemia, a hematologic malignancy, a solid tumor cancer, prostate cancer, breast cancer, liver cancer, a brain tumor, or diabetes.

19. The compound of claim 1, wherein the compound has the structure of Formula (II-C):



20. The compound of claim 1, wherein the compound has the structure of Formula (II-F) or Formula (II-H):



21. The compound of claim 1, wherein the compound is selected from the group consisting of:

