

The ability of $\gamma\delta$ T-cells to recognize an antigen on diseased cells directly and to exhibit inherent ability to kill tumor cells renders $\gamma\delta$ T-cells an attractive therapeutic tool. Adoptive transfer of V γ 9V δ 2 T cells has yielded limited objective clinical responses for investigational treatment of cancer (Kondo et al, Cytotherapy, 10:842- 856, 2008; Lang et al, Cancer Immunology, Immunotherapy: CII, 60: 1447-1460, 2011; Nagamine et al, 2009; Nicol et al, British Journal of Cancer, 105:778-786, 2011; Wilhelm et al, Blood. 2003 Jul 1;102(1):200-6), indicating the need to isolate and test clinically new $\gamma\delta$ T-cell populations.

The ability to selectively expand $\gamma\delta$ T-cell subset populations having potent anti-tumor activity with improved purity and in clinically-relevant levels is highly desirable. Although antibodies and cytokine cocktails have been used to propagate a more diverse set of $\gamma\delta$ T cells, activation of specific $\gamma\delta$ T-cell subsets to sufficient purity and clinically-relevant levels, was not achieved (Dokouhaki et al, 2010; Kang et al, 2009; Lopez et al, 2000; Kress, 2006). Therefore, clinically-relevant methods of expanding specific $\gamma\delta$ T cell subsets *ex vivo*, and the cells produced thereby, are greatly needed.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on June 21, 2017 is named ADC-0002-WO_SL.txt and is 207,179 Kilobytes in size.

SUMMARY OF THE INVENTION

The present invention, in one aspect, provides *ex vivo* methods for producing enriched $\gamma\delta$ T-cell populations. The enriched $\gamma\delta$ T-cell populations can be produced from isolated mixed cell populations by a method comprising contacting the mixed cell population, or a purified fraction thereof, with one or more agents:

- i) which selectively expand δ 1 T-cells by binding to an epitope specific of a δ 1 TCR,
- ii) which selectively expand δ 2 T-cells by binding to an epitope specific of a δ 2 TCR,
- iii) which selectively expand δ 1 and δ 4 T-cells by binding to an epitope specific of a δ 1 TCR and a δ 4 TCR, or

FIG. 4 depicts a graph illustrating growth of $\gamma\delta 1$ and $\gamma\delta 2$ lymphocytes isolated from colon adenocarcinoma metastasis to liver (TIL 1) and renal tumors (TIL 2) and have been shown to express CCR4 and CCR7.

FIG. 5 depicts a graph illustrating $\gamma\delta$ T-cells growth in serum-containing and serum-free media.

FIG. 6 depicts a graph illustrating anti- $\gamma\delta$ TCR antibody blocking experiments with 5A6.E9, B1, TS8.2, 15D, B3, B6, TS-1, $\gamma 3.20$, IMMU510, or 11F2.

FIG. 7 depicts a graph illustrating anti- $\gamma\delta$ TCR antibody blocking experiments with 5A6.E9, B1, TS8.2, 15D, B3, B6, TS-1, $\gamma 3.20$, IMMU510, or 11F2.

FIG. 8 depicts a competition experiment with anti-TCR V $\delta 1$ TS-1 antibody.

FIG. 9 depicts a competition experiment with anti-TCR V $\delta 1$ TS8.2 antibody.

FIG. 10 depicts a graph illustrating activation and expansion of $\delta 1$ T-cells from PBMC.

FIG. 11 depicts a graph illustrating activation and expansion of $\delta 2$ T-cells from PBMC.

FIG. 12 depicts a graph illustrating fold expansion of $\delta 1$ T-cells from PBMC.

FIG. 13 depicts a graph illustrating fold expansion of $\delta 2$ T-cells from PBMC.

FIG. 14 depicts the human V $\delta 1$ amino acid sequence (SEQ ID NO: 133). CDR1, CDR3, and CDR3 regions are underlined. FIG. 14 also discloses the nucleotide sequence as SEQ ID NO: 132.

FIG. 15 depicts the human V $\delta 2$ amino acid sequence (SEQ ID NO: 135). CDR1, CDR3, and CDR3 regions are underlined. FIG. 15 discloses SEQ ID NOS 135, 134, 137, and 136, respectively, in order of appearance.

FIG. 16 depicts activation of PBMCs with TS-1 or TS8.2 antibodies resulting in significant and specific expansion of V $\delta 1^+$ T cells. **(A)** MAbs were directly coated (1 $\mu\text{g/mL}$) (TS1, TS8.2, B6, and 15D) or captured (0.1 $\mu\text{g/mL}$) with goat-anti-mouse Fc (5 $\mu\text{g/mL}$) (TS1Fc, TS8.2Fc, B6Fc and 15DFc) in 24-well plates. PBMCs were plated at 10^6 cells/mL in RPMI with 10% FBS and 100 IU/mL IL-2. On day 7, cells were transferred to new plates without antibodies and further expanded until day 14. Culture media was replenished every 2-3 days. Data depicts expansion of V $\delta 1^+$ cells over 14 days; **(B)** Same culture as A, showing percentage of V $\delta 1^+$ cells on day 14 and day 0 (d0); **(C)** V $\delta 1^+$ cells were expanded from isolated PBMCs from a different donor as follows. MAbs were directly coated (1 $\mu\text{g/mL}$) (TS1, TS8.2, B6 and 15D) or captured (0.1 $\mu\text{g/mL}$) with goat-anti-mouse Fc (5 $\mu\text{g/mL}$) (TS1Fc, TS8.2Fc, B6Fc and 15DFc) in 24-well plates. PBMCs were plated at 10^6 cells/mL in RPMI with 10% FBS and 100 IU/mL IL-2. Cells were transferred to new plate without

to the V δ 1J1 chain at the selected positions. BE-13 refers to δ 1J region from a T cell leukemia cell line expressing δ 1 γ 8 TCR (DSMZ accession #ACC 396). Replacement of a single amino acid residue at position Lys120 by Thr or Ala in the V δ 1J1 and V δ 1J2 region completely abolished binding of TS-1 and TS8.2 MAbs indicating that this amino acid in the V δ 1J1 and V δ 1J2 region is
5 contributes to the binding of TS-1 and TS8.2. Replacement of a single amino acid residue at position Thr120 by Lys in the V δ 1J3 region resulted in a gain of binding of TS-1 and TS8.2 MAbs further indicating that this amino acid in the V δ 1J1 and V δ 1J2 region contributes to the binding of TS-1 and TS8.2. FIG. 24 discloses SEQ ID NOS 138-143, respectively, in order of appearance.

FIG. 25 depicts that point mutation in V δ 1J1 from Lys120 to Thr or Ala resulted in the loss
10 of binding by TS-1 and TS8.2 MAbs.

FIG. 26 depicts of human V δ 1 protein sequence (SEQ ID NO: 144) and the six mutated human V δ 1 sequences (SEQ ID NOS 145-150, respectively, in order of appearance) based on the differences between human V δ 1 and the bovine V δ 1 amino acid sequence (GenBank:AFP25162.1). Mutation are shown in bold.

FIG. 27 depicts loss of binding of TS-1 and TS8.2 antibodies to the human V δ 1 mutated chains.
15

FIG. 28 depicts binding of the B6 MAb to different V δ 2/ γ chain pairings (γ 3, γ 8, γ 9).

FIG. 29 depicts the protein sequence alignment of human (IMGT Human TRDV2) (SEQ ID NO: 151) and Rhesus monkey (GenBank: AY190028.1) (SEQ ID NO: 152) V δ 2 variable regions
20 and the changes made in V δ 2 CDR1 (G35S), CDR2 (D65G) and CDR3 (C104S).

FIG. 30 depicts loss of binding of 15D to V δ 2 (G35S) mutated at CDR1.

FIG. 31 depicts characterization of highly enriched V δ 1+, V δ 2+ and $\alpha\beta$ T cell cultures using TS8.2, B6 and IP26 antibodies for δ 1, δ 2 and $\alpha\beta$, respectively. The populations derived from PBMC were expanded with a combination of TS-1 and TS8.2 to expand δ 1 cells, or 15D and B6 to expand
25 δ 2 cells. The expanded culture was depleted of $\alpha\beta$ T cells using IP26 MicroBeads. Positively selected $\alpha\beta$ T cells were also collected ($\alpha\beta$ enriched).

FIG. 32 depicts cytotoxicity of V δ 1+, V δ 2+ populations toward solid tumor (BxPC3, SKMEL5) and plasmacytoma (RPMI8226) cell lines.

FIG. 33 depicts heavy-chain framework and complementarity determining region amino acid sequences of δ 1-specific MAbs. FIG. 33 discloses SEQ ID NOS 153-179, respectively, in order of
30 appearance.

FIG. 34 depicts light-chain framework and complementarity determining region amino acid

sequences of the δ 1-specific MAbs described in **FIG. 33**. FIG. 34 discloses SEQ ID NOS 180-206, respectively, in order of appearance.

FIG. 35 depicts heavy-chain framework and complementarity determining region amino acid sequences of δ 2-specific MAbs. FIG. 35 discloses SEQ ID NOS 207-215, respectively, in order of appearance.

FIG. 36 depicts light-chain framework and complementarity determining region amino acid sequences of δ 2-specific MAbs. FIG. 36 discloses SEQ ID NOS 216-225, respectively, in order of appearance.

FIG. 37 depicts cross-reactivity of the indicated δ 1-specific antibodies with other $\gamma\delta$ TCRs as determined by ELISA.

FIG. 38 depicts an alignment of V δ 1 deduced amino acid sequence between human (SEQ ID NO: 133) and dolphin (SEQ ID NO: 226).

FIG. 39 depicts the percentage of proliferating V δ 1 cells (A) or V δ 2 cells (B) obtained by a $\gamma\delta$ T cell expansion method described herein.

FIG. 40 depicts the fold expansion of V δ 1 cells (A) or V δ 2 cells (B) from an isolated mixed cell population obtained by a $\gamma\delta$ T cell expansion method described herein.

FIG. 41 depicts phenotype data from an isolated mixed cell population obtained by a $\gamma\delta$ T cell expansion method described herein. Cell phenotypes are determined by CD45RA and CD27 expression using flow cytometry analysis.

FIG. 42 illustrates δ 1 T cells activated and expanded according to $\gamma\delta$ T cell expansion methods described herein can show robust cytotoxicity against tumor cell lines.

FIG. 43 illustrates V δ 1 fold expansion and purity on day 19 achieved by a method of the present invention.

FIG. 44 illustrates V δ 1 fold expansion and accumulative doubling time on day 19 achieved by a method of the present invention.

FIG. 45 illustrates V δ 1 fold expansion and accumulative doubling time from day 14 to day 19 achieved by a method of the present invention.

FIG. 46 illustrates expression of an anti-CD20 chimeric antigen receptor (CAR) on δ 1 T cells. Expanded cells were stained with V δ 1 specific antibody (R9.12) and anti-Rituximab FITC conjugate on Day 7 post-transduction with CAR retroviral construct. Up to 35% of V δ 1 cells were CD20 CAR+.

FIG. 47A-B depicts epitope binding specificity data for δ -1 specific $\gamma\delta$ T cell activators.

FIG. 48 depicts epitope binding specificity data for δ -2 specific $\gamma\delta$ T cell activators.

determining spatial conformation of amino acids with methods that include, e.g., x-ray crystallography and 2-dimensional nuclear magnetic resonance. Some epitope mapping methods, such as, x-ray analyses of crystals of antigen:antibody complexes can provide atomic resolution of the epitope. In other cases, computational combinatorial methods for epitope mapping can be employed to model a potential epitope based on the sequence of the antibody, such as TS-1 antibody or TS8.2 antibody. In such cases, the antigen binding portion of the antibody is sequenced, and computation models are used to reconstruct and predict a potential binding site of the antibody.

In some cases the disclosure provides a method of determining an epitope of a $\gamma\delta$ T-cell receptor, comprising: (a) preparing a library of epitopes from the $\gamma\delta$ T-cell receptor; (b) contacting the library of epitopes with an antibody; and (b) identifying the amino acid sequence of at least one epitope in the library of epitopes that is bound by the antibody. In some cases, the antibody is selected from the group consisting of, TS8.2, 15D, B6, TS-1, and R9.12, antibodies. In one instance, the antibody is attached to a solid support. The library of epitopes can comprise sequences that correspond to continuous and discontinuous epitopes of a T-cell receptor, such as a γ TCR or a δ TCR. In some cases, the library of epitopes comprises fragments from a $\gamma\delta$ T-cell receptor ranging from about 10 amino acids to about 30 amino acids in length, from about 10 amino acids to about 20 amino acids in length, or from about 5 amino acids to about 12 amino acids in length. In some cases, the antibody is labeled and the label is a radioactive molecule, a luminescent molecule, a fluorescent molecule, an enzyme, or biotin.

δ 1 Epitope Bins

In some embodiments, the epitope, that is (e.g., specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 47-70 of human V δ 1 (SKEMIFLIRQGSDEQNA) (SEQ ID NO: 17) and J1 (TDKLIFGKGTRVTVEP) (SEQ ID NO: 18) or J2 (LTAQLFFGKGTQLLIVEP) (SEQ ID NO: 19), wherein the activating agent does not bind an epitope containing a K120T mutation in J1 or J2. This epitope is referred to herein as a Bin 1 δ 1 epitope. Exemplary activating agents that bind a Bin 1 δ 1 epitope include but are not limited to TS-1; and δ 1-18. A human V δ 1 J region can vary due to (D)(J) recombination, an exemplary V δ 1 J region of a δ 1 chain of a $\gamma\delta$ TCR having the Bin 1 δ 1 epitope is SKEMIFLIRQGSDEQNAKSGRYSVNFKKAASVALTISALQLEDSAKYFCALGTGVRGLQD TDKLIFGKGTRVTVEP (SEQ ID NO: 20); and another exemplary V δ 1 J region Bin 1 δ 1 epitope of

a $\delta 1$ chain of a $\gamma\delta$ TCR is

SKEMIFLIRQGSDEQNAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGEAPSAWGK
HLTAQLFFGKGTLIVEP (SEQ ID NO: 21).

In some cases, activating agents that bind a Bin 1 $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR
5 having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVE
P (SEQ ID NO: 22); but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
10 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDSWDTRQMFFGTGIK
LFVEP (SEQ ID NO: 23).

The activating agents that bind a Bin 1 $\delta 1$ epitope can also bind a $\delta 1$ chain of a $\gamma\delta$ TCR
having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
15 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGEAPSAWGKHLTAQLFFGKGTLI
VEP (SEQ ID NO: 24).

In some embodiments, the epitope, that is (*e.g.*, specifically) recognized by an activating
agent of interest is an epitope comprised of amino acids 47-70 of human V $\delta 1$ (SKEMIFLIRQ
GSDEQNA) (SEQ ID NO: 17) and J1 (TDKLIFGKGTRVTVEP) (SEQ ID NO: 18), wherein the
20 activating agent does not bind an epitope containing a K120T mutation in J1. This epitope is
referred to herein as a Bin 1b $\delta 1$ epitope. Exemplary activating agents that bind a Bin 1b $\delta 1$ epitope
include but are not limited to $\delta 1$ -37. An exemplary V $\delta 1$ J region of a $\delta 1$ chain of a $\gamma\delta$ TCR having
the Bin 1b $\delta 1$ epitope is

SKEMIFLIRQGSDEQNAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQD
25 TDKLIFGKGTRVTVEP (SEQ ID NO: 20).

In some cases, activating agents that bind a Bin 1b $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR
having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVE
30 P (SEQ ID NO: 22); but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDSWDTRQMFFGTGIK
LFVEP (SEQ ID NO: 23).

5 The activating agents that bind a Bin 1b $\delta 1$ epitope also do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGEAPSAWGKHLTAQLFFGKGQTQLI
VEP (SEQ ID NO: 24).

10 In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 11-21 of human V $\delta 1$ (VSMPVRKAVTL) (SEQ ID NO: 26). This epitope is referred to herein as a Bin 2 $\delta 1$ epitope. Exemplary activating agents that bind a Bin 2 $\delta 1$ epitope include but are not limited to and $\delta 1$ -285.

In some cases, activating agents that bind a Bin 2 $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

15 AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQDKLIFGKGTRVTVE
P (SEQ ID NO: 22); but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQVQRAMSSQLGEAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQDKLIFGKGTRVTVE
20 PRSQPHTKPSVFVMKNGTNVACLVEF (SEQ ID NO: 25).

In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 11-21 of human V $\delta 1$ (VSMPVRKAVTL) (SEQ ID NO: 26), wherein the activating agent that binds this epitope does not bind an epitope containing a mutation of R16 in V $\delta 1$, such as an R16N mutation. This epitope is referred to herein
25 as a Bin 2b $\delta 1$ epitope. Exemplary activating agents that bind a Bin 2b $\delta 1$ epitope include but are not limited to R9.12.

In some cases, activating agents that bind a Bin 2b $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
30 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQDKLIFGKGTRVTVE

P (SEQ ID NO: 22); but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVNKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVE
P (SEQ ID NO: 27); and/or do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

5 AQKVTQVQGRAMSSQLGEAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVE
PRSQPHTKPSVFMKNGTNVACLVEF (SEQ ID NO: 25).

In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 11-21 of human V $\delta 1$ (VSMPVRKAVTL) (SEQ ID NO: 26), wherein the activating agent that binds this epitope also binds (cross-reacts) with $\delta 3$, $\delta 4$, and $\delta 5$ $\gamma\delta$ TCRs. This epitope is referred to herein as a Bin 2c $\delta 1$ epitope. Exemplary activating agents that bind a Bin 2c $\delta 1$ epitope include but are not limited to $\delta 1$ -39.

In some cases, activating agents that bind a Bin 2c $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

15 AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVE
P (SEQ ID NO: 22); but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQVQGRAMSSQLGEAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVE
20 PRSQPHTKPSVFMKNGTNVACLVEF (SEQ ID NO: 25).

In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 80-95 of human V $\delta 1$ (FKKAAKSVALTISALQ) (SEQ ID NO: 28) or 70 to 95 of human V $\delta 1$ (AKSGRYSVNFKKAAKSVALTISALQ) (SEQ ID NO: 29). This epitope is referred to herein as a Bin 3 $\delta 1$ epitope. Exemplary activating agents that bind a Bin 3 $\delta 1$ epitope include but are not limited to $\delta 1$ -08; and $\delta 1$ -23.

In some cases, activating agents that bind a Bin 3 $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
30 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVE

P (SEQ ID NO: 22)

but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQVQRAMSSQLGEAVTLSCQYETSLSWYDIFWYKQLPSGEMTFLIHQISSDQ
NAKNGRYSVNFQERHKFISLTISALQLEDSAKYFCALGTGVRGLQDQDKLIFGKGTRVTVEP
5 RSQPHTKPSVFMKNGTNVACLVKEFYPKD (SEQ ID NO: 30).

In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 1-11 of human V $\delta 1$ (AQKVTQAQSSV) (SEQ ID NO: 227) and J1 or J2. This epitope is referred to herein as a Bin 4 $\delta 1$ epitope. In some cases, the Bin 4 $\delta 1$ epitope binding activating agent does not bind an epitope containing a K120T
10 mutation in J1 or J2. Exemplary activating agents that bind a Bin 4 $\delta 1$ epitope include but are not limited to $\delta 1$ -35; and $\delta 1$ -203.

In some cases, activating agents that bind a Bin 4 $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
15 NAKSGRYSVNFKKAASVALTISALQLEDSAKYFCALGTGVRGLQDQDKLIFGKGTRVTVE
P (SEQ ID NO: 22); but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAASVALTISALQLEDSAKYFCALGTGVRGLQDSWDTRQMFFGTGIK
LFVEP (SEQ ID NO: 23).

20 In some cases, activating agents that bind a Bin 4 $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAASVALTISALQLEDSAKYFCALGEAPSAWGKHLTAQLFFGKGQTQLI
VEP (SEQ ID NO: 24).

25 In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 28-47 of human V $\delta 1$ (SWWSYYIFWYKQLPS) (SEQ ID NO: 31) and J1. This epitope is referred to herein as a Bin 5 $\delta 1$ epitope. Exemplary activating agents that bind a Bin 5 $\delta 1$ epitope include but are not limited to $\delta 1$ -113; $\delta 1$ -155; $\delta 1$ -183; $\delta 1$ -191; $\delta 1$ -278; and $\delta 1$ -282. An exemplary V $\delta 1$ J1 region of a $\delta 1$ chain of a
30 $\gamma\delta$ TCR having the Bin 5 $\delta 1$ epitope is

SWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVEP (SEQ ID NO: 32).

In some cases, activating agents that bind a Bin 5 $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

5 A Q K V T Q A Q S S V S M P V R K A V T L N C L Y E T S W W S Y Y I F W Y K Q L P S K E M I F L I R Q G S D E Q
N A K S G R Y S V N F K K A A K S V A L T I S A L Q L E D S A K Y F C A L G T G V R G L Q D T D K L I F G K G T R V T V E
P (SEQ ID NO: 22); but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

10 A Q K V T Q A Q S S V S M P V R K A V T L N C L Y E T S W W S Y Y I F W Y K Q L P S K E M I F L I R Q G S D E Q
N A K S G R Y S V N F K K A A K S V A L T I S A L Q L E D S A K Y F C A L G E A P S A W G K H L T A Q L F F G K G T Q L I
V E P (SEQ ID NO: 24).

In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 21-28 of human V $\delta 1$ (LNCLYETS) (SEQ ID NO: 33) and J1. This epitope is referred to herein as a Bin 6 $\delta 1$ epitope. Exemplary activating agents that bind a Bin 6 $\delta 1$ epitope include but are not limited to TS 8.2 and $\delta 1$ -143. An exemplary V $\delta 1$ J1 region of a $\delta 1$ chain of a $\gamma\delta$ TCR having the Bin 6 $\delta 1$ epitope is

15 L N C L Y E T S W W S Y Y I F W Y K Q L P S K E M I F L I R Q G S D E Q N A K S G R Y S V N F K K A A K S V A L T I S A L
Q L E D S A K Y F C A L G T G V R G L Q D T D K L I F G K G T R V T V E P (SEQ ID NO: 34).

In some cases, activating agents that bind a Bin 6 $\delta 1$ epitope bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

20 A Q K V T Q A Q S S V S M P V R K A V T L N C L Y E T S W W S Y Y I F W Y K Q L P S K E M I F L I R Q G S D E Q
N A K S G R Y S V N F K K A A K S V A L T I S A L Q L E D S A K Y F C A L G T G V R G L Q D T D K L I F G K G T R V T V E
P (SEQ ID NO: 22); but do not bind a $\delta 1$ chain of a $\gamma\delta$ TCR having the sequence of:

25 A Q K V T Q A Q S S V S M P V R K A V T L N C L Y E T S W W S Y Y I F W Y K Q L P S K E M I F L I R Q G S D E Q
N A K S G R Y S V N F K K A A K S V A L T I S A L Q L E D S A K Y F C A L G T G V R G L Q D S W D T R Q M F F G T G I K
L F V E P (SEQ ID NO: 23).

In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 47-70 of human V $\delta 1$ (SKEMIFLIRQGSDEQNA) (SEQ ID NO: 17) and J1 or J2. This epitope is referred to herein as a Bin 7 $\delta 1$ epitope. Exemplary activating agents that bind a Bin 7 $\delta 1$ epitope include but are not

30 limited to $\delta 1$ -149; $\delta 1$ -253, and $\delta 1$ -257. An exemplary V $\delta 1$ J region of a $\delta 1$ chain of a $\gamma\delta$ TCR

having the Bin 7 δ 1 epitope is

SKEMIFLIRQGSDEQNAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQD
TDKLIFGKGTRVTVEP (SEQ ID NO: 20); and another exemplary V δ 1 J region Bin 7 δ 1 epitope of
a δ 1 chain of a $\gamma\delta$ TCR is

5 SKEMIFLIRQGSDEQNAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGEAPSAWGK
HLTAQLFFGKGQTQLIVEP (SEQ ID NO: 21).

In some cases, activating agents that bind a Bin 7 δ 1 epitope bind a δ 1 chain of a $\gamma\delta$ TCR
having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
10 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQD
TDKLIFGKGTRVTVEP (SEQ ID NO: 22); but do not bind a δ 1 chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDSWDTRQMFFGTGIK
LFVEP (SEQ ID NO: 23).

15 In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating
agent of interest is an epitope comprised of amino acids 70-80 of human V δ 1 (AKSGRYSVNF)
(SEQ ID NO: 35) and J1 or J2. This epitope is referred to herein as a Bin 8 δ 1 epitope. Exemplary
activating agents that bind a Bin 8 δ 1 epitope include but are not limited to δ 1-192. An exemplary
V δ 1 J region of a δ 1 chain of a $\gamma\delta$ TCR having the Bin 8 δ 1 epitope is
20 AKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQD
TDKLIFGKGTRVTVEP (SEQ ID NO: 36); and another exemplary V δ 1 J region Bin 8 δ 1 epitope of a δ 1 chain of a $\gamma\delta$ TCR is
AKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGEAPSAWGKHLTAQLFFGKGQTQLIV
EP (SEQ ID NO: 37).

25 In some cases, activating agents that bind a Bin 8 δ 1 epitope bind a δ 1 chain of a $\gamma\delta$ TCR
having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQD
TDKLIFGKGTRVTVEP (SEQ ID NO: 22); but do not bind a δ 1 chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
30 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDSWDTRQMFFGTGIK

LFVEP (SEQ ID NO: 23).

In some embodiments, the epitope that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 80-95 of human V δ 1

(FKKAAKSVALTISALQ) (SEQ ID NO: 28). This epitope is referred to herein as a Bin 9 δ 1

5 epitope. Exemplary activating agents that bind a Bin 9 δ 1 epitope include but are not limited to δ 1-201.

In some cases, activating agents that bind a Bin 9 δ 1 epitope bind a δ 1 chain of a $\gamma\delta$ TCR having the sequence of:

AQKVTQAQSSVSMRPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ

10 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQDKLIFGKGTRVTVEP (SEQ ID NO: 22); but do not bind a δ 1 chain of a $\gamma\delta$ TCR having the sequence of

AQKVTQVQGRAMSSQLGEAVTLSCQYETSLSWYDIFWYKQLPSGEMTFLIHQISSDQ

NAKNGRYSVNFQERHKFISLTISALQLEDSAKYFCALGTGVRGLQDQDKLIFGKGTRVTVEP (SEQ ID NO: 38).

15 The δ 1-specific antibodies described herein selectively bind δ 1-containing $\gamma\delta$ -TCRs over δ 2-containing $\gamma\delta$ -TCRs. As such, the foregoing δ 1-specific antibodies do not bind, *e.g.*, the sequence of and/or a $\gamma\delta$ -TCR comprising the sequence of:

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGP

20 GFKDNFQGDIDIAKNLAVLKILAPSERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ ID NO: 39).

δ 2 Epitope Bins

In some embodiments, the epitope, that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 83-94 of human V δ 2 (AKNLAVLKILAP)

(SEQ ID NO: 40). This epitope is referred to herein as a Bin 1 δ 2 epitope. In some cases, the

25 activating agent that binds the Bin 1 δ 2 epitope does not bind an epitope containing a mutation of K90 in V δ 2, such as an K90N mutation. Exemplary activating agents that bind a Bin 1 δ 2 epitope include but are not limited to δ 2-17, and B6.

In some cases, activating agents that bind a Bin 1 δ 2 epitope bind a δ 2 chain of a $\gamma\delta$ TCR having the sequence of

30 AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGP

GFKDNFQGDIDIAKNLAVLKILAPSERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ ID NO: 39); but do not bind a $\delta 2$ chain of a $\gamma\delta$ TCR having the sequence of:

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGP
GFKDNFQGDIDFLNNQAVLNILEASERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ
5 ID NO: 41).

In some embodiments, the epitope, that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 28-38 of human V $\delta 2$ (EAIGNYY) (SEQ ID NO: 42). This epitope is referred to herein as a Bin 2 $\delta 2$ epitope. In some cases, the activating agent that binds the Bin 2 $\delta 2$ epitope does not bind an epitope containing a mutation of G35 in V $\delta 2$, such
10 as an G35S mutation. Exemplary activating agents that bind a Bin 2 $\delta 2$ epitope include but are not limited to 15D.

In some cases, activating agents that bind a Bin 2 $\delta 2$ epitope bind a $\delta 2$ chain of a $\gamma\delta$ TCR having the sequence of:

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGP
15 GFK.DNFQGDIDIAKNLAVLKILAPSERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ ID NO: 39); but do not bind a $\delta 2$ chain of a $\gamma\delta$ TCR having the sequence of:

AIELVPEHQTVPVSIGVPATLRCSMKGDSISNYTTFWYRRTPGNTMTLIYREGGTYG
PGFEDNLQGEIDFLNNQAVLNILEASERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ
ID NO: 43).

20 In some embodiments, the epitope, that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 72-83 of human V $\delta 2$ (KDNFQGDIDIA) (SEQ ID NO: 44). This epitope is referred to herein as a Bin 3 $\delta 2$ epitope. Exemplary activating agents that bind a Bin 3 $\delta 2$ epitope include but are not limited to $\delta 2$ -32.

25 In some cases, activating agents that bind a Bin 3 $\delta 2$ epitope bind a $\delta 2$ chain of a $\gamma\delta$ TCR having the sequence of:

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGP
GFK.DNFQGDIDIAKNLAVLKILAPSERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ
ID NO: 39); but do not bind a $\delta 2$ chain of a $\gamma\delta$ TCR having the sequence of:

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGP
30 GFEDNLQGEIDFLNNQAVLNILEASERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ

ID NO: 45).

In some embodiments, the epitope, that is (*e.g.*, specifically) recognized by an activating agent of interest is an epitope comprised of amino acids 1-27 of human V δ 2 (AIELVPEHQTPVPSIGVPATLRCSMKG) (SEQ ID NO: 46). This epitope is referred to herein as a Bin 4 δ 2 epitope. Exemplary activating agents that bind a Bin 4 δ 2 epitope include but are not limited to δ 2-14; δ 2-22; δ 2-30; δ 2-31; δ 2-36; and δ 2-37.

In some cases, activating agents that bind a Bin 4 δ 2 epitope bind a δ 2 chain of a $\gamma\delta$ TCR having the sequence of:

AIELVPEHQTPVPSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGP
GFKDNFQGDIDIAKNLAVLKILAPSERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ
ID NO: 39); but do not bind a δ 2 chain of a $\gamma\delta$ TCR having the sequence of:

AVTLVPQNPQARSVSVGESVTLRCSMKGDSISNYYTFWYRRTPGNTMTLIYREGGTY
GPGFEDNLQGEIDFLNNQAVLNILEASERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP
(SEQ ID NO: 47).

The δ 2-specific antibodies described herein selectively bind δ 2-containing $\gamma\delta$ -TCRs over δ 1-containing $\gamma\delta$ -TCRs. As such, the foregoing δ 2-specific antibodies do not bind, *e.g.*, the sequence of and/or a $\gamma\delta$ -TCR comprising the sequence of:

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDQDKLIFGKGTRVTVE
P (SEQ ID NO: 22).

Generally, the δ 1- and δ 2-specific antibodies described herein recognize a conformational epitope in the context of a $\gamma\delta$ -TCR. In some cases, the δ 1- and δ 2-specific antibodies described herein are specific for one or more pairs of $\gamma\delta$ 1- or $\gamma\delta$ 2- TCRs respectively. For example, in some cases, the δ 1- specific antibodies described herein are specific for a $\gamma\delta$ 1-TCR. In some cases, the δ 1- specific antibodies described herein bind $\gamma\delta$ 1-TCR but not $\gamma\delta$ 2-TCR.

Methods of Treatment

Pharmaceutical compositions containing a non-engineered, enriched $\gamma\delta$ T-cell population, an engineered, enriched $\gamma\delta$ T-cell population, and/or admixtures thereof, as described herein may be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, the

compositions can be administered to a subject already suffering from a disease or condition in an amount sufficient to cure or at least partially arrest the symptoms of the disease or condition. A non-engineered, enriched $\gamma\delta$ T-cell population, an engineered, enriched $\gamma\delta$ T-cell population, and/or admixtures thereof, can also be administered to lessen a likelihood of developing, contracting, or worsening a condition. Effective amounts of a population of a non-engineered, enriched $\gamma\delta$ T-cell population, an engineered, enriched $\gamma\delta$ T-cell population, and/or admixtures thereof, for therapeutic use can vary based on the severity and course of the disease or condition, previous therapy, the subject's health status, weight, and/or response to the drugs, and/or the judgment of the treating physician.

A non-engineered, enriched $\gamma\delta$ T-cell population, an engineered, enriched $\gamma\delta$ T-cell population, and /or admixtures thereof, of the disclosure can be used to treat a subject in need of

TCR chains were amplified and cloned out of TS8.2 and B6 activated populations. PBMCs cultures were harvested after 14 days of specific activation with $\delta 1$ specific activating mAbs TS8.2 and the $\delta 2$ specific MAb B6. PBMCs cultures were lysed in 350 μ l RLT buffer (Qiagen,), Total RNA was purified from 5×10^5 cell using the Qiagen RNeasy Mini Kit (Qiagen, catalog number 74106), following the manufacturer's instructions. In brief, 350 μ l of 70% Ethanol was added to the cell lysate to provide ideal binding conditions. The lysate was then loaded onto the RNeasy silica membrane spin column. Contaminants were washed away. Concentrated RNA was eluted in 50 μ l water.

One step RT PCR: Full length delta chains were amplified using the 5' $\delta 1$ (5' GCCCAGAAGGTTACTCAAGCCCAGTC3') (SEQ ID NO: 48) and 5' $\delta 2$ (5' GCCATTGAGTTGGTGCCTGAACACC 3') (SEQ ID NO: 49) primers, in combination with 3' human C δ (5' TTACAAGAAAAATAACTTGGCAGTCAAGAGAAA 3') (SEQ ID NO: 50) primer. A 800 bp PCR fragments of the full length delta chain were cloned into the pCI expression vector (Promega catalog number E1841). Similarly, full length gamma chains were amplified using the 5' primers $\gamma 2, \gamma 3, \gamma 4$ (5' TCTTCCAACCTTGAAGGGAGAACGAAGTC 3') (SEQ ID NO: 51) $\gamma 5$ (5' TCTTCCAACCTTGAAGGGGGAACGA 3') (SEQ ID NO: 52) , $\gamma 8$ (5' TCTTCCAACCTTGAAGGGAGAACAAAGTC 3') (SEQ ID NO: 53) $\gamma 9$ (5' GCAGGTCACCTAGAGCAACCTCAAATTTCC 3') (SEQ ID NO: 54) in combination with a single 3' Cgamma primer (5' GGAAGAAAAATAGTGGGCTTGGGGGAA 3') (SEQ ID NO: 13)

The delta and gamma transcripts were amplified from 100 ng total RNA using reverse transcriptase polymerase chain reaction (RT-PCR). A total of five RT-PCR reactions were run for each sample activated with the TS8.2 MAb ($\delta 1, \gamma 2\text{-}\gamma 4, \gamma 5, \gamma 8$ and $\gamma 9$) and two RT-PCR reactions were run for each sample activated with the B6 MAb ($\delta 2$, and $\gamma 9$). The QIAGEN One Step RT-PCR kit was used for amplification, (Qiagen, Inc.). This kit provides a blend of Sensiscript and Omniscript Reverse Transcriptases, HotStarTaq DNA Polymerase, dNTP mix, buffer and Q-Solution, a novel additive that enables efficient amplification of "difficult" (e.g., GC-rich) templates. Reaction mixtures were prepared that included 5 μ l of RNA, 0.5 of 100 μ M of either delta or gamma chains primers (custom synthesized by IDT), 5 μ l of 5 $\times$ RT-PCR buffer, 1 μ l dNTPs, 1 μ l of enzyme

mix containing reverse transcriptase and DNA polymerase, and 0.4 µl of ribonuclease inhibitor RNasin (1 unit). The reaction mixture contains all of the reagents required for both reverse transcription and PCR. The thermal cycler program was RT step 50° C. for 30 minutes 95° C. for 15 minutes followed by 30 cycles of (95° C. for 30 seconds, 48° C. for 30 seconds, 72° C. for 1.0 minutes). There was then a final incubation at 72° C. for 10 minutes. The extracted PCR products were directly cloned into pCI expression vector. Nucleotide sequences were analyzed using IMGT to identify germline V, D and J gene members with the highest sequence homology.

Examples of sequences of Vδ1 chains activated by TS8.2 CDR3 (junctional diversity is indicated by bold text):

TS8.2 activated clone1 Vδ1, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKS
GRYSVNFKKA AKSVALTISALQLEDSAKYF**CALGPVVIPKGKLS**FGKGTRVTVEP (SEQ ID
NO: 55)

TS8.2 activated clone2 Vδ1, D3, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKS
GRYSVNFKKA AKSVALTISALQLEDSAKYF**CALGELCLGDTYTDKLIF**GKGTRVTVEP
(SEQ ID NO: 56)

TS8.82 activated clone 3 Vδ1, D2+D3, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKS
GRYSVNFKKA AKSVALTISALQLEDSAKYF**CALGDPKVYWGCTDKLIF**GKGTRVTVEP
(SEQ ID NO: 57)

TS8.2 activated clone 4 Vδ1, D3, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQDAKS
GRYSVNFKKA AKSVALTISALQLEDSAKYF**CALGTGVRGLQDADKLIF**GKGTRVTVEP
(SEQ ID NO: 58)

TS8.2 activated clone 5 Vδ1 D1+D3, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKS
GRYSVNFKKA AKSVALTISALQLEDSAKYF**CALLLGDTSFYTDKLIF**GKGTRVTVEP (SEQ
ID NO: 59)

TS8.2 activated clone 6 Vδ1 D1+D3, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKS
GRYSVNFKKA AKSVALTISALQLEDSAKYF**CAALLPFLPSDWGIPVTDKLIF**GKGTRVTVE
P (SEQ ID NO: 60)

TS8.2 activated clone 7 Vδ1, D2 and D3, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKS
GRYSVNFKKAASKSVALTISALQLEDSAKYFCALGTGVRGLQD**TDKL**IFGTGTRVTVEP
(SEQ ID NO: 61)

5 TS8.2 activated clone 8 V81, D2 and D3, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKS
GRYSVNFKKAASKSVALTISALQLEDSAKYFCAL**GGLSSLDLGD**TDNHY**TDKL**IFGKGTRVTVEP
(SEQ ID NO: 62)

10 TS8.2 activated clone 9 V81, D2 and D3, J1

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQNAKS
GRYSVNFKKAASKSVALTISALQLEDSAKYFCAL**GHPRSLMGVYTDKL**IFGKGTRVTVEP
(SEQ ID NO: 63)

15 **Examples of Gamma chains activated by TS8.2 Mab:**

B8 TS8.82 activated Clone-1 γ 4, J1/J2

20 SSNLEGRTKSVIRQTGSSAEITCDLAEGSTGYIHWYHLHQEGKAPQRLLYYDSYTSSVVLES
SPGKYDTYGSTRKNLRIILRNLIENDSGVYY**CATWDDGKKL**FGSGTTLVVT (SEQ ID NO:
64)

B8 TS8.82 activated Clone-2 γ 9, JP1

25 AGHLEQPQISSTKTLSKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYY**CALWDDTRLGKKIKV**FAPGTKLIIT (SEQ ID
NO: 65)

TS8.2 activated Clone-3 γ 4, J1

30 SSNLEGRTKSVIRQTGSSAEITCDLAEGSTGYIHWYHLHQEGKAPQRLLYYDSYTSSVVLES
SPGKYDTYGSTRKNLRMILRNLIENDSGVYY**CATWSDKKL**FGSGTTLVVT (SEQ ID NO:
66)

TS8.2 activated Clone-4 γ 9, J1/J2

35 AGHLEQPQISSTKTLSKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYY**CALWIHKKL**FGSGTTLVVT (SEQ ID NO: 67)

TS8.2 activated Clone 5 γ 2, J1

40 SSNLEGRTKSVIRQTGSSAENTCDLAEGSNGYIHWYHLHQEGKTPQRLQYYDSYNSKVVLES
GVSPGKYTYASTRNNLRLILRNLIENDSGVYY**CATWDCHYKKL**FGSGTTLVVT (SEQ ID
NO: 68)

TS8.2 activated Clone 6 γ 3, JP2

45 SSNLEGRTKSVTRQTGSSAEITCDLTVTNTFYIHWYHLHQEGKAPQRLLYYDVSTARDVLESG
LSPGKYTYHTPRRWSWILRLQNLIENDSGVYY**CATWDRRWIKT**FAKGTKLIVTSP (SEQ ID
NO: 69)

- TS8.2 activated Clone 7 γ 2, JP1
SSNLEGRTKSVIRQTGSSAEITCDLAEGSNGYIHWYHLHQEGKAPQRLQYYDSYNSKVVLES
GVSPGKYITYASTRNNLRLILRNLIENDSGVYYCATWDGLDATCGVDTTGWFKIFAEGT
KLIVTSP (SEQ ID NO: 70)
- 5 TS8.2 activated Clone 8 γ 4, J1/J2
SSNLEGRTKSVTRPTGSSAVITCDLPVENAVYTHWYHLHQEGKAPRRLLYYDSYTSSVVLES
GISPGKYDTYGSTRKNLRMILRNLIENDSGVYYCATKKLFGSGTTLVVT (SEQ ID NO: 71)
- 10 TS8.2 activated Clone 9 γ 9, JP1
AGHLEQPQISSTKTLSTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYYCALWEIASQLGKKIKVFGPGTKLIIT (SEQ
ID NO: 72)
- 15 TS8.2 activated Clone 10 γ 3, J1
SSNLGGRTKSVTRQTGSSAEITCDLTVTNTFYIHWYHLHQEGKAPQRLLYYDVSTARDVLES
GLSPGKYTYHTPRRWSWILRLQNLIENDSGVYYCATWDRYKYKKLFGSGTTLVVT (SEQ
ID NO: 73)
- 20 TS8.2 activated Clone 11 γ 3, JP2
SSNLEGRTKSVTRQTGSSAEITCDLTVTNTFYIHWYHLHQEGKAPQRLLYYDVSTARDVLESG
LSPGKYTYHTPRRWSWILRLQNLIENDSGVYYPNSSDWIKTFAGTKLIVTSP (SEQ ID NO:
74)
- 25 TS8.2 activated Clone 12 γ 4, J2
SSNLEGRTKSVTRPTRSSAEITCDLAERNFTFYIHWYHLHQEGKAPQRLQYYDSYTSSVVLESG
ISPGKYDTYGSTRKNLRMILRNLIENDSGVYYCATRDVPNYKKLFGSGTTLVVT (SEQ ID
NO: 75)
- 30 TS8.2 activated Clone 13 γ 2, JP1
SSNLEGRTKSVIRQTGSSAEITCDLAEGSNGYIHWYHLHQEGKAPQRLQYYDSYNSKVVLES
GVSPGKYITYASTRNNLRLILRNLIENDSGVYYCATWDGRVSYTTGWFKIFAEGTKLIVTS
P (SEQ ID NO: 76)
- 35 TS8.2 activated Clone 14 γ 4, J1/J2
SSNLEGRTKSVIRQTGSSAEITCDLAEGSTGYIHWYHLHQEGKAPQRLLYYDSYTSSVVLESGI
SPGKYDTYGSTRKNLRMILRNLIENDSGVYYCATWDKGRKLFGSGTTLVVT (SEQ ID NO:
77)
- 40 TS8.2 activated Clone 15 γ 9, J1
AGHLEQPQISSTKTLSTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYYCALWETHYKKLFGSGTTLVVT (SEQ ID NO:
78)
- TS8.2 activated Clone 16 γ 2, J1

SSNLEGRTKSVIRQTGSSAEITCDLAEGSNGYIHWYHQQEGKAPQRLQYYDSYNSKVVLES
GVSPGKYTYASTRNNLRLILRNLIENDSGVYYCATWDGRYKKLFGSGTTLVVT (SEQ ID
NO: 79)

5 TS8.2 activated Clone 17 γ 4, JP1

SSNLEGRTKSVIRQTGSSAEITCDLAEGSTGYIHWYHQQEGKAPQRLLYYDSYTSSVVLES
SPGKYDTYGSTRKNLRMILRNLIENDSGVYYCATWGTTGWFKIFAEGTKLIVTSP (SEQ ID
NO: 80)

10 **B8(Buffy coat 8)- B6 activated clones:**

B6 activated Clone 82, D3,J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPGFKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDKVLGVPTASYTDNKLIFGKGTRVTVEP
15 (SEQ ID NO: 81)

B6 activated Clone 82, D3,J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMSFIYREKDIYGPGFKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDTVGLPYDKLIFGKGTRVTVEP (SEQ ID
20 NO: 82)

B6 activated Clone 82, D2+D3, J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPGFKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDILTVLGDNRDTDKLIFGKGTRVTVEP (SEQ
25 ID NO: 83)

B6 activated Clone 82, D3,J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPGFKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDVVGEGGADKLIFGKGTRVTVEP (SEQ ID
30 NO: 84)

B6 activated Clone 82, D3,J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPGFKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDTVGGGEYTDKLIFGKGTRVTVEP (SEQ
35 ID NO: 85)

B6 activated Clone 82, D3,J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPGFKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDTVGTGDIRTYTDKLIFGKGTRVTVEP
40 (SEQ ID NO: 86)

B6 activated Clone 82, D2+D3,J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPGFKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDSLTTGSGSLTDKLIFGKGTRVTVEP (SEQ
45 ID NO: 87)

B6 activated Clone 82, D3, J3

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPFGKD
NFQGDIDIAKNLAVLKILPPSERDEGSYYCACDTGGYSSWDTRQMFFGTGIKLFVEP (SEQ
ID NO: 88)

5

B6 activated Clone 82, D2+D3, J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPFGKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDPLKTLGTYTDKLIFGKGTRVTVEP (SEQ
ID NO: 89)

10

B6 activated Clone 82, D1+D3, J1

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPFGKD
NFQGDIDIAKNLAVLKILAPSERDEGSYYCACDAVIAGGSFTDKLIFGKGTRVTVEP (SEQ
ID NO: 90)

15

B6 activated Gamma chains:

B6 activated clone γ 9, JP1

20 AGHLEQPQISSTKTLSKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYYCALWEDQELGKKIKVFGPGTKLIIT (SEQ ID
NO: 91)

B6 activated clone γ 9, JP1

25 AGHLEQPQISSTKTLSKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYYCALWAYPPELGKKIKVFGPGTKLIIT (SEQ
ID NO: 92)

B6 activated clone γ 9, JP1

30 AGHLEQPQISSTKTLSKTARLECAVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYYCALWEVQELGKKIKVFGPGTKLIIT (SEQ ID
NO: 93)

B6 activated clone γ 9, JP1

35 AGHLEQPQISSTKTLSKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYYCALWEVLQELGKKIKVFGPGTKLIIT (SEQ
ID NO: 94)

B6 activated clone γ 9, JP1

40 AGHLEQPQISSTKTLSKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKESGI
PSGKFEVDRIPESTSTLTIHNVEKQDIATYYCALWEVRELGKKIKVFGPGTKLIIT (SEQ ID
NO: 95)

B6 activated clone γ 9,

JP1AGHLEQPQISSTKTLSKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKE

SGIPSGKFEVD RIPETSTSTLTIHNVEKQDIATYYCALWRELGKKIKVFGPGTKLIIT (SEQ ID NO: 96)

B6 activated clone γ 9.

5 JP1AGHLEQPQISSTKTL SKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKE
SGIPSGKFEVD RIPETSTSTLTIHNVEKQDIATYYCALWEAQELGKKIKVFGPGTKLIIT
(SEQ ID NO: 97)

B6 activated clone γ 9.

10 JP1AGHLEQPQISSTKTL SKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKE
SGIPSGKFEVD RIPETSTSTLTIHNVEKQDIATYYCALWETELGKKIKVFGPGTKLIIT (SEQ
ID NO: 98)

B6 activated clone γ 9.

15 JP1AGHLEQPQISSTKTL SKTARLECVVSGITISATSVYWYRERPGEVIQFLVSISYDGTVRKE
SGIPSGKFEVD RIPETSTSTLTIHNVEKQDIATYYCALWEEVELGKKIKVFGPGTKLIIT
(SEQ ID NO: 99)

B6 activated clone γ 9, JP1

20 AGHLEQPQISSTKTL SKTARLECVVSGITISATSVYWYRERPGEVIQFPVSISYDGTVRKESGI
PSGKFEVD RIPETSTSTLTIHNVEKQDIATYYCALWEPPQSLGKKIKVFGPGTKLIIT (SEQ
ID NO: 100)

**Example 34. Cryopreservation of $\gamma\delta$ T-Cells in Freeze Media to Generate Cell Banks
for Further Processing**

25 Non-modified $\gamma\delta$ T-Cells will be formulated in freezing media and placed in cryogenic storage
units such as liquid nitrogen freezers (-195°C) or ultra-low temperature freezers (-65°C , -80°C or -120°C)
for long-term storage. The freeze media will contain dimethyl sulfoxide (DMSO), sodium
chloride (NaCl), dextrose, dextran sulfate or hydroxyethyl starch (HES) with physiological pH
buffering agents ranging between 6.5 and 7.5.

30 The cryopreserved $\gamma\delta$ T-cells will be thawed and further processed by stimulation with
antibodies, proteins, peptides, and cytokines. The cryopreserved $\gamma\delta$ T-Cells will be thawed and
genetically modified as previously described in this application. The engineered $\gamma\delta$ T-cells will be
further cryopreserved to generate cell banks in quantities of 10, 100, 200 vials at 10^6 to 10^8 cells per
mL in freeze media. The non-engineered and/or engineered $\gamma\delta$ T-cells will be further cryopreserved
35 to generate cell banks in quantities of 10, 100, 200, 500, and 1,000 vials or bags at 10^6 to 10^8 cells per
mL in freeze media.

HEK-293 cells were cultured in 90-mm cell culture plates in a DMEM medium (Thermo Fisher Scientific) with 10% fetal calf serum (HyClone™ FetalClone™ I Serum (U.S.) - GE Healthcare Life Sciences). For every construct 1×10^6 HEK-293 cells were transfected with 10 µg of plasmid using 293 Fectin as transfection reagent. Supernatants were harvested 72 hours post transfection

81 specific activators- monoclonal antibodies differed in their binding pattern to the chimeras and a distinct binding pattern could be identified for each monoclonal antibody, as tested in ELISA assay.

Polystyrene microtiter plates (Greiner bio-one # 655061) were coated overnight at room temperature with 1 µg/well of Goat anti human FC specific antibody (Jackson ImmunoResearch # 109-005-098) in ELISA coating buffer (50 mM Na₂CO₃, 0.05% NaH₃, pH 9.6). The plates were washed three times with washing buffer (PBS 0.05% (v/v) Tween 20) and then blocked for 1 h at room temperature (RT) with 100 µl/well of ELISA blocking buffer (3%BSA in PBST). Binding of soluble TCRs to blocked plates was done by adding 100µl of each supernatant and incubation for 1 hour at Room temperature. Each antibody was diluted to 1 µg/ml in ELISA blocking buffer, and tested for binding to each γδ TCR chimeric molecule.

Detection of binding was done by adding 100 µl/well of HRP conjugated secondary antibody: Goat anti mouse FC specific (Jackson ImmunoResearch Laboratories, Inc. Peroxidase-AffiniPure Goat Anti-Mouse IgG, Fcγ Fragment Specific # 115-035-008) diluted to 1:10,000 in blocking buffer. The amount of bound antibodies was detected by the use of TMB substrate (Thermo Fisher #34029)Color development was measured spectrophotometrically at 450nm after 10 minutes. All assays were done in duplicate.

Sequence of all constructs

δ1-D3-J1 FC- Fully human TCR

gccagaagggtactcaagcccagtcacgtatccatgccagtgaggaaagcagtcacacctgaactgcctgtatgaaacaagttggt
ggatcatattatatttttggtacaagcaactccagcaaagagatgattttccttattcgccagggttctgatgaacagaatgcaaaaagtggtcgcta
ttctgtcaactcaagaaagcagcgaaatccgtcgccttaaccatttcagccttacagctagaagattcagcaaagtacttttgtgctcttgggacgg
gggtgaggggactccaggacaccgataaactcatctttgaaaaaggaacccgtgtgactgtggaaccaa (SEQ ID NO: 101)

AQKVTQAQSSVSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAASVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVE
P (SEQ ID NO: 22)

5 Dolphin TRD isolate 5R1D8

gcccagaaagtactcaagtcagcgagccatgtccagtcagctaggggagggcggtcaccttgagctgtcagtatgaaacaagcttg
agctggtagcatatttttggataagcagcttccagtgagagatgactttccttattcatcagatttcttctgacaaaaatgcaagaatggccgct
attctgtaaaccttcaggaaagacataaattcatcagcctcaccatttcagccttactgggtggaagattctgcaaaactacttctgtgctctccgggagc
gcgtgtccacgtcggtttattcaatacgcgcaataagccactgctattcggaaggaacctatctgaacgttgaaccaa (SEQ ID NO:
10 102)

AQKVTQVQRAMSSQLGEAVTLSCQYETSLSWYDIFWYKQLPSGEMTFLIHQISSDQ
NAKNGRYSVNFQERHKFISLTISALLVEDSANYFCALRERVVHVVLFNTRNKPLLFGKGTYL
NVEP (SEQ ID NO: 103)

15

Set of dolphin/human chimera -soluble TCRs were cloned -Human sequence in blue

Dolphin/Human 1-11

gcccagaaagtactcaagtcagcgagccatgtccatgccagtgaggaaagcagtcaccctgaactgcctgtatgaaacaagttggt
20 ggtagcatattatatttttggtagcaagcaacttccagcaagagatgattttccttattcgccaggggttctgatgaacagaatgcaaaaagtggtcgcta
ttctgtcaacttcaagaaagcagcgaaatccgtcgcttaaccatttcagccttacagctagaagattcagcaaaagtactttgtgctcttgggacgg
gggtgaggggactccaggacaccgataaactcatctttgaaaaggaacccgtgtgactgtggaaccaa (SEQ ID NO: 104)

AQKVTQVQRAMSMPVRKAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDE
25 QNAKSGRYSVNFKKAASVALTISALQLEDSAKYFCALGTGVRGLQDQTDKLIFGKGTRVTVEP (SEQ ID NO: 105)

Dolphin/Human 1-21

gcccagaaagtactcaagtcagcgagccatgtccagtcagctaggggagggcggtcaccttgactgcctgtatgaaacaagttggt
30 ggtagcatattatatttttggtagcaagcaacttccagcaagagatgattttccttattcgccaggggttctgatgaacagaatgcaaaaagtggtcgcta

ttctgtcaacttcaagaaagcagcgaaatccgtcgccttaaccatttcagccttacagctagaagattcagcaaagtacttttgctcttgggacgg
gggtgaggggactccaggacaccgataaactcatctttggaaaaggaacccgtgtgactgtggaaccaa (SEQ ID NO: 106)

AQKVTQVQRAMSSQLGEAVTLNCLYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
5 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDTDKLIFGKGTRVTVE
P (SEQ ID NO: 107)

Dolphin/Human 1-28

gcccagaaagtactcaagtccagcgagccatgtccagtcagctaggggagggcggtcaccttgagctgtcagtatgaaacaagctgg
10 tggatcatattatatttttggtacaagcaactccagcaaaagagatgattttccttattcgccaggggttctgatgaacagaatgcaaaaagtggcgcg
attctgtcaacttcaagaaagcagcgaaatccgtcgccttaaccatttcagccttacagctagaagattcagcaaagtacttttgctcttgggacg
ggggtgaggggactccaggacaccgataaactcatctttggaaaaggaacccgtgtgactgtggaaccaa (SEQ ID NO: 108)

AQKVTQVQRAMSSQLGEAVTLSCQYETSWWSYYIFWYKQLPSKEMIFLIRQGSDEQ
15 NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDTDKLIFGKGTRVTVE
P (SEQ ID NO: 109)

Dolphin/Human 1-47

gcccagaaagtactcaagtccagcgagccatgtccagtcagctaggggagggcggtcaccttgagctgtcagtatgaaacaagcttg
20 agctggtacgatatttttggtataagcagcttccagtaaagagatgattttccttattcgccaggggttctgatgaacagaatgcaaaaagtggcgcg
tattctgtcaacttcaagaaagcagcgaaatccgtcgccttaaccatttcagccttacagctagaagattcagcaaagtacttttgctcttgggacg
ggggtgaggggactccaggacaccgataaactcatctttggaaaaggaacccgtgtgactgtggaaccaa (SEQ ID NO: 110)

AQKVTQVQRAMSSQLGEAVTLSCQYETSLSWYDIFWYKQLPSKEMIFLIRQGSDEQ
NAKSGRYSVNFKKAAKSVALTISALQLEDSAKYFCALGTGVRGLQDTDKLIFGKGTRVTVE
25 P (SEQ ID NO: 111)

Dolphin/Human 1-70

gcccagaaagtactcaagtccagcgagccatgtccagtcagctaggggagggcggtcaccttgagctgtcagtatgaaacaagcttg
agctggtacgatatttttggtataagcagcttccagtgagagatgactttccttattcatcagatttctgatgaacagaatgcaaaaagtggcgcg

attctgtcaactcaagaaagcagcgaatccgtcgccttaaccatttcagccttacagctagaagattcagcaaagtactttgtgctcttgggacg
ggggtgaggggactccaggacaccgataaactcatctttgaaaaggaacccgtgtgactgtggaaccaa (SEQ ID NO: 112)

AQKVTQVQGRAMSSQLGEAVTLSCQYETSLSWYDIFWYKQLPSGEMTFLIHQISDEQ
5 NAKSGRYSVNFKKA AKSVALTISALQLEDSAKYFCALGTGVRGLQDTDKLIFGKGTRVTVE
P (SEQ ID NO: 113)

Dolphin/human 1-80

gcccagaaagtactcaagtccagcagccatgtccagtcagctaggggagggcggtcaccttgagctgtcagtatgaaacaagcttg
10 agctggtacgatatttttggataagcagcttccagtgagagatgactttccttattcatcagatttcttctgacaaaaatgcaaagaatggccgct
attctgtaaactttaagaaagcagcgaatccgtcgccttaaccatttcagccttacagctagaagattcagcaaagtactttgtgctcttgggacgg
gggtgaggggactccaggacaccgataaactcatctttgaaaaggaacccgtgtgactgtggaaccaa (SEQ ID NO: 114)

AQKVTQVQGRAMSSQLGEAVTLSCQYETSLSWYDIFWYKQLPSGEMTFLIHQISSDQ
15 NAKNGRYSVNFKKA AKSVALTISALQLEDSAKYFCALGTGVRGLQDTDKLIFGKGTRVTVE
P (SEQ ID NO: 115)

Dolphin/human 1-95

gcccagaaagtactcaagtccagcagccatgtccagtcagctaggggagggcggtcaccttgagctgtcagtatgaaacaagcttg
20 agctggtacgatatttttggataagcagcttccagtgagagatgactttccttattcatcagatttcttctgacaaaaatgcaaagaatggccgct
attctgtaaactttcaggaaagacataaattcatcagcctcaccatttcagccttacagctagaagattcagcaaagtactttgtgctcttgggacgg
gggtgaggggactccaggacaccgataaactcatctttgaaaaggaacccgtgtgactgtggaaccaa (SEQ ID NO: 116)

AQKVTQVQGRAMSSQLGEAVTLSCQYETSLSWYDIFWYKQLPSGEMTFLIHQISSDQ
NAKNGRYSVNFQERHKFISLTISALQLEDSAKYFCALGTGVRGLQDTDKLIFGKGTRVTVEP
25 (SEQ ID NO: 38)

RESULTS

Functional TCR genes are assembled from separate V, D and J region segments by genetic
recombination. When the TCR delta chain gene is assembled a D gene segment is juxtaposed to a J
30 segment and this is followed by rearrangement of a V segment to the assembled downstream DJ
segment. The immunoglobulin-like fold of the TCR chains positions three loops or complementarity

Bin6: (aa 21-28+ J1 region): TS8.2; δ 1-143
 Bin7: (aa 47-70+ J1/J2 region): δ 1-149; δ 1-253, and δ 1-257
 Bin8: (aa 70-80) + J1/J2: δ 1-192
 Bin9: (aa 80-95): δ 1-201.

5

Similar to the δ 1 epitope mapping: based on Linguiti G. et al., 2016. Genomic and expression analyses of *Tursiops truncatus* T cell receptor gamma (TRG) and alpha/delta (TRA/TRD) loci reveal a similar basic public; repertoire in dolphin and human. BMC Genomics. A set of 5 recombinant chimeric δ 2 chains were cloned to produce human/dolphin δ 2-FC TCR chains. Dolphin residues comprised amino acids 28-94, 44-94, 72-94 and, 82-94 respectively, according to IMGT unique numbering for T cell receptor variable domains (Lefranc et al., Developmental and Comparative Immunology 27 (2003) 55–77), as illustrated in **Fig.15**.

Transfection of δ 2/ γ 9-FC constructs

15 Human TRGV9-JP-FC

GCAGGTCACCTAGAGCAACCTCAAATTTCCAGTACTAAAACGCTGTCAAAAACAGCCCGCCTGGAATGTGTGGTGTCTGGAATAACAAT
 TTCTGCAACATCTGTATATTGGTATCGAGAGAGACCTGGTGAAGTCATACAGTTCCTGGTGTCCATTTTCATATGACGGCACTGTCAGAA
 AGGAATCCGGCATTCCGTCAGGCAAATTTGAGGTGGATAGGATACCTGAAACGTCTACATCCACTCTCACCATTACAAATGTAGAGAAA
 CAGGACATAGCTACCTACTACTGTGCCTTGTGGGAGGTAAAGCAAGAGTTGGGCAAAAAAATCAAGGTATTGGTCCCGGAACAAAGCT
 TATCATTACAG (SEQ ID NO: 117)

20 AGHLEQPPQISSTKTLTKTARLECVVSGITISATSVYWRERPEVQFLVSISYDGTVRKESGIPSGKFEVDRIPESTSTLTIHNVK
 QDIATYYCALWEVKQELGKKIKVFGPGTKLIIT (SEQ ID NO: 118)

25 Human V δ 21J1-FC

GCCATTGAGTTGGTGCCTGAACACCAAACAGTGCCTGTGTCAATAGGGGTCCCTGCCACCCTCAGGTGCTCCATGAAAGG
 AGAAGCGATCGGTAACCTACTATATCAACTGGTACAGGAAGACCCAAGGTAACACAATGACTTTTCATATACCGAGAAAAGG
 ACATCTATGGCCCTGGTTTCAAAGACAATTTCCAAGGTGACATTGATATTGCAAAGAACCCTGGCTGTACTTAAGATACTT
 GCACCATCAGAGAGAGATGAAGGGTCTTACTACTGTGCCTGTGACCCTCTTGGCGGACCCCCGATAAACTCATCTTTGG
 AAAAGGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 119)

30 AIELVPEHQTPVPSIGVPATLRCSMKGEAIGNYYINWYRKTQGNMTFTIYREKDIYGPFGKDNFQGDIDIAKNLAVLKIL
 APSEDEGSYYACDPLGGPPDKLIFGKGTRVTVEP (SEQ ID NO: 39)

35 Dolphin V δ 2J1-FC

GCTGTACGTTGGTGCCTCAAAAACCAAGCAAGGAGTGTGTCTGTGGGGGAATCTGTACCCCTCAGGTGCTCCATGAAAGG
 AGACTCCATCAGTAACTATTATACCTTCTGGTACAGGAGAACACCGGGTAACACAATGACTCTCATATACCGAGAAGGGG
 GCACATATGGCCCTGGTTTCAAGACAACCTCCAAGGTGAAATTGATTTTTTAAACAACCAGGCTGTGCTGAATATCCTG
 GAGGCATCAGAGAGAGATGAAGGATCTTACTACTGTGCCTGTGACCCTCTTGGCGGACCCCCGATAAACTCATCTTTGG
 AAAA
 40 GGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 120)

AVTLVPQNQARSVSVGESVTLRCSMKGDSISNYTFWYRRTPGNTMTLIYREGGTYGPFGEDNLQGEIDFLNNQAVLNIL
 EASERDEGSYYACDPLGGPPDKLIFGKGTRVTVEP (SEQ ID NO: 47)

Human/Dolphin Vδ2J1-FC (28-94)

5 GCCATTGAGTTGGTGCCTGAACACCAAACAGTGCCTGTGTCAATAGGGGTCCCTGCCACCCTCAGGTGCTCCATGAAAGG
AGACTCCATCAGTAACTATTATACCTTCTGGTACAGGAGAACACCGGGTAACACAATGACTCTCATATACCGAGAAGGGG
GCACATATGGCCCTGGTTTCGAAGACAACCTCCAAGGTGAAATTGATTTTTTAAACAACCAGGCTGTGCTGAATATCCTG
GAGGCATCAGAGAGAGATGAAGGATCTTACTACTGTGCCTGTGACCCTCTTGGCGGACCCCCCGATAAACTCATCTTTGG
AAAA
GGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 121)

10 AIELVPEHQTVPVSIGVPATLRCSMKGDSISNYTTFWYRRTPGNTMTLIYREGGTYGPGFEDNLQGEIDFLNNQAVLNIL
EASERDEGSYYC**CACDPLGGPPDKLIF**GKGTRVTVEP (SEQ ID NO: 43)

Human/Dolphin Vδ2J1-FC (44-94)

15 GCCATTGAGTTGGTGCCTGAACACCAAACAGTGCCTGTGTCAATAGGGGTCCCTGCCACCCTCAGGTGCTCCATGAAAGG
AGAAGCGATCGGTAACTACTATATCAACTGGTACAGGAGAACACCGGGTAACACAATGACTTTCATATACCGAGAAAGGG
GCACATATGGCCCTGGTTTCGAAGACAACCTCCAAGGTGAAATTGATTTTTTAAACAACCAGGCTGTGCTGAATATCCTG
GAGGCATCAGAGAGAGATGAAGGATCTTACTACTGTGCCTGTGACCCTCTTGGCGGACCCCCCGATAAACTCATCTTTGG
AAAAGGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 122)

20 AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRRTPGNTMTLIYREGGTYGPGFEDNLQGEIDFLNNQAVLNIL
EASERDEGSYYC**CADPLGGPPDKLIF**GKGTRVTVEP (SEQ ID NO: 123)

Human/ Dolphin Vδ2J1-FC (72-94)

25 GCCATTGAGTTGGTGCCTGAACACCAAACAGTGCCTGTGTCAATAGGGGTCCCTGCCACCCTCAGGTGCTCCATGAAAGG
AGAAGCGATCGGTAACTACTATATCAACTGGTACAGGAAGACCCAAGGTAACACAATGACTTTCATATACCGAGAAAAGG
ACATCTATGGCCCTGGTTTCGAAGACAACCTCCAAGGTGAAATTGATTTTTTAAACAACCAGGCTGTGCTGAATATCCTG
GAGGCATCAGAGAGAGATGAAGGATCTTACTACTGTGCCTGTGACCCTCTTGGCGGACCCCCCGATAAACTCATCTTTGG
AAAAGGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 124)

30 AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNMTMTFIYREKDIYGPGFEDNLQGEIDFLNNQAVLNIL
EASERDEGSYYC**CADPLGGPPDKLIF**GKGTRVTVEP (SEQ ID NO: 45)

Human Dolphin Vδ2J1-FC (82-94)

35 GCCATTGAGTTGGTGCCTGAACACCAAACAGTGCCTGTGTCAATAGGGGTCCCTGCCACCCTCAGGTGCTCCATGAAAGG
AGAAGCGATCGGTAACTACTATATCAACTGGTACAGGAAGACCCAAGGTAACACAATGACTTTCATATACCGAGAAAAGG
ACATCTATGGCCCTGGTTTCAAAGACAATTTCCAAGGTGACATTGATTTTTTAAACAACCAGGCTGTGCTGAATATCCTG
GAGGCATCAGAGAGAGATGAAGGATCTTACTACTGTGCCTGTGACCCTCTTGGCGGACCCCCCGATAAACTCATCTTTGG
AAAAGGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 125)

40 AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNMTMTFIYREKDIYGPGFKDNFQGDIDFLNNQAVLNIL
EASERDEGSYY**CACDPLGGPPDKLIF**GKGTRVTVEP (SEQ ID NO: 41)

Human Vδ21J1-FC (CDR1-S)

45 GCCATTGAGTTGGTGCCTGAACACCAAACAGTGCCTGTGTCAATAGGGGTCCCTGCCACCCTCAGGTGCTCCATGAAAGG
AGAAGCGATCAGTAACTACTATATCAACTGGTACAGGAAGACCCAAGGTAACACAATGACTTTCATATACCGAGAAAAGG
ACATCTATGGCCCTGGTTTCAAAGACAATTTCCAAGGTGACATTGATATTGCAAAGAACCTGGCTGTACTTAAGATACTT
GCACCATCAGAGAGAGATGAAGGGTCTTACTACTGTGCCTGTGACCCTCTTGGCGGACCCCCCGATAAACTCATCTTTGG
AAAAGGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 126)

50 AIELVPEHQTVPVSIGVPATLRCSMKGEA**IS**NYINWYRKTQGNMTMTFIYREKDIYGPGFKDNFQGDIDIAKNLAVLKIL
APSERDEGSYYC**CADPLGGPPDKLIF**GKGTRVTVEP (SEQ ID NO: 127)

Human Vδ21J1-FC (CDR2-G)

GCCATTGAGTTGGTGCCTGAACACCAAACAGTGCCTGTGTCAATAGGGGTCCCTGCCACCCTCAGGTGCTCCATGAAAGG
AGAAGCGATCGGTAAGTACTATATCAACTGGTACAGGAAGACCCAAGGTAACACAATGACTTTCATATACCGAGAAAAGG
GCATCTATGGCCCTGGTTTCAAAGACAATTTCCAAGGTGACATTGATATTGCAAAGAACCTGGCTGTACTTAAGATACTT
GCACCATCAGAGAGAGATGAAGGGTCTTACTACTGTGCCTGTGACCCTCTTGGCGGACCCCCCGATAAACTCATCTTTGG
AAAAGGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 128)

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKGIYGPFGKDNFQGDIDIAKNLAVLKIL
APSERDEGSYYCACDPLGGPPDKLIFGKGTRVTVEP (SEQ ID NO: 129)

Human Vδ21J1-FC (CDR3-S)

GCCATTGAGTTGGTGCCTGAACACCAAACAGTGCCTGTGTCAATAGGGGTCCCTGCCACCCTCAGGTGCTCCATGAAAGG
AGAAGCGATCGGTAAGTACTATATCAACTGGTACAGGAAGACCCAAGGTAACACAATGACTTTCATATACCGAGAAAAGG
ACATCTATGGCCCTGGTTTCAAAGACAATTTCCAAGGTGACATTGATATTGCAAAGAACCTGGCTGTACTTAAGATACTT
GCACCATCAGAGAGAGATGAAGGGTCTTACTACTGTGCCTCTGACCCTCTTGGCGGACCCCCCGATAAACTCATCTTTGG
AAAAGGAACCCGTGTGACTGTGGAACCAA (SEQ ID NO: 130)

AIELVPEHQTVPVSIGVPATLRCSMKGEAIGNYYINWYRKTQGNTMTFIYREKDIYGPFGKDNFQGDIDIAKNLAVLKIL
APSERDEGSYYCASDPLGGPPDKLIFGKGTRVTVEP (SEQ ID NO: 131)

ELISA results are shown in **Fig. 48**, identifying the following epitope bins:

Bin	1	2	3	4
Residues	83-94	28-38	72-83	1-27
mAbs	δ2-17	15D	δ 2-32	δ 2-14
	B6			δ 2-22
				δ 2-30
				δ 2-31
				δ 2-36
				δ 2-37

* * *

While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is

CLAIMS

WHAT IS CLAIMED IS:

1. An *ex vivo* method for producing an enriched $\delta 1$ $\gamma\delta$ T-cell population from an isolated mixed cell population without prior selection of $\gamma\delta$ T-cells and/or depletion of non- $\gamma\delta$ T-cells, and under
5 culture conditions that do not comprise IL-4 and wherein said cell population has not been exposed to exogenous IL-4 prior to expansion, said method comprising directly contacting the mixed cell population with at least one antibody that selectively expands $\delta 1$ T-cells without significant expansion of $\delta 2$ T-cells by binding to an epitope specific to a $\delta 1$ variable region of a $\delta 1$ TCR to
10 provide clinically-relevant levels of an enriched $\delta 1$ $\gamma\delta$ T cell population, wherein the at least one antibody comprises a heavy chain variable region/light chain variable region (HCVR/LCVR) sequence pair selected from the group consisting of SEQ ID NOs: 153/180, 154/181, 155/182, 156/183, 157/184, 158/185, 159/186, 160/187, 161/188, 162/189, 163/190, 164/191, 165/192, 166/193, 167/194, 168/195, 169/196, 170/197, 171/198, 172/199, 173/200, 174/201, 175/202, 176/203, 177/204, 178/205, and 179/206.
15
2. The method of claim 1, wherein the percentage of $\delta 1$ T cells in the enriched $\gamma\delta$ T cell population is greater than 60% $\delta 1$ T-cells of total viable cells.
3. The method of any one of the preceding claims, wherein said method does not comprise
20 expanding the enriched $\delta 1$ $\gamma\delta$ T-cell population by an antigen presenting cell or an aminophosphate.
4. The method of claim 1, wherein said at least one antibody that selectively expands $\delta 1$ T-cells is immobilized on a surface.
- 25 5. The method of claim 1, wherein said isolated mixed cell population is selected from a peripheral blood sample, a cord blood sample or a tumor.
6. The method of claim 1, wherein said enriched $\delta 1$ $\gamma\delta$ T-cell population comprise polyclonal TCR diversity.

30

7. The method of claim 1, wherein the enriched $\delta 1$ $\gamma\delta$ T-cell population is further formulated for administration to a subject.

8. The method of claim 1, wherein the enriched $\delta 1$ $\gamma\delta$ T-cell population includes a therapeutically effective amount of $\gamma\delta$ T-cells.

9. The method of claim 1, wherein said $\delta 1$ $\gamma\delta$ T-cell population is further engineered to stably express one or more tumor recognition moieties.

10. The method of claim 1, further comprising:
directly contacting at least a portion of the enriched $\delta 1$ $\gamma\delta$ T-cell population with antigen presenting cells (APCs) in a second $\gamma\delta$ T-cell expansion, thereby producing a second enriched $\gamma\delta$ T-cell population,

wherein the second enriched $\gamma\delta$ T-cell population comprises a clinically relevant number of $\gamma\delta$ T-cells.

11. The method of claim 1, further comprising:
directly contacting at least a portion of the enriched $\delta 1$ $\gamma\delta$ T-cell population with one or more agents that (a) expand $\gamma\delta$ T-cells, and/or (b) deplete $\alpha\beta$ T-cells.

12. The method of claim 1, wherein an antibody that selectively expands $\delta 1$ T-cells is an antibody that specifically recognizes the $\delta 1$ chain variable region and i) the CDR3 loop created by the restricted juxtaposition of V(D)J1 and V(D)J2 but not V(D)J3 segments, or iii) the CDR3 loop created by the juxtaposition of V(D)J1, V(D)J2 and V(D)J3 segments.

13. The method of claim 1, wherein the antibody binds to an epitope comprising residues Arg71, Asp72 and Lys120 of the $\delta 1$ variable region.

14. The method of claim 1, wherein the antibody has reduced binding to a mutant $\delta 1$ TCR polypeptide comprising a mutation at K120 at the delta J1 and delta J2.

15. The method of claim 1, wherein the epitope is selected from a $\delta 1$ TCR Bin 1 $\delta 1$ epitope, Bin 1b $\delta 1$ epitope, Bin 2 $\delta 1$ epitope, Bin 2b $\delta 1$ epitope, Bin 2c $\delta 1$ epitope, Bin 3 $\delta 1$ epitope, Bin 4 $\delta 1$ epitope, Bin 5 $\delta 1$ epitope, Bin 6 $\delta 1$ epitope, Bin 7 $\delta 1$ epitope, Bin 8 $\delta 1$ epitope, or Bin 9 $\delta 1$ epitope of
5 a human $\delta 1$ TCR.