

CLAIMS

What is claimed is:

1. A method of manufacturing an immune cell or a population thereof, comprising subjecting the immune cell to a small compound treatment comprising at least one of dexamethasone, lenalidomide, AQX-1125, or a derivative or an analogue thereof, thereby obtaining an immune cell having enhanced post-thaw cytotoxicity as compared to a counterpart immune cell without the same small compound treatment.
2. The method of claim 1, wherein the immune cell is a derivative effector immune cell differentiated from an induced pluripotent stem cell (iPSC), wherein the effector immune cell comprises: a derivative CD34 cell, a derivative hematopoietic stem and progenitor cell, a derivative hematopoietic multipotent progenitor cell, a derivative T cell progenitor, a derivative NK cell progenitor, a derivative T cell, a derivative NKT cell, a derivative NK cell, a derivative B cell, or a derivative effector cell having one or more functional features that are not present in a counterpart primary T, NK, NKT, and/or B cell.
3. The method of claim 2, wherein the iPSC comprises at least one of the following edits:
 - (i) a first chimeric antigen receptor (CAR) having a first targeting specificity;
 - (ii) CD38 knockout;
 - (iii) HLA-I deficiency and/or HLA-II deficiency, in comparison to its native counterpart cell;
 - (iv) introduced expression of HLA-G or non-cleavable HLA-G, or knockout of one or both of CD58 and CD54;
 - (v) a CD16 or a variant thereof;
 - (vi) a second CAR having a second targeting specificity;
 - (vii) a signaling complex comprising a partial or full peptide of a cell surface expressed exogenous cytokine and/or a receptor thereof;
 - (viii) at least one of the genotypes listed in Table 2;
 - (ix) deletion or reduced expression in at least one of B2M, CIITA, TAP1, TAP2, Tapasin, NLRC5, CIITA, RFXANK, RFX5, RFXAP, TCR α or β constant region, NKG2A, NKG2D, CD25, CD69, CD44, CD56, CIS, CBL-B, SOCS2, PD1, CTLA4, LAG3, TIM3, and TIGIT, in comparison to its native counterpart cell; or

- (x) introduced or increased expression in at least one of HLA-E, 41BBL, CD3, CD4, CD8, CD16, CD47, CD113, CD131, CD137, CD80, PDL1, A_{2A}R, antigen-specific TCR, Fc receptor, an antibody or fragment thereof, a checkpoint inhibitor, an engager, and surface triggering receptor for coupling with bi- or multi-specific or universal engagers; and wherein the effector immune cell differentiated from the iPSC comprises the same one or more edits as the iPSC.
4. The method of claim 3, wherein the first CAR comprises:
- (i) an ectodomain comprising at least one antigen recognition region, a transmembrane domain, and an endodomain comprising at least one signaling domain; and wherein the at least one signaling domain is originated from a cytoplasmic domain of a signal transducing protein specific to T and/or NK cell activation or functioning;
 - (ii) an antigen recognition domain that specifically binds an antigen associated with a disease, a pathogen, a liquid tumor, or a solid tumor; or
 - (iii) an antigen recognition domain that is specific to:
 - (a) any one of CD19, BCMA, CD20, CD22, CD38, CD123, HER2, CD52, EGFR, GD2, MICA/B, MSLN, VEGF-R2, PSMA and PDL1; or
 - (b) any one of ADGRE2, carbonic anhydrase IX (CAIX), CCR1, CCR4, carcinoembryonic antigen (CEA), CD3, CD5, CD7, CD8, CD10, CD20, CD22, CD30, CD33, CD34, CD38, CD41, CD44, CD44V6, CD49f, CD56, CD70, CD74, CD99, CD123, CD133, CD138, CDS, CLEC12A, an antigen of a cytomegalovirus (CMV) infected cell, epithelial glycoprotein2 (EGP 2), epithelial glycoprotein-40 (EGP-40), epithelial cell adhesion molecule (EpCAM), EGFRvIII, receptor tyrosine-protein kinases erb- B2,3,4, EGFR, EGFR-VIII, ERBB folate-binding protein (FBP), fetal acetylcholine receptor (AChR), folate receptor-a, Ganglioside G2 (GD2), Ganglioside G3 (GD3), human Epidermal Growth Factor Receptor 2 (HER-2), human telomerase reverse transcriptase (hTERT), ICAM-1, Integrin B7, Interleukin-13 receptor subunit alpha-2 (IL-13R α 2), κ -light chain, kinase insert domain receptor (KDR), Lewis A (CA19.9), Lewis Y (LeY), L1 cell adhesion molecule (L1-CAM), LILRB2, melanoma antigen family A 1 (MAGE-A1), MICA/B, Mucin 1 (Muc-1), Mucin 16 (Muc-16), Mesothelin (MSLN), NKCSI, NKG2D ligands, c-Met, cancer-testis antigen NY-ESO-1, oncofetal antigen

(h5T4), PRAME, prostate stem cell antigen (PSCA), PRAME prostate-specific membrane antigen (PSMA), tumor-associated glycoprotein 72 (TAG-72), TIM-3, TRBC1, TRBC2, vascular endothelial growth factor R2 (VEGF- R2), and Wilms tumor protein (WT-1).

5. The method of claim 3, wherein the first CAR is comprised in a bi-cistronic construct co-expressing:

- (1) a partial or full-length peptide of a cell surface expressed exogenous cytokine or a receptor thereof, wherein the exogenous cytokine or receptor thereof comprises:
 - (a) at least one of IL2, IL4, IL6, IL7, IL9, IL10, IL11, IL12, IL15, IL18, IL21, or its respective receptor;
 - (b) at least one of:
 - (i) co-expression of IL15 and IL15R α by using a self-cleaving peptide;
 - (ii) a fusion protein of IL15 and IL15R α ;
 - (iii) an IL15/IL15R α fusion protein with intracellular domain of IL15R α truncated or eliminated;
 - (iv) a fusion protein of IL15 and membrane bound Sushi domain of IL15R α ;
 - (v) a fusion protein of IL15 and IL15R β ;
 - (vi) a fusion protein of IL15 and common receptor γ C, wherein the common receptor γ C is native or modified; and
 - (vii) a homodimer of IL15R β ;
- (2) an antibody or fragment thereof;
- (3) an engager; or
- (4) a checkpoint inhibitor.

6. The method of claim 1, wherein the small compound treatment of the immune cell is prior to or subsequent to cryopreservation of the immune cell.

7. The method of claim 1, wherein the method further comprises cryopreserving the immune cell subjected to the small compound treatment.

8. The method of claim 7, wherein the cryopreservation is free or substantially free of one or more small compounds of the treatment.

9. The method of claim 1, wherein the enhanced post-thaw cytotoxicity comprises enhanced *in vivo* efficacy of immune cells thawed after cryogenic preservation, and wherein the post-thaw immune cells having the small compound treatment comprise at least one of the following characteristics:
- (i) enhanced ability in tumor control, tumor clearance, and/or reducing tumor relapse;
 - (ii) improved tumor penetration; or
 - (iii) enhanced ability in migrating to bone marrow and/or to tumor sites,
- as compared to post-thaw counterpart immune cells without the same small compound treatment.
10. The method of claim 1, wherein the small compound treatment:
- (i) comprises dexamethasone, or a derivative or an analog thereof;
 - (ii) is free or essentially free of cytokine IL7, optionally, wherein the immune cell under the treatment is a T cell;
 - (iii) is free or essentially free of cytokine IL2 and/or cytokine IL15, optionally, wherein the immune cell under the treatment is an NK cell;
 - (iv) comprises dexamethasone, but does not comprise cytokine IL7;
 - (v) is free or essentially free of cytokines;
 - (vi) is during cell culturing and/or prior to or subsequent to cryopreservation;
 - (vii) is during immune cell expansion after differentiating the cell from iPSC; and/or
 - (viii) lasts between about 1 to about 12 days, or between about 3 to about 6 days, prior to cryopreservation.
11. The method of claim 10, wherein the dexamethasone is present at a concentration range between about 10nM to about 20μM.
12. A cell or a population thereof, wherein:
- (i) the cell is an immune cell that has been subjected to a small compound treatment comprising at least one of dexamethasone, lenalidomide, AQX-1125, and a derivative or an analogue thereof; and
 - (ii) the immune cell comprises enhanced post-thaw cytotoxicity as compared to a counterpart immune cell without the same small compound treatment.
13. The cell or the population thereof of claim 12, wherein:

- (iii) the immune cell is a derivative effector immune cell differentiated from an induced pluripotent stem cell (iPSC); and
- (iv) the effector immune cell comprises: a derivative CD34 cell, a derivative hematopoietic stem and progenitor cell, a derivative hematopoietic multipotent progenitor cell, a derivative T cell progenitor, a derivative NK cell progenitor, a derivative T cell, a derivative NKT cell, a derivative NK cell, a derivative B cell, or a derivative effector cell having one or more functional features that are not present in a counterpart primary T, NK, NKT, and/or B cell.

14. The cell or the population thereof of claim 13, wherein the iPSC comprises at least one of the following edits:

- (i) a first chimeric antigen receptor (CAR) having a first targeting specificity;
- (ii) CD38 knockout;
- (iii) HLA-I deficiency and/or HLA-II deficiency, in comparison to its native counterpart cell;
- (iv) introduced expression of HLA-G or non-cleavable HLA-G, or knockout of one or both of CD58 and CD54;
- (v) a CD16 or a variant thereof;
- (vi) a second CAR having a second targeting specificity;
- (vii) a signaling complex comprising a partial or full peptide of a cell surface expressed exogenous cytokine and/or a receptor thereof;
- (viii) at least one of the genotypes listed in Table 2;
- (ix) deletion or reduced expression in at least one of B2M, CIITA, TAP1, TAP2, Tapasin, NLRC5, CIITA, RFXANK, RFX5, RFXAP, TCR α or β constant region, NKG2A, NKG2D, CD25, CD69, CD44, CD56, CIS, CBL-B, SOCS2, PD1, CTLA4, LAG3, TIM3, and TIGIT, in comparison to its native counterpart cell; or
- (x) introduced or increased expression in at least one of HLA-E, 41BBL, CD3, CD4, CD8, CD16, CD47, CD113, CD131, CD137, CD80, PDL1, A_{2A}R, antigen-specific TCR, Fc receptor, an antibody or fragment thereof, a checkpoint inhibitor, an engager, and surface triggering receptor for coupling with bi- or multi- specific or universal engagers; and wherein the effector immune cell differentiated from the iPSC comprises the same one or more edits as the iPSC.

15. The cell or the population thereof of claim 14, wherein the first and second CARs independently comprise:

- (i) an ectodomain comprising at least one antigen recognition region, a transmembrane domain, and an endodomain comprising at least one signaling domain; and wherein the at least one signaling domain is originated from a cytoplasmic domain of a signal transducing protein specific to T and/or NK cell activation or functioning;
- (ii) an antigen recognition domain that specifically binds an antigen associated with a disease, a pathogen, a liquid tumor, or a solid tumor; or
- (iii) an antigen recognition domain that is specific to:
 - (a) any one of CD19, BCMA, CD20, CD22, CD38, CD123, HER2, CD52, EGFR, GD2, MICA/B, MSLN, VEGF-R2, PSMA and PDL1; or
 - (b) any one of ADGRE2, carbonic anhydrase IX (CAIX), CCR1, CCR4, carcinoembryonic antigen (CEA), CD3, CD5, CD7, CD8, CD10, CD20, CD22, CD30, CD33, CD34, CD38, CD41, CD44, CD44V6, CD49f, CD56, CD70, CD74, CD99, CD123, CD133, CD138, CDS, CLEC12A, an antigen of a cytomegalovirus (CMV) infected cell, epithelial glycoprotein2 (EGP 2), epithelial glycoprotein-40 (EGP-40), epithelial cell adhesion molecule (EpCAM), EGFRvIII, receptor tyrosine-protein kinases erb- B2,3,4, EGFR, EGFR-VIII, ERBB folate-binding protein (FBP), fetal acetylcholine receptor (AChR), folate receptor-a, Ganglioside G2 (GD2), Ganglioside G3 (GD3), human Epidermal Growth Factor Receptor 2 (HER-2), human telomerase reverse transcriptase (hTERT), ICAM-1, Integrin B7, Interleukin-13 receptor subunit alpha-2 (IL-13R α 2), κ -light chain, kinase insert domain receptor (KDR), Lewis A (CA19.9), Lewis Y (LeY), L1 cell adhesion molecule (L1-CAM), LILRB2, melanoma antigen family A 1 (MAGE-A1), MICA/B, Mucin 1 (Muc-1), Mucin 16 (Muc-16), Mesothelin (MSLN), NKCSI, NKG2D ligands, c-Met, cancer-testis antigen NY-ESO-1, oncofetal antigen (h5T4), PRAME, prostate stem cell antigen (PSCA), PRAME prostate-specific membrane antigen (PSMA), tumor-associated glycoprotein 72 (TAG-72), TIM-3, TRBC1, TRBC2, vascular endothelial growth factor R2 (VEGF-R2), and Wilms tumor protein (WT-1).

16. The cell or the population thereof of claim 14, wherein the first CAR is comprised in a bi-cistronic construct co-expressing:

- (1) a partial or full-length peptide of a cell surface expressed exogenous cytokine or a receptor thereof, wherein the exogenous cytokine or receptor thereof comprises:
 - (a) at least one of IL2, IL4, IL6, IL7, IL9, IL10, IL11, IL12, IL15, IL18, IL21, or its respective receptor;
 - (b) at least one of:
 - (i) co-expression of IL15 and IL15R α by using a self-cleaving peptide;
 - (ii) a fusion protein of IL15 and IL15R α ;
 - (iii) an IL15/IL15R α fusion protein with intracellular domain of IL15R α truncated or eliminated;
 - (iv) a fusion protein of IL15 and membrane bound Sushi domain of IL15R α ;
 - (v) a fusion protein of IL15 and IL15R β ;
 - (vi) a fusion protein of IL15 and common receptor γ C, wherein the common receptor γ C is native or modified; and
 - (vii) a homodimer of IL15R β ;
- (2) an antibody or fragment thereof; or
- (3) a checkpoint inhibitor.

17. The cell or the population thereof of claim 12, wherein the small compound treatment of the immune cell is prior to cryopreservation of the immune cell.

18. The cell or the population thereof of claim 12, wherein the small compound treated immune cell is:

- (i) comprised in a pre-cryopreservation medium;
- (ii) comprised in a cryopreservation medium;
- (iii) in cryopreservation; or
- (iv) post-thaw from cryopreservation.

19. The cell or the population thereof of claim 12, wherein the cryopreservation is free or substantially free of one or more small compounds of the treatment.

20. The cell or the population thereof of claim 12, wherein the enhanced post-thaw cytotoxicity comprises enhanced *in vivo* efficacy of immune cells thawed after cryogenic

preservation, and wherein the post-thaw immune cells having the small compound treatment prior to the cryogenic preservation comprise at least one of the following characteristics:

- (i) enhanced ability in tumor control, tumor clearance, and/or reducing tumor relapse;
- (ii) improved tumor penetration; or
- (iii) enhanced ability in migrating to bone marrow and/or to tumor sites,

as compared to post-thaw counterpart immune cells without the same small compound treatment.

21. The cell or the population thereof of claim 12, wherein the small compound treatment:

- (i) comprises dexamethasone;
- (ii) is free or essentially free of cytokine IL7, optionally, wherein the immune cell under the treatment is a T cell;
- (iii) is free or essentially free of cytokine IL2 and/or cytokine IL15, optionally, wherein the immune cell under the treatment is an NK cell;
- (iv) comprises dexamethasone, but does not comprise cytokine IL7;
- (v) is free or essentially free of cytokines;
- (vi) is during cell culturing and/or prior to or subsequent to cryopreservation;
- (vii) is during immune cell expansion after differentiating the cell from iPSC; and/or
- (viii) lasts between about 1 to about 12 days, or between about 3 to about 6 days, prior to cryopreservation.

22. The cell or the population thereof of claim 21, wherein the dexamethasone is present at a concentration range between about 10nM to about 20μM.

23. The cell or the population thereof of claim 12, wherein the immune cell comprises one or more differentially expressed genes comprising at least one of:

- (i) SPOCK2, PTGDS, IL7R, LCNL1, RASGRP2, SMAP2, IL6ST, IL-7R, and IL2RA up-regulation; or
- (ii) JCHAIN, KLF3, KLRB1, IGFBP4, NUCB2, CSF2RB, and CXCR6 down-regulation,

as compared to a counterpart immune cell without the same small compound treatment.

24. The cell or the population thereof of claim 12, wherein the immune cell is comprised in a medium, wherein the medium:

- (i) comprises dexamethasone;
- (ii) comprises lenalidomide;

- (iii) comprises AQX-1125;
- (iv) comprises dexamethasone and lenalidomide;
- (v) comprises dexamethasone, but not cytokine IL7, and optionally, wherein the immune cell is a T cell;
- (vi) comprises dexamethasone, but not cytokine IL2 or cytokine IL15, and optionally, wherein the immune cell is an NK cell; or
- (vii) comprises dexamethasone and is free or essentially free of cytokines.

25. A method of manufacturing an immune cell or a population thereof, wherein the method comprises:

- (a) differentiating a genetically engineered iPSC to obtain the immune cell, wherein the iPSC comprises at least one of the following edits:
 - (i) a first chimeric antigen receptor (CAR) having a first targeting specificity;
 - (ii) CD38 knockout;
 - (iii) HLA-I deficiency and/or HLA-II deficiency, in comparison to its native counterpart cell;
 - (iv) introduced expression of HLA-G or non-cleavable HLA-G, or knockout of one or both of CD58 and CD54;
 - (v) a CD16 or a variant thereof;
 - (vi) a second CAR having a second targeting specificity;
 - (vii) a signaling complex comprising a partial or full peptide of a cell surface expressed exogenous cytokine and/or a receptor thereof;
 - (viii) at least one of the genotypes listed in Table 2;
 - (ix) deletion or reduced expression in at least one of B2M, CIITA, TAP1, TAP2, Tapasin, NLRC5, CIITA, RFXANK, RFX5, RFXAP, TCR α or β constant region, NKG2A, NKG2D, CD25, CD69, CD44, CD56, CIS, CBL-B, SOCS2, PD1, CTLA4, LAG3, TIM3, and TIGIT, in comparison to its native counterpart cell; or
 - (x) introduced or increased expression in at least one of HLA-E, 41BBL, CD3, CD4, CD8, CD16, CD47, CD113, CD131, CD137, CD80, PDL1, A_{2A}R, antigen-specific TCR, Fc receptor, an antibody or fragment thereof, a checkpoint inhibitor, an engager, and surface triggering receptor for coupling with bi- or multi- specific or universal engagers; and
- wherein the immune cell differentiated from the iPSC comprises the same one or more edits as the iPSC; and

- (b) subjecting the immune cell to a small compound treatment comprising at least one of dexamethasone, lenalidomide, AQX-1125, or a derivative or an analogue thereof,

thereby obtaining an immune cell having enhanced post-thaw cytotoxicity as compared to a counterpart immune cell without the same small compound treatment.

26. The method of claim 25, wherein the method further comprises: (c) cryopreserving the treated immune cell from step (b).

27. The method of claim 25, further comprising genomically engineering a clonal iPSC to knock in a polynucleotide encoding the first CAR, and optionally:

- (i) to knock out CD38;
- (ii) to knock out B2M and CIITA;
- (iii) to knock out one or both CD58 and CD54; and/or
- (iv) to introduce expression of HLA-G or non-cleavable HLA-G, a CD16 or a variant thereof, a second CAR, and/or a partial or full peptide of a cell surface expressed exogenous cytokine or a receptor thereof.

28. The method of claim 27, wherein the genomic engineering comprises targeted deletion, insertion, or in/del, and wherein the genomic engineering is carried out by CRISPR, ZFN, TALEN, homing nuclease, homology recombination, or any other functional variation of these methods.

29. The method of claim 25, wherein the immune cell differentiated from the induced pluripotent stem cell (iPSC) comprises: a derivative CD34 cell, a derivative hematopoietic stem and progenitor cell, a derivative hematopoietic multipotent progenitor cell, a derivative T cell progenitor, a derivative NK cell progenitor, a derivative T cell, a derivative NKT cell, a derivative NK cell, a derivative B cell, or a derivative effector cell having one or more functional features that are not present in a counterpart primary T, NK, NKT, and/or B cell.

30. The method of claim 26, wherein the method further comprises (d) thawing the cryopreserved immune cell from step (c).

31. A composition for therapeutic use comprising the immune cell of any one of claims 12-23, and one or more therapeutic agents.

32. The composition of claim 31, wherein the one or more therapeutic agents comprise a peptide, a cytokine, a checkpoint inhibitor, a mitogen, a growth factor, a small RNA, a dsRNA (double stranded RNA), mononuclear blood cells, feeder cells, feeder cell components or replacement factors thereof, a vector comprising one or more polynucleic acids of interest, an antibody, a chemotherapeutic agent or a radioactive moiety, or an immunomodulatory drug (IMiD).
33. The composition of claim 32, wherein;
- (i) the checkpoint inhibitor comprises:
 - (a) one or more antagonists to checkpoint molecules comprising PD-1, PDL-1, TIM-3, TIGIT, LAG-3, CTLA-4, 2B4, 4-1BB, 4-1BBL, A2aR, BATE, BTLA, CD39, CD47, CD73, CD94, CD96, CD160, CD200, CD200R, CD274, CEACAM1, CSF-1R, Foxp1, GARP, HVEM, IDO, EDO, TDO, LAIR-1, MICA/B, NR4A2, MAFB, OCT-2, Rara (retinoic acid receptor alpha), TLR3, VISTA, NKG2A/HLA-E, or inhibitory KIR;
 - (b) one or more of atezolizumab, avelumab, durvalumab, ipilimumab, IPH4102, IPH43, IPH33, lirimumab, monalizumab, nivolumab, pembrolizumab, and their derivatives or functional equivalents; or
 - (c) at least one of atezolizumab, nivolumab, and pembrolizumab; or
 - (ii) the therapeutic agents comprise one or more of venetoclax, azacitidine, and pomalidomide.
34. The composition of claim 32, wherein the antibody comprises:
- (a) an anti-CD20, an anti-HER2, an anti-CD52, an anti-EGFR, an anti-CD123, an anti-GD2, an anti-PDL1, and/or an anti-CD38 antibody;
 - (b) one or more of rituximab, veltuzumab, ofatumumab, ublituximab, ocaratuzumab, obinutuzumab, trastuzumab, pertuzumab, alemtuzumab, certuximab, dinutuximab, avelumab, daratumumab, isatuximab, MOR202, 7G3, CSL362, elotuzumab, and their humanized or Fc modified variants or fragments and their functional equivalents and biosimilars; or
 - (c) daratumumab, and wherein the derivative hematopoietic cells comprise derivative NK cells or derivative T cells comprising a CD38 knockout, and optionally expression of CD16 or a variant thereof.

35. Therapeutic use of the composition of any one of the claims 31-34 by introducing the composition to a subject suitable for adoptive cell therapy, wherein the subject has an autoimmune disorder, a hematological malignancy, a solid tumor, cancer, or a virus infection.
36. A method of treating a disease or a condition comprising:
- (i) thawing one or more units of cryopreserved immune cells manufactured according to any one of claims 26-29; and
 - (ii) administering to a subject a composition comprising the post-thaw immune cells of step (i).
37. The method of claim 36, wherein the immune cells are iPSC derived NK cells, iPSC derived T cells, or iPSC derived effector cells having one or more functional features that are not present in counterpart primary T, NK, NKT, and/or B cells.