

-1-

QUATERNARY AMMONIUM COMPOUNDS AS TACHYKININ ANTAGONISTS

This invention relates to quaternary ammonium compounds. More particularly, this invention relates to 1-(2-acylimidazol-1-ylalkyl)quinuclidinium salts and to processes for the preparation of, intermediates used in the preparation of, compositions containing and the uses of, such salts.

The present compounds are antagonists of tachykinins, including NKA (neurokinin A), NKB (neurokinin B) and Substance P, acting at the human neurokinin-1(NK₁), neurokinin-2 (NK₂) and neurokinin-3 (NK₃) receptors.


These compounds are particularly useful as dual NK₁ and NK₂ receptor antagonists and can therefore be used for treating an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis or Crohn's disease, a disease caused by Helicobacter pylori or another urease-positive Gram negative bacteria, a urogenital tract disorder such as incontinence, impotence, hyperreflexia or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis, urticaria, eczematoid dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a proliferative disorder such as a cancer or a disorder involving fibroblast proliferation, a vasospastic disease such as angiogenesis, angina or Reynaud's disease, a fibrosing or collagen disease such as atherosclerosis, scleroderma or eosinophilic fascioliasis, reflux sympathetic dystrophy such as shoulder/hand syndrome, an addiction disorder such as alcoholism, a stress-related somatic disorder, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn,

-2-

herpetic neuralgia or post-herpetic neuralgia, a neuropathological disorder such as Alzheimer's disease or multiple sclerosis, a disorder related to immune enhancement or suppression such as systemic lupus erythematosus, a
 5 rheumatic disease such as fibrositis, emesis, cough, acute or chronic pain, migraine, an ophthalmic disease such as proliferative retinopathy, influenza or a cold.

EP-A-680962 and EP-A-0739891 disclose heterocyclic compounds which are non-peptide antagonists of NKA and that are useful for the treatment
 10 of diseases such as asthma. EP-A-0591040 discloses quaternary compounds with tachykinin antagonist activity.

The present invention provides compounds of the formula:



wherein

R is phenyl, C₃-C₇ cycloalkyl or heteroaryl, each of which being optionally benzo- or C₃-C₇ cycloalkyl-fused and optionally substituted, including in the benzo- or C₃-C₇ cycloalkyl-fused portion, by from 1 to 3 substituents each
 25 independently selected from C₁-C₄ alkyl, fluoro(C₁-C₄)alkyl, C₁-C₄ alkoxy, fluoro(C₁-C₄)alkoxy, phenoxy, C₂-C₄ alkanoyl, halo, C₁-C₄ alkoxycarbonyl, C₃-C₇ cycloalkyl, -S(O)_m(C₁-C₄ alkyl), cyano, -NR²R³, -S(O)_mNR²R³, -NR⁴(C₁-C₄ alkanoyl) and -CONR²R³, or R is 2,3-dihydrobenzo[b]furanyl or chromanyl;

30 R¹ is H or C₁-C₆ alkyl;

-3-

R^2 and R^3 are either each independently selected from H and C_1-C_6 alkyl, or when taken together, represent C_4-C_6 alkylene;

5 R^4 is H or C_1-C_6 alkyl;

W is a direct link, methylene or ethylene;

X is unbranched C_2-C_4 alkylene;

10

Y is phenyl, naphthyl, benzyl, pyridyl, thienyl or C_3-C_7 cycloalkyl, each of which being optionally substituted by from 1 to 3 substituents each independently selected from C_1-C_4 alkyl, fluoro(C_1-C_4)alkyl, C_1-C_4 alkoxy, fluoro(C_1-C_4)alkoxy, halo and cyano;

15

Ar is phenyl, naphthyl, benzyl, thienyl, benzo[b]thienyl or indolyl, each of which being optionally substituted by from 1 to 3 substituents each independently selected from C_1-C_4 alkyl, fluoro(C_1-C_4)alkyl, C_1-C_4 alkoxy, fluoro(C_1-C_4)alkoxy, halo and cyano, or Ar is 1,3-benzodioxolan-4 or 5-yl or 1,4-benzodioxan-5 or
20 6-yl;

m is 0, 1 or 2;

Z^A is a pharmaceutically acceptable anion;

25

and "heteroaryl", used in the definition of R, means thienyl or a 5- or 6-membered ring heteroaryl group containing either from 1 to 4 nitrogen heteroatoms, or 1 or 2 nitrogen heteroatom(s) and 1 oxygen or sulphur heteroatom,

30

-4-

with the proviso that when W is a direct link and R is optionally fused and optionally substituted heteroaryl, said heteroaryl is linked by a ring carbon atom to the carbonyl group.

5

In the above definitions, "halo" means fluoro, chloro, bromo or iodo and alkyl and alkoxy groups having three or more carbon atoms, alkanoyl groups having four carbon atoms and alkylene groups having two or more carbon atoms (except where stated) may be unbranched- or branched-chain.

10

Z^A is a pharmaceutically acceptable anion such as chloride, bromide, nitrate, methanesulphonate, para-toluenesulphonate, benzenesulphonate, hydrogen sulphate or sulphate.

15

Preferably, Z^A is chloride or methanesulphonate.

Most preferably, Z^A is methanesulphonate.

20

A compound of the formula (I) contains one or more asymmetric carbon atoms and therefore exists in two or more stereoisomeric forms. The present invention includes the individual stereoisomers of the compounds of the formula (I) and mixtures thereof.

25

Separation of diastereoisomers may be achieved by conventional techniques, e.g. by fractional crystallisation, chromatography or H.P.L.C. of a stereoisomeric mixture of a compound of the formula (I) or a suitable salt or derivative thereof. An individual enantiomer of a compound of the formula (I) may also be prepared from a corresponding optically pure intermediate or by resolution, such as by H.P.L.C. of the corresponding racemate using a suitable chiral support or by fractional crystallisation of the diastereoisomeric salts formed by reaction of the corresponding racemate with a suitable optically active acid.

30

-5-

Preferably, R is phenyl which is optionally benzo- or C₃-C₇ cycloalkyl-fused and optionally substituted, including in the benzo- or C₃-C₇ cycloalkyl-fused portion, by 1, 2 or 3 substituents each independently selected from C₁-C₄ alkyl, fluoro(C₁-C₄)alkyl, C₁-C₄ alkoxy, fluoro(C₁-C₄)alkoxy, phenoxy and halo, or R is 2,3-dihydrobenzo[b]furanyl.

More preferably, R is phenyl which is optionally benzo- or C₃-C₇ cycloalkyl-fused and optionally substituted, including in the benzo- or C₃-C₇ cycloalkyl-fused portion, by 1, 2 or 3 substituents each independently selected from methyl, ethyl, trifluoromethyl, methoxy, isopropoxy, trifluoromethoxy, phenoxy, fluoro and chloro, or R is 2,3-dihydrobenzo[b]furanyl.

Yet more preferably, R is phenyl, naphthyl or tetrahydronaphthyl, each of which being optionally substituted by 1, 2 or 3 substituents each independently selected from methyl, ethyl, trifluoromethyl, methoxy, isopropoxy, trifluoromethoxy, phenoxy, fluoro and chloro, or R is 2,3-dihydrobenzo[b]furanyl.

Yet further preferably, R is phenyl, 3,5-dimethylphenyl, 2,3-dimethylphenyl, 2-trifluoromethoxyphenyl, 2-methoxy-3-methylphenyl, 2,3-dihydrobenzo[b]furan-7-yl, naphth-2-yl, 4-fluoro-3-trifluoromethylphenyl, 1,2,3,4-tetrahydronaphth-5-yl, 1,2,3,4-tetrahydronaphth-6-yl, 5-chloro-2-methoxyphenyl, 2-methoxyphenyl, 2-trifluoromethylphenyl, 2-isopropoxyphenyl, 2-ethylphenyl, 2-phenoxyphenyl or 3,5-bis(trifluoromethyl)phenyl.

Most preferably, R is 2,3-dimethylphenyl, naphth-2-yl, 1,2,3,4-tetrahydronaphth-5-yl or 2-methoxyphenyl.

Preferably, R¹ is H.

-6-

Preferably, W is a direct link or methylene.

Most preferably, W is a direct link.

5 Preferably, X is 1,2-ethylene.

Preferably, Y is phenyl, naphthyl or cyclohexyl, each of which being optionally substituted by 1, 2 or 3 C₁-C₄ alkyl substituents.

10 More preferably, Y is phenyl, 3,5-dimethylphenyl, cyclohexyl or naphth-2-yl.

Most preferably, Y is phenyl.

Preferably, Ar is phenyl optionally substituted by 1, 2 or 3 halo substituents.


15 More preferably, Ar is phenyl substituted by 1 or 2 chloro substituents.

Most preferably, Ar is 3,4-dichlorophenyl.

Preferred examples of compounds of the formula (I) are compounds of the formula:

20

25



(IA)

wherein

- 1) R-W- is 3,5-dimethylphenyl, Y is phenyl and Z^A is CH₃SO₃⁻;
- 30 2) R-W- is 2,3-dimethylphenyl, Y is phenyl and Z^A is CH₃SO₃⁻;

-7-


- 3) R-W- is 2-trifluoromethoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 4) R-W- is 2-methoxy-3-methylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 5) R-W- is 2,3-dihydrobenzo[b]furan-7-yl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 5 6) R-W- is naphth-2-yl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 7) R-W- is 4-fluoro-3-trifluoromethylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 8) R-W- is 1,2,3,4-tetrahydronaphth-5-yl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 9) R-W- is 1,2,3,4-tetrahydronaphth-6-yl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 10) R-W- is 5-chloro-2-methoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 10 11) R-W- is 2-methoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 12) R-W- is 2-trifluoromethylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 13) R-W- is 2-isopropoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 14) R-W- is 2-ethylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 15) R-W- is 2-phenoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 15 16) R-W- is benzyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - 17) R-W- is 3,5-bis(trifluoromethyl)phenyl, Y is phenyl and Z^A is Cl^- ;
 - 18) R-W- is 2-methoxyphenyl, Y is cyclohexyl and Z^A is CH_3SO_3^- ;
 - 19) R-W- is 4-fluoro-3-trifluoromethylphenyl, Y is cyclohexyl and Z^A is CH_3SO_3^- ;
 - 20 20) R-W- is 2-methoxyphenyl, Y is 3,5-dimethylphenyl and Z^A is CH_3SO_3^- ; or
 - 21) R-W- is 2-methoxyphenyl, Y is naphth-2-yl and Z^A is CH_3SO_3^- ;
- or an alternative pharmaceutically acceptable salt of any thereof (re Z^A).

Particularly preferred examples of the compounds of the formula (I) are


- 25 4-phenyl-1-(3(S)-[3,4-dichlorophenyl]-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)-imidazol-1-yl]butyl)quinuclidinium methanesulphonate and 4-phenyl-1-(3(R)-[3,4-dichlorophenyl]-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate.

- 30 All the compounds of the formula (I) can be prepared by reaction of a compound of the formula:

-8-



10 wherein R, R¹, Ar, W and X are as previously defined for a compound of the formula (I), Z is a suitable leaving group capable of forming a pharmaceutically acceptable anion (Z^A) and Z¹ is a suitable leaving group, with a compound of the formula:



wherein Y is as previously defined for a compound of the formula (I), said process being followed by either (a), where Z¹ is a suitable leaving group, exchange for a pharmaceutically acceptable anion (Z^A), or (b), optionally, where Z^A is a pharmaceutically acceptable anion, exchange for another pharmaceutically acceptable anion.

20

Preferred examples of Z are C₁-C₄ alkanesulphonyloxy, benzenesulphonyloxy, para-toluenesulphonyloxy, chloro, bromo and iodo.

An example of Z¹ is trifluoromethanesulphonyloxy.

25 Preferably, the leaving group in the compound of the formula (II) forms a pharmaceutically acceptable anion (Z/Z^A), e.g. methanesulphonyloxy/methanesulphonate, and therefore anion exchange at the end of the process is unnecessary.

-9-

It is possible to exchange pharmaceutically acceptable anions (re Z^A) in the work-up procedure, e.g. methanesulphonate may be exchanged to chloride by treatment of the isolated compound or the crude mixture with aqueous

5 hydrochloric acid solution.


The reaction of the compounds (II) and (III) is generally carried out in a suitable solvent, e.g. acetonitrile, at elevated temperatures, preferably at the reflux temperature thereof.

The starting materials of the formula (II) can be prepared as shown in


10 Scheme 1.

-10-


Scheme 1




-11-




$\text{CH}_3\text{SO}_2\text{Cl}$, acid acceptor (e.g. NEt_3)




Imidazole, optional additional acid acceptor (e.g. K_2CO_3)




aqueous mineral acid (e.g. 5N HCl)



Reduction (e.g. NaBH_4)



-12-



- 5 wherein R, R¹, Ar, W, X, Z and Z¹ are as previously defined for a compound of the formula (II) and L and L¹ are suitable leaving groups, e.g. chloro, bromo,

-13-

iodo, methanesulphonyloxy, trifluoromethanesulphonyloxy, benzenesulphonyloxy and para-toluenesulphonyloxy.


5 Suitable reaction conditions, reagents and solvents for carrying out any one of the steps shown in Scheme 1 will be well-known to those skilled in the art with reference to the Preparations herein.

 With regard to the last step in the reaction sequence, a compound of the formula (XV) can be converted to a compound of the formula (II) using
10 conventional conditions. For example, an alcohol of the formula (XV) can be converted to a compound of the formula (II) where Z is methanesulphonyloxy by treatment with methanesulphonyl chloride, triethylamine and dichloromethane, and a compound of the formula (II) wherein Z¹ is trifluoromethanesulphonyloxy may be prepared by treating an alcohol of the
15 formula (XV) with trifluoromethanesulphonic anhydride, optionally in the presence of a suitable acid acceptor, and in a suitable solvent, e.g. dichloromethane.


 The compounds of the formula (XII) can also be prepared as shown in Scheme 2:

-14-

Scheme 2



-15-



wherein R^1 and Ar are as previously defined for a compound of the formula (XII)
 5 and L^2 and L^3 are suitable leaving groups, e.g. as previously defined for L and L^1 .


-16-

Suitable reaction conditions, reagents and solvents for carrying out any one of the steps shown in Scheme 2 will be well-known to those skilled in the art with reference to the Preparations herein.

The compounds of the formula (XV) can also be prepared as shown in Scheme 3:


-17-

Scheme 3




-18-


MeSO₂Cl, acid acceptor (e.g. NEt₃)




Imidazole, optional additional acid acceptor (e.g. K₂CO₃)



R-W-COCl, acid acceptor
(e.g. triethylamine)



Deprotection




-19-

wherein R, R¹, Ar, W and X are as previously defined for a compound of the formula (XV), L⁴ and L⁵ are suitable leaving groups, e.g. as defined for L and L¹, and P is a suitable protecting group.

5 Examples of suitable protecting groups (P) together with methods for their removal can be found in the publication "Protective Groups in Organic Synthesis", T.W. Greene and P.G.M. Wuts, Second Edition, Wiley-Interscience. A preferred example of P is tetrahydropyran-2-yl that can be removed using
10 Amberlyst 15 (trade mark) ion exchange resin or methanol saturated with hydrogen chloride gas.

 Suitable reaction conditions, reagents and solvents for carrying out any one of the steps shown in Scheme 3 will be well known to those skilled in the art with reference to the Preparations herein.


 The compounds of the formula (XXXI) can also be prepared by reacting
15 a compound of the formula (XXIX) with a compound of the formula:



wherein R and W are as previously defined for a compound of the formula
20 (XXXI), optionally in the presence of an additional acid acceptor, e.g. potassium carbonate.

 The compounds of the formula (XXXII) may be prepared as shown in Scheme 4.

-20-

Scheme 4

5

wherein R and W are as previously defined for a compound of the formula (XXXII).

-21-

The compounds of the formula (III) can be prepared by a similar method to that described in Chem.Ber., 108, 3475 (1975).

Alternatively, the compounds of the formula (III) where Y is cyclohexyl
5 optionally substituted as previously defined for the definition of Y for a compound of the formula (I) can be prepared by catalytic hydrogenation of a compound of the formula:



(IIIA)

10

wherein Y is phenyl optionally substituted as previously defined for the above definition of Y. The reduction can be carried out under a hydrogen atmosphere using a suitable catalyst, e.g. rhodium-on-alumina, and in a suitable solvent, e.g. acetic acid.


15

The compounds of the formula (IIIA) may be prepared by similar methods to those described in Chem.Ber., 108, 3475 (1975) and J.Org.Chem., 22, 1484 (1957).


The compounds of the formula (III) can also be prepared as shown in Scheme 5:

-22-

Scheme 5



-23-



wherein Y is as previously defined for a compound of the formula (III).

- 5 Suitable reaction conditions, reagents and solvents for carrying out any – one of the steps shown in Scheme 5 will be well-known to those skilled in the art with reference to the Preparations herein.

-24-

All of the above reactions and the preparations of novel starting materials used in the preceding methods are conventional and appropriate reagents and reaction conditions for their performance or preparation as well as
5 procedures for isolating the desired products will be well known to those skilled in the art with reference to literature precedents and the Examples and Preparations hereto.

The affinity of the compounds of the formula (I) for the human NK₁ receptor can be determined in vitro by determining their ability to inhibit [³H]-
10 Substance P binding to membranes prepared from the human IM9 cell line expressing the human NK₁ receptor using a modification of the method described in McLean, S. *et al*, J. Pharm.Exp.Ther., 267, 472-9 (1993) in which whole cells were used.

The affinity of the compounds of formula (I) for the human NK₂ receptor
15 can be determined in vitro by determining their ability to compete with [³H]-NKA (neurokinin A) for binding to membranes prepared from Chinese hamster ovary cells expressing the cloned human NK₂ receptor. In this method, washed Chinese hamster ovary cell membranes are prepared as described for the previous method where IM9 cells are used instead. The membranes are
20 incubated (90 min, 25°C) with [³H]-NKA and with a range of concentrations of the test compound. Non-specific binding is determined in the presence of 10µM NKA.

The NK₁ receptor antagonist activity of the compounds of the formula (I) can be determined in vitro by testing their ability to antagonise the contractile
25 effects of Substance P in de-epithelialised guinea pig tracheal strips. Tissues can be prepared from guinea pigs (350-600g) which are killed by stunning and exsanguination. The excised trachea is cleared of connective tissue and opened longitudinally, opposite the trachealis muscle band. The epithelial layer can then be removed by rubbing the inner surface of the trachea with a cotton
30 bud. Strips of approximately 4 cartilage bands wide are cut and mounted under 1g tension in organ baths containing Krebs solution (composition: NaCl 118mM,

-25-

KCl 4.6mM, NaHCO₃ 25mM, KH₂PO₄ 1.4mM, MgSO₄ 1.2mM, CaCl₂ 2.5mM, glucose 11mM) at 37°C and gassed with 95% O₂/5% CO₂. The potential action of Substance P on the NK₂ receptor population found in this tissue can be prevented by the inclusion of the selective NK₂ receptor antagonist $\pm$ SR-48,968 (1 μ M) in the Krebs buffer solution. Additionally, indomethacin (3 μ M) is added to remove the influence of endogenous prostanoids. Tension changes of the tissue in response to cumulative addition of the agonist Substance P are recorded isometrically. The potency of the compounds of the formula (I) can be assessed by the magnitude of shift induced in the Substance P dose response curve, using standard Schild analysis, following 30 minutes incubation of the compound with the tissue.

The de-epithelialised guinea pig trachea strip preparation may also be used to evaluate the NK₂ receptor antagonist activity of the compounds of the formula (I) *in vitro* by using the selective NK₂ receptor agonist [β -Ala⁸]NKA₍₄₋₁₀₎ as the contractile agent. For such studies, strips are prepared and mounted in organ baths as described above, using Krebs solution of the following composition: NaCl 118mM, KCl 4.6mM, NaHCO₃ 25mM, KH₂PO₄ 1.4mM, MgSO₄ 1.2mM, CaCl₂ 2.5mM, glucose 11mM, indomethacin 3 μ M. The potency of the compounds may be assessed by the magnitude of the shift induced in the [β -Ala⁸]NKA₍₄₋₁₀₎ dose response curve, using standard Schild analysis, following 30 minutes incubation of the compound with the tissue.

The compounds of the formula (I) can be tested for NK₃ receptor antagonist activity, *in vitro*, by testing their ability to antagonise the contractile effects of the selective NK₃ receptor agonist senktide in the guinea-pig ileum using the method of Maggi *et al*, Br.J.Pharmacol., 101, 996-1000 (1990).

For human use, the compounds of the formula (I) can be administered alone, but will generally be administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice.

-26-

For example, they can be administered orally or sublingually in the form of tablets containing such excipients as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents.

They can be injected parenterally, for example, intravenously, intramuscularly or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood.

For oral and parenteral administration to human patients, the daily dosage level of the compounds of the formula (I) will be from 0.01 to 20mg/kg (in single or divided doses).

Thus tablets or capsules of the compounds will contain from 1mg to 1.0g of active compound for administration singly or two or more at a time, as appropriate. The physician in any event will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited and such are within the scope of this invention.

The compounds of formula (I) can also be administered intranasally or by inhalation and are conveniently delivered in the form of a dry powder inhaler or an aerosol spray presentation from a pressurised container or a nebuliser with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane (HFA 134A [trade mark]) or 1,1,1,2,3,3,3-heptafluoropropane (HFA 227EA [trade mark]), carbon dioxide or other suitable gas. In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurised container or nebuliser may contain a solution or suspension of the active compound, e.g.

-27-

using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g. sorbitan trioleate. Capsules and cartridges
5 (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated to contain a powder mix of a compound of the formula (I) and a suitable powder base such as lactose or starch.

Aerosol formulations are preferably arranged so that each metered dose or "puff" of aerosol contains from 20 μ g to 1000 μ g of a compound of formula (I)
10 for delivery to the patient. The overall daily dose with an aerosol will be in the range of from 20 μ g to 20mg which may be administered in a single dose or, more usually, in divided doses throughout the day.

Alternatively, the compounds of the formula (I) can be administered in the form of a suppository or pessary, or they may be applied topically in the
15 form of a lotion, solution, cream, ointment or dusting powder. For example, they can be incorporated into a cream consisting of an aqueous emulsion of polyethylene glycols or liquid paraffin, or they can be incorporated, at a concentration of from 1 to 10% by weight, into an ointment consisting of a white wax or white soft paraffin base together with such stabilisers and preservatives
20 as may be required. The compounds of the formula (I) may also be transdermally administered by the use of a skin patch.

It is to be appreciated that reference to treatment includes curative, palliative or prophylactic treatment.

Thus the invention further provides:-

- 25 (i) a pharmaceutical composition comprising a compound of the formula (I) together with a pharmaceutically acceptable diluent or carrier;
- (ii) a compound of the formula (I) or a pharmaceutically acceptable composition thereof, for use as a medicament;
- (iii) the use of a compound of the formula (I), or of a pharmaceutically
30 acceptable composition thereof, for the manufacture of a medicament for

-28-

the treatment of a disease by producing an antagonist effect on a tachykinin receptor or on a combination of tachykinin receptors;

- (iv) use as in (iii) where the antagonist effect is on the human NK₁ and NK₂ tachykinin receptors;
- 5 (v) use as in (iii) or (iv) where the disease is an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel
- 10 disease, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis or Crohn's disease, a disease caused by Helicobacter pylori or another urease-positive Gram negative bacteria, a urogenital tract disorder such as incontinence, hyperreflexia or cystitis, a
- 15 pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, emesis, cough, migraine or acute or chronic pain;
- 20 (vi) a method of treatment of a human to treat a disease by producing an antagonist effect on a tachykinin receptor or on a combination of tachykinin receptors, which comprises treating said human with an effective amount of a compound of the formula (I) or with a pharmaceutically acceptable composition thereof;
- 25 (vii) a method as in (vi) where the antagonist effect is on the human NK₁ and NK₂ tachykinin receptors;
- (viii) a method as in (vi) or (vii) where the disease is an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a

-29-

central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis or Crohn's disease, a disease caused by Helicobacter pylori or another urease-positive Gram negative bacteria, a urogenital tract disorder such as incontinence, hyperreflexia or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, emesis, cough, migraine or acute or chronic pain; and

(ix) a compound of the formula (II), (XI), (XII), (XIII), (XIV), (XV), (XXIII), (XXX) or (XXXI).


The following Examples illustrate the preparation of the compounds of the formula (I):-

THIS SHEET HAS BEEN
REPLACED

-30-

Example 14-Phenyl-1-(3-[3,4-dichlorophenyl]-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate

5



1-Methanesulphonyloxy-3-(3,4-dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]-butane (0.71g) (see Preparation 74) and 4-phenylquinuclidine (0.32g) (see J. Org. Chem., 22, 1484, (1957)) were dissolved in acetonitrile (10ml) and the mixture heated under reflux for 4 hours. The solvent was removed under reduced pressure and the resulting residue dissolved in dichloromethane before removal of the solvent under reduced pressure. The residue was chromatographed on silica gel eluting with a solvent gradient of 95:5 changing to 85:15, by volume, dichloromethane : methanol to give the product as a white foam. This was then triturated with diethyl ether, filtered and dried at room temperature under reduced pressure to give 4-phenyl-1-(3-[3,4-dichlorophenyl]-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate (0.71g) as a white solid.


$^1\text{H-NMR}$ (CDCl_3): δ = 7.62 (2H, s), 7.16-7.41 (10H, m), 7.11 (1H, s), 4.66-4.86 (2H, m), 3.52-3.81 (7H, m), 3.32-3.47 (1H, m), 2.91-3.08 (1H, m), 2.82 (3H, s), 2.22-2.50 (14H, m) ppm.

Found: C, 61.53; H, 6.17; N, 6.03. $\text{C}_{36}\text{H}_{41}\text{Cl}_2\text{N}_3\text{O}_4\text{S}$ requires C, 61.70; H, 6.18; N, 5.99%.

-31-

Examples 2-16





The compounds of the following tabulated Examples (Table 1) of the general formula :












were prepared by a similar method to that of Example 1 using the appropriate mesylate (see Preparations 75-84 and 86-90) and 4-phenylquinuclidine as the starting materials.

TABLE 1

Example no.	Starting material Prep.no.	R-W-	Analytical Data
2	75		¹ H-NMR (CDCl ₃): δ = 7.03-7.41 (13H, m), 4.80-4.97 (2H, m), 3.59-3.84 (7H, m), 3.38-3.50 (1H, m), 2.97-3.10 (1H, m), 2.84 (3H, s), 2.40-2.59 (2H, m), 2.23-2.34 (9H, m), 2.09 (3H, s) ppm. Found: C, 61.70; H, 6.01; N, 6.00. C ₃₆ H ₄₁ Cl ₂ N ₃ O ₄ S. 1.00 mol H ₂ O requires C, 61.70; H, 6.18; N, 6.00%.
3	76		¹ H-NMR (CDCl ₃): δ = 7.21-7.59 (13H, m), 7.11 (1H, s), 4.89 (1H, dd), 4.75 (1H, dd), 3.55-3.85 (7H, m), 3.30-3.42 (1H, m), 2.99-3.11 (1H, m), 2.84 (3H, s), 2.21-2.62 (8H, m) ppm. Found: C, 55.35; H, 4.86; N, 5.48. C ₃₅ H ₃₆ Cl ₂ F ₃ N ₃ O ₅ S. 0.4 mol CH ₂ Cl ₂ requires C, 55.03; H, 4.93; N, 5.43%.

4	77		¹ H-NMR (CDCl ₃): δ = 7.21-7.46 (10H, m), 7.01-7.16 (3H, m), 4.72-4.92 (2H, m), 3.52-3.85 (10H, m), 3.32-3.49 (1H, m), 3.00-3.12 (1H, m), 2.82 (3H, s), 2.42-2.56 (2H, m), 2.20-2.38 (9H, m) ppm. Found: C, 59.25; H, 5.86; N, 5.67. C ₃₆ H ₄₁ Cl ₂ N ₃ O ₅ S. 0.5 mol H ₂ O. 0.3 mol CH ₂ Cl ₂ requires C, 59.29; H, 5.84; N, 5.71%.
5	78		¹ H-NMR (CDCl ₃): δ = 7.60 (1H, d), 7.21-7.42 (10H, m), 7.09 (1H, s), 6.89 (1H, t), 4.82 (1H, dd), 4.57-4.71 (3H, m), 3.58-3.84 (7H, m), 3.41-3.52 (1H, m), 3.22 (2H, t), 3.02-3.14 (1H, m), 2.81 (3H, s), 2.20-2.55 (8H, m) ppm. Found: C, 60.32; H, 5.66; N, 5.88. C ₃₆ H ₃₉ Cl ₂ N ₃ O ₅ S. 0.3 mol CH ₂ Cl ₂ requires C, 60.38; H, 5.53; N, 5.82%.
6	79		¹ H-NMR (CDCl ₃): δ = 8.79 (1H, s), 8.11 (1H, d), 8.00 (1H, d), 7.89 (2H, t), 7.51-7.64 (2H, m), 7.48 (1H, s), 7.19-7.40 (9H, m), 4.71-4.90 (2H, m), 3.59-3.80 (7H, m), 3.03-3.51 (1H, m), 2.99-3.09 (1H, m), 2.88 (3H, s), 2.42-2.55 (2H, m), 2.22-2.38 (6H, m) ppm. Found: C, 64.08; H, 5.68; N, 5.90. C ₃₈ H ₃₉ Cl ₂ N ₃ O ₄ S 0.5 mol H ₂ O requires C, 63.95; H, 5.65; N, 5.89%.
7	80		¹ H-NMR (CDCl ₃): δ = 8.41-8.52 (2H, m), 7.51 (1H, s), 7.11-7.42 (10H, m), 4.72-4.89 (2H, m), 3.60-3.89 (7H, m), 3.31-3.46 (1H, m), 2.97-3.09 (1H, m), 2.85 (3H, s), 2.42-2.59 (2H, m), 2.21-2.39 (6H, m) ppm. Found: C, 56.15; H, 4.88; N, 5.77. C ₃₅ H ₃₅ Cl ₂ F ₄ N ₃ O ₄ S. 0.5 mol H ₂ O requires C, 56.08; H, 4.84; N, 5.61%.

8	81		¹ H-NMR (CDCl ₃): δ = 7.05-7.44 (13H, m), 4.80-4.94 (2H, m), 3.61-3.89 (7H, m), 3.38-3.71 (1H, m), 3.01-3.12 (1H, m), 2.76-2.91 (5H, m), 2.22-2.69 (10H, m), 1.63-1.86 (4H, m) ppm. Found: C, 64.11; H, 6.30; N, 6.00. C ₃₈ H ₄₃ Cl ₂ N ₃ O ₄ S requires C, 64.40; H, 6.12; N, 5.93%.
9	82		¹ H-NMR (CDCl ₃): δ = 7.78-7.86 (2H, m), 7.21-7.40 (9H, m), 7.11-7.16 (2H, m), 4.80 (1H, dd), 4.68 (1H, dd), 3.60-3.85 (7H, m), 3.37-3.49 (1H, m), 3.02-3.11 (1H, m), 2.79-2.90 (7H, m), 2.40-2.51 (2H, m), 2.22-2.36 (6H, m), 1.82 (4H, br. s) ppm. Found: C, 63.90; H, 6.25; N, 6.05. C ₃₈ H ₄₃ Cl ₂ N ₃ O ₄ S. 0.25 mol H ₂ O requires C, 63.99; H, 6.15; N, 5.89%.
10	83		¹ H-NMR (CDCl ₃): δ = 7.24-7.46 (11H, m), 7.08 (1H, s), 6.91 (1H, d), 4.89 (1H, dd), 4.72 (1H, dd), 3.64-3.88 (10H, m), 3.15-3.46 (1H, m), 3.07-3.17 (1H, m), 2.88 (3H, s), 2.46-2.56 (2H, m), 2.26-2.40 (6H, m) ppm.
11	84		¹ H-NMR (CDCl ₃): δ = 6.94-7.49 (14H, m), 4.90 (1H, dd), 4.70 (1H, dd), 3.61-3.84 (10H, m), 3.36-3.49 (1H, m), 3.09-3.20 (1H, m), 2.82 (3H, s), 2.22-2.59 (8H, m) ppm. Found: C, 58.80; H, 5.68; N, 5.88. C ₃₅ H ₃₉ Cl ₂ N ₃ O ₅ S. 0.5 mol H ₂ O requires C, 58.87; H, 5.38; N, 6.06%.
12	86		¹ H-NMR (CDCl ₃): δ = 7.72 (1H, d), 7.55-7.66 (2H, m), 7.20-7.49 (10H, m), 7.09 (1H, s), 4.94 (1H, dd), 4.72 (1H, dd), 3.60-3.85 (7H, m), 3.30-3.42 (1H, m), 3.05-3.18 (1H, m), 2.86 (3H, s), 2.36-2.59 (2H, m), 2.21-2.32 (6H, m) ppm. Found: C, 57.41; H, 5.13; N, 5.79. C ₃₅ H ₃₆ Cl ₂ F ₃ N ₃ O ₄ S. 0.5 mol H ₂ O requires C, 57.46; H, 5.10; N, 5.74%.


13	87		¹ H-NMR (CDCl ₃): δ = 7.20-7.46 (10H, m), 6.91-7.10 (4H, m), 4.83 (1H, dd), 4.43-4.68 (2H, m), 3.66-3.90 (7H, m), 3.32-3.48 (1H, m), 3.15-3.31 (1H, m), 2.42-2.58 (2H, m), 2.22-2.36 (6H, m), 1.11-1.21 (6H, m) ppm. Found: C, 61.88; H, 5.97; N, 6.11. C ₃₇ H ₄₃ Cl ₂ N ₃ O ₅ S. 0.25 mol H ₂ O requires C, 61.96; H, 6.11; N, 5.86%.
14	88		¹ H-NMR (CDCl ₃): δ = 7.15-7.41 (13H, m), 7.06 (1H, s), 4.80-4.91 (2H, m), 3.59-3.86 (7H, m), 3.34-3.49 (1H, m), 3.00-3.12 (1H, m), 2.83 (3H, s), 2.40-2.66 (4H, m), 2.21-2.35 (6H, m), 1.15 (3H, t) ppm. Found: C, 62.45; H, 6.11; N, 6.13. C ₃₆ H ₄₁ Cl ₂ N ₃ O ₄ S. 0.2 mol CH ₂ Cl ₂ requires C, 62.14; H, 5.96; N, 6.01%.
15	89		¹ H-NMR (CDCl ₃): δ = 6.83-7.53 (19H, m), 4.79 (1H, dd), 4.55 (1H, dd), 3.43-3.72 (7H, m), 3.03-3.30 (2H, m), 2.81 (3H, s), 2.09-2.49 (8H, m) ppm. Found: C, 63.02; H, 5.55; N, 5.72. C ₄₀ H ₄₁ Cl ₂ N ₃ O ₅ S. 0.25 mol CH ₂ Cl ₂ requires C, 62.95; H, 5.45; N, 5.47%.
16	90		¹ H-NMR (CDCl ₃): δ = 7.08-7.40 (15H, m), 4.74 (1H, dd), 4.61 (1H, dd), 4.46 (1H, d), 4.29 (1H, d), 3.48-3.71 (7H, m), 3.14-3.28 (1H, m), 2.88-3.04 (1H, m), 2.79 (3H, s), 2.15-2.39 (8H, m) ppm. Found: C, 61.41; H, 5.68; N, 6.19. C ₃₅ H ₃₉ Cl ₂ N ₃ O ₄ S. 0.2 mol CH ₂ Cl ₂ requires C, 61.66; H, 5.79; N, 6.13%.

-35-

Example 17

4-Phenyl-1-(3-[3,4-dichlorophenyl]-4-{2-[3,5-bis(trifluoromethyl)benzoyl]imidazol-1-yl}butyl)quinuclidinium chloride

5



1-Methanesulphonyloxy-3-(3,4-dichlorophenyl)-4-{2-[3,5-bis(trifluoromethyl)benzoyl]imidazol-1-yl}butane (0.7g) (see Preparation 85) and 4-phenylquinuclidine (0.32g) (see J. Org. Chem., 22, 1484,(1957)) were dissolved in acetonitrile (10ml) and the mixture heated under reflux for 18 hours. The solvent was removed under reduced pressure and the resulting residue dissolved in dichloromethane and washed twice with 2N aqueous hydrochloric acid solution. The organic phase was then dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 95:5 changing to 90:10, by volume, dichloromethane : methanol to give 4-phenyl-1-(3-[3,4-dichlorophenyl]-4-{2-[3,5-bis(trifluoromethyl)benzoyl]imidazol-1-yl}butyl)quinuclidinium chloride (0.37g) as a white foam.

¹H-NMR (CDCl₃): δ = 8.69 (2H, s), 8.04 (1H, s), 7.54 (1H, s), 7.14-7.41 (9H, m), 4.76-4.96 (2H, m), 3.70-3.98 (7H, m), 3.80-3.93 (1H, m), 3.10-3.21 (1H, m), 2.49-2.62 (2H, m), 2.23-2.40 (6H, m) ppm.


Found: C, 56.34; H, 4.53; N, 5.55. C₃₅H₃₂Cl₂F₆N₃O. 1.00 mol H₂O requires C, 56.12; H, 4.58; N, 5.61%.

-36-

Example 18

4-Phenyl-1-(3(S)-[3,4-dichlorophenyl]-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate

5



1-Methanesulphonyloxy-3(S)-[3,4-dichlorophenyl]-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butane (0.82g) (see Preparation 105) and 4-phenylquinuclidine (0.35g) (see J. Org. Chem., 22, 1484, (1957)) were dissolved in acetonitrile (10ml) and the mixture heated under reflux for 4 hours and then left to stand at room temperature overnight. The solvent was removed under reduced pressure and the residue chromatographed on silica gel eluting with a solvent gradient of 9:1 changing to 8:2, by volume, dichloromethane : methanol to give the product as a white foam. This was dissolved in dichloromethane, filtered and the solvent removed from the filtrate to give 4-phenyl-1-(3(S)-[3,4-dichlorophenyl]-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate (0.7g) as a white powder.

¹H-NMR (CDCl₃): δ = 7.05-7.44 (13H, m), 4.80-4.94 (2H, m), 3.61-3.89 (7H, m), 3.38-3.71 (1H, m), 3.01-3.12 (1H, m), 2.76-2.91 (5H, m), 2.22-2.69 (10H, m), 1.63-1.86 (4H, m) ppm.


Found: C, 63.81; H, 6.20; N, 5.99. C₃₈H₄₃Cl₂N₃O₄ requires C, 64.40; H, 6.12; N, 5.93%.

-37-

Example 19

4-Phenyl-1-(3(R)-[3,4-dichlorophenyl]-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate

5



This compound was prepared in an analogous fashion to Example 18 using 1-methanesulphonyloxy-3(R)-(3,4-dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-

10 naphthoyl)imidazol-1-yl]butane (see Preparation 106) and 4-phenylquinuclidine (0.36g) (see J. Org. Chem., 22, 1484 (1957)) as the starting materials.


¹H-NMR (CDCl₃): δ = 7.05-7.44 (13H, m), 4.80-4.94 (2H, m), 3.61-3.89 (7H, m), 3.38-3.71 (1H, m), 3.01-3.12 (1H, m), 2.76-2.91 (5H, m), 2.22-2.69 (10H, m), 1.63-1.86 (4H, m) ppm.

15 Found: C, 62.69; H, 6.07; N, 5.87. C₃₈H₄₃Cl₂N₃O₄S. 1.00 mol H₂O requires C, 62.80; H, 6.25; N, 5.78%.

-38-

Example 20

4-Cyclohexyl-1-(3-[3,4-dichlorophenyl]-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate



1-Methanesulphonyloxy-3-(3,4-dichlorophenyl)-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butane (0.59g) (see Preparation 84) and 4-cyclohexylquinuclidine (0.27g) (see Preparation 1) were dissolved in acetonitrile (8ml) and heated under reflux for 3.5 hours. The solvent was removed under reduced pressure and the resulting residue dissolved in dichloromethane and the solvent removed under reduced pressure. The residue was chromatographed on silica gel eluting with a solvent gradient of 95:5 changing to 85:15, by volume, dichloromethane : methanol to give 4-cyclohexyl-1-(3-[3,4-dichlorophenyl]-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate (0.69g) as a white foam.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.21-7.49 (6H, m), 6.94-7.08 (3H, m), 4.87 (1H, dd), 4.71 (1H, dd), 3.77 (3H, s), 3.33-3.70 (8H, m), 2.91-3.06 (1H, m), 2.83 (3H, s), 2.30-2.49 (2H, m), 1.60-1.89 (11H, m), 0.80-1.28 (6H, m) ppm.


Found: C, 56.76; H, 6.31; N, 5.63. $\text{C}_{35}\text{H}_{45}\text{Cl}_2\text{N}_3\text{O}_5\text{S}$. 0.5mol CH_2Cl_2 . 0.5 mol H_2O requires C, 56.64; H, 6.38; N, 5.66%.

-39-

Example 21

4-Cyclohexyl-1-(3-[3,4-dichlorophenyl]-4-[2-(3-trifluoromethyl-4-fluorobenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate

5



- 1-Methanesulphonyloxy-3-(3,4-dichlorophenyl)-4-[2-(3-trifluoromethyl-4-fluorobenzoyl)imidazol-1-yl]butane (0.77g) (see Preparation 80) and 4-cyclohexylquinuclidine (0.31g) (see Preparation 1) were dissolved in acetonitrile (10ml) and heated under reflux for 5 hours. The solvent was removed under reduced pressure, the resulting residue dissolved in dichloromethane and the solvent removed under reduced pressure. The residue was chromatographed on silica gel eluting with a solvent gradient of 95:5 changing to 80:20, by volume, dichloromethane : methanol to give 4-cyclohexyl-1-(3-[3,4-dichlorophenyl]-4-[2-(3-trifluoromethyl-4-fluorobenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate (0.69g) as a white foam.


-40-

¹H-NMR (CDCl₃): δ = 8.42-8.50 (2H, m), 7.55 (1H, s), 7.22-7.32 (3H, m), 7.11-7.20 (2H, m), 4.80-4.85 (2H, m), 3.34-3.76 (8H, m), 2.90-2.98 (1H, m), 2.85 (3H, s), 2.30-2.50 (2H, m), 1.75-1.81 (6H, m), 1.61-1.72 (5H, m), 1.02-1.27 (4H, m), 0.81-0.95 (2H, m) ppm.

Found: C, 56.14; H, 5.62; N, 5.70. C₃₅H₄₁Cl₂F₄N₃O₄S requires C, 56.30; H, 5.53; N, 5.63%.

Example 22

10 4-(3,5-Dimethylphenyl)-1-(3-[3,4-dichlorophenyl]-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate



15 1-Methanesulphonyloxy-3-(3,4-dichlorophenyl)-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butane (0.55g) (see Preparation 84) and 4-(3,5-dimethylphenyl)quinuclidine (0.28g) (see Preparation 12) were dissolved in acetonitrile (10ml) and heated under reflux for 2.5 hours. The solvent was removed under reduced pressure, the resulting residue dissolved in dichloromethane and the solvent removed under reduced pressure. The
20 residue was chromatographed on silica gel eluting with a solvent gradient of 95:5 changing to 90:10, by volume, dichloromethane : methanol to give 4-(3,5-dimethylphenyl)-1-(3-[3,4-dichlorophenyl]-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate (0.69g) as a white foam.


-41-

¹H-NMR (CDCl₃): δ = 7.28-7.48 (5H, m), 7.23 (1H, d), 6.92-7.07 (3H, m), 6.88 (1H, s), 6.82 (2H, s), 4.89 (1H, dd), 4.71 (1H, dd), 3.53-3.80 (10H, m), 3.34-3.47 (1H, m), 3.01-3.12 (1H, m), 2.82 (3H, s), 2.42-2.51 (2H, m), 2.20-2.34 (12H, m) ppm.

5

Example 23

4-(2-Naphthyl)-1-(3-[3,4-dichlorophenyl]-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate



10

1-Methanesulphonyloxy-3-(3,4-dichlorophenyl)-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butane (0.55g) (see Preparation 84) and 4-(2-naphthyl)quinuclidine (0.31g) (see Preparation 7) were dissolved in acetonitrile (10ml) and heated under reflux for 2.5 hours. The solvent was removed under reduced pressure and the resulting residue dissolved in dichloromethane and the solvent removed under reduced pressure. The residue was chromatographed on silica gel eluting with a solvent gradient of 95:5 changing to 85:15, by volume, dichloromethane : methanol to give 4-(2-naphthyl)-1-(3-[3,4-dichlorophenyl]-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate (0.65g) as a white foam.

20


¹H-NMR (CDCl₃): δ = 7.71-7.86 (3H, m), 7.62 (1H, s), 7.23-7.51 (8H, m), 7.17 (1H, s), 6.92-7.07 (3H, m), 4.89 (1H, dd), 4.70 (1H, dd), 3.61-3.86 (10H, m), 3.34-3.66 (1H, m), 3.03-3.16 (1H, m), 2.84 (3H, s), 2.30-2.55 (8H, m) ppm.

The following Preparations describe the preparation of certain starting materials used in the syntheses of the compounds of the preceding Examples.

5

Preparation 1

4-Cyclohexylquinuclidine




- 10 4-Phenylquinuclidine (5g) (see J. Org. Chem., 22, 1484, (1957)) was dissolved in glacial acetic acid (25ml), 5% w/w rhodium-on-alumina (2g) was added and the mixture hydrogenated for 5 days at 345kPa (50psi). The mixture was filtered through a short column of Arbacel (trade mark) filter aid and the pad washed with methanol. The filtrate was collected and the solvent removed under reduced pressure to give a residue which
- 15 was dissolved in water. The pH was adjusted to >10 by addition of 0.88 aqueous ammonia solution. The aqueous mixture was extracted with ethyl acetate (x3) and the organic layers combined, washed with brine, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give 4-cyclohexylquinuclidine (4.7g) as a pale pink solid.

20

¹H-NMR (CDCl₃): δ = 2.75-2.96 (6H, m), 1.60-1.85 (5H, m), 1.06-1.45 (9H, m), 1.80-1.98 (3H, m) ppm.


-43-

Preparation 2Ethyl 2-(1-benzylpiperidin-4-ylidene)-2-cyanoacetate

N-Benzylpiperidin-4-one (25g), ethyl cyanoacetate (16.4g), glacial acetic acid (6ml) and ammonium acetate (2.54g) were heated together in toluene (200ml), with removal of water using a Dean and Stark apparatus, for 90 minutes. The mixture was cooled, a further amount of toluene (100ml) added and the solution washed sequentially with water (100ml) and brine (100ml) at which point a red oily product precipitated. The phases were separated and the oily product dissolved in dichloromethane. The toluene and dichloromethane solutions were combined and the solvents removed under reduced pressure to give a residue. The residue was dissolved in dichloromethane and washed sequentially with water and saturated aqueous sodium hydrogen carbonate solution before removal of the solvent under reduced pressure. The crude product was chromatographed on silica gel eluting with 98:2, by volume, dichloromethane : methanol to give the title compound (30.8g) as an oil.

20 $^1\text{H-NMR}$ (CDCl_3): δ = 7.24-7.34 (5H, m), 4.22-4.32 (2H, q), 3.54 (2H, s), 3.12-3.16 (2H, m), 2.79 (2H, d), 2.66 (2H, d), 2.56 (2H, d), 1.32 (3H, t) ppm.

-44-

Preparation 3Ethyl 2-cyano-2-(1-benzyl-4-(2-naphthyl)piperidin-4-yl)ethanoate

2-Bromonaphthalene (26g) was dissolved in anhydrous diethyl ether (100ml) and 1/5 of this solution added to a vigorously stirred mixture of magnesium turnings (3.3g) and 2-3 iodine crystals under a nitrogen atmosphere. Gentle heating was applied to initiate formation of the Grignard reagent since it proved difficult to maintain spontaneous refluxing. The remainder of the 2-bromonaphthalene solution was added in 4 portions, allowing the refluxing to subside after each addition, and the mixture finally heated under reflux for 1 hour after which time two dark organic phases had formed. The mixture was cooled in an ice-bath and anhydrous tetrahydrofuran (50ml) added followed by dropwise addition of a solution of ethyl 2-(1-benzylpiperidin-4-ylidene)-2-cyanoacetate (12g) (see Preparation 2) in tetrahydrofuran (100ml). The single phase mixture was stirred at 0°C for 30 minutes before being left to stand at room temperature overnight. The mixture was then poured into saturated aqueous ammonium chloride solution (450ml) and extracted twice with diethyl ether. The organic extracts were combined and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 4:0 changing to 4:1, by volume, dichloromethane : diethyl ether to give the title compound (5.9g) as an oil.


-45-

$^1\text{H-NMR}$ (CDCl_3): δ = 7.72-7.90 (4H, m), 7.46-7.57 (3H, m), 7.20-7.38 (5H, m), 3.85 (2H, q), 3.71 (1H, s), 3.37 (2H, s), 2.60-2.80 (4H, m), 2.15-2.40 (4H, m), 0.80 (3H, t) ppm.

5

Preparation 4

Ethyl 2-(1-benzyl-4-(2-naphthyl)piperidin-4-yl)ethanoate



10

A solution of potassium hydroxide (10g) in water (30ml) was added to a solution of ethyl 2-cyano-2-(1-benzyl-4-(2-naphthyl)piperidin-4-yl)ethanoate (5.9g) (see Preparation 3) in ethanol (40ml) resulting in formation of a suspension which was stirred at room temperature overnight and then heated on a steam bath until a clear solution was formed. The solution was divided between two microwave vessels and microwaved at 690kPa (100psi)/100% power for 5 hours. The solutions were then combined, the solvent removed under reduced pressure and the residue azeotroped 4 times with toluene to remove any residual water. The residue was dissolved in ethanol (250ml), cooled in an ice-bath and the solution saturated with hydrogen chloride gas at which stage a precipitate was formed. The mixture was left to stand at room temperature for 3 days, filtered and the solvent removed from the filtrate under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 98:2 changing to 96:4, by volume, dichloromethane : methanol to give the hydrochloride of the title compound. The hydrochloride salt was partitioned between

20

-46-

dichloromethane and saturated aqueous sodium carbonate solution, the organic phase separated, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give the title compound (3.6g) as an oil.


5

$^1\text{H-NMR}$ (CDCl_3): δ = 7.70-7.85 (4H, m), 7.40-7.52 (3H, m), 7.20-7.32 (5H, m), 3.78 (2H, q), 3.41 (2H, s), 2.58-2.70 (4H, m), 2.30-2.50 (4H, m), 2.10-2.20 (2H, m), 0.84 (3H, t) ppm.

10

Preparation 5

4-(2-Naphthyl)-4-(2-hydroxyethyl)-N-benzylpiperidine



- 15 Ethyl 2-(1-benzyl-4-(2-naphthyl)piperidin-4-yl)ethanoate (5.8g) (see Preparation 4) was dissolved in anhydrous diethyl ether (100ml), cooled in an ice-bath and lithium aluminium hydride (0.57g) added, portionwise. The mixture was stirred at room temperature for 30 minutes, water (0.8ml) was then carefully added followed by 2N aqueous sodium hydroxide solution (0.8ml) and further water (1.6ml). The mixture was
- 20 stirred for 20 minutes and the resulting granular precipitate removed by filtration. The solvent was removed from the filtrate under reduced pressure to give a solid which was dissolved in dichloromethane, dried over anhydrous sodium sulphate and the solvent again removed under reduced pressure to yield 4-(2-naphthyl)-4-(2-hydroxyethyl)-N-benzylpiperidine (2.2g) as a white solid.


-47-

The granular precipitate was triturated with dichloromethane and filtered. The filtrate was washed with water, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to yield a second crop of 4-(2-naphthyl)-4-(2-hydroxyethyl)-N-benzylpiperidine (2.7g) as a white solid.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.75-7.86 (3H, m), 7.70 (1H, s), 7.43-7.50 (3H, m), 7.20-7.35 (5H, m), 3.35-3.45 (4H, m), 2.60-2.70 (2H, m), 2.25-2.40 (4H, m), 1.92-2.04 (4H, m), 0.90 (1H, br. s) ppm.

Preparation 6

4-(2-Naphthyl)-N-benzylquinuclidinium 4-methylphenylsulphonate



15

4-(2-Naphthyl)-4-(2-hydroxyethyl)-N-benzylpiperidine (4.9g) (see Preparation 5) was dissolved in pyridine (30ml) and cooled in an ice-bath before addition of 4-methylphenylsulphonyl chloride (3.0g). The mixture was left at 0°C for 16 hours before removal of the solvent under reduced pressure. The residue was suspended in 10% w/w aqueous potassium carbonate solution (120ml) and extracted with toluene (2x130ml). The combined organic phases were washed with 10% w/w aqueous potassium carbonate solution, dried over anhydrous potassium carbonate for 10 minutes and filtered. The filtrate was collected and the volume reduced to ~60ml under reduced pressure to yield

20

-48-


a suspension. This was then heated at 90°C for 6 hours, left to stand at room temperature overnight and the resulting precipitate filtered off. The precipitate was washed sequentially with toluene and diethyl ether and dried to yield 4-(2-naphthyl)-N-benzylquinuclinium 4-methylphenylsulphonate (3.5g) as a white solid.

¹H-NMR (CDCl₃): δ = 7.90 (2H, d), 7.78 (3H, m), 7.30-7.65 (9H, m), 7.16 (2H, d), 4.89 (2H, s), 3.80-3.95 (6H, m), 2.31 (3H, s), 2.13-2.29 (6H, m) ppm.

10

Preparation 7

4-(2-Naphthyl)quinuclidine



15 4-(2-Naphthyl)-N-benzylquinuclinium 4-methylphenylsulphonate (3.5g) (see Preparation 6) was dissolved in methanol (35ml), 10% w/w palladium-on-carbon (0.5g) was added and the mixture hydrogenated for 40 hours at 345kPa (50psi). The mixture was then filtered through a short column of Arbacel (trade mark) filter aid and the solvent removed from the filtrate under reduced pressure to give a residue. This was
20 partitioned between diethyl ether and 1N aqueous sodium hydroxide solution. The two phases were separated and the aqueous phase extracted twice with diethyl ether. The organic phases were combined and the solvent removed under reduced pressure. The resulting residue was dissolved in ethyl acetate and the solution washed sequentially

-49-

with 0.88 aqueous ammonia solution and brine before being dried over anhydrous sodium sulphate. The solvent was again removed under reduced pressure yield 4-(2-naphthyl)quinuclidine (1.56 g) as a solid.


5

$^1\text{H-NMR}$ (CDCl_3): δ = 7.80 (3H, d), 7.63 (1H, s), 7.40-7.52 (3H, m), 3.00-3.15 (6H, m), 1.82-1.96 (6H, m) ppm.

Preparation 8

10

Ethyl 2-cyano-2-(1-benzyl-4-(3,5-dimethylphenyl)piperidin-4-yl)ethanoate



1,3-Dimethyl-5-bromobenzene (15.6g) was dissolved in anhydrous diethyl ether (60ml) and added to a vigorously stirred mixture of magnesium turnings (2.2g) and 2-3 iodine crystals under a nitrogen atmosphere. Gentle heating was applied to initiate formation of the Grignard reagent and once the spontaneous refluxing had subsided the mixture was heated under reflux for a further 30 minutes. The mixture was cooled in an ice-bath and a solution of ethyl 2-(1-benzylpiperidin-4-ylidene)-2-cyanoacetate (8.0g) (see Preparation 2) in anhydrous tetrahydrofuran (80ml) added over 20 minutes. The mixture was stirred at 0°C for 15 minutes before being left to stand at room temperature for 3 days. The mixture was then poured into saturated aqueous ammonium chloride solution - (300ml) and extracted twice with diethyl ether. The organic extracts were combined and

-50-

the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 3:0 changing to 3:1, by volume, dichloromethane : diethyl ether to give the title compound (4.7g) as an oil.


5

$^1\text{H-NMR}$ (CDCl_3): δ = 7.20-7.35 (5H, m), 6.91 (3H, s), 3.94 (2H, q), 3.61 (1H, s), 3.40 (2H, s), 2.40-2.70 (4H, m), 2.30 (6H, s), 2.10-2.22 (4H, m), 1.00 (3H, t) ppm.

Preparation 9

10

Ethyl 2-(1-benzyl-4-(3,5-dimethylphenyl)piperidin-4-yl)ethanoate



A solution of potassium hydroxide (3.4g) in water (20ml) was added to a solution of ethyl 2-cyano-2-(1-benzyl-4-(3,5-dimethylphenyl)piperidin-4-yl)ethanoate (4.7g) (see Preparation 8) in ethanol (24ml) and the mixture microwaved at 345kPa (100psi)/100% power for 2.5 hours. The solvent was removed under reduced pressure and the residue azeotroped with toluene to remove any residual water. The residue was dissolved in ethanol (100ml), cooled in an ice-bath and the solution saturated with hydrogen chloride gas. The mixture was left to stand at room temperature overnight before re-saturating the solution with hydrogen chloride gas and leaving to stand for a further 24 hours. The solution was filtered and the solvent removed from the filtrate under reduced pressure to give a residue which was dissolved in saturated aqueous sodium carbonate solution and extracted twice with dichloromethane. The organic phases were combined, dried over


-51-

anhydrous sodium sulphate and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 97:3 changing to 90:10, by volume, dichloromethane : methanol to give the title compound (2.0g) as an oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.20-7.35 (5H, m), 6.90 (2H, s), 6.82 (1H, s), 3.80-3.90 (2H, q), 3.43 (2H, s), 2.50-2.63 (4H, m), 2.20-2.40 (10H, m), 1.92-2.08 (2H, m), 0.95-1.03 (3H, t) ppm.

Preparation 10

4-(3,5-Dimethylphenyl)-4-(2-hydroxyethyl)-N-benzylpiperidine




Ethyl 2-(1-benzyl-4-(3,5-dimethylphenyl)piperidin-4-yl)ethanoate (2.0g) (see Preparation 9) was dissolved in anhydrous diethyl ether (50ml), cooled in an ice-bath and lithium aluminium hydride (0.21g) added portionwise. The mixture was stirred at room temperature for 45 minutes, water (0.3ml) was carefully added, followed by 2N aqueous sodium hydroxide solution (0.3ml) and further water (0.6ml). The mixture was stirred for 20 minutes and the resulting granular precipitate removed by filtration and washed with diethyl ether. The solvent was removed from the filtrate under reduced pressure to give a solid which was dissolved in dichloromethane, dried over anhydrous sodium sulphate and the solvent again removed under reduced pressure to yield 4-(3,5-dimethylphenyl)-4-(2-hydroxyethyl)-N-benzylpiperidine (1.67g) as an oil.

-52-

$^1\text{H-NMR}$ (CDCl_3): $\delta = 7.20-7.35$ (5H, m), 6.90 (2H, s), 6.83 (1H, s), 3.32-3.43 (4H, m), 2.54-2.64 (2H, m), 2.13-2.37 (11H, m), 1.80-1.90 (4H, m) ppm.

5

Preparation 114-(3,5-Dimethylphenyl)-N-benzylquinuclidinium 4-methylphenylsulphonate

10

4-(3,5-Dimethylphenyl)-4-(2-hydroxyethyl)-N-benzylpiperidine (1.6g) (see Preparation 10) was dissolved in pyridine (10ml), cooled in an ice-bath and 4-methylphenylsulphonyl chloride (1.0g) added in 4 portions over 10 minutes. The mixture was left at 0°C for 16 hours before removal of the solvent under reduced

15 pressure. The residue was suspended in 10% w/w aqueous potassium carbonate solution (30ml) and extracted with toluene (50ml). The organic phase was washed with 10% w/w aqueous potassium carbonate solution, dried over anhydrous potassium carbonate and filtered. The filtrate was collected and the volume reduced to ~12ml under reduced pressure to yield a suspension which was then heated at 90°C for 3 hours, allowed to

20 cool room temperature and the volume reduced to ~5ml at which point the product began to precipitate out of solution. Diethyl ether (30ml) was added, the precipitate filtered, washed with diethyl ether and dried to yield 4-(3,5-dimethylphenyl)-N-benzylquinuclidinium 4-methylphenylsulphonate (1.59g) as a white solid.


-53-

$^1\text{H-NMR}$ (CDCl_3): δ = 7.96 (2H, d), 7.58 (2H, d), 7.30-7.40 (3H, m), 7.16 (2H, d), 6.96 (1H, s), 6.75 (2H, s), 4.86 (2H, d), 3.72-3.82 (6H, m), 2.34 (3H, s), 2.25 (6H, s), 2.05-2.12 (6H, m) ppm.

5

Preparation 12

4-(3,5-Dimethylphenyl)quinuclidine



10

4-(3,5-Dimethylphenyl)-N-benzylquinuclidinium 4-methylphenylsulphonate (1.4g) (see Preparation 11) was dissolved in methanol (14ml), 10% w/w palladium-on-carbon (0.2g) was added and the mixture hydrogenated for 40 hours at 345kPa (50psi). The mixture was then filtered through a short column of Arbocel (trade mark) filter aid and the solvent removed under reduced pressure to give a residue. This was partitioned between diethyl ether and 1N aqueous sodium hydroxide solution. The two phases were separated and the aqueous phase extracted twice with diethyl ether. The organic phases were combined and the solvent removed under reduced pressure. The resulting residue was dissolved in ethyl acetate and the solution washed with brine before being dried over anhydrous sodium sulphate. The solvent was again removed under reduced pressure yield 4-(3,5-dimethylphenyl)quinuclidine (0.59g) as a white solid.


20

$^1\text{H-NMR}$ (CDCl_3): δ = 6.89 (2H, s), 6.83 (1H, s), 2.95-3.05 (6H, m), 2.33 (6H, s), 1.70-1.81 (6H, m) ppm.

-54-

Preparation 132-(3,4-Dichlorophenyl)-3-(1,3-dioxolan-2-yl)propanenitrile

5




- Sodium hydride (60% w/w dispersion in mineral oil) (4.73g) was suspended in tetrahydrofuran (70ml) under a nitrogen atmosphere and the mixture cooled in an ice-bath. A solution of 3,4-dichlorophenylacetonitrile (20g) in tetrahydrofuran (80ml) was added dropwise over 35 minutes, the mixture allowed to warm to room temperature and left to stir for 16 hours. 2-Bromomethyl-1,3-dioxolane (19.71g) and tetra-n-butylammonium iodide (2g) were added and the resulting mixture was heated at reflux for 4 hours. The mixture was cooled and partitioned between ethyl acetate and water.
- The organic layer was separated and washed with brine. The organic solvents were removed under reduced pressure to yield a brown oil which was chromatographed on silica gel using 80:20, by volume, ethyl acetate : hexane as the eluent to yield the product as an orange mobile oil. This oil was then dissolved in methanol, cooled in ice and the precipitate that formed was filtered off, washed with methanol and dried under reduced pressure to yield 2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propanenitrile (15.8g) as a white solid.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.40-7.50 (2H, m), 7.20-7.25 (1H, dd), 4.95 (1H, dd), 3.82-4.05 (5H, m), 2.30-2.40 (1H, m), 2.05-2.15 (1H, m) ppm.

-55-

Preparation 142-(3,4-Dichlorophenyl)-3-(1,3-dioxolan-2-yl)propan-1-al

5



2-(3,4-Dichlorophenyl)-3-(1,3-dioxolan-2-yl)propanenitrile (64g) (see Preparation 13) was suspended in anhydrous toluene (500ml) and cooled to -70°C under a nitrogen atmosphere. Diisobutylaluminium hydride (200ml of a 1.5M solution in toluene) was then added over 50 minutes, by which time a clear solution was achieved. The mixture was stirred at -70°C for a further 30 minutes and then allowed to warm slowly to -20°C . Water (24ml) was carefully added (exothermic reaction) before pouring the mixture into a 15% (by weight) aqueous solution of citric acid (1800ml). Toluene (100ml) was added and the mixture was vigorously stirred for 1 hour. The resulting emulsion was filtered through a short column of Arbacel (trade mark) filter aid to give two clear phases which were separated. The organic phase was washed with brine, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give 2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propan-1-al as a yellow oil (53.5g)


20

$^1\text{H-NMR}$ (CDCl_3): δ = 9.90 (1H, s), 7.10-7.50 (3H, m), 4.89 (1H, t), 3.70-4.00 (5H, m), 1.45-1.55 (1H, m), 2.20-2.10 (1H, m) ppm.

-56-

Preparation 152-(3,4-Dichlorophenyl)-3-(1,3-dioxolan-2-yl)propan-1-ol

5



Sodium borohydride (5g) was added in two portions, over 40 minutes, to an ice-cooled solution of 2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propan-1-al (53.5g) (see
10 Preparation 14) in ethanol (300ml). The mixture was stirred for a further 30 minutes before removing the solvent under reduced pressure to give a residue. This was suspended in water (200ml), cooled to 0°C and the mixture acidified (pH<6) with glacial acetic acid. Dichloromethane was added and the aqueous phase basified (pH>8) by addition of solid sodium carbonate. A further amount of dichloromethane (200ml)
15 was added and the organic and aqueous phases separated. The aqueous phase was further extracted with dichloromethane (200ml). The organic phases were combined, washed with brine, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give 2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propan-1-ol (54.6g) as a yellow oil.


20

¹H-NMR (CDCl₃): δ = 7.30-7.45 (2H, m), 7.09 (1H, dd), 4.79 (1H, t), 3.70-4.00 (6H, m), 2.97-3.08 (1H, m), 1.96-2.09 (3H, m) ppm.

-57-

Preparation 161-Methanesulphonyloxy-2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propane

5




Methanesulphonyl chloride (5.5g) was added over 10 minutes to an ice-cooled solution of 2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propan-1-ol (12g) (see Preparation 15) and triethylamine (5.7g) in dichloromethane (100ml). The mixture was stirred for 30 minutes before further addition of dichloromethane (50ml). The solution was then washed sequentially with water (3x 50ml) and brine (50ml) before drying over anhydrous sodium sulphate. Removal of the solvent under reduced pressure gave 1-methanesulphonyloxy-2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propane as a yellow oil (15.6g) which solidified upon standing.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.41 (1H, d), 7.34 (1H, d), 7.10 (1H, dd), 4.75 (1H, t), 4.26-4.43 (2H, m), 3.90-4.00 (2H, m), 3.80-3.87 (2H, m), 3.30 (1H, m), 2.90 (3H, s), 2.00-2.10 (2H, m) ppm.

-58-

Preparation 17

1-(Imidazol-1-yl)-2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propane

5


1-Methanesulphonyloxy-2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propane (15.5g) (see Preparation 16) and imidazole (9g) were dissolved in anhydrous acetonitrile (100ml) and the mixture heated at reflux for 90 hours. The solvent was removed under reduced pressure, the residue dissolved in dichloromethane (100ml) and the solvent again removed under reduced pressure. The residue was dissolved in dichloromethane (300ml) and washed with sufficient aqueous sodium carbonate solution to ensure that the aqueous phase reached pH14. The two phases were separated and the aqueous phase extracted with dichloromethane (100ml). The organic phases were then combined and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel, eluting with a solvent gradient of 99:1 changing to 95:5, by volume, dichloromethane : methanol to give 1-(imidazol-1-yl)-2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propane (11.1g) as an orange oil.

15

¹H-NMR (CDCl₃): δ = 6.95-7.40 (4H, m), 6.85 (1H, dd), 6.69 (1H, s), 4.69 (1H, m), 4.15-4.25 (1H, m), 3.70-4.10 (5H, m), 3.15-3.25 (1H, m), 1.90-2.10 (2H, m) ppm.

20

-59-


Preparation 183-(3,4-Dichlorophenyl)-4-(imidazol-1-yl)butan-1-al

5N Aqueous hydrochloric acid solution (100ml) was added to an ice-cooled solution of 1-(imidazol-1-yl)-2-(3,4-dichlorophenyl)-3-(1,3-dioxolan-2-yl)propane (11g) (see Preparation 17) in tetrahydrofuran (100ml). The mixture was allowed to warm slowly to room temperature and then left to stand for 24 hours. The tetrahydrofuran was removed under reduced pressure and the acidic phase extracted with dichloromethane (2x 100ml). The organic phases were combined, washed with brine and dried over anhydrous sodium sulphate. The solvent was removed under reduced pressure to give the crude product which was chromatographed on silica gel eluting with a solvent gradient of 97.5:2.5 changing to 95:5, by volume, dichloromethane : methanol, to give 3-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-al (4.4g) as a viscous oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 9.70 (1H, s), 7.38 (1H, d), 7.20-7.30 (2H, m), 7.01 (1H, s), 6.89 (1H, dd), 6.71 (1H, s), 4.00-4.22 (2H, m), 3.60 (1H, m), 2.72-2.92 (2H, m) ppm.

-60-

Preparation 19


3-(3,4-Dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol

5

Sodium borohydride (0.52g) was added in three portions, over 5 minutes, to an ice-cooled solution of 3-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-al (3.3g) (see Preparation 18) in ethanol (25ml). The mixture was stirred for a further hour before
10 removing the solvent under reduced pressure to give a residue. The residue was then suspended in water (50ml), cooled to 0°C and the mixture first acidified to pH1 with 2N aqueous hydrochloric acid solution and then basified to pH14 by addition of solid sodium carbonate before being extracted with ethyl acetate (3x 200ml). The organic
15 phases were combined, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to yield 3-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol (2.84g) as a cream solid.

¹H-NMR (CDCl₃): δ = 7.35 (1H, d), 7.15-7.30 (2H, m), 6.95 (1H, s), 6.89 (1H, d), 6.70 (1H, s), 4.00-4.25 (2H, m), 3.60-3.70 (1H, m), 3.40-3.50 (1H, m), 3.15-3.30 (1H, m),
20 2.10 (1H, br. s), 1.75-2.00 (2H, m) ppm.

-61-


Preparation 204-Cyano-4-(3,4-dichlorophenyl)but-1-ene

3,4-Dichlorophenylacetonitrile (80g) was dissolved in anhydrous tetrahydrofuran (800ml) under a nitrogen atmosphere, cooled to -70°C , and lithium di-isopropylamide (320ml of a 1.5M solution in cyclohexane) added. The mixture was stirred at -70°C for 10 30 minutes, allyl bromide (63g) added over 15 minutes and the mixture stirred for a further 30 minutes. 2N Aqueous hydrochloric acid solution (600ml) was then added and the mixture extracted twice with diethyl ether. The organic extracts were combined and the solvent removed under reduced pressure. The resulting residue was dissolved in 15 dichloromethane, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give a red mobile oil. This was chromatographed on silica gel eluting with 4:1, by volume, hexane : dichloromethane to give 4-cyano-4-(3,4-dichlorophenyl)but-1-ene (94.8g) as an oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.40-7.51 (2H, m), 7.14-7.21 (1H, m), 5.70-5.95 (1H, m), 5.13-20 5.27 (2H, m), 3.84 (1H, t), 2.56-2.70 (2H, m) ppm.

-62-

Preparation 21


4-(3,4-Dichlorophenyl)-4-formylbut-1-ene

4-Cyano-4-(3,4-dichlorophenyl)but-1-ene (13g) (see Preparation 20) was dissolved in anhydrous toluene (100ml) and cooled to -70°C under a nitrogen atmosphere.

Diisobutylaluminium hydride (50ml of a 1.5M solution in toluene) was added to the solution over 30 minutes, the mixture stirred at -70°C for a further 30 minutes and then allowed to warm slowly to -10°C. Water (6ml) was carefully added (exothermic reaction) before pouring the mixture into a 15% (by weight) aqueous solution of citric acid (500ml), followed by further addition of toluene (300ml) and vigorous stirring of the mixture for 30 minutes. The two phases were separated, the organic phase washed with brine, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give 4-(3,4-dichlorophenyl)-4-formylbut-1-ene as an oil (14g).

¹H-NMR (CDCl₃): δ = 9.68 (1H, s), 7.48 (1H, d), 7.31 (1H, s), 7.03 (1H, d), 5.61-5.77 (1H, m), 4.94-5.13 (2H, m), 3.56-3.65 (1H, m), 2.77-2.91 (1H, m), 2.40-2.54 (1H, m) ppm.

-63-

Preparation 224-(3,4-Dichlorophenyl)-5-hydroxypent-1-ene

5

Sodium borohydride (2.2g) was added in four portions, over 10 minutes, to an ice-cooled solution of 4-(3,4-dichlorophenyl)-4-formylbut-1-ene (13g) (see Preparation 21) in ethanol (100ml). The mixture was stirred for a further 30 minutes before removing the solvent under reduced pressure to give a residue. This was suspended in water
10 (50ml), cooled to 0°C and the mixture acidified (pH<6) with 2N aqueous hydrochloric acid solution. The mixture was extracted three times with dichloromethane, the organic phases combined and the solvent removed under reduced pressure. Dichloromethane (200ml) was then added, the mixture stirred and filtered and the solvent removed from the filtrate under reduced pressure to give an oil. This was chromatographed on silica
15 gel eluting with dichloromethane to give 4-(3,4-dichlorophenyl)-5-hydroxypent-1-ene (7g) as a orange oil.

¹H-NMR (CDCl₃): δ = 7.24-7.45 (2H, m), 7.06 (1H, dd), 5.61-5.75 (1H, m), 4.96-5.10 (2H, m), 3.70-3.90 (2H, m), 2.80-2.91 (1H, m), 2.30-2.55 (2H, m), 1.32 (1H, t) ppm.

-64-

Preparation 234-(3,4-Dichlorophenyl)-5-(methanesulphonyloxy)pent-1-ene

A solution of methanesulphonyl chloride (2.3g) in dichloromethane (10ml) was added to an ice-cooled solution of 4-(3,4-dichlorophenyl)-5-hydroxypent-1-ene (4.3g) (see Preparation 22) and triethylamine (2.5g) in dichloromethane (50ml) over 10 minutes. The mixture was stirred for a further 30 minutes before being washed sequentially with water (3x 25ml) and brine (25ml), then dried over anhydrous sodium sulphate. Removal of the solvent under reduced pressure gave 4-(3,4-dichlorophenyl)-5-(methanesulphonyloxy)pent-1-ene (5.4g) as an oil.

15

$^1\text{H-NMR}$ (CDCl_3): δ = 7.42 (1H, d), 7.30 (1H, s), 7.04 (1H, dd), 5.56-5.72 (1H, m), 5.01-5.10 (2H, m), 4.22-4.39 (2H, m), 3.05-3.15 (1H, m), 2.88 (3H, s), 2.33-2.60 (2H, m) ppm.

20

Preparation 244-(3,4-Dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene (and hydrochloride salt)

-65-

4-(3,4-Dichlorophenyl)-5-(methanesulphonyloxy)pent-1-ene (5.4g) (see Preparation 23) and imidazole (3.6g) were dissolved in anhydrous acetonitrile (40ml) and the mixture heated under reflux for 100 hours. The solvent was removed under reduced pressure, the residue dissolved in dichloromethane and the solvent removed under reduced pressure. The residue was dissolved in dichloromethane (100ml) and washed with sufficient aqueous sodium carbonate solution to ensure that the aqueous phase reached pH14. The two phases were separated and the aqueous phase extracted with dichloromethane (100ml). The organic phases were combined and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 100:0 changing to 98:2, by volume, dichloromethane : methanol, to give 4-(3,4-dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene (3.45g) as an orange oil.

¹H-NMR (CDCl₃): δ = 7.35 (1H, d), 7.15-7.30 (2H, m), 6.99 (1H, s), 6.84 (1H, d), 6.69 (1H, s), 5.55-5.70 (1H, m), 5.07 (2H, d), 3.95-4.22 (2H, m), 2.95-3.10 (1H, m), 2.32-2.45 (2H, m) ppm.

The hydrochloride salt was prepared by dissolving the free base in dichloromethane, treating the solution with hydrogen chloride gas and removal of the solvent under reduced pressure to give 4-(3,4-dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene hydrochloride as a foam.

-66-

Preparation 25(alternative method to Preparation 18)3-(3,4-Dichlorophenyl)-4-(imidazol-1-yl)butan-1-al

5



4-(3,4-Dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene hydrochloride (3.6g) (see Preparation 24) was dissolved in a mixture of acetonitrile (50ml) and water (20ml) and osmium tetroxide (4ml of a 0.05M solution in toluene) added. The mixture was stirred for 30 minutes, sodium periodate (5.3g) added and stirring continued for 2 hours. A further portion of acetonitrile (20ml) was added and stirring continued for 16 hours before removal of the organic solvent under reduced pressure to give an aqueous suspension. This was basified to pH>7 by the addition of solid sodium carbonate. The mixture was then extracted with dichloromethane, the organic phase dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give a dark oil which was chromatographed on silica gel eluting with a solvent gradient of 98:2 changing to 92.5:7.5, by volume, dichloromethane : methanol to give 3-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-al (2.28g) as a colourless, viscous oil.

20

$^1\text{H-NMR}$ (CDCl_3): δ = 9.70 (1H, s), 7.38 (1H, d), 7.20-7.30 (2H, m), 7.01 (1H, s), 6.89 (1H, dd), 6.71 (1H, s), 4.00-4.22 (2H, m), 3.60 (1H, m), 2.72-2.92 (2H, m) ppm.

-67-


Preparation 262-(3,4-Dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butanenitrile

To a mixture of sodium hydride (60% w/w dispersion in mineral oil) (19.24g) and anhydrous tetrahydrofuran (450ml) at 0°C under a nitrogen atmosphere was added a solution of 3,4-dichlorophenylacetonitrile (89.5g) in anhydrous tetrahydrofuran (450ml), dropwise over 40 minutes. After a further 30 minutes, a solution of 2-(2-bromoethoxy)tetrahydropyran (100g) in anhydrous tetrahydrofuran (100ml) was added. The mixture was allowed to warm to room temperature and then stirred for 14 hours. 30% w/w Aqueous ammonium chloride solution (500ml) was added and the mixture extracted with diethyl ether (2 x 400ml). The organic extracts were combined, washed with water (2 x 400ml), dried over anhydrous magnesium sulphate and the solvent removed under reduced pressure. The residue was chromatographed on silica gel eluting with a solvent gradient of 1:9 changing to 1:1, by volume, diethyl ether : hexane to give 2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)-butanenitrile (51g) as an oil and as a mixture of diastereomers.

20

¹H-NMR (CDCl₃) (mixture of 2 diastereomers): δ = 7.25-7.50 (2H, m), 7.20-7.25 (1H, m), 4.50-4.60 (1H, m), 4.00-4.10 (1H, m), 2.80-2.95 (2H, m), 2.40-2.65 (2H, m), 2.05-2.30 (2H, m), 1.50-1.90 (6H, m) ppm.

-68-

Preparation 272-(3,4-Dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butan-1-al


2-(3,4-Dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butanenitrile (20.2g) (see Preparation 26) was dissolved in anhydrous toluene (300ml) and cooled to -78°C under a nitrogen atmosphere. Diisobutylaluminium hydride (85.6ml of a 1.0M solution in toluene) was then added dropwise. The mixture was stirred at -78°C for 1.5 hours and then allowed to warm slowly to -40°C . Water (100ml) and saturated aqueous ammonium chloride solution (50ml) were carefully added (exothermic reaction) and the mixture allowed to warm to 10°C before further addition of water (100ml) and saturated aqueous ammonium chloride solution (50ml). A 10% w/w aqueous solution of Rochelle's salt (potassium sodium tartrate tetrahydrate) (400ml) was added and the mixture extracted with diethyl ether. The organic phase was dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give a crude product. This was chromatographed on silica gel eluting with 98:2, by volume, dichloromethane : methanol to give 2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butan-1-al (17.02g) as a yellow oil and as a mixture of diastereomers.

$^1\text{H-NMR}$ (CDCl_3) (mixture of two diastereomers): δ = 9.70 (1H, s), 7.44 (1H, d), 7.32 (1H, m), 7.06 (1H, m), 4.55 (0.5H, t), 4.46 (0.5H, t), 3.20-3.90 (6H, m), 2.35-2.50 (1H, m), 1.45-2.00 (6H, m) ppm.

-69-

Preparation 282-(3,4-Dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butan-1-ol

5




Sodium borohydride (2.03g) was added portionwise to a solution of 2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butan-1-al (17.02g) (see Preparation 27) in 2-propanol (250ml). The mixture was stirred at room temperature overnight and glacial acetic acid (4ml) carefully added, followed by addition of water (2ml). The solvent was removed under reduced pressure to give a residue which was dissolved in dichloromethane and washed sequentially with water, dilute aqueous sodium hydrogen carbonate solution and brine. The solution was dried over anhydrous sodium sulphate and the solvent removed under reduced pressure. The crude product was chromatographed on silica gel eluting with 1:1, by volume, ethyl acetate : hexane to give 2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butan-1-ol (14.7g) as a colourless oil and as a mixture of diastereomers.

¹H-NMR (CDCl₃) (mixture of two diastereomers): δ = 7.30-7.40 (2H, m), 7.09 (1H, d), 4.55 (0.5H, t), 4.46 (0.5H, t), 3.24-3.82 (6H, m), 2.90-3.00 (1H, m), 1.45-2.10 (9H, m) ppm.

-70-

Preparation 291-Methanesulphonyloxy-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane

5




Methanesulphonyl chloride (2.4ml) was added dropwise to an ice-cooled solution of 2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butan-1-ol (8.13g) (see Preparation 28) and triethylamine (5.32ml) in dichloromethane (100ml) under a nitrogen atmosphere. The mixture was stirred for 10 minutes at 0°C and then 90 minutes at room temperature before being washed twice with water. The organic phase was separated, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give the crude product. This was chromatographed on silica gel eluting with 95:5, by volume, dichloromethane : diethyl ether to give 1-methanesulphonyloxy-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane (8.85g) as a yellow oil and as a mixture of diastereomers.

¹H-NMR (CDCl₃) (mixture of two diastereomers): δ = 7.42 (1H, d), 7.32 (1H, dd), 7.09 (1H, m), 4.30-4.50 (3H, m), 3.20-3.80 (5H, m), 2.89 (3H, s), 2.05-2.15 (1H, m), 1.50-1.95 (7H, m) ppm.

-71-

Preparation 301-(Imidazol-1-yl)-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane

5




1-Methanesulphonyloxy-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane (10g) (see Preparation 29), imidazole (2.07g) and potassium carbonate (7.65g) were dissolved in anhydrous acetonitrile (50ml) and heated under reflux under a nitrogen atmosphere for 3 days. The organic solvent was then removed under reduced pressure and the resulting aqueous suspension partitioned between dichloromethane and water. The organic phase was separated, washed sequentially with water and brine, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure. The crude product was chromatographed on silica gel eluting with 95:5, by volume, dichloromethane : methanol to give 1-(imidazol-1-yl)-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane (3g) as an oil and as a mixture of diastereomers.

¹H-NMR (CDCl₃) (mixture of two diastereomers): δ = 7.35 (1H, d), 7.18-7.28 (2H, m), 6.99 (1H, s), 6.88 (1H, m), 6.69 (1H, s), 4.49 (0.5H, t), 4.40 (0.5H, t), 4.00-4.25 (2H, m), 3.10-3.85 (5H, m), 1.45-2.10 (8H, m) ppm.

-72-

Preparation 313-(3,4-Dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butyl 3,5-dimethylbenzoate


5



3,5-Dimethylbenzoyl chloride (1.33g) was added dropwise to a suspension of 3-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol (0.75g) (see Preparation 19) and triethylamine (1.1g) in anhydrous acetonitrile (15ml) and the solution stirred at room temperature for 120 hours. The mixture was then mixed with water (30ml) and dichloromethane (50ml), the organic phase separated and washed sequentially with water and brine. The solvent was removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 100:0 changing to 99 : 1, by volume, dichloromethane : methanol to give 3-(3,4-dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butyl 3,5-dimethylbenzoate (0.94 g) as a yellow foam.

¹H-NMR (CDCl₃): δ = 7.69 (2H, s), 7.49 (2H, s), 7.10-7.27 (5H, m), 6.92 (1H, dd), 6.81 (1H, s), 4.76 (1H, dd), 4.51 (1H, dd), 4.10-4.30 (2H, m), 3.32 (1H, m), 2.35 (6H, s), 2.29 (6H, s), 2.19 (2H, m) ppm.

-73-

Preparation 322,3-Dihydrobenzo[b]furan-7-oic acid

5

N,N,N',N'-Tetramethylethylenediamine (38ml) was dissolved in hexane (300ml), cooled in an ice-bath and n-butyllithium (100ml of a 2.5M solution in hexane) added. The mixture was stirred at 0°C for 15 minutes before adding 2,3-dihydrobenzo[b]furan (30g) dropwise over 30 minutes. The mixture was allowed to warm to room temperature over 30 minutes, stirred at room temperature for a further 4 hours, poured onto an excess of solid carbon dioxide and left to stand for 3 days by which time the solvent had evaporated off. The residue was partitioned between ethyl acetate (1L) and 4N aqueous hydrochloric acid solution (240ml), the layers separated and the aqueous layer extracted with ethyl acetate (500ml). The organic extracts were combined, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure. The residue was then triturated with diethyl ether to give 2,3-dihydrobenzo[b]furan-7-oic acid as a white solid (21g)

¹H-NMR (CDCl₃): δ = 7.75 (1H, d), 7.31 (1H, d), 6.88 (1H, t), 4.69 (2H, t), 3.20 (2H, t) ppm.

20

-74-

Preparation 331,2,3,4-Tetrahydro-5-naphthoic acid

5




10% w/w Palladium-on-carbon (10g) was added to a solution of 1-naphthoic acid (33.4g) in glacial acetic acid (150ml) and the mixture hydrogenated at 345kPa (50psi) and 85°C for 4 days. The warm mixture was filtered through a short column of Arbocel (trade mark) filter aid and the pad washed with glacial acetic acid (150ml). Water (1.5L) was added to the filtrate and the resulting precipitate filtered off and washed with water. The precipitate was dissolved in dichloromethane, the solution dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give an oil which was crystallised from ethyl acetate to give 1,2,3,4-tetrahydro-5-naphthoic acid (2.94g) as a white solid (m.p. 148-150°C).

¹H-NMR (CDCl₃): δ = 7.85 (1H, d), 7.28 (1H, d), 7.16 (1H, m), 3.15 (2H, br. s), 2.84 (2H, br. s), 1.72-1.90 (4H, m) ppm.

-75-

Preparation 341,2,3,4-Tetrahydro-6-naphthoic acid

5



1,2,3,4-Tetrahydro-6-naphthaldehyde (2.0g) was suspended in 0.5M aqueous sodium hydroxide solution (125ml), stirred vigorously and a 70% w/w solution of tertiary butyl hydroperoxide in water (10.3ml) added. The mixture was heated at about 70°C for 4 hours and then left to stand at room temperature for 3 days. A further quantity of a 70% w/w solution of tertiary butyl hydroperoxide in water (10ml) was added and the mixture heated at about 70°C for a further 24 hours. The mixture was cooled, diethyl ether (100ml) added and the phases separated. The aqueous phase was acidified to pH 1 with 2N aqueous hydrochloric acid solution and extracted with diethyl ether (2x 100ml). The organic phases were combined, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 1:1 changing to 1:0, by volume, diethyl ether : pentane to give 1,2,3,4-tetrahydro-6-naphthoic acid (0.62 g) as a white solid.


20

¹H-NMR (CDCl₃): δ = 7.78-7.86 (2H, m), 7.14 (1H, d), 2.78-2.87 (4H, br. s), 1.79-1.88 (4H, br. s) ppm.

-76-


Preparation 352,3-Dihydrobenzo[b]furan-7-oyl chloride

5




2,3-Dihydrobenzo[b]furan-7-oic acid (3g) (see Preparation 32) was suspended in anhydrous dichloromethane (30ml) and oxalyl chloride (3.5g) added, followed by addition of dimethylformamide (3 drops). The mixture was stirred at room temperature for 2.5 hours, the solvent then removed under reduced pressure, the resulting residue dissolved in dichloromethane and the solvent removed under reduced pressure. The residue was again dissolved in dichloromethane and the solvent removed under reduced pressure to give 2,3-dihydrobenzo[b]furan-7-oyl chloride as a pink solid (3.3g).

15

Preparation 361,2,3,4-Tetrahydro-5-naphthoyl chloride

This compound was prepared by an analogous method to that used in Preparation 35 using 1,2,3,4-tetrahydro-5-naphthoic acid (see Preparation 33) as the starting material.

-77-

Preparation 371,2,3,4-Tetrahydro-6-naphthoyl chloride


5

This compound was prepared by an analogous method to that used in Preparation 35 using 1,2,3,4-tetrahydro-6-naphthoic acid (see Preparation 34) as the starting material.

10

Preparations 38 - 46






The compounds of the following tabulated preparations (Table 2), of the general formula:







15

were prepared by a similar method to that of Preparation 31 using 3-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol (see Preparation 19) and the appropriate acid chloride starting material.

TABLE 2


Prep. no.	Acid Chloride starting material	R-W-	Analytical Data
38	2,3-dimethylbenzoyl chloride		¹ H-NMR (CDCl ₃): δ = 7.47 (1H, d), 7.33 (1H, d), 7.22-7.29 (3H, m), 6.96-7.15 (5H, m), 6.82 (1H, s), 4.83 (1H, dd), 4.64 (1H, dd), 4.27-4.36 (1H, m), 4.09-4.20 (1H, m), 3.32-3.44 (1H, m), 2.41 (3H, s), 2.31 (6H, s), 2.15-2.25 (2H, m), 2.15 (3H, s) ppm.
39	2-trifluoromethoxybenzoyl chloride		¹ H-NMR (CDCl ₃): δ = 7.82 (1H, d), 7.20-7.65 (9H, m), 7.10 (1H, s), 6.93 (1H, dd), 6.79 (1H, s), 4.86 (1H, dd), 4.53 (1H, dd), 4.30-4.42 (1H, m), 4.12-4.28 (1H, m), 3.28-3.42 (1H, m), 2.10-2.30 (2H, m) ppm.
40	2-methoxy-3-methylbenzoyl chloride		¹ H-NMR (CDCl ₃): δ = 7.48 (1H, d), 7.24-7.36 (4H, m), 7.18 (1H, d), 6.95-7.09 (4H, m), 6.80 (1H, s), 4.86 (1H, dd), 4.56 (1H, dd), 4.27-4.36 (1H, m), 4.10-4.22 (1H, m), 3.76 (3H, s), 3.66 (3H, s), 3.31-3.43 (1H, m), 2.30 (3H, s), 2.28 (3H, s), 2.08-2.18 (2H, m) ppm.
41	See Prep. no. 35		¹ H-NMR (CDCl ₃): δ = 7.58 (1H, d), 7.50 (1H, d), 7.20-7.36 (4H, m), 7.08 (1H, s), 6.98 (1H, d), 6.72-6.90 (3H, m), 4.50-4.81 (6H, m), 4.27-4.38 (1H, m), 4.03-4.15 (1H, m), 3.35-3.49 (1H, m), 3.05-3.28 (4H, m), 2.04-2.23 (2H, m) ppm.
42 (footnote (a))	2-naphthoyl chloride		¹ H-NMR (CDCl ₃): δ = 8.82 (1H, s), 8.46 (1H, s), 8.15 (1H, d), 7.74-7.99 (8H, m), 7.43-7.64 (4H, m), 7.26 (1H, s), 7.17 (1H, s), 6.98 (1H, d), 6.86 (1H, s), 4.89 (1H, dd), 4.58 (1H, dd), 4.39-4.48 (1H, m), 4.25-4.30 (1H, m), 3.42-3.52 (1H, m), 2.20-2.38 (2H, m) ppm.

43 (footnote (b))	3- trifluoro- methyl-4- fluoro- benzoyl chloride		¹ H-NMR (CDCl ₃): δ = 8.49-8.62 (2H, m), 8.20 (1H, d), 8.04-8.09 (1H, m), 7.12-7.32 (5H, m), 6.90 (1H, d), 6.82 (1H, s), 4.81 (1H, dd), 4.26-4.51 (3H, m), 3.29-3.40 (1H, m), 2.20-2.29 (2H, m) ppm.
44 (footnote (a))	See Prep. No. 36		¹ H-NMR (CDCl ₃): δ = 7.50 (1H, d), 6.95-7.38 (10H, m), 4.83 (1H, dd), 4.63 (1H, m), 4.30-4.35 (1H, m), 4.08-4.16 (1H, m), 3.30-3.40 (1H, m), 3.00 (2H, br. s), 2.63-2.84 (6H, m), 2.10-2.25 (2H, m), 1.67-1.83 (8H, m) ppm.
45 (footnote (c))	See Prep. No. 37		¹ H-NMR (CDCl ₃): δ = 7.79-7.88 (2H, m), 7.52-7.62 (2H, m), 7.01-7.34 (5H, m), 6.90 (1H, dd), 6.79 (1H, s), 4.76 (1H, dd), 4.51 (1H, dd), 4.10-4.32 (2H, m), 3.27-3.40 (1H, m), 2.68-2.91 (8H, m), 2.11-2.26 (2H, m), 1.72-1.91 (8H, m) ppm.
46	2- methoxy- 5-chloro- benzoyl chloride		¹ H-NMR (CDCl ₃): δ = 7.62 (1H, s), 7.32-7.43 (4H, m), 7.24 (1H, s), 7.07 (1H, s), 7.02 (1H, dd), 6.90 (2H, t), 6.77 (1H, s), 4.87 (1H, dd), 4.49 (1H, dd), 4.30-4.35 (1H, m), 4.11-4.20 (1H, m), 3.86 (3H, s), 3.72 (3H, s), 3.32-3.41 (1H, m), 2.10-2.25 (2H, m) ppm.

Footnotes

- 5 (a) chromatographed on silica gel eluting with a solvent gradient of 1:2 changing to 1:0, by volume, ethyl acetate : pentane.
- (b) chromatographed on silica gel eluting with diethyl ether.
- (c) chromatographed on silica gel eluting with a solvent gradient of 1:2 changing to 2:1, by volume, ethyl acetate : pentane.

-80-

Preparation 473-(3,4-Dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butan-1-ol


3-(3,4-Dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butyl 3,5-dimethylbenzoate (0.93g) (see Preparation 31) was dissolved in methanol (10ml) and 2N aqueous sodium hydroxide solution (2ml) added. A thick gum formed, additional methanol was added (50ml) and the resulting suspension stirred at room temperature overnight. The methanol was then removed under reduced pressure and the residue partitioned between dichloromethane (50ml) and water (10ml). The organic phase was separated, washed with brine and the solvent removed under reduced pressure to give the crude product which was chromatographed on silica gel eluting with a solvent gradient of 99:1 changing to 95 : 5, by volume, dichloromethane : methanol, to give 3-(3,4-dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butan-1-ol (0.67 g) as a white foam.

¹H-NMR (CDCl₃): δ = 7.70 (2H, s), 7.10-7.32 (4H, m), 6.89 (1H, d), 6.72 (1H, s), 4.83 (1H, dd), 4.40 (1H, dd), 3.70-3.80 (1H, m), 3.50-3.62 (1H, m), 3.28-3.40 (1H, m), 2.40 (6H, s), 1.85-2.07 (3H, m), ppm.

-81-

Preparations 48 - 53





The compounds of the following tabulated preparations (Table 3) of the general
 5 formula :




were prepared by a similar method to that of Preparation 47 using the appropriate
 10 starting materials (see Preparations 38-41, 44 and 46).

TABLE 3

Prep. no.	Starting material Prep. No.	R-W-	Analytical Data
48	38		¹ H-NMR (CDCl ₃): δ = 7.31 (1H, d), 7.10-7.28 (4H, m), 7.06 (1H, s), 6.92 (1H, dd), 6.75 (1H, s), 4.90 (1H, dd), 4.50 (1H, dd), 4.70-4.81 (1H, m), 3.50-3.64 (1H, m), 3.30-3.43 (1H, m), 2.29 (3H, s), 2.18 (3H, s), 1.90-2.05 (2H, m), 1.80 (1H, t) ppm.
49	39		¹ H-NMR (CDCl ₃): δ = 7.20-7.64 (6H, m), 7.09 (1H, s), 6.91 (1H, d), 6.72 (1H, s), 4.96 (1H, dd), 4.39 (1H, dd), 3.69-3.83 (1H, m), 3.51-3.67 (1H, m), 3.28-3.41 (1H, m), 1.87-2.08 (2H, m), 1.74 (1H, t) ppm.

50	40		¹ H-NMR (CDCl ₃): δ = 7.18-7.37 (4H, m), 7.05-7.10 (2H, m), 6.93 (1H, dd), 6.72 (1H, s), 4.92 (1H, dd), 4.42 (1H, dd), 3.75 (1H, m), 3.70 (3H, s), 3.50-3.62 (1H, m), 3.31-3.45 (1H, m), 2.32 (3H, s), 1.86-2.02 (3H, m) ppm.
51	41		¹ H-NMR (CDCl ₃): δ = 7.64 (1H, d), 7.11-7.38 (3H, m), 7.06 (1H, s), 6.85-6.96 (2H, m), 6.67 (1H, s), 4.89 (1H, dd), 4.60-4.75 (2H, m), 4.35 (1H, dd), 3.63-3.75 (1H, m), 3.49-3.61 (1H, m), 3.18-3.41 (3H, m), 1.84-2.06 (2H, m), 1.72 (1H, t) ppm.
52	44		¹ H-NMR (CDCl ₃): δ = 7.10-7.35 (5H, m), 7.06 (1H, s), 6.92 (1H, dd), 6.74 (1H, s), 4.90 (1H, dd), 4.50 (1H, dd), 3.70-3.82 (1H, m), 3.53-3.64 (1H, m), 3.30-3.42 (1H, m), 2.70-2.86 (4H, m), 1.70-2.04 (6H, m) ppm.
53	46		¹ H-NMR (CDCl ₃): δ = 7.30-7.42 (3H, m), 7.19 (1H, s), 7.03 (1H, s), 6.92 (2H, t), 6.69 (1H, s), 4.91 (1H, dd), 4.35 (1H, dd), 3.50-3.80 (4H, m), 3.50-3.63 (1H, m), 3.29-3.40 (1H, m), 1.89-2.01 (2H, m), 1.72 (1H, t) ppm.

-83-

Preparation 543-(3,4-Dichlorophenyl)-4-[2-(2-naphthoyl)imidazol-1-yl]butan-1-ol

5

3-(3,4-Dichlorophenyl)-4-[2-(2-naphthoyl)imidazol-1-yl]butyl 2-naphthoate (1.25g)
(see Preparation 42) was dissolved in 1,4-dioxane (15ml), 1N aqueous sodium
hydroxide solution (4ml) added and the mixture stirred at room temperature overnight.


- 10 The dioxane was then removed under reduced pressure and the residue partitioned
between ethyl acetate and water. The organic phase was separated, washed sequentially
with 1N aqueous sodium hydroxide solution, water and brine and the solvent removed
under reduced pressure to give the crude product which was chromatographed on silica
gel eluting with a solvent gradient of 99:1 changing to 95 : 5, by volume,
15 dichloromethane : methanol, to give 3-(3,4-dichlorophenyl)-4-[2-(2-
naphthoyl)imidazol-1-yl]butan-1-ol (0.8 g) as a gum.

¹H-NMR (CDCl₃): δ = 8.82 (1H, s), 8.16 (1H, d), 7.99 (1H, d), 7.89 (2H, t), 7.50-7.63
(2H, m), 7.26 (1H, d), 7.16 (1H, d), 6.90 (1H, dd), 6.76 (1H, s), 4.90 (1H, dd), 4.43 (1H,
20 dd), 3.69-3.83 (1H, m), 3.51-3.68 (1H, m), 3.32-3.45 (1H, m), 1.86-2.08 (3H, m) ppm.

-84-

Preparations 55 - 56

The compounds of the following tabulated preparations (Table 4) of the general
 5 formula:



were prepared by a similar method to that of Preparation 54 using the appropriate
 10 starting material compounds (see Preparations 43 and 45).

TABLE 4

Prep. no.	Starting material Prep. No.	R	Analytical Data
55 (footnote (a))	43		¹ H-NMR (CDCl ₃): δ = 8.49-8.62 (2H, m), 7.23-7.32 (2H, m), 7.11 (2H, s), 6.86 (1H, d), 6.80 (1H, s), 4.85 (1H, dd), 4.47 (1H, dd), 3.70-3.81 (1H, m), 3.52-3.63 (1H, m), 3.29-3.39 (1H, m), 1.89-2.09 (2H, m), 1.71 (1H, br. s) ppm.
56	45		¹ H-NMR (CDCl ₃): δ = 7.81-7.91 (2H, m), 7.05-7.32 (4H, m), 6.88 (1H, dd), 6.71 (1H, s), 4.85 (1H, dd), 4.37 (1H, dd), 3.69-3.80 (1H, m), 3.51-3.66 (1H, m), 3.07-3.42 (1H, m), 2.70-2.92 (4H, m), 1.76-2.05 (6H, m) ppm.

-85-


Footnote

(a) chromatographed on silica gel eluting with a solvent gradient of 1:0 changing to 0:1, by volume, diethyl ether : ethyl acetate.

5

Preparation 57

1-[2-(2-Methoxybenzoyl)imidazol-1-yl]-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane



1-(Imidazol-1-yl)-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane (0.86g) (see Preparation 30) and triethylamine (1.3ml) were dissolved in acetonitrile (15ml), under a nitrogen atmosphere, the solution cooled in an ice-bath and 2-methoxybenzoyl chloride (1.4ml) added, dropwise. The mixture was allowed to warm to room temperature slowly and then stirred for 2 days. The acetonitrile was removed under reduced pressure and the residue dissolved in dichloromethane (20ml) and washed sequentially with water and brine. The organic phase was then dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with 97:3, by volume, dichloromethane : methanol to give 1-[2-(2-methoxybenzoyl)imidazol-1-yl]-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane (1.1g) as an oil and as a mixture of diastereomers.

15

20


-86-

$^1\text{H-NMR}$ (CDCl_3) (mixture of two diastereomers): $\delta = 6.76\text{--}7.52$ (9H, m), $4.10\text{--}4.90$ (3H, m), $3.11\text{--}3.90$ (8H, m), $1.45\text{--}2.25$ (6H, m), 1.26 (1H, t), 1.06 (1H, t)ppm.

5

Preparations 58 - 62






The compounds of the following tabulated preparations (Table 5) of the general formula:



10 were prepared by a similar method to that of Preparation 57 using the appropriate starting materials.

-87-


TABLE 5

Prep. no.	Acid Chloride Starting material	R-W-	Analytical Data
58	3,5-bis(trifluoromethyl)-benzoyl chloride		¹ H-NMR (CDCl ₃) (mixture of two diastereomers): δ = 8.71 (2H, d), 8.06 (1H, s), 7.13-7.32 (3H, m), 6.98 (1H, d), 6.85-6.95 (1H, m), 4.40-4.80 (3H, m), 3.60-3.85 (2H, m), 3.20-3.50 (3H, m), 1.43-2.18 (8H, m) ppm.
59	2-trifluoromethyl-benzoyl chloride		¹ H-NMR (CDCl ₃) (mixture of two diastereomers): δ = 7.75 (1H, dd), 7.60 (2H, m), 7.43 (2H, m), 7.24 (1H, t), 7.09 (1H, s), 6.97 (1H, m), 6.86 (1H, d), 4.80-4.90 (1H, m), 4.40-4.64 (2H, m), 3.55-3.83 (2H, m), 3.20-3.50 (3H, m), 1.90-2.18 (2H, m), 1.43-1.86 (6H, m) ppm.
60	2-isopropoxy-benzoyl chloride		¹ H-NMR (CDCl ₃) (mixture of two diastereomers): δ = 6.75-7.58 (9H, m), 4.38-4.91 (4H, m), 3.20-3.80 (5H, m), 1.15-2.13 (14H, m) ppm.
61	2-ethylbenzoyl chloride		¹ H-NMR (CDCl ₃) (mixture of two diastereomers): δ = 6.87-7.43 (9H, m), 4.40-4.83 (3H, m), 3.20-3.82 (5H, m), 2.62-2.75 (2H, m), 1.10-2.10 (11H, m) ppm.
62	2-phenoxybenzoyl chloride		¹ H-NMR (CDCl ₃) (mixture of two diastereomers): δ = 6.90-7.52 (12H, m), 6.84 (1H, d), 6.67 (1H, d), 4.69-4.80 (1H, m), 4.29-4.48 (2H, m), 3.12-3.79 (5H, m), 1.41-2.08 (8H, m) ppm.

-88-

Preparation 631-Ethoxymethylimidazole


5



Imidazole (20g) was dissolved in tetrahydrofuran (700ml) under a nitrogen atmosphere, cooled to -70°C and n-butyllithium (117.5ml of a 2.5M solution in hexane) added dropwise, over 15 minutes. The mixture was allowed to warm to -20°C and stirred at -20°C for 30 minutes before the dropwise addition of chloromethyl ethyl ether (30.5g). The mixture was allowed to warm to room temperature and stirred for a further hour. The solvent was removed under reduced pressure to give a residue which was triturated with dichloromethane and filtered through a short pad of Arbocel (trade mark) filter aid. The solvent was removed from the filtrate under reduced pressure to give a mobile oil which was distilled under reduced pressure (0.7millibars, 0.53mmHg) to give 1-ethoxymethylimidazole (20.8g) as an oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.61 (1H, s), 7.09 (2H, d), 5.29 (2H, s), 3.49 (2H, q), 1.21 (3H, t) ppm.

-89-

Preparation 641-Ethoxymethyl-2-phenacetylimidazole


5

1-Ethoxymethylimidazole (11.98g) (see Preparation 63) was dissolved in tetrahydrofuran (400ml) under a nitrogen atmosphere, cooled to -70°C and n-butyl lithium (40ml of a 2.5M solution in hexane) added, dropwise, over 5 minutes. The mixture was stirred at -70°C for 1 hour and chlorotrimethylsilane (10.83g) added, dropwise. The mixture was stirred at -70°C for 1 hour before being allowed to warm to room temperature and stirred for a further 3 hours. Phenacetyl chloride (14.68g) was added and stirring continued for 18 hours before removing the solvent under reduced pressure. The resulting residue was dissolved in dichloromethane and washed sequentially with water, saturated aqueous sodium hydrogen carbonate solution and brine before drying over anhydrous sodium sulphate. The solvent was removed from the organic phase under reduced pressure. The crude product was chromatographed on silica gel eluting with 95:5, by volume, dichloromethane : methanol to give 1-ethoxymethyl-2-phenacetylimidazole (8.23g) as an oil.

20

$^1\text{H-NMR}$ (CDCl_3): δ = 7.19-7.39 (7H, m), 5.72 (2H, s), 4.43 (2H, s), 3.48 (2H, q), 1.14 (3H, t) ppm.

-90-

Preparation 652-Phenacetylimidazole

5

- 1-Ethoxymethyl-2-phenylacetylimidazole (8.23g) (see Preparation 64) was dissolved in ethanol (200ml), 2N aqueous hydrochloric acid solution (200ml) added and the resulting suspension heated under reflux for 30 minutes. A further quantity of ethanol (50ml) was added to dissolve the remaining suspended material and the mixture heated under reflux for a further 6 hours before being left to stand at room temperature overnight. The organic solvent was removed under reduced pressure and the resulting aqueous suspension basified to pH 9 by addition of a saturated aqueous sodium hydrogen carbonate solution. The mixture was then extracted (three times) with dichloromethane.
- 15 The organic extracts were combined, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give 2-phenacetylimidazole (6.29g) as a yellow solid.


¹H-NMR (CDCl₃): δ = 7.19-7.39 (7H, m), 4.40 (2H, s) ppm.

-91-

Preparation 66

1-[2-Phenacetyl-1-imidazol-1-yl]-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane

5




- 1-Methanesulfonyloxy-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane
 10 (0.57g) (see Preparation 29) and 2-phenacetyl-1-imidazole (0.27g) (see Preparation 65)
 were dissolved in acetonitrile (20ml). Potassium carbonate (0.41g) was added and the
 mixture heated under reflux, under a nitrogen atmosphere, for 18 hours. The mixture
 was cooled, the solvent removed under reduced pressure and the residue partitioned
 between dichloromethane and water. The organic phase was separated, washed with
 15 brine, dried over anhydrous sodium sulphate and the solvent removed under reduced
 pressure. The resulting residue was then chromatographed on silica gel eluting with
 99:1, by volume, dichloromethane : methanol to give 1-[2-phenacetyl-1-imidazol-1-yl]-2-
 (3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane (0.23g) as an oil and as a
 mixture of diastereomers.

-92-

¹H-NMR (CDCl₃) (mixture of two diastereomers): δ = 7.22-7.38 (6H, m), 7.05-7.12 (2H, m), 6.71-6.84 (2H, m), 4.64-4.76 (1H, m), 4.28-4.50 (4H, m), 3.10-3.79 (5H, m), 1.42-2.05 (8H, m) ppm.

5

Preparation 673-(3,4-Dichlorophenyl)-4-[2-(2-methoxybenzoyl)imidazol-1-yl]butan-1-ol

10

15 1-[2-(2-Methoxybenzoyl)imidazol-1-yl]-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane (1.1 g) (see Preparation 57) was dissolved in methanol (20ml), Amberlyst 15 (trade mark) ion-exchange resin (0.11 g) added and the mixture stirred at room temperature for 5 days. The resin was removed by filtration through a short column of Arbocel (trade mark) filter aid and the solvent removed from the filtrate under reduced pressure to give a residue. This was chromatographed on silica gel eluting with 97:3, by volume, dichloromethane : methanol to give 3-(3,4-dichlorophenyl)-4-[2-(2-methoxybenzoyl)-imidazol-1-yl]butan-1-ol (0.31 g) as a white foam.


-93-

$^1\text{H-NMR}$ (CDCl_3): δ = 7.19-7.48 (4H, m), 6.91-7.06 (4H, m), 6.67 (1H, s), 4.93 (1H, dd), 4.36 (1H, dd), 3.52-3.80 (2H, m), 3.80 (3H, s), 3.30-3.41 (1H, m), 1.90-2.03 (2H, m), 1.72 (1H, br. s) ppm.

5

Preparation 68

3-(3,4-Dichlorophenyl)-4-[2-[3,5-bis(trifluoromethyl)benzoyl]imidazol-1-yl]-
butan-1-ol




10

3-(3,4-Dichlorophenyl)-4-[2-[3,5-bis(trifluoromethyl)benzoyl]imidazol-1-yl]butan-1-ol
was prepared in an analogous fashion to the compound of Preparation 67 using the
15 compound of Preparation 58 as the starting material.

$^1\text{H-NMR}$ (CDCl_3): δ = 8.76 (2H, s), 8.08 (1H, s), 7.30 (1H, d), 7.19 (2H, d), 6.85-6.92 (2H, m), 4.88 (1H, dd), 4.50 (1H, dd), 3.71-3.81 (1H, m), 3.51-3.63 (1H, m), 3.30-3.42 (1H, m), 1.90-2.10 (2H, m), 1.71 (1H, t) ppm.

-94-

Preparation 693-(3,4-Dichlorophenyl)-4-[2-(2-trifluoromethylbenzoyl)imidazol-1-yl]butan-1-ol

5

1-[2-(2-Trifluoromethylbenzoyl)imidazol-1-yl]-2-(3,4-dichlorophenyl)-4-(tetrahydropyran-2-yloxy)butane (0.68g) (see Preparation 59) was dissolved in methanol (15ml), which had been previously saturated with hydrogen chloride gas, and left to stand for 4 hours. The solvent was then removed under reduced pressure to give a residue which was partitioned between dichloromethane and saturated aqueous sodium hydrogen carbonate solution (the pH of the aqueous layer was kept at 8-9). The organic phase was separated and the aqueous phase extracted (twice) with dichloromethane. The organic phases were then combined, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give 3-(3,4-dichlorophenyl)-4-[2-(2-trifluoromethylbenzoyl)imidazol-1-yl]butan-1-ol (0.41g) as a white foam.


$^1\text{H-NMR}$ (CDCl_3): δ = 7.77 (1H, d), 7.54-7.69 (2H, m), 7.47 (1H, m), 7.32 (1H, d), 7.22 (1H, d), 7.07 (1H, s), 6.91 (1H, dd), 6.74 (1H, s), 4.98 (1H, dd), 4.42 (1H, dd), 3.70-3.81 (1H, m), 3.52-3.67 (1H, m), 3.28-3.41 (1H, m), 1.90-2.09 (2H, m), 1.77 (1H, br. s) ppm.

20

-95-

Preparations 70 - 73

The compounds of the following tabulated preparations (Table 6) of the general
 5 formula:





were prepared by a similar method to that of Preparation 69 using the appropriate
 10 tetrahydropyranyl protected starting materials (see Preparations 60-62 and 66)

TABLE 6

Prep. no.	Starting material Prep. no.	R-W-	Analytical Data
70 (footnote (a))	60		¹ H-NMR (CDCl ₃): δ = 7.31-7.46 (3H, m), 7.24 (1H, d), 6.91-7.04 (4H, m), 6.85 (1H, s), 4.95 (1H, dd), 4.55 (1H, q), 4.30 (1H, dd), 3.71-3.83 (1H, m), 3.52-3.66 (1H, m), 3.31-3.44 (1H, m), 1.90-2.05 (2H, m), 1.22 (6H, d) ppm.
71 (footnote (a))	61		¹ H-NMR (CDCl ₃): δ = 7.19-7.47 (6H, m), 7.09 (1H, s), 6.91 (1H, dd), 6.76 (1H, s), 4.91 (1H, dd), 4.51 (1H, dd), 3.71-3.84 (1H, m), 3.52-3.66 (1H, m), 3.30-3.44 (1H, m), 2.70 (2H, q), 1.90-2.09 (2H, m), 1.86 (1H, br. s), 1.24 (3H, t) ppm.

-96-


72	62		¹ H-NMR (CDCl ₃): δ = 6.90-7.59 (12H, m), 6.79 (1H, dd), 6.56 (1H, s), 4.86 (1H, dd), 4.17 (1H, dd), 3.62-3.76 (1H, m), 3.48-3.59 (1H, m), 3.15-3.27 (1H, m), 1.82-1.95 (2H, m), 1.76 (1H, br. t) ppm.
73	66		¹ H-NMR (CDCl ₃): δ = 7.22-7.38 (6H, m), 7.04-7.10 (2H, m), 6.76 (1H, dd), 6.64 (1H, s), 4.81 (1H, dd), 4.40 (2H, dd), 4.21 (1H, dd), 3.60-3.70 (1H, m), 3.44-3.55 (1H, m), 3.10-3.21 (1H, m), 1.80-1.91 (2H, m), 1.67 (1H, t) ppm.

Footnote

- 5 (a) The product was purified by chromatography on silica gel eluting with 98:2, by volume, dichloromethane : methanol.

Preparation 74

1-Methanesulphonyloxy-3-(3,4-dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butane




-97-

3-(3,4-Dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]butan-1-ol (0.63g) (see Preparation 47) and triethylamine (0.20g) were dissolved in dichloromethane (12ml), cooled in an ice-bath and methanesulphonyl chloride (0.19g) added. The mixture was stirred for 30 minutes, dichloromethane (35ml) added and the solution washed sequentially with water (2x 30ml) and brine (30ml). The organic phase was dried over anhydrous sodium sulphate and the solvent removed under reduced pressure. The residue was dissolved in acetonitrile and the solvent removed under reduced pressure to give 1-methanesulphonyloxy-3-(3,4-dichlorophenyl)-4-[2-(3,5-dimethylbenzoyl)imidazol-1-yl]-butane (0.72g) as an oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.68 (2H, s), 7.10-7.32 (4H, m), 6.91 (1H, dd), 6.83 (1H, s), 4.73 (1H, dd), 4.53 (1H, dd), 4.15-4.24 (1H, m), 4.00-4.10 (1H, m), 3.26-3.36 (1H, m), 2.92 (3H, s), 2.40 (6H, s), 2.05-2.28 (2H, m) ppm.





Preparations 75 - 90







The compounds of the following tabulated preparations (Table 7) of the general formula:








were prepared by a similar method to that of Preparation 74 using the appropriate alcohol starting material (see Preparations 48-56 and 67-73).


TABLE 7

Prep. no.	Starting material Prep. No.	R	Analytical Data
75	48		¹ H-NMR (CDCl ₃): δ = 7.38 (1H, d), 7.21-7.29 (2H, m), 7.15 (2H, d), 7.09 (1H, s), 6.98 (1H, d), 6.84 (1H, s), 4.82 (1H, dd), 4.65 (1H, dd), 4.20-4.28 (1H, m), 4.02-4.11 (1H, m), 3.30-3.43 (1H, m), 2.96 (3H, s), 2.31 (3H, s), 2.08-2.28 (2H, m), 2.19 (3H, s) ppm.
76	49		¹ H-NMR (CDCl ₃): δ = 7.50-7.60 (2H, m), 7.31-7.44 (3H, m), 7.26 (1H, dd), 7.10 (1H, s), 6.95 (1H, dd), 6.79 (1H, s), 4.89 (1H, dd), 4.48 (1H, dd), 4.20-4.28 (1H, m), 4.03-4.11 (1H, m), 3.28-3.39 (1H, m), 2.05-2.30 (2H, m), 1.98 (3H, s) ppm.
77	50		¹ H-NMR (CDCl ₃): δ = 7.17-7.41 (4H, m), 6.96-7.11 (3H, m), 6.80 (1H, s), 4.86 (1H, dd), 4.52 (1H, dd), 4.20-4.29 (1H, m), 4.02-4.11 (1H, m), 3.70 (3H, s), 3.30-3.44 (1H, m), 2.95 (3H, s), 2.31 (3H, s), 2.05-2.29 (2H, m) ppm.
78	51		¹ H-NMR (CDCl ₃): δ = 7.64 (1H, d), 7.20-7.39 (3H, m), 7.09 (1H, s), 6.86-6.99 (2H, m), 6.72 (1H, s), 4.63-4.85 (3H, m), 4.45 (1H, dd), 4.12-4.23 (1H, m), 4.00-4.09 (1H, m), 3.20-3.40 (3H, m), 2.93 (3H, s), 2.02-2.18 (2H, m) ppm.
79	54		¹ H-NMR (CDCl ₃): δ = 8.81 (1H, s), 8.14 (1H, d), 8.00 (1H, d), 7.89 (2H, t), 7.50-7.64 (2H, m), 7.18-7.31 (3H, m), 6.95 (1H, d), 6.86 (1H, s), 4.82 (1H, dd), 4.56 (1H, dd), 4.02-4.29 (2H, m), 3.31-3.43 (1H, m), 2.94 (3H, s), 2.06-2.32 (2H, m) ppm.

80	55		¹ H-NMR (CDCl ₃): δ = 8.44-8.58 (2H, m), 7.25-7.35 (2H, m), 7.17 (2H, s), 6.90 (2H, m), 4.75 (1H, dd), 4.59 (1H, dd), 4.21-4.30 (1H, m), 4.03-4.11 (1H, m), 3.27-3.39 (1H, m), 2.98 (3H, s), 2.08-2.33 (2H, m) ppm.
81	52		¹ H-NMR (CDCl ₃): δ = 7.37 (1H, d), 7.05-7.31 (5H, m), 6.99 (1H, dd), 6.84 (1H, s), 4.80 (1H, dd), 4.65 (1H, dd), 4.05-4.30 (2H, m), 3.30-3.40 (1H, m), 2.97 (3H, s), 2.65-2.85 (4H, m), 2.10-2.30 (2H, m), 1.70-1.85 (4H, m) ppm.
82	56		¹ H-NMR (CDCl ₃): δ = 7.79-7.89 (2H, m), 7.10-7.35 (4H, m), 6.91 (1H, dd), 6.81 (1H, s), 4.72 (1H, dd), 4.51 (1H, dd), 4.17-4.27 (1H, m), 4.00-4.10 (1H, m), 3.06-3.39 (1H, m), 2.79-2.99 (7H, m), 2.03-2.29 (2H, m), 1.75-1.87 (4H, m) ppm.
83	53		¹ H-NMR (CDCl ₃): δ = 7.31-7.42 (3H, m), 7.21 (1H, d), 7.07 (1H, s), 6.99 (1H, dd), 6.91 (1H, d), 6.75 (1H, s), 4.83 (1H, dd), 4.48 (1H, dd), 4.16-4.28 (1H, m), 4.02-4.11 (1H, m), 3.75 (3H, s), 3.28-3.39 (1H, m), 2.92 (3H, s), 2.02-2.28 (2H, m) ppm.
84	67		¹ H-NMR (CDCl ₃): δ = 7.35-7.50 (3H, m), 7.24 (1H, d), 6.95-7.07 (4H, m), 6.72 (1H, s), 4.89 (1H, dd), 4.49 (1H, dd), 4.19-4.28 (1H, m), 4.05-4.14 (1H, m), 3.79 (3H, s), 3.30-3.41 (1H, m), 2.94 (3H, s), 2.06-2.29 (2H, m), ppm.
85	68		¹ H-NMR (CDCl ₃): δ = 8.79 (2H, s), 8.07 (1H, s), 7.32 (1H, d), 7.20 (2H, d), 6.91 (2H, d), 4.80 (1H, dd), 4.61 (1H, dd), 4.20-4.29 (1H, m), 4.04-4.12 (1H, m), 3.27-3.40 (1H, m), 2.93 (3H, s), 2.08-2.30 (2H, m) ppm.

86	69		¹ H-NMR (CDCl ₃): δ = 7.76 (1H, d), 7.58-7.68 (2H, m), 7.49 (1H, d), 7.39 (1H, d), 7.26 (1H, d), 7.10 (1H, s), 6.95 (1H, dd), 6.80 (1H, s), 4.91 (1H, dd), 4.50 (1H, dd), 4.22-4.30 (1H, m), 4.05-4.15 (1H, m), 3.30-3.41 (1H, m), 2.98 (3H, s), 2.07-2.30 (2H, m) ppm.
87	70		¹ H-NMR (CDCl ₃): δ = 7.32-7.46 (3H, m), 7.24 (1H, s), 6.92-7.07 (4H, m), 6.71 (1H, s), 4.87 (1H, dd), 4.40-4.56 (2H, m), 4.18-4.27 (1H, m), 4.04-4.11 (1H, m), 3.30-3.42 (1H, m), 2.94 (3H, s), 2.07-2.27 (2H, m), 1.21 (6H, d) ppm.
88	71		¹ H-NMR (CDCl ₃): δ = 7.21-7.45 (6H, m), 7.08 (1H, s), 7.96 (1H, dd), 6.85 (1H, s), 4.82 (1H, dd), 4.63 (1H, dd), 4.20-4.30 (1H, m), 4.03-4.11 (1H, m), 3.30-3.42 (1H, m), 2.94 (3H, s), 2.68 (2H, q), 2.06-2.31 (2H, m), 1.21 (3H, t) ppm.
89	72		¹ H-NMR (CDCl ₃): δ = 6.90-7.59 (12H, m), 6.83 (1H, dd), 6.63 (1H, s), 4.80 (1H, dd), 3.96-4.32 (3H, m), 3.15-3.25 (1H, m), 2.90 (3H, s), 2.00-2.21 (2H, m) ppm.
90	73		¹ H-NMR (CDCl ₃): δ = 7.22-7.37 (6H, m), 7.10 (2H, dd), 6.81 (1H, dd), 6.70 (1H, s), 4.76 (1H, dd), 4.25-4.50 (3H, m), 4.10-4.18 (1H, m), 3.91-4.03 (1H, m), 3.10-3.21 (1H, m), 2.89 (3H, s), 1.94-2.15 (2H, m) ppm.

-101-

Preparation 914(S)-(3,4-Dichlorophenyl)-5-hydroxypent-1-ene


5

4(S)-Benzyl-3-(2(S)-(3,4-dichlorophenyl)pent-4-en-1-oyl)oxazolidin-2-one (30.32g) (see Bioorganic and Medicinal Chemistry Letters, 3, 319, (1993)) was dissolved in anhydrous tetrahydrofuran (400ml), cooled in an ice-bath and lithium aluminium hydride (5.7g) carefully added in two portions (exothermic reaction). The mixture was stirred at 0°C for 20 minutes and then stirred at room temperature for 1 hour before being cooled to 0°C. Water (6ml) was carefully added followed by 2N aqueous sodium hydroxide solution (6ml) and further water (12ml). The mixture was stirred vigorously for 40 minutes and the resulting precipitate removed by filtration and washed with diethyl ether. The filtrate was collected and the organic and aqueous phases separated.

15 The solvent was then removed from the organic phase under reduced pressure to give a yellow oil which was chromatographed on silica gel eluting with a solvent gradient of 4:1 changing to 4:0, by volume, dichloromethane : hexane to give 4(S)-(3,4-dichlorophenyl)-5-hydroxypent-1-ene (11.74g) as an oil.

20 ¹H-NMR (CDCl₃): δ = 7.24-7.45 (2H, m), 7.06 (1H, dd), 5.61-5.75 (1H, m), 4.96-5.10 (2H, m), 3.70-3.90 (2H, m), 2.80-2.91 (1H, m), 2.30-2.55 (2H, m), 1.32 (1H, t) ppm.

-102-


Preparation 924(R)-(3,4-Dichlorophenyl)-5-hydropent-1-ene

This compound was prepared by an analogous method to that used to prepare the compound of Preparation 91 using the appropriate starting material (see Bioorganic and Medicinal Chemistry Letters, 3, 319(1993)).

10 $^1\text{H-NMR}$ (CDCl_3): δ = 7.24-7.45 (2H, m), 7.06 (1H, dd), 5.61-5.75 (1H, m), 4.96-5.10 (2H, m), 3.70-3.90 (2H, m), 2.80-2.91 (1H, m), 2.30-2.55 (2H, m), 1.32 (1H, t) ppm.

Preparation 934(S)-(3,4-Dichlorophenyl)-5-(methanesulphonyloxy)pent-1-ene

15




-103-

Methanesulphonyl chloride (3.55g) was added dropwise to an ice-cooled solution of 4(S)-(3,4-dichlorophenyl)-5-hydroxypent-1-ene (6.0g) (see Preparation 91) and triethylamine (3.48g) in dichloromethane (100ml) and the mixture stirred at room temperature for 1 hour. The solvent was then removed under reduced pressure and the residue partitioned between diethyl ether and water. The two phases were separated and the aqueous layer further extracted with diethyl ether. The organic phases were then combined and washed sequentially with water, dilute aqueous citric acid solution, brine, saturated aqueous sodium hydrogen carbonate solution and brine before drying over anhydrous sodium sulphate. Removal of the solvent under reduced pressure gave 4(S)-(3,4-dichlorophenyl)-5-(methanesulphonyloxy)pent-1-ene (8.25g) as an oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.42 (1H, d), 7.30 (1H, s), 7.04 (1H, dd), 5.56-5.72 (1H, m), 5.01-5.10 (2H, m), 4.22-4.39 (2H, m), 3.05-3.15 (1H, m), 2.88 (3H, s), 2.33-2.60 (2H, m) ppm.

Preparation 94

4(R)-(3,4-Dichlorophenyl)-5-(methanesulphonyloxy)pent-1-ene




This compound was prepared by an analogous method to that used to prepare the compound of Preparation 93 using 4(R)-(3,4-dichlorophenyl)-5-hydroxypent-1-ene (see Preparation 92) as the starting material.

-104-

¹H-NMR (CDCl₃): δ = 7.42 (1H, d), 7.30 (1H, s), 7.04 (1H, dd), 5.56-5.72 (1H, m), 5.01-5.10 (2H, m), 4.22-4.39 (2H, m), 3.05-3.15 (1H, m), 2.88 (3H, s), 2.33-2.60 (2H, m) ppm.

5

Preparation 95

4(S)-(3,4-Dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene

10

4(S)-(3,4-Dichlorophenyl)-5-(methanesulphonyloxy)pent-1-ene (8.25g) (see Preparation 93) and imidazole (5.3g) were dissolved in anhydrous acetonitrile (80ml) and the mixture heated under reflux for 6 days. The solvent was removed under reduced pressure to give a residue which was partitioned between diethyl ether and aqueous saturated sodium hydrogen carbonate solution (pH of the aqueous phase was kept >7). The phases were separated and the aqueous phase extracted with diethyl ether. The combined organic extracts were then washed with water, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel, eluting with a solvent gradient of 98:2 changing to 95:5, by volume, dichloromethane : methanol to give 4(S)-(3,4-dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene (4.78g) as an oil.


20

¹H-NMR (CDCl₃): δ = 7.35 (1H, d), 7.15-7.30 (2H, m), 6.99 (1H, s), 6.84 (1H, d), 6.69 (1H, s), 5.55-5.70 (1H, m), 5.07 (2H, d), 3.95-4.22 (2H, m), 2.95-3.10 (1H, m), 2.32-2.45 (2H, m) ppm.

25

-105-

Preparation 96

4(R)-(3,4-Dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene


This compound was prepared by an analogous method to that used to prepare the compound of Preparation 95 using 4(R)-(3,4-dichlorophenyl)-5-(methanesulfonyloxy)pent-1-ene (see Preparation 94) as the starting material.

10

$^1\text{H-NMR}$ (CDCl_3): δ = 7.35 (1H, d), 7.15-7.30 (2H, m), 6.99 (1H, s), 6.84 (1H, d), 6.69 (1H, s), 5.55-5.70 (1H, m), 5.07 (2H, d), 3.95-4.22 (2H, m), 2.95-3.10 (1H, m), 2.32-2.45 (2H, m) ppm.

15

Preparation 97

3(S)-(3,4-Dichlorophenyl)-4-(imidazol-1-yl)butan-1-al


-106-

4(S)-(3,4-Dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene (4.75g) (see Preparation 95) was dissolved in a mixture of acetonitrile (75ml), water (13ml) and 1N aqueous hydrochloric acid solution (17ml). Osmium tetroxide (3.4ml of a 0.05M solution in toluene) was added and the mixture stirred for 20 minutes. Sodium periodate (5.3g) was then added together with a further quantity of acetonitrile (30ml) and stirring continued overnight. Water (ca. 100ml) was added and the acetonitrile removed under reduced pressure to give an aqueous suspension which was basified to pH>7 by the addition of solid sodium carbonate. The mixture was then extracted (three times) with ethyl acetate and the combined organic extracts washed sequentially with water and brine before being dried over anhydrous sodium sulphate. Removal of the solvent under reduced pressure gave 3(S)-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-al (4.3g) as an oil.

¹H-NMR (CDCl₃): δ = 9.70 (1H, s), 7.38 (1H, d), 7.20-7.30 (2H, m), 7.01 (1H, s), 6.89 (1H, dd), 6.71 (1H, s), 4.00-4.22 (2H, m), 3.60 (1H, m), 2.72-2.92 (2H, m) ppm.

Preparation 98

3(R)-(3,4-Dichlorophenyl)-4-(imidazol-1-yl)butan-1-al




This compound was prepared by an analogous method to that used to prepare the compound of Preparation 97 using 4(R)-(3,4-dichlorophenyl)-5-(imidazol-1-yl)pent-1-ene (see Preparation 96) as the starting material.

-107-


$^1\text{H-NMR}$ (CDCl_3): δ = 9.70 (1H, s), 7.38 (1H, d), 7.20-7.30 (2H, m), 7.01 (1H, s), 6.89 (1H, dd), 6.71 (1H, s), 4.00-4.22 (2H, m), 3.60 (1H, m), 2.72-2.92 (2H, m) ppm.

5

Preparation 993(S)-(3,4-Dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol

- 10 Sodium borohydride (0.68g) was added carefully to an ice-cooled solution of 3(S)-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-al (4.3g) (see Preparation 97) in ethanol (40ml) and the mixture stirred at room temperature for one hour before removing the solvent under reduced pressure to give a residue. The residue was suspended in water (30ml), cooled in an ice-bath and the mixture first acidified to pH1 with 2N aqueous
- 15 hydrochloric acid solution and then basified to pH14 by addition of 2N aqueous sodium hydroxide solution. The resulting suspension was filtered and the residue washed with water, dried and then crystallised from acetonitrile to give 3(S)-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol (1.37g) as a cream solid.
- 20 $^1\text{H-NMR}$ (CDCl_3): δ = 7.35 (1H, d), 7.15-7.30 (2H, m), 6.95 (1H, s), 6.89 (1H, d), 6.70 (1H, s), 4.00-4.25 (2H, m), 3.60-3.70 (1H, m), 3.40-3.50 (1H, m), 3.15-3.30 (1H, m), 2.10 (1H, br. s), 1.75-2.00 (2H, m) ppm.

-108-

Preparation 1003(R)-(3,4-Dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol

5

This compound was prepared by an analogous method to that used to prepare the compound of Preparation 99 using 3(R)-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-al (see Preparation 98) as the starting material.


- 10 ¹H-NMR (CDCl₃): δ = 7.35 (1H, d), 7.15-7.30 (2H, m), 6.95 (1H, s), 6.89 (1H, d), 6.70 (1H, s), 4.00-4.25 (2H, m), 3.60-3.70 (1H, m), 3.40-3.50 (1H, m), 3.15-3.30 (1H, m), 2.10 (1H, br. s), 1.75-2.00 (2H, m) ppm.

-109-

Preparation 101

3(S)-(3,4-Dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl
1,2,3,4-tetrahydro-5-naphthoate

5



1,2,3,4-Tetrahydro-5-naphthoyl chloride (1.70g) (see Preparation 36) was added dropwise to a suspension of 3(S)-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol (0.90g) (see Preparation 99) and triethylamine (1.45g) in anhydrous acetonitrile (15ml) and the solution stirred at room temperature for 4 days. The solvent was then removed under reduced pressure to give a residue which was partitioned between ethyl acetate and saturated aqueous sodium hydrogen carbonate solution. The phases were separated and the aqueous phase further extracted with ethyl acetate. The organic phases were combined, washed sequentially with water and brine, dried over anhydrous sodium sulphate and the solvent removed under reduced pressure to give a residue which was chromatographed on silica gel eluting with a solvent gradient of 2:1 changing to 0:1, by volume, pentane : ethyl acetate to give 3(S)-(3,4-dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl 1,2,3,4-tetrahydro-5-naphthoate (1.38 g) as a gum.

20


-110-

$^1\text{H-NMR}$ (CDCl_3): δ = 7.50 (1H, d), 6.95-7.38 (10H, m), 4.83 (1H, dd), 4.63 (1H, m), 4.30-4.35 (1H, m), 4.08-4.16 (1H, m), 3.30-3.40 (1H, m), 3.00 (2H, br. s), 2.63-2.84 (6H, m), 2.10-2.25 (2H, m), 1.67-1.83 (8H, m) ppm.

5

Preparation 102

3(R)-(3,4-Dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl
1,2,3,4-tetrahydro-5-naphthoate



10

This compound was prepared by an analogous method to that used to prepare the compound of Preparation 101 using 3(R)-(3,4-dichlorophenyl)-4-(imidazol-1-yl)butan-1-ol (see Preparation 100) as the starting material.

15


$^1\text{H-NMR}$ (CDCl_3): δ = 7.50 (1H, d), 6.95-7.38 (10H, m), 4.83 (1H, dd), 4.63 (1H, m), 4.30-4.35 (1H, m), 4.08-4.16 (1H, m), 3.30-3.40 (1H, m), 3.00 (2H, br. s), 2.63-2.84 (6H, m), 2.10-2.25 (2H, m), 1.67-1.83 (8H, m) ppm.

-111-

Preparation 103

3(S)-(3,4-Dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butan-1-ol

5



- 3(S)-(3,4-Dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl
 10 1,2,3,4-tetrahydro-5-naphthoate (1.38 g) (see Preparation 101) was dissolved in
 methanol (40ml), 1N aqueous sodium hydroxide solution (5ml) added and the resulting
 mixture stirred at room temperature overnight. The methanol was then removed under
 reduced pressure and the residue partitioned between ethyl acetate and water. The
 organic phase was separated, washed sequentially with saturated aqueous sodium
 15 hydrogen carbonate solution and brine, dried over anhydrous sodium sulphate and the
 solvent removed under reduced pressure to give 3(S)-(3,4-dichlorophenyl)-4-[2-
 (1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butan-1-ol (0.82 g) as a gum.


- ¹H-NMR (CDCl₃): δ = 7.10-7.35 (5H, m), 7.06 (1H, s), 6.92 (1H, dd), 6.74 (1H, s),
 20 4.90 (1H, dd), 4.50 (1H, dd), 3.70-3.82 (1H, m), 3.53-3.64 (1H, m), 3.30-3.42 (1H, m),
 2.70-2.86 (4H, m), 1.70-2.04 (6H, m) ppm.

-112-

Preparation 104

3(R)-(3,4-Dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butan-1-ol

5



This compound was prepared by an analogous method to that used to prepare the
 10 compound of Preparation 103 using 3(R)-(3,4-dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-
 5-naphthoyl)imidazol-1-yl]butyl 1,2,3,4-tetrahydro-5-naphthoate (see Preparation 102)
 as the starting material.


¹H-NMR (CDCl₃): δ = 7.10-7.35 (5H, m), 7.06 (1H, s), 6.92 (1H, dd), 6.74 (1H, s),
 15 4.90 (1H, dd), 4.50 (1H, dd), 3.70-3.82 (1H, m), 3.53-3.64 (1H, m), 3.30-3.42 (1H, m),
 2.70-2.86 (4H, m), 1.70-2.04 (6H, m) ppm.

-113-

Preparation 105

1-Methanesulphonyloxy-3(S)-(3,4-dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butane

5



3(S)-(3,4-Dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butan-1-ol (0.68g) (see Preparation 103) and triethylamine (0.22g) were dissolved in
 10 dichloromethane (15ml), the solution cooled in an ice-bath and methanesulphonyl chloride (0.22g) added. The mixture was stirred for 4 hours before removal of the solvent under reduced pressure to give a residue which was partitioned between ethyl acetate and water. The organic phase was separated and washed with saturated aqueous sodium hydrogen carbonate solution before being dried over anhydrous sodium
 15 sulphate. The solvent was removed under reduced pressure to give 1-methanesulphonyloxy-3(S)-(3,4-dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butane (0.82g) as an oil.


¹H-NMR (CDCl₃): δ = 7.37 (1H, d), 7.05-7.31 (5H, m), 6.99 (1H, dd), 6.84 (1H, s),
 20 4.80 (1H, dd), 4.65 (1H, dd), 4.05-4.30 (2H, m), 3.30-3.40 (1H, m), 2.97 (3H, s), 2.65-
 2.85 (4H, m), 2.10-2.30 (2H, m), 1.70-1.85 (4H, m) ppm.

-114-

Preparation 106

1-Methanesulphonyloxy-3(R)-(3,4-dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butane

5



This compound was prepared by an analogous method to that used to prepare the compound of Preparation 105 using 3(R)-(3,4-dichlorophenyl)-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butan-1-ol (see Preparation 104) as the starting material.

$^1\text{H-NMR}$ (CDCl_3): δ = 7.37 (1H, d), 7.05-7.31 (5H, m), 6.99 (1H, dd), 6.84 (1H, s), 4.80 (1H, dd), 4.65 (1H, dd), 4.05-4.30 (2H, m), 3.30-3.40 (1H, m), 2.97 (3H, s), 2.65-2.85 (4H, m), 2.10-2.30 (2H, m), 1.70-1.85 (4H, m) ppm.

-115-

PHARMACOLOGICAL DATA

The affinities of the following compounds of the Examples for the human NK₁ and NK₂ receptors were determined by the methods described on page 24, lines 8 to 22, of the description, and the results are presented in the Table below.


Example No.	NK1 Binding (IC ₅₀)	NK2 Binding (IC ₅₀)
Example 2	0.6 nM	4 nM
Example 5	8 nM	11 nM
Example 6	9 nM	96 nM
Example 18	123 nM	3 nM
Example 19	2 nM	24 nM

These data show that the compounds are dual NK₁ and NK₂ receptor antagonists.

-116-

CLAIMS

1. A compound of the formula:



wherein


R is phenyl, C₃-C₇ cycloalkyl or heteroaryl, each of which being optionally benzo- or C₃-C₇ cycloalkyl-fused and optionally substituted, including in the benzo- or C₃-C₇ cycloalkyl-fused portion, by from 1 to 3 substituents each independently selected from C₁-C₄ alkyl, fluoro(C₁-C₄)alkyl, C₁-C₄ alkoxy, fluoro(C₁-C₄)alkoxy, phenoxy, C₂-C₄ alkanoyl, halo, C₁-C₄ alkoxycarbonyl, C₃-C₇ cycloalkyl, -S(O)_m(C₁-C₄ alkyl), cyano, -NR²R³, -S(O)_mNR²R³, -NR⁴(C₁-C₄ alkanoyl) and -CONR²R³, or R is 2,3-dihydrobenzo[b]furanyl or chromanyl;

R¹ is H or C₁-C₆ alkyl;

R² and R³ are either each independently selected from H and C₁-C₆ alkyl, or when taken together, represent C₄-C₆ alkylene;

R⁴ is H or C₁-C₆ alkyl;

W is a direct link, methylene or ethylene;



-117-

X is unbranched C₂-C₄ alkylene;

Y is phenyl, naphthyl, benzyl, pyridyl, thienyl or C₃-C₇ cycloalkyl, each of which being optionally substituted by from 1 to 3 substituents each independently selected from C₁-C₄ alkyl, fluoro(C₁-C₄)alkyl, C₁-C₄ alkoxy, fluoro(C₁-C₄)alkoxy, halo and cyano;

Ar is phenyl, naphthyl, benzyl, thienyl, benzo[b]thienyl or indolyl, each of which being optionally substituted by from 1 to 3 substituents each independently selected from C₁-C₄ alkyl, fluoro(C₁-C₄)alkyl, C₁-C₄ alkoxy, fluoro(C₁-C₄)alkoxy, halo and cyano, or Ar is 1,3-benzodioxolan-4 or 5-yl or 1,4-benzodioxan-5 or 6-yl;

m is 0, 1 or 2;

Z^A is a pharmaceutically acceptable anion;

and "heteroaryl", used in the definition of R, means thienyl or a 5- or 6-membered ring heteroaryl group containing either from 1 to 4 nitrogen heteroatoms, or 1 or 2 nitrogen heteroatom(s) and 1 oxygen or sulphur heteroatom,


with the proviso that when W is a direct link and R is optionally fused and optionally substituted heteroaryl, said heteroaryl is linked by a ring carbon atom to the carbonyl group.



* C L M S *


-118-

2. A compound as claimed in claim 1 wherein R is phenyl which is optionally benzo- or C₃-C₇ cycloalkyl-fused and optionally substituted, including in the benzo- or C₃-C₇ cycloalkyl-fused portion, by 1, 2 or 3 substituents each independently selected from C₁-C₄ alkyl, fluoro(C₁-C₄)alkyl, C₁-C₄ alkoxy, fluoro(C₁-C₄)alkoxy, phenoxy and halo, or R is 2,3-dihydrobenzo[b]furanyl.
3. A compound as claimed in claim 1 or 2 wherein R is phenyl which is optionally benzo- or C₃-C₇ cycloalkyl-fused and optionally substituted, including in the benzo- or C₃-C₇ cycloalkyl-fused portion, by 1, 2 or 3 substituents each independently selected from methyl, ethyl, trifluoromethyl, methoxy, isopropoxy, trifluoromethoxy, phenoxy, fluoro and chloro, or R is 2,3-dihydrobenzo[b]furanyl.
4. A compound as claimed in claim 1, 2 or 3 wherein R is phenyl, naphthyl or tetrahydronaphthyl, each of which being optionally substituted by 1, 2 or 3 substituents each independently selected from methyl, ethyl, trifluoromethyl, methoxy, isopropoxy, trifluoromethoxy, phenoxy, fluoro and chloro, or R is 2,3-dihydrobenzo[b]furanyl.
5. A compound as claimed in any one of claims 1 to 4 wherein R is phenyl, 3,5-dimethylphenyl, 2,3-dimethylphenyl, 2-trifluoromethoxyphenyl, 2-methoxy-3-methylphenyl, 2,3-dihydrobenzo[b]furan-7-yl, naphth-2-yl, 4-fluoro-3-trifluoromethylphenyl, 1,2,3,4-tetrahydronaphth-5-yl, 1,2,3,4-tetrahydronaphth-6-yl, 5-chloro-2-methoxyphenyl, 2-methoxyphenyl, 2-trifluoromethylphenyl, 2-isopropoxyphenyl, 2-ethylphenyl, 2-phenoxyphenyl or 3,5-bis(trifluoromethyl)phenyl.
6. A compound as claimed in any one of claims 1 to 5 wherein R is 2,3-dimethylphenyl, naphth-2-yl, 1,2,3,4-tetrahydronaphth-5-yl or 2-methoxyphenyl.




-119-

7. A compound as claimed in any one of claims 1 to 6 wherein R^1 is H.
8. A compound as claimed in any one of claims 1 to 7 wherein W is a direct link or methylene.
9. A compound as claimed in any one of claims 1 to 8 wherein W is a direct link.
10. A compound as claimed in any one of claims 1 to 9 wherein X is 1,2-ethylene.
11. A compound as claimed in any one of claims 1 to 10 wherein Y is phenyl, naphthyl or cyclohexyl, each of which being optionally substituted by 1, 2 or 3 C_1 - C_4 alkyl substituents.
12. A compound as claimed in any one of claims 1 to 11 wherein Y is phenyl, 3,5-dimethylphenyl, cyclohexyl or naphth-2-yl.
13. A compound as claimed in any one of claims 1 to 12 wherein Y is phenyl.
14. A compound as claimed in any one of claims 1 to 13 wherein Ar is phenyl optionally substituted by 1, 2 or 3 halo substituents.
15. A compound as claimed in any one of claims 1 to 14 wherein Ar is phenyl substituted by 1 or 2 chloro substituents.
16. A compound as claimed in any one of claims 1 to 15 wherein Ar is 3,4-dichlorophenyl.



-120-

17. A compound as claimed in any one of claims 1 to 16 wherein Z^A is chloride, bromide, nitrate, methanesulphonate, para-toluenesulphonate, benzenesulphonate, hydrogen sulphate or sulphate.
18. A compound as claimed in any one of claims 1 to 17 wherein Z^A is chloride or methanesulphonate.
19. A compound as claimed in any one of claims 1 to 18 wherein Z^A is methanesulphonate.
20. A compound as claimed in claim 1 wherein R^1 is H, X is $-\text{CH}_2\text{CH}_2-$ and Ar is 3,4-dichlorophenyl, and wherein
- R-W- is 3,5-dimethylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2,3-dimethylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2-trifluoromethoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2-methoxy-3-methylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2,3-dihydrobenzo[b]furan-7-yl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is naphth-2-yl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 4-fluoro-3-trifluoromethylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 1,2,3,4-tetrahydronaphth-5-yl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 1,2,3,4-tetrahydronaphth-6-yl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 5-chloro-2-methoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2-methoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2-trifluoromethylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2-isopropoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2-ethylphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is 2-phenoxyphenyl, Y is phenyl and Z^A is CH_3SO_3^- ;
 - R-W- is benzyl, Y is phenyl and Z^A is CH_3SO_3^- ;




* C L M S *

-121-


R-W- is 3,5-bis(trifluoromethyl)phenyl, Y is phenyl and Z^A is Cl^- ;
 R-W- is 2-methoxyphenyl, Y is cyclohexyl and Z^A is CH_3SO_3^- ;
 R-W- is 4-fluoro-3-trifluoromethylphenyl, Y is cyclohexyl and Z^A is CH_3SO_3^- ;
 R-W- is 2-methoxyphenyl, Y is 3,5-dimethylphenyl and Z^A is CH_3SO_3^- ; or
 R-W- is 2-methoxyphenyl, Y is naphth-2-yl and Z^A is CH_3SO_3^- ;
 or wherein Z^A is an alternative pharmaceutically acceptable anion in respect of any thereof.

21. A compound as claimed in claim 1 which is 4-phenyl-1-(3(S)-[3,4-dichlorophenyl]-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate or 4-phenyl-1-(3(R)-[3,4-dichlorophenyl]-4-[2-(1,2,3,4-tetrahydro-5-naphthoyl)imidazol-1-yl]butyl)quinuclidinium methanesulphonate.


22. A process for the preparation of a compound of the formula (I) as claimed in claim 1 wherein R, R^1 , W, X, Y, Ar and Z^A are as defined in claim 1 comprising reaction of a compound of the formula:



wherein R, R^1 , Ar, W and X are as previously defined for a compound of the formula (I) in this claim, Z is a leaving group capable of forming a pharmaceutically acceptable anion (Z^A) and Z^1 is a leaving group, with a compound of the formula:



-122-



(III)

wherein Y is as previously defined for a compound of the formula (I), said process being followed by either (a), where Z^1 is a leaving group, exchange for a pharmaceutically acceptable anion (Z^A), or (b), optionally, where Z^A is a pharmaceutically acceptable anion, exchange for another pharmaceutically acceptable anion.


23. A process as claimed in claim 22 wherein Z is C_1 - C_4 alkanesulphonyloxy, benzenesulphonyloxy, para-toluenesulphonyloxy, chloro, bromo or iodo, and Z^1 is trifluoromethanesulphonyloxy.

24. A process as claimed in claim 22 wherein Z is methanesulphonyloxy and Z^A is methanesulphonate.

25. A pharmaceutical composition comprising a compound of the formula (I) as claimed in any one of claims 1 to 21, together with a pharmaceutically acceptable diluent or carrier.

26. A compound of the formula (I) or a pharmaceutically acceptable composition thereof, as claimed in any one of claims 1 to 21 and 25, respectively, for use as a medicament.

27. The use of a compound of the formula (I), or of a pharmaceutically acceptable composition thereof, as claimed in any one of claims 1 to 21 and 25, respectively, for the manufacture of a medicament for the treatment of a disease by producing an antagonist effect on a tachykinin receptor or on a combination of tachykinin receptors.



-123-

28. Use as claimed in claim 27 where the antagonist effect is on the human NK_1 and NK_2 tachykinin receptors.

29. Use as claimed in claim 27 or 28 where the disease is an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis or Crohn's disease, a disease caused by Helicobacter pylori or another urease-positive Gram negative bacteria, a urogenital tract disorder such as incontinence, hyperreflexia or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, emesis, cough, migraine or acute or chronic pain.

30. A method of treatment of a human to treat a disease by producing an antagonist effect on a tachykinin receptor or on a combination of tachykinin receptors, which comprises treating said human with an effective amount of a compound of the formula (I) or with a pharmaceutically acceptable composition thereof, as claimed in any one of claims 1 to 21 and 25, respectively.



31. A method as claimed in claim 30 where the antagonist effect is on the human NK_1 and NK_2 tachykinin receptors.









-124-

32. A method as claimed in claim 30 or 31 where the disease is an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis or Crohn's disease, a disease caused by Helicobacter pylori or another urease-positive Gram negative bacteria, a urogenital tract disorder such as incontinence, hyperreflexia or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, emesis, cough, migraine or acute or chronic pain.

33. A compound of the formula:




-125-




* C L H S *

-126-



(XXX) ; or



(XXXI) :

wherein R, R¹, W, X and Ar are as defined in claim 1, Z is a leaving group capable of forming a pharmaceutically acceptable anion (Z^A), Z¹ is a leaving group and P is a protecting group.

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced

