


Ibandronate polymorph A

The present invention relates to a new polymorph crystal form of 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt monohydrate (Ibandronate) with the following formula



Formula 1

5

and a process for its preparation.

Ibandronate is one of the most potent anti-resorptive drugs that directly inhibit osteoclast activity and present an effective pharmacologic alternative for controlling hypercalcemia. Ibandronate binds to hydroxyapatite in calcified bone, rendering it
 10 resistant to hydrolytic dissolution by phosphatases, thereby inhibiting both normal and abnormal bone resorption. This drug increases bone mass and decreases the risk of fractures and is therefore particularly well adapted to bone and calcium metabolic diseases such as for instance osteoporosis or Paget's disease (EP-A 0252504).

It has been found that Ibandronate may exist in various polymorphs.

15 One polymorph of Ibandronate hereinafter also designated as polymorph A was identified as being thermodynamically more stable, whereas a second polymorph of Ibandronate, hereinafter also designated as polymorph B, is easier to separate within the production process.

20 The ability of a substance to exist in more than one crystal form is defined as polymorphism and these different crystal forms are known as "polymorph modifications" or "polymorphs". Polymorphism can influence many aspects of solid state properties of a drug. Different crystal modifications of a substance may differ considerably from one to another in different physical properties which may influence directly their solubility for instance. Polymorphism is found in several organic compounds.

An exhaustive treatment of polymorphism in pharmaceutical and molecular crystals is given e.g. by H.G. Brittain in *Polymorphism in Pharmaceutical Solids*, H.G. Brittain or, Marcel Dekker Inc., New York, 1999 and in *Solid-State Chemistry of Drugs*, SSCI Inc., West Lafayette; Indiana, 1999

Object of the embodiments disclosed herein is to specifically isolate and characterize Ibandronate polymorph A and to develop a process for the preparation of Ibandronate polymorph A.

The object has been achieved with the identification of crystalline polymorph A of Ibandronate and with a process for its preparation as discussed herein.

In a first aspect, there is provided a crystalline polymorph of 3-(N-methyl-N-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid, monosodium salt monohydrate (Ibandronate) which is characterized by an X-ray powder diffraction pattern having characteristic peaks expressed in angle 2-theta at approximately

Angle 2-theta
10.2°
11.5°
15.7°
19.4°
26.3°

In a second aspect, there is provided a crystalline Ibandronate polymorph which is characterized by an IR absorption spectrum having characteristic peaks expressed in cm^{-1} at approximately 3678 cm^{-1} , 3164 cm^{-1} , 2854 cm^{-1} , 1377 cm^{-1} , 1288 cm^{-1} , 1157 cm^{-1} , 1094 cm^{-1} , 1069 cm^{-1} , 1035 cm^{-1} , 966 cm^{-1} , 951 cm^{-1} , 933 cm^{-1} , 903 cm^{-1} , 760 cm^{-1} and 723 cm^{-1} .

In a third aspect, there is provided a crystalline Ibandronate polymorph which is characterized by a characteristic peak expressed in cm^{-1} at approximately 1460 cm^{-1} in a vibrational Raman spectrum.

In a fourth aspect, there is provided an Ibandronate containing at least 80% of a crystalline polymorph as characterized above.

In a fifth aspect, there is provided a process for the preparation of a crystalline Ibandronate polymorph as characterized above comprising crystallizing a 3-(N-methyl-N-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt or a monohydrate, a polymorph or a polymorph mixture thereof at a temperature of 50°C to 70°C in a polar solvent.

In a sixth aspect, there is provided a process for the preparation of a crystalline Ibandronate polymorph as characterized above comprising tempering a moist 3-(N-methyl-N-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt or a monohydrate, a polymorph or a polymorph mixture thereof at a temperature of 30°C to 90°C.

In a seventh aspect, there is provided pharmaceutical compositions comprising a crystalline polymorph as defined above or obtained as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

In an eighth aspect, there is provided a crystalline polymorph as defined above or obtained as defined above for use as therapeutic active substance.

Unless otherwise indicated, the following definitions are set forth to define the meaning and scope of the various terms used and described in the description.

The term "Ibandronate polymorph A" refers to the polymorph crystal form of 3-(N-methyl-N-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt monohydrate as defined in claims and description of the present invention.

The term "crystalline polymorph" refers to a crystal modification which can be characterized by analytical methods such as e.g. X-ray powder diffraction, IR spectroscopy and Raman spectroscopy.

The term IR means infrared.

Description of figures:

Figure 1: shows an X-ray diffraction pattern of crystalline Ibandronate polymorph A as obtained in example 1.

Figure 2: shows an IR-spectrum of crystalline Ibandronate polymorph A

Figure 3: shows a Raman spectrum of crystalline Ibandronate polymorph A

Figure 4: shows an X-ray diffraction pattern of crystalline Ibandronate polymorph B as obtained in Ref. example 2.

Figure 5: shows an IR-spectrum of crystalline Ibandronate polymorph B

Figure 6: shows a Raman spectrum of crystalline Ibandronate polymorph B.

The crystalline Ibandronate polymorph A of the present invention can be characterized by an X-ray powder diffraction pattern having characteristic peaks expressed in angle 2-theta at approximately:

Angle 2-theta	Intensity [%]
10.2°	0.6
11.5°	0.7
15.7°	1.6
19.4°	1.9
26.3°	3.3

5

Angle 2-theta means an incertitude of ± 0.2 (expressed in degrees), 2-theta represent the reflexion angle according to the Bragg law. A reflection at a glancing angle θ gives rise to a reflection at an angle 2θ to the direction of the incident beam.

The crystalline Ibandronate polymorph A as described above can further be
10 characterized by the x-ray powder diffraction pattern as shown in Figure 1.

The crystalline Ibandronate polymorph A can furthermore be characterized by its IR absorption spectrum having characteristic peaks expressed in cm^{-1} at approximately:

5

10

Bands [cm ⁻¹]
3678
3164
2854
1377
1288
1157
1094
1069
1035
966
951
933
903
760
723

The term approximately means in this context that the cm⁻¹ value can vary by about ± 4 cm⁻¹.

- 15 The crystalline Ibandronate polymorph A can further be characterized by the IR absorption spectrum shown in figure 2.

The crystalline Ibandronate polymorph A as described above can furthermore be described by Raman vibrational spectroscopy. The Raman spectrum has the following characteristic bands expressed in cm⁻¹ at approximately:

Bands [cm^{-1}]
2950
2927
2889
2851
1460
1443
1308
1137
1056
1024
954
904
839
761
678

5

10 The term approximately means in this context that the cm^{-1} value can vary by $\pm 8 \text{ cm}^{-1}$.

The crystalline Ibandronate polymorph A can further be characterized by the Raman spectrum shown in Figure 3.

15 The crystalline Ibandronate polymorph A described above is further characterized by a solubility in water of about 278 g/l at 25°C.

The process according to the present invention is characterized by a crystallization of 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt or a monohydrate, a polymorph or a mixture of polymorphs at a temperature of 50°C to 70°C in a polar solvent.

20 As polar solvent preferably water is used.

Preferably the crystallization takes place in water at a temperature of about 60°C.

A polar aprotic solvent may be added for inducing the crystallization.

A suitable polar aprotic solvent is acetone.

Expediently the crystallization temperature is maintained for 15 minutes to 120 minutes.

Starting product for the process of the present invention can either be 3-(*N*-methyl-
5 *N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt e.g. obtained from a process as outlined in reference example 1, or crystalline Ibandronate polymorph B e.g. obtained according to reference example 2, or from a mixture of crystalline Ibandronate polymorph B with crystalline Ibandronate polymorph A.

The starting product can either be dissolved in the polar solvent at about room
10 temperature and then be warmed to the crystallization temperature or alternatively be dissolved at a higher temperature and then be cooled to the crystallization temperature as indicated above.

Ethanol residues which may be present from the manufacturing process (according to reference example 1) can easily be removed by known methods e.g. by distilling the
15 ethanol off as azeotrope.

Crystallization as a rule occurs spontaneously but can also be initiated by addition of crystals of Ibandronate polymorph A.

The crystal suspension so obtained is as a rule cooled under stirring until crystallization is complete before the filtration takes place.

20 The whole crystallization process may be controlled regarding temperature, heating and cooling periods with equipment common to the skilled practitioner.

Separation of the desired polymorph can be effected by common filtration techniques. As a rule the precipitate is washed with the polar solvent used for the crystallization, preferably with a mixture of water and acetone in a ratio of about 1:1
25 (V/V).

Drying of the crystalline Ibandronate polymorph A preferably happens at a temperature from 40°C to 80°C for about 9 hours to 72 hours either at normal or at reduced pressure.

The crystalline Ibandronate polymorph according to the present invention can be
30 obtained with a content of the crystalline Ibandronate polymorph A of at least 80%.

Alternatively the crystalline Ibandronate polymorph A can be obtained by tempering a moist 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt e.g. obtained from a process as outlined in reference example 1, or a crystalline Ibandronate polymorph B e.g. obtained according to reference example 2, or
5 from a mixture of crystalline Ibandronate polymorph B with crystalline Ibandronate polymorph A at a temperature of 30°C to 90°C.

Ethanol residues which may be present from the manufacturing process (according to reference example 1) can easily be removed by known methods e.g. by distilling the ethanol off as azeotrope.

10 The term "moist" in this context means that the starting products contain some, as a rule about 10% water.

The crystalline Ibandronate polymorph A of the present invention can be used as a pharmaceutically active compound which acts as an efficient anti-resorptive drug that directly inhibits osteoclasts activity and thus also increases bone mass. Accordingly this
15 polymorph can be used for the treatment and/or prevention of diseases which are associated with bone and calcium metabolic diseases such as for instance osteoporosis or Paget's disease.

The invention therefore also relates to pharmaceutical compositions comprising a crystalline Ibandronate polymorph A as defined above and a pharmaceutically acceptable
20 carrier and/or adjuvant.

The invention also encompasses a crystalline Ibandronate polymorph A as described above for use as therapeutically active substances.

The following examples serve as illustration of the invention.

Examples

X-ray powder diffraction measurement

The x-ray powder diffraction patterns of the individual crystalline Ibandronate polymorphs A and B were recorded with a Bruker D8 Advance AXS diffractometry (geometry: Bragg-Brentano; radiation: CuK α in the angle range $2\theta = 2^\circ$ to 40° ; Cu secondary monochromator; step scan with 0.02° and a measuring time of e.g. 4.0 s per step). The samples weighing approximately 500 mg were prepared into carrier wells and exposed to CuK α radiation. Radiation diffracted at the lattice levels is converted into electronic signals by a scintillation counter and processed by the software package "Diffrac plus". The x-ray powder diffraction patterns of the individual crystalline Ibandronate polymorphs A and B are shown in Figure 1 and 4.

IR measurement method

The IR spectra of the individual crystalline Ibandronate polymorphs A and B were recorded as film of Nujol suspension consisting of approx. 15 mg of sample in approx. 15 mg of Nujol between two sodium chloride plates. Measurements were carried out with an FT-IR spectrometer (IFS55 (Bruker) or equivalent instrument) in transmittance mode (resolution 4 cm^{-1} , detector: DTGS). The IR spectra of the individual crystalline Ibandronate polymorphs A and B are shown in Figure 2 and 5.

Raman measurement method

The Raman spectra of the individual crystalline Ibandronate polymorphs A and B were recorded as sample of about 20 mg of the powder and were filled into a glass tube (shortened NMR tube). The samples were measured with the FT-Raman of Nicolet coupled to a Magna 860 (Nicolet) in 90° scattering arrangement, detector: InGaAs. Measurement parameters resolution 8 cm^{-1} ; laser power 0.95W, No of scan 300. The FT-Raman spectra of the individual crystalline Ibandronate polymorphs A and B are shown in Figure 3 and 6.

Solubility measurement

The solubility of the individual crystalline Ibandronate polymorphs A and B was measured for various solutions. Approximately 10 g of the corresponding polymorph A or B were suspended in three different buffer solutions at pH 2, pH 4 (Titrisol-buffer, citrate/HCl), at pH 7 (methenamine buffer, HCl) or in water. The suspension was stirred

for 24 hours at 25°C and was kept additionally 24 hours without stirring at the same temperature. Solubility was calculated by titration according to the following method.

- The residue was filtered, 2 ml of the filtrate taken, 5 ml of Titriplex III solution were added and diluted with water to 100 ml. 2 ml of this solution were added to
- 5 approximately 0.1 ml of xylenol orange indicator and the pH was adjusted to 6.5 by adding small portions of methenamine buffer solution or 0.1 M hydrochloric acid. The solution is titrated immediately with Th-DCTA-xylenol orange complex until the color changes from yellow to reddish-violet. The endpoint is determined photometrically.

Results are gathered in the following table

10

	Polymorph A solubility in [g/l]	Polymorph B solubility in [g/l]*
pH 2	280	274
pH 4	276	278
pH 7	292	299
water	278	279

*under at least partial conversion to polymorph A

Reference example 1

- 15 Preparation of 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt

- 250 g (1.19 mol) *N*-methyl-*N*-pentyl- β -alanine hydrochloride, 233 g (2.84 mol) phosphorous acid, 151 ml (1.65 mol) phosphorous oxychloride and 900 ml diethyl carbonate were heated stepwise to 80°C. After 2 hours reaction time under continued heating the mixture was cooled to 60°C and 1733 ml demineralized water were added,
- 20 followed by azeotropic distillation of diethyl carbonate / water at 90 to 101°C. 358 ml demineralized water were added, the mixture was refluxed and water was distilled off. 316 ml demineralized water were added and water was distilled off twice. Finally 2040 ml demineralized water were added and the residue was cooled to 24°C. The pH was adjusted at 23°C with sodium hydroxide solution (50%) to 4.4. Thereafter, 1100 ml

- 10 -

ethanol were added to start crystallization. The suspension was stirred for 8 hours at 21 to 22°C. Then the solid was separated, washed with 344 ml cold ethanol/demineralized water (7/5 V/V), subsequently with 344 ml acetone / demineralized water (5/2 V/V) and dried at 60°C. 315.6 g (73.7 %) of the title product were obtained in the form of colorless
5 crystals.

Assay (complexometric titration): 100.6 % (calculated on the anhydrous and solvent free basis)

Residual solvents: 2.3% ethanol (GC)
3.9 % water (KF)

10

Reference example 2

Preparation of crystalline Ibandronate polymorph B

55 g of 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt (obtained according to reference example 1) were dissolved in 240 ml
15 demineralized water. 75 ml water were distilled off. After filtration, the remaining solution was brought to 35°C and 190 ml acetone were added in 20 minutes. Thereafter, the mixture was cooled until $\leq 25^{\circ}\text{C}$ under stirring, up to the complete crystallization. The product was separated and washed with a mixture of acetone / demineralized water in a ratio of 1:1 (V/V). The product was dried in vacuum from 150 to 20 mbar for 12
20 hours at 40°C and 24 hours at 60°C.

Yield: 81 %

Identified with x-ray powder diffraction as crystalline Ibandronate polymorph B.
(Figure 4)

25

Example 1

Preparation of crystalline Ibandronate polymorph A.

150 g of 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid
5 monosodium salt (obtained according to reference example 1) were dissolved in 390 ml
demineralized water at about 70-90°C. 205 ml water were distilled off. After filtration the
solution was cooled to 60°C and stirred for 45 minutes. Crystallization was initiated by
means of crystalline Ibandronate polymorph A. After crystallization the suspension was
cooled to about 20-25°C under stirring until the crystallization was complete. The
10 product was separated and washed with a mixture of 50 ml acetone/demineralized water
in a ratio of 1:1 (V/V). The product was dried for 48 hours in a vacuum from 150 to 20
mbar at 60°C.

Yield: 75 %

Identified with x-ray powder diffraction as crystalline Ibandronate polymorph A.
15 (Figure 1)

Assay (complexometric titration): 101.0 % (calculated on the anhydrous and
solvent free basis)

Example 2

20 Preparation of crystalline Ibandronate polymorph A

150 g of crystalline Ibandronate polymorph B (obtained according to reference
example 2) were dissolved in 185 ml demineralized water at about 90°C. The solution was
then cooled to 60°C and stirred for 30 min. Crystallization was initiated by means of
crystalline Ibandronate polymorph A. The suspension was cooled to about 20-25°C until
25 the crystallization was complete. The product was separated and washed with a mixture
of 50 ml acetone/demineralized water in a ratio of 1:1 (V/V). The product was dried for
48 hours in vacuum from 150 to 20 mbar at 60 °C.

Yield: 80 %

Identified with x-ray powder diffraction as crystalline Ibandronate polymorph A.
30 (Figure 1)

Example 3

Preparation of crystalline Ibandronate polymorph A

15 150 g of 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt (obtained according to reference example 1), were dissolved in 390 ml demineralized water at about 70-90°C. 205 ml water were distilled off. After filtration the filtrate was cooled to 60°C and stirred for 45 minutes. Crystallization was initiated by means of crystalline Ibandronate polymorph A. After crystallization a mixture of demineralized water/acetone (290 ml/518 ml) previously heated to 50°C was added under
10 stirring. Thereafter, the suspension was cooled to about 20-25°C under stirring, until the crystallization was complete. The product was separated and washed with a mixture of 50 ml acetone/demineralized water in a ratio of 1:1 (V/V). The product was dried for 48 hours in a vacuum from 150 to 20 mbar at 60°C.

Yield: 85 %

15 Identified with x-ray powder diffraction as crystalline Ibandronate polymorph A. (Figure 1)

Assay (complexometric titration): 101.0 % (calculated on the anhydrous and solvent free basis)

20

Example 4

Preparation of crystalline Ibandronate polymorph A

100 g of crystalline Ibandronate polymorph B (obtained according to reference example 2) were dissolved in 304 ml demineralized water, warmed to 60°C and it was filtrated. To the filtrate which was kept at 55°C, 347 ml acetone were added dropwise
25 during 1 hour. The mixture was maintained at this temperature under stirring for 2 hours, then cooled to 15-20°C. The product was isolated by filtration and washed with 120 ml of a solvent mixture of acetone/demineralized water 1:1 (V/V). The product was dried in a vacuum from 150 to 20 mbar at 40°C for 14 hours and then at 60°C for 24 hours.

30 Yield: 88 %

Identified with x-ray powder diffraction as crystalline Ibandronate polymorph A.
(Figure 1)

Example 5

5 Preparation of crystalline Ibandronate polymorph A

100 g crystalline Ibandronate polymorph B (obtained according to reference example 2) were dissolved in 304 ml demineralized water, warmed to 60°C and the solution was then filtrated. To the filtrate which was kept at 55°C 347 ml acetone were added dropwise during 1 hour. In the course of the acetone addition crystals of
10 Ibandronate polymorph A were added in order to initiate crystallization. The mixture was maintained at this temperature under stirring for 2 hours then cooled to 15-20°C. The product was isolated by filtration and washed with 120 ml of a solvent mixture of acetone/demineralized water (1:1). The product was dried in a vacuum from 150 to 20 mbar at 40°C for 14 hours and then at 60°C for 24 hours.

15 Yield: 90 %

Identified with x-ray powder diffraction as crystalline Ibandronate polymorph A.
(Figure 1)

Example 6

20 Preparation of crystalline Ibandronate polymorph A (tempering)

30 g freshly precipitated Ibandronate obtained according to reference example 2, but directly used after filtration without drying and verified as polymorph B (moisture content about 10%, dried sample verified as polymorph B) were heated for 45 min at 60°C and 60 mbar until acetone is evaporated, followed by tempering at 60°C and 900
25 mbar for 19 hours and drying at 60°C and 40 mbar for 9 hours under rotary motion.

Yield: quantitative

Identified with x-ray powder diffraction as crystalline Ibandronate polymorph A.
(content polymorph B: 19 %)

Example 7

Preparation of crystalline Ibandronate polymorph A

5 30 g crystalline Ibandronate polymorph B obtained according to reference example
2 were suspended in a mixture of 95.4 ml demineralized water and 103.4 ml acetone at
20°C. The suspension was heated to 60°C in 1 hour, stirred for 15 min., cooled to 20°C in
1 hour and stirred for 5 min. at 20°C. This cycle was repeated twice. The suspension was
stirred for 17 hours. The product was separated and dried for 18 hours in a vacuum from
10 150 to 20 mbar at 60°C.

Identification with x-ray-powder diffraction as crystalline Ibandronate polymorph
A (content polymorph B: 23 %).

Claims

1. A crystalline polymorph of 3-(N-methyl-N-pentyl) amino-1-hydroxypropane-1, 1-diphosphonic acid, monosodium salt monohydrate (Ibandronate) which is characterized by an X-ray powder diffraction pattern having characteristic peaks
- 5 expressed in angle 2-theta at approximately

Angle 2-theta
10.2°
11.5°
15.7°
19.4°
26.3°

2. A crystalline polymorph of claim 1, characterized by the x-ray powder diffraction pattern shown in Figures 1.
3. A crystalline Ibandronate polymorph which is characterized by an IR absorption
- 10 spectrum having characteristic peaks expressed in cm^{-1} at approximately 3678 cm^{-1} , 3164 cm^{-1} , 2854 cm^{-1} , 1377 cm^{-1} , 1288 cm^{-1} , 1157 cm^{-1} , 1094 cm^{-1} , 1069 cm^{-1} , 1035 cm^{-1} , 966 cm^{-1} , 951 cm^{-1} , 933 cm^{-1} , 903 cm^{-1} , 760 cm^{-1} and 723 cm^{-1} .
4. A crystalline polymorph of claim 3, characterized by the IR absorption spectrum shown in Figure 2.
- 15 5. A crystalline Ibandronate polymorph which is characterized by a characteristic peak expressed in cm^{-1} at approximately 1460 cm^{-1} in a vibrational Raman spectrum.
6. A crystalline polymorph of claim 5 which is characterized by the characteristic peaks expressed in cm^{-1} at approximately 2950 cm^{-1} , 2927 cm^{-1} , 2889 cm^{-1} , 2851 cm^{-1} , 1460 cm^{-1} , 1443 cm^{-1} , 1308 cm^{-1} , 1137 cm^{-1} , 1056 cm^{-1} , 1024 cm^{-1} , 954 cm^{-1} , 904 cm^{-1} , 839
- 20 cm^{-1} , 761 cm^{-1} and 678 cm^{-1} in a vibrational Raman spectrum.
7. A crystalline polymorph of claim 5 and 6, characterized by the vibrational Raman spectra shown in Figure 3.
8. Ibandronate containing at least 80% of a crystalline polymorph as characterized by claims 1 to 7.

9. A process for the preparation of a crystalline Ibandronate polymorph as characterized by the claims 1 to 8 comprising crystallizing a 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1,1-diphosphonic acid monosodium salt or a monohydrate, a polymorph or a polymorph mixture thereof at a temperature of 50°C to 70°C in a polar solvent.

10. Process according to claim 9, characterized in that the polar solvent is water.

11. Process according to claims 9 or 10, characterized in that acetone as a polar aprotic solvent is added for inducing the crystallization.

12. A process for the preparation of a crystalline Ibandronate polymorph as characterized by the claims 1 to 8 comprising tempering a moist 3-(*N*-methyl-*N*-pentyl) amino-1-hydroxypropane-1, 1-diphosphonic acid monosodium salt or a monohydrate, a polymorph or a polymorph mixture thereof at a temperature of 30°C to 90°C.

13. Pharmaceutical compositions comprising a crystalline polymorph according to any of claims 1 to 8 or obtained according to a process of claims 9 to 12 and a pharmaceutically acceptable carrier and/or adjuvant.

14. A crystalline polymorph according to any of claims 1 to 8 or obtained according to a process of claims 9 to 12 for use as therapeutic active substance.