

CLAIMS

1. A humanized antibody targeting c-Met or antigen-binding fragment thereof, comprising a heavy chain variable region and a light chain variable region, wherein:

5 (1) the heavy chain variable region has a sequence shown in SEQ ID NO: 7, or a sequence that has the same CDRs 1-3 as SEQ ID NO: 7 and an identity of greater than 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% to SEQ ID NO: 7; and/or

(2) the light chain variable region has a sequence shown in SEQ ID NO: 8, or a sequence that has the same CDRs 1-3 as SEQ ID NO: 8 and an identity of greater than 80%, 85%, 90%,
10 95%, 96%, 97%, 98% or 99% to SEQ ID NO: 8.

2. The antibody or antigen-binding fragment thereof according to claim 1, wherein the humanized antibody or antigen-binding fragment thereof is selected from the group consisting of monoclonal antibody, bispecific antibody, multispecific antibody, recombinant protein
15 comprising the antigen-binding fragment, Fab fragment, F(ab') fragment, F(ab')₂ fragment, Fv fragment, dAb, Fd, and single chain antibody (scFv).

3. The antibody or antigen-binding fragment thereof according to claim 1, further comprising a constant region of an immunoglobulin, wherein the immunoglobulin is selected
20 from the group consisting of IgG1, IgG2, IgG3 and IgG4.

4. The antibody or antigen-binding fragment thereof according to claim 1, wherein:

(1) the heavy chain variable region has an amino acid sequence shown in SEQ ID NO: 7;
and/or

25 (2) the light chain variable region has an amino acid sequence shown in SEQ ID NO: 8.

5. The antibody or antigen-binding fragment thereof according to claim 4, wherein the

humanized antibody targeting c-Met comprises a heavy chain and a light chain, and

(1) the heavy chain has an amino acid sequence shown in SEQ ID NO: 9; and/or

(2) the light chain has an amino acid sequence shown in SEQ ID NO:10.

5 6. An isolated polynucleotide encoding the antibody or antigen-binding fragment thereof according to claims 1-5.

7. A nucleic acid construct comprising the polynucleotide according to claim 6.

10 8. The nucleic acid construct according to claim 7, wherein the nucleic acid construct is an expression vector, and the polynucleotide is operably linked to a regulatory sequence that permits expression of a polypeptide encoded by the polynucleotide in a host cell or cell-free expression system.

15 9. A host cell comprising the polynucleotide according to claim 6 or the nucleic acid construct according to claim 7 or 8, wherein the host cell is preferably selected from the group consisting of a prokaryotic cell, eukaryotic cell, yeast cell, mammalian cell and E. coli cell; and more preferably, the host cell is selected from the group consisting of a CHO cell, NS0 cell, Sp2/0 cell and BHK cell.

20 10. A method for producing the antibody or antigen-binding fragment thereof according to any one of claims 1 to 5, comprising culturing the host cell according to claim 9 under a condition allowing the expression of the nucleic acid construct according to claim 8, and recovering the produced expressed protein from the culture.

25 11. An antibody-drug conjugate targeting c-Met, comprising the antibody or antigen-binding fragment thereof according to any one of claims 1 to 5 coupled with one or more therapeutic











agents.











12. The antibody-drug conjugate according to claim 11, wherein the therapeutic agent is selected from the group consisting of a cytotoxic molecule, immunoenhancer and radioisotope, the cytotoxic molecule includes but is not limited to a tubulin inhibitor or DNA damage agent; further preferably, the tubulin inhibitor includes but is not limited to a cytotoxic molecule of dolastatins and auristatins and a cytotoxic molecule of maytansines; the DNA damaging agent includes but is not limited to calicheamicins, duocarmycins, pyrrolbenzodiazepine derivatives (PBD), camptothecins and derivatives thereof, SN-38 and Dxd; more preferably, the cytotoxic molecule of auristatins includes but is not limited to MMAE, MMAF, and a derivative thereof, and the cytotoxic molecule of maytansines includes but is not limited to DM1, DM4, and a derivative thereof.


13. The antibody-drug conjugate according to claim 12, wherein the antibody-drug conjugate has a structural formula of $A-(L-U)_n$, wherein A represents the antibody or antigen-binding fragment thereof according to any one of claims 1 to 5; U is an active drug unit; L is any linking group, and L is covalently linked to A and U, respectively; n is an integer selected from 1, 2, 3, 4, 5, 6, 7 or 8; and A is linked to 1, 2, 3, 4, 5, 6, 7 or 8 of U through one or more L.

14. The antibody-drug conjugate according to claim 13, wherein L is covalently linked to the amino residue or thiol residue on A; preferably, L is covalently linked to the thiol residue on A; more preferably, L is covalently linked to the thiol residue formed after the interchain disulfide bond on A is opened.













15. The antibody-drug conjugate according to claim 14, wherein L comprises a cleavable linker and a non-cleavable linker.

















18. The antibody-drug conjugate according to claim 13 or claim 17, wherein U has a structure selected from the group consisting of:









19. The antibody-drug conjugate according to claim 13, wherein the antibody-drug conjugate has a structure selected from the group consisting of:




wherein, n in ADC-1 is an integer selected from 1, 2, 3, 4, 5, 6, 7 and 8; or




wherein, n in ADC-2 is an integer selected from 1, 2, 3 and 4; or




wherein, n in ADC-3 is an integer selected from 1, 2, 3 and 4; or




wherein, n in ADC-4 is an integer selected from 1 and 2; or



wherein, n in ADC-5 is an integer selected from 1, 2, 3, 4, 5, 6, 7 and 8; or



wherein, n in ADC-6 is an integer selected from 1, 2, 3, 4, 5, 6, 7 and 8; or



ADC-7

wherein, n in ADC-7 is an integer selected from 1, 2, 3, 4, 5, 6, 7 and 8.

20. A pharmaceutical composition comprising the antibody or antigen-binding fragment thereof according to any one of claims 1-5 and/or the antibody-drug conjugate according to any one of claims 11-19, and a pharmaceutically acceptable carrier.

21. Use of the antibody or antigen-binding fragment thereof according to any one of claims 1-5, the polynucleotide according to claim 6, the nucleic acid construct according to claim 7 or 8, the antibody-drug conjugate according to any one of claims 11-19 or the pharmaceutical composition according to claim 20 in the manufacture of a medicament for treating or preventing cancer.

22. The use according to claim 21, wherein the cancer is a c-Met positive cancer.

23. The use according to claim 22, wherein the c-Met positive cancer includes lung cancer and gastric cancer.

24. A kit comprising a container, a preparation placed in the container and an optional instruction; wherein the preparation comprises the antibody or antigen-binding fragment thereof according to any one of claims 1-5, the antibody-drug conjugate according to any one of claims 11-19 or a pharmaceutically acceptable salt or solvate thereof, or the pharmaceutical composition

according to claim 20.

25. Use of the antibody or antigen-binding fragment thereof according to any one of claims 1-5, the antibody-drug conjugate according to any one of claims 11-19, the polynucleotide according to claim 6, the nucleic acid construct according to claim 7 or 8, or the pharmaceutical composition according to claim 20 in the manufacture of a medicament for treating or preventing cancer.

26. The use according to claim 25, wherein the cancer is a solid tumor; further, the solid tumor includes lung cancer and gastric cancer.

27. A recombinant protein comprising the antibody or antigen-binding fragment according to any one of claims 1-5.

28. The recombinant protein according to claim 27, wherein the recombinant protein is a bispecific antibody or a multispecific antibody.

29. Use of the antibody or antigen-binding fragment according to any one of claims 1-5 in the manufacture of a recombinant protein.