
PYRIDAZINONES AND METHODS OF USE THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of priority to U.S. Provisional Patent Application No. 62/732,728, filed September 18, 2018; and U.S. Provisional Patent Application No. 62/780,553, filed December 17, 2018; each of which is incorporated by reference herein in its entirety.

BACKGROUND

Proteinuria is a condition in which an excessive amount of protein in the blood leaks into the urine. Proteinuria can progress from a loss of 30 mg of protein in the urine over a 24-hour period (called microalbuminuria) to >300 mg/day (called macroalbuminuria), before reaching levels of 3.5 grams of protein or more over a 24-hour period, or 25 times the normal amount. Proteinuria occurs when there is a malfunction in the kidney's glomeruli, causing fluid to accumulate in the body (edema). Prolonged protein leakage has been shown to result in kidney failure. Nephrotic Syndrome (NS) disease accounts for approximately 12% of prevalent end stage renal disease cases at an annual cost in the United States of more than \$3 billion. Approximately 5 out of every 100,000 children are diagnosed with NS every year and 15 out of every 100,000 children are living with it today. For patients who respond positively to treatment, the relapse frequency is extremely high. Ninety % of children with Nephrotic Syndrome will respond to treatment, however, an estimated 75% will relapse. There is a need for more effective methods of treating, or reducing risk of developing, kidney disease, e.g., proteinuria.

Mammalian TRP channel proteins form six-transmembrane cation-permeable channels that may be grouped into six subfamilies on the basis of amino acid sequence homology (TRPC, TRPV, TRPM, TRPA, TRPP, and TRPML). Recent studies of TRP channels indicate that they are involved in numerous fundamental cell functions and are considered to play an important role in the pathophysiology of many diseases. Many TRPs are expressed in kidney along different parts of the nephron and growing evidence suggest that these channels are involved in hereditary, as well as acquired kidney disorders. TRPC6, TRPM6, and TRPP2 have been implicated in hereditary focal

segmental glomerulosclerosis (FSGS), hypomagnesemia with secondary hypocalcemia (HSH), and polycystic kidney disease (PKD), respectively.


TRPC5 has also been reported to contribute to the mechanisms underlying regulation of innate fear responses. (J Neurosci. 2014 Mar 5; 34(10): 3653–3667).

Hence, there is a need for additional inhibitors of TRPC5 or TRPC4 or both.

SUMMARY

This invention is based, at least in part, on the discovery that Transient Receptor Potential Cation Channel, subfamily C, member 5 (TRPC5), activity abolishes actin stress fibers and diminishes focal adhesion formation, rendering a motile, migratory podocyte phenotype.

One aspect of the invention is compounds that are antagonists of TRPC5 or TRPC4 or both. In some embodiments, the compound of the invention is a compound of structural formula I:



(I), or a pharmaceutically acceptable salt thereof;

wherein:

“---” is a single bond or a double bond

X¹ is CH or N;

when “---” is a double bond, X² is CH or N;

when “---” is a single bond, X² is N(CH₃),

when X¹ is CH, X² is N or N(CH₃);

Y is -O-, -N(CH₃)-, -N(CH₂CH₂OH)-, cyclopropan-1,1-diyl, or -CH(CH₃)-

Q is 2-trifluoromethyl-4-fluorophenyl, 2-difluoromethyl-4-fluorophenyl, 2-trifluoromethylphenyl, 2-methyl-4-fluorophenyl, 2-chloro-4-fluorophenyl, 2-chlorophenyl, 1-(benzyl)-4-methylpiperidin-3-yl, 4-trifluoromethylpyridin-3-yl, 2-

trifluoromethyl-6-fluorophenyl, 2-trifluoromethyl-3-cyanophenyl, 2-ethyl-3-fluorophenyl, 2-chloro-3-cyanophenyl, 2-trifluoromethyl-5-fluorophenyl, or 2-difluoromethylphenyl;


R^3 is hydrogen, $-\text{CH}_2\text{OH}$, $-\text{CH}(\text{OH})-\text{CH}_2\text{OH}$, $-\text{NH}_2$, $-\text{CH}(\text{OH})\text{CH}_3$, $-\text{OCH}_3$, or $-\text{NH}-(\text{CH}_2)_2\text{OH}$; and when “---” is a double bond, R^4 is absent;

and when “---” is a single bond, R^3 and R^4 are taken together to form $=\text{O}$; and

each of R^5 and R^6 is independently hydrogen or $-\text{CH}_3$,

provided that if X^1 is N, X^2 is N, Y is $-\text{O}-$ or $-\text{N}(\text{CH}_3)-$, and Q is 2-trifluoromethylphenyl, then at least one of R^3 , R^5 , and R^6 is not hydrogen.

In some embodiments, the compound of the invention is represented by structural formula II:



(II), or a pharmaceutically acceptable salt thereof; wherein:

R^1 is chloro, $-\text{CF}_3$, $-\text{CHF}_2$, or $-\text{CH}_3$;

R^2 is hydrogen or fluoro; and

R^3 is hydrogen, $-\text{NH}_2$, $-\text{CH}_2\text{OH}$, or $\text{CH}(\text{OH})-\text{CH}_2\text{OH}$.

In one aspect, the invention features a pharmaceutical composition comprising a compound of the invention and a pharmaceutically acceptable carrier.

In one aspect, the invention relates to methods of treating, or reducing the risk of developing, a disease or condition selected from kidney disease, pulmonary arterial hypertension, anxiety, depression, cancer, diabetic retinopathy, or pain, comprising administering to a subject in need thereof a therapeutically effective amount of the compound or the composition. In some embodiments, the disease is kidney disease, anxiety, depression, cancer, or diabetic retinopathy. In some embodiments, the disease or

condition is kidney disease selected from Focal Segmental Glomerulosclerosis (FSGS), Diabetic nephropathy, Alport syndrome, hypertensive kidney disease, nephrotic syndrome, steroid-resistant nephrotic syndrome, minimal change disease, membranous nephropathy, idiopathic membranous nephropathy, membranoproliferative glomerulonephritis (MPGN), immune complex-mediated MPGN, complement-mediated MPGN, Lupus nephritis, postinfectious glomerulonephritis, thin basement membrane disease, mesangial proliferative glomerulonephritis, amyloidosis (primary), c1q nephropathy, rapidly progressive GN, anti-GBM disease, C3 glomerulonephritis, hypertensive nephrosclerosis, or IgA nephropathy. In some embodiments, the kidney disease is proteinuric kidney disease. In some embodiments, the kidney disease is microalbuminuria or macroalbuminuria kidney disease. In some embodiments, the disease or condition to be treated is pulmonary arterial hypertension. In some embodiments, the disease or condition to be treated is pain selected from neuropathic pain and visceral pain.

In some embodiments, the disease or condition is cancer selected from chemoresistant breast carcinoma, adriamycin-resistant breast cancer, chemoresistant colorectal cancer, medulloblastoma, and tumor angiogenesis.

In some embodiments, the disease or condition to be treated is transplant-related FSGS, transplant-related nephrotic syndrome, transplant-related proteinuria, cholestatic liver disease, polycystic kidney disease, autosomal dominant polycystic kidney disease (ADPKD), obesity, insulin resistance, Type II diabetes, prediabetes, metabolic syndrome, non-alcoholic fatty liver disease (NAFLD), or non-alcoholic steatohepatitis (NASH).

The methods are effective for a variety of subjects including mammals, e.g., humans and other animals, such as laboratory animals, e.g., mice, rats, rabbits, or monkeys, or domesticated and farm animals, e.g., cats, dogs, goats, sheep, pigs, cows, or horses.

The invention provides several advantages. The prophylactic and therapeutic methods described herein are effective in treating kidney disease, e.g., proteinuria, and have minimal, if any, side effects. Further, methods described herein are effective to

identify compounds that treat or reduce risk of developing a kidney disease, anxiety, depression, or cancer.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features, objects, and advantages of the invention will be apparent from the detailed description, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows albumin excretion in PAN-injured rats treated with compound 100 or mizoribine.

Figure 2 shows urine protein creatinine ratio data in AT1R transgenic rats treated with compound 100 compared to vehicle, with AngII infusion.

Figure 3 shows the urine protein creatinine ratio data presented in Figure 2 expressed as percent of baseline.

DETAILED DESCRIPTION

Definitions

The term “acyl” is art-recognized and refers to a group represented by the general formula hydrocarbylC(O)-, preferably alkylC(O)-.

The term "acylamino" is art-recognized and refers to an amino group substituted with an acyl group and may be represented, for example, by the formula hydrocarbylC(O)NH-

The term "acyloxy" is art-recognized and refers to a group represented by the general formula hydrocarbylC(O)O-, preferably alkylC(O)O-.

The term "alkoxy" refers to an alkyl group, preferably a lower alkyl group, having an oxygen attached thereto. Representative alkoxy groups include methoxy, trifluoromethoxy, ethoxy, propoxy, tert-butoxy and the like.

The term "alkoxyalkyl" refers to an alkyl group substituted with an alkoxy group and may be represented by the general formula alkyl-O-alkyl.

The term "alkenyl", as used herein, refers to an aliphatic group containing at least one double bond and is intended to include both "unsubstituted alkenyls" and "substituted alkenyls", the latter of which refers to alkenyl moieties having substituents replacing a hydrogen on one or more carbons of the alkenyl group. Such substituents may occur on one or more carbons that are included or not included in one or more double bonds. Moreover, such substituents include all those contemplated for alkyl groups, as discussed below, except where stability is prohibitive. For example, substitution of alkenyl groups by one or more alkyl, carbocyclyl, aryl, heterocyclyl, or heteroaryl groups is contemplated.

An "alkyl" group or "alkane" is a straight chained or branched non-aromatic hydrocarbon which is completely saturated. Typically, a straight chained or branched alkyl group has from 1 to about 20 carbon atoms, preferably from 1 to about 10 unless otherwise defined. Examples of straight chained and branched alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, pentyl, hexyl, pentyl and octyl. A C₁-C₆ straight chained or branched alkyl group is also referred to as a "lower alkyl" group.

Moreover, the term "alkyl" (or "lower alkyl") as used throughout the specification, examples, and claims is intended to include both "unsubstituted alkyls" and "substituted alkyls", the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents, if not otherwise specified, can include, for example, a halogen (e.g., fluoro),

a hydroxyl, a carbonyl (such as a carboxyl, an alkoxycarbonyl, a formyl, or an acyl), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an alkoxy, a phosphoryl, a phosphate, a phosphonate, a phosphinate, an amino, an amido, an amidine, an imine, a cyano, a nitro, an azido, a sulfhydryl, an alkylthio, a sulfate, a sulfonate, a sulfamoyl, a sulfonamido, a sulfonyl, a heterocyclyl, an aralkyl, or an aromatic or heteroaromatic moiety. In preferred embodiments, the substituents on substituted alkyls are selected from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, halogen, carbonyl, cyano, or hydroxyl. In more preferred embodiments, the substituents on substituted alkyls are selected from fluoro, carbonyl, cyano, or hydroxyl. It will be understood by those skilled in the art that the moieties substituted on the hydrocarbon chain can themselves be substituted, if appropriate. For instance, the substituents of a substituted alkyl may include substituted and unsubstituted forms of amino, azido, imino, amido, phosphoryl (including phosphonate and phosphinate), sulfonyl (including sulfate, sulfonamido, sulfamoyl and sulfonate), and silyl groups, as well as ethers, alkylthios, carbonyls (including ketones, aldehydes, carboxylates, and esters), -CF₃, -CN and the like. Exemplary substituted alkyls are described below. Cycloalkyls can be further substituted with alkyls, alkenyls, alkoxy, alkylthios, aminoalkyls, carbonyl-substituted alkyls, -CF₃, -CN, and the like.

Unless otherwise specified, “alkylene” by itself or as part of another substituent refers to a saturated straight-chain or branched divalent group having the stated number of carbon atoms and derived from the removal of two hydrogen atoms from the corresponding alkane. Examples of straight chained and branched alkylene groups include -CH₂- (methylene), -CH₂-CH₂- (ethylene), -CH₂-CH₂-CH₂- (propylene), -C(CH₃)₂-, -CH₂-CH(CH₃)-, -CH₂-CH₂-CH₂-CH₂-, -CH₂-CH₂-CH₂-CH₂-CH₂- (pentylene), -CH₂-CH(CH₃)-CH₂-, and -CH₂-C(CH₃)₂-CH₂-.

The term “C_{x-y}” when used in conjunction with a chemical moiety, such as, acyl, acyloxy, alkyl, alkenyl, alkynyl, or alkoxy is meant to include groups that contain from x to y carbons in the chain. For example, the term “C_{x-y} alkyl” refers to substituted or unsubstituted saturated hydrocarbon groups, including straight-chain alkyl and branched-chain alkyl groups that contain from x to y carbons in the chain, including haloalkyl groups. Preferred haloalkyl groups include trifluoromethyl, difluoromethyl, 2,2,2-trifluoroethyl, and pentafluoroethyl. C₀ alkyl indicates a hydrogen where the group is in


a terminal position, a bond if internal. The terms “C_{2-y} alkenyl” and “C_{2-y} alkynyl” refer to substituted or unsubstituted unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively.

The term “alkylamino”, as used herein, refers to an amino group substituted with at least one alkyl group.

The term “alkylthio”, as used herein, refers to a thiol group substituted with an alkyl group and may be represented by the general formula alkylS-


The term “alkynyl”, as used herein, refers to an aliphatic group containing at least one triple bond and is intended to include both "unsubstituted alkynyls" and "substituted alkynyls", the latter of which refers to alkynyl moieties having substituents replacing a hydrogen on one or more carbons of the alkynyl group. Such substituents may occur on one or more carbons that are included or not included in one or more triple bonds. Moreover, such substituents include all those contemplated for alkyl groups, as discussed above, except where stability is prohibitive. For example, substitution of alkynyl groups by one or more alkyl, carbocyclyl, aryl, heterocyclyl, or heteroaryl groups is contemplated.

The term “amide”, as used herein, refers to a group



wherein each R^A independently represent a hydrogen or hydrocarbyl group, or two R^A are taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure.

The terms “amine” and “amino” are art-recognized and refer to both unsubstituted and substituted amines and salts thereof, e.g., a moiety that can be represented by




wherein each R^A independently represents a hydrogen or a hydrocarbonyl group, or two R^A are taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure.

The term “aminoalkyl”, as used herein, refers to an alkyl group substituted with an amino group.

The term “aralkyl”, as used herein, refers to an alkyl group substituted with an aryl group.

The term “aryl” as used herein include substituted or unsubstituted single-ring aromatic groups in which each atom of the ring is carbon. Preferably the ring is a 6- or 10-membered ring, more preferably a 6-membered ring. The term “aryl” also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings wherein at least one of the rings is aromatic, e.g., the other cyclic rings can be cycloalkyls, cycloalkenyls, aryls, heteroaryls, and/or heterocyclyls. Aryl groups include benzene, naphthalene, phenanthrene, phenol, aniline, and the like.

The term “carbamate” is art-recognized and refers to a group



wherein each R^A independently represent hydrogen or a hydrocarbonyl group, such as an alkyl group, or both R^A taken together with the intervening atom(s) complete a heterocycle having from 4 to 8 atoms in the ring structure.

The terms “carbocycle”, and “carbocyclic”, as used herein, refers to a saturated or unsaturated ring in which each atom of the ring is carbon. The term carbocycle includes both aromatic carbocycles and non-aromatic carbocycles. Non-aromatic carbocycles include both cycloalkane rings, in which all carbon atoms are saturated, and cycloalkene rings, which contain at least one double bond. “Carbocycle” includes 5-7 membered monocyclic and 8-12 membered bicyclic rings. Each ring of a bicyclic carbocycle may be selected from saturated, unsaturated and aromatic rings. Carbocycle includes bicyclic molecules in which one, two or three or more atoms are shared between the two rings. The term “fused carbocycle” refers to a bicyclic carbocycle in which each of the rings

shares two adjacent atoms with the other ring. Each ring of a fused carbocycle may be selected from saturated, unsaturated and aromatic rings. In an exemplary embodiment, an aromatic ring, e.g., phenyl, may be fused to a saturated or unsaturated ring, e.g., cyclohexane, cyclopentane, or cyclohexene. Any combination of saturated, unsaturated and aromatic bicyclic rings, as valence permits, is included in the definition of carbocyclic. Exemplary “carbocycles” include cyclopentane, cyclohexane, bicyclo[2.2.1]heptane, 1,5-cyclooctadiene, 1,2,3,4-tetrahydronaphthalene, bicyclo[4.2.0]oct-3-ene, naphthalene and adamantane. Exemplary fused carbocycles include decalin, naphthalene, 1,2,3,4-tetrahydronaphthalene, bicyclo[4.2.0]octane, 4,5,6,7-tetrahydro-1H-indene and bicyclo[4.1.0]hept-3-ene. “Carbocycles” may be substituted at any one or more positions capable of bearing a hydrogen atom.

A “cycloalkyl” group is a cyclic hydrocarbon which is completely saturated. “Cycloalkyl” includes monocyclic and bicyclic rings. Typically, a monocyclic cycloalkyl group has from 3 to about 10 carbon atoms, more typically 3 to 8 carbon atoms unless otherwise defined. The second ring of a bicyclic cycloalkyl may be selected from saturated, unsaturated and aromatic rings. Cycloalkyl includes bicyclic molecules in which one, two or three or more atoms are shared between the two rings. The term “fused cycloalkyl” refers to a bicyclic cycloalkyl in which each of the rings shares two adjacent atoms with the other ring. The second ring of a fused bicyclic cycloalkyl may be selected from saturated, unsaturated and aromatic rings. A “cycloalkenyl” group is a cyclic hydrocarbon containing one or more double bonds.

The term “carbocyclalkyl”, as used herein, refers to an alkyl group substituted with a carbocycle group.

The term “carbonate” is art-recognized and refers to a group $-\text{OCO}_2\text{-R}^{\text{A}}$, wherein R^{A} represents a hydrocarbyl group.

The term “carboxy”, as used herein, refers to a group represented by the formula $-\text{CO}_2\text{H}$.

The term “ester”, as used herein, refers to a group $-\text{C}(\text{O})\text{OR}^{\text{A}}$ wherein R^{A} represents a hydrocarbyl group.

The term “ether”, as used herein, refers to a hydrocarbyl group linked through an oxygen to another hydrocarbyl group. Accordingly, an ether substituent of a hydrocarbyl

group may be hydrocarbyl-O-. Ethers may be either symmetrical or unsymmetrical. Examples of ethers include, but are not limited to, heterocycle-O-heterocycle and aryl-O-heterocycle. Ethers include “alkoxyalkyl” groups, which may be represented by the general formula alkyl-O-alkyl.

The terms “halo” and “halogen” as used herein means halogen and includes chloro, fluoro, bromo, and iodo.

The terms “hetaralkyl” and “heteroaralkyl”, as used herein, refers to an alkyl group substituted with a hetaryl group.

The term “heteroalkyl”, as used herein, refers to a saturated or unsaturated chain of carbon atoms and at least one heteroatom, wherein no two heteroatoms are adjacent.

The terms “heteroaryl” and “hetaryl” include substituted or unsubstituted aromatic single ring structures, preferably 5- to 7-membered rings, more preferably 5- to 6-membered rings, whose ring structures include at least one heteroatom, preferably one to four heteroatoms, more preferably one or two heteroatoms. The terms “heteroaryl” and “hetaryl” also include polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings wherein at least one of the rings is heteroaromatic, e.g., the other cyclic rings can be cycloalkyls, cycloalkenyls, aryls, heteroaryls, and/or heterocyclyls. Heteroaryl groups include, for example, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, pyrazole, pyridine, pyrazine, pyridazine, and pyrimidine, and the like.

The term “heteroatom” as used herein means an atom of any element other than carbon or hydrogen. Preferred heteroatoms are nitrogen, oxygen, and sulfur.

The terms “heterocyclyl”, “heterocycle”, and “heterocyclic” refer to substituted or unsubstituted non-aromatic ring structures, preferably 3- to 10-membered rings, more preferably 3- to 7-membered rings, whose ring structures include at least one heteroatom, preferably one to four heteroatoms, more preferably one or two heteroatoms. The terms “heterocyclyl” and “heterocyclic” also include polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings wherein at least one of the rings is heterocyclic, e.g., the other cyclic rings can be cycloalkyls, cycloalkenyls, aryls, heteroaryls, and/or heterocyclyls. Heterocyclyl groups

include, for example, piperidine, piperazine, pyrrolidine, tetrahydropyran, tetrahydrofuran, morpholine, lactones, lactams, and the like.

The term “heterocyclalkyl” or “heterocycloalkyl”, as used herein, refers to an alkyl group substituted with a heterocycle group.

The term “hydrocarbyl”, as used herein, refers to a group that is bonded through a carbon atom that does not have a =O or =S substituent, and typically has at least one carbon-hydrogen bond and a primarily carbon backbone, but may optionally include heteroatoms. Thus, groups like methyl, ethoxyethyl, 2-pyridyl, and trifluoromethyl are considered to be hydrocarbyl for the purposes of this application, but substituents such as acetyl (which has a =O substituent on the linking carbon) and ethoxy (which is linked through oxygen, not carbon) are not. Hydrocarbyl groups include, but are not limited to aryl, heteroaryl, carbocycle, heterocyclyl, alkyl, alkenyl, alkynyl, and combinations thereof.

The term “hydroxyalkyl”, as used herein, refers to an alkyl group substituted with a hydroxy group.

The term “lower” when used in conjunction with a chemical moiety, such as, acyl, acyloxy, alkyl, alkenyl, alkynyl, or alkoxy is meant to include groups where there are ten or fewer non-hydrogen atoms in the substituent, preferably six or fewer. A “lower alkyl”, for example, refers to an alkyl group that contains ten or fewer carbon atoms, preferably six or fewer. In certain embodiments, acyl, acyloxy, alkyl, alkenyl, alkynyl, or alkoxy substituents defined herein are respectively lower acyl, lower acyloxy, lower alkyl, lower alkenyl, lower alkynyl, or lower alkoxy, whether they appear alone or in combination with other substituents, such as in the recitations hydroxyalkyl and aralkyl (in which case, for example, the atoms within the aryl group are not counted when counting the carbon atoms in the alkyl substituent).


The terms “polycyclyl”, “polycycle”, and “polycyclic” refer to two or more rings (e.g., cycloalkyls, cycloalkenyls, aryls, heteroaryls, and/or heterocyclyls) in which two or more atoms are common to two adjoining rings, e.g., the rings are “fused rings”. Each of the rings of the polycycle can be substituted or unsubstituted. In certain embodiments, each ring of the polycycle contains from 3 to 10 atoms in the ring, preferably from 5 to 7.

The term “silyl” refers to a silicon moiety with three hydrocarbyl moieties attached thereto.

The term “substituted” refers to moieties having substituents replacing a hydrogen on one or more carbons of the backbone. It will be understood that “substitution” or “substituted with” includes the implicit proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, e.g., which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc. As used herein, the term “substituted” is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and non-aromatic substituents of organic compounds. The permissible substituents can be one or more and the same or different for appropriate organic compounds. For purposes of this invention, the heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valences of the heteroatoms. Substituents can include any substituents described herein, for example, a halogen, a hydroxyl, a carbonyl (such as a carboxyl, an alkoxycarbonyl, a formyl, or an acyl), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an alkoxy, a phosphoryl, a phosphate, a phosphonate, a phosphinate, an amino, an amido, an amidine, an imine, a cyano, a nitro, an azido, a sulfhydryl, an alkylthio, a sulfate, a sulfonate, a sulfamoyl, a sulfonamido, a sulfonyl, a heterocyclyl, an aralkyl, or an aromatic or heteroaromatic moiety. In preferred embodiments, the substituents on substituted alkyls are selected from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, halogen, carbonyl, cyano, or hydroxyl. In more preferred embodiments, the substituents on substituted alkyls are selected from fluoro, carbonyl, cyano, or hydroxyl. It will be understood by those skilled in the art that substituents can themselves be substituted, if appropriate. Unless specifically stated as “unsubstituted,” references to chemical moieties herein are understood to include substituted variants. For example, reference to an “aryl” group or moiety implicitly includes both substituted and unsubstituted variants.

The term “sulfate” is art-recognized and refers to the group -OSO₃H, or a pharmaceutically acceptable salt thereof.

The term “sulfonamide” is art-recognized and refers to the group represented by the general formulae



wherein each R^{A} independently represents hydrogen or hydrocarbyl, such as alkyl, or both R^{A} taken together with the intervening atom(s) complete a heterocycle having from 4 to 8 atoms in the ring structure.

The term “sulfoxide” is art-recognized and refers to the group $-\text{S}(\text{O})-\text{R}^{\text{A}}$, wherein R^{A} represents a hydrocarbyl.

The term “sulfonate” is art-recognized and refers to the group SO_3H , or a pharmaceutically acceptable salt thereof.


The term “sulfone” is art-recognized and refers to the group $-\text{S}(\text{O})_2-\text{R}^{\text{A}}$, wherein R^{A} represents a hydrocarbyl.

The term “thioalkyl”, as used herein, refers to an alkyl group substituted with a thiol group.

The term “thioester”, as used herein, refers to a group $-\text{C}(\text{O})\text{SR}^{\text{A}}$ or $-\text{SC}(\text{O})\text{R}^{\text{A}}$ wherein R^{A} represents a hydrocarbyl.

The term “thioether”, as used herein, is equivalent to an ether, wherein the oxygen is replaced with a sulfur.

The term “urea” is art-recognized and may be represented by the general formula



wherein each R^{A} independently represents hydrogen or a hydrocarbyl, such as alkyl, or any occurrence of R^{A} taken together with another and the intervening atom(s) complete a heterocycle having from 4 to 8 atoms in the ring structure.

“Protecting group” refers to a group of atoms that, when attached to a reactive functional group in a molecule, mask, reduce or prevent the reactivity of the functional group. Typically, a protecting group may be selectively removed as desired during the course of a synthesis. Examples of protecting groups can be found in Greene and Wuts, *Protective Groups in Organic Chemistry*, 3rd Ed., 1999, John Wiley & Sons, NY and

Harrison et al., *Compendium of Synthetic Organic Methods*, Vols. 1-8, 1971-1996, John Wiley & Sons, NY. Representative nitrogen protecting groups include, but are not limited to, formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl ("CBZ"), tert-butoxycarbonyl ("Boc"), trimethylsilyl ("TMS"), 2-trimethylsilyl-ethanesulfonyl ("TES"), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl ("Fmoc"), nitro-veratryloxycarbonyl ("NVOC") and the like. Representative hydroxyl protecting groups include, but are not limited to, those where the hydroxyl group is either acylated (esterified) or alkylated such as benzyl and trityl ethers, as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers (e.g., TMS or TIPS groups), glycol ethers, such as ethylene glycol and propylene glycol derivatives and allyl ethers.

As used herein, a therapeutic that "prevents" or "reduces the risk of developing" a disease, disorder, or condition refers to a compound that, in a statistical sample, reduces the occurrence of the disease, disorder, or condition in the treated sample relative to an untreated control sample, or delays the onset or reduces the severity of one or more symptoms of the disorder or condition relative to the untreated control sample.

The term "treating" includes prophylactic and/or therapeutic treatments. The term "prophylactic or therapeutic" treatment is art-recognized and includes administration to the host of one or more of the subject compositions. If it is administered prior to clinical manifestation of the unwanted condition (e.g., disease or other unwanted state of the host animal) then the treatment is prophylactic (i.e., it protects the host against developing the unwanted condition), whereas if it is administered after manifestation of the unwanted condition, the treatment is therapeutic, (i.e., it is intended to diminish, ameliorate, or stabilize the existing unwanted condition or side effects thereof).

The phrases "conjoint administration" and "administered conjointly" refer to any form of administration of two or more different therapeutic compounds such that the second compound is administered while the previously administered therapeutic compound is still effective in the body (e.g., the two compounds are simultaneously effective in the patient, which may include synergistic effects of the two compounds). For example, the different therapeutic compounds can be administered either in the same formulation or in a separate formulation, either concomitantly or sequentially. In certain

embodiments, the different therapeutic compounds can be administered within one hour, 12 hours, 24 hours, 36 hours, 48 hours, 72 hours, or a week of one another. Thus, an individual who receives such treatment can benefit from a combined effect of different therapeutic compounds.

The term “prodrug” is intended to encompass compounds which, under physiologic conditions, are converted into the therapeutically active agents of the present invention. A common method for making a prodrug is to include one or more selected moieties which are hydrolyzed under physiologic conditions to reveal the desired molecule. In other embodiments, the prodrug is converted by an enzymatic activity of the host animal. For example, esters or carbonates (e.g., esters or carbonates of alcohols or carboxylic acids) are preferred prodrugs of the present invention. In certain embodiments, some or all of the compounds of the invention in a formulation represented above can be replaced with the corresponding suitable prodrug, e.g., wherein a hydroxyl in the parent compound is presented as an ester or a carbonate or carboxylic acid present in the parent compound is presented as an ester.

As used herein, “small molecules” refers to small organic or inorganic molecules of molecular weight below about 3,000 Daltons. In general, small molecules useful for the invention have a molecular weight of less than 3,000 Daltons (Da). The small molecules can be, e.g., from at least about 100 Da to about 3,000 Da (e.g., between about 100 to about 3,000 Da, about 100 to about 2500 Da, about 100 to about 2,000 Da, about 100 to about 1,750 Da, about 100 to about 1,500 Da, about 100 to about 1,250 Da, about 100 to about 1,000 Da, about 100 to about 750 Da, about 100 to about 500 Da, about 200 to about 1500, about 500 to about 1000, about 300 to about 1000 Da, or about 100 to about 250 Da).


In some embodiments, a “small molecule” refers to an organic, inorganic, or organometallic compound typically having a molecular weight of less than about 1000. In some embodiments, a small molecule is an organic compound, with a size on the order of 1 nm. In some embodiments, small molecule drugs of the invention encompass oligopeptides and other biomolecules having a molecular weight of less than about 1000.

An “effective amount” is an amount sufficient to effect beneficial or desired results. For example, a therapeutic amount is one that achieves the desired therapeutic effect. This amount can be the same or different from a prophylactically effective amount, which is an amount necessary to prevent onset of disease or disease symptoms. An effective amount can be administered in one or more administrations, applications or dosages. A therapeutically effective amount of a composition depends on the composition selected. The compositions can be administered from one or more times per day to one or more times per week; including once every other day. The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of the compositions described herein can include a single treatment or a series of treatments.

Compounds of the Invention

One aspect of the invention provides small molecule inhibitors of TRPC5.

In some embodiments, the compound of the invention is a compound of structural formula I:



(I), or a pharmaceutically acceptable salt thereof;

wherein:

“---” is a single bond or a double bond

X¹ is CH or N;

when “---” is a double bond, X² is CH or N;

when “---” is a single bond, X² is N(CH₃),

when X¹ is CH, X² is N or N(CH₃);

Y is -O-, -N(CH₃)-, -N(CH₂CH₂OH)-, cyclopropan-1,1-diyl, or -CH(CH₃)-;

Q is 2-trifluoromethyl-4-fluorophenyl, 2-difluoromethyl-4-fluorophenyl, 2-trifluoromethylphenyl, 2-methyl-4-fluorophenyl, 2-chloro-4-fluorophenyl, 2-chlorophenyl, 1-(benzyl)-4-methylpiperidin-3-yl, 4-trifluoromethylpyridin-3-yl, 2-trifluoromethyl-6-fluorophenyl, 2-trifluoromethyl-3-cyanophenyl, 2-ethyl-3-fluorophenyl, 2-chloro-3-cyanophenyl, 2-trifluoromethyl-5-fluorophenyl, or 2-difluoromethylphenyl;


R³ is hydrogen, -CH₂OH, -CH(OH)-CH₂OH, -NH₂, -CH(OH)CH₃, -OCH₃, or -NH-(CH₂)₂OH; and when “---” is a double bond, R⁴ is absent;

and when “---” is a single bond, R³ and R⁴ are taken together to form =O; and

each of R⁵ and R⁶ is independently hydrogen or -CH₃,

provided that if X¹ is N, X² is N, Y is -O- or -N(CH₃)-, and Q is 2-trifluoromethylphenyl, then at least one of R³, R⁵, and R⁶ is not hydrogen.

In some embodiments, the compound of the invention is a compound represented by structural formula II:



(II), or a pharmaceutically acceptable salt thereof; wherein:

R¹ is chloro, -CF₃, -CHF₂, or -CH₃;







R² is hydrogen or fluoro; and







R³ is hydrogen, -NH₂, -CH₂OH, or CH(OH)-CH₂OH.

In some embodiments, when R¹ is -CHF₂, R² is not hydrogen.

In some embodiments, the compound of the invention is selected from any one of the following compounds, or a pharmaceutically acceptable salt thereof:

Compound	Structure
100	 <chem>ClC1=CN=C(N1C2=NC=NC=C2Oc3ccc(cc3)C(F)(F)F)C(=O)NN</chem>
101	 <chem>ClC1=CN=C(N1C2=NC=NC=C2Oc3ccc(cc3)Cl)C(=O)NN</chem>
102	 <chem>ClC1=CN=C(N1C2=NC=NC=C2Oc3ccc(cc3)C(F)(F)F)C(=O)NN</chem>
103	 <chem>ClC1=CN=C(N1C2=NC=NC=C2Oc3ccc(cc3)C(F)(F)F)C(=O)NN</chem>
104	 <chem>ClC1=CN=C(N1C2=NC=NC=C2Oc3ccc(cc3)Cl)C(=O)NN</chem>
105	 <chem>ClC1=CN=C(N1C2=NC=NC=C2Oc3ccc(cc3)C(F)(F)F)C(=O)NN</chem>

Compound	Structure
106	
107	
108	
109	
110	
111	

Compound	Structure
112	
113	
114	
115	
116	
117	

Compound	Structure
117a	
118	
119	
120	
121	
122	

Compound	Structure
123	
124	
125	
126	
126a	
127	

Compound	Structure
128	
129	
130	
131	
132	
133	

Compound	Structure
133a	
134	
135	
136	
137	
138	







Compound	Structure
139	
140	







In some embodiments, the compound of the invention is selected from any one of the following compounds, or a pharmaceutically acceptable salt thereof:

Compound	Structure	Compound	Structure
100		104	
101		105	
102		112	

Compound	Structure	Compound	Structure
113		128	
114		134	
116		135	
124		137	
125			

In some embodiments, the compound of the invention is selected from any one of the following compounds, or a pharmaceutically acceptable salt thereof:

Compound	Structure
100	
101	
102	
104	
105	
114	

Compound	Structure
116	
124	
125	
128	
134	
135	

Compound	Structure
137	

In certain embodiments, the compounds of the invention may be racemic. In certain embodiments, the compounds of the invention may be enriched in one enantiomer. For example, a compound of the invention may have greater than 30% ee, 40% ee, 50% ee, 60% ee, 70% ee, 80% ee, 90% ee, or even 95% or greater ee.

The compounds of the invention have more than one stereocenter. Accordingly, the compounds of the invention may be enriched in one or more diastereomers. For example, a compound of the invention may have greater than 30% de, 40% de, 50% de, 60% de, 70% de, 80% de, 90% de, or even 95% or greater de. In certain embodiments, the compounds of the invention have substantially one isomeric configuration at one or more stereogenic centers, and have multiple isomeric configurations at the remaining stereogenic centers.

In certain embodiments, the enantiomeric excess of the stereocenter is at least 40% ee, 50% ee, 60% ee, 70% ee, 80% ee, 90% ee, 92% ee, 94% ee, 95% ee, 96% ee, 98% ee or greater ee.

As used herein, single bonds drawn without stereochemistry do not indicate the stereochemistry of the compound.

As used herein, hashed or bolded non-wedge bonds indicate relative, but not absolute, stereochemical configuration (e.g., do not distinguish between enantiomers of a given diastereomer).

As used herein, hashed or bolded wedge bonds indicate absolute stereochemical configuration.

In some embodiments, the invention relates to pharmaceutical composition comprising a compound of the invention and a pharmaceutically acceptable carrier. In certain embodiments, a therapeutic preparation or pharmaceutical composition of the compound of the invention may be enriched to provide predominantly one enantiomer of a compound. An enantiomerically

enriched mixture may comprise, for example, at least 60 mol percent of one enantiomer, or more preferably at least 75, 90, 95, or even 99 mol percent. In certain embodiments, the compound enriched in one enantiomer is substantially free of the other enantiomer, wherein substantially free means that the substance in question makes up less than 10%, or less than 5%, or less than 4%, or less than 3%, or less than 2%, or less than 1% as compared to the amount of the other enantiomer, *e.g.*, in the composition or compound mixture. For example, if a composition or compound mixture contains 98 grams of a first enantiomer and 2 grams of a second enantiomer, it would be said to contain 98 mol percent of the first enantiomer and only 2% of the second enantiomer.

In certain embodiments, a therapeutic preparation or pharmaceutical composition may be enriched to provide predominantly one diastereomer of the compound of the invention. A diastereomerically enriched mixture may comprise, for example, at least 60 mol percent of one diastereomer, or more preferably at least 75, 90, 95, or even 99 mol percent.

Methods of Treatment

The non-selective Ca^{2+} -permeable Transient Receptor Potential (TRP) channels act as sensors that transduce extracellular cues to the intracellular environment in diverse cellular processes, including actin remodeling and cell migration (Greka et al., *Nat Neurosci* 6, 837-845, 2003; Ramsey et al., *Annu Rev Physiol* 68, 619-647, 2006; Montell, *Pflugers Arch* 451, 19-28, 2005; Clapham, *Nature* 426, 517-524, 2003). Dynamic rearrangement of the actin cytoskeleton relies on spatiotemporally regulated Ca^{2+} influx (Zheng and Poo, *Annu Rev Cell Dev Biol* 23, 375-404, 2007); Brandman and Meyer, *Science* 322, 390-395, 2008); Collins and Meyer, *Dev Cell* 16, 160-161, 2009) and the small GTPases RhoA and Rac1 serve as key modulators of these changes (Etienne-Manneville and Hall, *Nature* 420, 629-635, 2002); Raftopoulou and Hall, *Dev Biol* 265, 23-32, 2004). RhoA induces stress fiber and focal adhesion formation, while Rac1 mediates lamellipodia formation (Etienne-Manneville and Hall, *Nature* 420, 629-635, 2002). The Transient Receptor Potential Cation Channel, subfamily C, member 5 (TRPC5) acts in concert with TRPC6 to regulate Ca^{2+} influx, actin remodeling, and cell motility in kidney podocytes and fibroblasts. TRPC5-mediated Ca^{2+} influx increases Rac1 activity, whereas TRPC6-mediated Ca^{2+} influx promotes RhoA activity. Gene silencing of TRPC6 channels abolishes stress fibers and diminishes focal contacts, rendering a motile, migratory cell phenotype. In contrast, gene

silencing of TRPC5 channels rescues stress fiber formation, rendering a contractile cell phenotype. The results described herein unveil a conserved signaling mechanism whereby TRPC5 and TRPC6 channels control a tightly regulated balance of cytoskeletal dynamics through differential coupling to Rac1 and RhoA.

Ca²⁺-dependent remodeling of the actin cytoskeleton is a dynamic process that drives cell migration (Wei et al., *Nature* 457, 901-905, 2009). RhoA and Rac1 act as switches responsible for cytoskeletal rearrangements in migrating cells (Etienne-Manneville and Hall, *Nature* 420, 629-635, 2002); Raftopoulou and Hall, *Dev Biol* 265, 23-32, 2004). Activation of Rac1 mediates a motile cell phenotype, whereas RhoA activity promotes a contractile phenotype (Etienne-Manneville and Hall, *Nature* 420, 629-635, 2002). Ca²⁺ plays a central role in small GTPase regulation (Aspenstrom et al., *Biochem J* 377, 327-337, 2004). Spatially and temporally restricted flickers of Ca²⁺ are enriched near the leading edge of migrating cells (Wei et al., *Nature* 457, 901-905, 2009). Ca²⁺-microdomains have thus joined local bursts in Rac1 activity (Gardiner et al., *Curr Biol* 12, 2029-2034, 2002; Machacek et al., *Nature* 461, 99-103, 2009) as critical events at the leading edge. To date, the sources of Ca²⁺-influx responsible for GTPase regulation remain largely elusive. TRP (Transient Receptor Potential) channels generate time and space-limited Ca²⁺ signals linked to cell migration in fibroblasts and neuronal growth cones⁰. Specifically, TRPC5 channels are known regulators of neuronal growth cone guidance¹ and their activity in neurons is dependent on PI3K and Rac1 activity (Bezzarides et al., *Nat Cell Biol* 6, 709-720, 2004).

Podocytes are neuronal-like cells that originate from the metanephric mesenchyme of the kidney glomerulus and are essential to the formation of the kidney filtration apparatus (Somlo and Mundel, *Nat Genet.* 24, 333-335, 2000; Fukasawa et al., *J Am Soc Nephrol* 20, 1491-1503, 2009). Podocytes possess an exquisitely refined repertoire of cytoskeletal adaptations to environmental cues (Somlo and Mundel, *Nat Genet* 24, 333-335, 2000; Garg et al., *Mol Cell Biol* 27, 8698-8712, 2007; Verma et al., *J Clin Invest* 116, 1346-1359, 2006; Verma et al., *J Biol Chem* 278, 20716-20723, 2003; Barletta et al., *J Biol Chem* 278, 19266-19271, 2003; Holzman et al., *Kidney Int* 56, 1481-1491, 1999; Ahola et al., *Am J Pathol* 155, 907-913, 1999; Tryggvason and Wartiovaara, *N Engl J Med* 354, 1387-1401, 2006; Schnabel and Farquhar, *J Cell Biol* 111, 1255-1263, 1990; Kurihara et al., *Proc Natl Acad Sci USA* 89, 7075-7079, 1992).

Early events of podocyte injury are characterized by dysregulation of the actin cytoskeleton (Faul et al., Trends Cell Biol 17, 428-437, 2007; Takeda et al., J Clin Invest 108, 289-301, 2001; Asanuma et al., Nat Cell Biol 8, 485-491, 2006) and Ca^{2+} homeostasis (Hunt et al., J Am Soc Nephrol 16, 1593-1602, 2005; Faul et al., Nat Med 14, 931-938, 2008). These changes are associated with the onset of proteinuria, the loss of albumin into the urinary space, and ultimately kidney failure (Tryggvason and Wartiovaara, N Engl J Med 354, 1387-1401, 2006). The vasoactive hormone Angiotensin II induces Ca^{2+} influx in podocytes, and prolonged treatment results in loss of stress fibers (Hsu et al., J Mol Med 86, 1379-1394, 2008). While there is a recognized link between Ca^{2+} influx and cytoskeletal reorganization, the mechanisms by which the podocyte senses and transduces extracellular cues that modulate cell shape and motility remain elusive. TRP Canonical 6 (TRPC6) channel mutations have been linked to podocyte injury (Winn et al., Science 308, 1801-1804, 2005; Reiser et al., Nat Genet 37, 739-744, 2005; Moller et al., J Am Soc Nephrol 18, 29-36, 2007; Hsu et al., Biochim Biophys Acta 1772, 928-936, 2007), but little is known about the specific pathways that regulate this process. Moreover, TRPC6 shares close homology with six other members of the TRPC channel family (Ramsey et al., Annu Rev Physiol 68, 619-647, 2006; Clapham, Nature 426, 517-524, 2003). TRPC5 channels antagonize TRPC6 channel activity to control a tightly regulated balance of cytoskeletal dynamics through differential coupling to distinct small GTPases.

Proteinuria

Proteinuria is a pathological condition wherein protein is present in the urine. Albuminuria is a type of proteinuria. Microalbuminuria occurs when the kidney leaks small amounts of albumin into the urine. In a properly functioning body, albumin is not normally present in urine because it is retained in the bloodstream by the kidneys. Microalbuminuria is diagnosed either from a 24-hour urine collection (20 to 200 $\mu\text{g}/\text{min}$) or, more commonly, from elevated concentrations (30 to 300 mg/L) on at least two occasions. Microalbuminuria can be a forerunner of diabetic nephropathy. An albumin level above these values is called macroalbuminuria. Subjects with certain conditions, e.g., diabetic nephropathy, can progress from microalbuminuria to macroalbuminuria and reach a nephrotic range ($>3.5 \text{ g}/24 \text{ hours}$) as kidney disease reaches advanced stages.

Causes of Proteinuria

Proteinuria can be associated with a number of conditions, including focal segmental glomerulosclerosis, IgA nephropathy, diabetic nephropathy, lupus nephritis, membranoproliferative glomerulonephritis, progressive (crescentic) glomerulonephritis, and membranous glomerulonephritis.

A. Focal Segmental Glomerulosclerosis (FSGS)

Focal Segmental Glomerulosclerosis (FSGS) is a disease that attacks the kidney's filtering system (glomeruli) causing serious scarring. FSGS is one of the many causes of a disease known as Nephrotic Syndrome, which occurs when protein in the blood leaks into the urine (proteinuria).

Very few treatments are available for patients with FSGS. Many patients are treated with steroid regimens, most of which have very harsh side effects. Some patients have shown to respond positively to immunosuppressive drugs as well as blood pressure drugs which have shown to lower the level of protein in the urine. To date, there is no commonly accepted effective treatment or cure and there are no FDA approved drugs to treat FSGS. Therefore, more effective methods to reduce or inhibit proteinuria are desirable.

B. IgA Nephropathy

IgA nephropathy (also known as IgA nephritis, IgAN, Berger's disease, and synpharyngitic glomerulonephritis) is a form of glomerulonephritis (inflammation of the glomeruli of the kidney). IgA nephropathy is the most common glomerulonephritis throughout the world. Primary IgA nephropathy is characterized by deposition of the IgA antibody in the glomerulus. There are other diseases associated with glomerular IgA deposits, the most common being Henoch-Schönlein purpura (HSP), which is considered by many to be a systemic form of IgA nephropathy. Henoch-Schönlein purpura presents with a characteristic purpuric skin rash, arthritis, and abdominal pain and occurs more commonly in young adults (16-35 yrs old). HSP is associated with a more benign prognosis than IgA nephropathy. In IgA nephropathy there is a slow progression to chronic renal failure in 25-30% of cases during a period of 20 years.

C. Diabetic Nephropathy

Diabetic nephropathy, also known as Kimmelstiel-Wilson syndrome and intercapillary glomerulonephritis, is a progressive kidney disease caused by angiopathy of capillaries in the kidney glomeruli. It is characterized by nephrotic syndrome and diffuse glomerulosclerosis. It is due to longstanding diabetes mellitus and is a prime cause for dialysis. The earliest detectable change in the course of diabetic nephropathy is a thickening in the glomerulus. At this stage, the kidney may start allowing more serum albumin than normal in the urine. As diabetic nephropathy progresses, increasing numbers of glomeruli are destroyed by nodular glomerulosclerosis and the amount of albumin excreted in the urine increases.

D. Lupus Nephritis

Lupus nephritis is a kidney disorder that is a complication of systemic lupus erythematosus. Lupus nephritis occurs when antibodies and complement build up in the kidneys, causing inflammation. It often causes proteinuria and may progress rapidly to renal failure. Nitrogen waste products build up in the bloodstream. Systemic lupus erythematosus causes various disorders of the internal structures of the kidney, including interstitial nephritis. Lupus nephritis affects approximately 3 out of 10,000 people.

E. Membranoproliferative Glomerulonephritis I/II/III

Membranoproliferative glomerulonephritis is a type of glomerulonephritis caused by deposits in the kidney glomerular mesangium and basement membrane thickening, activating complement and damaging the glomeruli. There are three types of membranoproliferative glomerulonephritis. Type I is caused by immune complexes depositing in the kidney and is believed to be associated with the classical complement pathway. Type II is similar to Type I, however, it is believed to be associated with the alternative complement pathway. Type III is very rare and it is characterized by a mixture of subepithelial deposits and the typical pathological findings of Type I disease.

F. Progressive (Crescentic) Glomerulonephritis

Progressive (crescentic) glomerulonephritis (PG) is a syndrome of the kidney that, if left untreated, rapidly progresses into acute renal failure and death within months. In 50% of cases, PG is associated with an underlying disease such as Goodpasture's syndrome, systemic lupus erythematosus, or Wegener granulomatosis; the remaining cases are idiopathic. Regardless of the underlying cause, PG involves severe injury to the kidney's glomeruli, with many of the glomeruli containing characteristic crescent-shaped scars. Patients with PG have hematuria, proteinuria, and occasionally, hypertension and edema. The clinical picture is consistent with nephritic syndrome, although the degree of proteinuria may occasionally exceed 3 g/24 hours, a range associated with nephrotic syndrome. Untreated disease may progress to decreased urinary volume (oliguria), which is associated with poor kidney function.

G. Membranous Glomerulonephritis

Membranous glomerulonephritis (MGN) is a slowly progressive disease of the kidney affecting mostly patients between ages of 30 and 50 years, usually Caucasian. It can develop into nephrotic syndrome. MGN is caused by circulating immune complex. Current research indicates that the majority of the immune complexes are formed via binding of antibodies to antigens in situ to the glomerular basement membrane. The said antigens may be endogenous to the basement membrane, or deposited from systemic circulation.

Measurement of Urine Protein Levels

Protein levels in urine can be measured using methods known in the art. Until recently, an accurate protein measurement required a 24-hour urine collection. In a 24-hour collection, the patient urinates into a container, which is kept refrigerated between trips to the bathroom. The patient is instructed to begin collecting urine after the first trip to the bathroom in the morning. Every drop of urine for the rest of the day is to be collected in the container. The next morning, the patient adds the first urination after waking and the collection is complete.

More recently, researchers have found that a single urine sample can provide the needed information. In the newer technique, the amount of albumin in the urine sample is compared with the amount of creatinine, a waste product of normal muscle breakdown. The measurement is

called a urine albumin-to-creatinine ratio (UACR). A urine sample containing more than 30 milligrams of albumin for each gram of creatinine (30 mg/g) is a warning that there may be a problem. If the laboratory test exceeds 30 mg/g, another UACR test should be performed 1 to 2 weeks later. If the second test also shows high levels of protein, the person has persistent proteinuria, a sign of declining kidney function, and should have additional tests to evaluate kidney function.

Tests that measure the amount of creatinine in the blood will also show whether a subject's kidneys are removing wastes efficiently. Too much creatinine in the blood is a sign that a person has kidney damage. A physician can use the creatinine measurement to estimate how efficiently the kidneys are filtering the blood. This calculation is called the estimated glomerular filtration rate, or eGFR. Chronic kidney disease is present when the eGFR is less than 60 milliliters per minute (mL/min).

TRPC5

TRPC is a family of transient receptor potential cation channels in animals. TRPC5 is subtype of the TRPC family of mammalian transient receptor potential ion channels. Three examples of TRPC5 are highlighted below in Table 1.

TABLE 1

The TRPC5 orthologs from three different species along with their GenBank Ref Seq Accession Numbers.			
Species	Nucleic Acid	Amino Acid	GeneID
<i>Homo sapiens</i>	NM_012471.2	NP_036603.1	7224
<i>Mus musculus</i>	NM_009428.2	NP_033454.1	22067
<i>Rattus norvegicus</i>	NM_080898.2	NP_543174.1	140933

Accordingly, in certain embodiments, the invention provides methods for treating, or the reducing risk of developing, a disease or condition selected from kidney disease, pulmonary arterial hypertension, anxiety, depression, cancer, diabetic retinopathy, or pain, comprising

administering to a subject in need thereof a therapeutically effective amount of a compound of the invention (e.g., a compound of structural formula I) or a pharmaceutical composition comprising said compound.

In some embodiments, the disease is kidney disease, anxiety, depression, cancer, or diabetic retinopathy.

In some embodiments, the disease or condition is kidney disease selected from Focal Segmental Glomerulosclerosis (FSGS), Diabetic nephropathy, Alport syndrome, hypertensive kidney disease, nephrotic syndrome, steroid-resistant nephrotic syndrome, minimal change disease, membranous nephropathy, idiopathic membranous nephropathy, membranoproliferative glomerulonephritis (MPGN), immune complex-mediated MPGN, complement-mediated MPGN, Lupus nephritis, postinfectious glomerulonephritis, thin basement membrane disease, mesangial proliferative glomerulonephritis, amyloidosis (primary), c1q nephropathy, rapidly progressive GN, anti-GBM disease, C3 glomerulonephritis, hypertensive nephrosclerosis, or IgA nephropathy. In some embodiments, the kidney disease is proteinuric kidney disease. In some embodiments, the kidney disease is microalbuminuria or macroalbuminuria kidney disease.

In some embodiments, the disease or condition to be treated is pulmonary arterial hypertension.

In some embodiments, the disease or condition to be treated is pain selected from neuropathic pain and visceral pain.

In some embodiments, the disease or condition is cancer selected from chemoresistant breast carcinoma, adriamycin-resistant breast cancer, chemoresistant colorectal cancer, medulloblastoma, and tumor angiogenesis.

The invention also provides methods of treating, or the reducing risk of developing, anxiety, or depression, or cancer, comprising administering to a subject in need thereof a therapeutically effective amount of a compound of the invention (e.g., a compound of Formula I), or a pharmaceutical composition comprising said compound.

In some embodiments, the disease or condition to be treated is transplant-related FSGS, transplant-related nephrotic syndrome, transplant-related proteinuria, cholestatic liver disease, polycystic kidney disease, autosomal dominant polycystic kidney disease (ADPKD), obesity,

insulin resistance, Type II diabetes, prediabetes, metabolic syndrome, non-alcoholic fatty liver disease (NAFLD), or non-alcoholic steatohepatitis (NASH).

Subjects to be Treated

In one aspect of the invention, a subject is selected on the basis that they have, or are at risk of developing, a kidney disease, pulmonary arterial hypertension, anxiety, depression, cancer, diabetic retinopathy, or pain. In another aspect, a subject is selected on the basis that they have, or are at risk of developing, kidney disease, anxiety, depression, cancer, or diabetic retinopathy. In another aspect of the invention, a subject is selected on the basis that they have, or are at risk of developing, pain, neuropathic pain, visceral pain, transplant-related FSGS, transplant-related nephrotic syndrome, transplant-related proteinuria, cholestatic liver disease, polycystic kidney disease, autosomal dominant polycystic kidney disease (ADPKD), obesity, insulin resistance, Type II diabetes, prediabetes, metabolic syndrome, non-alcoholic fatty liver disease (NAFLD), or non-alcoholic steatohepatitis (NASH).

Subjects that have, or are at risk of developing, proteinuria include those with diabetes, hypertension, or certain family backgrounds. In the United States, diabetes is the leading cause of end-stage renal disease (ESRD). In both type 1 and type 2 diabetes, albumin in the urine is one of the first signs of deteriorating kidney function. As kidney function declines, the amount of albumin in the urine increases. Another risk factor for developing proteinuria is hypertension. Proteinuria in a person with high blood pressure is an indicator of declining kidney function. If the hypertension is not controlled, the person can progress to full kidney failure. African Americans are more likely than Caucasians to have high blood pressure and to develop kidney problems from it, even when their blood pressure is only mildly elevated. Other groups at risk for proteinuria are American Indians, Hispanics/Latinos, Pacific Islander Americans, older adults, and overweight subjects.

In one aspect of the invention, a subject is selected on the basis that they have, or are at risk of developing proteinuria. A subject that has, or is at risk of developing, proteinuria is one having one or more symptoms of the condition. Symptoms of proteinuria are known to those of skill in the art and include, without limitation, large amounts of protein in the urine, which may cause it to look foamy in the toilet. Loss of large amounts of protein may result in edema, where


swelling in the hands, feet, abdomen, or face may occur. These are signs of large protein loss and indicate that kidney disease has progressed. Laboratory testing is the only way to find out whether protein is in a subject's urine before extensive kidney damage occurs.

The methods are effective for a variety of subjects including mammals, e.g., humans and other animals, such as laboratory animals, e.g., mice, rats, rabbits, or monkeys, or domesticated and farm animals, e.g., cats, dogs, goats, sheep, pigs, cows, or horses. In some embodiments, the subject is a mammal. In some embodiments, the subject is a human.

EXAMPLES

The invention is further described in the following examples, which do not limit the scope of the invention described in the claims.

Example 1. Synthesis of Compound 100



tert-butyl 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (400 mg, 1.48 mmol, 1 equiv.) and 4-fluoro-2-(trifluoromethyl)phenol (400.6 mg, 2.22 mmol, 1.5 equiv.) in acetonitrile (10 mL) was added DBU (451.5 mg, 2.97 mmol, 2.00 equiv.) at room temperature. The resulting mixture was stirred for 2 h at 80 °C. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The resulting

mixture was concentrated under reduced pressure. The resulting mixture was extracted with DCM (3 x 100 mL). The combined organic layers were washed with brine (3 x 100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE/EtOAc 2:1) to afford tert-butyl 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (110 mg, 17.94%) as a brown solid.

4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine

To a stirred solution of tert-butyl 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (110 mg, 0.27 mmol, 1 equiv.) in DCM (4 mL) was added TFA (1 mL, 13.46 mmol, 50.59 equiv.) at room temperature. The resulting mixture was stirred for 1 h at room temperature. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The mixture was basified to pH 8 with saturated NaHCO₃ (aq.). The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (DCM / MeOH 12:1) to afford 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (50 mg, 59.98%) as a brown solid.

4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


To a stirred solution of 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (50 mg, 0.16 mmol, 1 equiv.) in DIEA (2 mL) was added 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (47.5 mg, 0.19 mmol, 1.19 equiv.) at room temperature. The resulting mixture was stirred for 2 h at 100°C. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The residue was purified by Prep-TLC (PE/EtOAc 2:1) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (40 mg, 47.65%) as a brown solid.

4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (40 mg, 0.08 mmol, 1 equiv.) in DCM (4 mL) was added TFA (1 mL, 13.46 mmol, 177.00 equiv.) dropwise at room temperature. The resulting mixture was stirred for 1 h at room temperature. The reaction was

monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The mixture was basified to pH 8 with saturated NaHCO_3 (aq.). The resulting mixture was concentrated under reduced pressure. The crude product (40 mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30×150mm 5μm; Mobile Phase A: Water(10MMOL/L NH_4HCO_3), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 18% B to 47% B in 7 min; 220 nm; Rt: 6.22 min) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one(8.6 mg, 25.59%) as a white solid.

Example 2. Synthesis of Compound 140



Tert-butyl 4-bromo-5,6,7,8-tetrahydro-1,7-naphthyridine-7-carboxylate

To a solution of 4-bromo-5,6,7,8-tetrahydro-1,7-naphthyridine (250 mg, 1.173 mmol, 1 equiv.) in THF (10 mL, 123.430 mmol, 105.20 equiv.) were added Boc_2O (512.13 mg, 2.347 mmol, 2.00 equiv.) and TEA (474.90 mg, 4.693 mmol, 4 equiv.) at 25°C. The solution was stirred at 25°C for 2 hours. The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE/EA 5/1) to afford tert-butyl 4-bromo-5,6,7,8-tetrahydro-1,7-naphthyridine-7-carboxylate (210 mg, 57.15%) as a light yellow oil.

Tert-butyl 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridine-7-carboxylate

To a solution of tert-butyl 4-bromo-5,6,7,8-tetrahydro-1,7-naphthyridine-7-carboxylate (210 mg, 0.671 mmol, 1 equiv.) and 4-fluoro-2-(trifluoromethyl)phenol (241.52 mg, 1.341 mmol, 2 equiv.) in DMSO (10 mL) were added Cs_2CO_3 (873.86 mg, 2.682 mmol, 4 equiv.), 2-(dimethylamino)acetic acid (41.46 mg, 0.402 mmol, 0.6 equiv.) and CuI (76.62 mg, 0.402 mmol,

0.60 equiv). After stirring for 4 hours at 120°C under a nitrogen atmosphere, the resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC, eluted with PE/EA (5/1) to afford tert-butyl 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridine-7-carboxylate (100 mg, 36.17%) as a light yellow solid.

4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridine


To a solution of tert-butyl 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridine-7-carboxylate (150 mg, 0.364 mmol, 1 equiv.) in DCM (10 mL, 157.300 mmol, 432.46 equiv.) was added TFA (414.75 mg, 3.637 mmol, 10 equiv.) at 25°C. The solution was stirred at 25°C for 2 hours. The resulting mixture was concentrated under reduced pressure. The residue was used the next step.

4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

A mixture of 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridine (60 mg, 0.192 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (47.86 mg, 0.192 mmol, 1.00 equiv.) in DIEA (49.67 mg, 0.384 mmol, 2 equiv.) was stirred for 2 hours at 100°C under N₂ atmosphere. The residue was purified by Prep-TLC (PE/EA 1/1) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (100 mg, 99.15%) as a light yellow solid.

4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridin-7-yl]-2,3-dihydropyridazin-3-one

To a solution of 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (100 mg, 0.191 mmol, 1 equiv.) in DCM (10 mL, 157.300 mmol, 825.67 equiv.) was added TFA (217.23 mg, 1.905 mmol, 10.00 equiv.) at 25°C. The solution was stirred at 25°C for 2 hours. The crude product (150 mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Shield RP18 OBD Column 30*150mm, 5µm ; Mobile Phase A: Water (10 mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 20% B to 40% B in 7 min; 220 nm; Rt: 6.63 min) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5,6,7,8-tetrahydro-1,7-naphthyridin-7-yl]-2,3-dihydropyridazin-3-one (42.9 mg, 51.09%) as a white solid.

Example 3. Synthesis of Compound 120**2-Benzyl-5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridine**

To a stirred mixture of 2-benzyl-5-bromo-1,2,3,4-tetrahydro-2,6-naphthyridine (250 mg, 0.825 mmol, 1 equiv.) and 2-(dimethylamino)acetic acid (170.05 mg, 1.649 mmol, 2.00 equiv.) in DMSO (5 mL) were added 4-fluoro-2-(trifluoromethyl)phenol (89.10 mg, 0.495 mmol, 0.6 equiv.) and CuI (94.22 mg, 0.495 mmol, 0.6 equiv.) at room temperature. Then Cs₂CO₃ (1074.59 mg, 3.298 mmol, 4 equiv.) was added at room temperature. The final reaction mixture was irradiated with microwave radiation for 1 hours at 120°C. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The crude product was purified by reverse phase flash with the following conditions (Column: XBridge Prep OBD C18 Column 30×150mm 5μm; Mobile Phase A: Water (10mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 18% B to 35% B in 8 min; 220 nm; Rt: 7.12 min) to afford 2-benzyl-5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridine (180 mg, 54.25%) as a brown solid.

5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridine


To a stirred solution of 2-benzyl-5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridine (180 mg) in MeOH (10 mL) was added Pd/C (20 mg) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 5 hours at room temperature under hydrogen atmosphere. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (DCM/MeOH 12:1) to afford 5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridine (100mg) as a brown solid.

4-chloro-5-[5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridin-2-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

To a stirred solution of 5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridine (100 mg, 0.320 mmol, 1 equiv.) in DIEA (0.1 mL) was added 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (63.81 mg, 0.256 mmol, 0.8 equiv.) at room temperature. The resulting mixture was stirred for 1 hours at 90°C. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The residue was purified by Prep-TLC (DCM/MeOH; 12:1) to afford 4-chloro-5-[5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridin-2-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (130 mg, 77.34%) as a white solid.

4-chloro-5-[5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridin-2-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridin-2-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (107 mg, 0.204 mmol, 1 equiv.) in DCM (4 mL) was added TFA (1 mL) at room temperature. The resulting mixture was stirred for 1 hours at room temperature. The reaction was monitored by LCMS. The mixture was basified to pH 7 with saturated NaHCO₃ (aq.). The resulting mixture was concentrated under reduced pressure. The crude product (50 mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30×150mm 5μm; Mobile Phase A: Water (10 mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 30% B to 50% B in 8 min; 220 nm; Rt: 7.55 min) to afford 4-chloro-5-[5-[4-fluoro-2-(trifluoromethyl)phenoxy]-1,2,3,4-tetrahydro-2,6-naphthyridin-2-yl]-2,3-dihydropyridazin-3-one (60 mg, 66.78%) as a white solid.

Example 4. Synthesis of Compound 118**Ethyl 2-(benzylamino)propanoate**

To a stirred solution of benzaldehyde (8 g, 75.384 mmol, 1 equiv.) and TEA (7.63 g, 75.384 mmol, 1 equiv.) in DCE (100 mL, 1263.149 mmol, 16.76 equiv.) was added TEA (7.63 g, 75.384 mmol, 1 equiv.) and $\text{NaBH}(\text{OAc})_3$ (31.95 g, 150.767 mmol, 2 equiv.) in portions at room temperature under nitrogen atmosphere. The mixture was stirred at rt overnight. Desired product could be detected by LCMS. The resulting mixture was extracted with DCM (2 x 150 mL). The combined organic layers were washed with brine (1x90 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure to afford ethyl 2-(benzylamino)propanoate (12 g, 76.80%) as colorless oil.

Methyl 4-[benzyl(1-ethoxy-1-oxopropan-2-yl)amino]butanoate

To a stirred solution of ethyl 2-(benzylamino)propanoate (8 g, 38.596 mmol, 1 equiv.) and methyl 4-oxobutanoate (4.48 g, 38.596 mmol, 1.00 equiv.) in DCE (120 mL, 1515.779 mmol, 39.27 equiv.) was added TEA (3.91 g, 38.596 mmol, 1 equiv.) and $\text{NaBH}(\text{OAc})_3$ (16.36 g, 77.193 mmol, 2 equiv.) in portions at room temperature under nitrogen atmosphere. The mixture was stirred at rt overnight. Desired product could be detected by LCMS. The resulting mixture

was extracted with DCM (2 x 150 mL). The combined organic layers were washed with brine (1x90 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure to afford methyl 4-[benzyl(1-ethoxy-1-oxopropan-2-yl)amino]butanoate (10g, 84.29%) as colorless oil.

Methyl 1-benzyl-2-methyl-3-oxopiperidine-4-carboxylate

To a stirred solution of methyl 4-[benzyl(1-ethoxy-1-oxopropan-2-yl)amino]butanoate (8 g, 26.026 mmol, 1 equiv.) in Toluene (100 mL) was added t-BuOK (5.00 g, 52.051 mmol, 2 equiv.) in portions at room temperature under nitrogen atmosphere. The mixture was stirred at 80°C for 2 hours. Desired product was detected by LCMS. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (5:1 to 2:1) to afford methyl 1-benzyl-2-methyl-3-oxopiperidine-4-carboxylate (6.5 g, 95.57%) as a white solid.

7-Benzyl-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-ol

To a stirred solution of methyl 1-benzyl-2-methyl-3-oxopiperidine-4-carboxylate (6 g, 22.960 mmol, 1 equiv.) in EtOH (80 mL, 1377.083 mmol, 59.98 equiv.) was added t-BuONa (4.41 g, 45.921 mmol, 2 equiv.) and methanimidamide hydrochloride (3.70 g, 45.921 mmol, 2.00 equiv.) in portions at room temperature under nitrogen atmosphere. The mixture was stirred at 80°C for 2h. Desired product could be detected by LCMS. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (3:1 to 2:1) to afford 7-benzyl-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-ol (5 g, 85.29%) as a white solid.

Tert-Butyl 4-hydroxy-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a solution of 7-benzyl-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-ol (5 g, 19.583 mmol, 1 equiv.) in EtOH (60 mL, 1032.812 mmol, 52.74 equiv.) was added Boc₂O (8.55 g, 39.166 mmol, 2 equiv), CH₃COONa (1.81 g, 23.500 mmol, 1.2 equiv), Pd(OH)₂/C (275.01 mg, 1.958 mmol, 0.1 equiv.) under nitrogen atmosphere. The mixture was hydrogenated at room temperature for 2h under hydrogen atmosphere, filtered through a Celite pad and concentrated under reduced pressure to afford tert-butyl 4-hydroxy-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (4.5 g, 86.61%) as white solid.

Tert-butyl 4-chloro-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 4-hydroxy-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (4.5 g, 16.961 mmol, 1 equiv.) and PPh₃ (6.67 g, 25.442 mmol, 1.5 equiv.) in DCE (60 mL, 0.606 mmol, 0.04 equiv.) was added CCl₄ (5.22 g, 33.922 mmol, 2 equiv.) in portions at room temperature under nitrogen atmosphere. The mixture was stirred at 70°C for 2 hours.

Desired product could be detected by LCMS. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (7:1) to afford tert-butyl 4-chloro-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (4 g, 83.11%) as a white solid.

Tert-butyl 4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 4-chloro-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (4 g, 14.096 mmol, 1 equiv.) and 2-chloro-4-fluorophenol (2.07 g, 14.096 mmol, 1 equiv.) in DMF (50 mL) was added K₂CO₃ (3.90 g, 28.193 mmol, 2 equiv.) in portions at room temperature under nitrogen atmosphere. The mixture was stirred at 70 for 1h. Desired product could be detected by LCMS. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (1:1) to afford tert-butyl 4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (4 g, 72.05%) as a white solid.

4-(2-Chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine

To a stirred solution of tert-butyl 4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (4 g, 1 equiv.) in DCM (20 mL) was added TFA (4 mL) dropwise/ in portions at room temperature under nitrogen atmosphere. The mixture was stirred at rt for 2h. Desired product could be detected by LCMS. The resulting mixture was concentrated under reduced pressure to afford 4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (2.7 g, 90.51%) as off-white solid.

4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

To a stirred solution of 4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (1 g, 3.404 mmol, 1 equiv.) in DIEA (1 mL) was added 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (0.85 g, 3.404 mmol, 1 equiv.) in portions at room temperature under nitrogen atmosphere. The mixture was stirred at 100°C overnight. The desired product


could be detected by LCMS. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (1:1 to 1:2) to afford 4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (1 g, 58.01%) as a white solid.

4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (1 g, 1 equiv.) in DCM (10 mL) was added TFA (2 mL) dropwise at room temperature under nitrogen atmosphere. The mixture was stirred at rt for 1h. Desired product could be detected by LCMS. The resulting mixture was concentrated under reduced pressure to afford 4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (600 mg, 71.95%) as white solid.

4-chloro-5-[(8R)-4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (250 mg, 1 equiv.) was separated by prep chiral- HPLC (Column: CHIRALPAK IG, 20*250mm, 5 um; Mobile Phase A:Hex:DCM=3:1 (0.1%FA)--HPLC, Mobile Phase B: EtOH--HPLC; Flow rate: 20 mL/min; Gradient: 15 B to 15 B in 19 min; 220/254 nm ; RT1:13.016 ; RT2:16.004) to afford 4-chloro-5-[(8R)-4-(2-chloro-4-fluorophenoxy)-8-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (144 mg, 57.60%) as white solid.

Example 5. Synthesis of Compound 103**Tert-Butyl 4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate**

To a stirred solution of tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (800 mg, 2.966 mmol, 1 equiv.) and 2-(difluoromethyl)phenyl acetate (1104.26 mg, 5.932 mmol, 2.00 equiv.) in DMF (20 mL) were added K₂CO₃ (1229.72 mg, 8.898 mmol, 3 equiv.) in portions at 80°C under nitrogen atmosphere. The mixture was stirred for 2 hours. The reaction was monitored by LCMS. The reaction was quenched with Water at room temperature. The mixture was extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine (3 x 100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeOH in water, 10% to 50% gradient in 10 min; detector, UV 254 nm. to afford tert-butyl 4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (900 mg, 80.41%) as off-white solid.

4-[2-(Difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine

To a stirred solution of tert-butyl 4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (900 mg, 2.385 mmol, 1 equiv.) in DCM was added 3,3,3-trifluoropropanoic acid (3 mL, 6.00 equiv.) dropwise at room temperature. The mixture was stirred for 1.5 hours. The reaction was monitored by TLC (PE/EtOAc 10:1). The residue was basified to pH 8 with saturated NaHCO₃ (aq.). The mixture was concentrated under reduced pressure. The crude product (100 mg) was purified by Prep-HPLC with the following conditions


to afford 4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (329 mg, 49.75%) as off-white solid.

4-Chloro-5-[4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

To a stirred solution of 4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (328 mg, 1.183 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (169.05 mg, 0.679 mmol, 1.00 equiv.) were added DIEA (175.43 mg, 1.357 mmol, 2.00 equiv.) in portions at 70°C. The mixture was stirred for 2 hours at 70°C. The residue was purified by reverse flash chromatography with the following conditions: column, C18 silica gel; mobile phase, MeOH in water, 10% to 50% gradient in 10 min; detector, UV 254 nm. to afford 4-chloro-5-[4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (328 mg, 56.60%) as off-white solid.

4-Chloro-5-[4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (328 mg, 0.670 mmol, 1 equiv.) in DCM (10 mL) was added trifluoroacetic acid (3 mL) dropwise at room temperature. The mixture was concentrated under vacuum. The product was purified by Prep-HPLC to afford 4-chloro-5-[4-[2-(difluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (256.4 mg, 94.38%) as off-white solid.

Example 6. Synthesis of Compound 117 and 117a**Ethyl 4-[(1-phenylethyl)amino]pentanoate**

To a stirred solution of 1-phenylethan-1-amine (25 g, 206.300 mmol, 1 equiv.) and ethyl 4-oxopentanoate (29.74 g, 206.300 mmol, 1 equiv.) in DCE (400 mL, 5052.598 mmol, 24.49 equiv.) was added $\text{NaBH}(\text{OAc})_3$ (65.59 g, 309.449 mmol, 1.5 equiv.) in portions at 25°C under nitrogen atmosphere. The solution was stirred at 25°C for 2 hours. The reaction was quenched by the addition of H_2O (400 mL) at 0°C. The resulting mixture was extracted with DCM (3 x 200 mL). The combined organic layers were washed with saturated NaCl (aq.) (3 x 200 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The crude product was used to the next step.

Ethyl 4-[(2-ethoxy-2-oxoethyl)(1-phenylethyl)amino]pentanoate

To a stirred solution of ethyl 4-[(1-phenylethyl)amino]pentanoate (49 g, 196.508 mmol, 1 equiv.) and ethyl 2-oxoacetate (40.12 g, 392.990 mmol, 2.00 equiv.) in DCE (500 mL, 6315.747 mmol, 32.14 equiv.) was added $\text{NaBH}(\text{OAc})_3$ (62.47 g, 294.762 mmol, 1.5 equiv.) in portions at 25°C under nitrogen atmosphere. The solution was stirred at 25°C for 2 hours. The reaction was

quenched by the addition of H₂O (400 mL) at 0°C. The resulting mixture was extracted with DCM (3 x 200 mL). The combined organic layers were washed with saturated NaCl (aq.) (3 x 200 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The crude product ethyl 4-[(2-ethoxy-2-oxoethyl)(1-phenylethyl)amino]pentanoate (57 g, 86.47%) was used to the next step.

Ethyl 2-methyl-5-oxo-1-(1-phenylethyl)piperidine-4-carboxylate

To a solution of ethyl 4-[(2-ethoxy-2-oxoethyl)(1-phenylethyl)amino]pentanoate (57 g, 169.924 mmol, 1 equiv.) in Toluene (500 mL, 4699.452 mmol, 27.66 equiv.) was added t-BuOK (47.67 g, 424.810 mmol, 2.5 equiv.) in ports at 0°C. The mixture was stirred at 25°C for 2 hours. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EA (50/1 to 10/1) to afford ethyl 2-methyl-5-oxo-1-(1-phenylethyl)piperidine-4-carboxylate (29 g, 58.98%) as a yellow oil

7-(1-cyclohexylethyl)-6-methyl-decahydropyrido[3,4-d]pyrimidin-4-ol

To a solution of ethyl 2-methyl-5-oxo-1-(1-phenylethyl)piperidine-4-carboxylate (10 g, 34.557 mmol, 1 equiv.) and methanimidamide hydrochloride (4.17 g, 51.836 mmol, 1.50 equiv.) in EtOH (100 mL, 1721.353 mmol, 49.81 equiv.) was added EtONa (5.88 g, 86.393 mmol, 2.50 equiv.) in ports at 25°C. The mixture was stirred at 90°C for 2 hours. The residue was purified by silica gel column chromatography, eluted with DCM/MeOH (20/1 to 10/1) to afford 7-(1-cyclohexylethyl)-6-methyl-decahydropyrido[3,4-d]pyrimidin-4-ol (3.4 g, 34.96%) as a yellow solid.

Tert-Butyl 4-hydroxy-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a solution of 6-methyl-7-(1-phenylethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-ol (3.5 g, 12.994 mmol, 1 equiv), HCOONH₄ (4.10 g, 65.022 mmol, 5.00 equiv.) and Boc₂O (8.51 g, 38.983 mmol, 3 equiv.) in EtOH (50 mL, 860.677 mmol, 66.23 equiv.) was added Pd(OH)₂/C (0.36 g, 2.599 mmol, 0.2 equiv.) under nitrogen atmosphere. The mixture was hydrogenated at 70°C for 2 hours under hydrogen atmosphere using a hydrogen balloon, filtered through a Celite pad and concentrated under reduced pressure. To afford tert-butyl 4-hydroxy-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1.8 g, 52.21%) as a yellow solid.

Tert-Butyl 4-chloro-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a solution of tert-butyl 4-hydroxy-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1.8 g, 6.784 mmol, 1 equiv.) and PPh₃ (3.56 g, 13.569 mmol, 2 equiv.) in DCE (20

mL, 252.630 mmol, 37.24 equiv.) was added CCl₄ (3.13 g, 20.353 mmol, 3 equiv.) at 25°C. The mixture was stirred at 70°C for 3 hours. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EA (10/1 to 1/1) to afford tert-butyl 4-chloro-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1.1 g, 57.14%) as a yellow solid.

Tert-Butyl 4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a solution of tert-butyl 4-chloro-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1.1 g, 3.877 mmol, 1 equiv.) and 2-chloro-4-fluorophenol (0.85 g, 5.800 mmol, 1.50 equiv.) in DMF (15 mL, 193.826 mmol, 50.00 equiv.) was added K₂CO₃ (1.07 g, 7.753 mmol, 2 equiv.) at 25°C. The mixture was stirred at 70°C for 1 hour. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EA (10/1 to 5/1) to afford tert-butyl 4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1.2 g, 78.60%) as a yellow solid.

4-Chloro-5-[4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


A mixture of 4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (800 mg, 2.724 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (678.42 mg, 2.724 mmol, 1.00 equiv.) in DIEA (704.01 mg, 5.447 mmol, 2 equiv.) was stirred for 16 hours at 100°C under nitrogen atmosphere. The residue was purified by Prep-TLC (PE/EA 1/1) to afford 4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (530 mg, 38.43%) as a light yellow solid.

4-chloro-5-[(6R)-4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a solution of 4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (530 mg, 1.047 mmol, 1 equiv.) in DCM (20 mL, 314.601 mmol, 300.57 equiv.) was added TFA (1193.47 mg, 10.467 mmol, 10 equiv.) at 25°C. The solution was stirred at 25°C for 2 hours. The resulting mixture was concentrated under reduced pressure. The crude product (600 mg) was purified by Prep-HPLC

with the following conditions (Column: XBridge Shield RP18 OBD Column 30*150mm,5um ; Mobile Phase A: Water (10 mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 20% B to 40% B in 7 min; 220 nm; Rt: 6.63 min) to afford the racemate (200 mg). The residue (200 mg) was purified by Chiral-Prep-HPLC with the following conditions: Column: CHIRALPAK IE, 2*25cm,5um; Mobile Phase A: MTBE (0.1%FA)-HPLC, Mobile Phase B: IPA--HPLC; Flow rate: 18 mL/min; Gradient: 20 B to 20 B in 15 min; 220/254 nm. Although the two isomers were separated by this technique, the absolute orientation was not determined. The compound designated as 4-chloro-5-[(6S)-4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (60.9mg,13.78%) was obtained at 9.688 min as a white solid. The compound designated as 4-chloro-5-[(6R)-4-(2-chloro-4-fluorophenoxy)-6-methyl-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (61.5mg,13.92%) was obtained at 11.813 min as a white solid.

Example 7. Synthesis of Compound 134



Tert-butyl 2-chloro-4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of 2-(difluoromethyl)-4-fluorophenol (5.33 g, 32.879 mmol, 2.00 equiv.) and tert-butyl 2,4-dichloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (5 g, 16.438 mmol, 1 equiv.) in DMF (30 mL) was added NaHCO₃ (4.14 g, 49.282 mmol, 3.00 equiv.) at room temperature. The solution was stirred at 70°C for 0.5 hours. The mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 um, 330 g; Mobile Phase A: Water

(plus 10 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 70% B - 95% B gradient in 100 min; Detector: 254 nm. The fractions containing the desired product were collected at 92% B and concentrated under reduced pressure to afford tert-butyl 2-chloro-4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (2.100g) as off-white solid.

7-tert-Butyl 2-methyl 4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-2,7-dicarboxylate

To a solution of tert-butyl 2-chloro-4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (400 mg, 0.931 mmol, 1 equiv.) and TEA (188.34 mg, 1.861 mmol, 2 equiv.) in MeOH (15 mL, 370.484 mmol, 398.10 equiv.) was added Pd(PPh₃)₄ (107.54 mg, 0.093 mmol, 0.1 equiv.) in a pressure tank. The mixture was purged with nitrogen for 1 hours and then was pressurized to 10 atm with carbon monoxide at 100°C for 16 hours. The reaction mixture was cooled to room temperature and filtered to remove insoluble solids. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C₁₈, 20 - 40 μm , 330 g; Mobile Phase A: Water (plus 10 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 35% B - 65% B gradient in 20 min; Detector: 254 nm. The fractions containing the desired product were collected at 62% B and concentrated under reduced pressure to afford 7-tert-butyl 2-methyl 4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-2,7-dicarboxylate (100 mg, 23.70%) as colorless oil.

Tert-butyl 4-[2-(difluoromethyl)-4-fluorophenoxy]-2-(hydroxymethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of 7-tert-butyl 2-methyl 4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-2,7-dicarboxylate (100 mg, 0.221 mmol, 1 equiv.) in t-BuOH (6 mL, 63.139 mmol, 286.29 equiv.) was added NaBH₄ (16.69 mg, 0.441 mmol, 2 equiv.) at room temperature. The solution was stirred at 70°C for 3 hours. To the mixture was added water (3 mL). The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C₁₈, 20 - 40 μm , 330 g; Mobile Phase A: Water (plus 10 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 45% B - 80% B gradient in 20 min; Detector: 254 nm. The fractions containing the desired product were collected at 74% B and concentrated under reduced pressure to afford tert-

butyl 4-[2-(difluoromethyl)-4-fluorophenoxy]-2-(hydroxymethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (35 mg, 37.30%) as colorless oil.

[4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]methanol

To a stirred solution of tert-butyl 4-[2-(difluoromethyl)-4-fluorophenoxy]-2-(hydroxymethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (35 mg) in DCM (6 mg) was added TFA (1 mg) at room temperature. The solution was stirred at rt for 2 hours. The mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μ m, 330 g; Mobile Phase A: Water (plus 10 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 25% B - 55% B gradient in 20 min; Detector: 254 nm. The fractions containing the desired product were collected at 41% B and concentrated under reduced pressure to afford as [4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]methanol (20 mg) as colorless oil.

4-chloro-5-[4-[2-(difluoromethyl)-4-fluorophenoxy]-2-(hydroxymethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


Into a 25 mL round-bottom flask were added [4-[2-(difluoromethyl)-4-fluorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]methanol (20 mg, 0.061 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (15.31 mg, 0.061 mmol, 1 equiv.) at room temperature. To the mixture was added DIEA (15.89 mg, 0.123 mmol, 2 equiv.) at rt. The mixture was stirred at 90°C for 2 hours. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μ m, 330 g; Mobile Phase A: Water (plus 10 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 35% B - 70% B gradient in 20 min; Detector: 254 nm. The fractions containing the desired product were collected at 65% B and concentrated under reduced pressure to afford 4-chloro-5-[4-[2-(difluoromethyl)-4-fluorophenoxy]-2-(hydroxymethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (30 mg, 90.71%) as colorless oil.

4-chloro-5-[4-[2-(difluoromethyl)-4-fluorophenoxy]-2-(hydroxymethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[4-[2-(difluoromethyl)-4-fluorophenoxy]-2-(hydroxymethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (30 mg) in DCM (5 mL) was added TFA (1 mL) at room temperature. The solution was stirred at rt for 2 hours. The mixture was concentrated under reduced pressure. The crude product (30 mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30×150mm 5um; Mobile Phase A: undefined, Mobile Phase B: undefined; Flow rate: 60 mL/min; Gradient: 20% B to 40% B in 8 min; 220 nm; Rt: 7.22 min) to afford 4-chloro-5-[4-[2-(difluoromethyl)-4-fluorophenoxy]-2-(hydroxymethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (8.7 mg) as a white solid.

Compounds 128, 125, 114 were prepared by the methods and scheme described in this Example by using 2-trifluoromethylphenol, 4-fluoro-2-trifluoromethylphenol and 4-fluoro-2-chlorophenol respectively, in place of 2-(difluoromethyl)-4-fluorophenol in the first step of the synthesis.

Example 8. Synthesis of Compound 112



Tert-butyl 2-chloro-4-(2-chloro-4-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred mixture of tert-butyl 2,4-dichloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (800 mg, 2.630 mmol, 1 equiv.) and 2-chloro-4-fluorophenol (578.16 mg, 3.945 mmol, 1.50 equiv.) in DMF (15 mL) was added K₂CO₃ (726.99 mg, 5.260 mmol, 2.00 equiv.) in portions at rt under nitrogen atmosphere. The resulting mixture was stirred for 0.5 hours at 70°C under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to

cool down to rt. The resulting mixture was concentrated under reduced pressure. The resulting mixture was extracted with EtOAc (3 x 200 mL). The combined organic layers were washed with brine (2 x 100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (30/1 to 10/1) to afford tert-butyl 2-chloro-4-(2-chloro-4-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1 g, 91.78%) as a yellow oil.

Tert-Butyl 4-(2-chloro-4-fluorophenoxy)-2-[[4-methoxyphenyl)methyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred mixture of tert-butyl 2-chloro-4-(2-chloro-4-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (700 mg, 1.690 mmol, 1 equiv.) in THF (30 mL) was added 1-(4-methoxyphenyl)methanamine (1159.02 mg, 8.449 mmol, 5.00 equiv.) in portions at rt under nitrogen atmosphere. The resulting mixture was stirred for 16 hours at 60°C under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to cool down to rt. The resulting mixture was concentrated under reduced pressure. The resulting mixture was extracted with EtOAc (3 x 200 mL). The combined organic layers were washed with brine (2 x 100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse flash chromatography with the following conditions (Column, C18 silica gel; mobile phase, acetonitrile in water, 60% to 95% gradient in 20 min; detector, UV 220 nm) to afford tert-butyl 4-(2-chloro-4-fluorophenoxy)-2-[[4-methoxyphenyl)methyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (350 mg, 40.22%) as a yellow oil.

4-(2-chloro-4-fluorophenoxy)-N-[[4-methoxyphenyl)methyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-amine

To a stirred solution of tert-butyl 4-(2-chloro-4-fluorophenoxy)-2-[[4-methoxyphenyl)methyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (350 mg, 1 equiv.) in DCM (10 mL) was added TFA (1 mL) dropwise at rt. The reaction mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH₄HCO₃ (aq.). The resulting mixture was extracted with DCM (3 x 100 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was

concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (Column: XBridge Shield RP18 OBD Column, 5um, 19*150mm; Mobile Phase A: Water (10mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 25 mL/min; Gradient: 2% B to 32% B in 1 min; 220/254 nm; Rt: 7.08 min) to afford 4-(2-chloro-4-fluorophenoxy)-N-[(4-methoxyphenyl)methyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-amine (260 mg) as a yellow oil.

4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-2-[[4-(methoxyphenyl)methyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

Into a 50 mL round-bottom flask were added 4-(2-chloro-4-fluorophenoxy)-N-[(4-methoxyphenyl)methyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-amine (260 mg, 0.627 mmol, 1 equiv), 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (156.11 mg, 0.627 mmol, 1.00 equiv.) and DIEA (242.99 mg, 1.880 mmol, 3.00 equiv.) at rt under nitrogen atmosphere. The resulting mixture was stirred for 2 hours at 90°C under nitrogen atmosphere. The residue was purified by reverse flash chromatography with the following conditions (Column, C18 silica gel; mobile phase, acetonitrile in water, 50% to 85% gradient in 25 min; detector, UV 220 nm) to afford 4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-2-[[4-(methoxyphenyl)methyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (350 mg, 89.00%) as a yellow solid.


5-[2-amino-4-(2-chloro-4-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[4-(2-chloro-4-fluorophenoxy)-2-[[4-(methoxyphenyl)methyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (200 mg) in TFA (8 mL, 107.704 mmol, 328.23 equiv). The final reaction mixture was irradiated with microwave radiation for 2 hours at 80°C. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH₄HCO₃ (aq.). The resulting mixture was extracted with DCM (2 x 100 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30x150mm 5um; Mobile Phase A: Water (10mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 25% B to 40% B in 8 min; 220

nm; Rt: 7.35 min) to afford 5-[2-amino-4-(2-chloro-4-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2,3-dihydropyridazin-3-one (52.4 mg) as a yellow solid.

Compounds 113, 116, and 102 were prepared by the methods and scheme described in this Example by using 2-chlorophenol, 4-fluoro-2-trifluoromethylphenol, 2-trifluorophenol respectively, in place of 2-chloro-4-fluorophenol in the first step of the synthesis.

Example 9. Synthesis of Compounds 129 and 130



1-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]ethan-1-one

To a mixture of tert-butyl 2-chloro-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (600 mg, 1.340 mmol, 1 equiv.) and tributyl (1-ethoxyethenyl)stannane (967.80 mg, 2.680 mmol, 2.00 equiv.) in Toluene (10 mL) was added Pd(PPh₃)₄ (77.41 mg, 0.067 mmol, 0.05 equiv.) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 4 hours at 110 °C. The reaction was monitored by LCMS. This resulted in tert-butyl 2-(1-ethoxyethenyl)-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (700 mg, 108.06%) as a yellow oil. The crude resulting mixture was used in the next step directly without further purification.

To a stirred solution of tert-butyl 2-(1-ethoxyethenyl)-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1 g, 2.068 mmol, 1 equiv.) in DCM (5 mL) was added TFA (3.33 mL, 29.239 mmol, 21.70 equiv.) at room temperature. The resulting mixture was stirred for 2 hours at room temperature. The reaction was monitored by LCMS. The

resulting mixture was concentrated under reduced pressure. The mixture/residue was basified to pH 8 with saturated NaHCO_3 (aq.). The resulting mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μm , 330 g; Mobile Phase A: Water (plus 5 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 43% B - 55% B gradient in 20 min; Detector: 220 nm. The fractions containing the desired product were collected at 50% B and concentrated under reduced pressure to afford 1-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]ethan-1-one (750 mg, 102.06%) as a light yellow solid.

5-[2-Acetyl-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

Into a 50 mL round-bottom flask were added 1-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]ethan-1-one (750 mg, 2.111 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (525.81 mg, 2.111 mmol, 1.00 equiv.) at room temperature. To the above mixture was added DIEA (818.47 mg, 6.333 mmol, 3.00 equiv.). The resulting mixture was stirred for 2 hours at 100°C. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μm , 330 g; Mobile Phase A: Water (plus 5 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 60% B - 85% B gradient in 20 min; Detector: 220 nm. The fractions containing the desired product were collected at 80% B and concentrated under reduced pressure to afford 5-[2-acetyl-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (230 mg, 19.18%) as a light yellow oil.

4-Chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-(1-hydroxyethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

To a stirred solution of 5-[2-acetyl-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (230 mg, 0.405 mmol, 1 equiv.) in MeOH (10 mL) was added NaBH_4 (30.64 mg, 0.810 mmol, 2.00 equiv.) in portions at 0°C under nitrogen atmosphere. The resulting mixture was stirred for 2 hours at room temperature. The reaction was monitored by LCMS. The resulting mixture was concentrated

under reduced pressure. The residue was purified by Prep-TLC (PE/EtOAc 1/1) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-(1-hydroxyethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (120 mg, 51.99%) as a light yellow oil.


4-Chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-[(1S)-1-hydroxyethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one and 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-[(1R)-1-hydroxyethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-(1-hydroxyethyl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (120 mg, 0.211 mmol, 1 equiv.) in DCM (5 mL) was added TFA (2.00 mL, 17.541 mmol, 127.89 equiv.) dropwise at room temperature. The resulting mixture was stirred for 2 hours at room temperature. The reaction was monitored by LCMS. The residue was basified to pH 8 with saturated NaHCO₃ (aq.). The resulting mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C₁₈, 20 - 40 μ m, 330 g; Mobile Phase A: Water (plus 5 mM NH₄HCO₃); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 40% B - 80% B gradient in 25 min; Detector: 220 nm. The fractions containing the desired product were collected at 55% B and concentrated under reduced pressure. The crude product (50 mg) was purified by Chiral-Prep-HPLC with the following conditions (Column: CHIRALPAK IE, 2*25cm,5 μ m; Mobile Phase A:Hex (0.1%FA)--HPLC, Mobile Phase B: EtOH--HPLC; Flow rate: 16 mL/min; Gradient: 30 B to 30 B in 33 min; 220/254 nm ; RT1:26.219 ; RT2:29.589). Although the two isomers were separated by this technique, the absolute orientation was not determined. The compound designated as 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-[(1S)-1-hydroxyethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (27.1 mg) was obtained at 29.589 min as an off-white solid. The compound designated as 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-[(1R)-1-hydroxyethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (22.6 mg) was obtained at 26.219 min as an off-white solid.

Compound 119 was prepared by the methods and scheme described in this Example using tert-butyl 2-chloro-4-[4-fluoro-2-chlorophenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate as the starting material.

Compounds 122 and 123 were prepared by the methods and scheme described in this Example using tert-butyl 2-chloro-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate as the starting material. Again, the absolute orientation of these separated isomers was not determined and the designation as (*S*) or (*R*) was arbitrary.

Example 10. Synthesis of Compound 115



tert-Butyl 2-chloro-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 2,4-dichloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (2 g, 6.58 mmol, 1 equiv.) and 2-(trifluoromethyl)phenol (1.6 g, 9.86 mmol, 1.5 equiv.) in acetonitrile (20 mL) was added DBU (2.0 g, 13.15 mmol, 2 equiv.) at room temperature. The solution was stirred at rt for 4 hours. The mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE/EtOAc 10:1) to afford tert-butyl 2-chloro-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (700 mg, 24.77%) as colorless oil.

tert-butyl 2-([2-((tert-butyldimethylsilyl)oxy)ethyl]amino)-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a solution of tert-butyl 2-chloro-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 1.163 mmol, 1 equiv.) in THF (15 mL) was added (2-aminoethoxy)(tert-butyl)dimethylsilane (1019.89 mg, 5.816 mmol, 5.00 equiv.) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 16 h at 50 °C. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE/EtOAc 3/1) to afford tert-butyl 2-([2-[(tert-butyl)dimethylsilyl]oxy]ethyl)amino)-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (440 mg, 66.51%) as a light yellow oil.

2-([4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]amino)ethan-1-ol

To a stirred solution of tert-butyl 2-([2-[(tert-butyl)dimethylsilyl]oxy]ethyl)amino)-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (440 mg, 0.774 mmol, 1 equiv.) in DCM (10 mL) was added TFA (3 mL, 40.389 mmol, 52.20 equiv.) at room temperature. The resulting mixture was stirred for 2 h at room temperature. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was purified by reverse flash chromatography with the following conditions: column, C18 silica gel; mobile phase, ACN in water, 40% to 60% gradient in 15 min; detector, UV 254 nm to afford 2-([4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]amino)ethan-1-ol (220 mg) as light yellow oil.

4-chloro-5-[2-[(2-hydroxyethyl)amino]-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


Into a 50 mL round-bottom flask were added 2-([4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]amino)ethan-1-ol (220 mg, 0.621 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (154.66 mg, 0.621 mmol, 1.00 equiv.) at room temperature. To the above mixture was added DIEA (240.74 mg, 1.863 mmol, 3.00 equiv.). The resulting mixture was stirred for 2 h at 100 degrees C. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 um, 330 g; Mobile Phase A: Water (plus 5 mM NH₄HCO₃); Mobile Phase B: ACN; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 45% B - 60% B gradient in 20 min; Detector: 220 nm. The fractions containing the desired product were collected at 55% B and concentrated under

reduced pressure to afford 4-chloro-5-[2-[(2-hydroxyethyl)amino]-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (210 mg, 59.66%) as a yellow solid.

4-chloro-5-[2-[(2-hydroxyethyl)amino]-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[2-[(2-hydroxyethyl)amino]-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (200 mg, 0.353 mmol, 1 equiv.) in DCM (5 mL) was added TFA (2 mL) at room temperature. The resulting mixture was stirred for 1 h at room temperature. The reaction was monitored by LCMS. The mixture was basified to pH 8 with saturated NaHCO_3 (aq.). The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30×150mm 5μm; Mobile Phase A: undefined, Mobile Phase B: undefined; Flow rate: 60 mL/min; Gradient: 25% B to 50% B in 8 min; 220 nm; Rt: 7.67 min) to afford 4-chloro-5-[2-[(2-hydroxyethyl)amino]-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (106.3 mg) as a white solid.

Example 11. Synthesis of Compounds 138 and 139



7-Benzyl-2-chloro-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine

To a stirred solution of 4-fluoro-2-(trifluoromethyl)phenol (1469.32 mg, 8.158 mmol, 1.20 equiv.) and 7-benzyl-2,4-dichloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (2000 mg, 6.799 mmol, 1 equiv.) in DMF (20 mL) was added K₂CO₃ (1879.20 mg, 13.597 mmol, 2 equiv.) at room temperature. The solution was stirred at 70°C for 0.5 hours. The mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μ m, 330 g; Mobile Phase A: Water (plus 5 mM TFA); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 70% B - 95% B gradient in 20 min; Detector: 254 nm. The fractions containing the desired product were collected at 95% B and concentrated under reduced pressure to afford 7-benzyl-2-chloro-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (2331 mg, 78.31%) as an off-white solid.

7-Benzyl-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-one

A solution of 7-benzyl-2-chloro-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (2 g, 4.568 mmol, 1 equiv.) in HAc (10 mL, 174.515 mmol, 38.20 equiv.) and H₂O (1 mL, 55.508 mmol, 12.15 equiv.) was stirred for 10 hours at 140°C under N₂ atmosphere. The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE/EA 1/1) to afford 7-benzyl-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-one (530 mg, 27.67%) as a light yellow solid.

4-[4-Fluoro-2-(trifluoromethyl)phenoxy]-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-one

To a solution of 7-benzyl-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-one (530 mg, 1.264 mmol, 1 equiv.) in MeOH (10 mL, 246.989 mmol, 195.44 equiv.) was added Pd/C (268.98 mg, 2.528 mmol, 2 equiv.) under nitrogen atmosphere. The mixture was hydrogenated at room temperature for 4 hours under hydrogen atmosphere using a hydrogen balloon, filtered through a Celite pad and concentrated under reduced pressure. To afford 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-one (430 mg, 103.34%) as a light yellow solid.

4-Chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-oxo-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

A mixture of 4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-one (430 mg, 1.306 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (357.84 mg, 1.437 mmol, 1.1 equiv.) in DIEA (337.58 mg, 2.612 mmol, 2.00 equiv.) was stirred for 2 hours at 100°C under N₂ atmosphere. The residue was purified by Prep-TLC (PE/EA 1/1) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-oxo-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (210 mg, 29.67%) as a light yellow solid.

4-Chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1-methyl-2-oxo-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one and 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-methoxy-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

To a solution of 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-oxo-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (90 mg, 0.166 mmol, 1 equiv.) and NaHCO₃ (27.90 mg, 0.332 mmol, 2 equiv.) in DMF (10 mL, 129.218 mmol, 778.02 equiv.) was added CH₃I (47.15 mg, 0.332 mmol, 2.00 equiv.) dropwise at 0°C under nitrogen atmosphere. The mixture was stirred at 25°C for 16 hours. The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE/EA 0/1) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1-methyl-2-oxo-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (60 mg, 64.99%) and 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-methoxy-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (15 mg) as a light yellow solid.

4-Chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1-methyl-2-oxo-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one


To a solution of 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1-methyl-2-oxo-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (60 mg, 0.108 mmol, 1 equiv.) in DCM (10 mL, 157.300 mmol, 1457.41 equiv.) was added TFA (123.07 mg, 1.079 mmol, 10 equiv.) at 25°C. The resulting mixture was concentrated under reduced pressure. The crude product (100 mg) was purified by Prep-HPLC with the following

conditions (Column: XBridge Shield RP18 OBD Column 30*150mm,5um ; Mobile Phase A: Water (10 mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 20% B to 40% B in 7 min; 220 nm; Rt: 6.63 min) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-1-methyl-2-oxo-1H,2H,5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (29.3 mg, 57.54%) as a white solid.

4-Chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-methoxy-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a solution of 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-methoxy-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (15 mg, 0.027 mmol, 1 equiv.) in DCM (5 mL, 78.650 mmol, 2914.83 equiv.) was added TFA (30.77 mg, 0.270 mmol, 10 equiv.) at 25°C. The resulting mixture was concentrated under reduced pressure. The crude product (20 mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Shield RP18 OBD Column 30*150mm,5um ; Mobile Phase A: Water (10 mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 20% B to 40% B in 7 min; 220 nm; Rt: 6.63 min) to afford 4-chloro-5-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-2-methoxy-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (7.5 mg, 58.91%) as a white solid.

Example 12. Synthesis of Compound 110



Tert-Butyl 2-chloro-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 2,4-dichloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (2 g, 6.58 mmol, 1 equiv.) and 2-(trifluoromethyl)phenol (1.6 g, 9.86 mmol, 1.5 equiv.) in acetonitrile (20 mL) was added DBU (2.0 g, 13.15 mmol, 2 equiv.) at room temperature. The solution was stirred at rt for 4 hours. The mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (PE/EtOAc 10:1) to afford tert-butyl 2-chloro-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (700 mg, 24.77%) as colorless oil.

Tert-Butyl 2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a solution of tert-butyl 2-chloro-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1 g, 2.327 mmol, 1 equiv.) in MeOH (20 mL, 493.978 mmol, 212.32 equiv.) was added NaOMe (0.25 g, 0.005 mmol, 2 equiv.) at 25°C. The mixture was stirred at 25°C for 4 hours. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EA (10/1 to 1/1) to afford tert-butyl 2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (100 mg, 10.10%) as a light yellow solid.

2-Methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine

To a solution of tert-butyl 2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (100 mg, 0.235 mmol, 1 equiv.) in DCM (10 mL) was added TFA (268.03 mg, 2.351 mmol, 10 equiv.) at 25°C. The solution was stirred at 25°C for 4 hours. The resulting mixture was concentrated under reduced pressure. The crude product (150 mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Shield RP18 OBD Column 30*150mm,5um ; Mobile Phase A: Water (10 mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 20% B to 40% B in 7 min; 220 nm; Rt: 6.63 min) to afford 2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (80 mg, 104.62%) as a light yellow solid.

4-Chloro-5-[2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


A solution of 2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (80 mg, 0.246 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (61.26 mg, 0.246 mmol, 1 equiv.) in DIEA (63.57 mg, 0.492 mmol, 2.00 equiv.) was stirred for 2

hours at 100°C under nitrogen atmosphere. The residue was purified by silica gel column chromatography, eluted with PE/EA (5/1 to 1/1) to afford 4-chloro-5-[2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (120 mg, 90.71%) as a light yellow solid.

4-Chloro-5-[2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a solution of 4-chloro-5-[2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (120 mg, 0.223 mmol, 1 equiv.) in DCM (5 mL, 78.650 mmol, 352.56 equiv.) was added TFA (254.36 mg, 2.231 mmol, 10.00 equiv.) at 25°C. The resulting mixture was concentrated under reduced pressure. The crude product (150 mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Shield RP18 OBD Column 30*150mm, 5µm; Mobile Phase A: Water (10 mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 20% B to 40% B in 7 min; 220 nm; Rt: 6.63 min) to afford 4-chloro-5-[2-methoxy-4-[2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (24.1 mg, 23.81%) as a white solid.

Example 13. Synthesis of Compound 108



Tert-Butyl 4-(3-bromo-2-chlorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 1.854 mmol, 1 equiv.) and 3-bromo-2-chlorophenol (461.46 mg, 2.224 mmol, 1.20

equiv.) in DMF (10 mL) was added K_2CO_3 (512.38 mg, 3.707 mmol, 2 equiv.). The resulting mixture was stirred for 1h at 70°C. The mixture was purified by reverse flash chromatography with the following conditions: Column: (spherical C18, 20-40 μ m, 330g; Mobile Phase A: Water (5mM NH_4HCO_3), Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 20% B to 60% B in 55 min; 254 nm). The fractions containing the desired product were collected at 40% B and concentrated under reduced pressure. This resulted in tert-butyl 4-(3-bromo-2-chlorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (300 mg, 36.72%) as an off-white solid.

Tert-Butyl 4-(2-chloro-3-cyanophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 4-(3-bromo-2-chlorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (450 mg, 1.021 mmol, 1 equiv.) and zinc dicarbonitrile (143.87 mg, 1.225 mmol, 1.20 equiv.) in DMF (5 mL) was added $Pd(PPh_3)_4$ (117.99 mg, 0.102 mmol, 0.1 equiv.). The resulting mixture was stirred for 2 hours at 120°C under nitrogen atmosphere. The residue was purified by reverse flash chromatography with the following conditions: Column: spherical C18, 20-40 μ m, 180g; Mobile Phase A: Water (5mM NH_4HCO_3), Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 10% B to 60% B in 55 min; 254 nm. The fractions containing the desired product were collected at 40% B and concentrated under reduced pressure. This resulted in tert-butyl 4-(2-chloro-3-cyanophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (280 mg, 70.89%) as a light yellow solid.

2-Chloro-3-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yloxy]benzonitrile

To a stirred solution of tert-butyl 4-(2-chloro-3-cyanophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (100 mg, 0.259 mmol, 1 equiv.) in DCM (3 mL) was added TFA (1 mL). The resulting mixture was stirred for 2 hours at room temperature under air atmosphere. The resulting mixture was concentrated under reduced pressure. The mixture was basified to pH 7 with saturated NH_4HCO_3 (aq.). The mixture was purified by reverse flash chromatography with the following conditions: Column: spherical C18, 20-40 μ m, 180g; Mobile Phase A: Water (5mM NH_4HCO_3), Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 30% B to 60% B in 30 min; 254 nm). The fractions containing the desired product were collected at 45% B and concentrated under reduced pressure. This resulted in 2-chloro-3-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yloxy]benzonitrile (60 mg, 80.95%) as a light yellow oil.


2-Chloro-3-([7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy)benzonitrile

To a stirred solution of tert-butyl 4-(2-chloro-3-cyanophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (60 mg, 0.155 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (38.63 mg, 0.155 mmol, 1.00 equiv.) in DIEA (40.09 mg, 0.310 mmol, 2 equiv). The resulting mixture was stirred for hours at 100°C under air atmosphere. The residue was purified by Prep-TLC (PE/EtOAc 1:1) to afford 2-chloro-3-([7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy)benzonitrile (50 mg, 64.56%) as a light yellow solid.

2-Chloro-3-[[7-(5-chloro-6-oxo-1,6-dihydropyridazin-4-yl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy]benzonitrile

To a stirred solution of 2-chloro-3-([7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy)benzonitrile (50 mg, 0.100 mmol, 1 equiv.) in DCM (3 mL) was added TFA (1 mL). The resulting mixture was stirred for 2 hours at room temperature. The resulting mixture was concentrated under reduced pressure. The mixture was basified to pH 7 with saturated NH_4CO_3 (aq.). The crude product was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30×150mm 5μm; Mobile Phase A: Water (10mM NH_4HCO_3), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 20% B to 42% B in 8 min; 220 nm; Rt: 7.58 min) to afford 2-chloro-3-[[7-(5-chloro-6-oxo-1,6-dihydropyridazin-4-yl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy]benzonitrile (14.5 mg, 34.88%) as an off-white solid.

Example 14. Synthesis of Compound 111



Tert-Butyl 4-[3-bromo-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred mixture of tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (180 mg, 0.667 mmol, 1 equiv.) and 3-bromo-2-(trifluoromethyl)phenol (241.25 mg, 1.001 mmol, 1.50 equiv.) in DMF (10 mL) was added Cs₂CO₃ (434.86 mg, 1.335 mmol, 2.00 equiv.) in portions at rt under nitrogen atmosphere. The resulting mixture was stirred for 0.5 hours at 70°C under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to cool down to rt. The resulting mixture was concentrated under reduced pressure. The resulting mixture was extracted with EtOAc (3 x 200 mL). The combined organic layers were washed with brine (2 x 100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse flash chromatography with the following conditions (Column, C18 silica gel; mobile phase, acetonitrile in water, 40% to 85% gradient in 30 min; detector, UV 220 nm) to afford tert-butyl 4-[3-bromo-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (150 mg, 47.39%) as a yellow oil.

Tert-Butyl 4-[3-cyano-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred mixture of tert-butyl 4-[3-bromo-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (150 mg, 0.316 mmol, 1 equiv.) and Zn(CN)₂ (111.43 mg, 0.949 mmol, 3.00 equiv.) in DMF (8 mL) was added Pd(PPh₃)₄ (36.55 mg, 0.032 mmol, 0.1 equiv.) in portions at rt under nitrogen atmosphere. The final reaction mixture was irradiated with microwave radiation for 3 hours at 150°C. The reaction was monitored by LCMS. The residue was purified by reverse flash chromatography with the following conditions (Column, C18 silica gel; mobile phase, acetonitrile in water, 40% to 95% gradient in 30 min; detector, UV 220 nm) to afford tert-butyl 4-[3-cyano-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (70 mg, 52.65%) as a yellow oil.

3-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yloxy]-2-(trifluoromethyl)benzonitrile

To a stirred solution of tert-butyl 4-[3-cyano-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (70 mg) in DCM (10 mL) was added TFA (1 mL) dropwise at rt. The reaction mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified

to pH=8 with saturated NH_4HCO_3 (aq.). The resulting mixture was extracted with DCM (3 x 100 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse flash chromatography with the following conditions (Column, C18 silica gel; mobile phase, acetonitrile in water, 30% to 60% gradient in 20 min; detector, UV 220 nm) to afford 3-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yloxy]-2-(trifluoromethyl)benzonitrile (40 mg) as a yellow oil.

3-([7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy)-2-(trifluoromethyl)benzonitrile


Into a 25 mL round-bottom flask were added 3-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yloxy]-2-(trifluoromethyl)benzonitrile (40 mg, 0.125 mmol, 1 equiv), 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (62.22 mg, 0.250 mmol, 2.00 equiv.) and DIEA (48.42 mg, 0.375 mmol, 3.00 equiv.) at rt under nitrogen atmosphere. The resulting mixture was stirred for 16 hours at 90°C under nitrogen atmosphere. The residue was purified by Prep-TLC (PE/EtOAc=5/1) to afford 3-([7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy)-2-(trifluoromethyl)benzonitrile (50 mg, 75.12%) as a yellow oil.

3-[[7-(5-Chloro-6-oxo-1,6-dihydropyridazin-4-yl)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy]-2-(trifluoromethyl)benzonitrile

To a stirred solution of 3-([7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy)-2-(trifluoromethyl)benzonitrile (50 mg) in DCM (10 mL) was added TFA (1 mL) dropwise at rt. The reaction mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH_4HCO_3 (aq.). The resulting mixture was extracted with DCM (3 x 100 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30x150mm 5um; Mobile Phase A: Water (10mM NH_4HCO_3), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 25% B to 45% B in 8 min; 220 nm; Rt: 7.07 min) to afford 3-[[7-(5-chloro-6-oxo-1,6-dihydropyridazin-4-yl)-

5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]oxy]-2-(trifluoromethyl)benzonitrile (10.8 mg) as a white solid.

Example 15. Synthesis of Compounds 126 and 126a



N-[(1E)-1-[4-fluoro-2-(trifluoromethyl)phenyl]ethylidene]-4-methylbenzene-1-sulfonohydrazide

To a stirred solution of 1-[4-fluoro-2-(trifluoromethyl)phenyl]ethan-1-one (2 g, 9.702 mmol, 1 equiv.) in EtOH (40 mL) was added 4-methylbenzene-1-sulfonohydrazide (1.81 g, 9.719 mmol, 1.00 equiv.) in portions at rt under nitrogen atmosphere. The resulting mixture was stirred for 6 hours at 90°C under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to cool down to rt. The resulting mixture was concentrated under vacuum. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μ m, 330 g; Mobile Phase A: Water (plus 5 mM AcOH); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 45% B - 70% B gradient in 20 min; Detector: 220 nm. The fractions containing the desired product were collected at 60% B and concentrated under reduced pressure to afford N-[(1E)-1-[4-fluoro-2-

(trifluoromethyl)phenyl]ethylidene]-4-methylbenzene-1-sulfonohydrazide (2.5 g, 68.83%) as a white solid.

Tert-Butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethenyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred mixture of tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (750 mg, 2.781 mmol, 1 equiv.) and N-[(1E)-1-[4-fluoro-2-(trifluoromethyl)phenyl]ethylidene]-4-methylbenzene-1-sulfonohydrazide (2081.80 mg, 5.561 mmol, 2.00 equiv.) in 1,4-dioxane (20 mL) were added Pd(acetonitrile)₂Cl₂ (72.14 mg, 0.278 mmol, 0.10 equiv), Dppf (307.18 mg, 0.556 mmol, 0.2 equiv.) and t-BuOLi (489.71 mg, 6.117 mmol, 2.20 equiv.) in portions at rt under nitrogen atmosphere. The final reaction mixture was irradiated with microwave radiation for 2 hours at 100°C. The reaction was monitored by LCMS. The mixture was allowed to cool down to rt. The resulting mixture was filtered, the filter cake was washed with EtOAc (2x50 mL). The filtrate was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 um, 330 g; Mobile Phase A: Water (plus 5 mM AcOH); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 50% B - 90% B gradient in 30 min; Detector: 220 nm. The fractions containing the desired product were collected at 85% B and concentrated under reduced pressure to afford tert-butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethenyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (800 mg, 67.95%) as a brown oil.

Tert-Butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a solution of tert-butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethenyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (150 mg) in 30 mL MeOH was added Pd/C (10%, 30 mg) under nitrogen atmosphere in a 100 mL round-bottom flask. The mixture was hydrogenated at room temperature for 4 hours under hydrogen atmosphere using a hydrogen balloon, filtered through a celite pad and concentrated under reduced pressure. This resulted in tert-butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (150 mg) as a yellow oil.

4-[1-[4-Fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine

To a stirred solution of tert-butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (150 mg) in DCM (10 mL) was added TFA (1 mL)

dropwise at rt. The reaction mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH_4HCO_3 (aq.). The resulting mixture was extracted with DCM (3 x 50 mL). The combined organic layers were washed with brine (1x30 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μm , 120 g; Mobile Phase A: Water (plus 5 mM AcOH); Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 5% - 5% B, 10 min, 40% B - 58% B gradient in 15 min; Detector: 254 nm. The fractions containing the desired product were collected at 53% B and concentrated under reduced pressure to afford 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (100 mg) as a yellow oil.

4-chloro-5-(4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


Into a 50 mL round-bottom flask were added 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (100 mg, 0.307 mmol, 1 equiv), 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (91.88 mg, 0.369 mmol, 1.20 equiv.) and DIEA (119.19 mg, 0.922 mmol, 3.00 equiv.) at rt under nitrogen atmosphere. The resulting mixture was stirred for 2 hours at 90°C under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to cool down to rt. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μm , 120 g; Mobile Phase A: Water (plus 5 mM AcOH); Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 5% - 5% B, 10 min, 40% B - 60% B gradient in 15 min; Detector: 220 nm. The fractions containing the desired product were collected at 53% B and concentrated under reduced pressure to afford 4-chloro-5-(4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (120 mg, 72.57%) as a yellow oil.

4-Chloro-5-[4-[(1S)-1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one and 4-chloro-5-[4-[(1R)-1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-(4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (200 mg)

in DCM (10 mL) was added TFA (1 mL) dropwise at rt. The reaction mixture was stirred for 4 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH_4HCO_3 (aq.). The resulting mixture was extracted with DCM (3 x 100 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by Chiral-Prep-HPLC with the following conditions (Column: XBridge Prep Phenyl OBD Column 19x150mm 5um 13nm; Mobile Phase A: Mobile Phase B:Flow rate: 60 mL/min; Gradient: 20% B to 37% B in 8 min; 220 nm; Rt: 7.97 min). Although the two isomers were separated by this technique, the absolute orientation was not determined. The compound designated as 4-chloro-5-[4-[(1S)-1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (11.8 mg) was obtained at 1.819 min as an off-white solid. The compound designated as 4-chloro-5-[4-[(1R)-1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (13.5 mg) was obtained at 2.470 min as a white solid.

Example 16. Synthesis of Compound 133



tert-Butyl 4-[methyl[(3R,4R)-4-methylpiperidin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

Into a 25 mL round-bottom flask were added (3R,4R)-1-benzyl-N,4-dimethylpiperidin-3-amine (2.43 g, 0.011 mmol, 1.50 equiv.) and tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-

d]pyrimidine-7-carboxylate (2 g, 0.007 mmol, 1 equiv.) at room temperature. To the mixture was added DIEA (1.92 g, 0.015 mmol, 2.00 equiv.) at rt. The mixture was stirred at 100°C for 2 hours. The residue was purified by Prep-TLC (PE/EtOAc 1:1) to afford tert-butyl 4-[methyl[(3R,4R)-4-methylpiperidin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (670mg, 25.00%) as an off-white solid.

(3R,4R)-1-Benzyl-N,4-dimethyl-N-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]piperidin-3-amine

To a stirred solution of tert-butyl 4-[[[(3R,4R)-1-benzyl-4-methylpiperidin-3-yl](methyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (413 mg, 0.914 mmol, 1 equiv.) in DCM (10 mL) was added trifluoroacetic acid (3 mL, 0.026 mmol, 6.00 equiv.) dropwise at 0°C. The mixture was stirred for 2 hours at room temperature. The reaction was monitored by LCMS. The solution was concentrated under reduced pressure. The crude product (362mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Shield RP18 OBD Column, 5µm, 19*150mm; Mobile Phase A: Water (10mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B to 80% B in 25 min; 220 nm; Rt: 21.65 min) to afford (3R,4R)-1-benzyl-N,4-dimethyl-N-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]piperidin-3-amine (250 mg, 77.77%) as red oil.

5-(4-[[[(3R,4R)-1-Benzyl-4-methylpiperidin-3-yl](methyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


Into a 25 mL round-bottom flask were added (3R,4R)-1-benzyl-N,4-dimethyl-N-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]piperidin-3-amine (263 mg, 0.748 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (186.38 mg, 0.748 mmol, 1.00 equiv.) at room temperature. To the mixture was added DIEA (193.41 mg, 1.261 mmol, 2 equiv.) at rt. The mixture was stirred for 2 hours at 100°C. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 µm, 330 g; Mobile Phase A: Water (plus 5 mM NH₄HCO₃); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 45% B - 95% B gradient in 30 min; Detector: 254 nm. The fractions containing the desired product were collected at 85% B and concentrated under reduced pressure to afford 5-(4-[[[(3R,4R)-1-benzyl-4-methylpiperidin-3-yl](methyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (245mg, 58.04%) as an off-white solid.

5-(4-[[[(3R,4R)-1-Benzyl-4-methylpiperidin-3-yl](methyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2,3-dihydropyridazin-3-one

To a stirred solution of 5-(4-[[[(3R,4R)-1-benzyl-4-methylpiperidin-3-yl](methyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (88 mg, 1 equiv.) in DCM (10 mL) was added trifluoroacetic acid (3 mL, 0.026 mmol, 6.00 equiv.) dropwise at 0°C. The mixture was stirred for 2 hours at room temperature. The solution was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μ m, 330 g; Mobile Phase A: Water (plus 5 mM TFA); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 33% B - 95% B gradient in 30 min; Detector: 254 nm. The fractions containing the desired product were collected at 90% B and concentrated under reduced pressure to afford 5-(4-[[[(3R,4R)-1-benzyl-4-methylpiperidin-3-yl](methyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-4-chloro-2,3-dihydropyridazin-3-one (33.5 mg, 44.74%) as an off-white solid.

Compound 133a was prepared by the methods and scheme described in this example by using (3S,4S)-1-benzyl-N,4-dimethylpiperidin-3-amine in place of (3R,4R)-1-benzyl-N,4-dimethylpiperidin-3-amine.

Example 17. Synthesis of Compound 136



Tert-Butyl 4-[[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

A mixture of 4-fluoro-2-(trifluoromethyl)aniline (6.64 g, 37.074 mmol, 2 equiv), tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (5 g, 18.537 mmol, 1 equiv),

Pd(AcO)₂ (0.83 g, 3.707 mmol, 0.2 equiv), XantPhos (4.29 g, 7.415 mmol, 0.4 equiv.) and Cs₂CO₃ (12.08 g, 37.074 mmol, 2 equiv.) in 1,4-dioxane (80 mL) was stirred at 110°C for 16 hours. The reaction mixture was filtered and the filtrate was concentrated to give the crude product which was purified by silica gel column chromatography, eluted with PE:EA (20:1 to 1:2) to afford tert-butyl 4-[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (5.6 g, 73.26%) as a white solid.

Tert-Butyl 4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-methoxy-2-oxoethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred mixture of tert-butyl 4-[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (3 g, 7.275 mmol, 1 equiv.) and methyl 2-bromoacetate (2.23 g, 14.578 mmol, 2.00 equiv.) in DMF (30 mL) was added Cs₂CO₃ (4.74 g, 14.548 mmol, 2.00 equiv.) in portions at rt under nitrogen atmosphere. The resulting mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was extracted with EtOAc (3 x 400 mL). The combined organic layers were washed with brine (2x 200 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 um, 330 g; Mobile Phase A: Water (plus 5 mM TFA); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 55% B - 85% B gradient in 30 min; Detector: 220 nm. The fractions containing the desired product were collected at 79% B and concentrated under reduced pressure to afford tert-butyl 4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-methoxy-2-oxoethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 14.19%) as a yellow solid.

Tert-Butyl 4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-hydroxyethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-methoxy-2-oxoethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 1.032 mmol, 1 equiv.) in THF (50 mL) was added LiAlH₄ (78.34 mg, 2.064 mmol, 2.00 equiv.) in portions at -30°C under nitrogen atmosphere. The reaction mixture was stirred for 16 hours at rt. The reaction was monitored by LCMS. The reaction was quenched by the addition of Water (1 mL) at -30°C. The precipitated solids were collected by filtration and washed with MeOH (3x30 mL). The resulting mixture was concentrated under vacuum. The residue was purified by Prep-TLC

(PE/EA = 1/1) to afford tert-butyl 4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-hydroxyethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (100 mg, 21.23%) as a yellow oil.

4-[1-[4-Fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine

To a stirred solution of tert-butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (150 mg) in DCM (10 mL) was added TFA (1 mL) dropwise at rt. The reaction mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH_4HCO_3 (aq.). The resulting mixture was extracted with DCM (3 x 50 mL). The combined organic layers were washed with brine (1x30 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μm , 120 g; Mobile Phase A: Water (plus 5 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 40% B - 58% B gradient in 15 min; Detector: 220 nm. The fractions containing the desired product were collected at 53% B and concentrated under reduced pressure to afford 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (100 mg) as a yellow oil.

4-Chloro-5-(4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-hydroxyethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


Into a 50 mL round-bottom flask were added 2-[[4-fluoro-2-(trifluoromethyl)phenyl]([5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl)]amino]ethan-1-ol (40 mg, 0.112 mmol, 1 equiv), 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (55.92 mg, 0.224 mmol, 2.00 equiv.) and DIEA (43.53 mg, 0.337 mmol, 3.00 equiv.) at rt under nitrogen atmosphere. The resulting mixture was stirred for 2 hours at 90°C under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to cool down to rt. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 μm , 120 g; Mobile Phase A: Water (plus 5 mM NH_4HCO_3); Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 5% - 5% B, 10 min, 40% B - 60% B gradient in 15 min; Detector: 220 nm. The fractions containing the desired product were collected at 55% B and concentrated under reduced pressure to afford 4-chloro-5-(4-[[4-fluoro-2-

(trifluoromethyl)phenyl](2-hydroxyethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (50 mg, 78.28%) as a yellow oil.

4-Chloro-5-(4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-hydroxyethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-(4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-hydroxyethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (50 mg) in DCM (10 mL) was added TFA (1 mL) dropwise at rt. The reaction mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH_4HCO_3 (aq.). The resulting mixture was extracted with EtOAc (2 x 50 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30x150mm 5um; Mobile Phase A: undefined, Mobile Phase B: undefined; Flow rate: 60 mL/min; Gradient: 30% B to 45% B in 8 min; 220 nm; Rt: 7.6 min) to afford 4-chloro-5-(4-[[4-fluoro-2-(trifluoromethyl)phenyl](2-hydroxyethyl)amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2,3-dihydropyridazin-3-one (6.2 mg) as a white solid.

Example 18. Synthesis of Compound 132



Tert-Butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of t-BuONa (226.97 mg, 2.362 mmol, 2.00 equiv.) in DMSO (20 mL) was added Me₃SiI (472.57 mg, 2.362 mmol, 2.00 equiv.) in portions at 40°C under nitrogen atmosphere. The resulting mixture was stirred for 0.5 hours at 40°C under nitrogen atmosphere. Then tert-butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]ethenyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 1.181 mmol, 1 equiv.) in DMSO (5 mL) was dropwise at rt under nitrogen atmosphere. The resulting mixture was stirred for 1 hours at rt under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to cool down to rt. The resulting mixture was extracted with EtOAc (3 x 100 mL). The combined organic layers were washed with brine (2x 100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 um, 330 g; Mobile Phase A: Water (plus 5 mM NH₄HCO₃); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 55% B - 80% B gradient in 25 min; Detector: 220 nm. The fractions containing the desired product were collected at 73% B and concentrated under reduced pressure to afford tert-butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (240 mg, 46.46%) as a yellow oil.

4-[1-[4-Fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine

To a stirred solution of tert-butyl 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (240 mg, 0.549 mmol, 1 equiv.) in DCM (10 mL) was added TFA (1 mL, 13.463 mmol, 24.54 equiv.) dropwise at rt. The reaction mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH₄HCO₃ (aq.). The resulting mixture was extracted with EtOAc (2 x 100 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 um, 120 g; Mobile Phase A: Water (plus 5 mM AcOH); Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 5% - 5% B, 10 min, 33% B - 45% B gradient in 20 min; Detector: 254 nm. The fractions containing the desired product were collected at 40% B and concentrated under

reduced pressure to afford 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (150 mg, 81.05%) as a yellow oil.

4-Chloro-5-(4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


Into a 50 mL round-bottom flask were added 4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (150 mg, 0.445 mmol, 1 equiv), 4,5-dichloro-2-(oxan-2-yl)-1,2,3,6-tetrahydropyridazin-3-one (134.00 mg, 0.534 mmol, 1.20 equiv.) and DIEA (172.42 mg, 1.334 mmol, 3.00 equiv.) at rt under nitrogen atmosphere. The resulting mixture was stirred for 2 hours at 90°C under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to cool down to rt. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20 - 40 um, 330 g; Mobile Phase A: Water (plus 5 mM NH₄HCO₃); Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% - 5% B, 10 min, 40% B - 60% B gradient in 15 min; Detector: 220 nm. The fractions containing the desired product were collected at 54% B and concentrated under reduced pressure to afford 4-chloro-5-(4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (200 mg, 81.78%) as a yellow oil.

4-Chloro-5-(4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-(4-[1-[4-fluoro-2-(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (200 mg) in DCM (10 mL) was added TFA (2 mL) dropwise at rt. The reaction mixture was stirred for 2 hours at rt. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH=8 with saturated NH₄HCO₃ (aq.). The resulting mixture was extracted with EtOAc (2 x 100 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30x150mm 5um; Mobile Phase A: undefined, Mobile Phase B: undefined; Flow rate: 60 mL/min; Gradient: 30% B to 55% B in 8 min; 220 nm; Rt: 7.232 min) to afford 4-chloro-5-(4-[1-[4-fluoro-2-

(trifluoromethyl)phenyl]cyclopropyl]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2,3-dihydropyridazin-3-one (39.2 mg) as an off-white solid.

Example 19. Synthesis of Compound 109



Tert-Butyl 4-(2-bromo-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 1.854 mmol, 1 equiv.) and 2-bromo-3-fluorophenol (424.87 mg, 2.224 mmol, 1.20 equiv.) in DMF (10 mL) was added K_2CO_3 (512.38 mg, 3.707 mmol, 2 equiv.). The resulting mixture was stirred for 0.5 hours at 70°C. The mixture was allowed to cool down to room temperature. The reaction was quenched with Water at room temperature. The resulting mixture was extracted with EtOAc (2 x 100 mL). The combined organic layers were washed with brine (2x100 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (5:1) to afford tert-butyl 4-(2-bromo-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 63.58%) as a white solid.

Tert-Butyl 4-(2-ethenyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a solution of tert-butyl 4-(2-bromo-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 1.178 mmol, 1 equiv.) and pentamethyl-1,3,2-dioxaborolane (334.72 mg, 2.357 mmol, 2.00 equiv.) in H₂O (2 mL) and 1,4-dioxane (16 mL) were added K₂CO₃ (325.75 mg, 2.357 mmol, 2 equiv.) and Pd(PPh₃)₄ (68.09 mg, 0.059 mmol, 0.05 equiv). After stirring for overnight at 90°C under a nitrogen atmosphere, the resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (5:1) to afford tert-butyl 4-(2-ethenyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (250 mg, 57.12%) as a yellow oil.

Tert-Butyl 4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred solution of tert-butyl 4-(2-ethenyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (250 mg, 0.673 mmol, 1 equiv.) in MeOH (10 mL) was added Pd/C (100 mg, 0.940 mmol, 1.40 equiv). The resulting mixture was stirred for 2 hours at RT under hydrogen atmosphere. The resulting mixture was filtered, the filter cake was washed with MeOH (2x10 mL). The filtrate was concentrated under reduced pressure. The resulting mixture was concentrated under reduced pressure. This resulted in tert-butyl 4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (210 mg, 0.08%) as a black oil.

4-(2-Ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine


To a stirred solution of tert-butyl 4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (210 mg, 0.562 mmol, 1 equiv.) in DCM (3 mL) was added TFA (1 mL). The resulting mixture was stirred for 2 hours at room temperature under air atmosphere. The resulting mixture was concentrated under reduced pressure. The mixture was basified to pH 8 with saturated NH₄HCO₃ (aq.). The mixture was purified by reverse flash chromatography with the following conditions: Column: spherical C18, 20-40 µm, 180g; Mobile Phase A: Water (5mM NH₄HCO₃), Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 25% B to 60% B in 40 min; 254 nm). The fractions containing the desired product were collected at 40% B and concentrated under reduced pressure. This resulted in 4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (120 mg, 78.07%) as a light yellow oil.

4-Chloro-5-[4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one

To a stirred solution of 4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine (120 mg, 0.439 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (109.37 mg, 0.439 mmol, 1.00 equiv.) in DIEA (113.49 mg, 0.878 mmol, 2 equiv). The resulting mixture was stirred for 2 hours at 100°C under air atmosphere. The residue was purified by Prep-TLC (PE/EtOAc 1:1) to afford 4-chloro-5-[4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (100 mg, 46.87%) as a light yellow solid.

4-Chloro-5-[4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-[4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (100 mg, 0.206 mmol, 1 equiv.) in DCM (3 mL) was added TFA (1 mL). The resulting mixture was stirred for 2 hours at room temperature. The resulting mixture was concentrated under reduced pressure. The mixture was basified to pH 7 with saturated NH_4HCO_3 (aq.). The crude product was purified by Prep-HPLC with the following conditions (Column: XBridge Prep OBD C18 Column 30×150mm 5μm; Mobile Phase A: Water (10mM NH_4HCO_3), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 30% B to 50% B in 8 min; 220 nm; Rt: 7.27 min) to afford 4-chloro-5-[4-(2-ethyl-3-fluorophenoxy)-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2,3-dihydropyridazin-3-one (41.6 mg, 50.31%) as a white solid.

Example 20. Synthesis of Compound 127**Methyl 3-(methylamino)pyridine-4-carboxylate**

To a stirred solution of 3-(methylamino)pyridine-4-carboxylic acid (11 g, 72.296 mmol, 1 equiv.) in MeOH (500 mL, 12349.455 mmol, 170.82 equiv.) was added SOCl₂ (43.01 g, 361.478 mmol, 5 equiv.) dropwise at 0°C. The resulting mixture was stirred for 30 hours at 70°C. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under reduced pressure. The residue was dissolved in ethyl acetate (50 mL). The mixture basified to pH 8 with saturated NaHCO₃ (aq.). The resulting mixture was extracted with EtOAc (2 x 20 mL). The combined organic layers were washed with brine (1x30 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure to afford methyl 3-(methylamino)pyridine-4-carboxylate (9 g, crude) as a yellow solid.

Methyl 3-(N-methylacetamido)pyridine-4-carboxylate

To a stirred solution of methyl 3-(methylamino)pyridine-4-carboxylate (9 g, 54.158 mmol, 1 equiv.) in DCM (100 mL) were added Pyridine (21.42 g, 270.791 mmol, 5 equiv.) and acetyl chloride (6.38 g, 81.237 mmol, 1.5 equiv.) dropwise at room temperature. The resulting mixture was stirred for 2 hours at room temperature. The reaction was monitored by LCMS. The solution was basified to pH 8 with saturated NaHCO₃ (aq.). The resulting mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash with the following

conditions (Column: C18 Column 330 g; Mobile Phase A: Water (10 mM NH_4HCO_3), Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 10% B to 30% B in 25 min; 254/220 nm) to afford methyl 3-(N-methylacetamido)pyridine-4-carboxylate (8 g, 70.94%) as a brown liquid.

4-Hydroxy-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one

To a stirred solution of methyl 3-(N-methylacetamido)pyridine-4-carboxylate (6 g, 28.816 mmol, 1 equiv.) in dry 1,4-dioxane (100 mL) was added t-BuOK (6.47 g, 57.632 mmol, 2 equiv.) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 1 hours at 90°C under nitrogen atmosphere. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel column chromatography, eluted with DCM/MeOH (10:1) to afford 4-hydroxy-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one (4.5 g, 88.64%) as a orange solid.

4-Chloro-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one

To a stirred solution of 4-hydroxy-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one (4.5 g, 25.543 mmol, 1 equiv.) in dry 1,4-dioxane (100 mL) was added POCl_3 (3.92 g, 25.543 mmol, 1 equiv.) dropwise at room temperature. The resulting mixture was stirred for 16 hours at 90°C. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with DCM/MeOH (10:1) to afford 4-chloro-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one (2 g, 40.23%) as a red solid.

4-[[4-Fluoro-2-(trifluoromethyl)phenyl]amino]-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one

To a stirred solution of 4-chloro-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one (0.8 g, 4.111 mmol, 1 equiv.) in dry 1,4-dioxane (15 mL) were added Cs_2CO_3 (2.68 g, 8.221 mmol, 2 equiv), 4-fluoro-2-(trifluoromethyl)aniline (1.47 g, 8.221 mmol, 2.00 equiv), XantPhos (0.95 g, 1.644 mmol, 0.4 equiv.) and $\text{Pd}(\text{AcO})_2$ (0.18 g, 0.822 mmol, 0.2 equiv.) at room temperature under nitrogen atmosphere. The final reaction mixture was irradiated with microwave radiation for 4 hours at 110°C. The reaction was monitored by LCMS. The resulting mixture was extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine (1x100 mL), dried over anhydrous Na_2SO_4 . After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by reverse phase flash with the following conditions (Column: C18

Column 330 g; Mobile Phase A: Water (10 mM AcOH), Mobile Phase B: acetonitrile; Flow rate: 50 mL/min; Gradient: 20% B to 40% B in 40 min; 254/220 nm) to afford 4-[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one (1.1 g, 79.34%) as an off-white solid.

4-[[4-Fluoro-2-(trifluoromethyl)phenyl]amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one

To a stirred solution of 4-[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-1-methyl-1,2-dihydro-1,7-naphthyridin-2-one (1 g, 2.965 mmol, 1 equiv.) in THF (20 mL) was added PtO₂ (67.33 mg, 0.296 mmol, 0.10 equiv.) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 16 hours at room temperature under hydrogen atmosphere. The reaction was monitored by LCMS. The resulting mixture was filtered, the filter cake was washed with EtOAc (3x20 mL). The filtrate was concentrated under reduced pressure. The resulting mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash with the following conditions (Column: C18 Column 330 g; Mobile Phase A: Water (10 mM AcOH), Mobile Phase B: acetonitrile; Flow rate: 80mL/min; Gradient: 5% B to 20% B in 40 min; 254/220 nm) The fractions containing the desired product were collected at 16% B and concentrated under reduced pressure to afford 4-[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one (750 mg, 74.11%) as an off-white solid.

7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-4-[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one

To a stirred mixture of 4-[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one (750 mg, 2.197 mmol, 1 equiv.) and 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (1.09 g, 4.395 mmol, 2 equiv.) was added DIPEA (568.00 mg, 4.395 mmol, 2 equiv.) at room temperature. The resulting mixture was stirred for 2 hours at 100°C. The reaction was monitored by LCMS. The residue was dissolved in DMF (10 mL). The solution was purified by reverse phase flash with the following conditions (Column: C18 Column 330 g; Mobile Phase A: Water (10 mM FA), Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 30% B to 50% B in 40 min; 254/220 nm). The fractions containing the desired product were collected at 44% B and concentrated under reduced pressure to afford 7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-4-[[4-fluoro-2-


(trifluoromethyl)phenyl]amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one (1 g, 82.15%) as a yellow oil.

7-[5-Chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-4-[[4-fluoro-2-(trifluoromethyl)phenyl](methyl)amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one

To a stirred solution of 7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-4-[[4-fluoro-2-(trifluoromethyl)phenyl]amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one (800 mg, 1.444 mmol, 1 equiv.) in DMF (20 mL) were added Cs₂CO₃ (0.94 g, 2.888 mmol, 2 equiv.) and MeI (614.96 mg, 4.333 mmol, 3 equiv.) at room temperature. The resulting mixture was stirred for 16 hours at room temperature. The reaction was monitored by LCMS. The mixture was purified by reverse phase flash with the following conditions (Column: C18 Column 120 g; Mobile Phase A: Water (10 mM AcOH), Mobile Phase B: acetonitrile; Flow rate: 60 mL/min; Gradient: 40% B to 60% B in 40 min; 254/220 nm). The fractions containing the desired product were collected at 49% B and concentrated under reduced pressure to afford 7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-4-[[4-fluoro-2-(trifluoromethyl)phenyl](methyl)amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one (80 mg, 9.75%) as a yellow oil.

7-(5-Chloro-6-oxo-1,6-dihydropyridazin-4-yl)-4-[[4-fluoro-2-(trifluoromethyl)phenyl](methyl)amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one

To a stirred solution of 7-[5-chloro-1-(oxan-2-yl)-6-oxo-1,6-dihydropyridazin-4-yl]-4-[[4-fluoro-2-(trifluoromethyl)phenyl](methyl)amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one (80 mg, 0.141 mmol, 1 equiv.) in DCM (4.5 mL) was added TFA (0.5 mL, 6.732 mmol, 31.07 equiv.) dropwise at room temperature. The resulting mixture was stirred for 2 hours at room temperature. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The residue was basified to pH 8 with saturated NaHCO₃ (aq.). The solution was purified by reverse phase flash to afford 7-(5-chloro-6-oxo-1,6-dihydropyridazin-4-yl)-4-[[4-fluoro-2-(trifluoromethyl)phenyl](methyl)amino]-1-methyl-1,2,5,6,7,8-hexahydro-1,7-naphthyridin-2-one (40mg, 58.69%) as a white solid.

Example 21. Synthesis of Compounds 135 and 137**Tert-butyl 2-chloro-4-(4-fluoro-2-(trifluoromethyl)phenoxy)-5,8-dihydropyrido[3,4-*d*]pyrimidine-7(6H)-carboxylate.**

To a solution of *tert*-butyl 2,4-dichloro-5H,6H,7H,8H-pyrido[3,4-*d*]pyrimidine-7-carboxylate (5 g, 16.44 mmol) in DMF (50 mL) were added 4-fluoro-2-(trifluoromethyl)phenol (4.44 g, 24.66 mmol) and added K_2CO_3 (3.41 g, 24.66 mmol) at room temperature. The resulting mixture was stirred for 1 hours at 70 °C. After cooling to room temperature. A filtration was performed and the filtrate was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions (Column: Spherical C18, 20~40 μ m, 330 g; Mobile Phase A: Water (plus 5 mM NH_4HCO_3 ; Mobile Phase B: acetonitrile; Flow rate: 80 mL/min; Gradient: 5% in 10 min, 35%B to 45%B in 10 min; Detector: 254 nm/220 nm. The fractions containing desired product were collected at 44% B and concentrated under reduced

pressure to afford tert-butyl 2-chloro-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (6.2 g, 85%) as a white solid.

Tert-butyl 4-(4-fluoro-2-(trifluoromethyl)phenoxy)-2-vinyl-5,8-dihydropyrido[3,4-d]pyrimidine-7(6H)-carboxylate.

To a solution of tert-butyl 2-chloro-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 1.12 mmol) in dioxane (10 mL) were added 2-ethenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (344 mg, 2.23 mmol) and H₂O (0.5 mL, 27.75 mmol) K₂CO₃ (309 mg, 2.23 mmol) and Pd(PPh₃)₄ (129 mg, 0.11 mmol). After stirring for 2 hours at 95°C under a nitrogen atmosphere, the resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC, eluted with 17% ethyl acetate in petroleum ether to afford tert-butyl 2-ethenyl-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (490 mg, 99%) as a light yellow solid.

Tert-butyl 2-(1,2-dihydroxyethyl)-4-(4-fluoro-2-(trifluoromethyl)phenoxy)-5,8-dihydropyrido[3,4-d]pyrimidine-7(6H)-carboxylate.

To a solution of tert-butyl 2-ethenyl-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (400 mg, 0.91 mmol) in DCM (20 mL) were added 4-hydroxy-4-methylmorpholin-4-ium (323 mg, 2.73 mmol) and K₂OsO₄·2H₂O (34 mg, 0.091 mmol) at room temperature. After stirring for additional 1 hour, the resulting mixture was concentrated under reduced pressure and the residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20~40 μ m, 120 g; Mobile Phase A: Water (plus 5 mM NH₄HCO₃; Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 5% B in 10 min, 45%B to 65%B in 15 min; Detector: 254 nm and 220 nm. The fractions containing desired product were collected at 64% B and concentrated under reduced pressure to afford tert-butyl 2-(1,2-dihydroxyethyl)-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (280 mg, 65%) as a white solid.

1-(4-(4-fluoro-2-(trifluoromethyl)phenoxy)-5,6,7,8-tetrahydropyrido[3,4-d]pyrimidin-2-yl)ethane-1,2-diol.

To a stirred solution of tert-butyl 2-(1,2-dihydroxyethyl)-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (280 mg, 0.59 mmol) in DCM (4 mL) was added TFA (1 mL) at room temperature. The resulting mixture was

stirred for 1 hours at room temperature. The resulting mixture was concentrated under vacuum. The residue was dissolved into DCM (50 mL) and washed with saturated aqueous NaHCO₃ (20 mL). the organic layer was separated out and dried over anhydrous sodium sulfate. After filtration, the filtrate was concentrated under reduced pressure and the residue was purified by Prep-TLC with 8% methanol in dichloromethane to afford 1-[4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-2-yl]ethane-1,2-diol (180 mg, 82%) as a brown solid.

4-chloro-5-(2-(1,2-dihydroxyethyl)-4-(4-fluoro-2-(trifluoromethyl)phenoxy))-5,8-dihydropyrido[3,4-d]pyrimidin-7(6H)-yl)-2-(tetrahydro-2H-pyran-2-yl)pyridazin-3(2H)-one.


To a stirred solution of 2-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yloxy]benzaldehyde (180 mg, 0.71 mmol) in DIEA (0.5 mL) was added 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (176 mg, 0.71 mmol) at room temperature. The resulting mixture was stirred for 1 hours at 90°C. After cooling to ambient temperature, the resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC, eluted with 8% methanol in dichloromethane to afford 4-chloro-5-(2-(1,2-dihydroxyethyl)-4-(4-fluoro-2-(trifluoromethyl)phenoxy))-5,8-dihydropyrido[3,4-d]pyrimidin-7(6H)-yl)-2-(tetrahydro-2H-pyran-2-yl)pyridazin-3(2H)-one (140 mg, 43%) as a brown solid.

(S)-4-chloro-5-(2-(1,2-dihydroxyethyl)-4-(4-fluoro-2-(trifluoromethyl)phenoxy))-5,8-dihydropyrido[3,4-d]pyrimidin-7(6H)-yl)pyridazin-3(2H)-one and (R)-4-chloro-5-(2-(1,2-dihydroxyethyl)-4-(4-fluoro-2-(trifluoromethyl)phenoxy))-5,8-dihydropyrido[3,4-d]pyrimidin-7(6H)-yl)pyridazin-3(2H)-one

To a solution of 4-chloro-5-[2-(1,2-dihydroxyethyl)-4-[4-fluoro-2-(trifluoromethyl)phenoxy]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl]-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (150 mg, 0.27 mmol) in DCM (4 mL) was added TFA (1 mL) at room temperature. The resulting mixture was stirred for 1 hours at room temperature. The resulting mixture was concentrated under reduced pressure. The residue was purified by reverse phase flash chromatography with the following conditions: Column: Spherical C18, 20~40 um, 120 g; Mobile Phase A: Water (plus 5 mM NH₄HCO₃); Mobile Phase B: acetonitrile; Flow rate: 45 mL/min; Gradient: 5% B in 10 min, 45%B to 65%B in 15 min; Detector: 254nm and 220 nm. The fractions containing desired product were collected at 64%B and concentrated under reduced pressure to afford the racemic

product (130 mg) which was separated by Prep-Chiral-HPLC with the following conditions: Column: XBridge Prep OBD C18 Column 30 × 150 mm, 5 μm; Mobile Phase A: Hexane, Mobile Phase B: EtOH; Flow rate: 20 mL/min; Gradient: 35% B in 10 min; Detector: 254/220 nm). Although the two isomers were separated by this technique, the absolute orientation was not determined. The fractions containing desired product were collected and concentrated under reduced pressure to afford the product: The compound designated as (S)-4-chloro-5-(2-(1,2-dihydroxyethyl)-4-(4-fluoro-2-(trifluoromethyl)phenoxy)-5,8-dihydropyrido[3,4-d]pyrimidin-7(6H)-yl)pyridazin-3(2H)-one: retention time (4.97 min) (49.5 mg, 39%) as a white solid and The compound designated as (R)-4-chloro-5-(2-(1,2-dihydroxyethyl)-4-(4-fluoro-2-(trifluoromethyl)phenoxy)-5,8-dihydropyrido[3,4-d]pyrimidin-7(6H)-yl)pyridazin-3(2H)-one: retention time (8.05 min) (45.7 mg, 36%) as a white solid.

Example 22. Synthesis of Compound 131



Tert-butyl 4-[[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred mixture of tert-butyl 4-chloro-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (500 mg, 1.854 mmol, 1 equiv.) and 4-(trifluoromethyl)pyridin-3-amine (601.03 mg, 3.707 mmol, 2.0 equiv.) in 1,4-dioxane (5 mL) were added Pd(AcO)₂ (83.24 mg, 0.371 mmol, 0.2 equiv.) and Cs₂CO₃ (1207.95 mg, 3.707 mmol, 2.0 equiv.) and XantPhos (429.04 mg, 0.741 mmol, 0.4 equiv.) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 2 h at 110 degrees C under nitrogen atmosphere. The reaction was monitored by

LCMS. The mixture was allowed to cool down to room temperature. The resulting mixture was filtered, the filter cake was washed with DCM (3 x 2 mL). The filtrate was concentrated under reduced pressure. The resulting mixture was concentrated under reduced pressure. The crude product was purified by reverse phase flash with the following conditions (Column: C18, 120 g; Mobile Phase A: Water/0.05% NH₄HCO₃, Mobile Phase B: ACN; Flow rate: 45 mL/min; Gradient: 45%B to 65%B in 15 min; Detector, 254nm and 220 nm, the desired product were collected at 64%B) to afford tert-butyl 4-[[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (600mg, 81.86%) as a white solid.

Tert-butyl 4-[methyl[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate

To a stirred mixture of tert-butyl 4-[[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (1.32 g, 3.339 mmol, 1 equiv.) and Cs₂CO₃ (2.18 g, 6.677 mmol, 2.0 equiv.) in DMF (10 mL) was added CH₃I (0.95 g, 6.677 mmol, 2.0 equiv.) at 0°C under nitrogen atmosphere. The resulting mixture was stirred for 2 h at room temperature under nitrogen atmosphere. The reaction was monitored by LCMS. The crude product was purified by reverse phase flash with the following conditions (Column: C18, 120 g; Mobile Phase A: Water/0.05% NH₄HCO₃, Mobile Phase B: ACN; Flow rate: 45 mL/min; Gradient: 45%B to 65%B in 15 min; Detector, 254nm and 220 nm, the desired product were collected at 64%B) to afford tert-butyl 4-[methyl[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (400 mg, 29.26%) as a brown solid.

N-methyl-N-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]-4-(trifluoromethyl)pyridin-3-amine

To a stirred solution of tert-butyl 4-[methyl[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidine-7-carboxylate (220 mg, 0.537 mmol, 1 equiv.) in DCM (4 mL) was added TFA (1 mL) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 1 h at room temperature. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The mixture was basified to pH 8 with saturated NaHCO₃ (aq.). The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (DCM / MeOH 12:1) to afford N-methyl-N-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]-4-(trifluoromethyl)pyridin-3-amine (130 mg, 78.22%) as a brown solid.

4-chloro-5-(4-[methyl[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one


To a stirred solution of N-methyl-N-[5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-4-yl]-4-(trifluoromethyl)pyridin-3-amine (130 mg, 0.420 mmol, 1 equiv.) in DIEA (0.5 mg) was added 4,5-dichloro-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (104.69 mg, 0.420 mmol, 1.0 equiv.) at room temperature. The resulting mixture was stirred for 1 h at 90°C. The reaction was monitored by LCMS. The mixture was allowed to cool down to room temperature. The resulting mixture was concentrated under reduced pressure. The residue was purified by Prep-TLC (DCM/MeOH 12:1) to afford 4-chloro-5-(4-[methyl[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (100 mg, 45.58%) as a brown solid.

4-chloro-5-(4-[methyl[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2,3-dihydropyridazin-3-one

To a stirred solution of 4-chloro-5-(4-[methyl[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2-(oxan-2-yl)-2,3-dihydropyridazin-3-one (100 mg, 0.192 mmol, 1 equiv.) in DCM (4 mL) was added TFA (1 mL) at room temperature. The resulting mixture was stirred for 1 h at room temperature. The reaction was monitored by LCMS. The resulting mixture was concentrated under reduced pressure. The mixture was basified to pH 8 with saturated NaHCO₃ (aq.). The resulting mixture was concentrated under reduced pressure. The crude product (100 mg) was purified by Prep-HPLC with the following conditions (Column: XBridge Prep Phenyl OBD Column 19×150mm 5um 13nm ; Mobile phase A: water, 5mM NH₄HCO₃, Mobile phase B: Acetonitrile; Flow rate: 60 mL/min; Gradient: 35% B to 55% B in 8 min; 220 nm; Rt: 7.13 min) to afford 4-chloro-5-(4-[methyl[4-(trifluoromethyl)pyridin-3-yl]amino]-5H,6H,7H,8H-pyrido[3,4-d]pyrimidin-7-yl)-2,3-dihydropyridazin-3-one (52 mg, 61.99%) as a white solid.

Example 23. Synthesis of Intermediates

A. 2-(Difluoromethyl)-4-fluorophenyl acetate




4-Fluoro-2-formylphenyl acetate

To a solution of 5-fluoro-2-hydroxybenzaldehyde (10 g, 71.371 mmol, 1 equiv.) in Pyridine (100 mL, 1242.353 mmol, 17.41 equiv.) was added acetyl acetate (14.57 g, 0.143 mmol, 2 equiv.) at 25°C. The solution was stirred at 25°C for 30 min. The resulting solution was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EA (100/1 to 20/1) to afford 4-fluoro-2-formylphenyl acetate (12 g, 92.31%) as a light yellow oil.

2-(Difluoromethyl)-4-fluorophenyl acetate

To a solution of 4-fluoro-2-formylphenyl acetate (12 g, 65.880 mmol, 1 equiv.) in DCM (200 mL, 3146.009 mmol, 47.75 equiv.) was added DAST (21.24 g, 131.760 mmol, 2 equiv.) at 0°C. The solution was stirred at 25°C for 4 hours. The resulting solution was quenched with water (100 mL). The resulting mixture was extracted with DCM (100 mL x 2). The combined organic layers were washed with saturated NaCl aq. (100 mL x 2) dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EA (10/1 to 5/1) to afford 2-(difluoromethyl)-4-fluorophenyl acetate (10 g, 74.35%) as a light yellow oil.

B. 2-(Difluoromethyl)-4-fluorophenol**1-Bromo-2-(difluoromethyl)-4-fluorobenzene**

To a stirred solution of 2-bromo-5-fluorobenzaldehyde (10 g, 49.26 mmol, 1 equiv.) in DCM (60 mL) was added DAST (15.9 g, 98.52 mmol, 2 equiv). The resulting mixture was stirred for 2 hours at -10°C. The reaction was quenched with Water at -10°C. The resulting mixture was extracted with EtOAc (4 x 30 mL). The combined organic layers were washed with brine (2x 40 mL), dried over anhydrous Na₂SO₄. After filtration, the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EtOAc (6:1) to afford 1-bromo-2-(difluoromethyl)-4-fluorobenzene (8 g, 72.18%) as a light yellow oil.

2-[2-(Difluoromethyl)-4-fluorophenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

To a solution of 1-bromo-2-(difluoromethyl)-4-fluorobenzene (31 g, 137.773 mmol, 1 equiv.) and BPD (52.48 g, 206.664 mmol, 1.50 equiv.) in 1,4-dioxane (300 mL, 3541.225 mmol, 25.70 equiv.) were added AcOK (27.04 g, 275.546 mmol, 2 equiv.) and Pd(dppf)Cl₂•CH₂Cl₂ (5.63 g, 6.889 mmol, 0.05 equiv.) at 25°C under nitrogen atmosphere. The mixture was stirred at 90°C for 2 hours. The resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with PE/EA (10/1) to afford 2-[2-(difluoromethyl)-4-fluorophenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (30 g, 80.03%) as a light yellow oil. The reaction was monitored by TLC. The crude was used the next step directly.

2-(Difluoromethyl)-4-fluorophenol

To a solution of 2-[2-(difluoromethyl)-4-fluorophenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (50 g, 183.776 mmol, 1 equiv.) in MeOH (300 mL, 7409.673 mmol, 40.32 equiv.) and H₂O (100 mL, 5550.837 mmol, 30.20 equiv.) was added H₂O₂ (30%) (50 mL, 2146.131 mmol, 11.68 equiv.) dropwise at 0°C. The solution was stirred at 25°C for 3 hours. The resulting solution was concentrated under reduced pressure. The residue was diluted with EA (500 mL). The organic layer was washed with 3 x 200 mL of saturated NaCl (aq.). Combined organic layers was dried with anhydrous Na₂SO₄, concentrated under reduced pressure to afford 2-(difluoromethyl)-4-fluorophenol (25 g, 83.91%) as a light yellow oil.

Example 24. TRPC4 Activity Assay

ICLN-1694 cells (HEK-TREx hTRPC4) expressing TRPC4 were generated as follows. Commercially available HekTrex-293 cells were seeded at 0.7x10⁶ cells/well in a 1x6-well plate 24 hrs prior to transfection using 2 mL cell growth media containing no antibiotics (1x DMEM/high glucose (Hyclone #SH30022.02); 10% fetal bovine serum (Sigma) 2mM sodium pyruvate, 10 mM HEPES). The human codon-optimized TRPC4 coding sequence was cloned into pcDNA5/TO (Invitrogen; Cat No. V103320) using hygromycin as the resistance gene and the plasmid (SEQ ID NO:1) propagated using T-Rex-293 cells (Invitrogen; Cat No. R71007) following manufacturer's directions. On day 2, 2 µg of plasmid DNA plus 6 µl of Xtreme-GENE HP reagent in Optimem (200 µl total volume) was prepared and incubated for 15 min at room temperature. This plasmid solution was then gently overlayed dropwise onto each well and the plate was gently swirled to mix complex with the media for approximately 30 seconds.

Transfected cells were incubated at 37 °C in a 10% CO₂ incubator for 24 hrs. The transfected cells were harvested and transferred into 2 x 150mm dishes containing cell growth media with no antibiotics at 37 °C

The next day selection was initiated to generate a stable pool by adding cell growth media containing 150 µg/mL Hygromycin and 5 µg/mL Blasticidin and cells were allowed to grow. Media with the selection agent was changed every 1-2 days as needed to remove dead cells. After 7 days, the hygromycin concentration was reduced to 75 µg/mL and cells growth was allowed to continue.

Single clones were selected as follows. The stable pool was diluted to 10 cells/mL and seeded (100 µl/well) into 24 x 96 well plates (~1 cell/well) and allowed to grow for 7 days in cell growth media. Fresh media (100 µl) was added and the cells allowed to grow for another 1-2 weeks and then stored frozen or used immediately.

Compounds were made up to, or supplied as, a 10 mM stock solution generally using DMSO as the vehicle. 10-point dose response curves were generated using the Echo-550 acoustic dispenser. Compound source plates were made by serially diluting compound stocks to create 10mM, 1mM, and 0.1mM solutions in DMSO into Echo-certified LDV plates. The Echo then serially spotted 100% DMSO stock solutions into source dose response plates to generate a 4-fold dilution scheme. 100% DMSO was added to the spotted dose response plates to bring the final volume to 5µl. 300nl of the dose response stock plate was then spotted into pre-incubation and stimulation assay plates. 50µl of pre-incubation buffer and 100µl of stimulation buffer was then added to the plates resulting in a final assay test concentration range of 30µM to 0.0001µM with a final DMSO concentration of 0.3%.

ICLN-1694 cells (HEK-TREx hTRPC4) were plated onto 384 well, black pdl-coated microplates and maintained in cell growth media supplemented with 1 µg/mL tetracycline the day prior to use for experiments. TRPC4 expression was induced by the application of 1 µg/mL tetracycline at the time of plating. Media was removed from the plates and 10µl of 4µM of Fluo-4 AM (mixed with equal volume of Pluronic F-127) in EBSS (NaCl (142 mM), KCl (5.4 mM), glucose (10 mM), CaCl₂ (1.8 mM), MgCl₂ (0.8 mM), HEPES (10 mM), pH 7.4) is added to the cells. Cells were incubated at room temperature, protected from light, for 60-90 minutes. After the incubation period, the dye was removed and replaced with 10µl of EBSS. Cell, pre-incubation and stimulation plates were loaded onto the FLIPR-II and the assay was initiated.

The FLIPR measured a 10 second baseline and then added 10µl of 2X compounds (or controls). Changes in fluorescence were monitored for an additional 5 minutes. After a 5 minute pre-incubation, 20µl of 2X Englerin A (with 1X compound or controls) was added to the cell plate. The final Englerin A stimulation concentration in the assay was 100nM. After the Englerin A addition, changes in fluorescence were monitored for an additional 5 minutes.

Compound modulation of TRPC4 calcium response was determined as follows. After the Englerin A, fluorescence was monitored for a 5-minute period. The maximum relative fluorescence response (minus the control response of 1 µM of an internal control compound known to maximally block TRPC4 calcium response, the “REF INHIB” in the formula below) was captured and exported from the FLIPR.

Compound effect is calculated as % inhibition using the following formula:

$$\% \text{ inhibition} = \frac{RFU \text{ TEST AGENT} - \text{Plate Average RFU REF INHIB}}{\text{Plate Average RFU CONTROL} - \text{Plate Average RFU REF INHIB}} \times 100$$

wherein “RFU” is the relative fluorescent units.

The results of these assays are shown in Table 2, below, wherein “A” indicates an IC₅₀ of less than or equal to 50 nM; “B” an IC₅₀ of greater than 50 nM and less than or equal to 500 nM; “C” an IC₅₀ of greater than 500 nM and less than 1 µM; “D” an IC₅₀ of 1 µM or greater; and “NT” indicates that the compound was not tested.

Example 25. TRPC5 Activity Assay

ICLN-1633 cells (HEK-TREx hTRPC5) expressing TRPC5 were generated as follows. Commercially available HekTrex-293 cells were seeded at 0.7x10⁶ cells/well in a 1x6-well plate 24 hrs prior to transfection using 2 mL cell growth media containing no antibiotics (1x DMEM/high glucose (Hyclone #SH30022.02); 10% fetal bovine serum (Sigma) 2mM sodium pyruvate, 10 mM HEPES). The human TRPC5 coding sequence (NM_012471 with a silent T478C mutation) was cloned into pcDNA5/TO (Invitrogen; Cat No. V103320) using hygromycin as the resistance gene and the plasmid (SEQ ID NO:2) propagated using T-Rex-293 cells (Invitrogen; Cat No. R71007) following manufacturer’s directions. On day 2, 2 µg of

plasmid DNA plus 6 μ l of Xtreme-GENE HP reagent in Optimem (200 μ l total volume) was prepared and incubated for 15 min at room temperature. This plasmid solution was then gently overlaid dropwise onto each well and the plate was gently swirled to mix complex with the media for approximately 30 seconds. Transfected cells were incubated at 37 °C in a 10% CO₂ incubator for 24 hrs. The transfected cells were harvested and transferred into 2 x 150mm dishes containing cell growth media with no antibiotics at 37 °C

The next day selection was initiated to generate a stable pool by adding cell growth media containing 150 μ g/mL Hygromycin and 5 μ g/mL Blasticidin and cells were allowed to grow. Media with the selection agent was changed every 1-2 days as needed to remove dead cells. After 7 days, the hygromycin concentration was reduced to 75 μ g/mL and cells growth was allowed to continue.

Single clones were selected as follows. The stable pool was diluted to 10 cells/mL and seeded (100 μ l/well) into 24 x 96 well plates (~1 cell/well) and allowed to grow for 7 days in cell growth media. Fresh media (100 μ l) was added and the cells allowed to grow for another 1-2 weeks and then stored frozen or used immediately.

Compounds were made up to, or supplied as a 10 mM stock solution generally using DMSO as the vehicle. 10-point dose response curves were generated using the Echo-550 acoustic dispenser. Compound source plates were made by serially diluting compound stocks to create 10mM, 1mM, and 0.1mM solutions in DMSO into Echo certified LDV plates. The Echo then serially spotted 100% DMSO stock solutions into source dose response plates to generate a 4-fold dilution scheme. 100% DMSO was added to the spotted dose response plates to bring the final volume to 5 μ l. 300nl of the dose response stock plate was then spotted into pre-incubation and stimulation assay plates. 50 μ l of pre-incubation buffer and 100 μ l of stimulation buffer was then added to the plates resulting in a final assay test concentration range of 30 μ M to 0.0001 μ M with a final DMSO concentration of 0.3%.

Human ICLN-1633 cells expressing were plated onto 384 well, black PDL-coated microplates and maintained in TRPC5 growth media the day prior to use for experiments. TRPC5 expression was induced by the application of 1 μ g/mL tetracycline at the time of plating. Media is removed from the plates and 10 μ l of 4 μ M of Fluo-4 AM (mixed with equal volume of Pluronic F-127) in EBSS is added to the cells. Cells are incubated at room temperature, protected from light, for 60-90 minutes. After the incubation period, the dye is removed and replaced with

10µl of EBSS. Cell, pre-incubation and stimulation plates are loaded onto the FLIPR-II and the assay is initiated. The FLIPR measures a 10 second baseline and then adds 10µl of 2X compounds (or controls). Changes in fluorescence are monitored for an additional 5 minutes. After the 5 minute pre-incubation, 20µl of 2X Riluzole (with 1X compound or controls) is added to the cell plate. The final Riluzole stimulation concentration in the assay is 30µM. After the Riluzole addition, changes in fluorescence are monitored for an additional 5 minutes.

Compound modulation of TRPC5 calcium response was determined as follows. After the Englerin A, fluorescence was monitored for a 5-minute period. The maximum relative fluorescence response (minus the control response of 1 µM of an internal control compound known to maximally block TRPC5 calcium response, the “REF INHIB” in the formula below) was captured and exported from the FLIPR.

Compound effect is calculated as % inhibition using the following formula:

% *inhibition*

$$= \frac{RFU \text{ TEST AGENT} - \text{Plate Average } RFU \text{ REF INHIB}}{\text{Plate Average } RFU \text{ CONTROL} - \text{Plate Average } RFU \text{ REF INHIB}} \times 100$$

wherein “RFU” is the relative fluorescent units.

The results of these assays are shown in Table 2, below, wherein “A” indicates an IC₅₀ of less than or equal to 50 nM; “B” an IC₅₀ of greater than 50 nM and less than or equal to 500 nM; “C” an IC₅₀ of greater than 500 nM and less than 1 µM; “D” an IC₅₀ of 1 µM or greater; and “NT” indicates that the compound was not tested.







Table 2. TRPC4 and TRPC5 Activities of Exemplary Compounds







<u>Compound</u>	<u>TRPC5</u>	<u>TRPC4</u>
100	A	A
101	A	A
102	A	A
103	A	A
104	A	A
105	A	A
106	A	B
107	A	B
108	B	C
109	A	B
110	B	A
111	B	NT
112	A	B
113	A	B
114	A	A
115	B	B
116	A	A
117	A	B
117a	D	NT
118	B	B
119	A	A
120	A	A
121	A	NT
122	A	NT
123	B	NT
124	A	NT
125	A	NT
126	A	NT
126a	B	NT
127	B	NT
128	A	NT
129	A	NT
130	A	NT
131	A	NT
132	B	NT
133	B	NT
133a	C	NT
134	A	NT
135	A	NT
136	A	NT
137	A	NT
138	B	NT

<u>Compound</u>	<u>TRPC5</u>	<u>TRPC4</u>
139	B	NT
140	B	NT







¹H NMR and MS data for selected compounds is provided in the table below:

Compound	Structure	NMR	MS
100		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.57 (s, 1H), 8.04 (s, 1H), 7.79 (dd, J = 8.4, 3.0 Hz, 1H), 7.70 (s, 1H), 7.70 – 7.59 (m, 1H), 4.68 (s, 2H), 3.79 (t, J = 5.6 Hz, 2H), 2.98 (s, 2H).	442
101		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.54 (s, 1H), 8.03 (s, 1H), 7.63 (dd, J = 8.0, 1.4 Hz, 1H), 7.51 – 7.40 (m, 2H), 7.36 (td, J = 7.4, 2.2 Hz, 1H), 4.67 (s, 2H), 3.80 (t, J = 5.7 Hz, 2H), 3.03 (t, J = 5.7 Hz, 2H).	390
102		¹ H NMR (400 MHz, Methanol-d ₄) chemical shifts 8.00 (s, 1H), 7.79 – 7.67 (m, 2H), 7.44 (dd, J = 13.1, 7.9 Hz, 2H), 4.51 (s, 2H), 3.87 (t, J = 5.8 Hz, 2H), 2.93 (t, J = 5.6 Hz, 2H).	439
103		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.54 (s, 1H), 8.04 (s, 1H), 7.71 (d, J = 7.7 Hz, 1H), 7.65 (t, J = 7.9 Hz, 1H), 7.46 (t, J = 7.6 Hz, 1H), 7.40 (d, J = 8.2 Hz, 1H), 7.13 (t, J = 54.3 Hz, 1H), 4.67 (s, 2H), 3.79 (t, J = 5.7 Hz, 2H), 3.03 (t, J = 5.7 Hz, 2H).	406
104		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.97 (s, 1H), 8.55 (s, 1H), 8.03 (s, 1H), 7.66 (dd, J = 8.3, 3.0 Hz, 1H), 7.51 (dd, J = 8.9, 5.3 Hz, 1H), 7.42 – 7.24 (m, 1H), 4.67 (s, 2H), 3.80 (t, J = 5.9 Hz, 2H), 3.02 (s, 2H).	407






Compound	Structure	NMR	MS
105		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.55 (s, 1H), 8.04 (s, 1H), 7.51 (ddd, J = 14.3, 10.8, 6.8 Hz, 3H), 7.13 (t, J = 53.9 Hz, 1H), 4.67 (s, 2H), 3.79 (t, J = 5.7 Hz, 2H), 3.03 (t, J = 5.8 Hz, 2H).	424
106		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.61 (s, 1H), 8.03 (s, 1H), 7.95 (d, J = 8.4 Hz, 1H), 7.62 (d, J = 9.7 Hz, 1H), 7.41 (s, 1H), 4.69 (s, 2H), 3.80 (s, 2H), 2.98 (s, 2H).	442
107		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.91 (s, 1H), 8.79 (d, J = 4.9 Hz, 1H), 8.59 (s, 1H), 8.04 (s, 1H), 7.92 (d, J = 5.0 Hz, 1H), 4.70 (s, 2H), 3.81 (t, J = 5.7 Hz, 2H), 3.02 (s, 2H).	425
108		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.57 (s, 1H), 8.04 (s, 1H), 7.99 (dd, J = 7.8, 1.5 Hz, 1H), 7.86 (dd, J = 8.3, 1.5 Hz, 1H), 7.69 (t, J = 8.0 Hz, 1H), 4.69 (s, 2H), 3.81 (t, J = 5.7 Hz, 2H), 3.08 – 3.01 (m, 2H).	415
109		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.01 (s, 1H), 8.54 (s, 1H), 8.04 (s, 1H), 7.38 – 7.28 (m, 1H), 7.15 (t, J = 8.9 Hz, 1H), 7.06 (d, J = 8.2 Hz, 1H), 4.67 (s, 2H), 3.80 (t, J = 5.8 Hz, 2H), 3.03 (d, J = 5.8 Hz, 2H), 2.48 (d, J = 7.4 Hz, 2H), 1.05 (t, J = 7.5 Hz, 3H).	402
110		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.98 (s, 1H), 8.02 (s, 1H), 7.88 – 7.75 (m, 2H), 7.59 – 7.48 (m, 2H), 4.60 (s, 2H), 3.77 (t, J = 5.7 Hz, 2H), 3.74 (s, 3H), 2.89 (t, J = 5.6 Hz, 2H).	454

Compound	Structure	NMR	MS
111		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.01 (s, 1H), 8.60 (s, 1H), 8.13 (d, J = 7.4 Hz, 1H), 8.07 – 7.99 (m, 2H), 7.96 (d, J = 8.4 Hz, 1H), 4.70 (s, 2H), 3.81 (t, J = 5.6 Hz, 2H), 3.01 (s, 2H).	449
112		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.95 (s, 1H), 7.97 (s, 1H), 7.62 (dd, J = 8.5, 3.1 Hz, 1H), 7.45 (dd, J = 9.0, 5.3 Hz, 1H), 7.31 (td, J = 8.5, 2.9 Hz, 1H), 6.50 (s, 2H), 4.41 (s, 2H), 3.72 (d, J = 6.0 Hz, 2H), 2.80 (s, 2H).	423
113		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.95 (s, 1H), 7.97 (s, 1H), 7.58 (dd, J = 7.9, 1.5 Hz, 1H), 7.47 – 7.38 (m, 1H), 7.41 – 7.34 (m, 1H), 7.34 – 7.25 (m, 1H), 6.48 (s, 2H), 4.41 (s, 2H), 3.73 (t, J = 5.8 Hz, 2H), 2.81 (d, J = 5.7 Hz, 2H).	405
114		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.03 (s, 1H), 7.66 (dd, J = 8.4, 3.0 Hz, 1H), 7.51 (dd, J = 9.0, 5.3 Hz, 1H), 7.35 (td, J = 8.6, 3.0 Hz, 1H), 5.17 (t, J = 6.3 Hz, 1H), 4.67 (s, 2H), 4.32 (d, J = 6.3 Hz, 2H), 3.80 (t, J = 5.7 Hz, 2H), 3.00 (d, J = 6.4 Hz, 2H).	438
115		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.95 (s, 1H), 7.98 (s, 1H), 7.83 – 7.72 (m, 2H), 7.48 (dd, J = 18.0, 8.2 Hz, 2H), 6.82 (s, 1H), 4.54 (s, 1H), 4.44 (s, 2H), 3.73 (t, J = 5.6 Hz, 2H), 3.33 (m, 4H), 2.78 (s, 2H).	483
116		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.96 (s, 1H), 7.97 (s, 1H), 7.79 – 7.71 (m, 1H), 7.66 (td, J = 8.5, 3.0 Hz, 1H), 7.56 (dd, J = 9.2, 4.6 Hz, 1H), 6.54 (s, 2H), 4.41 (s, 2H), 3.72 (t, J = 5.8 Hz, 2H), 2.76 (t, J = 5.7 Hz, 2H).	457

Compound	Structure	NMR	MS
117*		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.04 (s, 1H), 8.56 (s, 1H), 8.02 (s, 1H), 7.67 (dd, J = 8.4, 2.5 Hz, 1H), 7.53 (dd, J = 9.2, 5.3 Hz, 1H), 7.41 – 7.31 (m, 1H), 4.75 (d, J = 18.1 Hz, 1H), 4.62 – 4.34 (m, 2H), 3.23 (dd, J = 17.0, 6.0 Hz, 1H), 2.81 (d, J = 17.1 Hz, 1H), 1.20 (d, J = 6.6 Hz, 3H).	422
117a*		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.04 (s, 1H), 8.56 (s, 1H), 8.02 (s, 1H), 7.67 (dd, J = 8.6, 2.5 Hz, 1H), 7.58 – 7.50 (m, 1H), 7.36 (dd, J = 9.9, 7.3 Hz, 1H), 4.76 (d, J = 18.1 Hz, 1H), 4.63 – 4.40 (m, 2H), 3.23 (dd, J = 17.3, 5.9 Hz, 1H), 2.81 (d, J = 17.0 Hz, 1H), 1.20 (d, J = 6.7 Hz, 3H).	422
118		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.96 (s, 1H), 8.55 (s, 1H), 8.02 (s, 1H), 7.66 (dd, J = 8.4, 3.0 Hz, 1H), 7.50 (dd, J = 9.0, 5.3 Hz, 1H), 7.35 (td, J = 8.5, 3.0 Hz, 1H), 5.06 (q, J = 6.7 Hz, 1H), 4.00 (dd, J = 14.1, 5.7 Hz, 1H), 3.68 (ddd, J = 14.4, 11.1, 4.2 Hz, 1H), 3.06 (ddd, J = 17.1, 11.2, 6.0 Hz, 1H), 2.98 – 2.88 (m, 1H), 1.60 (d, J = 6.8 Hz, 3H).	422
119		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.98 (s, 1H), 8.03 (s, 1H), 7.66 (dd, J = 8.4, 3.0 Hz, 1H), 7.51 (dd, J = 9.1, 5.3 Hz, 1H), 7.35 (td, J = 8.5, 3.0 Hz, 1H), 5.06 (d, J = 5.3 Hz, 1H), 4.67 (s, 2H), 4.48 (p, J = 6.4 Hz, 1H), 3.80 (t, J = 5.5 Hz, 2H), 2.99 (s, 2H), 1.22 (t, J = 6.4 Hz, 3H).	452
120		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 7.98 (s, 1H), 7.90 (d, J = 5.2 Hz, 1H), 7.72 (dd, J = 8.6, 3.1 Hz, 1H), 7.63 (td, J = 8.6, 3.1 Hz, 1H), 7.49 (dd, J = 9.1, 4.6 Hz, 1H), 7.00 (d, J = 5.2 Hz, 1H), 4.69 (s, 2H), 3.78 (t, J = 5.8 Hz, 2H), 2.95 (t, J = 5.9 Hz, 2H).	441

Compound	Structure	NMR	MS
121		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.01 (s, 1H), 8.60 (s, 1H), 8.04 (s, 1H), 7.83 (t, J = 9.2 Hz, 1H), 7.73 (d, J = 7.9 Hz, 1H), 7.67 – 7.57 (m, 1H), 4.71 (s, 2H), 3.81 (t, J = 5.8 Hz, 2H), 3.01 (s, 2H).	442
122		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.03 (s, 1H), 7.88 – 7.75 (m, 2H), 7.57 (d, J = 8.2 Hz, 1H), 7.51 (t, J = 7.6 Hz, 1H), 5.07 (d, J = 5.3 Hz, 1H), 4.67 (s, 2H), 4.50 (p, J = 6.2 Hz, 1H), 3.80 (t, J = 5.6 Hz, 2H), 2.96 (s, 2H), 1.23 (d, J = 6.5 Hz, 3H).	468
123		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.03 (s, 1H), 7.88 – 7.75 (m, 2H), 7.57 (d, J = 8.3 Hz, 1H), 7.51 (t, J = 7.7 Hz, 1H), 5.07 (d, J = 5.2 Hz, 1H), 4.67 (s, 2H), 4.50 (p, J = 6.4 Hz, 1H), 3.80 (t, J = 5.6 Hz, 2H), 2.96 (s, 2H), 1.23 (d, J = 6.6 Hz, 3H).	468
124		¹ H NMR (DMSO-d ₆) δ: 12.97 (br s, 1H), 8.49 (s, 1H), 8.01 (s, 1H), 7.15-7.24 (m, 2H), 7.05-7.12 (m, 1H), 4.63 (s, 2H), 3.78 (t, J=5.7 Hz, 2H), 3.00 (s, 2H), 2.06 (s, 3H)	388.3
125		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.03 (s, 1H), 7.81 – 7.74 (m, 1H), 7.74 – 7.60 (m, 2H), 5.19 (t, J = 6.3 Hz, 1H), 4.67 (s, 2H), 4.34 (d, J = 6.3 Hz, 2H), 3.79 (t, J = 5.6 Hz, 2H), 2.95 (s, 2H).	472
126*		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.98 (s, 1H), 9.02 (s, 1H), 7.97 (s, 1H), 7.63 (dd, J = 9.4, 2.6 Hz, 1H), 7.53 (p, J = 8.7 Hz, 2H), 4.71 (t, J = 7.1 Hz, 1H), 4.65 (s, 2H), 3.68 (t, J = 5.7 Hz, 2H), 3.08 (d, J = 16.5 Hz, 1H), 2.57 (s, 1H), 1.60 (d, J = 6.8 Hz, 3H).	454

Compound	Structure	NMR	MS
126a*		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.97 (s, 1H), 9.02 (s, 1H), 7.97 (s, 1H), 7.63 (d, J = 7.9 Hz, 1H), 7.60 – 7.49 (m, 2H), 4.72 (d, J = 6.9 Hz, 1H), 4.65 (s, 2H), 3.68 (s, 2H), 3.08 (d, J = 16.3 Hz, 1H), 2.58 (s, 1H), 1.60 (d, J = 6.8 Hz, 3H).	454
127		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.98 (s, 1H), 7.96 (s, 1H), 7.81 – 7.74 (m, 1H), 7.55 (d, J = 8.8 Hz, 1H), 7.35 – 7.27 (m, 1H), 5.85 (s, 1H), 4.52 (s, 2H), 3.39 (s, 3H), 3.39 – 3.34 (m, 2H), 3.09 (s, 3H), 1.95 (s, 2H).	484
128		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.01 (s, 1H), 8.03 (s, 1H), 7.88 – 7.75 (m, 2H), 7.56 (d, J = 8.3 Hz, 1H), 7.51 (t, J = 7.7 Hz, 1H), 5.19 (t, J = 6.3 Hz, 1H), 4.68 (s, 2H), 4.34 (d, J = 6.3 Hz, 2H), 3.80 (t, J = 5.7 Hz, 2H), 2.96 (s, 2H).	454
129*		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.03 (s, 1H), 7.78 (dd, J = 8.6, 2.9 Hz, 1H), 7.75 – 7.61 (m, 2H), 5.08 (d, J = 5.3 Hz, 1H), 4.67 (s, 2H), 4.50 (p, J = 6.5 Hz, 1H), 3.79 (t, J = 5.7 Hz, 2H), 2.94 (d, J = 6.1 Hz, 2H), 1.23 (d, J = 6.5 Hz, 3H).	486
130*		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.00 (s, 1H), 8.03 (s, 1H), 7.78 (dd, J = 8.6, 3.0 Hz, 1H), 7.67 (ddd, J = 19.1, 8.6, 3.8 Hz, 2H), 5.08 (d, J = 5.3 Hz, 1H), 4.67 (s, 2H), 4.50 (p, J = 6.5 Hz, 1H), 3.79 (t, J = 5.5 Hz, 2H), 2.95 (s, 2H), 1.23 (d, J = 6.6 Hz, 3H).	486
131		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.07 (s, 1H), 8.89 – 8.79 (m, 2H), 8.59 (s, 1H), 7.94 – 7.84 (m, 2H), 4.53 (s, 2H), 3.44 (d, J = 11.0 Hz, 2H), 3.41 (s, 3H), 1.96 (t, J = 5.6 Hz, 2H).	438

Compound	Structure	NMR	MS
132		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.94 (s, 1H), 8.89 (s, 1H), 8.00 (d, J = 8.8 Hz, 1H), 7.87 (s, 1H), 7.62 (d, J = 8.6 Hz, 2H), 4.58 (s, 2H), 3.48 (s, 2H), 2.27 (s, 2H), 1.84 (s, 2H), 1.47 (s, 2H).	466
133		¹ H NMR (400 MHz, Methanol-d ₄) chemical shifts 8.34 (s, 1H), 8.02 (s, 1H), 7.33 (q, J = 7.8 Hz, 4H), 7.25 (t, J = 7.0 Hz, 1H), 4.67 (d, J = 17.3 Hz, 1H), 4.58 (d, J = 17.2 Hz, 2H), 3.97 (d, J = 13.9 Hz, 1H), 3.55 (s, 2H), 3.47 (s, 1H), 3.39 (s, 3H), 2.94 (s, 1H), 2.81 (d, J = 15.8 Hz, 1H), 2.74-2.53 (m, 3H), 2.35 (s, 1H), 2.15 (s, 1H), 1.76 (s, 2H), 1.01 (d, J = 7.0 Hz, 3H).	480
133a		¹ H NMR (400 MHz, Methanol-d ₄) chemical shifts 8.34 (s, 1H), 8.02 (s, 1H), 7.42 – 7.21 (m, 5H), 4.62 (q, J = 17.5 Hz, 3H), 3.97 (d, J = 13.3 Hz, 1H), 3.55 (s, 2H), 3.47 (s, 1H), 3.39 (s, 3H), 3.15 (s, 1H), 2.94 (s, 1H), 2.87 – 2.61 (m, 3H), 2.35 (s, 1H), 2.15 (s, 1H), 1.76 (s, 2H), 1.01 (d, J = 7.0 Hz, 3H).	480
134		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.01 (s, 1H), 8.03 (s, 1H), 7.58 – 7.49 (m, 3H), 7.27 – 7.00 (m, 1H), 5.17 (t, J = 6.3 Hz, 1H), 4.66 (s, 2H), 4.34 (d, J = 6.3 Hz, 2H), 3.79 (t, J = 5.7 Hz, 2H), 2.99 (s, 2H).	454
135*		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.97 (s, 1H), 8.03 (s, 1H), 7.78 (dd, J = 8.5, 3.0 Hz, 1H), 7.75 – 7.61 (m, 2H), 5.12 (d, J = 5.7 Hz, 1H), 4.67 (s, 2H), 4.57 (t, J = 6.0 Hz, 1H), 4.37 (q, J = 5.8 Hz, 1H), 3.79 (t, J = 5.7 Hz, 2H), 3.57 (dt, J = 11.2, 5.7 Hz, 1H), 3.46 (dt, J = 10.8, 6.1 Hz, 1H), 2.96 (t, J = 5.6 Hz, 2H).	502

Compound	Structure	NMR	MS
136		¹ H NMR (400 MHz, Methanol-d ₄) chemical shifts 8.54 (s, 1H), 7.87 (s, 1H), 7.75 (s, 1H), 7.63 (dd, J = 8.8, 3.0 Hz, 1H), 7.54 (s, 1H), 4.57 (m, 3H), 3.88 (s, 2H), 3.51 (m, 3H), 2.05 (s, 1H), 1.98 (s, 1H).	485
137*		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.94 (s, 1H), 8.03 (s, 1H), 7.78 (dd, J = 8.5, 3.0 Hz, 1H), 7.75 – 7.61 (m, 2H), 5.12 (d, J = 5.7 Hz, 1H), 4.67 (s, 2H), 4.57 (t, J = 6.0 Hz, 1H), 4.37 (q, J = 5.7 Hz, 1H), 3.79 (t, J = 5.7 Hz, 2H), 3.57 (dt, J = 11.2, 5.7 Hz, 1H), 3.52 – 3.43 (m, 1H), 2.96 (t, J = 5.6 Hz, 2H).	502
138		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 13.05 (s, 1H), 8.11 (s, 1H), 7.78 (dd, J = 8.5, 3.1 Hz, 1H), 7.70 (td, J = 8.5, 3.2 Hz, 1H), 7.57 (dd, J = 9.1, 4.6 Hz, 1H), 4.65 (s, 2H), 3.66 (t, J = 5.7 Hz, 2H), 3.38 (s, 3H), 2.74 (t, J = 5.4 Hz, 2H).	472
139		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.84 (s, 1H), 8.02 (s, 1H), 7.79 (dd, J = 8.6, 2.9 Hz, 1H), 7.68 (dtd, J = 18.8, 9.0, 3.8 Hz, 2H), 4.60 (s, 2H), 3.77 (d, J = 5.6 Hz, 2H), 3.75 (s, 3H), 2.88 (t, J = 5.7 Hz, 2H).	472
140		¹ H NMR (400 MHz, DMSO-d ₆) chemical shifts 12.97 (s, 1H), 8.30 (d, J = 5.6 Hz, 1H), 8.02 (s, 1H), 7.83 (dd, J = 8.5, 3.2 Hz, 1H), 7.68 (td, J = 8.6, 3.1 Hz, 1H), 7.50 (dd, J = 9.1, 4.5 Hz, 1H), 6.58 (d, J = 5.6 Hz, 1H), 4.69 (s, 2H), 3.77 (t, J = 5.7 Hz, 2H), 2.98 (t, J = 5.8 Hz, 2H).	441

*Compound is a pure isomer separated from its opposite isomer by HPLC of the corresponding racemic mixture. Absolute orientation has not been determined and therefore designation of the specific orientation around a chiral center is arbitrary.

Example 26. Effect of Compound 100 on Puromycin Aminonucleoside (PAN)-Induced Glomerular Injury in Rats

Objective:

The objective of this study is to evaluate the dose-dependent effects of compound 100 on PAN induced glomerular kidney injury as indexed by albuminuria.

Methods:

Eighty (80), male Sprague-Dawley rats weighing approximately 125-150 g and approximately 5-6 weeks of age were acquired from Charles River. They were fed a standard chow diet (Harlan 8640), housed under standard conditions, and allowed to acclimate for at least 5 days prior to study inception.

On D-2, rats were placed into weight-matched treatment groups and were placed, individually housed, into metabolic cages for the balance of the study.

A 24 hour baseline (Day 0) urine was collected followed by a baseline blood collection via conscious tail venous puncture. Rats were then administered vehicle or test article.

Two (2) hours following administration of vehicle or test agent on Day 0, rats received an administration of (5 ml/kg, s.c.) vehicle (sterile saline) or puromycin aminonucleoside (PAN; challenge agent; 75mg/kg) dissolved in vehicle.

Intermittent (Day 4, 7 and 10) 24 hour urine volumes were determined and samples (4 samples/animal/time point; 0.5 ml/sample) were obtained. Additionally, intermittent (Day 4, 7 and 10) blood samples were collected via conscious tail venous puncture 2 hours $\pm$ 1 minute post-AM dose.

Immediately following the last blood collection, rats were anesthetized with isoflurane, tissues harvested, and animals sacrificed. Endpoint kidney weights and indices were obtained.

Urine samples were immediately flash-frozen in liquid N₂ and stored at -80°C until analyzed.

Whole blood samples collected on K₃EDTA were processed appropriately for the production of plasma for PK measurements.

Results:

As shown in Figure 1, treatment with compound 100 at 30 mg/kg once- (QD) or twice- (BID) daily resulted in reduced urinary albumin excretion on following injury with PAN. Significant reductions were seen at 7 and 10 days with BID dosing, and at 10 days with QD dosing of compound 100. Mizoribine, the positive control compound, was also efficacious in reducing albuminuria.

Conclusion:

Compound 100 is effective in reducing albuminuria in the PAN model of glomerular injury in rats.

Example 27. Compound 100 is Efficacious in the AT1R Transgenic Rat Model of FSGS

The AT1R transgenic rat model of FSGS is characterized by podocyte-specific expression of human AT1R. Males have been shown to have substantially worse pathology compared to females. The efficacy of TRPC5 inhibitors in the AT1R model has been demonstrated with a tool compound. *See Zhou et al., Science (2017), vol. 358 (6368), 1332-1336.*

In the present study, pathophysiology in AT1R transgenic rats was accelerated with unilateral nephrectomy (UniNX) and minipump AngII infusion. Compound 100 was dosed orally once daily at 3 mg/kg or 10 mg/kg, and the urine protein creatinine ratio was determined at -1, 0, 1, 2, and 3 weeks of treatment. Figure 2 shows the urine protein creatinine ratio over the course of the study for rats treated with compound 100 or those treated with vehicle. UniNX, AngII incept, and compound 100 incept occurred at the time points indicated. Figure 3 shows the same data presented as % of baseline.

The results show that compound 100 is efficacious in the AT1R transgenic rat model of FSGS.

INCORPORATION BY REFERENCE

All of the U.S. patents and U.S. and PCT published patent applications cited herein are hereby incorporated by reference.

EQUIVALENTS

The foregoing written specification is sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by examples provided, since the examples are intended as a single illustration of one aspect of the invention and other functionally equivalent embodiments are within the scope of the invention. Various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims. The advantages and objects of the invention are not necessarily encompassed by each embodiment of the invention.

SEQ ID NO:1 TRPC4 Plasmid Sequence

The DNA sequence of the TRPC4 plasmid used in Example 24 is included below.
Underlined nucleic acids represent those encoding human TRPC4.

GACGGATCGGGAGATCTCCCGATCCCCTATGGTGCACTCTCAGTACAATCTGCTCTG
ATGCCGCATAGTTAAGCCAGTATCTGCTCCCTGCTTGTGTGTTGGAGGTCGCTGAGT
AGTGCGCGAGCAAAATTTAAGCTACAACAAGGCAAGGCTTGACCGACAATTGCATG
AAGAATCTGCTTAGGGTTAGGCGTTTTGCGCTGCTTCGCGATGTACGGGCCAGATAT
ACGCGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTA
GTTCATAGCCCATATATGGAGTTCCGCGTTACATAAATTACGGTAAATGGCCCCGCCT
GGCTGACCGCCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCAT
GTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACT
GCCCATTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTC
AATGACGGTAAATGGCCCCGCCTGGCATTATGCCCAGTACATGACCTTATGGGACTTT
CCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTT
GGCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCC
ACCCCATGACGTCAATGGGAGTTTGTGTTTGGCACCAAAATCAACGGGACTTTCCAA
AATGTCGTAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGG
GAGGTCTATATAAGCAGAGCTCTCCCTATCAGTGATAGAGATCTCCCTATCAGTGAT
AGAGATCGTCGACGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCC
ACGCTGTTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCGGACTCTAGC
GTTTAACTTAAGCTTGGTACCGAGCTCGGATCCGCCACCATGGCCAGTTCTACTA
TAAGAGAAACGTGAATGCCCCTTACCGCGACAGAATCCCCCTGAGAATCGTGAGGG
CAGAGTCCGAGCTGAGCCCATCCGAGAAGGCCTACCTGAACGCCGTGGAGAAGGGC
GACTATGCCAGCGTGAAGAAGTCCCTGGAGGAGGCCGAGATCTACTTTAAGATCAA
CATCAATTGCATCGATCCTCTGGGCAGAACCGCCCTGCTGATCGCCATCGAGAACGA
GAATCTGGAGCTGATCGAGCTGCTGCTGAGCTTCAACGTGTATGTGGGCGATGCCCT
GCTGCACGCCATCAGGAAGGAGGTGGTGGGAGCAGTGGAGCTGCTGCTGAATCACA
AGAAGCCAAGCGGAGAGAAGCAGGTGCCACCTATCCTGCTGGACAAGCAGTTCTCC
GAGTTTACCCAGATATCACACCCATCATCCTGGCCGCCACACCAACAATTACGAG

ATCATCAAGCTGCTGGTGCAGAAGGGCGTGTCCGTGCCTCGCCACACGAGGTGCG
GTGCAACTGCGTGGAGTGCGTGAGCTCCTCTGACGTGGATTCTCTGAGGCACAGCCG
GAGCCGGCTGAACATCTATAAGGCCCTGGCCTCCCCATCTCTGATCGCCCTGAGCTC
CGAGGACCCCTTCCTGACCGCCTTTCAGCTGTCTTGGGAGCTGCAGGAGCTGAGCAA
GGTGGAGAACGAGTTTAAGAGCGAGTACGAGGAGCTGTCCAGACAGTGCAAGCAGT
TCGCCAAGGACCTGCTGGATCAGACACGCTCTAGCCGGGAGCTGGAGATCATCTG
AACTATAGGGACGATAATTCTCTGATCGAGGAGCAGAGCGGAAACGACCTGGCACG
CCTGAAGCTGGCCATCAAGTACCGGCAGAAGGAGTTCGTGGCCCAGCCTAATTGTC
AGCAGCTGCTGGCCTCCCGCTGGTATGATGAGTTTCCAGGATGGCGGAGAAGGCAC
TGGGCAGTGAAGATGGTGACCTGCTTCATCATCGGCCTGCTGTTCCCGTGTTACG
GTGTGCTACCTGATCGCCCCTAAGTCTCCACTGGGCCTGTTTATCCGGAAGCCTTTCA
TCAAGTTTATCTGCCACACCGCCAGCTATCTGACATTCCTGTTTCTGCTGCTGCTGGC
CTCCCAGCACATCGACAGATCTGATCTGAACAGGCAGGGCCACCCCTACCATCGT
GGAGTGGATGATCCTGCCATGGGTGCTGGGCTTCATCTGGGGCGAGATCAAGCAGA
TGTGGGACGGCGGCCTGCAGGACTACATCCACGATTGGTGGAACCTGATGGATTTTG
TGATGAATTCCCTGTACCTGGCCACAATCTCTCTGAAGATCGTGGCCTTCGTGAAGT
ATAGCGCCCTGAATCCCAGAGAGTCCTGGGACATGTGGCACCCCTACCCTGGTGGCA
GAGGCCCTGTTTCGCAATCGCCAACATCTTTTCCTCTCTGCGCCTGATCAGCCTGTTTA
CAGCCAATTCCCACCTGGGACCACTGCAGATCTCCCTGGGACGGATGCTGCTGGATA
TCCTGAAGTTCTGTATTATCTACTGCCTGGTGCTGCTGGCCTTCGCCAACGGCCTGAA
TCAGCTGTACTTCTACTATGAGGAGACCAAGGGCCTGACATGCAAGGGCATCCGCTG
TGAGAAGCAGAACAAATGCCTTCAGCACCCCTGTTTCGAGACACTGCAGTCTCTGTTCTG
GAGCATCTTTGGCCTGATCAACCTGTACGTGACCAATGTGAAGGCCAGCACGAGTT
CACAGAGTTTGTGGGCGCCACCATGTTTCGGCACATACAACGTGATCTCTCTGGTGGT
GCTGCTGAATATGCTGATCGCCATGATGAACAATAGCTATCAGCTGATCGCCGACCA
CGCCGATATCGAGTGGAAGTTCGCCCCGACCAAGCTGTGGATGTCCTACTTTGAGGA
GGGCGGCACCCTGCCCACACCTTTCAACGTGATCCCATCCCCAAGTCTCTGTGGTA
TCTGATCAAGTGGATCTGGACACACCTGTGCAAGAAGAAGATGCGCCGGAAGCCTG
AGAGCTTTGGCACCATCGGCGTGCGCACACAGCACAGAAGGGCAGCAGACAACCTG
CGCCGGCACCAACAGTACCAGGAAGTGATGCGCAATCTGGTGAAGCGGTATGTGGC
CGCCATGATCAGGGACGCAAAGACCGAGGAGGGACTGACAGAGGAGAACTTCAAG

GAGCTGAAGCAGGATATCAGCTCCTTCAGATTTGAGGTGCTGGGCCTGCTGAGGGG
CAGCAAGCTGTCCACCATCCAGTCCGCCAACGCCTCTAAGGAGTCTAGCAATTCTGC
CGACAGCGATGAGAAGAGCGACTCCGAGGGCAACTCTAAGGATAAGAAGAAGAAC
TTCAGCCTGTTTTGACCTGACCACACTGATCCACCCACGCAGCGCCGCAATCGCATCC
GAGCGGCACAACATCTCCAATGGCTCTGCCCTGGTGGTGCAGGAGCCACCAAGAGA
GAAGCAGAGGAAGGTGAACTTTGTGACAGATATCAAGAATTTCGGCCTGTTTTACA
GAAGGAGCAAGCAGAACGCCGCCGAGCAGAACGCCAATCAGATCTTCTCTGTGAGC
GAGGAGGTGGCAAGACAGCAGGCAGCAGGACCACTGGAGAGGAATATCCAGCTGG
AGAGCCGGGGACTGGCAAGCAGGGGCGACCTGTCCATCCCAGGACTGTCTGAGCAG
TGCGTGCTGGTGGACCACAGGGAGCGGAACACCGATACACTGGGACTGCAAGTGGG
CAAGCGGGTGTGCCCTTTCAAGAGCGAGAAGGTCGTGGTGGAGGACACCGTGCCCA
TCATCCCTAAGGAGAAGCACGCCAAGGAGGAGGATTCCTCTATCGACTACGATCTG
AATCTGCCAGACACCGTGACACACGAGGATTATGTGACCACAAGGCTGTGAGCGGC
CGCTCTAGAGGGCCCGTTTAAACCCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGC
CAGCCATCTGTTGTTTGCCCCCTCCCCCGTGCCCTTCCTTGACCCTGGAAGGTGCCACTC
CCACTGTCCTTTTCCTAATAAAAATGAGGAAATTGCATCGCATTGTCTGAGTAGGTGTC
ATTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGAC
AATAGCAGGCATGCTGGGGATGCGGTGGGCTCTATGGCTTCTGAGGCGGAAAGAAC
CAGCTGGGGCTCTAGGGGGTATCCCCACGCGCCCTGTAGCGGCGCATTAAAGCGCGG
CGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCCG
CTCCTTTTCGCTTTCTTCCCTTCCTTTCTCGCCACGTTTCGCCGGCTTTCCCCGTCAAGCT
CTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCC
AAAAAAGTTGATTAGGGTGATGGTTACGTAGTGGGCCATCGCCCTGATAGACGGTT
TTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAACTG
GAACAACACTCAACCCTATCTCGGTCTATTCTTTTGATTTATAAGGGATTTTGCCGAT
TTCGGCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAATTAATT
CTGTGGAATGTGTGTCTAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGCAGGCAG
AAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAGTCCCCAG
GCTCCCCAGCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCATA
GTCCCGCCCTAACTCCGCCCATCCCGCCCTAACTCCGCCAGTTCCGCCATTCTC
CGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCTCTGCCTC

TGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAA
GCTCCCGGGAGCTTGTATATCCATTTTCGGATCTGATCAGCACGTGATGAAAAAGCC
TGAACTCACCGCGACGTCTGTGCGAGAAGTTTCTGATCGAAAAGTTCGACAGCGTCTC
CGACCTGATGCAGCTCTCGGAGGGCGAAGAATCTCGTGCTTTCAGCTTCGATGTAGG
AGGGCGTGGATATGTCCTGCGGGTAAATAGCTGCGCCGATGGTTTCTACAAAGATCG
TTATGTTTATCGGCACTTTGCATCGGCCGCGCTCCCGATTCCGGAAGTGCTTGACATT
GGGGAATTCAGCGAGAGCCTGACCTATTGCATCTCCCGCCGTGCACAGGGTGTAC
GTTGCAAGACCTGCCTGAAACCGAACTGCCCCGCTGTTCTGCAGCCGGTTCGCGGAGG
CCATGGATGCGATCGCTGCGGCCGATCTTAGCCAGACGAGCGGGTTCGGCCCATTTCG
GACCGCAAGGAATCGGTCAATACACTACATGGCGTGATTTTCATATGCGCGATTGCTG
ATCCCCATGTGTATCACTGGCAAACCTGTGATGGACGACACCGTCAGTGCGTCCGTCTG
CGCAGGCTCTCGATGAGCTGATGCTTTGGGCCGAGGACTGCCCCGAAGTCCGGCAC
CTCGTGACGCGGATTTTCGGCTCCAACAATGTCCTGACGGACAATGGCCGCATAACA
GCGGTCATTGACTGGAGCGAGGCGATGTTTCGGGGATTCCCAATACGAGGTCGCCAA
CATCTTCTTCTGGAGGCCGTGGTTGGCTTGTATGGAGCAGCAGACGCGCTACTTCGA
GCGGAGGCATCCGGAGCTTGCAGGATCGCCGCGGCTCCGGGCGTATATGCTCCGCA
TTGGTCTTGACCAACTCTATCAGAGCTTGGTTGACGGCAATTTTCGATGATGCAGCTT
GGGCGCAGGGTCGATGCGACGCAATCGTCCGATCCGGAGCCGGGACTGTGCGGCGT
ACACAAATCGCCCCGAGAAGCGCGGCCGTCTGGACCGATGGCTGTGTAGAAGTACT
CGCCGATAGTGGAACCGACGCCCCAGCACTCGTCCGAGGGCAAAGGAATAGCACG
TGCTACGAGATTTTCGATTCCACCGCCGCTTCTATGAAAGGTTGGGCTTCGGAATCG
TTTTCCGGGACGCCGGCTGGATGATCCTCCAGCGCGGGGATCTCATGCTGGAGTTCT
TCGCCCCACCCAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCA
TCACAAATTTACAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAA
ACTCATCAATGTATCTTATCATGTCTGTATACCGTCGACCTCTAGCTAGAGCTTGGCG
TAATCATGGTCATAGCTGTTTCCTGTGTGAAATTGTTATCCGCTCACAATTCCACACA
ACATACGAGCCGGAAGCATAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTAA
CTCACATTAATTGCGTTGCGCTCACTGCCCCGCTTTCAGTCGGGAAACCTGTCGTGC
CAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTTCGTATTGGGCG
CTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTTCGTTTCGGCTGCGGCGAGCG
GTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGC

AGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCC
GCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGA
CGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTCC
CCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTG
TCCGCCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCTGTAGGTATC
TCAGTTCGGTGTAGGTGCTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTC
AGCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGAC
ACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTAT
GTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAG
AACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGG
TAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTTGGTTTTTTGTTTGCAA
GCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTAC
GGGGTCTGACGCTCAGTGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAGATT
ATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAAT
CTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGC
ACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTG
TAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCG
CGAGACCCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAG
GGCCGAGCGCAGAAGTGGTCCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTG
TTGCCGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGCGCAACGTTGTTGC
CATTGCTACAGGCATCGTGGTGTACGCTCGTCGTTTGGTATGGCTTCATTCAGCTCC
GGTTCCCAACGATCAAGGCGAGTTACATGATCCCCATGTTGTGCAAAAAAGCGGTT
AGCTCCTTCGGTCCCTCCGATCGTTGTCAGAAGTAAGTTGGCCGCAGTGTTATCACTC
ATGGTTATGGCAGCACTGCATAATTCTCTTACTGTTCATGCCATCCGTAAGATGCTTTT
CTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGA
GTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAA
AAGTGCTCATCATTGGAACGTTCTTCGGGGCGAAAACTCTCAAGGATCTTACCGC
TGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAAGTATCTTCAGCATCTTT
TACTTTACCAAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAAATGCCGCAAAAA
AGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAATATT
ATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTA

GAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGACG
TC

SEQ ID NO:2 TRPC5 Plasmid Sequence

The DNA sequence of the TRPC5 plasmid used in Example 25 is included below.
Underlined nucleic acids represent those encoding human TRPC5.

GACGGATCGGGAGATCTCCCGATCCCCTATGGTGCACTCTCAGTACAATCTGCTCTG
ATGCCGCATAGTTAAGCCAGTATCTGCTCCCTGCTTGTGTGTTGGAGGTCGCTGAGT
AGTGCGCGAGCAAAATTTAAGCTACAACAAGGCAAGGCTTGACCGACAATTGCATG
AAGAATCTGCTTAGGGTTAGGCGTTTTGCGCTGCTTCGCGATGTACGGGCCAGATAT
ACGCGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTA
GTTTCATAGCCCATATATGGAGTTCCGCGTTACATAAATTACGGTAAATGGCCCCGCT
GGCTGACCGCCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCAT
GTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACT
GCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTC
AATGACGGTAAATGGCCCCGCTGGCATTATGCCCAGTACATGACCTTATGGGACTTT
CCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTT
GGCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCC
ACCCCATTGACGTCAATGGGAGTTTGTTTTGGCACCAAAATCAACGGGACTTTCCAA
AATGTCGTAACAACCTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGG
GAGGTCTATATAAGCAGAGCTCTCCCTATCAGTGATAGAGATCTCCCTATCAGTGAT
AGAGATCGTCGACGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCC
ACGCTGTTTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCGGACTCTAGC
GTTTAAACTTAAGCCCAAGCTGGCTAGACCGCCATGGCCCAACTGTACTACAAAAA
GGTCAACTACTCACCGTACAGAGACCGCATCCCCCTGCAAATTGTGAGGGCTGAGA
CAGAGCTCTCTGCAGAGGAGAAGGCCTTCCTCAATGCTGTGGAGAAGGGGGACTAT
GCCACTGTGAAGCAGGCCCTTCAGGAGGCTGAGATCTACTATAATGTTAACATCAAC
TGCATGGACCCCTTGGGCCGGAGTGCCCTGCTCATTGCCATTGAGAACGAGAACCTG
GAGATCATGGAGCTACTGCTGAACCACAGCGTGTATGTGGGTGATGCATTGCTCTAT

GCCATACGCAAGGAAGTGGTGGGCGCTGTGGAGCTTCTGCTCAGCTACAGGCGGCC
CAGCGGAGAGAAGCAGGTCCCCACTCTGATGATGGACACGCAGTTCTCTGAATTCA
CACCGGACATCACTCCCATCATGCTGGCTGCCCACACCAACAACCTACGAAATCATCA
AACTGCTTGTCCAAAAACGGGTCACTATCCCACGGCCCCACCAGATCCGCTGCAACT
GTGTGGAGTGTGTGTCTAGTTCAGAGGTAGACAGCCTGCGCCACTCTCGCTCCCGAC
TGAACATCTATAAGGCTCTGGCAAGCCCCCTCACTCATTGCCTTATCAAGTGAGGACC
CCATCCTAACTGCCTTCCGTCTGGGCTGGGAGCTCAAGGAGCTCAGCAAGGTGGAG
AATGAGTTCAAGGCCGAGTATGAGGAGCTCTCTCAGCAGTGCAAGCTCTTTGCCAAA
GACCTGCTGGACCAAGCTCGGAGCTCCAGGGAACCTGGAGATCATCCTCAACCATCG
AGATGACCACAGTGAAGAGCTTGACCCCTCAGAAGTACCATGACCTGGCCAAGTTGA
AGGTGGCAATCAAATACCACCAGAAAAGAGTTTGTGCTCAGCCCAACTGCCAACAG
TTGCTTGCCACCCTGTGGTATGATGGCTTCCCTGGATGGCGGCGGAAACACTGGGTA
GTCAAGCTTCTAACCTGCATGACCATTGGGTTCCTGTTTCCCATGCTGTCTATAGCCT
ACCTGATCTCACCCAGGAGCAACCTTGGGCTGTTTCATCAAGAAACCCTTTATCAAGT
TTATCTGCCACACAGCATCCTATTTGACCTTCCTCTTTATGCTTCTCCTGGCTTCTCAG
CACATTGTGAGGACAGACCTTCATGTACAGGGGCCTCCCCCAACTGTCGTGGAATGG
ATGATATTGCCTTGGGTTCTAGGTTTCATTTGGGGTGAGATTAAGGAAATGTGGGAT
GGTGGATTTACTGAATACATCCATGACTGGTGGAACTGATGGATTTTGCAATGAAC
TCCCTCTACCTGGCAACTATTTCCCTGAAGATTGTGGCCTATGTCAAGTATAATGGTT
CTCGTCCAAGGGAGGAATGGGAAATGTGGCACCCGACTCTGATTGCGGAAGCACTC
TTGCAATATCCAACATTTTAAGTTCGTTGCGTCTCATATCCCTGTTACAGCCAACT
CCCCTTAGGACCTCTGCAGATCTCTTTGGGACGCATGCTGCTTGATATCCTCAAATT
CCTCTTTATCTACTGCCTGGTACTACTAGCTTTTGCCAATGGACTGAACCAGCTTTAC
TTCTATTATGAAACCAGAGCTATCGATGAGCCTAACAACTGCAAGGGGATCCGATGT
GAGAAACAGAACAAATGCCTTCTCCACGCTCTTTGAGACTCTTCAGTCACTCTTCTGG
TCTGTATTTGGCCTTTTAAATCTATATGTCACCAATGTGAAAGCCAGACACGAATTC
ACCGAGTTTGTAGGAGCTACCATGTTTGGAACATACAATGTCATCTCCCTGGTAGTG
CTGCTGAACATGCTGATTGCTATGATGAACAACTCCTATCAGCTTATTGCCGATCAT
GCTGATATCGAGTGGAAGTTTGCAAGGACGAAGCTCTGGATGAGTTACTTTGATGAA
GGTGGCACCTTGCCACCTCCTTTCAACATCATCCCCAGCCCCAAGTCATTTCTATACC
TTGGTAACTGGTTCAACAACACCTTCTGCCCCAAAAGAGACCCTGACGGTAGACGG

AGAAGGCGCAACTTGAGAAGTTTCACAGAACGCAATGCTGACAGCCTGATACAAAA
TCAACATTATCAGGAAGTTATCAGGAATTTAGTCAAAAGATATGTGGCTGCTATGAT
AAGAAATTCCAAAACACATGAGGGACTTACAGAAGAAAATTTAAGGAATTAAAGC
AAGACATCTCCAGCTTTCGGTATGAAGTGCTTGACCTCTTGGGAAATAGAAAACATC
CAAGGAGCTTTTCCACTAGCAGCACTGAACTGTCTCAGAGAGACGATAATAATGAT
GGCAGTGGTGGGGCTCGGGCCAAATCCAAGAGTGTCTCTTTTAATTTAGGCTGCAAG
AAAAAGACTTGCCATGGGCCACCTCTCATCAGAACCATGCCAAGGTCCAGTGGTGC
CCAAGGAAAGTCAAAAGCTGAGTCATCAAGCAAACGCTCCTTCATGGGTCCTTCTCT
CAAGAACTGGGTCTCCTATTCTCCAAATTTAATGGTCATATGTCTGAACCCAGTTC
AGAGCCAATGTACACAATTTCTGATGGAATTGTTACGCAGCACTGTATGTGGCAGGA
CATCAGATATTCTCAGATGGAGAAAGGGAAAGCAGAGGCCTGTTCTCAAAGTGAAA
TTAACCTCAGTGAGGTAGAATTAGGTGAAGTCCAGGGCGCTGCTCAGAGCAGTGAA
TGCCCTCTAGCCTGTTCCAGCTCTCTTCACTGTGCATCCAGCATCTGCTCCTCAAATT
CTAAACTTTTAGACTCCTCAGAGGATGTATTTGAAACTTGGGGAGAGGCTTGTGACT
TGCTCATGCACAAATGGGGTGATGGACAGGAAGAACAAGTTACAACCTCGCCTCTAA
TGACTCGAGTCTAGAGGGCCCGTTTAAACCCGCTGATCAGCCTCGACTGTGCCTTCT
AGTTGCCAGCCATCTGTTGTTTGCCCCCTCCCCCGTGCCCTTCCTTGACCCTGGAAGGTG
CCACTCCCCTGTCCTTTCCTAATAAAATGAGGAAATTGCATCGCATTGTCTGAGTA
GGTGTCAATTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGG
GAAGACAATAGCAGGCATGCTGGGGATGCGGTGGGCTCTATGGCTTCTGAGGCGGA
AAGAACCAGCTGGGGCTCTAGGGGGTATCCCCACGCGCCCTGTAGCGGCGCATTA
GCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTA
GCGCCCGCTCCTTTTCGCTTTCTTCCCTTCCTTTCTCGCCACGTTCCGCGGCTTTCCCCG
TCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTC
GACCCCAAAAACTTGATTAGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAG
ACGGTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCC
AAACTGGAACAACACTCAACCTATCTCGGTCTATTCTTTTGATTTATAAGGGATTTT
GCCGATTTTCGGCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAA
TTAATTCTGTGGAATGTGTGTCAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGCA
GGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAGTC
CCCAGGCTCCCCAGCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAA

CCATAGTCCCCGCCCTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCCGCCC
ATTCTCCGCCCCATGGCTGACTAATTTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTC
TGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTG
CAAAAAGCTCCCGGGAGCTTGTATATCCATTTTCGGATCTGATCAGCACGTGATGAA
AAAGCCTGAACTCACCGCGACGTCTGTCGAGAAGTTTCTGATCGAAAAGTTCGACA
GCGTCTCCGACCTGATGCAGCTCTCGGAGGGCGAAGAATCTCGTGCTTTCAGCTTCG
ATGTAGGAGGGCGTGGATATGTCCTGCGGGTAAATAGCTGCGCCGATGGTTTCTACA
AAGATCGTTATGTTTATCGGCACCTTTCATCGGCCGCGCTCCCGATTCCGGAAGTGC
TTGACATTGGGGAATTCAGCGAGAGCCTGACCTATTGCATCTCCCGCCGTGCACAGG
GTGTACGTTGCAAGACCTGCCTGAAACCGAACTGCCCCGCTGTTCTGCAGCCGGTGC
CGGAGGCCATGGATGCGATCGCTGCGGCCGATCTTAGCCAGACGAGCGGGTTCGGC
CCATTCGGACCGCAAGGAATCGGTCAATACACTACATGGCGTGATTTTCATATGCGCG
ATTGCTGATCCCCATGTGTATCACTGGCAAACGTGTGATGGACGACACCGTCAGTGCG
TCCGTCGCGCAGGCTCTCGATGAGCTGATGCTTTGGGCCGAGGACTGCCCCGAAGTC
CGGCACCTCGTGACGCGGATTTTCGGCTCCAACAATGTCCTGACGGACAATGGCCGC
ATAACAGCGGTCATTGACTGGAGCGAGGCGATGTTTCGGGGATTCCCAATACGAGGT
CGCCAACATCTTCTTCTGGAGGCCGTGGTTGGCTTGTATGGAGCAGCAGACGCGCTA
CTTCGAGCGGAGGCATCCGGAGCTTGCAGGATCGCCGCGGCTCCGGGCGTATATGC
TCCGCATTGGTCTTGACCAACTCTATCAGAGCTTGGTTGACGGCAATTTTCGATGATG
CAGCTTGGGCGCAGGGTTCGATGCGACGCAATCGTCCGATCCGGAGCCGGGACTGTC
GGGCGTACACAAATCGCCCCGAGAAGCGCGGCCGTCTGGACCGATGGCTGTGTAGA
AGTACTCGCCGATAGTGGAACCGACGCCCCAGCACTCGTCCGAGGGCAAAGGAAT
AGCACGTGCTACGAGATTTTCGATTCCACCGCCGCCTTCTATGAAAGGTTGGGCTTCG
GAATCGTTTTCCGGGACGCCGGCTGGATGATCCTCCAGCGCGGGGATCTCATGCTGG
AGTTCTTCGCCCACCCCAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCA
ATAGCATCACAAATTTACAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTT
GTCCAAACTCATCAATGTATCTTATCATGTCTGTATAACGTCGACCTCTAGCTAGAGC
TTGGCGTAATCATGGTCATAGCTGTTTCTGTGTGAAATTGTTATCCGCTCACAATTC
CACACAACATACGAGCCGGAAGCATAAAGTGTAAGCCTGGGGTGCCTAATGAGTG
AGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCCGCTTTCAGTCGGGAAACCTG
TCGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGGCGGTTTGCCTATT

GGGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTTCGGCTGCGGC
GAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGAT
AACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAA
AGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAA
ATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCG
TTCCCCCTGGAAGCTCCCTCGTGCCTCTCCTGTTCCGACCCTGCCGCTTACCGGAT
ACCTGTCCGCCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCTGTAG
GTATCTCAGTTCGGTGTAGGTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCC
CGTTCAGCCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGT
AAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGA
GGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTA
GAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAG
TTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTTGGTTTTTTGTTTG
CAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTT
CTACGGGGTCTGACGCTCAGTGGAACGAAAACCTACGTTAAGGGATTTTGGTCATGA
GATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAATAAATGAAGTTTTAAAT
CAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTG
AGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGT
CGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGAT
ACCGCGAGACCCACGCTACCCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCG
GAAGGGCCGAGCGCAGAAGTGGTCCCTGCAACTTTATCCGCCTCCATCCAGTCTATTA
ATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTTCGCAACGTTG
TTGCCATTGCTACAGGCATCGTGGTGTCACGCTCGTCGTTTGGTATGGCTTCATTAG
CTCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCAAAAAAGC
GGTTAGCTCCTTCGGTCCTCCGATCGTTGTCAGAAGTAAGTTGGCCGCAGTGTTATC
ACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATGCCATCCGTAAGATG
CTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCG
ACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAA
CTTTAAAAGTGCTCATTCATTGAAAAACGTTCTTCGGGGCGAAAACCTCTCAAGGATCT
TACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAACTGATCTTCAG
CATCTTTTACTTTCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCG

CAAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTC
AATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAAT
GTATTTAGAAAAATAAACAAATAGGGGTTCGCGCACATTTCCCCGAAAAGTGCCA
CCTGACGTC