

CLAIMS

1. A pharmaceutical composition comprising a bispecific single chain antibody construct that binds to a target cell surface antigen selected from CD19 and CD33, via a first binding domain and to the T cell surface antigen CD3 via a second binding domain, a β -cyclodextrin and a buffer, wherein the bispecific single chain antibody construct comprises a sequence selected from the group consisting of SEQ ID NO: 100 and 174,

wherein the binding domains have pairs of VH regions and VL regions in the format of a single chain antibody (scFv), and wherein the bispecific single chain antibody construct optionally comprises a domain for extending serum half-life which binds to Fc domains,

wherein the buffer is selected from the group consisting of potassium phosphate, acetic acid/sodium acetate, citric acid/sodium citrate, succinic acid/sodium succinate, tartaric acid/sodium tartrate, histidine/histidine HCl, glycine, Tris, glutamate, acetate and mixtures thereof,

wherein the composition is characterized by a reduction or prevention of the formation of protein aggregates (high molecular weight species (HMWS)), and

wherein the β -cyclodextrin is sulfobutylether- β -cyclodextrin.

2. The composition according to claim 1, wherein the β -cyclodextrin is present in a concentration in the range of 0.1% to 20% (w/v).

3. The composition according to claim 1, wherein the bispecific single chain antibody construct is present in a concentration range of 0.1-5mg/ml.

4. The composition according to claim 1, wherein the buffer is selected from the group consisting of potassium phosphate, citric acid/sodium citrate, succinic acid, histidine, glutamate, acetate and combinations thereof.

5. The composition according to claim 1, wherein the pH of the composition is in the range of 4-7.5.

6. The composition according to claim 1, further comprising one or more excipients selected from the group consisting of sucrose, trehalose, mannitol, sorbitol, arginine, lysine, polysorbate 20, polysorbate 80, poloxamer 188, pluronic and combinations thereof.

7. The composition according to claim 1, further comprising one or more preservatives.

8. The composition according to claim 7, wherein the one or more preservative is selected from the group consisting of benzyl alcohol, chlorobutanol, phenol, meta-cresol, methylparaben, phenoxyethanol, propylparaben and thimerosal.

9. The composition according to claim 1, wherein the first binding domain comprises the amino acid sequence selected from the group consisting of SEQ ID NO: 99, 109, 119, 128, 137, 146, 155, 164, 173, 183, 185 and 187, and the second binding domain comprises the amino acid sequence selected from the group consisting of SEQ ID NO: 9, 18, 27, 36, 45, 54, 63, 72, 81, 179 and 90.

10. The composition according to claim 9, wherein the composition is free of preservatives, wherein the first and second binding domains of the antibody comprise the amino acid sequence set forth in SEQ ID NO 100 and the antibody is present at a concentration of about 0.5 mg/ml, wherein the β -cyclodextrin is sulfobutylether- β -cyclodextrin sodium salt and is present at a concentration of about 1% (w/v), wherein the buffer is potassium phosphate and is present at a concentration of about 10 mM, and wherein the composition further comprises sucrose at a concentration of about 8% (w/v) and polysorbate 80 at a concentration of about 0.01% (w/v) at a pH of about 6.0.

11. The composition according to claim 2, wherein the β -cyclodextrin is present in a concentration in the range of 0.5% to 2% (w/v).

12. The composition according to claim 11, wherein the β -cyclodextrin is present in a concentration in the range of 0.8% to 1.5% (w/v).

13. The composition according to claim 3, wherein the bispecific single chain antibody construct is present in a concentration range of 0.2-2.5 mg/ml.

14. The composition according to claim 13, wherein the bispecific single chain antibody construct is present in a concentration range of 0.25-1.0 mg/ml.

15. The composition according to claim 1, wherein the amino acid sequence of the bispecific single chain antibody construct is SEQ ID NO: 174.

16. The composition according to claim 9, wherein the cyclodextrin is sulfobutylether- β -cyclodextrin sodium salt in a concentration of about 1% (w/v) and the buffer is potassium phosphate in concentration of about 10 mM, and wherein the composition further comprises mannitol in a concentration of about 4%, sucrose in a concentration of about 2% (w/v), and polysorbate 80 in concentration of about 0.01% (w/v) at a pH of about 6.0.