

COMPOUNDS, COMPOSITIONS AND METHODS

CROSS-REFERENCE TO RELATED PATENT APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application number 60/373,454, filed April 17, 2002; and of U.S. Provisional Patent Application number 60/410,682, filed September 13, 2002. Each of these applications is incorporated herein by reference for all purposes.

FIELD OF THE INVENTION

[0002] This invention relates to compounds which are inhibitors of the mitotic kinesin KSP and are useful in the treatment of cellular proliferative diseases, for example cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders, fungal disorders, and inflammation.

BACKGROUND OF THE INVENTION

[0003] Among the therapeutic agents used to treat cancer are the taxanes and vinca alkaloids, which act on microtubules. Microtubules are the primary structural element of the mitotic spindle. The mitotic spindle is responsible for distribution of replicate copies of the genome to each of the two daughter cells that result from cell division. It is presumed that disruption of the mitotic spindle by these drugs results in inhibition of cancer cell division, and induction of cancer cell death. However, microtubules form other types of cellular structures, including tracks for intracellular transport in nerve processes. Because these agents do not specifically target mitotic spindles, they have side effects that limit their usefulness.

[0004] Improvements in the specificity of agents used to treat cancer is of considerable interest because of the therapeutic benefits which would be realized if the side effects associated with the administration of these agents could be reduced. Traditionally, dramatic improvements in the treatment of cancer are associated with identification of therapeutic agents acting through novel mechanisms. Examples of this include not only the taxanes, but also the camptothecin class of topoisomerase I inhibitors. From both of these perspectives, mitotic kinesins are attractive targets for new anti-cancer agents.

[0005] Mitotic kinesins are enzymes essential for assembly and function of the mitotic spindle, but are not generally part of other microtubule structures, such as in nerve processes. Mitotic kinesins play essential roles during all phases of mitosis. These enzymes are "molecular motors" that transform energy released by hydrolysis of ATP into mechanical force which drives the directional movement of cellular cargoes along microtubules. The catalytic domain sufficient for this task is a compact structure of approximately 340 amino acids. During mitosis, kinesins organize microtubules into the bipolar structure that is the mitotic spindle. Kinesins mediate movement of chromosomes along spindle microtubules, as well as structural changes in the mitotic spindle associated with specific phases of mitosis. Experimental perturbation of mitotic kinesin function causes malformation or dysfunction of the mitotic spindle, frequently resulting in cell cycle arrest and cell death.

[0006] Among the mitotic kinesins which have been identified is KSP. KSP belongs to an evolutionarily conserved kinesin subfamily of plus end-directed microtubule motors that assemble into bipolar homotetramers consisting of antiparallel homodimers. During mitosis KSP associates with microtubules of the mitotic spindle. Microinjection of antibodies directed against KSP into human cells prevents spindle pole separation during prometaphase, giving rise to monopolar spindles and causing mitotic arrest and induction of programmed cell death. KSP and related kinesins in other, non-human, organisms, bundle antiparallel microtubules and slide them relative to one another, thus forcing the two spindle poles apart. KSP may also mediate in anaphase B spindle elongation and focussing of microtubules at the spindle pole.

[0007] Human KSP (also termed HsEg5) has been described (Blangy, et al., *Cell*, 83:1159-69 (1995); Whitehead, et al., *Arthritis Rheum.*, 39:1635-42 (1996); Galgio et al., *J. Cell Biol.*, 135:339-414 (1996); Blangy, et al., *J Biol. Chem.*, 272:19418-24 (1997); Blangy, et al., *Cell Motil Cytoskeleton*, 40:174-82 (1998); Whitehead and Rattner, *J. Cell Sci.*, 111:2551-61 (1998); Kaiser, et al., *JBC* 274:18925-31 (1999); GenBank accession numbers: X85137, NM004523 and U37426), and a fragment of the KSP gene (TRIP5) has been described (Lee, et al., *Mol Endocrinol.*, 9:243-54 (1995); GenBank accession number L40372). *Xenopus* KSP homologs (Eg5), as well as *Drosophila* KLP61 F/KRP1 30 have been reported.


[0008] Mitotic kinesins, including KSP, are attractive targets for the discovery and development of novel antimitotic chemotherapeutics. Accordingly, it is an object of the present invention to provide compounds, compositions and methods useful in the inhibition

of KSP.

SUMMARY OF THE INVENTION

[0009] In accordance with the objects outlined above, the present invention provides compounds that can be used to treat cellular proliferative diseases. The compounds are KSP inhibitors, particularly human KSP inhibitors. The present invention also provides compositions comprising such compounds, and methods utilizing such compounds or compositions, which can be used to treat cellular proliferative diseases.

[0010] In one aspect, the invention relates to methods for treating cellular proliferative diseases, and for treating disorders by inhibiting the activity of KSP. The methods employ compounds represented by Formula I:



Formula I

wherein:

R_1 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-;

R_2 and $R_{2'}$ are independently chosen from hydrogen, optionally substituted alkyl-, optionally substituted alkoxy, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-; or R_2 and $R_{2'}$ taken together form an optionally substituted 3- to 7-membered ring;

R_{12} is selected from the group consisting of optionally substituted imidazolyl, optionally substituted imidazoliny, $-NHR_4$; $-N(R_4)(COR_3)$; $-N(R_4)(SO_2R_{3a})$; and -

$N(R_4)(CH_2R_{3b})$;

R_3 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, optionally substituted heteroaralkyl-, $R_{15}O$ - and $R_{17}NH$ -;

R_{3a} is chosen from optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, optionally substituted heteroaralkyl-, and $R_{17}NH$ -;

R_{3b} is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heterocycl-, and optionally substituted heteroaralkyl-;

R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, acyl, optionally substituted alkyl-, optionally substituted alkoxy, halogen, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl-, alkylsulfonamido-, alkylthio-, carboxyalkyl-, carboxamido-, aminocarbonyl-, optionally substituted aryl and optionally substituted heteroaryl-;

R_{15} is chosen from optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-; and

R_{17} is hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-,

optionally substituted heteroaryl-, or optionally substituted hetero-aralkyl-, including single stereoisomers, mixtures of stereoisomers;

a pharmaceutically acceptable salt of a compound of Formula I;

a pharmaceutically acceptable solvate of a pharmaceutically acceptable solvate of a compound of Formula I; or

a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I.

[0011] According to one embodiment, when either R_2 or $R_{2'}$ is hydrogen, the other is not hydrogen. In another embodiment, R_2 and $R_{2'}$ are each hydrogen; and R_1 is chosen from optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl, provided however, that R_1 is not substituted phenyl.

[0012] In one aspect, the invention relates to methods for treating cellular proliferative diseases and other disorders that can be treated by inhibiting KSP by the administration of a therapeutically effective amount of a compound of Formula I; a pharmaceutically acceptable salt of a compound of Formula I; or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I thereof. Such diseases and disorders include cancer, hyperplasia, restenosis, cardiac hypertrophy, immune disorders, fungal disorders and inflammation.

[0013] In another aspect, the invention relates to compounds useful in inhibiting KSP kinesin. The compounds have the structures shown above in Formula I; a pharmaceutically acceptable salt of a compound of Formula I; or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I. The invention also relates to pharmaceutical compositions containing a therapeutically effective amount of a compound of Formula I; a pharmaceutically acceptable salt of a compound of Formula I; or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I, admixed with at least one pharmaceutical excipient. In another aspect, the composition further comprises a chemotherapeutic agent other than a compound of the present invention.

[0014] In an additional aspect, the present invention provides methods of screening for compounds that will bind to a KSP kinesin, for example compounds that will displace or compete with the binding of a compound of the invention. The methods comprise combining a labeled compound of the invention, a KSP kinesin, and at least one candidate agent and determining the binding of the candidate agent to the KSP kinesin.

[0015] In a further aspect, the invention provides methods of screening for modulators of KSP kinesin activity. The methods comprise combining a compound of the invention, a KSP kinesin, and at least one candidate agent and determining the effect of the candidate agent on the KSP kinesin activity.

BRIEF DESCRIPTION OF THE FIGURES

The invention will be better understood by reference to the following description taken in conjunction with the accompanying drawings.

Figures 1 and 2 represent a XRPD and MDSC scan, respectively, of a salt prepared according to Preparation A.

Figure 3 represents an XRPD scan of a salt prepared according to Preparation B.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0016] As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise. The following abbreviations and terms have the indicated meanings throughout:

Ac	=	acetyl
Boc	=	t-butyloxy carbonyl
Bu	=	butyl
c-	=	cyclo
CBZ	=	carbobenzoxy = benzyloxycarbonyl
DCM	=	dichloromethane = methylene chloride = CH_2Cl_2
DIEA	=	N,N-diisopropylethylamine
DMF	=	N,N-dimethylformamide
DMSO	=	dimethyl sulfoxide
Et	=	ethyl
HBTU	=	O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate
HMDS	=	hexamethyldisilazane
HOAc	=	acetic acid
IPA	=	isopropyl alcohol
Me	=	methyl
Ph	=	phenyl
Py	=	pyridine
rt	=	room temperature
sat'd	=	saturated
s-	=	secondary
t-	=	tertiary
TEA	=	triethylamine
TFA	=	trifluoroacetic acid
THF	=	tetrahydrofuran
Tf	=	triflate

[0017] **Alkyl** is intended to include linear, branched, or cyclic aliphatic hydrocarbon structures and combinations thereof, which structures may be saturated or unsaturated. **Lower-alkyl** refers to alkyl groups of from 1 to 5 carbon atoms, preferably from 1 to 4 carbon atoms. Examples of lower-alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, s-and t-butyl and the like. Preferred alkyl groups are those of C₂₀ or below. More preferred alkyl groups are those of C₁₃ or below. **Cycloalkyl** is a subset of alkyl and includes cyclic aliphatic hydrocarbon groups of from 3 to 13 carbon atoms. Examples of cycloalkyl groups include c-propyl, c-butyl, c-pentyl, norbornyl, adamantyl and the like. **Cycloalkyl-alkyl-** is another subset of alkyl and refers to cycloalkyl attached to the parent structure through a non-cyclic alkyl. Examples of cycloalkyl-alkyl- include cyclohexylmethyl, cyclopropylmethyl, cyclohexylpropyl, and the like. In this application, alkyl includes alkanyl, alkenyl and alkynyl residues; it is intended to include vinyl, allyl, isoprenyl and the like. **Alkylene-, alkenylene-, and alkynylene-** are other subsets of alkyl, including the same residues as alkyl, but having two points of attachment within a chemical structure. Examples of alkylene include ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂-), dimethylpropylene (-CH₂C(CH₃)₂CH₂-) and cyclohexylpropylene (-CH₂CH₂CH(C₆H₁₃)-). Likewise, examples of alkenylene include ethenylene (-CH=CH-), propenylene (-CH=CH-CH₂-), and cyclohexylpropenylene (-CH=CHCH(C₆H₁₃)-). Examples of alkynylene include ethynylene (-C≡C-) and propynylene (-CH≡CH-CH₂-). When an alkyl residue having a specific number of carbons is named, all geometric isomers having that number of carbons are intended to be encompassed; thus, for example, "butyl" is meant to include n-butyl, sec-butyl, isobutyl and t-butyl; "propyl" includes n-propyl, isopropyl, and c-propyl.

[0018] **Alkoxy or alkoxyl** refers to an alkyl group, preferably including from 1 to 8 carbon atoms, of a straight, branched, or cyclic configuration, or a combination thereof, attached to the parent structure through an oxygen (i.e., the group alkyl-O-). Examples include methoxy-, ethoxy-, propoxy-, isopropoxy-, cyclopropyloxy-, cyclohexyloxy- and the like. **Lower-alkoxy** refers to alkoxy groups containing one to four carbons.

[0019] **Acyl** refers to groups of from 1 to 8 carbon atoms of a straight, branched, or cyclic configuration or a combination thereof, attached to the parent structure through a carbonyl functionality. Such groups may be saturated or unsaturated, and aliphatic or aromatic. One or more carbons in the acyl residue may be replaced by nitrogen, oxygen or sulfur as long as the point of attachment to the parent remains at the carbonyl. Examples

include acetyl, benzoyl, propionyl, isobutyryl, t-butoxycarbonyl, benzyloxycarbonyl and the like. **Lower-acyl** refers to acyl groups containing one to four carbons.

[0020] Amino refers to the group -NH_2 . The term "substituted amino" refers to the group -NHR or -NRR where each R is independently selected from the group: optionally substituted alkyl, optionally substituted alkoxy, optionally substituted amino carbonyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, acyl, alkoxycarbonyl, sulfanyl, sulfinyl and sulfonyl, e.g., diethylamino, methylsulfonylamino, furanyl-oxy-sulfonamino.

[0021] Aminocarbonyl- refers to the group $\text{-NR}^c\text{COR}^b$, $\text{-NR}^c\text{CO}_2\text{R}^b$, or $\text{-NR}^c\text{CONR}^b\text{R}^c$, where

R^b is H or optionally substituted $\text{C}_1\text{-C}_6$ alkyl, aryl, heteroaryl, aryl- $\text{C}_1\text{-C}_4$ alkyl-, or heteroaryl- $\text{C}_1\text{-C}_4$ alkyl- group; and

R^c is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; and

where each optionally substituted R^b group is independently unsubstituted or substituted with one or more substituents independently selected from $\text{C}_1\text{-C}_4$ alkyl, aryl, heteroaryl, aryl- $\text{C}_1\text{-C}_4$ alkyl-, heteroaryl- $\text{C}_1\text{-C}_4$ alkyl-, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{-OC}_1\text{-C}_4$ alkyl, $\text{-OC}_1\text{-C}_4$ alkylphenyl, $\text{-C}_1\text{-C}_4$ alkyl-OH, $\text{-OC}_1\text{-C}_4$ haloalkyl, halogen, -OH , -NH_2 , $\text{-C}_1\text{-C}_4$ alkyl- NH_2 , $\text{-N(C}_1\text{-C}_4\text{ alkyl)(C}_1\text{-C}_4\text{ alkyl)}$, $\text{-NH(C}_1\text{-C}_4\text{ alkyl)}$, $\text{-N(C}_1\text{-C}_4\text{ alkyl)(C}_1\text{-C}_4\text{ alkylphenyl)}$, $\text{-NH(C}_1\text{-C}_4\text{ alkylphenyl)}$, cyano, nitro, oxo (as a substituent for heteroaryl), $\text{-CO}_2\text{H}$, $\text{-C(O)OC}_1\text{-C}_4\text{ alkyl}$, $\text{-CON(C}_1\text{-C}_4\text{ alkyl)(C}_1\text{-C}_4\text{ alkyl)}$, $\text{-CONH(C}_1\text{-C}_4\text{ alkyl)}$, -CONH_2 , $\text{-NHC(O)(C}_1\text{-C}_4\text{ alkyl)}$, -NHC(O)(phenyl) , $\text{-N(C}_1\text{-C}_4\text{ alkyl)C(O)(C}_1\text{-C}_4\text{ alkyl)}$, $\text{-N(C}_1\text{-C}_4\text{ alkyl)C(O)(phenyl)}$, $\text{-C(O)C}_1\text{-C}_4\text{ alkyl}$, $\text{-C(O)C}_1\text{-C}_4\text{ phenyl}$, $\text{-C(O)C}_1\text{-C}_4\text{ haloalkyl}$, $\text{-OC(O)C}_1\text{-C}_4\text{ alkyl}$, $\text{-SO}_2\text{(C}_1\text{-C}_4\text{ alkyl)}$, $\text{-SO}_2\text{(phenyl)}$, $\text{-SO}_2\text{(C}_1\text{-C}_4\text{ haloalkyl)}$, $\text{-SO}_2\text{NH}_2$, $\text{-SO}_2\text{NH(C}_1\text{-C}_4\text{ alkyl)}$, $\text{-SO}_2\text{NH(phenyl)}$, $\text{-NHSO}_2\text{(C}_1\text{-C}_4\text{ alkyl)}$, $\text{-NHSO}_2\text{(phenyl)}$, and $\text{-NHSO}_2\text{(C}_1\text{-C}_4\text{ haloalkyl)}$.

[0022] Antimitotic refers to a drug for inhibiting or preventing mitosis, for example, by causing metaphase arrest. Some antitumour drugs block proliferation and are considered antimitotics.

[0023] Aryl and heteroaryl mean a 5- or 6-membered aromatic or heteroaromatic ring containing 0 or 1-4 heteroatoms, respectively, selected from O, N, or S; a bicyclic 9- or 10-membered aromatic or heteroaromatic ring system containing 0 or 1-4 (or more) heteroatoms, respectively, selected from O, N, or S; or a tricyclic 12- to 14-membered aromatic or heteroaromatic ring system containing 0 or 1-4 (or more) heteroatoms,

respectively, selected from O, N, or S. The aromatic 6- to 14-membered carbocyclic rings include, e.g., phenyl, naphthyl, indanyl, tetralinyl, and fluorenyl and the 5- to 10-membered aromatic heterocyclic rings include, e.g., imidazolyl, pyridinyl, indolyl, thienyl, benzopyranonyl, thiazolyl, furanyl, benzimidazolyl, quinolinyl, isoquinolinyl, quinoxalinyl, pyrimidinyl, pyrazinyl, tetrazolyl and pyrazolyl.

[0024] **Aralkyl-** refers to a residue in which an aryl moiety is attached to the parent structure via an alkyl residue. Examples include benzyl, phenethyl, phenylvinyl, phenylallyl and the like. **Heteroaralkyl-** refers to a residue in which a heteroaryl moiety is attached to the parent structure via an alkyl residue. Examples include furanylmethyl, pyridinylmethyl, pyrimidinylethyl and the like.

[0025] **Aralkoxy-** refers to the group -O-aralkyl. Similarly, **heteroaralkoxy-** refers to the group -O-heteroaralkyl; **aryloxy-** refers to the group -O-aryl; **acyloxy-** refers to the group -O-acyl; **heteroaryloxy-** refers to the group -O-heteroaryl; and **heterocyclyloxy-** refers to the group -O-heterocyclyl (i.e., aralkyl, heteroaralkyl, aryl, acyl, heterocyclyl, or heteroaryl is attached to the parent structure through an oxygen).

[0026] **Carboxyalkyl-** refers to the group -alkyl-COOH.

[0027] **Carboxamido** refers to the group -CONR^bR^c, where

R^b is H or optionally substituted C₁-C₆ alkyl, aryl, heteroaryl, aryl-C₁-C₄ alkyl-, or heteroaryl-C₁-C₄ alkyl- group; and

R^c is hydrogen or C₁-C₄ alkyl; and

where each optionally substituted R^b group is independently unsubstituted or substituted with one or more substituents independently selected from C₁-C₄ alkyl, aryl, heteroaryl, aryl-C₁-C₄ alkyl-, heteroaryl-C₁-C₄ alkyl-, C₁-C₄ haloalkyl, -OC₁-C₄ alkyl, -OC₁-C₄ alkylphenyl, -C₁-C₄ alkyl-OH, -OC₁-C₄ haloalkyl, halogen, -OH, -NH₂, -C₁-C₄ alkyl-NH₂, -N(C₁-C₄ alkyl)(C₁-C₄ alkyl), -NH(C₁-C₄ alkyl), -N(C₁-C₄ alkyl)(C₁-C₄ alkylphenyl), -NH(C₁-C₄ alkylphenyl), cyano, nitro, oxo (as a substituent for heteroaryl), -CO₂H, -C(O)OC₁-C₄ alkyl, -CON(C₁-C₄ alkyl)(C₁-C₄ alkyl), -CONH(C₁-C₄ alkyl), -CONH₂, -NHC(O)(C₁-C₄ alkyl), -NHC(O)(phenyl), -N(C₁-C₄ alkyl)C(O)(C₁-C₄ alkyl), -N(C₁-C₄ alkyl)C(O)(phenyl), -C(O)C₁-C₄ alkyl, -C(O)C₁-C₄ phenyl, -C(O)C₁-C₄ haloalkyl, -OC(O)C₁-C₄ alkyl, -SO₂(C₁-C₄ alkyl), -SO₂(phenyl), -SO₂(C₁-C₄ haloalkyl), -SO₂NH₂, -SO₂NH(C₁-C₄ alkyl), -SO₂NH(phenyl), -NHSO₂(C₁-C₄ alkyl), -NHSO₂(phenyl), and -NHSO₂(C₁-C₄ haloalkyl).

[0028] **Halogen** or **halo** refers to fluorine, chlorine, bromine or iodine. Fluorine,

chlorine and bromine are preferred. Dihaloaryl, dihaloalkyl, trihaloaryl etc. refer to aryl and alkyl substituted with the designated plurality of halogens (here, 2, 2 and 3, respectively), but not necessarily a plurality of the same halogen; thus 4-chloro-3-fluorophenyl is within the scope of dihaloaryl.

[0029] **Heterocyclyl** means a cycloalkyl or aryl residue in which one to four of the carbons is replaced by a heteroatom such as oxygen, nitrogen or sulfur. Examples of heterocycles that fall within the scope of the invention include azetidiny, imidazolinyl, pyrrolidinyl, pyrazolyl, pyrrolyl, indolyl, quinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, benzofuranyl, benzodioxanyl, benzodioxyl (commonly referred to as methylenedioxyphenyl, when occurring as a substituent), tetrazolyl, morpholinyl, thiazolyl, pyridinyl, pyridazinyl, piperidinyl, pyrimidinyl, thienyl, furanyl, oxazolyl, oxazolinyl, isoxazolyl, dioxanyl, tetrahydrofuranyl and the like. "N-heterocyclyl" refers to a nitrogen-containing heterocycle. The term heterocyclyl encompasses heteroaryl, which is a subset of heterocyclyl. Examples of N-heterocyclyl residues include azetidiny, 4-morpholinyl, 4-thiomorpholinyl, 1-piperidinyl, 1-pyrrolidinyl, 3-thiazolidinyl, piperazinyl and 4-(3,4-dihydrobenzoxazinyl). Examples of substituted heterocyclyl include 4-methyl-1-piperazinyl and 4-benzyl-1-piperidinyl.

[0030] A **leaving group or atom** is any group or atom that will, under the reaction conditions, cleave from the starting material, thus promoting reaction at a specified site. Suitable examples of such groups unless otherwise specified are halogen atoms, mesyloxy, p-nitrobenzensulphonyloxy and tosyloxy groups.

[0031] **Optional or optionally** means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstances occurs and instances in which it does not. For example, "optionally substituted alkyl" includes "alkyl" and "substituted alkyl" as defined herein. It will be understood by those skilled in the art with respect to any group containing one or more substituents that such groups are not intended to introduce any substitution or substitution patterns that are sterically impractical and/or synthetically non-feasible and/or inherently unstable.

[0032] **Substituted alkoxy** refers to alkoxy wherein the alkyl constituent is substituted (i.e., -O-(substituted alkyl)). One preferred substituted alkoxy group is "polyalkoxy" or -O-(optionally substituted alkylene)-(optionally substituted alkoxy), and includes groups such as -OCH₂CH₂OCH₃, and residues of glycol ethers such as

polyethyleneglycol, and $-O(CH_2CH_2O)_xCH_3$, where x is an integer of about 2-20, preferably about 2-10, and more preferably about 2-5. Another preferred substituted alkoxy group is hydroxyalkoxy or $-OCH_2(CH_2)_yOH$, where y is an integer of about 1-10, preferably about 1-4.

[0033] Substituted- alkyl, aryl, and heteroaryl, which includes the substituted alkyl, aryl and heteroaryl moieties of any group containing an optionally substituted alkyl, aryl and heteroaryl moiety (e.g., alkoxy, aralkyl and heteroaralkyl), refer respectively to alkyl, aryl, and heteroaryl wherein one or more (up to about 5, preferably up to about 3) hydrogen atoms are replaced by a substituent independently selected from the group:

$-R^a$, $-OR^b$, $-O(C_1-C_2 \text{ alkyl})O-$ (as an aryl substituent), $-SR^b$, $-NR^bR^c$, halogen, cyano, nitro, $-COR^b$, $-CO_2R^b$, $-CONR^bR^c$, $-OCOR^b$, $-OCO_2R^b$, $-OCONR^bR^c$, $-NR^cCOR^b$, $-NR^cCO_2R^b$, $-NR^cCONR^bR^c$, $-CO_2R^b$, $-CONR^bR^c$, $-NR^cCOR^b$, $-SOR^a$, $-SO_2R^a$, $-SO_2NR^bR^c$, and $-NR^cSO_2R^a$,

where R^a is an optionally substituted C_1-C_6 alkyl, aryl, heteroaryl, aryl- C_1-C_4 alkyl-, or heteroaryl- C_1-C_4 alkyl- group,

R^b is H or optionally substituted C_1-C_6 alkyl, aryl, heteroaryl, aryl- C_1-C_4 alkyl-, or heteroaryl- C_1-C_4 alkyl- group;

R^c is hydrogen or C_1-C_4 alkyl;

where each optionally substituted R^a group and R^b group is independently unsubstituted or substituted with one or more substituents independently selected from C_1-C_4 alkyl, aryl, heteroaryl, aryl- C_1-C_4 alkyl-, heteroaryl- C_1-C_4 alkyl-, C_1-C_4 haloalkyl, $-OC_1-C_4$ alkyl, $-OC_1-C_4$ alkylphenyl, $-C_1-C_4$ alkyl-OH, $-OC_1-C_4$ haloalkyl, halogen, -OH, $-NH_2$, $-C_1-C_4$ alkyl- NH_2 , $-N(C_1-C_4 \text{ alkyl})(C_1-C_4 \text{ alkyl})$, $-NH(C_1-C_4 \text{ alkyl})$, $-N(C_1-C_4 \text{ alkyl})(C_1-C_4 \text{ alkylphenyl})$, $-NH(C_1-C_4 \text{ alkylphenyl})$, cyano, nitro, oxo (as a substituent for heteroaryl), $-CO_2H$, $-C(O)OC_1-C_4 \text{ alkyl}$, $-CON(C_1-C_4 \text{ alkyl})(C_1-C_4 \text{ alkyl})$, $-CONH(C_1-C_4 \text{ alkyl})$, $-CONH_2$, $-NHC(O)(C_1-C_4 \text{ alkyl})$, $-NHC(O)(\text{phenyl})$, $-N(C_1-C_4 \text{ alkyl})C(O)(C_1-C_4 \text{ alkyl})$, $-N(C_1-C_4 \text{ alkyl})C(O)(\text{phenyl})$, $-C(O)C_1-C_4 \text{ alkyl}$, $-C(O)C_1-C_4 \text{ phenyl}$, $-C(O)C_1-C_4 \text{ haloalkyl}$, $-OC(O)C_1-C_4 \text{ alkyl}$, $-SO_2(C_1-C_4 \text{ alkyl})$, $-SO_2(\text{phenyl})$, $-SO_2(C_1-C_4 \text{ haloalkyl})$, $-SO_2NH_2$, $-SO_2NH(C_1-C_4 \text{ alkyl})$, $-SO_2NH(\text{phenyl})$, $-NHSO_2(C_1-C_4 \text{ alkyl})$, $-NHSO_2(\text{phenyl})$, and $-NHSO_2(C_1-C_4 \text{ haloalkyl})$.

[0034] Sulfanyl refers to the groups: $-S$ -(optionally substituted alkyl), $-S$ -(optionally substituted aryl), $-S$ -(optionally substituted heteroaryl), and $-S$ -(optionally substituted heterocyclyl).

[0035] Sulfinyl refers to the groups: $-S(O)-H$, $-S(O)$ -(optionally substituted alkyl),

-S(O)-(optionally substituted aryl), -S(O)-(optionally substituted heteroaryl), -S(O)-(optionally substituted heterocyclyl); and -S(O)-(optionally substituted amino).

[0036] Sulfonyl refers to the groups: -S(O₂)-H, -S(O₂)-(optionally substituted alkyl), -S(O₂)-(optionally substituted aryl), -S(O₂)-(optionally substituted heteroaryl), -S(O₂)-(optionally substituted heterocyclyl), -S(O₂)-(optionally substituted alkoxy), -S(O₂)-(optionally substituted aryloxy), -S(O₂)-(optionally substituted heteroaryloxy), -S(O₂)-(optionally substituted heterocyclyloxy); and -S(O₂)-(optionally substituted amino).

[0037] Pharmaceutically acceptable salts refers to those salts that retain the biological effectiveness of the free compound and that are not biologically or otherwise undesirable, formed with a suitable acid or base, and includes pharmaceutically acceptable acid addition salts and base addition salts. **Pharmaceutically acceptable acid addition salts** include those derived from inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and those derived from organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

[0038] Pharmaceutically acceptable base addition salts include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium, and magnesium salts. Base addition salts also include those derived from pharmaceutically acceptable organic non-toxic bases, including salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine.

[0039] Protecting group has the meaning conventionally associated with it in organic synthesis, i.e. a group that selectively blocks one or more reactive sites in a multifunctional compound such that a chemical reaction can be carried out selectively on another unprotected reactive site and such that the group can readily be removed after the selective reaction is complete. A variety of protecting groups are disclosed, for example, in T.H. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, Third Edition, John Wiley & Sons, New York (1999), which is incorporated herein by reference in its entirety. For example, a

hydroxy protected form is where at least one of the hydroxyl groups present in a compound is protected with a hydroxy protecting group. Likewise, amines and other reactive groups may similarly be protected.

[0040] **Solvate** refers to the compound formed by the interaction of a solvent and a compound of Formula I or salt thereof. Suitable solvates of the compounds of the Formula I are pharmaceutically acceptable solvates, such as hydrates, including monohydrates and hemi-hydrates.

[0041] Many of the compounds described herein contain one or more asymmetric centers (e.g. the carbon to which R_2 and R_2' are attached where R_2 differs from R_2') and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)-. The present invention is meant to include all such possible isomers, including racemic mixtures, optically pure forms and intermediate mixtures. Optically active (R)- and (S)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers. Likewise, all tautomeric forms and rotational isomers are also intended to be included.

[0042] When desired, the R- and S-isomers may be resolved by methods known to those skilled in the art, for example by formation of diastereoisomeric salts or complexes which may be separated, for example, by crystallization; via formation of diastereoisomeric derivatives which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic oxidation or reduction, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, such as silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be required to liberate the desired enantiomeric form. Alternatively, specific enantiomer may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer to the other by asymmetric transformation.

Compounds of the Present Invention


[0043] The present invention is directed to a class of novel compounds, which can be described as benzopyran-4-ones or chromen-4-ones, that are inhibitors of one or more mitotic kinesins. By inhibiting mitotic kinesins, but not other kinesins (e.g., transport kinesins), specific inhibition of cellular proliferation is accomplished. While not intending to be bound by any theory, the present invention capitalizes on the finding that perturbation of mitotic kinesin function causes malformation or dysfunction of mitotic spindles, frequently resulting in cell cycle arrest and cell death. According to one embodiment of the invention, the compounds described herein inhibit the mitotic kinesin, KSP. In another embodiment, the compounds inhibit the mitotic kinesin, KSP, as well as modulating one or more of the human mitotic kinesins selected from the group consisting of HSET (see, U.S. Patent No. 6,361,993, which is incorporated herein by reference); MCAK (see, U.S. Patent No. 6,331,424, which is incorporated herein by reference); CENP-E (see, PCT Publication No. WO 99/13061, which is incorporated herein by reference); Kif4 (see, U.S. Patent No. 6,440,684, which is incorporated herein by reference); MKLP1 (see, U.S. Patent No. 6,448,025, which is incorporated herein by reference); Kif15 (see, U.S. Patent No. 6,355,466, which is incorporated herein by reference); Kid (see, U.S. Patent No. 6,387,644, which is incorporated herein by reference); Mpp1, CMKrp, KinI-3 (see, U.S. Patent No. 6,461,855, which is incorporated herein by reference); Kip3a (see, PCT Publication No. WO 01/96593, which is incorporated herein by reference); Kip3d (see, U.S. Patent No. 6,492,151, which is incorporated herein by reference); and RabK6.

[0044] The methods of inhibiting a human KSP kinesin comprise contacting an inhibitor of the invention with a kinesin, particularly a human kinesin, preferably human KSP or fragments and variants thereof. The inhibition can be of the ATP hydrolysis activity of the KSP kinesin and/or the mitotic spindle formation activity, such that the mitotic spindles are disrupted. Meiotic spindles may also be disrupted.

[0045] An object of the present invention is to develop inhibitors of mitotic kinesins, in particular KSP and especially human KSP, for the treatment of disorders associated with cell proliferation. Traditionally, dramatic improvements in the treatment of cancer, one type of cellular proliferative disorder, have been associated with identification of therapeutic agents acting through novel mechanisms. Examples of this include not only the taxane class of agents that appear to act on microtubule formation, but also the camptothecin class of topoisomerase I inhibitors. The compounds, compositions and methods described herein can

differ in their selectivity and are preferably used to treat diseases of cellular proliferation, including, but not limited to cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders, fungal disorders and inflammation.

[0046] Accordingly, the present invention relates to methods employing compounds represented by Formula I:



Formula I

wherein:

R_1 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-;

R_2 and $R_{2'}$ are independently chosen from hydrogen, optionally substituted alkyl-, optionally substituted alkoxy, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-; or R_2 and $R_{2'}$ taken together form an optionally substituted 3- to 7-membered ring;

R_{12} is selected from the group consisting of optionally substituted imidazolyl, optionally substituted imidazolyl-, $-NHR_4$; $-N(R_4)(COR_3)$; $-N(R_4)(SO_2R_{3a})$; and $-N(R_4)(CH_2R_{3b})$;

R_3 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, optionally substituted heteroaralkyl-, $R_{15}O$ - and $R_{17}NH$ -;

R_{3a} is chosen from optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, optionally substituted heteroaralkyl-, and $R_{17}NH$ -;

R_{3b} is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heterocycl-, and optionally substituted heteroaralkyl-;

R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, acyl, optionally substituted alkyl-, optionally substituted alkoxy, halogen, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl-, alkylsulfonamido-, alkylthio-, carboxyalkyl-, carboxamido-, aminocarbonyl-, optionally substituted aryl and optionally substituted heteroaryl-;

R_{15} is chosen from optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-, and


R_{17} is hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, or optionally substituted heteroaralkyl-, including single stereoisomers, mixtures of stereoisomers;

a pharmaceutically acceptable salt of a compound of Formula I;

a pharmaceutically acceptable solvate of a pharmaceutically acceptable solvate of a compound of Formula I;

or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I.

[0047] When R_{12} is an imidazole, R_{12} has the formula:




wherein

R_9 is chosen from hydrogen, optionally substituted C_1 - C_8 alkyl, optionally substituted aryl, optionally substituted aryl- C_1 - C_4 -alkyl -, optionally substituted heteroaryl- C_1 - C_4 -alkyl -,

optionally substituted aryl-C₁-C₄-alkoxy -, optionally substituted heteroaryl-C₁-C₄-alkoxy -, optionally substituted heteroaryl-; and R₁₃ and R_{13'} are independently hydrogen, optionally substituted C₁-C₈ alkyl, optionally substituted aryl, or optionally substituted aryl-C₁-C₄-alkyl-

[0048] When R₁₂ is an imidazoline, R₁₂ has the formula



wherein

R₉ is chosen from hydrogen, optionally substituted C₁-C₈ alkyl, optionally substituted aryl, optionally substituted aryl-C₁-C₄-alkyl -, and optionally substituted heteroaryl-; and R₁₀, R_{10'}, R₁₄, and R_{14'} are independently chosen from hydrogen, optionally substituted C₁-C₈ alkyl, optionally substituted aryl, and optionally substituted aryl-C₁-C₄-alkyl -.

[0049] In one embodiment, R₁ is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl-;

R₂ and R_{2'} are independently chosen from hydrogen, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl-; or R₂ and R_{2'} taken together form an optionally substituted 3- to 7-membered ring, provided that if either R₂ or R_{2'} is hydrogen, then the other is not hydrogen;

R₁₂ is selected from the group consisting of optionally substituted imidazolyl, optionally substituted imidazoliny, -NHR₄; -N(R₄)(COR₃); -N(R₄)(SO₂R_{3a}); and -N(R₄)(CH₂R_{3b});

R₃ is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, R₁₅O- and R₁₇-NH-;

R_{3a} is chosen from optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, and R_{17} -NH-;

R_{3b} is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl-;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heterocycl-, and optionally substituted heteroaralkyl-;

R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, acyl, optionally substituted alkyl, optionally substituted alkoxy, halogen, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, optionally substituted aryl and optionally substituted heteroaryl-;

R_{15} is chosen from optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl-; and

R_{17} is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl, including single stereoisomers, mixtures of stereoisomers;

a pharmaceutically acceptable salt of a compound of Formula I;

a pharmaceutically acceptable solvate of a pharmaceutically acceptable solvate of a compound of Formula I; or

a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I.

[0050] In another embodiment, R_2 and R_2' are hydrogen; and

R_1 is chosen from optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl, provided however, that R_1 is not substituted phenyl;

R_{12} is selected from the group consisting of optionally substituted imidazolyl, optionally substituted imidazolyl-, $-NHR_4$; $-N(R_4)(COR_3)$; $-N(R_4)(SO_2R_{3a})$; and $-N(R_4)(CH_2R_{3b})$;

R_3 is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, optionally substituted

heteroaralkyl, $R_{15}O-$ and $R_{17}NH-$;

R_{3a} is chosen from optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, and $R_{17}NH-$;

R_{3b} is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl-;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heterocycl-, and optionally substituted heteroaralkyl-;

R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, acyl, optionally substituted alkyl, optionally substituted alkoxy, halogen, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, optionally substituted aryl and optionally substituted heteroaryl-;

R_{15} is chosen from optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl-; and

R_{17} is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl, including single stereoisomers, mixtures of stereoisomers;

a pharmaceutically acceptable salt of a compound of Formula I;


a pharmaceutically acceptable solvate of a pharmaceutically acceptable solvate of a compound of Formula I; or

a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I.

In a particularly preferred embodiment, where R_2 differs from R_2' , the stereogenic center to which R_2 and R_2' are attached is of the R configuration.


Nomenclature

[0051] The compounds of Formula I can be named and numbered in the manner (e.g., using AutoNom version 2.1 or ISIS-DRAW, each of which utilizes the IUPAC system of nomenclature) described below. For example, the compound:



i.e., the compound according to Formula I where R_1 is benzyl, R_2 is propyl (particularly i - propyl), $R_{2'}$ is hydrogen; R_{12} is $-N(R_4)(COR_3)$; R_3 is 3,4-dimethylphenyl-; R_4 is 3-aminopropyl-; R_5 , R_6 , and R_8 are hydrogen; and R_7 is cyano is named N-(3-amino-propyl)-N-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-3,4-dimethyl-benzamide.

[0052] Likewise, the compound:



i.e., the compound according to Formula I where R_1 is 3-methoxy-benzyl, R_2 is propyl (particularly i - propyl), $R_{2'}$ is hydrogen; R_{12} is substituted imidazoline; R_5 , R_6 , and R_8 are hydrogen; R_7 is chloro; R_9 is methylenedioxyphenyl-; R_{10} , $R_{10'}$, R_{14} , and $R_{14'}$ are hydrogen can be named 2-[1-(2-benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-7-chloro-3-(3-methoxy-benzyl)-chromen-4-one.

Synthesis of the Compounds of Formula I

[0053] The compounds of Formula I can be prepared by following the procedures described with reference to the Reaction Schemes below or utilizing techniques well known in the art. See, for example, Hirao et al. (1984) Synthesis 1076-1078 and Coppola et al. (1981) Synthesis 523-526, which are incorporated herein by reference.

[0054] Unless specified otherwise, the terms "solvent", "inert organic solvent" or "inert solvent" mean a solvent inert under the conditions of the reaction being described in

conjunction therewith [including, for example, benzene, toluene, acetonitrile, tetrahydrofuran ("THF"), dimethylformamide ("DMF"), chloroform, methylene chloride (or dichloromethane), diethyl ether, methanol, pyridine and the like]. Unless specified to the contrary, the solvents used in the reactions of the present invention are inert organic solvents.

[0055] The term "q.s." means adding a quantity sufficient to achieve a stated function, e.g., to bring a solution to the desired volume (i.e., 100%).

[0056] In general, esters of carboxylic acids may be prepared by conventional esterification procedures, for example alkyl esters may be prepared by treating the required carboxylic acid with the appropriate alkanol, generally under acidic conditions. Likewise, amides may be prepared using conventional amidation procedures, for example amides may be prepared by treating the relevant carboxylic acid with the appropriate amine. Alternatively, a lower-alkyl ester such as a methyl ester of the acid may be treated with an amine to provide the required amide, optionally in presence of trimethylaluminum following the procedure described in Tetrahedron Lett. 48, 4171-4173, (1977). Carboxyl groups may be protected as alkyl esters, for example methyl esters, which esters may be prepared and removed using conventional procedures, one convenient method for converting carbomethoxy to carboxyl is to use aqueous lithium hydroxide.

[0057] The salts and solvates of the compounds mentioned herein may as required be produced by methods conventional in the art. For example, if an inventive compound is an acid, a desired base addition salt can be prepared by treatment of the free acid with an inorganic or organic base, such as an amine (primary, secondary, or tertiary); an alkali metal or alkaline earth metal hydroxide; or the like. Illustrative examples of suitable salts include organic salts derived from amino acids such as glycine and arginine; ammonia; primary, secondary, and tertiary amines; such as ethylenediamine, and cyclic amines, such as cyclohexylamine, piperidine, morpholine, and piperazine; as well as inorganic salts derived from sodium, calcium, potassium, magnesium, manganese, iron, copper, zinc, aluminum, and lithium.

[0058] If a compound is a base, a desired acid addition salt may be prepared by any suitable method known in the art, including treatment of the free base with an inorganic acid, such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like, or with an organic acid, such as acetic acid, maleic acid, succinic acid, mandelic acid, fumaric acid, malonic acid, pyruvic acid, oxalic acid, glycolic acid, salicylic acid, pyranosidyl acid, such as glucuronic acid or galacturonic acid, alpha-hydroxy acid, such as citric acid or

tartaric acid, amino acid, such as aspartic acid or glutamic acid, aromatic acid, such as benzoic acid or cinnamic acid, sulfonic acid, such as p-toluenesulfonic acid, methanesulfonic acid, ethanesulfonic acid, or the like.

[0059] Isolation and purification of the compounds and intermediates described herein can be effected, if desired, by any suitable separation or purification procedure such as, for example, filtration, extraction, crystallization, column chromatography, thin-layer chromatography or thick-layer chromatography, or a combination of these procedures. Specific illustrations of suitable separation and isolation procedures can be had by reference to the examples hereinbelow. However, other equivalent separation or isolation procedures can, of course, also be used.

Brief Description Of Reaction Schemes

[0060] Reaction Scheme 1 illustrates a synthesis of compounds of formula 109, an intermediate in the synthesis of compounds of Formula I.

[0061] Reaction Scheme 2 illustrates a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(COR_3)$.

[0062] Reaction Scheme 3 illustrates a synthesis of compounds of Formula I wherein R_7 is $-OH$.

[0063] Reaction Scheme 4 illustrates a synthesis of compounds of Formula I wherein R_7 is $-OCH_3$.

[0064] Reaction Scheme 5 illustrates another synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(COR_3)$.

[0065] Reaction Scheme 6 illustrates a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(SO_2R_{3a})$.

[0066] Reaction Scheme 7 illustrates a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(CH_2R_{3b})$.

[0067] Reaction Scheme 8 illustrates a synthesis of compounds of Formula I wherein R_{12} is optionally substituted imidazolyl.

[0068] Reaction Scheme 9 illustrates another synthesis of compounds of Formula I wherein R_{12} is optionally substituted imidazolyl.

[0069] Reaction Scheme 10 illustrates a synthesis of compounds of Formula I wherein R_{12} is optionally substituted imidazoliny.

[0070] Reaction Scheme 11 illustrates a second synthesis of compounds of Formula I

wherein R_{12} is optionally substituted imidazoliny.

[0071] Reaction Scheme 12 illustrates a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(COR_3)$ wherein R_3 is $-OR_{15}$.

[0072] Reaction Scheme 13 illustrates a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(COR_3)$ wherein R_3 is $-NHR_{17}$.


[0073] Reaction Scheme 14 illustrates a synthesis of compounds of Formula 1407 which can be used as an intermediate in the synthesis of compounds of Formula I.

[0074] Reaction Scheme 15 illustrates a synthesis of compounds of Formula 1505 which can be used as an intermediate in the synthesis of compounds of Formula I.

Starting Materials

[0075] The optionally substituted compounds of Formula 101 are commercially available, e.g., from Aldrich Chemical Company, Milwaukee, WI. Other reactants are likewise commercially available or may be readily prepared by those skilled in the art using commonly employed synthetic methodology.

Reaction Scheme 1



Preparation of Compounds of Formula 103

[0076] Referring to Reaction Scheme 1, Step 1, about an equivalent of ethyl chloroformate is added over about one minute to a 0-5°C solution of a compound of Formula 101 (preferably wherein the amino protecting group, PG, is a Boc group) and a base such as triethylamine in a nonpolar, aprotic solvent such as THF. After about 15 minutes, a mixture of an excess of dimethylhydroxylamine hydrochloride (preferably about 1.2 equivalents) and a base such as triethylamine in a nonpolar, aprotic solvent such as THF is added over about 5 minutes. The product, a compound of Formula 103, is isolated and used without further purification.

Preparation of Compounds of Formula 105

[0077] Referring to Reaction Scheme 1, Step 2, a Grignard reagent is prepared by

mixing a compound of formula R_1CH_2Br (generally about 3 equivalents) and magnesium turnings in a nonpolar, aprotic solvent such as diethyl ether. After about 1.5 hours, the Grignard reaction is generally complete. A solution of a compound of Formula 103 in a nonpolar, aprotic solvent such as ether, is added to the Grignard reagent. The temperature should be monitored and not allowed to exceed $\sim 30^\circ C$. The product, a compound of Formula 105, is isolated and purified.

Preparation of Compounds of Formula 107

[0078] Referring to Reaction Scheme 1, Step 3, lithium bis(trimethylsilyl)amide (about 3.3 equivalents) is added slowly over ~ 3 minutes to a $-78^\circ C$ solution of a compound of Formula 105 in a nonpolar, aprotic solvent such as THF. The reaction solution temperature should be monitored and the addition of base conducted at a rate sufficient to prevent the temperature from exceeding about $-54^\circ C$. After the addition is complete, the resulting solution is maintained at $-78^\circ C$ for about 30 minutes. An acid chloride of Formula 106 (preferably, neat) is then added. The reaction solution is maintained at $-78^\circ C$ for about 30 minutes. The product is isolated and used without further purification.

[0079] A mixture of the above crude product, a base such as potassium carbonate, and a polar, aprotic solvent such as DMF is maintained at about room temperature for about 30 minutes. The product, a compound of Formula 107 is isolated and purified.

Preparation of Compounds of Formula 109

[0080] Referring to Reaction Scheme 1, Step 4, optionally, the protecting group, PG, may be removed from the amine. One of skill in the art will appreciate that the conditions for removal of the protecting group will vary with different protecting groups. Such conditions are well known in the art and can be found, e.g., in Greene et al. *supra*. When PG is Boc, it may be removed by treatment of a compound of Formula 107 with a mixture of aqueous TFA (preferably TFA:H₂O, 97.5:2.5) at room temperature. The product, a compound of Formula 109, is isolated and purified.


Preparation of Optically Active Compounds


[0081] In compounds of the invention where R_2 differs from R_2' , a particular stereo configuration (such as the (R) isomer) may be preferred at the stereogenic center to which R_2

and R_2 are attached. The optically active compound can be prepared by methods known in the art. For example, an amine of Formula 109 is dissolved in an inert organic solvent (such as IPA) and warmed to 60°C. In a separate vessel, a resolving agent (such as dibenzoyl-D-tartaric acid) is dissolved, preferably in the same warm solvent, and then quickly added (with agitation) to the warm amine solution. The reaction mixture is left to crystallize by cooling to room temperature over 16 hours under continuing agitation. The desired isomer, e.g., the (R) isomer of a compound of Formula 109, is isolated and purified.

[0082] For the sake of brevity in the remaining description of the synthesis of compounds of Formula I, it should be understood that either single isomer or a mixture of isomers may be employed to give the corresponding product.

Reaction Scheme 2





Preparation of Formula 203

[0083] Referring to Reaction Scheme 2, Step 1, to a solution of a compound of Formula 109 is added successively a slight excess (preferably about 1.2 equivalents) of an aldehyde comprising R_4 (i.e., a compound having the formula $R_4\text{CHO}$ where $R_4\text{CH}_2$ is equivalent to R_4 and R_4 is as described above or is a protected precursor to such a substituent, e.g., (3-oxo-propyl)-carbamic acid *tert*-butyl ester) and a reducing agent such as sodium triacetoxyborohydride. The resulting mixture is stirred for several hours. The product, a compound of Formula 203 is isolated and purified.


Preparation of Formula 205

[0084] Referring to Reaction Scheme 2, Step 2, to a solution of a compound of Formula 203 and an amine base such as diisopropylethylamine in a nonpolar, aprotic solvent such as dichloromethane is added an R_3 acyl chloride (such as $\text{Cl-C(O)-}R_3$ where R_3 is as described above). The resulting solution is stirred under nitrogen at room temperature for several hours. The product, a compound of Formula 205 is isolated and purified.

Preparation of Formula 207

[0085] Optionally, any protecting groups on compounds of Formula 205 are then removed. For example, if R_4 comprises a protected amine wherein the protecting group is a Boc group, the Boc group can be removed by treatment of the compound of Formula 205 with an acid such as trifluoroacetic acid in a nonpolar, aprotic solvent such as dichloromethane, while maintaining the reaction at about room temperature. The reaction is monitored e.g., by TLC. Upon completion, the product, a compound of Formula 207 is isolated and purified.

W0882


Reaction Scheme 3**Preparation of Compounds of Formula 303**

[0086] Referring to Reaction Scheme 3, Step 1, to a solution of a compound of Formula 301 in a nonpolar, aprotic solvent such as DMF is added sodium hydride. The resulting solution was stirred at about 45°C for about 5 minutes, then allyl alcohol (about 1.4 equivalents) is added via pipette. The resulting solution is stirred at about 45°C for about 12 hours and then cooled to room temperature. The product, a compound of Formula 303, is isolated and used without further purification.

Preparation of Compounds of Formula 305


[0087] Referring to Reaction Scheme 3, Step 2, to a room temperature solution of a compound of Formula 303, in an aprotic solvent such as acetonitrile, is added morpholine followed by $Pd(PPh_3)_4$. The resulting solution is stirred for about 5 minutes. The product, a compound of Formula 305, is isolated and purified.


Reaction Scheme 4



[0088] Referring to Reaction Scheme 4, a compound of Formula 401 is dissolved in 0.5 M sodium methoxide in methanol and heated to about 70°C. The temperature is maintained at about 70°C for about 12 hours and then cooled to room temperature. The product, a compound of Formula 403, is isolated and purified.

Reaction Scheme 5





Preparation of Compounds of Formula 503

[0089] Referring to Reaction Scheme 5, Step 1, to a solution of a compound of Formula 501 and a base such as triethylamine in a nonpolar, aprotic solvent such as CH_2Cl_2 at room temperature is added about an equivalent of an acid chloride of formula $\text{Cl}-(\text{CO})-\text{CH}_2\text{R}_1$ over about 5 minutes. After about 30 minutes, the product, a compound of Formula 503, is isolated and used without any further purification.

Preparation of Compounds of Formula 505

[0090] Referring to Reaction Scheme 5, Step 2, AlCl_3 (about 1.3 equivalents) is added slowly over about 15 minutes to a compound of Formula 503 at about 140°C . After gas evolution has ceased, the reaction mixture is cooled to room temperature. The product, a compound of Formula 505, is isolated and purified.

Preparation of Compounds of Formula 507

[0091] Referring to Reaction Scheme 5, Step 3, a solution of a compound of Formula

505 and an amino acid of Formula 506 (preferably about 1.1 equivalents) in which the amine group has been suitably protected with a protecting group PG (preferably a Boc group), a coupling reagent such as HBTU (preferably about 1.2 equivalents), a base such as TEA, and a nonpolar, aprotic solvent such as CH_2Cl_2 is maintained at room temperature for about 5 hours. The product, a compound of Formula 507, is isolated and purified.

Preparation of Compounds of Formula 509

[0092] Referring to Reaction Scheme 5, Step 4, a mixture of ester of Formula 507 and a base such as potassium carbonate in a polar, aprotic solvent such as DMF is heated at about 140°C . After about 30 mins, the product, a compound of Formula 509, is isolated and purified.

Preparation of Compounds of Formula 511

[0093] Referring to Reaction Scheme 5, Step 5, the amine protecting group PG is then removed. When PG is Boc, this may be accomplished by treatment of a compound of formula 509 with aqueous acid (preferably, 97.5:2.5 TFA: H_2O) at room temperature for about one hour. The free amine is isolated and used without further purification.


[0094] The resulting product; an aldehyde comprising R_4 (i.e., a compound having the formula $\text{R}_4\text{-CHO}$ where $\text{R}_4\text{-CH}_2\text{-}$ is equivalent to R_4 and R_4 is as described above or is a protected precursor to such a substituent, e.g., (3-oxo-propyl)-carbamic acid *tert*-butyl ester; preferably, about 1.45 equivalents); a reducing reagent, such as $\text{Na(OAc)}_3\text{BH}$; and a nonpolar, aprotic solvent such as CH_2Cl_2 is maintained at room temperature for about 3 hours. The product, a compound of Formula 511, is isolated and purified.

[0095]

Preparation of Compounds of Formula 513


[0096] Referring to Reaction Scheme 5, Step 6, to a solution of a compound of Formula 511, a base such as diisopropylethylamine, and a nonpolar, aprotic solvent such as CH_2Cl_2 at room temperature is added an acid chloride of the formula R_3COCl (preferably about 2 equivalents). After about 2.5 h, the product, a compound of Formula 513, is isolated and purified.

Reaction Scheme 6



[0097] Referring to Reaction Scheme 6, to a solution of a compound of Formula 203 and an amine base such as diisopropylethylamine in a nonpolar, aprotic solvent such as dichloromethane is added a compound having the formula $\text{Cl-S(O)}_2\text{-R}_{3a}$ or $\text{O-(S(O)}_2\text{-R}_{3a})_2$ where R_{3a} is as described above. The resulting solution is stirred under nitrogen at room temperature for several hours. The product, a compound of Formula 603 is isolated and purified.


Reaction Scheme 7



[0098] Referring to Reaction Scheme 7, to a solution of a compound of Formula 203 and an amine base such as diisopropylethylamine in a nonpolar, aprotic solvent such as dichloromethane is added a compound having the formula $\text{X-CH}_2\text{-R}_{3b}$ where R_{3b} is as described above and X is Br, Cl, mesylate, or tosylate. The resulting solution is stirred under nitrogen at room temperature or with heat for several hours. The product, a compound of

Formula 703 is isolated and purified.

Reaction Scheme 8



Preparation of Formula 803

[0099] Referring to Reaction Scheme 8, Step 1, to an optionally substituted compound of Formula 109 dissolved in a polar, aprotic solvent (such as DMF) in the presence of a base (such as potassium carbonate) is added one equivalent of an optionally substituted suitably protected aldehyde wherein such aldehyde further comprises a leaving group,

preferably, a halide (such as bromoacetaldehyde dimethylacetal). The solution is heated at reflux, monitoring completion of the reaction (e.g., by TLC). The reaction mixture is cooled and the corresponding, optionally substituted compound of Formula 803 is isolated and purified.


Preparation of Formula 805

[00100] Referring to Reaction Scheme 8, Step 2, to an optionally substituted compound of Formula 803 in an inert solvent (such as dichloromethane) in the presence of about 1.5 molar equivalents of an amine base (such as triethylamine) is added about 1.5 molar equivalents of an R_9 acid chloride, such as, $Cl-C(O)-R_9$, where R_9 is as described herein. The reaction takes place, with stirring, at room temperature over a period of 4 to 24 hours. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 805 is isolated and purified.

Preparation of Formula 807


[00101] Referring to Reaction Scheme 8, Step 3, a solution of a compound of Formula 805 and an excess of ammonium acetate in acetic acid is heated at reflux for 1-4 hours. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 807 is isolated and purified.

Reaction Scheme 9




903

Step 2




905

Step 3



907

Step 4



Preparation of Formula 903

[00102] Referring to Reaction Scheme 9, Step 1, a suspension of a compound of Formula 109, an alpha-haloketone reagent of the Formula $R_{13}(\text{CO})\text{CH}_2\text{X}$ wherein X is a halide and R_{13} is as described herein, and about an equivalent of a base, such as potassium carbonate in a polar, aprotic solvent such as DMF is stirred at room temperature. The reaction is diluted with water and the resulting compound, a compound of Formula 903, typically a solid, is used in the subsequent step without purification. Where the resulting compound is not a solid, it is isolated using standard procedures and then used in the subsequent step.

Preparation of Formula 905

[00103] Referring to Reaction Scheme 9, Step 2, a solution of the compound of Formula 903, about an equivalent of an amine base, such as triethylamine and about an equivalent of an acid chloride (such as a compound of Formula $R_9\text{-COCl}$) in an organic solvent such as methylene chloride is stirred at room temperature for several hours. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 905 is isolated and purified.

Preparation of Formula 907


[00104] Referring to Reaction Scheme 9, Step 3, a solution of a compound of Formula 905 and an excess of ammonium acetate in acetic acid is heated at reflux using a Dean-Stark trap and condenser. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 907 is isolated and purified.


Preparation of Formula 909

[00105] Referring to Reaction Scheme 9, Step 4, when R_{13} comprises a protected aminoalkyl group, the amino protected group may be removed. For example, when the amino group is protected as the corresponding phthalimide, the protecting group is removed as follows. A solution of a compound of Formula 907 and an excess of anhydrous hydrazine in a polar, protic solvent such as ethanol is heated at reflux. The reaction is cooled to about 5°C and any precipitate is filtered off. The filtrate is concentrated in vacuo and purified to yield a

compound of Formula 909. One of skill in the art will appreciate that other conditions may be used to remove other protecting groups.

Reaction Scheme 10





Preparation of Formula 1003

[00106] Referring to Reaction Scheme 10, Step 1, reductive amination of amines of Formula 109 with an optionally substituted, aldehyde-containing carbamic acid ester gives urethane intermediates. More specifically, to a solution of a compound of Formula 109 and an equivalent of a suitably protected aldehyde (Seki *et. al. Chem. Pharm. Bull.* 1996, 44, 2061) in dichloromethane is added a slight excess of a reducing agent, such as sodium triacetoxyborohydride. The resultant cloudy mixture is maintained at ambient temperature. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 1003 is isolated and used in the subsequent step without purification.

Preparation of Formula 1005

[00107] Referring to Reaction Scheme 10, Step 2, the amino protecting group PG is then removed. When PG is a Boc protecting group, this may be accomplished by the treatment of a solution of a compound of Formula 1003 in a nonpolar, aprotic solvent such as dichloromethane with a strong acid such as trifluoroacetic acid. The resultant solution is maintained at ambient temperature overnight and concentrated under reduced pressure. The residue is isolated to give a compound of Formula 1005 which was used in the subsequent step without purification.

Preparation of Formula 1007


[00108] Referring to Reaction Scheme 10, Step 3, to a solution of a compound of Formula 1005 in a nonpolar, aprotic solvent such as dichloromethane is added an excess, preferably about two equivalents of an amine base such as triethylamine, followed by about

an equivalent or slight excess of an acid chloride of the formula $R_9\text{-CO-Cl}$. The resultant solution is stirred at ambient temperature for about 3 hours. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 1007 is isolated and purified.

Preparation of Formula 1009

[00109] Referring to Reaction Scheme 10, Step 4, a solution of a compound of Formula 1007 in an excess of phosphorus oxychloride is heated at reflux. After 8 hours, the reaction mixture is allowed to cool to ambient temperature and concentrated under reduced pressure. The corresponding compound of Formula 1009 is isolated and purified.

Reaction Scheme 11




Preparation of Formula 1109

[00110] As an alternative to Steps 3 and 4 of Reaction Scheme 10, acylation of primary amines of Formula 1005, followed by acetic acid mediated cyclization, can proceed without isolation of the intermediate amides to provide the target compound of Formula 1109. This route is shown in Reaction Scheme 11.

[00111] More specifically, to a solution of a compound of Formula 1005 in a nonpolar, aprotic solvent such as dichloromethane is added an excess, preferably about two equivalents of an amine base, such as triethylamine, followed by about an equivalent of an acid chloride of the formula $R_9\text{-CO-Cl}$. The resultant solution is stirred at ambient temperature for 2 hours, then evaporated under reduced pressure. The resultant solid is treated with glacial acetic acid, then the resultant suspension is heated at reflux for about 48 hours. The reaction is cooled to


ambient temperature then evaporated under reduced pressure. The corresponding compound of Formula 1109 is isolated and purified.

Reaction Scheme 12




[00112] Referring to Reaction Scheme 12, a compound of Formula 203 is reacted with a slight excess of a compound of the formula $R_{15}O(CO)Cl$ in the presence of a base such as triethylamine in a nonpolar, aprotic solvent such as dichloromethane. The product, a compound of Formula 1203 is isolated and purified.

Reaction Scheme 13



[00113] Referring to Reaction Scheme 13, a compound of Formula 203 is treated with a slight excess of an isocyanate $R_{17}-N=C=O$ in the presence of a base, such as triethylamine, in a nonpolar, aprotic solvent, such as dichloromethane. The product, a compound of Formula 1303, is isolated and purified.

Reaction Scheme 14



Preparation of Compounds of Formula 1403

[00114] Referring to Reaction Scheme 14, Step 1, a nonpolar, aprotic solvent, such as THF, and an excess of a solution of an optionally substituted vinyl magnesium bromide in a nonpolar, aprotic solvent (and more preferably, about three equivalents of a 1.0 M solution of an optionally substituted vinyl magnesium bromide in THF) is cooled to -78°C while stirring under a nitrogen atmosphere. The mixture is treated dropwise with a solution of a compound of Formula 1401 in a nonpolar, aprotic solvent, such as THF over about 30 min. After the mixture is stirred for 30 min at -78°C , the cooling bath is removed and the reaction mixture is warmed slowly to room temperature overnight (about 15 h). The product, a compound of Formula 1403, is isolated and purified.

Preparation of Compounds of Formula 1405


[00115] Referring to Reaction Scheme 14, Step 2, to a solution of a compound of Formula 1403 in an anhydrous, nonpolar, aprotic solvent, such as acetonitrile under an inert atmosphere, such as argon, is added about an equivalent of a compound of the Formula R₁'-X wherein R₁' is an optionally substituted vinyl, optionally substituted aryl, or optionally substituted heteroaryl and X is I, Br, or -OTf, and a base such as triethylamine followed by

palladium (II) acetate (preferably, about 0.025 equivalents). The resulting solution is heated to about 80 °C. After about 15 h, the reaction mixture is allowed to cool to room temperature. The product, a compound of Formula 1405, is isolated and immediately purified.

Preparation of Compounds of Formula 1407

[00116] To a solution of a compound of Formula 1405 in a nonpolar, aprotic solvent such as ethyl acetate under nitrogen is added 10 wt % palladium on carbon. The nitrogen is replaced with a balloon of hydrogen and the flask is purged. After 3 h, the reaction flask is purged with nitrogen and filtered through a pad of celite (rinsing with a solvent such as ethyl acetate). The product, a compound of Formula 1407 is isolated and purified.

Reaction Scheme 15



Preparation of Compounds of Formula 1503

[00117] Referring to Reaction Scheme 15, Step 1, about one equivalent of carbonyldiimidazole is added slowly to a room temperature solution of a compound of Formula 1501 (preferably wherein the amino protecting group PG is Boc) in a nonpolar, aprotic solvent such as THF. After about one hour, the product, a compound of Formula 1503, is isolated and used without further purification.

Preparation of Compounds of Formula 1505

[00118] Referring to Reaction Scheme 15, Step 2, a Grignard reagent is prepared from a compound of Formula R_1CH_2Br and magnesium turnings in a nonpolar, aprotic solvent such as THF. A solution of a compound of Formula 1503 in a nonpolar, aprotic solvent such as THF is cooled to about 0-5°C. The solution of the Grignard reagent is then added via syringe to the 0-5°C solution of the compound of Formula 1503. The temperature is monitored by internal thermometer and is not allowed to exceed about 15°C. The reaction mixture is maintained at about 0-5°C for about one hour. The product, a compound of Formula 1505, is isolated and purified.

Preferred Processes and Last Steps

[00119] A compound of Formula I is optionally contacted with a pharmaceutically acceptable acid or base to form the corresponding acid or base addition salt.

[00120] A pharmaceutically acceptable acid addition salt of a compound of Formula I is optionally contacted with a base to form the corresponding free base of Formula I. A pharmaceutically acceptable base addition salt of a compound of Formula I is optionally contacted with an acid to form the corresponding free acid of Formula I.

Preferred Compounds

Preferred R_1 when Either R_2 or $R_{2'}$ is Not Hydrogen

[00121] When considering the compounds of Formula I, in a preferred embodiment when either one or both R_2 or $R_{2'}$ is not hydrogen (more preferably, either one of R_2 or $R_{2'}$ is not hydrogen), R_1 is selected from hydrogen, optionally substituted C_1 - C_8 alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted aryl- C_1 - C_4 -alkyl-, and optionally substituted heteroaryl- C_1 - C_4 -alkyl- (more preferably optionally substituted aryl and optionally substituted aryl- C_1 - C_4 -alkyl-). In a more preferred embodiment R_1 is selected from hydrogen, optionally substituted C_1 - C_4 alkyl, optionally substituted phenyl- C_1 - C_4 -alkyl-, optionally substituted naphthalenylmethyl, optionally substituted phenyl, and naphthyl. More preferably, R_1 is optionally substituted phenyl- C_1 - C_4 -alkyl- or optionally substituted heteroaryl- C_1 - C_4 -alkyl-.

[00122] In a most preferred embodiment R_1 is naphthyl, phenyl, bromophenyl, chlorophenyl, methoxyphenyl, ethoxyphenyl, tolyl, dimethylphenyl, chlorofluorophenyl,

methylchlorophenyl, ethylphenyl, phenethyl, benzyl, chlorobenzyl, methylbenzyl, methoxybenzyl, cyanobenzyl, hydroxybenzyl, dichlorobenzyl, dimethoxybenzyl, or naphthalenylmethyl. More preferably, R_1 is benzyl, cyanobenzyl, methoxybenzyl, or naphthalenylmethyl. Most preferably, R_1 is benzyl.

Preferred R_1 when R_2 and R_2' Are Hydrogen

[00123] In an embodiment wherein R_2 and R_2' are both hydrogen, preferably R_1 is chosen from optionally substituted aryl, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl, and optionally substituted heteroaryl- C_1 - C_4 -alkyl-, provided however, that R_1 is not substituted phenyl. More preferably, R_1 is optionally substituted aryl- C_1 - C_4 -alkyl- or optionally substituted heteroaryl- C_1 - C_4 -alkyl-. More preferably, when R_2 and R_2' are both hydrogen, R_1 is selected from optionally substituted phenyl- C_1 - C_4 -alkyl, and optionally substituted naphthalenylmethyl. In a more preferred embodiment wherein R_2 and R_2' are both hydrogen, R_1 is chosen from benzyl, chlorobenzyl, methylbenzyl, methoxybenzyl, cyanobenzyl, hydroxybenzyl, dichlorobenzyl, dimethoxybenzyl, and naphthalenylmethyl. More preferably, R_1 is benzyl, cyanobenzyl, methoxybenzyl, or naphthalenylmethyl. More preferably, R_1 is benzyl.

Preferred R_2

[00124] When considering the compounds of Formula I and as will be appreciated by those skilled in the art, the compounds described herein possess a potentially chiral center at the carbon to which R_2 and R_2' are attached. The R_2 and R_2' groups may be the same or different; if different, the compound is chiral (i.e., has a stereogenic center). When R_2 and R_2' are different, in preferred embodiments R_2' is hydrogen and R_2 is other than hydrogen. The invention contemplates the use of pure enantiomers and mixtures of enantiomers, including racemic mixtures, although the use of a substantially optically pure enantiomer will generally be preferred. The term "substantially optically pure" or "enantiomerically pure" means having at least about 95% of the described enantiomer with no single impurity greater than about 1% and preferably, at least about 97.5% enantiomeric excess. In a preferred embodiment, the stereogenic center to which R_2 and R_2' are attached is of the R configuration.

[00125] In one embodiment, R_2 is optionally substituted C_1 - C_4 alkyl, and R_2' is hydrogen or optionally substituted C_1 - C_4 alkyl. More preferably, R_2' is hydrogen and R_2 is optionally substituted C_1 - C_4 alkyl. In a most preferred embodiment R_2 is chosen from

methyl, ethyl, propyl (particularly, *c*-propyl or *i*-propyl), butyl (particularly, *t*-butyl), methylthioethyl, methylthiomethyl, aminobutyl, (CBZ)aminobutyl, cyclohexylmethyl, benzyloxymethyl, methylsulfinylethyl, methylsulfinylmethyl, and hydroxymethyl, and R_2 is hydrogen. Especially preferred is when R_2 is hydrogen and R_2 is ethyl or propyl (particularly, *c*-propyl or *i*-propyl). More preferably, R_2 is *i*-propyl. More preferred is the embodiment wherein the stereogenic center to which R_2 and R_2 is attached is of the *R* configuration.

[00126] In another embodiment, both R_2 and R_2 are hydrogen.

Preferred R_3 Groups When R_{12} is $-N(R_4)(COR_3)$

[00127] When considering the compounds of Formula I wherein R_{12} is $-N(R_4)(COR_3)$, in a preferred embodiment R_3 is selected from optionally substituted C_1 - C_8 alkyl, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl, optionally substituted aryl, $R_{15}O$ - and $R_{17}NH$ -, R_{15} is chosen from optionally substituted C_1 - C_8 alkyl and optionally substituted aryl, and R_{17} is chosen from hydrogen, optionally substituted C_1 - C_8 alkyl and optionally substituted aryl. Preferred R_3 are optionally substituted C_1 - C_8 alkyl (e.g., C_1 - C_8 alkyl substituted with lower-alkoxy), optionally substituted heteroaryl, and optionally substituted aryl.

[00128] In a more preferred embodiment, when R_3 is not $R_{17}NH$ - or $R_{15}O$ -, R_3 is chosen from phenyl; phenyl substituted with one or more of the following substituents: halo, C_1 - C_4 alkyl, C_1 - C_4 alkyl substituted with hydroxy (e.g., hydroxymethyl), C_1 - C_4 alkoxy, C_1 - C_4 alkyl substituted with C_1 - C_4 alkoxy, nitro, formyl, carboxy, cyano, methylenedioxy, ethylenedioxy, acyl (e.g., acetyl), *N*-acyl (e.g., *N*-acetyl) or trifluoromethyl; benzyl; phenoxymethyl-; halophenoxymethyl-; phenylvinyl-; heteroaryl-; heteroaryl- substituted with C_1 - C_4 alkyl or C_1 - C_4 alkyl substituted with halo (e.g., CF_3); C_1 - C_4 alkyl substituted with C_1 - C_4 alkoxy- and benzyloxymethyl-.

[00129] In a most preferred embodiment, when R_3 is not $R_{17}NH$ - or $R_{15}O$ -, R_3 is chosen from phenyl, halophenyl, dihalophenyl, cyanophenyl, halo(trifluoromethyl)phenyl, hydroxymethylphenyl, methoxyphenyl, ethoxyphenyl, carboxyphenyl, ethylphenyl, tolyl, methylenedioxyphenyl, ethylenedioxyphenyl, methoxychlorophenyl, dihydro-benzodioxinyl, methylhalophenyl, trifluoromethylphenyl, bis(trifluoromethyl)phenylbenzyl, furanyl, C_1 - C_4 alkyl substituted furanyl, trifluoromethylfuranyl, C_1 - C_4 alkyl substituted trifluoromethylfuranyl, benzofuranyl, thiophenyl, C_1 - C_4 alkyl substituted thiophenyl,

benzothiophenyl, benzothiadiazolyl, pyridinyl, indolyl, methylpyridinyl, trifluoromethylpyridinyl, pyrrolyl, quinolinyl, picolinyl, pyrazolyl, C₁-C₄ alkyl substituted pyrazolyl, N-methyl pyrazolyl, C₁-C₄ alkyl substituted N-methyl pyrazolyl, C₁-C₄ alkyl substituted pyrazinyl, C₁-C₄ alkyl substituted isoxazolyl, benzoisoxazolyl, morpholinomethyl, methylthiomethyl, methoxymethyl, N-methyl imidazolyl, and imidazolyl. Yet more preferably, R₃ is tolyl, halophenyl, halomethylphenyl, hydroxymethylphenyl, methylenedioxyphenyl, formylphenyl or cyanophenyl.

[00130] In a more preferred embodiment, when R₃ is R₁₇NH-, R₁₇ is chosen from hydrogen, C₁-C₄ alkyl; cyclohexyl; phenyl; and phenyl substituted with halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, or C₁-C₄ alkylthio.

[00131] In a most preferred embodiment, when R₃ is R₁₇NH-, R₁₇ is hydrogen isopropyl, butyl, cyclohexyl, phenyl, bromophenyl, dichlorophenyl, methoxyphenyl, ethylphenyl, tolyl, trifluoromethylphenyl or methylthiophenyl.

[00132] In an embodiment, wherein R₃ is R₁₅O-, R₁₅ is chosen from optionally substituted C₁-C₈ alkyl and optionally substituted aryl.

Preferred R_{3a} Groups when R₁₂ is -N(R₄)(SO₂R_{3a})

[00133] Preferably, when R₁₂ is -N(R₄)(SO₂R_{3a}), R_{3a} is chosen from C₁-C₁₃ alkyl; phenyl; naphthyl; phenyl substituted with halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, cyano, nitro, methylenedioxy, or trifluoromethyl; biphenyl and heteroaryl. More preferably, R_{3a} is chosen from phenyl substituted with halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, cyano, nitro, methylenedioxy, or trifluoromethyl and naphthyl.

Preferred R_{3b} Groups when R₁₂ is -N(R₄)(CH₂R_{3b})

[00134] Preferably, when R₁₂ is -N(R₄)(CH₂R_{3b}), R_{3b} is chosen from C₁-C₁₃ alkyl; substituted C₁-C₄ alkyl; phenyl; naphthyl; phenyl substituted with carboxy, alkoxycarbonyl, cyano, halo, C₁-C₄ alkyl-, C₁-C₄ alkoxy, nitro, methylenedioxy, or trifluoromethyl; biphenyl, benzyl; and heterocyclyl.

[00135] Most preferably, R_{3b} is chosen from halophenyl, polyhalophenyl, methylhalophenyl, tolyl, dimethylphenyl, methoxyphenyl, dimethoxyphenyl, cyanophenyl, trifluoromethylphenyl, trifluoromethoxyphenyl, bis(trifluoromethyl)phenyl, carboxyphenyl, t-butylphenyl, methoxycarbonylphenyl, piperidinyl and naphthyl.

Preferred R₄ Groups when R₁₂ is -NHR₄, -N(R₄)(COR₃), or -N(R₄)(CH₂R_{3b})

[00136] In a preferred embodiment when R₁₂ is -NHR₄, -N(R₄)(COR₃), or -N(R₄)(CH₂R_{3b}), R₄ is chosen from hydrogen, optionally substituted C₁-C₁₃ alkyl, optionally substituted aryl, optionally substituted aryl-C₁-C₄-alkyl-, optionally substituted heterocyclyl, and optionally substituted heteroaryl-C₁-C₄-alkyl- (preferably hydrogen or optionally substituted C₁-C₁₃ alkyl).

[00137] More preferably, R₄ is chosen from hydrogen, C₁-C₄ alkyl; cyclohexyl; phenyl substituted with hydroxyl, C₁-C₄ alkoxy or C₁-C₄ alkyl; benzyl; heteroarylmethyl-; heteroarylethyl-; heteroarylpropyl-; and R₁₆-alkylene-, wherein R₁₆ is hydroxyl, di(C₁-C₄ alkyl)amino-, (C₁-C₄ alkyl)amino-, amino, C₁-C₄ alkoxy-, or N-heterocyclyl-, particularly pyrrolidino, piperidino or imidazolyl.

[00138] More preferably, R₄ is R₁₆-alkylene-, wherein R₁₆ is amino, C₁-C₄ alkylamino-, di(C₁-C₄ alkyl)amino-, C₁-C₄ alkoxy-, hydroxyl, or N-heterocyclyl. Preferably R₁₆ is amino.

[00139] In a most preferred embodiment when R₁₂ is -NHR₄, -N(R₄)(COR₃), or -N(R₄)(CH₂R_{3b}), R₄ is chosen from hydrogen, methyl, ethyl, propyl, butyl, cyclohexyl, carboxyethyl, carboxymethyl, methoxyethyl, hydroxyethyl, hydroxypropyl, dimethylaminoethyl, dimethylaminopropyl, diethylaminoethyl, diethylaminopropyl, aminopropyl, methylaminopropyl, 2,2-dimethyl-3-(dimethylamino)propyl, 1-cyclohexyl-4-(diethylamino)butyl, aminoethyl, aminobutyl, aminopentyl, aminohexyl, aminoethoxyethyl, isopropylaminopropyl, diisopropylaminoethyl, 1-methyl-4-(diethylamino)butyl, (t-Boc)aminopropyl, hydroxyphenyl, benzyl, methoxyphenyl, methylmethoxyphenyl, dimethylphenyl, tolyl, ethylphenyl, (oxopyrrolidinyl)propyl, (methoxycarbonyl)ethyl, benzylpiperidinyl, pyridinylethyl, pyridinylmethyl, morpholinylethyl morpholinylpropyl, piperidinyl, azetidinylmethyl, azetidinylethyl, azetidinypropyl pyrrolidinylethyl, pyrrolidinylpropyl, piperidinylmethyl, piperidinylethyl, imidazolylpropyl, imidazolylethyl, (ethylpyrrolidinyl)methyl, (methylpyrrolidinyl)ethyl, (methylpiperidinyl)propyl, (methylpiperazinyl)propyl, furanylmethyl and indolylethyl.

[00140] More preferably, R₄ is aminoethyl, aminopropyl, aminobutyl, aminopentyl, aminohexyl, methylaminoethyl, methylaminopropyl, methylaminobutyl, methylaminopentyl, methylaminohexyl, dimethylaminoethyl, dimethylaminopropyl, dimethylaminobutyl, dimethylaminopentyl, dimethylaminohexyl, ethylaminoethyl, ethylaminopropyl, ethylaminobutyl, ethylaminopentyl, ethylaminohexyl, diethylaminoethyl, diethylaminopropyl,


diethylaminobutyl, diethylaminopentyl, or diethylaminoethyl, most preferably aminopropyl.

Preferred R_4 Groups when R_{12} is $--N(R_4)(SO_2R_{3a})$

[00141] Preferably, when R_{12} is $--N(R_4)(SO_2R_{3a})$, R_4 is chosen from C_1 - C_4 alkyl, cyclohexyl; phenyl substituted with hydroxyl, C_1 - C_4 alkoxy or C_1 - C_4 alkyl; benzyl; heteroarylmethyl-; heteroarylethyl-; heteroarylpropyl-; heteroarylethyl-; heteroarylpropyl- and R_{16} -alkylene-, wherein R_{16} is hydroxyl, di(C_1 - C_4 alkyl)amino-, (C_1 - C_4 alkyl)amino-, amino, C_1 - C_4 alkoxy-, or N-heterocyclyl-, particularly pyrrolidino, piperidino or imidazolyl.

R_{12} is an Imidazole

[00142] Preferably, when R_{12} is an imidazole, R_{12} has the formula:




wherein

R_9 is chosen from hydrogen, optionally substituted C_1 - C_8 alkyl, optionally substituted aryl, optionally substituted aryl- C_1 - C_4 -alkyl -, optionally substituted heteroaryl- C_1 - C_4 -alkyl -, optionally substituted aryl- C_1 - C_4 -alkoxy -, optionally substituted heteroaryl- C_1 - C_4 -alkoxy -, optionally substituted heteroaryl-; and R_{13} and $R_{13'}$ are independently hydrogen, optionally substituted C_1 - C_8 alkyl, optionally substituted aryl, or optionally substituted aryl- C_1 - C_4 -alkyl - (preferably optionally substituted aryl). More preferably, R_9 is phenyl substituted with C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy-, and/or halo (especially C_1 - C_4 -alkyl and/or halo); phenyl; or benzyl. Yet more preferably, R_9 is tolyl; halophenyl; or halomethylphenyl.

[00143] In a preferred embodiment, R_{13} is hydrogen and $R_{13'}$ is substituted C_1 - C_4 alkyl. More preferably, R_{13} is hydrogen and $R_{13'}$ is aminomethyl, aminoethyl, aminopropyl, acetylamino-methyl, acetylaminoethyl, benzyloxycarbonylamino-methyl or benzyloxycarbonylamino-ethyl.

R₁₂ is an Imidazoline

[00144] Preferably, when R₁₂ is an imidazoline, R₁₂ has the formula



wherein R₉ is chosen from hydrogen, optionally substituted C₁-C₈ alkyl, optionally substituted aryl, optionally substituted aryl-C₁-C₄-alkyl -, and optionally substituted heteroaryl-; and R₁₀, R_{10'}, R₁₄, and R_{14'} are independently chosen from hydrogen, optionally substituted C₁-C₈ alkyl, optionally substituted aryl, and optionally substituted aryl-C₁-C₄-alkyl -. More preferably, R₉ is methylenedioxyphenyl; phenyl; phenyl substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, and/or halo; or benzyl. In a preferred embodiment, R₉ is methylenedioxyphenyl-; phenyl; or phenyl substituted with methoxy, halo and/or methyl (preferably halo and/or methyl, including tolyl), more preferably methylenedioxyphenyl or said substituted phenyls. In another preferred embodiment, R₁₀, R_{10'}, R_{14'}, and R₁₄ are independently hydrogen or optionally substituted alkyl (preferably optionally substituted C₁-C₄ alkyl). More preferably, R₁₀ and R_{10'} are independently selected from the group consisting of hydrogen or optionally substituted C₁-C₄ alkyl (and more particularly, methyl or aminoalkyl-) and R_{14'} and R₁₄ are hydrogen.

Preferred R₅, R₆, R₇, and R₈ Groups

[00145] When considering the compounds of Formula I, in other preferred embodiments R₅, R₆, R₇ and R₈ are independently chosen from hydrogen; acyl, alkyl; alkyl substituted with alkyl, alkoxy, halo, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, lower-alkylaminocarbonyl- (e.g. methylaminocarbonyl- or ethylaminocarbonyl-), di(lower-alkyl)aminocarbonyl- (e.g. dimethylaminocarbonyl- or diethylaminocarbonyl-), aryl, or heteroaryl; alkoxy; alkoxy substituted with alkyl, acyl, alkoxy, halo, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, lower-alkylaminocarbonyl- (e.g. methylaminocarbonyl- or

ethylaminocarbonyl-), di(lower-alkyl)aminocarbonyl- (e.g. dimethylaminocarbonyl- or diethylaminocarbonyl-), aryl, or heteroaryl; halogen; hydroxyl; nitro; cyano; dialkylamino; alkylsulfonyl; alkylsulfonamido; alkylthio; carboxyalkyl; carboxamido; amidocarbonyl; aryl; aryl substituted with alkyl, acyl, alkoxy, halo, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, lower-alkylaminocarbonyl- (e.g. methylaminocarbonyl- or ethylaminocarbonyl-), di(lower-alkyl)aminocarbonyl- (e.g. dimethylaminocarbonyl- or diethylaminocarbonyl-), aryl, or heteroaryl; heteroaryl or heteroaryl substituted with alkyl, acyl, alkoxy, halo, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, lower-alkylaminocarbonyl- (e.g. methylaminocarbonyl- or ethylaminocarbonyl-), di(lower-alkyl)aminocarbonyl- (e.g. dimethylaminocarbonyl- or diethylaminocarbonyl-), aryl, or heteroaryl.

[00146] More preferably, R_5 , R_6 , R_7 , and R_8 are independently chosen from hydrogen, amino, alkylamino, hydroxyl, halogen (particularly chloro and fluoro), C_1 - C_4 alkyl (particularly methyl), C_1 - C_4 haloalkyl (particularly trifluoromethyl), C_1 - C_4 alkoxy (particularly methoxy), C_1 - C_4 haloalkoxy and cyano. More preferably, R_5 , R_6 , R_7 , and R_8 are methoxy, hydrogen, cyano, or halo (especially Cl, F). Further preferred for each of the specific substituents: R_5 is amino, alkylamino, trifluoromethyl, hydrogen or halo; R_6 is hydrogen, alkyl (particular, methyl) or halo; R_7 is hydrogen, halo, alkyl (particular, methyl), alkoxy (particularly, methoxy), cyano, or trifluoromethyl; and R_8 is hydrogen or halo. Still further preferred are the compounds where only one of R_5 , R_6 , R_7 , and R_8 is not hydrogen, especially R_7 . More preferred are the compounds where R_5 , R_6 , and R_8 are hydrogen and R_7 is cyano, methoxy or halogen (especially Cl, F).

Preferred Salt Forms

[00147] Preferred compounds will generally be capable of forming acid addition salts (i.e., will comprise a site which reacts with a pharmaceutically acceptable acid to form an acid addition salt.) The present invention includes pharmaceutically acceptable acid addition salts of the compounds of Formula I. Acid addition salts of the present compounds are prepared in a standard manner in a suitable solvent from the parent compound and an excess of an acid, such as hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, maleic, succinic or methanesulfonic. Preferred salt forms are hydrochloric, phosphoric, and oxalic acid salts with the hydrochloric acid salt form being especially preferred.

[00148] The salts and/or solvates of the compounds of the formula (I) which are not pharmaceutically acceptable may be useful as intermediates in the preparation of pharmaceutically acceptable salts and/or solvates of compounds of formula (I) or the compounds of the formula (I) themselves, and as such form another aspect of the present invention.

Preferred Subgenus

[00149] In a particularly preferred subgenus of compounds of Formula I, R_1 is benzyl, halobenzyl, methoxybenzyl-, cyanobenzyl, or naphthalenylmethyl-; R_2 is ethyl or propyl; R_2 is hydrogen; R_5 is hydrogen; R_6 is hydrogen; R_7 is halo, cyano, methoxy or hydrogen; R_8 is hydrogen; and R_{12} is $-NR_4(COR_3)$ wherein R_3 is optionally substituted aryl (preferably, halophenyl, halomethylphenyl-, methylenedioxyphenyl-, methoxyphenyl-, ethoxyphenyl-, cyanophenyl- or phenyl substituted with lower-acyl or lower-alkylaminocarbonyl-, e.g. methylaminocarbonyl- or ethylaminocarbonyl-, or di(lower-alkyl)aminocarbonyl-, e.g. dimethylaminocarbonyl- or diethylaminocarbonyl-; and R_4 is R_{16} -alkylene- wherein R_{16} is hydroxyl, di(C_1 - C_4)alkylamino-, (C_1 - C_4 alkyl)amino-, amino, pyrrolidino, piperidino, imidazolyl and morpholino (more preferably in such embodiments, R_1 is benzyl, halobenzyl, methoxybenzyl, cyanobenzyl, or naphthalenylmethyl; and R_2 is propyl (especially i- or c-propyl).

[00150] In another particularly preferred subgenus of compounds of Formula I, R_1 is benzyl, halobenzyl, methoxybenzyl-, cyanobenzyl, or naphthalenylmethyl-; R_2 is ethyl or propyl; R_2 is hydrogen; R_5 is hydrogen; R_6 is hydrogen; R_7 is halo, cyano, methoxy or hydrogen; R_8 is hydrogen; R_{12} is $-NR_4(CH_2R_{3b})$ wherein R_4 is R_{16} -alkylene- wherein R_{16} is hydroxyl, di(C_1 - C_4)alkylamino-, (C_1 - C_4 alkyl)amino-, amino, pyrrolidino, piperidino, imidazolyl or morpholino.; and R_{3b} is optionally substituted aryl.

[00151] In a particularly preferred subgenus of compounds of Formula I, R_1 is benzyl, halobenzyl, methoxybenzyl, cyanobenzyl, or naphthalenylmethyl; R_2 is chosen from ethyl or propyl; R_2 is hydrogen; R_5 is hydrogen; R_6 is hydrogen; R_7 is halo, cyano, methoxy or hydrogen; R_8 is hydrogen; and R_{12} is optionally substituted imidazolyl of the above formula wherein R_{10} , $R_{10'}$, R_{14} and $R_{14'}$ are independently hydrogen or optionally substituted alkyl (preferably optionally substituted C_1 - C_4 alkyl); and R_9 is optionally substituted phenyl (preferably, halophenyl, halomethylphenyl, tolyl, or methylenedioxyphenyl). More preferably

in such embodiments R_1 is benzyl, methoxybenzyl, or cyanobenzyl; R_2 is propyl (especially i- or c- propyl); and R_{16} is amino.

[00152] In a particularly preferred subgenus of compounds of Formula I, R_1 is benzyl, halobenzyl, methoxybenzyl, cyanobenzyl, or naphthalenylmethyl; R_2 is chosen from ethyl or propyl; $R_{2'}$ is hydrogen; R_5 is hydrogen; R_6 is hydrogen; R_7 is halo, cyano, methoxy or hydrogen; R_8 is hydrogen; and R_{12} is optionally substituted imidazole of the above formula wherein R_{13} is hydrogen and $R_{13'}$ is hydrogen or optionally substituted alkyl (preferably optionally substituted C_1 - C_4 alkyl); and R_9 is optionally substituted aryl (preferably, halophenyl, halomethylphenyl, tolyl, or methylenedioxyphenyl). More preferably, R_{13} is hydrogen and $R_{13'}$ is aminomethyl, aminoethyl aminopropyl, acetyl-amino-methyl, acetyl-aminoethyl, benzyloxycarbonylamino-methyl or Benzyloxycarbonylamino-ethyl. More preferably in such embodiments R_1 is benzyl, methoxybenzyl, or cyanobenzyl; R_2 is propyl (especially i- or c- propyl); and R_{16} is amino.

[00153] When R_{12} is $-N(R_4)(SO_2R_{3a})$, R_1 is most preferably chosen from C_1 - C_4 alkyl, benzyl, substituted benzyl and substituted phenyl; R_2 is C_1 - C_4 alkyl; $R_{2'}$ is hydrogen; R_{3a} is chosen from substituted phenyl and naphthyl; R_4 is R_{16} -alkylene-; R_7 is hydrogen, fluoro, methyl or chloro; R_5 , R_6 and R_8 are hydrogen; and R_{16} is chosen from hydroxyl, di(C_1 - C_4)amino, (C_1 - C_4 alkyl)amino, amino, pyrrolidino, piperidino, imidazolyl and morpholino.

[00154] When R_{12} is $-NHR_4$ or $-N(R_4)(CH_2R_{3b})$, R_1 is preferably chosen from hydrogen, optionally substituted C_1 - C_4 alkyl, optionally substituted benzyl, optionally substituted phenyl, and optionally substituted naphthalenylmethyl; R_2 is optionally substituted C_1 - C_4 alkyl and $R_{2'}$ is hydrogen; R_{3b} is chosen from optionally substituted alkyl; optionally substituted phenyl; biphenyl, optionally substituted aralkyl; and optionally substituted heterocyclyl; and R_4 is chosen from hydrogen, optionally substituted C_1 - C_4 alkyl; cyclohexyl; optionally substituted phenyl; optionally substituted benzyl; heterocyclyl; heteroaryl-methyl; heteroarylethyl; and heteroarylpropyl. More preferably, R_4 is R_{16} -alkylene-, wherein R_{16} is hydroxyl, di(C_1 - C_4)alkylamino-, (C_1 - C_4 alkyl)amino-, amino, C_1 - C_4 alkoxy-, or N-heterocyclyl.

[00155] When R_{12} is $-NHR_4$ or $-N(R_4)(CH_2R_{3b})$, R_1 is most preferably chosen from C_1 - C_4 alkyl, optionally substituted benzyl, and optionally substituted phenyl (more preferably optionally substituted benzyl, e.g., benzyl, cyanobenzyl); R_2 is optionally substituted C_1 - C_4

alkyl (more preferably propyl, especially i- or c-propyl); R_2 is hydrogen; R_{3b} is chosen from optionally substituted phenyl, optionally substituted heterocyclyl and naphthyl; R_4 is chosen from hydrogen, optionally substituted benzyl, optionally substituted heterocyclyl and R_{16} -alkylene-; R_6 and R_7 are chosen from halo, cyano, methoxy or hydrogen; R_5 and R_8 are hydrogen; and R_{16} is chosen from di(C_1 - C_4 alkylamino)-, (C_1 - C_4 alkyl)amino-, amino, pyrrolidinyl, piperidinyl, imidazolyl and morpholinyl.

[00156] In a particularly preferred subgenus of compounds of Formula I, R_1 is benzyl, halobenzyl (especially Cl-benzyl and F-benzyl), methoxybenzyl-, cyanobenzyl, or naphthalenylmethyl-; R_2 is ethyl or propyl; R_2 is hydrogen; R_5 is hydrogen; R_6 is hydrogen; R_7 is halo, cyano, methoxy or hydrogen; R_8 is hydrogen; and R_{12} is $-NHR_4$ wherein R_4 is hydrogen (more preferably in such embodiments, R_1 is benzyl, halobenzyl, cyanobenzyl; and R_2 is propyl, especially i- propyl or c-propyl).

[00157] When R_{3b} is present, it is most preferably chosen from phenyl substituted with one or more halo, methyl, methoxy, cyano, trifluoromethyl, trifluoromethoxy, carboxy, and or methoxycarbonyl groups [e.g., halophenyl, polyhalophenyl, tolyl, dimethylphenyl, methoxyphenyl, dimethoxyphenyl, cyanophenyl, trifluoromethylphenyl, trifluoromethoxyphenyl, bis(trifluoromethyl)phenyl, carboxyphenyl, t-butylphenyl, methoxycarbonylphenyl]; piperidinyl and naphthyl.

[00158] Particularly preferred compounds include:

N-(3-Amino-propyl)-N-{1-[3-(3-cyano-benzyl)-7-hydroxy-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-3-fluoro-4-methyl-benzamide;

4-Acetyl-N-(3-amino-propyl)-N-[1-(3-benzyl-7-methoxy-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-benzamide;

2-{1-[4-(2-Amino-ethyl)-2-p-tolyl-imidazol-1-yl]-2-methyl-propyl}-3-benzyl-4-oxo-4H-chromene-7-carbonitrile;

3-Benzyl-2-[1-(4,4-dimethyl-2-p-tolyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-4-oxo-4H-chromene-7-carbonitrile;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-{1-[7-chloro-3-(3-cyano-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-amide;

N-(3-Amino-propyl)-N-{1-[7-chloro-3-(3-cyano-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methyl-benzamide;

N-(3-Amino-propyl)-N-[1-(7-chloro-3-naphthalen-1-ylmethyl-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-2-methoxy-acetamide;

4-Acetyl-N-(3-amino-propyl)-N-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-benzamide;

N-(3-Amino-propyl)-N-{1-[7-chloro-3-(3-cyano-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-3-fluoro-4-methyl-benzamide;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

3-Benzyl-2-[1-(4,4-dimethyl-2-p-tolyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-7-hydroxy-chromen-4-one;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-3-fluoro-4-methyl-benzamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-methoxy-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-3-fluoro-4-methyl-benzamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-methoxy-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-methyl-benzamide;

3-Benzyl-2-[1-(4,4-dimethyl-2-p-tolyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-7-methoxy-chromen-4-one;

3-Benzyl-7-fluoro-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,4-dimethyl-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-chromen-4-one;

3-Benzyl-2-[1-(4,4-dimethyl-2-p-tolyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-7-fluoro-chromen-4-one;

3-Benzyl-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,4-dimethyl-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-7-cyano-chromen-4-one;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-methyl-benzamide;

4-Acetyl-N-(3-amino-propyl)-N-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-benzamide;

N-(3-Amino-propyl)-3-fluoro-N-{1-[7-fluoro-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methyl-benzamide;

4-Acetyl-N-(3-amino-propyl)-N-{1-[7-chloro-3-(3-cyano-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-benzamide;

(2-{1-[1-(3-Benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-2-p-tolyl-1H-

imidazol-4-yl}-ethyl)-carbamic acid benzyl ester;

2-[1-(2-Benzo[1,3]dioxol-5-yl-4,4-dimethyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-7-cyano-chromen-4-one;

4-Acetyl-N-(3-amino-propyl)-N-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-benzamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-3-fluoro-4-methyl-benzamide;

N-(3-Amino-propyl)-N-{1-[7-fluoro-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methoxy-benzamide;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-{1-[7-fluoro-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-amide;

2-{1-[4-(2-Amino-ethyl)-2-p-tolyl-imidazol-1-yl]-2-methyl-propyl}-3-benzyl-7-chloro-chromen-4-one;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-hydroxy-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-3-fluoro-4-methyl-benzamide;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-hydroxy-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-methyl-benzamide;

N-(3-Amino-propyl)-N-{1-[7-fluoro-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-2-methoxy-acetamide;

N-(3-Amino-propyl)-N-{1-[7-fluoro-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methyl-benzamide;

N-(3-Amino-propyl)-N-[(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-cyclopropyl-methyl]-4-methyl-benzamide;

2,3-Dihydro-benzo[1,4]dioxine-6-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

N-(3-Amino-propyl)-3-fluoro-N-{1-[7-cyano-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methyl-benzamide;

3-Benzyl-7-chloro-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-chromen-4-one;

3-Benzyl-7-fluoro-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-chromen-4-one;

2-[1-(4-Aminomethyl-2-p-tolyl-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one;

3-Benzyl-7-methoxy-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-chromen-4-one;

2-[1-(2-Benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one;

4-Acetyl-N-(3-amino-propyl)-N-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-benzamide;

4-Acetyl-N-(3-amino-propyl)-N-{1-[7-cyano-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-benzamide;

3-Benzyl-7-chloro-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-chromen-4-one;

3-Benzyl-7-fluoro-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-chromen-4-one;

2-[1-(2-Benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-7-fluoro-chromen-4-one;

N-(3-Amino-propyl)-N-{1-[7-cyano-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methyl-benzamide;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-{1-[7-cyano-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-amide;

3-(2-{1-[2-(3-Fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-7-methoxy-4-oxo-4H-chromen-3-ylmethyl)-benzonitrile;

3-{7-Chloro-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-4-oxo-4H-chromen-3-ylmethyl}-benzonitrile;

3-{7-Methoxy-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-4-oxo-4H-chromen-3-ylmethyl}-benzonitrile;

3-{7-Fluoro-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-4-oxo-4H-chromen-3-ylmethyl}-benzonitrile;

N-(3-Amino-propyl)-N-{1-[7-chloro-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-3-fluoro-4-methyl-benzamide;

N-(3-Amino-propyl)-N-{1-[3-(3-cyano-benzyl)-7-methoxy-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methyl-benzamide;

N-(3-Amino-propyl)-N-{1-[3-(3-cyano-benzyl)-7-methoxy-4-oxo-4H-chromen-2-yl]-

2-methyl-propyl}-3-fluoro-4-methyl-benzamide;

3-{2-[1-(2-Benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-7-fluoro-4-oxo-4H-chromen-3-ylmethyl}-benzonitrile;

3-Benzyl-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-4-oxo-4H-chromene-7-carbonitrile;

2-[1-(2-Benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-4-oxo-4H-chromene-7-carbonitrile;

3-(7-Chloro-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-4-oxo-4H-chromen-3-ylmethyl)-benzonitrile;

3-{2-[1-(2-Benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-7-methoxy-4-oxo-4H-chromen-3-ylmethyl}-benzonitrile;

3-(7-Fluoro-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-4-oxo-4H-chromen-3-ylmethyl)-benzonitrile;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-{1-[7-cyano-3-(3-cyano-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-amide;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-{1-[3-(3-cyano-benzyl)-7-methoxy-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-amide;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-{1-[3-(3-cyano-benzyl)-7-fluoro-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-amide;

N-(3-Amino-propyl)-N-{1-[3-(3-cyano-benzyl)-7-fluoro-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methyl-benzamide;

Benzo[1,3]dioxole-5-carboxylic acid (3-amino-propyl)-{1-[7-chloro-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-amide;

N-(3-Amino-propyl)-N-{1-[7-cyano-3-(3-cyano-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-4-methyl-benzamide;

3-Benzyl-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-4-oxo-4H-chromene-7-carbonitrile;

2-[1-(2-Benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-7-methoxy-chromen-4-one;

N-(3-Amino-propyl)-N-{1-[7-chloro-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-2-methoxy-acetamide;

N-(3-Amino-propyl)-N-{1-[7-cyano-3-(3-cyano-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl}-3-fluoro-4-methyl-benzamide;

N-(3-Amino-propyl)-N-[1-[3-(3-cyano-benzyl)-7-fluoro-4-oxo-4H-chromen-2-yl]-2-methyl-propyl]-3-fluoro-4-methyl-benzamide;

2-{1-[4-(2-Amino-ethyl)-2-p-tolyl-imidazol-1-yl]-2-methyl-propyl}-7-chloro-3-(3-methoxy-benzyl)-chromen-4-one;

2-{1-[4-(2-Amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-3-benzyl-7-fluoro-chromen-4-one;

2-{1-[4-(2-Amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-3-benzyl-4-oxo-4H-chromene-7-carbonitrile;

2-{1-[4-(2-Amino-ethyl)-2-p-tolyl-imidazol-1-yl]-2-methyl-propyl}-3-benzyl-7-fluoro-chromen-4-one;

2-{1-[4-(2-Amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-7-chloro-3-(3-methoxy-benzyl)-chromen-4-one;

2-{1-[4-(2-Amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-3-benzyl-7-chloro-chromen-4-one;

3-Benzyl-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-7-methoxy-chromen-4-one;

3-(2-{1-[4-(2-Amino-ethyl)-2-p-tolyl-imidazol-1-yl]-2-methyl-propyl}-7-chloro-4-oxo-4H-chromen-3-ylmethyl)-benzonitrile;

3-(2-{1-[4-(2-Amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-7-chloro-4-oxo-4H-chromen-3-ylmethyl)-benzonitrile;

2-{1-[4-(2-Amino-ethyl)-2-p-tolyl-imidazol-1-yl]-2-methyl-propyl}-7-fluoro-3-(3-methoxy-benzyl)-chromen-4-one;

2-{1-[4-(2-Amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-3-(3-cyano-benzyl)-4-oxo-4H-chromene-7-carbonitrile;

N-(3-Amino-propyl)-3-fluoro-N-[1-[7-hydroxy-3-(3-methoxy-benzyl)-4-oxo-4H-chromen-2-yl]-2-methyl-propyl]-4-methyl-benzamide;

2-{1-[4-(2-Amino-ethyl)-2-p-tolyl-imidazol-1-yl]-2-methyl-propyl}-3-(3-methoxy-benzyl)-4-oxo-4H-chromene-7-carbonitrile;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-ethoxy-benzamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-6-trifluoromethyl-nicotinamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-

propyl]-6-trifluoromethyl-nicotinamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-isonicotinamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-cyano-benzamide;

4-Acetylamino-N-(3-amino-propyl)-N-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-benzamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-6-trifluoromethyl-nicotinamide;

Benzo[1,2,3]thiadiazole-5-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

Benzo[1,2,3]thiadiazole-5-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

4-Acetylamino-N-(3-amino-propyl)-N-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-benzamide;

Benzo[1,2,3]thiadiazole-5-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-nicotinamide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-methoxy-benzamide;

Benzo[1,2,3]thiadiazole-5-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-pyrazine-2-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

N-(3-Amino-propyl)-N-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-3-dimethylamino-benzamide;

2-[1-(2-Benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-3-(3-methoxy-benzyl)-4-oxo-4H-chromene-7-carbonitrile;

7-Chloro-3-(3-methoxy-benzyl)-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-chromen-4-one;

7-Chloro-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-3-(3-methoxy-benzyl)-chromen-4-one;

2-[1-(2-Benzo[1,3]dioxol-5-yl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-7-chloro-3-(3-methoxy-benzyl)-chromen-4-one;

7-Fluoro-3-(3-methoxy-benzyl)-2-[2-methyl-1-(2-p-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-chromen-4-one;

7-Fluoro-2-{1-[2-(3-fluoro-4-methyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-3-(3-methoxy-benzyl)-chromen-4-one;

2-{1-[4-(2-Amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-7-fluoro-3-(3-methoxy-benzyl)-chromen-4-one;

2-{1-[4-(2-Acetyl-amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-3-(3-methoxy-benzyl)-4-oxo-4H-chromene-7-carboxylic acid amide;

3-(2-{1-[4-(2-Amino-ethyl)-2-(3-fluoro-4-methyl-phenyl)-imidazol-1-yl]-2-methyl-propyl}-7-fluoro-4-oxo-4H-chromen-3-ylmethyl)-benzonitrile;

N-{1-[1-(3-Benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-2-p-tolyl-1H-imidazol-4-ylmethyl}-acetamide;

Benzo[b]thiophene-2-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

1-Methyl-1H-indole-2-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-tert-Butyl-2-methyl-2H-pyrazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

2,5-Dimethyl-2H-pyrazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-tert-Butyl-2-methyl-2H-pyrazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

2,5-Dimethyl-2H-pyrazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

2,5-Dimethyl-2H-pyrazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-2H-pyrazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-2H-pyrazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

Furan-2-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-

2-yl)-2-methyl-propyl]-amide;

Furan-2-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

2,5-Dimethyl-furan-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

2,5-Dimethyl-furan-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-thiophene-2-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-thiophene-2-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-isoxazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-2-trifluoromethyl-furan-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-isoxazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-isoxazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

Benzo[c]isoxazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

Benzo[c]isoxazole-3-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

1-Methyl-1H-pyrrole-2-carboxylic acid (3-amino-propyl)-[1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

1-Methyl-1H-imidazole-4-carboxylic acid (3-amino-propyl)-[(R)-1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

N-(3-Amino-propyl)-N-[(R)-1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-3-dimethylamino-benzamide;

5-Methyl-2-trifluoromethyl-furan-3-carboxylic acid (3-amino-propyl)-[(R)-1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

5-Methyl-isoxazole-3-carboxylic acid (3-amino-propyl)-[(R)-1-(3-benzyl-7-cyano-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

1-Methyl-1H-imidazole-4-carboxylic acid (3-amino-propyl)-[(R)-1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

1-Methyl-1H-pyrrole-2-carboxylic acid (3-amino-propyl)-[(R)-1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide;

Benzo[c]isoxazole-3-carboxylic acid (3-amino-propyl)-[(R)-1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide; and

5-Methyl-isoxazole-3-carboxylic acid (3-amino-propyl)-[(R)-1-(3-benzyl-7-fluoro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-amide.

Utility, Testing and Administration

General Utility

[00159] Once made, the compounds of the invention find use in a variety of applications involving alteration of mitosis. As will be appreciated by those skilled in the art, mitosis may be altered in a variety of ways; that is, one can affect mitosis either by increasing or decreasing the activity of a component in the mitotic pathway. Stated differently, mitosis may be affected (e.g., disrupted) by disturbing equilibrium, either by inhibiting or activating certain components. Similar approaches may be used to alter meiosis.

[00160] In a preferred embodiment, the compounds of the invention are used to inhibit mitotic spindle formation, thus causing prolonged cell cycle arrest in mitosis. By "inhibit" in this context is meant decreasing or interfering with mitotic spindle formation or causing mitotic spindle dysfunction. By "mitotic spindle formation" herein is meant organization of microtubules into bipolar structures by mitotic kinesins. By "mitotic spindle dysfunction" herein is meant mitotic arrest and monopolar spindle formation.

[00161] The compounds of the invention are useful to bind to, and/or inhibit the activity of, a mitotic kinesin, KSP. In a preferred embodiment, the KSP is human KSP, although the compounds may be used to bind to or inhibit the activity of KSP kinesins from other organisms. In this context, "inhibit" means either increasing or decreasing spindle pole separation, causing malformation, i.e., splaying, of mitotic spindle poles, or otherwise causing morphological perturbation of the mitotic spindle. Also included within the definition of KSP for these purposes are variants and/or fragments of KSP. See U.S. Patent 6,437,115, hereby incorporated by reference in its entirety. The compounds of the invention have been shown to have specificity for KSP. However, the present invention includes the use of the compounds to bind to or modulate other mitotic kinesins.

[00162] The compounds of the invention are used to treat cellular proliferation diseases. Such disease states which can be treated by the compounds, compositions and methods provided herein include, but are not limited to, cancer (further discussed below), autoimmune disease, fungal disorders, arthritis, graft rejection, inflammatory bowel disease, cellular proliferation induced after medical procedures, including, but not limited to, surgery, angioplasty, and the like. Treatment includes inhibiting cellular proliferation. It is appreciated that in some cases the cells may not be in an abnormal state and still require treatment. Thus, in one embodiment, the invention herein includes application to cells or individuals afflicted or subject to impending affliction with any one of these disorders or states.

[00163] The compounds, compositions and methods provided herein are particularly deemed useful for the treatment of cancer including solid tumors such as skin, breast, brain, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that may be treated by the compounds, compositions and methods of the invention include, but are not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor (nephroblastoma), lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrosarcoma)

exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma (pinealoma), glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma (serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma), granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia (acute and chronic), acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma (malignant lymphoma); Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above identified conditions.

Testing

[00164] For assay of KSP-modulating activity, generally either KSP or a compound according to the invention is non-diffusably bound to an insoluble support having isolated sample receiving areas (e.g., a microtiter plate, an array, etc.). The insoluble support may be made of any composition to which the sample can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports may be solid or porous and of any convenient shape. Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharides, nylon or nitrocellulose, Teflon™, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the sample is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the sample and is

nondiffusable. Preferred methods of binding include the use of antibodies (which do not sterically block either the ligand binding site or activation sequence when the protein is bound to the support), direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the sample, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

[00165] The compounds of the invention may be used on their own to inhibit the activity of a mitotic kinesin, particularly KSP. In one embodiment, a compound of the invention is combined with KSP and the activity of KSP is assayed. Kinesin (including KSP) activity is known in the art and includes one or more kinesin activities. Kinesin activities include the ability to affect ATP hydrolysis; microtubule binding; gliding and polymerization/depolymerization (effects on microtubule dynamics); binding to other proteins of the spindle; binding to proteins involved in cell-cycle control; serving as a substrate to other enzymes, such as kinases or proteases; and specific kinesin cellular activities such as spindle pole separation.

[00166] Methods of performing motility assays are well known to those of skill in the art. (See e.g., Hall, et al. (1996), *Biophys. J.*, 71: 3467-3476, Turner et al., 1996, *Anal Biochem.* 242 (1):20-5; Gittes et al., 1996, *Biophys. J.* 70(1): 418-29; Shirakawa et al., 1995, *J. Exp. Biol.* 198: 1809-15; Winkelmann et al., 1995, *Biophys. J.* 68: 2444-53; Winkelmann et al., 1995, *Biophys. J.* 68: 72S.)

[00167] Methods known in the art for determining ATPase hydrolysis activity also can be used. Preferably, solution based assays are utilized. U.S. Patent 6,410,254, hereby incorporated by reference in its entirety, describes such assays. Alternatively, conventional methods are used. For example, P_i release from kinesin can be quantified. In one preferred embodiment, the ATPase hydrolysis activity assay utilizes 0.3 M PCA (perchloric acid) and malachite green reagent (8.27 mM sodium molybdate II, 0.33 mM malachite green oxalate, and 0.8 mM Triton X-100). To perform the assay, 10 μ L of the reaction mixture is quenched in 90 μ L of cold 0.3 M PCA. Phosphate standards are used so data can be converted to mM inorganic phosphate released. When all reactions and standards have been quenched in PCA, 100 μ L of malachite green reagent is added to the relevant wells in e.g., a microtiter plate. The mixture is developed for 10-15 minutes and the plate is read at an absorbance of 650 nm. If phosphate standards were used, absorbance readings can be converted to mM P_i and

plotted over time. Additionally, ATPase assays known in the art include the luciferase assay.

[00168] ATPase activity of kinesin motor domains also can be used to monitor the effects of agents and are well known to those skilled in the art. In one embodiment ATPase assays of kinesin are performed in the absence of microtubules. In another embodiment, the ATPase assays are performed in the presence of microtubules. Different types of agents can be detected in the above assays. In a preferred embodiment, the effect of a agent is independent of the concentration of microtubules and ATP. In another embodiment, the effect of the agents on kinesin ATPase can be decreased by increasing the concentrations of ATP, microtubules or both. In yet another embodiment, the effect of the agent is increased by increasing concentrations of ATP, microtubules or both.

[00169] Compounds that inhibit the biochemical activity of KSP in vitro may then be screened in vivo. In vivo screening methods include assays of cell cycle distribution, cell viability, or the presence, morphology, activity, distribution, or number of mitotic spindles. Methods for monitoring cell cycle distribution of a cell population, for example, by flow cytometry, are well known to those skilled in the art, as are methods for determining cell viability. See for example, U.S. Patent 6,437,115, hereby incorporated by reference in its entirety. Microscopic methods for monitoring spindle formation and malformation are well known to those of skill in the art (see, e.g., Whitehead and Rattner (1998), J. Cell Sci. 111:2551-61; Galgio et al, (1996) J. Cell Biol., 135:399-414), each incorporated herein by reference in its entirety.

[00170] The compounds of the invention inhibit the KSP kinesin. One measure of inhibition is IC_{50} , defined as the concentration of the compound at which the activity of KSP is decreased by fifty percent relative to a control. Preferred compounds have IC_{50} 's of less than about 1 mM, with preferred embodiments having IC_{50} 's of less than about 100 μ M, with more preferred embodiments having IC_{50} 's of less than about 10 μ M, with particularly preferred embodiments having IC_{50} 's of less than about 1 μ M, and especially preferred embodiments having IC_{50} 's of less than about 100 nM, and with the most preferred embodiments having IC_{50} 's of less than about 10 nM. Measurement of IC_{50} is done using an ATPase assay such as described herein.

[00171] Another measure of inhibition is K_i . For compounds with IC_{50} 's less than 1 μ M, the K_i or K_d is defined as the dissociation rate constant for the interaction of the compounds described herein with KSP. Preferred compounds have K_i 's of less than about

100 μM , with preferred embodiments having K_i 's of less than about 10 μM , and particularly preferred embodiments having K_i 's of less than about 1 μM and especially preferred embodiments having K_i 's of less than about 100 nM, and with the most preferred embodiments having K_i 's of less than about 10 nM.

[00172] The K_i for a compound is determined from the IC_{50} based on three assumptions and the Michaelis-Menten equation. First, only one compound molecule binds to the enzyme and there is no cooperativity. Second, the concentrations of active enzyme and the compound tested are known (i.e., there are no significant amounts of impurities or inactive forms in the preparations). Third, the enzymatic rate of the enzyme-inhibitor complex is zero. The rate (i.e., compound concentration) data are fitted to the equation:

$$V = V_{\max} E_0 \left[1 - \frac{(E_0 + I_0 + K_d) - \sqrt{(E_0 + I_0 + K_d)^2 - 4 E_0 I_0}}{2 E_0} \right]$$

where V is the observed rate, $V_{\max}$ is the rate of the free enzyme, I_0 is the inhibitor concentration, E_0 is the enzyme concentration, and K_d is the dissociation constant of the enzyme-inhibitor complex.

[00173] Another measure of inhibition is GI_{50} , defined as the concentration of the compound that results in a decrease in the rate of cell growth by fifty percent. Preferred compounds have GI_{50} 's of less than about 1 mM; those having a GI_{50} of less than about 20 μM are more preferred; those having a GI_{50} of less than about 10 μM more so; those having a GI_{50} of less than about 1 μM more so; those having a GI_{50} of less than about 100 nM more so; and those having a GI_{50} of less than about 10 nM even more so. Measurement of GI_{50} is done using a cell proliferation assay such as described herein. Compounds of this class were found to inhibit cell proliferation.

[00174] In vitro potency of small molecule inhibitors is determined, for example, by assaying human ovarian cancer cells (SKOV3) for viability following a 72-hour exposure to a 9-point dilution series of compound. Cell viability is determined by measuring the absorbance of formazon, a product formed by the bioreduction of MTS/PMS, a commercially available reagent. Each point on the dose-response curve is calculated as a percent of untreated control cells at 72 hours minus background absorption (complete cell kill).

[00175] Anti-proliferative compounds that have been successfully applied in the clinic to treatment of cancer (cancer chemotherapeutics) have GI_{50} 's that vary greatly. For example,

in A549 cells, paclitaxel GI_{50} is 4 nM, doxorubicin is 63 nM, 5-fluorouracil is 1 μ M, and hydroxyurea is 500 μ M (data provided by National Cancer Institute, Developmental Therapeutic Program, <http://dtp.nci.nih.gov/>). Therefore, compounds that inhibit cellular proliferation, irrespective of the concentration demonstrating inhibition, may be useful.

[00176] To employ the compounds of the invention in a method of screening for compounds that bind to KSP kinesin, the KSP is bound to a support, and a compound of the invention is added to the assay. Alternatively, the compound of the invention is bound to the support and KSP is added. Classes of compounds among which novel binding agents may be sought include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled in vitro protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

[00177] The determination of the binding of the compound of the invention to KSP may be done in a number of ways. In a preferred embodiment, the compound is labeled, for example, with a fluorescent or radioactive moiety, and binding is determined directly. For example, this may be done by attaching all or a portion of KSP to a solid support, adding a labeled test compound (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the amount of the label is that present on the solid support.

[00178] By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g., radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

[00179] In some embodiments, only one of the components is labeled. For example, the kinesin proteins may be labeled at tyrosine positions using ^{125}I , or with fluorophores. Alternatively, more than one component may be labeled with different labels; using ^{125}I for the proteins, for example, and a fluorophor for the antimetabolic agents.

[00180] The compounds of the invention may also be used as competitors to screen for additional drug candidates. "Candidate agent" or "drug candidate" or grammatical equivalents as used herein describe any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity. They may be capable of directly or indirectly altering the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular proliferation protein binding and/or activity is screened. Screens of this sort may be performed either in the presence or absence of microtubules. In the case where protein binding or activity is screened, preferred embodiments exclude molecules already known to bind to that particular protein, for example, polymer structures such as microtubules, and energy sources such as ATP. Preferred embodiments of assays herein include candidate agents which do not bind the cellular proliferation protein in its endogenous native state termed herein as "exogenous" agents. In another preferred embodiment, exogenous agents further exclude antibodies to KSP.

[00181] Candidate agents can encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

[00182] Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such

as acylation, alkylation, esterification, and/or amidification to produce structural analogs.

[00183] Competitive screening assays may be done by combining KSP and a drug candidate in a first sample. A second sample comprises a compound of the present invention, KSP and a drug candidate. This may be performed in either the presence or absence of microtubules. The binding of the drug candidate is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of a drug candidate capable of binding to KSP and potentially inhibiting its activity. That is, if the binding of the drug candidate is different in the second sample relative to the first sample, the drug candidate is capable of binding to KSP.

[00184] In a preferred embodiment, the binding of the candidate agent to KSP is determined through the use of competitive binding assays. In this embodiment, the competitor is a binding moiety known to bind to KSP, such as an antibody, peptide, binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as between the candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

[00185] In one embodiment, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to KSP for a time sufficient to allow binding, if present. Incubations may be performed at any temperature which facilitates optimal activity, typically between 4 and 40°C.

[00186] Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

[00187] In a preferred embodiment, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent is binding to KSP and thus is capable of binding to, and potentially inhibiting, the activity of KSP. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

[00188] In an alternative embodiment, the candidate agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the

competitor may indicate the candidate agent is bound to KSP with a higher affinity. Thus, if the candidate agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate the candidate agent is capable of binding to KSP.

[00189] Inhibition is tested by screening for candidate agents capable of inhibiting the activity of KSP comprising the steps of combining a candidate agent with KSP, as above, and determining an alteration in the biological activity of KSP. Thus, in this embodiment, the candidate agent should both bind to KSP (although this may not be necessary), and alter its biological or biochemical activity as defined herein. The methods include both in vitro screening methods and in vivo screening of cells for alterations in cell cycle distribution, cell viability, or for the presence, morphology, activity, distribution, or amount of mitotic spindles, as are generally outlined above.

[00190] Alternatively, differential screening may be used to identify drug candidates that bind to the native KSP, but cannot bind to modified KSP.

[00191] Positive controls and negative controls may be used in the assays. Preferably all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the binding of the agent to the protein. Following incubation, all samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

[00192] A variety of other reagents may be included in the screening assays. These include reagents like salts, neutral proteins, e.g., albumin, detergents, etc which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

Administration

[00193] Accordingly, the compounds of the invention are administered to cells. By "cells" herein is meant any cell in which mitosis or meiosis can be altered. By "administered" herein is meant administration of a therapeutically effective dose of a compound of the invention to a cell either in cell culture or in a patient. By "therapeutically effective dose" herein is meant a dose that produces the effects for which it is administered. The exact dose

will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is known in the art, adjustments for systemic versus localized delivery, route of administration, age, body weight, general health, sex, diet, time of administration, nature of the formulation, drug interaction, and the precise condition requiring treatment and its severity may be necessary, and will be ascertainable with routine experimentation by those skilled in the art. However, an effective amount of a compound of Formula I for the treatment of neoplastic growth (typically by intravenous administration), for example colon or breast carcinoma, will generally be in the range of 0.1 to 100 (including 1 to 100) mg/m² of surface area of the recipient per dose on a once a week to once a month schedule and usually in the range of 2 to 30 mg/m² of surface area of the recipient per dose on a once a week to once a month schedule. An effective amount of a salt, solvate, or solvate of a salt of a compound of Formula I may be determined as a proportion of the effective amount of the compound of Formula I per se. It is envisaged that similar dosages would be appropriate for treatment of the other conditions referred to herein.

[00194] A "patient" for the purposes of the present invention includes both humans and other animals, particularly mammals, and other organisms. Thus the methods are applicable to both human therapy and veterinary applications. In the preferred embodiment the patient is a mammal, and in the most preferred embodiment the patient is human.

[00195] Compounds of the invention having the desired pharmacological activity may be administered, preferably as a pharmaceutically acceptable composition comprising an pharmaceutical excipient, to a patient, as described herein. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways as discussed below. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt.%.

[00196] The agents may be administered alone or in combination with other treatments, i.e., radiation, or other chemotherapeutic agents such as the taxane class of agents that appear to act on microtubule formation or the camptothecin class of topoisomerase I inhibitors. When used, other chemotherapeutic agents may be administered before, concurrently, or after administration of a compound of the present invention. In one aspect of the invention, a compound of the present invention is co-administered with one or more other chemotherapeutic agents. By "co-administer" it is meant that the present compounds are administered to a patient such that the present compounds as well as the co-administered

compound may be found in the patient's bloodstream at the same time, regardless when the compounds are actually administered, including simultaneously.

[00197] The administration of the compounds and compositions of the present invention can be done in a variety of ways, including, but not limited to, orally, subcutaneously, intravenously, intranasally, transdermally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the compound or composition may be directly applied as a solution or spray.

[00198] Pharmaceutical dosage forms include a compound of formula I or a pharmaceutically acceptable salt or solvate thereof, and one or more pharmaceutical excipients. As is known in the art, pharmaceutical excipients are secondary ingredients which function to enable or enhance the delivery of a drug or medicine in a variety of dosage forms (e.g.: oral forms such as tablets, capsules, and liquids; topical forms such as dermal, ophthalmic, and otic forms; suppositories; injectables; respiratory forms and the like). Pharmaceutical excipients include inert or inactive ingredients, synergists or chemicals that substantively contribute to the medicinal effects of the active ingredient. For example, pharmaceutical excipients may function to improve flow characteristics, product uniformity, stability, taste, or appearance, to ease handling and administration of dose, for convenience of use, or to control bioavailability. While pharmaceutical excipients are commonly described as being inert or inactive, it is appreciated in the art that there is a relationship between the properties of the pharmaceutical excipients and the dosage forms containing them.

[00199] Pharmaceutical excipients suitable for use as carriers or diluents are well known in the art, and may be used in a variety of formulations. See, e.g., Remington's Pharmaceutical Sciences, 18th Edition, A. R. Gennaro, Editor, Mack Publishing Company (1990); Remington: The Science and Practice of Pharmacy, 20th Edition, A. R. Gennaro, Editor, Lippincott Williams & Wilkins (2000); Handbook of Pharmaceutical Excipients, 3rd Edition, A. H. Kibbe, Editor, American Pharmaceutical Association, and Pharmaceutical Press (2000); and Handbook of Pharmaceutical Additives, compiled by Michael and Irene Ash, Gower (1995), each of which is incorporated herein by reference for all purposes.

[00200] Oral solid dosage forms such as tablets will typically comprise one or more pharmaceutical excipients, which may for example help impart satisfactory processing and compression characteristics, or provide additional desirable physical characteristics to the tablet. Such pharmaceutical excipients may be selected from diluents, binders, glidants,

lubricants, disintegrants, colors, flavors, sweetening agents, polymers, waxes or other solubility-retarding materials.

[00201] Compositions for intravenous administration will generally comprise intravenous fluids, i.e., sterile solutions of simple chemicals such as sugars, amino acids or electrolytes, which can be easily carried by the circulatory system and assimilated. Such fluids are prepared with water for injection USP.

[00202] Fluids used commonly for intravenous (IV) use are disclosed in Remington, the Science and Practice of Pharmacy [full citation previously provided], and include:

alcohol (e.g., in dextrose and water ("D/W") [e.g., 5% dextrose] or dextrose and water [e.g., 5% dextrose] in normal saline solution ("NSS"); e.g. 5% alcohol);

synthetic amino acid such as Aminosyn, FreAmine, Travasol, e.g., 3.5 or 7; 8.5; 3.5, 5.5 or 8.5 % respectively;

ammonium chloride e.g., 2.14%;

dextran 40, in NSS e.g., 10% or in D5/W e.g., 10%;

dextran 70, in NSS e.g., 6% or in D5/W e.g., 6%;

dextrose (glucose, D5/W) e.g., 2.5-50%;

dextrose and sodium chloride e.g., 5-20% dextrose and 0.22-0.9% NaCl;

lactated Ringer's (Hartmann's) e.g., NaCl 0.6%, KCl 0.03%, CaCl_2 0.02%;

lactate 0.3%;

mannitol e.g., 5%, optionally in combination with dextrose e.g., 10% or NaCl e.g., 15 or 20%;

multiple electrolyte solutions with varying combinations of electrolytes, dextrose, fructose, invert sugar Ringer's e.g., NaCl 0.86%, KCl 0.03%, CaCl_2 0.033%;

sodium bicarbonate e.g., 5%;

sodium chloride e.g., 0.45, 0.9, 3, or 5%;

sodium lactate e.g., 1/6 M; and

sterile water for injection

The pH of such fluids may vary, and will typically be from 3.5 to 8 such as known in the art.

[00203] The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. All


publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.


EXAMPLES

[00204] All anhydrous solvents were purchased from Aldrich Chemical Company in SureSeal® containers. Reagents were added and aqueous extractions performed with single or multichannel pipettors. Filtrations were performed using Whatman/Polyfiltronics 24 well, 10 mL filtration blocks. Evaporation of volatile materials from the array was performed with a Labconco Vortex-Evaporator or by sweeping with a 4 x 6 nitrogen manifold.

Example 1

Synthesis of Compounds





a) 3-Phenyl-propionic acid 3-chloro-phenyl ester

[00205] To a solution of 3-chlorophenol (1, 50.1 g, 0.3 mol), triethylamine (TEA, 85 mL), and CH_2Cl_2 (500 mL) at 23°C was added hydrocinnamoyl chloride (2, 31 mL, 0.3 mol) over 5 minutes. After 30 minutes, the reaction mixture was concentrated in vacuo. The crude slurry was then dissolved in 10:1 hexanes:EtOAc (300 mL) and washed with 1 N NaOH (100 mL) and brine (100 mL). The organic layer was dried (MgSO_4) and filtered through a plug of silica gel (10:1 hexanes:EtOAc rinse). The eluent was concentrated to afford 78 g of a slightly yellow oil, which was used without any further purification.

b) 1-(4-Chloro-2-hydroxy-phenyl)-3-phenyl-propan-1-one

[00206] AlCl_3 (52 g, 0.39 mol) was added slowly over 15 minutes to ester 3 (78 g, 0.3 mol) at 140°C . After an additional 15 minutes gas evolution ceased and the reaction mixture was poured into a 1L beaker and allowed to cool to RT. The resulting solid was dissolved in CH_2Cl_2 (100 mL) and slowly quenched with 1 N HCl (200 mL). This mixture was diluted with EtOAc (600 mL) and the layers were separated. The organic layer was washed with brine (100 mL), dried (MgSO_4), filtered, and concentrated. The crude oil was dissolved in 20:1 hexanes:EtOAc (500 mL) and flushed through a plug of silica gel (100% hexanes; 20:1 hexanes:EtOAc rinse). The filtrate was concentrated to provide a slightly brown oil, which was purified by flash column chromatography (50:1 hexanes:EtOAc; 40:1 hexanes:EtOAc; 30:1 hexanes:EtOAc; 20:1 hexanes:EtOAc) to yield 31.2 g (40%) of 4 as a white solid.

c) 2-tert-Butoxycarbonylamino-3-methyl-butyric acid 5-chloro-2-(3-phenyl-propionyl)-phenyl ester

[00207] A solution of phenol 4 (23.18g, 89.1 mmol), BOC-D-Valine (5, 21.29 g, 98.05

mmol), HBTU (40.57 g, 107 mmol), TEA (37 mL, 265 mmol), and CH_2Cl_2 (155 mL) were maintained at 23°C for 5 hours. The reaction mixture was diluted with EtOAc (500 mL) and washed with saturated aqueous NH_4Cl (2 x 100 mL) and brine (100 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (100% hexanes; 50:1 hexanes:EtOAc; 40:1 hexanes:EtOAc; 30:1 hexanes:EtOAc; 20:1 hexanes:EtOAc; 10:1 hexanes:EtOAc) to yield 35.7 g (87%) of **6** as a yellow oil.

d) [1-(3-Benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-carbamic acid tert-butyl ester

[00208] A mixture of ester **6** (35.55 g, 79.17 mmol), K_2CO_3 (21.8 g, 158.4 mmol), and DMF (264 mL) was placed into a 140°C oil bath. After 30 mins, the reaction mixture was quenched with H_2O (300 mL) and extracted with Et_2O (3 x 200 mL). The organic layer was washed with brine (100 mL), dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 3.5 g (10.2%) of **7**.

e) 2-(1-Amino-2-methyl-propyl)-3-benzyl-7-chloro-chromen-4-one

[00209] Chromenone **7** (1.8 g, 4.18 mmol) and TFA: H_2O (97.5:2.5, 30 mL) was maintained at 23°C for 1h. The reaction mixture was concentrated. The residue was dissolved in EtOAc (100 mL) and washed with 1 N NaOH (25 mL) and brine (25 mL). The organic layer was dried (MgSO_4), filtered, and concentrated to provide a colorless oil, which was used without further purification.

f) N-{2-[1-(3-Benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propylamino]-1,1-dimethyl-ethyl}-4-methyl-benzamide


[00210] Chromenone **8** (72 mg, 0.22 mmol), aldehyde **9** (65 mg, 0.32 mmol), $\text{Na(OAc)}_3\text{BH}$ (184 mg, 0.87 mmol), and CH_2Cl_2 (1 mL) was maintained at 23°C for 3 h. The reaction mixture was diluted with EtOAc (20 mL) and washed with 1 N NaOH (5 mL) and brine (5 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (3:1 hexanes:EtOAc) to yield 80 mg (70%) of **10** as a white solid.

g) 3-Benzyl-7-chloro-2-[1-(4,4-dimethyl-2-p-tolyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-chromen-4-one

[00211] Chromenone **10** (80 mg, 0.15 mmol), POCl₃ (0.1 mL, 1.1 mmol), and PhMe (1 mL) were heated to 110°C. After 3.5h, an additional portion of POCl₃ (0.1 mL, 1.1 mmol) was added. After 1h, the reaction mixture was diluted with EtOAc (20 mL) and washed with 1 N NaOH (10 mL) and brine (10 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (20:1 CHCl₃:MeOH) to yield 50 mg (65%) of **11** as a white solid.

Example 2

Synthesis of Compounds



a) Preparation of {3-[1-(3-Benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propylamino]-propyl}-carbamic acid tert-butyl ester

[00212] Chromenone **8** (420 mg, 1.24 mmol), aldehyde **12** (280 mg, 1.6 mmol), NaCN(OAc)₃BH (790 mg, 3.7 mmol), and CH₂Cl₂ (4.1 mL) was maintained at 23°C for 3 h. The reaction mixture was diluted with EtOAc (20 mL) and washed with 1 N NaOH (5 mL)

and brine (5 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (5:1 hexanes:EtOAc; 3:1 hexanes:EtOAc) to yield 460 mg (75%) of 13 as a viscous oil.


b) Preparation of {3-[[1-(3-Benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-(4-methyl-benzoyl)-amino]-propyl}-carbamic acid tert-butyl ester

[00213] To a solution of chromenone 13 (1.3 g, 2.6 mmol), diisopropylethylamine (DIEA, 1.8 mL), and CH₂Cl₂ (7.5 mL) at 23°C was added *p*-toluoyl chloride (0.7 mL, 5.22 mmol). After 2.5 h, the reaction mixture was diluted with EtOAc (100 mL) and washed with saturated aqueous NaHCO₃ (2 x 20 mL) and brine (20 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (3:1 hexanes:EtOAc) to yield 1.43 g (89%) of 14 as a colorless oil.

c) Preparation of N-(3-Amino-propyl)-N-[1-(3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-methyl-benzamide


[00214] Chromenone 14 (1.43 g, 2.32 mmol) and TFA:H₂O (97.5:2.5, 30 mL) was maintained at 23°C for 1h. The reaction mixture was concentrated. The residue was dissolved in EtOAc (100 mL) and washed with 1 N NaOH (25 mL) and brine (25 mL). The organic layer was dried (MgSO₄), filtered, and concentrated to provide a white solid which was deemed >95% pure by ¹H NMR and LCMS analysis.

[00215] Using procedures analogous to those set forth above in Example 2, the following compound was prepared.

Structure	LRMS (MH) <i>m/z</i>
	471.2

Example 3

Synthesis of compounds



[00216] Carbonyldiimidazole (9.14 g, 56.37 mmol) was added slowly to a r.t. solution of BOC-D-Valine (5, 12.25 g, 56.37 mmol) and THF (185 mL). After 1 h, the solution was washed with 50% aqueous NaCl (200 mL), followed by brine (2 x 200 mL). The organic layer was dried (MgSO_4), filtered, and concentrated to provide a white solid, which was used without further purification.


[00217] (2-Bromoethyl)benzene (2.51 mL, 18.38 mmol), magnesium turnings (477 mg, 19.62 mmol), and THF (20 mL) were heated to 60°C for 1 hour, then allowed to cool to r.t. A solution of 16 (2.0 g, 9.19 mmol) and THF (20 mL) was cooled to $0-5^\circ\text{C}$. The solution of the phenethyl magnesium chloride was then added via syringe to the $0-5^\circ\text{C}$ solution of the valine

imidazole. The temperature was monitored by internal thermometer and was not allowed to exceed 15°C. The reaction mixture was maintained at 0-5°C for 1 hour. The reaction mixture was quenched with saturated aqueous NH₄Cl (20 mL), and diluted with EtOAc (100 mL). The layers were separated and the organic layer was washed with brine (30 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc; 5:1 hexanes:EtOAc) to yield 1.15 g (41%) of 17. LRMS (MH-*t*BuOCO) *m/z* 206.1.

[00218] LHMDS (1.M in THF, 9.53 mL, 3.3 equiv) was added slowly via syringe to a -78°C solution of ketone 17 (882 mg, 2.89 mmol). After the addition was complete the resulting light orange solution was maintained at -78°C for 40 mins. Neat 4-chloro-2-fluorobenzoyl chloride (18, 460 µL, 2.89 mmol) (assumed density to be 1.20 g/mL) was added dropwise via syringe. The reaction solution turned to an orange color and was maintained for 40 mins. The reaction solution was quenched with saturated aqueous NH₄Cl (20 mL), and diluted with EtOAc (50 mL). The layers were separated and the organic layer was washed with brine (30 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 1.15 g (86%) of 19. LRMS (MH-HF) *m/z* 442.1.


[00219] A mixture of 19 (1.15 g, 2.49 mmol), K₂CO₃ (420 mg, 3.04 mmol), and DMF (12 mL) was maintained at r.t. for 30 mins. The yellow reaction solution was quenched with brine (50 mL), and diluted with Et₂O (50 mL). The layers were separated and the organic layer was washed with brine (2 x 50 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 20. LRMS (MH) *m/z* 442.1.

[00220] Using methods analogous to those described above in Examples 2 and 3, the compounds shown below were prepared.



R ₇	R	R'	Stereochemistry (R/S)
Cl	H	H	3/1
Cl	-CH ₂ CMe ₂ N=C(<i>p</i> -tol)-		3/1
Cl	H ₂ N(CH ₂) ₃ -	<i>p</i> -toluoyl	3/1
Cl	H ₂ N(CH ₂) ₃ -	MeOCH ₂ C(O)-	3/1
Cl	H ₂ N(CH ₂) ₃ -	<i>p</i> -Br-phenyl-	3/1
Cl	H ₂ N(CH ₂) ₃ -	<i>p</i> -toluoyl	R
Cl	H ₂ N(CH ₂) ₃ -	<i>p</i> -toluoyl	S
Cl	-CH ₂ CMe ₂ N=C(<i>p</i> -tol)-		R
Cl	-CH ₂ CMe ₂ N=C(<i>p</i> -tol)-		S
F	H	H	3/1
F	H ₂ N(CH ₂) ₃ -	<i>p</i> -toluoyl	3/1
F	H ₂ N(CH ₂) ₃ -	MeOCH ₂ C(O)-	3/1
F	-CH ₂ CMe ₂ N=C(<i>p</i> -tol)-		3/1

Example 4**Alternative Methods for Preparation of Compounds**



[00221] Ethylchloroformate (11.0 mL, 115 mmol) was added over 1 minute to a 0-5°C solution of BOC-D-Valine (5, 25.0 g, 115 mmol), triethylamine (16.0 mL, 115 mmol), and THF (145 mL) under N₂. The internal temperature of the reaction solution rose to 9°C. After 15 mins, a mixture of dimethylhydroxylamine hydrochloride (13.46 g, 138 mmol), triethylamine (32.0 mL, 230 mmol), and THF (110 mL) was added over 5 minutes. The internal temperature rose to 17°C. Upon completion of addition, the ice/H₂O bath was removed and the reaction solution maintained at 23°C for 1 hour. The reaction solution was then concentrated. The crude residue was dissolved in EtOAc (200 mL) and washed with 1 N HCl (200 mL) and brine (100 mL). The organic layer was dried (MgSO₄), filtered, and concentrated to provide 30 g (~100%) of 21 as a colorless oil, which was used without further purification.


[00222] (2-Bromoethyl)benzene (38.0 mL, 273 mmol), magnesium turnings (7.0 g, 289 mmol), and Et₂O (500 mL) were mixed in a 1L round-bottom flask equipped with a reflux condenser at 23°C under a N₂ atmosphere. After ~10 mins the reaction mixture begins to exotherm and the reaction mixture was allowed to progress to reflux with intermittent cooling with an ice/H₂O bath. After 1.5 hour, the Grignard reaction was complete and the solution had cooled to 23°C. A solution of 21 (18.0 g, 82.7 mmol) and Et₂O (200 mL) was added via cannula to the 20°C solution of the phenethylmagnesium bromide. The temperature was

monitored by internal thermometer and was not allowed to exceed -30°C . The reaction mixture temperature was monitored by an internal thermometer and regulated ($20-30^{\circ}\text{C}$) with an ice/ H_2O bath. After 1 h at 23°C , the reaction mixture was quenched by pouring into 1 N HCl (300 mL). The layers were separated and the organic layer was washed with brine (100 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 13.4 g (53%) of **17**. LRMS (MH-*t*BuOCO) m/z 206.1.

[00223] Lithium bis(trimethylsilyl)amide (LHMDS, 1.0 M in THF, 94.0 mL, 3.3 equiv) was added slowly over ~ 3 minutes via syringe to a -78°C solution of ketone **17** (8.74 g, 28.62 mmol) and THF (100 mL). The reaction solution temperature was monitored by an internal thermometer, and addition of the base was done at a rate sufficient to prevent the temperature from exceeding -54°C . After the addition was complete the resulting solution was maintained at -78°C for 30 mins. Neat 4-chloro-2-fluorobenzoyl chloride (**18**, 4.58 mL, 28.62 mmol) (assumed density to be 1.20 g/mL) was added drop-wise over ~ 1 minute via syringe (temperature rose from -78°C to -59°C). The reaction solution was maintained at -78°C for 30 mins. The reaction solution was quenched with 1 N HCl (100 mL). The layers were separated and the organic layer was washed with brine (100 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was used without further purification.

[00224] A mixture of the above crude product, K_2CO_3 (4.75 g, 34.34 mmol) and DMF (100 mL) was maintained at 23°C for 30 mins. The reaction mixture was quenched by addition of Et_2O (200 mL) and brine (200 mL). The layers were separated and the organic layer was washed with brine (2 x 200 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue absorbed onto silica gel (CH_2Cl_2) and was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 11.5 g (91% for 2 steps) of **20** as a white solid. LRMS (MH) m/z 442.1.

Example 5



[00225] Methylmagnesium chloride (82.5 mL, 3.0 M in THF, 247 mmol) was added over 5 mins to a 0-5°C solution of 21 (17.95 g, 69.0 mmol) and THF (200 mL) under an N₂ atmosphere. The reaction mixture temperature was monitored by an internal thermometer, and addition of the Grignard was done at a rate sufficient to prevent the temperature from exceeding 19°C. Upon complete addition the cooling bath was removed and the reaction mixture was maintained at 23°C for 2 hours. The reaction mixture was then quenched with 1 N HCl (100 mL). The layers were separated and the organic layer was washed with brine (100 mL). The organic layer was dried (MgSO₄), filtered, and concentrated to provide 12.5 g (84%) of 22 as a white solid (>95% pure by ¹H NMR), which was used without further purification.

[00226] LHMDS (128 mL, 1.0 M in THF) was added via syringe over 3 minutes to a -78°C solution of ketone 22 (12.5 g, 58.1 mmol) and THF (200 mL). The reaction solution temperature was monitored by an internal thermometer, and addition of the base was done at a rate sufficient to prevent the temperature from exceeding -58°C. After 30 minutes, a solution of α -bromo-*m*-tolunitrile (23, 12.5 g, 63.9 mmol) and THF (50 mL) was added over 30 seconds (temperature rose from -78 to -60°C). The cooling bath was immediately replaced by a ice/H₂O bath and the reaction solution was maintained at ~0°C for 20 minutes. The reaction solution was quenched with 1 N HCl (100 mL). The layers were separated and the organic layer was washed with brine (100 mL). The organic layer was dried (MgSO₄),


filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 11.6 g (60%) of **24**. Inspection of the ^1H NMR revealed that **24** was only ~80% pure, but no further purification was performed. LRMS (MH-*t*BuOCO) m/z 231.1.

[00227] Lithium bis(trimethylsilyl)amide (LHMDS, 1.0 M in THF, 28.4 mL, 3.3 equiv) was added slowly via syringe to a -78°C solution of ketone **24** (2.84 g, 8.6 mmol) and THF (40 mL). The reaction solution temperature was monitored by an internal thermometer, and addition of the base was done at a rate sufficient to prevent the temperature from exceeding -48°C . After the addition was complete the resulting solution was maintained at -78°C for 30 mins. Neat 4-chloro-2-fluorobenzoyl chloride (**18**, 1.38 mL, 8.6 mmol) (assumed density to be 1.20 g/mL) was added dropwise via syringe. The reaction solution turned to an orange color and was maintained for 30 mins. The reaction solution was quenched with 1 N HCl (20 mL). The layers were separated and the organic layer was washed with brine (20 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was used without further purification.

[00228] A mixture of the above crude product, K_2CO_3 (1.43 g, 10.34 mmol) and DMF (43 mL) was maintained at 23°C for 1 hour. The reaction mixture was quenched by addition of Et_2O (100 mL) and brine (200 mL). The layers were separated and the organic layer was washed with brine (2 x 200 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue absorbed onto silica gel (CH_2Cl_2) and was purified by flash column chromatography (5:1 hexanes:EtOAc) to yield 2.7 g (67% for 2 steps) of **25**. Inspection of the ^1H NMR revealed that **25** was only ~80% pure, but no further purification was performed. LRMS (MH) m/z 467.1.

[00229] Chromenone **25** (2.71 g, 5.80 mmol) and TFA: H_2O (97.5:2.5, 25 mL) was maintained at 23°C for 1h. The reaction mixture was concentrated. The residue was dissolved in EtOAc (100 mL) and washed with 1 N NaOH (25 mL) and brine (25 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (1:1 hexanes:EtOAc; 1:2 hexanes:EtOAc; 1:4 hexanes:EtOAc) to yield 1.20 g (56%) of **26**. LRMS (MH) m/z 367.1.


Example 6



[00230] Lithium bis(trimethylsilyl)amide (LHMDS, 1.0 M in THF, 23.5 mL, 3.3 equiv) was added slowly via syringe to a -78°C solution of ketone 17 (2.18 g, 7.14 mmol) and THF (20 mL). The reaction solution temperature was monitored by an internal thermometer, and addition of the base was done at a rate sufficient to prevent the temperature from exceeding -50°C. After the addition was complete the resulting solution was maintained at -78°C for 30 mins. Neat 2,4-difluorobenzoyl chloride (27, 1.05 mL, 8.57 mmol) was added dropwise via syringe. The reaction solution turned to an orange color and was maintained for 30 mins. The reaction solution was quenched with 1 N HCl (20 mL). The layers were separated and the organic layer was washed with brine (20 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 2.16 g (68%) of 28. Inspection of the ¹H NMR revealed that 28 was only ~85% pure, but no further purification was performed.

[00231] A mixture of 28 (2.16 g, 4.9 mmol), K₂CO₃ (812 mg, 5.86 mmol), and DMF (24 mL) was maintained at 23°C for 1 hour. The reaction mixture was quenched with brine (50 mL), and diluted with Et₂O (50 mL). The layers were separated and the organic layer was washed with brine (2 x 50 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 1.25 g (60%) of 29. LRMS (MH) *m/z* 426.2.


Example 7



[00232] Lithium bis(trimethylsilyl)amide (LHMDS, 1.0 M in THF, 47 mL, 3.3 equiv) was added slowly via syringe to a -78°C solution of ketone 17 (4.37 g, 14.31 mmol) and THF (55 mL). The reaction solution temperature was monitored by an internal thermometer, and addition of the base was done at a rate sufficient to prevent the temperature from exceeding -50°C. After the addition was complete the resulting solution was maintained at -78°C for 30 mins. A solution of 4-cyano-2-fluorobenzoyl chloride (30, 1.38 mL, 8.6 mmol) and THF (5 mL) was added rapidly via syringe (temperature rose from -78 to -50°C). The reaction solution turned to a dark red color and was maintained for 30 mins. The reaction solution was quenched with 1 N HCl (20 mL). The layers were separated and the organic layer was washed with brine (20 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was used without further purification.


[00233] A mixture of the above crude product, K₂CO₃ (2.40 g, 17.17 mmol) and DMF (70 mL) was maintained at 23°C of 1 hour. The reaction mixture was quenched by addition of Et₂O (200 mL) and brine (200 mL). The layers were separated and the organic layer was washed with brine (2 x 200 mL). The organic layer was dried (MgSO₄), filtered, and concentrated. The resulting residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to yield 4.95 g (80% for 2 steps) of 31. LRMS (MH) *m/z* 433.2.

Example 8



[00234] 7-Fluoro-Chromenone 32 (96 mg, 0.19 mmol) was dissolved in 0.5 M sodium methoxide in methanol (10 mL) and heated to 70°C. The temperature was maintained at 70°C for 12 hours and then cooled to room temperature. The solvent was removed under reduced pressure. To the remaining residue was added EtOAc (20 mL) and water (20 mL). The layers were separated and the aqueous phase was extracted with additional EtOAc (3 x 15 mL). The organic phases were combined, washed with brine (25 mL) and dried (Na₂SO₄). Concentration *in vacuo* gave an amorphous white solid which was purified using flash column chromatography (20:1 DCM:MeOH) to provide 92 mg (94%) of 33. LRMS (MH) *m/z* 509.3.


Example 9



[00235] Compound 29 (608 mg, 1.43 mmol) was dissolved into a solution of 0.5 M sodium methoxide in methanol (40 mL). The mixture was heated to reflux for 3 hours (or until LC-MS indicated no starting material left). Then the solvent was evaporated. The residue was dissolved in dichloromethane (200 mL) and water (200 mL) was added, in which pH of the mixed solution was

adjusted to 9 by adding 2N HCl solution. The collected organic layers were dried by sodium sulfate. After evaporation of solvents, the final compound 34 (580 mg, 92%) was dried under vacuum.

Example 10



[00236] To a solution of 32 (56 mg, 0.11 mmol) in DMF (5 mL) was added NaH (5 mg of 60% dispersion in mineral oil, 0.15 mmol). The resulting solution was stirred at 45°C for 5 mins., then allyl alcohol was added (10 μ L, 0.15 mmol) via pipette. The resulting solution was stirred at 45°C for 12 hours and then cooled to room temperature. EtOAc (50 mL) and water (15 mL) were added and the layers were separated. The aqueous phase was extracted with EtOAc (3 x 15 mL). The organic phases were combined, washed with water (2 x 30 mL) and brine (2 x 30 mL), dried (Na_2SO_4) and concentrated under reduced pressure to yield 35 as an amorphous solid, which was used without further purification (54 mg, 91%). LRMS (MH) m/z 535.0.

[00237] To a room temperature solution of 35 (54 mg, 0.10 mmol) and MeCN (5 mL) was added morpholine (44 μ L, 0.50 mmol), followed by $\text{Pd(PPh}_3)_4$ (5 mg, 10%). The

resulting solution was stirred for 5 minutes and then concentrated under reduced pressure. The remaining residue was purified by flash column chromatography using stepwise elution (20:1 DCM:MeOH, 10:1 DCM:MeOH) to provide 41 mg of 36 as an off-white solid (81%). LRMS (MH) m/z 495.2.

Example 11

Preparation of 3-Benzyl-7-chloro-2-[2-methyl-1-(2-*p*-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-chromen-4-one

a) {2-[1-(3-Benzyl-7-chloro-4-oxo-4*H*-chromen-2-yl)-2-methyl-propylamino]-ethyl}-carbamic acid *tert*-butyl ester

[00238] To a solution of 2-(1-amino-2-methyl-propyl)-3-benzyl-7-chloro-chromen-4-one (6.0 g, 18 mmol) and (3-oxoethyl)carbamic acid *tert*-butyl ester (3.6 g, 23 mmol) in methylene chloride (150 mL) was added sodium triacetoxyborohydride (7.4 g, 35 mmol). The reaction mixture was stirred at room temperature for 16 hours, at which time it was diluted with 1 N sodium hydroxide (150 mL) and stirred vigorously for 2 hours. The organic layer was washed with 1 N sodium hydroxide (100 mL) and brine (100 mL), dried over magnesium sulfate, and concentrated. The residue was purified by flash chromatography (0→50% ethyl acetate/hexanes) to give 5.7 g (67%) of the title compound. MS(ES+) m/e 485 [M+H]⁺.

b) 2-[1-(2-Amino-ethylamino)-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one

[00239] A solution of {2-[1-(3-benzyl-7-chloro-4-oxo-4*H*-chromen-2-yl)-2-methyl-propylamino]-ethyl}-carbamic acid *tert*-butyl ester (5.7 g, 12 mmol) in 4:1 methylene chloride/trifluoroacetic acid (250 mL) was maintained at room temperature for 1.5 hours, at which time it was concentrated. The residue was dissolved in methylene chloride (200 mL), washed with 10% sodium carbonate, saturated sodium bicarbonate, and brine, dried over magnesium sulfate, and concentrated. The residue (4.2 g, 91% yield) was used without further purification.

c) *N*-{2-[1-(3-Benzyl-7-chloro-4-oxo-4*H*-chromen-2-yl)-2-methyl-propylamino]-ethyl}-4-methyl-benzamide


[00240] To a cooled (0°C) solution of 2-[1-(2-amino-ethylamino)-2-methyl-propyl]-3-

benzyl-7-chloro-chromen-4-one (4.1 g, 11 mmol) and triethylamine (2.2 mL, 17 mmol) in methylene chloride (100 mL) was added a solution of *p*-toluyl chloride (1.7 g, 11 mmol) in methylene chloride (20 mL). The reaction was maintained at 0 °C for 2 hours, at which time it was diluted with ether (250 mL). The resultant solution was washed with 1 N hydrogen chloride (2 x 200 mL), saturated sodium bicarbonate (200 mL) and brine (150 mL), dried over magnesium sulfate, and concentrated. The residue was purified by flash chromatography (0→60% ethyl acetate/hexanes) to give 2.8 g (50%) of the title compound. MS(ES+) *m/e* 503 [M+H]⁺.

d) 3-Benzyl-7-chloro-2-[2-methyl-1-(2-*p*-tolyl-4,5-dihydro-imidazol-1-yl)-propyl]-chromen-4-one

[00241] A mixture of *N*-{2-[1-(3-benzyl-7-chloro-4-oxo-4*H*-chromen-2-yl)-2-methyl-propylamino]-ethyl}-4-methyl-benzamide (2.8 g, 5.5 mmol) and phosphorus oxychloride (9.4 mL, 100 mmol) in toluene (60 mL) was heated at 85 °C for 7 hours, then heated at reflux for 1 hour. The reaction was concentrated and the residual phosphorus oxychloride removed by toluene azeotrope. The residue was diluted with ethyl acetate (100 mL) and washed with saturated sodium bicarbonate (100 mL) and brine (100 mL), dried over magnesium sulfate, and concentrated. The resultant residue was purified by flash chromatography (0→8% methanol/methylene chloride) to give 1.4 g (50%) of the title compound. MS(ES+) *m/e* 485 [M+H]⁺.

[00242] Following procedures analogous to those set forth herein in Examples 1, 3, and/or 4, the following compounds were prepared:



R ₇	R ₁	R ₉	[M+H] ⁺
----------------	----------------	----------------	--------------------

CN	3-MeO-Ph- CH ₂ -	3,4-piperonyl	536
Cl	3-MeO-Ph- CH ₂ -	4-Me-Ph	515
Cl	3-MeO-Ph- CH ₂ -	3-F-4-Me-Ph	533
Cl	3-MeO-Ph- CH ₂ -	3,4-piperonyl	545
F	3-MeO-Ph- CH ₂ -	4-Me-Ph	499
F	3-MeO-Ph- CH ₂ -	3-F-4-Me-Ph	517

Example 12**Preparation of 2-[(R)-1-(4-Aminomethyl-2-p-tolyl-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one**

a) 2-[(R)-1-(3-Phthalimido-2-oxo-propylamino)-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one

[00243] To 2-((R)-1-amino-2-methyl-propyl)-3-benzyl-7-chloro-chromen-4-one (0.5 g, 1.5 mMol) and K₂CO₃ (0.21 g, 1.5 mMol) in DMF (10 mL) was added N-(3-bromo-2-oxopropyl)-phthalimide (0.45 g, 1.5 mMol) (Nair et al.; J. Org. Chem.; 40; 1975; 1745). The reaction was stirred at RT for 3 h, concentrated under vacuum, taken up in EtOAc, washed with water, brine, dried (Na₂SO₄) and evaporated to give the title compound (0.88 g, 100%) as a yellow solid: MS (ES) m/e 543.2 (M + H)⁺.

b) 2-[(R)-1-[N-Toluoyl-(3-phthalimido-2-oxo-propyl)amino]-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one

[00244] To 2-[(R)-1-(3-phthalimido-2-oxo-propylamino)-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one (0.88 g, 1.5 mMol) in CH₂Cl₂ (10 mL) was added Et₃N (0.23 mL, 1.6 mMol) and toluoyl chloride (0.21 mL, 1.6 mMol). The reaction was stirred at RT for 18 h,

WO 03/

concentrated under vacuum, taken up in EtOAc, washed with 1 N HCl, brine, dried (MgSO₄) and evaporated to dryness. Purification by flash chromatography on silica gel (5-15% EtOAc/hexane) followed by trituration with pet. ether, filtration and drying under vacuum gave the title compound (0.89 g, 90%) as a white solid: MS (ES) m/e 661.2 (M + H)⁺.

c) N-{1-[(R)-1-(3-Benzyl-7-chloro-4-oxo-chromen-2-yl)-2-methyl-propyl]-2-p-tolyl-1H-imidazol-4-ylmethyl}-phthalimide


[00245] To 2-{(R)-1-[N-toluoyl-(3-phthalimido-2-oxo-propyl)amino]-2-methyl-propyl}-3-benzyl-7-chloro-chromen-4-one (0.88 g, 1.3 mMol) and NH₄OAc (5.13 g, 66.8 mMol) was added HOAc (15 mL). The reaction was stirred and heated to reflux (155 °C oil bath) for 2.5 h, cooled to RT and concentrated under vacuum. The remaining residue was taken up in EtOAc, washed with water, brine, dried (MgSO₄) and evaporated to dryness. Purification by flash chromatography on silica gel (50% EtOAc/hexane) followed by trituration with (1:2) Et₂O/ pet. ether, filtration and drying under vacuum gave the title compound (0.51 g, 61%) as a tan solid: ¹H NMR (400 MHz, CDCl₃) δ 8.11 (d, J = 8.6 Hz, 1 H), 7.89 (2d, 2 H), 7.73 (2d, 2 H), 7.53 (d, J = 1.8 Hz, 1 H), 7.52 (s, 1 H), 7.39 (d, J = 8.0 Hz, 2 H), 7.36 (dd, 1 H), 7.32 (d, J = 8.0 Hz, 2 H), 7.14 (m, 3 H), 6.74 (d, J = 1.5 Hz, 1 H), 6.72 (s, 1 H), 5.04 (d, J = 8.8 Hz, 1 H), 4.94 (s, 2 H), 4.01 (d, J = 15.5 Hz, 1 H), 2.63 (m, 1 H), 2.46 (s, 3 H), 2.44 (d, J = 15.5 Hz, 1 H), 0.95 (d, J = 6.6 Hz, 3 H), 0.29 (d, J = 6.7 Hz, 3 H); MS (ES) m/e 642.0 (M + H)⁺.

d) 2-[(R)-1-(4-Aminomethyl-2-p-tolyl-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one

[00246] To N-{1-[(R)-1-(3-benzyl-7-chloro-4-oxo-3,4-dihydro-chromen-2-yl)-2-methyl-propyl]-2-p-tolyl-1H-imidazol-4-ylmethyl}-phthalimide (0.50 g, 0.78 mMol) in EtOH (15 mL) was added hydrazine monohydrate (0.12 mL, 2.5 mMol). The reaction was stirred at RT for 72 h, filtered through a pad of Celite[®] to remove the insoluble precipitate, rinsed with EtOH and evaporated to dryness. Purification by flash chromatography on silica gel [5 - 10% (5% NH₄OH in MeOH)/CH₂Cl₂] gave the title compound (364 mg, 91%) as a white solid foam: ¹H NMR (400 MHz, CDCl₃) δ 8.14 (d, J = 8.6 Hz, 1 H), 7.63 (d, J = 1.8 Hz, 1 H), 7.37 (m, 6 H), 7.15 (m, 3 H), 6.71 (d, J = 1.7 Hz, 1 H), 6.69 (s, 1 H), 5.00 (d, J = 10.9 Hz, 1

H), 4.07 (d, $J = 15.5$ Hz, 1 H), 3.92 (d, 1 H), 3.88 (d, 1 H), 2.72 (br s, 2 H), 2.65 (m, 1 H), 2.49 (s, 3 H), 2.48 (d, $J = 15.5$ Hz, 1 H), 0.93 (d, $J = 6.6$ Hz, 3 H), 0.30 (d, $J = 6.7$ Hz, 3 H); MS (ES) m/e 512.2 ($M + H$)⁺.


[00247] Following procedures analogous to those set forth above, the following compounds were prepared:



R ₇	R ₁	R ₉	[M+H] ⁺
F	Ph-CH ₂ -	4-Me-Ph	510
F	Ph-CH ₂ -	3-F-4-Me-Ph	528
CN	Ph-CH ₂ -	3-F-4-Me-Ph	535
Cl	Ph-CH ₂ -	3-F-4-Me-Ph	544
Cl	3-MeO-Ph-CH ₂ -	4-Me-Ph	556
Cl	3-MeO-Ph-CH ₂ -	3-F-4-Me-Ph	574
CN	Ph-CH ₂ -	4-Me-Ph	517
Cl	3-CN-Ph-CH ₂ -	4-Me-Ph	551
Cl	3-CN-Ph-CH ₂ -	3-F-4-Me-Ph	569
F	3-MeO-Ph-CH ₂ -	4-Me-Ph	540
CN	3-MeO-Ph-CH ₂ -	4-Me-Ph	547
CN	3-CN-Ph-CH ₂ -	3-F-4-Me-Ph	560
F	3-MeO-Ph-CH ₂ -	3-F-4-Me-Ph	588
F	3-CN-Ph-CH ₂ -	3-F-4-Me-Ph	553

Example 13**Preparation of N-{1-[(R)-1-(3-Benzyl-7-chloro-4-oxo-3,4-dihydro-chromen-2-yl)-2-methyl-propyl]-2-p-tolyl-1H-imidazol-4-ylmethyl}-acetamide**

[00248] To 2-[(R)-1-(4-aminomethyl-2-p-tolyl-imidazol-1-yl)-2-methyl-propyl]-3-benzyl-7-chloro-chromen-4-one (0.17 g, 0.33 mMol) in CH_2Cl_2 (5 mL) was added with stirring, pyridine (27 μL , 0.33 mMol) and Ac_2O (63 μL , 0.67 mMol). After stirring at RT for 4 h the reaction was concentrated under vacuum. The remaining residue was triturated with (1:1) Et_2O /pet. ether, filtered and dried under vacuum to give the title compound (163mg, 89%) as a white solid: ^1H NMR (400 MHz, CDCl_3) δ 8.14 (d, $J = 8.6$ Hz, 1 H), 7.59 (d, $J = 1.8$ Hz, 1 H), 7.43 (m, 6 H), 7.17 (m, 3 H), 6.84 (br s, 1 H), 6.71 (d, $J = 2.2$ Hz, 1 H), 6.69 (s, 1 H), 5.07 (d, $J = 10.9$ Hz, 1 H), 4.50 (2d, 1 H), 4.43 (2d, 1 H), 4.07 (d, $J = 15.5$ Hz, 1 H), 2.66 (m, 1 H), 2.51 (s, 3 H), 2.48 (d, $J = 15.5$ Hz, 1 H), 2.00 (s, 3 H), 0.95 (d, $J = 6.6$ Hz, 3 H), 0.34 (d, $J = 6.7$ Hz, 3 H); MS (ES) m/e 554.4 ($\text{M} + \text{H}$) $^+$.


Example 14

[00249] A mixture of 4-cyano-2-fluorobenzoic acid (1, 25 g), DMF (0.1 mL), and thionyl chloride (50 mL) in a 250 mL round-bottom flask equipped with a reflux condenser under N_2 atmosphere was heated to reflux in a 90°C oil bath for 30 mins. The reflux

condenser was replaced with a distillation head and the excess thionyl chloride was distilled from the reaction pot. The oil bath temperature was increased to 120°C to facilitate distillation. After 2 hours, the reaction vessel was cooled to 23°C and placed under reduced pressure (~25 Torr). The reaction vessel was then heated to 90°C for 20 mins, then to 130°C for 90 minutes to remove any excess remaining thionyl chloride. The reaction vessel was then cooled to 23°C and placed under high vacuum (~0.2 Torr). The reaction vessel was equipped with a clean distillation head and collection flask. The vessel was placed into a 137°C oil bath and the product was distilled from the reaction pot (b.p. = 107°C @ ~0.2 Torr). Upon cooling the distillate solidified, providing 2 as a white solid in close to quantitative yield.

[00250] A mixture of 4-chloro-2-fluorobenzoic acid (3, 25 g), DMF (0.1 mL), and thionyl chloride (50 mL) in a 250 mL round-bottom flask equipped with a reflux condenser under N₂ atmosphere was heated to reflux in a 90°C oil bath for 30 mins. The reflux condenser was replaced with a distillation head and the excess thionyl chloride was distilled from the reaction pot. The oil bath temperature was increased to 120°C to facilitate distillation. After 2 hours, the reaction vessel was cooled to 23°C and placed under reduced pressure (~25 Torr). The reaction vessel was then heated to 90°C for 20 mins, then to 120°C for 90 minutes to remove any excess remaining thionyl chloride. The reaction vessel was then cooled to 23°C and then equipped with a clean distillation head and collection flask. The vessel was placed into a 160°C oil bath and the product was distilled from the reaction pot (b.p. = 124°C @ ~25 Torr), furnishing 4 as a colorless oil in near quantitative yield.


Example 15



[00251] Piperonyloyl Chloride (1, 7.8 g, 42 mmol) was added to a solution of 2-amino-2-methyl-1-propanol (4 ml, 42 mmol), CH_2Cl_2 (200 mL), and triethylamine (11.7 ml, 84 mmol) at room temperature. The reaction solution was concentrated after 45 minutes, and the resulting residue was diluted with EtOAc (40 mL) and washed with brine (30 mL). The organic layer was dried (MgSO_4), filtered, and concentrated to yield a light brown oil. The crude material (2, 8.8g) was used without further purification.

[00252] Tetrapropylammonium perruthenate (TPAP, 638 mg, 1.8 mmol) was added portion-wise to a solution of 2 (8.6 g, 36.3 mmol), CH_2Cl_2 (73 ml, 2mL/mmol), 4-methyl-morpholine N-oxide (6.4 g, 54.5 mmol), and molecular sieves, 4Å activated powder (18g, 500mg/mmol) at 0°C under N_2 . The reaction was allowed to warm to r.t. after 15 minutes. After 1 hour the reaction was complete (TLC) and was filtered through silica, eluted with EtOAc (100 mL), and the filtrate was concentrated. This yielded 7 g of off-white solid (3). The material was recrystallized: EtOAc (30 mL), MeOH (10 mL), and Hexanes (1 mL) were added portion wise with heating and sonication. This was brought to a boil after which hexane (100 mL) was added while cooling. Crystals immediately began to precipitate as the solution was cooled. The mixture was filtered and the crystals were washed with hexane (10 mL) to provide 4.88 g (57 %) of 3 as off-white fluffy crystals.

Example 16




A mixture of 2-amino-2-methyl-1-propanol (4 ml, 42 mmol), CH_2Cl_2 (200 mL), 4-acetylbenzoic acid (1, 6.9 g, 42 mmol), EDC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide HCl, 12.1 g, 63 mmol), HOBT (N-hydroxybenzotriazole H_2O , 6.4 g, 42 mmol), and Hunig's base (diisopropylethylamine, 22 mL, 126 mmol) was stirred at room temperature for 18 hours. Upon completion (TLC, LC/MS) the reaction mixture was concentrated, and the resulting residue was diluted with EtOAc (50 mL) and washed with NaHCO_3 (2 x 40 mL) and

brine (40 mL). The organic layer was dried (MgSO_4), filtered, and the filtrate was concentrated. The crude material was purified by flash column chromatography (4:1:1 hexanes:EtOAc: CH_2Cl_2 ; 1:1:1 hexanes:EtOAc: CH_2Cl_2 ; 1:1 EtOAc: CH_2Cl_2) to remove bisacylated material. Alcohol 2 was obtained in 54 % yield (5.36 g).


Tetrapropylammonium perruthenate (TPAP, 387 mg, 1.1 mmol) was added portion-wise to a solution of 2 (5.36 g, 22.8 mmol), CH_2Cl_2 (46 mL, 2mL/mmol), 4-methyl-morpholine N-oxide (4 g, 34.2 mmol), and molecular sieves, 4Å activated powder (11.4g, 500mg/mmol) at 0°C under N_2 . The reaction was allowed to warm to r.t. after 15 minutes. After 1 hour the reaction was complete (TLC) and was filtered through silica, which was eluted with EtOAc (100 mL), and the filtrate was concentrated. This yielded 4.4 g of light pink solid (3). The material was recrystallized: 1:1 Hexanes:EtOAc (5 mL), CH_2Cl_2 (20 mL), and MeOH (10 mL) were added portion-wise with heating and sonication. This was brought to a boil after which hexane (100 mL) was added while cooling. Crystals immediately began to precipitate as the solution was cooled. The mixture was filtered and the crystals were washed with hexane (10 mL), affording 2.46 g (46 %) of 3 as off-white fluffy crystals.

Example 17




[00253] A solution of 4-acetyl benzoic acid (500 mg) in oxalyl chloride (5 mL) was heated to reflux for 2 hours. Any remaining oxalyl chloride was evaporated by rot-vap, and the residue was dried by vacuum. The yield of product was quantitative.

[00254] Example 18



((R)-1-Isopropyl-2-oxo-but-3-enyl)-carbamic acid *tert*-butyl ester

[00255] Tetrahydrofuran (THF, 100 mL) and a 1.0M solution of vinyl magnesium bromide in THF (360 mL, 360 mmol, 3.1 equiv) was cooled to $-78\text{ }^{\circ}\text{C}$ while stirring under a nitrogen atmosphere. The mixture was treated dropwise with a solution of [(*R*)-(methoxymethyl-carbamoyl)-methyl-propyl]-carbamic acid *tert*-butyl ester (30.3 g, 116 mmol, 1 equiv) in THF (50 mL) over 30 min. After the resultant dark yellow mixture was stirred for 30 min at $-78\text{ }^{\circ}\text{C}$, the cooling bath was removed and the reaction mixture was warmed slowly to room temperature overnight (15 h). The reaction mixture was poured slowly into an ice-chilled solution of 1N aqueous hydrochloric acid (700 mL) and then warmed to room temperature. The organics were extracted with (3 x 600 mL) ethyl acetate, dried over sodium sulfate, filtered, and concentrated in vacuo. Purification by flash column chromatography (5-10% ethyl acetate/hexanes) provided the product as a white solid (16.8 g, 64%). ESMS $[\text{M}+\text{H}]^+$: 228.4.




[(*R*)-(E)-1-Isopropyl-4-(3-methoxy-phenyl)-2-oxo-but-3-enyl]-carbamic acid *tert*-butyl ester

[00256] To a solution of ((*R*)-1-Isopropyl-2-oxo-but-3-enyl)-carbamic acid *tert*-butyl ester (13.54 g, 59.6 mmol) in dry acetonitrile (150 mL) under argon, was added 3-iodoanisole (13.96 g, 59.6 mmol), triethylamine (9.1 mL, 65.6 mmol) followed by palladium (II) acetate (335 mg, 1.49 mmol). The resulting clear yellow solution was heated to $80\text{ }^{\circ}\text{C}$. Upon heating, the reaction darkened and the precipitation of palladium black occurred. After 15 h,

the reaction mixture was allowed to cool to room temperature, quenched with water (150 mL) and diluted with ether (150 mL). The ether layer was washed with brine (100 mL) and the combined aqueous layers were extracted with ether (two 50 mL portions). The extracts were dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue was immediately purified by silica gel chromatography (9:1 hexanes/EtOAc) to provide 17.6 g (88%) of [(*R*)-(*E*)-1-Isopropyl-4-(3-methoxy-phenyl)-2-oxo-but-3-enyl]-carbamic acid *tert*-butyl ester as a yellow oil. MS(ES+) *m/e* 334.0 [*M*+*H*]⁺.

[(*R*)-(*Z*)-4-(3-Cyano-phenyl)-1-isopropyl-2-oxo-but-3-enyl]-carbamic acid *tert*-butyl ester


[00257] Following the procedure described for [(*R*)-(*E*)-1-isopropyl-4-(3-methoxy-phenyl)-2-oxo-but-3-enyl]-carbamic acid *tert*-butyl ester with 3-iodobenzonitrile (5.50 g, 24.0 mmol, 1 equiv) afforded the title compound as a yellow solid (7.4 g of ~90% purity material). ESMS [*M*+*H*]⁺: 329.2.



[(*R*)-(*E*)-1-Isopropyl-4-(3-methoxy-phenyl)-2-oxo-butyl]-carbamic acid *tert*-butyl ester

[00258] To a solution of [(*R*)-(*E*)-1-Isopropyl-4-(3-methoxy-phenyl)-2-oxo-but-3-enyl]-carbamic acid *tert*-butyl ester (17.6 g, 52.9 mmol) in ethyl acetate (450 mL) under nitrogen was added 10 wt % palladium on carbon (300 mg). The nitrogen was replaced with a balloon of hydrogen and the flask was purged. After 3 h, the reaction flask was purged with nitrogen and filtered through a pad of celite (rinsing with ethyl acetate). The filtrate was concentrated under reduced pressure and the residue was purified by silica gel chromatography (9:1 hexanes/EtOAc) to provide 16.2 g (91%) of [(*R*)-(*E*)-1-Isopropyl-4-(3-

methoxy-phenyl)-2-oxo-butyl]-carbamic acid *tert*-butyl ester as a colorless oil. MS(ES+) *m/e* 336.4 [M+H]⁺. [α]_D²⁰ = +19.1 (*c* = 0.755, MeOH). ¹H NMR (400 MHz, CDCl₃) δ 7.21 (m, 1H), 6.80-6.77 (m, 2H), 6.75 (s, 1H), 5.13 (d, *J* = 8.4 Hz, 1H), 4.28 (dd, *J* = 8.8, 4.4 Hz, 1H), 3.81 (s, 3H), 2.93-2.88 (m, 2H), 2.85-2.76 (m, 2H), 2.14 (m, 1H), 1.46 (s, 9H), 1.00 (d, *J* = 6.8 Hz, 3H), 0.75 (d, *J* = 6.8 Hz, 3H).




[(*R*)-4-(3-Cyano-phenyl)-1-isopropyl-2-oxo-butyl]-carbamic acid *tert*-butyl ester

[00259] Palladium on carbon (740 mg of 10% wt/wt Pd/C) was added to a degassed solution of [(*R*)-(Z)-4-(3-cyano-phenyl)-1-isopropyl-2-oxo-but-3-enyl]-carbamic acid *tert*-butyl ester (7.4 g, 22.5 mmol, 1 equiv) in methanol (200 mL) at room temperature. The reaction mixture was thoroughly degassed and backfilled with hydrogen gas from a balloon. Hydrogenation proceeded at atmospheric pressure for 2.5 h. The (degassed) reaction mixture was then diluted with diethyl ether (300 mL), filtered through Celite, and washed with additional ether (2 x 100 mL). Upon concentration in vacuo, the residue was purified by flash column chromatography (20% ethyl acetate-hexanes) to provide the ketone product as a white solid (5.9 g, 79%). ESMS [M+H]⁺: 331.2. ¹H NMR (400 MHz, CDCl₃) δ 7.51 (m, 2H), 7.46 (m, 1H), 7.41 (m, 1H), 5.07 (m, 1H), 4.24 (m, 1H), 2.98 (m, 2H), 2.86 (m, 2H), 2.11 (m, 1H), 1.45 (s, 9H), 0.98 (d, 3H, *J* = 6.76 Hz), 0.74 (d, 3H, *J* = 6.78 Hz). [α]_D = +24.74 (*c* = 0.95, CH₃OH).

Example 19

[00260] Following procedures analogous to those set forth above, the following compounds were prepared:



R ₇	R ₁	R ₃	[M+H] ⁺
F	3-MeO-Ph-CH ₂ -	3-F-4-Me-Ph-	549
F	3-MeO-Ph-CH ₂ -	4-MeO-Ph-	547
F	3-MeO-Ph-CH ₂ -	4-Me-Ph-	531
F	3-MeO-Ph-CH ₂ -	3,4-piperonyl-	561
F	3-MeO-Ph-CH ₂ -	Methoxymethyl-	485
CN	3-MeO-Ph-CH ₂ -	4-Ac-Ph-	566
CN	3-MeO-Ph-CH ₂ -	4-Me-Ph-	538
CN	3-MeO-Ph-CH ₂ -	3,4-piperonyl-	568
CN	3-MeO-Ph-CH ₂ -	3-F-4-Me-Ph-	556
Cl	3-MeO-Ph-CH ₂ -	3-F-4-Me-Ph-	565

Cl	3-MeO-Ph- CH ₂ -	4-Me-Ph-	547
Cl	3-MeO-Ph- CH ₂ -	Methoxymethyl-	501
Cl	3-MeO-Ph- CH ₂ -	3,4-piperonyl-	577
MeO-	3-MeO-Ph- CH ₂ -	3-F-4-Me-Ph-	561
MeO-	3-MeO-Ph- CH ₂ -	4-Me-Ph-	543
MeO-	3-MeO-Ph- CH ₂ -	3,4-piperonyl	573
OH	3-MeO-Ph- CH ₂ -	3-F-4-Me-Ph-	547
CN	Ph-CH ₂ -	4-OEt-Ph-	538.2
CN	Ph-CH ₂ -	6-trifluoromethyl-3-pyridyl	563.2
CN	Ph-CH ₂ -	2-benzo[b]thiophene	550.2
CN	Ph-CH ₂ -	3-(5- <i>t</i> -butyl-2-methyl-2H-pyrazole)	554.4
CN	Ph-CH ₂ -	3-(2,5-dimethyl-2H-pyrazole)	512.4
F	Ph-CH ₂ -	6-trifluoromethyl-3-pyridyl	556.0
F	Ph-CH ₂ -	2-furyl	477.0
F	Ph-CH ₂ -	3-(5- <i>t</i> -butyl-2-methyl-2H-pyrazole)	547.2
Cl	Ph-CH ₂ -	4-CN-Ph-	529.2
Cl	Ph-CH ₂ -	4-AcNH-Ph-	561.4
Cl	Ph-CH ₂ -	6-trifluoromethyl-3-pyridyl-	573.0
Cl	Ph-CH ₂ -	5-benzo[1,2,3]thiadiazole-	562.2
Cl	Ph-CH ₂ -	2-furyl-	494.4
Cl	Ph-CH ₂ -	3-(2,5-dimethyl-2H-pyrazole)-	522.0
F	Ph-CH ₂ -	4-pyridyl-	488.4
F	Ph-CH ₂ -	3-(2,5-dimethyl-2H-pyrazole)-	505.4
F	Ph-CH ₂ -	4-AcNH-Ph-	544.4
F	Ph-CH ₂ -	5-benzo[1,2,3]thiadiazole-	545.2

CN	Ph-CH ₂ -	2-(1-methyl-1H-indole)-	547.4
Cl	Ph-CH ₂ -	3-pyridyl-	505.2
F	Ph-CH ₂ -	4-OMe-Ph-	517.2
CN	Ph-CH ₂ -	3-(2,5-dimethyl-furan)-	512.2
F	Ph-CH ₂ -	3-(2,5-dimethyl-furan)-	505.2
CN	Ph-CH ₂ -	2-(5-methyl-thiophene)-	514.2
F	Ph-CH ₂ -	2-(5-methyl-thiophene)-	507.0
CN	Ph-CH ₂ -	2-(1-methyl-1H-pyrrole)-	497.4
CN	Ph-CH ₂ -	5-benzo[1,2,3]thiadiazole-	550.0
Cl	Ph-CH ₂ -	6-Me-3-pyridyl-	518.2
Cl	Ph-CH ₂ -	2-(1-methyl-1H-pyrrole)-	506.0
Cl	Ph-CH ₂ -	2-(5-methyl-pyrazine)-	520.2
Cl	Ph-CH ₂ -	3-(5-methyl-isoxazole)-	508.2
Cl	Ph-CH ₂ -	3-benzo[c]isoxazole-	544.2
CN	Ph-CH ₂ -	4-(1-methyl-1H-imidazole)-	498.1
CN	Ph-CH ₂ -	3-N(CH ₃) ₂ -Ph-	537.4
CN	Ph-CH ₂ -	3-(5-methyl-2-trifluoromethyl-furan)-	566.4
CN	Ph-CH ₂ -	3-(5-methyl-isoxazole)-	499.6
F	Ph-CH ₂ -	4-(1-methyl-1H-imidazole)-	491.4
F	Ph-CH ₂ -	2-(1-methyl-1H-pyrrole)-	490.4
F	Ph-CH ₂ -	3-benzo[c]isoxazole-	528.4
F	Ph-CH ₂ -	3-(5-methyl-isoxazole)-	492.4

Example 20

[00261] A pharmaceutical composition for intravenous administration is prepared in the following manner.

1 mg/mL (as free base) IV solution with the vehicle being pH 5.0, 50 mM sodium acetate buffer containing 3.5% (w/v) mannitol:

Composition*	Unit Formula (mg/mL)
Compound of Example 2	
(free base)	1.000
Glacial Acetic Acid	1.081
Sodium Acetate Trihydrate	4.355
Mannitol, pyrogen free	35.000
Water for Injection (WFI)	q.s. to 1 mL

*All components other than the active compound are USP or Ph. Eur.

[00262] A suitable compounding vessel is filled to approximately 75% of the bulk solution volume with WFI. The glacial acetic acid (1.081 g), sodium acetate trihydrate (4.355 g), mannitol (35.000 g), and active (1.000 g) are weighed and individually added to the compounding vessel. After the additions, the ingredients are dissolved in the mixture by stirring with a mixer. The pH of the bulk solution is measured and adjusted to 5.0 with 5N NaOH or 5N glacial acetic acid. The solution is brought to its final volume (1 liter) with WFI. Where the active compound is a pharmaceutically acceptable salt, hydrate or salt of a hydrate, e.g., a monochloride salt or a hydrate of a monochloride salt, the amount of the active equivalent to the free base is added.

Example 21

Preparation of a hydrochloride salt of N-(3-Aminopropyl)-N-[(R)-1-3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl]-2-methyl-propyl]-4-methyl-benzamide

[00263] In one embodiment, a hydrochloric acid salt of N-(3-Aminopropyl)-N-[(R)-1-3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl]-2-methyl-propyl]-4-methyl-benzamide, such as prepared in accordance with Example 2, is formed in the following manner. An HCl salt is suitably prepared by reacting the free base (e.g. dissolved in a suitable solvent, such as one or more of TBME, THF, and ethylacetate) with hydrochloric acid, suitably aqueous hydrochloric acid or otherwise in the presence of water. The mixture is suitably seeded with HCl salt to assist crystallization. The resulting salt may be isolated using conventional techniques such as filtration and drying (suitably washing the salt with a solvent, such as one or more of TBME, THF, and ethylacetate).

[00264] In a particular embodiment, the salt is prepared as follows:

1. Dissolve the free base into 10 volumes of TBME and 5 volumes of THF, dissolution suitably conducted at room temperature;
2. Add aqueous HCl, e.g. 6.0 M or 12.0 M, suitably 1.1 eq, e.g. dropwise;
3. Suitably seed the solution with crystals of HCl salt, e.g. prepared according to preparation B below;
4. Suitably stir until crystals form;
5. Isolate the crystals, suitably by filtration, washing with a suitable solvent such as TBME, and drying.

[00265] In more particular embodiments a salt is prepared as follows:

Preparation A

[00266] To a vial fitted with a stir bar add 500 mg of N-(3-Aminopropyl)-N-[(R)-1-3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl]-2-methyl-propyl]-4-methyl-benzamide, and 3 ml TBME, and dissolve the free base in the TBME at room temperature. Bubble in HCl gas for about 30 seconds until a white solid forms. Filter and rinse the solids twice with 500 μ l TBME. Transfer the solids to a test tube, add 4 ml TBME and a stir bar, and stir at room temperature (the salt does not dissolve). Place the test tube containing the mixture in an Argonaut RS10 heating block (Argonaut Technologies, Foster City, California), or an equivalent, and temperature cycle as follows:

1. 30 min at room temperature
2. ramp to 40C over 30 min and hold at 40C for 30 min
3. ramp to room temperature over 30 min and hold at room temperature for 30 min
4. ramp to 10C over 30 min and hold at 10C for 30 min
5. ramp to room temperature over 30 min and hold at room temperature for 30 min
6. repeat 2 to 5 as desired

By 20 hours cycling some solvent evaporates, a white powder forms on the sides and a melt-like solid forms in the bottom of the test tube. Add 2 ml TBME. After 15 days, filter the light yellow solids and wash twice with about 0.5 ml TBME then twice with 1 ml hexane.

Dry in a vacuum oven at room temperature for 12-24 hours to obtain about 440 mg HCl salt.

[00267] Figures 1 and 2 represent a XRPD and MDSC scan, respectively, of a salt prepared according to Preparation A.

Preparation B

[00268] Dissolve 4.23 g of N-(3-Aminopropyl)-N-[(R)-1-3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-methyl-benzamide in 5 volumes (21 ml) of THF. Add 10 volumes (42 ml) of TBME (tert-butyl methyl ether), then 1.1 eq of 12N HCl (750 μ l) in one portion. Seed with HCl salt such as prepared in Preparation A (e.g. in 3 additions of about 10 mg each staggered over about 10 minutes). Stir the mixture overnight, e.g. for about 12-24 hours. Filter the mixture, wash the solids with TBME, and dry the solids in a vacuum oven at room temperature overnight, e.g. for about 12-24 hours, to yield about 4.3 g of HCl salt.

[00269] Figure 3 represents an XRPD scan of a salt prepared according to Preparation B.

[00270] The present invention includes crystalline N-(3-Aminopropyl)-N-[(R)-1-3-benzyl-7-chloro-4-oxo-4H-chromen-2-yl)-2-methyl-propyl]-4-methyl-benzamide hydrochloride salt having characteristic X-ray diffraction peaks at about 4.8, 9.7, 14.7, 17.9, 18.3, 20.1, 20.9, 22.5, 23.2, 23.8, 26.1 and 26.9 degrees 2 theta.

Example 22

Inhibition of Cellular Viability in Tumor Cell Lines Treated with KSP Inhibitors.

Materials and Solutions:

- Cells: SKOV3, Ovarian Cancer (human).
- Media: Phenol Red Free RPMI + 5% Fetal Bovine Serum + 2mM L-glutamine.
- Colorimetric Agent for Determining Cell Viability: Promega MTS tetrazolium compound.
- Control Compound for max cell kill: Topotecan, 1 μ M.

Procedure: Day 1 - Cell Plating:

[00271] Adherent SKOV3 cells are washed with 10mLs of PBS followed by the addition of 2mLs of 0.25% trypsin and incubation for 5 minutes at 37°C. The cells are rinsed from the flask using 8 mL of media (phenol red-free RPMI+ 5%FBS) and transferred to fresh flask. Cell concentration is determined using a Coulter counter and the appropriate volume of

cells to achieve 1000 cells/100 μ L is calculated. 100 μ L of media cell suspension (adjusted to 1000 cells/100 μ L) is added to all wells of 96-well plates, followed by incubation for 18 to 24 hours at 37°C, 100% humidity, and 5% CO₂, allowing the cells to adhere to the plates.

Procedure: Day 2 – Compound Addition:

[00272] To one column of the wells of an autoclaved assay block are added an initial 2.5 μ L of test compound(s) at 400X the highest desired concentration. 1.25 μ L of 400X (400 μ M) Topotecan is added to other wells (optical density's from these wells are used to subtract out for background absorbance of dead cells and vehicle). 500 μ L of media without DMSO are added to the wells containing test compound, and 250 μ L to the Topotecan wells. 250 μ L of media + 0.5% DMSO is added to all remaining wells, into which the test compound(s) are serially diluted. By row, compound-containing media is replica plated (in duplicate) from the assay block to the corresponding cell plates. The cell plates are incubated for 72 hours at 37°C, 100% humidity, and 5% CO₂.

Procedure: Day 4 – MTS Addition and OD Reading:

[00273] The plates are removed from the incubator and 40 μ L MTS / PMS is added to each well. Plates are then incubated for 120 minutes at 37°C, 100% humidity, 5%CO₂, followed by reading the ODs at 490nm after a 5 second shaking cycle in a ninety-six well spectrophotometer.

Data Analysis

[00274] The normalized % of control (absorbance- background) is calculated and an XLfit is used to generate a dose-response curve from which the concentration of compound required to inhibit viability by 50% is determined. The compounds of the present invention show activity when tested by this method as described above.

Example 23

Enantiomer Separation

[00275] An enriched 3:1 R:S mixture of chromenone enantiomers was separated into its pure enantiomers by chiral chromatography with the following conditions: Column -

Chiralpak AD, 250 x 4.6 mm (Diacel Inc.). Sample – 22.5 mg/ml in 1:1 *i*-PrOH:hexanes. Conditions – 40 min at isocratic 50% *i*-PrOH in Hexanes, (S)-enantiomer elutes at 18.35 min, (R)-enantiomer elutes at 26.87 min. The (R)-enantiomer was significantly more potent than the (S)-enantiomer of the compound of Example 2.

Example 24

Monopolar Spindle Formation following Application of a KSP Inhibitor

[00276] Human tumor cells Skov-3 (ovarian) were plated in 96-well plates at densities of 4,000 cells per well, allowed to adhere for 24 hours, and treated with various concentrations of the chromenone compounds for 24 hours. Cells were fixed in 4% formaldehyde and stained with antitubulin antibodies (subsequently recognized using fluorescently-labeled secondary antibody) and Hoechst dye (which stains DNA).

[00277] Visual inspection revealed that the compounds caused cell cycle arrest in the prometaphase stage of mitosis. DNA was condensed and spindle formation had initiated, but arrested cells uniformly displayed monopolar spindles, indicating that there was an inhibition of spindle pole body separation. Microinjection of anti-KSP antibodies also causes mitotic arrest with arrested cells displaying monopolar spindles.

Example 25

Inhibition of Cellular Proliferation in Tumor Cell Lines Treated with KSP Inhibitors.

[00278] Cells were plated in 96-well plates at densities from 1000-2500 cells/well of a 96-well plate and allowed to adhere/grow for 24 hours. They were then treated with various concentrations of drug for 48 hours. The time at which compounds are added is considered T_0 . A tetrazolium-based assay using the reagent 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulphophenyl)-2H-tetrazolium (MTS) (Patent No. 5,185,450) (see Promega product catalog #G3580, CellTiter 96® AQueous One Solution Cell Proliferation Assay) was used to determine the number of viable cells at T_0 and the number of cells remaining after 48 hours compound exposure. The number of cells remaining after 48 hours was compared to the number of viable cells at the time of drug addition, allowing for calculation of growth inhibition.

[00279] The growth over 48 hours of cells in control wells that had been treated with vehicle only (0.25% DMSO) is considered 100% growth and the growth of cells in wells with compounds is compared to this.

[00280] A GI_{50} was calculated by plotting the concentration of compound in μM vs the percentage of cell growth in treated wells. The GI_{50} calculated for the compounds is the estimated concentration at which growth is inhibited by 50% compared to control, i.e., the concentration at which:

$$100 \times [(Treated_{48} - T_0) / (Control_{48} - T_0)] = 50$$

wherein $Treated_{48}$ is the value at 48 hours for the treated cells and $Control_{48}$ is the value at 48 hours for the control population.

[00281] All concentrations of compounds are tested in duplicate and controls are averaged over 12 wells. A very similar 96-well plate layout and GI_{50} calculation scheme is used by the National Cancer Institute (see Monks, et al., J. Natl. Cancer Inst. 83:757-766 (1991)). However, the method by which the National Cancer Institute quantitates cell number does not use MTS, but instead employs alternative methods.

[00282] Compounds of Examples 1-13 above inhibited cell proliferation in human ovarian tumor cell lines (SKOV-3).

Example 26

Calculation of IC_{50} :

[00283] Measurement of a compound's IC_{50} for KSP activity uses an ATPase assay. The following solutions are used: Solution 1 consists of 3 mM phosphoenolpyruvate potassium salt (Sigma P-7127), 2 mM ATP (Sigma A-3377), 1 mM IDTT (Sigma D-9779), 5 μM paclitaxel (Sigma T-7402), 10 ppm antifoam 289 (Sigma A-8436), 25 mM Pipes/KOH pH 6.8 (Sigma P6757), 2 mM $MgCl_2$ (VWR JT400301), and 1 mM EGTA (Sigma E3889). Solution 2 consists of 1 mM NADH (Sigma N8129), 0.2 mg/ml BSA (Sigma A7906), pyruvate kinase 7U/ml, L-lactate dehydrogenase 10 U/ml (Sigma P0294), 100 nM KSP motor domain, 50 $\mu g/ml$ microtubules, 1 mM DTT (Sigma D9779), 5 μM paclitaxel (Sigma T-7402), 10 ppm antifoam 289 (Sigma A-8436), 25 mM Pipes/KOH pH 6.8 (Sigma P6757), 2 mM $MgCl_2$ (VWR JT4003-01), and 1 mM EGTA (Sigma E3889). Serial dilutions (8-12 two-fold dilutions) of the compound are made in a 96-well microtiter plate (Corning Costar 3695) using Solution 1. Following serial dilution each well has 50 μl of Solution 1. The reaction is started by adding 50 μl of solution 2 to each well. This may be done with a multichannel pipettor either manually or with automated liquid handling devices. The microtiter plate is then transferred to a microplate absorbance reader and multiple absorbance


readings at 340 nm are taken for each well in a kinetic mode. The observed rate of change, which is proportional to the ATPase rate, is then plotted as a function of the compound concentration. For a standard IC_{50} determination the data acquired is fit by the following four parameter equation using a nonlinear fitting program (e.g., Grafit 4):

$$y = \frac{\text{Range}}{1 + \left(\frac{x}{IC_{50}} \right)^s} + \text{Background}$$

where y is the observed rate and x is the compound concentration.

What is claimed is:

1. A compound having the structure:



wherein:

R_1 is chosen from optionally substituted aryl- C_1 - C_4 -alkyl- and optionally substituted heteroaryl- C_1 - C_4 -alkyl-;

R_2 and $R_{2'}$ are independently chosen from hydrogen, optionally substituted alkyl-, optionally substituted alkoxy, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-; or R_2 and $R_{2'}$ taken together form an optionally substituted 3- to 7-membered ring;

R_{12} is selected from the group consisting of optionally substituted imidazolyl, optionally substituted imidazolyl-, $-NHR_4$; $-N(R_4)(COR_3)$; $-N(R_4)(SO_2R_{3a})$; and $-N(R_4)(CH_2R_{3b})$;

R_3 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, optionally substituted heteroaralkyl-, $R_{15}O$ - and $R_{17}NH$ -;

R_{3a} is chosen from optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, optionally substituted heteroaralkyl-, and $R_{17}NH$ -;

R_{3b} is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted

heteroaralkyl;

R₄ is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heterocycl-, and optionally substituted heteroaralkyl-;

R₅, R₆, R₇ and R₈ are independently chosen from hydrogen, acyl, optionally substituted alkyl-, optionally substituted alkoxy, halogen, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl-, alkylsulfonamido-, alkylthio-, carboxyalkyl-, carboxamido-, aminocarbonyl-, optionally substituted aryl and optionally substituted heteroaryl; and

R₁₅ is chosen from optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-,

R₁₇ is hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, or optionally substituted heteroaralkyl-, including single stereoisomers, mixtures of stereoisomers;

a pharmaceutically acceptable salt of a compound of Formula I;

a pharmaceutically acceptable solvate of a pharmaceutically acceptable solvate of a compound of Formula I;

or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I.

2. A compound according to claim 1, wherein if either R₂ or R₂' is hydrogen, then the other is not hydrogen.
3. A compound according to claim 1, wherein R₅, R₆, R₇ and R₈ are each independently selected from hydrogen, amino, alkylamino, hydroxyl, halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy and cyano.
4. A compound according to claim 3, wherein R₅, R₆, R₇ and R₈ are each independently selected from hydrogen, cyano, methoxy, and halogen.

5. A compound according to claim 4, wherein R_5 , R_6 , and R_8 are each hydrogen and R_7 is cyano, methoxy or halogen.
6. A compound according to claim 1, wherein R_2 is optionally substituted C_1 - C_4 alkyl and R_2 is hydrogen or optionally substituted C_1 - C_4 alkyl.
7. A compound according to claim 6, wherein R_2 is hydrogen and R_3 is optionally substituted C_1 - C_4 alkyl.
8. A compound according to claim 7, wherein R_2 is hydrogen and R_2 is ethyl or propyl.
9. A compound according to claim 8, wherein R_2 is i-propyl.
10. A compound according to claim 1, wherein R_1 is optionally substituted phenyl- C_1 - C_4 -alkyl- and optionally substituted naphthalenylmethyl.
11. A compound according to claim 10, wherein R_1 is benzyl, chlorobenzyl, methylbenzyl, methoxybenzyl, cyanobenzyl, hydroxybenzyl, dichlorobenzyl, dimethoxybenzyl, or naphthalenylmethyl.
12. A compound according to claim 11, wherein R_1 is benzyl-, cyanobenzyl-, methoxybenzyl-, or naphthalenylmethyl.
13. A compound according to claim 12, wherein R_1 is benzyl.
14. A compound according to claim 1, wherein R_{12} is $-N(R_4)(COR_3)$ and R_3 is selected from optionally substituted alkyl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, optionally substituted heteroaryl-, optionally substituted aryl-, $R_{15}O$ - and $R_{17}NH$ -; R_{15} is chosen from optionally substituted alkyl and optionally substituted aryl; and R_{17} is chosen from

hydrogen, optionally substituted alkyl and optionally substituted aryl.

15. A compound according to claim 14, wherein R_3 is chosen from phenyl; phenyl substituted with one or more of the following substituents: halo, C_1 - C_4 alkyl, C_1 - C_4 alkyl substituted with hydroxyl, C_1 - C_4 alkoxy, C_1 - C_4 alkyl substituted with C_1 - C_4 alkoxy, nitro, formyl, carboxy, cyano, methylenedioxy, ethylenedioxy, acyl, -N-acyl, or trifluoromethyl; benzyl; phenoxymethyl-; halophenoxymethyl-; phenylvinyl-; heteroaryl-; heteroaryl- substituted with C_1 - C_4 alkyl or C_1 - C_4 alkyl substituted with halo; C_1 - C_4 alkyl substituted with C_1 - C_4 alkoxy- and benzyloxymethyl-.
16. A compound according to claim 15, wherein R_3 is chosen from phenyl, halophenyl, dihalophenyl, cyanophenyl, halo(trifluoromethyl)phenyl, hydroxymethylphenyl, methoxyphenyl, ethoxyphenyl, carboxyphenyl, ethylphenyl, tolyl, methylenedioxyphenyl, ethylenedioxyphenyl, methoxychlorophenyl, dihydro-benzodioxinyl, methylhalophenyl, trifluoromethylphenyl, bis(trifluoromethyl)phenylbenzyl, furanyl, C_1 - C_4 alkyl substituted furanyl, trifluoromethylfuranyl, C_1 - C_4 alkyl substituted trifluoromethylfuranyl, benzofuranyl, thiophenyl, C_1 - C_4 alkyl substituted thiophenyl, benzothiophenyl, benzothiadiazolyl, pyridinyl, indolyl, methylpyridinyl, trifluoromethylpyridinyl, pyrrolyl, quinoliny, picoliny, pyrazolyl, C_1 - C_4 alkyl substituted pyrazolyl, N-methyl pyrazolyl, C_1 - C_4 alkyl substituted N-methyl pyrazolyl, C_1 - C_4 alkyl substituted pyrazinyl, C_1 - C_4 alkyl substituted isoxazolyl, benzoisoxazolyl, morpholinomethyl, methylthiomethyl, methoxymethyl, N-methyl imidazolyl, and imidazolyl.
17. A compound according to claim 16, wherein R_3 is tolyl, halophenyl, halomethylphenyl, hydroxymethylphenyl, methylenedioxyphenyl, formylphenyl or cyanophenyl.
18. A compound according to claim 1, wherein R_{12} is $-N(R_4)(SO_2R_{3a})$ and R_{3a} is chosen from C_1 - C_{13} alkyl; phenyl; naphthyl; phenyl substituted with halo, C_1 - C_4 alkyl-, C_1 - C_4 alkoxy, cyano, nitro, methylenedioxy, or trifluoromethyl; biphenyl; and heteroaryl.


19. A compound according to claim 18, wherein R_{3a} is chosen from phenyl substituted with halo, C_1 - C_4 alkyl-, C_1 - C_4 alkoxy, cyano, nitro, methylenedioxy, or trifluoromethyl; and naphthyl.
20. A compound according to claim 1, wherein R_{12} is $-N(R_4)(CH_2R_{3b})$ and R_{3b} is chosen from C_1 - C_{13} alkyl; substituted C_1 - C_4 alkyl; phenyl; naphthyl; phenyl substituted with carboxy, alkoxycarbonyl, cyano, halo, C_1 - C_4 alkyl-, C_1 - C_4 alkoxy, nitro, methylenedioxy, or trifluoromethyl; biphenyl, benzyl; and heterocyclyl.
21. A compound according to claim 20, wherein R_{3b} is chosen from phenyl substituted with one or more halo, methyl, methoxy, cyano, trifluoromethyl, trifluoromethoxy, carboxy, and or methoxycarbonyl groups; piperidinyl; and naphthyl.
22. A compound according to claim 21, wherein R_{3b} is chosen from halophenyl, polyhalophenyl, tolyl, dimethylphenyl, methoxyphenyl, dimethoxyphenyl, cyanophenyl, trifluoromethylphenyl, trifluoromethoxyphenyl, bis(trifluoromethyl)phenyl, carboxyphenyl, t-butylphenyl, methoxycarbonylphenyl, piperidinyl and naphthyl.
23. A compound according to claim 1, wherein R_{12} is $-N(R_4)(CH_2R_{3b})$ and R_4 is chosen from hydrogen, optionally substituted C_1 - C_{13} alkyl, optionally substituted aryl, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heterocyclyl, and optionally substituted heteroaryl- C_1 - C_4 -alkyl- (preferably hydrogen or optionally substituted C_1 - C_{13} alkyl).
24. A compound according to claim 23, wherein R_4 is chosen from hydrogen, C_1 - C_4 alkyl; cyclohexyl; phenyl substituted with hydroxyl, C_1 - C_4 alkoxy or C_1 - C_4 alkyl; benzyl; heteroarylmethyl-; heteroarylethyl-; heteroarylpropyl-; and R_{16} -alkylene-, wherein R_{16} is hydroxyl, di(C_1 - C_4 alkyl)amino-, (C_1 - C_4 alkyl)amino-, amino, C_1 - C_4 alkoxy-, or N-heterocyclyl-, particularly pyrrolidino, piperidino or imidazolyl.

25. A compound according to claim 24, wherein R_4 is R_{16} -alkylene- wherein R_{16} is hydroxyl, di(C_1 - C_4 alkyl)amino-, (C_1 - C_4 alkyl)amino-, amino, C_1 - C_4 alkoxy, or N-heterocyclyl.
26. A compound according to claim 1, wherein R_{12} is $-NH(R_4)$ and R_4 is chosen from hydrogen, optionally substituted C_1 - C_{13} alkyl, optionally substituted aryl, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heterocyclyl, and optionally substituted heteroaryl- C_1 - C_4 -alkyl- (preferably hydrogen or optionally substituted C_1 - C_{13} alkyl).
27. A compound according to claim 26, wherein R_4 is R_{16} -alkylene- wherein R_{16} is hydroxyl, di(C_1 - C_4 alkyl)amino-, (C_1 - C_4 alkyl)amino-, amino, C_1 - C_4 alkoxy, or N-heterocyclyl.
28. A compound according to claim 1, wherein R_{12} is $-N(R_4)(COR_3)$ and R_4 is chosen from hydrogen, C_1 - C_4 alkyl; cyclohexyl; phenyl substituted with hydroxyl, C_1 - C_4 alkoxy or C_1 - C_4 alkyl; benzyl; heteroarylmethyl-; heteroarylethyl-; heteroarylpropyl-; and R_{16} -alkylene-, wherein R_{16} is hydroxyl, di(C_1 - C_4 alkyl)amino-, (C_1 - C_4 alkyl)amino-, amino, C_1 - C_4 alkoxy-, or N-heterocyclyl-, particularly pyrrolidino, piperidino or imidazolyl.
29. A compound according to claim 28, wherein R_4 is R_{16} -alkylene-, and R_{16} is C_1 - C_4 alkoxy, nitro, amino, alkylamino, dialkylamino, hydroxyl, or N-heterocyclyl.
30. A compound according to claim 29, wherein R_4 is R_{16} -alkylene-, and R_{16} is amino.
31. A compound according to claim 29, wherein R_4 is chosen from hydrogen, methyl, ethyl, propyl, butyl, cyclohexyl, carboxyethyl, carboxymethyl, methoxyethyl-, hydroxyethyl-, hydroxypropyl-, dimethylaminoethyl-, dimethylaminopropyl-, diethylaminoethyl-, diethylaminopropyl-, aminopropyl-, methylaminopropyl-, 2,2-dimethyl-3-(dimethylamino)propyl-, 1-cyclohexyl-4-(diethylamino)butyl-, aminoethyl-, aminopropyl-, aminobutyl-, aminopentyl-, aminohexyl-, aminoethoxyethyl-, isopropylaminopropyl-, diisopropylaminoethyl-, 1-methyl-4-(diethylamino)butyl-, (t-Boc)aminopropyl-, hydroxyphenyl-

, benzyl, methoxyphenyl-, methylmethoxyphenyl-, dimethylphenyl-, tolyl, ethylphenyl-, (oxopyrrolidinyl)propyl-, (methoxycarbonyl)ethyl-, benzylpiperidinyl-, pyridinylethyl-, pyridinylmethyl-, morpholinylethyl-, morpholinylpropyl-, piperidinyl-, azetidinylmethyl-, azetidinyethyl-, azetidinypropyl-, pyrrolidinylethyl-, pyrrolidinylpropyl-, piperidinylmethyl-, piperidinylethyl-, imidazolylpropyl-, imidazolylethyl-, (ethylpyrrolidinyl)methyl-, (methylpyrrolidinyl)ethyl-, (methylpiperidinyl)propyl-, (methylpiperazinyl)propyl-, furanylmethyl- and indolylethyl.

32. A compound according to claim 31, wherein R_4 is aminoethyl-, aminopropyl-, aminobutyl-, aminopentyl-, aminoethyl-, methylaminoethyl-, methylaminopropyl-, methylaminobutyl-, methylaminopentyl-, methylaminoethyl-, dimethylaminoethyl-, dimethylaminopropyl-, dimethylaminobutyl-, dimethylaminopentyl-, dimethylaminoethyl-, ethylaminoethyl-, ethylaminopropyl-, ethylaminobutyl-, ethylaminopentyl-, ethylaminoethyl-, diethylaminoethyl-, diethylaminopropyl-, diethylaminobutyl-, diethylaminopentyl-, or diethylaminoethyl.

33. A compound according to claim 1, wherein R_{12} is an imidazole and R_{12} has the formula:




wherein

R_9 is chosen from hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl- C_1 - C_4 -alkyl-, optionally substituted aryl- C_1 - C_4 -alkoxy, optionally substituted heteroaryl- C_1 - C_4 -alkoxy, optionally substituted heteroaryl; and

R_{13} and $R_{13'}$ are independently hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally

substituted aryl-, or optionally substituted aryl-C₁-C₄-alkyl.

34. A compound according to claim 33, wherein R₉ is phenyl substituted with C₁-C₄-alkyl, C₁-C₄-alkoxy-, and/or halo; phenyl; or benzyl.
35. A compound according to claim 34, wherein R₉ is tolyl; halophenyl; or halomethylphenyl.
36. A compound according to claim 33, wherein R₁₃ is hydrogen and R_{13'} is substituted C₁-C₄ alkyl.
37. A compound according to claim 36, wherein R₁₃ is hydrogen and R_{13'} is aminomethyl-, aminoethyl-, aminopropyl-, acetylamino-methyl-, acetylaminoethyl-, benzyloxycarbonylamino-methyl- or benzyloxycarbonylamino-ethyl-.
38. A compound according to claim 1, wherein R₁₂ is an imidazoline and R₁₂ has the formula



wherein,

R₉ is chosen from hydrogen, optionally substituted C₁-C₈ alkyl-, optionally substituted aryl-, optionally substituted aryl-C₁-C₄-alkyl-, and optionally substituted heteroaryl; and

R₁₀, R_{10'}, R₁₄, and R_{14'} are independently chosen from hydrogen, optionally substituted C₁-C₈ alkyl-, optionally substituted aryl-, and optionally substituted aryl-C₁-C₄-alkyl.

39. A compound according to claim 38, wherein R₉ is methylenedioxyphenyl; phenyl;


C₁-C₄ alkyl-, C₁-C₄ alkoxy-, and/or halo-substituted phenyl; or benzyl.

40. A compound according to claim 39, wherein R₉ is methylenedioxyphenyl; phenyl; tolyl; methoxyphenyl; halophenyl; or halomethylphenyl.

41. A compound according to claim 38, wherein R₁₀, R_{10'}, R_{14'}, and R₁₄ are independently hydrogen or optionally substituted C₁-C₄ alkyl.

42. A compound according to claim 41, wherein R_{14'} and R₁₄ are hydrogen.

43. A compound having the structure:



wherein

R₁ is benzyl, halobenzyl, methoxybenzyl-, cyanobenzyl, or naphthalenylmethyl-;

R₂ is ethyl or propyl;

R_{2'} is hydrogen;

R₅ is hydrogen;

R₆ is hydrogen;

R₇ is halo, cyano, methoxy or hydrogen;

R₈ is hydrogen; and

R₁₂ is -NR₄(COR₃) wherein R₃ is optionally substituted aryl and R₄ is R₁₆-alkylene- wherein R₁₆ is hydroxyl, di(C₁-C₄)alkylamino-, (C₁-C₄ alkyl)amino-, amino, pyrrolidino,

piperidino, imidazolyl and morpholino.

44. A compound according to claim 43, wherein

R_1 is benzyl, halobenzyl, methoxybenzyl-, cyanobenzyl, or naphthalenylmethyl-;

R_2 is propyl;

R_2 is hydrogen;

R_5 is hydrogen;


R_6 is hydrogen;

R_7 is halo, cyano, methoxy or hydrogen;

R_8 is hydrogen; and

R_{12} is $-NR_4(COR_3)$ wherein R_4 is aminopropyl and R_3 is tolyl.

45. A compound having the structure:



wherein

R_1 is benzyl, halobenzyl, methoxybenzyl-, cyanobenzyl, or naphthalenylmethyl;

R_2 is ethyl or propyl;

R_2 is hydrogen;


R_5 is hydrogen;

R_6 is hydrogen;

R_7 is halo, cyano, methoxy or hydrogen;


R_8 is hydrogen; and

R_{12} is optionally substituted imidazolyl having the structure



wherein R_{10} , $R_{10'}$, R_{14} and $R_{14'}$ are independently hydrogen or optionally substituted C_1 - C_4 alkyl; and R_9 is optionally substituted phenyl.

46. A compound having the structure:



wherein

R_1 is benzyl, halobenzyl, methoxybenzyl-, cyanobenzyl, or naphthalenylmethyl;

R_2 is ethyl or propyl;

$R_{2'}$ is hydrogen;


R_3 is hydrogen;

R_4 is hydrogen;

R_5 is halo, cyano, methoxy or hydrogen;

R_6 is hydrogen; and

R_{12} is optionally substituted imidazole, having the structure



wherein R_{13} is hydrogen and $R_{13'}$ is hydrogen or optionally substituted alkyl; and R_9 is optionally substituted aryl.

47. A compound according to claim 1 wherein R_2 and $R_{2'}$ are each attached to a stereogenic center having an R-configuration, or a pharmaceutically acceptable salt or solvate thereof.
48. A compound according to claim 1 that is a hydrochloric, phosphoric, or oxalic acid salt.
49. A composition comprising a pharmaceutical excipient and a compound, salt, or solvate thereof of any one of claims 1, 2, 3-9, or 10-48.
50. A composition according to claim 49, wherein said composition further comprises a chemotherapeutic agent other than a compound of Formula I or a pharmaceutical salt or solvate thereof.
51. A composition according to claim 50, wherein said composition further comprises a taxane.
52. A composition according to claim 50, wherein said composition further comprises a vinca alkaloid.
53. A composition according to claim 50, wherein said composition further comprises a topoisomerase I inhibitor.

54. A compound according to any one of claims 1 to 53 for use in the treatment of a cellular proliferative disease.
55. A compound according to claim 54, wherein said disease is selected from the group consisting of cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders, and inflammation.
56. A compound according to any one of claims 1 to 53 for use in the treatment of a disorder associated with KSP kinesin activity.
57. Use of a compound according to any one of claims 1 to 53 or a pharmaceutically acceptable salt or solvate thereof in the manufacture of a medicament for the treatment of a cellular proliferative disease.
58. Use of a compound according to claim 57, wherein said disease is selected from the group consisting of cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders, and inflammation.
59. Use of a compound according to claim 57 or 58 in the manufacture of a medicament for the treatment of a disorder associated with KSP kinesin activity.