


POLYSUBSTITUTED DERIVATIVES OF 2-ARYL-6-PHENYL-
IMIDAZO[1,2-a]PYRIDINES, AND PREPARATION AND THERAPEUTIC
USE THEREOF

Abstract

Compounds of formula (I):



in which: R_1 represents: a phenyl group or a naphthyl group, optionally substituted by one or more atoms or groups; X represents from 1 to 4 substituents, identical to or different from one another, chosen from halogen, (C_1-C_{10}) alkyl, halo (C_1-C_{10}) alkyl, (C_1-C_{10}) alkoxy, NRaRb, cyano, nitro, it being possible for the (C_1-C_{10}) alkyl to be optionally substituted by halogen or an NRaRb or hydroxyl group; R_2 and R_3 represent, independently of one another, a hydrogen atom or a (C_1-C_{10}) alkyl group, this group being optionally substituted by an Rf group, or an aryl group, the aryl being optionally substituted by one or more substituents chosen from a halogen or a (C_1-C_{10}) alkyl, halo (C_1-C_{10}) alkyl, (C_1-C_{10}) alkoxy, halo (C_1-C_{10}) alkoxy, NRaRb, hydroxyl, nitro or cyano group; R_2 and X can together form, with the carbon atoms which carry them, a carbon ring of 5 to 7 carbon atoms; R_4 represents a hydrogen atom, a (C_1-C_{10}) alkyl group, optionally substituted by an Rf group, or an optionally substituted aryl group, in the form of the base or of an addition salt with an acid.

Therapeutic use and process of synthesis.