RECOMBINANT NEWCASTLE DISEASE VIRUS AND PREPARATION METHOD THEREFOR, RECOMBINANT PLASMID, AND USE THEREOF

Technical Field

[0001] The present application in general relates to the field of biotechnologies, in particular to a recombinant Newcastle disease virus and a preparation method therefor, a recombinant plasmid, and use thereof.

Background

[0002] Major means for treating cancers include surgeries, radiotherapies, chemotherapies, immunotherapies, monoclonal antibody therapies, and viral vaccines.

[0003] Malignant tumours are one of the major diseases endangering human's health. According to the data in 2015 Global Cancer Statistics, there were about 14.1 million new cancer cases and up to 8.2 million cancer deaths across the world in 2015. It is predicted that by 2025, the number of new cancer cases across the world every year will reach up to 24.49 million. For a long time, the methods adopted by people to treat cancers mainly include traditional surgical resections, radiotherapies, and chemotherapies. These methods can control the development of tumours to some extent, but showing limited efficacy for patients with advanced tumour spread, and at the same time, causing severe trauma to the normal cells of a human body. Therefore, there is an urgent need of a new method for treating a malignant tumour. An oncolytic virus therapy is a novel tumour treatment method by which a tumour is killed by selectively infecting tumour cells with viruses. As an important part of tumour immunity, it has a great advantage as compared with other tumour immunotherapies. The oncolytic virus therapy has undergone three stages of development, and has achieved great results. At present, the anti-tumour effect can be enhanced by carrying exogenous genes, among which T-vec carrying GM-CSF exhibits a good efficacy and was approved by the US FDA in 2015 for the local treatment of an unresectable melanoma that recurred for the first time. In December of

the same year, T-vec was further approved by the European Union, indicating a great development potential of oncolytic viruses.

[0004] In 1950s, it was found that Newcastle disease viruses (NDV) were capable of inhibiting the metastasis of gastric cancer. Therefore, NDV can be used as an emerging tumour bio-therapeutic factor, and its application to tumour treatment has been studied deeper and deeper due to its high targeting property and the maturity of molecular biotechnologies. As a single negative-stranded RNA virus, NDV encodes six types of structural proteins (L, NP, P, HN, F, and M).

[0005] Compared with other oncolytic viruses, the Newcastle disease virus has a natural targeting property against tumour cells, and broad-spectrum anti-tumour properties, showing greater advantages in the anti-tumour aspect. However, how to improve its oncolytic effect and to ensure its safety is still a huge challenge for us to be confronted with.

Summary

[0006] An object of the present application is to provide a recombinant Newcastle disease virus and a preparation method therefor, a recombinant plasmid, and use thereof.

[0007] The present application provides a recombinant Newcastle disease virus obtained by replacing an F protein of a Newcastle disease virus lasota~~Losata~~ with an F protein of a Newcastle disease virus mesogenic strain.

[0008] Optionally, according to the recombinant Newcastle disease virus mentioned above, a coding DNA of an F protein of the recombinant Newcastle disease virus is as set forth in SEQ ID NO: 1 at positions 7274-8935 from a 5'-terminus.

[0009] Optionally, according to the recombinant Newcastle disease virus mentioned above, the recombinant Newcastle disease virus is obtained by replacing

an F gene in a genome of the Newcastle disease virus ~~Losata~~lasota with an F gene of the Newcastle disease virus mesogenic strain. Preferably, a DNA corresponding to the F gene of the recombinant Newcastle disease virus is as set forth in SEQ ID NO: 1 at the positions 7274-8935 from the 5'-terminus. More preferably, a DNA corresponding to a genome of the recombinant Newcastle disease virus is as set forth in SEQ ID NO: 1.

[0010] Optionally, according to the recombinant Newcastle disease virus mentioned above, a DNA corresponding to a genome of the recombinant Newcastle disease virus further comprises an exogenous gene, which is selected from one or more of a group consisting of DR5, TRAIL, hIL2, P53, PD1, CD, and mIL12.

[0011] Preferably, the exogenous gene is DR5, TRAIL, hIL2, P53, PD1, CD, or mIL12.

[0012] Preferably, the exogenous gene is mIL12-hIL2, P53-hIL2, PD1-hIL2, or DR5-TRAIL.

[0013] Preferably, the DR5 is as set forth in SEQ ID NO: 2, the TRAIL is as set forth in SEQ ID NO: 3, the hIL2 is as set forth in SEQ ID NO: 4, the P53 is as set forth in SEQ ID NO: 5, the PD1 is as set forth in SEQ ID NO: 6, the CD is as set forth in SEQ ID NO: 7, and the mIL12 is as set forth in SEQ ID NO: 8.

[0014] Preferably, the exogenous gene is located between a P gene and an M gene of the recombinant Newcastle disease virus.

[0015] The present application further provides a recombinant plasmid obtained by replacing an F gene in a pBr~~Losata~~lasota plasmid, wherein the F gene of the recombinant plasmid is as set forth in SEQ ID NO: 1 at positions 7274-8935 from a 5'-terminus.

[0016] Optionally, according to the recombinant plasmid mentioned above,

the recombinant plasmid is a DNA molecule plasmid as set forth in SEQ ID NO: 1.

[0017] Optionally, according to the recombinant plasmid mentioned above, the recombinant plasmid further comprises an exogenous gene, which is selected from one or more of a group consisting of DR5, TRAIL, hIL2, P53, PD1, CD, and mIL12.

[0018] Preferably, the exogenous gene is DR5, TRAIL, hIL2, P53, PD1, CD, or mIL12.

[0019] Preferably, the exogenous gene is mIL12-hIL2, P53-hIL2, PD1-hIL2, or DR5-TRAIL.

[0020] Preferably, the DR5 is as set forth in SEQ ID NO: 2, the TRAIL is as set forth in SEQ ID NO: 3, the hIL2 is as set forth in SEQ ID NO: 4, the P53 is as set forth in SEQ ID NO: 5, the PD1 is as set forth in SEQ ID NO: 6, the CD is as set forth in SEQ ID NO: 7, and the mIL12 is as set forth in SEQ ID NO: 8.

[0021] Preferably, the exogenous gene is located between a P gene and an M gene of a genome of the recombinant Newcastle disease virus.

[0022] The present application further provides a preparation method for a recombinant Newcastle disease virus, comprising transfecting the recombinant plasmid mentioned above to cells or cell line for culturing to obtain the recombinant Newcastle disease virus.

[0023] Optionally, according to the preparation method mentioned above, the recombinant plasmid and a helper plasmid are co-transfected to cells or cell line for culturing. The cell is a mammalian cell. The helper plasmid is, for example, a pBL-N plasmid, a pBL-P plasmid, or a pBL-L plasmid.

[0024] The present application further provides use of the recombinant Newcastle disease virus mentioned above or the recombinant plasmid mentioned

above in preparation of a drug. The drug has the following function(s) of (a) and/or (b) and/or (c): (a) treating a tumour; (b) inhibiting tumour cell proliferation; and (c) killing tumour cells.

[0025] The present application further provides a drug, comprising the recombinant Newcastle disease virus mentioned above, and /or the recombinant plasmid mentioned above. The drug has the following function(s) of (a) and/or (b) and/or (c): (a) treating a tumour; (b) inhibiting tumour cell proliferation; and (c) killing tumour cells.

[0026] Optionally, the tumour is selected from one or more of a group consisting of a liver cancer, a breast cancer, a non-small cell lung cancer, a melanoma, a neuroblastoma, a lung cancer, a pancreatic cancer, a thyroid cancer, a kidney cancer, a glioma, a myosarcoma, an esophageal cancer, a uterine cancer, ~~a breast cancer,~~ and a colorectal cancer.

[0027] Optionally, the tumour cell is selected from one or more of a group consisting of a liver cancer cell, a breast cancer cell, a non-small cell lung cancer cell, a melanoma cell, a neuroblastoma cell, a lung cancer cell, a pancreatic cancer cell, a thyroid cancer cell, a kidney cancer cell, a glioma cell, a myosarcoma cell, an esophageal cancer cell, an uterine cancer cell, ~~a breast cancer cell,~~ and a colorectal cancer cell.

[0028] In the present application, virus strains having a stronger effect of promoting cell fusion are screened out; the structural protein genes of the strains are acquired; structural protein genes of ~~Losata~~lasota are replaced by means of a reverse genetic manipulation technique and a molecular cloning technique; and exogenous genes are further inserted to successfully construct and rescue recombinant chimeric viruses. The recombinant chimeric viruses are continuously passaged in chicken embryos for 10 generations. The results of a proliferation stability experiment have demonstrated that the recombinant Newcastle disease virus could proliferate

continuously and stably in the chicken embryos, and the effects of the recombinant chimeric virus on killing tumour cells are systematically compared.

[0029] The recombinant chimeric virus according to the present application has good safety and is capable of effectively inhibiting tumour cells, promoting tumour cell apoptosis, and effectively treating a tumour.

Brief Description of the Drawings

[0030] FIG. 1 shows the results of a virus strain screening test according to Example 1;

[0031] FIG. 2 shows the results of an exogenous gene expression position determination test according to Example 2;

[0032] FIG. 3A to FIG. 3E show the killing effects of a recombinant chimeric virus against tumour cells as detected in a cell fusion experiment according to Example 4;

[0033] FIG. 4 shows the killing effect of a recombinant chimeric virus against HepG-2 cells as detected by an MTT method according to Example 5;

[0034] FIG. 5 shows the killing effect of the recombinant chimeric virus against Caki-1 cells as detected by the MTT method according to Example 5;

[0035] FIG. 6 shows the killing effect of the recombinant chimeric virus against A549 cells as detected by the MTT method according to Example 5;

[0036] FIG. 7 shows the killing effect of the recombinant chimeric virus against SK-MEL-5 cells as detected by the MTT method according to Example 5;

[0037] FIG. 8 shows the killing effect of the recombinant chimeric virus against LN229 cells as detected by an MTT method according to Example 5;

[0038] FIG. 9 shows the killing effect of the recombinant chimeric virus against liver cancer cells as detected by an Annexin V/PI method according to Example 5;

[0039] FIG. 10 shows the statistics of tumour volume after H22 tumour-bearing mice are treated with a recombinant chimeric virus according to

Example 6;

[0040] FIG. 11 shows the statistics of tumour weight after the H22 tumour-bearing mice are treated with the recombinant chimeric virus according to Example 6;

[0041] FIG. 12 shows the statistics of survival rate of the H22 tumour-bearing mice treated with the recombinant chimeric virus according to Example 6; and

[0042] FIG. 13 shows the pathological observations of tumour tissues after the H22 tumour-bearing mice are treated with the recombinant chimeric virus according to Example 6.

Detailed Description of Embodiments

[0043] Specific embodiments of the present application will be illustrated below in further detail in combination with accompanying drawings and examples for a better understanding of the solutions of the present application as well as the advantages thereof in various aspects. Nevertheless, the specific embodiments and examples described below are merely for an illustrative purpose, and are not intended to limit the present application.

[0044] The experimental methods in the examples below are conventional methods, unless otherwise stated. Test materials used in the examples below are purchased from common biochemical reagent stores, unless otherwise stated. For quantitative tests in the examples below, three repeated experiments are set up, and the results are averaged. Unless otherwise specified in the following examples, the technical means used in these examples are conventional means well-known to a person skilled in the art or common commercially-available instruments and reagents. References can be made to *Molecular Cloning: A Laboratory Manual* (Third Edition) (Science Press), *Microbiological Experiments* (Fourth Edition) (Higher Education Press), and the manufacturer's instructions for use for the corresponding instruments

and reagents, etc.

[0045] A reference mentioning "pBr~~Losata~~lasota plasmid" is "Identification of Optimal Insertion Site in Recombinant Newcastle Disease Virus (rNDV) Vector Expressing Foreign Gene to Enhance Its Anti-Tumour Effect" (Ziye Pan, PLOS ONE.; Harbin Boxiang Medical Technology Development Co., Ltd.).

[0046] A reference mentioning "pBL-N plasmid", "pBL-P plasmid", and "pBL-L plasmid" is "Genetically engineered Newcastle disease virus expressing interleukin 2 is a potential drug candidate for cancer immunotherapy" (Fuliang Bai, Immunology letters; Harbin Boxiang Medical Technology Development Co., Ltd.).

[0047] The pBr~~Losata~~lasota plasmid contains the NP, P, M, F, HN, and L genes of the Newcastle disease virus, with PmeI and SacII digestion recognition sites between the P gene and the M gene. The pBr~~Losata~~lasota, pBL-N, pBL-P, and pBL-L plasmids are co-transfected to mammalian cells and cultured (with the pBL-N, pBL-P, and pBL-L plasmids as helpers, and the pBr~~Losata~~lasota plasmid providing a virus whole-genome) to obtain a virus strain ~~Losata~~lasota.

[0048] The exogenous genes involved herein include a death receptor gene *DR5*, a tumour necrosis factor-related apoptosis-inducing ligand gene *TRAIL*, a human interleukin gene *2hIL2*, a cancer suppressor gene *P53*, a programmed death receptor 1 gene *PD1*, a suicide gene *CD*, and a murine interleukin gene *12mIL12*.

[0049] pMD19-T: purchased from TaKaRa Engineering (Dalian) Co., Ltd. (Dalian TaKaRa Bio Inc.).

[0050] BHK-21 cells (baby hamster kidney cells): purchased from ATCC.

[0051] Liver cancer cells HepG2, kidney cancer cells Caki-1, non-small cells lung cancer cells A549, neuroblastoma cells LN229, melanoma cells SK-MEL5, liver cancer cells HCCLM3, melanoma cells SK-MEL2, pancreatic cancer cells

ASPC-1, thyroid cancer cells FTC133, kidney cancer cells 786-O, neuroblastoma cells U87-MG, myosarcoma cells HT1080, esophageal cancer cells KYSE-150, uterine cancer cells Hela, breast cancer cells MCF7, and colorectal cancer cells HCT116 were purchased from the Cell Resource Center of Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences.

[0052] High-glucose DMEM medium, trypsin, new born calf serum (NBCS), and fetal bovine serum (FBS): purchased from GIBCO.

[0053] SPF chicken embryo: purchased from Harbin Veterinary Research Institute, Chinese Academy of Agricultural Sciences.

[0054] Balb/c mice (Kunming mice): purchased from Vital River Laboratories, Beijing.

[0055] Example 1: Virus strain screening

[0056] Chicken embryo fibroblasts in a logarithmic growth phase were digested with trypsin and collected, and then seeded into a 6-well plate. Three virulent Newcastle disease virus strains F48E9, Turkey/US(ND)/43084/92 (TkND) and Game fowl/US(CA)/211472/02 (CA02) and one mesogenic strain Anhinga (the strains were donated by Southeast Poultry Research Laboratory and stored in our laboratory, and a reference mentioning the virus strains included "Evaluation of Newcastle disease virus chimeras expressing the Hemagglutinin-Neuraminidase protein of velogenic strains in the context of a mesogenic recombinant virus backbone". Virus Research, Volume 129, Issue 2, November 2007, Pages 182-190) infected cells at 0.1 MOI respectively, and a control group was a virus-free SPF chicken embryo allantoic fluid. After 24 h, the 6-well plate were taken out and observed under an inverted fluorescence microscope for a cell fusion effect. By analysing a syncytial formation capability, virus strains having strong fusion capability and weak virulence were screened out for subsequent experiments. The results were shown in FIG. 1. Among

the four virus strains, F48E9 showed the strongest fusion-promoting capability, and the strain Anhinga shows a stronger fusion-promoting capability and the weakest virulence.

[0057] Example 2: Preparation of recombinant chimeric virus

[0058] I. Construction of recombinant plasmid (taking F gene replacement as an example)

[0059] 1. The virus strain Anhinga with a stronger fusion-promoting effect and weaker virulence was screened out, and the sequence of its F gene was acquired from NCBI (GenBank: AAS67135.1), and overlap PCR primers were designed.

[0060] 2. Taking a pAnh plasmid (the involved reference was "Evaluation of Newcastle disease virus chimeras expressing the Hemagglutinin-Neuraminidase protein of velogenic strains in the context of a mesogenic recombinant virus backbone") as a template, an F gene fragment was amplified by PCR by using F gene specific primers, and linked to pMD19-T; the linking product was transformed into Escherichia coli DH5α; the plasmid was extracted and identified; and the correctly identified pMD19-T-F plasmid was used for the next experiment.

[0061] 3. With the PCR technique, the target fragments A and B were amplified by using the pBr~~Losata~~lasota plasmid as a template (primers for the fragment A: 5' CACGTG AAAGCGCCAGAGAAGATTCCCGGGA 3'; P2: 5' TTTGGGGCCCATCTTGCACCTGGAGGGCGCCAACCGG 3'; P3: 5' TCCAGGTGCAAGATGGGCCCCAAACCCCCCACCGGAA 3'. B fragment primers: P6: 5' CCTCATCTGTGT TCATGTTTTTGTGGTGGCTCTCATC 3'; P7: 5' ACAAAAACATGAACACAGATGAGGAACGAAGGTTTCC 3'; and P8: 5' GGCCTGAGAGGCCACGCGTCGCCGCGGGCCGGTT 3'). The target fragment F was amplified by using the pMD19-T-F plasmid as a template, and the fragments A, F and B were extracted by gel. Then, the fragment A-F-B of interest was amplified by

the overlap PCR method using the purified fragments A, F and B as templates, and linked to a pMD19-T vector; the linking product was transformed into Escherichia coli DH5α; plasmids were extracted and identified; and the correctly identified plasmids were named as pMD19-T-A-F-B, which carry two digestion sites PmlI and SfiI.

[0062] 4. The plasmids pMD19-T-A-F-B from step 3 were digested using restriction enzymes PmlI and SfiI, and the cleaved product was extracted by gel.

[0063] 5. The pBr~~Losata~~lasota plasmid was digested using the restriction enzymes PmlI and SfiI, and the vector was extracted by gel.

[0064] 6. The digested product from step 4 and the vector from step 5 were linked to obtain a recombinant chimeric plasmid pBr~~Losata~~lasota-F. PCR and double digestion identification were performed on the recombinant chimeric plasmid; the correctly identified plasmid samples were sub-packaged and sent to Shanghai Sangon for sequencing. The sequencing results were aligned using sequence analysis software DNAMAN; and the plasmid with the sequencing result consistent with a sequence of interest was named as pBr~~Losata~~lasota-F. The nucleotide sequence of the recombinant chimeric plasmid pBr~~Losata~~lasota-F was as set forth in SEQ ID NO:1 of the sequence listing (the nucleotide sequence at positions 7274-8935 from the 5'-terminus in SEQ ID NO: 1 was the nucleotide sequence of the F gene).

[0065] Other recombinant chimeric plasmids were constructed in the same way: pBr~~Losata~~lasota-HN (HN gene sequence, GenBank: AAS67136.1) pBr~~Losata~~lasota-P (P gene sequence in GenBank: AAS67133.1), pBr~~Losata~~lasota-M (M gene sequence in GenBank: AAS67134.1).

[0066] II. Determination of optimal insertion position for exogenous gene

[0067] 1. Restriction Enzyme cutting –sites were respectively introduced to the NP/P, P/M, M/F, and F/HN sites of the pBr~~Losata~~lasota-F plasmid using a

molecular biology method. Recombinant plasmids pBr~~Losata~~lasota-F-NP/P (AscI, SfiI), pBr~~Losata~~lasota-F-P/M (SacII, PmeI), pBr~~Losata~~lasota-F-M/F (HpaI, MluI), and pBr~~Losata~~lasota-F-F/HN (HpaI, MluI) were obtained.

[0068] 2. RFP gene fragments carrying different restriction enzyme cutting sites were amplified by PCR using specific primers for a red fluorescent protein RFP gene (red fluorescent protein RFP gene sequence in GenBank: MG719763.1), and linked to a pMD19-T vector; the linking product was transformed into Escherichia coli DH5α; and the plasmid was extracted and identified as carrying different restriction enzyme cutting sites.

[0069] 3. The plasmid from step 2 was digested using a restriction enzyme, and the cleaved product was extracted by gel.

[0070] 4. The recombinant plasmid from step 1 was digested using a restriction enzyme, and the vector was extracted by gel.

[0071] 5. The digested product from step 3 and the vector from step 4 were linked to obtain recombinant plasmids pBr~~Losata~~lasota-F-NP/P-RFP, pBr~~Losata~~lasota-F-P/M-RFP, pBr~~Losata~~lasota-F-M/F-RFP, and pBr~~Losata~~lasota-F-F/HN-RFP.

[0072] 6. BHK-21 cells were co-transfected with the recombinant plasmid from step 5 and the helper plasmid to obtain recombinant virus r~~Losata~~lasota-F-NP/P-RFP, r~~Losata~~lasota-F-P/M-RFP, r~~Losata~~lasota-F-M/F-RFP, and r~~Losata~~lasota-F-F/HN-RFP.

[0073] 7. MCF7 cells in the logarithmic growth phase were digested with trypsin and then collected, and seeded into a 6-well plate. The cells were respectively infected with the recombinant chimeric viruses r~~Losata~~lasota-F-NP/P-RFP, r~~Losata~~lasota-F-P/M-RFP, r~~Losata~~lasota-F-M/F-RFP, and r~~Losata~~lasota-F-F/HN-RFP at 1 MOI, the parental virus r~~Losata~~lasota was used as a control, and the control group

was virus-free SPF chicken embryo allantoic fluid. After 24 h, the 6-well plate was taken out, and the average fluorescence intensity of the cells infected with respective virus was detected by the flow cytometry. The results were shown in FIG. 2, showing the strongest red fluorescence intensity between P and M sites, which indicates the highest RFP expression level between the P and M sites.

[0074] III. Construction of recombinant plasmid inserted with exogenous gene (taking PD1 gene insertion as an example)

[0075] 1. Monocytes were isolated from human peripheral blood cells, and RNA was extracted and reverse-transcribed into cDNA. According to the human PD1 sequence (AY238517.1) that has been published in GenBank, PCR primers P1 and P2 were designed using Primer Premier 5.0 software to clone the complete PD1 gene. Pme I and Sac II digestion sites were designed at both ends of the primers to match the pBr~~Losata~~lasota-F vector plasmid.

[0076] 2. The PCR product from step 1 was linked to pMD19-T and transformed into Escherichia coli; plasmids were extracted and identified; and the correctly identified pMD19-T-PD1 plasmids were used for the next experiment.

[0077] 3. The pMD19-PD1 plasmids from step 2 were digested using restriction enzymes Pme I and Sac II, and the cleaved product was extracted by gel.

[0078] 4. The pBr~~Losata~~lasota-F plasmids were digested using restriction enzymes Pme I and Sac II, and the vector was extracted by gel.

[0079] 5. The cleaved product from step 3 was linked to the vector from step 4 to obtain a recombinant chimeric plasmid pBr~~Losata~~lasota(F)-PD1, with the PD1 gene inserted between the P and M genes of the plasmid. PCR as well as Pme I and Sac II double digestion identification were performed on the recombinant chimeric plasmid; the correctly identified plasmid samples were sub-packaged and sent to Shanghai Sangon for sequencing. The sequencing results were aligned using

sequence analysis software DNAMAN; and the sequenced result was consistent with a sequence of interest. The PD1 nucleotide sequence of the recombinant chimeric plasmid pBr~~Losata~~lasota(F)-PD1 was as set forth in SEQ ID NO. 6 in the sequence listing.

[0080] Other recombinant chimeric plasmids inserted with exogenous genes were constructed using the same method. The constructed recombinant chimeric plasmids were specifically as follows: pBr~~Losata~~lasota(F)-DR5 (a DR5 sequence as set forth in SEQ ID NO: 2), pBr~~Losata~~lasota(F)-TRAIL (a TRAIL sequence as set forth in SEQ ID NO: 3), pBr~~Losata~~lasota(F)-hIL2 (a hIL2 sequence as set forth in SEQ ID NO: 4), pBr~~Losata~~lasota(F)-P53 (a P53 sequence as set forth in SEQ ID NO:5), pBr~~Losata~~lasota(F)-CD (a CD sequence as set forth in SEQ ID NO: 7), pBr~~Losata~~lasota(F)-mIL12 (a mIL12 sequence as set forth in SEQ ID NO: 8), pBr~~Losata~~lasota(F)-DR5-TRAIL (i.e., a DR5-TRAIL combination as the exogenous gene), pBr~~Losata~~lasota(F)-mIL12-hIL2 (i.e., a mIL12-hIL2 combination as the exogenous gene), pBr~~Losata~~lasota(F)-P53-hIL2 (i.e., a P53-hIL2 combination as the exogenous gene), and pBr~~Losata~~lasota(F)-PD1-hIL2 (i.e., a PD1-hIL2 combination as the exogenous gene).

[0081] IV. Preparation of recombinant chimeric virus

[0082] 1. BHK-21 cells were co-transfected respectively with the plasmid pBr~~Losata~~lasota, the recombinant chimeric plasmids pBr~~Losata~~lasota-HN, pBr~~Losata~~lasota-F, pBr~~Losata~~lasota-P, pBr~~Losata~~lasota-M, pBr~~Losata~~lasota(F)-DR5, pBr~~Losata~~lasota(F)-TRAIL, pBr~~Losata~~lasota(F)-hIL2, pBr~~Losata~~lasota(F)-P53, pBr~~Losata~~lasota(F)-PD1, pBr~~Losata~~lasota(F)-CD, pBr~~Losata~~lasota(F)-mIL12, pBr~~Losata~~lasota(F)-DR5-TRAIL, pBr~~Losata~~lasota(F)-mIL12-hIL2, pBr~~Losata~~lasota(F)-P53-hIL2 and pBr~~Losata~~lasota(F)-PD1-hIL2, together with the helper plasmids pBL-NP, pBL-P and pBL-L (about 1×10^6 cells transfected with 2 μg of recombinant chimeric plasmids, 1 μg of pBL-N plasmids, 0.5 μg of pBL-P plasmids and 0.25 μg of pBL-L plasmids). The transfected cells were placed in a 5%

CO_2 environment at 37℃ for static culture for 72 h.

[0083] 2. The transfected cells obtained in step 1 were repeatedly frozen and thawed three times, centrifuged to collect the cell supernatant, then seeded into the allantoic cavities of 9-11-day-old SPF chicken embryos. The chicken embryo allantoic cavities were cultured for 72 h in an environment at 37°C, and the chicken embryo allantoic fluid was collected for HA detection.

[0084] 3. The HA-positive chicken embryo allantoic liquid obtained in step 2 was passaged, seeded into the allantoic cavities of fresh 9-11-day-old SPF chicken embryos, and cultured for 72 h in an environment at 37°C. The chicken embryo allantoic liquid was collected for HA detection. The HA-positive chicken embryo allantoic liquid was passaged for 10 generations, and the proliferation stability of the recombinant viruses was detected.

[0085] 4. The chicken embryo allantoic liquid obtained from step 3 was taken for mass reproduction of the viruses, then seeded into the allantoic cavities of new 9-11-day-old SPF chicken embryos, and cultured for 72 h in an environment at 37°C. The chicken embryo allantoic liquid was collected for HA detection. The allantoic liquid with the HA titer greater than 2^9 was selected, mixed and sub-packaged for later use.

[0086] 5. The allantoic liquid was collected from each group for Western Blot to detect the expression of each exogenous gene. The results showed that, compared with the r~~Losata~~lasota group, the recombinant chimeric viruses expressing the exogenous gene was capable of stably expressing the exogenous gene.

[0087] In the following description, unless otherwise specified, the recombinant chimeric viruses, the chimeric viruses and the recombinant viruses refer to the recombinant chimeric viruses prepared by the above methods. r~~Losata~~lasota-HN, r~~Losata~~lasota-F, r~~Losata~~lasota-P, r~~Losata~~lasota-M,

rLosatalasota(F)-DR5, rLosatalasota(F)-TRAIL, rLosatalasota(F)-hIL2, rLosatalasota(F)-P53, rLosatalasota(F)-PD1, rLosatalasota(F)-CD, rLosatalasota(F)-mIL12, rLosatalasota(F)-DR5-TRAIL, rLosatalasota(F)-mIL12-hIL2, rLosatalasota(F)-P53-hIL2 and rLosatalasota(F)-PD1-hIL2 respectively refer to the recombinant chimeric viruses prepared with the above methods by using the recombinant chimeric plasmids pBrLosatalasota-HN, pBrLosatalasota-F, pBrLosatalasota-NP, pBrLosatalasota-M, pBrLosatalasota(F)-DR5, pBrLosatalasota(F)-TRAIL, pBrLosatalasota(F)-hIL2, pBrLosatalasota(F)-P53, pBrLosatalasota(F)-PD1, pBrLosatalasota(F)-CD, pBrLosatalasota(F)-mIL12, pBrLosatalasota(F)-DR5-TRAIL, pBrLosatalasota(F)-mIL12-hIL2, pBrLosatalasota(F)-P53-hIL2 and pBrLosatalasota(F)-PD1-hIL2.

[0088] In the following description, unless otherwise specified, the parental viruses refer to the Newcastle disease viruses rLosatalasota and Anhinga.

[0089] Example 3: Determination of virulence and pathogenicity of recombinant chimeric viruses

[0090] I. Determination of EID50

[0091] The rescued recombinant chimeric viruses or parental viruses were serially diluted 10 times with sterilized PBS respectively. Chicken embryos were taken out from an incubator at 37°C. Six 9-11-day-old SPF chicken embryos were seeded with the viruses at each dilution (100 μL for each), and incubated in the incubator at 37°C for 4-6 days. Each chicken embryo needed to be detected by HA assay. The number of embryos infected at each dilution was recorded. The EID50 of each of the recombinant chimeric viruses or parental viruses was calculated according to a Reed and Muench method.

[0092] II. Determination of TCID50

[0093] 1. DF-1 cells were seeded into a 96-well microplate by using a

DMEM medium containing 10% FBS and 1% antibiotics, and held in a 5% CO_2 incubator at 37°C overnight.

[0094] 2. Before the recombinant chimeric viruses and the parental viruses were seeded, the original cell culture medium added in step 1 was discarded, and 180 μL of fresh DMEM culture medium containing 10% allantoic fluid and 1% antibiotics.

[0095] 3. 20 μL of the recombinant chimeric viruses or parental viruses was seeded into the top row of cell wells; the mixture was gently pipetted up and mixed; and 20 μL of the mixture was pipetted to the lower cell wells for 10-fold serial dilution. Three replicates were set up for each virus.

[0096] 4. In the 5% CO_2 cell incubator at 37°C, the recombinant chimeric viruses or parental viruses were seeded and incubated for 1 h; then, the culture medium added in step 3 was discarded, and the residue was washed with PBS; and then, 200 μL of fresh DMEM culture medium containing 10% allantoic fluid and 1% antibiotics was added.

[0097] 5. The culture continued in a 5% CO_2 cell incubator at 37°C. After 72 h, the wells with cytopathy were observed under an optical inverted microscope, the number of wells with cytopathy was recorded, and the titer of the respective virus was calculated according to the Reed and Muench method.

[0098] III. Determination of MDT

[0099] The recombinant chimeric viruses or parental viruses were serially diluted 10 times with sterilized PBS respectively. The viruses subjected to doubling dilution were seeded to six 10-day-old SPF chicken embryos (100 μL for each) via allantoic cavities, and were continuously cultured for 7 days in the incubator at 37°C. The death time and death number of the chicken embryos seeded at each dilution were checked and recorded every 12 hours. Finally, the death times of all chicken embryos

seeded at the highest dilution ratio of the viruses are averaged as the MDT of the viruses in discussion.

[00100] IV. Determination of ICPI

[00101] According to the OIE standard, the recombinant chimeric viruses or parental viruses were first subjected to serial 10-fold dilution with sterilized PBS, and then intracerebrally seeded to ten 1-day-old SPF chicks (50 μL for each). Observations were continuously conducted for 10 days to record the daily morbidity and mortality of the chicks, and the intracerebral pathogenicity index of the viruses was calculated according to the Reed and Muench method.

[00102] The results of the measured virulence and pathogenicity of the recombinant chimeric viruses in Example 3 were shown in Table 1. The ICPI value of the parental virus strain Anhinga was 1.47; the ICPI values of r~~Losata~~lasota-HN, r~~Losata~~lasota-P and r~~Losata~~lasota-M were significantly lower than that of Anhinga, and were 0.05, 0.11, and 0.17 respectively; and the ICPI value of r~~Losata~~lasota-F was ~~1.59~~1.17, which is also lower than that of Anhinga. The results showed that, compared with the parental virus strain Anhinga, the HN, P, M and F replacements could reduce the pathogenicity of Anhinga in chickens. r~~Losata~~lasota-F as a recombinant virus showed higher controllability and safety than Anhinga.

Table 1 Comparison of virus titer, virulence, and pathogenicity

Virus	HA	$TCID_{50}$	EID_{50}	MDT	ICPI
r~~Losata~~lasota	2^9	2.09×10^7	6.67×10^9	>120 h	0.00
Anhinga	2^7	3.24×10^7	4.27×10^8	82 h	1.47
r~~Losata~~lasota-HN	2^7	1.13×10^7	1×10^9	>120 h	0.05
r~~Losata~~lasota-P	2^9	1.1×10^8	3.56×10^9	>120 h	0.11
r~~Losata~~lasota-F	2^9	1.62×10^8	6.67×10^9	96h	1.17
r~~Losata~~lasota- M	2^9	1.16×10^8	2.24×10^9	>120 h	0.17
r~~Losata~~lasota(F)-DR5	2^9	3.67×10^7	3.78×10^9	94h	1.11
r~~Losata~~lasota(F)-TRAIL	2^9	4.6×10^7	2.29×10^9	90h	1.23
r~~Losata~~lasota(F)-hIL2	2^{10}	1.71×10^8	6.67×10^9	96h	1.07
r~~Losata~~lasota(F)-P53	2^9	3.92×10^7	3.37×10^9	96h	1.20
r~~Losata~~lasota(F)-PD1	2^{10}	1.92×10^8	4.92×10^9	94h	1.12
r~~Losata~~lasota(F)-CD	2^9	4.11×10^7	2.12×10^9	98h	1.07
r~~Losata~~lasota(F)-mIL12	2^9	3.16×10^7	3.52×10^9	98h	1.06
r~~Losata~~lasota(F)-DR5-TRAIL	2^9	3.16×10^7	3.61×10^9	92h	1.20
r~~Losata~~lasota(F)-mIL12-hI	2^9	1.61×10^8	4.67×10^9	96h	1.17

L2					
r~~Losata~~lasota(F)-P53-hIL2	2^9	4.11×10^7	3.92×10^9	94h	1.21
r~~Losata~~lasota(F)-PD1-hIL2	2^{10}	1.58×10^8	5×10^9	98h	1.11

[00103] Example 4: Killing effect of recombinant chimeric virus against tumour cells as detected in cell fusion experiment

[00104] The tumour cells refer to liver cancer cells HepG2, kidney cancer cells Caki-1, non-small cell lung cancer cells A549, melanoma cells SK-MEL-5, or neuroblastoma cells LN229.

[00105] I. Infection of tumour cells with recombinant chimeric viruses

[00106] Tumour cells in the logarithmic growth phase were digested with trypsin and then collected, and seeded into a 6-well plate. The cells were infected with the recombinant chimeric viruses r~~Losata~~lasota-HN, r~~Losata~~lasota-F, r~~Losata~~lasota-P, and r~~Losata~~lasota-M at 1 MOI respectively, the parental virus r~~Losata~~lasota was used as a control, and a control group was a virus-free SPF chicken embryo allantoic fluid. After 24 h, the 6-well plate was taken out, and observed under the inverted fluorescent microscope for a cell fusion effect. The results could be found in FIG. 3A to FIG. 3E.

[00107] The results showed that, compared with the parental viruses, the recombinant chimeric viruses r~~Losata~~lasota-HN, r~~Losata~~lasota-P, and r~~Losata~~lasota-M produced more syncytia than r~~Losata~~lasota, but without significant difference; and the chimeric virus r~~Losata~~lasota-F produced more syncytia than r~~Losata~~lasota, with an extremely significant difference.

[00108] Example 5: Effect of recombinant chimeric virus on tumour cells

[00109] The tumour cells refer to liver cancer cells HepG2, kidney cancer cells Caki-1, non-small cell lung cancer cells A549, melanoma cells SK-MEL-5, or neuroblastoma cells LN229.

[00110] I. Detection of inhibitory effect of recombinant chimeric virus on

tumour cells by MTT method

[00111] Tumour cells in the logarithmic growth phase were digested by trypsin, and then, prepared into a cell suspension of 2×10^4 cells/mL by using the DMEM or RPMI-1640 culture medium containing 10% calf serum. The cell suspension was seeded in a 96-well plate (200 μl per well), placed in a 5% CO_2 environment at 37°C for static culture for 24 h; the culture supernatant was discarded; and the residue was washed with a PBS buffer and then grouped. 100 μL of chimeric viruses at 0.01 MOI, 0.1 MOI, 1 MOI, and 10 MOI were added to each group of experimental wells, respectively; 100 μL of r~~Losata~~lasota at 0.01 MOI, 0.1 MOI, 1 MOI, and 10 MOI was added to the control group, respectively; and 100 μL of DMEM was added to negative control wells. After 1 h of virus infection, the virus solution was discarded, 200 μL of PBS was added to each well for washing once; and a cell maintenance medium containing 5% serum was added for culture. After culturing for 24 h and 48 h respectively, 20 μL of MTT solution (5 mg/mL) was added to each well and incubated for 4 h; the culture solution was discarded; 150 μL of DMSO was added to each well; and after horizontal oscillating shake for 10 min, the OD value at 490 nm was measured with a microplate reader, and the cell growth inhibition rate was calculated.

[00112] Inhibition rate = (OD value of negative control treated well − OD value of experimental treated well) / OD value of negative control treated well × 100%.

[00113] The inhibition rate against HepG-2 cells could be found in FIG. 4; the inhibition rate against Caki-1 cells could be found in FIG. 5; the inhibition rate against A549 cells could be found in FIG. 6; the inhibition rate against SK-MEL-5 cells could be found in FIG. 7; and the inhibition rate against LN229 cells could be found in FIG. 8.

[00114] The results showed that both the parental virus r~~Losata~~lasota and the

chimerical viruses r~~Losata~~lasota-HN, r~~Losata~~lasota-F, r~~Losata~~lasota-P, and r~~Losata~~lasota-M showed an inhibitory effect on the above cells; the inhibitory effect of the chimeric virus ~~Losata~~lasota-F is significantly higher than that of the parental virus r~~Losata~~lasota; the inhibitory effects of other chimeric viruses r~~Losata~~lasota-HN, r~~Losata~~lasota-NP, and r~~Losata~~lasota-M are higher than that of the parental virus r~~Losata~~lasota, without significant difference; and the killing effects of the recombinant viruses on the tumour cells were positively correlated with time and dose.

[00115] II. Killing effect of recombinant chimeric virus against liver cancer cells as detected by Annexin V/PI method

[00116] HepG2 tumour cells in the logarithmic growth phase were collected and prepared into a cell suspension of 1-10×10^4 cells/mL (containing 0.001% trypsin). The cell suspension was mixed well and added to a 6-well plate, and after the cells grew to 60-80%, group experiments were carried out.

[00117] HepG2 cells added with the virus-free SPF chicken embryo allantoic fluid were taken as a null cell control group (control); HepG2 cells added with 1 MOI r~~Losata~~lasota were taken as a control group; and HepG2 cells added with 1 MOI recombinant chimeric viruses were taken as experiment groups.

[00118] After 12 hours of infection, the cells (0.5-1×10^6 cells) were digested with 0.25% trypsin, and washed twice with PBS; 500 uL of binding buffer was pipetted with a pipette to gently suspend the cells; 10 μL of FITC-labeled Annexin-V was added and mixed well; 5 μL of PI was then added and mixed well; immediately after 5-15 minutes of reaction in the dark, quantitative detection was carried out by flow cytometry; and at the same time, the HepG2 cells without AnnexinV-FITC and PI added were set up as a negative control.

[00119] The results could be found in FIG. 9 and showed that both the

parental viruses and the recombinant chimeric viruses could induce the apoptosis of tumour cells; the effect of the virus r~~Losata~~lasota-F was significantly stronger than that of the virus r~~Losata~~lasota; and the effects of other recombinant chimeric viruses were not significantly different from that of the virus r~~Losata~~lasota.

[00120] Example 6: Therapeutic effect of recombinant chimeric virus on tumours and virus safety detection

[00121] I. Therapeutic effect of recombinant chimeric virus on tumours

[00122] 1. Establishment of Balb/c mice H22 liver cancer animal model

[00123] Kunming mice of 6 weeks old were used for experiments. H22 mice liver cancer cells were seeded into the abdominal cavities of Humming mice, and after 7 days, the ascites occurred, and the mice were sacrificed by neck breakage. The ascites containing H22 cells was extracted under a sterile condition; a proper amount of PBS was added to formulate a cancer cell suspension; and the cells were counted and measured in viability, with the viable cell rate up to 95%. The cell density was adjusted to 10^6 cells/mL for later use. Subcutaneous injection was performed on each Balb/c mouse at the right groin at a dose of 0.2 mL, approximately containing 2×10^5 tumour cells. After 8-12 days, when a solid tumour with a diameter of 5-8 mm was formed, the modeling was successful, and the subsequent experiments could be carried out. The mice with the tumour diameter of 5-8 mm were selected, from which individuals with greatly varied forms and sizes were removed, and the remaining mice were taken as model mice.

[00124] 2. Therapeutic effect of recombinant chimeric virus on tumours

[00125] The model mice were randomly divided into six groups, with 15 mice in each group. They were treated as follows:

[00126] an r~~Losata~~lasota-HN group: 0.2 mL of r~~Losata~~lasota-HN virus

solution (containing 10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 14-day treatment;

[00127] an r~~Losata~~lasota-F group: 0.2 mL of r~~Losata~~lasota-F virus solution (containing 10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 14-day treatment;

[00128] an r~~Losata~~lasota-P group: 0.2 mL of r~~Losata~~lasota-P virus solution (containing 10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 14-day treatment;

[00129] an r~~Losata~~lasota-M group: 0.2 mL of r~~Losata~~lasota-M virus solution (containing 10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 14-day treatment;

[00130] an r~~Losata~~lasota group: 0.2 mL of r~~Losata~~lasota virus solution (containing 10^7 pfu viruses) prepared in Example 2 was injected into the tumours of the model mice every day for 14-day treatment;

[00131] an allantoic fluid group (Model): 0.2 mL of allantoic fluid was injected to the tumour of each model mice every day, and treated for 14 days.

[00132] The tumour volume was measured every other day. After the treatment was completed, the mice were sacrificed, the tumours were removed and measured in terms of volume and weight (the results were shown in FIG. 10 and FIG. 11). The survival conditions of the remaining mice were observed after the treatment was completed, and the control group included normal mice without any treatment (FIG. 12). The results showed that, compared with the allantoic fluid group, both the parental viruses and the chimeric viruses showed significant inhibitory effects on the tumours; the anti-tumour effects of r~~Losata~~lasota-P and r~~Losata~~lasota-M were higher than that of the r~~Losata~~lasota-HN group, without significant difference; and the inhibitory effect of r~~Losata~~lasota-F is significantly higher than those of the above

three types of chimeric viruses. Compared with the allantoic fluid group, both the parental viruses and the chimeric viruses could significantly improve the survival rate of the mice; and compared with the r~~Losata~~lasota group, only r~~Losata~~lasota-F could significantly improve the survival rate of the mice.

[00133] 3. Observation of pathological sections of tumours

[00134] To observe the tumour tissue cell morphology of NDV chimeric viruses and the control group after treatment, tumour tissues were acquired from each group of mice after treatment, and were prepared into tumour tissue paraffin sections following the conventional method. The morphological characteristics of tumour tissues in each treatment group were observed under a microscope, and then compared and analysed.

[00135] The results showed that after treatment, the tumour tissues from the model group were dense in structure with intact cell morphology and vigorous growth; and the tumour tissues from the r~~Losata~~lasota virus control group undergone slight necrosis; the r~~Losata~~lasota-HN, r~~Losata~~lasota-P, and r~~Losata~~lasota-M groups exhibited incomplete cell structure, accompanied with lymphocytes infiltration to a certain extent; and the r~~Losata~~lasota-F group exhibited loose tumour tissues and incomplete cell structure, accompanied with massive lymphocytes infiltration. (See FIG. 13)

[00136] II. Safety detection of recombinant chimeric viruses

[00137] Acute toxicity test

[00138] Health Kunming mice of 4-6 weeks old were selected, 20 mice in total including half female and half male, and the mice were divided into two groups, 10 mice in each group. The mice in the control group were fed normally. Each mouse in the experiment groups was intraperitoneally injected with 1×10^4 pfu of r~~Losata~~lasota-F, and was observed for 48 hours after injection. If the mice under test

experienced respiratory depression, unsteady limbs, paralysis symptoms, convulsions, fur trembles and other adverse reactions, and death, these mice were determined to be positive.

[00139] Subacute toxicity test

[00140] Health Kunming mice of 4-6 weeks old were selected, 20 mice in total including half female and half male, and the mice were divided into two groups, 10 mice in each group. Each mouse in the experiment groups was intraperitoneally injected with 1×10^4 pfu of r~~Losata~~lasota-F, and the mice in the control group were fed normally. The responses of the mice were continuously recorded within 4 weeks, and the observations focused on aspects such as water intake, food intake, fur colour and body weight. If everything is normal without any adverse reaction or death, these mice were determined to be negative.

[00141] The results showed that the recombinant chimeric viruses had no adverse effect on the normal growth of mice, and were safe and reliable.

[00142] Example 7: Effects of recombinant chimeric viruses expressing exogenous genes on tumour cells

[00143] The tumour cells included liver cancer cells HCCLM3, melanoma cells SK-MEL2, lung cancer cells A549, pancreatic cancer cells ASPC-1, thyroid cancer cells FTC133, kidney cancer cells 786-O, neuroblastoma cells U87-MG, myosarcoma cells HT1080, esophageal cancer cells KYSE-150, uterine cancer cells Hela, breast cancer cells MCF7, and colorectal cancer cells HCT116.

[00144] I. Detection of inhibitory effect of recombinant chimeric virus on tumour cells by MTT method

[00145] Tumour cells in the logarithmic growth phase were digested by trypsin, and then prepared into a cell suspension at 2×10^4 cells/mL by using a DMEM,

F12 or RPMI-1640 culture medium containing 10% calf serum. The cell suspension was seeded in a 96-well plate (200 μl per well), placed in a 5% CO_2 environment at 37°C for static culture for 24 h; the culture supernatant was discarded; and the residue was washed with a PBS buffer and then grouped. 100 μL of 0.1 MOI chimeric viruses expressing exogenous genes were added to each group of experimental wells; 0.1 MOI r~~Losata~~lasota -F was added to the control group; and 100 μL of DMEM was added to the negative control wells. After 1 h of virus infection, the virus solution was discarded, 200 μL of PBS was added to each well for washing once; and a cell maintenance medium containing 5% serum was added for culture. After culturing for 24 h, 20 μL of MTT solution (5 mg/mL) was added to each well and incubated for 4 h; the culture solution was discarded; 150 μL of DMSO was added to each well; and after horizontal oscillating shake for 10 min, the OD value at 490 nm was measured with a microplate reader, and the cell growth inhibition rate was calculated (the results were shown in Table 2).

[00146] Inhibition rate = (OD value of negative control treated well − OD value of experimental treated well) / OD value of negative control treated well * 100%.

Table 2 Inhibitory effects of recombinant viruses expressing different exogenous genes on different tumour cells (inhibition rate %)

	r~~Losata~~lasota-F	r~~Losata~~lasota(F)-DR5	r~~Losata~~lasota(F)-TRAIL	r~~Losata~~lasota(F)-hIL2	r~~Losata~~lasota(F)-P53	r~~Losata~~lasota(F)-PD1	r~~Losata~~lasota(F)-CD	r~~Losata~~lasota(F)-mIL12	r~~Losata~~lasota(F)-DR5-TRAIL	r~~Losata~~lasota(F)-mIL12-hIL2	r~~Losata~~lasota(F)-P53-hIL2	r~~Losata~~lasota(F)-PD1-hIL2
HCCLM3	27.39	32.55	27.50	42.13	35.83	32.43	30.17	33.43	55.56	57.73	50.95	55.04
SK-MEL2	21.39	32.55	32.49	49.83	31.95	30.44	31.67	38.43	48.06	60.73	56.95	60.04
U87-MG	19.39	24.69	25.88	41.83	27.96	40.87	37.25	28.44	33.18	42.83	46.95	43.04
786-O	15.89	19.56	17.83	26.84	22.95	23.43	31.25	23.46	38.18	42.84	36.97	38.56
FITC133	18.89	21.55	20.34	26.84	35.45	28.43	21.25	22.93	33.78	38.23	34.45	33.04
HT1080	13.47	18.33	19.62	25.98	23.46	25.87	32,25	32.93	29.12	31.88	39.95	35.64
Hela	11.96	28.09	17.84	30.22	10.45	12.86	9.63	15.67	33.10	32.83	24.95	29.94
ASPC-1	11.22	19.84	17.11	21.99	29.66	15.49	16.62	19.79	26.81	25.77	32.54	24.87
A549	15.41	17.26	18.98	16.14	26.22	22.39	20.16	25.9	24.69	28.12	29.95	30.66
KYSE-150	11.28	15.04	17.56	24.65	17.45	14.31	15.09	21.93	25.18	30.56	26.03	21.87
MCF7	9.8	17.63	16.56	25.66	14.45	29.74	15.09	14.43	21.89	26.31	25.53	33.28
HCT116	12.25	15.58	19.52	13.16	16.08	26.57	20.02	18.93	22.59	27.09	29.12	33.96

[00147] The results showed that r~~Losata~~lasota(F)-DR5,

rLosatalasota(F)-TRAIL, rLosatalasota(F)-hIL2, rLosatalasota(F)-P53, rLosatalasota(F)-PD1, rLosatalasota(F)-CD, rLosatalasota(F)-mIL12, rLosatalasota(F)-DR5-TRAIL, rLosatalasota(F)-mIL12-hIL2, rLosatalasota(F)-P53-hIL2 and rLosatalasota(F)-PD1-hIL2a all showed inhibitory effects on the above cells, and showed higher inhibition rate against HCCLM3, SK-MEL2, FTC133, 786-O, U87-MG, and HT1080 than against A549, ASPC-1, KYSE-150, Hela, MCF7, and HCT116. Also, different virus strains did not exhibit completely identical cytotoxicity against A549, ASPC-1, KYSE-150, Hela, MCF7, and HCT116. Among these 12 types of tumour cells, rLosatalasota(F)-DR5, rLosatalasota(F)-TRAIL, and rLosatalasota(F)-DR5-TRAIL showed higher inhibition rate against Hela cells; rLosatalasota(F)-P53 and rLosatalasota(F)-P53-hIL2 showed higher inhibition rate against A549 and ASPC-1 cells; rLosatalasota(F)-hIL2, rLosatalasota(F)-mIL12 and rLosatalasota(F)-mIL12-hIL2 showed higher inhibition rate against A549, KYSE-150, and Hela cells; rLosatalasota(F)-CD showed higher inhibition rate against A549 cells; and rLosatalasota(F)-PD1 and rLosatalasota(F)-PD1-hIL2 showed higher inhibition rate against A549, MCF7 and HCT116 cells.

[00148] Example 8: Therapeutic effect of recombinant chimeric virus expressing exogenous gene on tumours

[00149] 1. Balb/c 4T1 breast tumour-bearing mice were established according to the method described in Example 6.

[00150] 2. Therapeutic effect of recombinant chimeric virus on tumours

[00151] The model mice were randomly divided into 13 groups, with 10 mice in each group. These groups were treated as follows respectively:

[00152] an rLosatalasota(F)-DR5 group: 0.1 mL of rLosatalasota(F)-DR5 virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected

into the tumours of the model mice every day for 21-day treatment;

[00153] an r~~Losata~~lasota(F)-TRAIL group: 0.1 mL of r~~Losata~~lasota(F)-TRAIL virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 21-day treatment;

[00154] an r~~Losata~~lasota(F)-hIL2 group: 0.1 mL of r~~Losata~~lasota(F)-hIL2 virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 21-day treatment;

[00155] an r~~Losata~~lasota(F)-P53 group: 0.1 mL of r~~Losata~~lasota(F)-P53 virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 21-day treatment;

[00156] an r~~Losata~~lasota(F)-PD1 group: 0.1 mL of r~~Losata~~lasota(F)-PD1 virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 ~~–~~was injected into the tumours of the model mice every day for 21-day treatment;

[00157] an r~~Losata~~lasota(F)-CD group: 0.1 mL of r~~Losata~~lasota(F)-CD virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 ~~–~~was injected into the tumours of the model mice every day for 21-day treatment;

[00158] an r~~Losata~~lasota(F)-mIL12 group: 0.1 mL of r~~Losata~~lasota(F)-mIL12 virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 ~~–~~was injected into the tumours of the model mice every day for 21-day treatment;

[00159] an r~~Losata~~lasota(F)-DR5-TRAIL group: 0.1 mL of r~~Losata~~lasota(F)-DR5-TRAIL virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 21-day treatment;

[00160] an r~~Losata~~lasota(F)-mIL12-hIL2 group: 0.1 mL of r~~Losata~~lasota(F)-mIL12-hIL2 virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 21-day treatment;

[00161] an r~~Losata~~lasota(F)-P53-hIL2 group: 0.1 mL of r~~Losata~~lasota(F)-P53-hIL2 virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 21-day treatment;

[00162] an r~~Losata~~lasota(F)-PD1-hIL2 group: 0.1 mL of r~~Losata~~lasota(F)-PD1-hIL2 virus solution (containing 5×10^7 pfu viruses) prepared in Example ~~1~~ 2 was injected into the tumours of the model mice every day for 21-day treatment;

[00163] an r~~Losata~~lasota-F group: 0.1 mL of r~~Losata~~lasota-F virus solution (containing 5×10^7 pfu viruses) prepared in Example 2 was injected into the tumours of the model mice every day for 21-day treatment;

[00164] an allantoic fluid group (Model): 0.1 mL of allantoic fluid was injected to the tumour of each model mice every day, and treated for 21 days.

[00165] The tumour volume was measured every other day. After the treatment was completed, the mice were sacrificed, the tumours were removed and measured in weight (the results as shown in Table 3). The results showed that compared with the allantoic fluid group, r~~Losata~~lasota-F and the recombinant chimeric viruses expressing the exogenous genes exhibited significant inhibitory effects on the tumours; the anti-tumour effects of r~~Losata~~lasota(F)-DR5, r~~Losata~~lasota(F)-TRAIL and r~~Losata~~lasota(F)-CD are higher than that of the r~~Losata~~lasota-F group, without significant difference; the anti-tumour effects of r~~Losata~~lasota(F)-P53 and r~~Losata~~lasota(F)-mIL12 are significantly higher than that of

the r~~Losata~~lasota-F treatment group; and the anti-tumour effects of r~~Losata~~lasota(F)-hIL2 and r~~Losata~~lasota(F)-PD1 showed extremely significant differences from the r~~Losata~~lasota-F group. Compared with a single-gene treatment group, a dual-gene treatment group showed a further enhanced inhibitory effect against tumours. The r~~Losata~~lasota(F)-mIL12-hIL2 and r~~Losata~~lasota(F)-PD1-hIL2 groups exhibited the best effect, and the tumours in several mice almost regressed after the treatment was completed.

Table 3 Average volume and tumour weight of recombinant chimeric viruses expressing exogenous genes after the completion of treatment

	Average volume before treatment	Average volume after the completion of treatment	Average tumour weight after the completion of treatment
Model	108.66mm^3	1886.53mm^3	1.33 g
r~~Losata~~lasota-F	123.30mm^3	1152.25mm^3	1.01 g
r~~Losata~~lasota(F)-DR5	146.88mm^3	846.69mm^3	0.86 g
r~~Losata~~lasota(F)-TRAIL	116.88mm^3	759.84mm^3	0.79 g
r~~Losata~~lasota(F)-hIL2	143.59mm^3	516.74mm^3	0.51 g
r~~Losata~~lasota(F)-P53	142.58mm^3	607.42mm^3	0.62 g
r~~Losata~~lasota(F)-PD1	133.55mm^3	508.665mm^3	0.46 g
r~~Losata~~lasota(F)-CD	107.38mm^3	992.64mm^3	0.94 g
r~~Losata~~lasota(F)-mIL12	127.98mm^3	544.22mm^3	0.67 g
r~~Losata~~lasota(F)-DR5-TRAIL	115.26mm^3	462.54mm^3	0.44 g
r~~Losata~~lasota(F)-mIL12-hIL2	113.73mm^3	309.82mm^3	0.25 g
r~~Losata~~lasota(F)-P53-hIL2	127.29mm^3	401.58mm^3	0.38 g
r~~Losata~~lasota(F)-PD1-hIL2	130.43mm^3	289.28mm^3	0.18 g

[00166] Example 9: Safety detection of recombinant chimeric viruses expressing exogenous genes on tumours

[00167] Healthy SPF Balb/c mice of 6 weeks old were selected and grouped, 10 mice in each group. The mice in the control group were fed normally. Each mouse in the experiment groups was intraperitoneally injected with 5×10^8 pfu (10 times the therapeutic dose) recombinant chimeric virus (r~~Losata~~lasota(F)-DR5, r~~Losata~~lasota(F)-TRAIL, r~~Losata~~lasota(F)-hIL2, r~~Losata~~lasota(F)-P53, r~~Losata~~lasota(F)-PD1, r~~Losata~~lasota(F)-CD, r~~Losata~~lasota(F)-mIL12,

r~~Losata~~lasota(F)-DR5-TRAIL, r~~Losata~~lasota(F)-mIL12-hIL2, r~~Losata~~lasota(F)-P53-hIL2, or r~~Losata~~lasota(F)-PD1-hIL2), and then observed for 30 days. As a result, the mice with obvious adverse reactions such as listlessness occurred, ruffled fur and death were determined as positive. After the observation was completed, the serum of mice in each group was collected to detect AST, ALT and other related indicators.

[00168] The results showed that on the second day after injection, the fur of several mice in the experiment groups ruffled, and their food and water intake were not affected. After one week of continuous injections, the fur of the mice returned to normal; and after continuous observation for one month, none of the mice in each experimental group showed obvious adverse reactions such as listlessness, ruffled fur and death. After the observation was completed, the serum of mice in each group was collected to detect AST, ALT, BUN, CREA and other related indicators by ELISA. The results showed that compared with normal mice, the activities of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) as well as the concentrations of blood urea nitrogen (BUN) and serum creatinine (CREA) in each experiment group slightly increased, without significant difference. The results indicated that the chimeric virus treatment did not induce liver and kidney damage and serious toxic or side effects.

[00169] Finally, it should be noted that the above-mentioned examples are obviously examples given for the purpose of clearly illustrating the present application, and are not intended to limit the embodiments. For a person skilled in the art, other different forms of changes or variations can also be made based on the above description. It is unnecessary or impossible to exhaustively list all the embodiments herein. Furthermore, any obvious changes or variations derived herefrom shall still fall within the protection scope of the present application.

Sequence Listing

<110> JIANGSU KANIONREAL BIOMEDICAL TECHNOLOGY CO., LTD

<120> RECOMBINANT NEWCASTLE DISEASE VIRUS AND PREPARATION METHOD THEREFOR, RECOMBINANT PLASMID, AND USE THEREOF

<130> 191071P2

<160> 8

<170> SIPOSequenceListing 1.0

<210> 1
<211> 18391
<212> DNA
<213> Artificial Sequence

<220>
<221> gene
<222> (7274)..(8935)

<400> 1
ctggcgtaat agcgaagagg cccgcaccga tcgcccttcc caacagttgc gtagcctgaa 60
tggcgaatgg gacgcgccct gtagcggcgc attaagcgcg gcgggtgtgg tggttacgcg 120
cagcgtgacc gctacacttg ccagcgccct agcgcccgct cctttcgctt tcttcccttc 180
ctttctcgcc acgttcgccg gctttccccg tcaagctcta aatcgggggc tccctttagg 240
gttccgattt agtgctttac ggcacctcga ccccaaaaaa cttgattagg gtgatggttc 300
acgtagtggg ccatcgccct gatagacggt ttttcgccct ttgacgttgg agtccacgtt 360
ctttaatagt ggactcttgt tccaaactgg aacaacactc aaccctatct cggtctattc 420
ttttgattta taagggattt gccgatttc ggcctattgg ttaaaaaatg agctgattta 480
acaaaaattt aacgcgaatt ttaacaaaat attaacgttt acaatttcag gtggcacttt 540
tcggggaaat gtgcgcggaa cccctatttg tttatttttc taaatacatt caaatatgta 600
tccgctcatg agacaataac cctgataaat gcttcaataa tattgaaaaa ggaagagtat 660
gagtattcaa catttccgtg tcgcccttat tccctttttt gcggcatttt gccttcctgt 720
ttttgctcac ccagaaacgc tggtgaaagt aaaagatgct gaagatcagt tgggtgcacg 780
agtgggttac atcgaactgg atctcaacag cggtaagatc cttgagagtt ttcgccccga 840
agaacgtttt ccaatgatga gcacttttaa agttctgcta tgtggcgcgg tattatcccg 900
tattgacgcc gggcaagagc aactcggtcg ccgcatacac tattctcaga atgacttggt 960
tgagtactca ccagtcacag aaaagcatct tacggatggc atgacagtaa gagaattatg 1020
cagtgctgcc ataagcatga gtgataacac tgcggccaac ttacttctga caacgatcgg 1080
aggaccgaag gagctaaccg ctttttttca caacatgggg gatcatgtaa ctcgccttga 1140
tcgttgggaa ccggagctga atgaagccat accaaacgac gagcgtgaca ccacgatgcc 1200
tgtagcaatg gcaacaacgt tgcgcaaact attaactggc gaactactta ctctagcttc 1260
ccggcaacaa ttaatagact ggatggaggc ggataaagtt gcaggaccac ttctgcgctc 1320
ggcccttccg gctggctggt ttattgctga taaatctgga gccggtgagc gtgggtctcg 1380
cggtatcatt gcagcactgg ggccagatgg taagccctcc cgtatcgtag ttatctacac 1440
gacgggcagt caggcaacta tggatgaacg aaatagacag atcgctgaga taggtgcctc 1500
actgattaag cattggtaac tgtcagacca agtttactca tatatacttt agattgattt 1560
aaaacttcat ttttaattta aaaggatcta ggtgaagatc ctttttgata atctcatgac 1620
caaaatccct taacgtgagt tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa 1680

aggatcttct tgagatcctt tttttctgcg cgtaatctgc tgcttgcaaa caaaaaaacc 1740
accgctacca gcggtggttt gtttgccgga tcaagagcta ccaactcttt ttccgaaggt 1800
aactggcttc agcagagcgc agataccaaa tactgtcctt ctagtgtagc cgtagttagg 1860
ccaccacttc aagaactctg tagcaccgcc tacatacctc gctctgctaa tcctgttacc 1920
agtggctgct gccagtggcg ataagtcgtg tcttaccggg ttggactcaa gacgatagtt 1980
accggataag gcgcagcggt cgggctgaac gggggggttcg tgcacacagc ccagcttgga 2040
gcgaacgacc tacaccgaac tgagatacct acagcgtgag cattgagaaa gcgccacgct 2100
tcccgaaggg agaaaggcgg acaggtatcc ggtaagcggc agggtcggaa caggagagcg 2160
cacgagggag cttccagggg ggaacgcctg gtatctttat agtcctgtcg ggtttcgcca 2220
cctctgactt gagcgtcgat ttttgtgatg ctcgtcaggg gggccgagcc tatggaaaaa 2280
cgccagcaac gcggcctttt tacggttcct ggccttttgc tggccttttg ctcacatgtt 2340
ctttcctgcg ttatcccctg attctgtgga taaccgtatt accgcctttg agtgagctga 2400
taccgctcgc cgcagccgaa cgaccgagcg cagcgagtca gtgagcgagg aagcggaaga 2460
gcgcccaata cgcaaaccgc ctctccccgc gcgttggccg attcattaat gcagctggca 2520
cgacaggttt cccgactgga aagcgggcag tgagcgcaac gcaattaatg tgagttacct 2580
cactcattag gcaccccagg ctttacactt tatgcttccg gctcctatgt tgtgtggaat 2640
tgtgagcgga taacaatttc acacaggaaa cagctatgac catgattacg ccaagctcgg 2700
aagcggccgc taatacgact cactataggg accaaacaga gaatccgtaa gttacgataa 2760
aaggcgaagg agcaattgaa gtcgcacggg tagaaggtgt gaatctcgag tgcgagcccg 2820
aagcacaaac tcgagaaagc cttctgccaa catgtcttcc gtatttgatg agtacgaaca 2880
gctcctcgcg gctcagactc gccccaatgg agctcatgga gggggagaaa aagggagtac 2940
cttaaaagta gacgtcccgg tattcactct taacagtgat gacccagaag atagatggag 3000
ctttgtggta ttctgcctcc ggattgctgt tagcgaagat gccaacaaac cactcaggca 3060
aggtgctctc atatctcttt tatgctccca ctcacaggta atgaggaacc atgttgccct 3120
tgcagggaaa cagaatgaag ccacattggc cgtgcttgag attgatggct ttgccaacgg 3180
cacgccccag ttcaacaata ggagtggagt gtctgaagag agagcacaga gatttgcgat 3240
gatagcagga tctctccctc gggcatgcag caacggaacc ccgttcgtca cagccggggc 3300
cgaagatgat gcaccagaag acatcaccga taccctggag aggatcctct ctatccaggc 3360
tcaagtatgg gtcacagtag caaaagccat gactgcgtat gagactgcag atgagtcgga 3420
aacaaggcga atcaataagt atatgcagca aggcagggtc caaaagaaat acatcctcta 3480
ccccgtatgc aggagcacaa tccaactcac gatcagacag tctcttgcag tccgcatctt 3540
tttggttagc gagctcaaga gaggccgcaa cacggcaggt ggtacctcta cttattataa 3600
cctggtaggg gacgtagact catacatcag gaataccggg cttactgcat tcttcttgac 3660
actcaagtac ggaatcaaca ccaagacatc agcccttgca cttagtagcc tctcaggcga 3720
catccagaag atgaagcagc tcatgcgttt gtatcggatg aaaggagata atgcgccgta 3780
catgacatta cttggtgata gtgaccagat gagctttgcg cctgccgagt atgcacaact 3840
ttactccttt gccatgggta tggcatcagt cctagataaa ggtactggga aataccaatt 3900
tgccagggac tttatgagca catcattctg gagacttgga gtagagtacg ctcaggctca 3960
gggaagtagc attaacgagg atatggctgc cgagctaaag ctaaccccag cagcaaggag 4020
gggcctggca gctgctgccc aacgggtctc cgaggagacc agcagcatag acatgcctac 4080
tcaacaagtc ggagtcctca ctgggcttag cgaggggggg tcccaagctc tacaaggcgg 4140
atcgaataga tcgcaagggc aaccagaagc cggggatggg gagacccaat tcctggatct 4200
gatgagagcg gtagcaaata gcatgaggga ggcgccaaac tctgcacagg gcactcccca 4260
atcggggcct cccccaactc ctgggccatc ccaagataac gacaccgact gggggtattg 4320
atggacaaaa cccagcctgc ttccacaaaa acatcccaat gccctcaccc gtagtcgacc 4380
cctcgatttg cggctctata tgaccacacc ctcaaacaaa catccccctc tttcctccct 4440
ccccctgctg tacaactccg cacgccctag ataccacagg cacaatgcgg ctcactaaca 4500
atcaaaacag agccgaggga attagaaaaa agtacgggta gaagagggat attcagagat 4560
cagggcaagt ctcccgagtc tctgctctct cctctacctg atagaccagg acaaacatgg 4620
ccacctttac agatgcagag atcgacgagc tatttgagac aagtggaact gtcattgaca 4680
acataattac agcccagggt aaaccagcag agactgttgg aaggagtgca atcccacaag 4740
gcaagaccaa ggtgctgagc gcagcatggg agaagcatgg gagcatccag ccaccggcca 4800

gtcaagacaa ccccgatcga caggacagat ctgacaaaca accatccaca cccgagcaaa 4860
cgaccccgca tgacagcccg ccggccacat ccgccgacca gccccccacc caggccacag 4920
acgaagccgt cgacacacag ctcaggaccg gagcaagcaa ctctctgctg ttgatgcttg 4980
acaagctcag caataaatcg tccaatgcta aaaagggccc atggtcgagc ccccaagagg 5040
ggaatcacca acgtccgact caacagcagg ggagtcaacc cagtcgcgga aacagtcagg 5100
aaagaccgca gaaccaagtc aaggccgccc ctggaaacca gggcacagac gtgaacacag 5160
catatcatgg acaatgggag gagtcacaac tatcagctgg tgcaacccct catgctctcc 5220
gatcaaggca gagccaagac aatacccttg tatctgcgga tcatgtccag ccacctgtag 5280
actttgtgca agcgatgatg tctatgatgg aggcgatatc acagagagta agtaaggttg 5340
actatcagct agatcttgtc ttgaaacaga catcctccat ccctatgatg cggtccgaaa 5400
tccaacagct gaaaacatct gttgcagtca tggaagccaa cttgggaatg atgaagattc 5460
tggatcccgg ttgtgccaac atttcatctc tgagtgatct acgggcagtt gcccgatctc 5520
acccggtttt agtttcaggc cctggagacc cctctcccta tgtgacacaa ggaggcgaaa 5580
tggcacttaa taaactttcg caaccagtgc cacatccatc tgaattgatt aaacccgcca 5640
ctgcatgcgg gcctgatata ggagtggaaa aggacactgt ccgtgcattg atcatgtcac 5700
gcccaatgca cccgagttct tcagccaagc tcctaagcaa gttagatgca gccgggtcga 5760
tcgaggaaat caggaaaatc aagcgccttg ctctaaatgg ctaattacta ctgccacacg 5820
tagcgggtcc ctgtccactc ggcatcacac ggaatctgca ccgagttccc ccccgcagac 5880
ccaaggtcca actctccaag cggcaatcct ctctcgcttc ctcagcccca ctgaatgatc 5940
gcgtaaccgt aattaatcta gctacattta agattaagaa aaaatacggg tagaattgga 6000
gtgccccaat tgtgccaaga tggactcatc taggacaatt gggctgtact ttgattctgc 6060
ccattcttct agcaacctgt tagcatttcc gatcgtccta caagacacag gagatgggaa 6120
gaagcaaatc gccccgcaat ataggatcca gcgccttgac ttgtggactg atagtaagga 6180
ggactcagta ttcatcacca cctatggatt catctttcaa gttgggaatg aagaagccac 6240
tgtcggcatg atcgatgata aacccaagcg cgagttactt tccgctgcga tgctctgcct 6300
aggaagcgtc ccaaataccg gagaccttat tgagctggca agggcctgtc tcactatgat 6360
agtcacatgc aagaagagtg caactaatac tgagagaatg gttttctcag tagtgcaggc 6420
accccaagtg ctgcaaagct gtagggttgt ggcaaacaaa tactcatcag tgaatgcagt 6480
caagcacgtg aaagcgccag agaagattcc cgggagtgga accctagaat acaaggtgaa 6540
ctttgtctcc ttgactgtgg taccgaagaa ggatgtctac aagatcccag ctgcagtatt 6600
gaaggtttct ggctcgagtc tgtacaatct tgcgctcaat gtcactatta atgtggaggt 6660
agacccgagg agtcctttgg ttaaatctct gtctaagtct gacagcggat actatgctaa 6720
cctcttcttg catattggac ttatgaccac cgtagatagg aaggggaaga aagtgacatt 6780
tgacaagctg gaaaagaaaa taaggagcct tgatctatct gtcgggctca gtgatgtgct 6840
cgggccttcc gtgttggtaa aagcaagagg tgcacggact aagcttttgg caccttttctt 6900
ctctagcagt gggacagcct gctatcccat agcaaatgct tctcctcagg tggccaagat 6960
actctggagt caaaccgcgt gcctgcggag cgttaaaatc attatccaag caggtaccca 7020
acgcgctgtc gcagtgaccg ccgaccacga ggttacctct actaagctgg agaaggggca 7080
caccttgcc aaatacaatc cttttaagaa ataagctgcg tctctgagat tgcgctccgc 7140
ccactcaccc agatcatcat gacacaaaaa actaatctgt cttgattatt tacagttagt 7200
ttacctgtct atcaagttag aaaaaacacg ggtagaagat tctggatccc ggttggcgcc 7260
ctccaggtgc aagatgggcc ccaaaccccc caccggaacc ccagcgcctc tggtgctgat 7320
cgcccggacc gcgctggcgt tgggctgtgt ctgtccggcg ggctctcttg acggcagacc 7380
tcttgcagct gcagggattg tggtaacgag agataaagca gtcaatatat acacttcatc 7440
tcaaacgggg tcaatcatag tcaagttact cccaaatatg cccaaggaca aggaggcgtg 7500
cgcaaaagcc ccattagagg crtacaatag aacactgacc actttactca ctcctcttgg 7560
ygactccatc cgsaggatac aagggtctgc aactacatct agaggaagga gacagaaacg 7620
ttttgtaggt gctatcattg gcagtatagc tcttggggtt gcgacagctg cacaagtaac 7680
agcagctgca gctctgatac aagccaacca gaacgctgcc aacatcctcc ggcttaagga 7740
gagcattgct gcaaccaatg aagctgtgca cgaggtcact gacggattat cacaactagc 7800
gatggcgatt gggaagatgc agcagtttgt taatgaccag tttaataata cggcgcgaga 7860
attggactgc atcaaaatta cacaacaggt tggtgtcgaa ctcaatttgt atctaactga 7920

actgactaca gtattcgggc cacaaatcac ttcccctgct ttaactcagc taactatcca 7980
ggcactttat aatttagctg gtggcaatat gaattactta ttgactaagt taggtgtagg 8040
gaacaatcaa cttagctcat taattagtag tggcctgatc actggcaacc ccattttata 8100
tgactcacag acccaactct taggcataca gataaatgta ccctcagtcg ggagcctaaa 8160
taatatgcgt gccacctact tggagacctt atccgtaagc acaaccaggg ggttcgcctc 8220
agcacttgtc ccgaaagttg tgacgcaagt tggttctgtg atagaagaac ttgacacctc 8280
atattgtata gaatctaatc tggatttata ttgtacaagg atagtgacat tccccatgtc 8340
tcccggcatt tattcctgtc tgagcggtaa tacgtcagct tgtatgtatt caaagactga 8400
gggtgcactc actacaccat acatagctct caagggctca gttattgcta attgcaagat 8460
gattacatgt agatgtgcag acccccccagg tatcatatcg caaaattacg gagaagctgt 8520
gtccctaata gataaacatt catgtaatgt cttatcccta gacggaataa ccctgaggct 8580
cagtggggaa tttgatgcga cctatcaaaa gaacatctta atactagatt cccaggtcat 8640
cgtgacaggc aatctcgata tatcaactga acttgggaat gtcaacaact cgataagcag 8700
tgctctggac aaattagcgg aaagtaacag caagttaaac aaagtcaatg tcaacctaac 8760
tagcacatct gctctcatta cttatattgt tctagctgtc atatctcttg ttttcggcgt 8820
aattagcctg attctagcgt gctgcttgat gtataaacaa aaagcacaac aaaagacctt 8880
actatggctt gggaacaata ccctcgatca gatgagagcc accacaaaaa catgaacaca 8940
gatgaggaac gaaggtttcc ctaatagtaa tttgtgtgaa agttctggta gtctgtcagt 9000
taagaaaaaa tacgggtaga aggttaaccg gcccgcggcg acgcgtggcc tgagaggcct 9060
tcagagagtt aagaaaaaac taccggttgt agatgaccaa aggacgatat acgggtagaa 9120
cggtaagaga ggccgcccct caattgcgag ccaggcttca caacctccgt tctaccgctt 9180
caccgacaac agtcctcaat catggaccgc gccgttagcc aagttgcgtt agagaatgat 9240
gaaagagagg caaaaaatac atggcgcttg atattccgga ttgcaatctt attcttaaca 9300
gtagtgacct tggctatatc tgtagcctcc cttttatata gcatgggggc tagcacacct 9360
agcgatcttg taggcatacc gactaggatt tccagggcag aagaaaagat tacatctaca 9420
cttggttcca atcaagatgt agtagatagg atatataagc aagtggccct tgagtctccg 9480
ttggcattgt taaatactga gaccacaatt atgaacgcaa taacatctct ctcttatcag 9540
attaatggag ctgcaaacaa cagtgggtgg ggggcaccta tccatgaccc agattatata 9600
ggggggatag gcaaagaact cattgtagat gatgctagtg atgtcacatc attctatccc 9660
tctgcatttc aagaacatct gaattttatc ccggcgccta ctacaggatc aggttgcact 9720
cgaataccct catttgacat gagtgctacc cattactgct acacccataa tgtaatattg 9780
tctggatgca gagatcactc acattcatat cagtatttag cacttggtgt gctccggaca 9840
tctgcaacag ggagggtatt cttttctact ctgcgttcca tcaacctgga cgacacccaa 9900
aatcggaagt cttgcagtgt gagtgcaact cccctgggtt gtgatatgct gtgctcgaaa 9960
gtcacggaga cagaggaaga agattataac tcagctgtcc ctacgcggat ggtacatggg 10020
aggttagggt tcgacggcca gtaccacgaa aaggacctag atgtcacaac attattcggg 10080
gactgggtgg ccaactaccc aggagtaggg ggtggatctt ttattgacag ccgcgtatgg 10140
ttctcagtct acggagggtt aaaacccaat tcacccagtg acactgtaca ggaagggaaa 10200
tatgtgatat acaagcgata caatgacaca tgcccagatg agcaagacta ccagattcga 10260
atggccaagt cttcgtataa gcctggacgg tttggtggga aacgcataca gcaggctatc 10320
ttatctatca aggtgtcaac atccttaggc gaagacccgg tactgactgt accgcccaac 10380
acagtcacac tcatgggggc cgaaggcaga attctcacag tagggacatc tcatttcttg 10440
tatcaacgag ggtcatcata cttctctccc gcgttattat atcctatgac agtcagcaac 10500
aaaacagcca ctcttcatag tccttataca ttcaatgcct tcactcggcc aggtagtatc 10560
ccttgccagg cttcagcaag atgccccaac tcgtgtgtta ctggagtcta tacagatcca 10620
tatcccctaa tcttctatag aaaccacacc ttgcgagggg tattcgggac aatgcttgat 10680
ggtgtacaag caagacttaa ccctgcgtct gcagtattcg atagcacatc ccgcagtcgc 10740
attactcgag tgagttcaag cagtaccaaa gcagcataca caacatcaac ttgttttaaa 10800
gtggtcaaga ctaataagac ctattgtctc agcattgctg aaatatctaa tactctcttc 10860
ggagaattca gaatcgtccc gttactagtt gagatcctca aagatgacgg ggttagagaa 10920
gccaggtctg gctagttgag tcaattataa aggagttgga aagatggcat tgtatcacct 10980
atcttctgcg acatcaagaa tcaaaccgaa tgccggcgcg tgctcgaatt ccatgttgcc 11040

agttgaccac aatcagccag tgctcatgcg atcagattaa gccttgtcaa tagtctcttg 11100
attaagaaaa aatgtaagtg gcaatgagat acaaggcaaa acagctcatg gtaaataata 11160
cgggtaggac atggcgagct ccggtcctga aggggcagag catcagatta tcctaccaga 11220
gtcacacctg tcttcaccat tggtcaagca caaactactc tattactgga aattaactgg 11280
gctaccgctt cctgatgaat gtgacttcga ccacctcatt ctcagccgac aatggaaaaa 11340
aatacttgaa tcggcctctc ctgatactga gagaatgata aaactcggaa gggcagtaca 11400
ccaaactctt aaccacaatt ccagaataac cggagtgctc caccccaggt gtttagaaga 11460
actggctaat attgaggtcc cagattcaac caacaaattt cggaagattg agaagaagat 11520
ccaaattcac aacacgagat atggagaact gttcacaagg ctgtgtacgc atatagagaa 11580
gaaactgctg gggtcatctt ggtctaacaa tgtcccccgg tcagaggagt tcagcagcat 11640
tcgtacggat ccggcattct ggtttcactc aaaatggtcc acagccaagt ttgcatggct 11700
ccatataaaa cagatccaga ggcatctgat ggtggcagct aggacaaggt ctgcggccaa 11760
caaattggtg atgctaaccc ataaggtagg ccaagtcttt gtcactcctg aacttgtcgt 11820
tgtgacgcat acgaatgaga acaagttcac atgtcttacc caggaacttg tattgatgta 11880
tgcagatatg atggagggca gagatatggt caacataata tcaaccacgg cggtgcatct 11940
cagaagctta tcagagaaaa ttgatgacat tttgcggtta atagacgctc tggcaaaaga 12000
cttgggtaat caagtctacg atgttgtatc actaatggag ggatttgcat acggagctgt 12060
ccagctactc gagccgtcag gtacatttgc aggagatttc ttcgcattca acctgcagga 12120
gcttaaagac attctaattg gcctcctccc caatgatata gcagaatccg tgactcatgc 12180
aatcgctact gtattctctg gtttagaaca gaatcaagca gctgagatgt tgtgtctgtt 12240
gcgtctgtgg ggtcacccac tgcttgagtc ccgtattgca gcaaaggcag tcaggagcca 12300
aatgtgcgca ccgaaaatgg tagactttga tatgatcctt caggtactgt ctttcttcaa 12360
gggaacaatc atcaacgggt acagaaagaa gaatgcaggt gtgtggccgc gagtcaaagt 12420
ggatacaata tatgggaagg tcattgggca actacatgca gattcagcag agatttcaca 12480
cgatatcatg ttgagagagt ataagagttt atctgcactt gaatttgagc catgtataga 12540
atatgaccct gtcaccaacc tgagcatgtt cctaaaagac aaggcaatcg cacaccccaa 12600
cgataattgg cttgcctcgt ttaggcggaa ccttctctcc gaagaccaga agaaacatgt 12660
aaaagaagca acttcgacta atcgcctctt gatagagttt ttagagtcaa atgattttga 12720
tccatataaa gagatggaat atctgacgac ccttgagtac cttagagatg acaatgtggc 12780
agtatcatac tcgctcaagg agaaggaagt gaaagttaat ggacggatct tcgctaagct 12840
gacaaagaag ttaaggaact gtcaggtgat ggcggaaggg atcctagccg atcagattgc 12900
acctttcttt cagggaaatg gagtcattca ggatagcata tccttgacca agagtatgct 12960
agcgatgagt caactgtctt ttaacagcaa taagaaacgt atcactgact gtaaagaaag 13020
agtatcttca aaccgcaatc atgatccgaa aagcaagaac cgtcggagag ttgcaacctt 13080
cataacaact gacctgcaaa agtactgtct taattggaga tatcagacaa tcaaattgtt 13140
cgctcatgcc atcaatcagt tgatgggcct acctcacttc ttcgaatgga ttcacctaag 13200
actgatggac actacgatgt tcgtaggaga ccctttcaat cctccaagtg accctactga 13260
ctgtgacctc tcaagagtcc ctaatgatga catatatatt gtcagtgcca gaggggtat 13320
cgaaggatta tgccagaagc tatggacaat gatctcaatt gctgcaatcc aacttgctgc 13380
agctagatcg cattgtcgtg ttgcctgtat ggtacagggt gataatcaag taatagcagt 13440
aacgagagag gtaagatcag acgactctcc ggagatggtg ttgacacagt tgcatcaagc 13500
cagtgataat ttcttcaagg aattaattca tgtcaatcat ttgattggcc ataatttgaa 13560
ggatcgtgaa accatcaggt cagacacatt cttcatatac agcaaacgaa tcttcaaaga 13620
tggagcaatc ctcagtcaag tcctcaaaaa ttcatctaaa ttagtgctag tgtcaggtga 13680
tctcagtgaa aacaccgtaa tgtcctgtgc caacattgcc tctactgtag cacggctatg 13740
cgagaacggg cttcccaaag acttctgtta ctatttaaac tatataatga gttgtgtgca 13800
gacatacttt gactctgagt tctccatcac caacaattcg caccccgatc ttaatcagtc 13860
gtggattgag gacatctctt ttgtgcactc atatgttctg actcctgccc aattaggggg 13920
actgagtaac cttcaatact caaggctcta cactagaaat atcggtgacc cggggactac 13980
tgcttttgca gagatcaagc gactagaagc agtgggatta ctgagtccta acattatgac 14040
taatatctta actaggccgc ctgggaatgg agattgggcc agtctgtgca acgacccata 14100
ctctttcaat tttgagactg ttgcaagccc aaatattgtt cttaagaaac atacgcaaag 14160

agtcctattt gaaacttgtt caaatccctt attgtctgga gtgcacacag aggataatga 14220
ggcagaagag aaggcattgg ctgaattctt gcttaatcaa gaggtgattc atccccgcgt 14280
tgcgcatgcc atcatggagg caagctctgt aggtaggaga aagcaaattc aagggcttgt 14340
tgacacaaca aacaccgtaa ttaagattgc gcttactagg aggccattag gcatcaagag 14400
gctgatgcgg atagtcaatt attctagcat gcatgcaatg ctgtttagag acgatgtttt 14460
ttcctccagt agatccaacc acccccttagt ctcttctaat atgtgttctc tgacactggc 14520
agactatgca cggaatagaa gctggtcacc tttgacggga ggcaggaaaa tactgggtgt 14580
atctaatcct gatacgatag aactcgtaga gggtgagatt cttagtgtaa gcggagggtg 14640
tacaagatgt gacagcggag atgaacaatt tacttggttc catcttccaa gcaatataga 14700
attgaccgat gacaccagca agaatcctcc gatgagggta ccatatctcg ggtcaaagac 14760
acaggagagg agagctgcct cacttgcaaa aatagctcat atgtcgccac atgtaaaggc 14820
tgccctaagg gcatcatccg tgttgatctg ggcttatggg gataatgaag taaattggac 14880
tgctgctctt acgattgcaa aatctcggtg taatgtaaac ttagagtatc ttcggttact 14940
gtccccttta cccacggctg ggaatcttca acatagacta gatgatggta taactcagat 15000
gacattcacc cctgcatctc tctacagggt gtcaccttac attcacatat ccaatgattc 15060
tcaaaggctg ttcactgaag aaggagtcaa agaggggaat gtggtttacc aacagatcat 15120
gctcttgggt ttatctctaa tcgaatcgat ctttccaata acaacaacca ggacatatga 15180
tgagatcaca ctgcacctac atagtaaatt tagttgctgt atcagagaag cacctgttgc 15240
ggttcctttc gagctacttg gggtggtacc ggaactgagg acagtgacct caaataagtt 15300
tatgtatgat cctagccctg tatcggaggg agactttgcg agacttgact tagctacttt 15360
caagagttat gagcttaatc tggagtcata tcccacgata gagctaatga acattctttc 15420
aatatccagc gggaagttga ttggccagtc tgtggtttct tatgatgaag atacctccat 15480
aaagaatgac gccataatag tgtatgacaa tacccgaaat tggatcagtg aagctcagaa 15540
ttcagatgtg gtccgcctat ttgaatatgc agcacttgaa gtgctcctcg actgttctta 15600
ccaactctat tacctgagag taagaggcct agacaatatt gtcttatata tgggtgattt 15660
atacaagaat atgccaggaa ttctactttc caacattgca gctacaatat ctcatcccgt 15720
cattcattca aggttacatg cagtgggcct ggtcaaccat gacggatcac accaacttgc 15780
agatacggat tttatcgaaa tgtctgcaaa actattagta tcttgcaccc gacgtgtgat 15840
ctccggctta tattcaggaa ataagtatga tctgctgttc ccatctgtct tagatgataa 15900
cctgaatgag aagatgcttc agctgatatc ccggttatgc tgtctgtaca cggtactctt 15960
tgctacaaca agagaaatcc cgaaaataag aggcttaact gcagaagaga aatgttcaat 16020
actcactgag tatttactgt cggatgctgt gaaaccatta cttagtcccg atcaagtgag 16080
ctctatcatg tctcctaaca taattacatt cccagctaat ctgtactaca tgtctcggaa 16140
gagcctcaat ttgatcaggg aaagggagga cagggatact atcctggcgt tgttgttccc 16200
ccaagagcca ttattagagt tcccttctgt gcaagatatt ggtgctcgag tgaaagatcc 16260
attcacccga caacctgcgg catttttgca agagttagat ttgagtgctc cagcaaggta 16320
tgacgcattc acacttagtc agattcatcc tgaactcaca tctccaaatc cggaggaaga 16380
ctacttagta cgatacttgt tcagagggat agggactgca tcttcctctt ggtataaggc 16440
atctcatctc ctttctgtac ccgaggtaag atgtgcaaga cacgggaact ccttatactt 16500
agctgaaggg agcggagcca tcatgagtct tctcgaactg catgtaccac atgaaactat 16560
ctattacaat acgctctttt caaatgagat gaaccccccg caacgacatt tcgggccgac 16620
cccaactcag tttttgaatt cggttgttta taggaatcta caggcggagg taacatgcaa 16680
agatggattt gtccaagagt tccgtccatt atggagagaa aatacagagg aaagtgacct 16740
gacctcagat aaagtagtgg ggtatattac atctgcagtg ccctacagat ctgtatcatt 16800
gctgcattgt gacattgaaa ttcctccagg gtccaatcaa agcttactag atcaactagc 16860
tatcaattta tctctgattg ccatgcattc tgtaagggag ggcggggtag taatcatcaa 16920
agtgttgtat gcaatgggat actactttca tctactcatg aacttgtttg ctccgtgttc 16980
cacaaaagga tatattctct ctaatggtta tgcatgtcga ggagatatgg agtgttacct 17040
ggtatttgtc atgggttacc tgggcgggcc tacatttgta catgaggtgg tgaggatggc 17100
gaaaactctg gtgcagcggc acggtacgct tttgtctaaa tcagatgaga tcacactgac 17160
caggttattc acctcacagc ggcagcgtgt gacagacatc ctatccagtc ctttaccaag 17220
attaataaag tacttgagga agaatattga cactgcgctg attgaagccg ggggacagcc 17280

cgtccgtcca ttctgtgcgg agagtctggt gagcacgcta gcgaacataa ctcagataac 17340
ccagatcatc gctagtcaca ttgacacagt tatccggtct gtgatatata tggaagctga 17400
gggtgatctc gctgacacag tatttctatt tacccccttac aatctctcta ctgacgggaa 17460
aaagaggaca tcacttaaac agtgcacgag acagatccta gaggttacaa tactaggtct 17520
tagagtcgaa aatctcaata aaataggcga tataatcagc ctagtgctta aaggcatgat 17580
ctccatggag gaccttatcc cactaaggac atacttgaag catagtacct gccctaaata 17640
tttgaaggct gtcctaggta ttaccaaact caaagaaatg tttacagaca cttctgtact 17700
gtacttgact cgtgctcaac aaaaattcta catgaaaact ataggcaatg cagtcaaagg 17760
atattacagt aactgtgact cttaacgaaa atcacatatt aataggctcc ttttttggcc 17820
aattgtattc ttgttgattt aatcatatta tgttagaaaa aagttgaacc ctgactcctt 17880
aggactcgaa ttcgaactca aataaatgtc ttaaaaaaag gttgcgcaca attattcttg 17940
agtgtagtct cgtcattcac caaatctttg tttggtttgg tggccggcat ggtcccagcc 18000
tcctcgctgg cgccggctgg gcaacattcc gaggggaccg tcccctcggt aatggcgaat 18060
gggacgcggc cgatccggct gctaacaaag cccgaaagga agctgagttg gctgctgcca 18120
ccgctgagca ataactagca taacccccttg gggcctctaa acgggtcttg aggggttttt 18180
tgctgaaagg aggaactata tccggatcgg ccgatccggc tgctaacaaa gcccgaaagg 18240
aagctgagtt ggctgctgcc accgctgagc aataactagc ataacccctt ggggcctcta 18300
aacgggtctt gaggggtttt ttgctgaaag gaggaactat atccggatgg ccgccaccgg 18360
tgggccttgc agcacatccc cccttcgcca g 18391

<210> 2
<211> 1236
<212> DNA
<213> Homo sapiens

<400> 2
atggaacaac ggggacagaa cgccccggcc gcttcggggg cccggaaaag gcacggccca 60
ggacccaggg aggcgcgggg agccaggcct gggctccggg tccccaagac ccttgtgctc 120
gttgtcgccg cggtcctgct gttggtctca gctgagtctg ctctgatcac ccaacaagac 180
ctagctcccc agcagagagc ggccccacaa caaaagaggt ccagcccctc agagggattg 240
tgtccacctg gacaccatat ctcagaagac ggtagagatt gcatctcctg caaatatgga 300
caggactata gcactcactg gaatgacctc cttttctgct tgcgctgcac caggtgtgat 360
tcaggtgaag tggagctaag tccctgcacc acgaccagaa acacagtgtg tcagtgcgaa 420
gaaggcacct tccgggaaga agattctcct gagatgtgcc ggaagtgccg cacagggtgt 480
cccagaggga tggtcaaggt cggtgattgt acaccctgga gtgacatcga atgtgtccac 540
aaagaatcag gcatcatcat aggagtcaca gttgcagccg tagtcttgat tgtggctgtg 600
tttgtttgca agtctttact gtggaagaaa gtccttcctt acctgaaagg catctgctca 660
ggtggtggtg gggaccctga gcgtgtggac agaagctcac aacgacctgg ggctgaggac 720
aatgtcctca atgagatcgt gagtatcttg cagcccaccc aggtccctga gcaggaaatg 780
gaagtccagg agccagcaga gccaacaggt gtcaacatgt tgtcccccgg ggagtcagag 840
catctgctgg aaccggcaga agctgaaagg tctcagagga ggaggctgct ggttccagca 900
aatgaaggtg atcccactga gactctgaga cagtgcttcg atgactttgc agacttggtg 960
ccctttgact cctgggagcc gctcatgagg aagttgggcc tcatggacaa tgagataaag 1020
gtggctaaag ctgaggcagc gggccacagg gacaccttgt acacgatgct gataaagtgg 1080
gtcaacaaaa ccgggcgaga tgcctctgtc cacaccctgc tggatgcctt ggagacgctg 1140
ggagagagac ttgccaagca gaagattgag gaccacttgt tgagctctgg aaagttcatg 1200
tatctagaag gtaatgcaga ctctgccatg tcctaa 1236

<210> 3
<211> 624
<212> DNA
<213> Homo sapiens

<400> 3
atggagacag acacactcct gctatgggta ctgctgctct gggttccagg atccactggt 60
acctctgagg aaaccatttc tacagttcaa gaaaagcaac aaaatatttc tcccctagtg 120
agagaaagag gtcctcagag agtagcagct cacataactg ggaccagagg aagaagcaac 180
acattgtctt ctccaaactc caagaatgaa aaggctctgg gccgcaaaat aaactcctgg 240
gaatcatcaa ggagtgggca ttcattcctg agcaacttgc acttgaggaa tggtgaactg 300
gtcatccatg aaaaagggtt ttactacatc tattcccaaa catactttcg atttcaggag 360
gaaataaaag aaaacacaaa gaacgacaaa caaatggtcc aatatattta caaatacaca 420
agttatcctg accctatatt gttgatgaaa agtgctagaa atagttgttg gtctaaagat 480
gcagaatatg gactctattc catctatcaa gggggaatat ttgagcttaa ggaaaatgac 540
agaatttttg tttctgtaac aaatgagcac ttgatagaca tggaccatga agccagtttt 600
ttcggggcct ttttagttgg ctaa 624

<210> 4
<211> 462
<212> DNA
<213> Homo sapiens

<400> 4
atgtacagga tgcaactcct gtcttgcatt gcactaagtc ttgcacttgt cacaaacagt 60
gcacctactt caagttctac aaagaaaaca cagctacaac tggagcattt actgctggat 120
ttacagatga ttttgaatgg aattaataat tacaagaatc ccaaactcac caggatgctc 180
acatttaagt tttacatgcc caagaaggcc acagaactga aacatcttca gtgtctagaa 240
gaagaactca aacctctgga ggaagtgcta aatttagctc aaagcaaaaa ctttcactta 300
agacccaggg acttaatcag caatatcaac gtaatagttc tggaactaaa gggatctgaa 360
acaacattca tgtgtgaata tgctgatgag acagcaacca ttgtagaatt tctgaacaga 420
tggattacct tttgtcaaag catcatctca acactgactt aa 462

<210> 5
<211> 1182
<212> DNA
<213> Homo sapiens

<400> 5
atggaggagc cgcagtcaga tcctagcgtc gagccccctc tgagtcagga aacattttca 60
gacctatgga aactacttcc tgaaaacaac gttctgtccc ccttgccgtc ccaagcaatg 120
gatgatttga tgctgtcccc ggacgatatt gaacaatggt tcactgaaga cccaggtcca 180
gatgaagctc ccagaatgcc agaggctgct ccccccgtgg cccctgcacc agcagctcct 240
acaccggcgg cccctgcacc agccccctcc tggcccctgt catcttctgt cccttcccag 300
aaaacctacc agggcagcta cggtttccgt ctgggcttct tgcattctgg gacagccaag 360
tctgtgactt gcacgtactc ccctgccctc aacaagatgt tttgccaact ggccaagacc 420
tgccctgtgc agctgtgggt tgattccaca cccccgcccg gcacccgcgt ccgcgccatg 480
gccatctaca agcagtcaca gcacatgacg gaggttgtga ggcgctgccc ccaccatgag 540
cgctgctcag atagcgatgg tctggccccc cctcagcatc ttatccgagt ggaaggaaat 600
ttgcgtgtgg agtatttgga tgacagaaac acttttcgac atagtgtggt ggtgccctat 660
gagccgcctg aggttggctc tgactgtacc accatccact acaactacat gtgtaacagt 720
tcctgcatgg gcggcatgaa ccggaggccc atcctcacca tcatcacact ggaagactcc 780
agtggtaatc tactgggacg gaacagcttt gaggtgcgtg tttgtgcctg tcctgggaga 840
gaccggcgca cagaggaaga gaatctccgc aagaaagggg agcctcacca cgagctgccc 900
ccagggagca ctaagcgagc actgcccaac aacaccagct cctctcccca gccaaagaag 960
aaaccactgg atggagaata tttcaccctt cagatccgtg ggcgtgagcg cttcgagatg 1020

ttccgagagc tgaatgaggc cttggaactc aaggatgccc aggctgggaa ggagccaggg 1080
gggagcaggg ctcactccag ccacctgaag tccaaaaagg gtcagtctac ctcccgccat 1140
aaaaaactca tgttcaagac agaagggcct gactcagact ga 1182

<210> 6
<211> 504
<212> DNA
<213> Homo sapiens

<400> 6
atggagacag acacactcct gctatgggta ctgctgctct gggttccagg atccactggt 60
ccaggatggt tcttagactc cccagacagg ccctggaacc cccccacctt ctccccagcc 120
ctgctcgtgg tgaccgaagg ggacaacgcc accttcacct gcagcttctc caacacatcg 180
gagagcttcg tgctaaactg gtaccgcatg agccccagca accagacgga caagctggcc 240
gccttccccg aggaccgcag ccagcccggc caggactgcc gcttccgtgt cacacaactg 300
cccaacgggc gtgacttcca catgagcgtg gtcagggccc ggcgcaatga cagcggcacc 360
tacctctgtg gggccatctc cctggccccc aaggcgcaga tcaaagagag cctgcgggca 420
gagctcaggg tgacagagag aagggcagaa gtgcccacag cccaccccag cccctcaccc 480
aggccagccg gccagttcca atga 504

<210> 7
<211> 1284
<212> DNA
<213> Escherichia coli

<400> 7
atgtcgaata acgctttaca aacaattatt aacgcccggt taccaggcga agaggggctg 60
tggcagattc atctgcagga cggaaaaatc agcgccattg atgcgcaatc cggcgtgatg 120
cccataactg aaaacagcct ggatgccgaa caaggtttag ttataccgcc gtttgtggag 180
ccacatattc acctggacac cacgcaaacc gccggacaac cgaactggaa tcagtccggc 240
acgctgtttg aaggcattga acgctgggcc gagcgcaaag cgttattaac ccatgacgat 300
gtgaaacaac gcgcatggca aacgctgaaa tggcagattg ccaacggcat tcagcatgtg 360
cgtacccatg tcgatgtttc ggatgcaacg ctaactgcgc tgaaagcaat gctggaagtg 420
aagcaggaag tcgcgccgtg gattgatctg caaatcgtcg ccttccctca ggaagggatt 480
ttgtcgtatc ccaacggtga agcgttgctg gaagaggcgt tacgcttagg ggcagatgta 540
gtgggggcga ttccgcattt tgaatttacc cgtgaatacg gcgtggagtc gctgcataaa 600
accttcgccc tggcgcaaaa atacgaccgt ctcatcgacg ttcactgtga tgagatcgat 660
gacgagcagt cgcgctttgt cgaaaccgtt gctgccctgg cgcaccatga aggcatgggc 720
gcgcgagtca ccgccagcca caccacggca atgcactcct ataacggggc gtatacctca 780
cgcctgttcc gcttgctgaa aatgtccggt attaactttg tcgccaaccc gctggtcaat 840
attcatctgc aaggacgttt cgatacgtat ccaaaacgtc gcggcatcac gcgcgttaaa 900
gagatgctgg agtccggcat taacgtctgc tttggtcacg atgatgtctt cgatccgtgg 960
tatccgctgg gaacggcgaa tatgctgcaa gtgctgcata tggggctgca tgtttgccag 1020
ttgatgggct acgggcagat taacgatggc ctgaatttaa tcacccacca cagcgcaagg 1080
acgttgaatt tgcaggatta cggcattgcc gccggaaaca gcgccaacct gattatcctg 1140
ccggctgaaa atgggtttga tgcgctgcgc cgtcaggttc cggtacgtta ttcggtacgt 1200
ggcggcaagg tgattgccag cacacaaccg gcacaaacca ccgtatatct ggagcagcca 1260
gaagccatcg attacaaacg ttga 1284

<210> 8
<211> 1623
<212> DNA

<213> Mus musculus

<400> 8
atgtgtcaat cacgctacct cctctttttg gccacccttg ccctcctaaa ccacctcagt 60
ttggccaggg tcattccagt ctctggacct gccaggtgtc ttagccagtc ccgaaacctg 120
ctgaagacca cagatgacat ggtgaagacg gccagagaaa agctgaaaca ttattcctgc 180
actgctgaag acatcgatca tgaagacatc acacgggacc aaaccagcac attgaagacc 240
tgtttaccac tggaactaca caagaacgag agttgcctgg ctactagaga gacttcttcc 300
acaacaagag ggagctgcct gcccccacag aagacgtctt tgatgatgac cctgtgcctt 360
ggtagcatct atgaggactt gaagatgtac cagacagagt tccaggccat caacgcagca 420
cttcagaatc acaaccatca gcagatcatt ctagacaagg gcatgctggt ggccatcgat 480
gagctgatgc agtctctgaa tcataatggc gagactctgc gccagaaacc tcctgtggga 540
gaagcagacc cttacagagt gaaaatgaag ctctgcatcc tgcttcacgc cttcagcacc 600
cgcgtcgtga ccatcaacag ggtgatgggc tatctgagct ccgccgttcc tggagtaggg 660
gtacctggag tgggcggatc tatgtgggag ctggagaaag acgtttatgt tgtagaggtg 720
gactggactc ccgatgcccc tggagaaaca gtgaacctca cctgtgacac gcctgaagaa 780
gatgacatca cctggacctc agaccagaga catggagtca taggctctgg aaagaccctg 840
accatcactg tcaaagagtt tctagatgct ggccagtaca cctgccacaa aggaggcgag 900
actctgagcc actcacatct gctgctccac aagaaggaaa atggaatttg gtccactgaa 960
attttaaaaa atttcaaaaa caagactttc ctgaagtgtg aagcaccaaa ttactccgga 1020
cggttcacgt gctcatggct ggtgcaaaga aacatggact tgaagttcaa catcaagagc 1080
agtagcagtc cccccgactc tcgggcagtg acatgtggaa tggcgtctct gtctgcagag 1140
aaggtcacac tggaccaaag ggactatgag aagtattcag tgtcctgcca ggaggatgtc 1200
acctgcccaa ctgccgagga gaccctgccc attgaactgg cgttggaagc acggcagcag 1260
aataaatatg agaactacag caccagcttc ttcatcaggg acatcatcaa accagacccg 1320
cccaagaact tgcagatgaa gcctttgaag aactcacagg tggaggtcag ctgggagtac 1380
cctgactcct ggagcactcc ccattcctac ttctccctca agttctttgt tcgaatccag 1440
cgcaagaaag aaaagatgaa ggagacagag gaggggtgta accagaaagg tgcgttcctc 1500
gtagagaaga catctaccga agtccaatgc aaaggcggga atgtctgcgt gcaagctcag 1560
gatcgctatt acaattcctc atgcagcaag tgggcatgtg ttccctgcag ggtccgatcc 1620
tag 1623

What is claimed is:

1. A recombinant Newcastle disease virus, being obtained by replacing an F protein of a Newcastle disease virus ~~Losata~~lasota with an F protein of a Newcastle disease virus mesogenic strain.

2. The recombinant Newcastle disease virus according to claim 1, wherein a coding DNA of an F protein of the recombinant Newcastle disease virus is as set forth in SEQ ID NO: 1 at positions 7274-8935 from a 5'-terminus.

3 The recombinant Newcastle disease virus according to claim 1, being obtained by replacing an F gene in a genome of the Newcastle disease virus ~~Losata~~lasota with an F gene of the Newcastle disease virus mesogenic strain, wherein

preferably, a DNA corresponding to the F gene of the recombinant Newcastle disease virus is as set forth in SEQ ID NO: 1 at the positions 7274-8935 from the 5'-terminus; and

more preferably, a DNA corresponding to a genome of the recombinant Newcastle disease virus is as set forth in SEQ ID NO: 1.

4. The recombinant Newcastle disease virus according to claim 1, wherein a DNA corresponding to a genome of the recombinant Newcastle disease virus further comprises an exogenous gene, which is selected from one or more of a group consisting of DR5, TRAIL, hIL2, P53, PD1, CD, and mIL12;

preferably, the exogenous gene is DR5, TRAIL, hIL2, P53, PD1, CD, or mIL12;

preferably, the exogenous gene is mIL12-hIL2, P53-hIL2, PD1-hIL2, or DR5-TRAIL;

preferably, the DR5 is as set forth in SEQ ID NO: 2, the TRAIL is as set forth in SEQ ID NO: 3, the hIL2 is as set forth in SEQ ID NO: 4, the P53 is as set forth in SEQ ID NO: 5, the PD1 is as set forth in SEQ ID NO: 6, the CD is as set forth

in SEQ ID NO: 7, and the mIL12 is as set forth in SEQ ID NO: 8; and

preferably, the exogenous gene is located between a P gene and an M gene of the recombinant Newcastle disease virus.

5. A recombinant plasmid, being obtained by replacing an F gene in a pBr~~Losata~~lasota plasmid, wherein the F gene of the recombinant plasmid is as set forth in SEQ ID NO: 1 at positions 7274-8935 from a 5'-terminus.

6. The recombinant plasmid according to claim 5, wherein the recombinant plasmid is a DNA molecule plasmid as set forth in SEQ ID NO: 1.

7. The recombinant plasmid according to claim 5, further comprising an exogenous gene, which is selected from one or more of a group consisting of DR5, TRAIL, hIL2, P53, PD1, CD, and mIL12, wherein

preferably, the exogenous gene is DR5, TRAIL, hIL2, P53, PD1, CD, or mIL12;

preferably, the exogenous gene is mIL12-hIL2, P53-hIL2, PD1-hIL2, or DR5-TRAIL;

preferably, the DR5 is as set forth in SEQ ID NO: 2, the TRAIL is as set forth in SEQ ID NO: 3, the hIL2 is as set forth in SEQ ID NO: 4, the P53 is as set forth in SEQ ID NO: 5, the PD1 is as set forth in SEQ ID NO: 6, the CD is as set forth in SEQ ID NO: 7, and the mIL12 is as set forth in SEQ ID NO: 8; and

preferably, the exogenous gene is located between a P gene and an M gene of a genome of the recombinant Newcastle disease virus.

8. A preparation method for a recombinant Newcastle disease virus, comprising transfecting the recombinant plasmid according to any one of claims 5 to 7 to cells or cell line for culturing to obtain the recombinant Newcastle disease virus.

9. The preparation method according to claim 8, wherein

the recombinant plasmid according to any one of claims 5 to 7 and a helper plasmid are co-transfected to cells or cell line for culturing; and

the cell is a mammalian cell.

10. Use of the recombinant Newcastle disease virus according to any one of claims 1 to 4 or the recombinant plasmid according to any one of claims 5 to 7 in preparation of a drug having the following function(s) of (a) and/or (b) and/or (c):

(a) treating a tumour;

(b) inhibiting tumour cell proliferation; and

(c) killing tumour cells.

11. The use according to claim 10, wherein

the tumour is selected from one or more of a group consisting of a liver cancer, a breast cancer, a non-small cell lung cancer, a melanoma, a neuroblastoma, a lung cancer, a pancreatic cancer, a thyroid cancer, a kidney cancer, a glioma, a myosarcoma, an esophageal cancer, a uterine cancer, ~~a breast cancer,~~ and a colorectal cancer; and

the tumour cell is selected from one or more of a group consisting of a liver cancer cell, a breast cancer cell, a non-small cell lung cancer cell, a melanoma cell, a neuroblastoma cell, a lung cancer cell, a pancreatic cancer cell, a thyroid cancer cell, a kidney cancer cell, a glioma cell, a myosarcoma cell, an esophageal cancer cell, a uterine cancer cell, ~~a breast cancer cell,~~ and a colorectal cancer cell.

12. A drug, comprising

the recombinant Newcastle disease virus according to any one of claims 1 to 4, and/or

the recombinant plasmid according to any one of claims 5 to 7, wherein

the drug has the following function(s) of (a) and/or (b) and/or (c):

(a) treating a tumour;

(b) inhibiting tumour cell proliferation; and

(c) killing tumour cells.