PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 405/12 | A1

(11) International Publication Number: WO 99/55700

(43) International Publication Date: 4 November 1999 (04.11.99)

(21) International Application Number: PCT/US99/09092

(22) International Filing Date: 27 April 1999 (27.04.99)

(30) Priority Data:
60/083,493 29 April 1998 (29.04.98) US

(71) Applicants *(for all designated States except US)*: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). LONZA LTD. [CH/CH]; Muenchensteinerstrasse 38, CH–4002 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: MATSON, Patricia, Ann [US/US]; R.R.1, Box 147U, 10 Mill Street, Poolville, NY 13332 (US). GODLEWSKI, Michael, Selden [US/US]; 2327 State Highway 23, South Plymouth, NY 13844 (US). QUORROZ, Daniel [–/–]; – (**). GUGGISBERG, Yves [–/–]; – (**).

(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217–1087 (US).


(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.

(54) Title: PROCESS FOR MAKING 1,3–DISUBSTITUTED–4–OXOCYCLIC UREAS

(57) Abstract

A process for making 1,3–disubstituted–4–oxocyclic ureas of general formula (I) wherein R_1, R_2, and R_3 are independently selected from the group consisting of nil, Cl, F, Br, NH_2, NO_2, COOH, CH_3SO_2, NH, SO_3H, OH, alkoxy, alkyl, alkoxycarbonyl, hydroxyalkyl, carboxyalkyl, and acyloxy; R_4 is selected from the group consisting of a substituted or unsubstituted alkyl, alkenyl, alkynyl, alkylacyl, and heteroalkyl; and A is a substituted or unsubstituted, saturated or unsaturated, straight–chain or branched alkyl or alkenyl amino group comprised of 1–7 carbon atoms; or A is a substituted or unsubstituted, saturated or unsaturated heterocycle having 5, 6, or 7 members containing at least one nitrogen, and R_4 is attached to this nitrogen; wherein said 1,3–disubstituted–4–oxocyclic urea is made without isolation of intermediates and comprising the steps: Ia) reacting a 1–substituted–4–oxocyclic urea with a carbon chain containing at least two leaving groups in the presence of a mild base and a solvent to form an adduct containing at least one leaving group, and; Ib) condensing the adduct with an amine to form a 1,3–disubstituted–4–oxocyclic urea, and; II) recovering said 1,3–disubstituted–4–oxocyclic urea, are disclosed. This method is particularly preferred for making 1[[[5–(4–Chlorophenyl)–2–furanyl]methylene]amino]–3–[4–(4–methyl–1–piperazinyl)butyl)–2,4–imidazolidinedione.



(I)