

26 AUG 2003

DESCRIPTION

GENES CODING FOR FLAVONE SYNTHASES

Technical Field

5 The present invention relates to the control and
utilization of biosynthesis of flavones, which have
effects on flower color, protection from ultraviolet ray,
symbiosis with microorganisms, etc. in plants, by a
genetic engineering technique. More specifically, it
10 relates to genes encoding proteins with activity of
synthesizing flavones from flavanones, and to their
utilization.

Background Art

15 The abundance of different flower colors is one of
the pleasant aspects of life that enriches human minds
and hearts. It is expected to increase food production
to meet future population increase by the means of
accelerating the growth of plants through symbiosis with
microorganisms, or by increasing the number of nitrogen-
20 fixing leguminous bacteria, thus improving the plant
productivity as a result of increasing the content of
nitrogen in the soil. Elimination or reduction of the
use of agricultural chemicals is also desirable to
achieve more environmentally friendly agriculture, and
25 this requires improvement of the soil by the above-
mentioned biological means, as well as higher resistance
of plants against microbial infection. Another desired
goal is to obtain plants with high protective functions
against ultraviolet rays as a means of protecting the
30 plants from the destruction of the ozone layer.

 "Flavonoid" is a general term for a group of
compounds with a C6-C3-C6 carbon skeleton, and they are
widely distributed throughout plant cells. Flavonoids
are known to have such functions as attracting insects
35 and other pollinators, protecting plant from ultraviolet
rays, and participating in interaction with soil
microorganisms (BioEssays, 16 (1994), Koes et al., p.123;

26 AUG 2003

Trends in Plant Science, 1 (1997), Shirley, B.W., p.377).

Of flavonoids, flavone plays an important role in interaction of plants with microorganisms, especially in legumes, where they participate in the initial steps of the symbiosis with leguminous bacteria (Plant Cell, 7 (1995), Dixon and Paiva, p.1085; Annu. Rev. Phytopathol., 33 (1995), Spaink, p.345). Flavones in petals play a role in recognition by insects and act as copigments which form complexes with anthocyanins. (Gendai Kagaku, (May, 1998), Honda and Saito, p.25; Prog. Chem. Org. Natl. Prod., 52 (1987), Goto, T., p.114). It is known that when flavone forms a complex with anthocyanin, the absorption maximum of the anthocyanin shifts toward the longer wavelength, i.e. toward blue.

The biosynthesis pathways for flavonoids have been widely studied (Plant Cell, 7 (1995), Holton and Cornish, p.1071), and the genes for all of the enzymes involved in the biosynthesis of anthocyanidin 3-glucoside and flavonol, for example, have been isolated. However, the genes involved in the biosynthesis of flavones have not yet been isolated. The enzymes that synthesize flavones include those belonging to the dioxygenase family that depends on 2-oxoglutaric acid (flavone synthase I) and monooxygenase enzymes belonging to the cytochrome P450 family (flavone synthase II). These groups of enzymes are completely different enzymes with no structural homology.

It has been reported that in parsley, 2-oxoglutaric acid-dependent dioxygenase catalyzes a reaction which produces apigenin, a flavone, from naringenin, a flavanone (Z. Naturforsch., 36c (1981), Britsch et al., p.742; Arch. Biochem. Biophys., 282 (1990), Britsch, p.152). The other type, flavone synthase II, is known to exist in snapdragon (Z. Naturforsch., 36c (1981), Stotz and Forkmann, p.737) and soybean (Z. Naturforsch., 42c (1987), Kochs and Grisebach, p.343; Planta, 171 (1987), Kochs et al., p.519). A correlation has been recently

reported between a gene locus and flavone synthase II activity in the petals of gerbera (Phytochemistry, 49 (1998), Martens and Forkmann, p.1953). However, there are no reports that the genes for these flavone synthases I and II were isolated or that flavone synthase II was highly purified.

The properties of a cytochrome P450 protein, which had licodione-synthesizing activity that was induced when cultured cells of licorice (*Glycyrrhiza echinata*) were treated with an elicitor, were investigated. The protein is believed to catalyze the hydroxylation of 2-position of liquiritigenin which is a 5-deoxyflavanone, followed by non-enzymatic hemiacetal ring opening to produce licodione (Plant Physiol., 105 (1994), Otani et al., p.1427). For cloning of licodione synthase, a cDNA library was prepared from elicitor-treated *Glycyrrhiza* cultured cells, and 8 gene fragments encoding cytochrome P450 were cloned (Plant Science, 126 (1997), Akashi et al., p.39).

From these fragments there were obtained two different full-length cDNA sequences, each encoding a cytochrome P450, which had been unknown until that time. Specifically, they were CYPGe-3 (cytochrome P450 No.CYP81E1) and CYPGe-5 (cytochrome P450 No.CYP93B1, hereinafter indicated as CYP93B1) (Plant Physiol., 115 (1997), Akashi et al., p.1288). By further expressing the CYP93B1 cDNA in a system using cultured insect cells, the protein derived from the gene was shown to catalyze the reaction synthesizing licodione from liquiritigenin, a flavanone, and 2-hydroxynaringenin from naringenin, also a flavanone.

2-Hydroxynaringenin was converted to apigenin, a flavone, by acid treatment with 10% hydrochloric acid (room temperature, 2 hours). Also, eriodictyol was converted to luteolin, a flavone, by reacting eriodictyol with microsomes of CYP93B1-expressing yeast followed by acid treatment. It was therefore demonstrated that the

26 AUG 2003

cytochrome P450 gene encodes the function of flavanone 2-hydroxylase activity (FEBS Lett., 431 (1998), Akashi et al., p.287). Here, production of apigenin from naringenin required CYP93B1 as well as another unknown enzyme, so that it was concluded that a total of two enzymes were necessary.

However, no genes have yet been identified for enzymes with activity of synthesizing flavones (such as apigenin) directly from flavanones (such as naringenin) without acid treatment. Thus, despite the fact that flavones have numerous functions in plants, no techniques have yet been reported for controlling their biosynthesis in plants, and improving the biofunctions in which flavones are involved, such as flower color. The discovery of an enzyme which by itself can accomplish synthesis of flavones from flavanones and acquisition of its gene, and introduction of such a gene into plants, would be more practical and industrially applicable than the introduction into a plant of genes for two enzymes involved in the synthesis of flavones from flavanones.

Disclosure of the Invention

It is an aim of the present invention to provide flavone synthase genes, preferably flavone synthase II genes, and more preferably genes for flavone synthases with activity of synthesizing flavones directly from flavanones. The obtained flavone synthase genes may be introduced into plants and over-expressed to alter flower colors.

Moreover, in the petals of flowers that naturally contain large amounts of flavones, it is expected that controlling expression of the flavone synthase genes by an antisense method or a cosuppression method can also alter flower colors. Also, expression of the flavone synthase genes in the appropriate organs, in light of the antibacterial activity of flavones and their interaction with soil microorganisms, will result in an increase in the antibacterial properties of plants and improvement in

26 AUG 2003

the nitrogen fixing ability of legumes due to promoted symbiosis with rhizosphere microorganisms, as well as a protective effect against ultraviolet rays and light.

5 The present invention therefore provides genes encoding proteins that can synthesize flavones directly from flavanones. The genes are, specifically, genes encoding flavone synthase II that can synthesize flavones from flavanones by a single-enzyme reaction (hereinafter referred to as "flavone synthase II").

10 More specifically, the present invention provides genes encoding P450 proteins having the amino acid sequences listed as SEQ.ID. No. 2, 4 or 8 of the Sequence Listing and possessing activity of synthesizing flavones from flavanones, or genes encoding proteins having amino
15 acid sequences modified by additions or deletions of one or more amino acids and/or a substitution with different amino acids in said amino acid sequence, and possessing activity of synthesizing flavones from flavanones.

20 The invention further provides a gene encoding proteins having amino acid sequences with at least 55% identity with the amino acid sequences listed as SEQ.ID. No. 2, 4 or 8 of the Sequence Listing and possessing activity of synthesizing flavones from flavanones.

25 The invention still further provides genes encoding proteins possessing activity of synthesizing flavones from flavanones, and hybridizing with all or a part of the nucleotide sequences listed as SEQ.ID. No. 1, 3 or 7 of the Sequence List under the conditions of 5 x SSC, 50°C.

30 The invention still further provides a vector, particularly an expression vector, containing any one of the aforementioned genes.

The invention still further provides a host transformed with the aforementioned vector.

35 The invention still further provides a protein encoded by any of the aforementioned genes.

The invention still further provides a process for

26 AUG 2003

producing the aforementioned protein which is characterized by culturing or growing the aforementioned host, and collecting the protein with flavone-synthesizing activity from the host.

5 The invention still further provides a plant into which any one of the aforementioned genes has been introduced, or progenies of the plant or a tissue thereof, such as cut flowers, which exhibit the same properties.

10 The invention still further provides a method of altering amounts and compositions of flavonoid using the aforementioned genes; a method of altering amounts of flavones using the aforementioned genes; a method of altering flower colors using the aforementioned genes; a
15 method of bluing the color of flowers using the aforementioned genes; a method of reddening the color of flowers using the aforementioned genes; a method of modifying the photosensitivity of plants using the
20 aforementioned genes; and a method of controlling the interaction between plants and microbes using the aforementioned genes.

Brief Description of the Drawings

Fig. 1 is a chromatogram showing the results of HPLC analysis of products obtained from a substrate,
25 naringenin, using proteins encoded by CYP93B1 and TFNS5.

A and B: Obtained by adding a crude enzyme fraction of CYP93B1-expressing yeast.

C and D: Obtained by adding a crude enzyme fraction of TFNS5-expressing yeast.

30 A and C: Direct products obtained by addition of enzyme fraction.

B: Products obtained by acid treatment after reaction of A.

35 D: Products obtained by acid treatment after reaction of C.

Embodiments for Carrying out the Invention

Flavanone 2-hydroxylase encoded by the *Glycyrrhiza*

26 AUG 2003

CYP93B1 gene produces 2-hydroxyflavanones from flavanones as the substrates, and the products are converted to flavones by acid treatment. The present inventors viewed that it would be possible to obtain a gene encoding a flavone synthase II, which was an object of the invention, by using the *Glycyrrhiza*-derived cDNA, CYP93B1 for screening of a cDNA library of, for example, a flower containing a large amount of flavones, to thus obtain cDNA encoding proteins with activity of synthesizing flavones directly from flavanones as substrates.

According to the invention, a cDNA library of snapdragon which contains a large amount of flavones is screened using the *Glycyrrhiza*-derived cDNA, CYP93B1 as a probe, to obtain cDNA encoding a novel cytochrome P450 (see Example 1). The snapdragon cDNA, ANFNS2, obtained in this manner and the *Glycyrrhiza* CYP93B1 cDNA were then used as a mixed probe to obtain TFNS5, a cDNA encoding a novel cytochrome P450, from a cDNA library of *torenia* flower petals (see Example 2).

The *torenia*-derived cDNA was expressed in yeast and reacted with naringenin, a flavanone, as a substrate which resulted in production not of 2-hydroxynaringenin but rather of the flavone apigenin, without acid treatment (see Example 3). In other words, this enzyme directly produced flavones from flavanones without acid treatment, and its gene was confirmed to be a flavone synthase II which had never been cloned. The amino acid sequence encoded by the snapdragon-derived ANFNS2 of Example 1 exhibited high identity of 77% with the flavone synthase II encoded by TFNS5, and it exhibited the enzyme activity of flavone synthase II (Example 4). In addition, since an amino acid sequence encoded by *perilla*-derived cDNA also exhibited high identity of 76% and 75% with TFNS5 and ANFNS2, respectively (Example 8), it is speculated that the protein encoded by this cDNA also possesses the same enzymatic activity as the flavone synthases encoded by TFNS5 and ANFNS2.

26 AUG 2003

The genes of the present invention may be, for example, one encoding the amino acid sequences listed as SEQ.ID. No. 2, 4 or 8 of the Sequence Listing. However, it is known that proteins whose amino acid sequences are modified by additions or deletions of multiple amino acids and/or substitutions with different amino acids can maintain the same enzyme activity as the original protein. Consequently, proteins having the amino acid sequences listed as SEQ.ID. No. 2, 4 or 8 of the Sequence Listing wherein the amino acid sequence is modified by additions or deletions of one or more amino acids and/or substitutions with different amino acids, and genes encoding those proteins, are also encompassed by the present invention so long as they maintain the activity of producing flavones directly from flavanones.

The present invention also relates to genes that have the nucleotide sequences listed as SEQ.ID. Nos. 1, 3 and 7 and nucleotide sequences encoding the amino acid sequences listed therein, or that hybridize with portions of their nucleotide sequences under conditions of 5 x SSC, 50°C, for example, providing they encode proteins possessing activity of producing flavones from flavanones. The suitable hybridization temperature will differ depending on nucleotide sequences and the length of nucleotide sequences, and for example, when the probe used is a DNA fragment comprising 18 bases coding for 6 amino acids, the temperature is preferably not higher than 50°C.

A gene selected by such hybridization may be a naturally derived one, such as a plant-derived gene, for example, a gene derived from snapdragon, torenia or perilla; it may also be a gene from another plant, such as gentian, verbena, chrysanthemum, iris, or the like. A gene selected by hybridization may be cDNA or genomic DNA.

The invention also relates to genes encoding proteins that have amino acid sequences with identity of

at least 55%, preferably at least 70%, such as 80% or greater and even 90% or greater, with any one of the amino acid sequences listed as SEQ.ID. Nos. 2, 4 or 8 of the Sequence Listing, and that possess activity of synthesizing flavones from flavanones.

A gene with the natural nucleotide sequence can be obtained by screening of a cDNA library, for example, as demonstrated in detail in the examples. DNA encoding enzymes with modified amino acid sequences can be synthesized using common site-directed mutagenesis or a PCR method, using DNA with a natural nucleotide sequence as a starting material. For example, a DNA fragment into which a modification is to be introduced may be obtained by restriction enzyme treatments of natural cDNA or genomic DNA and then used as a template for site-directed mutagenesis or PCR using a primer having the desired mutation introduced therein, to obtain a DNA fragment having the desired modification introduced therein. Mutation-introduced DNA fragments may then be linked to a DNA fragment encoding another portion of a target enzyme.

Alternatively, in order to obtain DNA encoding an enzyme consisting of a shortened amino acid sequence, for example, DNA encoding an amino acid sequence which is longer than the aimed amino acid sequence, such as the full length amino acid sequence, may be cut with desired restriction endonucleases, and if the DNA fragment obtained thereby does not encode the entire target amino acid sequence, it may be linked with synthesized DNA comprising the rest of the sequence.

Thus obtained genes may be expressed in an expression system using *E. coli* or yeast and its enzyme activity measured to confirm that the obtained gene encodes flavone synthase. By expressing the gene, it is also possible to obtain the flavone synthase protein as the gene product. Alternatively, it is also possible to obtain a flavone synthase protein even using antibodies for a full or a partial amino acid sequence listed as

26 AUG 2003

SEQ.ID. No. 2, 4 or 8, and such antibodies may be used for cloning of a flavone synthase gene in another organism.

Consequently, the invention also relates to
5 recombinant vectors, and especially expression vectors, containing the aforementioned genes, and to hosts transformed by these vectors. The hosts used may be prokaryotic or eukaryotic organisms. Examples of
10 prokaryotic organisms that may commonly be used as hosts include bacteria belonging to the genus *Escherichia*, such as *Escherichia coli*, and microorganisms belonging to the genus *Bacillus*, such as *Bacillus subtilis*.

Examples of eukaryotic hosts that may be used include lower eukaryotic organisms, for example,
15 eukaryotic microorganisms, for example, Eumycota such as yeast and filamentous fungi. As yeast there may be mentioned microorganisms belonging to the genus *Saccharomyces*, such as *Saccharomyces cerevisiae*, and as filamentous fungi there may be mentioned microorganisms
20 belonging to the genus *Aspergillus*, such as *Aspergillus oryzae* and *Aspergillus niger* and microorganisms belonging to the genus *Penicillium*. Animal cells and plant cells may also be used, the animal cells being cell lines from mice, hamsters, monkeys or humans. Insect cells, such as
25 silkworm cells, or the adult silkworms themselves, may also be used.

The expression vectors of the invention will include expression regulating regions such as a promoter and a terminator, a replication origin, etc., depending on the
30 type of hosts into which they are to be introduced. Examples of promoters for bacterial expression vectors which may be used include conventional promoters such as *trc* promoter, *tac* promoter, *lac* promoter, etc., examples of yeast promoters that may be used include
35 glyceraldehyde-3-phosphate dehydrogenase promoter, PH05 promoter, etc., and examples of filamentous fungi promoters that may be used include amylase promoter,

trpC, etc. Examples of animal cell host promoters that may be used include viral promoters such as SV40 early promoter, SV40 late promoter, etc.

5 The expression vector may be prepared according to a conventional method using restriction endonucleases, ligases and the like. The transformation of a host with an expression vector may also be carried out according to conventional methods.

10 The hosts transformed by the expression vector may be cultured, cultivated or raised, and the target protein may be recovered and purified from the cultured product, etc. according to conventional methods such as filtration, centrifugal separation, cell crushing, gel filtration chromatography, ion-exchange chromatography
15 and the like.

The present specification throughout discusses flavone synthases II derived from snapdragon, torenia and perilla that are capable of synthesizing flavones directly from flavanones, and it is also known that the
20 cytochrome P450 genes constitute a superfamily (DNA and Cell Biology, 12 (1993), Nelson et al., p.1) and that cytochrome P450 proteins within the same family have 40% or greater identity in their amino acid sequences while cytochrome P450 proteins within a subfamily have 55% or
25 greater identity in their amino acid sequences, and their genes hybridize to each other (Pharmacogenetics, 6 (1996), Nelson et al., p.1).

For example, a gene for flavonoid 3',5'-hydroxylase, which was a type of cytochrome P450 and participated in
30 the pathway of flavonoid synthesis, was first isolated from petunias (Nature, 366 (1993), Holton et al., p.276), and the petunia flavonoid 3',5'-hydroxylase gene was used as a probe to easily isolate a flavonoid 3',5'-hydroxylase gene from gentian (Plant Cell Physiol., 37
35 (1996), Tanaka et al., p.711), prairie-gentian, bellflower (WO93/18155 (1993), Kikuchi et al.), lavender, torenia and verbena (Shokubutsu no Kagaku Chosetsu, 33

26 AUG 2003

(1998), Tanaka et al., p.55).

Thus, a part or all of any of the flavone synthase II genes of the invention derived from snapdragon, torenia or perilla, which are capable of synthesizing flavones directly from flavanones, can be used as a probe, in order to obtain flavone synthase II genes capable of synthesizing flavones directly from flavanones, from different species of plants. Furthermore, by purifying the snapdragon-, torenia- or perilla-derived flavone synthase II enzymes described in this specification which can synthesize flavones directly from flavanones, and obtaining antibodies against the enzymes by conventional methods, it is possible to obtain different flavone synthase II proteins that react with the antibodies, and obtain genes coding for those proteins.

Consequently, the present invention is not limited merely to snapdragon-, torenia- or perilla-derived genes for flavone synthases II capable of synthesizing flavones directly from flavanones, but further relates to flavone synthases II derived from numerous other plants, which are capable of synthesizing flavones directly from flavanones. The sources for such flavone synthase II genes may be, in addition to snapdragon, torenia and perilla described here, also gentian, verbena, chrysanthemum, iris, commelina, centaurea, salvia, nemophila and the like, although the scope of the invention is not limited to these plants.

The invention still further relates to plants whose colors are modified by introducing a gene or genes for flavone synthases II that can synthesize flavones directly from flavanones, and to progenies of the plants or their tissues, which may also be in the form of cut flowers. By using the flavone synthases II or their genes which have been cloned according to the invention, it is possible to produce flavones in plant species or varieties that otherwise produce little or absolutely no

26 AUG 2003

flavones. By expressing the flavone synthase II gene or the genes in flower petals, it is possible to increase the amount of flavones in the flower petals, thus allowing the colors of the flowers to be modified toward the blue, for example.

Conversely, by repressing synthesis of flavones in flower petals, it is possible to modify the colors of the flowers toward the red, for example. However, flavones have myriad effects on flower colors, and the changes in flower colors are therefore not limited to those mentioned here. With the current level of technology, it is possible to introduce a gene into a plant and express the gene in a constitutive or tissue-specific manner, while it is also possible to repress the expression of a target gene by an antisense method or a cosuppression method.

As examples of transformable plants there may be mentioned rose, chrysanthemum, carnation, snapdragon, cyclamen, orchid, prairie-gentian, freesia, gerbera, gladiolus, baby's breath, kalanchoe, lily, pelargonium, geranium, petunia, torenia, tulip, rice, barley, wheat, rapeseed, potato, tomato, poplar, banana, eucalyptus, sweet potato, soybean, alfalfa, lupin, corn, etc., but there is no limitation to these.

Because flavones have various physiological activities as explained above, they can impart new physiological activity or economic value to plants. For example, by expressing the gene to produce flavones in roots, it is possible to promote growth of microorganisms that are beneficial for the plant, and thus promote growth of the plant. It is also possible to synthesize flavones that exhibit physiological activity in humans, animals or insects.

Examples

The invention will now be explained in further detail by way of the following examples. Unless

26 AUG 2003

otherwise specified, the molecular biological methods were carried out according to Molecular Cloning (Sambrook et al., 1989).

Example 1. Cloning of snapdragon flavone synthase II

5 gene

RNA was extracted from about 5 g of young buds of a Yellow Butterfly snapdragon (commercial name by Sakata-no-Tane, KK.), and polyA⁺ RNA was obtained by an Oligotex. This polyA⁺ RNA was used as a template to
10 prepare a cDNA library using a Lambda ZAPII cDNA Library Synthesis Kit (Stratagene) by the method recommended by Stratagene (Stratagene Instruction Manual, Revision #065001). The cDNA library was screened using the full length cDNA CYP93B1 as the probe. The screening and
15 detection of positive clones were carried out using a DIG-DNA-labeling and detection kit (Boehringer) based on the method recommended by the same company, under a low stringent condition.

Specifically, a hybridization buffer (5 x SSC, 30%
20 formamide, 50 mM sodium phosphate buffer (pH 7.0), 1% SDS, 2% blocking reagent (Boehringer), 0.1% lauroylsarcosine, 80 µg/ml salmon sperm DNA) was used for prehybridization at 42°C for 2 hours, after which the DIG-labeled probe was added and the mixture was kept
25 overnight. The membrane was rinsed in 5 x SSC rinsing solution containing 1% SDS at 65°C for 1.5 hours. One positive clone was obtained, and it was designated as ANFNS1. Upon determining the nucleotide sequence at the
30 5' end of ANFNS1 it was expected that ANFNS1 encodes a sequence with high identity with the flavanone 2-hydroxylase encoded by licorice CYP93B1, and it was assumed that it encoded a P450 with a function similar to that of flavanone 2-hydroxylase.

However, a comparison with the amino acid sequence
35 of flavanone 2-hydroxylase encoded by CYP93B1 suggested that the cDNA of ANFNS1 is not a full-length cDNA, lacking the portion corresponding to approximately 65

26 AUG 2003

amino acid residues from the initiating methionine. The ANFNS1 cDNA was therefore used as a probe for-rescreening of the snapdragon cDNA library, to obtain cDNA (ANFNS2) which was believed to include the full-length amino acid sequence. The protein encoded by ANFNS2 obtained here exhibited 53% identity on the amino acid level with flavanone 2-hydroxylase encoded by snapdragon CYP93B1. The nucleotide sequence of ANFNS2 is listed as SEQ.ID. No.1, and the amino acid sequence deduced therefrom is listed as SEQ.ID. No.2.

Example 2. Cloning of torenia flavone synthase II gene

RNA was extracted from approximately 2 g of buds of a torenia variety (variety name: Sunrenive, Variety Registration Application No.: 7433 according to the Seeds and Seedlings Law, by Suntory Ltd.) and the polyA⁺ RNA was obtained with an Oligotex. The polyA⁺ RNA was used as a template to prepare a cDNA library using a Lambda ZAPII cDNA Library Synthesis Kit (Stratagene) by the method recommended by Stratagene as mentioned in Example 1. The cDNA library was screened using a mixture of the aforementioned CYP93B1 cDNA and ANFNS1 cDNA as the probes. The screening and detection of positive clones were carried out under the low stringent conditions as described in Example 1.

One positive clone was obtained, and was designated as TFNS5. Upon determining the full nucleotide sequence of TFNS5 cDNA, it was found that the protein encoded by TFNS5 cDNA exhibited 52% identity on the amino acid level with flavanone-2-hydroxylase encoded by snapdragon CYP93B1. This TFNS5 cDNA also had high identity of 77% with the protein encoded by ANFNS2, the snapdragon-derived cDNA obtained in Example 1. The determined nucleotide sequence is listed as SEQ.ID. No.3, and the amino acid sequence deduced therefrom is listed as SEQ.ID. No.4.

Example 3. Expression of torenia flavone synthase II
gene in yeast

The following experiment was conducted in order to detect the enzyme activity of the protein encoded by TFNS5, the torenia cDNA obtained in Example 2. Parts of the outside of the translated region of the gene were modified to introduce restriction enzyme sites therein to prepare a sense primer (5'-
5 AAATAGGATCCAAGCatgGACACAGTCTTAA-3'; underline = BamHI site; lowercase letters: initiation codon) (SEQ.ID. No.5)
10 and an antisense primer (5'-
CCCTTCTAGAtcaAGCACCCGATATTGTGGCCGGG-3'; underline = XbaI site; lowercase letters: termination codon) (SEQ.ID. No.6) were used with KOD polymerase (Toyobo) for PCR
15 reaction. The PCR conditions were 98°C for one minute, 20 cycles of (98°C for 15 seconds, 55°C for 10 seconds, 74°C for 30 seconds), followed by 74°C for 10 minutes.

After introducing the resultant PCR product into the EcoRV site of pBluescriptII SK(-) (Stratagene), it was
20 digested with restriction enzymes BamHI and XbaI and introduced at the BamHI-XbaI sites of the yeast expression vector pYES2 (Invitrogen). The resultant plasmid was then introduced into BJ2168 yeast (Nihon Gene). The enzyme activity was measured by the method
25 described by Akashi et al. (FEBS Lett., 431 (1998), Akashi et al., p.287). The transformed yeast cells were cultured in 20 ml of selective medium (6.7 mg/ml amino acid-free yeast nitrogen base (Difco), 20 mg/ml glucose, 30 µg/ml leucine, 20 µg/ml tryptophan and 5 mg/ml
30 casamino acid), at 30°C for 24 hours.

After harvesting the yeast cells with centrifugation, the harvested yeast cells were cultured at 30°C for 48 hours in an expressing medium (10 mg/ml yeast extract, 10 mg/ml peptone, 2 µg/ml hemin, 20 mg/ml
35 galactose). After collecting the yeast cells, they were washed by suspending in water and collecting them. Glass beads were used for 10 minutes of disrupting, after which

the cells were centrifuged at 8000 x g for 10 minutes. The supernatant was further centrifuged at 15,000 x g for 10 minutes to obtain a crude enzyme fraction.

5 A mixture of 15 μ g of (R,S)-naringenin (dissolved in 30 μ l of 2-methoxyethanol), 1 ml of crude enzyme solution and 1 mM NADPH (total reaction mixture volume: 1.05 ml) was reacted at 30°C for 2 hours. After terminating the reaction by addition of 30 μ l of acetic acid, 1 ml of ethyl acetate was added and mixed therewith. After 10 centrifugation, the ethyl acetate layer was dried with an evaporator. The residue was dissolved in 100 μ l of methanol and analyzed by HPLC. The analysis was carried out according to the method described by Akashi et al. The acid treatment involved dissolution of the 15 evaporator-dried sample in 150 μ l of ethanol containing 10% hydrochloric acid, and stirring for 30 minutes. This was diluted with 1.3 ml of water, 800 μ l of ethyl acetate was further added and mixed therewith, and after centrifugation, the ethyl acetate layer was recovered. 20 This was then dried, dissolved in 200 μ l of methanol, and analyzed by HPLC.

The yeast expressing licorice CYP93B1 produced 2-hydroxynaringenin from naringenin, but yielded no apigenin (Fig. 1, A). Only upon acid treatment of the 25 reaction mixture, apigenin was yielded from 2-hydroxynaringenin (Fig. 1, B). In contrast, the yeast expressing torenia TFNS5 yielded apigenin from naringenin without acid treatment of the reaction mixture (Fig. 1, C). This demonstrated that TFNS5 encodes a flavone 30 synthase II.

Example 4. Expression of snapdragon flavone synthase II gene in yeast

An approximately 1400 bp DNA fragment obtained by digesting ANFNS2 cDNA with BamHI and SphI, an 35 approximately 350 bp DNA fragment obtained by digesting the same with SphI and BamHI, and pYES2 digested with BamHI and XhoI were ligated to obtain a plasmid, which

was then introduced into yeast by the same method as described in Example 3. The resultant recombinant yeast cells were used to measure the flavone synthesis activity by the same method as in Example 3. The yeast expressing
5 the snapdragon-derived ANFNS2 produced apigenin without acid treatment, thus demonstrating that ANFNS2 encodes a flavone synthase II.

Example 5. Construction of expression vector in plants

A plant expression vector was constructed to
10 introduce TFNS5, the torenia cDNA obtained in Example 2, into plants. After digesting pBE2113-GUS (Plant Cell Physiol., 37 (1996), Mitsuhashi et al., p.49) with SacI, a blunting kit (Takara) was used to blunt the ends, after
15 which a XhoI linker (Toyobo) was inserted. The resulting plasmid was then digested with HindIII and EcoRI, and an approximately 3 kb DNA fragment was recovered. The DNA fragment was linked to the HindIII/EcoRI site of the binary vector pBINPLUS to prepare pBE2113'. Vector
20 pBINPLUS used here was obtained by modifying the binary vector Bin19 (Nucl. Acids Res., 12 (1984), Bevan, p.8711), which is widely used for gene introduction into plants using *Agrobacterium* cells, in the manner reported by van Engelen et al. (Transgenic Research, 4 (1995), van Engelen et al., p.288).

25 The TFNS5 cDNA was cut out of SK(-) vector by cleavage with BamHI/XhoI, and an approximately 1.7 kb fragment thus obtained was ligated to the BamHI/XhoI sites of the aforementioned binary vector pBE2113'. The construct thus obtained, pSPB441, expresses TFNS5 cDNA in
30 the sense direction under the control of 35S cauliflower mosaic virus promoter having a double repeat of the enhancer sequence (Plant cell Physiol., 37 (1996), Mitsuhashi et al., p.49).

Example 6. Alteration of torenia flower color

35 A torenia variety (variety name: Sunrenive, Variety Registration Application No.: 7433 according to the Seeds and Seedlings Law, by Suntory Ltd.) was transformed with

pSPB441 constructed in Example 5 above, according to the method of Aida et al. (Breeding Science, 45 (1995), Aida et al., p.71). Over 95% of the obtained transformants showed alteration of the flower color from the dark purple of the parent strain to a light purple. The left and the right flower petal colors of four flower petals were measured. While the flower petal color of the parent strain was Number 89A according to the Royal Horticultural Society Color Chart, the typical flower petal colors of the transformants were 82C, 87D, 87C, 88D, 91A, etc. These results indicated that introduction of TFNS5 into plants can alter flower colors.

In the transformed individuals, the amount of flavones ranged 1/5 to 1/10 that of the host, while the amount of anthocyanins was reduced to about 1/3 that of the host. Also detected were flavanones (naringenin, eriodictyol and pentahydroxyflavanone) which are flavone biosynthesis precursors that were not detected in the host.

20 Example 7. Expression of flavone synthase in petunias

Plasmid pSPB441 was introduced into a petunia variety (variety name: Revolution Violet Mini, Variety Registration Application No.: 9217, according to the Seeds and Seedlings Law, by Suntory Ltd.) according to the method of Napoli et al. (Plant Cell, 2, (1990), Napoli et al., p.279). Changes in flower colors occurred in two of the resultant transformants, where the flower colors were lighter than the parent strain. The flower color of the parent strain was Number 88A according to the Royal Horticultural Society Color Chart, whereas 87A in the transformants. Also, while no flavones were detected in the parent strain, a flavone, luteolin, was detected in the transformed strains.

30 Example 8. Cloning of perilla flavone synthase II gene

35 The method described in Example 1 was used to screen a cDNA library prepared from leaves of red perilla (*Perilla frutescens*) according to the method of Gong et

26 AUG 2003

al. (Plant Mol. Biol., 35, (1997), Gong et al., p.915)
using λ gt10 (Stratagene) as the vector. Culturing, DNA
preparation and subcloning of the resulting phage clones
#3 were carried out according to the method of Gong et
5 al. (Plant Mol. Biol., 35, (1997), Gong et al., p.915),
and the nucleotide sequence was determined and listed as
SEQ.ID. No.7. The deduced amino acid sequence encoded by
this nucleotide sequence was listed as SEQ.ID. No.8.
This amino acid sequence showed 76% and 75% identity with
10 TFNS5 and ANFNS2, respectively. It also showed 52%
identity with CYP93B1.

Example 9. Expression of perilla flavone synthase II
gene in yeast

The phage clone #3 obtained in Example 8 was used as
15 a template for PCR by the method described in Example 3,
using Lambda Arm Primer (Stratagene). The amplified DNA
fragment was subcloned at the EcoRV site of pBluescript
KS(-). A clone with the initiation codon of perilla
flavone synthase II cDNA on the SalI site side of
20 pBluescript KS(-) was selected, and was designated as
pFS3. The nucleotide sequence of the cDNA insert of pFS3
was determined, and PCR was conducted to confirm the
absence of errors.

An approximately 1.8 kb DNA fragment obtained by
25 digesting pFS3 with SalI and XbaI was ligated with pYES2
digested with XhoI and XbaI (Example 3) to obtain a
plasmid which was designated as pYFS3, and this was
introduced into yeast BJ2168 by the method described in
Example 3. When the flavone synthase activity of this
30 recombinant yeast was measured by the method described in
Example 3, production of apigenin from naringenin was
confirmed, indicating that the perilla phage clone #3
cDNA encodes a protein with flavone synthase II activity.

Industrial Applicability

35 It is possible to alter flower colors by linking
cDNA of the invention to an appropriate plant expression
vector and introducing it into plants to express or

26 AUG 2003

inhibit expression of flavone synthases. Furthermore, by expressing the flavone synthase genes not only in petals but also in entire plants or their appropriate organs, it is possible to increase the resistance against
5 microorganisms of plants or to improve the nitrogen fixing ability of legumes by promoting association with rhizosphere microorganisms, as well as to improve the protective effects of plants against ultraviolet rays and light.

26 AUG 2003

CLAIMS

1. An isolated gene encoding a flavone synthase II with activity of directly synthesizing flavones from flavanones.

2. An isolated gene according to claim 1 which encodes a protein having the amino acid sequence listed as SEQ.ID. No. 2, 4 or 8 of the Sequence Listing and showing activity of synthesizing flavones from flavanones, or an isolated gene encoding a protein having one of these amino acid sequences wherein the amino acid sequence has been modified by additions or deletions of one or more amino acids and/or one or more substitution with different amino acids, and possessing activity of synthesizing flavones from flavanones.

3. An isolated gene according to any one of claims 1 to 2, which has at least 55% identity with the amino acid sequence listed as SEQ.ID. No. 2, 4 or 8 of the Sequence Listing and possesses activity of synthesizing flavones from flavanones.

4. An isolated gene according to any one of claims 1 to 3, which hybridizes with all or a part of any one of the base sequence listed as SEQ.ID. No. 1, 3 and 7 of the Sequence Listing under conditions of 5 x SSC, 50°C, and which encodes a protein possessing activity of synthesizing flavones from flavanones.

5. A vector comprising an isolated gene according to any one of claims 1 to 4.

6. A host transformed by a vector according to claim 5.

7. A protein encoded by a gene according to any one of claims 1 to 4.

8. A method of producing a protein with flavone-synthesizing activity, which is characterized by culturing or growing a host according to claim 6 and recovering said protein from said host.

9. A plant into which a gene according to any one

of claims 1 to 4 has been introduced, or progenies of said plant or a tissue thereof, which exhibits the same properties.

5 10 . A cut flower from a plant or a progeny thereof having the same properties, according to claim 9 .

11 . A method of altering a composition flavonoids and/or its amount using a gene according to any one of claims 1 to 4 .

10 12 . A method of altering the amount of a flavone using a gene according to any one of claims 1 to 4 .

13 . A method of altering the color of a flower using a gene according to any one of claims 1 to 4 .

14 . A method of bluing the color of a flower using a gene according to any one of claims 1 to 4 .

15 15 . A method of reddening the color of a flower using a gene according to any one of claims 1 to 4 .

16 . A method of altering the photosensitivity of a plant using a gene according to any one of claims 1 to 4 .

20 17 . A method of controlling the interaction between a plant and microorganisms using a gene according to any one of claims 1 to 4 .

26 AUG 2003

SEQUENCE LISTING

<110> SUNTORY LIMITED

<120> Gene coding for flavone synthesizing enzyme

<130> 993009

<160> 8

<210> 1

<211> 1724

<212> DNA

<213> Antirrhinum majus

<220>

<223> Nucleotide sequence encoding a protein having an activity to directly convert flavanone to flavone

<400> 1

```

gctttacaca cacacacaca cacacacaca caaacaata atg tct aca ctt gtc      54
                                     Met Ser Thr Leu Val
                                     1           5
tac agc aca ctc ttc atc ctc tca acc ctc ctc ctc acc ctc cta acc      102
Tyr Ser Thr Leu Phe Ile Leu Ser Thr Leu Leu Leu Thr Leu Leu Thr
          10           15           20
cgc acc cgc cgc aag acc cgc ccg ccc ggc cca tta gcc ctc ccc tta      150
Arg Thr Arg Arg Lys Thr Arg Pro Pro Gly Pro Leu Ala Leu Pro Leu
          25           30           35
ata ggc cac tta cac ctc ctc ggc cca aag ctc cac cac acc ttc cac      198
Ile Gly His Leu His Leu Leu Gly Pro Lys Leu His His Thr Phe His
          40           45           50
caa ttc tcc caa cgc tac ggc ccg ctc atc cag ctc tac ctc ggc tcc      246
Gln Phe Ser Gln Arg Tyr Gly Pro Leu Ile Gln Leu Tyr Leu Gly Ser
          55           60           65

```

26 AUG 2003

gtc cca tgc gtc gtc gct tcc acg ccc gaa ctc gcc cgc gaa ttc ctc	294
Val Pro Cys Val Val Ala Ser Thr Pro Glu Leu Ala Arg Glu Phe Leu	
70 75 80 85	
aag acg cac gaa ctc gac ttc tcg tcc cgc aag cac tcc acc gcc atc	342
Lys Thr His Glu Leu Asp Phe Ser Ser Arg Lys His Ser Thr Ala Ile	
90 95 100	
gac atc gtc acg tac gac tcc tcg ttc gcc ttc gcg ccg tac ggg ccg	390
Asp Ile Val Thr Tyr Asp Ser Ser Phe Ala Phe Ala Pro Tyr Gly Pro	
105 110 115	
tac tgg aaa ttc atc aag aaa tta tgt act tac gag cta ctg ggt gcc	438
Tyr Trp Lys Phe Ile Lys Lys Leu Cys Thr Tyr Glu Leu Leu Gly Ala	
120 125 130	
cgg aac ttg agc cat ttc cag ccc att aga gct ttg gag gtc aac agt	486
Arg Asn Leu Ser His Phe Gln Pro Ile Arg Ala Leu Glu Val Asn Ser	
135 140 145	
ttc ttg aga att ttg tac gag aaa aca gag cag aaa cag agt gtt aat	534
Phe Leu Arg Ile Leu Tyr Glu Lys Thr Glu Gln Lys Gln Ser Val Asn	
150 155 160 165	
gtg act gag gag ctt gtg aag ctg acg agt aat gtg atc agt aac atg	582
Val Thr Glu Glu Leu Val Lys Leu Thr Ser Asn Val Ile Ser Asn Met	
170 175 180	
atg ttg ggg atc agg tgt tcg ggg acg gaa ggg gag gcg gag gtg gcg	630
Met Leu Gly Ile Arg Cys Ser Gly Thr Glu Gly Glu Ala Glu Val Ala	
185 190 195	
agg acg gtg ata agg gag gtg acg cag ata ttt ggg gag ttt gat gtg	678
Arg Thr Val Ile Arg Glu Val Thr Gln Ile Phe Gly Glu Phe Asp Val	
200 205 210	
tcg gag att gtt tgg ttt tgt aag aat ttg gat ctg cag ggg att agg	726
Ser Glu Ile Val Trp Phe Cys Lys Asn Leu Asp Leu Gln Gly Ile Arg	
215 220 225	
aag agg tcg gag gat att agg agg agg tat gat gct ttg ttg gag aag	774
Lys Arg Ser Glu Asp Ile Arg Arg Arg Tyr Asp Ala Leu Leu Glu Lys	
230 235 240 245	
att att agt gat agg gag agg ttg agg ttg agg ggg ggt ggt ggt gga	822
Ile Ile Ser Asp Arg Glu Arg Leu Arg Leu Arg Gly Gly Gly Gly Gly	
250 255 260	

26 AUG 2003

ggg ggt gga gag gtg aag gat ttt ttg gat atg ttg ttg gat gtg atg	870
Gly Gly Gly Glu Val Lys Asp Phe Leu Asp Met Leu Leu Asp Val Met	
265 270 275	
gag agt gag aaa tcg gag gtg gag ttt acg agg gag cat ctc aaa gct	918
Glu Ser Glu Lys Ser Glu Val Glu Phe Thr Arg Glu His Leu Lys Ala	
280 285 290	
ttg att ctg gat ttc ttc act gcc ggt aca gac aca aca gca atc aca	966
Leu Ile Leu Asp Phe Phe Thr Ala Gly Thr Asp Thr Thr Ala Ile Thr	
295 300 305	
aca gaa tgg gca ata gca gaa ctc att agc aat cca aat gta ctc aaa	1014
Thr Glu Trp Ala Ile Ala Glu Leu Ile Ser Asn Pro Asn Val Leu Lys	
310 315 320 325	
aaa gct caa gaa gag atg gac aaa gtc ata gga tca caa agg ttg ttg	1062
Lys Ala Gln Glu Glu Met Asp Lys Val Ile Gly Ser Gln Arg Leu Leu	
330 335 340	
caa gaa tcc gac gcc cct aac ttg cct tac ctc aac gcg atc ata aaa	1110
Gln Glu Ser Asp Ala Pro Asn Leu Pro Tyr Leu Asn Ala Ile Ile Lys	
345 350 355	
gaa acg ttc cgt ctc cac cct cca atc ccc atg ctc act aga aaa tca	1158
Glu Thr Phe Arg Leu His Pro Pro Ile Pro Met Leu Thr Arg Lys Ser	
360 365 370	
att tct gac gtt gtg gtc aac ggg tac acg atc cct gcc aaa acg cta	1206
Ile Ser Asp Val Val Val Asn Gly Tyr Thr Ile Pro Ala Lys Thr Leu	
375 380 385	
ttg ttt gtc aac ctt tgg tcc atg gga agg aat cct aac tac tgg gaa	1254
Leu Phe Val Asn Leu Trp Ser Met Gly Arg Asn Pro Asn Tyr Trp Glu	
390 395 400 405	
aat ccg atg gag ttc cga ccc gag agg ttt ctc gag aaa ggg acc ggg	1302
Asn Pro Met Glu Phe Arg Pro Glu Arg Phe Leu Glu Lys Gly Thr Gly	
410 415 420	
tcg ata gac gtt aaa ggg cag cat ttc gag ttg ctg ccg ttt ggc acg	1350
Ser Ile Asp Val Lys Gly Gln His Phe Glu Leu Leu Pro Phe Gly Thr	
425 430 435	
ggc agg cgg ggc tgc ccg ggg atg ttg tta ggc atg cag gag ttg ttt	1398
Gly Arg Arg Gly Cys Pro Gly Met Leu Leu Gly Met Gln Glu Leu Phe	
440 445 450	

26 AUG 2003

```

agt att atc ggg gct atg gtg cag tgc ttc gat tgg aaa ctg ccc gat      1446
Ser Ile Ile Gly Ala Met Val Gln Cys Phe Asp Trp Lys Leu Pro Asp
      455              460              465

ggt gtg aag tcg gtc gac atg acc gag cgg ccc ggg ttg acg gct cca      1494
Gly Val Lys Ser Val Asp Met Thr Glu Arg Pro Gly Leu Thr Ala Pro
470              475              480              485

cgt gcc aat gat ttg gtg tgc caa ttg gtg cca cgg att gac ccg gtc      1542
Arg Ala Asn Asp Leu Val Cys Gln Leu Val Pro Arg Ile Asp Pro Val
      490              495              500

gtt gtc tcc gga ccg tgaaccttaa ggtagtatcg ataatctggt taatt      1592
Val Val Ser Gly Pro
      505

aaattgttat ttgttgtag gatttgattt ttgttatgta tgattatgcg tggattaaga      1652
taagcctgca aggacaaatt ccctttcttt gattgatgtc aatgagtttg tgtcaaaaaa      1712
aaaaaaaaaa aa      1724

```

<210> 2

<211> 506

<212> PRT

<213> Antirrhinum majus

<220>

<223> Amino acid sequence of a protein having an activity to
directly convert flavanone to flavone

<400> 2

```

Met Ser Thr Leu Val Tyr Ser Thr Leu Phe Ile Leu Ser Thr Leu Leu
  1              5              10              15
Leu Thr Leu Leu Thr Arg Thr Arg Arg Lys Thr Arg Pro Pro Gly Pro
      20              25              30
Leu Ala Leu Pro Leu Ile Gly His Leu His Leu Leu Gly Pro Lys Leu
      35              40              45
His His Thr Phe His Gln Phe Ser Gln Arg Tyr Gly Pro Leu Ile Gln
      50              55              60
Leu Tyr Leu Gly Ser Val Pro Cys Val Val Ala Ser Thr Pro Glu Leu
      65              70              75              80
Ala Arg Glu Phe Leu Lys Thr His Glu Leu Asp Phe Ser Ser Arg Lys
      85              90              95

```

26 AUG 2003

```

His Ser Thr Ala Ile Asp Ile Val Thr Tyr Asp Ser Ser Phe Ala Phe
      100                      105                      110
Ala Pro Tyr Gly Pro Tyr Trp Lys Phe Ile Lys Lys Leu Cys Thr Tyr
      115                      120                      125
Glu Leu Leu Gly Ala Arg Asn Leu Ser His Phe Gln Pro Ile Arg Ala
      130                      135                      140
Leu Glu Val Asn Ser Phe Leu Arg Ile Leu Tyr Glu Lys Thr Glu Gln
      145                      150                      155                      160
Lys Gln Ser Val Asn Val Thr Glu Glu Leu Val Lys Leu Thr Ser Asn
      165                      170                      175
Val Ile Ser Asn Met Met Leu Gly Ile Arg Cys Ser Gly Thr Glu Gly
      180                      185                      190
Glu Ala Glu Val Ala Arg Thr Val Ile Arg Glu Val Thr Gln Ile Phe
      195                      200                      205
Gly Glu Phe Asp Val Ser Glu Ile Val Trp Phe Cys Lys Asn Leu Asp
      210                      215                      220
Leu Gln Gly Ile Arg Lys Arg Ser Glu Asp Ile Arg Arg Arg Tyr Asp
      225                      230                      235                      240
Ala Leu Leu Glu Lys Ile Ile Ser Asp Arg Glu Arg Leu Arg Leu Arg
      245                      250                      255
Gly Gly Gly Gly Gly Gly Gly Gly Glu Val Lys Asp Phe Leu Asp Met
      260                      265                      270
Leu Leu Asp Val Met Glu Ser Glu Lys Ser Glu Val Glu Phe Thr Arg
      275                      280                      285
Glu His Leu Lys Ala Leu Ile Leu Asp Phe Phe Thr Ala Gly Thr Asp
      290                      295                      300
Thr Thr Ala Ile Thr Thr Glu Trp Ala Ile Ala Glu Leu Ile Ser Asn
      305                      310                      315                      320
Pro Asn Val Leu Lys Lys Ala Gln Glu Glu Met Asp Lys Val Ile Gly
      325                      330                      335
Ser Gln Arg Leu Leu Gln Glu Ser Asp Ala Pro Asn Leu Pro Tyr Leu
      340                      345                      350
Asn Ala Ile Ile Lys Glu Thr Phe Arg Leu His Pro Pro Ile Pro Met
      355                      360                      365
Leu Thr Arg Lys Ser Ile Ser Asp Val Val Val Asn Gly Tyr Thr Ile
      370                      375                      380
Pro Ala Lys Thr Leu Leu Phe Val Asn Leu Trp Ser Met Gly Arg Asn
      385                      390                      395                      400

```

26 AUG 2003

Pro Asn Tyr Trp Glu Asn Pro Met Glu Phe Arg Pro Glu Arg Phe Leu
 405 410 415
 Glu Lys Gly Thr Gly Ser Ile Asp Val Lys Gly Gln His Phe Glu Leu
 420 425 430
 Leu Pro Phe Gly Thr Gly Arg Arg Gly Cys Pro Gly Met Leu Leu Gly
 435 440 445
 Met Gln Glu Leu Phe Ser Ile Ile Gly Ala Met Val Gln Cys Phe Asp
 450 455 460
 Trp Lys Leu Pro Asp Gly Val Lys Ser Val Asp Met Thr Glu Arg Pro
 465 470 475 480
 Gly Leu Thr Ala Pro Arg Ala Asn Asp Leu Val Cys Gln Leu Val Pro
 485 490 495
 Arg Ile Asp Pro Val Val Val Ser Gly Pro
 500 505

<210> 3

<211> 1730

<212> DNA

<213> Torenia hybrida

<220>

<223> Nucleotide sequence encoding a protein having an
 activity to directly convert flavanone to flavone

<400> 3

atcgaaaccg ctatatcatt acatttacaa cagcgctaaa aaaatatata taaagc	56
atg gac aca gtc tta atc aca ctc tac acc gcc ctg ttc gtc atc acc	104
Met Asp Thr Val Leu Ile Thr Leu Tyr Thr Ala Leu Phe Val Ile Thr	
1 5 10 15	
acc acc ttc ctc ctc ctc ctc cgc cga agg gga cca ccg tct ccg ccc	152
Thr Thr Phe Leu Leu Leu Leu Arg Arg Arg Gly Pro Pro Ser Pro Pro	
20 25 30	
ggt cct ctc tcc cta ccc ata att ggc cac ctc cac ctc ctc ggc cca	200
Gly Pro Leu Ser Leu Pro Ile Ile Gly His Leu His Leu Leu Gly Pro	
35 40 45	
aga ctc cac cac acg ttc cat gaa ttc tca ctc aaa tac ggc cca ttg	248
Arg Leu His His Thr Phe His Glu Phe Ser Leu Lys Tyr Gly Pro Leu	
50 55 60	

atc cag ctc aag ctc ggc tcg atc ccg tgc gtc gtg gcc tcg acg ccc	296
Ile Gln Leu Lys Leu Gly Ser Ile Pro Cys Val Val Ala Ser Thr Pro	
65 70 75 80	
gag ctc gcg aga gag ttt ctt aag acg aac gag ctc gcg ttc tcc tct	344
Glu Leu Ala Arg Glu Phe Leu Lys Thr Asn Glu Leu Ala Phe Ser Ser	
85 90 95	
cgc aag cac tct acg gcc ata gac atc gtc acc tac gac tcg tcc ttt	392
Arg Lys His Ser Thr Ala Ile Asp Ile Val Thr Tyr Asp Ser Ser Phe	
100 105 110	
gct ttc tct ccg tac gga ccc tac tgg aag tac atc aag aaa ctg tgt	440
Ala Phe Ser Pro Tyr Gly Pro Tyr Trp Lys Tyr Ile Lys Lys Leu Cys	
115 120 125	
acc tac gag ctg ctc gga gcg agg aac ctc gga cac ttt cag ccc att	488
Thr Tyr Glu Leu Leu Gly Ala Arg Asn Leu Gly His Phe Gln Pro Ile	
130 135 140	
agg aat ctc gag gtc agg tcc ttt ctg cag ctt ctg atg cac aag agc	536
Arg Asn Leu Glu Val Arg Ser Phe Leu Gln Leu Leu Met His Lys Ser	
145 150 155 160	
ttt aag ggc gag agt gtg aat gtg aca gac gag ctg gtg agg ctg acg	584
Phe Lys Gly Glu Ser Val Asn Val Thr Asp Glu Leu Val Arg Leu Thr	
165 170 175	
agc aat gtg ata tcc cac atg atg ctg agc ata agg tgc tcg gaa gat	632
Ser Asn Val Ile Ser His Met Met Leu Ser Ile Arg Cys Ser Glu Asp	
180 185 190	
gaa ggc gat gct gag gcg gcg aga aca gtg ata cgc gag gtg acg cag	680
Glu Gly Asp Ala Glu Ala Ala Arg Thr Val Ile Arg Glu Val Thr Gln	
195 200 205	
ata ttt ggg gaa ttc gat gtt acg gac ata ata tgg ttt tgc aag aaa	728
Ile Phe Gly Glu Phe Asp Val Thr Asp Ile Ile Trp Phe Cys Lys Lys	
210 215 220	
ttc gat ctg cag ggg ata aag aag agg tca gag gat att cag agg agg	776
Phe Asp Leu Gln Gly Ile Lys Lys Arg Ser Glu Asp Ile Gln Arg Arg	
225 230 235 240	
tat gat gct ttg ctc gag aag att att agt gat aga gag aga tcg agg	824
Tyr Asp Ala Leu Leu Glu Lys Ile Ile Ser Asp Arg Glu Arg Ser Arg	
245 250 255	

26 AUG 2003

agg caa aat cgt gat aag cat ggt ggc ggt aac aat gag gag gcc aag	872
Arg Gln Asn Arg Asp Lys His Gly Gly Gly Asn Asn Glu Glu Ala Lys	
260 265 270	
gat ttt ctt gat atg ttg ctt gat gtg atg gag agt ggg gac acg gag	920
Asp Phe Leu Asp Met Leu Leu Asp Val Met Glu Ser Gly Asp Thr Glu	
275 280 285	
gtc aaa ttc act aga gag cat ctc aag gct ttg att ctg gat ttc ttc	968
Val Lys Phe Thr Arg Glu His Leu Lys Ala Leu Ile Leu Asp Phe Phe	
290 295 300	
acg gcc ggt acg gac aca aca gcc ata gcc acc gag tgg gcc atc gcc	1016
Thr Ala Gly Thr Asp Thr Thr Ala Ile Ala Thr Glu Trp Ala Ile Ala	
305 310 315 320	
gag ctc atc aac aac ccg aac gtc ttg aag aag gcc caa gaa gaa ata	1064
Glu Leu Ile Asn Asn Pro Asn Val Leu Lys Lys Ala Gln Glu Glu Ile	
325 330 335	
tcc cgg atc atc gga acc aag cgg atc gta caa gaa tcc gac gcc cca	1112
Ser Arg Ile Ile Gly Thr Lys Arg Ile Val Gln Glu Ser Asp Ala Pro	
340 345 350	
gac cta ccc tac ctc cag gcc atc atc aag gag acg ttc cgg ctc cac	1160
Asp Leu Pro Tyr Leu Gln Ala Ile Ile Lys Glu Thr Phe Arg Leu His	
355 360 365	
cca ccg atc ccg atg ctc tcg cgt aag tcc acc tcc gat tgc acg gtc	1208
Pro Pro Ile Pro Met Leu Ser Arg Lys Ser Thr Ser Asp Cys Thr Val	
370 375 380	
aac ggc tac aaa atc caa gcc aag agc ctc ttg ttc gtg aac ata tgg	1256
Asn Gly Tyr Lys Ile Gln Ala Lys Ser Leu Leu Phe Val Asn Ile Trp	
385 390 395 400	
tcc atc ggt cga aac cct aat tac tgg gaa agc cct atg gag ttc agg	1304
Ser Ile Gly Arg Asn Pro Asn Tyr Trp Glu Ser Pro Met Glu Phe Arg	
405 410 415	
ccc gag cgg ttc ttg gag aag gga cgc gag tcc atc gac gtc aag ggc	1352
Pro Glu Arg Phe Leu Glu Lys Gly Arg Glu Ser Ile Asp Val Lys Gly	
420 425 430	
cag cac ttt gag ctc ttg cct ttt ggg acg ggc cgc agg ggc tgt ccc	1400
Gln His Phe Glu Leu Leu Pro Phe Gly Thr Gly Arg Arg Gly Cys Pro	
435 440 445	

26 AUG 2003

```

ggt atg ttg ctg gct ata caa gag gtg gtc agc atc att ggg acc atg      .448
Gly Met Leu Leu Ala Ile Gln Glu Val Val Ser Ile Ile Gly Thr Met
      450              455              460
ggt cag tgc ttc gac tgg aaa ttg gca gat ggt tgc ggc aat aat gtg      1496
Val Gln Cys Phe Asp Trp Lys Leu Ala Asp Gly Ser Gly Asn Asn Val
      465              470              475              480
gac atg acc gaa cgg tct gga ttg acc gct ccg aga gcg ttc gat ctg      1544
Asp Met Thr Glu Arg Ser Gly Leu Thr Ala Pro Arg Ala Phe Asp Leu
      485              490              495
ggt tgc cgg ttg tat cca cgg gtt gac ccg gcc aca ata tgc ggt gct      1592
Val Cys Arg Leu Tyr Pro Arg Val Asp Pro Ala Thr Ile Ser Gly Ala
      500              505              510              512
tgatgtagta gggtagggcg cgtgttggtg tttatcttt cggttttgtt ctgttagtat      1652
tattatggtc tgtgttgaag cctcaaggat tttaaaaaaa aaaaaaaaaa aaaaaaaaaa      1712
aaaaaaaaaa aaaaaaaaaa                                         1730

```

<210> 4

<211> 512

<212> PRT

<213> *Torenia hybrida*

<220>

<223> Amino acid sequence of a protein having an activity to
directly convert flavanone to flavone

<400> 4

```

Met Asp Thr Val Leu Ile Thr Leu Tyr Thr Ala Leu Phe Val Ile Thr
  1              5              10              15
Thr Thr Phe Leu Leu Leu Leu Arg Arg Arg Gly Pro Pro Ser Pro Pro
      20              25              30
Gly Pro Leu Ser Leu Pro Ile Ile Gly His Leu His Leu Leu Gly Pro
      35              40              45
Arg Leu His His Thr Phe His Glu Phe Ser Leu Lys Tyr Gly Pro Leu
      50              55              60
Ile Gln Leu Lys Leu Gly Ser Ile Pro Cys Val Val Ala Ser Thr Pro
      65              70              75              80
Glu Leu Ala Arg Glu Phe Leu Lys Thr Asn Glu Leu Ala Phe Ser Ser
      85              90              95

```

26 AUG 2003

Arg Lys His Ser Thr Ala Ile Asp Ile Val Thr Tyr Asp Ser Ser Phe
100 105 110
Ala Phe Ser Pro Tyr Gly Pro Tyr Trp Lys Tyr Ile Lys Lys Leu Cys
115 120 125
Thr Tyr Glu Leu Leu Gly Ala Arg Asn Leu Gly His Phe Gln Pro Ile
130 135 140
Arg Asn Leu Glu Val Arg Ser Phe Leu Gln Leu Leu Met His Lys Ser
145 150 155 160
Phe Lys Gly Glu Ser Val Asn Val Thr Asp Glu Leu Val Arg Leu Thr
165 170 175
Ser Asn Val Ile Ser His Met Met Leu Ser Ile Arg Cys Ser Glu Asp
180 185 190
Glu Gly Asp Ala Glu Ala Ala Arg Thr Val Ile Arg Glu Val Thr Gln
195 200 205
Ile Phe Gly Glu Phe Asp Val Thr Asp Ile Ile Trp Phe Cys Lys Lys
210 215 220
Phe Asp Leu Gln Gly Ile Lys Lys Arg Ser Glu Asp Ile Gln Arg Arg
225 230 235 240
Tyr Asp Ala Leu Leu Glu Lys Ile Ile Ser Asp Arg Glu Arg Ser Arg
245 250 255
Arg Gln Asn Arg Asp Lys His Gly Gly Gly Asn Asn Glu Glu Ala Lys
260 265 270
Asp Phe Leu Asp Met Leu Leu Asp Val Met Glu Ser Gly Asp Thr Glu
275 280 285
Val Lys Phe Thr Arg Glu His Leu Lys Ala Leu Ile Leu Asp Phe Phe
290 295 300
Thr Ala Gly Thr Asp Thr Thr Ala Ile Ala Thr Glu Trp Ala Ile Ala
305 310 315 320
Glu Leu Ile Asn Asn Pro Asn Val Leu Lys Lys Ala Gln Glu Glu Ile
325 330 335
Ser Arg Ile Ile Gly Thr Lys Arg Ile Val Gln Glu Ser Asp Ala Pro
340 345 350
Asp Leu Pro Tyr Leu Gln Ala Ile Ile Lys Glu Thr Phe Arg Leu His
355 360 365
Pro Pro Ile Pro Met Leu Ser Arg Lys Ser Thr Ser Asp Cys Thr Val
370 375 380
Asn Gly Tyr Lys Ile Gln Ala Lys Ser Leu Leu Phe Val Asn Ile Trp
385 390 395 400

26 AUG 2003

Ser Ile Gly Arg Asn Pro Asn Tyr Trp Glu Ser Pro Met Glu Phe Arg
405 410 415
Pro Glu Arg Phe Leu Glu Lys Gly Arg Glu Ser Ile Asp Val Lys Gly
420 425 430
Gln His Phe Glu Leu Leu Pro Phe Gly Thr Gly Arg Arg Gly Cys Pro
435 440 445
Gly Met Leu Leu Ala Ile Gln Glu Val Val Ser Ile Ile Gly Thr Met
450 455 460
Val Gln Cys Phe Asp Trp Lys Leu Ala Asp Gly Ser Gly Asn Asn Val
465 470 475 480
Asp Met Thr Glu Arg Ser Gly Leu Thr Ala Pro Arg Ala Phe Asp Leu
485 490 495
Val Cys Arg Leu Tyr Pro Arg Val Asp Pro Ala Thr Ile Ser Gly Ala
500 505 510

<210> 5
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 5
aaataggatc caagcatgga cacagtctta a

31

<210> 6
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 6
cccttctaga tcaagcaccc gatattgtgg ccggg

35

SEQUENCE LISTING

<210> 7

<211> 1770

<212> DNA

<213> *Perilla frutescens*

<220>

<223> Nucleotide sequence encoding a protein having an
activity to directly convert flavanone to flavone

<400> 7

tgctcgacgga gcaagtggaa atg gca ctg tac gcc gcc ctc ttc ctc ctg tcc	53
Met Ala Leu Tyr Ala Ala Leu Phe Leu Leu Ser	
1 5 10	
gcc gcc gtg gtc cgc tcc gtt ctg gat cga aaa cgc ggg cgg ccg ccc	101
Ala Ala Val Val Arg Ser Val Leu Asp Arg Lys Arg Gly Arg Pro Pro	
15 20 25	
tac cct ccc ggg ccg ttc cct ctt ccc atc atc ggc cac tta cac ctc	149
Tyr Pro Pro Gly Pro Phe Pro Leu Pro Ile Ile Gly His Leu His Leu	
30 35 40	
ctc ggg ccg aga ctc cac caa acc ttc cac gat ctg tcc caa cgg tac	197
Leu Gly Pro Arg Leu His Gln Thr Phe His Asp Leu Ser Gln Arg Tyr	
45 50 55	
ggg ccc tta atg cag ctc cgc ctc ggg tcc atc cgc tgc gtc att gct	245
Gly Pro Leu Met Gln Leu Arg Leu Gly Ser Ile Arg Cys Val Ile Ala	
60 65 70 75	
gcc tcg ccg gag ctc gcc aag gaa tgc ctc aag aca cac gag ctc gtc	293
Ala Ser Pro Glu Leu Ala Lys Glu Cys Leu Lys Thr His Glu Leu Val	
80 85 90	
ttc tcc tcc cgc aaa cac tcc acc gcc att gat atc gtc acc tac gat	341
Phe Ser Ser Arg Lys His Ser Thr Ala Ile Asp Ile Val Thr Tyr Asp	
95 100 105	

26 AUG 2003

tca tcc ttc gct ttc tct ccc tac ggg cct tac tgg aaa ttc atc aag	389
Ser Ser Phe Ala Phe Ser Pro Tyr Gly Pro Tyr Trp Lys Phe Ile Lys	
110 115 120	
aaa tta tgc acc tac gag ctg ctc ggg gcc cga aat ctc gcc cac ttt	437
Lys Leu Cys Thr Tyr Glu Leu Leu Gly Ala Arg Asn Leu Ala His Phe	
125 130 135	
cag ccc atc agg act ctc gaa gtc aag tct ttc ctc caa att ctt atg	485
Gln Pro Ile Arg Thr Leu Glu Val Lys Ser Phe Leu Gln Ile Leu Met	
140 145 150 155	
cgc aag ggt gaa tcg ggg gag agc ttc aac gtg act gag gag ctc gtg	533
Arg Lys Gly Glu Ser Gly Glu Ser Phe Asn Val Thr Glu Glu Leu Val	
160 165 170	
aag ctg acg agc aac gtc ata tcg cat atg atg ctg agc ata cgg tgt	581
Lys Leu Thr Ser Asn Val Ile Ser His Met Met Leu Ser Ile Arg Cys	
175 180 185	
tca gag acg gag tcg gag gcg gag gcg gcg agg acg gtg att cgg gag	629
Ser Glu Thr Glu Ser Glu Ala Glu Ala Ala Arg Thr Val Ile Arg Glu	
190 195 200	
gtc acg cag ata ttt ggg gag ttc gac gtc tcc gac atc ata tgg ctt	677
Val Thr Gln Ile Phe Gly Glu Phe Asp Val Ser Asp Ile Ile Trp Leu	
205 210 215	
tgt aag aac ttc gat ttc caa ggt ata agg aag cgg tcc gag gat atc	725
Cys Lys Asn Phe Asp Phe Gln Gly Ile Arg Lys Arg Ser Glu Asp Ile	
220 225 230 235	
cag agg aga tat gat gct ctg ctg gag aag atc atc acc gac aga gag	773
Gln Arg Arg Tyr Asp Ala Leu Leu Glu Lys Ile Ile Thr Asp Arg Glu	
240 245 250	
aag cag agg cgg acc cac ggc ggc ggt ggc ggc ggc ggg gaa gcc aag	821
Lys Gln Arg Arg Thr His Gly Gly Gly Gly Gly Gly Gly Glu Ala Lys	
255 260 265	
gat ttt ctt gac atg ttc ctc gac ata atg gag agc ggg aaa gcc gaa	869
Asp Phe Leu Asp Met Phe Leu Asp Ile Met Glu Ser Gly Lys Ala Glu	
270 275 280	
gtt aaa ttc acg agg gag cat ctc aaa gct ttg att ctg gat ttc ttc	917
Val Lys Phe Thr Arg Glu His Leu Lys Ala Leu Ile Leu Asp Phe Phe	
285 290 295	

26 AUG 2003

acc gcc ggc acc gac acg acg gcg atc gtg tgt gaa tgg gcg ata gca	965
Thr Ala Gly Thr Asp Thr Thr Ala Ile Val Cys Glu Trp Ala Ile Ala	
300 305 310 315	
gaa gtg atc aac aat cca aat gtg ttg aag aaa gct caa gaa gag att	1013
Glu Val Ile Asn Asn Pro Asn Val Leu Lys Lys Ala Gln Glu Glu Ile	
320 325 330	
gcc aac atc gtc gga ttc gac aga att ctg caa gaa tcc gac gcc cca	1061
Ala Asn Ile Val Gly Phe Asp Arg Ile Leu Gln Glu Ser Asp Ala Pro	
335 340 345	
aat ctg ccc tac ctt caa gcc ctc atc aaa gaa aca ttc cgg ctc cac	1109
Asn Leu Pro Tyr Leu Gln Ala Leu Ile Lys Glu Thr Phe Arg Leu His	
350 355 360	
cct cca atc cca atg ctg gcg agg aaa tcg atc tcc gac tgc gtc atc	1157
Pro Pro Ile Pro Met Leu Ala Arg Lys Ser Ile Ser Asp Cys Val Ile	
365 370 375	
gac ggc tac atg att ccg gcc aac acg ctg ctc ttc gtc aac ctc tgg	1205
Asp Gly Tyr Met Ile Pro Ala Asn Thr Leu Leu Phe Val Asn Leu Trp	
380 385 390 395	
tcc atg ggg cgg aac cct aaa atc tgg gac tac ccg acg gcg ttc cag	1253
Ser Met Gly Arg Asn Pro Lys Ile Trp Asp Tyr Pro Thr Ala Phe Gln	
400 405 410	
ccg gag agg ttt ctg gag aag gaa aag gcc gcc atc gat gtt aaa ggg	1301
Pro Glu Arg Phe Leu Glu Lys Glu Lys Ala Ala Ile Asp Val Lys Gly	
415 420 425	
cag cat ttt gag ctg cta ccg ttc gga acg ggc agg aga ggc tgc cca	1349
Gln His Phe Glu Leu Leu Pro Phe Gly Thr Gly Arg Arg Gly Cys Pro	
430 435 440	
ggg atg ctt tta gcc att cag gag gtg gtc atc ata att ggg acg atg	1397
Gly Met Leu Leu Ala Ile Gln Glu Val Val Ile Ile Ile Gly Thr Met	
445 450 455	
att caa tgc ttc gat tgg aag ctg ccc gac ggc tcc ggc cat gtt gat	1445
Ile Gln Cys Phe Asp Trp Lys Leu Pro Asp Gly Ser Gly His Val Asp	
460 465 470 475	
atg gca gaa cgg cca ggg ctc acg gca ccg cga gag acc gat ttg ttt	1493
Met Ala Glu Arg Pro Gly Leu Thr Ala Pro Arg Glu Thr Asp Leu Phe	
480 485 490	

26 AUG 2003

tgc cgt gtg gtg ccg cga gtt gat ccg ttg gtt gtt tcc acc cag	1538
Cys Arg Val Val Pro Arg Val Asp Pro Leu Val Val Ser Thr Gln	
495 500 505	
tgatcacccc ctttaaattt attaatgata tattttttatt ttgagaaaaa ataaaaatgc	1598
taattgtttt gtttcatgat gtaattgtta attagtttct attgtgcgct gtcgcgtgct	1658
gcgtggctta agataagatt gtatcattgg tacctaggat gtattttcat tttcaataaa	1718
ttattttgtg ctgtgtatat taaaaaaaaa aaagaaaaaa aaaaaaaaaa aa	1770

<210> 8

<211>

<212> PRT

<213> Perilla frutescens

<220>

<223> Amino acid sequence of a protein having an activity to
directly convert flavanone to flavone

<400> 8

Met Ala Leu Tyr Ala Ala Leu Phe Leu Leu Ser Ala Ala Val Val Arg	
1 5 10 15	
Ser Val Leu Asp Arg Lys Arg Gly Arg Pro Pro Tyr Pro Pro Gly Pro	
20 25 30	
Phe Pro Leu Pro Ile Ile Gly His Leu His Leu Leu Gly Pro Arg Leu	
35 40 45	
His Gln Thr Phe His Asp Leu Ser Gln Arg Tyr Gly Pro Leu Met Gln	
50 55 60	
Leu Arg Leu Gly Ser Ile Arg Cys Val Ile Ala Ala Ser Pro Glu Leu	
65 70 75 80	
Ala Lys Glu Cys Leu Lys Thr His Glu Leu Val Phe Ser Ser Arg Lys	
85 90 95	
His Ser Thr Ala Ile Asp Ile Val Thr Tyr Asp Ser Ser Phe Ala Phe	
100 105 110	

26 AUG 2003

Ser	Pro	Tyr	Gly	Pro	Tyr	Trp	Lys	Phe	Ile	Lys	Lys	Leu	Cys	Thr	Tyr			
																115	120	125
Glu	Leu	Leu	Gly	Ala	Arg	Asn	Leu	Ala	His	Phe	Gln	Pro	Ile	Arg	Thr			
																130	135	140
Leu	Glu	Val	Lys	Ser	Phe	Leu	Gln	Ile	Leu	Met	Arg	Lys	Gly	Glu	Ser			
																145	150	155
Gly	Glu	Ser	Phe	Asn	Val	Thr	Glu	Glu	Leu	Val	Lys	Leu	Thr	Ser	Asn			
																165	170	175
Val	Ile	Ser	His	Met	Met	Leu	Ser	Ile	Arg	Cys	Ser	Glu	Thr	Glu	Ser			
																180	185	190
Glu	Ala	Glu	Ala	Ala	Arg	Thr	Val	Ile	Arg	Glu	Val	Thr	Gln	Ile	Phe			
																195	200	205
Gly	Glu	Phe	Asp	Val	Ser	Asp	Ile	Ile	Trp	Leu	Cys	Lys	Asn	Phe	Asp			
																210	215	220
Phe	Gln	Gly	Ile	Arg	Lys	Arg	Ser	Glu	Asp	Ile	Gln	Arg	Arg	Tyr	Asp			
																225	230	235
Ala	Leu	Leu	Glu	Lys	Ile	Ile	Thr	Asp	Arg	Glu	Lys	Gln	Arg	Arg	Thr			
																245	250	255
His	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Glu	Ala	Lys	Asp	Phe	Leu	Asp	Met			
																260	265	270
Phe	Leu	Asp	Ile	Met	Glu	Ser	Gly	Lys	Ala	Glu	Val	Lys	Phe	Thr	Arg			
																275	280	285
Glu	His	Leu	Lys	Ala	Leu	Ile	Leu	Asp	Phe	Phe	Thr	Ala	Gly	Thr	Asp			
																290	295	300
Thr	Thr	Ala	Ile	Val	Cys	Glu	Trp	Ala	Ile	Ala	Glu	Val	Ile	Asn	Asn			
																305	310	315
Pro	Asn	Val	Leu	Lys	Lys	Ala	Gln	Glu	Glu	Ile	Ala	Asn	Ile	Val	Gly			
																325	330	335
Phe	Asp	Arg	Ile	Leu	Gln	Glu	Ser	Asp	Ala	Pro	Asn	Leu	Pro	Tyr	Leu			
																340	345	350
Gln	Ala	Leu	Ile	Lys	Glu	Thr	Phe	Arg	Leu	His	Pro	Pro	Ile	Pro	Met			
																355	360	365
Leu	Ala	Arg	Lys	Ser	Ile	Ser	Asp	Cys	Val	Ile	Asp	Gly	Tyr	Met	Ile			
																370	375	380

26 AUG 2003

Pro	Ala	Asn	Thr	Leu	Leu	Phe	Val	Asn	Leu	Trp	Ser	Met	Gly	Arg	Asn	
385																400
					390					395						
Pro	Lys	Ile	Trp	Asp	Tyr	Pro	Thr	Ala	Phe	Gln	Pro	Glu	Arg	Phe	Leu	
					405					410					415	
Glu	Lys	Glu	Lys	Ala	Ala	Ile	Asp	Val	Lys	Gly	Gln	His	Phe	Glu	Leu	
					420					425					430	
Leu	Pro	Phe	Gly	Thr	Gly	Arg	Arg	Gly	Cys	Pro	Gly	Met	Leu	Leu	Ala	
					435					440					445	
Ile	Gln	Glu	Val	Val	Ile	Ile	Ile	Gly	Thr	Met	Ile	Gln	Cys	Phe	Asp	
					450					455					460	
Trp	Lys	Leu	Pro	Asp	Gly	Ser	Gly	His	Val	Asp	Met	Ala	Glu	Arg	Pro	
					465					470					475	480
Gly	Leu	Thr	Ala	Pro	Arg	Glu	Thr	Asp	Leu	Phe	Cys	Arg	Val	Val	Pro	
					485					490					495	
Arg	Val	Asp	Pro	Leu	Val	Val	Ser	Thr	Gln							
					500				505							