
~~WHAT IS~~ CLAIM~~ED I~~S:

1. A compound of formula (I)



(**I**)

or a pharmaceutically acceptable salt thereof, or a stereoisomer thereof,

wherein

R^1 and R^2 are each independently hydrogen, halogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, oxo, -CN, -NO_2, -OR^a, -SO_2R^a, -COR^a, -CO_2R^a, -$CONR^aR^b$, -C (=NR^a)NR^bR^c, -NR^aR^b, -NR^aCOR^b, -$NR^aCONR^bR^c$, -$NR^aCO_2R^b$, -$NR^aSONR^bR^c$, -$NR^aSO_2NR^bR^c$, or –$NR^aSO_2R^b$, each of said -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with halogen, hydroxy, -C_{1-8}alkyoxy, cycloalkyl, heterocyclyl, aryl, or heteroaryl;

R^a, R^b, and R^c are each independently hydrogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl;

n is 0, 1, 2, 3 or 4;

R^3 and R^4, at each of its occurrence, are independently halogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, -C_{1-8}alkyl-heterocyclyl, -C_{1-8}alkyl-cycloalkyl, aryl, heteroaryl, oxo, -CN, -NO_2, -OR^{3a}, -SO_2R^{3a}, -$SO_2NR^{3a}R^{3b}$, -COR^{3a}, -CO_2R^{3a}, -$CONR^{3a}R^{3b}$, -C (=NR^{3a})$NR^{3b}R^{3c}$, -$NR^{3a}R^{3b}$, -$NR^{3a}COR^{3b}$, -$NR^{3a}CONR^{3b}R^{3c}$, -$NR^{3a}CO_2R^{3b}$, -$NR^{3a}SONR^{3b}R^{3c}$, -$NR^{3a}SO_2NR^{3b}R^{3c}$, or –$NR^{3a}SO_2R^{3b}$, each of said -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, -C_{1-8}alkyl-heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents R^{3d}; or R^3 and R^4, when on adjacent carbon atoms of the phenyl ring, together with the two intervening carbon atoms to which they are attached, form a 5- to 8-membered ring comprising 0, 1 or 2 heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{3e};

or two R^4, when on adjacent carbon atoms of the phenyl ring, together with the two intervening carbon atoms to which they are attached, form a 5- to 8-membered ring comprising 0, 1 or 2 heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s);


R^{3a}, R^{3b}, and R^{3c} are each independently hydrogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl, each of said -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents R^{3e}; or

(R^{3a} and R^{3b}), (R^{3b} and R^{3c}), or (R^{3c} and R^{3a}), together with the atom (s) to which they are attached, form a 3- to 12-membered ring, said ring comprising 0, 1 or 2 additional heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{3e};

R^{3d} and R^{3e} are each independently halogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, oxo, -CN, -NO_2, -OR^{3f}, -SO_2R^{3f}, -$SO_2NR^{3f}R^{3g}$, -COR^{3f}, -CO_2R^{3f}, -$CONR^{3f}R^{3g}$, -C (=NR^{3f})$NR^{3g}R^{3h}$, -$NR^{3f}R^{3g}$, -$NR^{3f}COR^{3g}$, -$NR^{3f}CONR^{3g}R^{3h}$, -$NR^{3f}CO_2R^{3f}$, -$NR^{3f}SONR^{3f}R^{3g}$, -$NR^{3f}SO_2NR^{3g}R^{3h}$, or –$NR^{3f}SO_2R^{3g}$, each of said -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents selected from halogen, -C_{1-8}alkyl, -OR^{3i}, -$NR^{3i}R^{3j}$, cycloalkyl, heterocyclyl, aryl, or heteroaryl;

R^{3f}, R^{3g}, R^{3h}, R^{3i}, and R^{3j} are each independently hydrogen, -C_{1-8}alkyl, C_{1-8}alkoxy-C_{1-8}alkyl-, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl;

L^1 is a single bond, alkylene, cycloalkylene, *1-O-alkylene-**1, *1-alkylene-O-**1, *1-NH-alkylene-**1, *1-alkylene-NH-**1, *1-NHC (O)-**1, *1-C (O)NH-**1, alkenylene, or alkynylene; wherein *1 refers to the position attached to Cy1, and **1 refers to the position attached to the pyrrolo[2, 3-b]pyrazine ring;



is a fused heterocyclyl, fused heteroaryl, fused aryl, fused cycloalkyl, fused cycloalkenyl, or fused cycloalkynyl, 

is optionally substituted with R^6;

p is 0, 1, 2, 3, 4; 

is optionally substituted with R^5, m is 0, 1, 2, 3, 4;

R^5 at each of its occurrence, is each independently halogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, -C_{1-8}alkyl-cycloalkyl, heterocyclyl, aryl, heteroaryl, oxo, -CN, -NO_2, -OR^{5a}, -SO_2R^{5a}, -$SO_2NR^{5a}R^{5b}$, -$POR^{5a}R^{5b}$, -COR^{5a}, -CO_2R^{5a}, -$CONR^{5a}R^{5b}$, -C (=NR^{5a})$NR^{5b}R^{5c}$, -$CH_2CONR^{5a}R^{5b}$, -$CH_2CH_2CONR^{5a}R^{5b}$, -$CH_2CH_2CH_2CONR^{5a}R^{5b}$, -$NR^{5a}R^{5b}$, -$CH_2NR^{5a}R^{5b}$, -$CH_2CH_2NR^{5a}R^{5b}$, -$CH_2CH_2CH_2NR^{5a}R^{5b}$, -$NR^{5a}COR^{5b}$, -$NR^{5a}CONR^{5b}R^{5c}$, -$NR^{5a}CO_2R^{5b}$, -$NR^{5a}SONR^{5b}R^{5c}$, -$NR^{5a}SO_2NR^{5b}R^{5c}$, or -$NR^{5a}SO_2R^{5b}$, each of said -$C_{1-8}$alkyl, -$C_{2-8}$alkenyl, -$C_{2-8}$alkynyl, cycloalkyl, -$C_{1-8}$alkyl-cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents R^{5d}; or two R^5, together with the atoms to which they are attached (provided that the valence theory is met), form a 4-, 5-, 6-, 7- or 8-membered ring, said ring comprising 0, 1 or 2 heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{5e};

R^{5a}, R^{5b} and R^{5c} are each independently hydrogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, -C_{1-8}alkyl-C_{1-8}alkoxy, cycloalkyl, -C_{1-8}alkyl-heterocyclyl, -C_{1-8}alkyl-aryl, heterocyclyl, aryl, or heteroaryl, each of said -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, -C_{1-8}alkyl-C_{1-8}alkoxy, cycloalkyl, -C_{1-8}alkyl-heterocyclyl, -C_{1-8}alkyl-aryl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents R^{5e};

(R^{5a} and R^{5b}), (R^{5b} and R^{5c}), or (R^{5c} and R^{5a}), together with the atom (s) to which they are attached, form a 3- to 12-membered ring, said ring comprising 0, 1 or 2 additional heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{5e};

R^{5d} and R^{5e} are each independently hydrogen, halogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, oxo, -CN, -NO_2, -CF_3, -OR^{5f}, -SO_2R^{5f}, -$SO_2NR^{5f}R^{5g}$, -$POR^{5f}R^{5g}$, -COR^{5f}, -CO_2R^{5f}, -$CONR^{5f}R^{5g}$, -C (=NR^{5h})$NR^{5f}R^{5g}$, -$NR^{5f}R^{5g}$, -$NR^{5f}COR^{5g}$, -$NR^{5h}CONR^{5f}R^{5g}$, -$NR^{5f}CO_2R^{5h}$, -$NR^{5h}SONR^{5f}R^{5g}$, -$NR^{5h}SO_2NR^{5f}R^{5g}$, or –$NR^{5f}SO_2R^{5g}$, each of said -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents selected from halogen, -C_{1-8}alkyl, -OR^{5i}, -$NR^{5i}R^{5j}$, cycloalkyl, heterocyclyl, aryl, or heteroaryl;

R^{5f}, R^{5g}, R^{5h}, R^{5i}, and R^{5j} are each independently hydrogen, oxo, -C_{1-8}alkyl, -C_{1-8}alkoxy, hydroxy, C_{1-8}alkoxy-C_{1-8}alkyl-, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl;

R^6 at each of its occurrence, is each independently hydrogen, halogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, oxo, -CN, -NO_2, -OR^{6a}, -SO_2R^{6a}, -$SO_2NR^{6a}R^{6b}$, -COR^{6a}, -CO_2R^{6a}, -$CONR^{6a}R^{6b}$, -C (=NR^{6a})$NR^{6b}R^{6c}$, -$CH_2CONR^{6a}R^{6b}$, -CH_2 $CH_2CONR^{6a}R^{6b}$, -$CH_2CH_2CH_2CONR^{6a}R^{6b}$, -$NR^{6a}R^{6b}$, -$CH_2NR^{6a}R^{6b}$, -$CH_2$ $CH_2NR^{6a}R^{6b}$, -$CH_2CH_2CH_2NR^{6a}R^{6b}$, -$NR^{6a}COR^{6b}$, -$NR^{6a}CONR^{6b}R^{6c}$, -$NR^{6a}CO_2R^{6b}$, -$NR^{6a}SONR^{6b}R^{6c}$, -$NR^{6a}SO_2NR^{6b}R^{6c}$, or –$NR^{6a}SO_2R^{6b}$, each of said -$C_{1-8}$alkyl, -$C_{2-8}$alkenyl, -$C_{2-8}$alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents R^{6d}; or two R^6, together with the atom to which they are attached (provided that the valence theory is met), form a 4-, 5-, 6-, 7- or 8-membered ring, said ring comprising 0, 1 or 2 heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{6e};


R^{6a}, R^{6b}, and R^{6c} are each independently hydrogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, C_{1-8}alkoxy-C_{1-8}alkyl-, cycloalkyl, heterocyclyl, aryl, or heteroaryl, each of said -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents R^{6e};

(R^{6a} and R^{6b}), (R^{6b} and R^{6c}), or (R^{6c} and R^{6a}), together with the atom (s) to which they are attached, form a 3- to 12-membered ring, said ring comprising 0, 1 or 2 additional heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{6e};

R^{6d} and R^{6e} are each independently hydrogen, halogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, oxo, -CN, -NO_2, -OR^{6f}, -SO_2R^{6f}, -$SO_2NR^{6f}R^{6g}$, -COR^{6f}, -CO_2R^{6f}, -$CONR^{6f}R^{6g}$, -C (=NR^{6h})$NR^{6f}R^{6g}$, -$NR^{6f}R^{6g}$, -$NR^{6f}COR^{6g}$, -$NR^{6h}CONR^{6f}R^{6g}$, -$NR^{6f}CO_2R^{6h}$, -$NR^{6h}SONR^{6f}R^{6g}$, -$NR^{6h}SO_2NR^{6f}R^{6g}$, or –$NR^{6f}SO_2R^{6g}$, each of said -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents selected from halogen, -C_{1-8}alkyl, -OR^{6i}, -$NR^{6i}R^{6j}$, cycloalkyl, heterocyclyl, aryl, or heteroaryl;

R^{6f}, R^{6g}, R^{6h}, R^{6i}, and R^{6j} are each independently hydrogen, oxo, -C_{1-8}alkyl, C_{1-8}alkoxy-C_{1-8}alkyl-, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl.


2. The compound of claim 1, wherein R^1 and R^2 are each hydrogen or -C_{1-8}alkyl ~~(preferably hydrogen or methyl)~~; and/or L^1 is a single bond.


3. The compound of claim 1 or 2, wherein R^3 is –C_{1-8}alkyl, -C_{1-8}alkyl-heterocyclyl or -C_{1-8}alkyl-cycloalkyl, wherein said –C_{1-8}alkyl, -C_{1-8}alkyl-heterocyclyl or -C_{1-8}alkyl-cycloalkyl is optionally substituted with at least one substituents R^{3d}; R^{3d} is independently selected from -C_{1-8}alkyl or -OR^{3f}; R^{3f} is each independently hydrogen, -C_{1-8}alkyl; preferably R^3 is 

; or

R^3 is -$CONR^{3a}R^{3b}$; R^{3a} and R^{3b} are each independently hydrogen or -C_{1-8}alkyl; each of said -C_{1-8}alkyl is substituted with at least one substituents selected from hydrogen, -OR^{3f}, CN, 3- to-7 membered heterocyclyl comprising 1 or 2 heteroatoms selected from nitrogen, oxygen; R^{3f} is selected from hydrogen or -C_{1-8}alkyl; or

R^3 is -$CONR^{3a}R^{3b}$ or -$NR^{3a}R^{3b}$; R^{3a} and R^{3b} together with the nitrogen atom to which they are attached is 4- to 12-membered ring comprising 1 or 2 additional nitrogen or oxygen heteroatoms as ring, said ring is optionally substituted with at least one substituents R^{3e}; R^{3e} each is selected from oxo, -C_{1-8}alkyl, -OR^{3f}, -$NR^{3f}R^{3g}$, said-C_{1-8}alkyl is optionally substituted by at least one halogen, wherein R^{3f} and R^{3g} are each independently hydrogen, or -C_{1-8}alkyl; or


R^3 is selected from 














or 







.





~~4. The compound of claim 1 or 2, wherein R^3 is -$CONR^{3a}R^{3b}$; R^{3a} and R^{3b} are each independently hydrogen or -C_{1-8}alkyl (preferably methyl, ethyl, propyl, butyl, pentyl or hexyl); each of said -C_{1-8}alkyl is substituted with at least one substituents selected from hydrogen, -OR^{3f}, CN, 3- to-7 membered heterocyclyl comprising 1 or 2 heteroatoms selected from nitrogen, oxygen (for example, piperazinyl, piperidinyl, piperidinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydropyranyl, or pyrrolidinyl); R^{3f} is selected from hydrogen or -C_{1-8}alkyl (e.g., methyl, ethyl, propyl, butyl, pentyl or hexyl).~~



~~5. The compound of claim 1 or 2, wherein R^3 is -$CONR^{3a}R^{3b}$ or -$NR^{3a}R^{3b}$; R^{3a} and R^{3b} together with the nitrogen atom to which they are attached is 4- to 12-membered ring comprising 1 or 2 additional nitrogen or oxygen heteroatoms as ring member (e.g., monocyclic 3- to 8-membered ring or bicyclic spiro 7- to 12-membered ring), said ring is optionally substituted with at least one substituents R^{3e}; R^{3e} each is selected from oxo, -C_{1-8}alkyl, -OR^{3f}, -$NR^{3f}R^{3g}$, said-C_{1-8}alkyl is optionally substituted by at least one halogen, wherein R^{3f} and R^{3g} are each independently hydrogen, or -C_{1-8}alkyl.~~

64. The compound of any one of claims 1-53, wherein R^3 is selected from 

( or ),, , , , , ( or ), , , , ( or ),

, , , , ( or ),

 ( or ) or  .











~~7. The compound of any claims 1-5, wherein R^3 is selected from~~  ~~or~~  ~~.~~

85. The compound of any one of claims 1-74, wherein n is 0, 1 or 2; R^4 is selected from halogen, -C_{1-8}alkyl (preferably methyl), halogen, CN, -OR^{3a} or -$NR^{3a}CONR^{3b}R^{3c}$; said -C_{1-8}alkyl is optionally substituted with at least one substituents R^{3d}; R^{3a}, R^{3b} and R^{3c} are each independently hydrogen, -C_{1-8}alkyl (preferably methyl); R^{3d} is each independently halogen or -C_{1-8}alkyl.

96. The compound of claim 85, wherein n is 1, R^4 is selected from -C_{1-8}alkyl ~~(preferably methyl)~~, halogen, CN, OR^{3a} or -$NR^{3a}CONR^{3b}R^{3c}$; said -C_{1-8}alkyl is optionally substituted with at least one substituents R^{3d}; R^{3a}, R^{3b} and R^{3c} are each independently hydrogen, or -C_{1-8}alkyl (preferably methyl); R^{3d} is each independently halogen or -C_{1-8}alkyl; or


n is 2, and R^4 is halogen; or


n is 1 or 2, and R^4 is selected from methyl, F, OH, CN, -CHF_2 or -$NHCOCH_3$;or

n is 0, and R^3 is selected from , , , , , , , ,  or ;or ~~.~~

~~10. The compound of claim 8, wherein n is 2, and R^4 is halogen.~~


~~11. The compound of claim 8, wherein n is 1 or 2, and R^4 is selected from methyl, F, OH, CN, -CHF_2 or -$NHCOCH_3$.~~

~~12. The compound of any claims 1-8, wherein n is 0, and R^3 is selected from~~  ~~,~~




~~or~~


~~13~~7. The compound of any one of claims 1-~~6~~9, ~~11~~, wherein n is 1, and R^3 is selected from

 or ;

and R^4 is methyl, F, OH, CN, $-CHF_2$ or $-NHCOCH_3$; or



n is 2, and R^3 is selected from , and R^4 is F.






~~14. The compound of any claims 1-8, 10-11, wherein n is 2, and R^3 is selected from~~





~~, and R^4 is F.~~




~~15~~8. The compound of any one of claims 1-2, wherein R^3 and R^4, when on adjacent carbon atoms of the phenyl ring, together with the two intervening carbon atoms to which they are attached, form a 5- to 8-membered ring comprising 0, 1 or 2 heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{3e}; each R^{3e} is independently selected from $-C_{1-8}$alkyl or oxo.

~~16~~9. The compound of claim ~~15~~8, wherein  is , X=NH or O, and R^{3e} is $-C_{1-8}$alkyl; or


wherein  is ; and n=2, two R^4, when on adjacent carbon atoms of the phenyl ring, together with the two intervening carbon atoms to which they are attached, form a 5-membered ring comprising 2 heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s); wherein  is  or .

~~17. The compound of any claims 1-2, wherein n=2, two R^4 when on adjacent carbon atoms of the phenyl ring, together with the two intervening carbon atoms to which they are attached, form a 5- to 8-membered ring comprising 0, 1 or 2 heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s).~~


~~18. The compound of claim 17, wherein~~  ~~is~~  ~~; and two R^4, when on adjacent carbon atoms of the phenyl ring, together with the two intervening carbon atoms to which they are attached, form a 5-membered ring comprising 2 heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s).~~

~~19. The compound of claim 18, wherein~~  ~~is~~  ~~or~~  ~~.~~

~~20. The compound of any claims 1-19, L^1 is a single bond.~~

~~21~~10. The compound of any one of claims 1-~~20~~9, wherein  is benzo fused heterocyclyl, benzo fused heteroaryl, benzo fused cycloalkyl, benzo fused cycloalkenyl, benzo fused cycloalkynyl;

wherein the benzo fused heterocyclyl is indolinyl, isoindolinyl, benzopyranyl, dihydrothiazolopyrimidinyl, tetrahydroquinolyl, tetrahydroisoquinolyl, dihydrobenzofuranyl, dihydrobenzoxazinyl, dihydrobenzoimidazolyl, tetrahydrobenzothienyl, tetrahydrobenzofuranyl, benzodioxolyl, benzodioxonyl, chromanyl, chromenyl, octahydrochromenyl, dihydrobenzodioxynyl, dihydrobenzoxezinyl, dihydrobenzodioxepinyl, dihydrothienodioxynyl, dihydrobenzooxazepinyl, tetrahydrobenzooxazepinyl, dihydrobenzoazepinyl, tetrahydrobenzoazepinyl, isochromanyl, or chromanyl;and/or ~~wherein~~ the benzo fused heteroaryl is benzisoxazolyl, benzodiazolyl, benzofuranyl, benzofurazanyl, benzofuryl, benzoimidazolyl, benzoisothiazolyl, benzothiadiazolyl, benzothiazolyl, benzothienyl, benzothiophenyl, benzotriazolyl, benzoxadiazolyl, benzoxazolyl, indazolyl, indolyl, isobenzofuryl, isoindolyl, isoquinolinyl (or isoquinolyl), phthalazinyl, pteridinyl, purinyl, quinazolinyl, quinolinyl (or quinolyl), or quinoxalinyl; in some examples, indazolyl is e.g., 1H-indazol-4-yl, 2H-indazol-4-yl, 1H-indazol-5-yl, 2H-indazol-5-yl, 1H-indazol-7-yl; benzodiazolyl is e.g., 1H-benzo[d]imidazol-4-yl, 1H-1, 3-benzodiazol-5-yl or 1H-benzo[d]imidazol-5-yl, 1H-benzo[d]imidazol-4-yl, 1H-benzo[d]imidazol-6-yl; benzooxazolyl is e.g., benzo[d]oxazol-6-yl; benzooxadiazolyl is e.g., benzo[c][1, 2, 5]oxadiazol-4-yl; or

wherein  is a bicyclic fused heteroaryl selected from furopyridinyl, furopyrrolyl, imidazopyridinyl, imidazopyridyl, imidazothiazolyl, indolizinyl, naphthyridinyl, pyrazinopyridazinyl, pyrazolopyridinyl, pyrazolopyrimidinyl, pyrazolopyridyl, pyrazolotriazinyl,


pyridazolopyridyl, pyrrolopyridinyl, thiazolopyridyl, thienopyrazinyl, thienopyrazolyl, thienopyridyl, thienopyrrolyl, thienothienyl, or triazolopyridyl; in some example, imidazopyridinyl is e.g., imidazo[1, 2-a]pyridin-5-yl or imidazo[1, 5-a]pyridin-5-yl.

22. The compound of claim 21, wherein the benzo fused heterocyclyl is indolinyl, isoindolinyl, benzopyranyl, dihydrothiazolopyrimidinyl, tetrahydroquinolyl, tetrahydroisoquinolyl, dihydrobenzofuranyl, dihydrobenzoxazinyl, dihydrobenzoimidazolyl, tetrahydrobenzothienyl, tetrahydrobenzofuranyl, benzodioxolyl, benzodioxonyl, chromanyl, chromenyl, octahydrochromenyl, dihydrobenzodioxynyl, dihydrobenzoxezinyl, dihydrobenzodioxepinyl, dihydrothienodioxynyl, dihydrobenzooxazepinyl, tetrahydrobenzooxazepinyl, dihydrobenzoazepinyl, tetrahydrobenzoazepinyl, isochromanyl, or chromanyl.




23. The compound of claim 21 or22, wherein the benzo fused heterocyclyl is indolinyl, isoindolinyl, tetrahydroisoquinolyl, dihydrobenzofuranyl, dihydrobenzoimidazolyl, tetrahydrobenzooxazepinyl, tetrahydrobenzoazepinyl, or isochromanyl.




2411. The compound of claim 2310, wherein tetrahydroisoquinolinyl is selected from 1, 2, 3, 4-tetrahydroisoquinolinyl, 1, 2, 3, 4-tetrahydroisoquinolin-6-yl or 1, 2, 3, 4-tetrahydroisoquinolin-7-yl; tetrahydrobenzooxazepinyl is selected from 2, 3, 4, 5-tetrahydrobenzooxazepinyl, 2, 3, 4, 5-tetrahydrobenzo[f][1, 4]oxazepin-8-yl, or 2, 3, 4, 5-tetrahydrobenzo[f][1, 4]oxazepin-7-yl; tetrahydrobenzoazepinyl is selected from 2, 3, 4, 5-tetrahydrobenzoazepinyl, 2, 3, 4, 5-tetrahydro-1H-benzo[c]azepin-7-yl and 2, 3, 4, 5-tetrahydro-1H-benzo[d]azepin-7-yl; isoindolinyl is selected from isoindolin-5-yl; isochromanyl is selected from isochroman-5-yl; dihydroisobenzofuranyl is selected from 1, 3-dihydroisobenzofuran-4-yl.

25. The compound of claim 21, wherein the benzo fused heteroaryl is benzisoxazolyl, benzodiazolyl, benzofuranyl, benzofurazanyl, benzofuryl, benzoimidazolyl, benzoisothiazolyl, benzothiadiazolyl, benzothiazolyl, benzothienyl, benzothiophenyl, benzotriazolyl, benzoxadiazolyl, benzoxazolyl, indazolyl, indolyl, isobenzofuryl, isoindolyl, isoquinolinyl (or isoquinolyl), phthalazinyl, pteridinyl, purinyl, quinazolinyl, quinolinyl (or quinolyl), or quinoxalinyl; in some examples, indazolyl is e.g., 1H-indazol-4-yl, 2H-indazol-4-yl, 1H-indazol-5-yl, 2H-indazol-5-yl, 1H-indazol-7-yl; benzodiazolyl is e.g., 1H-benzo[d]imidazol-4-yl, 1H-1, 3-benzodiazol-5-yl or 1H-benzo[d]imidazol-5-yl, 1H-benzo[d]imidazol-4-yl, 1H-benzo[d]imidazol-6-yl; benzooxazolyl is e.g., benzo[d]oxazol-6-yl; benzooxadiazolyl is e.g., benzo[e][1, 2, 5]oxadiazol-4-yl.




26. The compound of claim 21, wherein 

is a bicyclic fused heteroaryl selected from furopyridinyl, furopyrrolyl, imidazopyridinyl, imidazopyridyl, imidazothiazolyl, indolizinyl, naphthyridinyl, pyrazinopyridazinyl, pyrazolopyridinyl, pyrazolopyrimidinyl, pyrazolopyridyl, pyrazolotriazinyl, pyridazolopyridyl, pyrrolopyridinyl, thiazolopyridyl, thienopyrazinyl, thienopyrazolyl, thienopyridyl, thienopyrrolyl, thienothienyl, or triazolopyridyl; in some example, imidazopyridinyl is e.g., imidazo[1, 2-a]pyridin-5-yl or imidazo[1, 5-a]pyridin-5-yl.

2712. The compound of claim 2110, wherein  is  or , p=0, 1, 2 or 3; m=0, 1, 2, 3 or 4; R^6 and R^5 are defined for Formula (I).

2813. The compound of any one of claims 2110-2712, wherein  is  or  (preferably), m=1, 2, 3 or 4;

each R^5 is independently selected from -C_{1-8}alkyl, -C_{1-8}alkyl-cycloalkyl, heterocyclyl or heteroaryl -$CONR^{3a}R^{3b}$, $CH_2CONR^{5a}R^{5b}$, -CH_2 $CH_2CONR^{5a}R^{5b}$, -$NR^{5a}R^{5b}$, -$CH_2NR^{5a}R^{5b}$, -CH_2 $CH_2NR^{5a}R^{5b}$, said -C_{1-8}alkyl, -C_{1-8}alkyl-cycloalkyl or heterocyclyl or heteroaryl is optionally substituted with at least one R^{5d}, and R^{5a}, R^{5b} and R^{5d} are defined for Formula (I); wherein R^6 is selected from -C_{1-8}alkyl, -OR^{6a} or -$NR^{6a}R^{6b}$, said -C_{1-8}alkyl is optionally substituted with at least one substituents R^{6d}, R^{6a} and R^{6b} are each hydrogen or -C_{1-8}alkyl; R^{6d} is each independently hydrogen, halogen or -C_{1-8}alkyl. ~~R^6 and R^5 are defined for Formula (I) (preferably R^5 and R^6 are independently -C_{1-8}alkyl (e.g., methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-methyl-1-propyl, 1-methylpropyl or t-butyl).~~

14. The compound of any one of claims 10-12, wherein  is selected from  or , R^5 is selected from methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-methyl-1-propyl, 1-methylpropyl or t-butyl; .R^6 is selected from -CH_3, -OCH_3, -$NHCH_3$, -CHF_2, or -C $(CH_3)_2OH$.

~~29. The compound of any claims 27-28, wherein R^6 is selected from -C_{1-8}alkyl, -OR^{6a} or -$NR^{6a}R^{6b}$, said -C_{1-8}alkyl is optionally substituted with at least one substituents R^{6d}, R^{6a} and R^{6b} are each hydrogen or -C_{1-8}alkyl; R^{6d} is each independently hydrogen, halogen or -C_{1-8}alkyl; preferably R^6 is selected from -CH_3, -OCH_3, -$NHCH_3$, -CHF_2, or -C $(CH_3)_2OH$.~~


~~30. The compound of any claims 27-29, wherein each R^5 is independently selected from -C_{1-8}alkyl, -C_{1-8}alkyl-cycloalkyl, heterocyclyl or heteroaryl -$CONR^{3a}R^{3b}$, $CH_2CONR^{5a}R^{5b}$, -CH_2 $CH_2CONR^{5a}R^{5b}$, -$NR^{5a}R^{5b}$, -$CH_2NR^{5a}R^{5b}$, -CH_2-$CH_2NR^{5a}R^{5b}$, said -C_{1-8}alkyl, -C_{1-8}alkyl-~~


~~cycloalkyl or heterocyclyl or heteroaryl is optionally substituted with at least one R^{5d}, and R^{5a}, R^{5b} and R^{5d} are defined for Formula (I).~~

~~31~~15. The compound of any one of claims ~~27~~12-~~30~~13, wherein R^5 is selected from ~~C_{1-8}alkyl (e.g., methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-methyl-1-propyl, 1-methylpropyl or t-butyl)~~, -C_{1-8}alkyl-cycloalkyl, heterocyclyl or heteroaryl, said -C_{1-8}alkyl, -C_{1-8}alkyl-cycloalkyl, heteroaryl or heterocyclyl is optionally substituted with at least one substituents R^{5d}; R^{5d} is selected from -C_{1-8}alkyl, -OR^{5f}, -$NR^{5f}R^{5g}$, -CO_2R^{5f}, heteroaryl, heterocyclyl, -SO_2R^{5f}, -$POR^{5f}R^{5g}$, -$CONR^{5f}R^{5g}$, oxo or -CF_3; R^{5f} and R^{5g} are each independently hydrogen, -C_{1-8}alkyl, -C_{1-8}alkoxy or hydroxy; or





R^5 is -$NR^{5a}R^{5b}$, -$CH_2NR^{5a}R^{5b}$, -CH_2 $CH_2NR^{5a}R^{5b}$; R^{5a} and R^{5b} together with the nitrogen atom to which they are attached form a 4- to 7- membered ring, said ring comprising 0, 1 or 2 additional heteroatoms independently selected from nitrogen, oxygen, said ring is optionally substituted with at least one substituents R^{5e}; R^{5e} is each independently selected from C_{1-8}alkyl, OR^{5f}; R^{5f} is each independently selected from hydrogen or -C_{1-8}alkyl; or

R^5 is selected from -$CONR^{5a}R^{5b}$, -$CH_2CONR^{5a}R^{5b}$, -$CH_2CH_2CONR^{5a}R^{5b}$, wherein R^{5a} is hydrogen, methyl, ethyl; R^{5b} is selected from -C_{1-8}alkyl, cycloalkyl, -C_{1-8}alkyl-heterocyclyl, -C_{1-8}alkyl-aryl, heterocyclyl, aryl, said -C_{1-8}alkyl, cycloalkyl, -C_{1-8}alkyl-heterocyclyl, -C_{1-8}alkyl-aryl, heterocyclyl, aryl is optionally substituted with at least one substituents R^{5e}, R^{5e} is selected from halogen, -CH_2OH or OR^{5f}, R^{5f} is selected from hydrogen, cycloalkyl or -C_{1-8}alkyl.

~~32~~16. The compound of claim ~~31~~15, wherein, R^5 is methyl, ethyl, propyl, butyl, pentyl or hexyl, -CH_2OH, -C_2H_4OH, $C_2H_4NHCH_3$, -$C_2H_4OCH_3$, 




33. The compound of claim 30, wherein R^5 is $-NR^{5a}R^{5b}$, $-CH_2NR^{5a}R^{5b}$, $-CH_2CH_2NR^{5a}R^{5b}$; R^{5a} and R^{5b} together with the nitrogen atom to which they are attached form a 4- to 7- membered ring, said ring comprising 0, 1 or 2 additional heteroatoms independently selected from nitrogen, oxygen, said ring is optionally substituted with at least one substituents R^{5e}; R^{5e} is each independently selected from C_{1-8}alkyl (preferably methyl, ethyl), OR^{5f}; R^{5f} is each independently selected from hydrogen or C_{1-8}alkyl (e.g., methyl, ethyl).


34. The compound of claim 33, wherein R^5 is    or .

35. The compound of claim 30, wherein R^5 is selected from $-CONR^{5a}R^{5b}$, $-CH_2CONR^{5a}R^{5b}$, $-CH_2CH_2CONR^{5a}R^{5b}$, wherein R^{5a} and R^{5b} are each independently hydrogen, $-C_{1-8}$alkyl, C_{1-8}alkoxy-C_{1-8}alkyl-, cycloalkyl, heterocyclyl, $-C_{1-8}$alkyl-heterocyclyl, $-C_{1-8}$alkyl-aryl, -aryl, or heteroaryl, each of said $-C_{1-8}$alkyl, C_{1-8}alkoxy-C_{1-8}alkyl-, cycloalkyl, heterocyclyl, $-C_{1-8}$alkyl-heterocyclyl, $-C_{1-8}$alkyl-aryl, -aryl, or heteroaryl is optionally substituted with at least one substituents R^{5e}; R^{5e} is independently hydrogen, OR^{5f}, $-C_{1-8}$alkyl, heterocyclyl, $-C_{1-8}$alkyl-aryl, or C_{1-8}alkoxy-C_{1-8}alkyl-; R^{5f} is each independently selected from hydrogen, $-C_{1-8}$alkyl (e.g., methyl, ethyl) and cycloalkyl.


36. The compound of claim 35, wherein R^5 is selected from $-CONR^{5a}R^{5b}$, $-CH_2CONR^{5a}R^{5b}$, $-CH_2CH_2CONR^{5a}R^{5b}$, wherein R^{5a} is hydrogen or $-C_{1-8}$alkyl (e.g., methyl, ethyl); R^{5b} is selected from $-C_{1-8}$alkyl, cycloalkyl, $-C_{1-8}$alkyl-heterocyclyl, $-C_{1-8}$alkyl-aryl, heterocyclyl, aryl, said $-C_{1-8}$alkyl, cycloalkyl, $-C_{1-8}$alkyl-heterocyclyl, $-C_{1-8}$alkyl-aryl, heterocyclyl, aryl is optionally substituted with at least one substituents R^{5e}, R^{5e} is selected from halogen, $-CH_2OH$ or OR^{5f}, R^{5f} is selected from hydrogen, cycloalkyl or $-C_{1-8}$alkyl.

37. The compound of claim 36, wherein R^5 is selected from






~~38~~17. The compound of claim 15~~35 or 36~~, wherein R^5 is -$CONR^{3a}R^{3b}$, $CH_2CONR^{5a}R^{5b}$ or -CH_2 $CH_2CONR^{5a}R^{5b}$, R^{5a} is hydrogen or methyl; and R^{5b} is selected from ~~cycloalkyl, such as~~ cycloalkyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexenyl, cyclohexadienyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, cycloundecyl, and cyclododecyl groups~~; said cycloalkyl is optionally substituted with OH, CH_3, OCH_3; preferably R^5 is~~




~~39. The compound of claim 35 or 36, wherein R^5 is $CONR^{5a}R^{5b}$, $CH_2CONR^{5a}R^{5b}$ or CH_2 $CH_2CONR^{5a}R^{5b}$, R^{5a} is hydrogen or methyl; and R^{5b} is selected from C_{1-8}alkyl, cycloalkyl or C_{1-8}alkyl-heterocyclyl, said C_{1-8}alkyl, cycloalkyl or C_{1-8}alkyl-heterocyclyl is optionally substituted with at least one substituents R^{5e}, R^{5e} is selected from cycloalkyl, C_{1-8}alkyl, CH_2OH or OR^{5f}, OR^{5f} is selected from hydrogen or C_{1-8}alkyl. For example, R^5 is selected from~~



~~or~~

~~40~~18. The compound of claim ~~35 or 36~~17, wherein R^5 is -$CONR^{5a}R^{5b}$, $CH_2CONR^{5a}R^{5b}$ or -CH_2 $CH_2CONR^{5a}R^{5b}$, R^{5a} is hydrogen or methyl; and R^{5b} is heterocyclyl selected from~~(such as~~








~~)~~ or ~~aryl (such as~~ phenyl~~)~~, said heterocyclyl or phenyl~~aryl~~ is optionally substituted with at least one substituents R^{5e} selected from halogen, -C_{1-8}alkyl or -OR^{5f}; R^{5f} is each independently hydrogen or -C_{1-8}alkyl; or

R^5 is selected from -$CONR^{5a}R^{5b}$, $CH_2CONR^{5a}R^{5b}$, -$CH_2CH_2CONR^{5a}R^{5b}$, wherein R^{5a} and R^{5b} together with the nitrogen atom to which they are attached, form a 4- to 7-membered ring,




said ring comprising 0, 1 or 2 additional heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{5e}; R^{5e} is each independently halogen, -C_{1-8}alkyl, -OR^{5f}, R^{5f} is each independently hydrogen or -C_{1-8}alkyl.




~~41. The compound of claim 30, wherein R^5 is selected from -$CONR^{5a}R^{5b}$, $CH_2CONR^{5a}R^{5b}$, -$CH_2CH_2CONR^{5a}R^{5b}$, wherein R^{5a} and R^{5b} together with the nitrogen atom to which they are attached, form a 4- to 7-membered ring, said ring comprising 0, 1 or 2 additional heteroatoms independently selected from nitrogen, oxygen or optionally oxidized sulfur as ring member (s), said ring is optionally substituted with at least one substituents R^{5e}; R^{5e} is each independently halogen, -C_{1-8}alkyl, -OR^{5f}, R^{5f} is each independently hydrogen or -C_{1-8}alkyl.~~

~~42~~19. The compound of any one of claims 1-~~41~~18, wherein R^5 is selected from methyl, ethyl, propyl, butyl, pentyl or hexyl, -CH_2OH, -C_2H_4OH, $C_2H_4NHCH_3$, -$C_2H_4OCH_3$, , .







~~43~~20. The compound of claim ~~28~~13, wherein  is  or , $(R^5)_{m-1}$, m=2, 3 or 4, R^5 is each independently halogen, -C_{1-8}alkyl, -C_{2-8}alkenyl, -C_{2-8}alkynyl, cycloalkyl, -C_{1-8}alkyl-cycloalkyl, heterocyclyl, aryl, heteroaryl, oxo, -CN, -NO_2, -OR^{5a},






















$-SO_2R^{5a}$, $-SO_2NR^{5a}R^{5b}$, $-POR^{5a}R^{5b}$, $-COR^{5a}$, $-CO_2R^{5a}$, $-CONR^{5a}R^{5b}$, $-C(=NR^{5a})NR^{5b}R^{5c}$, $-CH_2CONR^{5a}R^{5b}$, $-CH_2CH_2CONR^{5a}R^{5b}$, $-CH_2CH_2CH_2CONR^{5a}R^{5b}$, $-NR^{5a}R^{5b}$, $-CH_2NR^{5a}R^{5b}$, $-CH_2CH_2NR^{5a}R^{5b}$, $-CH_2CH_2CH_2NR^{5a}R^{5b}$, $-NR^{5a}COR^{5b}$, $-NR^{5a}CONR^{5b}R^{5c}$, $-NR^{5a}CO_2R^{5b}$, $-NR^{5a}SONR^{5b}R^{5c}$, $-NR^{5a}SO_2NR^{5b}R^{5c}$, or $-NR^{5a}SO_2R^{5b}$, each of said $-C_{1-8}$alkyl, $-C_{2-8}$alkenyl, $-C_{2-8}$alkynyl, cycloalkyl, $-C_{1-8}$alkyl-cycloalkyl, heterocyclyl, aryl, or heteroaryl is optionally substituted with at least one substituents R^{5d}; R^6, R^{5a}, R^{5b}, R^{5c} and R^{5d} are defined for Formula (I).

























~~44~~21. The compound of claim ~~43~~20, wherein  is  or , m=2, R^5 is each independently $-C_{1-8}$alkyl or $-CONR^{5a}R^{5b}$, R^{5b} and R^{5d} are hydrogen or $-C_{1-8}$alkyl ~~(e.g., methyl, ethyl, propyl, butyl, pentyl or hexyl)~~; R^6 is selected from $-C_{1-8}$alkyl, $-OR^{6a}$ or $-NR^{6a}R^{6b}$, said $-C_{1-8}$alkyl is optionally substituted with at least one substituents R^{6d}, R^{6a} and R^{6b} are each hydrogen or $-C_{1-8}$alkyl; R^{6d} is each independently hydrogen, halogen or $-C_{1-8}$alkyl~~; preferably R^6 is selected from $-CH_3$, $-OCH_3$, $-NHCH_3$, $-CHF_2$, or $-C(CH_3)_2OH$~~.






















~~45~~22. The compound of claim ~~44~~21, wherein  is  or ; R^6 is selected from $-CH_3$, $-OCH_3$, $-NHCH_3$, $-CHF_2$, or $-C(CH_3)_2OH$.






















~~46~~23. The compound of claim 1, selected from:






















		
1	2	3
		
4	5	6
















		
7A	7B	8
		
9	10	11
		
12	13	14
		
15	16	17
		
18	19	20
		
21	22	23
		
24	25	26

		
27	28	29
		
30	31	32
		
33	34	35
		
36	37	38
		
39	40	41
		
42	43	44
		
45	46	47
		
48	49	50

		
51	52	53
		
54	55	56
		
57	58	59
		
60	61	62
		
63	64	65
		
66	67	68
		
69	70	71

		
72	73	74
		
75	76	77
		
78	79	80
		
81	82	83
		
84	85	86
		
87	88	89
		
90	91	92

		
93	94	95
		
96	97	98
		
99	100	101
		
102	103	104
		
105	106	107
		
108	109	110
		
111	112	113

		
114	115	116
		
117	118	119
		
120	121	122
		
123	124	125
		
126	127	128

4724. A pharmaceutical composition comprising the compound of any one of claims 1-4623 or a stereoisomer thereof, or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable excipient.

4825. A use of the compound of any one of claims 1-23 or a stereoisomer thereof, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament in ~~A method of~~ treating cancer~~, comprising administering a subject in need thereof the compound of any of claims 1-46 or a stereoisomer thereof, or a pharmaceutically acceptable salt thereof~~.