

[Name of Document] Description

[Title of Invention] Novel Nicotinamide Derivative or Salt Thereof

[Technical Field]

[0001]

5 The present invention relates to a nicotinamide derivative having Syk-inhibitory activity or a salt thereof.

[Background Art]

[0002]

10 Spleen Tyrosine Kinase (Syk), which is a non-receptor type intracellular tyrosine phosphatase, plays essential roles for activation of B cells and in an intracellular signaling system mediated by an Fc receptor. For example, Syk is associated with a FcεRI signal that is an immunoglobulin E receptor in mast cells, basophils, and other cells, and thus it regulates generation of inflammatory mediators, such as histamine or leukotrien, as
15 well as cytokine from these cells. At the same time, Syk plays a role in transmitting activation signals caused by stimulation of Fcγ receptor into monocytes, dendritic cells, and other cells (Non Patent Literatures 1 and 2). Moreover, it has been reported that Syk is also associated with cytokine signaling caused by integrin, IL-13, IL-15, etc. (Non Patent Literatures 3 and
20 4).

[0003]

 In the case of a B-cell, a signal is transmitted into the cell mediated by a BCR (a B-cell antigen receptor) expressed on the cell membrane, so that activation and differentiation of the cell is induced, resulting in generation of
25 an antibody. It has been reported that Syk is essential for such an activation and differentiation process (Non Patent Literature 5).

[0004]

 It is anticipated that it is possible to suppress various cell responses by inhibiting Syk (Non Patent Literatures 5 and 6).

30 [0005]

 In the case of a type I allergy, which is an immediate-type allergy reaction, for example, immunoglobulin E (IgE) binds to FcεRI, which is a high-affinity IgE receptor, and an allergen then binds thereto to promote activation of the FcεRI and the release of inflammatory mediator. As a result,
35 allergic symptoms are expressed. It is anticipated that inhibition of Syk activity will lead to the suppression of the activation of the FcεRI, and that it

will be useful for the treatment of representative type I allergy-related diseases such as bronchial asthma, allergic rhinitis, hives, and atopic dermatitis.

[0006]

5 Moreover, it is considered that inhibition of Syk activity leads to the suppression of the activation and/or maturation of immune B cells and generation of antibodies, and that such inhibition of Syk activity can also regulate immune reactions other than type I allergy. Accordingly, it is also anticipated that inhibition of Syk activity will be effective for autoimmune
10 diseases (rheumatoid arthritis, systemic lupus erythematosus, etc.), autoimmune hemolytic anemia, nephrotic syndrome, contact dermatitis, and the like. Furthermore, since inhibition of Syk activity also leads to the suppression of the activation of macrophages, it is anticipated that inhibition of Syk will be also effective for idiopathic thrombocytopenic purpura.

15 [0007]

 Further, inhibition of Syk activity suppresses not only immune and/or inflammatory diseases, but also activation and proliferation of lymphocytes, including B-cells as typical examples. Thus, it is anticipated that inhibition of Syk will be also effective for the treatment of various types of proliferative
20 diseases such as lymphoma and lymphocytic leukemia. Still further, since inhibition of Syk activity regulates proliferation and differentiation of bone marrow cells, it is anticipated that it will be also effective for acute myelocytic leukemia.

[0008]

25 On the other hand, Syk has been known to be involved in signaling mediated by integrin, which is a cell adhesion molecule. Since Syk is expressed in blood platelets and is involved in the activation thereof, an inhibitor of such Syk is anticipated to be effective as a therapeutic agent for diseases associated with the activation of blood platelets.

30 [0009]

 A large number of compounds having Syk-inhibitory activity have been reported (Patent Literatures 1 to 4). Useful compounds (Non Patent Literature 7) and compounds having Syk and/or JAK inhibitory activity (Patent Literatures 5 to 8) have been reported from clinical tests in which
35 rheumatoid arthritis and idiopathic thrombocytopenic purpura have been targeted.

[Prior Art Literatures]

[Patent Literature]

[0010]

[Patent Literature 1] International Publication WO00/75113

5 [Patent Literature 2] JP Patent Publication (Kokai) No. 2008-013499 A

[Patent Literature 3] International Publication WO07/120980

[Patent Literature 4] International Publication WO07/124221

[Patent Literature 5] International Publication WO09/026107

[Patent Literature 6] International Publication WO09/131687

10 [Patent Literature 7] International Publication WO09/136995

[Patent Literature 8] International Publication WO09/145856

[Non Patent Literature]

[0011]

15 [Non Patent Literature 1] The Journal of Biological Chemistry, Vol. 266, pp. 15790-15796, 1991

[Non Patent Literature 2] International Journal of Hematology, Vol. 75, No. 4, pp. 357-362, 2002

[Non Patent Literature 3] The Journal of Biological Chemistry, Vol. 270, pp. 16189-16197, 1995

20 [Non Patent Literature 4] The Journal of Immunology, Vol. 167, No. 11, pp. 6292-6302, 2001

[Non Patent Literature 5] Expert Opinion on Investigational Drugs, Vol. 13, No. 7, pp. 743-762, 2004

25 [Non Patent Literature 6] Expert Opinion on Therapeutic Targets, Vol. 9, No. 5, pp. 901-921, 2005

[Non Patent Literature 7] IDrugs, Vol. 12, No. 3, pp. 174-185, 2009

[Summary of Invention]

[Problem to be solved by the Invention]

[0012]

30 To date, various Syk inhibitors have been reported, but they have not been placed on the market yet. It has been desired to develop a compound and a pharmaceutical composition, which have excellent Syk-inhibitory activity.

[Means for Solution to Problem]


35 [0013]

As a result of intensive studies directed towards achieving the

aforementioned object, the present inventors have found that a nicotinamide derivative having a specific structure or a salt thereof has excellent Syk-inhibitory activity, thereby completing the present invention.

Specifically, the nicotinamide derivative of the present invention or a pharmaceutically acceptable salt thereof is characterized in that it is represented by the following formula (I):

[Formula 1]




(I)

wherein

R¹ is a substituent represented by the following formula (II-1), (III-1), or (IV-1):

[Formula 2]



(wherein

R³ represents a hydrogen atom or a C₁₋₆ alkyl, C₃₋₈ cycloalkyl, phenyl, pyridyl, or thienyl group, each of which optionally has at least one substituent,

R⁴ represents a hydrogen atom, a C₁₋₆ alkyl group, or a C₃₋₈ cycloalkyl group,

R⁵ represents a hydroxy group, a halogen atom, or a C₁₋₆ alkyl or C₁₋₆ alkoxy group, each of which optionally has at least one substituent,

n is an integer of 0-2,

R⁵ may be the same or different, and two R⁵s may, together with the carbon atom to which they bind, form a C₃₋₈ cycloalkane ring when n is 2,

X¹ represents an oxygen atom or -N(R⁶)- (wherein R⁶ represents a hydrogen atom or an acyl group),

"*" represents a binding site), and

R² represents a pyridyl, indazolyl, phenyl, pyrazolopyridyl, benzisoxazolyl, pyrimidinyl, or quinolyl group, each of which optionally has at least one substituent.

[0014]

In addition, the present invention provides a pharmaceutical composition comprising the above-described nicotinamide derivative or a salt thereof, particularly, a pharmaceutical composition for use in the treatment of a Syk-related disease, which comprises the above-described nicotinamide derivative or a salt thereof, and a pharmaceutical composition for use in the treatment of a disease selected from the group consisting of rheumatoid arthritis and idiopathic thrombocytopenic purpura, which comprises the above-described nicotinamide derivative or a salt thereof.

[0015]

From a further viewpoint, the present invention provides: use of the above-described nicotinamide derivative or a salt thereof for production of the above-described pharmaceutical composition; a method for treating a Syk-related disease, which comprises a step of administering a therapeutically effective amount of the above-described nicotinamide derivative or a salt thereof to mammals including a human; and a method for treating a disease selected from the group consisting of rheumatoid arthritis and idiopathic thrombocytopenic purpura, which comprises a step of administering a therapeutically effective amount of the above-described nicotinamide derivative or a salt thereof to mammals including a human.

[Advantageous Effects of Invention]

[0016]

The nicotinamide derivative of the present invention or a salt thereof has excellent Syk-inhibitory activity, and it is useful as a pharmaceutical composition for use in the treatment of a Syk-related disease.

[Description of Embodiments]

[0017]

Hereinafter, the compound of the present invention will be described in detail.

It is to be noted that the following definitions are applied in the present specification, unless otherwise specified.

The term "halogen atom" is used herein to mean a fluorine atom, a chlorine atom, a bromine atom or an iodine atom.

The term " C_{1-3} alkyl group" is used herein to mean a methyl, ethyl, propyl, or isopropyl group.

The term " C_{1-4} alkyl group" is used herein to mean a methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, isobutyl, or tert-butyl group.

The term "C₁₋₆ alkyl group" is used herein to mean a linear or branched C₁₋₆ alkyl group such as a methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, isobutyl, tert-butyl, pentyl, isopentyl, or hexyl group.

5 The term "C₂₋₄ alkyl group" is used herein to mean an ethyl, propyl, isopropyl, butyl, sec-butyl, isobutyl, or tert-butyl group.

The term "C₄₋₆ alkyl group" is used herein to mean a linear or branched C₄₋₆ alkyl group such as a butyl, sec-butyl, isobutyl, tert-butyl, pentyl, isopentyl, or hexyl group.

10 The term "C₃₋₈ cycloalkyl group" is used herein to mean a C₃₋₈ cycloalkyl group such as a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl group.

The term "ar-C₁₋₆ alkyl group" is used herein to mean an ar-C₁₋₆ alkyl group such as a benzyl, diphenylmethyl, trityl, phenethyl, or naphthylmethyl group.

15 [0018]

The term "C₁₋₃ alkoxy group" is used herein to mean a methoxy, ethoxy, propoxy, or isopropoxy group.

20 The term "C₁₋₆ alkoxy group" is used herein to mean a linear or branched C₁₋₆ alkyloxy group such as a methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy, or hexyloxy group.

The term "C₁₋₆ alkoxy C₁₋₆ alkyl group" is used herein to mean a C₁₋₆ alkoxy C₁₋₆ alkyl group such as a methoxymethyl or 1-ethoxyethyl group.

The term "C₁₋₃ alkylthio group" is used herein to mean a C₁₋₃ alkylthio group such as a methylthio, ethylthio, or propylthio group.

25 The term "C₁₋₆ alkylthio group" is used herein to mean a C₁₋₆ alkylthio group such as a methylthio, ethylthio, propylthio, butylthio, or pentylthio group.

30 The term "C₁₋₆ alkylsulfonyl group" is used herein to mean a C₁₋₆ alkylsulfonyl group such as a methylsulfonyl, ethylsulfonyl, or propylsulfonyl group.

The term "arylsulfonyl group" is used herein to mean a benzenesulfonyl, p-toluenesulfonyl, or naphthalenesulfonyl group.

35 The term "C₁₋₆ alkylsulfonyloxy group" is used herein to mean a C₁₋₆ alkylsulfonyloxy group such as a methylsulfonyloxy or ethylsulfonyloxy group.

The term "arylsulfonyloxy group" is used herein to mean a benzenesulfonyloxy or p-toluenesulfonyloxy group.

[0019]

The term " C_{2-12} alkanoyl group" is used herein to mean a linear or
5 branched C_{2-12} alkanoyl group such as an acetyl, propionyl, valeryl, isovaleryl, or pivaloyl group.

The term "aroyl group" is used herein to mean a benzoyl or naphthoyl group.

The term "acyl group" is used herein to mean a formyl group, a C_{2-12}
10 alkanoyl group, or an aroyl group.

The term " C_{1-6} alkoxycarbonyl group" is used herein to mean a linear or branched C_{1-6} alkoxycarbonyl group such as a methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tert-butoxycarbonyl, or 1,1-dimethylpropoxycarbonyl group.

15 The term "ar- C_{1-6} alkyloxycarbonyl group" is used herein to mean an ar- C_{1-6} alkyloxycarbonyl group such as a benzyloxycarbonyl or phenethyloxycarbonyl group.

The term "aryloxycarbonyl group" is used herein to mean a phenyloxycarbonyl or naphthyloxycarbonyl group.

20 [0020]

The term " C_{1-6} alkylamino group" is used herein to mean a linear or branched C_{1-6} alkylamino group such as a methylamino, ethylamino, propylamino, isopropylamino, butylamino, sec-butylamino, tert-butylamino, pentylamino, or hexylamino group.

25 The term "di- C_{1-6} alkylamino group" is used herein to mean a linear or branched di- C_{1-6} alkylamino group such as a dimethylamino, diethylamino, di propylamino, diisopropylamino, dibutylamino, di(tert-butyl)amino, dipentylamino, dihexylamino, (ethyl)(methyl)amino, or (methyl)(propyl)amino group.

30 The term "(di) C_{1-6} alkylamino group" is used herein to mean a C_{1-6} alkylamino group or di- C_{1-6} alkylamino group.

The term "silyl group" is used herein to mean a trimethylsilyl, triethylsilyl, or tributylsilyl group.

[0021]

35 The term " C_{3-8} cycloalkane" is used herein to mean a C_{3-8} cycloalkane ring such as a cyclopropane, cyclobutane, cyclopentane, or cyclohexane ring.

[0022]

The amino-protecting group includes all groups that can be used as protecting groups for ordinary amino groups. Examples of such an amino-protecting group include groups described in W. Greene et al.,
5 Protective Groups in Organic Synthesis, Vol. 4, pp. 696 to 926, 2007, John Wiley & Sons, INC. Specific examples include an ar-C₁₋₆ alkyl group, a C₁₋₆ alkoxy C₁₋₆ alkyl group, an acyl group, a C₁₋₆ alkoxycarbonyl group, an ar-C₁₋₆ alkyloxycarbonyl group, an aryloxycarbonyl group, a C₁₋₆ alkylsulfonyl group, an arylsulfonyl group, and a silyl group.

10 [0023]

The hydroxy-protecting group includes all groups that can be used as protecting groups for ordinary hydroxy groups. Examples of such a hydroxy-protecting group include groups described in W. Greene et al.,
Protective Groups in Organic Synthesis, Vol. 4, pp. 16 to 299, 2007, John
15 Wiley & Sons, INC. Specific examples include a C₁₋₆ alkyl group, an ar-C₁₋₆ alkyl group, a C₁₋₆ alkoxy C₁₋₆ alkyl group, acyl group, a C₁₋₆ alkoxycarbonyl group, an ar-C₁₋₆ alkyloxycarbonyl group, a C₁₋₆ alkylsulfonyl group, an arylsulfonyl group, a silyl group, a tetrahydrofuranyl group, and a tetrahydropyranyl group.

20 [0024]

The carboxyl-protecting group includes all groups that can be used as protecting groups for ordinary carboxyl groups. Examples of such a carboxyl-protecting group include groups described in W. Greene et al.,
Protective Groups in Organic Synthesis, Vol. 4, pp. 533 to 643, 2007, John
25 Wiley & Sons, INC. Specific examples include a C₁₋₆ alkyl group, an ar-C₁₋₆ alkyl group, a C₁₋₆ alkoxy C₁₋₆ alkyl group, an ar-C₁₋₆ alkyloxy C₁₋₆ alkyl group, and a silyl group.

[0025]

Examples of a leaving group include a halogen atom, a C₁₋₆
30 alkylsulfonyloxy group, and an arylsulfonyloxy group.

[0026]

Alcohols include methanol, ethanol, propanol, 2-propanol, butanol, and 2-methyl-2-propanol.

Aliphatic hydrocarbons include pentane, hexane, and cyclohexane.

35 Halogenated hydrocarbons include methylene chloride, chloroform, and dichloroethane.

Aromatic hydrocarbons include benzene, toluene, and xylene.

Glycols include ethylene glycol, propylene glycol, and diethylene glycol.

Ethers include diethyl ether, diisopropyl ether, dioxane, tetrahydrofuran, anisole, ethylene glycol dimethyl ether, diethylene glycol dimethyl ether, and diethylene glycol diethyl ether.

[0027]

Ketones include acetone, 2-butanone, and 4-methyl-2-pentanone.

Esters include methyl acetate, ethyl acetate, propyl acetate, and butyl acetate.

Amides include N,N-dimethylformamide, N,N-dimethylacetamide, and 1-methyl-2-pyrrolidone.

Nitriles include acetonitrile and propionitrile.

Sulfoxides include dimethyl sulfoxide.

[0028]

Salts of the compound represented by the formula (1) include generally known salts, namely, the salts of basic groups such as amino groups, and the salts of acidic groups such as hydroxy or carboxyl groups.

[0029]

Examples of the salts of basic groups include: salts with mineral acids such as hydrochloric acid, hydrobromic acid, nitric acid, and sulfuric acid; salts with organic carboxylic acids such as formic acid, acetic acid, citric acid, oxalic acid, fumaric acid, maleic acid, succinic acid, malic acid, tartaric acid, aspartic acid, trichloroacetic acid, and trifluoroacetic acid; and salts with sulfonic acids such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, mesitylenesulfonic acid, and naphthalenesulfonic acid.


[0030] Examples of the salts of acidic groups include: salts with alkaline metals such as sodium and potassium; salts with alkaline earth metals such as calcium and magnesium; ammonium salts; and salts with nitrogen-containing organic bases such as trimethylamine, triethylamine, tributylamine, pyridine, N,N-dimethyl aniline, N-methyl piperidine, N-methyl morpholine, diethylamine, dicyclohexylamine, procaine, dibenzylamine, N-benzyl- β -phenethylamine, 1-ephedrine, and N,N'-dibenzylethylenediamine.

Among the above-described salts, pharmaceutically acceptable salts are preferable.

[0031]

The nicotinamide derivative of the present invention is characterized in that it is represented by the following formula (I):

[Formula 3]




(I)

[0032]


R^1 is a substituent represented by the following formula (II-1), (III-1), or (IV-1):

[Formula 4]




(wherein R^3 , R^4 , R^5 , n , and X^1 have the same definitions as those described above), preferably a substituent represented by the following formula (II-2), (III-2), or (IV-2):

[Formula 5]



(wherein R^3 , R^4 , R^5 , n , and X^1 have the same definitions as those described above), and more preferably a substituent represented by the following formula (II-2) or (III-2):

[Formula 6]



(wherein R^3 , R^4 , R^5 , and n have the same definitions as those described above).

[0033]

R^3 is a hydrogen atom or a C_{1-6} alkyl, C_{3-8} cycloalkyl, phenyl, pyridyl, or thienyl group, each of which optionally has at least one substituent and preferably a C_{1-6} alkyl or C_{3-8} cycloalkyl group, each of which optionally has at least one substituent.

When R^3 is a C_{1-6} alkyl or C_{3-8} cycloalkyl group, each of which optionally has at least one substituent, toxicity of the compound can be further reduced.

R^3 is preferably a phenyl, pyridyl, or thienyl group, each of which optionally has at least one substituent. When R^3 is a phenyl, pyridyl, or thienyl group, each of which optionally has at least one substituent, pharmacological activity of the compound is improved.

When R^3 is a C_{1-6} alkyl group, it is preferably a C_{1-4} alkyl group. Preferred examples include methyl, ethyl, n-propyl, isopropyl, and isobutyl groups.

When R^3 is a C_{3-8} cycloalkyl group, it is preferably a cyclopropyl or cyclobutyl group and more preferably a cyclopropyl group.

[0034]

A substituent on a C_{1-6} alkyl, C_{3-8} cycloalkyl, phenyl, pyridyl, or thienyl group represented by R^3 is preferably selected from a substituent group α_{1-1} and more preferably selected from a substituent group α_{1-2} . Toxicity of the compound having such a substituent can be further reduced.

The substituent group α_{1-1} includes a halogen atom and C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkylthio, phenyl, and pyrazolyl groups, each of which has at least one halogen atom.

The substituent group α_{1-2} includes a halogen atom, a C_{1-6} alkyl group, and a C_{1-6} alkoxy group.

A halogen atom in each of the substituent groups α_{1-1} and α_{1-2} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

A C_{1-6} alkyl group in each of the substituent groups α_{1-1} and α_{1-2} is preferably a C_{1-3} alkyl group, more preferably a methyl group or an ethyl group, and further preferably a methyl group.

A C₃₋₈ cycloalkyl group in the substituent group α_{1-1} is preferably a cyclopropyl group or a cyclobutyl group and more preferably a cyclopropyl group.

5 A C₁₋₆ alkoxy group in each of the substituent groups α_{1-1} and α_{1-2} is preferably a C₁₋₃ alkoxy group, more preferably a methoxy group or an ethoxy group, and further preferably a methoxy group.

A C₁₋₆ alkylthio group in the substituent group α_{1-1} is preferably a C₁₋₃ alkylthio group, more preferably a methylthio group or an ethylthio group, and further preferably a methylthio group.

10 [0035]

When R³ is a C₁₋₆ alkyl group, the C₁₋₆ alkyl group is preferably unsubstituted or substituted with a halogen atom or a C₁₋₆ alkoxy group and more preferably unsubstituted or substituted with a C₁₋₆ alkoxy group.

15 When R³ is a C₃₋₈ cycloalkyl group, the C₃₋₈ cycloalkyl group is preferably unsubstituted or substituted with a halogen atom or a C₁₋₆ alkyl group.

When R³ is a thienyl group, the thienyl group is preferably unsubstituted or substituted with a C₁₋₆ alkyl group.

[0036]

20 R⁴ is a hydrogen atom, a C₁₋₆ alkyl group, or a C₃₋₈ cycloalkyl group, preferably a hydrogen atom or a C₁₋₃ alkyl group, more preferably a hydrogen atom, a methyl group, or an ethyl group, and further preferably a hydrogen atom or a methyl group.

[0037]

25 When R³ is a C₄₋₆ alkyl group, R⁴ is preferably a hydrogen atom. When R³ is a C₄₋₆ alkyl group, the C₄₋₆ alkyl group is preferably an isobutyl group. Toxicity of the compound having such a substituent can be further reduced.

30 When R³ is a C₁₋₃ alkyl or thienyl group, R⁴ is preferably a methyl group. Toxicity of the compound having such a substituent can be further reduced.

[0038]

R⁵ is a hydroxy group, a halogen atom, or a C₁₋₆ alkyl or C₁₋₆ alkoxy group, each of which optionally has at least one substituent.

35 Here, n is an integer of 0 to 2. When n is 2, R⁵ may be the same or different. Also, two R⁵s may, together with the carbon atom to which they

bind, form a C₃₋₈ cycloalkane ring. The C₃₋₈ cycloalkane ring is preferably a cyclopropane or cyclobutane ring and more preferably a cyclopropane ring.

Here, R⁵ is preferably a hydroxy group, a halogen atom, or a C₁₋₆ alkyl or C₁₋₆ alkoxy group, each of which optionally has at least one phenyl group and more preferably a hydroxy group or a halogen atom.

Here, n is an integer of 0-2, preferably an integer of 0 or 1, and more preferably an integer of 0.

[0039]

X¹ is an oxygen atom or -N(R⁶)- (wherein R⁶ has the same definition as that described above) and preferably an oxygen atom. When X¹ is an oxygen atom, pharmacological activity of the compound is improved.

R⁶ is a hydrogen atom or an acyl group, preferably a hydrogen atom or an acetyl group, and more preferably an acetyl group.

[0040]

R² is a pyridyl, indazolyl, phenyl, pyrazolopyridyl, benzisoxazolyl, pyrimidinyl, or quinolyl group, each of which optionally has at least one substituent, preferably a pyridyl or phenyl group, each of which optionally has at least one substituent, and more preferably a pyridyl group. When R² is a pyridyl or phenyl group, each of which optionally has at least one substituent, toxicity of the compound can be further reduced.

In addition, for idiopathic thrombocytopenic purpura (hereinafter also referred to as "ITP"), a pyridyl, phenyl, indazolyl, or pyrazolopyridyl group, each of which optionally has at least one substituent is preferable and an indazolyl or pyrazolopyridyl group, each of which optionally has at least one substituent is more preferable.

[0041]

A substituent that binds to a pyridyl, indazolyl, phenyl, pyrazolopyridyl, benzisoxazolyl, pyrimidinyl, or quinolyl group is preferably a substituent selected from a substituent group α_{2-1} .

The substituent group α_{2-1} includes: a halogen atom and C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, (di)C₁₋₆ alkylamino, acyl, pyrazolyl, triazolyl, morpholinyl, and pyrrolisyl groups, each of which optionally has at least one substituent selected from a substituent group β_{2-1} .

A halogen atom in the substituent group α_{2-1} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

A C₁₋₆ alkyl group in the substituent group α_{2-1} is preferably a C₁₋₃ alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

5 A C₃₋₈ cycloalkyl group in the substituent group α_{2-1} is preferably a cyclopropyl or cyclobutyl group and more preferably a cyclopropyl group.

A C₁₋₆ alkoxy group in the substituent group α_{2-1} is preferably a C₁₋₃ alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

10 A (di)C₁₋₆ alkylamino group in the substituent group α_{2-1} is a mono-C₁₋₆ alkylamino or di-C₁₋₆ alkylamino group and preferably a di-C₁₋₆ alkylamino group. Here, a C₁₋₆ alkyl group that binds to a nitrogen atom is preferably a C₁₋₃ alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

Acyl in the substituent group α_{2-1} is preferably an acetyl group.

15 [0042]

The substituent group β_{2-1} includes a halogen atom, oxo, and C₁₋₆ alkyl and C₁₋₆ alkoxy groups.

A halogen atom in the substituent group β_{2-1} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

20 A C₁₋₆ alkyl group in the substituent group β_{2-1} is preferably a C₁₋₃ alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

A C₁₋₆ alkoxy group in the substituent group β_{2-1} is preferably a C₁₋₃ alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

[0043]

When R² is pyridyl optionally having at least one substituent, a substituent that binds to the pyridyl group is preferably a substituent selected from the substituent group α_{2-1} , more preferably a substituent selected from a substituent group α_{3-1} , and further preferably a substituent selected from a substituent group α_{3-2} .

30 The substituent group α_{3-1} includes a halogen atom and C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, (di)C₁₋₆ alkylamino, acyl, pyrazolyl, triazolyl, morpholinyl, and pyrrolisyl groups, each of which optionally has at least one substituent selected from the following substituent group β_{3-1} . The preferred

35

ranges of the substituents are the same as those described for the substituent group α_{2-1} .

The substituent group β_{3-1} includes a halogen atom and C_{1-6} alkyl and C_{1-6} alkoxy groups. The preferred ranges of the substituents are the same as those described for the substituent group β_{2-1} .


The substituent group α_{3-2} includes a halogen atom and C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkoxy, (di) C_{1-6} alkylamino, pyrazolyl, triazolyl, and morpholinyl groups, each of which optionally has at least one substituent selected from the following substituent group β_{3-2} . The preferred ranges of the substituents are the same as those described for the substituent group α_{2-1} .

The substituent group β_{3-2} includes a halogen atom and a C_{1-6} alkyl group. The preferred ranges of the substituents are the same as those described for the substituent group β_{2-1} .

[0044]

When R^2 is pyridyl optionally having at least one substituent, a pyridyl group is preferably a substituent represented by the following formula (V-1) or (V-2):

[Formula 7]



(wherein R^7 , R^8 , R^9 , and R^{10} may be the same or different and are each a hydrogen atom or a substituent selected from the substituent group α_{3-2}), and more preferably a substituent represented by formula (V-1).

[0045]

R^7 is a hydrogen atom or a substituent selected from the substituent group α_{3-2} and preferably a hydrogen atom or a substituent selected from a substituent group α_{3-3} . Toxicity of a compound having such a substituent can be further reduced.

The substituent group α_{3-3} includes a halogen atom, a C_{1-6} alkyl group, and C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkoxy, pyrazolyl, and triazolyl groups, each of which optionally has at least one C_{1-6} alkyl group.

A halogen atom in the substituent group α_{3-3} is preferably a fluorine atom or chlorine atom and more preferably a fluorine atom.

A C₁₋₆ alkyl group in the substituent group α_{3-3} is preferably a C₁₋₃ alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

5 A C₃₋₈ cycloalkyl group in the substituent group α_{3-3} is preferably a cyclopropyl or cyclobutyl group and more preferably a cyclopropyl group.

A C₁₋₆ alkoxy group in the substituent group α_{3-3} is preferably a C₁₋₃ alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

[0046]

10 R⁸ is a hydrogen atom or a substituent selected from the substituent group α_{3-2} and preferably a hydrogen atom or a substituent selected from a substituent group α_{3-4} . Toxicity of a compound having such a substituent can be further reduced.

15 The substituent group α_{3-4} includes C₁₋₆ alkyl, C₁₋₆ alkoxy, and morpholinyl groups, each of which optionally has at least one halogen atom.

A halogen atom in the substituent group α_{3-4} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

20 A C₁₋₆ alkyl group in the substituent group α_{3-4} is preferably a C₁₋₃ alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

A C₁₋₆ alkoxy group in the substituent group α_{3-4} is preferably a C₁₋₃ alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

25 In addition, when R⁷ is a pyrazolyl or triazolyl group, R⁸ is preferably a hydrogen atom or a methyl group.

[0047]

30 R⁹ and R¹⁰ may be the same or different and are each a hydrogen atom or a substituent selected from the substituent group α_{3-2} and preferably a hydrogen atom or a substituent selected from a substituent group α_{3-5} . Toxicity of a compound having such a substituent can be further reduced.

The substituent group α_{3-5} includes C₁₋₆ alkyl and C₁₋₆ alkoxyl groups, each of which optionally has at least one halogen atom.

A halogen atom in the substituent group α_{3-5} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

A C₁₋₆ alkyl group in the substituent group α_{3-5} is preferably a C₁₋₃ alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

A C₁₋₆ alkoxy group in the substituent group α_{3-5} is preferably a C₁₋₃ alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

[0048]

When R² is an indazolyl group optionally having at least one substituent, a substituent that binds to the indazolyl group is preferably a substituent selected from the substituent group α_{2-1} , more preferably a substituent selected from a substituent group α_{4-1} , and further preferably a substituent selected from a substituent group α_{4-2} .

The substituent group α_{4-1} includes a halogen atom and C₁₋₆ alkyl, C₁₋₆ alkoxy, (di)C₁₋₆ alkylamino, and pyrrolsyl groups, each of which optionally has at least one substituent selected from a substituent group β_{4-1} . The preferred ranges of the substituents are the same as those described for the substituent group α_{2-1} .

The substituent group β_{4-1} includes a halogen atom, oxo, and a C₁₋₆ alkoxy group. The preferred ranges of the substituents are the same as those described for the substituent group β_{2-1} .


The substituent group α_{4-2} includes a halogen atom and C₁₋₆ alkyl, C₁₋₆ alkoxy, and (di)C₁₋₆ alkylamino groups, each of which optionally has at least one substituent selected from a substituent group β_{4-2} . The preferred ranges of the substituents are the same as those described for the substituent group α_{2-1} .

The substituent group β_{4-2} includes a halogen atom and a C₁₋₆ alkoxy group. The preferred ranges of the substituents are the same as those described for the substituent group β_{2-1} .

[0049]

When R² is an indazolyl group optionally having at least one substituent, the indazolyl group is preferably a substituent represented by the following formula (VI-1) or (VI-2):

[Formula 8]



(wherein R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} may be the same or different and are each a hydrogen atom or a substituent selected from the substituent group α_{4-2}).

5 [0050]

R^{11} and R^{14} are each a hydrogen atom or a substituent selected from the substituent group α_{4-2} and preferably a hydrogen atom or a substituent selected from a substituent group α_{4-3} . Toxicity of the compound having such a substituent can be further reduced.

10 The substituent group α_{4-3} includes C_{1-6} alkyl, C_{1-6} alkoxy, and (di) C_{1-6} alkylamino groups, each of which optionally has at least one halogen atom.

A halogen atom in the substituent group α_{4-3} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

15 A C_{1-6} alkyl group in the substituent group α_{4-3} is preferably a C_{1-3} alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

A C_{1-6} alkoxy group in the substituent group α_{4-3} is preferably a C_{1-3} alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

20 A (di) C_{1-6} alkylamino group in the substituent group α_{4-3} is a mono- C_{1-6} alkylamino or di- C_{1-6} alkylamino group and preferably a di- C_{1-6} alkylamino group. Here, a C_{1-6} alkyl group that binds to a nitrogen atom is preferably a C_{1-3} alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

25 [0051]

R^{12} and R^{15} are each a hydrogen atom or a substituent selected from the substituent group α_{4-2} and preferably a hydrogen atom or a substituent selected from a substituent group α_{4-4} . Toxicity of the compound having such a substituent can be further reduced.

30 The substituent group α_{4-4} includes a C_{1-6} alkyl group optionally having at least one substituent selected from a substituent group β_{4-4} .

A C₁₋₆ alkyl group in the substituent group α_{4-4} is preferably a C₁₋₃ alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

5 The substituent group β_{4-4} includes a halogen atom and a C₁₋₆ alkoxy group.

A halogen atom in the substituent group β_{4-4} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

10 A C₁₋₆ alkoxy group in the substituent group β_{4-4} is preferably a C₁₋₃ alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

[0052]

R¹³ and R¹⁶ are each preferably a hydrogen atom or a halogen atom. A halogen atom is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

15 [0053]

When R² is a phenyl group optionally having at least one substituent, a substituent that binds to the phenyl group is preferably a substituent selected from the substituent group α_{2-1} , more preferably a substituent selected from a substituent group α_{5-1} , and further preferably a substituent selected from a substituent group α_{5-2} .


20 The substituent group α_{5-1} includes a halogen atom and C₁₋₆ alkyl, C₁₋₆ alkoxy, acyl, and triazolyl groups, each of which optionally has at least one halogen atom. The preferred ranges of the substituents are the same as those described for the substituent group α_{2-1} .

25 The substituent group α_{5-2} includes a halogen atom, a C₁₋₆ alkoxy group, and C₁₋₆ alkyl and triazolyl groups, each of which optionally has at least one halogen atom. The preferred ranges of the substituents are the same as those described for the substituent group α_{2-1} .

[0054]

30 When R² is phenyl optionally having at least one substituent, a phenyl group is preferably a substituent represented by the following formula (VII-1):

[Formula 9]



(VII-1)

(wherein R^{17} , R^{18} , and R^{19} may be the same or different and are each a hydrogen atom or a substituent selected from the substituent group α_{5-2}).

[0055]

5 R^{17} and R^{18} are each a hydrogen atom or a substituent selected from the substituent group α_{5-2} and preferably a hydrogen atom or a substituent selected from a substituent group α_{5-3} . Toxicity of the compound having such a substituent can be further reduced.

10 The substituent group α_{5-3} includes a halogen atom, a C_{1-6} alkoxy group, a triazolyl group, and a C_{1-6} alkyl group optionally having at least one halogen atom.

A halogen atom in the substituent group α_{5-3} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

15 A C_{1-6} alkoxy group in the substituent group α_{5-3} is preferably a C_{1-3} alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

A C_{1-6} alkyl group in the substituent group α_{5-3} is preferably a C_{1-3} alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

20 [0056]

R^{19} is a hydrogen atom or a substituent selected from the substituent group α_{5-2} and preferably a hydrogen atom or a substituent selected from a substituent group α_{5-4} . Toxicity of the compound having such a substituent can be further reduced.

25 The substituent group α_{5-4} includes a halogen atom, a C_{1-6} alkoxy group, and a C_{1-6} alkyl group optionally having at least one halogen atom.

A halogen atom in the substituent group α_{5-4} is preferably a fluorine atom or a chlorine atom, and more preferably a fluorine atom.

30 A C_{1-6} alkoxy group in the substituent group α_{5-4} is preferably a C_{1-3} alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

A C₁₋₆ alkyl group in the substituent group α_{5-4} is preferably a C₁₋₃ alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

[0057]

- 5 When R² is a pyrazolopyridyl group optionally having at least one substituent, a substituent that binds to the pyrazolopyridyl group is preferably a substituent selected from the substituent group α_{2-1} and more preferably a substituent selected from a substituent group α_{6-1} .


10 The substituent group α_{6-1} includes a halogen atom and C₁₋₆ alkyl, C₁₋₆ alkoxy, and (di)C₁₋₆ alkylamino groups, each of which optionally has at least one substituent selected from a substituent group β_{6-1} . The preferred ranges of the substituents are the same as those described for the substituent group α_{2-1} .

15 The substituent group β_{6-1} includes a halogen atom and a C₁₋₆ alkoxy group. The preferred ranges of the substituents are the same as those described for the substituent group β_{2-1} .


[0058]

- 20 When R² is a pyrazolopyridyl group optionally having at least one substituent, the pyrazolopyridyl group is preferably a substituent represented by the following formula (VIII-1) or (VIII-2):

[Formula 10]



(VIII-1)



(VIII-2)

(wherein R²⁰, R²¹, R²², and R²³ may be the same or different and are each a hydrogen atom or a substituent selected from the substituent group α_{6-1}).

25 [0059]

R²⁰ is a hydrogen atom or a substituent selected from the substituent group α_{6-1} . Toxicity of the compound having such a substituent can be further reduced.

[0060]

- 30 R²¹ and R²³ are each a hydrogen atom or a substituent selected from the substituent group α_{6-1} and preferably a hydrogen atom or a substituent

selected from a substituent group α_{6-2} . Toxicity of the compound having such a substituent can be further reduced.

The substituent group α_{6-2} includes a C_{1-6} alkyl group optionally having at least one substituent selected from a halogen atom and a C_{1-6} alkoxy group.

A halogen atom in the substituent group α_{6-2} is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom.

A C_{1-6} alkoxy group in the substituent group α_{6-2} is preferably a C_{1-3} alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

A C_{1-6} alkyl group in the substituent group α_{6-2} is preferably a C_{1-3} alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

[0061]

R^{22} is a hydrogen atom or a substituent selected from the substituent group α_{6-1} and preferably a hydrogen atom, a halogen atom, or a C_{1-6} alkyl group. Here, a halogen atom is preferably a fluorine atom or a chlorine atom and more preferably a fluorine atom. In addition, a C_{1-6} alkyl group is preferably a C_{1-3} alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.

[0062]

When R^2 is a benzisoxazolyl group optionally having at least one substituent, a substituent that binds to the benzisoxazolyl group is preferably a substituent selected from the substituent group α_{2-1} and more preferably a C_{1-6} alkoxy group. Here, a C_{1-6} alkoxy group is preferably a C_{1-3} alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group.

[0063]

When R^2 is a pyrimidinyl group optionally having at least one substituent, a substituent that binds to the pyrimidinyl group is preferably a substituent selected from the substituent group α_{2-1} and more preferably C_{1-6} alkyl, C_{1-6} alkoxy, pyrazolyl, triazolyl, or morpholinyl. Here, a C_{1-6} alkoxy group is preferably a C_{1-3} alkoxy group, more preferably a methoxy or ethoxy group, and further preferably a methoxy group. A C_{1-6} alkyl group is preferably a C_{1-3} alkyl group, more preferably a methyl or ethyl group, and further preferably a methyl group.


[0064]

When R^2 is a quinolyl group optionally having at least one substituent, a substituent that binds to the quinolyl group is preferably a substituent selected from the substituent group α_{2-1} .

5 [0065]

The nicotinamide derivative of the present invention is represented preferably by the following formula (I-1) and more preferably by the following formula (I-2):

[Formula 11]




(wherein R^2 , R^3 , and R^4 are the same substituents as those described above and the preferred ranges thereof are also the same as those described above).

[0066]

15 The nicotinamide derivative of the present invention is represented preferably by the following formula (I-3) and more preferably by the following formula (I-4):

[Formula 12]




(wherein R^2 is the same substituent as that described above and the preferred range thereof is also the same as that described above).

[0067]

The nicotinamide derivative of the present invention is represented preferably by the following formula (I-5) and more preferably by the following formula (I-6):

25 [Formula 13]



(wherein R^2 is the same substituent as that described above and the preferred

range thereof is also the same as that described above).

[0068]

Preferred examples of the compound represented by the formula (1) of the present invention include the following compounds:

5 Example 2-1:

6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-10:

10 6-(((1R,2S)-2-amino-1-cyclobutyl propyl)amino)-5-fluoro-2-((1-methyl-1H-indazol-6-yl)amino)nicotinamide

Example 2-123:

6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxy ethyl)-1H-indazol-5-yl)amino)nicotinamide

Example 2-125:

15 6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxy ethyl)-1H-indazol-5-yl)amino)nicotinamide

Example 2-126:

6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxy ethyl)-1H-indazol-5-yl)amino)nicotinamide

20 Example 2-130:

(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-131:

25 6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-133:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-137:

30 (R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-138:

6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

35 Example 2-139:

6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-3

-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-142:

6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide

5 Example 2-148:

6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((7-chloro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-149:

10 6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((7-chloro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-159:

6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-17:

15 6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((5,6-dimethylpyridin-3-yl)amino)-5-fluoronicotinamide

Example 2-173:

6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

20 Example 2-18:

6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((5,6-dimethylpyridin-3-yl)amino)-5-fluoronicotinamide

Example 2-182:

25 6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-184:

6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-186:

30 6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-187:

6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

35 Example 2-188:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3-(dimethylamino)-7-fl

uoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-196:

6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((1,3-dimethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

5 Example 2-20:

6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide

Example 2-207:

10 6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide

Example 2-208:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1,3-dimethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-209:

15 6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-fluoro-1-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-210:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide

20 Example 2-211:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-213:

25 6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-214:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide

Example 2-218:

30 6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinamide

Example 2-23:

6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide

35 Example 2-230:

6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-2-((1-ethyl-1H-indazo

1-5-yl)amino)-5-fluoronicotinamide

Example 2-235:

6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((3-fluoro-2-morpholinopyridin-4-yl)amino)nicotinamide

5 Example 2-249:

6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide

Example 2-253:

10 6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide

Example 2-265:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide

Example 2-266:

15 6-(((3R,4S)-4-aminoheptan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide

Example 2-267:

6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide

20 Example 2-27:

6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((3-methoxyphenyl)amino)nicotinamide

Example 2-270:

25 6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide

Example 2-273:

6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)nicotinamide

Example 2-28:

30 6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide

Example 2-29:

6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3,4-dimethoxyphenyl)amino)-5-fluoronicotinamide

35 Example 2-31:

6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-tr

iazol-2-yl)pyridin-3-yl)amino)nicotinamide

Example 2-316:

6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide

5 Example 2-317:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide

Example 2-319:

10 6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide

Example 2-320:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide

Example 2-322:

15 6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide

Example 2-326:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

20 Example 2-328:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-329:

25 6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide

Example 2-330:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide

Example 2-332:

30 6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide

Example 2-362:

6-(((1R,2S)-2-amino-1-cyclopropyl-butyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

35 Example 2-37:

6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((1-methyl-1H-i

ndazol-6-yl)amino)nicotinamide

Example 2-375:

6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide

5 Example 2-376:

6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide

Example 2-377:

10 6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide

Example 2-378:

6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide

Example 2-38:

15 6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-6-yl)amino)nicotinamide

Example 2-381:

6-(((1S,2S)-2-amino-1-(pyridin-2-yl)propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

20 Example 2-39:

6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-6-yl)amino)nicotinamide

Example 2-4:

25 6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-40:

6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-404:

30 6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-(quinolin-5-ylamino)nicotinamide

Example 2-41:

6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((5-cyclopropylpyridin-3-yl)amino)-5-fluoronicotinamide

35 Example 2-410:

6-(((1R,2S)-2-amino-1-(2-fluorophenyl)

propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-413:

6-(((1R,2S)-2-amino-1-(3-fluorophenyl)

propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

5 Example 2-414:

6-(((1R,2S)-2-amino-1-(3-fluorophenyl)

propyl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide

Example 2-416:

6-(((1S,2S)-2-amino-1-(thiophen-2-yl)

10 propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-42:

6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyr
idin-4-yl)amino)nicotinamide

Example 2-434:

15 6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-(m-tolylamino)nicotinami
de

Example 2-437:

2-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-6-(((1R,2S)-2-aminocyclohexyl)a
mino)-5-fluoronicotinamide

20 Example 2-438:

2-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-6-(((1R,2S)-2-aminocyclohexyl)a
mino)-5-fluoronicotinamide

Example 2-439:

25 6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-chlorophenyl)amino)-5-fluoroni
cotinamide

Example 2-44:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-(m-tolylamino)n
icotinamide

Example 2-442:

30 2-((3-acetylphenyl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoroni
cotinamide

Example 2-454:

6-(((1R,2S)-2-amino-1-(3-fluorophenyl)

propyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicoti
35 namide

Example 2-46:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-methoxyphenyl)amino)nicotinamide

Example 2-47:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide

Example 2-472:

6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide

Example 2-475:

2-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoronicotinamide

Example 2-476:

2-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoronicotinamide

Example 2-478:

2-((3-acetylphenyl)amino)-6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoronicotinamide

Example 2-479:

2-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoronicotinamide

Example 2-48:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3,4-dimethoxyphenyl)amino)-5-fluoronicotinamide

Example 2-480:

2-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoronicotinamide

Example 2-481:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3,4-dimethylphenyl)amino)-5-fluoronicotinamide

Example 2-482:

2-((3-acetylphenyl)amino)-6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoronicotinamide

Example 2-5:

6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((2-methoxypyridin-4-yl)amino)nicotinamide

Example 2-508:

6-(((1S,2S)-2-amino-1-(thiophen-2-yl)propyl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide

Example 2-518:

5 2-((5-acetyl-6-methylpyridin-3-yl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoronicotinamide

Example 2-521:

6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide

10 Example 2-57:

6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((3-methoxybenzo[d]isoxazol-5-yl)amino)nicotinamide

Example 2-7:

15 6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide

Example 2-71:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide

20 Example 2-74:

6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-(2,2-difluoroethyl)-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide

Example 2-78:

25 (R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-8:

6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-(quinolin-6-ylamino)nicotinamide

Example 2-80:

30 6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide

Example 2-9:

6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide

35 [0069]


The compound represented by the formula (1) of the present invention

Examples of Syk-related diseases of the present invention include bronchial asthma, allergic rhinitis, hives, atopic dermatitis, rheumatoid arthritis, systemic lupus erythematosus, autoimmune hemolytic anemia, nephrotic syndrome, contact dermatitis, idiopathic thrombocytopenic purpura, lymphocytic leukemia, and acute myelocytic leukemia. Rheumatoid arthritis or idiopathic thrombocytopenic purpura is preferable. Idiopathic thrombocytopenic purpura is more preferable.

Next, a method for producing the compound of the present invention will be described.

[0072]

[Formula 14]



[0073]

(A1-1)

The compound of the formula [2] can be produced by hydrolyzing the compound of the formula [1] in the presence of a base and in the presence of
5 hydrogen peroxide.

[0074]

The solvent used in this reaction is not particularly limited, as long as it does not affect the reaction. Examples of such a solvent include aliphatic hydrocarbons, halogenated hydrocarbons, alcohols, glycols, ethers, ketones,
10 esters, amides, sulfoxides, aromatic hydrocarbons, and water. These substances may be used in combination.

Preferred solvents are alcohols and water.

[0075]

Examples of the base used in this reaction include: metal alkoxides
15 such as sodium methoxide, sodium ethoxide, potassium tert-butoxide, and sodium tert-butoxide; inorganic bases such as sodium hydroxide, potassium hydroxide, sodium hydrogencarbonate, sodium carbonate, potassium carbonate, sodium hydride, and potassium hydride; and organic bases such as triethylamine, diisopropylethylamine, and pyridine.

20 [0076]

The base may be used in a molar concentration 1 time or more, and preferably 1 to 10 times, higher than that of the compound of the formula [1].

The hydrogen peroxide may be used in a molar concentration 1 time or more, and preferably 1 to 10 times, higher than that of the compound of the
25 formula [1].

This reaction may be carried out at a temperature from 0°C to the boiling point of a solvent, and preferably from 10°C to 40°C, for 1 minute to 24 hours.

[0077]

30 (A1-2)

The compound of the formula [3] can be produced by deprotecting the compound of the formula [2] in the presence of an acid. This reaction can be carried out, for example, by the method described in W. Greene et al., Protective Groups in Organic Synthesis, Vol. 4, pp. 696 to 926, 2007, John
35 Wiley & Sons, INC.

[0078]

Examples of the acid used in this reaction include: inorganic acids such as hydrochloric acid, sulfuric acid, phosphoric acid, hydrogen chloride, and hydrogen bromide; organic carboxylic acids such as acetic acid, trichloroacetic acid, and trifluoroacetic acid; and organic sulfonic acids such as methanesulfonic acid and p-toluenesulfonic acid.

The acid may be used in a molar concentration 1 time or more, and preferably 1 to 5 times, higher than that of the compound of the formula [2]. In addition, such an acid may be used as a solvent.

[0079]


This reaction may be carried out in the coexistence of a solvent, as necessary. The solvent used is not particularly limited, as long as it does not affect the reaction. Examples of such a solvent include aliphatic hydrocarbons, halogenated hydrocarbons, alcohols, glycols, ethers, ketones, esters, amides, nitriles, sulfoxides, aromatic hydrocarbons, and water. These substances may be used in combination.

This reaction may be carried out at a temperature from 0°C to the boiling point of a solvent, and preferably from 10°C to 40°C, for 1 minute to 24 hours.

[0080]

[Production Method 2]

[Formula 15]



"wherein La and Lb may be the same or different, and a leaving group, Re, R^{a1}, R^{a2}, R^{b1}, R^{b2}, and R² have the same definitions as those described above."

[0081]

(A2-1)

The compound of the formula [6] can be produced by allowing the compound of the formula [4] to react with the compound of the formula [5] in the presence of a base.

The compound of the formula [4] can be produced by, for example, Production Method 3 described below.

A known example of the compound of the formula [4] is tert-butyl

((1R,2S)-1-amino-1-cyclobutylpropan-2-yl)carbamate.

A known example of the compound of the formula [5] is 2,6-dichloro-5-fluoronicotinonitrile.

[0082]

5 The solvent used in this reaction is not particularly limited, as long as it does not affect the reaction. Examples of such a solvent include aliphatic hydrocarbons, halogenated hydrocarbons, alcohols, glycols, ethers, ketones, esters, amides, nitriles, sulfoxides, aromatic hydrocarbons, and water. These substances may be used in combination.

10 Preferred solvents are amides and ethers.

[0083]

 Examples of the base used in this reaction include: inorganic bases such as sodium hydrogen carbonate, sodium carbonate, potassium carbonate, cesium carbonate, and tripotassium phosphate; and organic bases such as
15 pyridine, 4-(dimethylamino)pyridine, triethylamine, and diisopropyl ethylamine.

 The base may be used in a molar concentration 1 to 50 times, and preferably 1 to 5 times, higher than that of the compound of the formula [4].

[0084]

20 The compound of the formula [5] may be used in a molar concentration 1 to 50 times, and preferably 1 to 2 times, higher than that of the compound of the formula [4].

 This reaction may be carried out at a temperature from 0°C to the boiling point of a solvent, and preferably from 10°C to 40°C, for 1 minute to
25 24 hours.

[0085]

(A2-2)

 The compound of the formula [1] can be produced by allowing the compound of the formula [6] to react with the compound of the formula [7] in
30 the presence or absence of a base, in the presence of a palladium catalyst, and in the presence or absence of a ligand.

 A known example of the compound of the formula [7] is 5-fluoro-6-morpholinopyridin-3-amine.

[0086]

35 The solvent used in this reaction is not particularly limited, as long as it does not affect the reaction. Examples of the solvent include aliphatic

hydrocarbons, halogenated hydrocarbons, alcohols, glycols, ethers, ketones, esters, amides, nitriles, sulfoxides, aromatic hydrocarbons, and water. These substances may be used in combination.

Preferred solvents are ethers.

5 [0087]

Examples of the base used in this reaction as desired include: inorganic bases such as sodium hydrogen carbonate, sodium carbonate, potassium carbonate, cesium carbonate, and tripotassium phosphate; and organic bases such as pyridine, 4-(dimethylamino)pyridine, triethylamine, and
10 diisopropylethylamine.

The base may be used in a molar concentration 1 to 50 times, and preferably 1 to 5 times, higher than that of the compound of the formula [6].

[0088]

Examples of the palladium catalyst used in this reaction include:
15 metallic palladium such as palladium carbon and palladium black; inorganic palladium salts such as palladium chloride; organic palladium salts such as palladium acetate; organic palladium complexes such as tetrakis(triphenylphosphine)palladium (0), bis(triphenylphosphine)palladium (II) chloride, 1,1'-bis(diphenylphosphino)ferrocene-palladium (II) chloride,
20 and tris(dibenzylideneacetone)dipalladium (0); and polymer-bound organic palladium complexes such as polymer-supported bis(acetate)triphenylphosphine palladium (II) and polymer-supported di(acetate)dicyclohexylphenylphosphine palladium (II). These compounds may be used in combination.

25 The palladium catalyst may be used in a molar concentration 0.00001 to 1 time, and preferably 0.001 to 0.1 time, as high as that of the compound of the formula [6].

[0089]

Examples of the ligand used in this reaction as desired include:
30 trialkylphosphines such as trimethylphosphine and tri-tert-butylphosphine; tricycloalkylphosphines such as tricyclohexylphosphine; triarylphosphines such as triphenylphosphine and tritolylphosphine; trialkylphosphites such as trimethylphosphite, triethylphosphite, and tributylphosphite; tricycloalkylphosphites such as tricyclohexylphosphite; triarylphosphites
35 such as triphenylphosphite; imidazolium salts such as 1,3-bis(2,4,6-trimethylphenyl)imidazolium chloride; diketones such as

acetylacetone and octafluoroacetylacetone; amines such as trimethylamine, triethylamine, tripropylamine, and triisopropylamine; and 4,5-bis(diphenylphosphino)-9,9-dimethyl-xanthene, 1,1'-bis(diphenylphosphino)ferrocene,
 5 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl,
 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl,
 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl,
 2-(di-tert-butylphosphino)-2',4',6'-triisopropylbiphenyl, and
 2-(di-tert-butylphosphino)biphenyl. These compounds may be used in
 10 combination.

Such a ligand may be used in a molar concentration 0.00001 to 1 time, and preferably 0.001 to 0.5 time, as high as that of the compound of the formula [6].

[0090]


15 The compound of the formula [7] may be used in a molar concentration 1 to 50 times, and preferably 1 to 2 times, higher than that of the compound of the formula [6].

This reaction may be preferably carried out in an inert gas (e.g. nitrogen, argon) atmosphere at a temperature from 40°C to 170°C for 1 minute
 20 to 96 hours.

[0091]

[Production Method 3]

[Formula 16]



25 "wherein Le represents a leaving group, and Re, R^{a1}, R^{a2}, R^{b1}, and R^{b2} have the same definitions as those described above."

[0092]

(A3-1)

The compound of the formula [9] can be produced by allowing the
 30 compound of the formula [8] to react with a sulfonylchloride in the presence of a base.

A known example of the compound of the formula [8] is tert-butyl ((2S)-1-cyclopropyl-1-hydroxypropan-2-yl)carbamate.

[0093]

The solvent used in this reaction is not particularly limited, as long as it does not affect the reaction. Examples of the solvent include aliphatic hydrocarbons, halogenated hydrocarbons, ethers, ketones, esters, nitriles, sulfoxides, and aromatic hydrocarbons. These substances may be used in combination.

Preferred solvents are ethers.

[0094]

Examples of the sulfonyl chloride used in this reaction include methylsulfonyl chloride, ethylsulfonyl chloride, propylsulfonyl chloride, benzenesulfonyl chloride, p-toluenesulfonyl chloride, and naphthalenesulfonyl chloride.

Preferred sulfonyl chlorides include methylsulfonyl chloride and p-toluenesulfonyl chloride. Further, methylsulfonyl chloride is more preferable.

The sulfonyl chloride is used in a molar concentration of 1 time or more, and preferably 1 to 3 times, higher than that of the compound of the formula [8].

[0095]

Examples of the base used in this reaction include: inorganic bases such as sodium hydrogen carbonate, sodium carbonate, potassium carbonate, cesium carbonate, and tripotassium phosphate; and organic bases such as pyridine, 4-(dimethylamino)pyridine, triethylamine, and diisopropylethylamine.

The base may be used in a molar concentration 1 time or more, and preferably 1 to 3 times, higher than that of the compound of the formula [8].

This reaction may be carried out at a temperature from -78°C to the boiling point of a solvent, and preferably from 0°C to 80°C , for 1 minute to 24 hours.

[0096]

(A3-2)

The compound of the formula [10] can be produced by allowing the compound of the formula [9] to react with a phthalimide.

When the compound of the formula [9] is in the form of a diastereomeric mixture, the diastereomeric mixture may be separated in a step of isolating the compound of the formula [10].

[0097]

The solvent used in this reaction is not particularly limited, as long as it does not affect the reaction. Examples of the solvent include aliphatic hydrocarbons, halogenated hydrocarbons, alcohols, glycols, ethers, ketones, esters, amides, nitriles, sulfoxides, aromatic hydrocarbons, and water. These substances may be used in combination.

Preferred solvents are amides.

[0098]

Examples of the phthalimide used in this reaction include phthalimide sodium and phthalimide potassium. A preferred phthalimide is phthalimide potassium.

Such a phthalimide can also be produced in a reaction system.

Such a phthalimide is used in a molar concentration 1 time or more, and preferably 1 to 3 times, higher than that of the compound of the formula [9].

This reaction may be carried out at a temperature from 0°C to the boiling point of a solvent, and preferably from 0°C to 100°C, for 1 minute to 24 hours.

[0099]

(A3-3)


The compound of the formula [4] can be produced by deprotecting the compound of the formula [10]. This reaction can be carried out, for example, by the method described in W. Greene et al., Protective Groups in Organic Synthesis, the fourth edition, pp. 696 to 926, 2007, John Wiley & Sons, INC.

In this reaction, deprotection is preferably carried out using hydrazine.

[0100]

[Production Method 4]

[Formula 17]



"wherein R^a represents a C₁₋₆ alkyl, C₃₋₈ cycloalkyl, phenyl, pyridyl, or thienyl group, each of which optionally has at least one substituent, and Re, R^{b1}, R^{b2}, and Le have the same definitions as those described above."

[0101]

5 (A4-1)

The compound of the formula [12] can be produced by activating a carboxyl group of the compound of the formula [11] and then allowing the compound to react with an amine under basic conditions.

[0102]

10 The solvent used in this reaction is not particularly limited, as long as it does not affect the reaction. Examples of the solvent include aliphatic hydrocarbons, halogenated hydrocarbons, alcohols, glycols, ethers, ketones, esters, amides, nitriles, sulfoxides, aromatic hydrocarbons, and water. These substances may be used in combination.

15 Preferred solvents are halogenated hydrocarbons and ethers.

[0103]

Examples of a carboxyl activator used in this reaction include: carbodiimides such as N,N'-dicyclohexyl carbodiimide, N,N'-diisopropyl carbodiimide, and N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide; azide
20 phosphates such as diphenylphosphoryl azide; phosphoniums such as BOP reagents; carbonyldiimidazoles such as 1,1'-carbonyldiimidazole; and acid halides such as thionyl chloride.

[0104]

Examples of the base used in this reaction include: metal alkoxides
25 such as sodium methoxide, sodium ethoxide, potassium tert-butoxide, and sodium tert-butoxide; inorganic bases such as sodium hydroxide, potassium hydroxide, sodium hydrogen carbonate, sodium carbonate, potassium carbonate, sodium hydride, and potassium hydride; and organic bases such as triethylamine, diisopropylethylamine, and pyridine.

30 Preferred bases are organic bases.

Examples of the amine used in this reaction include methoxymethylamine.

The amine is used in a molar concentration 1 time or more, and preferably 1 to 3 times, higher than that of the compound of the formula [11].

35 The carboxyl activator is used in a molar concentration 1 time or more, and preferably 1 to 3 times, higher than that of the compound of the formula

[11].

The base is used in a molar concentration 1 time or more, and preferably 1 to 3 times, higher than that of the compound of the formula [11].

This reaction may be carried out at a temperature from 0°C to the boiling point of a solvent, and preferably from 0°C to 100°C, for 1 minute to 24 hours.

[0105]

(A4-2)

The compound of the formula [13] can be produced by allowing the compound of the formula [12] to react with a Grignard reagent.

[0106]

The solvent used in this reaction is not particularly limited, as long as it does not affect the reaction. Examples of the solvent include aliphatic hydrocarbons, halogenated hydrocarbons, ethers, aromatic hydrocarbons, and water. These substances may be used in combination.

Preferred solvents are ethers.

The Grignard reagent is used in a molar concentration 1 time or more, and preferably 1 to 5 times, higher than that of the compound of the formula [12].

This reaction may be carried out at a temperature from 0°C to the boiling point of a solvent, and preferably from 0°C to 100°C, for 1 minute to 24 hours.

[0107]

(A4-3)

The compound of the formula [14] can be produced by allowing the compound of the formula [13] to react with (R)-(+)-tert-butyl sulfinamide in the presence of an additive having the Lewis acid action and dehydrating action and then reducing a resulting imine.

(S)-(-)-tert-butyl sulfinamide may be used instead of (R)-(+)-tert-butyl sulfinamide.

[0108]

The solvent used in a series of reactions is not particularly limited, as long as it does not affect the reaction. Examples of such a solvent include aliphatic hydrocarbons, halogenated hydrocarbons, ethers, amides, nitriles, sulfoxides, and aromatic hydrocarbons. These substances may be used in combination.

Preferred solvents are halogenated hydrocarbons, aromatic hydrocarbons, and ethers.

Examples of the additive having the Lewis acid action and dehydrating action used in this reaction include: carboxylic acids such as acetic acid, citric acid, and formic acid; and metal alkoxides such as tetraethyl orthotitanate.

Preferred additives are acetic acid and tetraethyl orthotitanate.

The acid is used in a molar concentration 1 time or more, and preferably 1 to 10 times, higher than that of the compound of the formula [13].

10 (R)-(+)-tert-butyl sulfinamide is used in a molar concentration 1 time or more, and preferably 1 to 10 times, higher than that of the compound of the formula [13].

The reaction of generating an imine may be carried out at a temperature from 0°C to the boiling point of a solvent, and preferably from 15 0°C to 100°C, for 1 minute to 24 hours.

[0109]

Examples of a reducing agent to be used in a reaction of reducing an imine include boron hydrides such as sodium cyanoborohydride and sodium borohydride.

20 The boron hydride is used in a molar concentration 1 time or more, and preferably 1 to 10 times, higher than that of the compound of the formula [13].

The reaction of reducing an imine can be performed at -50°C to the boiling point of a solvent and preferably at -50°C to 100°C for 1 minute to 24 hours.

25 [0110]

(A4-4)

The compound of the formula [4]-1 can be produced through desulfinylation of the compound of the formula [14] under acidic conditions.

30 The solvent used in this reaction is not particularly limited, as long as it does not affect the reaction. Examples of the solvent include aliphatic hydrocarbons, halogenated hydrocarbons, alcohols, glycols, ethers, ketones, esters, amides, nitriles, sulfoxides, aromatic hydrocarbons, and water. These substances may be used in combination.

Preferred solvents are alcohols and ethers.

35 Examples of the acid used in this reaction include: inorganic acids such as hydrochloric acid, hydrogen bromide, and sulfuric acid; sulfonic acids

such as methanesulfonic acid and p-toluenesulfonic acid; carboxylic acids such as acetic acid, citric acid, and formic acid.

Preferred acids are inorganic acids such as hydrogen halide and sulfuric acid.


The inorganic acid is used in a molar concentration 1 time or more, and preferably 1 to 5 times, higher than that of the compound of the formula [14].

This reaction may be carried out at a temperature from -50°C to the boiling point of a solvent, and preferably from 0°C to 100°C , for 1 minute to 24 hours.

[0111]

[Production Method 5]

[Formula 18]



"wherein Re^1 and Re^2 may be the same or different, an amino protecting group, $\text{R}^{\text{a}1}$, $\text{R}^{\text{a}2}$, $\text{R}^{\text{b}1}$, $\text{R}^{\text{b}2}$, La , and Lb have the same definitions as those described above."

[0112]

(A5-1)

The compound of the formula [16] can be produced by deprotecting the compound of the formula [15].

[0113]

(A5-2)

The compound of the formula [17] can be produced by allowing the compound of the formula [16] to react with the compound of the formula [5] in accordance with Production Method 2.

[0114]

(A5-3)

The compound of the formula [6]-1 can be produced by deprotecting the compound of the formula [17] with the use of hydrazine or the like and then protecting an amino group.

5 [0115]

The compounds obtained by the above-described production methods can be induced to other compounds by subjecting them to well-known reactions such as condensation, addition, oxidation, reduction, dislocation, substitution, halogenation, dehydration or hydrolysis, or by combining these
10 reactions, as appropriate.

[0116]

When amino, hydroxy and/or carboxyl groups are present in the compounds obtained by the above-described production methods and the intermediates thereof, reactions can be carried out by replacing their
15 protecting groups with other groups, as appropriate. In addition, when two or more protecting groups are present, such protecting groups can be selectively deprotected by subjecting them to well-known reactions.

[0117]

Among compounds used in the above-described production methods,
20 those that can be in the form of salts can be used as salts. Examples of such salts are the same as the examples of the salt of the compound represented by the formula (I) of the present invention.

[0118]

When isomers (for example, optical isomers, geometric isomers,
25 tautomers, etc.) are present for the compounds used in the above-described production methods, these isomers can also be used. In addition, when solvates, hydrates, and various forms of crystals are present, these solvates, hydrates, and various forms of crystals can also be used.

[0119]

30 When the compound represented by the formula [1] of the present invention is used as a medicament, pharmaceutical additives commonly used in formulation of such a medicament, such as an excipient, a carrier, and a diluent, may be mixed into the compound of the present invention, as appropriate. The thus formulated medicament can be orally or parenterally
35 administered in the form of a tablet, a capsule, a powdered medicine, a syrup, a granule, a pill, a suspending agent, an emulsion, a liquid agent, a powdery

agent, a suppository, an eye drop, a nasal drop, an ear drop, a patch, an ointment, or an injection, according to ordinary methods. An administration method, a dosage, and a number of doses can be selected, as appropriate, depending on the age, body weight, and symptoms of a patient. In general, the medicament may be administered orally or parenterally (e.g. via injection, drip infusion, or administration into a rectal site) at a dosage from 0.01 to 1000 mg/kg to an adult per day, once or dividedly several times.

[Examples]

[0120]

The present invention is hereafter described with reference to the Reference Examples and the Examples, although the scope of the present invention is not limited thereto.

LC/MS analysis was conducted under the following conditions.

LC/MS analyzer: Waters SQD

Column: Waters BEHC18 1.7 μ m, 2.1 x 30 mm

Solvent: Liquid A: 0.1% formic acid-water

Liquid B: 0.1% formic acid-acetonitrile

Gradient cycle: 0.00 min (Liquid A/Liquid B = 95/5), 2.00 min (Liquid A/Liquid B = 5/95), 3.00 min (Liquid A/Liquid B = 5/95), 3.01 min (Liquid A/Liquid B = 100/0), 3.80 min (Liquid A/Liquid B = 100/0)

Flow rate: 0.5 mL/min (The column temperature was room temperature, and no temperature control was carried out.)

Ionization method: Electron Spray Ionization method (ESI positive and negative ion peaks were detected.)

UV detection: UV 254 nm


NMR spectra used herein are proton NMR spectra. NMR spectra were measured using a BRUKER AVANCE 300 (300 MHz spectrometer), and the δ value was expressed in ppm.

The carrier used for silica gel column chromatography is PSQ100B (spherical shape) (Fuji Silysia Chemical Ltd.), and the PLC glass plate used herein is a PLC glass plate silica gel 60 F₂₅₄ (Merck), unless otherwise specified.

The compound of the formula [1a] is a mixture of the compound of the formula [1b] and the compound of the formula [1c].

[0121]

[Formula 19]



[0122]


Abbreviations used in the Reference Examples and the Examples stand for the terms given below.

- 5 Boc: tert-butoxycarbonyl
 Bn: benzyl
 CDI: carbonyldiimidazole
 Cbz: benzyloxycarbonyl
 CHCl₃: chloroform
 10 CH₂Cl₂: dichloromethane
 dba: 1,3-dibenzylideneacetone
 DIAD: diisopropyl azodicarboxylate
 DIPEA: N,N-diisopropyl ethylamine
 DMAc: N,N-dimethylacetamide
 15 DMAP: N,N-dimethylaminopyridine
 DMF: N,N-dimethylformamide
 DMSO: dimethyl sulfoxide
 DMSO-d₆: deuterated dimethyl sulfoxide
 DPPA: diphenylphosphoryl azide
 20 Et: ethyl
 IPE: diisopropylether
 mCPBA: meta-chloroperoxybenzoic acid
 Me: methyl
 Ms: methanesulfonyl
 25 Ph: phenyl
 RT, rt: retention time
 TBAI: tetrabutylammonium iodide
 Tf: trifluoromethanesulfonyl
 TFA: trifluoroacetic acid
 30 THF: tetrahydrofuran
 Py: pyridine
 Xantphos: 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene

[0123]

Reference Example 1

[Formula 20]



[0124]

5 1st step

Hydrazine monohydrate (4.87 ml) was added to an EtOH (19 ml) solution containing 2-chloro-5-nitrobenzonitrile (1.83 g), followed by stirring for 0.5 hour under ice cooling. Water was added to the reaction solution, and a solid precipitate was collected by filtration and washed with IPE and ethyl acetate. A red solid of 5-nitro-1H-indazol-3-amine (1.45 g) was thus obtained.

MS (ESI m/z): 179(M+H)

RT (min): 0.77

[0125]

15 2nd step

5-nitro-1H-indazol-3-amine (254 mg) obtained in the 1st step and iodomethane (1 ml) were added to a DMF (3ml) suspension containing sodium hydride (60% in oil) (171 mg) under ice cooling, followed by stirring at room temperature for 1 hour. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with water and saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure, and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:1). A yellow solid of N,N,1-trimethyl-5-nitro-1H-indazol-3-amine (132 mg) and a yellow solid of N,N,2-trimethyl-5-nitro-2H-indazol-3-amine (72 mg) were thus obtained.

N,N,1-trimethyl-5-nitro-1H-indazol-3-amine

MS (ESI m/z): 221 (M+H)

RT (min): 1.24

30 N,N,2-trimethyl-5-nitro-2H-indazol-3-amine

MS (ESI m/z): 221 (M+H)

RT (min): 1.14

[0126]

3rd step

An MeOH (10 ml) solution containing N,N,1-trimethyl-5-nitro-1H-indazol-3-amine (132 mg) obtained in the 2nd step was prepared and subjected to a hydrogenation reaction (70°C; 50 bar; flow rate: 2 ml/min; 10%Pd/C) using H-cube™. The solvent was distilled
 5 away under reduced pressure. A red solid of N³,N³,1-trimethyl-1H-indazol-3,5-diamine (100 mg) was thus obtained.


MS (ESI m/z): 191 (M+H)

RT (min): 0.52

[0127]

10 Reference Example 2

[Formula 21]



[0128]

1st step

15 5-Nitro-1H-indazol-3-ol (112 mg) and iodomethane (0.5 ml) was added to a DMF (2 ml) suspension containing sodium hydride (60% in oil) (60 mg), followed by stirring at room temperature for 10 minutes. Water was added to the reaction solution, followed by extraction with chloroform. The organic layers were dried over anhydrous sodium sulfate. The solvent was distilled
 20 away under reduced pressure, and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:1). A yellow solid of 3-methoxy-1-methyl-5-nitro-1H-indazole (31 mg) was thus obtained.

3-Methoxy-1-methyl-5-nitro-1H-indazole

MS (ESI m/z): 208 (M+H)

25 RT (min): 1.33

[0129]

2nd step

The following compound was obtained as described in the 3rd step in Reference Example 1.

30 3-Methoxy-1-methyl-1H-indazol-5-amine


MS (ESI m/z): 178 (M+H)

RT (min): 0.44

[0130]

Reference Example 3

[Formula 22]



[0131]

1st step

5 3-Methoxy-5-nitro-1H-indazole (97 mg), 1-bromo-2-methoxyethane (70 μ l), and TBAI (2 mg) were added to a DMF (1 ml) suspension containing sodium hydride (60% in oil) (24 mg) under ice cooling, followed by stirring at 100°C for 1 hour. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with water and saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure, and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:1). A 3-methoxy-1-(2-methoxyethyl)-5-nitro-1H-indazole (50 mg) was thus obtained.

15 MS (ESI m/z): 252 (M+H)

RT (min): 1.40

[0132]

2nd step

 The following compound was obtained as described in the 3rd step in
20 Reference Example 1.


3-Methoxy-1-(2-methoxyethyl)-1H-indazol-5-amine

[0133]

Reference Example 4

 The following compound was obtained as described in Reference
25 Example 3.

[Formula 23]



1-(2-Methoxyethyl)-3-methyl-1H-indazol-5-amine


MS (ESI m/z): 206 (M+H)

30 RT (min): 0.79

[0134]

Reference Example 5

[Formula 24]



[0135]

5 1st step

A CH_2Cl_2 (10 ml) solution containing 2,2-difluoroethanol (5.0 g) and triethylamine (8.44 ml) was slowly added to a CH_2Cl_2 (10 ml) solution containing trifluoromethanesulfonic anhydride (10.2ml) at -78°C in a nitrogen atmosphere, followed by stirring for 45 minutes. The solvent was
 10 distilled away under reduced pressure. Colorless oily matter of 2,2-difluoroethyl trifluoromethane sulfonate (9.04 g) was thus obtained.

[0136]

2nd step

2,2-Difluoroethyl trifluoromethane sulfonate (2 ml) obtained in the 1st
 15 step and 5-nitroindazole (163 mg) were added to a DMF (2 ml) suspension containing sodium hydride (60% in oil) (44 mg) under ice cooling, followed by stirring at room temperature for 1 hour. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with water and saturated saline and dried over anhydrous sodium
 20 sulfate. The solvent was distilled away under reduced pressure, and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:1). 1-(2,2-Difluoroethyl)-5-nitro-1H-indazole (113 mg) was thus obtained.

MS (ESI m/z): 228 (M+H)

25 RT (min): 1.25

[0137]

3rd step

The following compound was obtained as described in the 3rd step in Reference Example 1.

30 1-(2,2-Difluoroethyl)-1H-indazol-5-amine


MS (ESI m/z): 228 (M+H)

RT (min): 1.18

[0138]

Reference Example 6

[Formula 25]



The following compound was obtained as described in Reference

5 Example 5.

[0139]

1st step

1-(2,2-Difluoroethyl)-4-nitro-1H-indazole

[0140]

10 2nd step

1-(2,2-Difluoroethyl)-1H-indazol-4-amine


MS (ESI m/z): 198 (M+H)

RT (min): 0.88

[0141]

15 Reference Example 7

[Formula 26]



The following compound was obtained as described in Reference
Example 5.

20 [0142]

1st step

1-(2,2-Difluoroethyl)-3-methoxy-5-nitro-1H-indazole

MS (ESI m/z): 258 (M+H)

RT (min): 1.40

25 [0143]


2nd step

1-(2,2-Difluoroethyl)-3-methoxy-1H-indazol-5-amine

[0144]

Reference Example 8

[Formula 27]



[0145]

- 5 The following compound was obtained as described in Reference Example 5.

1st step

1-(2,2-Difluoroethyl)-3-methyl-5-nitro-1H-indazole

[0146]

- 10 2nd step

1-(2,2-Difluoroethyl)-3-methyl-1H-indazol-5-amine


MS (ESI m/z): 212 ($M+H$)

RT (min): 0.49

[0147]

- 15 Reference Example 9

[Formula 28]



[0148]

The following compound was obtained as described in the 3rd step in

- 20 Reference Example 1.

 $N^3,N^3,1$ -trimethyl-2H-indazol-3,5-diamineMS (ESI m/z): 191 ($M+H$)


RT (min): 0.47

[0149]

- 25 Reference Example 10

The following compound was obtained with reference to WO2007/126841 A2.

[Formula 29]




1-Ethyl-4-nitro-1H-indazole

[0150]

Reference Example 11

5 The following compound was obtained as described in the 3rd step in Reference Example 1.

[Formula 30]



10 1-Ethyl-1H-indazol-4-amine

MS (ESI m/z): 162 (M+H)


RT (min): 0.92

[0151]

Reference Example 12

15 The following compound was obtained with reference to US2009/76275 A1.

[Formula 31]




1-(2-Methoxyethyl)-5-nitro-1H-indazole

20 [0152]

Reference Example 13

[Formula 32]



25 The following compound was obtained as described in the 3rd step in Reference Example 1.

1-(2-Methoxyethyl)-1H-indazol-5-amine


MS (ESI m/z): 192 (M+H)

RT (min): 0.39

[0153]

Reference Example 14

5 [Formula 33]



[0154]

1st step

Hydrazine monohydrate (19 ml) was added to an EtOH (15 ml) solution containing methyl 2-chloro-5-nitrobenzoate (10 g), followed by stirring at 90°C for 1 hour. The reaction solution was adjusted to room temperature. Water and concentrated hydrochloric acid (32ml) were added dropwise to the reaction solution. A solid precipitate was collected by filtration and washed with water. A brown solid of 5-nitro-1H-indazol-3-ol (5.42 g) was thus obtained.

MS (ESI m/z): 180 (M+H)

RT (min): 0.73

[0155]

2nd step

Ethyl chloroformate (5 ml) was added to a pyridine (30 ml) solution containing 5-nitro-1H-indazol-3-ol (5.42 g) obtained in the 1st step, followed by stirring at room temperature for 1.5 hours. Water and concentrated hydrochloric acid (32 ml) were added dropwise to the reaction solution and a solid precipitate was collected by filtration. The obtained residue was washed with water. A brown solid of ethyl 3-hydroxy-5-nitro-1H-indazole-1-carboxylate (7.5 g) was thus obtained.

MS (ESI m/z): 252 (M+H)

RT (min): 1.17

[0156]

3rd step

Iodomethane (10 ml) and cesium carbonate (4.89 g) were added to an acetone (20 ml) solution containing ethyl 3-hydroxy-5-nitro-1H-indazol-1-carboxylate (2.51 g) obtained in the 2nd step under ice cooling in a nitrogen atmosphere, followed by stirring at 80°C for 0.5 hours. An insoluble precipitate was removed by filtration and the solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 4:1). A white solid of ethyl 3-methoxy-5-nitro-1H-indazol-1-carboxylate (1.24 g) was thus obtained.

MS (ESI m/z): 266 (M+H)

RT (min): 1.45

[0157]

4th step

Potassium hydroxide (0.6 g) was added to an EtOH (20 ml) solution containing ethyl 3-methoxy-5-nitro-1H-indazol-1-carboxylate (1.24 g) obtained in the 3rd step, followed by stirring at room temperature for 0.5 hours. Water and concentrated hydrochloric acid (1 ml) were added dropwise to the reaction solution. A solid precipitate was collected by filtration and washed with water. A light yellow solid of 3-methoxy-5-nitro-1H-indazole (636 mg) was thus obtained.

MS (ESI m/z): 194 (M+H)

RT (min): 1.15

[0158]

5th step

3-methoxy-5-nitro-1H-indazole (100 mg) and iodoethane (0.1 ml) were added to a DMF (1 ml) suspension containing sodium hydride (60% in oil) (23 mg), followed by stirring at room temperature for 0.5 hours. Water was added to the reaction solution. A solid precipitate was collected by filtration and purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:1). 1-Ethyl-3-methoxy-5-nitro-1H-indazole (80 mg) was thus obtained.

MS (ESI m/z): 222 (M+H)

RT (min): 1.45

[0159]

6th step

The following compound was obtained as described in the 3rd step in

Reference Example 1.


1-Ethyl-3-methoxy-1H-indazol-5-amine

[0160]

Reference Example 15

5 The following compound was obtained with reference to Journal of Heterocyclic Chemistry, 1979, vol. 16, pp. 1599, 1600, and 1601.

[Formula 34]




7-Chloro-1-methyl-5-nitro-1H-indazole

10 [0161]

Reference Example 16

[Formula 35]



15 Tin (II) chloride (50 mg) was added to a ethanol (2 ml) solution containing 7-chloro-1-methyl-5-nitro-1H-indazole (30 mg), followed by stirring at 100°C for 0.5 hours. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 0:1). 7-Chloro-1-methyl-1H-indazol-5-amine (10 mg) was thus obtained.

20 MS (ESI m/z): 182 (M+H)


RT (min): 0.64

[0162]

Reference Example 17

25 The following compound was obtained with reference to EP1150962 B1, 2004.

[Formula 36]




3-Methoxybenz[d]isoxazol-5-amine

[0163]

30 Reference Example 18

[Formula 37]



[0164]

1st step

5 Sodium nitrate (1.7 g) was added to a concentrated sulfuric acid (7 ml) solution containing 2,3-difluorobenzoic acid (1.58 g), followed by stirring at room temperature for 0.5 hours. The reaction solution was poured into ice water, followed by extraction with ethyl acetate. The organic layers were dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure, the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 4:1). A light yellow solid of 2,3-difluoro-5-nitrobenzoic acid (1.61 g) was thus obtained.

MS (ESI m/z): 202 (M+H)

RT (min): 0.97

15 [0165]

2nd step

20 Oxalyl chloride (1 ml) and DMF (5 μ l) were added to a CH_2Cl_2 (1.6 ml) solution containing 2,3-difluoro-5-nitrobenzoic acid (1.61 g) obtained in the 1st step, followed by stirring at room temperature for 15 minutes. The reaction solution was poured into a liquid mixture of MeOH/Py (100 ml/1.28 ml) and the solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 4:1). A light yellow solid of methyl 2,3-difluoro-5-nitrobenzoate (1.71 g) was thus obtained.

25 [0166]

3rd step

Hydrazine monohydrate (1.91 ml) was added to an EtOH (40 ml) solution containing methyl 2,3-difluoro-5-nitrobenzoate (1.71 g) obtained in the 2nd step, followed by stirring at room temperature for 10 minutes. An

insoluble precipitate was removed by filtration and washed with EtOH.
7-Fluoro-5-nitro-1H-indazol-3-ol (956 mg) was thus obtained.

MS (ESI m/z): 198 (M+H)

RT (min): 0.90

5 [0167]

4th step

The following compound was obtained as described in the 1st step in
Reference Example 2.

7-Fluoro-3-methoxy-1-methyl-5-nitro-1H-indazole

10 MS (ESI m/z): 226 (M+H)

RT (min): 0.92

[0168]

5th step

The following compound was obtained as described in the 3rd step in
15 Reference Example 1.

7-Fluoro-3-methoxy-1-methyl-1H-indazol-5-amine


MS (ESI m/z): 196 (M+H)

RT (min): 0.61

[0169]

20 Reference Example 19

[Formula 38]



[0170]

1st step

25 Sodium chlorodifluoroacetate (4.75 g) and potassium carbonate (8.58 g) were added to a DMF (3 ml) solution containing ethyl 3-hydroxy-5-nitro-1H-indazol-1-carboxylate (1.56 g), followed by stirring at 80°C for 1 hour. The reaction solution was adjusted to room temperature and

ethyl acetate was added to remove an insoluble precipitate. The organic layers were washed with a saturated ammonium chloride aqueous solution, water, and saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure, and the obtained
5 residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 4:1). A yellow solid of ethyl 3-(difluoromethoxy)-5-nitro-1H-indazol-1-carboxylate (1.14 g) was thus obtained.

MS (ESI m/z): 302 (M+H)

10 RT (min): 1.53

[0171]

2nd step

Water (6 ml) and lithium hydroxide monohydrate (640 mg) were added to a THF (19 ml) solution containing ethyl
15 3-(difluoromethoxy)-5-nitro-1H-indazol-1-carboxylate (1.14 g) obtained in the 1st step, followed by reflux at 80°C for 3 hours. THF was distilled away under reduced pressure and a saturated ammonium chloride aqueous solution was added. An insoluble precipitate was removed by filtration. The obtained residue was washed with water, dissolved in ethyl acetate, and dried
20 over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. A yellow solid of 3-(difluoromethoxy)-5-nitro-1H-indazole (921 mg) was thus obtained.

MS (ESI m/z): 230 (M+H)

RT (min): 1.04

25 [0172]

3rd step

The following compound was obtained as described in the 2nd step in Reference Example 1.

3-(Difluoromethoxy)-1-methyl-5-nitro-1H-indazole

30 MS (ESI m/z): 244 (M+H)

RT (min): 1.51

[0173]

4th step

The following compound was obtained as described in the 3rd step in
35 Reference Example 1.

3-(Difluoromethoxy)-1-methyl-1H-indazol-5-amine


MS (ESI m/z): 214 (M+H)

RT (min): 0.59

[0174]

Reference Example 20

5 [Formula 39]



[0175]

The following compound was obtained as described in Reference Example 3.

10 1st step

3-(Difluoromethoxy)-1-(2-methoxyethyl)-5-nitro-1H-indazole

MS (ESI m/z): 288 (M+H)

RT (min): 1.55

[0176]

15 2nd step

3-(Difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-amine

MS (ESI m/z): 258 (M+H)


RT (min): 0.68

[0177]

20 Reference Example 21

The following compound was obtained with reference to WO2010/114971 A1.

[Formula 40]




25 2,3-Difluoro-5-nitrobenzonitrile

[0178]

Reference Example 22

[Formula 41]



[0179]

The following compound was obtained as described in Reference Example 1.

5 1st step

7-Fluoro-5-nitro-1H-indazol-3-amine

MS (ESI m/z): 197 (M+H)

RT (min): 0.93

[0180]

10 2nd step

7-Fluoro-N,N,1-trimethyl-5-nitro-1H-indazol-3-amine

MS (ESI m/z): 239 (M+H)

RT (min): 1.66

[0181]

15 3rd step

7-Fluoro-N³,N³,1-trimethyl-1H-indazol-3,5-diamine


MS (ESI m/z): 209 (M+H)

RT (min): 0.70

[0182]

20 Reference Example 23

[Formula 42]



1st step

Select flour (173 mg) and acetic acid (2.5 ml) were added to an acetonitrile (2.5 ml) solution containing 5-nitroindazole (615 mg), followed by microwave irradiation (Initiator TM, 150°C, 0.5 hours, 2.45 GHz, 0-240 W). The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:1). 3-Fluoro-5-nitro-1H-indazole (404 mg) was thus obtained.

30 [0183]

2nd step

Methyl iodide (41 μ l) and potassium carbonate (114 mg) were added to a 1,4-dioxane (2.5 ml) solution containing 3-fluoro-5-nitro-1H-indazole (100 mg), followed by stirring at 100°C for 2 hours. Ethyl acetate was added to the reaction solution. An insoluble precipitate was removed by filtration, the solvent was distilled away under reduced pressure, and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:1). 3-Fluoro-1-methyl-5-nitro-1H-indazole was thus obtained.

[0184]

3rd step

The following compound was obtained as described in the 3rd step in Reference Example 1.

3-Fluoro-1-methyl-1H-indazol-5-amine


MS (ESI m/z): 166 (M+H)

RT (min): 1.32

[0185]

Reference Example 24

[Formula 43]



[0186]

1st step

Potassium carbonate (200 mg) and 2-chloroethylmethyl ether (0.1 ml) were added to a DMF (1.5 ml) solution containing 4-nitro-1H-indazole (80 mg), followed by stirring at 60°C for 4 hours. An insoluble precipitate was collected by filtration and washed with ethyl acetate. A mixture of 1-(2-methoxyethyl)-4-nitro-1H-indazole and 2-(2-methoxyethyl)-4-nitro-2H-indazole was thus obtained.

1-(2-Methoxyethyl)-4-nitro-1H-indazole

MS (ESI m/z): 222 (M+H)

RT (min): 1.19

2-(2-Methoxyethyl)-4-nitro-2H-indazole

MS (ESI m/z): 222 (M+H)

RT (min): 1.12

[0187]

2nd step

Iron powder (170 mg), ammonium chloride (160 mg), and water (3 ml) were added to an EtOH (10 ml) solution containing the mixture of 1-(2-methoxyethyl)-4-nitro-1H-indazole and 2-(2-methoxyethyl)-4-nitro-2H-indazole obtained in the 1st step, followed by stirring at 80°C for 2 hours. Ethyl acetate was added to the reaction solution, insoluble matter was removed by filtration, and the solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel-alumina column chromatography.

1-(2-methoxyethyl)-1H-indazol-4-amine (49 mg) and 2-(2-methoxyethyl)-2H-indazol-4-amine (40 mg) were thus obtained.

1-(2-Methoxyethyl)-1H-indazol-4-amine

MS (ESI m/z): 192 (M+H)

RT (min): 0.72

2-(2-Methoxyethyl)-2H-indazol-4-amine


MS (ESI m/z): 192 (M+H)

RT (min): 0.53

[0188]

Reference Example 25

[Formula 44]



[0189]

1st step

An EtOH (12.5 ml) solution containing mucobromic acid (12.9 g) was added dropwise to a solution comprising sodium nitrate (13.1 g) and water (12.5 ml) at 50°C, followed by stirring for 0.5 hours. The reaction solution was adjusted to room temperature, a solid precipitate was collected by filtration. The obtained residue was washed with EtOH. A yellow solid of sodium (1,3-dioxopropan-2-ylidene)azinate (3.82 g) was thus obtained.

MS (ESI m/z): 116 (M-H)

RT (min): 0.31

[0190]

2nd step

Sodium (1,3-dioxopropan-2-ylidene)azinate (864 mg) obtained in the

1st step was added to an acetic acid (5.5 ml) solution containing 3-amino-5-hydroxypyrazole (495 mg), followed by stirring in a sealed tube at 90°C for 6 hours. The reaction solution was adjusted to room temperature and poured into water. A solid precipitate was collected by filtration.

5 5-Nitro-1H-pyrazolo[3,4-b]pyridin-3-ol (691 mg) was thus obtained.

MS (ESI m/z): 181 (M+H)

RT (min): 0.58

[0191]

3rd step

10 The following compound was obtained as described in Reference Example 2.

3-Methoxy-1-methyl-5-nitro-1H-pyrazolo[3,4-b]pyridine

MS (ESI m/z): 209 (M+H)

RT (min): 1.15

15 [0192]

4th step

The following compound was obtained as described in the 3rd step in Reference Example 1.

3-Methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-amine


20 MS (ESI m/z): 179 (M+H)

RT (min): 0.49

[0193]

Reference Example 26

[Formula 45]



[0194]

1st step

30 Sodium (1,3-dioxopropan-2-ylidene)azinate monohydrate (1.37 g) was added to an acetic acid (14 ml) solution containing pyrazole 3,5-diamine (2.41 g), followed by stirring in a sealed tube at 90°C for 6 hours. The reaction solution was adjusted to room temperature and poured into a saturated sodium hydrogen carbonate aqueous solution. A solid precipitate was collected by filtration. A liquid mixture of ethyl acetate/MeOH/THF (2/1/0.1) was added

to the obtained solid, insoluble matter was removed, and the organic layers were dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure, an orange-colored solid of 5-nitro-1H-pyrazolo[3,4-b]pyridin-3-amine (0.91 g) was thus obtained.

5 MS (ESI m/z): 180 (M+H)

RT (min): 0.55

[0195]

2nd step

The following compound was obtained as described in the 2nd step in

10 Reference Example 1.

N,N,1-trimethyl-5-nitro-1H-pyrazolo[3,4-b]pyridin-3-amine

MS (ESI m/z): 222 (M+H)

RT (min): 1.17

[0196]

15 3rd step

The following compound was obtained as described in the 3rd step in Reference Example 1.

N³,N³,1-trimethyl-1H-pyrazolo[3,4-b]pyridin-3,5-diamine


MS (ESI m/z): 192 (M+H)

20 RT (min): 0.49

[0197]

Reference Example 27

[Formula 46]



25 [0198]

1st step

Triethylamine (267 μ l), tert-butanol (230 μ l), and DPPA (413 μ l) were added to a toluene (5 ml) solution containing 5-fluoro-6-methoxynicotinic acid (275 mg), followed by reflux for 3 hours. The reaction solution was adjusted to room temperature and water was added, followed by extraction with ethyl acetate. Next, the organic layers were washed with saturated saline and dried over anhydrous sodium sulfate and the solvent was distilled away under reduced pressure. Then, the obtained residue was purified by

30

silica gel chromatography (n-hexane : ethyl acetate = 10:1 to 3:1). Colorless oily matter of tert-butyl (5-fluoro-6-methoxypyridin-3-yl)carbamate (279 mg) was thus obtained.

MS (ESI m/z): 243 (M+H)

5 RT (min): 1.46

[0199]

2nd step

TFA (2 ml) was added to tert-butyl (5-fluoro-6-methoxypyridin-3-yl)carbamate (279 mg) obtained in the 1st step, followed by stirring at room temperature for 1 hour. The solvent was distilled away under reduced pressure and a 5M sodium hydroxide aqueous solution was added to the obtained residue at 0°C so as to alkalify the mixture, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate and the solvent
15 was distilled away under reduced pressure. A light brown solid of 5-fluoro-6-methoxypyridin-3-amine (19 mg) was thus obtained.

MS (ESI m/z): 143 (M+H)


RT (min): 0.56

[0200]

20 Reference Example 28

The following compound was obtained with reference to EP1932845 A1, 2008 and EP2070929 A1, 2009.

[Formula 47]




25 4-Bromo-1H-pyrazolo[3,4-c]pyridine

[0201]

Reference Example 29

[Formula 48]



30 [0202]

1st step

The following compound was obtained as described in the 5th step in Reference Example 14.

4-Bromo-1-ethyl-1H-pyrazolo[3,4-c]pyridine

5 MS (ESI m/z): 226 (M+H)

RT (min): 1.12

[0203]

2nd step

tert-Butyl carbamate (90 mg), cesium carbonate (500 mg), Pd₂(dba)₃ (46 mg), and Xantphos (58 mg) were added to a dioxane solution (2 ml) containing 4-bromo-1-ethyl-1H-pyrazolo[3,4-c]pyridine (115 mg) obtained in the 1st step, followed by microwave irradiation (InitiatorTM, 130°C, 1 hour, 2.45 GHz, 0-240 W). Ethyl acetate was added to the reaction solution, insoluble matter was removed by filtration, and the solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 3:1). A white solid of tert-butyl (1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)carbamate (113 mg) was thus obtained.

MS (ESI m/z): 263 (M+H)

20 RT (min): 0.84

[0204]

3rd step

The following compound was obtained as described in the 2nd step in Reference Example 27.

25 1-Ethyl-1H-pyrazolo[3,4-c]pyridin-4-amine


MS (ESI m/z): 163 (M+H)

RT (min): 0.42

[0205]

Reference Example 30

30 [Formula 49]



[0206]

1st step

The following compound was obtained as described in the 1st step in Reference Example 3.

4-Bromo-1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridine

MS (ESI m/z): 256 (M+H)

5 RT (min): 1.03

[0207]

2nd step

The following compound was obtained as described in the 2nd step in Reference Example 29.

10 tert-Butyl (1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)carbamate

MS (ESI m/z): 293 (M+H)

RT (min): 0.87

[0208]

3rd step

15 The following compound was obtained as described in the 2nd step in Reference Example 27.


1-(2-Methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-amine

[0209]

Reference Example 31

20 The following compound was obtained with reference to WO2008/110863 A1, 2008 and WO2010/127855 A1, 2010.

[Formula 50]




6-Fluoro-4-nitro-1H-indazole

25 [0210]

Reference Example 32

[Formula 51]



[0211]

30 1st step

The following compound was obtained as described in the 5th step in Reference Example 14.

1-Ethyl-6-fluoro-4-nitro-1H-indazole

MS (ESI m/z): 210 (M+H)

5 RT (min): 1.35

[0212]

2nd step


The following compound was obtained as described in the 3rd step in Reference Example 1.

10 1-Ethyl-6-fluoro-1H-indazol-4-amine

[0213]

Reference Example 33

[Formula 52]



15 [0214]

1st step

The following compound was obtained as described in the 1st step in Reference Example 3.

6-Fluoro-1-(2-methoxyethyl)-4-nitro-1H-indazole

20 MS (ESI m/z): 240 (M+H)

RT (min): 1.31

[0215]

2nd step

The following compound was obtained as described in the 3rd step in


25 Reference Example 1.

6-Fluoro-1-(2-methoxyethyl)-1H-indazol-4-amine

[0216]

Reference Example 34

[Formula 53]



[0217]

1st step

The following compound was obtained as described in the 2nd step in

5 Reference Example 5.

4-Bromo-1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridine

MS (ESI m/z): 264 (M+H)

RT (min): 1.12

[0218]

10 The following compound was obtained as described in the 2nd and 3rd steps in Reference Example 29.

2nd step

tert-Butyl (1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)carbamate

MS (ESI m/z): 299 (M+H)

15 RT (min): 0.96

[0219]


3rd step

1-(2,2-Difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-amine

[0220]

20 Reference Example 35

[Formula 54]



[0221]

1st step

25 mCPBA (270 mg) was added to a CHCl_3 (5 ml) suspension containing

4-bromo-1H-pyrazolo[3,4-c]pyridine (197 mg), followed by stirring at room temperature for 10 minutes. IPE (10ml) was added to the reaction solution and a solid precipitate was collected by filtration. A white solid of 4-bromo-1H-pyrazolo[3,4-c]pyridine 6-oxide (148 mg) was thus obtained.

5 MS (ESI m/z): 214 (M+H)

RT (min): 0.56

[0222]

2nd step

10 Phosphorus oxychloride (2 ml) was added to 4-bromo-1H-pyrazolo[3,4-c]pyridine 6-oxide (148 mg) obtained in the 1st step, followed by stirring at room temperature for 3 hours. The reaction solution was poured into water, followed by extraction with ethyl acetate. An insoluble precipitate was removed and then the solvent was distilled away under reduced pressure. A yellow solid of 15 4-bromo-7-chloro-1H-pyrazolo[3,4-c]pyridine (126 mg) was thus obtained.

MS (ESI m/z): 232 (M+H)

RT (min): 1.09

[0223]

3rd step

20 Sodium methoxide (28% in MeOH) (3 ml) was added to 4-bromo-7-chloro-1H-pyrazolo[3,4-c]pyridine (126 mg) obtained in the 2nd step, followed by stirring at 90°C for 5 hours. A saturated ammonium chloride aqueous solution was added to the reaction solution and an insoluble precipitate was collected by filtration. The obtained residue was washed 25 with water and a yellow solid of 4-bromo-7-methoxy-1H-pyrazolo[3,4-c]pyridine (133 mg) was thus obtained.

MS (ESI m/z): 228 (M+H)

RT (min): 1.12

The following compound was obtained as described in Reference 30 Example 29.

[0224]

4th step

4-Bromo-7-methoxy-1-methyl-1H-pyrazolo[3,4-c]pyridine

MS (ESI m/z): 242 (M+H)

35 RT (min): 1.41

[0225]

The following compound was obtained as described in the 2nd and 3rd steps in Reference Example 29.

5th step

tert-Butyl (7-methoxy-1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)carbamate

5 MS (ESI m/z): 279 (M+H)

RT (min): 1.30

[0226]

6th step

7-Methoxy-1-methyl-1H-pyrazolo[3,4-c]pyridin-4-amine


10 MS (ESI m/z): 179 (M+H)

RT (min): 0.58

[0227]

Reference Example 36

[Formula 55]



15

[0228]

1st step

Potassium hydroxide (6.45 g) and iodine (15.6 g) were added to a DMF (60 ml) solution containing 5-nitroindazole (5 g), followed by stirring at 65°C for 1 hour. The reaction solution was poured into a saturated sodium hydrogen carbonate aqueous solution and a solid precipitate was collected by filtration. 3-Iodo-5-nitro-1H-indazole (8 g) was thus obtained.

20

[0229]

2nd step

The following compound was obtained as described in the 1st step in Reference Example 2.

25

3-Iodo-1-methyl-5-nitro-1H-indazole

MS (ESI m/z): 304 (M+H)

RT (min): 1.57

[0230]

3rd step

Pyrrolidine (90 mg), cesium carbonate (500 mg), $\text{Pd}_2(\text{dba})_3$ (46 mg),
 5 and Xantphos (58 mg) were added to a dioxane solution (4 ml) containing
 3-iodo-1-methyl-5-nitro-1H-indazole (303mg) obtained in the 1st step,
 followed by microwave irradiation (InitiatorTM, 160°C, 15 minutes, 2.45 GHz,
 0-240W). Then, ethyl acetate was added to the obtained residue, insoluble
 matter was removed by filtration, and the solvent was distilled away under
 10 reduced pressure. 1-Methyl-5-nitro-3-(pyrrolidine-1-yl)-1H-indazole (50
 mg) was thus obtained.

MS (ESI m/z): 247 (M+H)

RT (min): 1.37

[0231]

15 4th step

The following compound was obtained as described in the 3rd step in
 Reference Example 1.


1-Methyl-3-(pyrrolidine-1-yl)-1H-indazol-5-amine

[0232]

20 Reference Example 37

The following compound was obtained as described in the 3rd and 4th
 steps in Reference Example 36.

[Formula 56]



25 [0233]

1st step

1-(1-Methyl-5-nitro-1H-indazol-3-yl)pyrrolidin-2-one

MS (ESI m/z): 261 (M+H)

RT (min): 1.13

30 [0234]

2nd step

1-(5-Amino-1-methyl-1H-indazol-3-yl)pyrrolidin-2-one


MS (ESI m/z): 231 (M+H)

RT (min): 0.52

[0235]

Reference Example 38

5 [Formula 57]



[0236]

1st step

Di-tert-butyl carbonate (1.76 g), triethylamine (1.13 ml), and DMAP
 10 (10 mg) were added to a THF (8.2 ml) solution containing
 5-nitro-1H-indazol-3-amine (1.45 g) under ice cooling, followed by stirring
 for 20 minutes. A saturated ammonium chloride aqueous solution was added
 to the reaction solution, followed by extraction with ethyl acetate. The
 organic layers were extracted with water and saturated saline and dried over
 15 sodium sulfate and the solvent was distilled away under reduced pressure.
 The obtained residue was purified by silica gel chromatography (hexane :
 ethyl acetate = 1:0 to 3:2) and a yellow solid of tert-butyl
 (5-nitro-1H-indazol-3-yl)carbamate (1.8 g) was thus obtained.

[0237]

20 2nd step

Sodium hydride (60% in oil) (250 mg) and iodomethane (458 ml) were
 added to a DMF (5ml) solution containing tert-butyl
 (5-nitro-1H-indazol-3-yl)carbamate (680 mg) obtained in the 1st step under
 ice cooling, followed by stirring for 5 hours. Water was added to the
 25 reaction solution, followed by extraction with ethyl acetate and washing with
 water and saturated saline. The organic layers were dried over anhydrous
 sodium sulfate. Then, the solvent was distilled away under reduced pressure
 and the obtained residue was purified by silica gel chromatography
 (n-hexane : ethyl acetate = 1:0 to 5:1) and directly used in the subsequent
 30 reaction.

[0238]

3rd step

The following compound was obtained as described in the 2nd step in Reference Example 27.

5 1-Methyl-5-nitro-1H-indazol-3-amine

N,1-dimethyl-5-nitro-1H-indazol-3-amine

[0239]

4th step

The following compound was obtained as described in the 1st step in Reference Example 3.

N-(2-methoxyethyl)-1-methyl-5-nitro-1H-indazol-3-amine

MS (ESI m/z): 251 (M+H)

RT (min): 1.11

N-(2-methoxyethyl)-N,1-dimethyl-5-nitro-1H-indazol-3-amine

15 MS (ESI m/z): 265 (M+H)


RT (min): 1.29

[0240]


Reference Example 39

The following compound was obtained as described in the 3rd step in Reference Example 1.

[Formula 58]

N³-(2-methoxyethyl)-1-methyl-1H-indazol-3,5-diamine

[Formula 59]




25

N³-(2-methoxyethyl)-N,1-dimethyl-1H-indazol-3,5-diamine

[0241]

Reference Example 40

[Formula 60]



[0242]

1st step

di-tert-Butyl carbonate (1.76 g), triethylamine (1.13 ml), and DMAP
 5 (10 mg) were added to a THF (8.2 ml) solution containing
 5-nitro-1H-indazol-3-amine (1.45 g), followed by stirring for 20 minutes
 under ice cooling. A saturated ammonium chloride aqueous solution was
 added to the reaction solution, followed by extraction with ethyl acetate and
 washing with water and saturated saline. The obtained solution was dried
 10 over sodium sulfate and the solvent was distilled away under reduced pressure.
 The obtained residue was purified by silica gel chromatography (hexane :
 ethyl acetate = 1:0 to 3:2) and a yellow solid mixture (1.8 g) of tert-butyl
 3-((tert-butoxycarbonyl)amino)-5-nitro-1H-indazol-1-carboxylate, tert-butyl
 (5-nitro-1H-indazol-3-yl)carbamate, and tert-butyl
 15 3-amino-5-nitro-1H-indazol-1-carboxylate was thus obtained.

[0243]

2nd step

The following compound was obtained as described in the 2nd step in
 Reference Example 1.

20 tert-Butyl

3-((tert-butoxycarbonyl)(methyl)amino)-5-nitro-1H-indazol-1-carboxylate

MS (ESI m/z): 393 (M+H)

RT (min): 1.96

tert-Butyl methyl (1-methyl-5-nitro-1H-indazol-3-yl)carbamate

MS (ESI m/z): 307 (M+H)

RT (min): 1.59

tert-Butyl 3-(dimethylamino)-5-nitro-1H-indazol-1-carboxylate

5 MS (ESI m/z): 307 (M+H)

RT (min): 1.68

[0244]

3rd step

The following compound was obtained as described in the 3rd step in

10 Reference Example 1.

tert-Butyl

5-amino-3-((tert-butoxycarbonyl)(methyl)amino)-1H-indazol-1-carboxylate

MS (ESI m/z): 363 (M+H)

RT (min): 1.40

15 tert-Butyl (5-amino-1-methyl-1H-indazol-3-yl)(methyl)carbamate

MS (ESI m/z): 278 (M+H)

RT (min): 0.81

tert-Butyl 5-amino-3-(dimethylamino)-1H-indazol-1-carboxylate

MS (ESI m/z): 278 (M+H)


20 RT (min): 0.94

[0245]

Reference Example 41

The following compound was obtained as described in the 3rd step in
Reference Example 1.

25 [Formula 61]



[0246]

1st step

The following compound was obtained as described in the 1st step in

Reference Example 3.

tert-Butyl (1-(2-methoxyethyl)-5-nitro-1H-indazol-3-yl)carbamate

[0247]

2nd step

5 The following compound was obtained as described in the 2nd step in Reference Example 27.

1-(2-Methoxyethyl)-5-nitro-1H-indazol-3-amine

The following compound was obtained as described in the 2nd and 3rd steps in Reference Example 1.

10 [0248]

3rd step

1-(2-Methoxyethyl)-N,N-dimethyl-5-nitro-1H-indazol-3-amine

[0249]

4th step

15 1-(2-Methoxyethyl)-N³,N³-dimethyl-1H-indazol-3,5-diamine


MS (ESI m/z): 235 (M+H)

RT (min): 0.55

[0250]

Reference Example 42

20 [Formula 62]



Pyrazole (42 mg), cesium carbonate (340 mg), trans-N,N'-dimethylcyclohexan-1,2-diamine (74 mg), and copper iodide (50 mg) were added to a DMAc (2 ml) solution containing
 25 5-bromopyridin-3-amine (90 mg) in a nitrogen atmosphere, followed by stirring in a sealed tube at 150°C for 15 hours. The reaction solution was adjusted to room temperature and water was added, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under
 30 reduced pressure. The obtained residue was purified by silica gel chromatography (chloroform : MeOH = 1:0 to 10:1). A brown solid of 5-(pyrazol-1-yl)pyridin-3-amine (56.7 mg) was thus obtained.

MS (ESI m/z): 161 (M+H)

RT (min): 0.38


[0251]

Reference Example 43

The following compound was obtained with reference to US6133253

5 A1.

[Formula 63]




5-Bromo-6-methylpyridin-3-amine

[0252]

10 Reference Example 44

[Formula 64]



[0253]

1,2,3-Triazole (45 mg), cesium carbonate (260 mg),
 15 trans-N,N'-dimethylcyclohexan-1,2-diamine (76 mg), and copper iodide (50 mg) were added to a DMAc (2 ml) solution containing 5-bromo-6-methylpyridin-3-amine (100 mg) in a nitrogen atmosphere, followed by stirring in a sealed tube at 110°C for 8 hours. The reaction solution was adjusted to room temperature and water was added to the
 20 reaction solution, followed by extraction with ethyl acetate and washing with saturated saline. The organic layers were dried over anhydrous sodium sulfate. Next, the solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (chloroform : MeOH = 1:0 to 20:1). A white solid of
 25 6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-amine (12.8 mg) and brown oily matter of 6-methyl-5-(1H-1,2,3-triazol-1-yl)pyridin-3-amine (6.7 mg) were thus obtained.

6-Methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-amine

MS (ESI m/z): 176 (M+H)

30 RT (min): 0.44

$^1\text{H-NMR}$ (DMSO-d_6 , 300MHz) δ : 8.11 (s, 2H), 7.96 (d, 1H, $J = 2.7$ Hz), 7.25 (d, 1H, $J = 2.7$ Hz), 5.52 (br, 2H), 2.32 (s, 3H)

6-Methyl-5-(1H-1,2,3-triazol-1-yl)pyridin-3-amine


MS (ESI m/z): 176 ($M+H$)

5 RT (min): 0.20, 0.27

[0254]

Reference Example 45

[Formula 65]



10 [0255]

Cyclopropylboronic acid (360 mg), cesium carbonate (1.4 g), and $\text{Pd(PPh}_3)_4$ (166 mg) were added to a 1,4-dioxane/water (4.5 ml/0.5 ml) solution containing 5-bromopyridin-3-amine (500 mg) in a nitrogen atmosphere, followed by stirring in a sealed tube at 100°C for 5 hours. The reaction solution was adjusted to room temperature and water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate, the solvent was distilled away under reduced pressure, and the obtained residue was purified by silica gel chromatography (chloroform : MeOH = 1:0 to 30:1). A brown solid of 5-cyclopropyl pyridin-3-amine (314 mg) was thus obtained.


MS (ESI m/z): 135 ($M+H$)

RT (min): 0.39

[0256]

25 Reference Example 46

[Formula 66]



[0257]

1st step

1,2,3-triazole (340 mg) and cesium carbonate (1.74 g) were added to a DMAc (10 ml) solution containing methyl 2,6-dichloro-5-fluoronicotinate (1 g), followed by stirring at 70°C to 80°C for 1.5 hours. The reaction solution was adjusted to room temperature and water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. Next, the solvent was distilled away under reduced pressure. A mixture of methyl 2-chloro-5-fluoro-6-(2H-1,2,3-triazol-2-yl)nicotinate and methyl 2-chloro-5-fluoro-6-(1H-1,2,3-triazol-1-yl)nicotinate was thus obtained.

MS (ESI m/z): 257,259 (M+H)

RT (min): 1.07,1.13

[0258]

2nd step

The mixture of methyl 2-chloro-5-fluoro-6-(2H-1,2,3-triazol-2-yl)nicotinate and methyl 2-chloro-5-fluoro-6-(1H-1,2,3-triazol-1-yl)nicotinate obtained in the 1st step was dissolved in MeOH (5 ml) and ammonium formate (300 mg) and 10%Pd/C (200 mg) were added thereto, followed by reflux for 4.5 hours. The reaction solution was adjusted to room temperature and filtered through Celite, the filtrate was collected, and the solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 4:1). A white solid of methyl 5-fluoro-6-(2H-1,2,3-triazol-2-yl)nicotinate (250 mg) and a white solid of methyl 5-fluoro-6-(1H-1,2,3-triazol-1-yl)nicotinate (160 mg) were thus obtained.

Methyl 5-fluoro-6-(2H-1,2,3-triazol-2-yl)nicotinate

MS (ESI m/z): 223 (M+H)

RT (min): 0.90

Methyl 5-fluoro-6-(1H-1,2,3-triazol-1-yl)nicotinate


MS (ESI m/z): 223 (M+H)

RT (min): 0.95

[0259]

Reference Example 47

[Formula 67]



[0260]

1st step

A 5M potassium hydroxide aqueous solution (2ml) was added to a
 5 THF/MeOH (2 ml/2 ml) solution containing methyl
 5-fluoro-6-(2H-1,2,3-triazol-2-yl)nicotinate (250 mg), followed by stirring at
 room temperature for 1 hour. 6M hydrochloric acid was added to the
 reaction solution so as to acidify the reaction solution, followed by extraction
 with ethyl acetate. The organic layers were washed with saturated saline and
 10 dried over anhydrous sodium sulfate. The solvent was distilled away under
 reduced pressure and a white solid of
 5-fluoro-6-(2H-1,2,3-triazol-2-yl)nicotinic acid (198 mg) was thus obtained.
 MS (ESI m/z): 209 (M+H)

RT (min): 0.70

15 [0261]

2nd step

Triethylamine (158 μ l), tert-butanol, and DPPA (246 μ l) were added to
 a toluene (5 ml) solution containing
 5-fluoro-6-(2H-1,2,3-triazol-2-yl)nicotinic acid (198 mg) obtained in the 1st
 20 step, followed by stirring at 100°C for 1 hour. The reaction solution was
 adjusted to room temperature and the solvent was distilled away under
 reduced pressure. The obtained residue was purified by silica gel
 chromatography (n-hexane : ethyl acetate = 4:1 to 1:1) and a white solid of
 tert-butyl (5-fluoro-6-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)carbamate (94 mg)
 25 was thus obtained.

MS (ESI m/z): 180 (M+H)

RT (min): 0.64

[0262]

3rd step

30 TFA (1 ml) was added to tert-butyl
 (5-fluoro-6-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)carbamate (30 mg) obtained
 in the 2nd step, followed by stirring at room temperature for 1 hour. The
 solvent was distilled away under reduced pressure, a 5M sodium hydroxide
 aqueous solution was added to the obtained residue at 0°C so as to alkalify the

mixture, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and a light brown solid of 5-fluoro-6-(2H-1,2,3-triazol-2-yl)pyridin-3-amine (19 mg) was thus obtained.


MS (ESI m/z): 180 (M+H)

RT (min): 0.64

[0263]

Reference Example 48

[Formula 68]



[0264]

The following compound was obtained as described in Reference Example 47.

1st step

5-Fluoro-6-(1H-1,2,3-triazol-1-yl)nicotinic acid

MS (ESI m/z): 209 (M+H)

RT (min): 0.64

[0265]

2nd step

tert-Butyl (5-fluoro-6-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)carbamate

MS (ESI m/z): 280 (M+H)

RT (min): 1.20

3rd step

5-Fluoro-6-(1H-1,2,3-triazol-1-yl)pyridin-3-amine

MS (ESI m/z): 180 (M+H)


RT (min): 0.59

[0266]

Reference Example 49

The following compound was obtained with reference to Helvetica Chimica Acta, 1964, vol. 47, pp. 363, 376.

[Formula 69]




2,6-Dimethoxypyridin-4-amine

[0267]

Reference Example 50

5 [Formula 70]



N,N-dimethylglycine (1.27 g), copper iodide (1.88 g), potassium tert-butoxide (4.1 g), and 1H-1,2,3,-triazole (1.7 g) were added to a DMSO (25 ml) solution containing 5-bromo-6-methoxypyridin-3-amine (25 g), followed by stirring at 130°C for 2 hours. The reaction solution was adjusted to room temperature and water was added to the reaction solution. Then, 4M hydrochloric acid was added to adjust the pH to pH = 4, followed by extraction with ethyl acetate. Next, the obtained organic layers were dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:1). Yellow oily matter of 6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-amine (1 g) and a light yellow solid of 6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-amine (525 mg) was thus obtained.

20 6-Methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-amine

MS (ESI m/z): 192 (M+H)

RT (min): 0.58

¹H-NMR (CDCl₃, 300MHz) δ: 7.87 (s, 2H), 7.77 (d, 1H, J = 2.4Hz), 7.39 (d, 1H, J = 2.4Hz), 3.98 (s, 3H), 3.53 (br, 2H)

25 6-Methoxy-5-(1H-1,2,3-triazol-2-yl)pyridin-3-amine

MS (ESI m/z): 192 (M+H)


RT (min): 0.56

¹H-NMR (CDCl₃, 300MHz) δ: 8.36-8.33 (m, 1H), 7.82 (s, 1H), 7.77-7.72 (m, 2H), 3.98 (s, 3H), 3.60 (br, 2H)

30 [0268]

Reference Example 51

[Formula 71]



[0269]

5 1st step

2-methoxy ethanol (423 μ l) and sodium hydride (60% in oil) (196 mg) were added to a THF (10 ml) solution containing methyl 2,6-dichloro-5-fluoronicotinate (1g) under ice cooling in a nitrogen atmosphere, followed by stirring for 1.5 hours. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and yellow oily matter of methyl 2-chloro-5-fluoro-6-(2-methoxyethoxy)nicotinate (1.07 g) was thus obtained.

15 MS (ESI m/z): 264 (M+H)

RT (min): 1.41

[0270]

2nd step

10%Pd/C (200 mg) and ammonium formate (200 mg) were added to an MeOH (10 ml) solution containing methyl 2-chloro-5-fluoro-6-(2-methoxyethoxy)nicotinate (1.07g) obtained in the 1st step, followed by reflux for 1 hour. Next, insoluble matter was removed using Celite. The solvent was distilled away under reduced pressure. Then, the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:5). A white solid of methyl 5-fluoro-6-(2-methoxyethoxy)nicotinate (264 mg) was thus obtained.

MS (ESI m/z): 230 (M+H)

RT (min): 1.25

[0271]

3rd step

A 5M sodium hydroxide aqueous solution (2 ml) was added to a THF/MeOH (2 ml/2 ml) solution containing methyl
 5 5-fluoro-6-(2-methoxyethoxy)nicotinate (264 mg) obtained in the 2nd step, followed by stirring at room temperature for 1 hour. Next, 6M hydrochloric acid (2 ml) was added to the reaction solution at 5°C or less so as to acidify the reaction solution, followed by extraction with ethyl acetate. The organic
 10 layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and a white solid of 5-fluoro-6-(2-methoxyethoxy)nicotinic acid (197 mg) was thus obtained.

MS (ESI m/z): 216 (M+H)

RT (min): 0.95

15 [0272]

4th step

The following compound was obtained as described in the 2nd step in Reference Example 47.

tert-Butyl (5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)carbamate

20 MS (ESI m/z): 287 (M+H)

RT (min): 1.47

[0273]

5th step

The following compound was obtained as described in the 3rd step in
 25 Reference Example 47.

5-Fluoro-6-(2-methoxyethoxy)pyridin-3-amine


MS (ESI m/z): 187 (M+H)

RT (min): 0.67

[0274]

30 Reference Example 52

[Formula 72]



[0275]

1st step

The following compound was obtained as described in the 1st step in Reference Example 51.

2-(2-Methoxyethoxy)-5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridine

5 MS (ESI m/z): 266 (M+H)

RT (min): 1.18

[0276]

2nd step

10 10% Pd/C (200 mg) and ammonium formate (200 mg) were added to an MeOH/ethyl acetate (5 ml/5 ml) solution containing 2-(2-methoxyethoxy)-5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridine (286 mg) obtained in the 1st step, followed by reflux for 1 hour. Insoluble matter was removed using Celite and the solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel chromatography
15 (n-hexane : ethyl acetate = 1:0 to 1:5) and a white solid of 6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-amine (240 mg) was thus obtained.


MS (ESI m/z): 236 (M+H)

RT (min): 0.69

20 [0277]

Reference Example 53

[Formula 73]



[0278]

25 1st step

Cesium carbonate (24.4 g), 1,2,3-triazole (4.35 ml), 2,2,6,6-tetramethyl-3,5-heptanedione (3.89 ml), and copper iodide (7.1 g) were added to a DMSO (100 ml) solution containing 3-iodo-5-nitropyridin-2-ol (10 g) in a nitrogen atmosphere, followed by
30 stirring at 155°C for 1 hour. The reaction solution was adjusted to room temperature and water was added to the reaction solution. An insoluble precipitate was removed by filtration. 1M hydrochloric acid (110ml) was added to the resulting filtrate so as to acidify the reaction solution, followed

by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate and the solvent was distilled away under reduced pressure. Ethyl acetate was added to the obtained residue and insoluble matter was collected by filtration. A yellow solid mixture (4.86 g) of 5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridin-2-ol and 5-nitro-3-(1H-1,2,3-triazol-1-yl)pyridin-2-ol was thus obtained.

5-Nitro-3-(2H-1,2,3-triazol-2-yl)pyridin-2-ol

MS (ESI m/z): 208 (M+H)

RT (min): 0.69

5-Nitro-3-(1H-1,2,3-triazol-1-yl)pyridin-2-ol

MS (ESI m/z): 208 (M+H)

RT (min): 0.63

[0279]

2nd step

Thionyl chloride (19 ml) and DMF (4 ml) were added to a mixture (5.32 g) of 5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridin-2-ol and 5-nitro-3-(1H-1,2,3-triazol-1-yl)pyridin-2-ol, followed by reflux for 1 hour. The reaction solution was poured into ice water, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 10:1). A light yellow solid of 2-chloro-5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridine (0.72 g) and a yellow solid of 2-chloro-5-nitro-3-(1H-1,2,3-triazol-1-yl)pyridine (0.63 g) were thus obtained.

2-Chloro-5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridine

MS (ESI m/z): 226 (M+H)

RT (min): 1.18

2-Chloro-5-nitro-3-(1H-1,2,3-triazol-1-yl)pyridine

MS (ESI m/z): 226 (M+H)


RT (min): 0.92

[0280]

Reference Example 54

The following compound was obtained as described in Reference Example 52.

[Formula 74]



[0281]

1st step

2-Ethoxy-5-nitro-3-(1H-1,2,3-triazol-1-yl)pyridine

5 MS (ESI m/z): 236 (M+H)

RT (min): 1.21

2nd step

6-Ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-amine

MS (ESI m/z): 206 (M+H)


10 RT (min): 0.78

¹H-NMR (CDCl₃, 300MHz) δ: 8.39 (s, 1H), 7.83-7.80 (m, 1H), 7.77 (d, 1H, J = 2.7Hz), 7.72 (d, 1H, J = 2.7Hz), 4.43 (q, 2H, J = 7.2Hz), 3.60 (br, 2H), 1.40 (t, 3H, J = 7.2Hz)

[0282]

15 Reference Example 55

[Formula 75]



[0283]

20 The following compound was obtained as described in the 2nd and 3rd steps in Reference Example 47.

1st step

tert-Butyl (5-fluoro-6-methylpyridin-3-yl)carbamate

MS (ESI m/z): 227 (M+H)

RT (min): 1.32

25 [0284]

2nd step

5-Fluoro-6-methylpyridin-3-amine


MS (ESI m/z): 127 (M+H)

RT (min): 0.23, 0.29

30 [0285]

Reference Example 56

[Formula 76]



[0286]

Triethylamine (0.99 ml), benzyl alcohol, and DPPA (1.53 ml) were added to a toluene (10 ml) solution containing 5-fluoro-6-(2H-1,2,3-triazol-2-yl)nicotinic acid (920 mg), followed by stirring at 100°C for 1 hour. The reaction solution was adjusted to room temperature and the solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 4:1 to 1:1). A white solid of benzyl (5-fluoro-6-methylpyridin-3-yl)carbamate (94 mg) was thus obtained.

MS (ESI m/z): 261 (M+H)

RT (min): 1.35

[0287]

The following compound was obtained as described in the 2nd step in Reference Example 46.

5-Fluoro-6-methylpyridin-3-amine


MS (ESI m/z): 127 (M+H)

RT (min): 0.23,0.29

[0288]

Reference Example 57

[Formula 77]



[0289]

The following compound was obtained as described in Reference Example 53.

1st step

5-Nitro-3-(pyrazol-1-yl)pyridin-2-ol

MS (ESI m/z): 207 (M+H)

RT (min): 0.88

2nd step

2-Chloro-5-nitro-3-(pyrazol-1-yl)pyridine

MS (ESI m/z): 225, 227 (M+H)

RT (min): 1.17

5 [0290]

The following compound was obtained as described in Reference Example 52.

3rd step

2-Ethoxy-5-nitro-3-(pyrazol-1-yl) pyridine

10 MS (ESI m/z): 235 (M+H)

RT (min): 1.52

[0291]

4th step

6-Ethoxy-5-(pyrazol-1-yl)pyridin-3-amine


15 MS (ESI m/z): 205 (M+H)

RT (min): 0.93

[0292]

Reference Example 58

[Formula 78]



[0293]

1st step

25 Methylamine (9.8M in MeOH, 450 μ l) was added to an MeOH (5 ml) solution containing methyl 2,6-dichloro-5-fluoronicotinate (0.5 g) under ice cooling, followed by stirring at 70°C for 3.5 hours. The solvent and methylamine were distilled away under reduced pressure and methyl 2-chloro-5-fluoro-6-(methylamino)nicotinate was thus obtained.

MS (ESI m/z): 219, 221 (M+H)

RT (min): 1.20

30 [0294]

2nd step

The following compound was obtained as described in the 2nd step in Reference Example 46.

Methyl 5-fluoro-6-(methylamino)nicotinate

MS (ESI m/z): 185 (M+H)

RT (min): 0.88

[0295]

5 The following compound was obtained as described in the 1st step in Reference Example 47.

3rd step

5-fluoro-6-(methylamino)nicotinic acid

MS (ESI m/z): 171 (M+H)

10 RT (min): 0.50

[0296]

The following compound was obtained as described in Reference Example 56.

4th step

15 Benzozy(5-fluoro-6-(methylamino)pyridin-3-yl)carbamate

MS (ESI m/z): 276 (M+H)

RT (min): 1.04

5th step

3-Fluoro-N²-methylpyridin-2,5-diamine


20 MS (ESI m/z): 142 (M+H)

RT (min): 0.22

[0297]

Reference Example 59

[Formula 79]



25

1st step

tert-Butyl carbamate (215 mg), cesium carbonate (1.14 g), Pd₂(dba)₃ (240mg), and Xantphos (303 mg) were added to a toluene solution (9 ml) containing 2-chloro-3-fluoro-4-iodopyridine (450 mg), followed by stirring at 100°C for 3 hours. The reaction solution was adjusted to room temperature and water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline, dried over anhydrous sodium sulfate. Next, the solvent was distilled away under

30

reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 9:1 to 3:1). A yellow oily matter of tert-butyl (2-chloro-3-fluoropyridin-4-yl)carbamate (533 mg) was thus obtained.

5 MS (ESI m/z): 247, 249 (M+H)

RT (min): 1.46

[0298]

2nd step

The following compound was obtained as described in the 2nd step in

10 Reference Example 27.

2-Chloro-3-fluoropyridin-4-amine

MS (ESI m/z): 147, 149 (M+H)

RT (min): 0.72

[0299]

15 3rd step

Morpholine (3 ml) was added to 2-chloro-3-fluoropyridin-4-amine (133 mg) obtained in the 2nd step, followed by microwave irradiation (InitiatorTM, 180°C, 20 minutes, 2.45GHz, 0-240W). A saturated ammonium chloride aqueous solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with a saturated ammonium chloride aqueous solution and saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. A light brown solid of 3-fluoro-2-morpholinopyridin-4-amine (153 mg) was thus obtained.

25 3-Fluoro-2-morpholinopyridin-4-amine


MS (ESI m/z): 198 (M+H)

RT (min): 0.43

[0300]

Reference Example 60

30 [Formula 80]



1st step

Lithium N,N-diisopropyl amide (2M THF/ethylbenzene/heptane

solution) (2.9 ml) was mixed with THF (20 ml), and a THF (5 ml) solution containing 2-chloro-5-fluoropyridine (500 mg) was added to the mixture in a nitrogen atmosphere at -75°C , followed by stirring for 3 hours. Subsequently, a THF (5 ml) solution containing iodine (1.16 g) was added to the mixture, followed by stirring at -75°C for 1 hour. Next, water/THF (2 ml/8 ml), water (10 ml), and a 3M sodium thiosulfate aqueous solution were added to the reaction solution at -75°C , -50°C , and -35°C , respectively. The reaction solution was adjusted to room temperature, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 20:1 to 10:1). A white solid of 2-chloro-5-fluoro-4-iodopyridine (457 mg) was thus obtained.

2-Chloro-5-fluoro-4-iodopyridine

$^1\text{H-NMR}$ (CDCl_3 , 300MHz) δ : 8.14 (s, 1H), 7.77 (d, 1H, $J = 4.3\text{Hz}$)

[0301]

2nd step

tert-Butyl carbamate (960 mg), cesium carbonate (5.06 g), $\text{Pd}_2(\text{dba})_3$ (1.07 g), and Xantphos (1.35 g) were added to a toluene solution (40 ml) containing 2-chloro-5-fluoro-4-iodopyridine (2 g), followed by stirring at 100°C for 3 hours. The reaction solution was adjusted to room temperature and water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 9:1 to 4:1). A yellow oily matter of tert-butyl (2-chloro-5-fluoropyridin-4-yl)carbamate (1.53 g) was thus obtained.

tert-Butyl (2-chloro-5-fluoropyridin-4-yl)carbamate

MS (ESI m/z): 247, 249 ($M+H$)

RT (min): 1.64

[0302]

3rd step

The following compound was obtained as described in the 2nd step in Reference Example 27.

2-Chloro-5-fluoropyridin-4-amine

MS (ESI m/z): 147, 149 (M+H)

RT (min): 0.68

[0303]

4th step

5 Morpholine (3 ml) was added to 2-chloro-5-fluoropyridin-4-amine (262 mg) obtained in the 2nd step, followed by microwave irradiation (InitiatorTM, 235°C, 2 hours, 2.45 GHz, 0-240W). A saturated ammonium chloride aqueous solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with a
10 saturated ammonium chloride aqueous solution and saturated saline and dried over anhydrous sodium sulfate. Next, the solvent was distilled away under reduced pressure and a light brown solid of 5-fluoro-2-morpholinopyridin-4-amine (311 mg) was thus obtained.

5-Fluoro-2-morpholinopyridin-4-amine


15 MS (ESI m/z): 198 (M+H)

RT (min): 0.40

[0304]

Reference Example 61

[Formula 81]



20

1st step

Cesium carbonate (24.4 g), 1,2,3-triazole (4.35 ml), 2,2,6,6-tetramethyl-3,5-heptanedione (3.89 ml), and copper iodide (7.1 g) were added to a DMSO (100 ml) solution containing
25 3-iodo-5-nitropyridin-2-ol (10 g) in a nitrogen atmosphere, followed by stirring at 155°C for 1 hour. The reaction solution was adjusted to room temperature and water was added to the reaction solution. An insoluble precipitate was removed by filtration. 1M hydrochloric acid (110 ml) was added to the resulting filtrate, followed by extraction with ethyl acetate. The
30 organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure, a small amount of ethyl acetate was added to the obtained residue, and insoluble

matter was collected by filtration. A yellow solid mixture (4.86 g) of
 5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridin-2-ol and
 5-nitro-3-(1H-1,2,3-triazol-1-yl) pyridin-2-ol was thus obtained.

5-Nitro-3-(2H-1,2,3-triazol-2-yl)pyridin-2-ol

5 MS (ESI m/z): 208 (M+H)

RT (min): 0.69

5-Nitro-3-(1H-1,2,3-triazol-1-yl) pyridin-2-ol

MS (ESI m/z): 208 (M+H)

RT (min): 0.63

10 [0305]

2nd step

2,2-difluoro-2-(fluorosulfonyl)acetic acid (189 μ l) was added to an
 acetonitrile (2 ml) solution containing the mixture (105 mg) of
 5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridin-2-ol and
 15 5-nitro-3-(1H-1,2,3-triazol-1-yl) pyridin-2-ol obtained in the 1st step,
 followed by stirring at room temperature for 21 hours. Subsequently, sodium
 sulfate (100 mg) and 2,2-difluoro-2-(fluorosulfonyl)acetic acid (126 μ l) were
 added to the solution, followed by stirring at 70°C for 7 hours. The reaction
 solution was adjusted to room temperature and a saturated sodium hydrogen
 20 carbonate solution was added to the reaction solution, followed by extraction
 with ethyl acetate. The organic layers were washed with saturated saline and
 dried over anhydrous sodium sulfate. The solvent was distilled away under
 reduced pressure and the obtained residue was purified by silica gel
 chromatography (n-hexane : ethyl acetate = 1:0 to 4:1). A light yellow solid
 25 of 2-(difluoromethoxy)-5-nitro-3-(2H-1,2,3-triazol-2-yl)pyridine (15 mg) was
 thus obtained.


MS (ESI m/z): 258 (M+H)

RT (min): 1.29

[0306]

30 Reference Example 62

[Formula 82]



1st step

The following compound was obtained as described in the 3rd step in Reference Example 1.

6-(Difluoromethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-amine

5 MS (ESI m/z): 228 (M+H)


RT (min): 0.93

[0307]

Reference Example 63

10 The following compound was obtained with reference to WO2009/90548 A2, 2009.

[Formula 83]




3-(2H-1,2,3-triazol-2-yl)aniline

[0308]

15 Reference Example 64

The following compound was obtained with reference to Tetrahedron, 2011, vol. 67, # 2, pp. 289-292, WO2009/90548 A2, 2009.

[Formula 84]




20 3-(1H-1,2,3-triazol-1-yl) aniline

[0309]

Reference Example 65

[Formula 85]



25 4-Methyl-1H-pyrazole (114 mg), cesium carbonate (753 mg), trans-N,N'-dimethylcyclohexan-1,2-diamine (164 mg), and copper iodide (110 mg) were added to a DMAc (5 ml) solution containing

5-bromopyridin-3-amine (200 mg) in a nitrogen atmosphere, followed by microwave irradiation (InitiatorTM, 170°C, 0.5 hours, 2.45 GHz, 0-240 W). Water was added to the reaction solution and the reaction solution was filtered through Celite and the filtrate was extracted with ethyl acetate. The organic layers were washed with saturated saline and dried over sodium sulfate. The solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 4:1 to 0:1). 5-(4-Methylpyrazol-1-yl)pyridin-3-amine (173 mg) was thus obtained.


MS (ESI m/z): 175 (M+H)

10 RT (min): 0.54

[0310]

Reference Example 66

[Formula 86]



15 The following compound was obtained as described in Reference Example 65.

5-(3,5-Dimethylpyrazol-1-yl)pyridin-3-amine


MS (ESI m/z): 189 (M+H)

RT (min): 0.60

20 [0311]

Reference Example 67

[Formula 87]



25 [0312]

The following compound was obtained as described in Reference Example 65.

5-(4-Chloropyrazol-1-yl)pyridin-3-amine


MS (ESI m/z): 195, 197 (M+H)

RT (min): 0.66

[0313]

Reference Example 68

[Formula 88]



1st step

A sodium ethoxide solution (28% in MeOH) (1 ml) was added to an MeOH (3 ml) solution containing 3-bromo-2-chloro-5-nitropyridine (100 mg), followed by stirring at room temperature for 0.5 hours. A saturated ammonium chloride aqueous solution was added to the reaction solution and MeOH was distilled away under reduced pressure. An insoluble precipitate was washed with water and a white solid of 3-bromo-2-methoxy-5-nitropyridine (69 mg) was thus obtained.

MS (ESI m/z): 233, 235 (M+H)

15 RT (min): 1.40

[0314]

2nd step

Pd(PPh₃)₄ (60 mg) and tributyl (1-ethoxyvinyl)tin were added to a DMAc (3 ml) solution containing 3-bromo-2-methoxy-5-nitropyridine (69 mg) obtained in the 1st step, followed by microwave irradiation (InitiatorTM, 180°C, 10 minutes, 2.45 GHz, 0-240 W). Saturated sodium hydrogen carbonate was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 9:1 to 3:7). A light yellow solid of 1-(2-methoxy-5-nitropyridin-3-yl)ethanone (72 mg) was thus obtained.

MS (ESI m/z): 197 (M+H)

RT (min): 1.14

30 [0315]

3rd step

The following compound was obtained as described in the 3rd step in Reference Example 1.

1-(5-Amino-2-methoxypyridin-3-yl)ethanone

MS (ESI m/z): 167 (M+H)


RT (min): 0.60

[0316]

5 Reference Example 69

The following compound was obtained with reference to US2006/79522 A1.

[Formula 89]




10 3-Bromo-2-methyl-5-nitropyridine

[0317]

Reference Example 70

[Formula 90]



15 The following compound was obtained as described in the 2nd and 3rd steps in Reference Example 68.

1st step

1-(2-Methyl-5-nitropyridin-3-yl)ethanone

MS (ESI m/z): 181 (M+H)

20 RT (min): 0.90

[0318]

2nd step

1-(5-amino-2-methylpyridin-3-yl)ethanone


MS (ESI m/z): 151 (M+H)

25 RT (min): 0.28

[0319]

Reference Example 71

[Formula 91]



The following compound was obtained as described in Reference Example 68.

2-Methylamino-3-bromo-5-nitropyridine

5 MS (ESI m/z): 288, 290 (M+H)

RT (min): 1.36

1-(2-Methylamino-5-nitropyridin-3-yl)ethanone

MS (ESI m/z): 196 (M+H)

RT (min): 1.09

10 1-(5-Amino-2-methylaminopyridin-3-yl)ethanone


MS (ESI m/z): 166 (M+H)

RT (min): 0.32

[0320]

Reference Example 72

15 [Formula 92]



The following compound was obtained as described in Reference Example 68.

4-(3-Bromo-5-nitropyridin-2-yl)morpholine

20 MS (ESI m/z): 288, 290 (M+H)

RT (min): 1.36

1-(2-Morpholino-5-nitropyridin-3-yl)ethanone

MS (ESI m/z): 252 (M+H)

RT (min): 1.04

25 1-(5-Amino-2-morpholinopyridin-3-yl)ethanone


MS (ESI m/z): 222 (M+H)

RT (min): 0.53

[0321]

Reference Example 73

[Formula 93]



1st step

Pyrazole (130 mg) and cesium carbonate (610 mg) were added to a
 5 DMAc (10 ml) solution containing 5-bromo-2-chloropyrimidine (300 mg),
 followed by stirring at 120°C for 0.5 hours. The reaction mixture was
 adjusted to room temperature and water was added to the mixture. Next, the
 organic layers were collected, washed with saturated saline, and dried over
 anhydrous sodium sulfate. The solvent was distilled away under reduced
 10 pressure and a yellow solid of 5-bromo-2-(pyrazol-1-yl)pyrimidine (440 mg)
 was thus obtained.


MS (ESI m/z): 226 (M+H)

RT (min): 0.93

[0322]

15 Reference Example 74

[Formula 94]



The following compound was obtained as described in Reference
 Example 73.

20 5-Bromo-2-(2H-1,2,3-triazol-2-yl)pyrimidine

¹H-NMR: ¹H-NMR (CDCl₃) δ: 8.93 (2H, s), 8.01 (2H, s).

MS (ESI m/z): 227 (M+H)


RT (min): 0.68

[0323]

25 Reference Example 75

The following compound was obtained with reference to Chemische
 Berichte, 1967, vol. 100, #11 pp. 3485–3494.

[Formula 95]




4-methyl-2H-1,2,3-triazole

[0324]

Reference Example 76

5 [Formula 96]



The following compound was obtained as described in Reference Example 50.

10 5-(4-Methyl-2H-1,2,3-triazol-2-yl)pyridin-3-amine

MS (ESI m/z): 176 (M+H)

RT (min): 0.56

5-(4-Methyl-1H-1,2,3-triazol-1-yl)pyridin-3-amine


MS (ESI m/z): 176 (M+H)

15 RT (min): 0.53

[0325]

Reference Example 77

[Formula 97]




20 The following compound was obtained with reference to Chemische Berichte, 1967, vol. 100, #11 pp. 3485–3494.

4,5-Dimethyl-2H-1,2,3-triazole

[0326]

Reference Example 78

25 [Formula 98]



The following compound was obtained as described in Reference Example 50.

5-(4,5-Dimethyl-2H-1,2,3-triazol-2-yl)pyridin-3-amine

5 MS (ESI m/z): 190 (M+H)

RT (min): 0.62

5-(4,5-Dimethyl-1H-1,2,3-triazol-1-yl)pyridin-3-amine


MS (ESI m/z): 190 (M+H)

RT (min): 0.56

10 [0327]

Reference Example 79

[Formula 99]



[0328]

15 1st step

CDI (185.6 g) was added to a dichloromethane (2000 ml) solution containing N-(tert-butoxycarbonyl)-L-alanine (200 g) at 5°C or less, followed by stirring for 1 hour. Subsequently, triethylamine (115.8 g) and N-methoxy-N-methylamine hydrochloride (111.7 g) were added to the solution, followed by stirring at 15°C or less for 1.5 hours. Dichloromethane (230 ml) was added to the reaction solution. The organic layers were washed with a

20% sodium hydroxide aqueous solution and the solvent was distilled away under reduced pressure. Heptane was added to the obtained residue for suspension and a solid was collected by filtration. A white solid of (S)-tert-butyl (1-(methoxy(methyl)amino)-1-oxopropan-2-yl)carbamate (237.8 g) was thus obtained.

[0329]

2nd step

Magnesium turnings (7 g) and THF (30 ml) were introduced into a reaction container in a nitrogen atmosphere and dibromoethane (10 μ l) was added to the reaction container. After foaming was confirmed, a THF (150 ml) solution containing bromocyclobutane (40 g) was added dropwise at 70°C or less for 0.5 hours. Next, (S)-tert-butyl (1-(methoxy(methyl)amino)-1-oxopropan-2-yl)carbamate (27.9 g) obtained in the 1st step was added, followed by stirring at 50°C or less for 2 hours. The reaction solution was adjusted to room temperature and poured into a 10% citric acid aqueous solution (300 ml) so as to separate the organic layers. The obtained organic layers were dried over anhydrous sodium hydrogen sulfate and the solvent was distilled away under reduced pressure. A white solid of (S)-tert-butyl (1-cyclobutyl-3-oxobutan-2-yl)carbamate (36 g) was thus obtained.

[0330]

3rd and 4th steps

Tetraethyl orthotitanate (90.3 g), (S)-tert-butyl (3-oxobutan-2-yl)carbamate (36 g) obtained in the 2nd step and (R)-(+)-tert-butyl sulfinamide (23 g) were added to toluene (90 ml) in a nitrogen atmosphere, followed by stirring at 80°C or less for 7 hours. Sodium borohydride (11.95 g) was added thereto at -30°C or less, followed by stirring for 4 hours. Subsequently, the obtained mixture was adjusted to room temperature and MeOH was added so as to separate the organic layers. Next, the organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. Brown oily matter of tert-butyl ((2S)-1-((tert-butylsulfinyl)amino)-1-cyclobutylpropan-2-yl)carbamate (38 g) was thus obtained.

[0331]

5th step

4M hydrogen chloride/1,4-dioxane (100 ml) was added to an MeOH (900 ml) solution containing tert-butyl ((2S)-1-((tert-butylsulfinyl)amino)-1-cyclobutylpropan-2-yl)carbamate (38 g) obtained in the 4th step under ice cooling, followed by stirring for 30 minutes.

5 The reaction solution was adjusted to room temperature and poured into a 1M sodium hydroxide aqueous solution (600 ml), followed by extraction with ethyl acetate. The solvent was distilled away under reduced pressure and used in the subsequent step.

[0332]

10 6th step

DIPEA (17 ml) was added to a DMF (30 ml) solution containing tert-butyl ((1R,2S)-1-amino-1-cyclobutylpropan-2-yl)carbamate and 2,6-dichloro-5-fluoronicotinonitrile (16.2 g) obtained in the 5th step at 70°C, followed by stirring for 3 hours. The reaction solution was adjusted to room temperature and water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and the organic layers were dried over anhydrous sodium sulfate. The obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 10:1 to 1:1). A white solid of tert-butyl ((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate (16.5 g) was thus obtained.

¹H-NMR (CDCl₃) δ: 7.29 (1H, d, J = 9.9 Hz), 5.59 (1H, Br), 4.84 (1H, Br), 4.36-4.24 (1H, m), 3.90-3.76 (1H, m), 2.54-2.43 (1H, m), 2.05-1.81 (6H, m), 1.45 (9H, s), 1.14 (3H, d, J = 6.9 Hz).

25 MS (ESI m/z): 384 (M+H)


RT (min): 1.83

[0333]

Reference Example 80

The following compound was obtained as described in Reference Example 79.

30 [Formula 100]



tert-Butyl

((2S)-1-((tert-butylsulfinyl)imino)-1-(2-fluorophenyl)propan-2-yl)carbamate

MS (ESI m/z): 273 (M+H-Boc)

5 RT (min): 1.54

tert-Butyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate

¹H-NMR (CDCl₃) δ: 8.09 (1H, s), 7.36-7.22 (1H, m), 7.07-6.92 (3H, m), 5.00

10 (1H, d, J = 5.9 Hz), 4.32-4.24 (2H, m), 1.50 (9H, s), 1.14 (3H, d, J = 6.6 Hz)

MS (ESI m/z): 323 (M+H-Boc)


RT (min): 1.83

[0334]

Reference Example 81

15 The following compound was obtained as described in Reference Example 79.

[Formula 101]



(S)-tert-butyl (1-(3-fluorophenyl)-1-oxopropan-2-yl)carbamate

MS (ESI m/z): 168 (M+H-Boc)

RT (min): 1.52

5 tert-Butyl

((1R,2S)-1-((R)-1,1-dimethylethylsulfonamide)-1-(3-fluorophenyl)propan-2-yl)carbamate

MS (ESI m/z): 273 (M+H-Boc)

RT (min): 1.51

10 tert-Butyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate

¹H-NMR (CDCl₃) δ: 7.92 (1H, s), 7.24-7.17 (2H, m), 7.13-7.06 (2H, m), 5.38 (1H, d, J = 4.6 Hz), 4.39-4.25 (2H, m), 1.47 (9H, s), 1.16 (3H, d, J = 6.6 Hz)

15 MS (ESI m/z): 323 (M+H-Boc)


RT (min): 1.83

[0335]

Reference Example 82

The following compound was obtained as described in Reference Example 79.

[Formula 102]



(S)-tert-butyl (1-oxo-1-(pyridin-2-yl)propan-2-yl)carbamate

MS (ESI m/z): 151 (M+H-Boc)

25 RT (min): 1.31

tert-Butyl

((2S)-1-((tert-butylsulfinyl)imino)-1-(pyridin-2-yl)propan-2-yl)carbamate

MS (ESI m/z): 354 (M+H)

RT (min): 1.49

tert-Butyl

((1S,2S)-1-((R)-1,1-dimethylethylsulfonamide)-1-(pyridin-2-yl)propan-2-yl)carbamate

5 MS (ESI m/z): 356 (M+H)

RT (min): 1.13

tert-Butyl

((1S,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(pyridin-2-yl)propan-2-yl)carbamate

10 ¹H-NMR (CDCl₃) δ: 8.60 (1H, d, J = 4.6 Hz), 7.73 (1H, dd, J = 7.3, 7.3 Hz), 7.41 (1H, d, J = 7.9 Hz), 7.33-7.28 (2H, m), 5.49 (1H, s), 5.41-5.35 (1H, m), 4.29-4.19 (1H, m), 1.45 (9H, s), 1.05 (3H, d, J = 6.6 Hz).

MS (ESI m/z): 406 (M+H)


RT (min): 1.53

15 [0336]

Reference Example 83

The following compound was obtained as described in Reference Example 79.

[Formula 103]



20

(S)-tert-butyl (1-(4-methoxyphenyl)-1-oxopropan-2-yl)carbamate

MS (ESI m/z): 180 (M+H-Boc)

RT (min): 1.48

tert-Butyl

25 ((2S)-1-((tert-butylsulfinyl)imino)-1-(4-methoxyphenyl)propan-2-yl)carbamate

MS (ESI m/z): 283 (M+H-Boc)

RT (min): 1.65

tert-Butyl

((1R,2S)-1-((R)-1,1-dimethylethylsulfonamide)-1-(4-methoxyphenyl)propan-2-yl)carbamate

5 MS (ESI m/z): 385 (M+H)

RT (min): 1.50

tert-Butyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate

10 ¹H-NMR (CDCl₃) δ: 7.89 (1H, s), 7.17 (2H, t, J = 8.4 Hz), 6.88 (2H, t, J = 8.4 Hz), 4.98 (1H, d, J = 5.3 Hz), 4.34-4.20 (2H, m), 3.80 (3H, s), 1.48 (9H, s), 1.10 (3H, d, J = 6.6 Hz).

MS (ESI m/z): 433 (M+H)


RT (min): 1.81

15 [0337]

Reference Example 84

The following compound was obtained as described in Reference Example 79.

[Formula 104]



tert-Butyl

((1R,2S)-1-((R)-1,1-dimethylethylsulfonamide)-1-(4-(trifluoromethyl)phenyl)propan-2-yl)carbamate

MS (ESI m/z): 323 (M+H-Boc)

25 RT (min): 1.71

tert-Butyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(4-(trifluorome

thyl)phenyl)propan-2-yl)carbamate

$^1\text{H-NMR}$ (CDCl_3) δ : 8.21 (1H, s), 7.60 (2H, d, $J = 8.4$ Hz), 7.38 (2H, d, $J = 8.4$ Hz), 5.04 (1H, d, $J = 5.3$ Hz), 4.38-4.26 (2H, m), 1.49 (9H, s), 1.13 (3H, d, $J = 6.6$ Hz).

5 MS (ESI m/z): 373 ($M+H$)


RT (min): 1.91

[0338]

Reference Example 85

10 The following compound was obtained as described in Reference Example 79.

[Formula 105]



(S)-tert-butyl (3-oxo-4-phenylbutan-2-yl)carbamate

15 MS (ESI m/z): 164 ($M+H-\text{Boc}$)

RT (min): 1.48

tert-Butyl

((2S,3R)-3-((R)-1,1-dimethylethylsulfinamide)-4-phenylbutan-2-yl)carbamate

MS (ESI m/z): 269 ($M+H-\text{Boc}$)

20 RT (min): 1.63

tert-Butyl

((2S,3R)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate

25 $^1\text{H-NMR}$ (CDCl_3) δ : 7.25-7.15 (5H, m), 6.13 (1H, s), 4.69 (1H, s), 4.56-4.50 (1H, m), 4.00-3.90 (1H, m), 2.99 (1H, dd, $J = 13.9, 5.4$ Hz), 2.76 (1H, dd, $J = 13.9, 5.4$ Hz), 1.45 (9H, s), 1.18 (3H, d, $J = 6.6$ Hz).

MS (ESI m/z): 319 ($M+H-\text{Boc}$)


RT (min): 1.79

[0339]

Reference Example 86

The following compound was obtained as described in Reference
5 Example 79.

[Formula 106]



(S)-tert-butyl (1-oxo-1-(thiophen-2-yl)propan-2-yl)carbamate

10 MS (ESI m/z): 156 (M+H-Boc)

RT (min): 1.36

tert-Butyl

((2S)-1-((tert-butylsulfinyl)imino)-1-(thiophen-2-yl)propan-2-yl)carbamate

MS (ESI m/z): 381 (M+Na)

15 RT (min): 1.69

tert-Butyl ((1S,2S)-1-((R)-1,1-dimethylethylsulfinamide)-1-(thiophen-2-yl)
propan-2-yl)carbamate

MS (ESI m/z): 261 (M+H-Boc)

RT (min): 1.52

20 tert-Butyl

((1S,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(thiophen-2-yl)
propan-2-yl)carbamate

¹H-NMR (CDCl₃) δ: 7.87 (1H, s), 7.28-7.24 (1H, m), 7.06 (1H, d, J = 3.0 Hz),
7.00 (1H, dd, J = 5.3, 3.0 Hz), 5.37 (1H, d, J = 4.6 Hz), 4.52 (1H, s), 4.33-4.24
25 (1H, m), 1.47 (9H, s), 1.18 (3H, d, J = 6.6 Hz)

MS (ESI m/z): 409 (M-H)


RT (min): 1.78

[0340]

Reference Example 87

The following compound was obtained as described in Reference Example 79.

[Formula 107]



(S)-tert-butyl (1-(3-chlorothiophen-2-yl)-1-oxopropan-2-yl)carbamate

MS (ESI m/z): 190 (M+H-Boc)

RT (min): 1.59

10 tert-Butyl

((2S)-1-((tert-butylsulfinyl)imino)-1-(3-chlorothiophen-2-yl)propan-2-yl)carbamate

MS (ESI m/z): 293 (M+ H-Boc)

RT (min): 1.94

15 tert-Butyl

((1S,2S)-1-(3-chlorothiophen-2-yl)-1-((R)-1,1-dimethylethylsulfinamide)propan-2-yl)carbamate

MS (ESI m/z): 295 (M+H-Boc)

RT (min): 1.59

20 tert-Butyl

((1S,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(3-chlorothiophen-2-yl)propan-2-yl)carbamate

MS (ESI m/z): 433 (M-H)


RT (min): 1.92

25 [0341]

Reference Example 88

The following compound was obtained as described in Reference Example 79.

[Formula 108]



tert-Butyl

((1R,2S)-1-(3,4-difluorophenyl)-1-((R)-1,1-dimethylethylsulfonamide)propan-2-yl)carbamate

MS (ESI m/z): 291 (M+H-Boc)

RT (min): 1.58

tert-Butyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate

¹H-NMR (CDCl₃) δ: 8.18 (1H, s), 6.80-6.72 (3H, m), 4.93 (1H, d, J = 5.4 Hz), 4.32-4.21 (2H, m), 1.50 (9H, s), 1.16 (3H, d, J = 6.6 Hz).

MS (ESI m/z): 341 (M+H-Boc)


RT (min): 1.83

[0342]

Reference Example 89

The following compound was obtained as described in Reference Example 79.

[Formula 109]



tert-Butyl

((1R,2S)-1-(3,5-difluorophenyl)-1-((R)-1,1-dimethylethylsulfinamide)propan-2-yl)carbamate

5 MS (ESI m/z): 291 (M+H-Boc)

RT (min): 1.59

tert-Butyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(3,5-difluorophenyl)propan-2-yl)carbamate

10 ¹H-NMR (CDCl₃) δ: 8.20 (1H, s), 6.82-6.71 (3H, m), 4.95 (1H, d, J = 5.4 Hz), 4.32-4.18 (2H, m), 1.50 (9H, s), 1.13 (3H, d, J = 6.6 Hz).

MS (ESI m/z): 341 (M+H-Boc)


RT (min): 1.83

[0343]

15 Reference Example 90

The following compound was obtained as described in Reference Example 79.

[Formula 110]



(S)-benzyl (1-cyclohexyl-1-oxopropan-2-yl)carbamate

MS (ESI m/z): 290 (M+H)

RT (min): 1.82

5 Benzyl ((1R,2S)-1-amino-1-cyclohexylpropan-2-yl)carbamate

MS (ESI m/z): 291 (M+ H-Boc)

RT (min): 1.00

Benzyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclohexylpropan-2-yl)carbamate

10

MS (ESI m/z): 445 (M-H)


RT (min): 1.91

[0344]

Reference Example 91

15

[Formula 111]



[0345]

1st step

The following compound was obtained as described in the 1st step in Reference Example 79.

(S)-tert-butyl (4-methyl-3-oxopentan-2-yl)carbamate

2nd step

5 Sodium borohydride (3.4 g) was added to an MeOH/isopropanol (20 ml/20 ml) solution containing (S)-tert-butyl (4-methyl-3-oxopentan-2-yl)carbamate (16 g) obtained in the 1st step, followed by stirring at room temperature for 30 minutes. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic
10 layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and yellow oily matter of tert-butyl ((2S)-3-hydroxy-4-methylpentan-2-yl)carbamate (11.5 g) was thus obtained.

MS (ESI m/z): 218 (M+H)

15 RT (min): 1.22

[0346]

3rd step

p-Nitrobenzoic acid (10.6 g), triphenylphosphine (20.8 g), and DIAD (42 ml) were added to a THF (50 ml) solution containing tert-butyl
20 ((2S)-3-hydroxy-4-methylpentan-2-yl)carbamate (11.5 g) obtained in the 2nd step, followed by stirring at room temperature for 15 hours. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (hexane : ethyl acetate = 1:0 to 5:1). Yellow oily matter of (2S)-2-((tert-butoxycarbonyl)amino)-4-methylpentan-3-yl
25 4-nitrobenzoate (10 g) was thus obtained.

MS (ESI m/z): 367 (M+H)

RT (min): 1.87

4th step

A 1M lithium hydroxide aqueous solution (30 ml) was added to an
30 MeOH (10 ml) solution containing (2S)-2-((tert-butoxycarbonyl)amino)-4-methylpentan-3-yl 4-nitrobenzoate (10 g) obtained in the 3rd step, followed by stirring at room temperature for 1 hour. The solvent was distilled away under reduced pressure, followed by extraction with ethyl acetate. The organic layers were washed with a
35 saturated sodium hydrogen carbonate aqueous solution and saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away

under reduced pressure. Yellow oily matter of tert-butyl ((2S)-3-hydroxy-4-methylpentan-2-yl)carbamate (5.3 g) was thus obtained.

MS (ESI m/z): 218 (M+H)

RT (min): 1.22

5 [0347]

5th step

Phthalimide (4.3 g), triphenylphosphine (9.6 g), and DIAD (19.2 ml) were added to a THF (40 ml) solution containing tert-butyl ((2S)-3-hydroxy-4-methylpentan-2-yl)carbamate (5.3 g) obtained in the 4th step, followed by stirring at room temperature for 16 hours. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (hexane: ethyl acetate = 1:0 to 5:1). Yellow oily matter of tert-butyl ((2S,3R)-3-(1,3-dioxoisindolin-2-yl)-4-methylpentan-2-yl)carbamate (3 g) was thus obtained.

¹H-NMR (CDCl₃) δ: 7.89-7.81 (2H, m), 7.75-7.71 (2H, m), 4.44-4.24 (2H, m), 2.77-2.58 (1H, m), 1.46 (9H, s), 1.12 (6H, dd, J = 24.1, 6.6 Hz), 0.86 (3H, d, J = 6.6 Hz).

MS (ESI m/z): 347 (M+H)

20 RT (min): 1.73

6th step

An ethanol (10 ml) solution containing tert-butyl ((2S,3R)-3-(1,3-dioxoisindolin-2-yl)-4-methylpentan-2-yl)carbamate (3 g) obtained in the 5th step and hydrazine monohydrate (12.9 ml) was stirred at 80°C to 90°C for 48 hours. The solvent was distilled away under reduced pressure and an insoluble precipitate was removed. The obtained oily matter was used in the subsequent step.

tert-Butyl

((2S,3R)-3-(1,3-dioxoisindolin-2-yl)-4-methylpentan-2-yl)carbamate

30 MS (ESI m/z): 217 (M+H)

RT (min): 0.75

[0348]

7th step

The following compound was obtained as described in the 6th step in Reference Example 79.

tert-Butyl

((2S,3R)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate

¹H-NMR (CDCl₃) δ: 7.32 (1H, d, J = 9.9 Hz), 5.24 (1H, d, J = 8.3 Hz), 4.74 (1H, d, J = 7.6 Hz), 4.27-3.91(2H, m), 1.95-1.77 (1H, m), 1.44 (9H, s), 1.19-0.88 (9H, m).


MS (ESI m/z): 371 (M+H)

RT (min): 1.71

[0349]

Reference Example 92

[Formula 112]



[0350]

1st step

Triethylamine (59 ml) and methanesulfonylchloride (25 ml) were added to a THF solution (200 ml) containing (R)-tert-butyl (1-hydroxy-4-methylpentan-2-yl)carbamate (46.2 g) in an ice bath, followed by stirring for 0.5 hours. A saturated sodium hydrogen carbonate aqueous solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were dried over anhydrous sodium sulfate and the solvent was distilled away under reduced pressure. A white solid of (R)-2-((tert-butoxycarbonyl)amino)-4-methylpentyl methanesulfonate (60 g) was thus obtained.

2nd step

Potassium phthalimide (47 g) was added to a DMF (100 ml) solution containing (R)-2-((tert-butoxycarbonyl)amino)-4-methylpentyl methanesulfonate (60 g) obtained in the 1st step, followed by stirring at 70°C for 2 hours. The reaction solution was adjusted to room temperature and added dropwise to a saturated sodium hydrogen carbonate aqueous solution (1500 ml), followed by extraction with ethyl acetate. The organic layers were dried over anhydrous sodium sulfate and the solvent was distilled away

under reduced pressure. (R)-tert-butyl
 (1-(1,3-dioxoindolin-2-yl)-4-methylpentan-2-yl)carbamate was thus obtained.
 MS (ESI m/z): 347 (M+H)

RT (min): 1.65

5 [0351]

3rd step

4M hydrogen chloride/1,4-dioxane (100 ml) was added to a
 CHCl₃/MeOH (80 ml/40 ml) solution containing (R)-tert-butyl
 (1-(1,3-dioxoindolin-2-yl)-4-methylpentan-2-yl)carbamate obtained in the
 10 2nd step, followed by stirring at room temperature for 6 hours. The solvent
 was distilled away under reduced pressure and a yellow solid of
 (R)-2-(2-amino-4-methylpentyl)isoindolin-1,3-dione hydrochloride (21 g) was
 thus obtained.

MS (ESI m/z): 247 (M+H)

15 RT (min): 0.75

4th step

The following compound was obtained as described in the 7th step in
 Reference Example 91.

(R)-2-chloro-6-((1-(1,3-dioxoindolin-2-yl)-4-methylpentan-2-yl)amino)-5-flu
 20 oronicotinonitrile

MS (ESI m/z): 401 (M+H)

RT (min): 1.73

[0352]

5th step

25 The following compound was obtained as described in the 6th step in
 Reference Example 91.

(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-chloro-5-fluoronicotinonitrile

MS (ESI m/z): 271 (M+H)

RT (min): 0.96

30 6th step

The following compound was obtained as described in the 1st step in
 Reference Example 38.

(R)-tert-butyl

(2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-4-methylpentyl)carbamate

35 ¹H-NMR (CDCl₃, 300MHz) δ: 7.27 (d, 1H, J = 9.3Hz), 5.74 (d, 1H, J = 5.9Hz),
 4.79(br, 1H), 4.42-4.24 (m, 1H), 3.42-3.22 (m, 2H), 1.72-1.30 (m, 12H),

1.00-0.92 (m, 6H)

MS (ESI m/z): 371 (M+H)


RT (min): 1.81

[0353]

5 Reference Example 93

The following compound was obtained as described in Reference Example 91.

[Formula 113]



10 [0354]

(R)-tert-butyl (1-(1,3-dioxoisindolin-2-yl)-3-phenylpropan-2-yl)carbamate

MS (ESI m/z): 281 (M+H-Boc)

RT (min): 1.64

(R)-2-(2-amino-3-phenyl propyl)isoindolin-1,3-dione

15 MS (ESI m/z): 281 (M+H)

RT (min): 0.85

(R)-2-chloro-6-((1-(1,3-dioxoisindolin-2-yl)-3-phenylpropan-2-yl)amino)-5-fluoronicotinonitrile

MS (ESI m/z): 435 (M+H)

20 RT (min): 1.70

(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-2-chloro-5-fluoronicotinonitrile

MS (ESI m/z): 305 (M+H)

RT (min): 0.99

25 (R)-tert-butyl (2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-3-phenyl propyl)carbamate

¹H-NMR (CDCl₃) δ: 7.37-7.18 (6H, m), 6.55 (1H, d, J = 5.3 Hz), 4.83-4.70 (1H, m), 4.41-4.26 (1H, m), 3.43-3.21 (2H, m), 3.10 (1H, dd, J = 13.5, 5.0 Hz), 2.73 (1H, dd, J = 13.7, 8.8 Hz), 1.42 (9H, s).

30 MS (ESI m/z): 405 (M+H)


RT (min): 1.78

[0355]

Reference Example 94

The following compound was obtained as described in Reference
5 Example 91.

[Formula 114]



[0356]

(S)-tert-butyl

10 (1-(1,3-dioxoisindolin-2-yl)-3-(pyrazol-1-yl)propan-2-yl)carbamate

MS (ESI m/z): 371 (M+H)

RT (min): 1.34

(S)-2-(2-amino-3-(pyrazol-1-yl) propyl)isoindolin-1,3-dione

MS (ESI m/z): 271 (M+H)

15 RT (min): 0.57

(S)-2-chloro-6-((1-(1,3-dioxoisindolin-2-yl)-3-(pyrazol-1-yl)propan-2-yl)amino)-5-fluoronicotinonitrile

MS (ESI m/z): 425 (M+H)

RT (min): 1.38

20 (S)-6-((1-amino-3-(pyrazol-1-yl)propan-2-yl)amino)-2-chloro-5-fluoronicotinonitrile

MS (ESI m/z): 295 (M+H)

RT (min): 0.75

(S)-tert-butyl

25 (2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-3-(pyrazol-1-yl)propyl)carbamate

MS (ESI m/z): 395 (M+H)


RT (min): 1.42

[0357]

30 Reference Example 95

The following compound was synthesized with reference to WO2010/097248.

[Formula 115]




5 tert-Butyl ((3R,4R)-4-aminotetrahydro-2H-pyran-3-yl)carbamate

[0358]

Reference Example 96

The following compound was obtained as described in Reference Example 91.

10 [Formula 116]



[0359]

(2S)-2-((tert-butoxycarbonyl)amino)-1-cyclopropyl-butyl 4-nitrobenzoate

MS (ESI m/z): 379 (M+H)

15 RT (min): 1.80

tert-Butyl

((1R,2S)-1-cyclopropyl-1-(1,3-dioxoisindolin-2-yl)butan-2-yl)carbamate

MS (ESI m/z): 359 (M+H)

RT (min): 1.69

20 tert-Butyl ((1R,2S)-1-amino-1-cyclopropylbutan-2-yl)carbamate

MS (ESI m/z): 229 (M+H)

RT (min): 0.79

tert-Butyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate

¹H-NMR (CDCl₃) δ: 7.32 (1H, d, J = 9.9 Hz), 6.22 (1H, Br), 4.60 (1H, Br), 3.84-3.62 (2H, m), 1.79-1.57 (2H, m), 1.48 (9H, t, J = 9.2 Hz), 1.01 (3H, t, J = 7.3 Hz), 0.95-0.83 (1H, m), 0.70-0.35 (4H, m)

MS (ESI m/z): 383 (M+H).


RT (min): 1.79

[0360]

Reference Example 97

The following compound was obtained as described in Reference Example 91.

[Formula 117]



[0361]

(S)-tert-butyl (3-oxohexan-2-yl)carbamate

MS (ESI m/z): 216 (M+H)

RT (min): 1.37

((2S)-3-hydroxyhexan-2-yl)carbamate

MS (ESI m/z): 218 (M+H)

RT (min): 1.27

(2S)-2-((tert-butoxycarbonyl)amino)hexan-3-yl 4-nitrobenzoate

MS (ESI m/z): 367 (M+H)

RT (min): 1.86

tert-Butyl ((2S)-3-hydroxyhexan-2-yl)carbamate

MS (ESI m/z): 218 (M+H)

RT (min): 1.27

tert-Butyl ((2S,3R)-3-(1,3-dioxoisindolin-2-yl)hexan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.88-7.79 (m, 2H), 7.76-7.65 (m, 2H), 4.62-4.42 (m, 1H), 4.33-4.00 (m, 2H), 2.40-2.20 (m, 1H), 1.81-1.62 (m, 1H), 1.44 (s, 9H), 1.35-1.20 (m, 2H), 1.11 (d, 3H, J = 6.6Hz), 0.89 (t, 3H, J =

7.3 Hz)

MS (ESI m/z): 347 (M+H)

RT (min): 1.70

tert-Butyl ((2S,3R)-3-aminohexan-2-yl)carbamate

5 MS (ESI m/z): 217 (M+H)

RT (min): 0.79

7th step

tert-Butyl

((2S,3R)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)hexan-2-yl)carba

10 mate

¹H-NMR (CDCl₃, 300MHz) δ: 7.29 (d, 1H, J = 9.3Hz), 5.76 (d, 1H, J = 7.3Hz), 4.67 (d, 1H, J = 6.6Hz), 4.36-4.20 (m, 1H), 3.96-3.80 (m, 1H), 1.70-1.29 (m, 13H), 1.17 (d, 3H, J = 6.6Hz), 0.94 (t, 3H, J = 7.3Hz)

MS (ESI m/z): 371 (M+H)


15 RT (min): 1.78

[0362]

Reference Example 98

The following compound was obtained as described in Reference Example 91.

20 [Formula 118]



[0363]

tert-Butyl ((2S)-3-hydroxypentan-2-yl)carbamate

MS (ESI m/z): 204 (M+H)

25 RT (min): 1.12

(2S)-2-((tert-butoxycarbonyl)amino)pentan-3-yl 4-nitrobenzoate

MS (ESI m/z): 353 ($M+H$)

RT (min): 1.75

tert-Butyl ((2*S*)-3-hydroxypentan-2-yl)carbamate

30 MS (ESI m/z): 204 (M+H)

RT (min): 1.13

tert-Butyl ((2S,3R)-3-(1,3-dioxoisindolin-2-yl)pentan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.84 (dd, 2H, J = 3.3,5.4Hz), 7.72 (dd, 2H, J = 3.3,5.4Hz), 4.60-4.50 (m, 1H), 4.35-4.20 (m, 1H), 4.10-3.95 (m, 1H),
 5 2.38-2.17 (m, 1H), 1.93-1.80 (m, 1H), 1.43 (s, 9H), 1.11 (d, 3H, J = 6.6Hz),
 0.86 (d, 3H, J = 7.3Hz)

MS (ESI m/z): 333 (M+H)

RT (min): 1.56

tert-Butyl ((2S,3R)-3-aminopentan-2-yl)carbamate

10 MS (ESI m/z): 203 (M+H)

RT (min): 0.69

tert-Butyl

((2S,3R)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate

15 ¹H-NMR (CDCl₃, 300MHz) δ: 7.29 (d, 1H, J = 9.9Hz), 5.76 (d, 1H, J = 6.6 Hz),
 4.68 (d, 1H, J = 6.6Hz), 4.26-4.14 (m, 1H), 3.98-3.84 (m, 1H), 1.80-1.62 (m,
 1H), 1.49-1.36(m,10H), 1.17 (d, 3H, J = 7.2Hz), 0.97 (t, 3H, J = 7.7Hz)

MS (ESI m/z): 357 (M+H)


RT (min): 1.67

20 [0364]

Reference Example 99

The following compound was obtained as described in Reference Example 91.

[Formula 119]



25

[0365]

tert-Butyl ((2S)-1-cyclopropyl-1-hydroxypropan-2-yl)carbamate

MS (ESI m/z): 216 (M+H)

RT (min): 1.14

30 (2S)-2-((tert-butoxycarbonyl)amino)-1-cyclopropylpropyl-4-nitrobenzoate

MS (ESI m/z): 365 (M+H)

RT (min): 1.76

tert-Butyl ((2S)-1-cyclopropyl-1-hydroxypropan-2-yl)carbamate

MS (ESI m/z): 216 (M+H)

5 RT (min): 1.14

tert-Butyl

((1R,2S)-1-cyclopropyl-1-(1,3-dioxindolin-2-yl)propan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.87-7.68 (m, 4H), 4.62 (br, 1H), 4.45-4.28 (m, 1H), 3.31(dd, 1H, J = 10.7, 6.8Hz), 2.25-1.75 (m, 1H), 1.40 (s, 9H), 1.18 (t, 3H, J = 6.9Hz), 0.85-0.72 (m, 1H), 0.52-0.38 (m, 2H), 0.16-0.04 (m, 1H)

MS (ESI m/z): 345 (M+H)

RT (min): 1.60

tert-Butyl

15 ((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.32-7.28 (m, 1H), 6.20 (br, 1H), 4.90-4.74 (m, 1H), 4.12-3.98 (m, 1H), 3.68-3.50 (m, 1H), 1.44 (s, 9H), 1.27 (t, 3H, J = 3.3Hz), 0.98-0.85 (m, 1H), 0.73-0.40 (m, 4H)

MS (ESI m/z): 369 (M+H)


20 RT (min): 1.72

[0366]

Reference Example 100

The following compound was obtained as described in Reference Example 91.

25 [Formula 120]



[0367]

(2S)-2-((tert-butoxycarbonyl)amino)-1-cyclobutylpropyl 4-nitrobenzoate

MS (ESI m/z): 379 (M+H)

30 RT (min): 1.91

tert-Butyl

((1R,2S)-1-cyclobutyl-1-(1,3-dioxoisindolin-2-yl)propan-2-yl)carbamate

MS (ESI m/z): 359 (M+H)

RT (min): 1.71

5 tert-Butyl ((1R,2S)-1-amino-1-cyclobutylpropan-2-yl)carbamate

MS (ESI m/z): 229 (M+H)

RT (min): 0.85

tert-Butyl

((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate

10

MS (ESI m/z): 384 (M+H)


RT (min): 1.83

[0368]

Reference Example 101

15 The following compound was obtained as described in Reference Example 91.

[Formula 121]



[0369]

20 tert-Butyl ((2S)-4-cyclopropyl-3-hydroxybutan-2-yl)carbamate

MS (ESI m/z): 230 (M+H)

RT (min): 1.32

(3S)-3-((tert-butoxycarbonyl)amino)-1-cyclopropylbutan-2-yl)
4-nitrobenzoate

25 MS (ESI m/z): 379 (M+H)

RT (min): 1.89

tert-Butyl ((2S)-4-cyclopropyl-3-hydroxybutan-2-yl)carbamate

MS (ESI m/z): 230 (M+H)

RT (min): 1.32

tert-Butyl

((2S,3R)-4-cyclopropyl-3-(1,3-dioxoisindolin-2-yl)butan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.87-7.69 (m, 4H), 5.81-5.66 (m, 1H),
5.00-4.82 (m, 2H), 4.58-4.46 (br, 1H), 4.33-4.06 (m, 2H), 2.55-1.80 (m, 2H),
1.44 (s, 9H), 1.34-1.26 (m, 2H), 1.11 (d, 3H, J = 6.6Hz)

MS (ESI m/z): 359 (M+H)

RT (min): 1.70

tert-Butyl ((2S,3R)-3-amino-4-cyclopropylbutan-2-yl)carbamate

MS (ESI m/z): 229 (M+H)

RT (min): 0.89

tert-Butyl

((2S,3R)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-4-cyclopropylbutan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.29 (d, 1H, J = 9.9Hz), 5.94-5.74 (m, 1H),
5.06-4.95 (m, 2H), 4.62 (br, 1H), 4.34-4.25 (m, 1H), 3.96-3.87 (m, 1H),
2.17-2.08 (m, 2H), 1.78-1.67 (m, 1H), 1.55-1.46 (m, 2H), 1.44 (s, 9H), 1.18 (d,
3H, J = 7.3Hz)

MS (ESI m/z): 383 (M+H)


RT (min): 1.77

[0370]

Reference Example 102

The following compound was obtained as described in Reference Example 91.

[Formula 122]



[0371]

(S)-tert-butyl (3-oxoheptan-2-yl)carbamate

MS (ESI m/z): 230 (M+H)

RT (min): 1.53

tert-Butyl ((2S)-3-hydroxyheptan-2-yl)carbamate

MS (ESI m/z): 232 (M+H)

RT (min): 1.40

(2S)-2-((tert-butoxycarbonyl)amino)heptan-3-yl 4-nitrobenzoate

5 MS (ESI m/z): 381 (M+H)

RT (min): 1.96

tert-Butyl ((2S)-3-hydroxyheptan-2-yl)carbamate

MS (ESI m/z): 232 (M+H)

RT (min): 1.43

10 tert-Butyl ((2S,3R)-3-(1,3-dioxoisindolin-2-yl)heptan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.87-7.79 (m, 2H), 7.76-7.68 (m, 2H), 4.53(br,1H), 4.32-3.99 (m, 2H), 2.40-2.17 (m, 1H), 1.86-1.69 (m, 1H), 1.44 (s, 9H), 1.36-1.04 (m,7H), 0.83 (t, 3H, J = 7.2Hz)

MS (ESI m/z): 361 (M+H)

15 RT (min): 1.81

tert-Butyl ((2S,3R)-3-aminoheptan-2-yl)carbamate

MS (ESI m/z): 231 (M+H)

RT (min): 0.89

tert-Butyl

20 ((2S,3R)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)heptan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.29 (d, 1H, J = 9.9Hz), 5.74 (d, 1H, J = 7.3Hz), 4.68 (d, 1H, J = 6.6Hz), 4.34-4.18 (m, 1H), 3.97-3.80 (m, 1H), 1.71-1.22(m,15H), 1.17 (t, 3H, J = 6.6 Hz), 0.89 (t, 3H, J = 6.3Hz)

25 MS (ESI m/z): 385 (M+H)


RT (min): 1.87

[0372]

Reference Example 103

The following compound was obtained as described in Reference
30 Example 91.

[Formula 123]



[0373]

(S)-tert-butyl (5-methyl-3-oxohexan-2-yl)carbamate

MS (ESI m/z): 230 (M+H)

5 RT (min): 1.53

tert-Butyl ((2S)-3-hydroxy-5-methylhexan-2-yl)carbamate

MS (ESI m/z): 232 (M+H)

RT (min): 1.42

(2S)-2-((tert-butoxycarbonyl)amino)-5-methylhexan-3-yl 4-nitrobenzoate

10 MS (ESI m/z): 381 (M+H)

RT (min): 1.95

tert-Butyl ((2S)-3-hydroxy-5-methylhexan-2-yl)carbamate

MS (ESI m/z): 232 (M+H)

RT (min): 1.42

15 tert-Butyl

((2S,3R)-3-(1,3-dioxoisindolin-2-yl)-5-methylhexan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.86-7.77 (m, 2H), 7.75-7.66 (m, 2H), 4.55 (br, 1H), 4.32-4.12 (m, 2H), 2.48-2.30 (m, 1H), 1.51-1.36 (s, 10H), 1.32-1.22 (m, 1H), 1.11 (d, 3H, J = 6.6Hz), 0.92-0.84 (m, 6H)

20 MS (ESI m/z): 361 (M+H)

RT (min): 1.80

tert-Butyl ((2S,3R)-3-amino-5-methylhexan-2-yl)carbamate

MS (ESI m/z): 231 (M+H)

RT (min): 0.89

25 tert-Butyl

((2S,3R)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.29 (d, 1H, J = 9.9Hz), 5.69 (d, 1H, J = 7.9Hz), 4.67 (d, 1H, J = 6.6Hz), 4.46-4.28 (m, 1H), 3.96-3.80 (m, 1H), 1.70-1.32 (m, 12H), 1.16 (d, 3H, J = 6.6Hz), 0.94 (dd, 6H, J = 6.6, 2.0Hz)

30

MS (ESI m/z): 385 (M+H)


RT (min): 1.86

[0374]

Reference Example 104

- 5 The following compound was obtained as described in Reference Example 91.

[Formula 124]



tert-Butyl ((2S)-1-hydroxy-1-phenylpropan-2-yl)carbamate

- 10 MS (ESI m/z): 252 (M+H)

RT (min): 1.34

tert-Butyl

((1R,2S)-1-(1,3-dioxoisindolin-2-yl)-1-phenylpropan-2-yl)carbamate

MS (ESI m/z): 381 (M+H)

- 15 RT (min): 1.67

tert-Butyl ((1R,2S)-1-amino-1-phenylpropan-2-yl)carbamate

MS (ESI m/z): 251

RT (min): 0.86

tert-Butyl

- 20 ((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.95 (br, 1H), 7.42-7.19 (m, 6H), 5.04 (d, 1H, J = 6.3Hz), 4.37-4.20 (m, 2H), 1.49 (s, 9H), 1.13 (d, 3H, J = 6.3Hz)

MS (ESI m/z): 405 (M+H)


- 25 RT (min): 1.96

[0375]

Reference Example 105

The following compound was obtained as described in Reference Example 91.

[Formula 125]



[0376]

- 5 tert-Butyl ((2S)-1-(4-fluorophenyl)-1-hydroxypropan-2-yl)carbamate
 MS (ESI m/z): 270 (M+H)
 RT (min): 1.57
 (2S)-2-((tert-butoxycarbonyl)amino)-1-(4-fluorophenyl)propyl-4-nitrobenzoate
- 10 MS (ESI m/z): 419 (M+H)
 RT (min): 1.85
 tert-Butyl ((2S)-1-(4-fluorophenyl)-1-hydroxypropan-2-yl)carbamate
 MS (ESI m/z): 270 (M+H)
 RT (min): 1.57
- 15 tert-Butyl
 ((1R,2S)-1-(1,3-dioxoisindolin-2-yl)-1-(4-fluorophenyl)propan-2-yl)carbamate
 MS (ESI m/z): 399 (M+H)
 RT (min): 1.74
- 20 tert-Butyl ((1R,2S)-1-amino-1-(4-fluorophenyl)propan-2-yl)carbamate
 MS (ESI m/z): 269(M+H)
 RT (min): 0.89
 tert-Butyl
 ((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-(4-fluorophenyl)propan-2-yl)carbamate
- 25 ¹H-NMR (CDCl₃, 300MHz) δ: 8.07 (br, 1H), 7.25-7.16 (m, 3H), 7.09-6.98 (m, 2H), 4.99 (d, 1H, J = 5.9Hz), 4.36-4.16 (m, 2H), 1.50 (s, 9H), 1.12 (d, 3H, J = 6.6Hz)
 MS (ESI m/z): 423 (M+H)


RT (min): 1.81

[0377]

Reference Example 106

The following compound was obtained as described in Reference
5 Example 91.

[Formula 126]



[0378]

(S)-tert-butyl (2-oxopentan-3-yl)carbamate

10 MS (ESI m/z): 202 (M+H)

RT (min): 1.19

tert-Butyl ((3S)-2-hydroxypentan-3-yl)carbamate

MS (ESI m/z): 204 (M+H)

RT (min): 1.09

15 (3S)-3-((tert-butoxycarbonyl)amino)pentan-2-yl-4-nitrobenzoate

MS (ESI m/z): 353 (M+H)

RT (min): 1.75

tert-Butyl ((3S)-2-hydroxypentan-3-yl)carbamate

MS (ESI m/z): 204 (M+H)

20 RT (min): 1.09

tert-Butyl ((2R,3S)-2-(1,3-dioxoisindolin-2-yl)pentan-3-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.89-7.75 (m, 2H), 7.76-7.66 (m, 2H), 4.46 (d, 1H, J = 8.6Hz), 4.36-4.02 (m, 2H), 1.41 (s, 9H), 1.37-1.22 (m, 5H), 0.92 (t, 3H, J = 7.2)

25 MS (ESI m/z): 333 (M+H)

RT (min): 1.58

tert-Butyl ((2R,3S)-2-aminopentan-3-yl)carbamate

MS (ESI m/z): 203 (M+H)

RT (min): 0.69

30 tert-Butyl

((2R,3S)-2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)pentan-3-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.25 (d, 1H, J = 9.9Hz), 6.79 (d, 1H, J = 5.4Hz), 4.46 (d, 1H, J = 7.9Hz), 4.30-4.15 (m, 1H), 3.80-3.68 (m, 1H), 1.71-1.30(m,11H), 1.17 (d, 3H, J = 6.6Hz), 1.02 (t, 3H, J = 7.6Hz)

MS (ESI m/z): 357 (M+H)


RT (min): 1.72

[0379]

Reference Example 107

10 The following compound was obtained as described in Reference Example 91.

[Formula 127]



15 [0380]

(S)-tert-butyl (4-oxohexan-3-yl)carbamate

MS (ESI m/z): 216 (M+H)

RT (min): 1.36

tert-Butyl ((3S)-4-hydroxyhexan-3-yl)carbamate

20 MS (ESI m/z): 218 (M+H)

RT (min): 1.26

(4S)-4-((tert-butoxycarbonyl)amino)hexan-3-yl-4-nitrobenzoate

MS (ESI m/z): 367 (M+H)

RT (min): 1.85

25 tert-Butyl ((3S)-4-hydroxyhexan-3-yl)carbamate

MS (ESI m/z): 218 (M+H)

RT (min): 1.26

tert-Butyl ((3S,4R)-4-(1,3-dioxoisindolin-2-yl)hexan-3-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.89-7.78 (m, 2H), 7.76-7.66 (m, 2H), 4.46 (d,

1H, J = 8.6Hz), 4.36-3.90 (m, 2H), 2.39-2.15 (m, 1H), 1.96-1.76 (m, 1H), 1.67-1.40 (m, 10H), 1.34-1.16 (m, 1H), 0.96-0.80 (m, 6H)

MS (ESI m/z): 347 (M+H)

RT (min): 1.68

5 tert-Butyl ((3S,4R)-4-aminohexan-3-yl)carbamate

MS (ESI m/z): 217 (M+H)

RT (min): 0.75

tert-Butyl

((3S,4R)-4-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)hexan-3-yl)carbamate

10 ¹H-NMR (CDCl₃, 300MHz) δ: 7.28 (d, 1H, J = 9.9Hz), 5.80 (d, 1H, J = 7.9Hz), 4.43 (d, 1H, J = 8.6Hz), 4.29-4.05 (m, 1H), 3.74-3.60 (m, 1H), 1.78-1.27 (m, 13H), 1.00 (t, 3H, J = 7.7Hz), 0.96 (t, 3H, J = 7.5Hz)


MS (ESI m/z): 371 (M+H)

15 RT (min.): 1.77

[0381]

Reference Example 108

[Formula 128]



20 [0382]

1st step

The following compound was obtained as described in the 1st step in Reference Example 79.

(R)-tert-butyl

25 (3-methoxy-1-(methoxy(methyl)amino)-1-oxopropan-2-yl)carbamate

MS (ESI m/z): 263 (M+H)

RT (min): 1.03

The following compound was obtained as described in the 1st to 5th steps in Reference Example 91.

2nd step

(R)-tert-butyl (1-methoxy-3-oxobutan-2-yl)carbamate

MS (ESI m/z): 218 (M+H)

RT (min): 1.07

5 [0383]

3rd step

tert-Butyl ((2R)-3-hydroxy-1-methoxybutan-2-yl)carbamate

MS (ESI,m/z): 220 (M+H)

RT (min): 0.92

10 4th step

(3R)-3-((tert-butoxycarbonyl)amino)-4-methoxybutan-2-yl-4-nitrobenzoate

MS (ESI m/z): 369 (M+H)

RT (min): 1.67

[0384]

15 5th step

tert-Butyl ((2R)-3-hydroxy-1-methoxybutan-2-yl)carbamate

MS (ESI m/z): 220(M+H)

RT (min): 0.92

6th step

20 tert-Butyl

((2S,3S)-3-((1,3-dioxoisindolin-2-yl)-1-methoxybutan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.85-7.78 (m, 2H), 7.74-7.66 (m, 2H), 5.08-4.92 (m, 1H), 4.54-4.34 (m, 2H), 3.44-3.26 (m, 2H), 3.22 (s, 3H), 1.52 (d, 3H, J = 6.6Hz), 1.45 (s, 9H)

25 MS (ESI m/z): 349 (M+H)

RT (min): 1.50

[0385]

7th step

The following compound was obtained as described in the 3rd step in

30 Reference Example 47.

2-((2S,3S)-3-amino-4-methoxybutan-2-yl)isoindolin-1,3-dione


MS (ESI m/z): 249 (M+H),

RT (min): 0.64

[0386]

35 Reference Example 109

[Formula 129]



[0387]

1st step

The following compound was obtained as described in the 6th step in Reference Example 79.

2-chloro-6-(((2S,3S)-3-(1,3-dioxoisindolin-2-yl)-1-methoxybutan-2-yl)amino)-5-fluoronicotinonitrile

MS (ESI m/z): 403 (M+H),

RT (min): 1.59

2nd step

The following compound was obtained as described in the 6th step in Reference Example 91.

6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-chloro-5-fluoronicotinonitrile

MS (ESI m/z): 273 (M+H),

RT (min): 0.72

[0388]

3rd step

The following compound was obtained as described in the 1st step in Reference Example 38.

tert-Butyl

((2S,3S)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.31 (d, 1H, J = 9.6Hz), 6.10 (d, 1H, J = 7.6Hz), 5.17(d, 1H, J = 8.9Hz), 4.36-4.19 (m, 1H), 4.12-3.94 (m, 1H), 3.89 (s, 3H), 3.84-3.75 (m, 1H), 3.58-3.48 (m, 1H), 1.44 (s, 9H), 1.24 (d, 3H, J = 7.2Hz)

MS (ESI m/z): 373 (M+H)


RT (min): 1.60

[0389]

Reference Example 110

The following compound was obtained as described in Reference Example 108.

[Formula 130]



[0390]

1st step

(R)-tert-butyl

5 (3-ethoxy-1-(methoxy(methyl)amino)-1-oxopropan-2-yl)carbamate

$^1\text{H-NMR}$ (CDCl_3) δ : 5.38 (1H, d, $J = 8.6$ Hz), 4.90-4.78 (1H, m), 3.78 (3H, s), 3.72-3.43 (4H, m), 3.23 (3H, s), 1.46 (9H, s), 1.17 (3H, t, $J = 7.1$ Hz).

MS (ESI m/z): 117 ($M+H-\text{Boc}$)

RT (min): 1.17

10 2nd step

(R)-tert-butyl (1-ethoxy-3-oxobutan-2-yl)carbamate

$^1\text{H-NMR}$ (CDCl_3) δ : 5.55-5.45 (1H, br), 4.37-4.28 (1H, m), 3.92-3.41 (4H, m), 2.21 (3H, s), 1.47 (9H, d, $J = 7.3$ Hz), 1.17 (3H, q, $J = 7.3$ Hz).

MS (ESI m/z): 132 ($M+H-\text{Boc}$)

15 RT (min): 1.22

[0391]

3rd step

tert-Butyl ((2R)-3-hydroxy-1-ethoxybutan-2-yl)carbamate

MS (ESI, m/z): 234 ($M+H$)

20 RT (min): 1.07

4th step

(3R)-3-(((tert-butoxycarbonyl)amino)-4-ethoxybutan-2-yl)-4-nitrobenzoate

MS (ESI m/z): 383 ($M+H$)

RT (min): 1.71

25 [0392]

5th step

tert-Butyl ((2R)-3-hydroxy-1-ethoxybutan-2-yl)carbamate

MS (ESI m/z): 234 ($M+H$)

RT (min): 1.06

6th step

tert-Butyl

((2S,3S)-3-((1,3-dioxoisindolin-2-yl)-1-ethoxybutan-2-yl)carbamate

¹H-NMR (CDCl₃) δ: 7.86-7.81 (2H, m), 7.73-7.67 (2H, m), 5.10-4.95 (1H, m),
 4.54-4.35 (2H, m), 3.47-3.25 (4H, m), 1.46 (9H, s), 1.32 (3H, d, J = 6.3 Hz),
 1.02 (3H, t, J = 6.9 Hz).

MS (ESI m/z): 363 (M+H)

RT (min): 1.63

[0393]

10 7th step

2-((2S,3S)-3-amino-4-ethoxybutan-2-yl)isoindolin-1,3-dione

MS (ESI m/z): 263 (M+H),


RT (min): 0.77

[0394]

15 Reference Example 111

The following compound was obtained as described in Reference Example 109.

[Formula 131]



20 [0395]

1st step

2-Chloro-6-(((2S,3S)-3-(1,3-dioxoisindolin-2-yl)-1-ethoxybutan-2-yl)amino)-5-fluoronicotinonitrile

MS (ESI m/z): 417 (M+H)

25 RT (min): 1.41

2nd step

6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-2-chloro-5-fluoronicotinonitrile

MS (ESI m/z): 287 (M+H)

30 RT (min): 0.82

[0396]

3rd step

tert-Butyl

((2S,3S)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate

¹H-NMR (CDCl₃) δ: 7.31 (1H, d, J = 9.6 Hz), 6.14 (1H, d, J = 6.9 Hz), 5.25 (1H, d, J = 8.9 Hz), 4.33-4.19 (1H, dd, J = 7.9, 3.6 Hz), 4.11-3.93 (1H, m),
 5 3.79 (1H, dd, J = 9.9, 2.6 Hz), 3.62-3.43 (4H, m), 1.44 (9H, s), 1.31-1.19 (6H, m).

MS (ESI m/z): 387 (M+H)


RT (min): 1.71

[0397]

10 Reference Example 112

The following compound was obtained as described in Reference Example 108.

[Formula 132]



15 [0398]

(R)-tert-butyl

(1-(methoxy(methyl)amino)-4-(methylthio)-1-oxobutan-2-yl)carbamate

MS (ESI m/z): 293 (M+H)

RT (min): 1.24

20 (3R)-3-((tert-butoxycarbonyl)amino)-5-(methylthio)pentan-2-yl

4-nitrobenzoate

MS (ESI m/z): 399 (M+H)

RT (min): 1.78

tert-Butyl

25 ((3R,4S)-4-(1,3-dioxoisindolin-2-yl)-1-(methylthio)pentan-3-yl)carbamate

MS (ESI m/z): 379 (M+H)

RT (min): 1.60

2-((2S,3R)-3-amino-5-(methylthio)pentan-2-yl)isoindolin-1,3-dione

MS (ESI m/z): 279 (M+H)


RT (min): 0.75

[0399]

Reference Example 113

5 The following compound was obtained as described in Reference Example 109.

[Formula 133]



[0400]

10 1st step

2-Chloro-6-(((3R,4S)-4-(1,3-dioxoisindolin-2-yl)-1-(methylthio)pentan-3-yl)amino)-5-fluoronicotinonitrile

MS (ESI m/z): 433 (M+H)

RT (min): 1.67

15 2nd step

6-(((3R,4S)-4-amino-1-(methylthio)pentan-3-yl)amino)-2-chloro-5-fluoronicotinonitrile

MS (ESI m/z): 303 (M+H)

RT (min): 0.85

20 [0401]

3rd step

tert-Butyl

((2S,3R)-3-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate

25 MS (ESI m/z): 403 (M+H)


RT (min): 1.70

[0402]

Reference Example 114

30 The following compound was obtained as described in Reference Example 108.

[Formula 134]



[0403]

(2R)-2-((tert-butoxycarbonyl)amino)-1-cyclopropylpropyl-4-nitrobenzoate

MS (ESI m/z): 365 (M+H)

5 RT (min): 1.78

tert-Butyl

((1S,2R)-1-cyclopropyl-1-(1,3-dioxoisindolin-2-yl)propan-2-yl)carbamate

MS (ESI m/z): 399 (M+H)

RT (min): 1.73

10 2-((1S,2R)-2-amino-1-cyclopropylpropyl)isoindolin-1,3-dione

MS (ESI m/z): 299 (M+H)


RT (min): 0.73

[0404]

Reference Example 115

15 The following compound was obtained as described in Reference Example 109.

[Formula 135]



[0405]

20 2-Chloro-6-(((1S,2R)-1-cyclopropyl-1-(1,3-dioxoisindolin-2-yl)propan-2-yl)amino)-5-fluoronicotinonitrile

MS (ESI m/z): 400 (M+H)

RT (min): 1.74

25 6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-2-chloro-5-fluoronicotinonitrile

MS (ESI m/z): 270 (M+H)

RT (min): 0.84

tert-Butyl

((1S,2R)-2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate

MS (ESI m/z): 370 (M+H)


RT (min): 1.75

5 [0406]

Reference Example 116

The following compound was obtained with reference to Archiv der Pharmazie (Weinheim, Germany), 2004, vol. 337, #12 pp. 654-667.

[Formula 136]




(S,Z)-N-(2-((tert-butoxycarbonyl)amino)butylidene)-1-phenylmethaneamine oxide

[0407]

Reference Example 117

15 [Formula 137]



[0408]

1st step

20 Methylmagnesium bromide (3M diethyl ether solution, 0.86 ml) was added dropwise to a THF (5 ml) solution containing (S,Z)-N-(2-((tert-butoxycarbonyl)amino)butylidene)-1-phenylmethaneamine oxide (250 mg) at -50°C , followed by stirring at -50°C to -35°C for 2 hours. Further, methylmagnesium bromide (3M diethyl ether solution, 0.86 ml) was added dropwise to the reaction solution, followed by stirring at -45°C to -40°C for 1 hour. A saturated ammonium chloride aqueous solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 19:1 to 4:1). tert-Butyl ((3S,4R)-4-(benzyl(hydroxy)amino)pentan-3-yl)carbamate (39 mg) was thus obtained.

$^1\text{H-NMR}$ (CDCl_3 , 300MHz) δ : 7.39-7.18 (m, 5H), 6.70 (s, 1H), 4.43 (d, 1H, J =

10.2Hz), 4.11 (d, 1H, J = 13.9Hz), 4.10-3.97 (m, 1H), 3.64 (d, 1H, J = 13.9Hz), 2.78-2.68 (m, 1H), 1.47 (s, 9H), 1.44-1.26 (m, 2H), 1.03-0.94 (m, 9H)

[0409]

2nd step

5 An MeOH (20 ml) solution containing tert-butyl ((3S,4R)-4-(benzyl (hydroxy)amino)pentan-3-yl)carbamate (39 mg) was subjected to a hydrogenation reaction (45°C; 100 bar; flow rate: 1ml/min; 20% Pd(OH)₂/C) using H-cube™. Then, the solvent was distilled away under reduced pressure. Colorless oily matter of tert-butyl
10 ((3S,4R)-4-aminopentan-3-yl)carbamate (27 mg) was thus obtained.

3rd step

The following compound was obtained as described in the 7th step in Reference Example 417.

tert-Butyl

15 ((2R,3S)-2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)pentan-3-yl)carbamate


MS (ESI m/z): 357 (M+H), 355 (M-H)

[0410]

Reference Example 118

20 The following compound was obtained as described in Reference Example 117.

[Formula 138]



[0411]

25 tert-Butyl ((3S,4R)-4-(benzyl (hydroxy)amino)hexan-3-yl)carbamate

¹H-NMR (CDCl₃, 300MHz) δ: 7.40-7.20 (m, 5H), 5.88 (s, 1H), 4.62 (d, 1H, J = 9.6Hz), 4.07 (d, 1H, J = 13.9Hz), 4.01-3.88 (m, 1H), 3.73 (d, 1H, J = 13.9Hz), 2.59-2.50 (m, 1H), 1.69-1.32 (m, 4H), 1.45 (s, 9H), 1.05 (t, 3H, J = 7.6Hz), 0.98 (t, 3H, J = 7.3Hz)


30 tert-Butyl ((3S,4R)-4-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)hexan-3-yl)carbamate

MS (ESI m/z): 371 (M+H), 369 (M-H)

[0412]

Reference Example 119

[Formula 139]



[0413]

1st, 2nd, and 3rd steps

5 (S)-tert-butyl

(1-(methoxy(methyl)amino)-1-oxopropan-2-yl)carbamate (10 g) was added to (2-methyl-1-propen-1-yl)magnesium bromide (0.5 M in THF) (258.3 ml), followed by stirring at 50°C for 40 minutes. The reaction solution was adjusted to room temperature and poured into a 10% citric acid aqueous solution, followed by extraction with ethyl acetate. The organic layers were washed with water and saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. TFA (20 ml) was added to the obtained residue, followed by stirring at room temperature for 30 minutes. Next, the solvent was distilled away under reduced pressure. DMF (30 ml), potassium carbonate (13.8 g), and benzyl bromide (10.7 ml) were added to the obtained residue, followed by stirring at 70°C for 50 minutes. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with water and saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2). A yellow liquid of (S)-2-(dibenzylamino)-5-methyl-4-hexen-3-one (4.46 g) was thus obtained.

MS (ESI m/z): 308 (M+H)

25 RT (min): 1.56

[0414]

4th step

A methanol solution (10 ml) containing sodium borohydride (1.5 g) and (S)-2-(dibenzylamino)-5-methyl-4-hexen-3-one (4.2 g) obtained in the 3rd step were added to an MeOH (30 ml) solution containing cerium chloride (10 g), followed by stirring for 7 hours. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with water and saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2). A yellow liquid of (2S)-2-(dibenzylamino)-5-methyl-4-hexen-3-ol (3.5 g) was thus obtained.

MS (ESI m/z): 310 (M+H)

RT (min): 1.06

[0415]

5th step

A CH₂Cl₂ (5 ml) solution containing diiodomethane (3.5 ml) and a methylene chloride (5 ml) solution containing (2S)-2-amino-5-methyl-4-hexen-3-ol (2.7 g) obtained in the 4th step were added to a methylene chloride (40 ml) solution containing diethyl zinc (43.6 ml), followed by stirring at room temperature for 15 hours. Methanol and sodium hydrogen carbonate were added to the reaction solution, followed by filtration through Celite. The filtrate was extracted with ethyl acetate. The organic layers were washed with water and saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2). (2S)-2-(dibenzylamino)-1-(2,2-dimethylcyclopropyl)propan-1-ol (2.4 g) was thus obtained.

MS (ESI m/z): 324 (M+H)

RT (min): 1.13

[0416]

6th and 7th steps

Formic acid (2.4 ml) and 10%Pd/C (0.4 g) were added to an ethanol solution (60 ml) containing (1S,2S)-2-(dibenzylamino)-1-(2,2-dimethylcyclopropyl)propan-1-ol (2.4 g) obtained in the 5th step, followed by stirring at 90°C for 5 hours. The reaction solution was filtered through Celite and the filtrate was distilled

away under reduced pressure. Diisopropyl ethylamine (1.5 ml) and di-tert-butyl carbonate (1.75 g) were added to a THF (5 ml) solution containing the obtained residue, followed by stirring at room temperature for 40 minutes. Water and a 10% citric acid aqueous solution were added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with water and saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2). A yellow liquid of tert-butyl ((2S)-1-(2,2-dimethylcyclopropyl)-1-hydroxypropan-2-yl)carbamate (0.45 g) was thus obtained.

MS (ESI m/z): 244 (M+H)

RT (min): 1.36

[0417]

8th step

Phthalimide (0.147 g), triphenylphosphine (0.34 g), and diisopropyl azodicarboxylate (0.684 ml) were added to a THF (5 ml) solution containing tert-butyl

((2S)-1-(2,2-dimethylcyclopropyl)-1-hydroxypropan-2-yl)carbamate (0.20 g), followed by stirring at room temperature for 1.5 hours. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2). A yellow liquid of tert-butyl ((2S)-1-(2,2-dimethylcyclopropyl)-1-(1,3-dioxoisindolin-2-yl)propan-2-yl)carbamate (22.4 mg) was thus obtained.

MS (ESI m/z): 373 (M+H)

RT (min): 1.80

[0418]

9th and 10th steps

Hydrazine monohydrate (0.4 ml) was added to an ethanol solution (5 ml) containing tert-butyl ((2S)-1-(2,2-dimethylcyclopropyl)-1-(1,3-dioxoisindolin-2-yl)propan-2-yl)carbamate (22.4 mg), followed by stirring at 90°C for 3 hours. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under

reduced pressure. 2,6-Dichloro-5-fluoronicotinonitrile (15.3 mg) and diisopropyl ethylamine (0.1 ml) were added to the obtained residue, followed by stirring at 70°C for 50 minutes. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed
 5 with water and saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2). A yellow liquid of tert-butyl ((1R,2S)-1-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-1-((S)-2,2-dimet
 10 hylcyclopropyl)propan-2-yl)carbamate (6 mg) was thus obtained.


MS (ESI m/z): 397 (M+H)

RT (min): 1.89

[0419]

Reference Example 120

15 [Formula 140]



[0420]

1st and 2nd steps

A CH₂Cl₂ (125 ml) solution containing 1,4-hexadiene (8.0 g) was
 20 added to a solution comprising sodium hydrogen carbonate (12.6 g) and water (75 ml) at room temperature and mCPBA (16.4 g) was further added under ice cooling, followed by stirring for 1 hour and stirring at room temperature 10 hours. A 5% sodium thiosulfate aqueous solution was added to the reaction solution, followed by extraction with methylene chloride. The organic
 25 layers were washed with a saturated sodium hydrogen carbonate aqueous solution and saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. n-Butanol (20 ml) was added to the obtained residue. Next, (R)-(-)-1-phenylethylamine was added, followed by stirring at 90°C for 5 hours. The solvent was distilled away
 30 under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2). A yellow oily matter of (1S,6S)-6-(((R)-1-phenylethyl)amino)cyclohexyl-3-enol (2.6 g) was

thus obtained.

[0421]

3rd step

Potassium carbonate (1.9 g) and benzyl bromide (1.53 ml) were added
 5 to a DMF (5 ml) solution containing
 (1S,6S)-6-(((R)-1-phenylethyl)amino)cyclohexyl-3-enol (2.5 g) obtained in
 the 2nd step, followed by stirring at 90°C for 1 hour. Water was added to
 the reaction solution, followed by extraction with ethyl acetate. The organic
 layers were washed with water and saturated saline and dried over anhydrous
 10 sodium sulfate. The solvent was distilled away under reduced pressure and
 the obtained residue was purified by silica gel chromatography (n-hexane :
 ethyl acetate = 1:0 to 1:0.2). Yellow oily matter of (1S, 6S)-6-(benzyl
 ((R)-1-phenylethyl)amino)cyclohexan-3-ol (1.6 g) was thus obtained.


MS (ESI m/z): 308 (M+H)

15 RT (min): 0.98

[0422]

Reference Example 121

[Formula 141]



20

[0423]

1st step

Phthalimide (0.63 g), triphenylphosphine (1.3 g), and DIAD (1.9M in
 toluene solution) (2.6 ml) were added to a THF (50 ml) solution containing
 25 (1S, 6S)-6-(benzyl ((R)-1-phenylethyl)amino)cyclohexan-3-ol (1.2 g),
 followed by stirring at room temperature for 15 hours. The solvent was
 distilled away under reduced pressure and the obtained residue was purified
 by silica gel chromatography (hexane: ethyl acetate = 1:0 to 5:1). Yellow

oily matter of 2-((1R,6S)-6-(benzyl ((R)-1-phenylethyl)amino)cyclohex-3-en-1-yl)isoindolin-1,3-dione (1.5 g) was thus obtained.

MS (ESI m/z): 438 (M+H)

5 RT (min): 2.21

2nd step

A methylene chloride (3 ml) solution containing diiodomethane (0.2 ml) and a methylene chloride solution (3 ml) containing 2-((1R,6S)-6-(benzyl ((R)-1-phenylethyl)amino)-3-cyclohexen-1-yl)isoindolin-1,3-dione (0.36 g) 10 obtained in the 1st step were added to a methylene chloride solution (10 ml) containing diethyl zinc (1M in hexane) (2.47 ml), followed by stirring for 15 hours. Methanol and sodium hydrogen carbonate were added to the reaction solution. The reaction solution was filtered through Celite. The filtrate was extracted with ethyl acetate. The organic layers were washed with water 15 and saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2).

A 2-((3R,4S)-4-(benzyl ((R)-1-phenylethyl)amino)bicyclo[4.1.0]heptan-3-yl)isoindolin-1,3-dione 20 (0.20 g) was thus obtained.

MS (ESI m/z): 451 (M+H)

RT (min): 2.27

[0424]

3rd and 4th steps

25 Ammonium formate (0.084 g) and 10%Pd/C (0.1 g) were added to an ethanol solution (3 ml) containing 2-((3R,4S)-4-(benzyl ((R)-1-phenylethyl)amino)bicyclo[4.1.0]heptan-3-yl)isoindolin-1,3-dione (0.1 g) obtained in the 2nd step, followed by stirring 90°C for 8 hours. The reaction solution was filtered through Celite and the filtrate was distilled 30 away under reduced pressure. Diisopropyl ethylamine (0.2 ml) and di-tert-butyl carbonate (0.1 g) were added to a dimethylformamide solution (1 ml) containing the obtained residue, followed by stirring at room temperature for 45 minutes. Water and a 10% citric acid aqueous solution were added to the reaction solution, followed by extraction with ethyl acetate. The organic 35 layers were washed with water and saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure

and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 1:0 to 1:0.2). Yellow oily matter of tert-butyl (3S,4R)-4-(1,3-dioxoisindolin-2-yl)bicyclo[4.1.0]heptan-3-yl)carbamate (0.028 g) was thus obtained.

5 MS (ESI m/z): 357 (M+H)

RT (min): 1.81

[0425]

5th and 6th steps

tert-Butyl

(3S,4R)-4-(1,

10 3-dioxoisindolin-2-yl)bicyclo[4.1.0]heptan-3-yl)carbamate (0.028 g)

obtained in the 4th step and an ethanol (5 ml) solution containing hydrazine monohydrate (0.2 ml) was stirred at 90°C for 48 hours. The solvent was distilled away under reduced pressure and an insoluble precipitate was removed. Then, DMF (1 ml) containing 2,6-dichloro-5-fluoronicotinonitrile

15 (0.02 g) and DIPEA (0.1 ml) were added to the obtained oily matter, followed by stirring for 4 hours. The reaction solution was adjusted to room temperature and water was added to the reaction solution, followed by extraction with ethyl acetate and washing with saturated saline. The organic layers were dried over anhydrous sodium sulfate. The obtained residue was
20 purified by silica gel chromatography (n-hexane : ethyl acetate = 10:1 to 1:1).

A white solid of tert-butyl (3S,4R)-4-(6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)bicyclo[4.1.0]heptan-3-yl)carbamate (0.008 g) was thus obtained.

¹H-NMR (CDCl₃) δ: 7.23 (1H, d, J = 9.9 Hz), 6.39 (1H, br), 4.48 (1H, d, J =
25 7.9 Hz), 3.86-3.72 (1H, m), 3.47-3.33 (1H, m), 2.75-2.65 (1H, m), 2.33 (1H, dd, J = 12.6, 4.6 Hz), 1.80 (1H, td, J = 12.6, 4.6 Hz), 1.38 (9H, s), 1.29-0.69 (4H, m), 0.20-0.11 (1H, m)


MS (ESI m/z): 381 (M+H)

RT (min): 1.8

30 [0426]

Reference Example 122

[Formula 142]



[0427]

1st step

PPh₃ (412 mg), phthalimide (252 mg), and diethyl azodicarboxylate (40% in toluene solution) (0.712 ml) were added to a THF (5 ml) solution containing benzyl 3-((tert-butoxycarbonyl)amino)-4-hydroxypiperidin-1-carboxylate (500 mg), followed by stirring at room temperature for 4 hours. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane: ethyl acetate = 9:1 to 15:7). Colorless oily matter of benzyl 3-((tert-butoxycarbonyl)amino)-4-(1,3-dioxoisindolin-2-yl)piperidin-1-carboxylate (205 mg) was thus obtained.

2nd step

Ammonium formate (419 mg) and 10% Pd/C (84 mg) were added to an ethyl acetate/MeOH (4 ml/4 ml) solution containing benzyl 3-((tert-butoxycarbonyl)amino)-4-(1,3-dioxoisindolin-2-yl)piperidin-1-carboxylate (419 mg) obtained in the 1st step, followed by stirring at 60°C for 1 hour. The reaction solution was cooled to room temperature and insoluble matter was removed through Celite. Filter cake was washed with ethyl acetate and water. Subsequently, the filtrate was mixed with wash liquid and sodium chloride was added to the mixture. The organic layers were separated and washed with saturated saline. The organic layers were dried over anhydrous sodium sulfate and the solvent was distilled away under reduced pressure. A white solid of tert-butyl (4-(1,3-dioxoisindolin-2-yl)piperidin-3-yl)carbamate (280 mg) was thus obtained.

MS (ESI m/z): 346(M+H)

RT (min): 0.89

[0428]

3rd step

5 Sodium hydrogen carbonate (341 mg) and acetyl chloride (0.086 ml) were added to a THF/water (2 ml/2 ml) solution containing tert-butyl (4-(1,3-dioxoisindolin-2-yl)piperidin-3-yl)carbamate (280 mg) obtained in the 2nd step under ice cooling, followed by stirring at room temperature for 0.5 hours. Ethyl acetate was added to the reaction solution. The organic
10 layers were washed with saturated saline and dried over anhydrous sodium sulfate. Next, the solvent was distilled away under reduced pressure. A white solid of tert-butyl (1-acetyl-4-(1,3-dioxoisindolin-2-yl)piperidin-3-yl)carbamate (301 mg) was thus obtained.

15 MS (ESI m/z): 388(M+H)

RT (min): 1.17

4th step

Hydrazine monohydrate (1 ml) was added to an ethanol (5 ml) solution containing tert-butyl
20 (1-acetyl-4-(1,3-dioxoisindolin-2-yl)piperidin-3-yl)carbamate (301 mg) obtained in the 3rd step, followed by heating and stirring at 50°C for 0.5 hours. Ethyl acetate was added to the reaction solution. The reaction solution was washed with water and saturated saline and dried over anhydrous sodium sulfate. Next, the solvent was distilled away under reduced pressure.
25 Colorless oily matter of tert-butyl (1-acetyl-4-aminopiperidin-3-yl)carbamate (147 mg) was thus obtained.

MS (ESI m/z): 258(M+H)

RT (min): 0.50

[0429]

30 5th step

Triethylamine (0.096 ml) and 2,6-dichloro-5-fluoronicotinonitrile (109 mg) were added to a DMSO (2ml) solution containing tert-butyl (1-acetyl-4-aminopiperidin-3-yl)carbamate (147 mg) obtained in the 4th step, followed by stirring at room temperature for 1 hour. Ethyl acetate was added
35 to the reaction solution. The reaction solution was washed with water and saturated saline and dried over anhydrous sodium sulfate. Next, the solvent

was distilled away under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane: ethyl acetate = 4:1 to 2:3).

A light yellow solid of tert-butyl (1-acetyl-4-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)piperidin-3-yl)carbamate (91 mg) was thus obtained.

MS (ESI m/z): 412, 414 (M+H)


RT (min): 1.28

[0430]

Reference Example 123

The following compound was obtained with reference to Journal of Medicinal Chemistry, 2010, 53, 7107.

[Formula 143]




Benzyl 3-((tert-butoxycarbonyl)amino)-4-hydroxypiperidin-1-carboxylate

[0431]

Reference Example 124

[Formula 144]



[0432]

1st step

Triphenylphosphine (840 mg), diisopropyl azodicarboxylate (40% in toluene) (1.68 ml), and DPPA (0.86 ml) were added to a THF (9.3 ml) solution containing

benzyl 3-((tert-butoxycarbonyl)amino)-4-hydroxypiperidin-1-carboxylate (930 mg)

under ice cooling, followed by stirring at 50°C for 2 hours. The reaction solution was cooled to room temperature and water was added to the reaction

solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane : ethyl acetate = 9:1 to 17:3). Light yellow oily matter of benzyl 4-azide-3-((tert-butoxycarbonyl)amino)piperidin-1-carboxylate (780 mg) was thus obtained.

2nd step

Triphenylphosphine (820 mg) was added to a THF/water (7.8 ml/0.78 ml) solution containing benzyl 4-azide-3-((tert-butoxycarbonyl)amino)piperidin-1-carboxylate (780 mg) obtained in the 1st step, followed by heating and stirring at 80°C for 3 hours. The reaction solution was cooled to room temperature. Water and a 2M hydrochloric acid aqueous solution were added to the reaction solution so as to acidify the reaction solution. The reaction solution was washed with ethyl acetate. Next, the aqueous layers were collected and a 5M sodium hydroxide aqueous solution was added so as to alkalify the aqueous layers. The aqueous layers were subjected to extraction twice with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate and the solvent was distilled away under reduced pressure. Yellow oily matter of benzyl 4-amino-3-((tert-butoxycarbonyl)amino)piperidin-1-carboxylate (470 mg) was thus obtained.

[0433]

3rd step

Triethylamine (0.22 ml) and 2,6-dichloro-5-fluoronicotinonitrile (470 mg) were added to a DMSO (2.4 ml) solution containing benzyl 4-amino-3-((tert-butoxycarbonyl)amino)piperidin-1-carboxylate (470 mg) obtained in the 2nd step, followed by stirring at room temperature for 0.5 hours. Ethyl acetate and water were added to the reaction solution for extraction. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane: ethyl acetate = 19:1 to 15:3) and a white solid of benzyl

3-((tert-butoxycarbonyl)amino)-4-((6-chloro-5-cyano-3-fluoropyridin-2-yl)a

mino)piperidin-1-carboxylate (500 mg) was thus obtained.

MS (ESI m/z): 504, 506 (M+H)

RT (min): 1.88

[0434]

5 4th step

Benzyl

3-((tert-butoxycarbonyl)amino)-4-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)piperidin-1-carboxylate (50 mg) obtained in the 3rd step was mixed with TFA (1 ml), followed by stirring at room temperature for 15 minutes. TFA was distilled away under reduced pressure. Water (5 ml), a 5M sodium hydroxide aqueous solution (1 ml), and chloroform were added to the obtained residue and the organic layers were collected. Subsequently, the organic layers were washed with saturated saline and dried over anhydrous sodium sulfate and the solvent was distilled away under reduced pressure.

15 Next, DMF (1 ml) and phthalic anhydride (29 mg) were added to the obtained residue, followed by heating and stirring at 150°C for 1 hour. The reaction solution was cooled to room temperature. Ethyl acetate was added to the reaction solution. The reaction solution was washed with a saturated sodium hydrogen carbonate aqueous solution and saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. A white solid of benzyl 4-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-3-(1,3-dioxoisindolin-2-yl)piperidin-1-carboxylate (37 mg) was thus obtained.


MS (ESI m/z): 534, 536 (M+H)

25 RT (min): 1.75

[0435]

Reference Example 125

[Formula 145]



[0436]

1st step

3-Amino-3-methylpyridine (9 mg), cesium carbonate (45 mg),
 5 Pd₂(dba)₃ (10 mg), and Xantphos (12 mg) were added to a dioxane (3 ml)
 solution containing benzyl
 4-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)-3-(1,3-dioxoisindolin-2-yl)
 piperidin-1-carboxylate (37 mg), followed by microwave irradiation
 (InitiatorTM, 160°C, 10 minutes, 2.45 GHz, 0-240W). The obtained residue
 10 was filtered through Celite and filter cake was washed with ethyl acetate.
 Subsequently, the solvent was distilled away from the filtrate under reduced
 pressure. The obtained residue was purified by silica gel column
 chromatography (hexane : ethyl acetate = 4:1 to 1:1). Yellow oily matter of
 benzyl

15 4-((5-cyano-3-fluoro-6-((5-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-3-
 (1,3-dioxoisindolin-2-yl)piperidin-1-carboxylate (28 mg) was thus obtained.
 MS (ESI m/z): 606 (M+H)

RT (min): 1.30

2nd step

20 Ammonium formate (6 mg) and 10% Pd/C (6 mg) were added to an
 ethyl acetate/MeOH (1 ml/1 ml) solution containing benzyl
 4-((5-cyano-3-fluoro-6-((5-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-3-
 (1,3-dioxoisindolin-2-yl)piperidin-1-carboxylate (37 mg) obtained in the 1st
 step in a nitrogen atmosphere, followed by heating and stirring at 70°C for 1
 25 hour. Next, ammonium formate (30 mg) and 10% Pd/C (30 mg) were added
 to the solution, followed by heating and stirring at 70°C for 1 hour. The

reaction solution was cooled to room temperature and insoluble matter was removed by filtration through Celite. Filter cake was washed with ethyl acetate. The resulting filtrate was mixed with wash liquid. The obtained organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. A white solid of 6-((3-(1,3-dioxoisindolin-2-yl)piperidin-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinonitrile (15 mg) was thus obtained.

[0437]

10 3rd step

Sodium hydrogen carbonate (13 mg) and acetyl chloride (0.005 ml) were added to a THF/water (0.5 ml/0.5 ml) solution containing 6-((3-(1,3-dioxoisindolin-2-yl)piperidin-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinonitrile (15 mg) obtained in the 2nd step under ice cooling, followed by stirring at room temperature for 0.5 hours. Ethyl acetate was added to the reaction solution. The obtained organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. Then, the solvent was distilled away under reduced pressure. Yellow oily matter of 6-((1-acetyl-3-(1,3-dioxoisindolin-2-yl)piperidin-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinonitrile (12 mg) was thus obtained.

MS (ESI m/z): 514(M+H)

RT (min): 0.86

4th step

Hydrazine monohydrate (0.1 ml) was added to an EtOH (1 ml) solution containing 6-((1-acetyl-3-(1,3-dioxoisindolin-2-yl)piperidin-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinonitrile (12 mg) obtained in the 3rd step at room temperature, followed by heating and stirring at 50°C for 0.5 hours. Ethyl acetate was added to the reaction solution. The obtained organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. Next, ethyl acetate (1 ml) and 4M hydrochloric acid/1,4-dioxane (0.008 ml) were added to the obtained residue, followed by stirring at room temperature for 0.5 hours. The solvent was removed under reduced pressure and the obtained solid was washed with ethyl acetate. A yellow solid of 6-((1-acetyl-3-aminopiperidin-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl

)amino)nicotinonitrile hydrochloride (6 mg) was thus obtained.

MS (ESI m/z): 384 (M+H)


RT (min): 0.54

[0438]

5 Reference Example 126

The following compound was obtained with reference to J. Org. Chem., 1985, 50, 4154-4155 and Synth. Commun., 1992, 22, 3003-3012.

[Formula 146]




10 tert-Butyl ((1S,2S)-2-hydroxycyclohexyl)carbamate

[0439]

Reference Example 127

The following compound was obtained with reference to US 2003/0119855 A1.

15 [Formula 147]




tert-Butyl ((1S,2R)-2-aminocyclohexyl)carbamate

¹H NMR (CDCl₃, 300Mz) : 1.3-1.7 (17H, m), 2.9-3.0 (1H, m), 3.6-3.7 (1H, m), 4.9-5.0 (1H, m).

20 [0440]

Reference Example 128

[Formula 148]



Triphenylphosphine (14.3 g) was added to a THF (190 ml) solution containing tert-butyl ((1S,2S)-2-hydroxycyclohexyl)carbamate (10.0 g), followed by ice cooling. Diethyl azodicarboxylate (40% in toluene) (24.3 g)


and DPPA (15.3 g) were added dropwise to the reaction solution, followed by stirring at room temperature for 1 hour. The reaction solution was left overnight. The solvent was distilled away under reduced pressure. Water was added and then a 20% sodium hydroxide aqueous solution was added. Then, the organic layers were collected. Water (30 ml) was added to the obtained organic layers, followed by heating to 60°C. A THF (40 ml) solution containing triphenylphosphine (14.3 g) was added dropwise, followed by reflux for 2.5 hours. The solvent was distilled away under ordinary pressure. Toluene was added and the pH was adjusted with 3M hydrochloric acid to pH = 1 or less. Then, the resulting aqueous layers were collected, ethyl acetate was added, and the pH was adjusted with a 20% sodium hydroxide aqueous solution to pH 12. The organic layers were collected and dried over anhydrous sodium sulfate. Light yellow oily matter of tert-butyl ((1S,2R)-2-aminocyclohexyl)carbamate (5.24 g) was thus obtained.

[0441]

Reference Example 129

The following compound was obtained with reference to ChemCatChem, 2010, 2, 1215-1218 (optical resolution by lipase).


[Formula 149]



[0442]

Reference Example 130

[Formula 150]



[0443]


Potassium carbonate (3.62 g) and 2,6-dichloro-5-fluoronicotinonitrile (5.00 g) were added to a THF (50 ml) solution containing tert-butyl ((1S,2R)-2-aminocyclohexyl)carbamate (5.61 g), followed by reflux at 60°C for 8 hours. Then, the solvent was distilled away at 70°C. 1,4-dioxane (100 ml) was added to the resulting solution, followed by stirring at 100°C for 10 hours. The reaction solution was adjusted to room temperature. Ethyl

acetate and 2M hydrochloric acid were added to the reaction solution. The organic layers were collected. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 9:1 to 2:1). Diisopropyl ether was added to the obtained oily matter and a solid precipitate was collected by filtration. A white solid of tert-butyl ((1S,2R)-2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate (5.77 g) was thus obtained.

[0444]

Reference Example 131

[Formula 151]




Water (100 ml) was added to (1R, 2R)-2-aminocyclohexan-1-ol (47.8 g) and a THF (200 ml) solution containing di-tert-butyl dicarbonate (95.1 g) was added dropwise, followed by stirring at room temperature 3 hours. The organic layers were collected at 36°C, followed by cooling. Then, a solid precipitate was collected by filtration and washed with hexane and ethyl acetate. A white solid of tert-butyl ((1R,2R)-2-hydroxycyclohexyl)carbamate (73.7 g) was thus obtained.

[0445]

Reference Example 132

[Formula 152]



[0446]

1st step

Diethyl azodicarboxylate (40% in toluene) (36.4 g) and DPPA (23.0 g)

were added dropwise to a THF (190 ml) solution containing tert-butyl ((1R,2R)-2-hydroxycyclohexyl)carbamate (15.0 g) and triphenylphosphine (21.9 g) under ice cooling, followed by stirring at room temperature for 7 hours. The solvent of the reaction solution was distilled away under reduced pressure and the obtained residue was purified by silica gel chromatography (n-hexane : ethyl acetate = 4:1). Light yellow oily matter of tert-butyl ((1R,2S)-2-azide cyclohexyl)carbamate (21.1 g) was thus obtained.

2nd step

p-Toluene sulfonic acid monohydrate (13.3 g) was added to a 2-propanol (100 ml) solution containing tert-butyl ((1R,2S)-2-azide cyclohexyl)carbamate (21.1 g), followed by reflux for 40 minutes. After cooling, toluene and water were added, and aqueous layers were collected. Isopropyl acetate was added to the obtained aqueous layers. A 20% sodium hydroxide aqueous solution was added to adjust the pH to pH 12-13. Then, the organic layers were collected. The obtained organic layers were dried over anhydrous sodium sulfate and the solvent was distilled away under reduced pressure. (1R,2S)-2-azide cyclohexan-1-amine (8.7 g) was thus obtained.

[0447]

3rd step

(Liquid A)

Potassium carbonate (0.87 g) was added to a DMSO (5 ml) solution containing 2,6-dichloro-5-fluoronicotinonitrile (1.00 g) at room temperature. The solution was heated to 50°C and a DMSO (0.5 ml) solution containing (1R,2S)-2-azide cyclohexan-1-amine (0.73 g) was added to the solution. Subsequently, a DMSO (0.5 ml) solution containing (1R,2S)-2-azide cyclohexan-1-amine (0.22 g) was added, followed by stirring for 20 minutes. Further, a DMSO (0.5 ml) solution containing (1R,2S)-2-azide cyclohexan-1-amine (0.22 g) was added, followed by stirring for 20 minutes.

(Liquid B)

A toluene (25 ml) solution containing sodium carbonate (4.9 g) and (1R,2S)-2-azide cyclohexan-1-amine (7.5 g) was added dropwise to a DMSO (15 ml) solution containing 2,6-dichloro-5-fluoronicotinonitrile (7.3 g), followed by stirring at 45°C for 2.5 hours.

4th step

Water, 6M hydrochloric acid, and toluene were added to a mixture of

liquid A and liquid B. The organic layers were collected. The solvent was distilled away under reduced pressure. THF (25 ml) and water (30 ml) were added to the obtained residue, followed by heating to 60°C. A THF (25 ml) solution containing triphenylphosphine (11.4 g) was added dropwise, followed by reflux for 3 hours. After cooling, toluene, water and 6M hydrochloric acid were added, and aqueous layers were collected. Isopropyl acetate and a 20% sodium hydroxide aqueous solution were added to the obtained aqueous layers, and organic layers were collected. The solvent was distilled away under reduced pressure. A residue was thus obtained.

[0448]

5th step

An ethyl acetate (15 ml) solution containing di-tert-butyl dicarbonate (8.1 g) was added dropwise to an ethyl acetate (50 ml) solution containing the residue obtained in the 4th step, followed by stirring at room temperature for 40 minutes. Next, an ethyl acetate (5 ml) solution containing di-tert-butyl dicarbonate (0.81 g) was added dropwise, followed by stirring at room temperature for 9 hours. Water and ethyl acetate were added to the reaction solution, and organic layers were collected. The solvent was distilled away under reduced pressure. 2-Propanol was added to the obtained residue. The solvent was distilled away under reduced pressure. Water and seed crystals were added, followed by stirring. Further, 2-propanol and water were added and a solid precipitate was collected by filtration. A white solid of tert-butyl ((1S,2R)-2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate (7.5 g) was thus obtained.

[0449]

tert-Butyl

((1S,2R)-2-((6-chloro-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate obtained in the 5th step in Reference Example 132 was analyzed under the following conditions. It was confirmed that an optically-active substance was synthesized.

[0450]

<Chiral HPLC conditions>

Apparatus: SHIMAZU 10A series

Column: Daicel CHIRALPAK IC-3

Mobile phase: n-Hex/IPA/i-PrNH₂=95/5/0.1

Flow rate: 1.0 mL/min

Temperature: 40°C


Wavelength: 210 nm

Retention time (minute): (R,S) 7.6, (S,R) 8.7

[0451]

5 Reference Example 133

[Formula 153]



[0452]

1st step

- 10 7-Fluoro- N^3,N^3 ,1-trimethyl-1H-indazol-3,5-diamine (200 mg), cesium carbonate (625 mg), $\text{Pd}_2(\text{dba})_3$ (132 mg), and Xantphos (167 mg) were added to a toluene (11 ml) solution containing tert-butyl cis-2-(6-chloro-5-cyano-3-fluoropyridin-2-ylamino)cyclohexylcarbamate (354 mg), followed by stirring at 100°C for 6 hours in a nitrogen atmosphere.
- 15 The reaction mixture was cooled to room temperature, insoluble matter was removed by filtration, and filter cake was washed with ethyl acetate. Then, the solvent was distilled away under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane: ethyl acetate = 4:1 to 2:3). Brown oily matter of tert-butyl
- 20 ((1S,2R)-2-((5-cyano-6-((3-dimethylamino-7-fluoro-1-methyl-1H-indazole-5-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexylcarbamate (277 mg) was thus obtained.

MS (ESI m/z): 541 ($M+H$)

RT (min): 1.88

25 [0453]

Reference Example 134

The compounds shown in table 1 were obtained as described in Reference Example 133.

[0454]

30 [Table 1]

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-1		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((2-methoxypyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	471	1.35
Reference Example 134-2		tert-butyl ((1R,2S)-1-((5-cyano-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	501	1.94
Reference Example 134-3		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	515	1.42
Reference Example 134-4		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	490	1.38
Reference Example 134-5		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	508	1.80
Reference Example 134-6		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-6-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	494	1.76

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-7		tert-butyl ((2S,3R)-3-((5-cyano-6-((1-ethyl-1H-indazol-6-yl)amino)-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	510	1.86
Reference Example 134-8		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-4-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	495	1.83
Reference Example 134-9		tert-butyl ((2S,3R)-3-((5-cyano-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	461	1.10
Reference Example 134-10		tert-butyl ((2S,3R)-3-((5-cyano-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	489	1.24
Reference Example 134-11		tert-butyl ((2S,3R)-3-((5-cyano-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-heptan-2-yl)carbamate	489	1.25
Reference Example 134-12		tert-butyl ((1R,2S)-1-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	507	1.63

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-13		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(m-tolylamino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	454	1.98
Reference Example 134-14		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,5-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	468	2.08
Reference Example 134-15		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((3-methoxyphenyl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	470	1.93
Reference Example 134-16		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	500	1.92
Reference Example 134-17		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,4-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	500	1.80
Reference Example 134-18		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((3-methoxy-4-methylphenyl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	484	2.05

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-19		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	514	1.44
Reference Example 134-20		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-fluorophenyl)propan-2-yl)carbamate	580	1.60
Reference Example 134-21		tert-butyl ((2R,3S)-2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	514	1.48
Reference Example 134-22		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	507	1.12
Reference Example 134-23		tert-butyl ((2R,3S)-2-((5-cyano-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-3-yl)carbamate	461	1.10
Reference Example 134-24		tert-butyl ((3S,4R)-4-((5-cyano-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)hexan-3-yl)carbamate	475	1.16

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-25		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-6-yl)amino)pyridin-2-yl)amino)heptan-2-yl)carbamate	514	1.63
Reference Example 134-26		tert-butyl ((2R,3S)-2-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-6-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	486	1.50
Reference Example 134-27		tert-butyl ((2R,3S)-2-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	486	1.44
Reference Example 134-28		tert-butyl ((2R,3S)-2-((5-cyano-6-((5-cyclopropylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-3-yl)carbamate	473	1.20
Reference Example 134-29		tert-butyl ((2R,3S)-2-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	507	1.16
Reference Example 134-30		tert-butyl ((3S,4R)-4-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)hexan-3-yl)carbamate	521	1.19
Reference Example 134-31		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(m-tolylamino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	440	1.93

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-32		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,5-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	454	2.02
Reference Example 134-33		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((3-methoxyphenyl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	456	1.83
Reference Example 134-34		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	486	1.82
Reference Example 134-35		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,4-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	486	1.69
Reference Example 134-36		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((3-methoxy-4-methylphenyl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	470	1.94
Reference Example 134-37		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((4-fluorophenyl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	444	1.84
Reference Example 134-38		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,4-difluorophenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	462	1.87

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-39		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((4-fluorophenyl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	458	1.95
Reference Example 134-40		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,4-difluorophenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	476	1.98
Reference Example 134-41		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	475	1.76
Reference Example 134-42		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	489	1.87
Reference Example 134-43		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	463	1.75
Reference Example 134-44		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	477	1.84

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-45		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	491	1.90
Reference Example 134-46		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((5-fluoro-6-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	512	1.66
Reference Example 134-47		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((5-fluoro-6-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	512	1.59
Reference Example 134-48		tert-butyl ((2S,3R)-3-((6-((5-((1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)heptan-2-yl)carbamate	509	1.76
Reference Example 134-49		tert-butyl ((2S,3R)-3-((6-((5-((1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	509	1.74
Reference Example 134-50		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	530	1.64

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-51		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	544	1.75
Reference Example 134-52		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	532	1.76
Reference Example 134-53		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	510	1.71
Reference Example 134-54		tert-butyl ((2S,3R)-3-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	495	1.67
Reference Example 134-55		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-6-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	482	1.72
Reference Example 134-56		tert-butyl ((2S,3R)-3-((5-cyano-6-((1-ethyl-1H-indazol-6-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	496	1.80

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-57		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	482	1.70
Reference Example 134-58		tert-butyl ((2S,3R)-3-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	496	1.78
Reference Example 134-59		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	497	1.20
Reference Example 134-60		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((2-methoxypyridin-4-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	477	1.13
Reference Example 134-61		tert-butyl ((2S,3R)-3-((5-cyano-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	507	1.70
Reference Example 134-62		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	521	1.18
Reference Example 134-63		tert-butyl ((2S,3R)-3-((5-cyano-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	475	1.18

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-64		tert-butyl ((2S,3R)-3-((5-cyano-6-((5-cyclopropylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	487	1.22
Reference Example 134-65		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	540	1.81
Reference Example 134-66		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	524	1.67
Reference Example 134-67		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	538	1.77
Reference Example 134-68		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	519	1.73
Reference Example 134-69		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	533	1.83

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-70		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	507	1.72
Reference Example 134-71		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	521	1.80
Reference Example 134-72		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	507	1.72
Reference Example 134-73		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	584	1.85
Reference Example 134-74		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	540	1.83

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-75		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	459	1.74
Reference Example 134-76		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	568	1.70
Reference Example 134-77		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	524	1.69
Reference Example 134-78		tert-butyl ((1R,2S)-1-((5-cyano-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	538	1.79
Reference Example 134-79		tert-butyl ((1R,2S)-1-((5-cyano-6-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	537	1.96

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-80		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	582	1.80
Reference Example 134-81		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	438	1.79
Reference Example 134-82		tert-butyl ((1R,2S)-1-((5-cyano-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	552	1.89
Reference Example 134-83		tert-butyl ((1R,2S)-1-((5-cyano-6-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	551	2.07
Reference Example 134-84		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	516	1.53

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-85		tert-butyl ((2S,3S)-3-((5-cyano-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	530	1.72
Reference Example 134-86		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	566	1.62
Reference Example 134-87		tert-butyl ((2S,3S)-3-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	530	1.62
Reference Example 134-88		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	544	1.56
Reference Example 134-89		tert-butyl ((2S,3S)-3-((5-cyano-6-((1,3-dimethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	498	1.66
Reference Example 134-90		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	512	1.62

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-91		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-yl)amino)pyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	534	1.68
Reference Example 134-92		tert-butyl ((2S,3R)-3-((5-cyano-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	554	1.95
Reference Example 134-93		tert-butyl ((2S,3R)-3-((5-cyano-6-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	525	2.00
Reference Example 134-94		tert-butyl ((2S,3R)-3-((5-cyano-6-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)hexan-2-yl)carbamate	539	2.08
Reference Example 134-95		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	556	1.73

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-96		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	570	1.81
Reference Example 134-97		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	512	1.71
Reference Example 134-98		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	526	1.79
Reference Example 134-99		tert-butyl ((2S,3R)-3-((5-cyano-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	526	1.82
Reference Example 134-100		tert-butyl ((2S,3R)-3-((5-cyano-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)hexan-2-yl)carbamate	540	1.90
Reference Example 134-101		tert-butyl ((3R,4R)-4-((5-cyano-6-((1,3-dimethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	496	1.41

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-102		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((3-fluoro-1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	500	1.53
Reference Example 134-103		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	512	1.54
Reference Example 134-104		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	530	1.63
Reference Example 134-105		tert-butyl ((3R,4R)-4-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	525	1.48
Reference Example 134-106		tert-butyl ((3R,4R)-4-((5-cyano-6-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	543	1.64
Reference Example 134-107		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	510	1.38







Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-108		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	526	1.41
Reference Example 134-109		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	526	1.43
Reference Example 134-110		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	532	1.49
Reference Example 134-111		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((5-methylpyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	443	1.03
Reference Example 134-112		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	443	1.00
Reference Example 134-113		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	459	1.61

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-114		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	473	1.72
Reference Example 134-115		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	447	1.59
Reference Example 134-116		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	461	1.69
Reference Example 134-117		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	475	1.77
Reference Example 134-118		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	461	1.40
Reference Example 134-119		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	508	1.65

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-120		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	530	1.70
Reference Example 134-121		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	501	1.31
Reference Example 134-122		tert-butyl ((1S,2R)-2-((5-cyano-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	486	1.83
Reference Example 134-123		tert-butyl ((1S,2R)-2-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	494	1.71
Reference Example 134-124		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((3-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	532	1.33
Reference Example 134-125		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((5-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	532	1.19

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-126		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((3-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	530	1.53
Reference Example 134-127		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	530	1.44
Reference Example 134-128		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((3-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	546	1.71
Reference Example 134-129		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	546	1.63
Reference Example 134-130		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	546	1.82
Reference Example 134-131		tert-butyl ((1S,2R)-2-((5-cyano-6-((6-(difluoromethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	560	1.79

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-132		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	536	1.51
Reference Example 134-133		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	550	1.59
Reference Example 134-134		(R)-tert-butyl (2-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-4-methylpentyl)carbamate	550	1.63
Reference Example 134-135		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)heptan-2-yl)carbamate	564	1.69
Reference Example 134-136		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclohexylpropan-2-yl)carbamate	584	1.84

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-137		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclohexylpropan-2-yl)carbamate	570	1.91
Reference Example 134-138		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	542	1.45
Reference Example 134-139		tert-butyl ((3S,4R)-4-((5-cyano-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)hexan-3-yl)carbamate	544	1.50
Reference Example 134-140		tert-butyl ((2S,3S)-3-((5-cyano-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	508	1.58
Reference Example 134-141		tert-butyl ((2S,3S)-3-((5-cyano-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	522	1.68
Reference Example 134-142		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-fluorophenyl)propan-2-yl)carbamate	558	1.75

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-143		tert-butyl ((2R,3S)-2-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	447	1.58
Reference Example 134-144		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	463	1.67
Reference Example 134-145		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-fluorophenyl)propan-2-yl)carbamate	513	1.82
Reference Example 134-146		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	474	1.42
Reference Example 134-147		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	474	1.37
Reference Example 134-148		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	488	1.48

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-149		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	462	1.34
Reference Example 134-150		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	476	1.45
Reference Example 134-151		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	490	1.54
Reference Example 134-152		tert-butyl ((2R,3S)-2-((5-cyano-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	530	1.52
Reference Example 134-153		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	546	1.44
Reference Example 134-154		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	560	1.53

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-155		tert-butyl ((2R,3S)-2-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	530	1.52
Reference Example 134-156		tert-butyl ((3S,4R)-4-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-3-yl)carbamate	542	1.58
Reference Example 134-157		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	546	1.44
Reference Example 134-158		tert-butyl ((2S,3S)-3-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	558	1.54
Reference Example 134-159		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	524	1.66
Reference Example 134-160		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	524	1.69

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-161		tert-butyl ((1S,2R)-2-((5-cyano-6-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	544	1.75
Reference Example 134-162		tert-butyl ((1S,2R)-2-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	523	1.70
Reference Example 134-163		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	513	1.45
Reference Example 134-164		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	544	1.73
Reference Example 134-165		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	566	1.80
Reference Example 134-166		tert-butyl ((1R,2S)-1-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	559	1.78

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-167		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	460 (+H-Boc)	1.74
Reference Example 134-168		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	543	1.44
Reference Example 134-169		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	574	1.71
Reference Example 134-170		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	490 (+H-Boc)	1.72
Reference Example 134-171		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	596	1.77

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-172		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	560	1.77
Reference Example 134-173		tert-butyl ((1R,2S)-1-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	587	1.76
Reference Example 134-174		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate	542	1.61
Reference Example 134-175		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate	564	1.61
Reference Example 134-176		tert-butyl ((2S,3R)-3-((5-cyano-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate	535	1.26

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-177		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate	520	1.74
Reference Example 134-178		tert-butyl ((3R,4S)-4-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)bicyclo[4.1.0]heptan-3-yl)carbamate	505	1.66
Reference Example 134-179		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-(trifluoromethyl)phenyl)propan-2-yl)carbamate	612	1.85
Reference Example 134-180		tert-butyl ((1R,2S)-1-((5-cyano-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-(trifluoromethyl)phenyl)propan-2-yl)carbamate	591	2.03
Reference Example 134-181		tert-butyl ((1R,2S)-1-((5-cyano-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	523	1.93
Reference Example 134-182		tert-butyl ((1R,2S)-1-((5-cyano-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	553	1.89

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-183		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	522	1.68
Reference Example 134-184		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	544	1.74
Reference Example 134-185		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	515	1.41
Reference Example 134-186		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	500	1.86
Reference Example 134-187		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	508	1.75
Reference Example 134-188		(R)-tert-butyl (2-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-3-phenylpropyl)carbamate	548	1.60

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-189		(R)-tert-butyl 2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl)carbamate	562	1.51
Reference Example 134-190		(R)-tert-butyl 2-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl)carbamate	548	1.60
Reference Example 134-191		tert-butyl ((1R,2S)-1-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	507	1.63
Reference Example 134-192		tert-butyl ((3S,4R)-4-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)bicyclo[4.1.0]heptan-3-yl)carbamate	505	1.66
Reference Example 134-193		tert-butyl ((3S,4R)-4-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)bicyclo[4.1.0]heptan-3-yl)carbamate	520	1.70
Reference Example 134-194		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(quinolin-6-yl)amino)pyridin-2-yl)amino)-1-(4-(trifluoromethyl)phenyl)propan-2-yl)carbamate	581	1.63

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-195		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-((S)-2,2-dimethylcyclopropyl)propan-2-yl)carbamate	536	1.75
Reference Example 134-196		tert-butyl ((1S,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(pyridin-2-yl)propan-2-yl)carbamate	531	1.47
Reference Example 134-197		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	473	1.68
Reference Example 134-198		tert-butyl ((1R,2S)-1-((5-cyano-6-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	555	1.87
Reference Example 134-199		tert-butyl ((1S,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(pyridin-2-yl)propan-2-yl)carbamate	545	1.46
Reference Example 134-200		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(quinolin-5-ylamino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	477	1.33

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-201		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	562	1.73
Reference Example 134-202		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	548	1.79
Reference Example 134-203		tert-butyl ((1R,2S)-1-((5-cyano-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	541	1.91
Reference Example 134-204		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	562	1.70
Reference Example 134-205		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	548	1.79
Reference Example 134-206		tert-butyl ((1R,2S)-1-((5-cyano-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	541	1.89

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-207		tert-butyl ((1S,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	550	1.70
Reference Example 134-208		tert-butyl ((1S,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	536	1.77
Reference Example 134-209		(R)-tert-butyl (2-((5-cyano-3-fluoro-6-((2-methoxypyridin-4-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl)carbamate	ND	ND
Reference Example 134-210		(R)-tert-butyl (2-((5-cyano-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-3-phenylpropyl)carbamate	541	1.75
Reference Example 134-211		(R)-tert-butyl (2-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl)carbamate	555	1.20
Reference Example 134-212		(R)-tert-butyl (2-((5-cyano-3-fluoro-6-((6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl)carbamate	566	1.20

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-213		(R)-tert-butyl 2-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-3-phenylpropyl carbamate	577	1.58
Reference Example 134-214		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-phenylbutan-2-yl) carbamate	558	1.72
Reference Example 134-215		tert-butyl ((2S,3R)-3-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-phenylbutan-2-yl) carbamate	544	1.77
Reference Example 134-216		tert-butyl ((2S,3R)-3-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-phenylbutan-2-yl) carbamate	573	1.77
Reference Example 134-217		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-4-phenylbutan-2-yl) carbamate	580	1.75

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-218		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate	507	1.70
Reference Example 134-219		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate	551	1.43
Reference Example 134-220		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-(m-tolylamino)pyridin-2-yl)amino)cyclohexyl)carbamate	440	1.96
Reference Example 134-221		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-(p-tolylamino)pyridin-2-yl)amino)cyclohexyl)carbamate	440	1.96
Reference Example 134-222		tert-butyl ((1S,2R)-2-((5-cyano-6-((3,4-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	454	2.03
Reference Example 134-223		tert-butyl ((1S,2R)-2-((6-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	493	1.85

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-224		tert-butyl ((1S,2R)-2-((6-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	493	1.61
Reference Example 134-225		tert-butyl ((1S,2R)-2-((6-((3-chlorophenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	460 462	1.99
Reference Example 134-226		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((3-fluorophenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	444	1.90
Reference Example 134-227		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((4-fluorophenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	444	1.87
Reference Example 134-228		tert-butyl ((1S,2R)-2-((6-((3-acetylphenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	468	1.74
Reference Example 134-229		tert-butyl ((1S,2R)-2-((6-((3,5-bis(trifluoromethyl)phenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	561	2.16
Reference Example 134-230		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((3-(trifluoromethoxy)phenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	510	2.04







Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-231		tert-butyl ((1S,2R)-2-((6-((3-chloro-4-methylphenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	474	2.09
Reference Example 134-232		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((3-isopropoxyphenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	484	2.01
Reference Example 134-233		tert-butyl ((1S,2R)-2-((5-cyano-6-((3,4-difluorophenyl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	462	1.92
Reference Example 134-234		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((4-isopropylphenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	468	2.11
Reference Example 134-235		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((4-isopropoxyphenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	484	1.98
Reference Example 134-236		tert-butyl ((1S,2R)-2-((5-cyano-6-((3-ethylphenyl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	454	2.04

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-237		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	584	1.78
Reference Example 134-238		tert-butyl ((1R,2S)-1-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	577	1.77
Reference Example 134-239		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	513	1.73
Reference Example 134-240		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	578	1.74
Reference Example 134-241		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	590	1.71

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-242		(S)-tert-butyl 2-((5-cyano-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl)carbamate	545	1.00
Reference Example 134-243		(S)-tert-butyl 2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl)carbamate	552	1.23
Reference Example 134-244		(S)-tert-butyl 2-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl)carbamate	574	1.32
Reference Example 134-245		(S)-tert-butyl 2-((5-cyano-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl)carbamate	503	1.18
Reference Example 134-246		(S)-tert-butyl 2-((5-cyano-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl)carbamate	538	1.34
Reference Example 134-247		(S)-tert-butyl 2-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl)carbamate	567	1.31

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-248		(S)-tert-butyl 2-((5-cyano-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl carbamate	538	1.42
Reference Example 134-249		tert-butyl ((1S,2R)-2-((5-cyano-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	507	1.65
Reference Example 134-250		tert-butyl ((1S,2R)-2-((5-cyano-6-((5-(3,5-dimethyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	521	1.62
Reference Example 134-251		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-(m-tolylamino)pyridin-2-yl)amino)pentan-2-yl)carbamate	428	1.89
Reference Example 134-252		tert-butyl ((2S,3R)-3-((6-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	481	1.84
Reference Example 134-253		tert-butyl ((2S,3R)-3-((6-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	481	1.60

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-254		tert-butyl ((2S,3R)-3-((5-cyano-6-((3,4-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	442	1.95
Reference Example 134-255		tert-butyl ((2S,3R)-3-((6-((3-acetylphenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	456	1.72
Reference Example 134-256		tert-butyl ((1R,2S)-1-((6-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	493	1.84
Reference Example 134-257		tert-butyl ((1R,2S)-1-((6-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	493	1.61
Reference Example 134-258		tert-butyl ((1R,2S)-1-((5-cyano-6-((3,4-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	454	1.96
Reference Example 134-259		tert-butyl ((1R,2S)-1-((6-((3-acetylphenyl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	468	1.73
Reference Example 134-260		tert-butyl ((1S,2R)-2-((6-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	527 529	1.76

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-261		Mixture of tert-butyl (3-(benzyloxy)-2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate (1S,2S,3S),(1R,2R,3R)	614	1.78
Reference Example 134-262		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-(pyridin-3-ylamino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	441	1.23
Reference Example 134-263		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoropyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	459	1.64
Reference Example 134-264		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	489	1.79
Reference Example 134-265		tert-butyl ((1S,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	572	1.78
Reference Example 134-266		tert-butyl ((1S,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	536	1.82

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-267		tert-butyl ((1S,2S)-1-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	565	1.77
Reference Example 134-268		tert-butyl ((1S,2S)-1-(5-chlorothiophen-2-yl)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)propan-2-yl)carbamate	606	1.90
Reference Example 134-269		tert-butyl ((1S,2S)-1-(5-chlorothiophen-2-yl)-1-((5-cyano-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)propan-2-yl)carbamate	570	1.94
Reference Example 134-270		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	584	1.80
Reference Example 134-271		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	548	1.83

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-272		tert-butyl ((1S,2R)-2-((6-((5-acetyl-6-methoxypyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	499	1.76
Reference Example 134-273		tert-butyl ((1S,2R)-2-((6-((5-acetyl-6-(methylamino)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	498	1.50
Reference Example 134-274		tert-butyl ((1S,2R)-2-((6-((5-acetyl-6-morpholinopyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	554	1.63
Reference Example 134-275		tert-butyl ((1S,2R)-2-((6-((5-acetyl-6-methylpyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	483	1.43
Reference Example 134-276		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	507	1.64
Reference Example 134-277		tert-butyl ((1R,2S)-1-((6-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	527 529	1.74

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-278		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	495	1.64
Reference Example 134-279		tert-butyl ((2S,3R)-3-((6-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-cyano-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	515 517	1.74
Reference Example 134-280		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	521	1.75
Reference Example 134-281		(R)-tert-butyl (2-((5-cyano-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methylpentyl)carbamate	509	1.75
Reference Example 134-282		tert-butyl ((2S,3R)-3-((5-cyano-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	523	1.80
Reference Example 134-283		tert-butyl ((3R,4R)-4-((5-cyano-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	509	1.41

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-284		Mixture of tert-butyl (5-(benzyloxy)-2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate (1S,2R,5R),(1R,2S,5S)	614	1.80
Reference Example 134-285		Mixture of tert-butyl (2-(benzyloxy)-6-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate (1R,2R,6R),(1S,2S,6S)	614	1.80
Reference Example 134-286		Mixture of tert-butyl (2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-hydroxycyclohexyl)carbamate (1S,2R,5R),(1R,2S,5S)	614	1.80
Reference Example 134-287		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((5-(4-methyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	526	1.54
Reference Example 134-288		tert-butyl ((1S,2R)-2-((5-cyano-6-((5-(4,5-dimethyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	540	1.59
Reference Example 134-289		tert-butyl ((1R,2S)-1-((5-cyano-6-((5-(4,5-dimethyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	540	1.56

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-290		tert-butyl ((1R,2S)-1-((5-cyano-6-((1,3-dimethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	508	1.71
Reference Example 134-291		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((1-methyl-1H-indazol-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	494	1.70
Reference Example 134-292		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	544	1.74
Reference Example 134-293		tert-butyl ((1R,2S)-1-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3,5-difluorophenyl)propan-2-yl)carbamate	595	1.82
Reference Example 134-294		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-(3,5-difluorophenyl)propan-2-yl)carbamate	594	1.76
Reference Example 134-295		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3,5-difluorophenyl)propan-2-yl)carbamate	566	1.86


Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-296		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate	580	1.78
Reference Example 134-297		tert-butyl ((1R,2S)-1-((5-cyano-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate	595	1.82
Reference Example 134-298		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate	596	1.76
Reference Example 134-299		tert-butyl ((1R,2S)-1-((5-cyano-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate	564	1.85
Reference Example 134-300		Mixture of tert-butyl (2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-6-hydroxycyclohexyl)carbamate (1R,2R,6R),(1S,2S,6S)	614	1.80

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 134-301		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((2-methoxypyrimidin-5-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	458	1.49
Reference Example 134-302		tert-butyl ((1R,2S)-1-((5-cyano-3-fluoro-6-((2-morpholinopyrimidin-5-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	513	1.56
Reference Example 134-303		Mixture of tert-butyl (2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5,5-difluorocyclohexyl)carbamate (1S,2R),(1R,2S)	544	1.62
Reference Example 134-304		Mixture of tert-butyl (2-((5-cyano-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-6-fluorocyclohexyl)carbamate (1R,2R,6S),(1S,2S,6R)	614	1.80

[0455]

Reference Example 135

[Formula 154]



5 A 5N sodium hydroxide aqueous solution (0.512 ml) and a 30% hydrogen peroxide solution (0.29 ml) were added to a DMSO/EtOH (3 ml/3 ml) mixed solution containing tert-butyl ((1S,2R)-2-((5-cyano-6-((3-dimethylamino-7-fluoro-1-methyl-1H-indazole-5-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexylcarbamate (277 mg),
 10 followed by stirring at room temperature for 30 minutes. Water was added to the reaction solution and a solid was collected by filtration. A yellow solid of tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazole-5-yl)amino)-3-fluoropyridin-2-yl)amino) cyclohexyl)carbamate (283 mg)
 15 was thus obtained.

MS (ESI m/z): 559 (M+H)

RT (min): 1.72

[0456]

Reference Example 136

20 The compounds shown in table 2 were obtained as described in Reference Example 135.

[0457]

[Table 2]

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-1		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((2-methoxypyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	489	1.11
Reference Example 136-2		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	519	1.70
Reference Example 136-3		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	433	1.16
Reference Example 136-4		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	509	1.17
Reference Example 136-5		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	526	1.59
Reference Example 136-6		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-6-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	512	1.52

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-7		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((1-ethyl-1H-indazol-6-yl)amino)-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	528	1.68
Reference Example 136-8		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-4-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	514	1.71
Reference Example 136-9		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	461	1.10
Reference Example 136-10		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	489	1.24
Reference Example 136-11		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)heptan-2-yl)carbamate	489	1.25
Reference Example 136-12		tert-butyl ((1R,2S)-1-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	525	1.53

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-13		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(m-tolylamino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	472	1.89
Reference Example 136-14		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,5-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	486	1.98
Reference Example 136-15		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((3-methoxyphenyl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	488	1.78
Reference Example 136-16		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	518	1.76
Reference Example 136-17		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,4-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	518	1.61
Reference Example 136-18		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((3-methoxy-4-methylphenyl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	502	1.85

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-19		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	514	1.44
Reference Example 136-20		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-fluorophenyl)propan-2-yl)carbamate	580	1.60
Reference Example 136-21		tert-butyl ((2R,3S)-2-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	514	1.48
Reference Example 136-22		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	507	1.12
Reference Example 136-23		tert-butyl ((2R,3S)-2-((5-carbamoyl-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-3-yl)carbamate	461	1.10
Reference Example 136-24		tert-butyl ((3S,4R)-4-((5-carbamoyl-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)hexan-3-yl)carbamate	475	1.16

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-25		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-6-yl)amino)pyridin-2-yl)amino)heptan-2-yl)carbamate	514	1.63
Reference Example 136-26		tert-butyl ((2R,3S)-2-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-6-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	486	1.50
Reference Example 136-27		tert-butyl ((2R,3S)-2-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	486	1.44
Reference Example 136-28		tert-butyl ((2R,3S)-2-((5-carbamoyl-6-((5-cyclopropylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-3-yl)carbamate	473	1.20
Reference Example 136-29		tert-butyl ((2R,3S)-2-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	507	1.16
Reference Example 136-30		tert-butyl ((3S,4R)-4-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)hexan-3-yl)carbamate	521	1.19

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-31		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(m-tolylamino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	458	1.78
Reference Example 136-32		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,5-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	472	1.87
Reference Example 136-33		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((3-methoxyphenyl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	474	1.68
Reference Example 136-34		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	504	1.67
Reference Example 136-35		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,4-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	504	1.52
Reference Example 136-36		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((3-methoxy-4-methylphenyl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	488	1.79

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-37		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((4-fluorophenyl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	462	1.70
Reference Example 136-38		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,4-difluorophenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	480	1.76
Reference Example 136-39		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((4-fluorophenyl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	478	1.81
Reference Example 136-40		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,4-difluorophenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	494	1.86
Reference Example 136-41		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	493	1.61
Reference Example 136-42		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	507	1.72

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-43		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	481	1.61
Reference Example 136-44		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	495	1.69
Reference Example 136-45		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	509	1.77
Reference Example 136-46		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	530	1.56
Reference Example 136-47		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	530	1.47
Reference Example 136-48		tert-butyl ((2S,3R)-3-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)heptan-2-yl)carbamate	527	1.61






Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-49		tert-butyl ((2S,3R)-3-((6-((5-((1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	527	1.60
Reference Example 136-50		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	548	1.48
Reference Example 136-51		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	562	1.59
Reference Example 136-52		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	550	1.61
Reference Example 136-53		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	528	1.55

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-54		tert-butyl ((2S,3R)-3-((6-((5-((1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	513	1.53
Reference Example 136-55		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-6-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	500	1.53
Reference Example 136-56		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((1-ethyl-1H-indazol-6-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	514	1.61
Reference Example 136-57		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	500	1.52
Reference Example 136-58		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	514	1.60
Reference Example 136-59		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	497	1.20

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-60		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((2-methoxypyridin-4-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	477	1.13
Reference Example 136-61		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	507	1.70
Reference Example 136-62		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	521	1.18
Reference Example 136-63		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((5,6-dimethylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	475	1.18
Reference Example 136-64		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((5-cyclopropylpyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methylpentan-2-yl)carbamate	487	1.22
Reference Example 136-65		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	558	1.66

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-66		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	542	1.52
Reference Example 136-67		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	556	1.61
Reference Example 136-68		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	537	1.59
Reference Example 136-69		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	551	1.69
Reference Example 136-70		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	525	1.57

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-71		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	539	1.66
Reference Example 136-72		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	553	1.72
Reference Example 136-73		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	602	1.71
Reference Example 136-74		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	558	1.71
Reference Example 136-75		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	477	1.57

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-76		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	586	1.55
Reference Example 136-77		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	542	1.56
Reference Example 136-78		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	556	1.68
Reference Example 136-79		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	555	1.85
Reference Example 136-80		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	600	1.65

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-81		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	556	1.66
Reference Example 136-82		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	570	1.77
Reference Example 136-83		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	569	1.95
Reference Example 136-84		tert-butyl ((2S,3S)-3-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	516	1.53
Reference Example 136-85		tert-butyl ((2S,3S)-3-((5-carbamoyl-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	530	1.72

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-86		tert-butyl ((2S,3S)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	566	1.62
Reference Example 136-87		tert-butyl ((2S,3S)-3-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	530	1.62
Reference Example 136-88		tert-butyl ((2S,3S)-3-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	544	1.56
Reference Example 136-89		tert-butyl ((2S,3S)-3-((5-carbamoyl-6-((1,3-dimethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	516	1.49
Reference Example 136-90		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	572	1.84

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-91		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	543	1.89
Reference Example 136-92		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)hexan-2-yl)carbamate	557	1.97
Reference Example 136-93		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	574	1.59
Reference Example 136-94		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	588	1.67
Reference Example 136-95		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	530	1.59

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-96		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	544	1.67
Reference Example 136-97		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	544	1.70
Reference Example 136-98		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)hexan-2-yl)carbamate	558	1.78
Reference Example 136-99		tert-butyl ((3R,4R)-4-((5-carbamoyl-6-((1,3-dimethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	514	1.32
Reference Example 136-100		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((3-fluoro-1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	518	1.44
Reference Example 136-101		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	530	1.37

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-102		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	548	1.52
Reference Example 136-103		tert-butyl ((3R,4R)-4-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	543	1.24
Reference Example 136-104		tert-butyl ((3R,4R)-4-((5-carbamoyl-6-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	561	1.41
Reference Example 136-105		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	528	1.17
Reference Example 136-106		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	544	1.22
Reference Example 136-107		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	544	1.24

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-108		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	550	1.30
Reference Example 136-109		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((5-methylpyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	461	0.88
Reference Example 136-110		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	461	0.86
Reference Example 136-111		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	477	1.55
Reference Example 136-112		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	491	1.67
Reference Example 136-113		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	465	1.43

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-114		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	479	1.53
Reference Example 136-115		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	493	1.61
Reference Example 136-116		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	479	1.24
Reference Example 136-117		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	526	1.48
Reference Example 136-118		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	548	1.54
Reference Example 136-119		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	519	1.16

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-120		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	504	1.67
Reference Example 136-121		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	512	-
Reference Example 136-122		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((3-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	550	1.05
Reference Example 136-123		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((5-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	550	0.99
Reference Example 136-124		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((3-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	548	1.21
Reference Example 136-125		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	548	1.15

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-126		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((3-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	564	1.34
Reference Example 136-127		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-2-morpholinopyridin-4-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	564	1.26
Reference Example 136-128		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	564	1.67
Reference Example 136-129		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((6-(difluoromethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	578	1.67
Reference Example 136-130		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	536	1.51

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-131		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	550	1.59
Reference Example 136-132		(R)-tert-butyl (2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-4-methylpentyl)carbamate	550	1.63
Reference Example 136-133		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)heptan-2-yl)carbamate	564	1.69
Reference Example 136-134		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclohexylpropan-2-yl)carbamate	568	1.67
Reference Example 136-135		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclohexylpropan-2-yl)carbamate	554	1.75

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-137		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	542	1.45
Reference Example 136-137		tert-butyl ((3S,4R)-4-((5-carbamoyl-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)hexan-3-yl)carbamate	544	1.50
Reference Example 136-138		tert-butyl ((2S,3S)-3-((5-carbamoyl-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	508	1.58
Reference Example 136-139		tert-butyl ((2S,3S)-3-((5-carbamoyl-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	522	1.68
Reference Example 136-140		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-fluorophenyl)propan-2-yl)carbamate	558	1.75
Reference Example 136-141		tert-butyl ((2R,3S)-2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	465	1.44

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-142		tert-butyl ((2S,3S)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	481	1.33
Reference Example 136-143		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-fluorophenyl)propan-2-yl)carbamate	531	1.60
Reference Example 136-144		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	492	1.19
Reference Example 136-145		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	492	1.15
Reference Example 136-146		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	506	1.24
Reference Example 136-147		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	480	1.13

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-148		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-2-yl)carbamate	494	1.22
Reference Example 136-149		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	508	1.30
Reference Example 136-150		tert-butyl ((2R,3S)-2-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	530	1.52
Reference Example 136-151		tert-butyl ((2S,3S)-3-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	546	1.44
Reference Example 136-152		tert-butyl ((2S,3S)-3-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	560	1.53
Reference Example 136-153		tert-butyl ((2R,3S)-2-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-3-yl)carbamate	530	1.52

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-154		tert-butyl ((3S,4R)-4-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)hexan-3-yl)carbamate	542	1.58
Reference Example 136-155		tert-butyl ((2S,3S)-3-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methoxybutan-2-yl)carbamate	546	1.44
Reference Example 136-156		tert-butyl ((2S,3S)-3-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-ethoxybutan-2-yl)carbamate	558	1.54
Reference Example 136-157		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	542	1.54
Reference Example 136-158		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	542	1.54
Reference Example 136-159		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	562	1.59

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-160		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropyl)carbamate	541	1.51
Reference Example 136-161		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	531	1.30
Reference Example 136-162		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	562	1.56
Reference Example 136-163		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	584	1.64
Reference Example 136-164		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	577	1.60
Reference Example 136-165		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	578	1.61

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-166		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	561	1.29
Reference Example 136-167		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	592	1.54
Reference Example 136-168		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	608	1.59
Reference Example 136-169		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	614	1.62
Reference Example 136-170		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-(1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	578	1.60

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-171		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	607	1.57
Reference Example 136-172		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate	560	—
Reference Example 136-173		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate	582	—
Reference Example 136-174		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate	553	—
Reference Example 136-175		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-5-(methylthio)pentan-2-yl)carbamate	538	—
Reference Example 136-176		tert-butyl ((3R,4S)-4-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)bicyclo[4.1.0]heptan-3-yl)carbamate	523	1.50

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-177		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(4-(trifluoromethyl)phenyl)propan-2-yl)carbamate	630	1.71
Reference Example 136-178		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-(trifluoromethyl)phenyl)propan-2-yl)carbamate	609	1.90
Reference Example 136-179		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-phenylpropan-2-yl)carbamate	541	1.78
Reference Example 136-180		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	571	1.74
Reference Example 136-181		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	540	1.75

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-182		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	562	1.72
Reference Example 136-183		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	533	1.22
Reference Example 136-184		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,5-dimethoxyphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	518	1.72
Reference Example 136-185		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	526	1.58
Reference Example 136-186		(R)-tert-butyl (2-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-3-phenylpropyl)carbamate	548	1.60
Reference Example 136-187		(R)-tert-butyl (2-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl)carbamate	562	1.51

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-188		(R)-tert-butyl 2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl carbamate	548	1.60
Reference Example 136-189		tert-butyl ((1R,2S)-1-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	525	1.49
Reference Example 136-190		tert-butyl ((3S,4R)-4-((6-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)bicyclo[4.1.0]heptan-3-yl)carbamate	523	1.50
Reference Example 136-191		tert-butyl ((3S,4R)-4-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)bicyclo[4.1.0]heptan-3-yl)carbamate	538	1.53
Reference Example 136-192		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(quinolin-6-ylamino)pyridin-2-yl)amino)-1-(4-(trifluoromethyl)phenyl)propan-2-yl)carbamate	599	1.43
Reference Example 136-193		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-((S)-2,2-dimethylcyclopropyl)propan-2-yl)carbamate	554	1.62

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-194		tert-butyl ((1S,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(pyridin-2-yl)propan-2-yl)carbamate	549	1.30
Reference Example 136-195		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	491	1.51
Reference Example 136-196		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	573	1.72
Reference Example 136-197		tert-butyl ((1S,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(pyridin-2-yl)propan-2-yl)carbamate	563	1.29
Reference Example 136-198		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(quinolin-5-ylamino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	495	1.18
Reference Example 136-199		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	580	1.58

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-200		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	566	1.63
Reference Example 136-201		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	559	1.78
Reference Example 136-202		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	580	1.55
Reference Example 136-203		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	566	1.62
Reference Example 136-204		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	559	1.74
Reference Example 136-205		tert-butyl ((1S,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	568	1.53

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-206		tert-butyl ((1S,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	554	1.59
Reference Example 136-207		(R)-tert-butyl (2-((5-carbamoyl-6-((2,6-dimethoxypyridin-4-yl)amino)-3-fluoropyridin-2-yl)amino)-3-phenylpropyl)carbamate	541	1.75
Reference Example 136-208		(R)-tert-butyl (2-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl)carbamate	555	1.20
Reference Example 136-209		(R)-tert-butyl (2-((5-carbamoyl-3-fluoro-6-((6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-3-phenylpropyl)carbamate	566	1.20
Reference Example 136-210		(R)-tert-butyl (2-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-3-phenylpropyl)carbamate	577	1.58
Reference Example 136-211		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate	576	1.56

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-212		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate	562	1.60
Reference Example 136-213		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate	591	1.59
Reference Example 136-214		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate	598	1.60
Reference Example 136-215		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate	527	1.53
Reference Example 136-216		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-4-phenylbutan-2-yl)carbamate	569	1.23
Reference Example 136-217		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-(m-tolylamino)pyridin-2-yl)amino)cyclohexyl)carbamate	458	1.82

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-218		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-(p-tolylamino)pyridin-2-yl)amino)cyclohexyl)carbamate	458	1.82
Reference Example 136-219		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((3,4-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	472	1.89
Reference Example 136-220		tert-butyl ((1S,2R)-2-((6-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	511	1.73
Reference Example 136-221		tert-butyl ((1S,2R)-2-((6-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	511	1.47
Reference Example 136-222		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((3-chlorophenyl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	478 480	1.88
Reference Example 136-223		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((3-fluorophenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	462	1.78

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-224		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((4-fluorophenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	462	1.75
Reference Example 136-225		tert-butyl ((1S,2R)-2-((6-((3-acetylphenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	486	1.61
Reference Example 136-228		tert-butyl ((1S,2R)-2-((6-((3,5-bis(trifluoromethyl)phenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	580	2.09
Reference Example 136-227		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((3-(trifluoromethoxy)phenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	528	2.00
Reference Example 136-228		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((3-chloro-4-methylphenyl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	492 494	2.25
Reference Example 136-229		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((3-isopropoxyphenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	502	2.07

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-230		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((3,4-difluorophenyl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	480	1.87
Reference Example 136-231		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((4-isopropylphenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	486	2.00
Reference Example 136-232		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((4-isopropoxyphenyl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	502	1.84
Reference Example 136-233		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((3-ethylphenyl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	472	1.91
Reference Example 136-234		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	602	1.62

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-235		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	595	1.58
Reference Example 136-236		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	531	1.55
Reference Example 136-237		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-(3-fluorophenyl)propan-2-yl)carbamate	596	1.55
Reference Example 136-238		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-(4-methoxyphenyl)propan-2-yl)carbamate	608	1.50
Reference Example 136-239		(S)-tert-butyl (2-((5-carbamoyl-3-fluoro-6-((2-(2-methoxyethoxy)pyridin-4-yl)amino)pyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl)carbamate	545	1.00
Reference Example 136-240		(S)-tert-butyl (2-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl)carbamate	552	1.23

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-241		(S)-tert-butyl 2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl carbamate	574	1.32
Reference Example 136-242		(S)-tert-butyl 2-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methylpyridin-3-yl)amino)pyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl carbamate	503	1.18
Reference Example 136-243		(S)-tert-butyl 2-((5-carbamoyl-6-((1-ethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl carbamate	538	1.34
Reference Example 136-244		(S)-tert-butyl 2-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl carbamate	567	1.31
Reference Example 136-245		(S)-tert-butyl 2-((5-carbamoyl-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-3-(1H-pyrazol-1-yl)propyl carbamate	538	1.42
Reference Example 136-246		tert-butyl ((1S,2R)-2-((5-carbamoyl-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	525	1.49

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-247		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((5-(3,5-dimethyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	539	1.46
Reference Example 136-248		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-(m-tolylamino)pyridin-2-yl)amino)pentan-2-yl)carbamate	446	1.74
Reference Example 136-249		tert-butyl ((2S,3R)-3-((6-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	499	1.76
Reference Example 136-250		tert-butyl ((2S,3R)-3-((6-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	499	1.55
Reference Example 136-251		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((3,4-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	460	1.93
Reference Example 136-252		tert-butyl ((2S,3R)-3-((6-((3-acetylphenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	474	1.64

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-253		tert-butyl ((1R,2S)-1-((6-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	511	1.73
Reference Example 136-254		tert-butyl ((1R,2S)-1-((6-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	511	1.49
Reference Example 136-255		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3,4-dimethylphenyl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	472	1.83
Reference Example 136-256		tert-butyl ((1R,2S)-1-((6-((3-acetylphenyl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	486	1.62
Reference Example 136-257		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	545 547	1.65
Reference Example 136-258		Mixture of tert-butyl (3-(benzyloxy)-2-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate (1S,2S,3S), (1R,2R,3R)	632	1.64

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-259		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-(pyridin-3-ylamino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	459	1.09
Reference Example 136-260		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoropyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	477	1.51
Reference Example 136-281		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-methoxypyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	507	1.66
Reference Example 136-262		tert-butyl ((1S,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	590	1.63
Reference Example 136-263		tert-butyl ((1S,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	554	1.71
Reference Example 136-264		tert-butyl ((1S,2S)-1-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(thiophen-2-yl)propan-2-yl)carbamate	583	1.57

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-265		tert-butyl ((1S,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(5-chlorothiophen-2-yl)propan-2-yl)carbamate	624	1.75
Reference Example 136-266		tert-butyl ((1S,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(5-chlorothiophen-2-yl)propan-2-yl)carbamate	588	1.81
Reference Example 136-267		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-fluoro-6-morpholinopyridin-3-yl)amino)pyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	602	1.63
Reference Example 136-268		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(2-fluorophenyl)propan-2-yl)carbamate	566	1.71
Reference Example 136-269		tert-butyl ((1S,2R)-2-((6-((5-acetyl-6-methoxypyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	517	1.65
Reference Example 136-270		tert-butyl ((1S,2R)-2-((6-((5-acetyl-6-(methylamino)pyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	516	1.26

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-271		tert-butyl ((1S,2R)-2-((6-((5-acetyl-6-morpholinopyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	572	1.49
Reference Example 136-272		tert-butyl ((1S,2R)-2-((6-((5-acetyl-6-methylpyridin-3-yl)amino)-5-carbamoyl-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	501	1.23
Reference Example 136-273		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	525	1.50
Reference Example 136-274		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	545 547	1.63
Reference Example 136-275		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)pentan-2-yl)carbamate	513	1.49
Reference Example 136-276		tert-butyl ((2S,3R)-3-((5-carbamoyl-6-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)pentan-2-yl)carbamate	533 535	1.62

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-277		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclobutylpropan-2-yl)carbamate	539	1.60
Reference Example 136-278		(R)-tert-butyl (2-((5-carbamoyl-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-4-methylpentyl)carbamate	527	1.60
Reference Example 136-279		tert-butyl ((2S,3R)-3-((5-carbamoyl-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5-methylhexan-2-yl)carbamate	541	1.66
Reference Example 136-280		tert-butyl ((3R,4R)-4-((5-carbamoyl-3-fluoro-6-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)tetrahydro-2H-pyran-3-yl)carbamate	527	1.25
Reference Example 136-281		Mixture of tert-butyl (5-(benzyloxy)-2-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate (1S,2R,5R), (1R,2S,5S)	632	1.66
Reference Example 136-282		tert-butyl ((1S,2R,4S)-4-(benzyloxy)-2-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate	632	1.57

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-283		Mixture of tert-butyl (2-(benzyloxy)-6-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)cyclohexyl)carbamate (1R,2R,6R),(1S,2S,6S)	632	1.63
Reference Example 136-284		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((5-(4-methyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	526	1.54
Reference Example 136-285		tert-butyl ((1S,2R)-2-((5-carbamoyl-6-((5-(4,5-dimethyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)cyclohexyl)carbamate	540	1.59
Reference Example 136-286		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((5-(4,5-dimethyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	540	1.56
Reference Example 136-287		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1,3-dimethyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	526	1.52
Reference Example 136-288		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((1-methyl-1H-indazol-4-yl)amino)pyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	512	1.60

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-289		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-cyclopropylbutan-2-yl)carbamate	562	1.65
Reference Example 136-290		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3,5-difluorophenyl)propan-2-yl)carbamate	598	1.63
Reference Example 136-291		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3,5-difluorophenyl)propan-2-yl)carbamate	613	1.62
Reference Example 136-292		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-(3,5-difluorophenyl)propan-2-yl)carbamate	614	1.58
Reference Example 136-293		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3,5-difluorophenyl)propan-2-yl)carbamate	584	1.74


Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-294		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate	598	1.61
Reference Example 136-295		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate	613	1.62
Reference Example 136-296		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)pyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate	614	1.58
Reference Example 136-297		tert-butyl ((1R,2S)-1-((5-carbamoyl-6-((1-ethyl-1H-indazol-4-yl)amino)-3-fluoropyridin-2-yl)amino)-1-(3,4-difluorophenyl)propan-2-yl)carbamate	584	1.72
Reference Example 136-298		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((2-methoxypyrimidine-5-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	476	-

Reference Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Reference Example 136-299		tert-butyl ((1R,2S)-1-((5-carbamoyl-3-fluoro-6-((2-morpholinopyrimidine-5-yl)amino)pyridin-2-yl)amino)-1-cyclopropylpropan-2-yl)carbamate	531	1.37
Reference Example 136-300		Mixture of tert-butyl (2-((5-carbamoyl-3-fluoro-6-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)pyridin-2-yl)amino)-5,5-difluorocyclohexyl)carbamate (1S,2R),(1R,2S)	562	1.47

[0458]

Example 1 (Example 2-184)

[Formula 155]



5 A mixture of tert-butyl
 ((1S,2R)-2-((5-carbamoyl-6-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazole-5-yl)amino)-3-fluoropyridin-2-yl)amino) cyclohexyl)carbamate (283 mg)
 and TFA (3 ml) was stirred at room temperature for 30 minutes. The solvent
 was distilled away under reduced pressure. 4N hydrogen
 10 chloride/1,4-dioxane (0.253 ml) was added to an ethyl acetate (10 ml)
 suspension containing the obtained residue, followed by stirring at room
 temperature for 30 minutes. A solid was collected by filtration and washed
 with ethyl acetate. A light yellow solid of
 15 6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-7-fluoro-1-me
 thyl-1H-indazole-5-yl)amino)-5-fluoronicotinamide hydrochloride (241 mg)
 was thus obtained.

(Table 3 (Example 2-184) lists ¹H-NMR data and MS data.)

[0459]

Example 2

20 The compounds shown in table 3 were obtained as described in
 Example 1.

[0460]

[Table 3]

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-1		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	441	0.9	¹ H-NMR (MeOD) δ: 8.01 (1H, br s), 7.83 (1H, br s), 7.79 (1H, br s), 7.58 (1H, br s), 4.32-4.29 (1H, m), 3.91 (3H, s), 3.70-3.66 (1H, m), 3.38 (6H, br s), 1.79-1.50 (8H, m).
Example 2-2		6-(((2R,3S)-3-amino-1-cyclopropylbutan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	455	0.95	
Example 2-3		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	457	1.03	¹ H-NMR (MeOD) δ: 8.05 (1H, br s), 7.79 (1H, br s), 7.76 (1H, br s), 7.62 (1H, br s), 4.35-4.30 (1H, m), 3.94 (3H, s), 3.54-3.48 (1H, m), 3.43 (6H, s), 1.71-1.56 (2H, m), 1.39-1.27 (1H, m), 1.21 (3H, d, J = 6.6 Hz), 0.95 (3H, d, J = 6.6 Hz), 0.84 (3H, d, J = 6.6 Hz).
Example 2-4		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	429	0.89	
Example 2-5		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((2-methoxypyridin-4-yl)amino)nicotinamide	389	0.72	
Example 2-6		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide	419	1.07	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ HNMR
Example 2-7		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	433	0.79	
Example 2-8		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-(quinolin-6-ylamino)nicotinamide	0.78	409	
Example 2-9		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	426	1.01	
Example 2-10		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((1-methyl-1H-indazol-6-yl)amino)nicotinamide	412	0.96	
Example 2-11		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((1-ethyl-1H-indazol-6-yl)amino)-5-fluoronicotinamide	428	1.07	
Example 2-12		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-4-yl)amino)nicotinamide	414	1.05	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-13		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-2-methyl-2H-indazol-5-yl)amino)-5-fluoronicotinamide	441	0.85	
Example 2-14		6-(((2R,3S)-3-amino-1-cyclopropylbutan-2-yl)amino)-2-((3-(dimethylamino)-2-methyl-2H-indazol-5-yl)amino)-5-fluoronicotinamide	455	0.88	
Example 2-15		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((3-(dimethylamino)-2-methyl-2H-indazol-5-yl)amino)-5-fluoronicotinamide	457	0.94	
Example 2-16		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(dimethylamino)-2-methyl-2H-indazol-5-yl)amino)-5-fluoronicotinamide	429	0.82	
Example 2-17		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((5,6-dimethylpyridin-3-yl)amino)-5-fluoronicotinamide	361	0.62	
Example 2-18		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((5,6-dimethylpyridin-3-yl)amino)-5-fluoronicotinamide	389	0.77	¹ H-NMR (MeOD) δ : 9.34 (1H, s), 8.14 (1H, s), 7.82 (1H, d, J =11.7 Hz), 4.65-4.55 (1H, m), 3.58-3.49 (1H, m), 2.67 (3H, s), 2.47 (3H, s), 1.72-1.33 (6H, m), 0.98 (3H, d, 6.3 Hz), 0.91 (3H, d, 6.3 Hz).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-19		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((5,6-dimethylpyridin-3-yl)amino)-5-fluoronicotinamide	389	0.76	
Example 2-20		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide	448	0.98	
Example 2-21		6-(((2R,3S)-3-amino-1-cyclopropylbutan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide	462	1	
Example 2-22		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide	464	1.05	
Example 2-23		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide	436	0.94	
Example 2-24		2-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoronicotinamide	425	0.97	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-25		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-(m-tolylamino)nicotinamide	372	1.1	
Example 2-26		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3,5-dimethylphenyl)amino)-5-fluoronicotinamide	386	1.18	
Example 2-27		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((3-methoxyphenyl)amino)nicotinamide	388	1.05	
Example 2-28		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide	418	1.07	¹ H-NMR (DMSO-d ₆) δ: 11.58 (1H, m), 7.91-7.60 (5H, m), 6.90-6.40 (4H, m), 5.17-6.13 (1H, m), 4.47-4.33 (1H, m), 3.80 (3H, s), 3.74 (3H, s), 3.35-3.26 (1H, m), 2.74-2.60 (1H, m), 2.10-1.60 (6H, m), 1.17 (3H, d, J = 7.5 Hz).
Example 2-29		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3,4-dimethoxyphenyl)amino)-5-fluoronicotinamide	418	0.97	¹ H-NMR (DMSO-d ₆) δ: 11.35 (1H, s), 7.91-7.40 (5H, m), 7.25-6.77 (5H, m), 4.36-4.25 (1H, m), 3.80 (3H, s), 3.74 (3H, s), 3.25-3.35 (1H, m), 2.74-2.60 (1H, m), 2.10-1.62 (6H, m), 1.14 (3H, d, J = 7.5 Hz).
Example 2-30		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((3-methoxy-4-methylphenyl)amino)nicotinamide	402	1.13	
Example 2-31		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	414	0.86	¹ H-NMR (MeOD) δ: 9.30 (1H, d, J = 2.3 Hz), 8.83-8.78 (1H, m), 8.15 (2H, s), 7.87 (1H, d, J = 11.9 Hz), 4.72-4.63 (1H, m), 3.66-3.55 (1H, m), 2.84 (3H, s), 1.89-1.57 (2H, m), 1.31 (3H, d, J = 6.6 Hz), 0.98 (3H, t, J = 7.3 Hz).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-32		6-(((1R,2S)-2-amino-1-(4-fluorophenyl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	480	0.96	
Example 2-33		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	414	0.83	
Example 2-34		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	407	0.67	
Example 2-35		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((5,6-dimethylpyridin-3-yl)amino)-5-fluoronicotinamide	361	0.52	
Example 2-36		6-(((3R,4S)-4-aminohexan-3-yl)amino)-2-((5,6-dimethylpyridin-3-yl)amino)-5-fluoronicotinamide	375	0.64	
Example 2-37		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-6-yl)amino)nicotinamide	414	1.02	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-38		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-6-yl)amino)nicotinamide	414	1.01	
Example 2-39		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-6-yl)amino)nicotinamide	386	0.85	
Example 2-40		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-5-yl)amino)nicotinamide	386	0.83	¹ H-NMR (MeOD) δ: 7.94 (1H, s), 7.92-7.89 (1H, m), 7.74 (1H, d, J = 11.9 Hz), 7.53 (1H, d, J = 8.9 Hz), 7.42 (1H, dd, J = 8.9, 2.0 Hz), 4.36-4.24 (1H, m), 4.06 (3H, s), 3.46-3.33 (1H, m), 1.67-1.40 (2H, m), 1.28 (3H, d, J = 7.3 Hz), 0.68 (3H, t, J = 7.6 Hz).
Example 2-41		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((5-cyclopropylpyridin-3-yl)amino)-5-fluoronicotinamide	373	0.64	
Example 2-42		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	407	0.65	
Example 2-43		6-(((3R,4S)-4-aminohexan-3-yl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	421	0.71	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-44		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-(m-tolylamino)nicotinamide	358	1.03	¹ H-NMR (DMSO-d ₆) δ: 11.52 (1H, s), 7.95-7.05 (10H, m), 6.76 (1H, d, J = 7.3 Hz), 3.72-3.61 (1H, m), 3.50-3.42 (1H, m), 2.26 (3H, s), 1.26 (3H, d, J = 8.4 Hz), 1.21-1.08 (1H, m), 0.66-0.34 (4H, m)
Example 2-45		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3,5-dimethylphenyl)amino)-5-fluoronicotinamide	372	1.11	
Example 2-46		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-methoxyphenyl)amino)nicotinamide	374	0.98	¹ H-NMR (DMSO-d ₆) δ: 11.57 (1H, s), 7.95-7.50 (5H, m), 7.30-7.00 (5H, m), 6.64 (1H, d, J = 7.8 Hz), 3.74 (3H, s), 3.72-3.61 (1H, m), 3.50-3.42 (1H, m), 1.29 (3H, d, J = 8.4 Hz), 1.16-1.00 (1H, m), 0.60-0.25 (4H, m)
Example 2-47		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide	404	1	¹ H-NMR (DMSO-d ₆) δ: 11.65 (1H, s), 7.88-6.90 (7H, m), 6.68 (2H, d, J = 3.0 Hz), 6.17-6.13 (1H, m), 3.72 (6H, s), 3.72-3.61 (1H, m), 3.60-3.45 (1H, m), 1.29 (3H, d, J = 8.4 Hz), 1.16-1.00 (1H, m), 0.70-0.28 (4H, m)
Example 2-48		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3,4-dimethoxyphenyl)amino)-5-fluoronicotinamide	404	0.89	¹ H-NMR (DMSO-d ₆) δ: 11.31 (1H, s), 7.90-7.50 (5H, m), 7.20-6.65 (6H, m), 3.75 (3H, s), 3.73 (3H, s), 3.68-3.58 (1H, m), 3.60-3.45 (1H, m), 1.23 (3H, d, J = 8.4 Hz), 1.16-1.00 (1H, m), 0.64-0.25 (4H, m)
Example 2-49		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-methoxy-4-methylphenyl)amino)nicotinamide	388	1.06	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-50		6-(((2R,3S)-3-amino-1-cyclopropylbutan-2-yl)amino)-2-((1-ethyl-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide	456	1.1	
Example 2-51		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((1-ethyl-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide	458	1.16	
Example 2-52		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((1-ethyl-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide	430	1.01	¹ H-NMR (MeOD) δ: 7.92 (1H, d, J = 2.0 Hz), 7.75 (1H, d, J = 11.9 Hz), 7.37 (1H, d, J = 9.2 Hz), 7.29 (1H, dd, J = 9.2, 2.0 Hz), 4.25 (2H, q, J = 7.3 Hz), 4.24-4.18 (1H, m), 4.06 (3H, s), 3.48 (1H, dd, J = 6.9, 3.6 Hz), 1.76-1.56 (2H, m), 1.38 (3H, t, J = 7.3 Hz), 1.22 (3H, d, J = 6.6 Hz), 1.07 (3H, t, J = 7.3 Hz).
Example 2-53		6-(((2R,3S)-3-amino-1-cyclopropylbutan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide	492	1.11	
Example 2-54		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide	494	1.15	
Example 2-55		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide	466	1.03	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-56		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((3-methoxybenzo[d]isoxazol-5-yl)amino)nicotinamide	431	1.15	
Example 2-57		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((3-methoxybenzo[d]isoxazol-5-yl)amino)nicotinamide	403	0.87	
Example 2-58		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((4-fluorophenyl)amino)nicotinamide	362	0.98	
Example 2-59		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3,4-difluorophenyl)amino)-5-fluoronicotinamide	380	1.03	
Example 2-60		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((4-fluorophenyl)amino)nicotinamide	376	1.07	
Example 2-61		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3,4-difluorophenyl)amino)-5-fluoronicotinamide	394	1.13	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-62		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((5-fluoro-6-methoxypyridin-3-yl)amino)nicotinamide	393	0.94	¹ H-NMR (DMSO-d ₆) δ: 11.41 (1H, s), 8.02 (1H, dd, J = 2.4, 12.9 Hz), 7.95-7.65 (6H, m), 7.40-7.10 (2H, m), 3.91 (3H, s), 3.65-3.40 (2H, m), 1.25 (3H, d, J = 6.6 Hz), 1.14-1.10 (1H, m), 0.68-0.54 (1H, m), 0.48-0.22 (3H, m)
Example 2-63		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((5-fluoro-6-methoxypyridin-3-yl)amino)nicotinamide	407	1.02	¹ H-NMR (DMSO-d ₆) δ: 11.44 (1H, s), 8.18 (1H, dd, J = 2.1, 12.6 Hz), 7.99 (1H, d, J = 2.1 Hz), 7.91 (1H, d, J = 12.6 Hz), 7.85-7.60 (4H, m), 7.42-7.10 (1H, m), 6.88 (1H, d, J = 9.9 Hz), 4.37-4.25 (1H, m), 3.92 (3H, s), 3.40-3.25 (1H, m), 2.70-2.54 (1H, m), 2.10-1.62 (6H, m), 1.15 (3H, d, J = 6.6 Hz)
Example 2-64		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-methoxypyridin-3-yl)amino)nicotinamide	381	0.93	
Example 2-65		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-methoxypyridin-3-yl)amino)nicotinamide	395	1.02	¹ H-NMR (DMSO-d ₆) δ: 11.44 (1H, s), 8.08-7.97 (2H, m), 7.92 (1H, d, J = 12.6 Hz), 7.92-7.65 (4H, m), 7.40-7.16 (1H, m), 6.99 (1H, d, J = 8.4 Hz), 4.20-4.05 (1H, m), 3.91 (3H, s), 3.46-3.25 (1H, m), 1.64-1.48 (2H, m), 1.47-1.13 (5H, m), 0.85 (3H, t, J = 7.4 Hz)
Example 2-66		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-methoxypyridin-3-yl)amino)nicotinamide	409	1.08	¹ H-NMR (DMSO-d ₆) δ: 11.40 (1H, s), 8.08-7.97 (2H, m), 7.91 (1H, d, J = 12.6 Hz), 7.85-7.62 (4H, m), 7.40-7.10 (1H, m), 7.00 (1H, d, J = 8.4 Hz), 4.26-4.12 (1H, m), 3.90 (3H, s), 3.40-3.26 (1H, m), 1.66-1.48 (2H, m), 1.38-1.24 (1H, m), 1.17 (3H, d, J = 6.6 Hz), 0.89 (3H, d, J = 6.6 Hz), 0.76 (3H, d, J = 6.6 Hz)
Example 2-67		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((5-fluoro-6-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	430	0.9	¹ H-NMR (DMSO-d ₆) δ: 12.23 (1H, s), 8.58-8.47 (2H, m), 8.17 (2H, s), 8.03 (1H, d, J = 12.6 Hz), 8.02-7.80 (4H, m), 7.58-7.40 (1H, m), 7.11 (1H, d, J = 6.6 Hz), 4.37-4.22 (1H, m), 3.76-3.62 (1H, m), 2.02-1.80 (2H, m), 1.80-1.54 (4H, m), 1.52-1.34 (2H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-68		6-(((1R,2S)-2-amino-2-cyclohexylamino)-5-fluoro-2-((5-fluoro-6-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	430	0.86	¹ H-NMR (DMSO-d ₆) δ: 12.24 (1H, s), 8.67-8.48 (3H, m), 8.15-7.75 (6H, m), 7.60-7.40 (1H, m), 7.11 (1H, d, J = 6.6 Hz), 4.38-4.23 (1H, m), 3.75-3.50 (1H, m), 2.04-1.80 (2H, m), 1.80-1.54 (4H, m), 1.54-1.32 (2H, m)
Example 2-69		2-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoronicotinamide	427	1.02	
Example 2-70		2-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoronicotinamide	427	1.02	
Example 2-71		6-(((1R,2S)-2-amino-1-cyclopropylpropylamino)-2-((1-(2,2-difluoroethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	448	0.94	
Example 2-72		6-(((1R,2S)-2-amino-1-cyclopropylpropylamino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	441	0.9	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-73		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-ethyl-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide	442	1.01	
Example 2-74		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-(2,2-difluoroethyl)-3-methoxy-1H-indazol-5-yl)amino)-5-fluoronicotinamide	478	0.97	
Example 2-75		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide	448	0.97	
Example 2-76		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	448	0.87	¹ H-NMR (DMSO-d ₆) δ: 11.46 (1H, s), 8.10-7.66 (6H, m), 7.45-6.96 (3H, m), 3.74 (4H, t, J = 4.6 Hz), 3.69-3.68 (1H, m), 3.60-3.45 (1H, m), 3.24 (4H, t, J = 4.6 Hz), 1.23 (3H, d, J = 8.4 Hz), 1.13-1.00 (1H, m), 0.68-0.20 (4H, m)
Example 2-77		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	462	0.96	¹ H-NMR (DMSO-d ₆) δ: 11.49 (1H, s), 8.20-7.69 (7H, m), 7.40-6.75 (3H, m), 4.45-4.39 (m, 1H), 3.74 (4H, t, J = 4.6 Hz), 3.35-3.25 (4H, m), 2.74-2.60 (1H, m), 2.10-1.62 (6H, m), 1.14 (3H, d, J = 7.5 Hz)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-78		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	430	1.06	¹ H-NMR (MeOD) δ: 7.96 (1H, br s), 7.73 (1H, d, J = 11.9 Hz), 7.32 (1H, br s), 7.52 (1H, br s), 4.44-4.37 (1H, m), 4.05 (3H, s), 3.84 (3H, s), 3.18-2.98 (2H, m), 1.77-1.60 (2H, m), 1.46-1.35 (1H, m), 0.93 (6H, t, J = 7.0 Hz).
Example 2-79		(R)-6-((1-amino-3-cyclopropylpropan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	428	1.02	
Example 2-80		6-(((2S,3R)-2-amino-3-aminohexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	430	1	
Example 2-81		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	430	1	
Example 2-82		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	442	1.05	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-83		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	444	1.1	¹ H-NMR (MeOD) δ: 7.92 (1H, d, J = 1.3 Hz), 7.75 (1H, d, J = 11.9 Hz), 7.35 (1H, d, J = 9.2 Hz), 7.29 (1H, dd, J = 8.9, 1.7 Hz), 4.34-4.27 (1H, m), 4.04 (3H, s), 3.85 (3H, s), 3.46-3.41 (1H, m), 1.68-1.58 (2H, m), 1.51-1.34 (4H, m), 1.22 (3H, d, J = 7.3 Hz), 0.93 (3H, t, J = 6.9 Hz).
Example 2-84		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	416	0.93	
Example 2-85		6-(((3R,4S)-4-aminohexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	430	0.97	
Example 2-86		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	443	1.04	
Example 2-87		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	443	1	
Example 2-88		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	443	0.98	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-89		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	455	1.02	¹ H-NMR (MeOD) δ: 7.99 (1H, d, J = 9.2 Hz), 7.84 (1H, s), 7.77 (1H, d, J = 11.9 Hz), 7.63 (1H, d, J = 8.6 Hz), 4.29 (1H, dd, J = 9.9, 3.3 Hz), 4.14-4.05 (1H, m), 3.95 (3H, s), 3.40 (6H, s), 2.69-2.64 (1H, m), 2.13-2.02 (2H, m), 1.98-1.78 (4H, m), 1.18 (3H, d, J = 7.3 Hz).
Example 2-90		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	457	1.07	
Example 2-91		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	429	0.88	
Example 2-92		6-(((3R,4S)-4-aminohexan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	443	0.95	
Example 2-93		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	450	0.99	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HMR
Example 2-94		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	428	0.95	
Example 2-95		2-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoronicotinamide	413	0.94	
Example 2-96		6-(((2S,3R)-2-amino-4-aminohexan-3-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	414	1.01	
Example 2-97		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	414	0.97	
Example 2-98		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	426	1.02	
Example 2-99		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	428	1.08	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-100		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	400	0.93	
Example 2-101		6-(((3R,4S)-4-aminohexan-3-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	414	0.98	¹ H-NMR (MeOD) δ: 8.06 (1H, s), 7.83 (1H, d, J = 11.0 Hz), 7.80 (1H, d, J = 6.6 Hz), 7.38 (1H, t, J = 7.9 Hz), 7.19 (1H, d, J = 8.6 Hz), 4.45 (2H, q, J = 7.0 Hz), 4.31 (1H, td, J = 7.3, 3.3 Hz), 3.42-3.34 (1H, m), 1.81-1.67 (2H, m), 1.67-1.53 (2H, m), 1.47 (3H, t, J = 7.0 Hz), 1.08 (3H, t, J = 7.3 Hz), 0.93 (3H, t, J = 7.6 Hz).
Example 2-102		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-6-yl)amino)nicotinamide	400	0.93	
Example 2-103		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-2-((1-ethyl-1H-indazol-6-yl)amino)-5-fluoronicotinamide	414	0.99	
Example 2-104		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-5-yl)amino)nicotinamide	400	0.91	
Example 2-105		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	414	0.98	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-106		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoro-2-(quinolin-6-ylamino)nicotinamide	397	0.72	
Example 2-107		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoro-2-((2-methoxypyridin-4-yl)amino)nicotinamide	377	0.6	
Example 2-108		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide	407	1.01	
Example 2-109		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	421	0.72	
Example 2-110		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-2-((5,6-dimethylpyridin-3-yl)amino)-5-fluoronicotinamide	375	0.68	
Example 2-111		6-(((2S,3R)-2-amino-4-methylpentan-3-yl)amino)-2-((5-cyclopropylpyridin-3-yl)amino)-5-fluoronicotinamide	387	0.76	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-112		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	458	1.02	¹ H-NMR (DMSO-d ₆) δ: 11.56 (1H, s), 8.53 (1H, d, J = 2.7 Hz), 8.27 (1H, d, J = 2.7 Hz), 8.12 (2H, s), 7.93 (1H, d, J = 12.3 Hz), 7.85-7.55 (4H, m), 7.45-7.15 (1H, m), 7.03 (1H, d, J = 8.7 Hz), 4.34-4.26 (1H, m), 3.89 (3H, s), 3.40-3.20 (1H, m), 1.64-1.44 (2H, m), 1.34-1.12 (1H, m), 1.02 (3H, d, J = 7.2 Hz), 0.84 (3H, d, J = 6.0 Hz), 0.63 (3H, d, J = 5.7 Hz)
Example 2-113		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	442	0.89	¹ H-NMR (DMSO-d ₆) δ: 11.51 (1H, s), 8.39 (1H, d, J = 2.7 Hz), 8.26 (1H, d, J = 2.7 Hz), 8.12 (2H, s), 8.05-7.70 (5H, m), 7.40-7.17 (1H, m), 7.17 (1H, d, J = 8.7 Hz), 3.89 (3H, s), 3.68-3.54 (1H, m), 3.50-3.30 (1H, m), 1.14 (3H, d, J = 6.6 Hz), 1.08-0.94 (1H, m), 0.54-0.42 (1H, m), 0.42-0.30 (1H, m), 0.26-0.10 (2H, m)
Example 2-114		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	456	0.98	¹ H-NMR (DMSO-d ₆) δ: 11.56 (1H, s), 8.58 (1H, d, J = 2.3 Hz), 8.27 (1H, d, J = 2.3 Hz), 8.14 (2H, s), 7.92 (1H, d, J = 12.6 Hz), 7.85-7.55 (4H, m), 7.50-7.10 (1H, m), 6.91 (1H, d, J = 8.7 Hz), 4.40-4.28 (1H, m), 3.90 (3H, s), 3.40-3.26 (1H, m), 2.62-2.50 (1H, m), 1.96-1.80 (2H, m), 1.80-1.62 (4H, m), 1.06 (3H, d, J = 7.5 Hz)
Example 2-115		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)nicotinamide	437	0.95	
Example 2-116		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)nicotinamide	451	1.03	¹ H-NMR (DMSO-d ₆) δ: 11.44 (1H, s), 8.18 (1H, dd, J = 2.3, 12.6 Hz), 7.98 (1H, d, J = 2.3 Hz), 7.91 (1H, d, J = 12.6 Hz), 7.88-7.68 (4H, m), 7.40-7.12 (1H, m), 6.90 (1H, d, J = 8.7 Hz), 4.46-4.39 (2H, m), 4.37-4.25 (1H, m), 3.72-3.65 (2H, m), 3.40-3.26 (4H, m), 2.75-2.55 (1H, m), 2.15-1.62 (6H, m), 1.16 (3H, d, J = 6.6 Hz)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-117		6-(((2S,3R)-2-amino-3-ethylpentan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)nicotinamide	425	0.94	
Example 2-118		6-(((2S,3R)-2-amino-3-hexylpentan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)nicotinamide	439	1.02	
Example 2-119		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-(2-methoxyethoxy)pyridin-3-yl)amino)nicotinamide	453	1.08	
Example 2-120		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	445	0.81	
Example 2-121		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	432	0.84	
Example 2-122		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	416	0.87	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-123		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	472	0.94	¹ H-NMR (MeOD) δ: 8.10 (1H, s), 7.74 (1H, d, J = 11.9 Hz), 7.36 (1H, d, J = 8.6 Hz), 7.23 (1H, d, J = 9.2 Hz), 4.61-4.57 (1H, m), 4.34 (2H, t, J = 5.3 Hz), 4.06 (3H, s), 3.76 (2H, t, J = 5.3 Hz), 3.58-3.54 (1H, m), 3.13 (3H, s), 1.52-1.37 (8H, m).
Example 2-124		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	474	1.02	
Example 2-125		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	460	0.9	
Example 2-126		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	474	0.98	
Example 2-127		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	488	1.11	
Example 2-128		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	476	0.93	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-129		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	460	0.94	
Example 2-130		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide	458	1.02	
Example 2-131		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide	444	0.9	
Example 2-132		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide	458	0.99	
Example 2-133		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide	456	0.94	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-134		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide	472	1.05	
Example 2-135		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide	460	0.88	¹ H-NMR (MeOD) δ: 7.86 (1H, s), 7.77 (1H, d, J = 11.9 Hz), 7.52 (1H, br s), 7.52 (1H, br s), 4.62 (2H, t, J = 5.3 Hz), 4.44-4.38 (1H, m), 3.79 (2H, t, J = 5.3 Hz), 3.78-3.74 (1H, m), 3.65-3.55 (2H, m), 3.40 (3H, s), 3.28 (3H, s), 2.58 (3H, s), 1.21 (3H, d, J = 6.6 Hz).
Example 2-136		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide	444	0.88	
Example 2-137		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	464	1.03	
Example 2-138		6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	464	1.02	
Example 2-139		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	466	0.92	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-140		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	502	1.07	
Example 2-141		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	458	1.05	
Example 2-142		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	377	0.89	¹ H-NMR (DMSO-d ₆) δ: 11.84 (1H, s), 8.46-8.34 (1H, m), 8.20-8.11 (1H, m), 8.00-7.75 (5H, m), 7.46-7.28 (1H, m), 7.01 (1H, d, J = 7.2 Hz), 4.30-4.16 (1H, m), 3.60-3.40 (1H, m), 2.39 (3H, d, J = 2.7 Hz), 2.00-1.80 (2H, m), 1.79-1.54 (4H, m), 1.54-1.34 (2H, m)
Example 2-143		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-4-yl)amino)nicotinamide	427	1.06	
Example 2-144		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-4-yl)amino)nicotinamide	444	1.36	
Example 2-145		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-4-yl)amino)nicotinamide	415	1.07	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-146		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-4-yl)amino)nicotinamide	428	1.03	
Example 2-147		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-4-yl)amino)nicotinamide	432	0.95	
Example 2-148		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((7-chloro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	420	1.01	
Example 2-149		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((7-chloro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	436	0.97	
Example 2-150		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	446	1.06	
Example 2-151		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	448	1.12	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-152		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	434	1.04	
Example 2-153		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	448	1.06	
Example 2-154		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	446	1.07	
Example 2-155		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	462	1.17	
Example 2-156		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	462	1.15	
Example 2-157		6-(((2R,3S)-3-amino-1-cyclopropylbutan-2-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	460	1.13	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-158		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	460	1.13	
Example 2-159		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	450	1	
Example 2-160		6-(((3R,4S)-4-aminohexan-3-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	448	1.04	
Example 2-161		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	486	1	
Example 2-162		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	442	0.98	¹ H-NMR (DMSO-d ₆) δ: 11.63 (1H, s), 8.74 (1H, d, J = 2.4 Hz), 8.65-8.61 (1H, m), 8.22 (1H, d, J = 2.4 Hz), 8.00-7.97 (1H, m), 7.94 (1H, d, J = 12.3 Hz), 7.90-7.60 (4H, m), 7.40-7.20 (1H, m), 7.19 (1H, d, J = 8.7 Hz), 3.96 (3H, s), 3.80-3.66 (1H, m), 3.54-3.40 (1H, m), 1.15 (3H, d, J = 6.6 Hz), 1.10-0.96 (1H, m), 0.55-0.28 (3H, m), 0.20-0.10 (1H, m)
Example 2-163		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	456	1.07	¹ H-NMR (DMSO-d ₆) δ: 11.63 (1H, s), 8.75 (1H, d, J = 2.4 Hz), 8.66-8.62 (1H, m), 8.20 (1H, d, J = 2.4 Hz), 8.02-7.62 (6H, m), 7.40-7.12 (2H, m), 4.42 (2H, q, J = 6.9 Hz), 3.80-3.68 (1H, m), 3.56-3.40 (1H, m), 1.34 (3H, t, J = 6.9 Hz), 1.15 (3H, d, J = 6.6 Hz), 1.12-0.98 (1H, m), 0.54-0.30 (3H, m), 0.22-0.10 (1H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-164		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	455	1.2	
Example 2-165		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	500	1.04	
Example 2-166		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	456	1.07	
Example 2-167		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	470	1.17	¹ H-NMR (DMSO-d ₆) δ: 11.66 (1H, s), 8.95 (1H, d, J = 2.7 Hz), 8.68-8.65 (1H, m), 8.22 (1H, d, J = 2.7 Hz), 8.01-7.98 (1H, m), 7.93 (1H, d, J = 12.3 Hz), 7.85-7.45 (4H, m), 7.45-7.15 (1H, m), 6.93 (1H, d, J = 8.4 Hz), 4.67-4.46 (1H, m), 4.43 (2H, q, J = 6.9 Hz), 3.46-3.25 (1H, m), 2.66-2.50 (1H, m), 1.95-1.62 (6H, m), 1.35 (3H, t, J = 6.9 Hz), 1.08 (3H, d, J = 6.6 Hz)
Example 2-168		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	469	1.32	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-169		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	464	1.08	¹ H-NMR (MeOD) δ: 8.24 (1H, d, J = 1.3 Hz), 7.77 (1H, d, J = 11.9 Hz), 7.60-7.69 (1H, m), 7.44 (1H, d, J = 9.2 Hz), 7.31-7.29 (1H, m), 4.45-4.42 (1H, m), 3.93 (3H, s), 3.83-3.80 (1H, m), 1.87-1.67 (8H, m).
Example 2-170		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	452	1.04	
Example 2-171		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	464	1.06	
Example 2-172		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	480	1.19	
Example 2-173		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	468	1	
Example 2-174		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	508	1.14	¹ H-NMR (MeOD) δ: 8.21 (1H, d, J = 2.0 Hz), 7.77 (1H, d, J = 11.9 Hz), 7.48 (1H, d, J = 9.2 Hz), 7.35-7.25 (1H, m), 7.30-7.26 (1H, m), 4.45-4.42 (1H, m), 4.42 (2H, t, J = 5.3 Hz), 3.83-3.80 (1H, m), 3.78 (2H, t, J = 5.0 Hz), 3.28 (3H, s), 1.81-1.66 (8H, m).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-175		6-(((2S,3R)-2-amino-3-aminopentan-3-yl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	496	1.11	
Example 2-176		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	508	1.12	
Example 2-177		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	524	1.26	
Example 2-178		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	512	1.07	¹ H-NMR (MeOD) δ: 7.97 (1H, d, J = 1.3 Hz), 7.77 (1H, d, J = 11.9 Hz), 7.48 (1H, d, J = 9.2 Hz), 7.36 (1H, dd, J = 9.2, 2.0 Hz), 7.13 (1H, d, J = 73.3 Hz), 4.46-4.39 (1H, m), 4.42 (2H, t, J = 5.0 Hz), 3.87 (1H, dd, J = 10.2, 3.6 Hz), 3.79 (2H, t, J = 5.0 Hz), 3.72-3.62 (2H, m), 3.42 (3H, s), 3.28 (3H, s), 1.32 (3H, d, J = 7.3 Hz).
Example 2-179		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-5-fluoro-2-((1-methyl-1H-indazol-5-yl)amino)nicotinamide	416	0.93	
Example 2-180		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	430	1.04	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-181		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	466	0.99	
Example 2-182		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	430	1	
Example 2-183		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	444	0.95	
Example 2-184		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	459	1.1	¹ H-NMR (MeOD) δ: 7.61 (2H, m), 7.60 (1H, br s), 4.31-4.28 (1H, m), 4.04 (3H, br s), 3.78-3.77 (1H, m), 2.06 (6H, br s), 2.00-1.56 (6H, m).
Example 2-185		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	461	1.15	
Example 2-186		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	447	1.05	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-187		6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	461	1.13	
Example 2-188		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	459	1.07	¹ H-NMR (MeOD) δ: 7.81-7.77 (2H, m), 7.47 (1H, br s), 4.21-4.10 (1H, m), 4.07 (3H, br s), 3.67-3.58 (1H, m), 3.40-3.34 (6H, m), 1.35 (3H, d, J = 8.6 Hz), 1.15-1.00 (1H, m), 0.81-0.72 (1H, m), 0.66-0.57 (1H, m), 0.50-0.35 (2H, m).
Example 2-189		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	475	1.2	
Example 2-190		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	475	1.22	
Example 2-191		6-(((2R,3S)-3-amino-1-cyclopropylbutan-2-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	473	1.14	
Example 2-192		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	471	1.16	¹ H-NMR (MeOD) δ: 7.76 (1H, d, J = 13.2 Hz), 7.69 (1H, d, J = 16.2 Hz), 7.52 (1H, s), 4.39-4.36 (1H, m), 4.09 (3H, s), 3.67-3.58 (1H, m), 3.21-3.11 (6H, m), 2.70-2.55 (1H, m), 1.96-1.86 (4H, m), 1.32 (3H, s), 1.20 (2H, d, J = 7.3 Hz).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-193		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	463	1.02	¹ H-NMR (MeOD) δ: 7.77 (1H, d, J = 11.9 Hz), 7.64 (1H, d, J = 16.2 Hz), 7.49 (1H, d, J = 13.9 Hz), 4.37-4.32 (1H, m), 3.99 (3H, s), 3.67-3.69 (1H, m), 3.56-3.61 (2H, m), 3.41 (3H, s), 3.07 (6H, br s), 1.33 (3H, s).
Example 2-194		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	447	1.05	¹ H-NMR (MeOD) δ: 7.76 (1H, d, J = 12.6 Hz), 7.74 (1H, d, J = 13.0 Hz), 7.52 (1H, s), 4.42-4.36 (1H, m), 4.04 (3H, s), 3.42-3.36 (1H, m), 3.26 (6H, s), 1.73-1.63 (2H, m), 1.30 (3H, d, J = 6.6 Hz), 0.81 (3H, t, J = 7.6 Hz).
Example 2-195		6-(((3R,4S)-4-aminohexan-3-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	461	1.12	
Example 2-196		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((1,3-dimethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	416	0.9	
Example 2-197		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	430	0.88	¹ H-NMR (MeOD) δ: 9.41 (1H, d, J = 2.6 Hz), 8.98 (1H, s), 8.19 (2H, s), 7.91 (1H, d, J = 11.9 Hz), 3.85-3.62 (4H, m), 3.39 (3H, s), 2.94 (3H, s), 1.37 (3H, d, J = 6.6 Hz).
Example 2-198		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	452	0.93	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-199		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	472	1.17	
Example 2-200		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	443	1.17	
Example 2-201		6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((6-ethoxy-5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	457	1.25	
Example 2-202		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	474	0.95	
Example 2-203		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((6-(2-methoxyethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	488	1.02	
Example 2-204		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	430	0.92	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-205		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide.	444	1	¹ H-NMR (DMSO-d ₆) δ: 11.63 (1H, s), 8.83 (1H, d, J = 2.6 Hz), 8.64 (1H, s), 8.27 (1H, d, J = 2.6 Hz), 7.98 (1H, s), 7.94 (1H, d, J = 12.3 Hz), 7.86-7.50 (4H, m), 7.45-7.15 (1H, m), 7.05 (1H, d, J = 9.0 Hz), 4.47-4.32 (1H, m), 3.40-3.25 (1H, m), 3.96 (3H, s), 1.60-1.42 (2H, m), 1.40-1.10 (2H, m), 1.06 (3H, d, J = 6.6 Hz), 0.75 (3H, t, J = 7.7 Hz)
Example 2-206		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	444	1.01	
Example 2-207		6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((6-ethoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	458	1.09	
Example 2-208		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1,3-dimethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	414	0.79	¹ H-NMR (MeOD) δ: 7.78-7.63 (3H, m), 7.50-7.51 (1H, m), 4.24 (1H, t, J = 8.3, 4.4 Hz), 4.10-4.00 (1H, m), 4.05 (3H, s), 3.74-3.53 (3H, m), 2.60 (3H, s), 2.10-1.53 (3H, m)
Example 2-209		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-fluoro-1-methyl-1H-indazol-5-yl)amino)nicotinamide	418	0.87	
Example 2-210		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	430	0.84	¹ H-NMR (MeOD) δ: 8.10 (1H, d, J = 2.0 Hz), 7.75 (1H, d, J = 11.9 Hz), 7.31 (1H, d, J = 9.2 Hz), 7.21 (1H, dd, J = 8.6, 2.0 Hz), 4.40 (1H, t, J = 8.4, 4.2 Hz), 4.15-4.05 (4H, m), 3.98-3.55 (7H, m), 2.14-1.80 (2H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-211		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((7-fluoro-3-methoxy-1-methyl-1H-indazol-5-yl)amino)nicotinamide	448	0.93	¹ H-NMR (MeOD) δ : 8.29 (1H, d, J = 1.7 Hz), 8.25 (1H, d, J = 11.9 Hz), 7.63 (1H, dd, J = 11.9, 1.7 Hz), 4.93 (1H, td, J = 8.1, 4.6 Hz), 4.65-4.12 (1H, m), 2.65-2.37 (2H, m)
Example 2-212		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	443	0.81	
Example 2-213		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	461	0.9	
Example 2-214		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	428	0.75	¹ H-NMR (MeOD) δ : 9.26 (1H, d, J = 2.5 Hz), 8.88 (1H, d, J = 2.5 Hz), 8.20 (2H, s), 7.90 (1H, d, J = 11.9 Hz), 4.64-4.53 (1H, m), 4.14-4.04 (1H, m), 3.90-3.50 (4H, m), 2.90 (3H, s), 2.20-1.86 (2H, m)
Example 2-215		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	444	0.79	¹ H-NMR (MeOD) δ : 8.82 (1H, d, J = 2.6 Hz), 8.60 (1H, d, J = 1.0 Hz), 8.17 (1H, d, J = 2.6 Hz), 7.95 (1H, d, J = 1.0 Hz), 7.79 (1H, d, J = 11.9 Hz), 4.56-4.44 (1H, m), 4.12-3.96 (1H, m), 4.06 (3H, s), 3.80-3.60 (3H, m), 3.45-3.35 (1H, m), 2.12-1.82 (2H, m)
Example 2-216		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	444	0.79	¹ H-NMR (MeOD) δ : 8.38 (1H, d, J = 2.5 Hz), 8.25 (1H, d, J = 2.5 Hz), 8.02 (2H, s), 7.78 (1H, d, J = 11.9 Hz), 4.40-4.30 (1H, m), 4.08-3.95 (1H, m), 3.97 (3H, s), 3.80-3.67 (2H, m), 3.62-3.40 (2H, m), 2.10-1.76 (2H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-217		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	450	0.81	¹ H-NMR (MeOD) δ: 8.28 (1H, d, J = 2.3 Hz), 8.24 (1H, dd, J = 14.9, 2.3 Hz), 7.82 (1H, d, J = 11.9 Hz), 4.56-4.40 (1H, m), 4.18-3.65 (9H, m), 3.58-3.50 (4H, m), 2.20-1.83 (2H, m)
Example 2-218		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinamide	361	0.5	¹ H-NMR (MeOD) δ: 9.22 (1H, d, J = 2.0 Hz), 8.28-8.20 (2H, m), 7.89 (1H, d, J = 11.6 Hz), 4.60-4.45 (1H, m), 4.18-4.08 (1H, m), 4.06-3.97 (1H, m), 3.88-3.76 (3H, m), 2.53 (3H, s), 2.23-1.85 (2H, m)
Example 2-219		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((6-methylpyridin-3-yl)amino)nicotinamide	361	0.48	¹ H-NMR (MeOD) δ: 9.15 (1H, d, J = 2.3 Hz), 8.47 (1H, dd, J = 8.8, 2.3 Hz), 7.89 (1H, d, J = 11.9 Hz), 7.81 (1H, d, J = 8.8 Hz), 4.57-4.46 (1H, m), 4.18-4.08 (1H, m), 4.06-3.72 (4H, m), 2.73 (3H, s), 2.23-1.87 (2H, m)
Example 2-220		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	377	0.83	¹ H-NMR (DMSO-d ₆) δ: 11.92 (1H, s), 8.54 (1H, s), 8.15-7.70 (6H, m), 7.46-7.26 (2H, m), 3.74-3.60 (1H, m), 3.58-3.42 (1H, m), 2.43 (3H, d, J = 2.4 Hz), 1.30 (3H, d, J = 6.6 Hz), 1.20-1.06 (1H, m), 0.70-0.58 (1H, m), 0.56-0.38 (2H, m), 0.34-0.22 (1H, m)
Example 2-221		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	391	0.92	¹ H-NMR (DMSO-d ₆) δ: 11.91 (1H, s), 8.53 (1H, s), 8.31-8.20 (1H, m), 8.00-7.75 (5H, m), 7.47-7.27 (1H, m), 7.05 (1H, d, J = 8.7 Hz), 4.44-4.30 (1H, m), 3.42-3.26 (1H, m), 2.77-2.59 (1H, m), 2.44 (3H, d, J = 2.1 Hz), 2.13-1.65 (6H, m), 1.20 (3H, d, J = 6.6 Hz)
Example 2-222		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	365	0.8	¹ H-NMR (DMSO-d ₆) δ: 11.92 (1H, s), 8.60 (1H, s), 8.20-8.10 (1H, m), 8.06-7.75 (5H, m), 7.50-7.27 (1H, m), 7.13 (1H, d, J = 8.4 Hz), 4.35-4.21 (1H, m), 3.50-3.36 (1H, m), 2.43 (3H, d, J = 2.4 Hz), 1.80-1.50 (2H, m), 1.25 (3H, d, J = 6.6 Hz), 0.90 (3H, t, J = 7.4 Hz)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-223		6-(((2S,3R)-2-amino-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	379	0.89	¹ H-NMR (DMSO-d ₆) δ: 11.89 (1H, s), 8.53 (1H, s), 8.18-8.06 (1H, m), 8.04-7.78 (5H, m), 7.48-7.26 (1H, m), 7.13 (1H, d, J = 9.3 Hz), 4.28-4.14 (1H, m), 3.50-3.34 (1H, m), 2.41 (3H, d, J = 2.7 Hz), 1.68-1.54 (2H, m), 1.50-1.30 (2H, m), 1.24 (3H, d, J = 6.6 Hz), 0.86 (3H, t, J = 7.2 Hz)
Example 2-224		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	393	0.97	¹ H-NMR (DMSO-d ₆) δ: 11.87 (1H, s), 8.46 (1H, s), 8.16-8.06 (1H, m), 8.00-7.70 (5H, m), 7.48-7.24 (1H, m), 7.14 (1H, d, J = 7.8 Hz), 4.33-4.20 (1H, m), 3.46-3.30 (1H, m), 2.40 (3H, d, J = 2.7 Hz), 1.70-1.54 (2H, m), 1.44-1.30 (1H, m), 1.23 (3H, d, J = 6.6 Hz), 0.91 (3H, d, J = 6.6 Hz), 0.89 (3H, d, J = 6.6 Hz)
Example 2-225		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	379	0.72	
Example 2-226		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	426	0.84	
Example 2-227		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	448	0.9	
Example 2-228		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	419	0.63	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-229		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide	404	0.96	
Example 2-230		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	412	0.9	
Example 2-231		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-fluoro-2-morpholinopyridin-4-yl)amino)nicotinamide	450	0.65	¹ H-NMR (MeOD) δ: 8.37 (1H, dd, J = 6.9, 6.6 Hz), 7.96-7.90 (2H, m), 4.56-4.46 (1H, m), 4.17-4.08 (1H, m), 4.07-3.71 (8H, m), 3.61-3.55 (4H, m), 2.25-1.82 (2H, m)
Example 2-232		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((5-fluoro-2-morpholinopyridin-4-yl)amino)nicotinamide	450	0.57	¹ H-NMR (MeOD) δ: 8.02 (1H, d, J = 5.0 Hz), 7.99-7.90 (2H, m), 4.61-4.41 (1H, m), 4.22-4.10 (1H, m), 4.09-3.79 (6H, m), 3.72-3.49 (6H, m), 2.25-1.93 (2H, m)
Example 2-233		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-fluoro-2-morpholinopyridin-4-yl)amino)nicotinamide	448	0.74	¹ H-NMR (MeOD) δ: 8.33 (1H, dd, J = 6.6, 6.6 Hz), 7.99-7.93 (1H, m), 7.91 (1H, d, J = 11.9 Hz), 3.90-3.62 (6H, m), 3.60-3.54 (4H, m), 1.46 (3H, d, J = 6.6 Hz), 1.20-1.06 (1H, m), 0.84-0.73 (1H, m), 0.69-0.55 (2H, m), 0.48-0.37 (1H, m)
Example 2-234		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((5-fluoro-2-morpholinopyridin-4-yl)amino)nicotinamide	448	0.67	¹ H-NMR (MeOD) δ: 8.01 (1H, d, J = 4.6 Hz), 7.96-7.89 (2H, m), 4.22-4.16 (1H, m), 3.91-3.84 (4H, m), 3.74-3.64 (1H, m), 3.58-3.50 (4H, m), 1.42 (3H, d, J = 6.6 Hz), 1.20-1.07 (1H, m), 0.75-0.61 (2H, m), 0.57-0.48 (1H, m), 0.39-0.30 (1H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-235		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((3-fluoro-2-morpholinopyridin-4-yl)amino)nicotinamide	464	0.87	¹ H-NMR (MeOD) δ : 8.46 (1H, dd, J = 6.9, 6.6 Hz), 7.97-7.91 (2H, m), 4.63-4.50 (1H, m), 3.90-3.83 (4H, m), 3.62-3.56 (5H, m), 1.81-1.62 (2H, m), 1.56-1.44 (1H, m), 1.38 (3H, d, J = 6.6 Hz), 1.01 (3H, d, J = 6.6 Hz), 0.94 (3H, d, J = 6.6 Hz)
Example 2-236		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((5-fluoro-2-morpholinopyridin-4-yl)amino)nicotinamide	464	0.8	¹ H-NMR (MeOD) δ : 8.03 (1H, d, J = 5.3 Hz), 7.99-7.92 (2H, m), 4.55-4.46 (1H, m), 3.91-3.84 (4H, m), 3.66-3.47 (5H, m), 1.83-1.69 (2H, m), 1.38-1.51 (1H, m), 1.34 (3H, d, J = 6.6 Hz), 1.01 (3H, d, J = 6.6 Hz), 0.94 (3H, d, J = 5.9 Hz)
Example 2-237		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	464	1.04	
Example 2-238		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((6-(difluoromethoxy)-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	478	1.03	¹ H-NMR (DMSO-d ₆) δ : 11.87 (1H, s), 8.63-8.78 (1H, m), 8.36-8.32 (1H, m), 8.23 (2H, s), 8.02-7.92 (1H, m), 7.95-7.20 (6H, m), 6.94 (1H, d, J = 7.2 Hz), 4.24-4.10 (1H, m), 3.50-3.40 (1H, m), 1.85-1.65 (2H, m), 1.65-1.45 (4H, m), 1.40-1.15 (2H, m)
Example 2-239		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	413	0.84	
Example 2-240		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	415	0.93	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-241		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	401	0.82	
Example 2-242		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	415	0.89	
Example 2-243		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	413	0.83	¹ H-NMR (MeOD) δ: 8.73 (1H, d, J = 2.6 Hz), 8.09 (1H, d, J = 2.6 Hz), 7.76 (1H, d, J = 12.6 Hz), 4.83-4.71 (1H, m), 4.03 (3H, s), 3.57-3.50 (1H, m), 2.64 (3H, s), 1.26 (3H, d, J = 6.6 Hz), 1.05-1.00 (1H, m), 0.73-0.66 (1H, m), 0.61-0.52 (1H, m), 0.36-0.28 (2H, m).
Example 2-244		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	429	0.96	
Example 2-245		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	429	0.94	
Example 2-246		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	417	0.77	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-247		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-(((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	399	0.78	
Example 2-248		6-(((3R,4S)-4-aminohexan-3-yl)amino)-2-(((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	415	0.85	
Example 2-249		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-(((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	429	0.89	
Example 2-250		(R)-6-(((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-(((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	431	0.98	
Example 2-251		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-(((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	417	0.87	¹ H-NMR (MeOD) δ : 8.59 (1H, d, J = 2.6 Hz), 8.23 (1H, d, J = 2.6 Hz), 7.77 (1H, d, J = 11.9 Hz), 4.21-4.18 (1H, m), 4.07 (3H, s), 3.94 (3H, s), 3.51-3.45 (1H, m), 1.72-1.56 (2H, m), 1.24 (3H, d, J = 7.3 Hz), 1.02 (3H, t, J = 7.3 Hz).
Example 2-252		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-(((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	431	0.94	¹ H-NMR (MeOD) δ : 8.58 (1H, d, J = 2.6 Hz), 8.24 (1H, d, J = 2.6 Hz), 7.77 (1H, d, J = 11.9 Hz), 4.32-4.26 (1H, m), 4.06 (3H, s), 3.92 (3H, s), 3.49-3.40 (1H, m), 1.62-1.32 (4H, m), 1.24 (3H, d, J = 7.3 Hz), 0.93 (3H, t, J = 7.3 Hz).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-253		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	429	0.88	
Example 2-254		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	445	1	
Example 2-255		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	445	1	
Example 2-256		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	433	0.82	
Example 2-257		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	417	0.83	
Example 2-258		6-(((3R,4S)-4-aminohexan-3-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	431	0.89	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-259		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	436	0.89	
Example 2-260		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	450	0.95	
Example 2-261		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	450	1	
Example 2-262		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	464	1.03	
Example 2-263		6-(((1R,2S)-2-amino-1-cyclohexylpropyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	468	1.06	
Example 2-264		6-(((1R,2S)-2-amino-1-cyclohexylpropyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	454	1.11	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-265		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	442	0.87	¹ H-NMR (MeOD) δ: 7.96 (1H, s), 7.86 (1H, s), 7.74 (1H, d, J = 12.1 Hz), 7.56 (1H, d, J = 8.9 Hz), 7.42-7.32 (1H, m), 4.56 (2H, t, J = 5.1 Hz), 3.82 (2H, t, J = 4.8 Hz), 3.67-3.45 (2H, m), 3.28 (3H, s), 1.28 (3H, d, J = 6.9 Hz), 1.13-0.98 (1H, m), 0.80-0.55 (2H, m), 0.43 (2H, d, J = 4.5 Hz).
Example 2-266		6-(((3R,4S)-4-amino-3-hexyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	444	0.9	
Example 2-267		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide	408	0.92	¹ H-NMR (MeOD) δ: 7.76 (1H, d, J = 11.9 Hz), 6.75 (2H, d, J = 2.3 Hz), 6.19 (1H, t, J = 2.3 Hz), 4.51-4.44 (1H, m), 3.91-3.83 (1H, m), 3.77 (6H, s), 3.68-3.60 (1H, m), 3.43 (3H, s), 3.40-3.34 (1H, m), 1.39 (3H, d, J = 5.9 Hz).
Example 2-268		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide	422	0.99	
Example 2-269		6-(((1R,2S)-2-amino-1-(4-fluorophenyl)propyl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide	458	1.04	
Example 2-270		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	365	0.79	¹ H-NMR (DMSO-d ₆) δ: 11.89 (1H, s), 8.55 (1H, s), 8.20-8.02 (4H, m), 7.99 (1H, d, J = 12.6 Hz), 7.96-7.70 (1H, m), 7.46-7.20 (2H, m), 4.40-4.26 (1H, m), 3.32-3.17 (1H, m), 2.43 (3H, d, J = 2.7 Hz), 1.72-1.57 (2H, m), 1.26 (3H, d, J = 7.2 Hz), 0.89 (3H, t, J = 7.4 Hz).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-271		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	381	0.76	¹ H-NMR (DMSO-d ₆) δ: 11.78 (1H, s), 8.39 (1H, s), 8.10-8.02 (1H, m), 7.97 (1H, d, J = 12.3 Hz), 7.90-7.72 (4H, m), 7.50-7.25 (1H, m), 7.06 (1H, d, J = 8.7 Hz), 4.46-4.34 (1H, m), 3.60-3.40 (3H, m), 3.30 (3H, s), 2.39 (3H, d, J = 2.7 Hz), 1.26 (3H, d, J = 6.6 Hz)
Example 2-272		6-(((1R,2S)-2-amino-1-(4-fluorophenyl)propyl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	431	0.92	¹ H-NMR (DMSO-d ₆) δ: 11.81 (1H, s), 8.36 (1H, s), 8.02-7.80 (6H, m), 7.75 (1H, d, J = 7.8 Hz), 7.50-7.60 (2H, m), 7.48-7.26 (1H, m), 7.25-7.14 (1H, m), 6.38-6.28 (2H, m), 3.80-3.60 (1H, m), 2.43 (3H, d, J = 2.4 Hz), 1.26 (3H, d, J = 6.6 Hz)
Example 2-273		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((5-fluoro-6-methylamino)pyridin-3-yl)amino)nicotinamide	392	0.68	¹ H-NMR (DMSO-d ₆) δ: 11.23 (1H, s), 8.06-7.65 (7H, m), 7.45-7.19 (1H, m), 6.86 (1H, d, J = 6.6 Hz), 4.26-4.11 (3H, m), 3.40-3.22 (1H, m), 2.92 (3H, s), 1.90-1.75 (2H, m), 1.75-1.52 (4H, m), 1.50-1.33 (2H, m)
Example 2-274		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((5-fluoro-6-methylamino)pyridin-3-yl)amino)nicotinamide	392	0.67	
Example 2-275		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((5-fluoro-6-methylamino)pyridin-3-yl)amino)nicotinamide	406	0.77	¹ H-NMR (DMSO-d ₆) δ: 11.46 (1H, s), 8.50-8.22 (1H, m), 8.10-7.50 (6H, m), 7.45-7.12 (1H, m), 7.02-6.87 (1H, m), 4.46-4.32 (1H, m), 3.40-3.22 (1H, m), 2.97 (3H, s), 2.75-2.55 (1H, m), 2.07-1.60 (6H, m), 1.19 (3H, d, J = 6.6 Hz)
Example 2-276		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-methylamino)pyridin-3-yl)amino)nicotinamide	380	0.65	¹ H-NMR (DMSO-d ₆) δ: 11.40 (1H, s), 8.40-8.16 (1H, m), 8.10-7.65 (6H, m), 7.40-7.12 (1H, m), 7.07-6.92 (1H, m), 4.28-4.07 (1H, m), 3.40-3.25 (1H, m), 2.93 (3H, s), 1.80-1.62 (1H, m), 1.63-1.44 (1H, m), 1.21 (3H, d, J = 6.6 Hz), 0.86 (3H, t, J = 6.9 Hz)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-277		6-(((2S,3R)-2-amino-5-fluoro-2-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)nicotinamide	394	0.74	¹ H-NMR (DMSO-d ₆) δ: 11.44 (1H, s), 3.48-8.24 (1H, m), 8.15-7.60 (6H, m), 7.40-7.16 (1H, m), 7.10-6.94 (1H, m), 4.36-4.19 (1H, m), 3.40-3.25 (1H, m), 2.95 (3H, s), 1.70-1.48 (2H, m), 1.45-1.23 (2H, m), 1.21 (3H, d, J = 6.6 Hz), 0.83 (3H, t, J = 7.2 Hz)
Example 2-278		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-(methylamino)pyridin-3-yl)amino)nicotinamide	408	0.82	¹ H-NMR (DMSO-d ₆) δ: 11.21 (1H, s), 8.12-7.60 (6H, m), 6.98-6.88 (1H, m), 4.28-4.14 (1H, m), 3.40-3.25 (1H, m), 2.87 (3H, s), 1.64-1.49 (2H, m), 1.40-1.20 (1H, m), 1.16 (3H, d, J = 7.2 Hz), 0.89 (3H, d, J = 6.6 Hz), 0.76 (3H, d, J = 6.6 Hz)
Example 2-279		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	430	0.82	
Example 2-280		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	446	0.81	
Example 2-281		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-5-fluoro-2-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	460	0.88	
Example 2-282		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	430	0.84	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ HNMR
Example 2-283		6-(((3R,4S)-4-aminohexan-3-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	444	0.89	
Example 2-284		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	446	0.82	
Example 2-285		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	460	0.89	
Example 2-286		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	442	0.87	
Example 2-287		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	444	0.95	
Example 2-288		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	430	0.84	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-289		6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	444	0.91	
Example 2-290		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	442	0.86	
Example 2-291		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	458	0.98	
Example 2-292		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	458	0.97	
Example 2-293		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	456	0.93	
Example 2-294		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	446	0.81	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-295		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	460	0.87	
Example 2-296		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	430	0.81	
Example 2-297		6-(((3R,4S)-4-aminohexan-3-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	444	0.87	
Example 2-298		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	442	0.86	
Example 2-299		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-5-fluoro-2-((6-methoxy-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	442	0.86	
Example 2-300		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	462	0.94	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-301		6-(((1S,2R)-1-amino-1-cyclopropylpropan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	441	0.86	
Example 2-302		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	442	0.93	¹ H-NMR (MeOD) δ: 8.11 (1H, s), 7.86 (1H, d, J = 7.9 Hz), 7.82 (1H, t, J = 6.9 Hz), 7.39-7.34 (1H, m), 7.20 (1H, d, J = 7.9 Hz), 4.55 (2H, t, J = 5.3 Hz), 4.38-4.37 (1H, m), 3.90-3.87 (1H, m), 3.83 (2H, t, J = 5.3 Hz), 3.27 (3H, s), 1.93-1.75 (8H, m).
Example 2-303		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	444	1.01	
Example 2-304		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	430	0.89	
Example 2-305		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	444	0.97	
Example 2-306		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	442	0.9	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-307		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	458	1.05	
Example 2-308		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	458	1.03	
Example 2-309		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	446	0.85	
Example 2-310		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	460	0.92	
Example 2-311		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	430	0.88	
Example 2-312		6-(((1R,2S)-2-amino-1-phenylpropyl)amino)-5-fluoro-2-(quinolin-6-ylamino)nicotinamide	431	0.73	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-313		6-(((1R,2S)-2-amino-1-phenylpropyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	462	0.91	¹ H-NMR (MeOD) δ: 9.20 (1H, d, J = 2.3 Hz), 8.84 (1H, d, J = 2.3 Hz), 8.15 (2H, s), 7.89 (1H, d, J = 11.9 Hz), 7.45-7.26 (5H, m), 5.76 (1H, d, J = 6.6 Hz), 3.99-3.90 (1H, m), 2.87 (3H, s), 1.41 (3H, d, J = 6.6 Hz)
Example 2-314		6-(((1R,2S)-2-amino-1-phenylpropyl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	484	0.96	
Example 2-315		6-(((1R,2S)-2-amino-1-phenylpropyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	477	0.93	
Example 2-316		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	413	0.68	
Example 2-317		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	415	0.58	
Example 2-318		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	401	0.65	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-319		6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	415	0.73	
Example 2-320		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	413	0.66	
Example 2-321		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	429	0.81	
Example 2-322		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	429	0.8	
Example 2-323		6-(((2S,3S)-3-amino-1-methoxybutan-2-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	417	0.61	
Example 2-324		6-(((2S,3S)-3-amino-1-ethoxybutan-2-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	431	0.67	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-325		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	401	0.62	
Example 2-326		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	414	0.82	¹ H-NMR (MeOD) δ: 7.99 (1H, s), 7.92 (1H, d, J = 1.3 Hz), 7.75 (1H, d, J = 11.9 Hz), 7.56 (1H, d, J = 9.2 Hz), 7.44 (1H, dd, J = 8.9, 1.7 Hz), 4.47 (2H, q, J = 7.3 Hz), 4.28 (1H, td, J = 8.4, 4.6 Hz), 4.06 (1H, dd, J = 11.9, 4.6 Hz), 3.80-3.51 (3H, m), 3.46-3.39 (1H, m), 2.11-1.94 (1H, m), 1.89-1.80 (1H, m), 1.47 (3H, t, J = 7.3 Hz).
Example 2-327		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-methyl-1H-indazol-5-yl)amino)nicotinamide	458	0.81	
Example 2-328		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-(2,2-difluoroethyl)-3-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	464	0.86	¹ H-NMR (MeOD) δ: 7.74 (1H, d, J = 11.9 Hz), 7.63 (1H, dd, J = 8.9, 1.7 Hz), 7.60 (1H, s), 7.53 (1H, d, J = 8.6 Hz), 6.22 (1H, tt, J = 55.5, 3.6 Hz), 4.75 (2H, td, J = 14.7, 3.5 Hz), 4.23-4.14 (1H, m), 4.06-4.00 (1H, m), 3.69-3.59 (3H, m), 3.58-3.48 (1H, m), 2.54 (3H, s), 2.04-1.94 (1H, m), 1.86-1.78 (1H, m).
Example 2-329		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	474	0.86	
Example 2-330		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	414	0.82	¹ H-NMR (MeOD) δ: 8.08 (1H, s), 7.83 (1H, d, J = 12.6 Hz), 7.80 (1H, d, J = 7.9 Hz), 7.41-7.36 (1H, m), 7.20 (1H, d, J = 7.9 Hz), 4.51-4.42 (2H, m), 4.45-4.35 (1H, m), 4.12-4.08 (1H, m), 3.93-3.89 (2H, m), 3.69-3.53 (2H, m), 2.09-1.87 (2H, m), 1.48 (3H, t, J = 6.9 Hz).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-331		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	444	0.82	¹ H-NMR (MeOD) δ : 8.08 (1H, s), 7.82 (1H, d, J = 11.9 Hz), 7.71 (1H, d, J = 7.3 Hz), 7.37 (1H, t, J = 7.9 Hz), 7.23 (1H, d, J = 8.6 Hz), 4.55 (2H, t, J = 6.3 Hz), 4.39-4.32 (1H, m), 4.12-4.03 (1H, m), 3.90-3.83 (2H, m), 3.88-3.80 (2H, m), 3.64-3.55 (2H, m), 3.36 (3H, s), 2.07-1.99 (2H, m).
Example 2-332		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide	450	0.86	¹ H-NMR (MeOD) δ : 8.15 (1H, s), 7.83 (1H, d, J = 12.6 Hz), 7.80 (1H, d, J = 7.9 Hz), 7.42 (1H, t, J = 7.9 Hz), 7.24 (1H, d, J = 8.6 Hz), 6.48-6.05 (1H, m), 4.84-4.78 (2H, m), 4.42-4.34 (1H, m), 4.14-4.02 (1H, m), 3.92-3.85 (2H, m), 3.70-3.51 (2H, m), 1.93-1.84 (2H, m).
Example 2-333		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	415	0.73	
Example 2-334		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-methoxy-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)nicotinamide	431	0.77	
Example 2-335		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-pyrazolo[3,4-b]pyridin-5-yl)amino)-5-fluoronicotinamide	444	0.76	
Example 2-336		6-(((1R,2S)-2-amino-1-phenylpropyl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	478	0.93	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-337		6-(((1R,2S)-2-amino-1-(4-methoxyphenyl)propyl)amino)-5-fluoro-2-(quinolin-6-ylamino)nicotinamide	461	0.73	
Example 2-338		6-(((1R,2S)-2-amino-1-(4-methoxyphenyl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	492	0.85	¹ H-NMR (MeOD) δ: 9.09 (1H, d, J = 2.0 Hz), 8.85 (1H, d, J = 2.0 Hz), 8.15 (2H, s), 7.88 (1H, d, J = 11.9 Hz), 7.30 (2H, d, J = 8.6 Hz), 6.88 (2H, d, J = 8.6 Hz), 5.55 (1H, d, J = 6.6 Hz), 3.91-3.85 (1H, m), 3.75 (3H, s), 2.82 (3H, s), 1.39 (3H, d, J = 7.3 Hz)
Example 2-339		6-(((1R,2S)-2-amino-1-(4-methoxyphenyl)propyl)amino)-5-fluoro-2-((6-methoxy-5-(1H-1,2,3-triazol-1-yl)pyridin-3-yl)amino)nicotinamide	508	0.91	
Example 2-340		6-(((1R,2S)-2-amino-1-(4-methoxyphenyl)propyl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	514	0.96	
Example 2-341		6-(((1R,2S)-2-amino-1-(4-methoxyphenyl)propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	478	0.97	¹ H-NMR (MeOD) δ: 7.97 (1H, s), 7.87 (1H, d, J = 1.3 Hz), 7.75 (1H, d, J = 11.9 Hz), 7.50 (1H, d, J = 8.6 Hz), 7.36-7.27 (3H, m), 6.96 (2H, d, J = 8.6 Hz), 5.38 (1H, d, J = 5.7 Hz), 4.48 (2H, q, J = 7.2 Hz), 3.81-3.72 (4H, m), 1.49 (3H, t, J = 7.2 Hz), 1.30 (3H, d, J = 6.6 Hz)
Example 2-342		6-(((1R,2S)-2-amino-1-(4-methoxyphenyl)propyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	507	0.94	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-343		6-(((3R,4S)-4-amino-1-(methylthio)pentan-3-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	460	0.87	
Example 2-344		6-(((3R,4S)-4-amino-1-(methylthio)pentan-3-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	482	0.94	
Example 2-345		6-(((3R,4S)-4-amino-1-(methylthio)pentan-3-yl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide	453	0.72	
Example 2-346		6-(((3R,4S)-4-amino-1-(methylthio)pentan-3-yl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	438	1	
Example 2-347		2-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-6-(((3S,4R)-4-aminobicyclo[4.1.0]heptan-3-yl)amino)-5-fluoronicotinamide	423	0.87	
Example 2-348		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	466	0.91	¹ H-NMR (MeOD) δ: 8.14 (1H, d, J = 1.3 Hz), 7.77 (1H, d, J = 11.9 Hz), 7.45 (1H, d, J = 8.6 Hz), 7.31 (1H, dd, J = 8.6, 2.6 Hz), 7.31 (1H, d, J = 147.0 Hz), 4.44 (1H, td, J = 8.3, 4.0 Hz), 4.09 (1H, dd, J = 11.6, 4.3 Hz), 3.93 (3H, s), 3.86-3.74 (2H, m), 3.77-3.65 (2H, m), 2.04-2.00 (2H, m).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-349		6-(((2S,3R)-2-amino-3-yl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	466	1.08	
Example 2-350		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	478	1.09	
Example 2-351		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	452	0.98	
Example 2-352		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	510	0.94	¹ H-NMR (MeOD) δ: 8.08 (1H, s), 7.77 (1H, d, J = 11.9 Hz), 7.49 (1H, d, J = 8.6 Hz), 7.34-7.30 (1H, m), 7.32 (1H, d, J = 147.0 Hz), 4.43 (2H, t, J = 5.3 Hz), 4.40-4.37 (1H, m), 4.08 (1H, dd, J = 11.2, 4.6 Hz), 3.79 (2H, t, J = 6.3 Hz), 3.77-3.72 (2H, m), 3.72-3.65 (2H, m), 3.35 (3H, s), 2.09-1.98 (2H, m).
Example 2-353		6-(((2S,3R)-2-amino-3-yl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	510	1.1	
Example 2-354		6-(((1R,2S)-2-amino-1-(4-(trifluoromethyl)phenyl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	530	1.02	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-355		6-(((1R,2S)-2-amino-1-(4-(trifluoromethyl)phenyl)propyl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide	509	1.15	
Example 2-356		6-(((1R,2S)-2-amino-1-phenylpropyl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide	441	0.99	
Example 2-357		6-(((1R,2S)-2-amino-1-(4-methoxyphenyl)propyl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide	471	1	
Example 2-358		6-(((1R,2S)-2-amino-1-(cyclopropylbutyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	440	0.89	¹ H-NMR (DMSO-d ₆) δ: 11.87 (1H, s), 8.65 (1H, s), 8.41 (1H, s), 8.21 (2H, s), 7.97 (5H, m), 7.45-7.32 (1H, m), 7.26 (1H, d, J = 7.9 Hz), 3.75-3.65 (1H, m), 3.40-3.36 (1H, m), 2.60 (3H, m), 1.65-1.45 (2H, m), 1.11-0.90 (1H, m), 0.75 (3H, t, J = 7.3 Hz), 0.60-0.16 (4H, m)
Example 2-359		6-(((1R,2S)-2-amino-1-(cyclopropylbutyl)amino)-5-fluoro-2-((6-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	462	0.95	
Example 2-360		6-(((1R,2S)-2-amino-1-(cyclopropylbutyl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	433	0.72	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-361		6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-2-((3,5-dimethoxyphenyl)amino)-5-fluoronicotinamide	418	1.03	
Example 2-362		6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	426	0.95	
Example 2-363		(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	448	1.02	
Example 2-364		(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	462	0.96	
Example 2-365		(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	484	1.01	¹ H-NMR (MeOD) δ: 8.66 (1H, s), 7.90 (1H, dd, J = 14.4, 2.1 Hz), 7.72 (1H, d, J = 11.9 Hz), 7.21-7.10 (5H, m), 4.87 (2H, 溶媒ピークと重なっている), 3.92-3.86 (4H, m), 3.63-3.53 (4H, m), 3.19-2.82 (3H, m).
Example 2-366		2-((5-(1H-pyrazol-1-yl)pyridin-3-yl)amino)-6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-5-fluoronicotinamide	425	0.87	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-367		2-((5-((1H-pyrazol-1-yl)pyridin-3-yl)amino)-6-(((3R,4S)-4-aminobicyclo[4.1.0]heptan-3-yl)amino)-5-fluoronicotinamide	423	0.88	
Example 2-368		6-(((3R,4S)-4-aminobicyclo[4.1.0]heptan-3-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	438	0.9	
Example 2-369		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((1-ethyl-6-fluoro-1H-indazol-4-yl)amino)-5-fluoronicotinamide	430	1.04	
Example 2-370		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-ethyl-6-fluoro-1H-indazol-4-yl)amino)-5-fluoronicotinamide	432	0.93	¹ H-NMR (MeOD) δ: 8.10 (1H, s), 7.92 (1H, dd, J = 2.0, 12.9 Hz), 7.87 (1H, d, J = 11.9 Hz), 6.89 (1H, d, J = 7.9 Hz), 4.60-4.50 (1H, m), 4.38 (2H, q, J = 7.3 Hz), 4.20-4.18 (1H, m), 4.08-4.00 (2H, m), 3.97-3.66 (2H, m), 2.66-1.90 (2H, m), 1.45 (3H, t, J = 7.3 Hz).
Example 2-371		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((1-ethyl-6-fluoro-1H-indazol-4-yl)amino)-5-fluoronicotinamide	418	1.01	
Example 2-372		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-ethyl-6-fluoro-1H-indazol-4-yl)amino)-5-fluoronicotinamide	430	1.01	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-373		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((1-ethyl-6-fluoro-1H-indazol-4-yl)amino)-5-fluoronicotinamide	418	0.97	
Example 2-374		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	460	0.98	¹ H-NMR (MeOD) δ: 8.11 (1H, s), 7.94 (1H, dd, J = 12.6, 2.0 Hz), 7.86 (1H, d, J = 11.9 Hz), 6.88 (1H, d, J = 9.2 Hz), 4.49 (2H, t, J = 5.3 Hz), 4.48-4.42 (1H, m), 4.06-4.00 (1H, m), 3.80 (2H, t, J = 5.3 Hz), 3.27 (3H, s), 2.01-1.60 (8H, m).
Example 2-375		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	462	0.86	¹ H-NMR (MeOD) δ: 8.11 (1H, s), 7.92-7.84 (1H, m), 7.87 (1H, d, J = 13.2 Hz), 6.90 (1H, d, J = 8.6 Hz), 4.67-4.49 (1H, m), 4.49 (2H, t, J = 5.0 Hz), 4.19-4.11 (1H, m), 4.07-3.99 (2H, m), 3.87-3.62 (2H, m), 3.80 (2H, t, J = 5.0 Hz), 3.27 (3H, s), 2.02-1.87 (2H, m).
Example 2-376		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	448	0.95	¹ H-NMR (MeOD) δ: 8.11 (1H, s), 7.91 (1H, dd, J = 12.6, 1.9 Hz), 7.85 (1H, d, J = 11.9 Hz), 6.88 (1H, d, J = 8.6 Hz), 4.49 (2H, t, J = 5.0 Hz), 4.21-4.18 (1H, m), 3.80 (2H, t, J = 5.0 Hz), 3.61-3.45 (1H, m), 3.27 (3H, s), 2.20 (3H, d, J = 7.3 Hz), 1.64-1.54 (2H, m), 1.10 (3H, t, J = 7.6 Hz).
Example 2-377		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	460	0.95	¹ H-NMR (MeOD) δ: 8.10 (1H, s), 7.84 (1H, d, J = 12.0 Hz), 7.78 (1H, dd, J = 13.9, 2.0 Hz), 6.88 (1H, d, J = 10.0 Hz), 4.48 (2H, t, J = 5.3 Hz), 4.14-4.05 (1H, m), 3.80 (2H, t, J = 5.3 Hz), 3.79-3.74 (1H, m), 3.27 (3H, s), 1.45 (3H, d, J = 5.0 Hz), 1.17-1.11 (1H, m), 0.87-0.78 (1H, m), 0.67-0.60 (1H, m), 0.60-0.45 (2H, m).
Example 2-378		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((6-fluoro-1-(2-methoxyethyl)-1H-indazol-4-yl)amino)nicotinamide	448	0.93	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-379		6-(((1R,2S)-2-amino-1-(4-(trifluoromethyl)phenyl)propyl)amino)-5-fluoro-2-(quinolin-6-ylamino)nicotinamide	499	0.87	
Example 2-380		6-(((1R,2S)-2-amino-1-((S)-2,2-dimethylcyclopropyl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	454	1	
Example 2-381		6-(((1S,2S)-2-amino-1-(pyridin-2-yl)propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	449	0.88	
Example 2-382		6-(((1R,2S)-2-amino-1-(cyclopropylbutyl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	391	0.88	
Example 2-383		6-(((1R,2S)-2-amino-1-(cyclopropylbutyl)amino)-2-((3-(dimethylamino)-7-fluoro-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	473	1.04	
Example 2-384		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	443	0.71	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	1HMR
Example 2-385		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	445	0.59	
Example 2-386		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	445	0.76	
Example 2-387		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	431	0.65	
Example 2-388		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	443	0.68	
Example 2-389		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	459	0.78	
Example 2-390		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	459	0.76	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ HMR
Example 2-391		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	457	0.74	
Example 2-392		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	431	0.63	
Example 2-393		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	449	0.72	¹ H-NMR (MeOD) δ: 9.25 (1H, s), 9.00 (1H, m), 8.54 (1H, s), 7.95 (1H, d, J = 11.9 Hz), 6.38 (1H, t, J = 54.8, 2.9 Hz), 4.78-4.68 (2H, m), 4.56-4.51 (1H, m), 3.86-3.81 (1H, m), 1.96-1.55 (8H, m).
Example 2-394		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	451	0.62	
Example 2-395		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	451	0.82	
Example 2-396		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	437	0.7	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-397		6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	451	0.78	
Example 2-398		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	449	0.72	
Example 2-399		6-(((2S,3R)-2-aminoheptan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	465	0.83	
Example 2-400		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	465	0.82	
Example 2-401		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	463	0.79	
Example 2-402		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((1-(2,2-difluoroethyl)-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	437	0.66	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-403		6-(((1S,2S)-2-amino-1-(pyridin-2-yl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	463	0.85	
Example 2-404		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-(quinolin-5-ylamino)nicotinamide	395	0.68	
Example 2-405		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	415	0.78	
Example 2-406		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((1-ethyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)-5-fluoronicotinamide	427	0.75	
Example 2-407		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((7-methoxy-1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	429	0.91	
Example 2-408		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((7-methoxy-1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)amino)nicotinamide	429	0.89	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-409		6-(((1R,2S)-2-amino-1-(2-fluorophenyl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	480	0.92	
Example 2-410		6-(((1R,2S)-2-amino-1-(2-fluorophenyl)propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	466	0.98	
Example 2-411		6-(((1R,2S)-2-amino-1-(2-fluorophenyl)propyl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide	459	1.04	
Example 2-412		6-(((1R,2S)-2-amino-1-(3-fluorophenyl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	479	0.9	
Example 2-413		6-(((1R,2S)-2-amino-1-(3-fluorophenyl)propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	466	0.96	
Example 2-414		6-(((1R,2S)-2-amino-1-(3-fluorophenyl)propyl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide	459	1	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-415		6-(((1S,2S)-2-amino-1-(thiophen-2-yl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	468	0.88	
Example 2-416		6-(((1S,2S)-2-amino-1-(thiophen-2-yl)propyl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	454	0.94	
Example 2-417		(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-5-fluoro-2-((2-methoxypyridin-4-yl)amino)nicotinamide	411	0.74	
Example 2-418		(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-2-((2,6-dimethoxypyridin-4-yl)amino)-5-fluoronicotinamide	441	1.06	
Example 2-419		(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-5-fluoro-2-((2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	455	0.78	
Example 2-420		(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-5-fluoro-2-((6-morpholinopyridin-3-yl)amino)nicotinamide	466	0.79	¹ H-NMR (MeOD) δ : 8.81 (1H, s), 7.90 (1H, dd, J = 9.6, 2.3 Hz), 7.70 (1H, d, J = 11.9 Hz), 7.20 (1H, d, J = 9.9 Hz), 7.15-7.09 (5H, m), 4.87 (2H, 溶媒ピークと重なっている), 3.95-3.84 (4H, m), 3.71-3.63 (4H, m), 3.18-2.94 (2H, m), 2.87-2.74 (1H, m).

Example	Structure	Compound name	MS(ESI m/z): ($M+H$)	RT(min)	HNMR
Example 2-421		(R)-6-((1-amino-3-phenylpropan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	477	0.99	
Example 2-422		6-(((2R,3S)-3-amino-1-phenylbutan-2-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	476	0.96	
Example 2-423		6-(((2R,3S)-3-amino-1-phenylbutan-2-yl)amino)-2-((1-ethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	462	1.01	
Example 2-424		6-(((2R,3S)-3-amino-1-phenylbutan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	491	1	
Example 2-425		6-(((2R,3S)-3-amino-1-phenylbutan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	498	1.01	
Example 2-426		6-(((2R,3S)-3-amino-1-phenylbutan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	427	0.95	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-427		6-(((2R,3S)-3-amino-1-phenylbutan-2-yl)amino)-5-fluoro-2-((2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	469	0.79	
Example 2-428		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((1-methyl-3-(pyrrolidin-1-yl)-1H-indazol-5-yl)amino)nicotinamide	467	0.92	
Example 2-429		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((1-methyl-3-(pyrrolidin-1-yl)-1H-indazol-5-yl)amino)nicotinamide	455	0.89	
Example 2-430		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-methyl-3-(pyrrolidin-1-yl)-1H-indazol-5-yl)amino)nicotinamide	467	0.91	
Example 2-431		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((1-methyl-3-(2-oxopyrrolidin-1-yl)-1H-indazol-5-yl)amino)nicotinamide	481	0.87	
Example 2-432		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-methyl-3-(2-oxopyrrolidin-1-yl)-1H-indazol-5-yl)amino)nicotinamide	481	0.93	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-433		6-(((2R,3S)-3-amino-2-methyl-3-oxopent-2-yl)amino)-5-fluoro-2-((1-methyl-3-(2-oxopyrrolidin-1-yl)-1H-indazol-5-yl)amino)nicotinamide	469	0.86	
Example 2-434		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-(m-tolylamino)nicotinamide	358	1.05	¹ H-NMR (MeOD) δ: 7.74 (1H, d, J = 11.9 Hz), 7.42-7.36 (1H, m), 7.29 (1H, s), 7.18 (1H, dd, J = 7.9, 7.9 Hz), 6.87-6.81 (1H, m), 4.42-4.32 (1H, m), 3.86-3.78 (1H, m), 2.34 (3H, s), 1.94-1.52 (8H, m)
Example 2-435		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-(p-tolylamino)nicotinamide	358	1.04	¹ H-NMR (MeOD) δ: 7.72 (1H, d, J = 11.9 Hz), 7.40 (2H, d, J = 8.6 Hz), 7.12 (2H, d, J = 8.6 Hz), 4.34-4.25 (1H, m), 3.88-3.80 (1H, m), 2.30 (3H, s), 1.92-1.63 (8H, m)
Example 2-436		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3,4-dimethylphenyl)amino)-5-fluoronicotinamide	372	1.1	¹ H-NMR (MeOD) δ: 7.72 (1H, d, J = 11.9 Hz), 7.32-7.27 (1H, m), 7.24-7.20 (1H, m), 7.09-7.03 (1H, m), 4.37-4.29 (1H, m), 3.83-3.75 (1H, m), 2.26 (3H, s), 2.22 (3H, s), 1.89-1.63 (8H, m)
Example 2-437		2-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoronicotinamide	411	1.03	
Example 2-438		2-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoronicotinamide	411	0.89	¹ H-NMR (MeOD) δ: 8.89-8.85 (1H, m), 8.68 (1H, d, J = 1.0 Hz), 7.94 (1H, d, J = 1.3 Hz), 7.80 (1H, d, J = 11.9 Hz), 7.51-7.44 (1H, m), 7.36-7.31 (1H, m), 7.26-7.21 (1H, m), 4.75-4.67 (1H, m), 3.86-3.78 (1H, m), 1.88-1.45 (8H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-439		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-chlorophenyl)amino)-5-fluoronicotinamide	378 380	1.1	¹ H-NMR (MeOD) δ: 7.99-7.96 (1H, m), 7.77 (1H, d, J = 11.9 Hz), 7.29-7.17 (2H, m), 6.99-6.93 (1H, m), 4.46-4.36 (1H, m), 3.89-3.83 (1H, m), 2.03-1.54 (8H, m)
Example 2-440		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-fluorophenyl)amino)nicotinamide	362	1.07	¹ H-NMR (MeOD) δ: 7.80-7.70 (2H, m), 7.30-7.21 (1H, m), 7.10-7.04 (1H, m), 6.73-6.65 (1H, m), 4.42-4.33 (1H, m), 3.98-3.89 (1H, m), 1.97-1.61 (8H, m)
Example 2-441		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((4-fluorophenyl)amino)nicotinamide	362	1.02	¹ H-NMR (MeOD) δ: 7.74 (1H, d, J = 11.9 Hz), 7.54-7.47 (2H, m), 7.10-7.01 (2H, m), 4.33-4.24 (1H, m), 3.87-3.79 (1H, m), 1.92-1.52 (8H, m)
Example 2-442		2-((3-acetylphenyl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoronicotinamide	386	0.93	¹ H-NMR (MeOD) δ: 8.58-8.53 (1H, m), 7.78 (1H, d, J = 11.9 Hz), 7.68-7.62 (1H, m), 7.54-7.48 (1H, m), 7.47-7.39 (1H, m), 4.66-4.58 (1H, m), 3.85-3.76 (1H, m), 2.62 (3H, s), 1.94-1.52 (8H, m)
Example 2-443		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3,5-bis(trifluoromethyl)phenyl)amino)-5-fluoronicotinamide	480	1.33	¹ H-NMR (MeOD) δ: 8.19 (2H, s), 7.83 (1H, d, J = 11.9 Hz), 7.47 (1H, s), 4.60-4.51 (1H, m), 3.70-3.60 (1H, m), 1.96-1.50 (8H, m)
Example 2-444		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-(trifluoromethoxy)phenyl)amino)nicotinamide	428	1.19	¹ H-NMR (MeOD) δ: 7.85-7.75 (2H, m), 7.40-7.29 (2H, m), 6.90-6.84 (1H, m), 4.47-4.38 (1H, m), 3.88-3.80 (1H, m), 1.98-1.51 (8H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-445		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-chloro-4-methylphenyl)amino)-5-fluoronicotinamide	392 394	1.17	¹ H-NMR (MeOD) δ: 7.94-7.91 (1H, m), 7.75 (1H, d, J = 11.9 Hz), 7.21-7.09 (2H, m), 4.44-4.34 (1H, m), 3.88-3.81 (1H, m), 2.31 (3H, s), 1.99-1.55 (8H, m)
Example 2-446		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-isopropoxyphenyl)amino)nicotinamide	402	1.12	¹ H-NMR (MeOD) δ: 7.74 (1H, d, J = 11.9 Hz), 7.43-7.40 (1H, m), 7.22-7.12 (1H, m), 6.93-6.87 (1H, m), 6.59-6.53 (1H, m), 4.65-4.55 (1H, m), 4.40-4.32 (1H, m), 3.97-3.91 (1H, m), 2.12-1.53 (6H, m), 1.36-1.29 (6H, m)
Example 2-447		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3,4-difluorophenyl)amino)-5-fluoronicotinamide	380	1.06	¹ H-NMR (MeOD) δ: 7.90-7.78 (1H, m), 7.76 (1H, d, J = 11.9 Hz), 7.20-7.02 (1H, m), 7.10-7.03 (1H, m), 4.40-4.31 (1H, m), 3.87-3.82 (1H, m), 2.00-1.55 (8H, m)
Example 2-448		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((4-isopropylphenyl)amino)nicotinamide	386	1.19	¹ H-NMR (MeOD) δ: 7.72 (1H, d, J = 11.9 Hz), 7.46-7.40 (2H, m), 7.21-7.13 (2H, m), 4.33-4.25 (1H, m), 3.92-3.85 (1H, m), 2.92-2.80 (1H, m), 1.90-1.52 (8H, m), 1.24 (6H, d, J = 6.9 Hz)
Example 2-449		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((4-isopropoxyphenyl)amino)nicotinamide	402	1.1	¹ H-NMR (MeOD) δ: 7.71 (1H, d, J = 11.9 Hz), 7.41-7.34 (2H, m), 6.92-6.83 (2H, m), 4.62-4.47 (1H, m), 4.27-4.18 (1H, m), 3.82-3.77 (1H, m), 1.90-1.51 (8H, m), 1.30 (6H, d, J = 5.9 Hz)
Example 2-450		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-ethylphenyl)amino)-5-fluoronicotinamide	372	1.12	¹ H-NMR (MeOD) δ: 7.73 (1H, d, J = 11.9 Hz), 7.46 (1H, d, J = 7.9 Hz), 7.27-7.18 (2H, m), 6.86 (1H, d, J = 7.9 Hz), 4.43-4.34 (1H, m), 3.82-3.77 (1H, m), 2.63 (2H, q, J = 7.6 Hz), 1.92-1.60 (8H, m), 1.24 (3H, t, J = 7.4 Hz)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-451		6-(((1R,2S)-2-amino-1-(3-fluorophenyl)propyl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	502	0.96	
Example 2-452		6-(((1R,2S)-2-amino-1-(3-fluorophenyl)propyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	495	0.93	
Example 2-453		6-(((1R,2S)-2-amino-1-(3-fluorophenyl)propyl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	431	0.87	
Example 2-454		6-(((1R,2S)-2-amino-1-(3-fluorophenyl)propyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	496	0.93	
Example 2-455		6-(((1R,2S)-2-amino-1-(4-methoxyphenyl)propyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	508	0.92	
Example 2-456		6-(((1R,2S)-2-amino-1-(aminocyclohexyl)amino)-5-fluoro-2-((3-((2-methoxyethyl)(methyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	485	0.94	¹ H-NMR (MeOD) δ: 8.12 (1H, d, J = 9.9 Hz), 7.78 (1H, d, J = 7.5 Hz), 7.76 (1H, d, J = 3.5 Hz), 7.58 (1H, d, J = 9.9 Hz), 4.32-4.23 (1H, m), 3.92-3.85 (1H, m), 3.90 (3H, s), 3.77-3.53 (4H, m), 2.39 (3H, s), 3.34 (3H, s), 1.87-1.45 (8H, m).

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-457		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((3-((2-methoxyethyl)(methyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	487	0.83	
Example 2-458		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((3-((2-methoxyethyl)(methyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	473	0.89	
Example 2-459		6-(((2S,3R)-2-aminohexan-3-yl)amino)-5-fluoro-2-((3-((2-methoxyethyl)(methyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	487	0.95	
Example 2-460		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-((2-methoxyethyl)(methyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	485	0.9	
Example 2-461		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((3-((2-methoxyethyl)(methyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	473	0.86	
Example 2-462		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-((2-methoxyethyl)(methyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	471	0.87	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ HNMR
Example 2-463		6-(((2S,3R)-2-amino-3-ethylpentan-3-yl)amino)-5-fluoro-2-((3-((2-methoxyethyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	459	0.85	
Example 2-464		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-((2-methoxyethyl)amino)-1-methyl-1H-indazol-5-yl)amino)nicotinamide	471	0.85	
Example 2-465		(S)-6-((1-amino-3-(1H-pyrazol-1-yl)propan-2-yl)amino)-5-fluoro-2-((2-(2-methoxyethoxy)pyridin-4-yl)amino)nicotinamide	445	0.61	
Example 2-466		(S)-6-((1-amino-3-(1H-pyrazol-1-yl)propan-2-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	452	0.76	
Example 2-467		(S)-6-((1-amino-3-(1H-pyrazol-1-yl)propan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	474	0.82	
Example 2-468		(S)-6-((1-amino-3-(1H-pyrazol-1-yl)propan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	403	0.71	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-469		(S)-6-((1-amino-3-(1H-pyrazol-1-yl)propan-2-yl)amino)-5-fluoro-2-((5-fluoro-6-methylpyridin-3-yl)amino)nicotinamide	438	0.85	
Example 2-470		(S)-6-((1-amino-3-(1H-pyrazol-1-yl)propan-2-yl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	467	0.82	
Example 2-471		(S)-6-((1-amino-3-(1H-pyrazol-1-yl)propan-2-yl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	438	0.88	
Example 2-472		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide	425	0.91	¹ H-NMR (MeOD) δ : 9.35-9.31 (1H, m), 8.82-8.67 (2H, m), 8.27 (1H, s), 7.89 (1H, d, J = 11.9 Hz), 7.74 (1H, s), 4.74-4.64 (1H, m), 3.80-3.71 (1H, m), 2.20 (3H, s), 1.96-1.47 (6H, m)
Example 2-473		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((5-(3,5-dimethyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	439	0.91	¹ H-NMR (MeOD) δ : 8.87-8.78 (2H, m), 8.40 (1H, s), 7.88 (1H, d, J = 11.9 Hz), 6.21 (1H, s), 4.43-4.34 (1H, m), 3.74-3.64 (1H, m), 2.41 (3H, s), 2.30 (3H, s), 1.90-1.36 (6H, m)
Example 2-474		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-(m-tolylamino)nicotinamide	346	0.99	¹ H-NMR (MeOD) δ : 7.74 (1H, d, J = 12.2 Hz), 7.40-7.34 (1H, m), 7.26-7.17 (2H, m), 6.87 (1H, d, J = 7.6 Hz), 4.30-4.19 (1H, m), 3.58-3.47 (1H, m), 2.34 (3H, s), 1.82-1.55 (2H, m), 1.26 (3H, d, J = 6.9 Hz), 1.06 (3H, t, J = 7.3 Hz)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-475		2-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoronicotinamide	399	1	¹ H-NMR (MeOD) δ: 9.04-9.01 (1H, m), 7.98 (2H, s), 7.81 (1H, d, J = 11.9 Hz), 7.71-7.64 (1H, m), 7.47-7.38 (1H, m), 7.16-7.10 (1H, m), 4.83-4.73 (1H, m), 3.70-3.60 (1H, m), 1.87-1.60 (2H, m), 1.27 (3H, d, J = 6.6 Hz), 1.05 (3H, t, J = 7.3 Hz)
Example 2-476		2-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoronicotinamide	399	0.87	¹ H-NMR (MeOD) δ: 9.13-9.09 (1H, m), 8.64-8.59 (1H, m), 7.96-7.94 (1H, m), 7.81 (1H, d, J = 12.2 Hz), 7.52-7.44 (1H, m), 7.36-7.30 (1H, m), 7.24-7.13 (1H, m), 4.83-4.73 (1H, m), 3.82-3.68 (1H, m), 1.88-1.55 (2H, m), 1.31 (3H, d, J = 6.9 Hz), 0.94 (3H, t, J = 7.4 Hz)
Example 2-477		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3,4-dimethylphenyl)amino)-5-fluoronicotinamide	360	1.05	¹ H-NMR (MeOD) δ: 7.73 (1H, d, J = 11.9 Hz), 7.30-7.23 (1H, m), 7.19-7.14 (1H, m), 7.13-7.07 (1H, m), 4.24-4.14 (1H, m), 3.56-3.43 (1H, m), 2.27 (3H, s), 2.24 (3H, s), 1.78-1.55 (2H, m), 1.23 (3H, d, J = 6.6 Hz), 1.05 (3H, t, J = 7.4 Hz)
Example 2-478		2-((3-acetylphenyl)amino)-6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoronicotinamide	374	0.9	¹ H-NMR (MeOD) δ: 8.82-8.76 (1H, m), 7.79 (1H, d, J = 11.9 Hz), 7.69-7.63 (1H, m), 7.49-7.37 (2H, m), 4.68-4.58 (1H, m), 3.72-3.60 (1H, m), 2.65 (3H, s), 1.88-1.56 (2H, m), 1.33 (3H, d, J = 6.6 Hz), 1.00 (3H, t, J = 7.3 Hz)
Example 2-479		2-((3-(2H-1,2,3-triazol-2-yl)phenyl)amino)-6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoronicotinamide	411	1.01	¹ H-NMR (MeOD) δ: 8.79-8.75 (1H, m), 7.98 (2H, s), 7.80 (1H, d, J = 11.9 Hz), 7.70-7.65 (1H, m), 7.42 (1H, dd, J = 8.3 Hz, 8.3 Hz), 7.17-7.11 (1H, m), 3.96 (1H, dd, J = 4.0 Hz, 9.9 Hz), 3.86-3.74 (1H, m), 1.34 (3H, d, J = 6.9 Hz), 1.19-1.07 (1H, m), 0.78-0.57 (2H, m), 0.57-0.37 (2H, m)
Example 2-480		2-((3-(1H-1,2,3-triazol-1-yl)phenyl)amino)-6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoronicotinamide	411	0.89	¹ H-NMR (MeOD) δ: 9.03-9.00 (1H, m), 8.61 (1H, d, J = 1.0 Hz), 7.96 (1H, d, J = 1.0 Hz), 7.80 (1H, d, J = 12.2 Hz), 7.47 (1H, dd, J = 7.9 Hz, 8.3 Hz), 7.35-7.29 (1H, m), 7.19-7.12 (1H, m), 4.09 (1H, dd, J = 5.3 Hz, 9.6 Hz), 3.91-3.79 (1H, m), 1.39 (3H, d, J = 6.9 Hz), 1.20-1.04 (1H, m), 0.76-0.48 (2H, m), 0.48-0.38 (1H, m), 0.36-0.25 (1H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-481		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3,4-dimethylphenyl)amino)-5-fluoronicotinamide	372	1.07	¹ H-NMR (MeOD) δ: 7.71 (1H, d, J = 11.9 Hz), 7.24-7.17 (1H, m), 7.14-7.11 (1H, m), 7.09-7.04 (1H, m), 3.69-3.68 (1H, m), 3.63 (1H, dd, J = 4.5 Hz, 9.7 Hz), 2.25 (3H, s), 2.23 (3H, s), 1.33 (3H, d, J = 6.9 Hz), 1.17-1.02 (1H, m), 0.82-0.67 (2H, m), 0.50-0.37 (2H, m)
Example 2-482		2-((3-acetylphenyl)amino)-6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoronicotinamide	386	0.93	¹ H-NMR (MeOD) δ: 8.75-8.71 (1H, m), 7.77 (1H, d, J = 12.2 Hz), 7.68-7.62 (1H, m), 7.47-7.34 (2H, m), 3.96 (1H, dd, J = 5.1 Hz, 9.7 Hz), 3.83-3.70 (1H, m), 2.64 (3H, s), 1.41 (3H, d, J = 6.9 Hz), 1.22-1.07 (1H, m), 0.80-0.66 (1H, m), 0.64-0.52 (2H, m), 0.40-0.26 (1H, m)
Example 2-483		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	445 447	1	¹ H-NMR (DMSO-d ₆) δ: 11.96 (1H, s), 8.94 (1H, s), 8.83-8.47 (1H, m), 8.77 (1H, s), 8.04-7.66 (7H, m), 7.52-7.31 (1H, m), 7.01 (1H, d, J = 6.3 Hz), 4.40-4.26 (1H, m), 3.62-3.51 (1H, m), 1.93-1.28 (8H, m)
Example 2-484		Mixture of 6-((2-amino-6-(benzyloxy)cyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1S,2S,6S), (1R,2R,6R)	532	1.1	¹ H-NMR (DMSO-d ₆) δ: 11.75 (1H, s), 8.91-8.85 (1H, m), 8.33-8.27 (1H, m), 8.18 (2H, s), 7.99 (1H, d, J = 12.0 Hz), 7.90-7.72 (4H, m), 7.50-7.30 (1H, m), 7.30-7.12 (5H, m), 6.81 (1H, d, J = 7.8 Hz), 4.66-4.54 (1H, m), 4.44 (1H, d, J = 11.7 Hz), 4.35 (1H, d, J = 11.7 Hz), 3.71-3.56 (2H, m), 2.40 (3H, s), 1.95-1.72 (2H, m), 1.65-1.40 (4H, m)
Example 2-485		6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-5-fluoro-2-(pyridin-3-ylamino)nicotinamide	359	0.57	
Example 2-486		6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-5-fluoro-2-((5-fluoropyridin-3-yl)amino)nicotinamide	377	0.85	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HMR
Example 2-487		6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-5-fluoro-2-((5-fluoro-6-methoxypyridin-3-yl)amino)nicotinamide	407	0.96	
Example 2-488		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((3-(methylamino)-1H-indazol-5-yl)amino)nicotinamide	413	0.68	
Example 2-489		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((3-(methylamino)-1H-indazol-5-yl)amino)nicotinamide	401	0.66	
Example 2-490		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-(methylamino)-1H-indazol-5-yl)amino)nicotinamide	413	0.67	
Example 2-491		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((3-(methylamino)-1H-indazol-5-yl)amino)nicotinamide	401	0.62	
Example 2-492		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	427	0.8	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-493		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(dimethylamino)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	415	0.78	
Example 2-494		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3-(dimethylamino)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	427	0.79	
Example 2-495		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((3-(dimethylamino)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	415	0.73	
Example 2-496		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((1-methyl-3-(methylamino)-1H-indazol-5-yl)amino)nicotinamide	427	0.81	¹ H-NMR (MeOD) δ: 8.06 (1H, d, J = 9.2 Hz), 7.86 (1H, s), 7.77 (1H, d, J = 12.6 Hz), 7.54 (1H, d, J = 9.2 Hz), 4.35-4.28 (1H, m), 3.77-3.71 (1H, m), 3.75 (3H, s), 3.34 (3H, s), 1.85-1.48 (8H, m).
Example 2-497		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((1-methyl-3-(methylamino)-1H-indazol-5-yl)amino)nicotinamide	415	0.79	
Example 2-498		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-methyl-3-(methylamino)-1H-indazol-5-yl)amino)nicotinamide	427	0.79	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-499		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-methyl-3-(methylamino)-1H-indazol-5-yl)amino)nicotinamide	413	0.74	
Example 2-500		6-(((1R,2S)-2-aminocyclohexyl)amino)-2-((3-(dimethylamino)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	485	0.95	
Example 2-501		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-2-((3-(dimethylamino)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	487	0.84	¹ H-NMR (MeOD) δ: 8.00-7.45 (4H, m), 4.57-4.40 (1H, m), 4.28-4.00 (2H, m), 3.85-3.75 (2H, m), 3.73-3.66 (1H, m), 3.61-3.43 (4H, m), 3.35 (6H, s), 3.13 (3H, s), 2.10-1.77 (2H, m).
Example 2-502		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((3-(dimethylamino)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	473	0.92	
Example 2-503		6-(((2S,3R)-2-aminohexan-3-yl)amino)-2-((3-(dimethylamino)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	487	0.98	
Example 2-504		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((3-(dimethylamino)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	485	0.93	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ HMR
Example 2-505		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-2-((3-(dimethylamino)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	501	1.06	
Example 2-506		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3-(dimethylamino)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	499	1.01	
Example 2-507		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((3-(dimethylamino)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	473	0.88	
Example 2-508		6-(((1S,2S)-2-amino-1-(thiophen-2-yl)propyl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	490	0.89	¹ H-NMR (MeOD) δ: 8.55 (1H, s), 7.86-7.76 (2H, m), 7.39 (1H, d, J = 5.0 Hz), 7.23 (1H, d, J = 3.6 Hz), 7.03 (1H, dd, J = 5.0, 3.6 Hz), 5.80 (1H, d, J = 7.9 Hz), 3.88-3.84 (5H, m), 3.52-3.47 (4H, m), 1.49 (3H, d, J = 6.6 Hz).
Example 2-509		6-(((1S,2S)-2-amino-1-(thiophen-2-yl)propyl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	454	0.92	
Example 2-510		6-(((1S,2S)-2-amino-1-(thiophen-2-yl)propyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	483	0.87	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-511		6-(((1S,2S)-2-amino-1-(5-chlorothiophen-2-yl)propyl)amino)-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	524	1.04	
Example 2-512		6-(((1S,2S)-2-amino-1-(5-chlorothiophen-2-yl)propyl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	488	1.07	
Example 2-513		6-(((1R,2S)-2-amino-1-(2-fluorophenyl)propyl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide	502	0.96	
Example 2-514		6-(((1R,2S)-2-amino-1-(2-fluorophenyl)propyl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	466	1	
Example 2-515		2-((5-acetyl-6-methoxypyridin-3-yl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoronicotinamide	417	0.96	¹ H-NMR (MeOD) δ: 8.65 (1H, d, J = 3.0 Hz), 8.29 (1H, d, J = 2.6 Hz), 7.77 (1H, d, J = 11.9 Hz), 4.65-4.42 (1H, m), 4.05 (3H, s), 3.77-3.66 (1H, m), 2.65 (3H, s), 1.92-1.48 (8H, m)
Example 2-516		2-((5-acetyl-6-(methylamino)pyridin-3-yl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoronicotinamide	416	0.76	¹ H-NMR (MeOD) δ: 8.66 (1H, d, J = 2.3 Hz), 8.57 (1H, d, J = 2.6 Hz), 7.83 (1H, d, J = 11.9 Hz), 4.62-4.62 (1H, m), 3.74-3.65 (1H, m), 3.19 (3H, s), 2.69 (3H, s), 1.93-1.47 (8H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-517		2-((5-acetyl-6-morpholinopyridin-3-yl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoronicotinamide	472	0.90	¹ H-NMR (MeOD) δ: 8.62-8.57 (1H, m), 8.37 (1H, d, J = 2.6 Hz), 7.90 (1H, d, J = 11.9 Hz), 4.57-4.44 (1H, m), 3.90-3.84 (4H, m), 3.79-3.71 (1H, m), 3.40-3.30 (4H, m), 2.68 (3H, s), 1.98-1.50 (8H, m)
Example 2-518		2-((5-acetyl-6-methylpyridin-3-yl)amino)-6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoronicotinamide	401	0.74	¹ H-NMR (MeOD) δ: 9.40 (1H, d, J = 2.6 Hz), 8.73 (1H, d, J = 2.3 Hz), 7.91 (1H, d, J = 11.6 Hz), 4.63-4.54 (1H, m), 3.77-3.68 (1H, m), 2.83 (3H, s), 2.71 (3H, s), 2.04-1.54 (8H, m)
Example 2-519		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide	425	0.93	¹ H-NMR (MeOD) δ: 9.21-9.19 (1H, m), 8.83 (1H, s), 8.73 (1H, s), 8.27 (1H, s), 7.89 (1H, d, J = 11.9 Hz), 7.76 (1H, s), 3.98 (1H, dd, J = 4.6 Hz, 9.6 Hz), 3.82-3.70 (1H, m), 2.21 (3H, s), 1.43 (3H, d, J = 6.6 Hz), 1.25-1.09 (1H, m), 0.77-0.66 (1H, m), 0.65-0.48 (2H, m), 0.43-0.32 (1H, m)
Example 2-520		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	445	1.00	¹ H-NMR (MeOD) δ: 9.09-9.05 (1H, m), 9.04-8.94 (1H, m), 8.82-8.74 (1H, m), 8.67 (1H, s), 7.92 (1H, s), 7.89 (1H, d, J = 11.9 Hz), 3.96 (1H, dd, J = 5.3 Hz, 9.2 Hz), 3.78-3.64 (1H, m), 1.43 (3H, d, J = 6.6 Hz), 1.23-1.06 (1H, m), 0.80-0.30 (4H, m)
Example 2-521		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide	413	0.92	¹ H-NMR (MeOD) δ: 9.59 (1H, dd, J = 2.0 Hz, 2.3 Hz), 8.78-8.71 (2H, m), 8.28 (1H, s), 7.90 (1H, d, J = 11.9 Hz), 7.77 (1H, s), 4.78-4.71 (1H, m), 3.70-3.59 (1H, m), 2.21 (3H, s), 1.88-1.60 (2H, m), 1.34 (3H, d, J = 6.9 Hz), 1.01 (3H, t, J = 7.3 Hz)
Example 2-522		6-(((2S,3R)-2-aminopentan-3-yl)amino)-2-((5-(4-chloro-1H-pyrazol-1-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	433 435	0.99	¹ H-NMR (MeOD) δ: 9.39-9.34 (1H, m), 8.78-8.72 (1H, m), 8.72-8.66 (1H, m), 8.66-8.63 (1H, m), 7.93-7.84 (2H, m), 4.75-4.65 (1H, m), 3.68-3.55 (1H, m), 1.90-1.68 (2H, m), 1.32 (3H, d, J = 6.9 Hz), 1.02 (3H, t, J = 7.3 Hz)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-523		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide	439	1.03	¹ H-NMR (MeOD) δ: 9.60-9.55 (1H, m), 8.71-8.61 (2H, m), 8.32 (1H, s), 7.88 (1H, d, J = 11.9 Hz), 7.80 (1H, s), 4.78-4.71 (1H, m), 3.72-3.61 (1H, m), 2.78-2.61 (1H, m), 2.21 (3H, s), 2.16-1.66 (6H, m), 1.32 (3H, d, J = 6.6 Hz)
Example 2-524		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide	427	1.03	¹ H-NMR (MeOD) δ: 9.67-9.62 (1H, m), 8.89-8.83 (1H, m), 8.78-8.74 (1H, m), 8.30 (1H, s), 7.90 (1H, d, J = 11.9 Hz), 7.78 (1H, s), 3.41-3.28 (2H, m), 3.15-3.04 (1H, m), 2.21 (3H, s), 1.78-1.62 (2H, m), 1.57-1.42 (1H, m), 0.91 (3H, d, J = 6.3 Hz), 0.82 (3H, d, J = 6.3 Hz)
Example 2-525		6-(((2S,3R)-2-amino-5-methylhexan-3-yl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide	441	1.05	¹ H-NMR (MeOD) δ: 9.60-9.55 (1H, m), 8.79-8.70 (2H, m), 8.29 (1H, s), 7.90 (1H, d, J = 11.9 Hz), 7.79 (1H, s), 4.78-4.71 (1H, m), 3.73-3.61 (1H, m), 2.21 (3H, s), 1.82-1.68 (2H, m), 1.52-1.37 (1H, m), 1.36 (3H, d, J = 6.6 Hz), 0.92 (3H, d, J = 6.6 Hz), 0.81 (3H, d, J = 6.6 Hz)
Example 2-526		6-(((3R,4R)-3-aminotetrahydro-2H-pyran-4-yl)amino)-5-fluoro-2-((5-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)amino)nicotinamide	427	0.80	¹ H-NMR (MeOD) δ: 9.18-9.13 (1H, m), 8.75-8.66 (2H, m), 8.27 (1H, s), 7.88 (1H, d, J = 11.9 Hz), 7.73 (1H, s), 4.46-4.56 (1H, m), 4.14-4.02 (1H, m), 3.86-3.77 (2H, m), 3.73-3.62 (1H, m), 3.68-3.48 (1H, m), 2.21 (3H, s), 2.17-2.04 (1H, m), 2.00-1.88 (1H, m)
Example 2-527		Mixture of 6-((2-amino-5-hydroxycyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1R,2S,5S), (1S,2R,5R)	442	0.62	
Example 2-528		Mixture of 6-((2-amino-4-(benzyloxy)cyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1R,2S,4R), (1S,2R,4S)	532	1.16	¹ H-NMR (DMSO-d6) δ: 11.94 (1H, s), 8.73-8.67 (1H, m), 8.57-8.51 (1H, m), 8.04 (2H, s), 7.98 (1H, d, J = 12.6 Hz), 7.92-7.78 (4H, m), 7.46-7.30 (6H, m), 6.98 (1H, d, J = 6.6 Hz), 4.56-4.37 (3H, m), 3.84-3.60 (2H, m), 2.50 (3H, s), 2.18-2.00 (1H, m), 1.92-1.62 (4H, m), 1.40-1.45 (1H, m)

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-529		Mixture of 6-((2-amino-5-(benzyloxy)cyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1S,2R,5R), (1R,2S,5S)	532	1.08	¹ H-NMR (DMSO-d ₆) δ: 11.91 (1H, s), 8.72-8.67 (1H, m), 8.49-8.42 (1H, m), 8.16 (2H, s), 7.98 (1H, d, J = 12.6 Hz), 7.98-7.80 (4H, m), 7.50-7.28 (1H, m), 7.26-7.18 (5H, m), 6.94 (1H, d, J = 6.6 Hz), 4.60-4.48 (1H, m), 4.45 (1H, d, J = 12.6 Hz), 4.38 (1H, d, J = 12.6 Hz), 3.70-3.60 (2H, m), 2.64 (3H, s), 2.15-1.98 (1H, m), 1.95-1.45 (5H, m)
Example 2-530		Mixture of 6-((2-amino-3-(benzyloxy)cyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1R,2R,3R), (1S,2S,3S)	532	1.13	¹ H-NMR (DMSO-d ₆) δ: 11.93 (1H, s), 8.73-8.69 (2H, m), 8.69-8.65 (1H, m), 8.24 (2H, s), 8.13-7.92 (5H, m), 7.48-7.26 (6H, m), 6.82 (1H, d, J = 7.2 Hz), 4.84-4.71 (1H, m), 4.64 (1H, d, J = 11.1 Hz), 4.50 (1H, d, J = 11.1 Hz), 4.00-3.85 (1H, m), 3.38-3.25 (1H, m), 2.62 (3H, s), 2.22-2.08 (1H, m), 1.88-1.74 (1H, m), 1.60-1.37 (2H, m), 1.37-1.20 (1H, m)
Example 2-531		Mixture of 6-((2-amino-4-hydroxycyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1R,2S,4R), (1S,2R,4S)	442	0.73	
Example 2-532		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((5-(4-methyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	426	0.93	
Example 2-533		6-(((1S,2R)-2-aminocyclohexyl)amino)-2-((5-(4,5-dimethyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	440	0.99	
Example 2-534		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-2-((5-(4,5-dimethyl-2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)-5-fluoronicotinamide	440	0.99	


Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-535		6-(((1R,2S)-2-aminocyclohexyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-((2-methoxyethyl)amino)-1H-indazol-5-yl)amino)nicotinamide	515	0.88	
Example 2-536		6-(((2S,3R)-2-aminopentan-3-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-((2-methoxyethyl)amino)-1H-indazol-5-yl)amino)nicotinamide	503	0.86	
Example 2-537		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-((2-methoxyethyl)amino)-1H-indazol-5-yl)amino)nicotinamide	515	0.88	
Example 2-538		6-(((2R,3S)-3-aminopentan-2-yl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-3-((2-methoxyethyl)amino)-1H-indazol-5-yl)amino)nicotinamide	503	0.83	
Example 2-539		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	472	0.94	
Example 2-540		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-5-fluoro-2-((3-methoxy-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	486	1.04	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-541		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((3-(difluoromethoxy)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	466	1.09	
Example 2-542		(R)-6-((1-amino-4-methylpentan-2-yl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	510	1.12	
Example 2-543		6-(((1R,2S)-2-amino-1-cyclobutylpropyl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	522	1.11	
Example 2-544		6-(((2R,3S)-3-aminopentan-2-yl)amino)-2-((3-(difluoromethoxy)-1-(2-methoxyethyl)-1H-indazol-5-yl)amino)-5-fluoronicotinamide	497	0.97	
Example 2-545		6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-2-((1,3-dimethyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	426	0.91	
Example 2-546		6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-5-fluoro-2-((1-methyl-1H-indazol-4-yl)amino)nicotinamide	412	0.93	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-547		6-(((1R,2S)-2-amino-1-cyclopropylbutyl)amino)-2-((1-(2,2-difluoroethyl)-1H-indazol-4-yl)amino)-5-fluoronicotinamide	462	0.99	
Example 2-548		6-(((1R,2S)-2-amino-1-(3,5-difluorophenyl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	498	0.92	
Example 2-549		6-(((1R,2S)-2-amino-1-(3,5-difluorophenyl)propyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	513	0.95	
Example 2-550		6-(((1R,2S)-2-amino-1-(3,5-difluorophenyl)propyl)amino)-5-fluoro-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	514	0.95	
Example 2-551		6-(((1R,2S)-2-amino-1-(3,5-difluorophenyl)propyl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	484	1.01	
Example 2-552		6-(((1R,2S)-2-amino-1-(3,4-difluorophenyl)propyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide	498	0.96	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H-NMR
Example 2-553		6-(((1R,2S)-2-amino-1-(3,4-difluorophenyl)propyl)amino)-2-((3-(dimethylamino)-1-methyl-1H-indazol-5-yl)amino)-5-fluoronicotinamide	513	0.96	
Example 2-554		6-(((1R,2S)-2-amino-1-(3,4-difluorophenyl)propyl)amino)-2-((1-(2-methoxyethyl)-1H-indazol-5-yl)amino)nicotinamide	515	0.93	
Example 2-555		6-(((1R,2S)-2-amino-1-(3,4-difluorophenyl)propyl)amino)-2-((1-ethyl-1H-indazol-4-yl)amino)-5-fluoronicotinamide	484	1.02	
Example 2-556		Mixture of 6-((2-amino-3-hydroxycyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1R,2R,3R), (1S,1S,3S)	442	0.8	¹ H-NMR (DMSO-d ₆) δ: 11.89 (1H, s), 8.72-8.68 (1H, m), 8.59-8.54 (1H, m), 8.21 (2H, s), 8.02-7.75 (6H, m), 7.50-7.25 (1H, m), 6.63 (1H, d, J = 7.8 Hz), 4.82-4.70 (1H, m), 4.00-3.85 (1H, m), 3.10-2.96 (1H, m), 2.45 (3H, s), 2.00-1.20 (6H, m)
Example 2-557		Mixture of 6-((2-aminocyclohexyl)amino)-5-fluoro-2-((2-methoxypyrimidin-5-yl)amino)nicotinamide (1R,2S), (1S,2R)	376	0.78	
Example 2-558		Mixture of 6-((2-aminocyclohexyl)amino)-5-fluoro-2-((2-morpholinopyrimidin-5-yl)amino)nicotinamide (1R,2S), (1S,2R)	431	0.85	


Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	¹ H NMR
Example 2-559		Mixture of 6-((2-aminocyclohexyl)amino)-5-fluoro-2-((2-methylpyrimidin-5-yl)amino)nicotinamide (1R,2S), (1S,2R)	360	0.73	
Example 2-560		Mixture of 2-((2-(2H-1,2,3-triazol-2-yl)pyrimidin-5-yl)amino)-6-((2-aminocyclohexyl)amino)-5-fluoronicotinamide (1S,2R), (1R,2S)	413	0.76	
Example 2-561		Mixture of 2-((2-(1H-pyrazol-1-yl)pyrimidin-5-yl)amino)-6-((2-aminocyclohexyl)amino)-5-fluoronicotinamide (1S,2R), (1R,2S)	412	0.85	
Example 2-562		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((2-methoxypyrimidin-5-yl)amino)nicotinamide	376	0.75	
Example 2-563		6-(((1R,2S)-2-amino-1-cyclopropylpropyl)amino)-5-fluoro-2-((2-morpholinopyrimidin-5-yl)amino)nicotinamide	431	0.83	
Example 2-564		Mixture of 6-((2-amino-4,4-difluorocyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1R,2S), (1S,2R)	462	0.9	

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)	HNMR
Example 2-565		Mixture of 6-((2-amino-3-fluorocyclohexyl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (1R,2R,3S), (1S,2S,3R)	444	0.85	

[0461]

Example 3 (Example 4-1)

[Formula 156]



5 A 5M sodium hydroxide aqueous solution (0.038 ml) and a 30% hydrogen peroxide solution (0.008 ml) were added to a DMSO/EtOH (0.1 ml/0.1 ml) solution containing 6-((1-acetyl-3-aminopiperidin-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinonitrile hydrochloride (6 mg), followed by stirring at room temperature for 30 minutes. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layers were washed with saturated saline and dried over anhydrous sodium sulfate. The solvent was distilled away under reduced pressure. Ethyl acetate (0.5 ml) and 4M hydrochloric acid/1,4-dioxane (0.004 ml) were added to the obtained residue, followed by stirring at room temperature for 0.5 hours. The solvent was removed under reduced pressure. The obtained residue was washed with ethyl acetate. A yellow solid of 6-((1-acetyl-3-aminopiperidin-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinamide hydrochloride (3.2 mg) was thus obtained.

10

15

20 (Table 4 (Example 4-1) lists MS data.)





[0462]

Example 4

The compounds shown in table 4 were obtained as described in Example 3.

25 [0463]

[Table 4]

Example	Structure	Compound name	MS(ESI m/z): (M+H)	RT(min)
Example 4-1		Mixture of 6-((1-acetyl-3-aminopiperidin-4-yl)amino)-5-fluoro-2-((5-methylpyridin-3-yl)amino)nicotinamide (3S,4R) (3R,4S)	402	0.51
Example 4-2		Mixture of 6-((1-acetyl-3-aminopiperidin-4-yl)amino)-5-fluoro-2-((5-fluoro-6-morpholinopyridin-3-yl)amino)nicotinamide (3S,4R) (3R,4S)	491	0.8
Example 4-3		Mixture of 6-((1-acetyl-3-aminopiperidin-4-yl)amino)-5-fluoro-2-((6-methyl-5-(2H-1,2,3-triazol-2-yl)pyridin-3-yl)amino)nicotinamide (3S,4R) (3R,4S)	469	0.74
Example 4-4		Mixture of 6-((1-acetyl-3-aminopiperidin-4-yl)amino)-5-fluoro-2-((3-fluoro-2-morpholinopyridin-4-yl)amino)nicotinamide (3S,4R) (3R,4S)	491	0.65

[0464]

Next, the usefulness of representative compounds of the present invention will be described in the following Test Examples.

[0465]

5 Test Example 1: Syk Enzyme Assay

A glutathione S-transferase (GST)-linked full-length human Syk protein (Carna Biosciences), which had been generated using a Baculovirus expression system, was used in the Syk enzyme assay.

7.5 μ l of a reaction solution (0.83 nM Syk, 20 mM HEPES, 10 mM
10 MgCl_2 , 50 mM NaCl, 2 mM DTT, 0.05% BSA, pH 7.0) containing a Syk protein and a predetermined concentration of a test compound was shaken for 2 minutes, and it was then left at rest at room temperature for 13 minutes. Thereafter, 0.5 μ M substrate peptide (Biotin-EDPDYEWPSA-NH₂) and 5 μ L of a solution containing 67.5 μ M ATP were added to the reaction solution, and
15 the obtained mixture was then shaken for 2 minutes. The reaction solution was further left at rest at room temperature for 40 minutes, so as to carry out an enzyme reaction.

Thereafter, 50 μ L of a reaction termination solution [60 nM APC-SA, 0.45 μ g/mL Eu-PT66, 30 mM HEPES (pH 7.0), 150 mM KF, 30 mM EDTA,
20 0.15% BSA, 0.075% Tween20] containing Allophycocyanin-Streptavidin (APC-SA; PerkinElmer) and an Eu-W1024-labeled anti-phosphotyrosine PT66 antibody (Eu-PT66; PerkinElmer) was added to the reaction solution to terminate the enzyme reaction. At the same time, the reaction solution was left at rest at room temperature for 1 hour, so as to carry out an
25 antigen-antibody reaction. Thereafter, using EnVision (PerkinElmer), the level of time-resolved fluorescence was measured at 615 nm and 665 nm, so that the phosphorylation of the substrate peptide was measured.

Table 5 shows the results. The following are used in Table 5 to denote standards for evaluating IC₅₀ of Syk-inhibitory activity.

- 30 A: Up to 10 nM
B: 10 to 50 nM
C: 50 to 100 nM

The set of numbers (XYZ-xyz) given in each Example number column indicates the corresponding Example number (Example XYZ-xyz) in Table 5.

35 [0466]

[Table 5]

Example 2-1	A	Example 2-150	A	Example 2-197	B
Example 2-10	A	Example 2-151	A	Example 2-199	A
Example 2-100	A	Example 2-152	A	Example 2-2	A
Example 2-101	A	Example 2-153	A	Example 2-20	A
Example 2-102	C	Example 2-154	A	Example 2-200	B
Example 2-103	C	Example 2-155	A	Example 2-201	B
Example 2-104	C	Example 2-156	A	Example 2-202	C
Example 2-105	B	Example 2-157	A	Example 2-203	B
Example 2-106	B	Example 2-158	A	Example 2-204	B
Example 2-107	C	Example 2-159	A	Example 2-205	B
Example 2-108	C	Example 2-16	B	Example 2-206	B
Example 2-11	A	Example 2-160	A	Example 2-207	A
Example 2-111	B	Example 2-161	B	Example 2-208	A
Example 2-112	B	Example 2-162	B	Example 2-209	A
Example 2-113	B	Example 2-163	B	Example 2-21	A
Example 2-114	A	Example 2-164	B	Example 2-210	A
Example 2-115	B	Example 2-165	B	Example 2-211	A
Example 2-116	B	Example 2-166	B	Example 2-212	A
Example 2-117	C	Example 2-167	B	Example 2-213	A
Example 2-118	C	Example 2-168	B	Example 2-214	A
Example 2-119	B	Example 2-169	A	Example 2-215	B
Example 2-12	A	Example 2-17	A	Example 2-216	B
Example 2-120	A	Example 2-170	A	Example 2-217	C
Example 2-121	A	Example 2-171	A	Example 2-218	B
Example 2-122	B	Example 2-172	A	Example 2-219	B
Example 2-123	A	Example 2-173	A	Example 2-22	A
Example 2-124	A	Example 2-174	A	Example 2-220	B
Example 2-125	A	Example 2-175	A	Example 2-221	A
Example 2-126	A	Example 2-176	A	Example 2-222	B
Example 2-127	A	Example 2-177	A	Example 2-223	A
Example 2-128	B	Example 2-178	A	Example 2-224	A
Example 2-129	B	Example 2-179	B	Example 2-225	B
Example 2-13	A	Example 2-18	A	Example 2-226	C
Example 2-130	A	Example 2-180	B	Example 2-228	C
Example 2-131	A	Example 2-182	A	Example 2-229	B
Example 2-132	A	Example 2-183	C	Example 2-23	A
Example 2-133	A	Example 2-184	A	Example 2-230	B
Example 2-134	A	Example 2-185	A	Example 2-231	B
Example 2-135	B	Example 2-186	A	Example 2-233	B
Example 2-136	B	Example 2-187	A	Example 2-234	C
Example 2-137	A	Example 2-188	A	Example 2-235	A
Example 2-138	A	Example 2-189	A	Example 2-236	B
Example 2-139	A	Example 2-19	A	Example 2-237	B
Example 2-14	B	Example 2-190	A	Example 2-238	B
Example 2-140	B	Example 2-191	B	Example 2-239	B
Example 2-141	B	Example 2-192	A	Example 2-24	A
Example 2-142	B	Example 2-193	B	Example 2-240	A
Example 2-148	A	Example 2-194	A	Example 2-241	B
Example 2-149	A	Example 2-195	B	Example 2-242	B
Example 2-15	A	Example 2-196	A	Example 2-243	B

Example 2-244	B	Example 2-303	A	Example 2-351	A
Example 2-245	A	Example 2-304	A	Example 2-352	A
Example 2-248	C	Example 2-305	A	Example 2-353	A
Example 2-249	A	Example 2-306	A	Example 2-354	B
Example 2-25	A	Example 2-307	A	Example 2-355	A
Example 2-250	A	Example 2-308	A	Example 2-356	B
Example 2-251	B	Example 2-309	B	Example 2-357	B
Example 2-252	A	Example 2-31	A	Example 2-358	B
Example 2-253	A	Example 2-310	C	Example 2-359	C
Example 2-254	A	Example 2-311	A	Example 2-36	B
Example 2-255	A	Example 2-312	A	Example 2-360	B
Example 2-256	C	Example 2-313	A	Example 2-361	B
Example 2-257	C	Example 2-314	A	Example 2-362	A
Example 2-258	B	Example 2-315	B	Example 2-363	B
Example 2-259	C	Example 2-316	A	Example 2-364	C
Example 2-26	A	Example 2-317	A	Example 2-365	C
Example 2-260	B	Example 2-318	A	Example 2-366	B
Example 2-261	B	Example 2-319	A	Example 2-369	A
Example 2-262	B	Example 2-32	A	Example 2-37	A
Example 2-264	B	Example 2-320	A	Example 2-370	A
Example 2-265	A	Example 2-321	A	Example 2-371	A
Example 2-266	A	Example 2-322	A	Example 2-372	A
Example 2-267	A	Example 2-323	B	Example 2-373	A
Example 2-268	B	Example 2-324	B	Example 2-374	A
Example 2-269	A	Example 2-325	A	Example 2-375	A
Example 2-27	A	Example 2-326	A	Example 2-376	A
Example 2-270	B	Example 2-327	A	Example 2-377	A
Example 2-272	B	Example 2-328	A	Example 2-378	A
Example 2-273	A	Example 2-329	A	Example 2-38	A
Example 2-274	B	Example 2-33	B	Example 2-380	A
Example 2-275	B	Example 2-330	A	Example 2-381	A
Example 2-276	B	Example 2-331	A	Example 2-382	B
Example 2-277	B	Example 2-332	A	Example 2-383	A
Example 2-278	B	Example 2-333	B	Example 2-384	A
Example 2-279	B	Example 2-334	A	Example 2-385	B
Example 2-28	A	Example 2-335	B	Example 2-386	A
Example 2-286	B	Example 2-336	A	Example 2-387	B
Example 2-287	B	Example 2-337	C	Example 2-388	A
Example 2-288	B	Example 2-338	B	Example 2-389	B
Example 2-289	A	Example 2-339	C	Example 2-39	A
Example 2-29	A	Example 2-34	B	Example 2-390	A
Example 2-290	B	Example 2-341	A	Example 2-391	A
Example 2-291	B	Example 2-342	B	Example 2-393	A
Example 2-292	A	Example 2-343	B	Example 2-394	A
Example 2-293	A	Example 2-345	B	Example 2-395	A
Example 2-3	A	Example 2-346	A	Example 2-396	A
Example 2-30	A	Example 2-348	A	Example 2-397	A
Example 2-300	B	Example 2-349	A	Example 2-398	A
Example 2-301	C	Example 2-35	B	Example 2-399	A
Example 2-302	A	Example 2-350	A	Example 2-4	A

Example 2-40	A	Example 2-451	A	Example 2-502	B
Example 2-400	A	Example 2-452	A	Example 2-503	B
Example 2-401	A	Example 2-453	A	Example 2-504	B
Example 2-402	A	Example 2-454	A	Example 2-505	B
Example 2-403	B	Example 2-455	A	Example 2-506	B
Example 2-404	A	Example 2-456	A	Example 2-507	C
Example 2-405	A	Example 2-457	A	Example 2-508	A
Example 2-406	A	Example 2-458	A	Example 2-509	A
Example 2-407	B	Example 2-459	B	Example 2-51	A
Example 2-408	C	Example 2-46	A	Example 2-510	A
Example 2-409	A	Example 2-460	A	Example 2-511	B
Example 2-41	A	Example 2-461	B	Example 2-512	A
Example 2-410	A	Example 2-462	B	Example 2-513	B
Example 2-411	A	Example 2-463	B	Example 2-514	A
Example 2-412	B	Example 2-464	B	Example 2-515	B
Example 2-413	A	Example 2-469	B	Example 2-518	A
Example 2-414	A	Example 2-47	A	Example 2-519	A
Example 2-415	A	Example 2-472	A	Example 2-52	A
Example 2-416	A	Example 2-473	B	Example 2-520	A
Example 2-418	B	Example 2-474	A	Example 2-521	B
Example 2-42	B	Example 2-475	A	Example 2-522	B
Example 2-421	B	Example 2-476	A	Example 2-523	A
Example 2-422	B	Example 2-477	A	Example 2-524	A
Example 2-423	A	Example 2-478	A	Example 2-525	A
Example 2-424	A	Example 2-479	A	Example 2-526	A
Example 2-426	B	Example 2-48	A	Example 2-528	C
Example 2-427	B	Example 2-480	A	Example 2-53	A
Example 2-428	B	Example 2-481	A	Example 2-531	B
Example 2-429	A	Example 2-482	A	Example 2-532	A
Example 2-43	A	Example 2-483	A	Example 2-533	C
Example 2-430	B	Example 2-484	B	Example 2-534	C
Example 2-431	B	Example 2-485	B	Example 2-535	B
Example 2-434	A	Example 2-486	B	Example 2-536	B
Example 2-435	A	Example 2-488	A	Example 2-537	B
Example 2-436	A	Example 2-489	B	Example 2-539	A
Example 2-437	A	Example 2-49	A	Example 2-54	A
Example 2-438	A	Example 2-490	B	Example 2-540	A
Example 2-439	A	Example 2-491	B	Example 2-541	A
Example 2-44	A	Example 2-492	A	Example 2-542	A
Example 2-440	A	Example 2-493	A	Example 2-543	A
Example 2-441	A	Example 2-494	A	Example 2-544	B
Example 2-442	A	Example 2-495	A	Example 2-545	A
Example 2-444	A	Example 2-496	A	Example 2-546	A
Example 2-445	A	Example 2-497	C	Example 2-547	A
Example 2-446	A	Example 2-498	A	Example 2-548	B
Example 2-447	A	Example 2-499	B	Example 2-549	B
Example 2-448	B	Example 2-5	A	Example 2-55	A
Example 2-449	A	Example 2-50	A	Example 2-550	A
Example 2-45	A	Example 2-500	B	Example 2-551	B
Example 2-450	A	Example 2-501	B	Example 2-552	A

Example 2-553	A	Example 3-4	C
Example 2-554	A		
Example 2-555	A		
Example 2-563	C		
Example 2-564	B		
Example 2-57	A		
Example 2-58	B		
Example 2-59	B		
Example 2-6	A		
Example 2-60	B		
Example 2-61	B		
Example 2-62	B		
Example 2-63	B		
Example 2-64	B		
Example 2-65	B		
Example 2-66	B		
Example 2-67	B		
Example 2-68	B		
Example 2-69	B		
Example 2-7	A		
Example 2-70	A		
Example 2-71	A		
Example 2-72	A		
Example 2-73	A		
Example 2-74	A		
Example 2-75	A		
Example 2-76	B		
Example 2-77	B		
Example 2-78	A		
Example 2-79	A		
Example 2-8	A		
Example 2-80	A		
Example 2-81	B		
Example 2-82	A		
Example 2-83	A		
Example 2-84	A		
Example 2-85	B		
Example 2-86	A		
Example 2-87	A		
Example 2-89	A		
Example 2-9	A		
Example 2-90	B		
Example 2-91	B		
Example 2-92	B		
Example 2-96	A		
Example 2-97	C		
Example 2-98	A		
Example 2-99	A		
Example 3-1	B		
Example 3-3	A		

[0467]

Test Example 2: Selectivity of kinase inhibition

The concentrations of test compounds were adjusted to 100 nM. The test compounds were examined using Profiler Pro kits (Caliper) in terms of activity against each of 191 types of kinases excluding Syk. As a result, highly selective compounds (Example 2-1, Example 2-28, and Example 2-77) having kinase inhibitory rates of 75% or more with respect to only 0-1 type of kinases other than Syk and a compound (Example 2-76) having a kinase inhibitory rate of 75% or more with respect to 8 types of kinases were obtained.

[0468]

Test Example 3: TNF α generation assay

THP-1 cells (2×10^5 cells/ml), which were human monocytoid cells, were cultured in the presence of 10 ng/ml IFN- γ (Roche) for 2 days, and they were induced to differentiate into macrophage-like cells. The differentiation-induced THP-1 cells were recovered, and the cells (1×10^6 cell/ml) were then allowed to react with a predetermined concentration of test compound at room temperature for 30 minutes. On the other hand, 100 μ l of human IgG (10 μ g/ml, SIGMA-ALDRICH) diluted with PBS was added to a 96-well plate, and it was then incubated at room temperature overnight. Thereafter, the resultant was washed with PBS twice to produce a human IgG-coated plate. A cell fluid that contained a compound was plated on the obtained human IgG-coated plate (5×10^4 cells/well), and it was then cultured for 7 hours. Thereafter, the cultured solution was recovered, and the amount of TNF α secreted into the culture solution was then measured by the AlphaLISA method (PerkinElmer).

Table 6 shows the results. The following are used in Table 6 to denote standards for evaluating IC₅₀ of TNF α generation inhibitory activity.

A: Up to 65 nM

30 B: 65 to 130 nM

C: 130 to 200 nM

[0469]

[Table 6]

Example 2-1	B	Example 2-196	B	Example 2-28	B
Example 2-10	A	Example 2-20	B	Example 2-29	B
Example 2-100	C	Example 2-207	B	Example 2-302	C
Example 2-101	C	Example 2-208	A	Example 2-306	C
Example 2-106	C	Example 2-209	B	Example 2-31	B
Example 2-11	C	Example 2-21	C	Example 2-311	C
Example 2-114	C	Example 2-210	A	Example 2-316	B
Example 2-121	C	Example 2-211	B	Example 2-317	B
Example 2-122	C	Example 2-213	B	Example 2-319	B
Example 2-123	B	Example 2-214	B	Example 2-32	C
Example 2-125	B	Example 2-215	C	Example 2-320	B
Example 2-126	B	Example 2-216	C	Example 2-321	C
Example 2-127	C	Example 2-218	B	Example 2-322	B
Example 2-129	C	Example 2-219	C	Example 2-325	C
Example 2-130	B	Example 2-22	C	Example 2-326	A
Example 2-131	B	Example 2-220	C	Example 2-327	C
Example 2-132	C	Example 2-221	C	Example 2-328	A
Example 2-133	B	Example 2-222	C	Example 2-329	B
Example 2-134	C	Example 2-223	C	Example 2-33	C
Example 2-135	C	Example 2-224	C	Example 2-330	B
Example 2-136	C	Example 2-225	C	Example 2-332	B
Example 2-137	B	Example 2-229	C	Example 2-334	C
Example 2-138	A	Example 2-23	B	Example 2-348	C
Example 2-139	B	Example 2-230	B	Example 2-36	C
Example 2-141	C	Example 2-233	C	Example 2-362	B
Example 2-142	B	Example 2-235	B	Example 2-366	C
Example 2-148	B	Example 2-239	C	Example 2-37	B
Example 2-149	B	Example 2-24	C	Example 2-370	C
Example 2-150	C	Example 2-240	C	Example 2-372	C
Example 2-151	C	Example 2-242	C	Example 2-374	C
Example 2-152	C	Example 2-243	C	Example 2-375	B
Example 2-153	C	Example 2-249	B	Example 2-376	B
Example 2-159	B	Example 2-25	C	Example 2-377	B
Example 2-162	C	Example 2-251	C	Example 2-378	B
Example 2-17	B	Example 2-252	C	Example 2-38	B
Example 2-170	C	Example 2-253	B	Example 2-381	B
Example 2-172	C	Example 2-254	C	Example 2-39	A
Example 2-173	B	Example 2-255	C	Example 2-4	B
Example 2-174	C	Example 2-26	C	Example 2-40	A
Example 2-179	C	Example 2-265	B	Example 2-404	B
Example 2-18	B	Example 2-266	B	Example 2-406	C
Example 2-180	C	Example 2-267	B	Example 2-409	C
Example 2-182	B	Example 2-268	C	Example 2-41	B
Example 2-184	B	Example 2-269	B	Example 2-410	B
Example 2-185	C	Example 2-27	B	Example 2-413	B
Example 2-186	B	Example 2-270	B	Example 2-414	B
Example 2-187	B	Example 2-272	C	Example 2-415	C
Example 2-188	A	Example 2-273	B	Example 2-416	A
Example 2-19	C	Example 2-274	C	Example 2-42	B
Example 2-193	C	Example 2-275	C	Example 2-428	C

Example 2-43	C	Example 2-75	C
Example 2-434	B	Example 2-76	C
Example 2-436	C	Example 2-77	C
Example 2-437	B	Example 2-78	B
Example 2-438	A	Example 2-79	C
Example 2-439	B	Example 2-8	A
Example 2-44	B	Example 2-80	B
Example 2-440	C	Example 2-82	C
Example 2-441	C	Example 2-83	C
Example 2-442	B	Example 2-84	C
Example 2-45	C	Example 2-87	C
Example 2-454	B	Example 2-9	B
Example 2-46	B	Example 2-91	C
Example 2-47	B	Example 2-96	C
Example 2-472	A	Example 2-98	C
Example 2-474	C		
Example 2-475	B		
Example 2-476	B		
Example 2-477	C		
Example 2-478	B		
Example 2-479	B		
Example 2-48	B		
Example 2-480	B		
Example 2-481	B		
Example 2-482	B		
Example 2-49	C		
Example 2-496	C		
Example 2-5	B		
Example 2-508	B		
Example 2-51	C		
Example 2-518	B		
Example 2-519	C		
Example 2-52	C		
Example 2-521	B		
Example 2-522	C		
Example 2-539	C		
Example 2-54	C		
Example 2-546	C		
Example 2-57	B		
Example 2-58	C		
Example 2-6	C		
Example 2-66	C		
Example 2-67	C		
Example 2-69	C		
Example 2-7	B		
Example 2-70	C		
Example 2-71	A		
Example 2-72	C		
Example 2-73	C		
Example 2-74	B		

[0470]

Test Example 4: Antibody-dependent phagocytosis assay

THP-1 cells (2×10^5 cells/ml), which were human monocytoid cells, were cultured in the presence of 10 ng/ml IFN γ for 2 days, and they were induced to differentiate into macrophage-like cells. The differentiation-induced THP-1 cells were recovered, and the cells (5×10^4 cells/well) were then allowed to react with a predetermined concentration of test compound at room temperature for 30 minutes. Thereafter, Escherichia coli (Life Technologies) labeled with a pH-sensitive dye (pHrodo) was subjected to opsonization using an anti-Escherichia coli antibody (Life Technologies). Then, the resultant was added to the THP-1 cells obtained above, followed by incubation at 37°C for 3 hours. Opsonized Escherichia coli and cell-permeable fluorescent dye (Calcein AM) were simultaneously added thereto, followed by quantitative determination of phagocytosis of opsonized Escherichia coli in viable cells using an IN Cell Analyzer.

The test results are listed in Table 7 below. The following are used in Table 7 to denote standards for evaluating IC₅₀ upon phagocytosis inhibition.

A: Up to 1 μ M

B: 1 to 3 μ M

C: 3 to 6 μ M

[0471]

[Table 7]

25

Example 2-1	A	Example 2-184	A	Example 2-358	B
Example 2-2	A	Example 2-188	A	Example 2-365	A
Example 2-4	B	Example 2-192	B	Example 2-375	B
Example 2-13	A	Example 2-193	B	Example 2-376	A
Example 2-17	A	Example 2-194	B	Example 2-380	B
Example 2-18	A	Example 2-197	C	Example 2-393	B
Example 2-21	A	Example 2-205	B	Example 2-394	C
Example 2-22	A	Example 2-213	A	Example 2-397	B
Example 2-23	A	Example 2-214	B	Example 2-399	B
Example 2-28	A	Example 2-216	B	Example 2-405	B
Example 2-31	A	Example 2-217	B	Example 2-406	B
Example 2-36	B	Example 2-218	B	Example 2-409	B
Example 2-40	A	Example 2-221	A	Example 2-420	A
Example 2-44	A	Example 2-224	B	Example 2-427	B
Example 2-48	A	Example 2-231	B		
Example 2-50	A	Example 2-235	A		
Example 2-51	A	Example 2-236	B		
Example 2-52	A	Example 2-238	A		
Example 2-70	B	Example 2-252	A		
Example 2-66	A	Example 2-265	A		
Example 2-74	A	Example 2-267	B		
Example 2-76	B	Example 2-270	B		
Example 2-77	A	Example 2-272	A		
Example 2-85	A	Example 2-275	B		
Example 2-87	B	Example 2-291	B		
Example 2-89	B	Example 2-292	B		
Example 2-91	B	Example 2-302	B		
Example 2-96	A	Example 2-305	B		
Example 2-98	A	Example 2-306	B		
Example 2-100	B	Example 2-312	A		
Example 2-113	B	Example 2-313	C		
Example 2-114	B	Example 2-316	A		
Example 2-116	B	Example 2-317	B		
Example 2-121	B	Example 2-318	B		
Example 2-123	A	Example 2-319	A		
Example 2-124	A	Example 2-320	A		
Example 2-126	A	Example 2-321	B		
Example 2-127	A	Example 2-322	A		
Example 2-128	B	Example 2-323	B		
Example 2-135	B	Example 2-324	A		
Example 2-136	B	Example 2-325	A		
Example 2-142	A	Example 2-330	A		
Example 2-158	A	Example 2-331	B		
Example 2-162	B	Example 2-332	A		
Example 2-163	B	Example 2-338	A		
Example 2-166	B	Example 2-343	B		
Example 2-169	A	Example 2-346	B		
Example 2-170	B	Example 2-348	A		
Example 2-174	A	Example 2-349	B		
Example 2-178	B	Example 2-352	A		

[0472]

Test Example 5: Ames test

Four *Salmonella typhimurium* strains (TA100, TA1535, TA98, and TA1537) and one *Escherichia coli* strain (WP2uvrA) were used for the Ames
5 test.

A solution containing a test compound (0.1 ml) was added to a test tube. 0.1 M Na-phosphate buffer was added to the tube for no metabolic activation (S9(-)) or an S-9 mix (Kikkoman) (0.5 ml) was added to the tube for metabolic activation (S9(+)). Further, a precultured bacterial cell
10 suspension (0.1 ml) was added to the tube, followed by shaking at 37°C for 20 minutes. Thereafter, 2-ml top agar (a solution prepared by mixing 5 mM L-histidine and a 5 mM D-biotin preparation solution at a volume ratio of 99:1 in a Bacto™ Agar aqueous solution for salmonella or a solution prepared by
15 mixing a 5 mM L-tryptophan aqueous solution and a 5 mM D-biotin preparation solution at a volume ratio of 99:1 in a Bacto™ Agar aqueous solution for *Escherichia coli*) was added, followed by sufficient stirring. The content of the tube was poured onto a minimal glucose agar plate medium and cultured at 37°C for 48 hours.

Colony count was performed using an auto colony counter. The
20 average of colony counts for two plates was defined as the the measurement value.

When the average number of revertant colonies per plate for a test compound was at least two times that for a negative control (DMSO solvent alone) and increased in a dose-dependent manner, such test compound was
25 determined to yield a positive test result.

As a result, the following compounds were found to yield negative test results. .

[0473]

Example 2-1, Example 2-28, Example 2-48, Example 2-76, Example 2-77,
30 Example 2-184, Example 2-188, Example 2-213, Example 2-220, Example 2-221, Example 2-235, Example 2-192, Example 2-302, Example 2-358

[0474]

Test Example 6: Micronucleus test using culture cells

CHL cells (from Chinese hamster lung) were seeded on a 96 well plate
35 (5000 cells/well) and cultured at 37°C and 5% CO₂ for 24 hours. Thereafter, CHL cells were divided into a no metabolic activation (S9(-)) group and a

metabolic activation (S9(+)) group. Phosphate buffered saline (hereinafter abbreviated as PBS(-)) or S-9 mix (Kikkoman) was added to each group. Test substances were separately added, followed by culture at 37°C and 5% CO₂ for 6 hours. Then, each mixture was washed with PBS(-) and a culture solution (100 µl) was again added thereto, followed by culture at 37°C and 5% CO₂ for 18 hours. Cells were fixed with ethanol, followed by removal of the supernatant. PBS(-) containing 2 µg/mL Hoechst 33342 (Invitrogen) and 2 µg/mL CellMask (Invitrogen) (100 µL) was added for staining at room temperature for 30 minutes. Cells were washed with PBS(-), PBS(-) (100 µL) was added thereto, and cells having micronuclei were detected using an IN Cell Analyzer (GE). At least 1000 cells were analyzed per condition for calculation of the frequency of micronuclei. In addition, a cell toxicity test using CellTiter-Glo (Promega) was conducted at the same time as the micronucleus test.

The mutagenicity of each test compound was assessed according to the criteria described below. Dunnett's statistical analysis was conducted for a statistical significance test.

Positive: Statistically significant increase and dose relationship

Negative: No significant increase

False positive: Significant increase and no dose relationship or
Significant increase and strong cell toxicity (survival rate: 50% or less)

Compounds listed in Table 8 were tested according to the above standards. As a result, each compound was found to yield a negative test result.

[0475]

[Table 8]

Example 2-1	Example 2-154	Example 2-322
Example 2-2	Example 2-156	Example 2-325
Example 2-4	Example 2-158	Example 2-331
Example 2-13	Example 2-162	Example 2-332
Example 2-21	Example 2-163	Example 2-338
Example 2-22	Example 2-169	Example 2-346
Example 2-23	Example 2-170	Example 2-348
Example 2-24	Example 2-171	Example 2-349
Example 2-25	Example 2-172	Example 2-352
Example 2-30	Example 2-174	Example 2-358
Example 2-31	Example 2-184	Example 2-370
Example 2-41	Example 2-188	Example 2-373
Example 2-50	Example 2-192	Example 2-374
Example 2-52	Example 2-194	Example 2-376
Example 2-59	Example 2-197	Example 2-380
Example 2-63	Example 2-208	Example 2-393
Example 2-65	Example 2-214	Example 2-394
Example 2-76	Example 2-216	Example 2-397
Example 2-77	Example 2-217	Example 2-434
Example 2-78	Example 2-218	Example 2-435
Example 2-83	Example 2-223	Example 2-436
Example 2-85	Example 2-224	Example 2-439
Example 2-87	Example 2-235	Example 2-441
Example 2-89	Example 2-236	Example 2-444
Example 2-91	Example 2-238	Example 2-448
Example 2-96	Example 2-252	Example 2-473
Example 2-98	Example 2-265	Example 2-483
Example 2-100	Example 2-268	
Example 2-101	Example 2-270	
Example 2-113	Example 2-272	
Example 2-114	Example 2-275	
Example 2-116	Example 2-291	
Example 2-121	Example 2-292	
Example 2-123	Example 2-305	
Example 2-124	Example 2-306	
Example 2-126	Example 2-312	
Example 2-127	Example 2-316	
Example 2-135	Example 2-318	
Example 2-136	Example 2-319	
Example 2-142	Example 2-320	

[0476]

Test Example 7: Rat-type-II-collagen-induced arthritis

The compounds were tested to examine effects upon rat type II collagen arthritis. Equivalent volumes of 0.05 mol/L acetic acid in which a 3 mg/mL bovine type II collagen solution (Collagen Gijutsu Kenshu-Kai) had been dissolved and Freund's incomplete adjuvant (Wako Pure Chemical Industries, Ltd.) were mixed to prepare an emulsion. A portion of the emulsion (0.5 ml) was intradermally injected into the tail bases of 6- to 8-week-old female Lewis rats (Charles River Laboratories Japan, Inc.) (Day 0). Each rat was subjected to the same treatment on Day 7 after the initial inoculation so as to induce arthritis. Each test compound was orally administered from Day 7 to Day 20 once daily. At a given time during the period from Day 7 to Day 21, the rat hindlimb volume was determined using a plethysmometer (UGO BASILE), and the result was designated as an arthritis index. The following compound group significantly inhibited hindlimb swelling on Day 20 compared with the control group in the case of oral administration at 10 mg/kg/day (Student's t-test): Example 2-1, Example 2-76, and Example 2-184.

[0477]

Test Example 8: Mouse thrombocytopenia model

Test compounds were tested to examine effects upon mouse thrombocytopenia. Each test compound (50 mg/kg) was administered to 5- to 7-week-old female BALB/c mice (Charles River Laboratories Japan, Inc.). One hour thereafter, an anti-mouse CD41 (Integrin α IIb) antibody (SCB) (1 μ g (200 μ l)) was intravenously administered to each mouse so as to induce thrombocytopenia. Four hours after administration of the anti-CD41 antibody, blood sampling from the postcava was performed. The platelet count was determined using an automated hematology analyzer.

As a result, the groups to which the following compounds had been administered showed the improvement of platelet count (50% or more improvement) compared with the control group:

Example 2-1, Example 2-2, Example 2-18, Example 2-28, Example 2-48, Example 2-76, Example 2-77, Example 2-87, Example 2-89, Example 2-91, Example 2-100, Example 2-123, Example 2-142, Example 2-174, Example 2-184, Example 2-188, Example 2-192, Example 2-193, Example 2-213, Example 2-251, Example 2-252, Example 2-265, Example 2-267, Example


2-302, Example 2-326, Example 2-328, Example 2-330, Example 2-331, Example 2-348, Example 2-352, Example 2-358, Example 2-415, Example 2-456, Example 2-473, and Example 2-483.

[Name of Document] Claims

[Claim 1]

A nicotinamide derivative represented by the following formula (I) or a salt thereof:


5 [Formula 1]



wherein

R¹ is a substituent represented by the following formula (II-1), (III-1), or (IV-1):

10 [Formula 2]



(wherein

R³ represents a hydrogen atom, or a C₁₋₆ alkyl, C₃₋₈ cycloalkyl, phenyl, pyridyl or thienyl group, each of which optionally has at least one substituent,

15 R⁴ represents a hydrogen atom, C₁₋₆ alkyl group, or C₃₋₈ cycloalkyl group,

R⁵ represents a hydroxy group, a halogen atom, or a C₁₋₆ alkyl or C₁₋₆ alkoxy group, each of which optionally has at least one substituent,

n represents an integer of 0-2,

20 when n is 2, R⁵ may be the same or different, and two R⁵s may, together with a carbon atom to which they bind, form a C₃₋₈ cycloalkane ring, X¹ represents an oxygen atom or -N(R⁶)- (wherein R⁶ represents a hydrogen atom or acyl group, and


"*" represents a binding site)), and

25 R² represents a pyridyl, indazolyl, phenyl, pyrazolopyridyl, benzisoxazolyl, pyrimidinyl, or quinolyl group, each of which optionally has at least one substituent.

[Claim 2]

The nicotinamide derivative or a salt thereof according to claim 1, wherein R^1 is a substituent represented by the following formula (II-2), (III-2), or (IV-2):

[Formula 3]



(wherein R^3 , R^4 , R^5 , n , and X^1 have the same definitions as those described in claim 1).

[Claim 3]

The nicotinamide derivative or a salt thereof according to claim 1 or 2, wherein R^3 is a hydrogen atom, or a C_{1-6} alkyl, C_{3-8} cycloalkyl, phenyl, pyridyl or thienyl group, each of which optionally has at least one substituent selected from a substituent group α_{1-1}

substituent group α_{1-1} : a halogen atom and C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkylthio, phenyl, and pyrazolyl groups, each of which optionally has at least one halogen atom.

[Claim 4]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 3, wherein R^4 represents a hydrogen atom or a C_{1-3} alkyl group.

[Claim 5]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 4, wherein R^5 represents a hydroxy group, a halogen atom, or a C_{1-6} alkyl or C_{1-6} alkoxy group, each of which optionally has at least one phenyl group.

[Claim 6]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 5, wherein n is an integer of 0.

[Claim 7]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 6, wherein X^1 represents an oxygen atom.

[Claim 8]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 7, wherein R^2 represents a pyridyl, indazolyl, phenyl,

pyrazolopyridyl, benzisoxazolyl, pyrimidinyl, or quinolyl group, each of which optionally has at least one substituent selected from a substituent group α_{2-1}

substituent group α_{2-1} : a halogen atom and C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, (di)C₁₋₆ alkylamino, acyl, pyrazolyl, triazolyl, morpholinyl, and pyrrollysyl groups, each of which optionally has at least one substituent selected from a substituent group β_{2-1}

substituent group β_{2-1} : a halogen atom, oxo, and C₁₋₆ alkyl and C₁₋₆ alkoxy groups

10 [Claim 9]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 7, wherein R² represents a pyridyl group optionally having at least one substituent selected from a substituent group α_{3-1}


substituent group α_{3-1} : a halogen atom and C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, (di)C₁₋₆ alkylamino, acyl, pyrazolyl, triazolyl, morpholinyl, and pyrrollysyl groups, each of which optionally has at least one substituent selected from a substituent group β_{3-1}

substituent group β_{3-1} : a halogen atom and C₁₋₆ alkyl and C₁₋₆ alkoxy groups.

[Claim 10]

20 The nicotinamide derivative or a salt thereof according to any one of claims 1 to 7, wherein R² is a substituent represented by the following formula (V-1) or (V-2):

[Formula 4]



25 (wherein R⁷, R⁸, R⁹ and R¹⁰, which may be the same or different, are each a hydrogen atom or a substituent selected from a substituent group α_{3-2})
 substituent group α_{3-2} : a halogen atom and C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, (di)C₁₋₆ alkylamino, pyrazolyl, triazolyl, and morpholinyl groups, each of which optionally has at least one substituent selected from a substituent group β_{3-2}

substituent group β_{3-2} : a halogen atom and a C₁₋₆ alkyl group.

[Claim 11]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 7, wherein R^2 represents an indazolyl group optionally having at least one substituent selected from a substituent group α_{4-1}


substituent group α_{4-1} : a halogen atom and C_{1-6} alkyl, C_{1-6} alkoxy, (di) C_{1-6} alkylamino, and pyrrolsyl groups, each of which optionally has at least one substituent selected from a substituent group β_{4-1}

substituent group β_{4-1} : a halogen atom, oxo, and a C_{1-6} alkoxy group.

[Claim 12]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 7, wherein R^2 is a substituent represented by the following formula (VI-1) or (VI-2):

[Formula 5]



(wherein R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} , which may be the same or different, are each a hydrogen atom or a substituent selected from a substituent group α_{4-2})

substituent group α_{4-2} : a halogen atom and C_{1-6} alkyl, C_{1-6} alkoxy, and (di) C_{1-6} alkylamino groups, each of which optionally has at least one substituent selected from a substituent group β_{4-2}

substituent group β_{4-2} : a halogen atom and a C_{1-6} alkoxy group.

[Claim 13]


The nicotinamide derivative or a salt thereof according to any one of claims 1 to 7, wherein R^2 represents a phenyl group optionally having at least one substituent selected from a substituent group α_{5-1}

substituent group α_{5-1} : a halogen atom and C_{1-6} alkyl, C_{1-6} alkoxy, acyl, and triazolyl groups, each of which optionally has at least one halogen.

[Claim 14]

The nicotinamide derivative or a salt thereof according to any one of claims 1 to 7, wherein R^2 is a substituent represented by the following formula (VII-1):

[Formula 6]



(VII-1)

(wherein R^{17} , R^{18} , and R^{19} , which may be the same or different, are each a hydrogen atom or a substituent selected from a substituent group α_{5-2})
 substituent group α_{5-2} : a halogen atom a C_{1-6} alkoxy group; and C_{1-6} alkyl and
 5 triazolyl groups each of which optionally has at least one halogen atom.

[Claim 15]

The nicotinamide derivative or a salt thereof according to any one of
 claims 1 to 7, wherein R^2 represents a pyrazolopyridyl group optionally
 having at least one substituent selected from a substituent group α_{6-1}


10 substituent group α_{6-1} : a halogen atom and C_{1-6} alkyl, C_{1-6} alkoxy, and (di) C_{1-6}
 alkylamino groups, each of which optionally has at least one substituent
 selected from a substituent group β_{6-1}

substituent group β_{6-1} : a halogen atom and a C_{1-6} alkoxy group.


[Claim 16]

15 The nicotinamide derivative or a salt thereof according to any one of
 claims 1 to 7, wherein R^2 represents a substituent represented by the following
 formula (VIII-1) or (VIII-2):

[Formula 7]



(VIII-1)




(VIII-2)

20 (wherein R^{20} , R^{21} , R^{22} , and R^{23} , which may be the same or different, are each a
 hydrogen atom or a substituent selected from the substituent group α_{6-1}
 described in claim 15).

[Claim 17]

The nicotinamide derivative or a salt thereof according to claim 1,
 25 wherein the nicotinamide derivative is represented by the following formula
 (I-1)

[Formula 8]:




(wherein R^2 , R^3 , and R^4 have the same definitions as R^2 , R^3 , and R^4 described in claim 1).

[Claim 18]

- 5 The nicotinamide derivative or a salt thereof according to claim 1, wherein the nicotinamide derivative is represented by the following formula (I-3):

[Formula 9]




- 10 (wherein R^2 has the same definition as R^2 described in claim 1).

[Claim 19]

 The nicotinamide derivative or a salt thereof according to claim 1, wherein the nicotinamide derivative is represented by the following formula (I-5):

- 15 [Formula 10]



(wherein R^2 has the same definition as R^2 described in claim 1).

[Claim 20]

- A pharmaceutical composition, comprising the nicotinamide derivative or a salt thereof according to any one of claims 1-19.

[Claim 21]

 The pharmaceutical composition according to claim 20, which is used for treatment of Syk-related diseases.

[Claim 22]

- 25 The pharmaceutical composition according to claim 20, which is used for treatment of a disease selected from the group consisting of rheumatoid arthritis and idiopathic thrombocytopenic purpura.