


**Piperidinyl- and piperazinyl-alkylcarbamate
derivatives, preparation and therapeutic application
thereof**

5 The invention relates to piperidyl- and piperazinyl-alkylcarbamate derivatives and to the preparation and therapeutic application thereof.

The compounds of the invention correspond to the general formula (I):



(I)

10

in which

A represents a nitrogen atom or a group CR₂ in which R₂ represents a hydrogen or fluorine atom or a hydroxyl, cyano, trifluoromethyl, C₁₋₆-alkyl or C₁₋₆-alkoxy group;

15 n represents an integer equal to 2 or 3 and m

represents an integer equal to 2 when A represents a nitrogen atom;

n represents an integer equal to 1, 2 or 3 and m

represents an integer equal to 1 or 2 when A represents

20 a group CR₂;

B represents a covalent bond or a C₁₋₈-alkylene group;

R₁ represents a group chosen from phenyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl,

- oxazolyl, isoxazolyl, thiazolyl, isothiazolyl,
 imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl,
 naphthyl, quinolyl, tetrahydroquinolyl, isoquinolyl,
 tetrahydroisoquinolyl, phthalazinyl, quinazolinyl,
 5 quinoxalinyl, naphthyridinyl, cinnolyl,
 imidazopyrimidinyl, thienopyrimidinyl, benzofuranyl,
 benzothienyl, benzimidazolyl, benzoxazolyl,
 benzisoxazolyl, benzothiazolyl, benzisothiazolyl,
 indazolyl, pyrrolopyridyl, furopyridyl,
 10 dihydrofuropyridyl, thienopyridyl, dihydro-
 thienopyridyl, imidazopyridyl, pyrazolopyridyl,
 oxazolopyridyl, isoxazolopyridyl and thiazolopyridyl;
 the group R_1 optionally being substituted with one or
 more groups R' and/or R'' ;
- 15 R' represents a halogen atom or a cyano, nitro,
 hydroxyl, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -thioalkyl, C_{1-6} -
 fluoroalkyl, C_{1-6} -fluoroalkoxy, C_{1-6} -fluorothioalkyl,
 C_{3-7} -cycloalkyl, C_{3-7} -cycloalkyl- C_{1-6} -alkylene,
 azetidiny, piperidyl, pyrrolidinyl, morpholinyl,
 20 piperazinyl, azepinyl, NH_2 , NHR_6 , NR_6R_7 , NR_6COR_7 ,
 $NR_6SO_2R_7$, COR_6 , CO_2R_6 , SO_2R_6 , $SO_2NR_6R_7$ or $-O-(C_{1-6}$ -
 alkylene)-O- group;
- R'' represents a phenyl, imidazolyl, pyridyl, pyridaz-
 inyl, pyrazinyl or pyrimidinyl;
- 25 the group(s) R'' being optionally substituted with one
 or more groups R' , which may be identical to or
 different from each other;

R_3 represents a group of general formula CHR_4CONHR_5 in which

R_4 represents a hydrogen atom or a C_{1-6} -alkyl group and

R_5 represents a hydrogen atom or a C_{1-6} -alkyl, C_{3-7} -cycloalkyl or C_{3-7} -cycloalkyl- C_{1-6} -alkylene group;

R_6 and R_7 represent, independently of each other, a C_{1-6} -alkyl group.

Among the compounds of general formula (I), a first subgroup of compounds consists of compounds for which:

A represents a nitrogen atom; and/or

n represents an integer equal to 2 or 3 and m

represents an integer equal to 2; and/or

B represents a C_{1-8} -alkylene group, more particularly an ethyl or propyl; and/or

R_1 represents a group chosen from phenyl, pyridyl, pyrimidinyl, thiadiazolyl and naphthyl;

the group R_1 being optionally substituted with one or more groups R' and/or R'' ; and/or

R' represents a halogen atom, more particularly a chlorine, or a nitro or C_{1-6} -fluoroalkyl group, more particularly a trifluoromethyl; and/or

R'' represents a phenyl optionally substituted with one or more groups, which may be identical to or different from each other, chosen from a halogen atom, more particularly a chlorine, and a cyano, C_{1-6} -alkoxy, more particularly a methoxy, or C_{1-6} -fluoroalkoxy, more particularly a trifluoromethoxy, group; and/or

R_3 represents a group of general formula CHR_4CONHR_5 in which

R_4 represents a hydrogen atom and

R_5 represents a hydrogen atom or a C_{1-6} -alkyl group, more particularly a methyl or an ethyl, a C_{3-7} -cycloalkyl group, more particularly a cyclopropyl, or a C_{3-7} -cycloalkyl- C_{1-6} -alkylene group, more particularly a cyclopropyl-methylene.

Among the compounds of general formula (I), a second subgroup of compounds consists of compounds for which:

A represents a group CR_2 in which R_2 represents a hydrogen or fluorine atom or a hydroxyl group; and/or

m represents an integer equal to 1 or 2 and n represents an integer equal to 1 or 2; and/or

B represents a covalent bond or a C_{1-4} -alkylene group, more particularly a methyl, ethyl or n-propyl; and/or

R_1 represents a group chosen from phenyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, thiazolyl, quinolyl, isoquinolyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, furopyridyl, thienopyrimidinyl, imidazopyrimidinyl, benzothiazolyl, benzimidazolyl and benzoxazolyl;

the group R_1 being optionally substituted with one or more groups R' and/or R'' ; and/or

R' represents a halogen atom, more particularly a fluorine, a chlorine or a bromine, or a cyano group,

C_{1-6} -alkyl, more particularly a methyl, ethyl, n-propyl
 or isobutyl, a C_{1-6} -alkoxy, more particularly a methoxy,
 C_{1-6} -fluoroalkyl, more particularly a trifluoromethyl,
 C_{1-6} -fluoroalkoxy, more particularly a trifluoromethoxy,
 5 C_{3-7} -cycloalkyl, more particularly a cyclopropyl or
 cyclopentyl, pyrrolidinyl, NH_2 , NR_6R_7 or COR_6 ; and/or
 R'' represents a phenyl, imidazolyl or pyridyl; the
 group(s) R'' being optionally substituted with one or
 more groups R' , which may be identical to or different
 10 from each other, more particularly with one or more
 chlorine or fluorine atoms; and/or
 R_3 represents a group of general formula CHR_4CONHR_5 in
 which
 R_4 represents a hydrogen atom or a C_{1-6} -alkyl group,
 15 more particularly a methyl, and
 R_5 represents a hydrogen atom or a C_{1-6} -alkyl group,
 more particularly a methyl or an ethyl, C_{3-7} -cycloalkyl,
 more particularly a cyclopropyl, C_{3-7} -cycloalkyl- C_{1-6} -
 alkylene, more particularly a cyclopropyl-methylene;
 20 and/or
 R_6 and R_7 represent, independently of each other, a C_{1-6} -
 alkyl group, more particularly a methyl.

Among the compounds of general formula (I), a
 third subgroup of compounds consists of compounds for
 25 which:

A represents a group CR_2 in which R_2 represents a
 hydrogen atom; and/or
 m is equal to 2 and n is equal to 2; and/or

B represents an ethyl group; and/or

R₁ represents a group chosen from phenyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, thiazolyl, quinolyl, isoquinolyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, furopyridyl, thienopyrimidinyl, imidazopyrimidinyl, benzothiazolyl, benzimidazolyl and benzoxazolyl;

the group R₁ being optionally substituted with one or more groups R' and/or R''; and/or

10 R' represents a halogen atom, more particularly a fluorine, a chlorine or a bromine, or a cyano group, C₁₋₆-alkyl, more particularly a methyl, ethyl, n-propyl or isobutyl, a C₁₋₆-alkoxy, more particularly a methoxy, C₁₋₆-fluoroalkyl, more particularly a trifluoromethyl, 15 C₁₋₆-fluoroalkoxy, more particularly a trifluoromethoxy, C₃₋₇-cycloalkyl, more particularly a cyclopropyl or cyclopentyl, pyrrolidinyl, NH₂, NR₆R₇ or COR₆; and/or R'' represents a phenyl, imidazolyl or pyridyl;

the group(s) R'' being optionally substituted with one 20 or more groups R', which may be identical to or different from each other, more particularly with one or more chlorine or fluorine atoms; and/or


R₃ represents a group of general formula CHR₄CONHR₅ in which

25 R₄ represents a hydrogen atom and

R₅ represents a hydrogen atom or a C₁-C₆-alkyl group, more particularly a methyl or an ethyl; and/or

R_6 and R_7 represent, independently of each other, a C_{1-6} -alkyl group, more particularly a methyl.

A subject of the invention is also, among the compounds of general formula (I), compounds
 5 corresponding to the general formula (I'):



(I')

in which

A represents a nitrogen atom or a group CR_2 in which R_2 represents a hydrogen or fluorine atom or a hydroxyl,
 10 cyano, trifluoromethyl, C_{1-5} -alkyl or C_{1-5} -alkoxy group;
 n represents an integer equal to 2 or 3 and m represents an integer equal to 2 when A represents a nitrogen atom;
 n represents an integer equal to 1, 2 or 3 and m
 15 represents an integer equal to 1 or 2 when A represents a group CR_2 ;
 B represents a covalent bond or a C_{1-8} -alkylene group;
 R_1 represents a group chosen from phenyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl,
 20 thiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, naphthyl, quinolyl, tetrahydroquinolyl, isoquinolyl, tetrahydroisoquinolyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, cinnolyl, imidazopyrimidinyl, thienopyrimidinyl, benzofuranyl,

- benzothienyl, benzimidazolyl, benzoxazolyl,
 benzisoxazolyl, benzothiazolyl, benzisothiazolyl,
 indazolyl, pyrrolopyridyl, furopyridyl,
 dihydrofuropyridyl, thienopyridyl, dihydrothieno-
 5 pyridyl, imidazopyridyl, pyrazolopyridyl,
 oxazolopyridyl, isoxazolopyridyl and thiazolopyridyl;
 optionally substituted with one or more substituents
 chosen from a halogen atom and a cyano, nitro, C₁₋₅-
 alkyl, C₁₋₃-alkoxy, C₁₋₅-thioalkyl, C₁₋₅-fluoroalkyl,
 10 C₁₋₃-fluoroalkoxy, C₁₋₃-fluorothioalkyl, C₃₋₅-cycloalkyl,
 C₃₋₅-cycloalkyl-C₁₋₃-alkylene, piperidyl, pyrrolidinyl,
 morpholinyl, NH₂, NHR₆, NR₆R₇, NHCOR₆, COR₆, CO₂R₆, SO₂R₆,
 -O-(C₁₋₃-alkylene)-O-, phenyl, pyridyl or pyrimidinyl
 group;
- 15 the phenyl, pyridyl and pyrimidinyl groups possibly
 being substituted with one or more substituents chosen
 from a halogen atom and a cyano, nitro, hydroxyl, C₁₋₅-
 alkyl, C₁₋₃-alkoxy, C₁₋₅-thioalkyl, C₁₋₅-fluoroalkyl,
 C₁₋₃-fluoroalkoxy, C₁₋₃-fluorothioalkyl, C₃₋₅-cycloalkyl,
 20 C₃₋₅-cycloalkyl-C₁₋₃-alkylene, piperidyl, pyrrolidinyl,
 morpholinyl, NH₂, NHR₆, NR₆R₇, NHCOR₆, COR₆, CO₂R₆, SO₂R₆,
 or -O-(C₁₋₃-alkylene)-O- group;
- R₃ represents a group of general formula CHR₄CONHR₅ in
 which
- 25 R₄ represents a hydrogen atom or a C₁₋₃-alkyl group and
 R₅ represents a hydrogen atom or a C₁₋₃-alkyl, C₃₋₅-
 cycloalkyl or C₃₋₅-cycloalkyl-C₁₋₃-alkylene group;

R_6 and R_7 represent, independently of each other, a C_{1-3} -alkyl group.

The compounds of general formula (I) may comprise one or more asymmetric carbons. They may exist
5 in the form of enantiomers or diastereoisomers. These enantiomers and diastereoisomers, and also mixtures thereof, including racemic mixtures, form part of the invention.

The compounds of general formula (I) may
10 exist in the form of bases or of acid-addition salts. Such addition salts form part of the invention.

These salts are advantageously prepared with pharmaceutically acceptable acids, but the salts of other acids that are useful, for example, for the
15 purification or isolation of the compounds of formula (I) also form part of the invention.

The compounds of general formula (I) may be in the form of hydrates or solvates, i.e. in the form of associations or combinations with one or more water
20 molecules or with a solvent. Such hydrates and solvates also form part of the invention.

In the context of the invention, the following definitions apply:

- C_{t-z} in which t and z may take values from 1 to 8, a
25 carbon-based chain possibly containing from t to z carbon atoms, for example a C_{1-3} carbon-based chain which may contain from 1 to 3 carbon atoms,

- alkyl, a saturated, linear or branched aliphatic group, for example a C₁₋₃-alkyl group represents a linear or branched carbon-based chain of 1 to 3 carbon atoms, more particularly a methyl, ethyl, propyl or 1-methylethyl,
- alkylene, a saturated, linear or branched divalent alkyl group, for example a C₁₋₃-alkylene group represents a linear or branched divalent carbon-based chain of 1 to 3 carbon atoms, more particularly a methylene, ethylene, 1-methylethylene or propylene,
- cycloalkyl, a cyclic alkyl group, for example a C₃₋₅-cycloalkyl group represents a cyclic carbon-based group of 3 to 5 carbon atoms, more particularly a cyclopropyl, cyclobutyl or cyclopentyl,
- alkoxy, an -O-alkyl group containing a saturated, linear or branched aliphatic chain,
- thioalkyl, an -S-alkyl group containing a saturated, linear or branched aliphatic chain,
- fluoroalkyl, an alkyl group in which one or more hydrogen atoms have been replaced with a fluorine atom,
- fluoroalkoxy, an alkoxy group in which one or more hydrogen atoms have been replaced with a fluorine atom,
- fluorothioalkyl, a thioalkyl group in which one or more hydrogen atoms have been replaced with a fluorine atom,

- halogen atom, a fluorine, a chlorine, a bromine or an iodine.


The compounds of the invention may be prepared according to various methods, illustrated by
5 the schemes that follow.

Thus, a first method (Scheme 1) consists in reacting an amine of general formula (II), in which A, B, R₁, n and m are as defined above, with a carbonate of general formula (IIIa) in which Z represents a hydrogen
10 atom or a nitro group, R₄ is as defined above and R represents a methyl or ethyl group. The carbamate ester of general formula (Ia) thus obtained is then converted into a compound of general formula (I), via aminolysis using an amine of general formula R₅NH₂ in which R₅ is
15 as defined above. The aminolysis reaction may be performed in a solvent such as methanol or a mixture of solvents such as methanol and tetrahydrofuran or methanol and dioxane.

One variant for obtaining the compounds of
20 general formula (I) (Scheme 1) consists in reacting an amine of general formula (II), as defined above, with a carbonate of general formula (IIIb), in which Z represents a hydrogen atom or a nitro group and R₄ and R₅ are as defined above, in a solvent such as toluene or
25 dichloroethane, at a temperature of between 0°C and 80°C.

The carbonates of general formula (IIIa) and (IIIb) may be prepared according to any method

described in the literature, for example by reacting an alcohol of general formula $\text{HOCHR}_4\text{COOR}$, in which R represents a methyl or ethyl group, or $\text{HOCHR}_4\text{CONHR}_5$ in which R_4 and R_5 are as defined above, with phenyl chloroformate or 4-nitrophenyl chloroformate, in the presence of a base such as triethylamine or diisopropylethylamine.




10

A second method for obtaining the compounds of general formula (I) (Scheme 2) consists in reacting a derivative of general formula (IIa) in which Y represents a hydroxyl, mesylate or tosylate group or a chlorine, bromine or iodine atom and A, B, R_1 , n and m are as defined above, with an oxazolidinedione of general formula (IV) in which R_4 is as defined above, to

give the oxazolidinedione derivative of general formula (V). In the case where Y represents a hydroxyl group, the reaction may be performed according to the Mitsunobu conditions (Synthesis, 1981, 1-28), for example via the action of diethyl or diisopropyl azodicarboxylate in the presence of triphenylphosphine. In the case where Y represents a chlorine, bromine or iodine atom or a mesylate or tosylate group, the reaction may be performed in the presence of a base such as 1,1,3,3-tetramethylguanidine, sodium hydride or sodium tert-butoxide in a solvent such as tetrahydrofuran, acetonitrile or dimethylformamide, at a temperature of between 0°C and the reflux temperature of the solvent. The oxazolidinedione derivative of general formula (V) thus obtained is then converted into a compound of general formula (I), via aminolysis using an amine of general formula R_5NH_2 in which R_5 is as defined above.

20



The compounds of general formulae (I), (Ia), (II), (IIa) and (V), in which R_1 is substituted with a group R'' , may also be prepared by reacting the corresponding compounds of general formulae (I), (Ia), (II), (IIa) and (V), for which R_1 is substituted with a chlorine, bromine or iodine atom or with a triflate group in the position into which the group R'' is to be introduced, with an aryl- or heteroaryl-boronic acid derivative according to the Suzuki reaction conditions (Chem. Rev. 1995, 95, 2457-2483) or with an aryl- or heteroaryl-trialkylstannane derivative according to the Stille reaction conditions (Angew. Chem. Int. Ed. 1986, 25, 504-524).

The compounds of general formulae (II), (IIa) and (IV), when their preparation method is not described, are commercially available or are described in the literature, or alternatively may be prepared according to methods that are described therein or known to those skilled in the art.

The amines of general formula R_5NH_2 are commercially available.

According to another of its aspects, a subject of the invention is also the compounds of formula (Ia) in which n , m , A , B , R_1 and R_4 are as defined above and R represents a methyl or ethyl group. The compounds of formula (Ia) are useful as synthetic intermediates for the preparation of the compounds of formula (I).

According to another of its aspects, a subject of the invention is also the compounds of formula (V) in which n, m, A, B, R₁ and R₄ are as defined above, the following compounds being excluded:

- 5 * 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-2,4-oxazolidinedione
- * 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-5-methyl-2,4-oxazolidinedione
- * 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-
- 10 piperidyl]-5-ethyl-2,4-oxazolidinedione
- * 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-5-propyl-2,4-oxazolidinedione
- * 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-5-(1-methylethyl)-2,4-oxazolidinedione.

15 The compounds of formula (V) are useful as synthetic intermediates for the preparation of the compounds of formula (I).

The examples that follow illustrate the preparation of a number of compounds of the invention.

20 These examples are not limiting and are given merely to illustrate the invention. The microanalyses and the IR, NMR and/or LC/MS (Liquid Chromatography coupled to Mass Spectroscopy) spectra confirm the structures and purities of the compounds obtained.


25 m.p. (°C) represents the melting point in degrees Celsius.

The numbers indicated in parentheses in the example titles correspond to those in the first column of the table hereinbelow.

The IUPAC (International Union of Pure and Applied Chemistry) nomenclature has been used for the naming of the compounds in the examples that follow.

Example 1 (Compound 38)

2-(Methylamino)-2-oxoethyl 2-[4-(1-naphthyl)-1-piperazinyl]ethylcarbamate hydrochloride



1.1. 2-{2-[4-(1-Naphthyl)-1-piperazinyl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione

A suspension of 1.17 g (5.52 mmol) of 1-(1-naphthyl)piperazine (Tetrahedron Letters, 1998, 39(15), 2219-2222), 0.99 g (7.15 mol) of potassium carbonate and 1.68 g (6.62 mmol) of 2-(2-bromoethyl)-1*H*-isoindole-1,3(2*H*)-dione in 15 ml of *N,N*-dimethylformamide is heated at 90°C for 2 hours.

The mixture is allowed to cool to room temperature and is concentrated under reduced pressure. The residue is taken up in ethyl acetate and water, the aqueous phase is separated out and extracted three times with ethyl acetate, and the combined organic phases are washed with saturated aqueous sodium

chloride solution and dried over sodium sulfate. After evaporating off the solvent, the residue obtained is purified by chromatography on silica gel, eluting with a 40/60 mixture of ethyl acetate and cyclohexane.

5 1.14 g of pure product are thus obtained in the form of an oil.

1.2. 2-[4-(1-Naphthyl)-1-piperazinyl]ethanamine

0.17 ml (3.55 mmol) of hydrazine monohydrate
10 is added slowly at room temperature to a solution of 1.14 g (2.96 mmol) of 2-{2-[4-(1-naphthyl)-1-piperazinyl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione, prepared in step 1.1, in 15 ml of ethanol. The reaction mixture is then refluxed for 1 hour.

15 The mixture is allowed to cool to room temperature, the insoluble material is separated out by filtration and the filtrate is concentrated under reduced pressure. The residue is taken up in 20 ml of ether and stirred at room temperature for 20 minutes.
20 The insoluble material is again separated out and the filtrate is concentrated under reduced pressure. The residue obtained is purified by chromatography on silica gel, eluting with a 90/10/1 mixture of dichloromethane, methanol and 28% aqueous ammonia.

25 0.45 g of amine is thus obtained in the form of a colourless oil.

1.3. Ethyl [(phenyloxycarbonyl)oxy]acetate

32 ml (256 mmol) of phenyl chloroformate are added slowly at room temperature to a solution of 25 g (240 mmol) of ethyl glycolate and 55 ml (315 mmol) of diisopropylethylamine in 500 ml of toluene. Stirring is continued at room temperature for 2 hours.

The salt formed is separated out and the filtrate is concentrated under reduced pressure.

53.7 g of oily product are obtained, and are used without further purification in the following step.

1.4. Ethyl [(2,4-(1-naphthyl)-1-piperazinyl)ethyl]-amino)carbonyl]oxyacetate

A solution of 0.45 g (1.76 mmol) of 2-[4-(1-naphthyl)-1-piperazinyl]ethanamine, prepared in step 1.2, and 0.48 g (2.16 mmol) of ethyl [(phenyloxy-carbonyl)oxy]acetate, obtained in step 1.3, in 15 ml of toluene is heated at 50°C for 6 hours.

The mixture is allowed to cool to room temperature, the insoluble material is separated out by filtration and the filtrate is concentrated under reduced pressure. The residue is taken up in dichloromethane and water, the aqueous phase is separated out and extracted three times with dichloromethane, and the combined organic phases are washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. After evaporating off the solvent, the residue obtained is purified by chromatography on silica gel,

eluting with a 97/3 and then 93/7 mixture of dichloromethane and methanol. 0.49 g of pure product is thus obtained in the form of an oil.

5 1.5. 2-(Methylamino)-2-oxoethyl 2-[4-(1-naphthyl)-1-piperazinyl]ethylcarbamate

1.9 ml (3.75 mmol) of a solution (2M) of methylamine in tetrahydrofuran are added to a solution of 0.48 g (1.25 mmol) of ethyl [({2,4-(1-naphthyl)-1-
10 piperazinyl}ethyl}amino)carbonyl]oxyacetate, prepared in step 1.4, in 5 ml of methanol. Stirring is continued at room temperature for 1 hour.

After concentrating under reduced pressure, the residue obtained is purified by chromatography on
15 silica gel, eluting with a 97/3 mixture of dichloromethane and methanol. An oily residue is obtained, which is taken up with a solution of hydrochloric acid (5N) in isopropanol. The mixture is concentrated to dryness and the salt obtained is then recrystallized
20 from a 1/1 mixture of acetone and diisopropyl ether.

0.23 g of pure monohydrochloride product is thus obtained in the form of a white solid.


LC-MS: M+H = 371

m.p. (°C): 166°C

25 ¹H NMR (DMSO) δ (ppm): 2.55 (d, 3H); 3.20-3.75 (unresolved peak, 12H); 4.40 (s, 2H); 7.10 (d, 1H); 7.35-7.70 (unresolved peak, 4H); 7.80-8.05 (unresolved peak, 2H); 8.15 (dd, 1H); 10.90 (broad s, 1H).

Example 2 (Compound 157)**2-(Methylamino)-2-oxoethyl 3-[1-(1-isoquinolyl)-4-piperidyl]propylcarbamate hydrochloride**

5

**2.1. 2-[1-(1-Isoquinolyl)-4-piperidyl]ethanol**

4.17 g (25.50 mmol) of 1-chloroisoquinoline,
 10 3.62 g (28 mmol) of 2-(4-piperidyl)ethanol, 5.90 g
 (61.20 mmol) of sodium tert-butoxide and 0.476 g
 (0.765 mmol) of BINAP (2,2'-bis(diphenylphosphino)-
 1,1'-binaphthyl) suspended in 50 ml of toluene are
 introduced under an inert atmosphere. 0.233 g
 15 (0.255 mmol) of tris(dibenzylideneacetone)dipalladium
 is then added. The reaction mixture is then refluxed
 for 12 hours.

The salts are separated out by filtration
 through Celite and the filtrate is then concentrated
 20 under reduced pressure. The residue is taken up in
 ethyl acetate and water, the aqueous phase is separated
 out and extracted twice with ethyl acetate, and the
 combined organic phases are washed with saturated
 aqueous sodium chloride solution and dried over sodium
 25 sulfate. After evaporating off the solvent, the residue
 obtained is purified by chromatography on silica gel,

eluting with a 97/3 mixture of dichloromethane and methanol.

3.34 g of product are obtained in the form of a grey paste.

5

2.2. 1-[4-(2-Chloroethyl)-1-piperidyl]isoquinoline

2.20 ml (30.10 mmol) of thionyl chloride are added dropwise to a solution of 3.09 g (12.10 mmol) of 2-[1-(1-isoquinolyl)-4-piperidyl]ethanol, prepared in
10 step 2.1, in 30 ml of dichloromethane. The reaction mixture is then refluxed for 5 hours.

The mixture is concentrated to dryness under reduced pressure. The residue is taken up in 40 ml of dichloromethane and 30 ml of aqueous sodium hydroxide
15 solution (1M). The aqueous phase is separated out and extracted twice with dichloromethane, the combined organic phases are washed with saturated aqueous sodium chloride solution and dried over sodium sulfate, and the filtrate is concentrated under reduced pressure.
20 2.70 g of product are obtained in the form of a brown paste, which is used without further purification in the following step.

2.3. 3-[1-(1-Isoquinolyl)-4-piperidyl]propanenitrile

25 0.63 g (9.57 mmol) of potassium cyanide is added portionwise to a suspension of 2.63 g (9.57 mmol) of 1-[4-(2-chloroethyl)-1-piperidyl]isoquinoline, prepared in step 2.2, and 0.048 g (0.29 mmol) of

potassium iodide in 30 ml of dimethyl sulfoxide. The reaction mixture is maintained at about 120°C for 16 hours. The mixture is allowed to cool to room temperature and 90 ml of water and ethyl acetate are then added. The aqueous phase is separated out and extracted three times with ethyl acetate, the combined organic phases are washed with saturated aqueous sodium chloride solution and dried over sodium sulfate, and the filtrate is concentrated under reduced pressure.

10 The residue thus obtained is purified by chromatography on silica gel, eluting with a 30/70 mixture of ethyl acetate and cyclohexane.

0.77 g of product is obtained in the form of a yellow solid.

15 m.p. (°C): 141-143°C

2.4. 3-[1-(1-Isoquinolyl)-4-piperidyl]propanamine

A solution of 0.77 g (2.90 mmol) of 3-[1-(1-isoquinolyl)-4-piperidyl]propanenitrile, prepared in step 2.3, in 16 ml of tetrahydrofuran is added dropwise to a suspension of 0.22 g (5.80 mmol) of lithium aluminium hydride in 8 ml of tetrahydrofuran. The reaction mixture is then refluxed for 5 hours.

20

The reaction medium is cooled to about 0°C and 15 ml of aqueous sodium hydroxide solution (1M) are then added slowly. The mixture is stirred at room temperature for 30 minutes, followed by portionwise addition of wet sodium sulfate. The salts are separated

25

out by filtration through paper and the phases are then allowed to separate by settling. The aqueous phase is extracted three times with ethyl acetate, the combined organic phases are dried over sodium sulfate and the
5 filtrate is concentrated under reduced pressure. The residue thus obtained is purified by chromatography on silica gel, eluting with a 93/7/0.7 mixture of dichloromethane, methanol and 28% aqueous ammonia.

0.329 g of product is obtained in the form of
10 a yellow oil that crystallizes at room temperature.

2.5. Ethyl [{3-[1-(1-isoquinolyl)-4-piperidyl]propyl}-amino]carbonyloxyacetate

The process is performed as in Example 1
15 (step 1.4). Starting with 0.350 g (1.30 mmol) of 3-[1-(1-isoquinolyl)-4-piperidyl]propanamine, obtained in step 2.4, and 0.32 g (1.43 mmol) of ethyl [(phenyloxy-carbonyl)oxy]acetate, prepared in step 1.3 of Example 1, and after chromatography on silica gel, eluting with
20 a 20/80 and then 30/70 mixture of ethyl acetate and cyclohexane, 0.383 g of product is obtained in the form of a yellow paste.

2.6. 2-(Methylamino)-2-oxoethyl 3-[1-(1-isoquinolyl)-4-
25 piperidyl]propylcarbamate

The process is performed as described in Example 1 (step 1.5). Starting with 0.380 g (0.95 mmol) of ethyl [{3-[1-(1-isoquinolyl)-4-piperidyl]propyl}-

amino)carbonyl]oxyacetate, obtained in step 2.5, and 4.8 ml (9.51 mmol) of a solution (2M) of methylamine in tetrahydrofuran, and after chromatography on silica gel, eluting with a 98/2 and then 95/5 mixture of dichloromethane and methanol, 0.277 g of product is obtained in the form of an oil. This oily residue is then taken up in a solution of hydrochloric acid (5N) in isopropanol and the salt formed is filtered off and then washed with ethyl acetate.


After drying under vacuum at about 40°C, 0.204 g of hydrochloride is obtained in the form of an amorphous white solid.

LC-MS: M+H = 385

¹H NMR (DMSO+D₂O) δ (ppm): 1.30 (m, 2H); 1.40-1.80 (unresolved peak, 5H); 1.90 (broad d, 2H); 2.60 (s, 3H); 3.05 (m, 2H); 3.40 (t, 2H); 4.05 (broad d, 2H); 4.30 (s, 2H); 7.50 (d, 1H); 7.80 (m, 2H); 7.95 (t, 1H); 8.05 (d, 1H); 8.20 (d, 1H).

Example 3 (Compound 44)

2-Amino-2-oxoethyl 2-{1-[3-(trifluoromethyl)phenyl]-4-piperidyl}ethylcarbamate



3.1. 2-{1-[3-(Trifluoromethyl)phenyl]-4-piperidyl}-ethanol

The process is performed as described in Example 2 (step 2.1). Starting with 25.6 g (113.90 mmol) of 1-bromo-3-(trifluoromethyl)benzene, 17.66 g (136.60 mmol) of 2-(4-piperidyl)ethanol, 26.24 g (273 mmol) of sodium tert-butoxide, 2.12 g (3.41 mmol) of BINAP and 1.04 g (1.14 mmol) of tris-(dibenzylideneacetone)dipalladium, and after chromatography on silica gel, eluting with a 25/75 mixture of ethyl acetate and cyclohexane, 17.90 g of an orange-coloured oily residue are obtained. This residue is then taken up in 100 ml of methanol, a solution of 4.24 g of potassium hydroxide in 15 ml of methanol is then added and stirring is continued at room temperature for 1 hour. The mixture is concentrated under reduced pressure and the residue is then taken up in chloroform and aqueous hydrochloric acid solution (1N). The organic phase is separated out and dried over sodium sulfate, and the filtrate is concentrated under reduced pressure. 14 g of product are obtained in the form of a dark yellow oil, which is used without further purification in the following step.

3.2. 2-{1-[3-(Trifluoromethyl)phenyl]-4-piperidyl}ethyl methanesulfonate

A solution of 1 g (8.78 mmol) of mesyl chloride in 5 ml of dichloromethane is added dropwise

under an inert atmosphere to a solution of 2 g
(7.32 mmol) of 2-{1-[3-(trifluoromethyl)phenyl]-4-
piperidyl}ethanol, obtained in step 3.1, and 1.53 ml
(10.98 mmol) of triethylamine in 40 ml of dichloro-
5 methane, cooled to about 0°C. Stirring is continued at
0°C for 1 hour and then at room temperature for
2 hours.

Water is added to the reaction medium, the
aqueous phase is separated out and extracted three
10 times with dichloromethane, the combined organic phases
are washed with saturated aqueous sodium chloride
solution and dried over sodium sulfate, and the
filtrate is concentrated under reduced pressure.

2.5 g of product are thus obtained in the
15 form of an oil, which is used without further
purification in the following step.

3.3. 3-(2-{1-[3-(Trifluoromethyl)phenyl]-4-piperidyl}- ethyl)-1,3-oxazolidine-2,4-dione

20 A solution of 2.3 g (6.545 mmol) of 2-{1-[3-
(trifluoromethyl)phenyl]-4-piperidyl}ethyl methane-
sulfonate, prepared in step 3.2, 0.694 g (6.87 mmol) of
1,3-oxazolidine-2,4-dione (J. Med. Chem., 1991, 34,
1538-1544) and 1.5 g (13.09 mmol) of 1,1,3,3-tetra-
25 methylguanidine in 30 ml of tetrahydrofuran is refluxed
for 12 hours under an inert atmosphere.

The mixture is concentrated under reduced
pressure. The residue is taken up in dichloromethane

and water, the aqueous phase is separated out and extracted twice with dichloromethane, and the combined organic phases are washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. After
5 evaporating off the solvent, the residue obtained is purified by chromatography on silica gel, eluting with a 20/80 and then 40/60 mixture of ethyl acetate and cyclohexane.

1.61 g of pure product are obtained in the
10 form of an oil.

3.4. 2-Amino-2-oxoethyl 2-{1-[3-(trifluoromethyl)-phenyl]-4-piperidyl}ethylcarbamate

9.3 ml (64.82 mmol) of a solution of aqueous
15 ammonia (7N) in methanol are added to a solution of 0.77 g (2.16 mmol) of 3-(2-{1-[3-(trifluoromethyl)-phenyl]-4-piperidyl}ethyl)-1,3-oxazolidine-2,4-dione, obtained in step 3.3, in 10 ml of a 1/1 mixture of methanol and tetrahydrofuran. Stirring is continued at
20 room temperature for 24 hours.

After concentrating under reduced pressure, the residue obtained is purified by chromatography on silica gel, eluting with a 97/3 and then 95/5 mixture of dichloromethane and methanol, followed by
25 recrystallization from a mixture of ethyl acetate and diisopropyl ether.

0.370 g of pure product is obtained in the form of a white solid.


LC-MS: M+H = 374

m.p. (°C): 140-142°C

¹H NMR (CDCl₃) δ (ppm): 1.30-1.55 (unresolved peak, 5H);
 1.90 (broad d, 2H); 2.80 (t, 2H); 3.35 (q, 2H); 3.80
 5 (broad d, 2H); 4.60 (s, 2H); 4.90 (broad s, 1H); 5.55
 (broad s, 1H); 6.05 (broad s, 1H); 7.10 (m, 3H); 7.35
 (t, 1H).

Example 4 (Compound 47)

10 **2-(Methylamino)-2-oxoethyl 2-[1-(6-methyl-2-pyridyl)-4-piperidyl]ethylcarbamate hydrochloride**



15 4.1. 2-[1-(6-Methyl-2-pyridyl)-4-piperidyl]ethanol

1 g (7.74 mmol) of 2-(4-piperidyl)ethanol and
 0.987 g (7.74 mmol) of 2-chloro-6-methylpyridine are
 introduced into an autoclave. The mixture is then
 heated at 130°C for 17 hours.

20 The reaction mixture is allowed to cool to
 room temperature and is then taken up in chloroform and
 saturated aqueous sodium hydrogen carbonate solution.
 The aqueous phase is separated out and extracted twice
 with chloroform, the combined organic phases are washed
 25 with saturated aqueous sodium chloride solution and

dried over sodium sulfate, and the filtrate is concentrated under reduced pressure.

1.21 g of product are thus obtained in the form of an orange-coloured liquid, which is used without further purification in the following step.

4.2. 2-[1-(6-Methyl-2-pyridyl)-4-piperidyl]ethyl methanesulfonate

The process is performed as in Example 3 (step 3.2). Starting with 0.661 g (3 mmol) of 2-[1-(6-methyl-2-pyridyl)-4-piperidyl]ethanol, obtained in step 4.1, 0.378 g (3.30 mmol) of mesyl chloride and 0.63 ml (4.50 mmol) of triethylamine, 0.779 g of product is obtained in the form of an orange-coloured oil, which is used without further purification in the following step.

4.3. 3-{2-[1-(6-Methyl-2-pyridyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione

The process is performed according to the method described in Example 3 (step 3.3). Starting with 0.776 g (2.60 mmol) of 2-[1-(6-methyl-2-pyridyl)-4-piperidyl]ethyl methanesulfonate, obtained in step 4.2, 0.315 g (3.12 mmol) of 1,3-oxazolidine-2,4-dione and 0.65 ml (5.20 mmol) of 1,1,3,3-tetramethylguanidine, and after chromatography on silica gel, eluting with a 30/70 mixture of ethyl acetate and cyclohexane, 0.76 g

of pure product is obtained in the form of a yellow oil.

4.4. 2-(Methylamino)-2-oxoethyl 2-[1-(6-methyl-2-pyridyl)-4-piperidyl]ethylcarbamate

The process is performed according to the procedure described in Example 1 (step 1.5). Starting with 0.841 g (2.77 mmol) of 3-{2-[1-(6-methyl-2-pyridyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, obtained in step 4.3, and 6.9 ml (13.86 mmol) of a solution (2M) of methylamine in tetrahydrofuran, 0.598 g of pure product is obtained in the form of an oil, after chromatography on silica gel, eluting with a 95/5 mixture of dichloromethane and methanol. This oily residue is then taken up with a solution of hydrochloric acid (5N) in isopropanol and the salt formed is filtered off and then washed successively with acetone and then with ether.

After drying under vacuum at about 80°C, 0.492 g of hydrochloride is obtained in the form of a white powder.


LC-MS: M+H = 335

m.p. (°C): 95-100°C

¹H NMR (DMSO+D₂O) δ (ppm): 0.95-1.45 (unresolved peak, 4H); 1.60 (m, 1H); 1.80 (broad d, 2H); 2.40 (s, 3H); 2.60 (s, 3H); 2.90-3.20 (unresolved peak, 4H); 4.10 (broad d, 2H); 4.30 (s, 2H); 6.70 (d, 1H); 7.10 (d, 1H); 7.75 (dd, 1H).

Example 5 (Compound 154)

2-(Methylamino)-2-oxoethyl [1-(1-isoquinolyl)-4-piperidyl]methylcarbamate hydrochloride



5

5.1. [1-(1-Isoquinolyl)-4-piperidyl]methanol

The process is performed as described in Example 2 (step 2.1). Starting with 2.50 g (15.28 mmol) of 1-chloroisoquinoline, 1.94 g (136.6 mmol) of 4-piperidylmethanol, 3.53 g (36.67 mmol) of sodium tert-butoxide, 0.285 g (0.46 mmol) of BINAP and 0.140 g (0.15 mmol) of tris(dibenzylideneacetone)dipalladium, and after chromatography on silica gel, eluting with a 98/2/0.2 and then 95/5/0.5 mixture of dichloromethane, methanol and 28% aqueous ammonia, 2.40 g of pure product are obtained in the form of a viscous orange oil.

5.2. 3-{[1-(1-Isoquinolyl)-4-piperidyl]methyl}-1,3-oxazolidine-2,4-dione

A solution of 2.01 g (9.95 mmol) of diisopropyl azodicarboxylate (DIAD) in 5 ml of tetrahydrofuran is added dropwise, under an inert atmosphere, to a solution of 2.4 g (9.95 mmol) of [1-(1-isoquinolyl)-

25

4-piperidyl]methanol, prepared in step 5.1, 2.87 g
(10.94 mmol) of triphenylphosphine and 1.21 g
(11.93 mmol) of 1,3-oxazolidine-2,4-dione in 40 ml of
tetrahydrofuran, cooled to about -10°C , the temperature
5 of the reaction medium being maintained throughout
between -10°C and 0°C . Stirring is then continued at
 0°C for 1 hour and then at 25°C for 18 hours.

The mixture is concentrated under reduced
pressure and the residue is taken up in dichloromethane
10 and 10 ml of aqueous 5% sodium hydroxide solution. The
aqueous phase is separated out and then extracted twice
with dichloromethane. The organic phases are combined
and washed successively with aqueous hydrochloric acid
solution (1N) and then with saturated aqueous sodium
15 hydrogen carbonate solution and with saturated aqueous
sodium chloride solution. The organic phase is dried
over sodium sulfate and the filtrate is concentrated
under reduced pressure. The residue thus obtained is
purified by chromatography on silica gel, eluting with
20 a 99/1/0.1 and then 98/2/0.2 mixture of dichloro-
methane, methanol and 28% aqueous ammonia.

3.57 g of oxazolidinedione are thus obtained
in the form of an orange paste.

25 5.3. 2-(Methylamino)-2-oxoethyl [1-(1-isoquinolyl)-4-
piperidyl]methylcarbamate

The process is performed according to the
procedure described in Example 1 (step 1.5). Starting

with 3.57 g (10.97 mmol) of 3-([1-(1-isoquinolyl)-4-piperidyl]methyl)-1,3-oxazolidine-2,4-dione, obtained in step 5.2, and 27 ml (54.86 mmol) of a solution (2M) of methylamine in tetrahydrofuran, and after
5 chromatography on silica gel, eluting with a 95/5/0.5 mixture of dichloromethane, methanol and 28% aqueous ammonia, 0.90 g of pure product is obtained in the form of a yellow paste. This residue is then taken up in a solution of hydrochloric acid (5N) in isopropanol, and
10 the salt formed is filtered off and then washed with acetone.

After drying under vacuum at about 80°C, 0.728 g of hydrochloride is obtained in the form of a pale yellow solid.


15 LC-MS: M+H = 357

m.p. (°C): 208-212°C (decomposition)

¹H NMR (D₂O) δ (ppm): 1.55 (m, 2H); 1.95 (m, 3H); 2.70 (s, 3H); 3.20 (broad d, 2H); 3.45 (t, 2H); 4.10 (broad d, 2H); 4.50 (s, 2H); 7.35 (d, 1H); 7.55 (d, 1H); 7.70
20 (m, 1H); 7.90 (d, 2H); 8.20 (d, 1H).

Example 6 (Compound 158)

2-Amino-2-oxoethyl 2-[1-(6-fluoro-1-isoquinolyl)-4-piperidyl]ethylcarbamate



6.1. 2-[1-(6-Fluoro-1-isoquinolyl)-4-piperidyl]ethanol

The process is performed as described in
 5 Example 4 (step 4.1). Starting with 1.52 g (8.39 mmol)
 of 1-chloro-6-fluoroisoquinoline and 1.20 g (9.23 mmol)
 of 2-(4-piperidyl)ethanol, and after chromatography on
 silica gel, eluting with a 95/5/0.5 mixture of
 dichloromethane, methanol and 28% aqueous ammonia,
 10 0.90 g of pure product is obtained in the form of a
 yellow paste.

6.2. 2-[1-(6-Fluoro-1-isoquinolyl)-4-piperidyl]ethyl methanesulfonate

15 The process is performed as described in
 Example 3 (step 3.2). Starting with 1.47 g (5.36 mmol)
 of 2-[1-(6-fluoro-1-isoquinolyl)-4-piperidyl]ethanol,
 obtained in step 6.1, 0.675 g (5.89 mmol) of mesyl
 chloride and 1.13 ml (8.04 mmol) of triethylamine,
 20 1.80 g of product are obtained in the form of an oil,
 which is used without further purification in the
 following step.

6.3. 3-{2-[1-(6-Fluoro-1-isoquinolyl)-4-piperidyl]- 25 ethyl}-1,3-oxazolidine-2,4-dione

The process is performed according to the method described in Example 3 (step 3.3). Starting with 1.8 g (5.10 mmol) of 2-[1-(6-fluoro-1-isoquinolyl)-4-piperidyl]ethyl methanesulfonate, obtained in step 6.2, 5 0.62 g (6.13 mmol) of 1,3-oxazolidine-2,4-dione and 1.30 ml (10.21 mmol) of 1,1,3,3-tetramethylguanidine, and after chromatography on silica gel, eluting with a 40/60 mixture of ethyl acetate and cyclohexane, 1.34 g of pure product are obtained in the form of an 10 amorphous white solid.

6.4. 2-Amino-2-oxoethyl 2-[1-(6-fluoro-1-isoquinolyl)-4-piperidyl]ethylcarbamate

The process is performed according to the 15 procedure described in Example 3 (step 3.4). Starting with 0.597 g (1.67 mmol) of 3-{2-[1-(6-fluoro-1-isoquinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, obtained in Step 6.3, and 14.30 ml (100.20 mmol) of a solution (7M) of aqueous ammonia in methanol, and after 20 chromatography on silica gel, eluting with a 95/5 mixture of dichloromethane and methanol, followed by recrystallization from diisopropyl ether, 0.168 g of pure product is obtained in the form of a white solid.

LC-MS: M+H = 375


25 m.p. (°C): 135-139°C

¹H NMR (DMSO) δ (ppm): 1.20-1.70 (unresolved peak, 5H); 1.80 (broad d, 2H); 2.85 (t, 2H); 3.10 (broad d, 2H); 3.70 (broad d, 2H); 4.30 (s, 2H); 6.95-7.20 (unresolved

peak, 3H); 7.30 (d, 1H); 7.45 (td, 1H); 7.60 (dd, 1H);
8.10 (m, 2H).

Example 7 (Compound 172)

5 **2-(Methylamino)-2-oxoethyl 2-[1-(4-isoquinolyl)-4-piperidyl]ethylcarbamate hydrochloride**



10 7.1. 2-[1-(4-Isoquinolyl)-4-piperidyl]ethanol

The process is performed according to the protocol described in Example 2 (step 2.1). Starting with 1 g (4.81 mmol) of 4-bromoisoquinoline, 0.683 g (5.29 mmol) of 2-(4-piperidyl)ethanol, 1.11 g (11.50 mmol) of sodium tert-butoxide, 0.090 g (0.144 mmol) of BINAP and 0.044 g (0.048 mmol) of tris(dibenzylideneacetone)dipalladium, and after chromatography on silica gel, eluting with a 97/3 and then 95/5 mixture of dichloromethane and methanol,
15 0.810 g of product is obtained in the form of a viscous green liquid.

7.2. 3-{2-[1-(4-Isoquinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione

The procedure described in Example 5 (step 5.2) is used. Starting with 0.801 g (3.12 mmol) of 2-[1-(4-isoquinolyl)-4-piperidyl]ethanol, prepared in step 7.1, 0.902 g (3.44 mmol) of triphenylphosphine, 5 0.379 g (3.75 mmol) of 1,3-oxazolidine-2,4-dione and 0.632 g (3.12 mmol) of diisopropyl azodicarboxylate (DIAD), and after chromatography on silica gel, eluting with a 98/2 mixture of dichloromethane and methanol, 1 g of product is obtained in the form of a green
10 paste.

7.3. 2-(Methylamino)-2-oxoethyl 2-[1-(4-isoquinolyl)-4-piperidyl]ethylcarbamate

The procedure described in Example 1 (step 15 1.5) is used. Starting with 1 g (2.95 mmol) of 3-{2-[1-(4-isoquinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, obtained in step 7.2, and 7.40 ml (14.73 mmol) of a solution (2M) of methylamine in tetrahydrofuran, and after chromatography on silica gel, eluting with a
20 95/5 mixture of dichloromethane and methanol, 0.410 g of pure product is obtained in the form of an amorphous white solid. This oily residue is then taken up in a solution of hydrochloric acid (5N) in isopropanol and the salt formed is filtered off and then washed
25 successively with acetone and then with diisopropyl ether.

After drying under vacuum at about 90°C,
0.359 g of hydrochloride is obtained in the form of a
yellow solid.

LC-MS: M+H = 371


5 m.p. (°C): 205-210°C

¹H NMR (DMSO) δ (ppm): 1.30-1.70 (unresolved peak, 5H);
1.90 (broad d, 2H); 2.60 (d, 3H); 2.90 (t, 2H); 3.15
(m, 2H); 3.50 (broad d, 2H); 4.35 (s, 2H); 7.25 (broad
t, 1H); 7.80 (broad s, 1H); 8.0 (dd, 1H); 8.05-8.30
10 (unresolved peak, 3H); 8.45 (d, 1H); 9.45 (s, 1H).

Example 8 (Compound 126)

**2-(Methylamino)-2-oxoethyl 2-[1-(2-quinolyl)-4-
piperidyl]ethylcarbamate**

15



8.1. 2-[1-(2-Quinolyl)-4-piperidyl]ethanol

The process is performed as in Example 4
20 (step 4.1). Starting with 2 g (12.20 mmol) of 2-chloro-
quinoline and 1.58 g (12.20 mmol) of 2-(4-piperidyl)-
ethanol, and after chromatography on silica gel,
eluting with a 98/2/0.2 and then 95/5/0.5 mixture of
dichloromethane, methanol and 28% aqueous ammonia,

2.36 g of pure product are obtained in the form of a pale yellow oil that crystallizes at room temperature.

8.2. 3{2-[1-(2-Quinolyl)-4-piperidyl]ethyl-1,3-oxazolidine-2,4-dione

The process is performed according to the procedure described in Example 5 (step 5.2). Starting with 2.22 g (8.65 mmol) of 2-[1-(2-quinolyl)-4-piperidyl]ethanol, prepared in step 8.1, 2.50 g (9.52 mmol) of triphenylphosphine, 1.05 g (10.38 mmol) of 1,3-oxazolidine-2,4-dione and 1.75 g (8.65 mmol) of diisopropyl azodicarboxylate (DIAD), and after chromatography on silica gel, eluting with a 30/70 and then 40/60 mixture of ethyl acetate and cyclohexane, 2.63 g of product are obtained in the form of an amorphous white solid.

8.3. 2-(Methylamino)-2-oxoethyl 2-[1-(2-quinolyl)-4-piperidyl]ethylcarbamate

The process is performed according to the procedure described in Example 1 (step 1.5). Starting with 1.5 g (4.42 mmol) of 3{2-[1-(2-quinolyl)-4-piperidyl]ethyl-1,3-oxazolidine-2,4-dione, obtained in step 8.2, and 11 ml (22.10 mmol) of a solution (2M) of methylamine in tetrahydrofuran, and after chromatography on silica gel, eluting with a 98/2/0.2 and then 95/5/0.5 mixture of dichloromethane, methanol and 28% aqueous ammonia, followed by crystallization

from ethyl acetate, 0.405 g of pure product is obtained in the form of a white solid.

LC-MS: M+H = 371


m.p. (°C): 125-128°C

5 ¹H NMR (CDCl₃) δ (ppm): 1.20-1.60 (unresolved peak, 5H);
1.85 (broad d, 2H); 2.85 (d, 3H); 3.0 (broad t, 2H);
3.30 (broad q, 2H); 4.55 (broad d, 2H); 4.60 (s, 2H);
4.85 (broad s, 1H); 6.10 (broad s, 1H); 7.0 (d, 1H);
7.20 (t, 1H); 7.55 (m, 2H); 7.70 (d, 1H); 7.90 (d, 1H).

10

Example 9 (Compound 127)

2-Amino-2-oxoethyl 2-[1-(2-quinolyl)-4-piperidyl]-ethylcarbamate



15

The process is performed according to the procedure described in Example 3 (step 3.4). Starting with 1.14 g (3.36 mmol) of 3{2-[1-(2-quinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, described
20 in Example 8 (step 8.2), and 9.60 ml (67.20 mmol) of a solution (7M) of aqueous ammonia in methanol, and after chromatography on silica gel, eluting with a 95/5 mixture of dichloromethane and methanol, followed by

recrystallization from ethyl acetate, 0.360 g of pure product is obtained in the form of a white solid.

LC-MS: M+H = 357


m.p. (°C): 135-137°C

5 ¹H NMR (CDCl₃) δ (ppm): 1.15-1.70 (unresolved peak, 5H);
1.85 (broad d, 2H); 2.95 (t, 2H); 3.35 (q, 2H); 4.55
(broad d, 2H); 4.60 (s, 2H); 4.85 (broad s, 1H); 5.55
(broad s, 1H); 6.05 (broad s, 1H); 7.0 (d, 1H); 7.20
(t, 1H); 7.55 (t, 1H); 7.60 (d, 1H); 7.70 (d, 1H); 7.90
10 (d, 1H).

Example 10 (Compound 137)

2-Amino-2-oxoethyl 2-[1-(6-chloro-2-quinolyl)-4-piperidyl]ethylcarbamate

15



10.1. 2-[1-(6-Chloro-2-quinolyl)-4-piperidyl]ethanol

The process is performed as described in
20 Example 4 (step 4.1). Starting with 2 g (10.10 mmol) of
2,6-dichloroquinoline and 1.44 g (11.10 mmol) of 2-(4-
piperidyl)ethanol, and after chromatography on silica
gel, eluting with a 98/2 mixture of dichloromethane and
methanol, 2.54 g of pure product are obtained in the
25 form of a white solid.

10.2. 2-[1-(6-Chloro-2-quinolyl)-4-piperidyl]ethyl
methanesulfonate

The process is performed as described in
Example 3 (step 3.2). Starting with 2.49 g (8.56 mmol)
5 of 2-[1-(6-chloro-2-quinolyl)-4-piperidyl]ethanol,
obtained in step 10.1, 1.08 g (9.42 mmol) of mesyl
chloride and 1.81 ml (12.84 mmol) of triethylamine,
3.10 g of product are obtained in the form of an oil,
which is used without further purification in the
10 following step.

10.3. 3-{2-[1-(6-Chloro-2-quinolyl)-4-piperidyl]ethyl}-
1,3-oxazolidine-2,4-dione

The process is performed according to the
15 method described in Example 3 (step 3.3). Starting with
3 g (8.13 mmol) of 2-[1-(6-chloro-2-quinolyl)-4-
piperidyl]ethyl methanesulfonate, obtained in step
10.2, 1.09 g (10.8 mmol) of 1,3-oxazolidine-2,4-dione
and 2.30 ml (18 mmol) of 1,1,3,3-tetramethylguanidine,
20 and after chromatography on silica gel, eluting with a
98/2 mixture of dichloromethane and methanol, 2.57 g of
product are obtained.

10.4. 2-Amino-2-oxoethyl 2-[1-(6-chloro-2-quinolyl)-4-
25 piperidyl]ethylcarbamate

The process is performed according to the
procedure described in Example 3 (step 3.4). Starting
with 1.28 g (3.42 mmol) of 3-{2-[1-(6-chloro-2-

quinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, obtained in step 10.3, and 22.10 ml (154.08 mmol) of a solution (7M) of aqueous ammonia in methanol, and after crystallization from ethanol, 0.64 g of pure product is
 5 obtained in the form of a white solid.


LC-MS: M+H = 391

m.p. (°C): 189-191°C

¹H NMR (DMSO) δ (ppm): 1.10 (m, 2H); 1.40 (m, 2H); 1.60 (m, 1H); 1.80 (broad d, 2H); 2.90 (broad t, 2H); 3.05
 10 (m, 2H); 4.30 (s, 2H); 4.50 (broad d, 2H); 7.15 (m, 3H); 7.25 (d, 1H); 7.50 (m, 2H); 7.75 (d, 1H); 7.95 (d, 1H).

Example 11 (Compound 166)

15 **2-(Methylamino)-2-oxoethyl 2-[1-(3-isoquinolyl)-4-piperidyl]ethylcarbamate**



20 11.1. 2-[1-(3-Isoquinolyl)-4-piperidyl]ethanol

The process is performed as described in Example 4 (step 4.1). Starting with 1 g (6.11 mmol) of 3-chloroisoquinoline and 0.869 g (6.72 mmol) of 2-(4-piperidyl)ethanol, and after chromatography on silica
 25 gel, eluting with a 98/2 mixture of dichloromethane and

methanol, 0.34 g of pure product is obtained in the form of an oil.

11.2. 2-[1-(3-Isoquinolyl)-4-piperidyl]ethyl methane-
5 sulfonate

The process is performed as described in Example 3 (step 3.2). Starting with 0.34 g (1.33 mmol) of 2-[1-(3-isoquinolyl)-4-piperidyl]ethanol, obtained in step 11.1, 0.18 g (1.59 mmol) of mesyl chloride and
10 0.30 ml (1.99 mmol) of triethylamine, 0.44 g of product is obtained in the form of an oil, which is used without further purification in the following step.

11.3. 3-{2-[1-(3-Isoquinolyl)-4-piperidyl]ethyl}-1,3-
15 oxazolidine-2,4-dione

The process is performed according to the method described in Example 3 (step 3.3). Starting with 0.44 g (1.32 mmol) of 2-[1-(3-isoquinolyl)-4-piperidyl]ethyl methanesulfonate, obtained in step
20 11.2, 0.16 g (1.58 mmol) of 1,3-oxazolidine-2,4-dione and 0.30 g (2.63 mmol) of 1,1,3,3-tetramethylguanidine, and after chromatography on silica gel, eluting with a 98/2 mixture of dichloromethane and methanol, 0.25 g of product is obtained.

25

11.4. 2-(Methylamino)-2-oxoethyl 2-[1-(3-isoquinolyl)-
4-piperidyl]ethylcarbamate

The process is performed according to the procedure described in Example 1 (step 1.5). Starting with 0.24 g (0.71 mmol) of 3-{2-[1-(3-isoquinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, obtained in
 5 step 11.3, and 1.8 ml (3.53 mmol) of a solution (2M) of methylamine in tetrahydrofuran, and after chromatography on silica gel, eluting with a 98/2 and then 96/4 mixture of dichloromethane and methanol, followed by crystallization from diisopropyl ether,
 10 0.16 g of pure product is obtained in the form of a white solid.


LC-MS: M+H = 371

m.p. (°C): 156-158°C

¹H NMR (CDCl₃) δ (ppm): 1.20-1.70 (unresolved peak, 5H);
 15 1.85 (d, 2H); 2.90 (m, 5H); 3.30 (q, 2H); 4.40 (d, 2H);
 4.60 (s, 2H); 4.85 (broad s, 1H); 6.10 (broad s, 1H);
 6.80 (s, 1H); 7.30 (m, 1H); 7.60 (m, 2H); 7.80 (d, 1H);
 8.95 (s, 1H).

20 **Example 12 (Compound 128)**

2-(Methylamino)-2-oxoethyl 2-[4-fluoro-1-(2-quinolyl)-4-piperidyl]ethylcarbamate



12.1. *tert*-Butyl 4-hydroxy-4-(2-hydroxyethyl)-1-piperidinecarboxylate

A solution of 31.20 g (108.50 mmol) of *tert*-butyl 4-(2-ethoxy-2-oxoethyl)-4-hydroxy-1-piperidinecarboxylate (WO 02/16352) in 150 ml of tetrahydrofuran is added dropwise to a suspension of 4.12 g (108.50 mmol) of lithium aluminium hydride in 150 ml of tetrahydrofuran. The mixture is stirred at room temperature for 2 hours and is then worked up as described in Example 2 (step 2.4).

26 g of product are obtained in the form of a yellow oil, which is used without further purification in the following step.

12.2. *tert*-Butyl 4-(2-([*tert*-butyldiphenylsilyl]oxy)ethyl)-4-hydroxy-1-piperidinecarboxylate

A solution of 21.50 ml (82.50 mmol) of *tert*-butyldiphenylsilyl chloride in 15 ml of dichloromethane is added dropwise, under an inert atmosphere, to a solution of 18.4 g (75 mmol) of *tert*-butyl 4-hydroxy-4-(2-hydroxyethyl)-1-piperidinecarboxylate, obtained in step 12.1, and 11.60 ml (82.50 mmol) of triethylamine in 100 ml of dichloromethane, cooled to about 0°C. The mixture is allowed to warm to room temperature and stirring is then continued for 12 hours. Saturated aqueous ammonium chloride solution is added to the reaction medium. The aqueous phase is separated out and extracted twice with dichloromethane, the combined

organic phases are washed with saturated aqueous sodium chloride solution and dried over sodium sulfate, and the filtrate is concentrated under reduced pressure. The residue thus obtained is purified by chromatography
5 on silica gel, eluting with a 10/90 and then 20/80 mixture of ethyl acetate and cyclohexane.

33.48 g of product are thus obtained in the form of a yellow oil.

10 12.3. *tert*-Butyl 4-(2-([*tert*-butyl(diphenyl)silyl]oxy)-ethyl)-4-fluoro-1-piperidinecarboxylate

A solution of 1.70 ml (13.40 mmol) of 1,1'-[(trifluoro- λ^4 -sulfanyl)imino]diethane (DAST) in 10 ml of dichloromethane is added dropwise, under an inert
15 atmosphere, to a solution of 5 g (10.34 mmol) of *tert*-butyl 4-(2-([*tert*-butyl(diphenyl)silyl]oxy)ethyl)-4-hydroxy-1-piperidinecarboxylate, obtained in step 12.2, in 100 ml of dichloromethane, cooled to about 0°C. The mixture is allowed to warm to room temperature and
20 stirring is then continued for 12 hours. Saturated aqueous sodium hydrogen carbonate solution is added to the reaction medium. The aqueous phase is separated out and extracted three times with dichloromethane, and the combined organic phases are washed with saturated
25 aqueous sodium chloride solution and dried over sodium sulfate. The filtrate is concentrated under reduced pressure and the residue thus obtained is then purified by chromatography on silica gel, eluting with a 10/90

mixture of ethyl acetate and cyclohexane. 4.65 g of product are thus obtained in the form of an orange-coloured oil. 0.60 ml of a solution of osmium tetroxide (2.5%) in *tert*-butanol is added at room temperature to
5 a solution of 4.50 g of this oily residue and 1.25 g (10.7 mmol) of *N*-methyldmorpholine oxide (NMO) in a mixture of 8 ml of acetone and 6 ml of water. Stirring is continued for 21 hours. The residue is taken up in ethyl acetate and water, the aqueous phase is separated
10 out and extracted twice with ethyl acetate, and the combined organic phases are washed with saturated aqueous sodium chloride solution and dried over sodium sulfate. After evaporating off the solvent, the residue obtained is purified by chromatography on silica gel,
15 eluting with a 6/94 mixture of ethyl acetate and cyclohexane.

3.40 g of product are thus obtained in the form of a pale yellow oil.

20 12.4. 4-(2-([*tert*-butyl(diphenyl)silyl]oxy)ethyl)-4-fluoropiperidine

2.80 ml (37.06 mmol) of trifluoroacetic acid are added slowly to a solution of 3 g (6.17 mmol) of *tert*-butyl 4-(2-([*tert*-butyldiphenylsilyl]oxy)ethyl)-4-
25 fluoro-1-piperidinecarboxylate, obtained in step 12.3, in 20 ml of dichloromethane. Stirring is continued at room temperature for 5 hours. The reaction mixture is poured into a mixture of ice-water and 28% aqueous

ammonia. The phases are separated by settling, the aqueous phase is extracted twice with dichloromethane and the combined organic phases are washed with saturated aqueous sodium chloride solution, dried over sodium sulfate and concentrated under reduced pressure.

2.30 g of product are obtained in the form of a yellow oil, which is used without further purification in the following step.

10 12.5. 2-[4-(2-([*tert*-butyldiphenylsilyl]oxy)ethyl)-4-fluoro-1-piperidyl]quinoline

The process is performed as described in Example 2 (step 2.1). Starting with 1.18 g (5.68 mmol) of 2-bromoquinoline (Eur. J. Org. Chem. 2002, 4181-4184), 2.30 g (5.98 mmol) of 4-(2-([*tert*-butyl-(diphenyl)silyl]oxy)ethyl)-4-fluoropiperidine, obtained in step 12.4, 0.66 g (6.81 mmol) of sodium *tert*-butoxide, 0.149 g (0.239 mmol) of BINAP and 0.074 g (0.081 mmol) of tris(dibenzylideneacetone)dipalladium, and after chromatography on silica gel, eluting with a 10/90 mixture of ethyl acetate and cyclohexane, 2.15 g of pure product are obtained in the form of an orange-coloured oil.

25 12.6. 2-[4-Fluoro-1-(2-quinolyl)-4-piperidyl]ethanol

0.40 g (1.26 mmol) of *n*-tetrabutylammonium fluoride trihydrate is added to a solution of 2.15 g (4.19 mmol) of 2-[4-(2-([*tert*-butyldiphenylsilyl]-

oxy)ethyl)-4-fluoro-1-piperidyl]quinoline, obtained in step 12.5, in 20 ml of tetrahydrofuran. Stirring is continued at room temperature for 4 hours. The mixture is concentrated to dryness and the residue obtained is then purified by chromatography on silica gel, eluting with a 35/65 and then 40/60 mixture of ethyl acetate and cyclohexane.

0.61 g of product is obtained in the form of an orange-coloured oil.

10

12.7. 2-[4-Fluoro-1-(2-quinolyl)-4-piperidyl]ethyl methanesulfonate

The process is performed as described in Example 3 (step 3.2). Starting with 0.61 g (2.22 mmol) of 2-[4-fluoro-1-(2-quinolyl)-4-piperidyl]ethanol, obtained in step 12.6, 0.280 g (2.45 mmol) of mesyl chloride and 0.35 ml (2.45 mmol) of triethylamine, 0.80 g of product is obtained in the form of an orange-coloured oil, which is used without further purification in the following step.

20

12.8. 3-{2-[4-Fluoro-1-(2-quinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione

The process is performed according to the method described in Example 3 (step 3.3). Starting with 0.780 g (2.22 mmol) of 2-[4-fluoro-1-(2-quinolyl)-4-piperidyl]ethyl methanesulfonate, obtained in step 12.7, 0.27 g (2.66 mmol) of 1,3-oxazolidine-2,4-dione

25

and 0.51 g (4.43 mmol) of 1,1,3,3-tetramethylguanidine,
and after chromatography on silica gel, eluting with a
99/1 mixture of dichloromethane and methanol, 0.520 g
of pure product is obtained in the form of a beige-
5 coloured solid.

12.9. 2-(Methylamino)-2-oxoethyl 2-[4-fluoro-1-(2-
quinolyl)-4-piperidyl]ethylcarbamate

The process is performed according to the
10 procedure described in Example 1 (step 1.5). Starting
with 0.52 g (1.46 mmol) of 3-{2-[4-fluoro-1-(2-
quinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione,
obtained in step 12.8, and 3.6 ml (7.28 mmol) of a
solution (2M) of methylamine in tetrahydrofuran, and
15 after chromatography on silica gel, eluting with a 99/1
mixture of ethyl acetate and methanol, followed by
crystallization from diethyl ether, 0.390 g of pure
product is obtained in the form of a white solid.


LC-MS: M+H = 389

20 m.p. (°C): 147-149°C

¹H NMR (CDCl₃) δ (ppm): 1.70-2.10 (unresolved peak, 6H);
2.90 (d, 3H); 3.40 (broad t, 2H); 3.50 (q, 2H); 4.40
(broad d, 2H); 4.60 (s, 2H); 5.15 (broad s, 1H); 6.15
(broad s, 1H); 7.05 (d, 1H); 7.25 (t, 1H); 7.55 (t,
25 1H); 7.65 (d, 1H); 7.75 (d, 1H); 7.95 (d, 1H).

Example 13 (Compound 49)

2-(Methylamino)-2-oxoethyl 2-[1-(6-isobutyl-2-pyridyl)-4-piperidyl]ethylcarbamate



5

13.1. 2-[1-(6-Bromo-2-pyridyl)-4-piperidyl]ethanol

The process is performed as described in Example 4 (step 4.1). Starting with 30.20 g (127 mmol) of 2,6-dibromopyridine and 16.45 g (127 mmol) of 2-(4-piperidyl)ethanol, and after chromatography on silica gel, eluting with a 30/70 mixture of ethyl acetate and cyclohexane, 7 g of pure product are obtained in the form of an oil.

15

13.2. 2-[1-(6-Bromo-2-pyridyl)-4-piperidyl]ethyl methanesulfonate

The process is performed as described in Example 3 (step 3.2). Starting with 7 g (24.50 mmol) of 2-[1-(6-bromo-2-pyridyl)-4-piperidyl]ethanol, obtained in step 13.1, 2.50 ml (26.90 mmol) of mesyl chloride and 3.80 ml (26.90 mmol) of triethylamine, 8.68 g of product are obtained in the form of an oil, which is used without further purification in the following step.

25

13.3. 3-{2-[1-(6-Bromo-2-pyridyl)-4-piperidyl]ethyl}-
1,3-oxazolidine-2,4-dione

The process is performed according to the method described in Example 3 (step 3.3). Starting with
5 8.68 g (23.80 mmol) of 2-[1-(6-bromo-2-pyridyl)-4-piperidyl]ethyl methanesulfonate, obtained in step 13.2, 2.90 g (28.60 mmol) of 1,3-oxazolidine-2,4-dione and 6 ml (47.60 mmol) of 1,1,3,3-tetramethylguanidine, and after chromatography on silica gel, eluting with a
10 97/3 mixture of dichloromethane and methanol, 4.52 g of product are obtained in the form of an oil.

13.4. 3-{2-[1-(6-Isobutyl-2-pyridyl)-4-piperidyl]-ethyl}-1,3-oxazolidine-2,4-dione

15 2 g (5.43 mmol) of 3-{2-[1-(6-bromo-2-pyridyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, prepared in step 13.3, and 0.20 g (0.271 mmol) of dichlorobis(triphenylphosphine)palladium ($\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$) suspended in 10 ml of tetrahydrofuran are introduced
20 under an inert atmosphere. 22 ml (10.80 mmol) of a solution (0.5 M) of bromo(isobutyl)zinc in tetrahydrofuran are then added. Stirring is continued at room temperature for 17 hours. The reaction mixture is poured into water and ethyl acetate. The phases are
25 separated by settling, the aqueous phase is extracted twice with ethyl acetate, the combined organic phases are dried over sodium sulfate and the filtrate is concentrated under reduced pressure. The residue thus

obtained is purified by chromatography on silica gel, eluting with a 20/80 mixture of ethyl acetate and cyclohexane.

1.41 g of product are obtained in the form of
5 a white solid.

m.p. (°C): 94-96°C

13.5. 2-(Methylamino)-2-oxoethyl 2-[1-(6-isobutyl-2-pyridyl)-4-piperidyl]ethylcarbamate

10 The process is performed according to the procedure described in Example 1 (step 1.5). Starting with 0.72 g (2.08 mmol) of 3-{2-[1-(6-isobutyl-2-pyridyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, obtained in step 13.4, and 5.20 ml (10.40 mmol) of a
15 solution (2M) of methylamine in tetrahydrofuran, and after chromatography on silica gel, eluting with a 95/5/0.5 mixture of dichloromethane, methanol and 28% aqueous ammonia, followed by crystallization from diisopropyl ether, 0.540 g of pure product is obtained
20 in the form of a white solid.


LC-MS: M+H = 377

m.p. (°C): 97-99°C

¹H NMR (DMSO) δ (ppm): 0.85 (d, 6H); 1.05 (m, 2H); 1.20-1.60 (unresolved peak, 3H); 1.70 (broad d, 2H); 2.0 (m, 1H); 2.40 (d, 2H); 2.55 (d, 3H); 2.70 (broad t, 2H);
25 3.05 (broad q, 2H); 4.20 (broad d, 2H); 4.30 (s, 2H); 6.35 (d, 1H); 6.55 (d, 1H); 7.15 (broad t, 1H); 7.40 (dd, 1H); 7.75 (broad s, 1H).

Example 14 (Compound 58)

2-Amino-2-oxoethyl 2-[1-(6-phenyl-2-pyridyl)-4-piperidyl]methylcarbamate



5

14.1. 3-{2-[1-(6-Phenyl-2-pyridyl)-4-piperidyl]methyl}-1,3-oxazolidine-2,4-dione

0.20 g (0.56 mmol) of 3-{2-[1-(6-bromo-2-pyridyl)-4-piperidyl]methyl}-1,3-oxazolidine-2,4-dione,
 10 prepared according to the procedure described in Example 13 (steps 13.1, 13.2 and 13.3), 0.089 g (0.73 mmol) of phenylboronic acid and 0.480 g (2.25 mmol) of hydrated potassium phosphate suspended
 15 in 3 ml of 1,2-dimethoxyethane are introduced under an inert atmosphere. 0.040 g (0.0346 mmol) of tetrakis-(triphenylphosphine)palladium is then added. The reaction mixture is then maintained at about 85°C for 16 hours. The resulting mixture is concentrated under
 20 reduced pressure. The residue is taken up in ethyl acetate and water, the aqueous phase is separated out and extracted twice with ethyl acetate, the combined organic phases are dried over sodium sulfate and the filtrate is concentrated under reduced pressure. The
 25 residue thus obtained is purified by chromatography on

silica gel, eluting with a 40/60 mixture of ethyl acetate and cyclohexane.

0.175 g of product is obtained.

5 14.2. 2-Amino-2-oxoethyl 2-[1-(6-phenyl-2-pyridyl)-4-piperidyl]methylcarbamate

The process is performed according to the procedure described in Example 3 (step 3.4). Starting with 0.175 g (0.499 mmol) of 3-{2-[1-(6-phenyl-2-pyridyl)-4-piperidyl]methyl}-1,3-oxazolidine-2,4-dione,
10 obtained in step 14.1, and 2.5 ml (17.45 mmol) of a solution (7M) of aqueous ammonia in methanol, 0.070 g of pure product is obtained in the form of a white solid, after crystallization from ethyl acetate.


15 LC-MS: M+H = 369

m.p. (°C): 131-132°C

¹H NMR (CDCl₃) δ (ppm): 1.20-1.90 (unresolved peak, 5H);
2.90 (broad t, 2H); 3.20 (t, 2H); 4.50 (broad d, 2H);
4.60 (s, 2H); 5.0 (broad s, 1H); 5.55 (broad s, 1H);
20 6.15 (broad s, 1H); 6.65 (d, 1H); 7.10 (d, 1H); 7.35-
7.60 (m, 4H); 8.15 (dd, 2H).

Example 15 (Compound 130)

**2-(Methylamino)-2-oxoethyl 2-[1-(5-chloro-2-quinolyl)-
25 4-piperidyl]ethylcarbamate**



15.1. 2-[1-(5-Chloro-2-quinolyl)-4-piperidyl]ethanol

The process is performed as described in
 5 Example 4 (step 4.1). Starting with 4.78 g (24.14 mmol)
 of 2-chloro-5-chloroquinoline (J. Med. Chem., 2002, 45,
 3130-3137) and 3.43 g (26.55 mmol) of 2-(4-piperidyl)-
 ethanol, 7 g of product are obtained in the form of an
 oil, which is used without further purification in the
 10 following step.

15.2. 2-[1-(5-Chloro-2-quinolyl)-4-piperidyl]ethyl methanesulfonate

The process is performed as described in
 15 Example 3 (step 3.2). Starting with 7 g (24.07 mmol) of
 2-[1-(5-chloro-2-quinolyl)-4-piperidyl]ethanol,
 obtained in step 15.1, 3.31 g (28.89 mmol) of mesyl
 chloride and 5.10 ml (36.11 mmol) of triethylamine,
 8.70 g of product are obtained in the form of an oil,
 20 which is used without further purification in the
 following step.

15.3. 3-{2-[1-(5-Chloro-2-quinolyl)-4-piperidyl]ethyl}- 1,3-oxazolidine-2,4-dione

The process is performed according to the method described in Example 3 (step 3.3). Starting with 8.7 g (23.58 mmol) of 2-[1-(5-chloro-2-quinolyl)-4-piperidyl]ethyl methanesulfonate, obtained in step 15.2, 2.86 g (28.30 mmol) of 1,3-oxazolidine-2,4-dione and 5.43 g (47.17 mmol) of 1,1,3,3-tetramethylguanidine, and after chromatography on silica gel, eluting with a 99.5/0.5 mixture of dichloromethane and methanol, 6.40 g of product are obtained in the form of a white solid.

m.p. (°C): 136°C

15.4. 2-(Methylamino)-2-oxoethyl 2-[1-(5-chloro-2-quinolyl)-4-piperidyl]ethylcarbamate

15 The process is performed according to the procedure described in Example 1 (step 1.5). Starting with 6.40 g (17.12 mmol) of 3-{2-[1-(5-chloro-2-quinolyl)-4-piperidyl]ethyl}-1,3-oxazolidine-2,4-dione, obtained in step 15.3, and 60 ml (119.84 mmol) of a solution (2M) of methylamine in tetrahydrofuran, and after chromatography on silica gel, eluting with a 98/2 and then 96/4 mixture of dichloromethane and methanol, followed by crystallization from diisopropyl ether, 5.14 g of product are obtained in the form of a white solid.

LC-MS: M+H = 405

m.p. (°C): 158-162°C

^1H NMR (CDCl_3) δ (ppm): 1.10-1.80 (unresolved peak, 5H);
 1.9 (broad d, 2H); 2.90 (d, 3H); 3.0 (m, 2H); 3.30 (q,
 2H); 4.60 (m, 4H); 4.85 (broad s, 1H); 6.10 (broad s,
 1H); 7.05 (d, 1H); 7.25 (d, 1H); 7.40 (dd, 1H); 7.60
 5 (d, 1H); 8.30 (s, 1H).


Table 1 below illustrates the chemical
 structures and physical properties of a number of
 compounds according to the invention.



10 In this table:













- in the "base or salt" column, "base" represents a
 compound in free base form, whereas "HCl" represents
 a compound in hydrochloride form;
- OMe represents a methoxy group.













15













Table 1




























No	R ₁	m	n	A	B	R ₃	base or salt	m.p. (°C)
1.		2	2	N	(CH ₂) ₃	CH ₂ CONHCH ₃	base	131
2.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	417*
3.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	165-166
4.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	144-145
5.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	138-139












No	R ₁	m	n	A	B	R ₃	base or salt	m.p. (°C)
6.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	173-174
7.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	196-197
8.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	157-158
9.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	225-226
10.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	121-122
11.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	141-142
12.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	139-140
13.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	170-171
14.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	128-129
15.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	153-154
16.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	160-161
17.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	148-149
18.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	156-157
19.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	157-158
20.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	178-179
21.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	432*
22.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	158-159
23.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	180-181
24.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	185-186
25.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	193-194
26.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	153-154
27.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	125-126
28.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	193-194












No	R ₁	m	n	A	B	R ₃	base or salt	m.p. (°C)
29.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	176-177
30.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	160-161
31.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	156-157
32.		2	2	N	(CH ₂) ₂	CH ₂ CONH ₂	base	189-190
33.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	base	196-197
34.		2	3	N	(CH ₂) ₃	CH ₂ CONHCH ₃	base	100 (dec.)
35.		2	3	N	(CH ₂) ₃	CH ₂ CONHCH ₃	HCl	451*
36.		2	3	N	(CH ₂) ₃	CH ₂ CONHCH ₃	HCl	172-174
37.		2	3	N	(CH ₂) ₃	CH ₂ CONHCH ₃	base	118-122
38.		2	2	N	(CH ₂) ₂	CH ₂ CONHCH ₃	HCl	166
39.		2	2	N	(CH ₂) ₃	CH ₂ CONHCH ₃	HCl	167
40.		1	1	CH	CH ₂	CH ₂ CONH ₂	base	149-151
41.		1	1	CH	CH ₂	CH ₂ CONHCH ₃	base	137-139
42.		1	1	CH	CH ₂	CH ₂ CONHCH ₂ (cyclopropyl)	base	119-121
43.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	110-112
44.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	140-142
45.		2	2	CF	(CH ₂) ₂	CH ₂ CONHCH ₃	base	127-128
46.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	HCl	168-170
47.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	HCl	95-100
48.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	113-115
49.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	97-99
50.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	107-109
51.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	108-110









No	R ₁	m	n	A	B	R ₃	base or salt	m.p. (°C)
52.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	98-100
53.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	123-125
54.		2	2	CH	-	CH ₂ CONH ₂	base	169
55.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	148-149
56.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	131-132
57.		2	2	CH	-	CH ₂ CONH ₂	base	153
58.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	131-132
59.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	116-117
60.		2	2	CH	-	CH ₂ CONH ₂	base	190-191
61.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	155-156
62.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	155-156
63.		2	2	CH	-	CH ₂ CONH ₂	base	159-160
64.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	132
65.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	134-135
66.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	146-147
67.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	332*
68.		2	2	CH	-	CH ₂ CONH ₂	base	131-132
69.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	129-130
70.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	86-87
71.		2	2	CH	-	CH ₂ CONH ₂	base	58-59
72.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	387*
73.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	58-59
74.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	152-153
75.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	163
76.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	121-123
77.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	137-139
78.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	327*
79.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	162-163

No	R ₁	m	n	A	B	R ₃	base or salt	m.p. (°C)
80.		2	2	CH	-	CH ₂ CONH ₂	base	168
81.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	176
82.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	174-175
83.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	169-170
84.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	145-146
85.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	119-121
86.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	109-111
87.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	169-170
88.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	164-165
89.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	174-175
90.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	162-163
91.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	151-152
92.		2	2	CH	-	CH ₂ CONH ₂	base	195-196
93.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	229-230
94.		2	2	CH	-	CH ₂ CONH ₂	base	194-195
95.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	222-223
96.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	182
97.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	196-199
98.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	161-163
99.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	157-158
100.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	175-176
101.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	163-164
102.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	418*
103.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	190-191

No	R ₁	m	n	A	B	R ₃	base or salt	m.p. (°C)
104.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	165-166
105.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	323*
106.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	202-203
107.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	172-173
108.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	370*
109.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	177-178
110.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	202-203
111.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	185-186
112.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	165-166
113.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	126-127
114.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	309*
115.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	159-160
116.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	161-162
117.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	126-127
118.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	158-159
119.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	174-175
120.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	204-205
121.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	229-230
122.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	228-229
123.		2	2	CH	CH ₂	CH ₂ CONHCH ₃	base	153-157
124.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	206-212

No	R ₁	m	n	A	B	R ₃	base or salt	m.p.(°C)
125.		2	2	CH	CH ₂	CH(CH ₃)CONHCH ₃	base	129-130
126.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	126-128
127.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	135-137
128.		2	2	CF	(CH ₂) ₂	CH ₂ CONHCH ₃	base	147-149
129.		2	2	C(OH)	(CH ₂) ₂	CH ₂ CONHCH ₃	base	82-90
130.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	158-162
131.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	171-175
132.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	162-163
133.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	154-156
134.		2	2	CH	CH ₂	CH ₂ CONHCH ₃	base	147-151
135.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	207-208
136.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	155-157
137.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	189-191
138.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	163-165
139.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	135-137
140.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₂ CH ₃	base	114-116
141.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₂ (cyclopropyl)	base	95-100
142.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	191-193
143.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	149-151
144.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	146-150
145.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	171-173
146.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	140-142
147.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	160-162
148.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	HCl	148-150
149.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	HCl	219-221
150.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	130-132
151.		1	2	CH	CH ₂	CH ₂ CONH ₂	base	109-111
152.		1	2	CH	CH ₂	CH ₂ CONHCH ₃	base	99-102

No	R ₁	m	n	A	B	R ₃	base or salt	m.p. (°C)
153.		2	2	CH	-	CH ₂ CONHCH ₃	HCl	232-237
154.		2	2	CH	CH ₂	CH ₂ CONHCH ₃	HCl	208-212
155.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	HCl	132-136
156.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	HCl	120-124
157.		2	2	CH	(CH ₂) ₃	CH ₂ CONHCH ₃	HCl	385*
158.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	135-139
159.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	144-148
160.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	134-136
161.		2	2	CH	CH ₂	CH ₂ CONHCH ₃	HCl	387*
162.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	100-104
163.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	118-120
164.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	HCl	186-188
165.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	80-86
166.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	156-158
167.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	176-178
168.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	148-150
169.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	154-156
170.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	156-158
171.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	156-158
172.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	HCl	205-210
173.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	419*

No	R ₁	m	n	A	B	R ₃	base or salt	m.p. (°C)
174.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	105-106
175.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	129-130
176.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	112-113
177.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	180-181
178.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	244-245
179.		2	2	CH	(CH ₂) ₂	CH ₂ CONHCH ₃	base	146-150
180.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	169-171
181.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	137-138
182.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	208-209
183.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	183-184
184.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	172-173
185.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	187-188
186.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	191-192
187.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	222-223
188.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	149-150
189.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	167-168
190.		2	2	CH	CH ₂	CH ₂ CONH ₂	base	220-221
191.		2	2	CH	(CH ₂) ₂	CH ₂ CONH ₂	base	201-202

★ M+H (LC-MS)

dec. = decomposition of the product

The compounds of the invention underwent
 5 pharmacological trials to determine their inhibitory
 effect on the enzyme FAAH (Fatty Acid Amide Hydrolase).

The inhibitory activity was demonstrated in a
 radioenzymatic test based on measuring the product of
 hydrolysis (ethanolamine [1-³H]) of anandamide [ethanol-
 10 amine 1-³H] with FAAH (*Life Sciences* (1995), 56, 1999-
 2005 and *Journal of Pharmacology and Experimented
 Therapeutics* (1997), 283, 729-734). Thus, mouse brains
 (minus the cerebellum) are removed and stored at -80°C.

Membrane homogenates are prepared extemporaneously by homogenizing the tissues using a Polytron blender in 10 mM tris-HCl buffer (pH 8.0) containing 150 mM NaCl and 1 mM EDTA. The enzymatic reaction is then performed in 70 μ l of buffer containing fatty-acid-free bovine serum albumin (1 mg/ml). To test compounds at different concentrations, anandamide [ethanolamine 1- 3 H] (specific activity of 15-20 Ci/mmol) diluted to 10 μ M with cold anandamide, and the membrane preparation (400 μ g of frozen tissue per test) are successively added. After 15 minutes at 25°C, the enzymatic reaction is quenched by adding 140 μ L of chloroform/methanol (2:1). The mixture is stirred for 10 minutes and then centrifuged for 15 minutes at 3500 g. An aliquot (30 μ L) of the aqueous phase containing the ethanolamine [1- 3 H] is counted by liquid scintillation. Under these conditions, the most active compounds of the invention have IC₅₀ values (concentration that inhibits 50% of the control enzymatic activity of FAAH) of between 0.001 and 1 μ M.

Table 2 below shows the IC₅₀ values of a few compounds according to the invention.

Table 2

Compound No.	IC ₅₀
47	85 μ M

126	113 μ M
166	87 μ M

It is thus seen that the compounds according to the invention have inhibitory activity on the enzyme FAAH.

5 The *in vivo* activity of the compounds of the invention was evaluated in a test of analgesia.

 Thus, the intraperitoneal (i.p.) administration of PBQ (phenylbenzoquinone, 2 mg/kg in 0.9% sodium chloride solution containing 5% ethanol) to
10 male OF1 mice weighing 25 to 30 g, causes abdominal tractions, on average 30 torsions or contractions during the period of 5 to 15 minutes after injection. The test compounds are administered orally or intraperitoneally as a 0.5% suspension in Tween 80, 60
15 minutes or 120 minutes before the administration of PBQ. Under these conditions, the most powerful compounds of the invention produce a 35% to 70% reduction in the number of tractions induced with PBQ, within a dose range of between 1 and 30 mg/kg.

20 Table 3 below shows the results of the analgesia test for a few compounds according to the invention.

Table 3

Compound No.	Reduction in the number of tractions (%)
47	-63% (b)
126	-43% (b)
166	-62% (a)

(a) 1 mg/kg p.o. at 2 hours;

(b) 3 mg/kg p.o. at 1 hour

5

The enzyme FAAH (*Chemistry and Physics of Lipids*, (2000), 108, 107-121) catalyses the hydrolysis of the endogenous amide and ester derivatives of various fatty acids such as *N*-arachidonylethanolamine (anandamide), *N*-palmitoylethanolamine, *N*-oleoylethanol-
 10 amine, oleamide or 2-arachidonoylglycerol. These derivatives exert different pharmacological activities by interacting, inter alia, with the cannabinoid and vanilloid receptors.

15

The compounds of the invention block this degradation pathway and increase the tissue content of these endogenous substances. They may be used in this respect in the prevention and treatment of pathologies in which the endogenous cannabinoids, and/or any other
 20 substrates metabolized by the enzyme FAAH, are involved. Examples that may be mentioned include the following diseases and complaints:

pain, especially acute or chronic pain of neurogenic type: migraine, neuropathic pain including the forms associated with the herpes virus and diabetes, acute or chronic pain associated with inflammatory diseases:

- 5 arthritis, rheumatoid arthritis, osteoarthritis, spondylitis, gout, vascularitis, Crohn's disease, irritable bowel syndrome, acute or chronic peripheral pain, vertigo, vomiting, nausea, in particular post-10 chemotherapy nausea, eating disorders, in particular anorexia and cachexia of diverse nature, neurological and psychiatric pathologies: tremor, dyskinesia, dystonia, spasticity, compulsive and15 obsessive behaviour, Tourette's syndrome, all forms of depression and anxiety of any nature or origin, mood disorders, psychoses, acute and chronic neurodegenerative diseases: Parkinson's disease, Alzheimer's disease, senile20 dementia, Huntington's chorea, lesions associated with cerebral ischaemia and cranial and medullary trauma, epilepsy, sleeping disorders, including sleep apnoea, cardiovascular diseases, in particular hypertension,25 cardiac arrhythmia, arteriosclerosis, heart attack, cardiac ischaemia, renal ischaemia,

- cancers: benign skin tumours, papillomas and cerebral tumours, prostate tumours, cerebral tumours (glioblastomas, medullo-epitheliomas, medullo-blastomas, neuroblastomas, tumours of embryonic origin,
- 5 astrocytomas, astroblastomas, ependyomas, oligodendro-gliomas, plexus tumour, neuroepitheliomas, pineal gland tumours, ependymoblastomas, malignant meningiomas, sarcomatoses, malignant melanomas, schwennomas), immune system disorders, especially autoimmune
- 10 diseases: psoriasis, lupus erythematosus, connective tissue diseases, Sjögren's syndrome, ankylosing spondylitis, undifferentiated spondylitis, Behcet's disease, haemolytic autoimmune anaemias, multiple sclerosis, amyotrophic lateral sclerosis, amylosis,
- 15 graft rejection, diseases affecting the plasmocytic line,
- allergic diseases: immediate or delayed hypersensitivity, allergic rhinitis or allergic conjunctivitis, contact dermatitis,
- 20 parasitic, viral or bacterial infectious diseases: AIDS, meningitis, inflammatory diseases, especially articular diseases: arthritis, rheumatoid arthritis, osteoarthritis, spondylitis, gout, vascularitis, Crohn's disease, irritable bowel syndrome,
- 25 osteoporosis, ocular complaints: ocular hypertension, glaucoma,
- pulmonary complaints: respiratory pathway diseases, bronchospasms, coughing, asthma, chronic bronchitis,

chronic obstruction of the respiratory pathways,
emphysema,
gastrointestinal diseases: irritable bowel syndrome,
intestinal inflammatory disorders, ulcers, diarrhoea,
5 urinary incontinence and inflammation of the bladder.

The use of the compounds according to the
invention, in the form of the base, or a
pharmaceutically acceptable acid-addition salt, hydrate
or solvate, for the preparation of a medicinal product
10 for treating the pathologies mentioned above forms an
integral part of the invention.

A subject of the invention is also medicinal
products comprising a compound of formula (I), or an
acid-addition salt, or alternatively a pharmaceutically
15 acceptable hydrate or solvate of the compound of
formula (I). These medicinal products find their
therapeutic use especially in the treatment of the
pathologies mentioned above.

According to another of its aspects, the
20 present invention relates to pharmaceutical
compositions containing, as active principal, at least
one compound according to the invention. These
pharmaceutical compounds contain an effective dose of a
compound according to the invention, or a
25 pharmaceutically acceptable acid-addition salt, hydrate
or solvate of the said compound, and optionally one or
more pharmaceutically acceptable excipients.

The said excipients are chosen, according to the pharmaceutical form and the desired administration form, from the usual excipients known to those skilled in the art.

5 In the pharmaceutical compositions of the present invention for oral, sublingual, subcutaneous, intramuscular, intravenous, topical, local, intrathecal, intranasal, transdermal, pulmonary, ocular or rectal administration, the active principal of
10 formula (I) above, or the possible acid-addition salt, solvate or hydrate thereof, may be administered in a unit administration form, as a mixture with standard pharmaceutical excipients, to man and animals for the prophylaxis or treatment of the above disorders or
15 diseases.

 The appropriate unit administration forms comprise oral forms such as tablets, soft or hard gel capsules, powders, granules, chewing gums and oral solutions or suspensions, sublingual, buccal, intra-
20 tracheal, intraocular and intranasal administration forms, forms for administration by inhalation, subcutaneous, intramuscular or intravenous administration forms and rectal or vaginal administration forms. For topical administration, the compounds according to the
25 invention may be used in creams, ointments or lotions.

 By way of example, a unit administration form of a compound according to the invention in the form of a tablet may comprise the following components:

	Compound according to the invention	50.0 mg
	Mannitol	223.75 mg
	Sodium croscarmellose	6.0 mg
	Maize starch	15.0 mg
5	Hydroxypropylmethylcellulose	2.25 mg
	Magnesium stearate	3.0 mg


The said unit forms are dosed to allow a daily administration of from 0.01 to 20 mg of active principal per kg of body weight, depending on the presentation form.

There may be particular cases in which higher or lower doses are suitable, and such doses also form part of the invention. According to the usual practice, the dose that is suitable for each patient is determined by the doctor according to the mode of administration and the weight and response of the said patient.

According to another of its aspects, the invention also relates to a method for treating the pathologies mentioned above, which comprises the administration of an effective dose of a compound according to the invention, a pharmaceutically acceptable acid-addition salt thereof or a solvate or hydrate of the said compound.

CLAIMS

1. Compound corresponding to formula (I)



(I)

5

in which

- A represents a nitrogen atom or a group CR₂ in which R₂ represents a hydrogen or fluorine atom or a hydroxyl, cyano, trifluoromethyl, C₁₋₆-alkyl or C₁₋₆-alkoxy group;
 10 n represents an integer equal to 2 or 3 and m represents an integer equal to 2 when A represents a nitrogen atom;
 n represents an integer equal to 1, 2 or 3 and m represents an integer equal to 1 or 2 when A represents a group CR₂;
 15 B represents a covalent bond or a C₁₋₈-alkylene group;
 R₁ represents a group chosen from phenyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl,
 20 imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, naphthyl, quinolyl, tetrahydroquinolyl, isoquinolyl, tetrahydroisoquinolyl, phthalazinyl, quinazolinyl,

quinoxaliny, naphthyridiny, cinnoly,
 imidazopyrimidiny, thienopyrimidiny, benzofurany,
 benzothieny, benzimidazolyl, benzoxazolyl,
 benzisoxazolyl, benzothiazolyl, benzisothiazolyl,
 5 indazolyl, pyrrolopyridyl, furopyridyl,
 dihydrofuropyridyl, thienopyridyl, dihydro-
 thienopyridyl, imidazopyridyl, pyrazolopyridyl,
 oxazolopyridyl, isoxazolopyridyl and thiazolopyridyl;
 the group R_1 optionally being substituted with one or
 10 more groups R' and/or R'' ;
 R' represents a halogen atom or a cyano, nitro,
 hydroxyl, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -thioalkyl, C_{1-6} -
 fluoroalkyl, C_{1-6} -fluoroalkoxy, C_{1-6} -fluorothioalkyl,
 C_{3-7} -cycloalkyl, C_{3-7} -cycloalkyl- C_{1-6} -alkylene,
 15 azetidiny, piperidyl, pyrrolidiny, morpholiny,
 piperaziny, azepiny, NH_2 , NHR_6 , NR_6R_7 , NR_6COR_7 ,
 $NR_6SO_2R_7$, COR_6 , CO_2R_6 , SO_2R_6 , $SO_2NR_6R_7$ or $-O-(C_{1-6}$ -
 alkylene)-O- group;
 R'' represents a phenyl, imidazolyl, pyridyl or
 20 pyrimidiny;
 the group(s) R'' being optionally substituted with one
 or more groups R' , which may be identical to or
 different from each other;
 R_3 represents a group of general formula CHR_4CONHR_5 in
 25 which
 R_4 represents a hydrogen atom or a C_{1-6} -alkyl group and
 R_5 represents a hydrogen atom or a C_{1-6} -alkyl, C_{3-7} -
 cycloalkyl or C_{3-7} -cycloalkyl- C_{1-6} -alkylene group;

R_6 and R_7 represent, independently of each other, a C_{1-6} -alkyl group;

in the form of a base, an acid-addition salt, a hydrate or a solvate.

5 2. Compound of formula (I) according to Claim 1, characterized in that:

A represents a nitrogen atom;

n represents an integer equal to 2 or 3 and m

represents an integer equal to 2;

10 B represents a C_{1-8} -alkylene group;

R_1 represents a group chosen from phenyl, pyridyl, pyrimidinyl, thiadiazolyl and naphthyl;

the group R_1 being optionally substituted with one or more groups R' and/or R'' ;

15 R' represents a halogen atom or a nitro or C_{1-6} -fluoroalkyl group;

R'' represents a phenyl optionally substituted with one or more groups, which may be identical to or different from each other, chosen from a halogen atom or a cyano,

20 C_{1-6} -alkoxy or C_{1-6} -fluoroalkoxy group;

R_3 represents a group of general formula CHR_4CONHR_5 in which

R_4 represents a hydrogen atom and

R_5 represents a hydrogen atom or a C_{1-6} -alkyl, C_{3-7} -

25 cycloalkyl or C_{3-7} -cycloalkyl- C_{1-6} -alkylene group;

in the form of a base, an acid-addition salt, a hydrate or a solvate.

3. Compound of formula (I) according to Claim 1, characterized in that:

A represents a group CR_2 in which R_2 represents a hydrogen or fluorine atom or a hydroxyl group;

5 m represents an integer equal to 1 or 2 and n represents an integer equal to 1 or 2;

B represents a covalent bond or a C_{1-4} -alkylene group;

R_1 represents a group chosen from phenyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, thiazolyl, quinolyl, isoquinolyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, furopyridyl, thienopyrimidinyl, imidazopyrimidinyl, benzothiazolyl, benzimidazolyl and benzoxazolyl;

15 the group R_1 being optionally substituted with one or more groups R' and/or R'' ;

R' represents a halogen atom, a cyano or C_{1-6} -alkyl, a C_{1-6} -alkoxy, C_{1-6} -fluoroalkyl, C_{1-6} -fluoroalkoxy, C_{3-7} -cycloalkyl, pyrrolidinyl, NH_2 , NR_6R_7 or COR_6 group;

R'' represents a phenyl, imidazolyl or pyridyl;

20 the group(s) R'' being optionally substituted with one or more groups R' , which may be identical to or different from each other, more particularly with one or more chlorine or fluorine atoms;

R_3 represents a group of general formula CHR_4CONHR_5 in which

R_4 represents a hydrogen atom or a C_{1-6} -alkyl group, and

R_5 represents a hydrogen atom or a C_{1-6} -alkyl, C_{3-7} -cycloalkyl or C_{3-7} -cycloalkyl- C_{1-6} -alkylene group;

R_6 and R_7 represent, independently of each other, a C_{1-6} -alkyl group;

in the form of a base, an acid-addition salt, a hydrate or a solvate.

5 4. Compound of formula (I) according to Claim 1 or 3, characterized in that:

A represents a group CR_2 in which R_2 represents a hydrogen atom;

m is equal to 2 and n is equal to 2;

10 B represents an ethyl group;

R_1 represents a group chosen from phenyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, thiazolyl, quinolyl, isoquinolyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, furopyridyl, thienopyrimidinyl, imidazopyrimidinyl, benzothiazolyl, benzimidazolyl and benzoxazolyl;

the group R_1 being optionally substituted with one or more groups R' and/or R'' ;

20 R' represents a halogen atom or a cyano or C_{1-6} -alkyl, a C_{1-6} -alkoxy, C_{1-6} -fluoroalkyl, C_{1-6} -fluoroalkoxy, C_{3-7} -cycloalkyl, pyrrolidinyl, NH_2 , NR_6R_7 or COR_6 group;

R'' represents a phenyl, imidazolyl or pyridyl;

the group(s) R'' being optionally substituted with one or more groups R' , which may be identical to or
25 different from each other, more particularly with one or more chlorine or fluorine atoms;

R_3 represents a group of general formula CHR_4CONHR_5 in which

R_4 represents a hydrogen atom and


R_5 represents a hydrogen atom or a C_1 - C_6 -alkyl group;

R_6 and R_7 represent, independently of each other, a C_1 - C_6 -alkyl group;

5 in the form of the base, an acid-addition salt, a hydrate or a solvate.

5. Process for preparing a compound of formula (I) according to any one of Claims 1 to 4, comprising the step consisting in

10 converting the carbamate ester of general formula (Ia)




in which n , m , A , B , R_1 and R_4 are as defined in the
15 general formula (I) according to Claim 1 and R

represents a methyl or ethyl group,

by aminolysis using an amine of general formula R_5NH_2 in which R_5 is as defined in the general formula (I) according to Claim 1.


20 6. Process for preparing a compound of formula (I) according to any one of Claims 1 to 4, comprising the step consisting in converting the oxazolidinedione derivative of general formula (V)



in which n , m , A , B , R_1 and R_4 are as defined in the
 general formula (I) according to Claim 1,
 5 by aminolysis using an amine of general formula R_5NH_2 in
 which R_5 is as defined in the general formula (I)
 according to Claim 1.

7. Compound corresponding to the general
 formula (Ia)

10



in which
 A represents a nitrogen atom or a group CR_2 in which R_2
 15 represents a hydrogen or fluorine atom or a hydroxyl,
 cyano, trifluoromethyl, C_1-6 -alkyl or C_1-6 -alkoxy group;
 n represents an integer equal to 2 or 3 and m
 represents an integer equal to 2 when A represents a
 nitrogen atom;
 20 n represents an integer equal to 1, 2 or 3 and m
 represents an integer equal to 1 or 2 when A represents
 a group CR_2 ;

- B represents a covalent bond or a C₁₋₈-alkylene group;
 R₁ represents a group chosen from phenyl, pyridyl,
 pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl,
 oxazolyl, isoxazolyl, thiazolyl, isothiazolyl,
 5 imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl,
 naphthyl, quinolyl, tetrahydroquinolyl, isoquinolyl,
 tetrahydroisoquinolyl, phthalazinyl, quinazolinyl,
 quinoxalinyl, naphthyridinyl, cinnolyl,
 imidazopyrimidinyl, thienopyrimidinyl, benzofuranyl,
 10 benzothienyl, benzimidazolyl, benzoxazolyl,
 benzisoxazolyl, benzothiazolyl, benzisothiazolyl,
 indazolyl, pyrrolopyridyl, furopyridyl,
 dihydrofuropyridyl, thienopyridyl, dihydro-
 thienopyridyl, imidazopyridyl, pyrazolopyridyl,
 15 oxazolopyridyl, isoxazolopyridyl and thiazolopyridyl;
 the group R₁ optionally being substituted with one or
 more groups R' and/or R'';
- R' represents a halogen atom or a cyano, nitro,
 hydroxyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-thioalkyl, C₁₋₆-
 20 fluoroalkyl, C₁₋₆-fluoroalkoxy, C₁₋₆-fluorothioalkyl,
 C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-C₁₋₆-alkylene,
 azetidiny, piperidyl, pyrrolidinyl, morpholinyl,
 piperazinyl, azepinyl, NH₂, NHR₆, NR₆R₇, NR₆COR₇,
 NR₆SO₂R₇, COR₆, CO₂R₆, SO₂R₆, SO₂NR₆R₇ or -O-(C₁₋₆-
 25 alkylene)-O- group;
- R'' represents a phenyl, imidazolyl, pyridyl, pyrazinyl,
 pyridazinyl or pyrimidinyl;

the group(s) R'' being optionally substituted with one or more groups R' , which may be identical to or different from each other;


R_4 represents a hydrogen atom or a C_{1-6} -alkyl group;

5 R_6 and R_7 represent, independently of each other, a C_{1-6} -alkyl group;

R represents a methyl or ethyl group.

8. Compound corresponding to the general formula (V)

10



in which

A represents a nitrogen atom or a group CR_2 in which R_2 represents a hydrogen or fluorine atom or a hydroxyl, cyano, trifluoromethyl, C_{1-6} -alkyl or C_{1-6} -alkoxy group;

n represents an integer equal to 2 or 3 and m

represents an integer equal to 2 when A represents a nitrogen atom;

20 n represents an integer equal to 1, 2 or 3 and m

represents an integer equal to 1 or 2 when A represents a group CR_2 ;

B represents a covalent bond or a C_{1-8} -alkylene group;

- R_1 represents a group chosen from phenyl, pyridyl,
 pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl,
 oxazolyl, isoxazolyl, thiazolyl, isothiazolyl,
 imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl,
 5 naphthyl, quinolyl, tetrahydroquinolyl, isoquinolyl,
 tetrahydroisoquinolyl, phthalazinyl, quinazolinyl,
 quinoxalinyl, naphthyridinyl, cinnolyl,
 imidazopyrimidinyl, thienopyrimidinyl, benzofuranyl,
 benzothienyl, benzimidazolyl, benzoxazolyl,
 10 benzisoxazolyl, benzothiazolyl, benzisothiazolyl,
 indazolyl, pyrrolopyridyl, furopyridyl,
 dihydrofuropyridyl, thienopyridyl, dihydro-
 thienopyridyl, imidazopyridyl, pyrazolopyridyl,
 oxazolopyridyl, isoxazolopyridyl and thiazolopyridyl;
 15 the group R_1 optionally being substituted with one or
 more groups R' and/or R'' ;
 R' represents a halogen atom or a cyano, nitro,
 hydroxyl, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -thioalkyl, C_{1-6} -
 fluoroalkyl, C_{1-6} -fluoroalkoxy, C_{1-6} -fluorothioalkyl,
 20 C_{3-7} -cycloalkyl, C_{3-7} -cycloalkyl- C_{1-6} -alkylene,
 azetidiny, piperidyl, pyrrolidinyl, morpholinyl,
 piperazinyl, azepinyl, NH_2 , NHR_6 , NR_6R_7 , NR_6COR_7 ,
 $NR_6SO_2R_7$, COR_6 , CO_2R_6 , SO_2R_6 , $SO_2NR_6R_7$ or $-O-(C_{1-6}$ -
 alkylene)-O- group;
 25 R'' represents a phenyl, imidazolyl, pyridyl, pyrazinyl,
 pyridazinyl or pyrimidinyl;

the group(s) R" being optionally substituted with one or more groups R', which may be identical to or different from each other;

R₄ represents a hydrogen atom or a C₁₋₆-alkyl group;

5 R₆ and R₇ represent, independently of each other, a C₁₋₆-alkyl group;

the following compounds being excluded:

* 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-2,4-oxazolidinedione

10 * 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-5-methyl-2,4-oxazolidinedione

* 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-5-ethyl-2,4-oxazolidinedione

15 * 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-5-propyl-2,4-oxazolidinedione

* 3-[1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-piperidyl]-5-(1-methylethyl)-2,4-oxazolidinedione.

9. Compound of formula (I) according to any one of Claims 1 to 4, in the form of the base or of a
20 pharmaceutically acceptable acid-addition salt, hydrate or solvate, for its use as a medicament.

10. Pharmaceutical composition containing at least one compound of formula (I) according to any one of Claims 1 to 4, in the form of the base or of a
25 pharmaceutically acceptable acid-addition salt, hydrate or solvate, and optionally one or more pharmaceutically acceptable excipients.

11. Use of a compound of formula (I)
according to any one of Claims 1 to 4, in the form of
the base or of a pharmaceutically acceptable acid-
addition salt, hydrate or solvate, for the preparation
5 of a medicament for preventing or treating a pathology
in which the endogenous cannabinoids and/or any other
substrate metabolized by the enzyme FAAH are involved.

12. Use of a compound of formula (I)
according to any one of Claims 1 to 4, in the form of
10 the base or of a pharmaceutically acceptable acid-
addition salt, hydrate or solvate, for the preparation
of a medicament for preventing or treating acute or
chronic pain, vertigo, vomiting, nausea, eating
disorders, neurological and psychiatric pathologies,
15 acute or chronic neurodegenerative diseases, epilepsy,
sleeping disorders, cardiovascular diseases, renal
ischaemia, cancers, immune system disorders, allergic
diseases, parasitic, viral or bacterial infectious
diseases, inflammatory diseases, osteoporosis, ocular
20 complaints, pulmonary complaints, gastrointestinal
diseases or urinary incontinence.