


clinical trials in similar patient population for ALN-RSV01 are on-going
(www.clinicaltrials.gov).

Nevertheless, safe and effective treatment for RSV disease is needed urgently.

It has been found that the compounds of the present ~~invention~~disclosure belong to a new
chemical class of RSV inhibitors for the treatment or prophylaxis of RSV infection. The
compounds of the ~~invention~~disclosure are therefore useful in the treatment or prophylaxis of
RSV disease.

In a first aspect, there is provided a compound of formula (I)



Wherein

R¹ is hydrogen, halogen, or C₁₋₆alkyl;

R² is hydrogen, halogen, or C₁₋₆alkyl;

R³ is hydrogen, halogen, or C₁₋₆alkyl;

R⁴ is hydrogen, or C₁₋₆alkyl;

R⁵ is hydrogen, or halogen;

R⁶ is hydrogen, halogen, hydroxy, C₁₋₆alkoxy, carboxy, morpholinyl, or 4-C₀-alkylpiperazin-1-yl;

R⁷ is hydrogen, halogen, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylaminocarbonyl, diC₁₋₆alkylaminocarbonyl, C₁₋₆alkylsulfonyl, phenoxy, or hydroxy(CH₂)₂₋₆-O-;

R⁸ is hydrogen, halogen, or C₁₋₆alkoxy;

R⁹ is hydrogen, C₁₋₆alkyl, or =O;

R¹⁰ is hydrogen, or =O, provided that R⁹ and R¹⁰ are not =O simultaneously;

A is nitrogen, or -C-R¹¹, wherein R¹¹ is hydrogen, halogen, C₁₋₆alkyl, cycloalkyl, C₁₋₆alkoxy, trifluoromethyl, trifluoromethoxy, pyridinyloxy, C₁₋₆alkoxy(CH₂)₁₋₆-O-,

difluoromethoxy, cyano, nitro, amino, vinyl, acetylenyl, aminocarbonyl, hydroxy(CH₂)₂₋₆-O-, C₁₋₆alkylsulfinyl, C₁₋₆alkylsulfonyl, hydroxy(CH₂)₁₋₆, deuteratedC₁₋₆alkyl, carboxyl, C₁₋₆alkoxycarbonyl, hydroxy, difluoromethyl, -CH(hydroxy) C₁₋₆alkyl, or C₁₋₆alkylsulfanyl;

X is -CH₂-, -O-, -NH-, -CF₂-, -C(C₁₋₆alkyl)(OH)-, -S-, -C(=O)-, -C(=NOC₀₋₆alkyl)-, -S(=O)-, -S(O₂)- or -S(=O)(NH)-;

Y is -CH-, or nitrogen;

Q is hydrogen; halogen; C₁₋₆alkyl, unsubstituted or once or twice substituted by amino or hydroxy, provided that di-substitution is not on the same carbon; amino(CH₂)₂₋₆aminosulfonyl; 2-amino-4,5-dihydro-1,3-oxazol-4-yl(CH₂)₁₋₆; carboxy(CH₂)₁₋₆; phenylsulfonyl; piperidin-4-yl-carbonyl; 1*H*-pyrazol-3-yl; pyrrolidin-3-yl; piperidin-4-yl; amino(CH₂)₂₋₆-O-; or

NR¹²R¹³, wherein one of R¹² and R¹³ is hydrogen, C₁₋₆alkyl, or hydroxy(CH₂)₂₋₆, and the other one is {1-[amino(CH₂)₀₋₆]-3,3-difluorocyclobutyl}(CH₂)₁₋₆; guanidino(CH₂)₂₋₆; (*S*-C₁₋₆alkylsulfonimidoyl)(CH₂)₂₋₆; 2-oxa-6-aza-spiro[3.4]oct-8-yl; {3-[amino(CH₂)₀₋₆]tetrahydrofuran-3-yl}(CH₂)₁₋₆; 3-aminomethyl-1,1-dioxidothietan-3-ylmethyl; 3-amino-1,1-dioxidothietan-3-ylmethyl; 3-(aminomethyl)thietan-3-ylmethyl; (1,1-dioxidothiomorpholin-4-yl)ethyl; C₀₋₆alkyl(oxetanyl)N(CH₂)₂₋₆; 4,5-dihydro-1*H*-imidazol-2-yl; amino(CH₂)₂₋₆-O-(CH₂)₂₋₆; amino(CH₂)₂₋₁₀; amino(CH₂)₁₋₆difluoromethyl(CH₂)₁₋₆; amino(CH₂)₁₋₆difluoromethyldifluoromethyl(CH₂)₁₋₆; amino(CH₂)₁₋₆fluoromethyl(CH₂)₁₋₆; amino(CH₂)₁₋₆oxetanyl(CH₂)₀₋₆; amino(CH₂)₀₋₆oxetanyl(CH₂)₁₋₆; amino(CH₂)₂₋₆sulfanyl(CH₂)₂₋₆; amino(CH₂)₂₋₆sulfonyl(CH₂)₂₋₆; amino(CH₂)₀₋₆carbonyl(CH₂)₀₋₆; aminocycloalkyl(CH₂)₀₋₆; 2-aminodihydrooxazol-4-yl(CH₂)₁₋₆; 2-aminodihydrooxazol-5-yl(CH₂)₁₋₆; (2-amino-5-methyl-4,5-dihydro-1,3-oxazol-5-yl)methyl; aminophenyl; 4-aminotetrahydropyran-4-yl(CH₂)₁₋₆; azetidin-2-yl(CH₂)₁₋₆; azetidin-3-yl(CH₂)₀₋₆; azetidiny carbonyl; C₁₋₆alkoxy(CH₂)₂₋₆; C₁₋₆alkoxy(CH₂)₂₋₆amino(CH₂)₂₋₆; C₁₋₆alkyl; C₁₋₆alkylamino(CH₂)₂₋₆; C₁₋₆alkylaminocarbonyl(CH₂)₀₋₆; C₁₋₆alkylaminooxetanyl(CH₂)₁₋₆; C₁₋₆alkylcarbonyl; C₁₋₆alkylcarbonylamino(CH₂)₂₋₆; C₁₋₆alkylcarbonylamino(CH₂)₁₋₆oxetanyl(CH₂)₀₋₆; C₁₋₆alkylsulfinyl(CH₂)₂₋₆; C₁₋₆alkylsulfonyl; carboxy(CH₂)₁₋₆; cyano(CH₂)₁₋₆; diC₁₋₆alkylamino(CH₂)₂₋₆; diC₁₋₆alkylaminocarbonyl; difluoromethyl(CH₂)₁₋₆amino(CH₂)₂₋₆; hydrogen; hydroxy(CH₂)₂₋₁₀; hydroxy(CH₂)₂₋₆amino(CH₂)₂₋₆; hydroxy(CH₂)₁₋₆carbonyl; hydroxy(CH₂)₀₋₆oxetanyl(CH₂)₁₋₆; hydroxy(CH₂)₁₋₆oxetanyl(CH₂)₀₋₆; hydroxycycloalkyl; isoxazolyl; morpholin-2-yl(CH₂)₁₋₆; morpholin-4-yl(CH₂)₂₋₆; oxetanyl(CH₂)₀₋₆; *N*-oxetanylpiperidin-3-yl; oxo-piperidinylcarbonyl; phenylaminocarbonyl; phenyl(CH₂)₀₋₆aminooxetanyl(CH₂)₁₋₆; phenylcarbonyl; piperazinyl(CH₂)₂₋₆; piperidin-1-yl(CH₂)₂₋₆; piperidin-2-yl(CH₂)₁₋₆; piperidin-3-yl(CH₂)₀₋₆; piperidin-4-yl(CH₂)₀₋₆; piperidinylcarbonyl; pyrazinylcarbonyl; pyrazol-3-yl;

pyridazinylcarbonyl; pyridinyl(CH₂)₀₋₆carbonyl; pyridinylamino(CH₂)₂₋₆; pyrrolidin-3-yl, unsubstituted or 4-substituted by halogen; pyrrolidin-4-yl, unsubstituted or 3-substituted by hydroxy or C₁₋₆alkoxy; pyrrolidin-2-yl(CH₂)₁₋₆; pyrrolidinylcarbonyl; tetrahydrofuran-3-yl;

tetrahydropyran-4-yl; tetrazolyl(CH₂)₂₋₆; trifluoromethylcarbonylamino(CH₂)₁₋₆oxetanyl;

trifluoromethylsulfonyl; $-(CH_2)_{1-6}-\overset{\overset{R^{14}}{\diagup}}{\underset{\underset{R^{16}}{\diagdown}}{C}}-R^{15}$, wherein R¹⁴ is hydrogen, C₁₋₆alkyl or hydroxy(CH₂)₁₋₆; R¹⁵ is hydroxy, C₁₋₆alkyl, hydroxy(CH₂)₁₋₆ or amino; and R¹⁶ is C₁₋₆alkyl, trifluoromethyl, hydroxy(CH₂)₁₋₆, amino(CH₂)₁₋₆, aminocarboxy or carboxy(CH₂)₁₋₆;

$\overset{\overset{R^{17}}{\diagup}}{\underset{\underset{R^{19}}{\diagdown}}{C}}-R^{18}$, wherein R¹⁷ is hydrogen, C₁₋₆alkyl or hydroxy(CH₂)₁₋₆; R¹⁸ is hydroxy(CH₂)₁₋₆ or C₁₋₆alkyl; R¹⁹ is hydroxy(CH₂)₁₋₆, amino(CH₂)₁₋₆, carboxy or aminocarboxy(CH₂)₀₋₆;

or $\overset{\overset{O}{\parallel}}{\underset{\underset{R^{22}}{\diagdown}}{C}}-\overset{\overset{R^{20}}{\diagup}}{C}-R^{21}$, wherein R²⁰ is hydrogen or C₁₋₆alkyl; R²¹ is C₁₋₆alkyl; R²² is C₁₋₆alkoxy or amino;

R¹² and R¹³, with the nitrogen atom to which they are attached, may form a pyrrolidinyl, piperazinyl, piperidinyl, morpholinyl, azetidiny, diazepanyl or oxopyrrolidinyl ring; which may be unsubstituted, once or twice substituted by a group selected from halogen, C₁₋₆alkyl, C₁₋₆alkoxy, gemdimethyl, amino, aminocarbonyl, hydroxy, oxetanylamino, C₁₋₆alkylpiperazinyl, and amino(CH₂)₁₋₆;

R¹² and R¹³, with the nitrogen atom to which they are attached may form a bridge ring or a spiral ring selected from 2-oxa-6-aza-spiro[3.4]octan-6-yl, 2-oxa-5,7-diazaspiro[3.4]octan-6-one-5-yl, (4aS,7aR)-hexahydropyrrolo[3,4-b][1,4]oxazin-6(2H)-yl, 4,5,6,6a-tetrahydro-3aH-pyrrolo[3,4-d][1,3]oxazol-5-yl, 2-aza-bicyclo[2.1.1]hexan-2-yl, and 3-aza-bicyclo[3.1.0]hexan-3-yl; which may be unsubstituted or further substituted by amino;

and pharmaceutically acceptable salt and stereoisomers thereof.

In a second aspect, there is provided a compound as defined above for use as a therapeutically active substance.

In a third aspect, there is provided a pharmaceutical composition comprising a compound as defined above and a therapeutically inert carrier.

In a fourth aspect, there is provided the use of a compound as defined above for the preparation of a medicament for the treatment or prophylaxis of respiratory syncytial virus infection.

In a fifth aspect, there is provided a compound as defined above for the treatment or prophylaxis of respiratory syncytial virus infection.

As used herein, the term "C₀₋₆alkyl" alone or in combination signifies a chemical bond, or hydrogen, or saturated, linear- or branched chain alkyl group containing 1 to 6, preferably 1 to 4 carbon atoms, for example methyl, ethyl, propyl, isopropyl, 1-butyl, 2-butyl, *tert*-butyl and the like. Preferred "C₀₋₆alkyl" groups are chemical bond, hydrogen, methyl, ethyl, isopropyl, *tert*-butyl.

As used herein, the term "C₁₋₆alkyl" alone or in combination signifies a saturated, linear- or branched chain alkyl group containing 1 to 6, preferably 1 to 4 carbon atoms, for example methyl, ethyl, propyl, isopropyl, 1-butyl, 2-butyl, *tert*-butyl and the like. Preferred "C₁₋₆alkyl" groups are methyl, ethyl, isopropyl, *tert*-butyl.

As used herein, the term "C₂₋₆alkyl" alone or in combination signifies a saturated, linear- or branched chain alkyl group containing 2 to 6, preferably 2 to 4 carbon atoms, for example ethyl, propyl, isopropyl, 1-butyl, 2-butyl, *tert*-butyl and the like. Preferred "C₂₋₆alkyl" groups are ethyl, isopropyl, *tert*-butyl.

As used herein, the term "-(CH₂)₀₋₆-" signifies a chemical link, hydrogen, or a saturated, linear alkyl chain containing from 1 to 6 carbon atoms, preferably, the term signifies hydrogen or -(CH₂)₁₋₄.

As used herein, the term "-(CH₂)₁₋₆-" signifies a saturated, linear alkyl chain containing from 1 to 6 carbon atoms, preferably from 1 to 4 carbon atoms.

As used herein, the term "-(CH₂)₂₋₆-" signifies a saturated, linear alkyl chain containing from 2 to 6 carbon atoms, preferably from 2 to 4 carbon atoms.

As used herein, the term "-(CH₂)₂₋₁₀-" signifies a saturated, linear alkyl chain containing from 2 to 10 carbon atoms, preferably from 2 to 4 carbon atoms.

The term "cycloalkyl", alone or in combination, refers to a saturated carbon ring containing from 3 to 7 carbon atoms, preferably from 3 to 6 carbon atoms, for example,

(1,3,4,5-tetrahydro-2*H*-2-benzazepin-2-yl)quinolin-4-yl]propan-1-amine; [2-(1,3,4,5-tetrahydro-2*H*-2-benzazepin-2-yl)quinolin-4-yl]methanol; 2-(6-chloroquinolin-2-yl)-2,3,4,5-tetrahydro-1*H*-2-benzazepine; 3-[6-chloro-2-(1,3,4,5-tetrahydro-2*H*-2-benzazepin-2-yl)quinolin-4-yl]propane-1,2-diol; (4*S*)-4-{2-[2-(1,3,4,5-tetrahydro-2*H*-2-benzazepin-2-yl)quinolin-4-yl]ethyl}-4,5-dihydro-1,3-oxazol-2-amine; *N*-(2-aminoethyl)-2-(1,3,4,5-tetrahydro-2*H*-2-benzazepin-2-yl)quinoline-4-sulfonamide trifluoroacetate; 4-{4-[(2-aminoethyl)amino]-6-methylquinolin-2-yl}-1,3,4,5-tetrahydro-2*H*-1,4-benzodiazepin-2-one; *N*-[6-methyl-2-(1,2,3,5-tetrahydro-4*H*-1,4-benzodiazepin-4-yl)quinolin-4-yl]ethane-1,2-diamine; *N*-[2-(2,3-dihydro-1,4-benzoxazepin-4(5*H*)-yl)quinolin-4-yl]ethane-1,2-diamine; *N*-[(3-aminooxetan-3-yl)methyl]-2-[(5*E*)-5-(methoxyimino)-1,3,4,5-tetrahydro-2*H*-2-benzazepin-2-yl]-6-methylquinolin-4-amine and 2-(4-{[(3-aminooxetan-3-yl)methyl]amino}-6-methylquinazolin-2-yl)-1,2,3,4-tetrahydro-5*H*-2-benzazepin-5-one.

16. A compound according to any one of claims 1 to 15 for use as a therapeutically active substance.

17. A pharmaceutical composition comprising a compound in accordance with any one of claims 1 to 15 and a therapeutically inert carrier.

~~18. The use of a compound according to any one of claims 1 to 15 for the treatment or prophylaxis of respiratory syncytial virus infection.~~

~~18~~19. The use of a compound according to any one of claims 1 to 15 for the preparation of a medicament for the treatment or prophylaxis of respiratory syncytial virus infection.

~~19~~20. A compound according to any one of claims 1 to 15 for the treatment or prophylaxis of respiratory syncytial virus infection.

~~21. A method for the treatment or prophylaxis of respiratory syncytial virus infection, which method comprises administering an effective amount of a compound as defined in any one of claims 1 to 15.~~
