

TITLE

PYRIMIDINYLOXY BENZENE DERIVATIVES AS HERBICIDES

FIELD OF THE INVENTION

This invention relates to certain pyrimidinyloxy benzene derivatives, their *N*-oxides, salts and compositions, and methods of their use for controlling undesirable vegetation.


BACKGROUND OF THE INVENTION

The control of undesired vegetation is extremely important in achieving high crop efficiency. Achievement of selective control of the growth of weeds especially in such useful crops as rice, soybean, sugar beet, maize, potato, wheat, barley, tomato and plantation crops, among others, is very desirable. Unchecked weed growth in such useful crops can cause significant reduction in productivity and thereby result in increased costs to the consumer. The control of undesired vegetation in noncrop areas is also important. Many products are commercially available for these purposes, but the need continues for new compounds that are more effective, less costly, less toxic, environmentally safer or have different sites of action.

JP 61236766 A (Sumitomo, 1986) discloses certain carbon-linked pyrimidinyloxy benzene derivatives as herbicides. WO 94/17059 (Nippon Soda, 1994) discloses certain carbon linked pyrimidinyloxy benzene derivative as herbicides.

SUMMARY OF THE INVENTION

This invention is directed to compounds of Formula 1 (including all stereoisomers), (*N*-oxides, and salts thereof), agricultural compositions containing them and their use as herbicides:



Q is a 5- or 6-membered aromatic heterocyclic ring, bound to the remainder of

Formula 1 through a carbon atom, and optionally substituted with 1 to 4 R¹;

Z is O or S;

each R¹ is independently halogen, cyano, nitro, SF₅, CHO, C(=O)NH₂, C(=S)NH₂,

SO₂NH₂, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄

haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₄-C₈

alkylcycloalkyl, C₄-C₈ cycloalkylalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆

- haloalkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₃-C₇ cycloalkylcarbonyl, C₂-C₈ alkylaminocarbonyl, C₃-C₁₀ dialkylaminocarbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₃-C₆ cycloalkoxy, C₃-C₆ halocycloalkoxy, C₄-C₈ cycloalkylalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ alkoxyhaloalkyl, C₂-C₆ alkoxyalkoxy, C₂-C₄ alkylcarbonyloxy, C₂-C₆ cyanoalkyl, C₂-C₆ cyanoalkoxy, C₁-C₄ hydroxyalkyl, C₂-C₄ alkylthioalkyl, SO_nR^{1A}, Si(CH₃)₃ or B(-OC(R^{1B})₂C(R^{1B})₂O-); or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{1C}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{1C} on carbon atom ring members and R^{1D} on nitrogen atom ring members;
- R² is halogen, cyano, nitro, C₁-C₄ alkoxy, C₁-C₄ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, SO_nR^{2A}, C₁-C₄ haloalkyl or C₃-C₆ cycloalkyl;
- each R³ is independently halogen, cyano, hydroxy, nitro, amino, CHO, C(=O)NH₂, C(=S)NH₂, SO₂NH₂, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₄-C₈ alkylcycloalkyl, C₄-C₈ cycloalkylalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₃-C₇ cycloalkylcarbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₃-C₆ cycloalkoxy, C₃-C₆ halocycloalkoxy, C₄-C₈ cycloalkylalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ alkoxyhaloalkyl, C₂-C₆ alkoxyalkoxy, C₂-C₄ alkylcarbonyloxy, C₂-C₆ cyanoalkyl, C₂-C₆ cyanoalkoxy, C₂-C₄ alkylthioalkyl, Si(CH₃)₃, C≡CSi(CH₃)₃, C(=O)N(R^{3A})(R^{3B}), C(=NOR^{3C})H, C(=NR^{3D})H, SO_nR^{3E}; or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{3F}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members; or pyrimidinyloxy;
- m is 0, 1, 2 or 3;
- each n is independently 0, 1 or 2;
- each R^{1A}, R^{2A} and R^{3E} is independently C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkylamino or C₂-C₆ dialkylamino;

each R^{1B} is independently H or C₁-C₄ alkyl;

each R^{1C} is independently hydroxy, halogen, cyano, nitro, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₁-C₆ haloalkoxy;

each R^{1D} is independently cyano, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₂-C₆ alkylcarbonyl;

each R^{3A} is independently C₁-C₄ alkyl or C₁-C₄ haloalkyl;

each R^{3B} is independently H, C₁-C₄ alkyl or C₁-C₄ haloalkyl;

each R^{3C} is independently H or C₁-C₄ alkyl;

each R^{3D} is independently H, amino, C₁-C₄ alkyl or C₁-C₄ alkylamino;

each R^{3F} is independently hydroxy, halogen, cyano, nitro, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₁-C₆ haloalkoxy; and

each R^{3G} is independently cyano, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₂-C₆ alkylcarbonyl;

More particularly, this invention pertains to a compound of Formula **1** (including all stereoisomers), an *N*-oxide or a salt thereof. This invention also relates to a herbicidal composition comprising a compound of the invention (i.e. in a herbicidally effective amount) and at least one component selected from the group consisting of surfactants, solid diluents and liquid diluents. This invention further relates to a method for controlling the growth of undesired vegetation comprising contacting the vegetation or its environment with a herbicidally effective amount of a compound of the invention (e.g., as a composition described herein).

This invention also includes a herbicidal mixture comprising (a) a compound selected from Formula **1**, *N*-oxides, and salts thereof, and (b) at least one additional active ingredient selected from (b1) through (b16); and salts of compounds of (b1) through (b16).

DETAILS OF THE INVENTION

As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains,” “containing,” “characterized by” or any other variation thereof, are intended to cover a non-exclusive inclusion, subject to any limitation explicitly indicated. For example, a composition, mixture, process or method that comprises a list of elements is not necessarily limited to only those elements but may include other elements not expressly listed or inherent to such composition, mixture, process or method.

The transitional phrase “consisting of” excludes any element, step, or ingredient not specified. If in the claim, such would close the claim to the inclusion of materials other than those recited except for impurities ordinarily associated therewith. When the phrase “consisting of” appears in a clause of the body of a claim, rather than immediately following the preamble, it limits only the element set forth in that clause; other elements are not excluded from the claim as a whole.

The transitional phrase “consisting essentially of” is used to define a composition, process or method that includes materials, steps, features, components, or elements, in addition to those literally disclosed, provided that these additional materials, steps, features, components, or elements do not materially affect the basic and novel characteristic(s) of the claimed invention. The term “consisting essentially of” occupies a middle ground between “comprising” and “consisting of”.

Where applicants have defined an invention or a portion thereof with an open-ended term such as “comprising,” it should be readily understood that (unless otherwise stated) the description should be interpreted to also describe such an invention using the terms “consisting essentially of” or “consisting of.”

Further, unless expressly stated to the contrary, “or” refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B are true (or present).

Also, the indefinite articles “a” and “an” preceding an element or component of the invention are intended to be nonrestrictive regarding the number of instances (i.e. occurrences) of the element or component. Therefore “a” or “an” should be read to include one or at least one, and the singular word form of the element or component also includes the plural unless the number is obviously meant to be singular.

As referred to herein, the term “seedling”, used either alone or in a combination of words means a young plant developing from the embryo of a seed.

As referred to herein, the term “broadleaf” used either alone or in words such as “broadleaf weed” means dicot or dicotyledon, a term used to describe a group of angiosperms characterized by embryos having two cotyledons. As used herein, the term “alkylating agent” refers to a chemical compound in which a carbon-containing radical is bound through a carbon atom to a leaving group such as halide or sulfonate, which is displaceable by bonding of a nucleophile to said carbon atom. Unless otherwise indicated, the term “alkylating” does not limit the carbon-containing radical to alkyl; the carbon-containing radicals in alkylating agents include the variety of carbon-bound substituent radicals specified for Q, R¹ and R³.

In the above recitations, the term “alkyl”, used either alone or in compound words such as “alkylthio” or “haloalkyl” includes straight-chain or branched alkyl, such as, methyl, ethyl, *n*-propyl, *i*-propyl, or the different butyl, pentyl or hexyl isomers. “Alkenyl” includes straight-chain or branched alkenes such as ethenyl, 1-propenyl, 2-propenyl, and the different butenyl, pentenyl and hexenyl isomers. “Alkenyl” also includes polyenes such as 1,2-propadienyl and 2,4-hexadienyl. “Alkynyl” includes straight-chain or branched alkynes such as ethynyl, 1-propynyl, 2-propynyl and the different butynyl, pentynyl and hexynyl isomers.

“Alkoxy” includes, for example, methoxy, ethoxy, *n*-propyloxy, isopropyloxy and the different butoxy, pentoxy and hexyloxy isomers. “Alkoxyalkyl” denotes alkoxy substitution on alkyl. Examples of “alkoxyalkyl” include CH_3OCH_2 , $\text{CH}_3\text{OCH}_2\text{CH}_2$, $\text{CH}_3\text{CH}_2\text{OCH}_2$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OCH}_2$ and $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_2$. “Alkenyloxy” includes straight-chain or branched alkenyloxy moieties. Examples of “alkenyloxy” include $\text{H}_2\text{C}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)\text{CH}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)\text{CH}=\text{C}(\text{CH}_3)\text{CH}_2\text{O}$ and $\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{O}$. “Alkynyloxy” includes straight-chain or branched alkynyloxy moieties. Examples of “alkynyloxy” include $\text{HC}\equiv\text{CCH}_2\text{O}$, $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{O}$ and $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{CH}_2\text{O}$. “Alkylthio” includes branched or straight-chain alkylthio moieties such as methylthio, ethylthio, and the different propylthio, butylthio, pentylthio and hexylthio isomers. “Alkylthioalkyl” denotes alkylthio substitution on alkyl. Examples of “alkylthioalkyl” include CH_3SCH_2 , $\text{CH}_3\text{SCH}_2\text{CH}_2$, $\text{CH}_3\text{CH}_2\text{SCH}_2$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{SCH}_2$ and $\text{CH}_3\text{CH}_2\text{SCH}_2\text{CH}_2$. “Alkylthioalkoxy” denotes alkylthio substitution on alkoxy. “Cyanoalkyl” denotes an alkyl group substituted with one cyano group. Examples of “cyanoalkyl” include NCCH_2 , NCCH_2CH_2 and $\text{CH}_3\text{CH}(\text{CN})\text{CH}_2$. “Alkylamino”, “dialkylamino”, and the like, are defined analogously to the above examples.

“Cycloalkyl” includes, for example, cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The term “halogen”, either alone or in compound words such as “haloalkyl”, or when used in descriptions such as “alkyl substituted with halogen” includes fluorine, chlorine, bromine or iodine. Further, when used in compound words such as “haloalkyl”, or when used in descriptions such as “alkyl substituted with halogen” said alkyl may be partially or fully substituted with halogen atoms which may be the same or different. Examples of “haloalkyl” or “alkyl substituted with halogen” include F_3C , ClCH_2 , CF_3CH_2 and CF_3CCl_2 . The terms “haloalkoxy”, and the like, is defined analogously to the term “haloalkyl”. Examples of “haloalkoxy” include $\text{CF}_3\text{O}-$, $\text{CCl}_3\text{CH}_2\text{O}-$, $\text{HCF}_2\text{CH}_2\text{CH}_2\text{O}-$ and $\text{CF}_3\text{CH}_2\text{O}-$. “Alkylcarbonyl” denotes a straight-chain or branched alkyl moieties bonded to a $\text{C}(=\text{O})$ moiety. Examples of “alkylcarbonyl” include $\text{CH}_3\text{C}(=\text{O})-$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{C}(=\text{O})-$ and $(\text{CH}_3)_2\text{CHC}(=\text{O})-$. Examples of “alkoxycarbonyl” include $\text{CH}_3\text{OC}(=\text{O})-$, $\text{CH}_3\text{CH}_2\text{OC}(=\text{O})-$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OC}(=\text{O})-$, $(\text{CH}_3)_2\text{CHOC}(=\text{O})-$ and the different butoxy- or pentoxy carbonyl isomers.

The total number of carbon atoms in a substituent group is indicated by the “ $\text{C}_i\text{-C}_j$ ” prefix where *i* and *j* are numbers from 1 to 6. For example, $\text{C}_1\text{-C}_4$ alkylsulfonyl designates methylsulfonyl through butylsulfonyl; C_2 alkoxyalkyl designates CH_3OCH_2- ; C_3 alkoxyalkyl designates, for example, $\text{CH}_3\text{CH}(\text{OCH}_3)-$, $\text{CH}_3\text{OCH}_2\text{CH}_2-$ or $\text{CH}_3\text{CH}_2\text{OCH}_2-$; and C_4 alkoxyalkyl designates the various isomers of an alkyl group substituted with an alkoxy group containing a total of four carbon atoms, examples including $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2-$ and $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_2-$.

When a compound is substituted with a substituent bearing a subscript that indicates the number of said substituents can exceed 1, said substituents (when they exceed 1) are independently selected from the group of defined substituents, (e.g., $(R^3)_n$, n is 0, 1, 2 or 3). Further, when the subscript indicates a range, e.g. $(R)_{i-j}$, then the number of substituents may be selected from the integers between i and j inclusive. When a group contains a substituent which can be hydrogen, for example (when $m = 0$), then when this substituent is taken as hydrogen, it is recognized that this is equivalent to said group being unsubstituted. When a variable group is shown to be optionally attached to a position, (for example $(R^1)_n$ attached to Q wherein n may be 0, then hydrogen may be at the position even if not recited in the variable group definition. When one or more positions on a group are said to be “not substituted” or “unsubstituted”, then hydrogen atoms are attached to take up any free valency.

Unless otherwise indicated, a “ring” as a component of Formula 1 (e.g., substituent Q) is carbocyclic or heterocyclic. The term “ring member” refers to an atom or heteroatom forming the backbone of a ring. When a fully unsaturated carbocyclic ring satisfies Hückel’s rule, then said ring is also called an “aromatic ring”. “Saturated carbocyclic” refers to a ring having a backbone consisting of carbon atoms linked to one another by single bonds; unless otherwise specified, the remaining carbon valences are occupied by hydrogen atoms.

The terms “heterocyclic ring”, “heterocycle” denote a ring in which at least one atom forming the ring backbone is not carbon, e.g., nitrogen, oxygen or sulfur. Typically a heterocyclic ring contains no more than 4 nitrogens, no more than 2 oxygens and no more than 2 sulfurs. Unless otherwise indicated, a heterocyclic ring can be a saturated, partially unsaturated, or fully unsaturated ring. When a fully unsaturated heterocyclic ring satisfies Hückel’s rule, then said ring is also called a “heteroaromatic ring” or “aromatic heterocyclic ring”. Unless otherwise indicated, heterocyclic rings can be attached through any available carbon or nitrogen by replacement of a hydrogen on said carbon or nitrogen.


“Aromatic” indicates that each of the ring atoms is essentially in the same plane and has a p -orbital perpendicular to the ring plane, and that $(4n + 2) \pi$ electrons, where n is a positive integer, are associated with the ring to comply with Hückel’s rule.


The term “optionally substituted” in connection with the heterocyclic rings refers to groups which are unsubstituted or have at least one non-hydrogen substituent that does not extinguish the biological activity possessed by the unsubstituted analog. As used herein, the following definitions shall apply unless otherwise indicated. The term “optionally substituted” is used interchangeably with the phrase “substituted or unsubstituted” or with the term “(un)substituted.” Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and each substitution is independent of the other.


When Q is a 5- or 6-membered (nitrogen-containing) heterocyclic ring, it may be attached to the remainder of Formula 1 through any available carbon or nitrogen ring atom, unless otherwise described. As noted above, Q can be (among others) phenyl optionally substituted with one or more substituents selected from a group of substituents as defined in the Summary of the Invention. An example of phenyl optionally substituted with one to five substituents is the ring illustrated as U-1 in Exhibit 1, wherein R^V is R^1 as defined in the Summary of the Invention for Q and r is an integer (from 0 to 4).

As noted above, Q can be (among others) 5- or 6-membered aromatic heterocyclic ring, which may be saturated or unsaturated, optionally substituted with one or more substituents selected from a group of substituents as defined in the Summary of the Invention. Examples of a 5- or 6-membered unsaturated aromatic heterocyclic ring optionally substituted with from one or more substituents include the rings U-2 through U-61 illustrated in Exhibit 1 wherein R^V is any substituent as defined in the Summary of the Invention for Q (i.e. R^1) and r is an integer from 0 to 4, limited by the number of available positions on each U group. As U-29, U-30, U-36, U-37, U-38, U-39, U-40, U-41, U-42 and U-43 have only one available position, for these U groups r is limited to the integers 0 or 1, and r being 0 means that the U group is unsubstituted and a hydrogen is present at the position indicated by $(R^V)_r$.

Exhibit 1







Although R^v groups are shown in the structures U-1 through U-61, it is noted that they do not need to be present since they are optional substituents. Note that when R^v is H when attached to an atom, this is the same as if said atom is unsubstituted. The nitrogen atoms that require substitution to fill their valence are substituted with H or R^v . Note that when the attachment point between $(R^v)_r$ and the U group is illustrated as floating, $(R^v)_r$ can be attached to any available carbon atom or nitrogen atom of the U group. Note that when the attachment point on the U group is illustrated as floating, the U group can be attached to the remainder of Formula 1 through any available carbon or nitrogen of the U group by replacement of a hydrogen atom. Note that some U groups can only be substituted with less than 4 R^v groups (e.g., U-2 through U-47 and U-52 through U-61).

A wide variety of synthetic methods are known in the art to enable preparation of aromatic and nonaromatic heterocyclic rings and ring systems; for extensive reviews see the eight volume set of *Comprehensive Heterocyclic Chemistry*, A. R. Katritzky and C. W. Rees editors-in-chief, Pergamon Press, Oxford, 1984 and the twelve volume set of *Comprehensive Heterocyclic Chemistry II*, A. R. Katritzky, C. W. Rees and E. F. V. Scriven editors-in-chief, Pergamon Press, Oxford, 1996.

Compounds of this invention can exist as one or more stereoisomers. The various stereoisomers include enantiomers, diastereomers, atropisomers and geometric isomers. Stereoisomers are isomers of identical constitution but differing in the arrangement of their atoms in space and include enantiomers, diastereomers, cis-trans isomers (also known as geometric isomers) and atropisomers. Atropisomers result from restricted rotation about single bonds where the rotational barrier is high enough to permit isolation of the isomeric species. One skilled in the art will appreciate that one stereoisomer may be more active and/or may exhibit beneficial effects when enriched relative to the other stereoisomer(s) or when separated from the other stereoisomer(s). Additionally, the skilled artisan knows how to separate, enrich, and/or to selectively prepare said stereoisomers. The compounds of the invention may be present as a mixture of stereoisomers, individual stereoisomers or as an optically active form.

Compounds of Formula 1 typically exist in more than one form, and Formula 1 thus include all crystalline and non-crystalline forms of the compounds they represent. Non-crystalline forms include embodiments which are solids such as waxes and gums as well as embodiments which are liquids such as solutions and melts. Crystalline forms include

embodiments which represent essentially a single crystal type and embodiments which represent a mixture of polymorphs (i.e. different crystalline types). The term “polymorph” refers to a particular crystalline form of a chemical compound that can crystallize in different crystalline forms, these forms having different arrangements and/or conformations of the molecules in the crystal lattice. Although polymorphs can have the same chemical composition, they can also differ in composition due the presence or absence of co-crystallized water or other molecules, which can be weakly or strongly bound in the lattice. Polymorphs can differ in such chemical, physical and biological properties as crystal shape, density, hardness, color, chemical stability, melting point, hygroscopicity, suspensibility, dissolution rate and biological availability. One skilled in the art will appreciate that a polymorph of a compound of Formula 1 can exhibit beneficial effects (e.g., suitability for preparation of useful formulations, improved biological performance) relative to another polymorph or a mixture of polymorphs of the same compound of Formula 1. Preparation and isolation of a particular polymorph of a compound of Formula 1 can be achieved by methods known to those skilled in the art including, for example, crystallization using selected solvents and temperatures. For a comprehensive discussion of polymorphism see R. Hilfiker, Ed., *Polymorphism in the Pharmaceutical Industry*, Wiley-VCH, Weinheim, 2006.


One skilled in the art will appreciate that not all nitrogen-containing heterocycles can form *N*-oxides since the nitrogen requires an available lone pair for oxidation to the oxide; one skilled in the art will recognize those nitrogen-containing heterocycles which can form *N*-oxides. One skilled in the art will also recognize that tertiary amines can form *N*-oxides. Synthetic methods for the preparation of *N*-oxides of heterocycles and tertiary amines are very well known by one skilled in the art including the oxidation of heterocycles and tertiary amines with peroxy acids such as peracetic and *m*-chloroperbenzoic acid (MCPBA), hydrogen peroxide, alkyl hydroperoxides such as *t*-butyl hydroperoxide, sodium perborate, and dioxiranes such as dimethyldioxirane. These methods for the preparation of *N*-oxides have been extensively described and reviewed in the literature, see for example: T. L. Gilchrist in *Comprehensive Organic Synthesis*, vol. 7, pp 748–750, S. V. Ley, Ed., Pergamon Press; M. Tisler and B. Stanovnik in *Comprehensive Heterocyclic Chemistry*, vol. 3, pp 18–20, A. J. Boulton and A. McKillop, Eds., Pergamon Press; M. R. Grimmett and B. R. T. Keene in *Advances in Heterocyclic Chemistry*, vol. 43, pp 149–161, A. R. Katritzky, Ed., Academic Press; M. Tisler and B. Stanovnik in *Advances in Heterocyclic Chemistry*, vol. 9, pp 285–291, A. R. Katritzky and A. J. Boulton, Eds., Academic Press; and G. W. H. Cheeseman and E. S. G. Werstiuk in *Advances in Heterocyclic Chemistry*, vol. 22, pp 390–392, A. R. Katritzky and A. J. Boulton, Eds., Academic Press.


One skilled in the art recognizes that because in the environment and under physiological conditions salts of chemical compounds are in equilibrium with their corresponding nonsalt forms, salts share the biological utility of the nonsalt forms. Thus a

wide variety of salts of a compound of Formula 1 are useful for control of undesired vegetation (i.e. are agriculturally suitable). The salts of a compound of Formula 1 include acid-addition salts with inorganic or organic acids such as hydrobromic, hydrochloric, nitric, phosphoric, sulfuric, acetic, butyric, fumaric, lactic, maleic, malonic, oxalic, propionic, salicylic, tartaric, 4-toluenesulfonic or valeric acids. When a compound of Formula 1 contains an acidic moiety such as a carboxylic acid or phenol, salts also include those formed with organic or inorganic bases such as pyridine, triethylamine or ammonia, or amides, hydrides, hydroxides or carbonates of sodium, potassium, lithium, calcium, magnesium or barium. Accordingly, the present invention comprises compounds selected from Formula 1, *N*-oxides and agriculturally suitable salts thereof.

Embodiments of the present invention as described in the Summary of the Invention include (where Formula 1 as used in the following Embodiments includes *N*-oxides and salts thereof):

Embodiment 1. A compound of Formula 1 wherein Q is selected from





wherein r is 0, 1, 2 or 3; and s is 0 or 1.

Embodiment 2. A compound of Embodiment 1 wherein Q is selected from Q-1 through Q-42.

Embodiment 3. A compound of Embodiment 2 wherein Q is selected from Q-7 through Q-24.

Embodiment 4. A compound of Embodiment 3 wherein Q is selected from Q-16 and Q-18.

Embodiment 5. A compound of Embodiment 4 wherein Q is Q-16.

Embodiment 6. A compound of Embodiment 4 wherein Q is Q-18.

Embodiment 7. A compound of Embodiment 1 wherein Q is selected from Q-43 through Q-55.

Embodiment 8. A compound of Embodiment 7 wherein Q is selected from Q-43, Q-44, Q-45, Q-48, Q-49 and Q-50.

Embodiment 9. A compound of Embodiment 8 wherein Q is selected from Q-43, Q-44 and Q-45.

Embodiment 10. A compound of Embodiment 9 wherein Q is Q-43.

Embodiment 11. A compound of Embodiment 10 wherein Q is Q-45.

Embodiment 12. A compound of Formula 1 or any one of Embodiments 1 through 11 either alone or in combination, wherein Z is O.

Embodiment 13. A compound of Formula 1 or any one of Embodiments 1 through 12 either alone or in combination, wherein each R¹ is independently halogen, cyano, SF₅, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ cyanoalkyl, C₁-C₄ hydroxyalkyl, C₂-C₄ alkylthioalkyl or SO_nR^{1A}.

Embodiment 14. A compound of Embodiment 13 wherein each R¹ is independently halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₁-C₄ hydroxyalkyl, C₂-C₄ alkylthioalkyl or SO_nR^{1A}.

Embodiment 15. A compound of Embodiment 14 wherein each R¹ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy or SO_nR^{1A}.

Embodiment 16. A compound of Embodiment 15 wherein each R¹ is independently halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy.

Embodiment 17. A compound of Embodiment 16 wherein each R^1 is independently halogen, C_1 - C_4 haloalkyl or C_1 - C_4 haloalkoxy.

Embodiment 18. A compound of Embodiment 17 wherein each R^1 is independently halogen or C_1 - C_4 haloalkyl.

5 Embodiment 19. A compound of Embodiment 18 wherein each R^1 is independently F, Cl, Br, CF_3 , CHF_2 or CH_2F .

Embodiment 20. A compound of Formula 1 or any one of Embodiments 1 through 19 either alone or in combination, wherein r is 0, 1 or 2.

Embodiment 20a. A compound of Embodiment 20 wherein r is 1.

10 Embodiment 21. A compound of Formula 1 or any one of Embodiments 1 through 19 either alone or in combination, wherein s is 1.

Embodiment 21a. A compound of Formula 1 or any one of Embodiments 1 through 20a either alone or in combination, wherein when Q is Q-16 and r is 1 then R^1 is attached at the 5 position of the Q-16 ring.

15 Embodiment 21b. A compound of Formula 1 or any one of Embodiments 1 through 20a either alone or in combination, wherein when Q is Q-18 and r is 1 then R^1 is attached at the 3 position of the Q-18 ring.

Embodiment 22. A compound of Formula 1 or any one of Embodiments 1 through 21b either alone or in combination, wherein R^2 is halogen, C_1 - C_4 alkyl or C_1 - C_4 haloalkyl.

20

Embodiment 23. A compound of Embodiment 22 wherein R^2 is halogen or C_1 - C_4 alkyl.

Embodiment 24. A compound of Embodiment 23 wherein R^2 is halogen or CH_3 .

Embodiment 25. A compound of Embodiment 24 wherein R^2 is halogen.

25 Embodiment 26. A compound of Embodiment 25 wherein R^2 is F, Cl or Br.

Embodiment 27. A compound of Formula 1 or any one of Embodiments 1 through 26 either alone or in combination, wherein m is 0, 1 or 2.

Embodiment 28. A compound of Embodiment 27 wherein m is 0 or 1.

Embodiment 29. A compound of Embodiment 28 wherein m is 1.

30 Embodiment 30. A compound of Embodiment 27 wherein m is 0 (i.e. the 3-, 4-, 5- and 6-positions are unsubstituted by R^3).

Embodiment 31. A compound of Formula 1 or any one of Embodiments 1 through 30 either alone or in combination, wherein each R^3 is independently halogen, cyano, CHO, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 haloalkyl, C_2 - C_4 haloalkenyl, C_2 - C_4 haloalkynyl, C_3 - C_6 cycloalkyl, C_3 - C_6 halocycloalkyl, C_4 - C_8 alkylcycloalkyl, C_2 - C_6 alkylcarbonyl, C_2 - C_6 haloalkylcarbonyl, C_2 - C_6 alkoxy carbonyl, C_1 - C_4 alkoxy, C_3 - C_4 alkenyloxy, C_3 - C_4 alkynyloxy, C_1 - C_4 haloalkoxy, C_3 - C_4 haloalkenyloxy, C_3 - C_4 haloalkynyloxy, C_3 - C_6 cycloalkoxy,

35

C₃-C₆ halocycloalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₄ alkylcarbonyloxy, C₂-C₆ cyanoalkyl, C(=O)N(R^{3A})(R^{3B}), C(=NOR^{3C})H, SO_nR^{3E}; or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{3F}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members.

Embodiment 32. A compound of Embodiment 31 wherein each R³ is independently halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxycarbonyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ cyanoalkyl, SO_nR^{3E}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members.

Embodiment 33. A compound of Embodiment 32 wherein each R³ is independently halogen, cyano, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxycarbonyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₂-C₆ alkoxyalkyl or C₂-C₆ haloalkoxyalkyl.

Embodiment 34. A compound of Embodiment 33 wherein each R³ is independently halogen, cyano, C₁-C₄ alkyl or C₁-C₄ haloalkyl.

Embodiment 35. A compound of Embodiment 34 wherein each R³ is independently halogen or cyano.

Embodiment 36. A compound of Embodiment 35 wherein each R³ is independently halogen.

Embodiment 37. A compound of Formula 1 or any one of Embodiments 1 through 36 either alone or in combination, wherein R³ is attached to the remainder of Formula 1 at the 3-position.

Embodiment 38. A compound of Formula 1 or any one of Embodiments 1 through 37 either alone or in combination, wherein each R^{1A} is independently C₁-C₄ alkyl or C₁-C₄ haloalkyl.

Embodiment 39. A compound of Embodiment 38 wherein each R^{1A} is independently C₁-C₄ haloalkyl.

Embodiment 40. A compound of Formula 1 or any one of Embodiments 1 through 39 either alone or in combination, wherein each R^{3E} is independently C_1 - C_4 alkyl.

Embodiment 41. A compound of Formula 1 or any one of Embodiments 1 through 40 either alone or in combination, wherein each R^{3A} is independently C_1 - C_4 alkyl.

5 Embodiment 42. A compound of Formula 1 or any one of Embodiments 1 through 41 either alone or in combination, wherein each R^{3B} is independently H or C_1 - C_4 alkyl.

10 Embodiment 43. A compound of Formula 1 or any one of Embodiments 1 through 42 either alone or in combination, wherein each R^{3C} is independently H or C_1 - C_4 alkyl.

Embodiment 44. A compound of Formula 1 or any one of Embodiments 1 through 43 either alone or in combination, wherein each R^{3D} is independently H or C_1 - C_4 alkyl.

15 Embodiment 45. A compound of Formula 1 or any one of Embodiments 1 through 44 either alone or in combination, wherein each n is independently 0 or 2.

Embodiment 46. A compound of Embodiment 45 wherein n is 2.

Embodiment 47. A compound of Embodiment 45 wherein n is 0.


20 Embodiment 48. A compound of Formula 1 or any one of Embodiments 1 through 47 either alone or in combination, provided that i) when Q is 5-chloro-2-pyridinyl; Z is O; and R³ is 4 chloro, then R² is other than Cl or Br; ii) when Q is 4-CF₃-2-pyrimidinyl; Z is O; and m is 0, then R² is other than Cl or Br; and iii) when Q is 6-CF₃-2-pyridinyl; Z is O; and m is 0, then R² is other than Br.


Embodiments of the present invention as described in the Summary of the Invention and Embodiment AAA also include the following:


25 Embodiment 1P. A compound of Formula 1 (including all stereoisomers), *N*-oxides, and salts thereof, agricultural compositions containing them and their use as herbicides as described in the Summary of the Invention.

30 Embodiment 2P. A compound of Embodiment 1 wherein Q is a 5- or 6-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, and optionally substituted with 1 to 3 R^1 .

Embodiment 3P. A compound of Embodiment 2 wherein Q is selected from







r is 0, 1, 2 or 3; and

s is 0 or 1.

Embodiment 4P. A compound of any one of Embodiments 1 through 3 wherein Q is a 5-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, optionally substituted with R^1 , and is selected from Q-1 through Q-41.

Embodiment 5P. A compound of Embodiment 4 wherein Q is selected from Q-7 through Q-24.

Embodiment 6P. A compound of Embodiment 5 wherein Q is selected from Q-9, Q-11, Q-12, Q-16, Q-18, Q-22, Q-23, Q-24 and Q-25.

Embodiment 7P. A compound of Embodiment 6 wherein Q is selected from Q-11, Q-18 and Q-22.

Embodiment 8P. A compound of any one of Embodiments 1 through 3 wherein Q is a 6-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, optionally substituted with R^1 , and is selected from Q-42 through Q-54.


Embodiment 9P. A compound of Embodiment 8 wherein Q is selected from Q-42, Q-43, Q-44, Q-47, Q-48 and Q-49.

Embodiment 10P. A compound of Embodiment 9 wherein Q is selected from Q-42, Q-43, Q-47 and Q-48.

Embodiment 11P. A compound of Embodiment 10 wherein Q is selected from Q-42, Q-47 and Q-48.

Embodiment 12P. A compound of Embodiment 11 wherein Q is selected from Q-42.

Embodiment 13P. A compound of Embodiment 12 wherein Q is



Embodiment 14P. A compound of any one of Embodiments 1 through 3 wherein Q is selected from Q-7 through Q-24, Q-42, Q-43, Q-44, Q-47, Q-48 and Q-49.

Embodiment 15P. A compound of Embodiment 14 wherein Q is selected from Q-9, Q-11, Q-12, Q-16, Q-18, Q-22, Q-23, Q-24, Q-25, Q-42, Q-43, Q-47 and Q-48.

Embodiment 16P. A compound of Embodiment 1 wherein Q is phenyl substituted with 1 to 3 R¹.

Embodiment 17P. A compound of Embodiment 16 wherein Q is phenyl substituted with 1 to 2 R¹.

Embodiment 18P. A compound of Embodiment 17 wherein Q is phenyl substituted with 1 R¹ at the 3- or 4-positions (i.e. *meta* or *para* to the attachment of phenyl to the remainder of Formula 1).

Embodiment 19P. A compound of Embodiment 1 wherein when Q is phenyl substituted with 1 to 3 R¹, m is 1, 2 or 3.

Embodiment 20P. A compound of Embodiment 1 wherein when Q is phenyl substituted with 1 to 3 R¹, m is 1 or 2.

Embodiment 21P. A compound of Embodiment 1 wherein Q is other than phenyl substituted with 1 to 4 R¹.

Embodiment 22P. A compound of any one of Embodiments 1 through 21 wherein R¹ is halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₂-C₄ alkoxyalkyl, C₂-C₄ alkylthioalkyl or SO_nR^{1A}.

Embodiment 23P. A compound of Embodiment 22 wherein R¹ is halogen, cyano, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, or SCF₃.

Embodiment 24P. A compound of Embodiment 23 wherein R¹ is halogen, C₁-C₄ alkyl or C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy.

Embodiment 25P. A compound of Embodiment 24 wherein R¹ is halogen C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy.

Embodiment 26P. A compound of Embodiment 25 wherein R¹ is Cl, Br, CF₃ or OCF₃.

Embodiment 27P. A compound of any one of Embodiments 1 through 22 wherein each n is independently 0, 1 or 2.

Embodiment 28P. A compound of Embodiment 27 wherein each n is independently 0.

Embodiment 29P. A compound of Embodiment 28 wherein each n is independently 2.

5 Embodiment 30P. A compound of any one of Embodiments 1 through 29 wherein R² is halogen, C₁-C₄ alkyl or C₁-C₄ haloalkyl.

Embodiment 31P. A compound of Embodiment 30 wherein R² is halogen or C₁-C₄ alkyl.

Embodiment 32P. A compound of Embodiment 31 wherein R² is halogen or CH₃.

10 Embodiment 33P. A compound of Embodiment 32 wherein R² is halogen.

Embodiment 34P. A compound of Embodiment 33 wherein R² is F, Cl or Br.

Embodiment 35P. A compound of any one of Embodiments 1 through 34 wherein m is 0, 1 or 2.

Embodiment 36P. A compound of Embodiment 35 wherein m is 0 or 1.

15 Embodiment 37P. A compound of Embodiment 36 wherein m is 1.

Embodiment 38P. A compound of Embodiment 37 wherein m is 0 (i.e. the 3-, 4-, 5- and 6-positions of the benzene ring are unsubstituted by R³).

20 Embodiment 39P. A compound of any one of Embodiments 1 through 37 wherein each R³ is independently halogen, cyano, hydroxy, nitro, amino, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C(=O)N(R^{3A})(R^{3B}), C(=NOR^{3C})H, C(=N)(R^{3D})H, C₁-C₄ alkoxy, C₂-C₄ cyanoalkoxy, C₂-C₄ alkylcarbonyl, C₂-C₄ alkoxy carbonyl, C₂-C₄ alkylcarbonyloxy, C₂-C₄ alkoxyalkyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, SO_nR^{3E} or C₃-C₆ cycloalkyl.

25 Embodiment 40P. A compound of Embodiment 39 wherein each R³ is independently halogen, cyano, amino, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₂-C₄ alkoxy carbonyl, C₂-C₄ alkylcarbonyloxy, C₂-C₄ alkoxyalkyl or C₁-C₄ haloalkyl.

Embodiment 41P. A compound of Embodiment 40 wherein each R³ is independently halogen, cyano, amino or C₁-C₄ alkyl.

30 Embodiment 42P. A compound of Embodiment 41 wherein each R³ is independently cyano.

Embodiment 43P. A compound of any one of Embodiments 1 through 37 or 39 through 42 wherein each R³ is attached to the remainder of Formula 1 at the 3-, 4- or 6-position.

35 Embodiment 44P. A compound of Embodiments 43 wherein each R³ is attached to the remainder of Formula 1 at the 3- or 4-position.

Embodiment 45P. A compound of Embodiment 44 wherein R³ is attached to the remainder of Formula 1 at the 3-position.

Embodiment 46P. A compound of any one of Embodiments 1 through 22 or 27 or 29 through 45 wherein R^{1A} is C₁-C₄ alkyl or C₁-C₄ haloalkyl.

Embodiment 47P. A compound of Embodiment 46 wherein R^{1A} is C₁-C₄ haloalkyl.

Embodiment 48P. A compound of any one of Embodiments 1 through 37 or 39 wherein R^{3E} is C₁-C₄ alkyl.

Embodiment 49P. A compound of any one of Embodiments 1 through 37 or 39 wherein R^{3A} is C₁-C₄ alkyl.

Embodiment 50P. A compound of any one of Embodiments 1 through 37 or 39 wherein R^{3B} is H or C₁-C₄ alkyl.

Embodiment 51P. A compound of any one of Embodiments 1 through 37 or 39 wherein R^{3C} is H or C₁-C₄ alkyl.

Embodiment 52P. A compound of any one of Embodiments 1 through 37 or 39 wherein R^{3D} is H or C₁-C₄ alkyl.

Embodiment 53P. A compound of any one of Embodiments 1 through 52 wherein Z is O.

Embodiment 54P. A compound of any one of Embodiments 1 through 53 wherein when m is 1, R³ is positioned at the 3-, 5- or 6-positions (i.e. the 3-, 5- and 6-positions of the benzene ring).

Embodiment 55P. A compound of any one of Embodiments 1 through 53 wherein when m is 1, R³ is other than Cl at the 4-position.

Embodiments of this invention, including Embodiments 1-48 and 1P-55P above as well as any other embodiments described herein, can be combined in any manner, and the descriptions of variables in the embodiments pertain not only to the a compound of Formula 1 but also to the starting compounds and intermediate compounds useful for preparing the compounds of Formula 1. In addition, embodiments of this invention, including Embodiments 1-48 and 1P-55P above as well as any other embodiments described herein, and any combination thereof, pertain to the compositions and methods of the present invention.

Embodiment AAA. A compound of Formula 1 wherein

Q is a 5- or 6-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, and optionally substituted with 1 to 4 R¹; or Q is phenyl substituted with 1 to 4 R¹;

Z is O or S;

R¹ is halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₂-C₄ alkoxyalkyl, C₂-C₄ alkylthioalkyl, SO_nR^{1A}, C₂-C₆ dialkylamino, C₁-C₄ cyanoalkyl, C₁-C₄ hydroxyalkyl, CH(=NOH) or C₃-C₆ cycloalkyl; or unsubstituted phenyl; or unsubstituted pyridyl;

R² is halogen, cyano, nitro, C₁-C₄ alkoxy, C₁-C₄ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, SO_nR^{2A} or C₁-C₄ haloalkyl;

each R³ is independently halogen, cyano, hydroxy, nitro, amino, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C(=O)N(R^{3A})(R^{3B}), C(=NOR^{3C})H, C(=N)(R^{3D})H, C₁-C₄ alkoxy, C₂-C₄ cyanoalkoxy, C₂-C₄ alkylcarbonyl, C₂-C₄ alkoxy carbonyl, C₂-C₄ alkylcarbonyloxy, C₂-C₄ alkoxyalkyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, SO_nR^{3E} or C₃-C₆ cycloalkyl; or phenyl optionally substituted with cyano, halogen or C₁-C₄ alkyl;

m is 0, 1, 2 or 3;

each R^{1A}, R^{2A} and R^{3E} is independently C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkylamino or C₂-C₆ dialkylamino;

R^{3A} is C₁-C₄ alkyl or C₁-C₄ haloalkyl;

R^{3B} is H, C₁-C₄ alkyl or C₁-C₄ haloalkyl;

R^{3C} is independently H or C₁-C₄ alkyl;

R^{3D} is independently H or C₁-C₄ alkyl; and

n is 0, 1, or 2;

provided the compound of Formula 1 is other than 5-chloro-2-[(4'-methyl[1,1'-biphenyl]2-yl)oxy]-pyrimidine (CAS # 107492-74-0), 5-chloro-2-[(4'-chloro[1,1'-biphenyl]2-yl)oxy]-pyrimidine (CAS # 107492-72-8), 5-chloro-2-[(3'-chloro[1,1'-biphenyl]2-yl)oxy]-pyrimidine (CAS # 107492-76-2) and 5-chloro-2-[[3'-(trifluoromethyl)[1,1'-biphenyl]2-yl]oxy]-pyrimidine (CAS # 107492-75-1); and provided i) when Q is 5-chloro-2-pyridinyl; Z is O; and R³ is 4-chloro, then R² is other than Cl or Br; ii) when Q is 4-CF₃-2-pyrimidinyl; Z is O; and m is 0, then R² is other than Cl or Br; and iii) when Q is 6-CF₃-2-pyridinyl; Z is O; and m is 0, then R² is other than Br.

Embodiment AA. A compound of Embodiment AAA or a compound of Formula 1 as described in the Summary of the Invention wherein

Q is a 5- or 6-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, and optionally substituted with 1 to 4 R¹;

Z is O or S;

each R¹ is independently halogen, cyano, nitro, SF₅, CHO, C(=O)NH₂, C(=S)NH₂, SO₂NH₂, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₄-C₈ alkylcycloalkyl, C₄-C₈ cycloalkylalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₃-C₇ cycloalkylcarbonyl, C₂-C₈ alkylaminocarbonyl, C₃-C₁₀ dialkylaminocarbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₃-C₆ cycloalkoxy, C₃-C₆ halocycloalkoxy, C₄-C₈

cycloalkylalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ alkoxyhaloalkyl, C₂-C₆ alkoxyalkoxy, C₂-C₄ alkylcarbonyloxy, C₂-C₆ cyanoalkyl, C₂-C₆ cyanoalkoxy, C₁-C₄ hydroxyalkyl, C₂-C₄ alkylthioalkyl, SO_nR^{1A}, Si(CH₃)₃ or B(-OC(R^{1B})₂C(R^{1B})₂O-); or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{1C}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{1C} on carbon atom ring members and R^{1D} on nitrogen atom ring members;

R² is halogen, cyano, nitro, C₁-C₄ alkoxy, C₁-C₄ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, SO_nR^{2A}, C₁-C₄ haloalkyl or C₃-C₆ cycloalkyl;

each R³ is independently halogen, cyano, hydroxy, nitro, amino, CHO, C(=O)NH₂, C(=S)NH₂, SO₂NH₂, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₄-C₈ alkylcycloalkyl, C₄-C₈ cycloalkylalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxycarbonyl, C₃-C₇ cycloalkylcarbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₃-C₆ cycloalkoxy, C₃-C₆ halocycloalkoxy, C₄-C₈ cycloalkylalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ alkoxyhaloalkyl, C₂-C₆ alkoxyalkoxy, C₂-C₄ alkylcarbonyloxy, C₂-C₆ cyanoalkyl, C₂-C₆ cyanoalkoxy, C₂-C₄ alkylthioalkyl, Si(CH₃)₃, C≡CSi(CH₃)₃, C(=O)N(R^{3A})(R^{3B}), C(=NOR^{3C})H, C(=NR^{3D})H, SO_nR^{3E}; or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{3F}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members; or pyrimidinyloxy;

m is 0, 1, 2 or 3;

each n is independently 0, 1 or 2;

each R^{1A}, R^{2A} and R^{3E} is independently C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkylamino or C₂-C₆ dialkylamino;

each R^{1B} is independently H or C₁-C₄ alkyl;

each R^{1C} is independently hydroxy, halogen, cyano, nitro, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₁-C₆ haloalkoxy;

each R^{1D} is independently cyano, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₂-C₆ alkylcarbonyl;

each R^{3A} is independently C₁-C₄ alkyl or C₁-C₄ haloalkyl;

each R^{3B} is independently H, C₁-C₄ alkyl or C₁-C₄ haloalkyl;

5 each R^{3C} is independently H or C₁-C₄ alkyl;

each R^{3D} is independently H, amino, C₁-C₄ alkyl or C₁-C₄ alkylamino;

each R^{3F} is independently hydroxy, halogen, cyano, nitro, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₁-C₆ haloalkoxy; and

10 each R^{3G} is independently cyano, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₂-C₆ alkylcarbonyl;

Embodiment A. A compound of Embodiment AA wherein

Q is selected from Q-1 through Q-55 wherein r is 0, 1, 2 or 3; and s is 0 or 1;

15 each R¹ is independently halogen, cyano, SF₅, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxycarbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ cyanoalkyl, C₁-C₄ hydroxyalkyl, C₂-C₄ alkylthioalkyl or SO_nR^{1A};

20 R³ is independently halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₄-C₈ alkylcycloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxycarbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₃-C₆ cycloalkoxy, C₃-C₆ halocycloalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₄ alkylcarbonyloxy, C₂-C₆ cyanoalkyl, C(=O)N(R^{3A})(R^{3B}), C(=NOR^{3C})H, SO_nR^{3E}; or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{3F}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members;

Z is O; and

m is 0, 1 or 2.

35 Embodiment B. A compound of Embodiment A wherein

each R¹ is independently halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy,

C₃-C₄ haloalkynyloxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₁-C₄ hydroxyalkyl, C₂-C₄ alkylthioalkyl or SO_nR^{1A};

R² is halogen, C₁-C₄ alkyl or C₁-C₄ haloalkyl;

each R³ is independently halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ cyanoalkyl, SO_nR^{3E}; or a

5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members; and

m is 0 or 1.

Embodiment C1. A compound of Embodiment B wherein

Q is selected from Q-7 through Q-24;

each R¹ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy or SO_nR^{1A};

R² is halogen or C₁-C₄ alkyl;

each R³ is independently halogen, cyano, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl or C₂-C₆ haloalkoxyalkyl; and

each R^{1A} is independently C₁-C₄ alkyl or C₁-C₄ haloalkyl.

Embodiment C2. A compound of Embodiment B wherein

Q is selected from Q-43, Q-44, Q-45, Q-48, Q-49 and Q-50;

each R¹ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy or SO_nR^{1A};

R² is halogen or C₁-C₄ alkyl;

each R³ is independently halogen, cyano, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl or C₂-C₆ haloalkoxyalkyl; and

each R^{1A} is independently C₁-C₄ alkyl or C₁-C₄ haloalkyl.

Embodiment D1. A compound of Embodiment C1 wherein

Q is selected from Q-16 and Q-18;

each R¹ is independently halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy;

R² is halogen or CH₃; and

each R³ is independently halogen, cyano, C₁-C₄ alkyl or C₁-C₄ haloalkyl.

Embodiment D2. A compound of Embodiment C2 wherein

Q is selected from Q-43, Q-44 and Q-45;

each R¹ is independently halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy;

5 R² is halogen or CH₃; and

each R³ is independently halogen, cyano, C₁-C₄ alkyl or C₁-C₄ haloalkyl.

Specific embodiments include compounds of Formula 1 selected from the group consisting of:

5-chloro-2-[2-(5-chloro-2-pyridinyl)phenoxy]pyrimidine (Compound 1),
5-chloro-2-[2-[5-(fluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine (Compound 32),
2-[2-(3-bromo-5-isoxazolyl)phenoxy]-5-chloropyrimidine (Compound 12),
5-chloro-2-[2-[5-(trifluoromethyl)-2-pyridinyl]phenoxy]pyrimidine (Compound 27),
5-chloro-2-[3-chloro-2-(5-chloro-2-pyridinyl)phenoxy]pyrimidine (Compound 23),
4-[2-[(5-bromo-2-pyrimidinyl)oxy]phenyl]-2-(trifluoromethyl)pyrimidine (Compound 21),
2-[2-(2-bromo-5-thiazolyl)phenoxy]-5-(trifluoromethyl)pyrimidine (Compound 15),
5-chloro-2-[4-methyl-2-[2-(trifluoromethyl)-4-pyridinyl]phenoxy]pyrimidine (Compound 24),
5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine (Compound 35),
5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]phenoxy]pyrimidine (Compound 53),
5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine (Compound 55),
5-bromo-2-[2-[5-(difluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine (Compound 62),
5-chloro-2-[2-[3-(trifluoromethyl)-5-isoxazolyl]phenoxy]pyrimidine (Compound 63),
5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine (Compound 144),
5-bromo-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine (Compound 145),
5-chloro-2-[2-[5-(trifluoromethyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine (Compound 168) and

5-chloro-2-[2-[5-(trifluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine (Compound 200).

Embodiments of the present invention as described in the Summary of the Invention and Embodiment AAA also include the following:

Embodiment Ap. A compound of the Summary of the Invention wherein

Q is a 5- or 6-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, and optionally substituted with 1 to 3 R¹; or

Q is phenyl substituted with 1 to 3 R¹;

R¹ is halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₂-C₄ alkoxyalkyl, C₂-C₄ alkylthioalkyl or SO_nR^{1A};

each n is independently 0, 1 or 2

R² is halogen, C₁-C₄ alkyl or C₁-C₄ haloalkyl;

m is 0, 1 or 2;

each R³ is independently halogen, cyano, hydroxy, nitro, amino, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C(=O)N(R^{3A})(R^{3B}), C(=NOR^{3C})H, C(=N)(R^{3D})H, C₁-C₄ alkoxy, C₂-C₄ cyanoalkoxy, C₂-C₄ alkylcarbonyl, C₂-C₄ alkoxyalkyl, C₂-C₄ alkylcarbonyloxy, C₂-C₄ alkoxyalkyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, SO_nR^{3E} or C₃-C₆ cycloalkyl;

each R³ is attached to the remainder of Formula 1 at the 3-, 4- or 6-position;

R^{1A} is C₁-C₄ alkyl or C₁-C₄ haloalkyl;

R^{3E} is C₁-C₄ alkyl;

R^{3A} is C₁-C₄ alkyl;

R^{3B} is H or C₁-C₄ alkyl;

R^{3C} is H or C₁-C₄ alkyl; and

R^{3D} is H or C₁-C₄ alkyl.

Embodiment Bp. A compound of Embodiment A wherein

Q is selected from Q-1 through Q-54 (i.e. as described in Embodiment 3);

Z is O;

R¹ is halogen, cyano, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, or SCF₃;

R² is halogen or C₁-C₄ alkyl;

m is 0 or 1;

each R³ is independently halogen, cyano, amino, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₂-C₄ alkoxyalkyl, C₂-C₄ alkylcarbonyloxy, C₂-C₄ alkoxyalkyl or C₁-C₄ haloalkyl; and

each R³ is attached to the remainder of Formula 1 at the 3- or 4-position.

Embodiment Cp. A compound of Embodiment B wherein

Q is a 5-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, optionally substituted with R¹, and is selected from Q-1 through Q-41;

R¹ is halogen C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy;

R² is halogen or CH₃; and

each R³ is independently halogen, cyano, amino or C₁-C₄ alkyl.

Embodiment Dp. A compound of Embodiment C wherein

Q is 6-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, optionally substituted with R¹, and is selected from Q-42 through Q-54;

R¹ is halogen C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy;

R² is halogen or CH₃; and

each R³ is independently halogen, cyano, amino or C₁-C₄ alkyl.

Embodiment Ep. A compound of Embodiment D wherein

Q is selected from Q-42, Q-43, Q-44, Q-47, Q-48 and Q-49;

R¹ is Cl, Br, CF₃ or OCF₃;

R² is halogen; and

each R³ is independently cyano.

Specific embodiments include a compound of Formula 1 selected from the group consisting of:

5-chloro-2-[2-(5-chloro-2-pyridinyl)phenoxy]pyrimidine (Compound 1);

5-chloro-2-[2-[5-(fluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine (Compound 32);

2-[2-(3-bromo-5-isoxazolyl)phenoxy]-5-chloropyrimidine (Compound 12);

5-chloro-2-[[4'-(trifluoromethoxy)[1,1'-biphenyl]-2-yl]oxy]pyrimidine (Compound 42);

5-chloro-2-[2-[5-(trifluoromethyl)-2-pyridinyl]phenoxy]pyrimidine (Compound 27);

5-chloro-2-[3-chloro-2-(5-chloro-2-pyridinyl)phenoxy]pyrimidine (Compound 23);

4-[2-[(5-bromo-2-pyrimidinyl)oxy]phenyl]-2-(trifluoromethyl)pyrimidine (Compound 21);

2-[2-(2-bromo-5-thiazolyl)phenoxy]-5-(trifluoromethyl)pyrimidine (Compound 15);

and

5-chloro-2-[4-methyl-2-[2-(trifluoromethyl)-4-pyridinyl]phenoxy]pyrimidine (Compound 24).

This invention also relates to a method for controlling undesired vegetation comprising applying to the locus of the vegetation herbicidally effective amounts of the compounds of the invention (e.g., as a composition described herein). Of note as embodiments relating to methods of use are those involving the compounds of embodiments described above. Compounds of the invention are particularly useful for selective control of weeds in crops

such as wheat, barley, maize, soybean, sunflower, cotton, oilseed rape and rice, and specialty crops such as sugarcane, citrus, fruit and nut crops.

Also noteworthy as embodiments are herbicidal compositions of the present invention comprising the compounds of embodiments described above.

5 This invention also includes a herbicidal mixture comprising (a) a compound selected from Formula 1, *N*-oxides, and salts thereof, and (b) at least one additional active ingredient selected from (b1) photosystem II inhibitors, (b2) acetohydroxy acid synthase (AHAS) inhibitors, (b3) acetyl-CoA carboxylase (ACCase) inhibitors, (b4) auxin mimics and (b5) 5-enol-pyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, (b6) photosystem I
10 electron diverters, (b7) protoporphyrinogen oxidase (PPO) inhibitors, (b8) glutamine synthetase (GS) inhibitors, (b9) very long chain fatty acid (VLCFA) elongase inhibitors, (b10) auxin transport inhibitors, (b11) phytoene desaturase (PDS) inhibitors, (b12) 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors, (b13) homogentisate solanesyltransferase (HST) inhibitors, (b14) cellulose biosynthesis inhibitors, (b15) other
15 herbicides including mitotic disruptors, organic arsenicals, asulam, bromobutide, cinmethylin, cumyluron, dazomet, difenzoquat, dymron, etobenzanid, flurenol, fosamine, fosamine-ammonium, metam, methyldymron, oleic acid, oxaziclomefone, pelargonic acid and pyributicarb, and (b16) herbicide safeners; and salts of compounds of (b1) through (b16).

20 "Photosystem II inhibitors" (b1) are chemical compounds that bind to the D-1 protein at the Q_B-binding niche and thus block electron transport from Q_A to Q_B in the chloroplast thylakoid membranes. The electrons blocked from passing through photosystem II are transferred through a series of reactions to form toxic compounds that disrupt cell membranes and cause chloroplast swelling, membrane leakage, and ultimately cellular
25 destruction. The Q_B-binding niche has three different binding sites: binding site A binds the triazines such as atrazine, triazinones such as hexazinone, and uracils such as bromacil, binding site B binds the phenylureas such as diuron, and binding site C binds benzothiadiazoles such as bentazon, nitriles such as bromoxynil and phenyl-pyridazines such as pyridate. Examples of photosystem II inhibitors include ametryn, amicarbazone, atrazine,
30 bentazon, bromacil, bromofenoxim, bromoxynil, chlorbromuron, chloridazon, chlorotoluron, chloroxuron, cumyluron, cyanazine, daimuron, desmedipham, desmetryn, dimefuron, dimethametryn, diuron, ethidimuron, fenuron, fluometuron, hexazinone, ioxynil, isoproturon, isouron, lenacil, linuron, metamiltron, methabenzthiazuron, metobromuron, metoxuron, metribuzin, monolinuron, neburon, pentanochlor, phenmedipham, prometon,
35 prometryn, propanil, propazine, pyridafol, pyridate, siduron, simazine, simetryn, tebuthiuron, terbacil, terbutometon, terbutylazine, terbutryn and trietazine.

"AHAS inhibitors" (b2) are chemical compounds that inhibit acetohydroxy acid synthase (AHAS), also known as acetolactate synthase (ALS), and thus kill plants by

inhibiting the production of the branched-chain aliphatic amino acids such as valine, leucine and isoleucine, which are required for protein synthesis and cell growth. Examples of AHAS inhibitors include amidosulfuron, azimsulfuron, bensulfuron-methyl, bispyribac-sodium, cloransulam-methyl, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, diclosulam, ethametsulfuron-methyl, ethoxysulfuron, flazasulfuron, florasulam, flucarbazone-sodium, flumetsulam, flupyrsulfuron-methyl, flupyrsulfuron-sodium, foramsulfuron, halosulfuron-methyl, imazamethabenz-methyl, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, imazosulfuron, iodosulfuron-methyl (including sodium salt), iofensulfuron (2-iodo-*N*-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide), mesosulfuron-methyl, metazosulfuron (3-chloro-4-(5,6-dihydro-5-methyl-1,4,2-dioxazin-3-yl)-*N*-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-1-methyl-1*H*-pyrazole-5-sulfonamide), metosulam, metsulfuron-methyl, nicosulfuron, oxasulfuron, penoxsulam, primisulfuron-methyl, propoxycarbazone-sodium, propyrisulfuron (2-chloro-*N*-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-6-propylimidazo[1,2-*b*]pyridazine-3-sulfonamide), prosulfuron, pyrazosulfuron-ethyl, pyribenzoxim, pyriftalid, pyriminobac-methyl, pyrithiobac-sodium, rimsulfuron, sulfometuron-methyl, sulfosulfuron, thiencarbazone, thifensulfuron-methyl, triafamone (*N*-[2-[(4,6-dimethoxy-1,3,5-triazin-2-yl)carbonyl]-6-fluorophenyl]-1,1-difluoro-*N*-methylmethanesulfonamide), triasulfuron, tribenuron-methyl, trifloxysulfuron (including sodium salt), triflusulfuron-methyl and tritosulfuron.

“ACCCase inhibitors” (b3) are chemical compounds that inhibit the acetyl-CoA carboxylase enzyme, which is responsible for catalyzing an early step in lipid and fatty acid synthesis in plants. Lipids are essential components of cell membranes, and without them, new cells cannot be produced. The inhibition of acetyl CoA carboxylase and the subsequent lack of lipid production leads to losses in cell membrane integrity, especially in regions of active growth such as meristems. Eventually shoot and rhizome growth ceases, and shoot meristems and rhizome buds begin to die back. Examples of ACCCase inhibitors include alloxydim, butoxydim, clethodim, clodinafop, cycloxydim, cyhalofop, diclofop, fenoxaprop, fluazifop, haloxyfop, pinoxaden, profoxydim, propaquizafop, quizalofop, sethoxydim, tepraloxydim and tralkoxydim, including resolved forms such as fenoxaprop-P, fluazifop-P, haloxyfop-P and quizalofop-P and ester forms such as clodinafop-propargyl, cyhalofop-butyl, diclofop-methyl and fenoxaprop-P-ethyl.

Auxin is a plant hormone that regulates growth in many plant tissues. “Auxin mimics” (b4) are chemical compounds mimicking the plant growth hormone auxin, thus causing uncontrolled and disorganized growth leading to plant death in susceptible species. Examples of auxin mimics include aminocyclopyrachlor (6-amino-5-chloro-2-cyclopropyl-4-pyrimidinecarboxylic acid) and its methyl and ethyl esters and its sodium and potassium salts, aminopyralid, benazolin-ethyl, chloramben, clacyfos, clomeprop, clopyralid, dicamba,

2,4-D, 2,4-DB, dichlorprop, fluroxypyr, halauxifen (4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-2-pyridinecarboxylic acid), halauxifen-methyl (methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-2-pyridinecarboxylate), MCPA, MCPB, mecoprop, picloram, quinclorac, quinmerac, 2,3,6-TBA, triclopyr, and methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoro-2-pyridinecarboxylate.

“EPSP (5-enol-pyruvylshikimate-3-phosphate) synthase inhibitors” (b5) are chemical compounds that inhibit the enzyme, 5-enol-pyruvylshikimate-3-phosphate synthase, which is involved in the synthesis of aromatic amino acids such as tyrosine, tryptophan and phenylalanine. EPSP inhibitor herbicides are readily absorbed through plant foliage and translocated in the phloem to the growing points. Glyphosate is a relatively nonselective postemergence herbicide that belongs to this group. Glyphosate includes esters and salts such as ammonium, isopropylammonium, potassium, sodium (including sesquisodium) and trimesium (alternatively named sulfosate).

“Photosystem I electron diverters” (b6) are chemical compounds that accept electrons from Photosystem I, and after several cycles, generate hydroxyl radicals. These radicals are extremely reactive and readily destroy unsaturated lipids, including membrane fatty acids and chlorophyll. This destroys cell membrane integrity, so that cells and organelles “leak”, leading to rapid leaf wilting and desiccation, and eventually to plant death. Examples of this second type of photosynthesis inhibitor include diquat and paraquat.

“PPO inhibitors” (b7) are chemical compounds that inhibit the enzyme protoporphyrinogen oxidase, quickly resulting in formation of highly reactive compounds in plants that rupture cell membranes, causing cell fluids to leak out. Examples of PPO inhibitors include acifluorfen-sodium, azafenidin, benzfendizone, bifenox, butafenacil, carfentrazone, carfentrazone-ethyl, chlomethoxyfen, cinidon-ethyl, fluazolate, flufenpyr-ethyl, flumiclorac-pentyl, flumioxazin, fluoroglycofen-ethyl, fluthiacet-methyl, fomesafen, halosafen, lactofen, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, profluzol, pyraclostrobin, pyraflufen-ethyl, saflufenacil, sulfentrazone, thidiazimin, tiafenacil (methyl *N*-[2-[[[2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2*H*)-pyrimidinyl]-4-fluorophenyl]thio]-1-oxopropyl]-β-alaninate) and 3-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propyn-1-yl)-2*H*-1,4-benzoxazin-6-yl]dihydro-1,5-dimethyl-6-thioxo-1,3,5-triazine-2,4(1*H*,3*H*)-dione.

“GS (glutamine synthase) inhibitors” (b8) are chemical compounds that inhibit the activity of the glutamine synthetase enzyme, which plants use to convert ammonia into glutamine. Consequently, ammonia accumulates and glutamine levels decrease. Plant damage probably occurs due to the combined effects of ammonia toxicity and deficiency of amino acids required for other metabolic processes. The GS inhibitors include glufosinate and its esters and salts such as glufosinate-ammonium and other phosphinothricin

derivatives, glufosinate-P ((2*S*)-2-amino-4-(hydroxymethylphosphinyl)butanoic acid) and bilanaphos.

“VLCFA (very long chain fatty acid) elongase inhibitors” (b9) are herbicides having a wide variety of chemical structures, which inhibit the elongase. Elongase is one of the enzymes located in or near chloroplasts which are involved in biosynthesis of VLCFAs. In plants, very-long-chain fatty acids are the main constituents of hydrophobic polymers that prevent desiccation at the leaf surface and provide stability to pollen grains. Such herbicides include acetochlor, alachlor, anilofos, butachlor, cafenstrole, dimethachlor, dimethenamid, diphenamid, fenoxasulfone (3-[[[(2,5-dichloro-4-ethoxyphenyl)methyl]sulfonyl]-4,5-dihydro-5,5-dimethylisoxazole), fentrazamide, flufenacet, indanofan, mefenacet, metazachlor, metolachlor, naproanilide, napropamide, napropamide-M ((2*R*)-*N,N*-diethyl-2-(1-naphthalenyloxy)propanamide), pethoxamid, piperophos, pretilachlor, propachlor, propisochlor, pyroxasulfone, and thenylchlor, including resolved forms such as *S*-metolachlor and chloroacetamides and oxyacetamides.


“Auxin transport inhibitors” (b10) are chemical substances that inhibit auxin transport in plants, such as by binding with an auxin-carrier protein. Examples of auxin transport inhibitors include diflufenzopyr, naptalam (also known as *N*-(1-naphthyl)phthalamic acid and 2-[(1-naphthalenylamino)carbonyl]benzoic acid).

“PDS (phytoene desaturase inhibitors) (b11) are chemical compounds that inhibit carotenoid biosynthesis pathway at the phytoene desaturase step. Examples of PDS inhibitors include beflubutamid, diflufenican, fluridone, flurochloridone, flurtamone norflurzon and picolinafen.

“HPPD (4-hydroxyphenyl-pyruvate dioxygenase) inhibitors” (b12) are chemical substances that inhibit the biosynthesis of synthesis of 4-hydroxyphenyl-pyruvate dioxygenase. Examples of HPPD inhibitors include benzobicyclon, benzofenap, bicyclopyrone (4-hydroxy-3-[[2-[(2-methoxyethoxy)methyl]-6-(trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one), fenquinotrine (2-[[8-chloro-3,4-dihydro-4-(4-methoxyphenyl)-3-oxo-2-quinoxaliny]carbonyl]-1,3-cyclohexanedione), isoxachlortole, isoxaflutole, mesotrione, pyrasulfotole, pyrazolynate, pyrazoxyfen, sulcotrione, tefuryltrione, tembotrione, topramezone, 5-chloro-3-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-1-(4-methoxyphenyl)-2(1*H*)-quinoxalinone, 4-(2,6-diethyl-4-methylphenyl)-5-hydroxy-2,6-dimethyl-3(2*H*)-pyridazinone, 4-(4-fluorophenyl)-6-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-2-methyl-1,2,4-triazine-3,5(2*H*,4*H*)-dione, 5-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-2-(3-methoxyphenyl)-3-(3-methoxypropyl)-4(3*H*)-pyrimidinone, 2-methyl-*N*-(4-methyl-1,2,5-oxadiazol-3-yl)-3-(methylsulfinyl)-4-(trifluoromethyl)benzamide and 2-methyl-3-(methylsulfonyl)-*N*-(1-methyl-1*H*-tetrazol-5-yl)-4-(trifluoromethyl)benzamide.

HST (homogentisate solanesyltransferase) inhibitors (b13) disrupt a plant's ability to convert homogentisate to 2-methyl-6-solanyl-1,4-benzoquinone, thereby disrupting carotenoid biosynthesis. Examples of HST inhibitors include haloxydine, pyriclor, 3-(2-chloro-3,6-difluorophenyl)-4-hydroxy-1-methyl-1,5-naphthyridin-2(1*H*)-one, 7-(3,5-dichloro-4-pyridinyl)-5-(2,2-difluoroethyl)-8-hydroxypyrido[2,3-*b*]pyrazin-6(5*H*)-one and 4-(2,6-diethyl-4-methylphenyl)-5-hydroxy-2,6-dimethyl-3(2*H*)-pyridazinone.

HST inhibitors also include compounds of Formulae **A** and **B**.



wherein R^{d1} is H, Cl or CF_3 ; R^{d2} is H, Cl or Br; R^{d3} is H or Cl; R^{d4} is H, Cl or CF_3 ; R^{d5} is CH_3 , CH_2CH_3 or CH_2CHF_2 ; and R^{d6} is OH, or $-OC(=O)-i\text{-Pr}$; and R^{e1} is H, F, Cl, CH_3 or CH_2CH_3 ; R^{e2} is H or CF_3 ; R^{e3} is H, CH_3 or CH_2CH_3 ; R^{e4} is H, F or Br; R^{e5} is Cl, CH_3 , CF_3 , OCF_3 or CH_2CH_3 ; R^{e6} is H, CH_3 , CH_2CHF_2 or $C\equiv CH$; R^{e7} is OH, $-OC(=O)Et$, $-OC(=O)-i\text{-Pr}$ or $-OC(=O)-t\text{-Bu}$; and A^{e8} is N or CH.

Cellulose biosynthesis inhibitors (b14) inhibit the biosynthesis of cellulose in certain plants. They are most effective when using a pre-application or early post-application on young or rapidly growing plants. Examples of cellulose biosynthesis inhibitors include chlorthiamid, dichlobenil, flupoxam, indaziflam (*N*²-[(1*R*,2*S*)-2,3-dihydro-2,6-dimethyl-1*H*-inden-1-yl]-6-(1-fluoroethyl)-1,3,5-triazine-2,4-diamine), isoxaben and triaziflam.

Other herbicides (b15) include herbicides that act through a variety of different modes of action such as mitotic disruptors (e.g., flamprop-M-methyl and flamprop-M-isopropyl) organic arsenicals (e.g., DSMA, and MSMA), 7,8-dihydropteroate synthase inhibitors, chloroplast isoprenoid synthesis inhibitors and cell-wall biosynthesis inhibitors. Other herbicides include those herbicides having unknown modes of action or do not fall into a specific category listed in (b1) through (b14) or act through a combination of modes of action listed above. Examples of other herbicides include aclonifen, asulam, amitrole, bromobutide, cinmethylin, clomazone, cumyluron, cyclopyrimorate (6-chloro-3-(2-cyclopropyl-6-methylphenoxy)-4-pyridazinyl 4-morpholinecarboxylate), daimuron, difenzoquat, etobenzanid, fluometuron, flurenol, fosamine, fosamine-ammonium, dazomet, dymron, ipfencarbazone (1-(2,4-dichlorophenyl)-*N*-(2,4-difluorophenyl)-1,5-dihydro-*N*-(1-methylethyl)-5-oxo-4*H*-1,2,4-triazole-4-carboxamide), metam, methyldymron, oleic acid,

oxaziclomefone, pelargonic acid, pyributicarb and 5-[[2,6-difluorophenyl)methoxy]methyl]-4,5-dihydro-5-methyl-3-(3-methyl-2-thienyl)isoxazole.


“Herbicide safeners” (b16) are substances added to a herbicide formulation to eliminate or reduce phytotoxic effects of the herbicide to certain crops. These compounds protect crops from injury by herbicides but typically do not prevent the herbicide from controlling undesired vegetation. Examples of herbicide safeners include but are not limited to benoxacor, cloquintocet-mexyl, cumyluron, cyometrinil, cyprosulfamide, daimuron, dichlormid, dicyclonon, dimepiperate, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, isoxadifen-ethyl, mefenpyr-diethyl, mephenate, methoxyphenone, naphthalic anhydride, oxabetrinil, *N*-(aminocarbonyl)-2-methylbenzenesulfonamide and *N*-(aminocarbonyl)-2-fluorobenzenesulfonamide, 1-bromo-4-[(chloromethyl)sulfonyl]benzene, 2-(dichloromethyl)-2-methyl-1,3-dioxolane (MG 191), 4-(dichloroacetyl)-1-oxa-4-azospiro[4.5]decane (MON 4660).

The compounds of Formula 1 can be prepared by general methods known in the art of synthetic organic chemistry. One or more of the following methods and variations as described in Schemes 1–9 can be used to prepare the compounds of Formula 1. The definitions of Q, R¹, R² and R³ in the compounds of Formulae 1–11 below are as defined above in the Summary of the Invention unless otherwise noted. Compounds of Formulae 1A–1C, 2A–2F, 4A and 8A are various subsets of the compounds of Formula 1, 2 4 and 8 and all substituents for Formulae 1A–1C, 2A–2F, 4A and 8A are as defined above for Formula 1 unless otherwise noted.

One or more of the following methods and variations as described in Schemes 1–9 can be used to prepare the compounds of Formula 1. The definitions of Q, R¹, R² and R³ in the compounds of Formulae 1–11 below are as defined above in the Summary of the Invention unless otherwise noted.


As shown in Scheme 1 a compound of Formula 1 can be prepared by nucleophilic substitution by heating a compound of Formula 2 in a suitable solvent, such as acetonitrile, tetrahydrofuran or *N,N*-dimethylformamide in the presence of a base such as potassium or cesium carbonate, at temperatures ranging from 50 to 110 °C, with a compound of Formula 3 (where LG is halogen or SO₂Me). The reaction is typically conducted at temperatures ranging from 50 to 110 °C.

35




Alternatively, as shown in Scheme 2, boron compounds of Formula **5** or tin compounds of Formula **6** can be coupled with intermediates of Formula **4** under Suzuki or Stille conditions to give compounds of Formula **1**. Suzuki couplings typically are conducted in the presence of Pd(0) or Pd(II) salts, a suitable ligand, and a base. Suitable bases for this transformation include potassium carbonate or cesium carbonate, while Pd(II) salts such as Pd(OAc)₂ or PdCl₂ can be used in conjunction with ligands such as triphenylphosphine or 1,1'-bis(diphenylphosphino)ferrocene (dppf). Conditions for Suzuki couplings are well documented in the literature (see for example *Angewandte Chemie International Edition* **2006**, 45, 3484 and *Tetrahedron Letters* **2002**, 58(14), 2885). Boron intermediates of Formula **5** are commercially available or can be prepared from the corresponding halides or trifluoromethanesulfonates by methods known in the literature (see for example PCT Patent Publication WO 2007/043278, US Pat. No. 8,080,566, *Organic Letters* **2011**, 13(6), 1366 and *Organic Letters* **2012**, 14(2), 600). Stille couplings typically can be conducted in the presence of Pd(0) or a Pd(II) salt, a ligand and a Cu(I) salt such as copper(I) iodide. The reaction can be run in a solvent such as dioxane, 1,2-dimethoxyethane or toluene at a temperature ranging from ambient to reflux. For conditions and reagents employed in Stille couplings see *Chemical Reviews* **2007**, 107(1), 133–173.

Scheme 2




As shown in Scheme 3, a compound of Formula **2C** (i.e. a compound of Formula **2** where Z is O) can be prepared by deprotection of a compound of Formula **2B** (i.e. a compound of Formula **2A** wherein Z is O; and R^A is CH₃ or -C(=O)CH₃) with a suitable deprotecting agent. Suitable methoxy (i.e. when R^A is CH₃) deprotecting reagents such as BBr₃, AlCl₃ and HBr in acetic acid can be used in the presence of solvents such as toluene, dichloromethane and dichloroethane at a temperature of from -80 to 120 °C. Suitable acetoxy (i.e. when R^A is -C(=O)CH₃) deprotecting agents include potassium carbonate in methanol or ammonium acetate in aqueous methanol at room temperature can be used as discussed in Das, et al., *Tetrahedron* **2003**, 59, 1049–1054 and methods cited therein. Alternatively, a compound of Formula **2B** can be combined with Amberlyst 15[®] in methanol (as discussed in Das, et al. *Tet. Lett.* **2003**, 44, 5465–5468) or combined with sodium acetate in ethanol (as discussed in Narender, T., et al. *Synthetic Communications* **2009**, 39(11), 1949–1956) to obtain a compound of Formula **2C**. Other useful phenolic protecting groups suitable for use in preparing a compound of Formula **2C** can be found in Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 4th ed.; Wiley: Hoboken, New Jersey, 1991.

Scheme 3




An intermediate of Formula **2B** can be prepared as shown in Scheme 4 from an intermediate of Formula **7** by a variety of methods known to one skilled in the art. Compounds of Formula **2B** can be accessed by coupling precursors of Formula **7** wherein J is Br, Cl, I or trifluoromethanesulfonate with boronate or trialkyltin group-containing heterocycles (i.e compounds of Formula **5** or Formula **6** using the Suzuki conditions or the Stille conditions of Scheme 2). Alternatively, compounds of Formula **7** wherein J is a boronate or trialkyltin group may be coupled with halogen-substituted heterocycles Q-X using the methods shown in Scheme 2 to afford compounds of Formula **2B**. The skilled chemist will realize that with the prudent choice of groups X and J in reactions involving compounds of Formula **7** and Q-X can synthesize the intermediate **2B** utilizing various cross coupling procedures such as Kumada coupling, Hiyama coupling or Negishi coupling described in "Metal-Catalyzed Cross-Coupling Reactions", Eds. A. de Meijere and F. Diederich, Wiley-VCH, Weinheim, 2004, vols 1 and 2.

When J in Formula **7** is an alkene, alkyne, oxime, nitrile or ketone, various heterocycles can be prepared using methods described in Katritzky, *Advances in Heterocyclic Chemistry*, Vol. 1-104, Elsevier. In cases where regioisomeric mixtures are produced, the desired product can be isolated using routine separation techniques known in the art.




As shown in Scheme 5, a compound of Formula **4A** can be prepared by coupling of phenols of Formula **9** with a compound of Formula **3** under the nucleophilic substitution conditions described in Scheme 1.



As shown in Scheme 6, a compound of Formula **1B**, (i.e. a compound of Formula **1** where Z is O; and m is 1 at the 3-position) can be prepared by "C-H activation" of a compound of Formula **1A** (a compound of Formula **1** wherein Z is O; and m is 0). For


example, palladium(II) acetate along with either an *N*-halosuccinimide, $\text{PhI}(\text{OAc})_2$, *N*-fluoropyridinium tetrafluoroborate, or a lower alkyl boronic acid can be used to introduce the R^3 variable as I, Br, Cl, -OAc, F, and lower alkyl substituents respectively. These methods are detailed in reviews of selective activation of C-H bonds in *Chemical Reviews* **2010**, *110*, 575-1211 and references cited therein. Methods for "C-H activation" can also be found in Wencel-Delord et al., *Nature Chemistry* **2013**, *5*, 369-375 and a series of reviews of "C-H activation" in *Accounts of Chemical Research* **2012**, *45*, 777-958 and references cited therein. Iodides and bromides of Formula **1B** can then be further functionalized by various cross coupling procedures described in "Metal-Catalyzed Cross-Coupling Reactions", Eds A. de Meijere and F. Diederich, Wiley-VCH, Weinheim, 2004, vols 1 and 2.

Scheme 6



Chemistry based on "C-H activation" can also be used to prepare a compound of Formula **2D** (i.e. a compound of Formula **2** wherein Z is O; R^A is $-\text{C}(\text{O})\text{CH}_3$; and m is 1 at the 3-position) as shown in Scheme 7 utilizing palladium(II) acetate and (diacetoxyiodo)benzene as described above for Scheme 6. A compound of Formula **2D** can subsequently be converted via methods disclosed in Schemes 1 and 6 to provide a compound of Formula **1**.


Scheme 7



Similarly, chemistry based on "C-H activation" can be used to prepare a compound of Formulae **2F** (i.e. a compound of Formula **2A** wherein Z is S) as shown in Scheme 8. A compound of Formula **8** can first be converted to a compound of Formula **8A** (i.e. a compound of Formula **6** wherein the *ortho* "H" is X; and X is Br or I) by utilizing a stepwise introduction of substituents using "C-H activation". Iodides and bromides of Formula **8A** can then be further functionalized by copper mediated cross-coupling with thiourea as


described in Qi, Junsheng, *Chin. J. Chem.* **2010**, 28, 1441–1443 to provide the aryl thiol after acidic deprotection. Palladium catalyzed cross-coupling reactions of aryl halides can give protected thiols that can, in turn, be deprotected under either acidic conditions or basic conditions (e.g. cesium fluoride) to provide a compound of Formula **2F**. These conditions are discussed in Organ, Michael G., *Angew. Chem. Int. Ed.* **2012**, 51, 3314–3322 and the references cited therein. Also, relevant conditions can be found in Takashiro Itoh, *J. Org. Chem.* **2006**, 71, 2203–2206. A compound of Formula **2F** can then be converted via methods disclosed in Schemes 1 and 7 to provide a compound of Formula **1**.

Scheme 8



In Scheme 9, the phenol, **2E** is reacted with *N,N*-dimethyl thiocarbamoyl chloride in *N,N*-dimethylformamide in the presence of a strong tertiary amine base such as 1,4-diazabicyclo[2.2.2]octane or *N*-methylmorpholine for acidic phenols (for less-acidic phenols, prior deprotonation with sodium hydride may be advantageous) to form the *O*-aryl *N,N*-dimethylthiocarbamate of Formula **10**. Newman-Kwart rearrangement of a compound of Formula **10** at temperatures ranging from 200 to 300 °C provides the intermediate *S*-aryl dimethylthiocarbamate of Formula **11**. A one-pot deprotection of a compound of Formula **11** is readily achieved using 10% aqueous sodium hydroxide or methanolic potassium hydroxide to afford the corresponding aryl thiol. Subsequent reaction with a compound of Formula **3** at or slightly above room temperature provides the product **1C** (i.e. a compound of Formula **1** wherein Z is S). Methods for Newman-Kwart rearrangements are found in Lloyd-Jones, Guy C., *Synthesis* **2008**, 661-689.

Scheme 9



It is recognized by one skilled in the art that various functional groups can be converted into others to provide different a compound of Formula 1. For a valuable resource that illustrates the interconversion of functional groups in a simple and straightforward fashion, see Larock, R. C., *Comprehensive Organic Transformations: A Guide to Functional Group Preparations*, 2nd Ed., Wiley-VCH, New York, 1999. For example, intermediates for the preparation of a compound of Formula 1 may contain aromatic nitro groups, which can be reduced to amino groups, and then be converted via reactions well known in the art such as the Sandmeyer reaction, to various halides, providing a compound of Formula 1. The above reactions can also in many cases be performed in alternate order

It is recognized that some reagents and reaction conditions described above for preparing a compound of Formula 1 may not be compatible with certain functionalities present in the intermediates. In these instances, the incorporation of protection/deprotection sequences or functional group interconversions into the synthesis will aid in obtaining the desired products. The use and choice of the protecting groups will be apparent to one skilled in chemical synthesis (see, for example, Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 4th ed.; Wiley: Hoboken, New Jersey, 1991). One skilled in the art will recognize that, in some cases, after the introduction of a given reagent as it is depicted in any individual scheme, it may be necessary to perform additional routine synthetic steps not described in detail to complete the synthesis of a compound of Formula 1. One skilled in the art will also recognize that it may be necessary to perform a combination of the steps illustrated in the above schemes in an order other than that implied by the particular presented to prepare a compound of Formula 1.

One skilled in the art will also recognize that a compound of Formula 1 and the intermediates described herein can be subjected to various electrophilic, nucleophilic,

radical, organometallic, oxidation, and reduction reactions to add substituents or modify existing substituents.

Without further elaboration, it is believed that one skilled in the art using the preceding description can utilize the present invention to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative, and not limiting of the disclosure in any way whatsoever. Steps in the following Examples illustrate a procedure for each step in an overall synthetic transformation, and the starting material for each step may not have necessarily been prepared by a particular preparative run whose procedure is described in other Examples or Steps. Percentages are by weight except for chromatographic solvent mixtures or where otherwise indicated. Parts and percentages for chromatographic solvent mixtures are by volume unless otherwise indicated. ¹H NMR spectra are reported in ppm downfield from tetramethylsilane in CDCl₃; “s” means singlet, “d” means doublet, “t” means triplet, “q” means quartet, “m” means multiplet, “dd” means doublet of doublets, “dt” means doublet of triplets, and “bs” means broad singlet.

It is recognized by one skilled in the art that various functional groups can be converted into others to provide different compounds of Formula 1. For a valuable resource that illustrates the interconversion of functional groups in a simple and straightforward fashion, see Larock, R. C., *Comprehensive Organic Transformations: A Guide to Functional Group Preparations*, 2nd Ed., Wiley-VCH, New York, 1999. For example, intermediates for the preparation of compounds of Formula 1 may contain aromatic nitro groups, which can be reduced to amino groups, and then be converted via reactions well known in the art such as the Sandmeyer reaction, to various halides, providing compounds of Formula 1. The above reactions can also in many cases be performed in alternate order

It is recognized that some reagents and reaction conditions described above for preparing compounds of Formula 1 may not be compatible with certain functionalities present in the intermediates. In these instances, the incorporation of protection/deprotection sequences or functional group interconversions into the synthesis will aid in obtaining the desired products. The use and choice of the protecting groups will be apparent to one skilled in chemical synthesis (see, for example, Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 2nd ed.; Wiley: New York, 1991). One skilled in the art will recognize that, in some cases, after the introduction of a given reagent as it is depicted in any individual scheme, it may be necessary to perform additional routine synthetic steps not described in detail to complete the synthesis of compounds of Formula 1. One skilled in the art will also recognize that it may be necessary to perform a combination of the steps illustrated in the above schemes in an order other than that implied by the particular presented to prepare the compounds of Formula 1.

One skilled in the art will also recognize that compounds of Formula 1 and the intermediates described herein can be subjected to various electrophilic, nucleophilic,

radical, organometallic, oxidation, and reduction reactions to add substituents or modify existing substituents.

Without further elaboration, it is believed that one skilled in the art using the preceding description can utilize the present invention to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative, and not limiting of the disclosure in any way whatsoever. Steps in the following Examples illustrate a procedure for each step in an overall synthetic transformation, and the starting material for each step may not have necessarily been prepared by a particular preparative run whose procedure is described in other Examples or Steps. Percentages are by weight except for chromatographic solvent mixtures or where otherwise indicated. Parts and percentages for chromatographic solvent mixtures are by volume unless otherwise indicated. ¹H NMR spectra are reported in ppm downfield from tetramethylsilane at 500 MHz in CDCl₃ unless otherwise indicated; “s” means singlet, “d” means doublet, “t” means triplet, “q” means quartet, “m” means multiplet, “dd” means doublet of doublets and “dt” means doublet of triplets.

SYNTHESIS EXAMPLE 1

Synthesis of 3-[2-[(5-chloro-2-pyrimidinyl)oxy]phenyl]-5-isoxazolemethanol
(Compound 31)

Step A: Synthesis of 5-chloro-2-[2-[5-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]methyl]-3-isoxazolyl]phenoxy]pyrimidine

To a solution of 3-(2-methoxyphenyl)-5-isoxazolemethanol (prepared as described in *Bioorganic Med. Chem.* **2004**, *12*, 3965 (0.500 mg, 0.243 mmol) in tetrahydrofuran (25 mL) was added *t*-butyldiphenylsilyl chloride (0.804 mg, 2.92 mmol) followed by imidazole (0.199 mg, 2.92 mmol). After 2 h the solvent was removed under vacuum. Purification using chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes to afford the intermediate 5-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]methyl]-3-(2-methoxyphenyl)isoxazole and the material was taken on without further purification.

To a solution of 5-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]methyl]-3-(2-methoxyphenyl)isoxazole (3.10 g, 0.699 mmol) in dichloromethane (35 mL) at 0 °C was added a 1.0 M solution of boron tribromide (34.9 mL) and the reaction was stirred at this temperature for 1 h. The reaction was quenched with a saturated solution of sodium bicarbonate. The phases were separated, and the aqueous layer was washed with additional dichloromethane. The combined organic phases were combined, dried with MgSO₄ and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes afforded 2-[5-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]methyl]-3-isoxazolyl]phenol which was taken to the next step without further purification.

To a solution of 2-[5-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]methyl]-3-isoxazolyl]phenol (2.78 g, 6.47 mmol) in acetonitrile (60 mL) was added 2,5-dichloropyrimidine (1.15 g, 7.70 mmol) and potassium carbonate (2.24 g, 16.2 mmol) and the reaction was heated to 80 °C for 6 h. The reaction mixture was allowed to warm to room temperature and the solvent was removed under vacuum. Purification by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes afforded the title product (2.27 g).

¹H NMR δ 8.41 (s, 2H), 8.00–7.97 (m, 1H), 7.67–7.61 (m, 4H), 7.56–7.50 (m, 1H), 7.47–7.36 (m, 7H), 7.28–7.26 (m, 1H), 6.56 (t, 1H), 1.05 (s, 9H). MS (AP⁺) = 542.

10 Step B: Synthesis of 3-[2-[(5-chloro-2-pyrimidinyl)oxy]phenyl]-5-isoxazolemethanol

To a solution of 5-chloro-2-[5-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]methyl]-3-isoxazolyl]phenoxy]pyrimidine (i.e. the product from Step A) (2.27 g, 4.19 mmol) in tetrahydrofuran (15 mL) was added acetic acid (0.50 mL) followed by a solution of 75% tetrabutylammonium fluoride in water (2.9 mL) and the reaction was allowed to stir for 2 h.

15 The reaction was quenched with a saturated solution of sodium bicarbonate and the phases were partitioned and the aqueous phase was further washed with ethyl acetate. The combined organic phases were combined, dried over MgSO₄ and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes afforded the title product, a compound of the present invention (1.21 g).

20 ¹H NMR δ 8.45 (s, 2H), 7.99–7.95 (m, 1H), 7.56–7.52 (m, 1H), 7.42–7.38 (m, 1H), 7.28–7.25 (m, 1H), 6.64–6.61 (m, 1H), 4.77–4.73 (m, 2H).

SYNTHESIS EXAMPLE 2

Synthesis of 3-[2-[(5-chloro-2-pyrimidinyl)oxy]phenyl]-5-isoxazolecarboxaldehyde (Compound 33)

25 Step A: Synthesis of 3-[2-[(5-chloro-2-pyrimidinyl)oxy]phenyl]-5-isoxazolecarboxaldehyde

Pyridinium chlorochromate (263 mg, 1.22 mmol) and silica gel (200 mg) were combined and mixed as solids. This mixture was then added to a stirring solution of 3-[2-[(5-chloro-2-pyrimidinyl)oxy]phenyl]-5-isoxazolemethanol (i.e. the product obtained in Step A of Example 1) (309 mg, 1.02 mmol) in dichloromethane (5.0 mL) and the reaction was allowed to stir for 18 h. The solution was filtered to remove the silica gel and the organic phase was washed with a 1 M hydrochloric acid solution. The organic phase was dried with MgSO₄ and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes to afforded the title product, a compound of the present invention (0.307 g).

35 ¹H NMR δ 9.95 (s, 1H), 8.46 (s, 2H), 8.06–8.01 (m, 1H), 7.61–7.56 (m, 1H), 7.47–7.41 (m, 1H), 7.38 (s, 1H), 7.32–7.29 (m, 1H). MS (AP⁺) = 302.

SYNTHESIS EXAMPLE 3

Synthesis of 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine
(Compound 35)

Step A Synthesis of 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine

To a stirred solution of 3-[2-[(5-chloro-2-pyrimidinyl)oxy]phenyl]-5-isoxazolecarboxaldehyde (i.e. the product from Step A of Example 2) (100 mg, 0.332 mmol) in dichloromethane (3.0 mL) at -78°C was added Deoxo-Fluor[®] (161 mg, 0.729 mmol) and the reaction was allowed to return to ambient temperature. Upon consumption of the starting material as evidenced by thin-layer chromatography, the solvent was removed under vacuum. Purification by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes to afforded the title product, a compound of the present invention (36.3 mg).

^1H NMR δ 8.46 (s, 2H), 8.01–7.96 (m, 1H), 7.59–7.54 (m, 1H), 7.44–7.38 (m, 1H), 7.31–7.27 (m, 1H), 6.98–6.96 (s, 1H), 6.83–6.60 (m, 1H). MS (ESI⁺) = 324.

SYNTHESIS EXAMPLE 4

Synthesis of 2-[2-(3-bromo-5-isoxazolyl)phenoxy]-5-chloropyrimidine (Compound 12)

Step A: Synthesis of 3-bromo-5-(2-methoxyphenyl)isoxazole

To a solution of 1-ethynyl-2-methoxybenzene (0.78 g, 5.92 mmol) in dichloromethane (10 mL) was added dibromoformaldoxime (1.00 g, 4.93 mmol). The mixture was cooled to 0°C and potassium bicarbonate (1.48 g, 14.8 mmol) was added, followed by heating to 40°C for 18 h. Water was added to the reaction mixture, the phases separated, and the aqueous layer was again washed with dichloromethane. The combined organic phases were dried over MgSO_4 , concentrated under vacuum, and purified by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes to afford the title product, a compound of the present invention (1.04 g).

^1H NMR δ 7.94 (dd, 1H), 7.47–7.42 (m, 1H), 7.09 (dd, 1H), 7.02 (dd, 1H), 6.85 (s, 1H), 3.97 (s, 3H). MS (AP⁺) = 254.

Step B: Synthesis of 2-(3-bromo-5-isoxazolyl)phenol

To a solution of 3-bromo-5-(2-methoxyphenyl)isoxazole (i.e. the product from Step A) (0.50 g, 1.97 mmol) in dichloromethane (20mL) was added a 1 M solution of boron tribromide in dichloromethane (9.86 mmol) at -78°C and the solution was allowed to warm to room temperature and stir for 18 h. Dichloroethane (20 mL) was added, and reaction mixture was concentrated to remove the excess dichloromethane. Boron tribromide in dichloromethane (9.86 mmol) was again added and the reaction was heated to 80°C until completion as evidenced by thin-layer chromatography. The reaction mixture was allowed to cool to ambient temperature and quenched with a saturated solution of sodium bicarbonate. The phases were separated and the aqueous layer was again washed with

dichloromethane. The combined organic phases were dried with MgSO_4 , concentrated under vacuum, and purified by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes to afford the title product (0.395 g).

^1H NMR δ 7.76 (dd, 1H), 7.39–7.33 (m, 1H), 7.09–7.02 (m, 1H), 6.96–6.93 (m, 1H), 6.02 (s, 1H). MS (AP^-) = 238.

Step C: 2-[2-(3-bromo-5-isoxazolyl)phenoxy]-5-chloropyrimidine

To a solution of 2-(3-bromo-5-isoxazolyl)phenol (i.e. the product from Step B) (100 mg, 0.417 mmol) in acetonitrile (5 mL) was added 2,5-dichloropyrimidine (75.0 mg, 0.503 mmol) and potassium carbonate (288 mg, 2.08 mmol) then the solution was stirred at ambient temperature for 18 h. The reaction was then heated at 40 °C for 2 h followed by 80 °C for two hours. The solution was then cooled to ambient temperature, water was added, the phases were separated and the aqueous layer was again washed with dichloromethane. The combined organic phases were dried over MgSO_4 , concentrated under vacuum, and purified by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes to afford the title product, a compound of the present invention (122 mg).

^1H NMR δ 8.49 (s, 2H), 8.03 (dd, 1H), 7.58–7.53 (m, 1H), 7.43 (dt, 1H), 7.29 (dd, 1H), 6.74 (s, 1H). MS (AP^+) = 352.

SYNTHESIS EXAMPLE 5

Synthesis of 5-chloro-2-[2-[4-(trifluoromethyl)-2-pyridinyl]phenoxy]pyrimidine
(Compound 25)

Step A: Synthesis of 2-[4-(trifluoromethyl)-2-pyridinyl]phenol

2-Chloro-4-trifluoromethylpyridine (1.0 g, 5.5 mmol) and 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenol (1.57 g, 7.16 mmol) were combined in dimethoxyethane (18 mL) and water (1.8 mL). To this mixture were added sodium carbonate (2.28 g, 16.5 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.32 g, 0.27 mmol). The reaction was heated at 90 °C for 2.5 h and allowed to stir at 23 °C for 18 h. The mixture was diluted with water (20 mL) and dichloromethane (20 mL) and the layers separated. The aqueous layer was washed with dichloromethane (10 mL). The combined dichloromethane layers were washed with saturated aqueous sodium chloride solution (10 mL) and dried over sodium sulfate. After filtration the organic layer was evaporated and the solid thus obtained was triturated with hexanes (20 mL). The filtrate was concentrated to provide 1.18 g of the title compound as a yellow solid which was used in Step B without further purification.

^1H NMR δ 13.61 (s, 1H), 8.72 (d, 1H), 8.12 (s, 1H), 7.83 (d, 1H), 7.47 (m, 1H), 7.36 (s, 1H), 7.06 (d, 1H), 6.96 (t, 1H).

Step B Synthesis of 5-chloro-2-[2-[4-(trifluoromethyl)-2-pyridinyl]phenoxy]pyrimidine

2-[4-(Trifluoromethyl)-2-pyridinyl]phenol (i.e. the product from Step A) (0.20 g, 0.84 mmol) and 2,5-dichloropyrimidine (0.14 g, 0.92 mmol) were dissolved in acetonitrile (2 mL) and treated with powdered potassium carbonate (0.34 g, 2.5 mmol). The mixture was heated to 80 °C for 18 h. After cooling, the reaction mixture was diluted with water (10 mL) and ethyl acetate (10 mL) and the layers separated. The aqueous layer was washed with ethyl acetate (10 mL). The combined ethyl acetate solution was washed with saturated aqueous sodium chloride solution (10 mL) and dried over MgSO₄. The filtrate was evaporated under reduced pressure and subjected to chromatography through 12 g silica gel eluting with 10 to 20% ethyl acetate in hexanes. Appropriate fractions were pooled and evaporated to provide the title compound, a compound of the present invention (0.2 g) as a clear oil.

¹H NMR δ 8.75 (d, 1H), 8.39 (s, 2H), 7.91 (s, 1H), 7.88 (m, 1H), 7.54 (m, 1H), 7.44 (m, 1H), 7.37 (d, 1H), 7.28 (m, 1H).

SYNTHESIS EXAMPLE 6

Synthesis of 5-chloro-2-[4-methyl-2-[6-(trifluoromethyl)-3-pyridinyl]phenoxy]pyrimidine (Compound 22)

Step A: Synthesis of 2-(2-bromo-4-methylphenoxy)-5-chloropyrimidine

2-Bromo-4-methylphenol (280 mg, 1.5 mmol) and 2,5-dichloropyrimidine (246 mg, 1.65 mmol) were combined in 6 mL of acetonitrile under a nitrogen atmosphere. Powdered potassium carbonate (455 mg, 3.3 mmol) was added and the resulting mixture was heated at reflux for 6 h. The reaction mixture was cooled and diluted with de-ionized water and ethyl acetate. The aqueous layer was separated and extracted twice with ethyl acetate. The combined organic layers were washed with brine, dried over MgSO₄, filtered and concentrated under reduced pressure. The resulting residue was purified by medium pressure liquid chromatography on silica gel eluting with 0 to 15% ethyl acetate in hexanes to yield the title compound (270 mg).

¹H NMR δ 8.48 (s, 2H), 7.47 (d, 1H), 7.18 (m, 1H), 7.11 (m, 1H), 2.37 (s, 3H).

Step B Synthesis of 5-chloro-2-[4-methyl-2-[6-(trifluoromethyl)-3-pyridinyl]phenoxy]pyrimidine

A mixture of 2-(2-bromo-4-methylphenoxy)-5-chloropyrimidine (i.e. the product of Step A; 190 mg, 0.63 mmol), *B*-[6-(trifluoromethyl)-3-pyridinyl]-boronic acid (133 mg, 0.70 mmol), sodium carbonate (0.6 mL of 2 M aqueous solution, 1.26 mmol) and tetrakis(triphenylphosphine)palladium(0) (73 mg, 0.06 mmol) in toluene (9 mL) and ethanol (1 mL) was heated at 90 °C for 2 h. The reaction mixture was then concentrated under reduced pressure, and the residue was purified by medium pressure liquid chromatography

on silica gel eluted with 0 to 10% ethyl acetate in hexanes to yield the title compound, a compound of the present invention (190 mg).

^1H NMR δ 8.77 (d, 1H), 8.36 (s, 2H), 8.02 (m, 1H), 7.64 (d, 1H), 7.31 (m, 2H), 7.15 (d, 1H), 2.45 (s, 3H).

5

SYNTHESIS EXAMPLE 7

Synthesis of 5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]phenoxy]pyrimidine
(Compound 53)

Step A: Synthesis of 2-(3-difluoromethyl-5-isoxazolyl)phenol

To a solution of 25% sodium methoxide in methanol (5 mL) and tetrahydrofuran (10 mL), acetophenone (1 g, 7.3 mmol) and difluoroacetate (1 g, 8.1 mmol) in tetrahydrofuran (2 mL) was added and heated at 60 °C for 5 h. The reaction was cooled to room temperature and treated with 36% aq hydrochloric acid (4 mL) and stirred at 60 °C for 2 h. The reaction was quenched by adding water (15 mL) and the organic solvent was removed under vacuum. The precipitated product 2-difluoromethyl-4-chromenone (1.4 g) was filtered and dissolved in ethanol (5 mL). To this solution, hydroxylamine acetate (22 mmol) in water (5 mL) was added and the mixture was heated at 60 °C for 3 h. After cooling the reaction to ambient temperature 4,4-difluoro-1-(2-hydroxyphenyl)-butane-1,3-dione 3-oxime was precipitated with the addition of water (20 mL). This product was collected by filtration and suspended in acetic acid (5 mL) and 36% aqueous hydrochloric acid (1.8 mL) at room temperature and stirred at 80 °C for 15 min to obtain the title compound as a beige solid (800 mg).

^1H NMR δ 7.82 (m, 1 H), 7.36 (s, 1 H), 7.07 (m, 1 H), 6.95 (m, 2H), 6.82 (t, 1H), 6.05 (s, 1H). MS (ESI^+) = 212

Step B: Synthesis of 5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]phenoxy]-pyrimidine

To a solution of 2-(3-difluoromethyl-5-isoxazolyl)phenol (i.e. the product from Step A) (2.1 g, 9.71 mmol) in anhydrous *N,N*-dimethylformamide (8 mL) was added 2,5-dichloropyrimidine (1.5 g, 10.2 mmol) and potassium carbonate (2.9 g, 21.3 mmol). The reaction was heated at 90 °C for 1 h. The solution was cooled to ambient temperature and diluted with water. The phases were separated and the aqueous phase was washed with additional ethyl acetate. The organic phases were combined, dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 10% ethyl acetate in hexanes afforded the title compound, a compound of the present invention, as a solid (2.2 g).

^1H NMR δ 8.49 (s, 2H), 8.06 (m, 1H), 7.57 (m, 1H), 7.44 (m, 1H), 7.31 (m, 1H), 6.88 (s, 1H), 6.74 (t, 1H). MS (ESI^+) = 324

SYNTHESIS EXAMPLE 8

Synthesis of 5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine
(Compound 144)

Step A: Synthesis of 4,4-difluoro-1-(2-fluoro-6-methoxyphenyl)butane-1,3-dione

To a solution of 1-(2-fluoro-6-methoxyphenyl)ethanone (2.6 g, 15.5 mmol) and difluoroacetic acid ethyl ester (3.9 mL, 31.0 mmol) in anhydrous *N,N*-dimethylformamide at 0 °C was added sodium hydride (1.2 g, 31.0 mmol). The reaction mixture was heated at 80 °C for 1 h. The reaction was then cooled to 0 °C, diluted with ethyl acetate and acidified with 1 N aqueous hydrochloric acid. The phases were separated and the aqueous phase was washed with additional ethyl acetate. The organic phases were combined and dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 15% ethyl acetate in hexanes afforded the title compound (2.5 g). ¹H NMR δ 7.39 (m, 1H), 6.77 (m, 2H), 6.24 (s, 1H), 6.01 (t, 1 H), 3.87 (s, 3 H). MS (ESI⁺) = 247

Step B: Synthesis of 3-difluoromethyl-5-(2-fluoro-6-methoxyphenyl)isoxazole

A solution of 4,4-difluoro-1-(2-fluoro-6-methoxyphenyl)butane-1,3-dione (i.e. the product from Step A) (2.5 g, 10 mmol) and hydroxylamine hydrochloride (2.1 g, 30 mmol) in ethanol (25 mL) was stirred at 80 °C. After 1 h the solvent was removed under vacuum. The resulting residue was diluted with water and extracted with dichloromethane. The organic phase was dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 15% ethyl acetate in hexanes afforded the title compound (1.5 g).

¹H NMR δ 7.41 (m, 1 H), 6.69 - 6.98 (m, 4H), 3.93 (s, 3 H). MS (ESI⁺) = 244

Step C: Synthesis of 2-(3-difluoromethyl-5-isoxazolyl)-3-fluorophenol

To a solution of 3-difluoromethyl-5-(2-fluoro-6-methoxyphenyl)isoxazole (i.e. the product from Step B) (1.5 g, 6.2 mmol.) in dichloromethane (10 mL) at 0 °C was added a 1.0 M solution of boron tribromide in dichloromethane (31 mL, 31 mmol). The reaction mixture was warmed to ambient temperature and stirred for 6 h. The reaction was cooled to 0 °C and slowly quenched with a saturated aqueous solution of sodium bicarbonate. The biphasic mixture was stirred at room temperature for 1 h. The phases were separated and the aqueous phase was extracted with dichloromethane. The combined organic phases were dried and concentrated under vacuum. The crude residue was purified by chromatography on silica gel, eluting with 0 to 10% ethyl acetate in hexanes, to afford the title compound (980 mg).

¹H NMR δ 7.33 (m, 1 H), 6.66 - 6.99 (m, 4 H). MS (ESI⁺) = 230

Step D: Synthesis of 5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine

To a solution of 2-(3-difluoromethyl-5-isoxazolyl)-3-fluorophenol (i.e. the product from Step C) (120 mg, 0.5 mmol) in anhydrous *N,N*-dimethylformamide (2 mL) was added 2,5-dichloropyrimidine (85 mg, 0.57 mmol) and potassium carbonate (244 mg, 1.04 mmol). The reaction was heated at 80 °C for 4 h. The mixture was cooled to ambient temperature and diluted with water. The phases were separated and the aqueous phase was washed with additional ethyl acetate. The organic phases were combined, dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 15% ethyl acetate in hexanes afforded the title compound, a compound of the present invention, as a solid (110 mg).

¹H NMR δ 8.46 (s, 2H), 7.56 (m, 1H), 7.21 (m, 1H), 7.13 (m, 1H), 6.87 (m, 1 H), 6.74 (t, 1 H). MS (ESI⁺) = 342

SYNTHESIS EXAMPLE 9

Synthesis of 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine (Compound 55)

Step A: Synthesis of 4,4-difluoro-1-(2-fluoro-6-methoxyphenyl)butane-1,3-dione

A solution of 2-fluoro-6-methoxyacetophenone (6.83 g, 40.6 mmol) and ethyl difluoroacetate (7.45g, 60 mmol) in tetrahydrofuran (35 mL) was added dropwise to a solution of tetrahydrofuran (20 mL) and 25% sodium methoxide (10.2g, 47.2 mmol) over 15 minutes. The reaction was complete in 3h as determined by high pressure liquid chromatography. The reaction was partially concentrated under vacuum to remove most of the tetrahydrofuran and methanol, and then diluted with toluene and water. The aqueous phase was acidified with 37% hydrochloric acid (5g), followed by extraction with toluene. The combined organic phases were concentrated under vacuum to provide title compound (7.98 g).

¹H NMR δ 7.39 (td, 1 H) 6.72 - 6.81 (m, 2 H) 6.25 (d, 1 H) 5.87 - 6.14 (m, 1 H) 3.88 (s, 3 H).

Step B: Synthesis of 5-(difluoromethyl)-3-(2-fluoro-6-methoxyphenyl)-4*H*-isoxazol-5-ol

To a solution of 4,4-difluoro-1-(2-fluoro-6-methoxyphenyl)butane-1,3-dione (i.e. the product from Step A) (7.98 g, 32.4 mmol) in 35 mL methanol at room temperature was added a 50% hydroxylamine solution (2.78g, 42.1 mmol) and 1N sodium hydroxide solution (1.50 mL, 1.50 mmol) followed by heating to 65 °C for 2h. The reaction was cooled to room temperature and diluted with water and toluene. The phases were separated and the organic phase was concentrated under vacuum to provide the title compound (7.99 g).

¹H NMR δ 7.36 (td, 1 H) 6.73 - 6.82 (m, 2 H) 5.79 - 6.05 (m, 1 H) 3.88 (s, 3 H) 3.67 - 3.73 (m, 1 H) 3.47 - 3.51 (m, 1 H) 3.34 - 3.42 (m, 1 H).

Step C: Synthesis of 5-(difluoromethyl)-3-(2-fluoro-6-methoxyphenyl)isoxazole

To toluene (80 mL) was added 5-(difluoromethyl)-3-(2-fluoro-6-methoxyphenyl)-4H-isoxazol-5-ol (i.e. the product from Step B) (7.99 g, 30.6 mmol) followed by *p*-toluenesulfonic acid monohydrate (0.700 g, 3.68 mmol). The mixture was heated to a vigorous reflux (107 – 111 °C) for two hours at which point high pressure liquid chromatography determined the reaction was complete. The cooled reaction mixture was washed with a saturated sodium bicarbonate solution, followed by water. The organic phase was concentrated under vacuum to provide the title compound (7.44 g).

¹H NMR δ 7.40 (td, 1 H) 6.69 - 6.94 (m, 4 H) 3.88 (s, 3 H).

Step D: Synthesis of 2-[5-(difluoromethyl)-3-isoxazolyl]-3-fluorophenol

To a solution of 5-(difluoromethyl)-3-(2-fluoro-6-methoxyphenyl)isoxazole (i.e. the product from Step C) (3.72 g, 15.3 mmol) in dichloromethane (15 mL) at 3 °C was added a 1M solution of boron tribromide in dichloromethane (18.0 mL, 18 mmol) over 5 min. The reaction was then allowed to warm to room temperature. After 90 minutes it was determined the reaction was complete using high pressure liquid chromatography and the reaction was treated with a 10% aqueous solution of potassium bicarbonate (10 mL). The phases were separated and the organic phase was concentrated under vacuum. The resulting brown solid was triturated with a water/methanol solution (~2/1) providing the title compound (3.34 g).

¹H NMR δ 9.63 - 9.75 (m, 1 H) 7.33 (td, 1 H) 7.21 (ddd, 1 H) 6.71 - 6.96 (m, 3 H).

Step E: Synthesis of 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine

To a solution of 2-[5-(difluoromethyl)-3-isoxazolyl]-3-fluorophenol (i.e. the product from Step D) (1.61 g, 7.02 mmol) and 5-methyl-2-methylsulfonylpyrimidine (1.49 g, 7.72 mmol) in *N,N*-dimethylformamide (9 mL) was added potassium carbonate (4.24 g, 17.5 mmol) and the reaction was allowed to stir at room temperature for 24 h. The reaction was diluted with water and toluene, the phases were separated and the organic solvent was removed under vacuum. To the resulting oil was added 8 mL of methanol and a tan slurry formed, after further dilution with a methanol/water solution (20 mL), the precipitate was filtered providing the title compound, a compound of the present invention (2.24g).

¹H NMR δ 8.45 (s, 2 H) 7.54 (td, 1 H) 7.19 (ddd, 1 H) 7.14 (dt, 1 H) 6.88 (dt, 1 H) 6.61 - 6.85 (m, 1 H).

SYNTHESIS EXAMPLE 10

Preparation of 5-chloro-2-[3-cyano-2-[4-(trifluoromethyl)-2-pyridinyl]phenoxy]pyrimidine
(Compound 158)

A solution of 5-chloro-2-[2-[4-(trifluoromethyl)-2-pyridinyl]phenoxy]pyrimidine (i.e. the product of example 5, step B) (0.30 g, 0.853 mmols) in 4.27 mL of *N,N*-dimethylformamide under a nitrogen atmosphere was treated with copper(II) bromide (0.19 g, 0.853 mmols), palladium(II) acetate (9 mg, 0.0426 mmols) and potassium ferricyanide (0.06 g, 0.17 mmols). The mixture was heated at 130°C for 18 hours. The mixture was then cooled, diluted with diethyl ether and water, filtered thru a celite pad and rinsed with ethyl acetate and water. The phases were separated and the aqueous phase was extracted twice with diethyl ether. The combined organic phases were washed twice with water and saturated aqueous sodium chloride, dried over magnesium sulfate, and concentrated to provide 0.21g crude product. The crude product was purified with a 12 g Teledyne Isco silica gel column eluting with 10 to 30% EtOAc-Hexanes gradient to provide the title compound, a compound of the present invention as a solid (0.23 g).

^1H NMR δ 8.86 (d, 1H), 8.40 (s, 2H), 7.78 (d&s, 2H), 7.62 (t, 1H), 7.52 (d, 1H), 7.50 (d, 1H).

SYNTHESIS EXAMPLE 11

Synthesis of 5-chloro-2-[2-[5-(trifluoromethyl)-2-pyridinyl]phenoxy]pyrimidine (Compound 27)

Step A: Synthesis of 2-[5-(trifluoromethyl)-2-pyridinyl]phenol

2-Chloro-5-(trifluoromethyl)pyridine (1.0 g, 5.50 mmols) and 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenol (1.57 g, 7.16 mmols) were combined in 16 mL of 1,2-dimethoxyethane and 1.8 mL of de-ionized water under a nitrogen atmosphere. Solid sodium carbonate (2.28g, 16.5 mmols) and then tetrakis(triphenylphosphine) palladium (0) (0.32g, 0.27 mmols) were added. The reaction was heated at reflux approximately ninety minutes. The reaction was cooled, diluted with dichloromethane and filtered thru a celite pad, rinsing with dichloromethane and then de-ionized water. The phases were separated. The aqueous phase was extracted twice with dichloromethane. The combined organic phases were washed with saturated aqueous sodium chloride, dried over magnesium sulfate, filtered and concentrated to give a solid. A solid was filtered from hexanes to give 34 mg. A second crop was obtained from the filtrate from hexanes to yield 506 mg of a light orange-brown solid of the title compound.

^1H NMR δ 8.81 (s, 1H), 8.04 (m, 2H), 7.83 (d, 1H), 7.38 (t, 1H), 7.07 (d, 1H), 6.98 (t, 1H).

Step B: Synthesis of 5-chloro-2-[2-[5-(trifluoromethyl)-2-pyridinyl]phenoxy]-pyrimidine

A mixture of 2-[5-(trifluoromethyl)-2-pyridinyl]phenol (i.e. the product of step A) (0.20 g, 0.836 mmoles) and 2,5-dichloropyrimidine (0.14 g, 0.919 mmoles) in 2.0 mL of N,N-dimethylformamide was stirred under a nitrogen atmosphere. Powdered potassium carbonate (0.35 g, 2.51 mmoles) was added and the mixture was heated at 80 °C overnight. The reaction was cooled before diluting with de-ionized water and diethyl ether. The phases were separated. The aqueous phase was extracted twice with diethyl ether. The combined organic phases were washed three times with de-ionized water, dried over sodium sulfate, filtered and concentrated to 0.37 g of solid. A solid was filtered from hexanes and some diethyl ether to give 103 mg of the title compound, a compound of the present invention. ¹H NMR δ 8.84 (s, 1H), 8.40 (s, 2H), 7.92(d, 1H), 7.87 (s&d, 2H), 7.54 (t, 1H), 7.45 (t, 1H), 7.27 (d,1H).

SYNTHESIS EXAMPLE 12

Preparation of 5-chloro-2-[2-[5-(trifluoromethyl)-2-pyridinyl]-3-chlorophenoxy]pyrimidine (Compound 160)

5-Chloro-2-[2-[5-(trifluoromethyl)-2-pyridinyl]phenoxy]-pyrimidine (i.e. the product of example 11, step B) (0.14 g, 0.398 mmoles) was dissolved in 2 mL of acetic acid. Palladium acetate (0.01 g, 0.039 mmoles) and *N*-chlorosuccinimide (0.11 g, 0.796 mmoles) were added and the mixture was heated at 100 °C for three hours. The mixture was cooled to room temperature overnight and then diluted with toluene and ethyl acetate. The mixture was filtered thru a celite pad, rinsed with toluene and then ethyl acetate. The filtrate was washed twice with saturated aqueous sodium hydrogencarbonate, saturated aqueous sodium chloride, dried over magnesium sulfate and concentrated to a crude product. The crude product was purified with a 12 g Teledyne Isco silica gel column eluting with 10 to 30% EtOAc-Hexanes gradient to provide the title compound, a compound of the present invention as a solid (40 mg).

¹H NMR δ 8.82 (s, 1H), 8.41 (s, 2H), 7.93 (d, 1H), 7.52 (d, 1H), 7.47 (m, 2H), 7.19 (m, 1H).

SYNTHESIS EXAMPLE 13

Synthesis of 5-bromo-2-[2-[5-(difluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine (Compound 62)

Step A: Synthesis of 5-(difluoromethyl)-3-(2-methoxyphenyl)isoxazole

To a solution of acetophenone (3.0 g, 20 mmol) in methanol (15 mL) was added a 30% sodium methoxide in methanol solution (5.0 mL) and the reaction stirred for five minutes. Next ethyl difluoroacetate (2.97 g, 24 mmol) was added and the reaction was heated to reflux for 18 h. The reaction was allowed to cool to room temperature and the solvent was removed under vacuum. To the residue was added 1M hydrochloric acid and ethyl acetate,

the phases were separated, the organic phase was dried using magnesium sulfate and the solvent was removed under vacuum. Purification by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes afforded the desired product which was taken on directly. A solution of 4,4-difluoro-1-(2-methoxyphenyl)butane-1,3-dione from the previous step in ethanol (30 mL) was added dropwise to a solution of hydroxylamine hydrochloride (1.4 g, 20 mmol) in 1M sodium hydroxide (21 mL). The reaction was then heated to reflux for 2 h, followed by cooling to room temperature. The product was precipitated from solution by the addition of water and collected via vacuum filtration to afford the desired product which was taken on directly (AP+ 244, ^1H NMR δ ppm 7.79 (d, 1 H) 7.38 - 7.46 (m, 1 H) 6.93 - 7.04 (m, 2 H) 5.76 - 6.04 (m, 1 H) 3.88 (s, 3 H) 3.67 - 3.80 (m, 1 H) 3.49 - 3.55 (m, 1 H)). Next, a solution of 5-(difluoromethyl)-3-(2-methoxyphenyl)-4*H*-isoxazol-5-ol from the previous step was taken up in trifluoroacetic acid (20 mL) and heated to 70 °C for 18 h. The reaction was cooled to room temperature and the solvent was removed under vacuum. The resulting residue was dissolved in dichloromethane and washed with a saturated solution of sodium bicarbonate. The phases were separated, the organic phase was dried using magnesium sulfate and the solvent was removed under vacuum. Purification by chromatography on silica gel eluting with 0 to 100% ethyl acetate in hexanes afforded the title compound (2.0 g, AP+ = 226).

^1H NMR δ 7.91 (dd, 1 H) 7.42 - 7.50 (m, 1 H) 6.98 - 7.12 (m, 3 H) 6.66 - 6.93 (m, 1 H) 3.92 (s, 3 H).

Step B: Synthesis of 2-[5-(difluoromethyl)-3-isoxazolyl]phenol

To a solution of 5-(difluoromethyl)-3-(2-methoxyphenyl)isoxazole (i.e. the product of step A) (2.01 g, 8.92 mmol) in dichloromethane (50 mL) at 0 °C was added a 1M solution of boron tribromide in dichloromethane (13.3 mL, 13.3 mmol) and the reaction was allowed to warm to room temperature over three hours. The solvent was removed under vacuum and purified by chromatography on silica gel, eluting with 0 to 100% ethyl acetate in hexanes to afford the title compound (1.66g, AP- = 210).

^1H NMR δ 9.14 (s, 1 H) 7.51 (dd, 1 H) 7.39 (ddd, 1 H) 7.11 (dd, 1 H) 6.97 - 7.04 (m, 2 H) 6.70 - 6.95 (m, 1 H).

Step C: Synthesis of 5-bromo-2-[2-[5-(difluoromethyl)-3-isoxazolyl]phenoxy]-pyrimidine

To a solution of 2-[5-(difluoromethyl)-3-isoxazolyl]phenol (i.e. the product of step B) (427 mg, 2.01 mmol) and 5-bromo-2-chloro-pyrimidine (468 mg, 2.42 mmol) in acetonitrile (10 mL) was added potassium carbonate (695 mg, 5.03 mmol) and the reaction was heated to 80 °C for 18 h. The solvent was removed under vacuum and purified by chromatography on silica gel, eluting with 0 to 100% ethyl acetate in hexanes to afford the title compound, a compound of the present invention, as a solid (555 mg, mp = 88.9–92.8 °C).

¹H NMR δ 8.53 (s, 2 H) 7.98 (dd, 1 H) 7.53 - 7.60 (m, 1 H) 7.41 (td, 1 H) 7.28 (dd, 1 H) 6.97 (t, 1 H) 6.59 - 6.84 (m, 1 H).

SYNTHESIS EXAMPLE 14

Synthesis of 5-chloro-2-[2-[5-(trifluoromethyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine
(Compound 168)

Step A: Synthesis of 5-(difluoromethyl)-3-(2-methoxyphenyl)-4*H*-isoxazol-5-ol

To a solution of 2-fluoro-6-methoxyacetophenone (1.0 g, 5.9 mmol) in tetrahydrofuran (2 mL) was added a 30% sodium methoxide in methanol solution (1.4 mL). To this mixture was added dropwise a solution of ethyl trifluoroacetate (0.805 g, 6.49 mmol) in tetrahydrofuran (1 mL) and the reaction stirred at room temperature for 2 h. To the reaction was added 1M hydrochloric acid solution and ethyl acetate, the phases were separated and the aqueous phase was again washed with ethyl acetate. The combined organic phases were dried with magnesium sulfate and concentrated under vacuum to provide the desired product (AP- = 263) which was taken onto the next step directly. Next, to a solution of 4,4,4-trifluoro-1-(2-fluoro-6-methoxyphenyl)butane-1,3-dione (from the previous step) in ethanol (14 mL) was added 1M sodium hydroxide solution (7 mL) followed by hydroxylamine hydrochloride (410 mg, 5.9 mmol) and the reaction stirred overnight at room temperature. The solvent was removed under vacuum and the residue was purified by chromatography on silica gel, eluting with 0 to 100% ethyl acetate in hexanes and 0 to 20% methanol in dichloromethane to afford the product (AP+ = 280, ¹H NMR δ ppm 7.37 (td, 1 H) 6.74 - 6.83 (m, 2 H) 3.89 (s, 3 H) 3.79 (d, 1 H) 3.50 (dd, 1 H)) which was taken onto the next step directly. To a solution of 3-(2-fluoro-6-methoxy-phenyl)-5-(trifluoromethyl)-4*H*-isoxazol-5-ol (from the previous step) in dichloromethane (20 mL) at 0 °C was added a 1M solution of boron tribromide in dichloromethane (11.8 mL, 11.8 mmol) and the reaction was allowed to warm to room temperature over 2 h. The solvent was removed under vacuum. The residue was dissolved in dichloromethane and washed with a saturated solution of sodium bicarbonate, the aqueous phase was washed with dichloromethane. The combined organic phases were dried with magnesium sulfate, concentrated under vacuum and purified by chromatography on silica gel, eluting with 0 to 100% ethyl acetate in hexanes, providing the title compound (606 mg, AP- = 264).

¹H NMR δ 9.86 (s, 1 H) 7.32 (td, 1 H) 6.84 - 6.90 (m, 1 H) 6.68 (ddd, 1 H) 3.86 - 3.94 (m, 1 H) 3.75 (dd, 1 H) 3.56 (s, 1 H).

Step B: Synthesis of 5-chloro-2-[2-[5-(trifluoromethyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine

To a solution of 3-(2-fluoro-6-hydroxyphenyl)-5-(trifluoromethyl)-4*H*-isoxazol-5-ol (i.e. the product of step A) (606 mg, 2.29 mmol) in dimethylsulfoxide (15 mL) was added 5-chloro-2-methylsulfonyl-pyrimidine (527mg, 2.74mmol) followed by cesium carbonate

(1.1g, 3.43mmol) and the reaction stirred for 18 h. The reaction was partitioned between water and ethyl acetate, the phases were separated and the aqueous layer was again washed with ethyl acetate. The combined organic phases were dried with magnesium sulfate and concentrated under vacuum. The residue was purified by chromatography on silica gel, eluting with 0 to 100% ethyl acetate, to afford the title compound, a compound of the present invention, (198mg, AP+ = 360).

^1H NMR δ 8.46 (s, 2 H) 7.56 (td, 1 H) 7.21 (ddd, 1 H) 7.15 (dt, 1 H) 7.02 (dd, 1 H).

SYNTHESIS EXAMPLE 15

Synthesis of 5-chloro-2-[2-[3-(trifluoromethyl)-5-isoxazolyl]phenoxy]pyrimidine (Compound 63)

Step A: Synthesis of 2-(trifluoromethyl)-4*H*-1-benzopyran-4-one

2-Hydroxyacetophenone (10 g, 66.7 mmol) was dissolved in trifluoroacetic anhydride (19 ml, 133.3 mmol) and pyridine (10.8 mL, 133.3 mmol). The reaction mixture was heated at 70 °C and stirred for 12 h. After cooling the reaction mixture was diluted with 1 M hydrochloric acid and methylene chloride and washed with water. The organic phase was dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 5% ethyl acetate in hexanes afforded the title compound (10.5 g) as a pale yellow solid.

^1H NMR δ 8.21 (m, 1 H), 7.76 (m, 1H), 7.77 (d, 1H), 7.46 (m, 1H), 6.73 (s, 1H). MS (ESI⁺) = 215

Step B: Synthesis of 2-[3-(trifluoromethyl)-5-isoxazolyl]phenol

To a solution of 2-(trifluoromethyl)-4*H*-1-benzopyran-4-one (i.e. the product of step A) (10.5 g, 48.8 mmol) in ethanol (50 mL), hydroxylamine acetate (146 mmol) in water (50 mL) was added. The mixture was heated at 60 °C for 4 h. After cooling the reaction to ambient temperature 4,4,4-trifluoro-1-(2-hydroxyphenyl)-butane-1,3-dione 3-oxime was precipitated with the addition of water (200 mL).

^1H NMR δ 9.30 (s, 1H), 7.37 (m, 1H), 7.19 (m, 1H), 7.06 (m, 1H), 6.94 (m, 1H), 3.87 (d, 1H), 3.69 (d, 1H).

This product was collected by filtration and suspended in acetic acid (30 mL) and 36 % aqueous hydrochloric acid (10.8 mL) at room temperature. The mixture was stirred at 80 °C for 30 min to afford the title compound as a white solid (4.6 g).

^1H NMR δ 7.88 (m, 1H), 7.37 (m, 1H), 7.08 (m, 1H), 7.01 (s, 1H), 6.95 (m, 1H). MS (ESI⁺) = 230

Step C: Synthesis of 5-chloro-2-[2-[3-(trifluoromethyl)-5-isoxazolyl]phenoxy]pyrimidine

To a solution of 2-[3-(trifluoromethyl)-5-isoxazolyl]phenol (i.e. the product of step B) (2.2 g, 9.4 mmol) in anhydrous *N,N*-dimethylformamide (10 mL) was added 2,5-

dichloropyrimidine (1.5 g, 10.3 mmol) and potassium carbonate (2.9 g, 20.6 mmol). The reaction was heated at 80 °C for 2 h. The solution was cooled to ambient temperature and diluted with water. The phases were separated and the aqueous phase was washed with additional ethyl acetate. The organic phases were combined, dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 5% ethyl acetate in hexanes afforded the title compound, a compound of the present invention, as a solid (2.1 g).

¹H NMR δ 8.49 (s, 2H), 8.08 (m, 1H), 7.58 (m, 1H), 7.45 (m, 1H), 7.32 (m, 1H), 6.91 (s, 1H). MS (ESI⁺) = 342. Melting Point: 114-115 °C

SYNTHESIS EXAMPLE 16

Synthesis of 5-bromo-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine (Compound 145)

Step A: Synthesis of 4,4-difluoro-1-(2-fluoro-6-methoxyphenyl)butane-1,3-dione

To a solution of 1-(2-fluoro-6-methoxyphenyl)ethanone (2.6 g, 15.5 mmol) and difluoroacetic acid ethyl ester (3.9 mL, 31.0 mmol) in anhydrous *N,N*-dimethylformamide at 0 °C was added sodium hydride (1.2 g, 31.0 mmol). The reaction mixture was heated at 80 °C for 1 h. The reaction was then cooled down to 0 °C, diluted with ethyl acetate and acidified with 1 N aqueous hydrochloric acid. The phases were separated and the aqueous phase was washed with additional ethyl acetate. The organic phases were combined and dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 15% ethyl acetate in hexanes afforded the title compound (2.5 g).

¹H NMR δ 7.39 (m, 1H), 6.77 (m, 2H), 6.24 (s, 1H), 6.01 (t, 1 H), 3.87 (s, 3 H). MS (ESI⁺) = 247

Step B: Synthesis of 3-(difluoromethyl)-5-(2-fluoro-6-methoxyphenyl)isoxazole

A solution of 4,4-difluoro-1-(2-fluoro-6-methoxyphenyl)butane-1,3-dione (i.e. the product of step A) (2.5 g, 10 mmol) and hydroxylamine hydrochloride (2.1 g, 30 mmol) in ethanol (25 mL) was stirred at 80 °C. After 1 h the solvent was removed under vacuum. The resulting residue was diluted with water and extracted with dichloromethane. The organic phase was dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 15% ethyl acetate in hexanes afforded the title compound (1.5 g).

¹H NMR δ 7.41 (m, 1 H), 6.69 - 6.98 (m, 4H), 3.93 (s, 3 H). MS (ESI⁺) = 244

Step C: Synthesis of 2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenol

To a solution of 3-(difluoromethyl)-5-(2-fluoro-6-methoxyphenyl)isoxazole (i.e. the product of step B) (1.5 g, 6.2 mmol.) in dichloromethane (10 mL) at 0 °C was added a 1.0 M solution of boron tribromide in dichloromethane (31 mL, 31 mmol). The reaction mixture

was warmed to ambient temperature and stirred for 6 h. The reaction was cooled to 0 °C and slowly treated with a saturated aqueous solution of sodium bicarbonate. The biphasic mixture was stirred at room temperature for 1 h. The phases were separated and the aqueous phase was extracted with dichloromethane. The combined organic phases were dried and concentrated under vacuum. The crude residue was purified by chromatography on silica gel, eluting with 0 to 10% ethyl acetate in hexanes, to afford the title compound (980 mg).

^1H NMR δ 7.33 (m, 1 H), 6.66 - 6.99 (m, 4 H). MS (ESI+) = 230


Step D: Synthesis of 5-bromo-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine

To a solution of 2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenol (i.e. the product of step C) (229 mg, 1 mmol) in anhydrous *N,N*-dimethylformamide (2.5 mL) was added 5-bromo-2-chloropyrimidine (212 mg, 1.1 mmol) and potassium carbonate (304 mg, 2.2 mmol). The reaction was heated at 80 °C for 1 h. The solution was cooled to ambient temperature and diluted with water. The phases were separated and the aqueous layer was washed with additional ethyl acetate. The organic phases were combined, dried with magnesium sulfate and concentrated under vacuum. Purification by chromatography on silica gel eluting with 0 to 15% ethyl acetate in hexanes afforded the title compound, a compound of the present invention, as a solid (320 mg).

^1H NMR δ 8.54 (s, 2H), 7.54 (m, 1H), 7.20 (m, 1H), 7.13 (m, 1H), 6.86 (m, 1 H), 6.75 (t, 1 H). MS (ESI⁺) = 387

By the procedures described herein together with methods known in the art, the following compounds of Tables 1 to 1584 can be prepared. The following abbreviations are used in the Tables which follow: *t* means tertiary, *s* means secondary, *n* means normal, *i* means iso, *c* means cyclo, Me means methyl, Et means ethyl, Pr means propyl, Bu means butyl, *i*-Pr means isopropyl, Bu means butyl, *c*-Pr cyclopropyl, *c*-Bu means cyclobutyl, Ph means phenyl, OMe means methoxy, OEt means ethoxy, SMe means methylthio, SEt means ethylthio, NHMe methylamino, -CN means cyano, Py means pyridinyl, -NO₂ means nitro, tzl means triazol, pzl means pyrazol, izl means imidazole, odzl means oxadiazol, tdzl means thiadiazol and SO₂Me means methylsulfonyl.

Table 1



$R^2 = \text{Cl}$; $Z = \text{O}$; and $R^3 = \text{H}$ ($m = 0$);

and Q is:

Isoxazol-5-yl	5-CHO-isoxazol-3-yl	4-I-isothiazol-5-yl
3-F-isoxazol-5-yl	5-CN-isoxazol-3-yl	4-Me-isothiazol-5-yl
3-Cl-isoxazol-5-yl	5-CH ₂ CN-isoxazol-3-yl	4-Et-isothiazol-5-yl
3-Br-isoxazol-5-yl	5-OMe-isoxazol-3-yl	4-CF ₃ -isothiazol-5-yl
3-I-isoxazol-5-yl	5-OCF ₃ -isoxazol-3-yl	4-CHF ₂ -isothiazol-5-yl
3-Me-isoxazol-5-yl	5-Ph-isoxazol-3-yl	4-CHO-isothiazol-5-yl
3-Et-isoxazol-5-yl	4-F-isoxazol-3-yl	4-CN-isothiazol-5-yl
3-CF ₃ -isoxazol-5-yl	4-Cl-isoxazol-3-yl	4-OMe-isothiazol-5-yl
3-CHF ₂ -isoxazol-5-yl	4-Br-isoxazol-3-yl	4-OCF ₃ -isothiazol-5-yl
3-CHO-isoxazol-5-yl	4-I-isoxazol-3-yl	4-Ph-isothiazol-5-yl
3-CN-isoxazol-5-yl	4-Me-isoxazol-3-yl	Isothiazol-3-yl
3-OMe-isoxazol-5-yl	4-Et-isoxazol-3-yl	5-F-isothiazol-3-yl
3-OCF ₃ -isoxazol-5-yl	4-CF ₃ -isoxazol-3-yl	5-Cl-isothiazol-3-yl
3-Ph-isoxazol-5-yl	4-CHF ₂ -isoxazol-3-yl	5-Br-isothiazol-3-yl
4-F-isoxazol-5-yl	4-CHO-isoxazol-3-yl	5-I-isothiazol-3-yl
4-Cl-isoxazol-5-yl	4-CN-isoxazol-3-yl	5-Me-isothiazol-3-yl
4-Br-isoxazol-5-yl	4-OMe-isoxazol-3-yl	5-Et-isothiazol-3-yl
4-I-isoxazol-5-yl	4-OCF ₃ -isoxazol-3-yl	5-CF ₃ -isothiazol-3-yl
4-Me-isoxazol-5-yl	4-Ph-isoxazol-3-yl	5-CHF ₂ -isothiazol-3-yl
4-Et-isoxazol-5-yl	Isothiazol-5-yl	5-CHO-isothiazol-3-yl
4-CF ₃ -isoxazol-5-yl	3-F-isothiazol-5-yl	5-CN-isothiazol-3-yl
4-CHF ₂ -isoxazol-5-yl	3-Cl-isothiazol-5-yl	5-CH ₂ CN-isothiazol-3-yl
4-CHO-isoxazol-5-yl	3-Br-isothiazol-5-yl	5-OMe-isothiazol-3-yl
4-CN-isoxazol-5-yl	3-I-isothiazol-5-yl	5-OCF ₃ -isothiazol-3-yl
4-OMe-isoxazol-5-yl	3-Me-isothiazol-5-yl	5-Ph-isothiazol-3-yl
4-OCF ₃ -isoxazol-5-yl	3-Et-isothiazol-5-yl	4-F-isothiazol-3-yl
4-Ph-isoxazol-5-yl	3-CF ₃ -isothiazol-5-yl	4-Cl-isothiazol-3-yl
isoxazol-3-yl	3-CHF ₂ -isothiazol-5-yl	4-Br-isothiazol-3-yl
5-F-isoxazol-3-yl	3-CHO-isothiazol-5-yl	4-I-isothiazol-3-yl
5-Cl-isoxazol-3-yl	3-CN-isothiazol-5-yl	4-Me-isothiazol-3-yl
5-Br-isoxazol-3-yl	3-OMe-isothiazol-5-yl	4-Et-isothiazol-3-yl
5-I-isoxazol-3-yl	3-OCF ₃ -isothiazol-5-yl	4-CF ₃ -isothiazol-3-yl
5-Me-isoxazol-3-yl	3-Ph-isothiazol-5-yl	4-CHF ₂ -isothiazol-3-yl
5-Et-isoxazol-3-yl	4-F-isothiazol-5-yl	4-CHO-isothiazol-3-yl
5-CF ₃ -isoxazol-3-yl	4-Cl-isothiazol-5-yl	4-CN-isothiazol-3-yl
5-CHF ₂ -isoxazol-3-yl	4-Br-isothiazol-5-yl	4-OMe-isothiazol-3-yl

4-OCF ₃ -isothiazol-3-yl	3-CHO-isothiazol-4-yl	4-Et-oxazol-2-yl
4-Ph-isothiazol-3-yl	3-CN-isothiazol-4-yl	4-CF ₃ -oxazol-2-yl
Isoxazol-4-yl	3-OMe-isothiazol-4-yl	4-CHF ₂ -oxazol-2-yl
3-F-isoxazol-4-yl	3-OCF ₃ -isothiazol-4-yl	4-CHO-oxazol-2-yl
3-Cl-isoxazol-4-yl	3-Ph-isothiazol-4-yl	4-CN-oxazol-2-yl
3-Br-isoxazol-4-yl	5-F-isothiazol-4-yl	4-OMe-oxazol-2-yl
3-I-isoxazol-4-yl	5-Cl-isothiazol-4-yl	4-OCF ₃ -oxazol-2-yl
3-Me-isoxazol-4-yl	5-Br-isothiazol-4-yl	4-Ph-oxazol-2-yl
3-Et-isoxazol-4-yl	5-I-isothiazol-4-yl	Thiazol-2-yl
3-CF ₃ -isoxazol-4-yl	5-Me-isothiazol-4-yl	5-F-thiazol-2-yl
3-CHF ₂ -isoxazol-4-yl	5-Et-isothiazol-4-yl	5-Cl-thiazol-2-yl
3-CHO-isoxazol-4-yl	5-CF ₃ -isothiazol-4-yl	5-Br-thiazol-2-yl
3-CN-isoxazol-4-yl	5-CHF ₂ -isothiazol-4-yl	5-I-thiazol-2-yl
3-OMe-isoxazol-4-yl	5-CHO-isothiazol-4-yl	5-Me-thiazol-2-yl
3-OCF ₃ -isoxazol-4-yl	5-CN-isothiazol-4-yl	5-Et-thiazol-2-yl
3-Ph-isoxazol-4-yl	5-OMe-isothiazol-4-yl	5-CF ₃ -thiazol-2-yl
5-F-isoxazol-4-yl	5-OCF ₃ -isothiazol-4-yl	5-CHF ₂ -thiazol-2-yl
5-Cl-isoxazol-4-yl	5-Ph-isothiazol-4-yl	5-CHO-thiazol-2-yl
5-Br-isoxazol-4-yl	oxazol-2-yl	5-CN-thiazol-2-yl
5-I-isoxazol-4-yl	5-F-oxazol-2-yl	5-CH ₂ CN-thiazol-2-yl
5-Me-isoxazol-4-yl	5-Cl-oxazol-2-yl	5-OMe-thiazol-2-yl
5-Et-isoxazol-4-yl	5-Br-oxazol-2-yl	5-OCF ₃ -thiazol-2-yl
5-CF ₃ -isoxazol-4-yl	5-I-oxazol-2-yl	5-Ph-thiazol-2-yl
5-CHF ₂ -isoxazol-4-yl	5-Me-oxazol-2-yl	4-F-thiazol-2-yl
5-CHO-isoxazol-4-yl	5-Et-oxazol-2-yl	4-Cl-thiazol-2-yl
5-CN-isoxazol-4-yl	5-CF ₃ -oxazol-2-yl	4-Br-thiazol-2-yl
5-OMe-isoxazol-4-yl	5-CHF ₂ -oxazol-2-yl	4-I-thiazol-2-yl
5-OCF ₃ -isoxazol-4-yl	5-CHO-oxazol-2-yl	4-Me-thiazol-2-yl
5-Ph-isoxazol-4-yl	5-CN-oxazol-2-yl	4-Et-thiazol-2-yl
Isothiazol-4-yl	5-CH ₂ CN-oxazol-2-yl	4-CF ₃ -thiazol-2-yl
3-F-isothiazol-4-yl	5-OMe-oxazol-2-yl	4-CHF ₂ -thiazol-2-yl
3-Cl-isothiazol-4-yl	5-OCF ₃ -oxazol-2-yl	4-CHO-thiazol-2-yl
3-Br-isothiazol-4-yl	5-Ph-oxazol-2-yl	4-CN-thiazol-2-yl
3-I-isothiazol-4-yl	4-F-oxazol-2-yl	4-OMe-thiazol-2-yl
3-Me-isothiazol-4-yl	4-Cl-oxazol-2-yl	4-OCF ₃ -thiazol-2-yl
3-Et-isothiazol-4-yl	4-Br-oxazol-2-yl	4-Ph-thiazol-2-yl
3-CF ₃ -isothiazol-4-yl	4-I-oxazol-2-yl	Oxazol-5-yl
3-CHF ₂ -isothiazol-4-yl	4-Me-oxazol-2-yl	2-F-oxazol-5-yl

2-Cl-oxazol-5-yl	4-OCF ₃ -thiazol-5-yl	5-CF ₃ -thiazol-4-yl
2-Br-oxazol-5-yl	4-Ph-thiazol-5-yl	5-CHF ₂ -thiazol-4-yl
2-Me-oxazol-5-yl	Oxazol-4-yl	5-CN-thiazol-4-yl
2-CF ₃ -oxazol-5-yl	2-F-oxazol-4-yl	5-OMe-thiazol-4-yl
2-CHF ₂ -oxazol-5-yl	2-Cl-oxazol-4-yl	5-OCF ₃ -thiazol-4-yl
2-CN-oxazol-5-yl	2-Br-oxazol-4-yl	5-Ph-thiazol-4-yl
2-OMe-oxazol-5-yl	2-Me-oxazol-4-yl	1 <i>H</i> -izl-2-yl
2-OCF ₃ -oxazol-5-yl	2-CF ₃ -oxazol-4-yl	1-Me-1 <i>H</i> -izl-2-yl
2-Ph-oxazol-5-yl	2-CHF ₂ -oxazol-4-yl	4-F-1-Me-1 <i>H</i> -izl-2-yl
4-F-oxazol-5-yl	2-CN-oxazol-4-yl	4-Cl-1-Me-1 <i>H</i> -izl-2-yl
4-Cl-oxazol-5-yl	2-OMe-oxazol-4-yl	4-Br-1-Me-1 <i>H</i> -izl-2-yl
4-Br-oxazol-5-yl	2-OCF ₃ -oxazol-4-yl	1,4-di-Me-1 <i>H</i> -izl-2-yl
4-Me-oxazol-5-yl	2-Ph-oxazol-4-yl	4-CF ₃ -1-Me-1 <i>H</i> -izl-2-yl
4-CF ₃ -oxazol-5-yl	5-F-oxazol-4-yl	4-CHF ₂ -1-Me-1 <i>H</i> -izl-2-yl
4-CHF ₂ -oxazol-5-yl	5-Cl-oxazol-4-yl	4-CN-1-Me-1 <i>H</i> -izl-2-yl
4-CN-oxazol-5-yl	5-Br-oxazol-4-yl	4-OMe-1-Me-1 <i>H</i> -izl-2-yl
4-OMe-oxazol-5-yl	5-Me-oxazol-4-yl	4-OCF ₃ -1-Me-1 <i>H</i> -izl-2-yl
4-OCF ₃ -oxazol-5-yl	5-CF ₃ -oxazol-4-yl	4-Ph-1-Me-1 <i>H</i> -izl-2-yl
4-Ph-oxazol-5-yl	5-CHF ₂ -oxazol-4-yl	5-F-1-Me-1 <i>H</i> -izl-2-yl
Thiazol-5-yl	5-CN-oxazol-4-yl	5-Cl-1-Me-1 <i>H</i> -izl-2-yl
2-F-thiazol-5-yl	5-OMe-oxazol-4-yl	5-Br-1-Me-1 <i>H</i> -izl-2-yl
2-Cl-thiazol-5-yl	5-OCF ₃ -oxazol-4-yl	1,5-di-Me-1 <i>H</i> -izl-2-yl
2-Br-thiazol-5-yl	5-Ph-oxazol-4-yl	5-CF ₃ -1-Me-1 <i>H</i> -izl-2-yl
2-Me-thiazol-5-yl	Thiazol-4-yl	5-CHF ₂ -1-Me-1 <i>H</i> -izl-2-yl
2-CF ₃ -thiazol-5-yl	2-F-thiazol-4-yl	5-CN-1-Me-1 <i>H</i> -izl-2-yl
2-CHF ₂ -thiazol-5-yl	2-Cl-thiazol-4-yl	5-OMe-1-Me-1 <i>H</i> -izl-2-yl
2-CN-thiazol-5-yl	2-Br-thiazol-4-yl	5-OCF ₃ -1-Me-1 <i>H</i> -izl-2-yl
2-OMe-thiazol-5-yl	2-Me-thiazol-4-yl	5-Ph-1-Me-1 <i>H</i> -izl-2-yl
2-OCF ₃ -thiazol-5-yl	2-CF ₃ -thiazol-4-yl	1 <i>H</i> -izl-4-yl
2-Ph-thiazol-5-yl	2-CHF ₂ -thiazol-4-yl	1-Me-1 <i>H</i> -izl-4-yl
4-F-thiazol-5-yl	2-CN-thiazol-4-yl	2-F-1-Me-1 <i>H</i> -izl-4-yl
4-Cl-thiazol-5-yl	2-OMe-thiazol-4-yl	2-Cl-1-Me-1 <i>H</i> -izl-4-yl
4-Br-thiazol-5-yl	2-OCF ₃ -thiazol-4-yl	2-Br-1-Me-1 <i>H</i> -izl-4-yl
4-Me-thiazol-5-yl	2-Ph-thiazol-4-yl	1,2-di-Me-1 <i>H</i> -izl-4-yl
4-CF ₃ -thiazol-5-yl	5-F-thiazol-4-yl	2-CF ₃ -1-Me-1 <i>H</i> -izl-4-yl
4-CHF ₂ -thiazol-5-yl	5-Cl-thiazol-4-yl	2-CHF ₂ -1-Me-1 <i>H</i> -izl-4-yl
4-CN-thiazol-5-yl	5-Br-thiazol-4-yl	2-CN-1-Me-1 <i>H</i> -izl-4-yl
4-OMe-thiazol-5-yl	5-Me-thiazol-4-yl	2-OMe-1-Me-1 <i>H</i> -izl-4-yl

2-OCF ₃ -1-Me-1 <i>H</i> -izl-4-yl	4-Br-1-Me-1 <i>H</i> -pzl-3-yl	5-OCF ₃ -1-Me-1 <i>H</i> -pzl-4-yl
2-Ph-1-Me-1 <i>H</i> -izl-4-yl	1,4-di-Me-1 <i>H</i> -pzl-3-yl	5-Ph-1-Me-1 <i>H</i> -pzl-4-yl
5-F-1-Me-1 <i>H</i> -izl-4-yl	4-CF ₃ -1-Me-1 <i>H</i> -pzl-3-yl	1 <i>H</i> -pzl-5-yl
5-Cl-1-Me-1 <i>H</i> -izl-4-yl	4-CHF ₂ -1-Me-1 <i>H</i> -pzl-3-yl	1-Me-1 <i>H</i> -pzl-5-yl
5-Br-1-Me-1 <i>H</i> -izl-4-yl	4-CN-1-Me-1 <i>H</i> -pzl-3-yl	3-F-1-Me-1 <i>H</i> -pzl-5-yl
1,5-di-Me-1 <i>H</i> -izl-4-yl	4-OMe-1-Me-1 <i>H</i> -pzl-3-yl	3-Cl-1-Me-1 <i>H</i> -pzl-5-yl
5-CF ₃ -1-Me-1 <i>H</i> -izl-4-yl	4-OCF ₃ -1-Me-1 <i>H</i> -pzl-3-yl	3-Br-1-Me-1 <i>H</i> -pzl-5-yl
5-CHF ₂ -1-Me-1 <i>H</i> -izl-4-yl	4-Ph-1-Me-1 <i>H</i> -pzl-3-yl	1,3-di-Me-1 <i>H</i> -pzl-5-yl
5-CN-1-Me-1 <i>H</i> -izl-4-yl	5-F-1-Me-1 <i>H</i> -pzl-3-yl	3-CF ₃ -1-Me-1 <i>H</i> -pzl-5-yl
5-OMe-1-Me-1 <i>H</i> -izl-4-yl	5-Cl-1-Me-1 <i>H</i> -pzl-3-yl	3-CHF ₂ -1-Me-1 <i>H</i> -pzl-5-yl
5-OCF ₃ -1-Me-1 <i>H</i> -izl-4-yl	5-Br-1-Me-1 <i>H</i> -pzl-3-yl	3-CN-1-Me-1 <i>H</i> -pzl-5-yl
5-Ph-1-Me-1 <i>H</i> -izl-4-yl	1,5-di-Me-1 <i>H</i> -pzl-3-yl	3-OMe-1-Me-1 <i>H</i> -pzl-5-yl
1 <i>H</i> -izl-5-yl	5-CF ₃ -1-Me-1 <i>H</i> -pzl-3-yl	3-OCF ₃ -1-Me-1 <i>H</i> -pzl-5-yl
1-Me-1 <i>H</i> -izl-5-yl	5-CHF ₂ -1-Me-1 <i>H</i> -pzl-3-yl	3-Ph-1-Me-1 <i>H</i> -pzl-5-yl
2-F-1-Me-1 <i>H</i> -izl-5-yl	5-CN-1-Me-1 <i>H</i> -pzl-3-yl	4-F-1-Me-1 <i>H</i> -pzl-5-yl
2-Cl-1-Me-1 <i>H</i> -izl-5-yl	5-OMe-1-Me-1 <i>H</i> -pzl-3-yl	4-Cl-1-Me-1 <i>H</i> -pzl-5-yl
2-Br-1-Me-1 <i>H</i> -izl-5-yl	5-OCF ₃ -1-Me-1 <i>H</i> -pzl-3-yl	4-Br-1-Me-1 <i>H</i> -pzl-5-yl
1,2-di-Me-1 <i>H</i> -izl-5-yl	5-Ph-1-Me-1 <i>H</i> -pzl-3-yl	1,4-di-Me-1 <i>H</i> -pzl-5-yl
2-CF ₃ -1-Me-1 <i>H</i> -izl-5-yl	1 <i>H</i> -pzl-4-yl	4-CF ₃ -1-Me-1 <i>H</i> -pzl-5-yl
2-CHF ₂ -1-Me-1 <i>H</i> -izl-5-yl	1-Me-1 <i>H</i> -pzl-4-yl	4-CHF ₂ -1-Me-1 <i>H</i> -pzl-5-yl
2-CN-1-Me-1 <i>H</i> -izl-5-yl	3-F-1-Me-1 <i>H</i> -pzl-4-yl	4-CN-1-Me-1 <i>H</i> -pzl-5-yl
2-OMe-1-Me-1 <i>H</i> -izl-5-yl	3-Cl-1-Me-1 <i>H</i> -pzl-4-yl	4-OMe-1-Me-1 <i>H</i> -pzl-5-yl
2-OCF ₃ -1-Me-1 <i>H</i> -izl-5-yl	3-Br-1-Me-1 <i>H</i> -pzl-4-yl	4-OCF ₃ -1-Me-1 <i>H</i> -pzl-5-yl
2-Ph-1-Me-1 <i>H</i> -izl-5-yl	1,3-di-Me-1 <i>H</i> -pzl-4-yl	4-Ph-1-Me-1 <i>H</i> -pzl-5-yl
4-F-1-Me-1 <i>H</i> -izl-5-yl	3-CF ₃ -1-Me-1 <i>H</i> -pzl-4-yl	Thiophene-2-yl
4-Cl-1-Me-1 <i>H</i> -izl-5-yl	3-CHF ₂ -1-Me-1 <i>H</i> -pzl-4-yl	Thiophene-3-yl
4-Br-1-Me-1 <i>H</i> -izl-5-yl	3-CN-1-Me-1 <i>H</i> -pzl-4-yl	Furan-2-yl
1,4-di-Me-1 <i>H</i> -izl-5-yl	3-OMe-1-Me-1 <i>H</i> -pzl-4-yl	Furan-3-yl
4-CF ₃ -1-Me-1 <i>H</i> -izl-5-yl	3-OCF ₃ -1-Me-1 <i>H</i> -pzl-4-yl	1 <i>H</i> -pyrrol-2-yl
4-CHF ₂ -1-Me-1 <i>H</i> -izl-5-yl	3-Ph-1-Me-1 <i>H</i> -pzl-4-yl	1-Me-1 <i>H</i> -pyrrol-2-yl
4-CN-1-Me-1 <i>H</i> -izl-5-yl	5-F-1-Me-1 <i>H</i> -pzl-4-yl	1 <i>H</i> -pyrrol-3-yl
4-OMe-1-Me-1 <i>H</i> -izl-5-yl	5-Cl-1-Me-1 <i>H</i> -pzl-4-yl	1-Me-1 <i>H</i> -pyrrol-3-yl
4-OCF ₃ -1-Me-1 <i>H</i> -izl-5-yl	5-Br-1-Me-1 <i>H</i> -pzl-4-yl	[1,3,4]odzl-2-yl
4-Ph-1-Me-1 <i>H</i> -izl-5-yl	1,5-di-Me-1 <i>H</i> -pzl-4-yl	2-F-[1,3,4]odzl-5-yl
1 <i>H</i> -pzl-3-yl	5-CF ₃ -1-Me-1 <i>H</i> -pzl-4-yl	2-Cl-[1,3,4]odzl-5-yl
1-Me-1 <i>H</i> -pzl-3-yl	5-CHF ₂ -1-Me-1 <i>H</i> -pzl-4-yl	2-Br-[1,3,4]odzl-5-yl
4-F-1-Me-1 <i>H</i> -pzl-3-yl	5-CN-1-Me-1 <i>H</i> -pzl-4-yl	2-Me-[1,3,4]odzl-5-yl
4-Cl-1-Me-1 <i>H</i> -pzl-3-yl	5-OMe-1-Me-1 <i>H</i> -pzl-4-yl	2-CF ₃ -[1,3,4]odzl-5-yl

2-CHF ₂ -[1,3,4]odzl-5-yl	5-CHF ₂ -1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	5-Br-[1,2,4]odzl-3-yl
2-CN-[1,3,4]odzl-5-yl		5-Me-[1,2,4]odzl-3-yl
2-OMe-[1,3,4]odzl-5-yl	5-CN-1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	5-CF ₃ -[1,2,4]odzl-3-yl
2-OCF ₃ -[1,3,4]odzl-5-yl	5-OMe-1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	5-CHF ₂ -[1,2,4]odzl-3-yl
[1,3,4]tdzl-2-yl		5-CN-[1,2,4]odzl-3-yl
2-F-[1,3,4]tdzl-5-yl	5-OCF ₃ -1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	5-OMe-[1,2,4]odzl-3-yl
2-Cl-[1,3,4]tdzl-5-yl		5-OCF ₃ -[1,2,4]odzl-3-yl
2-Br-[1,3,4]tdzl-5-yl	5-Ph-1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	5-Ph-[1,2,4]odzl-3-yl
2-Me-[1,3,4]tdzl-5-yl	1 <i>H</i> -[1,2,4]tztl-5-yl	[1,2,4]tdzl-5-yl
2-CF ₃ -[1,3,4]tdzl-5-yl	1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	3-F-[1,2,4]tdzl-5-yl
2-CHF ₂ -[1,3,4]tdzl-5-yl	3-F-1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	3-Cl-[1,2,4]tdzl-5-yl
2-CN-[1,3,4]tdzl-5-yl	3-Cl-1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	3-Br-[1,2,4]tdzl-5-yl
2-OMe-[1,3,4]tdzl-5-yl	3-Br-1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	3-Me-[1,2,4]tdzl-5-yl
2-OCF ₃ -[1,3,4]tdzl-5-yl	1,3-di-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	3-CF ₃ -[1,2,4]tdzl-5-yl
4 <i>H</i> -[1,2,4]tztl-3-yl	3-CF ₃ -1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	3-CHF ₂ -[1,2,4]tdzl-5-yl
4-Me-4 <i>H</i> -[1,2,4]tztl-3-yl		3-CN-[1,2,4]tdzl-5-yl
3-F-4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl	3-CHF ₂ -1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	3-OMe-[1,2,4]tdzl-5-yl
3-Cl-4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl		3-OCF ₃ -[1,2,4]tdzl-5-yl
3-Br-4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl	3-CN-1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	3-Ph-[1,2,4]tdzl-5-yl
3,4-di-Me-4 <i>H</i> -[1,2,4]tztl-5-yl	3-OMe-1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	[1,2,4]tdzl-3-yl
3-CF ₃ -4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl		5-F-[1,2,4]tdzl-3-yl
	3-OCF ₃ -1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	5-Cl-[1,2,4]tdzl-3-yl
		5-Br-[1,2,4]tdzl-3-yl
3-CHF ₂ -4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl	3-Ph-1-Me-1 <i>H</i> -[1,2,4]tztl-5-yl	5-Me-[1,2,4]tdzl-3-yl
	[1,2,4]odzl-5-yl	5-CF ₃ -[1,2,4]tdzl-3-yl
3-CN-4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl	3-F-[1,2,4]odzl-5-yl	5-CHF ₂ -[1,2,4]tdzl-3-yl
3-OMe-4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl	3-Cl-[1,2,4]odzl-5-yl	5-CN-[1,2,4]tdzl-3-yl
	3-Br-[1,2,4]odzl-5-yl	5-OMe-[1,2,4]tdzl-3-yl
3-OCF ₃ -4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl	3-Me-[1,2,4]odzl-5-yl	5-OCF ₃ -[1,2,4]tdzl-3-yl
	3-CF ₃ -[1,2,4]odzl-5-yl	5-Ph-[1,2,4]tdzl-3-yl
3-Ph-4-Me-4 <i>H</i> -[1,2,4]tztl-5-yl	3-CHF ₂ -[1,2,4]odzl-5-yl	[1,2,3]odzl-5-yl
1 <i>H</i> -[1,2,4]tztl-3-yl	3-CN-[1,2,4]odzl-5-yl	4-F-[1,2,3]odzl-5-yl
1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	3-OMe-[1,2,4]odzl-5-yl	4-Cl-[1,2,3]odzl-5-yl
5-F-1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	3-OCF ₃ -[1,2,4]odzl-5-yl	4-Br-[1,2,3]odzl-5-yl
5-Cl-1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	3-Ph-[1,2,4]odzl-5-yl	4-Me-[1,2,3]odzl-5-yl
5-Br-1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	[1,2,4]odzl-3-yl	4-CF ₃ -[1,2,3]odzl-5-yl
1,5-di-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	5-F-[1,2,4]odzl-3-yl	4-CHF ₂ -[1,2,3]odzl-5-yl
5-CF ₃ -1-Me-1 <i>H</i> -[1,2,4]tztl-3-yl	5-Cl-[1,2,4]odzl-3-yl	4-CN-[1,2,3]odzl-5-yl

4-OMe-[1,2,3]odzl-5-yl	5-Cl-3 <i>H</i> -[1,2,4]tztl-3-yl	4-CN-1 <i>H</i> -[1,2,3]tztl-5-yl
4-OCF ₃ -[1,2,3]odzl-5-yl	5-Br-3 <i>H</i> -[1,2,4]tztl-3-yl	4-OMe-1 <i>H</i> -[1,2,3]tztl-5-yl
4-Ph-[1,2,3]odzl-5-yl	5-Me-3 <i>H</i> -[1,2,4]tztl-3-yl	4-OCF ₃ -1 <i>H</i> -[1,2,3]tztl-5-yl
[1,2,3]odzl-4-yl	5-CF ₃ -3 <i>H</i> -[1,2,4]tztl-3-yl	4-Ph-1 <i>H</i> -[1,2,3]tztl-5-yl
5-F-[1,2,3]odzl-4-yl	5-CHF ₂ -3 <i>H</i> -[1,2,4]tztl-3-yl	5-F-pyridin-2-yl
5-Cl-[1,2,3]odzl-4-yl	5-CN-3 <i>H</i> -[1,2,4]tztl-3-yl	5-Cl-pyridin-2-yl
5-Br-[1,2,3]odzl-4-yl	5-OMe-3 <i>H</i> -[1,2,4]tztl-3-yl	5-Br-pyridin-2-yl
5-Me-[1,2,3]odzl-4-yl	5-OCF ₃ -3 <i>H</i> -[1,2,4]tztl-3-yl	5-I-pyridin-2-yl
5-CF ₃ -[1,2,3]odzl-4-yl	5-Ph-3 <i>H</i> -[1,2,4]tztl-3-yl	5-Me-pyridin-2-yl
5-CHF ₂ -[1,2,3]odzl-4-yl	1 <i>H</i> -[1,2,3]tztl-4-yl	5-Et-pyridin-2-yl
5-CN-[1,2,3]odzl-4-yl	5-F-1 <i>H</i> -[1,2,3]tztl-4-yl	5-CF ₃ -pyridin-2-yl
5-OMe-[1,2,3]odzl-4-yl	5-Cl-1 <i>H</i> -[1,2,3]tztl-4-yl	5-CHF ₂ -pyridin-2-yl
5-OCF ₃ -[1,2,3]odzl-4-yl	5-Br-1 <i>H</i> -[1,2,3]tztl-4-yl	5-CHO-pyridin-2-yl
5-Ph-[1,2,3]odzl-4-yl	5-Me-1 <i>H</i> -[1,2,3]tztl-4-yl	5-CN-pyridin-2-yl
[1,2,3]tdzl-5-yl	5-CF ₃ -1 <i>H</i> -[1,2,3]tztl-4-yl	5-OMe-pyridin-2-yl
4-F-[1,2,3]tdzl-5-yl	5-CHF ₂ -1 <i>H</i> -[1,2,3]tztl-4-yl	5-OCF ₃ -pyridin-2-yl
4-Cl-[1,2,3]tdzl-5-yl	5-CN-1 <i>H</i> -[1,2,3]tztl-4-yl	5-N(Me) ₂ -pyridin-2-yl
4-Br-[1,2,3]tdzl-5-yl	5-OMe-1 <i>H</i> -[1,2,3]tztl-4-yl	5-Ph-pyridin-2-yl
4-Me-[1,2,3]tdzl-5-yl	5-OCF ₃ -1 <i>H</i> -[1,2,3]tztl-4-yl	3,5-di-Cl-pyridin-2-yl
4-CF ₃ -[1,2,3]tdzl-5-yl	5-Ph-1 <i>H</i> -[1,2,3]tztl-4-yl	3-Me-5-Cl-pyridin-2-yl
4-CHF ₂ -[1,2,3]tdzl-5-yl	2 <i>H</i> -[1,2,3]tztl-4-yl	3-CN-5-Cl-pyridin-2-yl
4-CN-[1,2,3]tdzl-5-yl	4-F-2 <i>H</i> -[1,2,3]tztl-5-yl	6-F-pyridin-2-yl
4-OMe-[1,2,3]tdzl-5-yl	4-Cl-2 <i>H</i> -[1,2,3]tztl-5-yl	6-Cl-pyridin-2-yl
4-OCF ₃ -[1,2,3]tdzl-5-yl	4-Br-2 <i>H</i> -[1,2,3]tztl-5-yl	6-Br-pyridin-2-yl
4-Ph-[1,2,3]tdzl-5-yl	4-Me-2 <i>H</i> -[1,2,3]tztl-5-yl	6-I-pyridin-2-yl
[1,2,3]tdzl-4-yl	4-CF ₃ -2 <i>H</i> -[1,2,3]tztl-5-yl	6-Me-pyridin-2-yl
5-F-[1,2,3]tdzl-4-yl	4-CHF ₂ -2 <i>H</i> -[1,2,3]tztl-5-yl	6-Et-pyridin-2-yl
5-Cl-[1,2,3]tdzl-4-yl	4-CN-2 <i>H</i> -[1,2,3]tztl-5-yl	6-CF ₃ -pyridin-2-yl
5-Br-[1,2,3]tdzl-4-yl	4-OMe-2 <i>H</i> -[1,2,3]tztl-5-yl	6-CHF ₂ -pyridin-2-yl
5-Me-[1,2,3]tdzl-4-yl	4-OCF ₃ -2 <i>H</i> -[1,2,3]tztl-5-yl	6-CHO-pyridin-2-yl
5-CF ₃ -[1,2,3]tdzl-4-yl	4-Ph-2 <i>H</i> -[1,2,3]tztl-5-yl	6-CN-pyridin-2-yl
5-CHF ₂ -[1,2,3]tdzl-4-yl	1 <i>H</i> -[1,2,3]tztl-5-yl	6-OMe-pyridin-2-yl
5-CN-[1,2,3]tdzl-4-yl	4-F-1 <i>H</i> -[1,2,3]tztl-5-yl	6-OCF ₃ -pyridin-2-yl
5-OMe-[1,2,3]tdzl-4-yl	4-Cl-1 <i>H</i> -[1,2,3]tztl-5-yl	6-N(Me) ₂ -pyridin-2-yl
5-OCF ₃ -[1,2,3]tdzl-4-yl	4-Br-1 <i>H</i> -[1,2,3]tztl-5-yl	6-Ph-pyridin-2-yl
5-Ph-[1,2,3]tdzl-4-yl	4-Me-1 <i>H</i> -[1,2,3]tztl-5-yl	3-F-pyridin-2-yl
3 <i>H</i> -[1,2,4]tztl-3-yl	4-CF ₃ -1 <i>H</i> -[1,2,3]tztl-5-yl	3-Cl-pyridin-2-yl
5-F-3 <i>H</i> -[1,2,4]tztl-3-yl	4-CHF ₂ -1 <i>H</i> -[1,2,3]tztl-5-yl	3-Br-pyridin-2-yl

3-I-pyridin-2-yl	3-OMe-pyridin-4-yl	5-F-pyridazin-2-yl
3-Me-pyridin-2-yl	3-OCF ₃ -pyridin-4-yl	5-Cl-pyridazin-2-yl
3-Et-pyridin-2-yl	3-N(Me) ₂ -pyridin-4-yl	5-Br-pyridazin-2-yl
3-CF ₃ -pyridin-2-yl	3-Ph-pyridin-4-yl	5-I-pyridazin-2-yl
3-CHF ₂ -pyridin-2-yl	3,5-di-Me-pyridin-4-yl	5-Me-pyridazin-2-yl
3-CHO-pyridin-2-yl	3,5-di-Cl-pyridin-4-yl	5-Et-pyridazin-2-yl
3-CN-pyridin-2-yl	6-F-pyridazin-3-yl	5-CF ₃ -pyridazin-2-yl
3-OMe-pyridin-2-yl	6-Cl-pyridazin-3-yl	5-CHF ₂ -pyridazin-2-yl
3-OCF ₃ -pyridin-2-yl	6-Br-pyridazin-3-yl	5-CHO-pyridazin-2-yl
3-N(Me) ₂ -pyridin-2-yl	6-I-pyridazin-3-yl	5-CN-pyridazin-2-yl
3-Ph-pyridin-2-yl	6-Me-pyridazin-3-yl	5-OMe-pyridazin-2-yl
5,6-di-Cl-pyridin-2-yl	6-Et-pyridazin-3-yl	5-OCF ₃ -pyridazin-2-yl
6-F-pyridin-3-yl	6-CF ₃ -pyridazin-3-yl	5-N(Me) ₂ -pyridazin-2-yl
6-Cl-pyridin-3-yl	6-CHF ₂ -pyridazin-3-yl	5-Ph-pyridazin-2-yl
6-Br-pyridin-3-yl	6-CHO-pyridazin-3-yl	5-F-pyrimidin-4-yl
6-I-pyridin-3-yl	6-CN-pyridazin-3-yl	5-Cl-pyrimidin-4-yl
6-Me-pyridin-3-yl	6-OMe-pyridazin-3-yl	5-Br-pyrimidin-4-yl
6-Et-pyridin-3-yl	6-OCF ₃ -pyridazin-3-yl	5-I-pyrimidin-4-yl
6-CF ₃ -pyridin-3-yl	6-N(Me) ₂ -pyridazin-3-yl	5-Me-pyrimidin-4-yl
6-CHF ₂ -pyridin-3-yl	6-Ph-pyridazin-3-yl	5-Et-pyrimidin-4-yl
6-CHO-pyridin-3-yl	4-Cl-pyridazin-3-yl	5-CF ₃ -pyrimidin-4-yl
6-CN-pyridin-3-yl	4-CN-pyridazin-3-yl	5-CHF ₂ -pyrimidin-4-yl
6-OMe-pyridin-3-yl	6-F-pyridazin-4-yl	5-CHO-pyrimidin-4-yl
6-OCF ₃ -pyridin-3-yl	6-Cl-pyridazin-4-yl	5-CN-pyrimidin-4-yl
6-N(Me) ₂ -pyridin-3-yl	6-Br-pyridazin-4-yl	5-OMe-pyrimidin-4-yl
6-Ph-pyridin-3-yl	6-I-pyridazin-4-yl	5-OCF ₃ -pyrimidin-4-yl
4,6-di-Cl-pyridin-3-yl	6-Me-pyridazin-4-yl	5-N(Me) ₂ -pyrimidin-4-yl
4-CN-6-Cl-pyridin-3-yl	6-Et-pyridazin-4-yl	5-Ph-pyrimidin-4-yl
3-F-pyridin-4-yl	6-CF ₃ -pyridazin-4-yl	2-F-pyrimidin-5-yl
3-Cl-pyridin-4-yl	6-CHF ₂ -pyridazin-4-yl	2-Cl-pyrimidin-5-yl
3-Br-pyridin-4-yl	6-CHO-pyridazin-4-yl	2-Br-pyrimidin-5-yl
3-I-pyridin-4-yl	6-CN-pyridazin-4-yl	2-I-pyrimidin-5-yl
3-Me-pyridin-4-yl	6-OMe-pyridazin-4-yl	2-Me-pyrimidin-5-yl
3-Et-pyridin-4-yl	6-OCF ₃ -pyridazin-4-yl	2-Et-pyrimidin-5-yl
3-CF ₃ -pyridin-4-yl	6-N(Me) ₂ -pyridazin-4-yl	2-CF ₃ -pyrimidin-5-yl
3-CHF ₂ -pyridin-4-yl	6-Ph-pyridazin-4-yl	2-CHF ₂ -pyrimidin-5-yl
3-CHO-pyridin-4-yl	4-Cl-pyridazin-4-yl	2-CHO-pyrimidin-5-yl
3-CN-pyridin-4-yl	4-CN-pyridazin-4-yl	2-CN-pyrimidin-5-yl

2-OMe-pyrimidin-5-yl	3-OMe-[1,2,4]triazin-6-yl	6-CN-[1,2,4]triazin-3-yl
2-OCF ₃ -pyrimidin-5-yl	3-CF ₃ -[1,2,4]triazin-6-yl	4-Cl-[1,3,5]triazin-2-yl
2-N(Me) ₂ -pyrimidin-5-yl	6-Cl-[1,2,4]triazin-5-yl	4-CF ₃ -phenyl
2-Ph-pyrimidin-5-yl	6-Me-[1,2,4]triazin-5-yl	4-OCF ₃ -phenyl
3-Cl-pyrazin-2-yl	6-OMe-[1,2,4]triazin-5-yl	3-OCF ₃ -phenyl
3-CN-pyrazin-2-yl	6-CN-[1,2,4]triazin-5-yl	3,5-di-OCF ₃ -phenyl
3-OMe-pyrazin-2-yl	6-Cl-[1,2,4]triazin-3-yl	3,5-di-Cl-phenyl
3-Cl-[1,2,4]triazin-6-yl	6-Me-[1,2,4]triazin-3-yl	
3-CN-[1,2,4]triazin-6-yl	6-OMe-[1,2,4]triazin-3-yl	

The present disclosure also includes Tables 2 through 1584. Each Table is constructed in the the same manner as Table 1 above, except that the row heading in Table 1 (i.e. “R² = Cl; Z = O; and R³ = H (m = 0)”) is replaced with the respective row heading shown below. For example, the first entry in Table 2 is a compound of Formula 1 wherein R¹ is H, R² is Cl, Z is O, R³ is H (m = 0), and Q is isoxazol-5-yl (i.e. unsubstituted isoxazole attached to the remainder of Formula 1 at the 5-position). The remainder of Table 2 is constructed in the same way, and hence the remainder of Tables 3 through 1584 is constructed the same way.

Table	Header Row	Table	Header Row
2	R ² = F, Z = O, R ³ = H (m = 0)	23	R ² = F, Z = O, R ³ = 3-Ph
3	R ² = F, Z = O, R ³ = 3-F	24	R ² = F, Z = S, R ³ = 3-F
4	R ² = F, Z = O, R ³ = 3-Cl	25	R ² = F, Z = S, R ³ = 3-Cl
5	R ² = F, Z = O, R ³ = 3-Br	26	R ² = F, Z = S, R ³ = 3-Br
6	R ² = F, Z = O, R ³ = 3-I	27	R ² = F, Z = S, R ³ = 3-I
7	R ² = F, Z = O, R ³ = 3-CN	28	R ² = F, Z = S, R ³ = 3-CN
8	R ² = F, Z = O, R ³ = 3-NO ₂	29	R ² = F, Z = S, R ³ = 3-NO ₂
9	R ² = F, Z = O, R ³ = 3-OMe	30	R ² = F, Z = S, R ³ = 3-OMe
10	R ² = F, Z = O, R ³ = 3-OCF ₃	31	R ² = F, Z = S, R ³ = 3-OCF ₃
11	R ² = F, Z = O, R ³ = 3-CF ₃	32	R ² = F, Z = S, R ³ = 3-CF ₃
12	R ² = F, Z = O, R ³ = 3-CHF ₂	33	R ² = F, Z = S, R ³ = 3-CHF ₂
13	R ² = F, Z = O, R ³ = 3-CH ₂ F	34	R ² = F, Z = S, R ³ = 3-CH ₂ F
14	R ² = F, Z = O, R ³ = 3-CHO	35	R ² = F, Z = S, R ³ = 3-CHO
15	R ² = F, Z = O, R ³ = 3-Me	36	R ² = F, Z = S, R ³ = 3-Me
16	R ² = F, Z = O, R ³ = 3-Et	37	R ² = F, Z = S, R ³ = 3-Et
17	R ² = F, Z = O, R ³ = 3-Ethynyl	38	R ² = F, Z = S, R ³ = 3-Ethynyl
18	R ² = F, Z = O, R ³ = 3-Ethenyl	39	R ² = F, Z = S, R ³ = 3-Ethenyl
19	R ² = F, Z = O, R ³ = 3-SO ₂ Me	40	R ² = F, Z = S, R ³ = 3-SO ₂ Me
20	R ² = F, Z = O, R ³ = 3-OAc	41	R ² = F, Z = S, R ³ = 3-OAc
21	R ² = F, Z = O, R ³ = 3- <i>c</i> -Pr	42	R ² = F, Z = S, R ³ = 3- <i>c</i> -Pr
22	R ² = F, Z = O, R ³ = 3- <i>i</i> -Pr	43	R ² = F, Z = S, R ³ = 3- <i>i</i> -Pr

Table	Header Row
44	$R^2 = F, Z = S, R^3 = 3\text{-Ph}$
45	$R^2 = F, Z = O, R^3 = 4\text{-F}$
46	$R^2 = F, Z = O, R^3 = 4\text{-Cl}$
47	$R^2 = F, Z = O, R^3 = 4\text{-Br}$
48	$R^2 = F, Z = O, R^3 = 4\text{-I}$
49	$R^2 = F, Z = O, R^3 = 4\text{-CN}$
50	$R^2 = F, Z = O, R^3 = 4\text{-NO}_2$
51	$R^2 = F, Z = O, R^3 = 4\text{-OMe}$
52	$R^2 = F, Z = O, R^3 = 4\text{-OCF}_3$
53	$R^2 = F, Z = O, R^3 = 4\text{-CF}_3$
54	$R^2 = F, Z = O, R^3 = 4\text{-CHF}_2$
55	$R^2 = F, Z = O, R^3 = 4\text{-CH}_2\text{F}$
56	$R^2 = F, Z = O, R^3 = 4\text{-CHO}$
57	$R^2 = F, Z = O, R^3 = 4\text{-Me}$
58	$R^2 = F, Z = O, R^3 = 4\text{-Et}$
59	$R^2 = F, Z = O, R^3 = 4\text{-Ethyneyl}$
60	$R^2 = F, Z = O, R^3 = 4\text{-Ethenyl}$
61	$R^2 = F, Z = O, R^3 = 4\text{-SO}_2\text{Me}$
62	$R^2 = F, Z = O, R^3 = 4\text{-OAc}$
63	$R^2 = F, Z = O, R^3 = 4\text{-c-Pr}$
64	$R^2 = F, Z = O, R^3 = 4\text{-i-Pr}$
65	$R^2 = F, Z = O, R^3 = 4\text{-Ph}$
66	$R^2 = F, Z = O, R^3 = 5\text{-F}$
67	$R^2 = F, Z = O, R^3 = 5\text{-Cl}$
68	$R^2 = F, Z = O, R^3 = 5\text{-Br}$
69	$R^2 = F, Z = O, R^3 = 5\text{-I}$
70	$R^2 = F, Z = O, R^3 = 5\text{-CN}$
71	$R^2 = F, Z = O, R^3 = 5\text{-NO}_2$
72	$R^2 = F, Z = O, R^3 = 5\text{-OMe}$
73	$R^2 = F, Z = O, R^3 = 5\text{-OCF}_3$
74	$R^2 = F, Z = O, R^3 = 5\text{-CF}_3$
75	$R^2 = F, Z = O, R^3 = 5\text{-CHF}_2$
76	$R^2 = F, Z = O, R^3 = 5\text{-CH}_2\text{F}$
77	$R^2 = F, Z = O, R^3 = 5\text{-CHO}$
78	$R^2 = F, Z = O, R^3 = 5\text{-Me}$
79	$R^2 = F, Z = O, R^3 = 5\text{-Et}$
80	$R^2 = F, Z = O, R^3 = 5\text{-Ethyneyl}$

Table	Header Row
81	$R^2 = F, Z = O, R^3 = 5\text{-Ethenyl}$
82	$R^2 = F, Z = O, R^3 = 5\text{-SO}_2\text{Me}$
83	$R^2 = F, Z = O, R^3 = 5\text{-OAc}$
84	$R^2 = F, Z = O, R^3 = 5\text{-c-Pr}$
85	$R^2 = F, Z = O, R^3 = 5\text{-i-Pr}$
86	$R^2 = F, Z = O, R^3 = 5\text{-Ph}$
87	$R^2 = F, Z = O, R^3 = 6\text{-F}$
88	$R^2 = F, Z = O, R^3 = 6\text{-Cl}$
89	$R^2 = F, Z = O, R^3 = 6\text{-Br}$
90	$R^2 = F, Z = O, R^3 = 6\text{-I}$
91	$R^2 = F, Z = O, R^3 = 6\text{-CN}$
92	$R^2 = F, Z = O, R^3 = 6\text{-NO}_2$
93	$R^2 = F, Z = O, R^3 = 6\text{-OMe}$
94	$R^2 = F, Z = O, R^3 = 6\text{-OCF}_3$
95	$R^2 = F, Z = O, R^3 = 6\text{-CF}_3$
96	$R^2 = F, Z = O, R^3 = 6\text{-CHF}_2$
97	$R^2 = F, Z = O, R^3 = 6\text{-CH}_2\text{F}$
98	$R^2 = F, Z = O, R^3 = 6\text{-CHO}$
99	$R^2 = F, Z = O, R^3 = 6\text{-Me}$
100	$R^2 = F, Z = O, R^3 = 6\text{-Et}$
101	$R^2 = F, Z = O, R^3 = 6\text{-Ethyneyl}$
102	$R^2 = F, Z = O, R^3 = 6\text{-Ethenyl}$
103	$R^2 = F, Z = O, R^3 = 6\text{-SO}_2\text{Me}$
104	$R^2 = F, Z = O, R^3 = 6\text{-OAc}$
105	$R^2 = F, Z = O, R^3 = 6\text{-c-Pr}$
106	$R^2 = F, Z = O, R^3 = 6\text{-i-Pr}$
107	$R^2 = F, Z = O, R^3 = 6\text{-Ph}$
108	$R^2 = F, Z = O, R^3 = 3,4\text{-di-F}$
109	$R^2 = F, Z = O, R^3 = 3,5\text{-di-F}$
110	$R^2 = F, Z = O, R^3 = 3,6\text{-di-F}$
111	$R^2 = F, Z = O, R^3 = 4,5\text{-di-F}$
112	$R^2 = F, Z = O, R^3 = 3,4\text{-di-Cl}$
113	$R^2 = F, Z = O, R^3 = 3,5\text{-di-Cl}$
114	$R^2 = F, Z = O, R^3 = 3,6\text{-di-Cl}$
115	$R^2 = F, Z = O, R^3 = 4,5\text{-di-Cl}$
116	$R^2 = F, Z = O, R^3 = 3,4\text{-di-Br}$
117	$R^2 = F, Z = O, R^3 = 3,5\text{-di-Br}$

Table	Header Row
118	$R^2 = F, Z = O, R^3 = 3,6\text{-di-Br}$
119	$R^2 = F, Z = O, R^3 = 4,5\text{-di-Br}$
120	$R^2 = F, Z = O, R^3 = 3,4\text{-di-CN}$
121	$R^2 = F, Z = O, R^3 = 3,5\text{-di-CN}$
122	$R^2 = F, Z = O, R^3 = 3,6\text{-di-CN}$
123	$R^2 = F, Z = O, R^3 = 4,5\text{-di-CN}$
124	$R^2 = F, Z = O, R^3 = 3,4\text{-di-Me}$
125	$R^2 = F, Z = O, R^3 = 3,5\text{-di-Me}$
126	$R^2 = F, Z = O, R^3 = 3,6\text{-di-Me}$
127	$R^2 = F, Z = O, R^3 = 4,5\text{-di-Me}$
128	$R^2 = F, Z = O, R^3 = 3,4\text{-di-OMe}$
129	$R^2 = F, Z = O, R^3 = 3,5\text{-di-OMe}$
130	$R^2 = F, Z = O, R^3 = 3,6\text{-di-OMe}$
131	$R^2 = F, Z = O, R^3 = 4,5\text{-di-OMe}$
132	$R^2 = F, Z = O, R^3 = 3,4\text{-di-CF}_3$
133	$R^2 = F, Z = O, R^3 = 3,5\text{-di-CF}_3$
134	$R^2 = F, Z = O, R^3 = 3,6\text{-di-CF}_3$
135	$R^2 = F, Z = O, R^3 = 4,5\text{-di-CF}_3$
136	$R^2 = F, Z = O, R^3 = 3\text{-CN, 4-Me}$
137	$R^2 = F, Z = O, R^3 = 3\text{-CN, 4-F}$
138	$R^2 = F, Z = O, R^3 = 3\text{-CN, 4-Br}$
139	$R^2 = F, Z = O, R^3 = 3\text{-CN, 4-OMe}$
140	$R^2 = F, Z = O, R^3 = 3\text{-CN, 4-CF}_3$
141	$R^2 = F, Z = O, R^3 = 3\text{-CN, 6-Me}$
142	$R^2 = F, Z = O, R^3 = 3\text{-CN, 6-F}$
143	$R^2 = F, Z = O, R^3 = 3\text{-CN, 6-Br}$
144	$R^2 = F, Z = O, R^3 = 3\text{-CN, 6-OMe}$
145	$R^2 = F, Z = O, R^3 = 3\text{-CN, 6-CF}_3$
146	$R^2 = Br, Z = O, R^3 = H (m = 0)$
147	$R^2 = Br, Z = O, R^3 = 3-F$
148	$R^2 = Br, Z = O, R^3 = 3-Cl$
149	$R^2 = Br, Z = O, R^3 = 3-Br$
150	$R^2 = Br, Z = O, R^3 = 3-I$
151	$R^2 = Br, Z = O, R^3 = 3-CN$
152	$R^2 = Br, Z = O, R^3 = 3-NO_2$
153	$R^2 = Br, Z = O, R^3 = 3-OMe$
154	$R^2 = Br, Z = O, R^3 = 3-OCF_3$

Table	Header Row
155	$R^2 = Br, Z = O, R^3 = 3-CF_3$
156	$R^2 = Br, Z = O, R^3 = 3-CHF_2$
157	$R^2 = Br, Z = O, R^3 = 3-CH_2F$
158	$R^2 = Br, Z = O, R^3 = 3-CHO$
159	$R^2 = Br, Z = O, R^3 = 3-Me$
160	$R^2 = Br, Z = O, R^3 = 3-Et$
161	$R^2 = Br, Z = O, R^3 = 3\text{-Ethynyl}$
162	$R^2 = Br, Z = O, R^3 = 3\text{-Ethenyl}$
163	$R^2 = Br, Z = O, R^3 = 3-SO_2Me$
164	$R^2 = Br, Z = O, R^3 = 3-OAc$
165	$R^2 = Br, Z = O, R^3 = 3\text{-}i\text{-Pr}$
166	$R^2 = Br, Z = O, R^3 = 3\text{-}i\text{-Pr}$
167	$R^2 = Br, Z = O, R^3 = 3-Ph$
168	$R^2 = Br, Z = S, R^3 = 3-F$
169	$R^2 = Br, Z = S, R^3 = 3-Cl$
170	$R^2 = Br, Z = S, R^3 = 3-Br$
171	$R^2 = Br, Z = S, R^3 = 3-I$
172	$R^2 = Br, Z = S, R^3 = 3-CN$
173	$R^2 = Br, Z = S, R^3 = 3-NO_2$
174	$R^2 = Br, Z = S, R^3 = 3-OMe$
175	$R^2 = Br, Z = S, R^3 = 3-OCF_3$
176	$R^2 = Br, Z = S, R^3 = 3-CF_3$
177	$R^2 = Br, Z = S, R^3 = 3-CHF_2$
178	$R^2 = Br, Z = S, R^3 = 3-CH_2F$
179	$R^2 = Br, Z = S, R^3 = 3-CHO$
180	$R^2 = Br, Z = S, R^3 = 3-Me$
181	$R^2 = Br, Z = S, R^3 = 3-Et$
182	$R^2 = Br, Z = S, R^3 = 3\text{-Ethynyl}$
183	$R^2 = Br, Z = S, R^3 = 3\text{-Ethenyl}$
184	$R^2 = Br, Z = S, R^3 = 3-SO_2Me$
185	$R^2 = Br, Z = S, R^3 = 3-OAc$
186	$R^2 = Br, Z = S, R^3 = 3\text{-}i\text{-Pr}$
187	$R^2 = Br, Z = S, R^3 = 3\text{-}i\text{-Pr}$
188	$R^2 = Br, Z = S, R^3 = 3-Ph$
189	$R^2 = Br, Z = O, R^3 = 4-F$
190	$R^2 = Br, Z = O, R^3 = 4-Cl$
191	$R^2 = Br, Z = O, R^3 = 4-Br$

Table	Header Row
192	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-I}$
193	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-CN}$
194	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-NO}_2$
195	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-OMe}$
196	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-OCF}_3$
197	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-CF}_3$
198	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-CHF}_2$
199	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-CH}_2\text{F}$
200	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-CHO}$
201	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-Me}$
202	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-Et}$
203	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-Ethynyl}$
204	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-Ethenyl}$
205	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-SO}_2\text{Me}$
206	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-OAc}$
207	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-}i\text{-Pr}$
208	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-}i\text{-Pr}$
209	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4\text{-Ph}$
210	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-F}$
211	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-Cl}$
212	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-Br}$
213	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-I}$
214	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-CN}$
215	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-NO}_2$
216	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-OMe}$
217	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-OCF}_3$
218	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-CF}_3$
219	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-CHF}_2$
220	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-CH}_2\text{F}$
221	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-CHO}$
222	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-Me}$
223	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-Et}$
224	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-Ethynyl}$
225	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-Ethenyl}$
226	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-SO}_2\text{Me}$
227	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-OAc}$
228	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$

Table	Header Row
229	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$
230	$R^2 = \text{Br}, Z = \text{O}, R^3 = 5\text{-Ph}$
231	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-F}$
232	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-Cl}$
233	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-Br}$
234	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-I}$
235	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-CN}$
236	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-NO}_2$
237	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-OMe}$
238	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-OCF}_3$
239	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-CF}_3$
240	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-CHF}_2$
241	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-CH}_2\text{F}$
242	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-CHO}$
243	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-Me}$
244	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-Et}$
245	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-Ethynyl}$
246	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-Ethenyl}$
247	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-SO}_2\text{Me}$
248	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-OAc}$
249	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
250	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
251	$R^2 = \text{Br}, Z = \text{O}, R^3 = 6\text{-Ph}$
252	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,4\text{-di-F}$
253	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,5\text{-di-F}$
254	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,6\text{-di-F}$
255	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4,5\text{-di-F}$
256	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,4\text{-di-Cl}$
257	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,5\text{-di-Cl}$
258	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,6\text{-di-Cl}$
259	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4,5\text{-di-Cl}$
260	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,4\text{-di-Br}$
261	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,5\text{-di-Br}$
262	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,6\text{-di-Br}$
263	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4,5\text{-di-Br}$
264	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,4\text{-di-CN}$
265	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,5\text{-di-CN}$

Table	Header Row
266	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,6\text{-di-CN}$
267	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4,5\text{-di-CN}$
268	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,4\text{-di-Me}$
269	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,5\text{-di-Me}$
270	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,6\text{-di-Me}$
271	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4,5\text{-di-Me}$
272	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,4\text{-di-OMe}$
273	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,5\text{-di-OMe}$
274	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,6\text{-di-OMe}$
275	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4,5\text{-di-OMe}$
276	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,4\text{-di-CF}_3$
277	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,5\text{-di-CF}_3$
278	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3,6\text{-di-CF}_3$
279	$R^2 = \text{Br}, Z = \text{O}, R^3 = 4,5\text{-di-CF}_3$
280	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Me}$
281	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-F}$
282	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Br}$
283	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-OMe}$
284	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-CF}_3$
285	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Me}$
286	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-F}$
287	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Br}$
288	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-OMe}$
289	$R^2 = \text{Br}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-CF}_3$
290	$R^2 = \text{Cl}, Z = \text{O}, R^3 = \text{H} (m = 0)$
291	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-F}$
292	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-Cl}$
293	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-Br}$
294	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-I}$
295	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-CN}$
296	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-NO}_2$
297	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-OMe}$
298	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-OCF}_3$
299	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-CF}_3$
300	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-CHF}_2$
301	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-CH}_2\text{F}$
302	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-CHO}$

Table	Header Row
303	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-Me}$
304	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-Et}$
305	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-Ethylyl}$
306	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-Ethenyl}$
307	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-SO}_2\text{Me}$
308	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-OAc}$
309	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-c-Pr}$
310	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-i-Pr}$
311	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 3\text{-Ph}$
312	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-F}$
313	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-Cl}$
314	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-Br}$
315	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-I}$
316	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-CN}$
317	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-NO}_2$
318	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-OMe}$
319	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-OCF}_3$
320	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-CF}_3$
321	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-CHF}_2$
322	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-CH}_2\text{F}$
323	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-CHO}$
324	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-Me}$
325	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-Et}$
326	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-Ethylyl}$
327	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-Ethenyl}$
328	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-SO}_2\text{Me}$
329	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-OAc}$
330	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-c-Pr}$
331	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-i-Pr}$
332	$R^2 = \text{Cl}, Z = \text{S}, R^3 = 3\text{-Ph}$
333	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 4\text{-F}$
334	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 4\text{-Cl}$
335	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 4\text{-Br}$
336	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 4\text{-I}$
337	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 4\text{-CN}$
338	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 4\text{-NO}_2$
339	$R^2 = \text{Cl}, Z = \text{O}, R^3 = 4\text{-OMe}$

Table	Header Row
340	$R^2 = Cl, Z = O, R^3 = 4-OCF_3$
341	$R^2 = Cl, Z = O, R^3 = 4-CF_3$
342	$R^2 = Cl, Z = O, R^3 = 4-CHF_2$
343	$R^2 = Cl, Z = O, R^3 = 4-CH_2F$
344	$R^2 = Cl, Z = O, R^3 = 4-CHO$
345	$R^2 = Cl, Z = O, R^3 = 4-Me$
346	$R^2 = Cl, Z = O, R^3 = 4-Et$
347	$R^2 = Cl, Z = O, R^3 = 4-Ethynyl$
348	$R^2 = Cl, Z = O, R^3 = 4-Ethenyl$
349	$R^2 = Cl, Z = O, R^3 = 4-SO_2Me$
350	$R^2 = Cl, Z = O, R^3 = 4-OAc$
351	$R^2 = Cl, Z = O, R^3 = 4-c-Pr$
352	$R^2 = Cl, Z = O, R^3 = 4-i-Pr$
353	$R^2 = Cl, Z = O, R^3 = 4-Ph$
354	$R^2 = Cl, Z = O, R^3 = 5-F$
355	$R^2 = Cl, Z = O, R^3 = 5-Cl$
356	$R^2 = Cl, Z = O, R^3 = 5-Br$
357	$R^2 = Cl, Z = O, R^3 = 5-I$
358	$R^2 = Cl, Z = O, R^3 = 5-CN$
359	$R^2 = Cl, Z = O, R^3 = 5-NO_2$
360	$R^2 = Cl, Z = O, R^3 = 5-OMe$
361	$R^2 = Cl, Z = O, R^3 = 5-OCF_3$
362	$R^2 = Cl, Z = O, R^3 = 5-CF_3$
363	$R^2 = Cl, Z = O, R^3 = 5-CHF_2$
364	$R^2 = Cl, Z = O, R^3 = 5-CH_2F$
365	$R^2 = Cl, Z = O, R^3 = 5-CHO$
366	$R^2 = Cl, Z = O, R^3 = 5-Me$
367	$R^2 = Cl, Z = O, R^3 = 5-Et$
368	$R^2 = Cl, Z = O, R^3 = 5-Ethynyl$
369	$R^2 = Cl, Z = O, R^3 = 5-Ethenyl$
370	$R^2 = Cl, Z = O, R^3 = 5-SO_2Me$
371	$R^2 = Cl, Z = O, R^3 = 5-OAc$
372	$R^2 = Cl, Z = O, R^3 = 5-c-Pr$
373	$R^2 = Cl, Z = O, R^3 = 5-i-Pr$
374	$R^2 = Cl, Z = O, R^3 = 5-Ph$
375	$R^2 = Cl, Z = O, R^3 = 6-F$
376	$R^2 = Cl, Z = O, R^3 = 6-Cl$

Table	Header Row
377	$R^2 = Cl, Z = O, R^3 = 6-Br$
378	$R^2 = Cl, Z = O, R^3 = 6-I$
379	$R^2 = Cl, Z = O, R^3 = 6-CN$
380	$R^2 = Cl, Z = O, R^3 = 6-NO_2$
381	$R^2 = Cl, Z = O, R^3 = 6-OMe$
382	$R^2 = Cl, Z = O, R^3 = 6-OCF_3$
383	$R^2 = Cl, Z = O, R^3 = 6-CF_3$
384	$R^2 = Cl, Z = O, R^3 = 6-CHF_2$
385	$R^2 = Cl, Z = O, R^3 = 6-CH_2F$
386	$R^2 = Cl, Z = O, R^3 = 6-CHO$
387	$R^2 = Cl, Z = O, R^3 = 6-Me$
388	$R^2 = Cl, Z = O, R^3 = 6-Et$
389	$R^2 = Cl, Z = O, R^3 = 6-Ethynyl$
390	$R^2 = Cl, Z = O, R^3 = 6-Ethenyl$
391	$R^2 = Cl, Z = O, R^3 = 6-SO_2Me$
392	$R^2 = Cl, Z = O, R^3 = 6-OAc$
393	$R^2 = Cl, Z = O, R^3 = 6-c-Pr$
394	$R^2 = Cl, Z = O, R^3 = 6-i-Pr$
395	$R^2 = Cl, Z = O, R^3 = 6-Ph$
396	$R^2 = Cl, Z = O, R^3 = 3,4-di-F$
397	$R^2 = Cl, Z = O, R^3 = 3,5-di-F$
398	$R^2 = Cl, Z = O, R^3 = 3,6-di-F$
399	$R^2 = Cl, Z = O, R^3 = 4,5-di-F$
400	$R^2 = Cl, Z = O, R^3 = 3,4-di-Cl$
401	$R^2 = Cl, Z = O, R^3 = 3,5-di-Cl$
402	$R^2 = Cl, Z = O, R^3 = 3,6-di-Cl$
403	$R^2 = Cl, Z = O, R^3 = 4,5-di-Cl$
404	$R^2 = Cl, Z = O, R^3 = 3,4-di-Br$
405	$R^2 = Cl, Z = O, R^3 = 3,5-di-Br$
406	$R^2 = Cl, Z = O, R^3 = 3,6-di-Br$
407	$R^2 = Cl, Z = O, R^3 = 4,5-di-Br$
408	$R^2 = Cl, Z = O, R^3 = 3,4-di-CN$
409	$R^2 = Cl, Z = O, R^3 = 3,5-di-CN$
410	$R^2 = Cl, Z = O, R^3 = 3,6-di-CN$
411	$R^2 = Cl, Z = O, R^3 = 4,5-di-CN$
412	$R^2 = Cl, Z = O, R^3 = 3,4-di-Me$
413	$R^2 = Cl, Z = O, R^3 = 3,5-di-Me$

Table	Header Row
414	$R^2 = Cl, Z = O, R^3 = 3,6\text{-di-Me}$
415	$R^2 = Cl, Z = O, R^3 = 4,5\text{-di-Me}$
416	$R^2 = Cl, Z = O, R^3 = 3,4\text{-di-OMe}$
417	$R^2 = Cl, Z = O, R^3 = 3,5\text{-di-OMe}$
418	$R^2 = Cl, Z = O, R^3 = 3,6\text{-di-OMe}$
419	$R^2 = Cl, Z = O, R^3 = 4,5\text{-di-OMe}$
420	$R^2 = Cl, Z = O, R^3 = 3,4\text{-di-CF}_3$
421	$R^2 = Cl, Z = O, R^3 = 3,5\text{-di-CF}_3$
422	$R^2 = Cl, Z = O, R^3 = 3,6\text{-di-CF}_3$
423	$R^2 = Cl, Z = O, R^3 = 4,5\text{-di-CF}_3$
424	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 4-Me}$
425	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 4-F}$
426	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 4-Br}$
427	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 4-OMe}$
428	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 4-CF}_3$
429	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 6-Me}$
430	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 6-F}$
431	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 6-Br}$
432	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 6-OMe}$
433	$R^2 = Cl, Z = O, R^3 = 3\text{-CN, 6-CF}_3$
434	$R^2 = I, Z = O, R^3 = H (m = 0)$
435	$R^2 = I, Z = O, R^3 = 3\text{-F}$
436	$R^2 = I, Z = O, R^3 = 3\text{-Cl}$
437	$R^2 = I, Z = O, R^3 = 3\text{-Br}$
438	$R^2 = I, Z = O, R^3 = 3\text{-I}$
439	$R^2 = I, Z = O, R^3 = 3\text{-CN}$
440	$R^2 = I, Z = O, R^3 = 3\text{-NO}_2$
441	$R^2 = I, Z = O, R^3 = 3\text{-OMe}$
442	$R^2 = I, Z = O, R^3 = 3\text{-OCF}_3$
443	$R^2 = I, Z = O, R^3 = 3\text{-CF}_3$
444	$R^2 = I, Z = O, R^3 = 3\text{-CHF}_2$
445	$R^2 = I, Z = O, R^3 = 3\text{-CH}_2\text{F}$
446	$R^2 = I, Z = O, R^3 = 3\text{-CHO}$
447	$R^2 = I, Z = O, R^3 = 3\text{-Me}$
448	$R^2 = I, Z = O, R^3 = 3\text{-Et}$
449	$R^2 = I, Z = O, R^3 = 3\text{-Ethyneyl}$
450	$R^2 = I, Z = O, R^3 = 3\text{-Ethenyl}$

Table	Header Row
451	$R^2 = I, Z = O, R^3 = 3\text{-SO}_2\text{Me}$
452	$R^2 = I, Z = O, R^3 = 3\text{-OAc}$
453	$R^2 = I, Z = O, R^3 = 3\text{-c-Pr}$
454	$R^2 = I, Z = O, R^3 = 3\text{-i-Pr}$
455	$R^2 = I, Z = O, R^3 = 3\text{-Ph}$
456	$R^2 = I, Z = S, R^3 = 3\text{-F}$
457	$R^2 = I, Z = S, R^3 = 3\text{-Cl}$
458	$R^2 = I, Z = S, R^3 = 3\text{-Br}$
459	$R^2 = I, Z = S, R^3 = 3\text{-I}$
460	$R^2 = I, Z = S, R^3 = 3\text{-CN}$
461	$R^2 = I, Z = S, R^3 = 3\text{-NO}_2$
462	$R^2 = I, Z = S, R^3 = 3\text{-OMe}$
463	$R^2 = I, Z = S, R^3 = 3\text{-OCF}_3$
464	$R^2 = I, Z = S, R^3 = 3\text{-CF}_3$
465	$R^2 = I, Z = S, R^3 = 3\text{-CHF}_2$
466	$R^2 = I, Z = S, R^3 = 3\text{-CH}_2\text{F}$
467	$R^2 = I, Z = S, R^3 = 3\text{-CHO}$
468	$R^2 = I, Z = S, R^3 = 3\text{-Me}$
469	$R^2 = I, Z = S, R^3 = 3\text{-Et}$
470	$R^2 = I, Z = S, R^3 = 3\text{-Ethyneyl}$
471	$R^2 = I, Z = S, R^3 = 3\text{-Ethenyl}$
472	$R^2 = I, Z = S, R^3 = 3\text{-SO}_2\text{Me}$
473	$R^2 = I, Z = S, R^3 = 3\text{-OAc}$
474	$R^2 = I, Z = S, R^3 = 3\text{-c-Pr}$
475	$R^2 = I, Z = S, R^3 = 3\text{-i-Pr}$
476	$R^2 = I, Z = S, R^3 = 3\text{-Ph}$
477	$R^2 = I, Z = O, R^3 = 4\text{-F}$
478	$R^2 = I, Z = O, R^3 = 4\text{-Cl}$
479	$R^2 = I, Z = O, R^3 = 4\text{-Br}$
480	$R^2 = I, Z = O, R^3 = 4\text{-I}$
481	$R^2 = I, Z = O, R^3 = 4\text{-CN}$
482	$R^2 = I, Z = O, R^3 = 4\text{-NO}_2$
483	$R^2 = I, Z = O, R^3 = 4\text{-OMe}$
484	$R^2 = I, Z = O, R^3 = 4\text{-OCF}_3$
485	$R^2 = I, Z = O, R^3 = 4\text{-CF}_3$
486	$R^2 = I, Z = O, R^3 = 4\text{-CHF}_2$
487	$R^2 = I, Z = O, R^3 = 4\text{-CH}_2\text{F}$

Table	Header Row
488	$R^2 = I, Z = O, R^3 = 4\text{-CHO}$
489	$R^2 = I, Z = O, R^3 = 4\text{-Me}$
490	$R^2 = I, Z = O, R^3 = 4\text{-Et}$
491	$R^2 = I, Z = O, R^3 = 4\text{-Ethyneyl}$
492	$R^2 = I, Z = O, R^3 = 4\text{-Ethenyl}$
493	$R^2 = I, Z = O, R^3 = 4\text{-SO}_2\text{Me}$
494	$R^2 = I, Z = O, R^3 = 4\text{-OAc}$
495	$R^2 = I, Z = O, R^3 = 4\text{-}i\text{-Pr}$
496	$R^2 = I, Z = O, R^3 = 4\text{-}i\text{-Pr}$
497	$R^2 = I, Z = O, R^3 = 4\text{-Ph}$
498	$R^2 = I, Z = O, R^3 = 5\text{-F}$
499	$R^2 = I, Z = O, R^3 = 5\text{-Cl}$
500	$R^2 = I, Z = O, R^3 = 5\text{-Br}$
501	$R^2 = I, Z = O, R^3 = 5\text{-I}$
502	$R^2 = I, Z = O, R^3 = 5\text{-CN}$
503	$R^2 = I, Z = O, R^3 = 5\text{-NO}_2$
504	$R^2 = I, Z = O, R^3 = 5\text{-OMe}$
505	$R^2 = I, Z = O, R^3 = 5\text{-OCF}_3$
506	$R^2 = I, Z = O, R^3 = 5\text{-CF}_3$
507	$R^2 = I, Z = O, R^3 = 5\text{-CHF}_2$
508	$R^2 = I, Z = O, R^3 = 5\text{-CH}_2\text{F}$
509	$R^2 = I, Z = O, R^3 = 5\text{-CHO}$
510	$R^2 = I, Z = O, R^3 = 5\text{-Me}$
511	$R^2 = I, Z = O, R^3 = 5\text{-Et}$
512	$R^2 = I, Z = O, R^3 = 5\text{-Ethyneyl}$
513	$R^2 = I, Z = O, R^3 = 5\text{-Ethenyl}$
514	$R^2 = I, Z = O, R^3 = 5\text{-SO}_2\text{Me}$
515	$R^2 = I, Z = O, R^3 = 5\text{-OAc}$
516	$R^2 = I, Z = O, R^3 = 5\text{-}i\text{-Pr}$
517	$R^2 = I, Z = O, R^3 = 5\text{-}i\text{-Pr}$
518	$R^2 = I, Z = O, R^3 = 5\text{-Ph}$
519	$R^2 = I, Z = O, R^3 = 6\text{-F}$
520	$R^2 = I, Z = O, R^3 = 6\text{-Cl}$
521	$R^2 = I, Z = O, R^3 = 6\text{-Br}$
522	$R^2 = I, Z = O, R^3 = 6\text{-I}$
523	$R^2 = I, Z = O, R^3 = 6\text{-CN}$
524	$R^2 = I, Z = O, R^3 = 6\text{-NO}_2$

Table	Header Row
525	$R^2 = I, Z = O, R^3 = 6\text{-OMe}$
526	$R^2 = I, Z = O, R^3 = 6\text{-OCF}_3$
527	$R^2 = I, Z = O, R^3 = 6\text{-CF}_3$
528	$R^2 = I, Z = O, R^3 = 6\text{-CHF}_2$
529	$R^2 = I, Z = O, R^3 = 6\text{-CH}_2\text{F}$
530	$R^2 = I, Z = O, R^3 = 6\text{-CHO}$
531	$R^2 = I, Z = O, R^3 = 6\text{-Me}$
532	$R^2 = I, Z = O, R^3 = 6\text{-Et}$
533	$R^2 = I, Z = O, R^3 = 6\text{-Ethyneyl}$
534	$R^2 = I, Z = O, R^3 = 6\text{-Ethenyl}$
535	$R^2 = I, Z = O, R^3 = 6\text{-SO}_2\text{Me}$
536	$R^2 = I, Z = O, R^3 = 6\text{-OAc}$
537	$R^2 = I, Z = O, R^3 = 6\text{-}i\text{-Pr}$
538	$R^2 = I, Z = O, R^3 = 6\text{-}i\text{-Pr}$
539	$R^2 = I, Z = O, R^3 = 6\text{-Ph}$
540	$R^2 = I, Z = O, R^3 = 3,4\text{-di-F}$
541	$R^2 = I, Z = O, R^3 = 3,5\text{-di-F}$
542	$R^2 = I, Z = O, R^3 = 3,6\text{-di-F}$
543	$R^2 = I, Z = O, R^3 = 4,5\text{-di-F}$
544	$R^2 = I, Z = O, R^3 = 3,4\text{-di-Cl}$
545	$R^2 = I, Z = O, R^3 = 3,5\text{-di-Cl}$
546	$R^2 = I, Z = O, R^3 = 3,6\text{-di-Cl}$
547	$R^2 = I, Z = O, R^3 = 4,5\text{-di-Cl}$
548	$R^2 = I, Z = O, R^3 = 3,4\text{-di-Br}$
549	$R^2 = I, Z = O, R^3 = 3,5\text{-di-Br}$
550	$R^2 = I, Z = O, R^3 = 3,6\text{-di-Br}$
551	$R^2 = I, Z = O, R^3 = 4,5\text{-di-Br}$
552	$R^2 = I, Z = O, R^3 = 3,4\text{-di-CN}$
553	$R^2 = I, Z = O, R^3 = 3,5\text{-di-CN}$
554	$R^2 = I, Z = O, R^3 = 3,6\text{-di-CN}$
555	$R^2 = I, Z = O, R^3 = 4,5\text{-di-CN}$
556	$R^2 = I, Z = O, R^3 = 3,4\text{-di-Me}$
557	$R^2 = I, Z = O, R^3 = 3,5\text{-di-Me}$
558	$R^2 = I, Z = O, R^3 = 3,6\text{-di-Me}$
559	$R^2 = I, Z = O, R^3 = 4,5\text{-di-Me}$
560	$R^2 = I, Z = O, R^3 = 3,4\text{-di-OMe}$
561	$R^2 = I, Z = O, R^3 = 3,5\text{-di-OMe}$

Table	Header Row
562	$R^2 = I, Z = O, R^3 = 3,6\text{-di-OMe}$
563	$R^2 = I, Z = O, R^3 = 4,5\text{-di-OMe}$
564	$R^2 = I, Z = O, R^3 = 3,4\text{-di-CF}_3$
565	$R^2 = I, Z = O, R^3 = 3,5\text{-di-CF}_3$
566	$R^2 = I, Z = O, R^3 = 3,6\text{-di-CF}_3$
567	$R^2 = I, Z = O, R^3 = 4,5\text{-di-CF}_3$
568	$R^2 = I, Z = O, R^3 = 3\text{-CN, 4-Me}$
569	$R^2 = I, Z = O, R^3 = 3\text{-CN, 4-F}$
570	$R^2 = I, Z = O, R^3 = 3\text{-CN, 4-Br}$
571	$R^2 = I, Z = O, R^3 = 3\text{-CN, 4-OMe}$
572	$R^2 = I, Z = O, R^3 = 3\text{-CN, 4-CF}_3$
573	$R^2 = I, Z = O, R^3 = 3\text{-CN, 6-Me}$
574	$R^2 = I, Z = O, R^3 = 3\text{-CN, 6-F}$
575	$R^2 = I, Z = O, R^3 = 3\text{-CN, 6-Br}$
576	$R^2 = I, Z = O, R^3 = 3\text{-CN, 6-OMe}$
577	$R^2 = I, Z = O, R^3 = 3\text{-CN, 6-CF}_3$
578	$R^2 = Me, Z = O, R^3 = H (m = 0)$
579	$R^2 = Me, Z = O, R^3 = 3\text{-F}$
580	$R^2 = Me, Z = O, R^3 = 3\text{-Cl}$
581	$R^2 = Me, Z = O, R^3 = 3\text{-Br}$
582	$R^2 = Me, Z = O, R^3 = 3\text{-I}$
583	$R^2 = Me, Z = O, R^3 = 3\text{-CN}$
584	$R^2 = Me, Z = O, R^3 = 3\text{-NO}_2$
585	$R^2 = Me, Z = O, R^3 = 3\text{-OMe}$
586	$R^2 = Me, Z = O, R^3 = 3\text{-OCF}_3$
587	$R^2 = Me, Z = O, R^3 = 3\text{-CF}_3$
588	$R^2 = Me, Z = O, R^3 = 3\text{-CHF}_2$
589	$R^2 = Me, Z = O, R^3 = 3\text{-CH}_2\text{F}$
590	$R^2 = Me, Z = O, R^3 = 3\text{-CHO}$
591	$R^2 = Me, Z = O, R^3 = 3\text{-Me}$
592	$R^2 = Me, Z = O, R^3 = 3\text{-Et}$
593	$R^2 = Me, Z = O, R^3 = 3\text{-Ethynyl}$
594	$R^2 = Me, Z = O, R^3 = 3\text{-Ethenyl}$
595	$R^2 = Me, Z = O, R^3 = 3\text{-SO}_2\text{Me}$
596	$R^2 = Me, Z = O, R^3 = 3\text{-OAc}$
597	$R^2 = Me, Z = O, R^3 = 3\text{-c-Pr}$
598	$R^2 = Me, Z = O, R^3 = 3\text{-i-Pr}$

Table	Header Row
599	$R^2 = Me, Z = O, R^3 = 3\text{-Ph}$
600	$R^2 = Me, Z = S, R^3 = 3\text{-F}$
601	$R^2 = Me, Z = S, R^3 = 3\text{-Cl}$
602	$R^2 = Me, Z = S, R^3 = 3\text{-Br}$
603	$R^2 = Me, Z = S, R^3 = 3\text{-I}$
604	$R^2 = Me, Z = S, R^3 = 3\text{-CN}$
605	$R^2 = Me, Z = S, R^3 = 3\text{-NO}_2$
606	$R^2 = Me, Z = S, R^3 = 3\text{-OMe}$
607	$R^2 = Me, Z = S, R^3 = 3\text{-OCF}_3$
608	$R^2 = Me, Z = S, R^3 = 3\text{-CF}_3$
609	$R^2 = Me, Z = S, R^3 = 3\text{-CHF}_2$
610	$R^2 = Me, Z = S, R^3 = 3\text{-CH}_2\text{F}$
611	$R^2 = Me, Z = S, R^3 = 3\text{-CHO}$
612	$R^2 = Me, Z = S, R^3 = 3\text{-Me}$
613	$R^2 = Me, Z = S, R^3 = 3\text{-Et}$
614	$R^2 = Me, Z = S, R^3 = 3\text{-Ethynyl}$
615	$R^2 = Me, Z = S, R^3 = 3\text{-Ethenyl}$
616	$R^2 = Me, Z = S, R^3 = 3\text{-SO}_2\text{Me}$
617	$R^2 = Me, Z = S, R^3 = 3\text{-OAc}$
618	$R^2 = Me, Z = S, R^3 = 3\text{-c-Pr}$
619	$R^2 = Me, Z = S, R^3 = 3\text{-i-Pr}$
620	$R^2 = Me, Z = S, R^3 = 3\text{-Ph}$
621	$R^2 = Me, Z = O, R^3 = 4\text{-F}$
622	$R^2 = Me, Z = O, R^3 = 4\text{-Cl}$
623	$R^2 = Me, Z = O, R^3 = 4\text{-Br}$
624	$R^2 = Me, Z = O, R^3 = 4\text{-I}$
625	$R^2 = Me, Z = O, R^3 = 4\text{-CN}$
626	$R^2 = Me, Z = O, R^3 = 4\text{-NO}_2$
627	$R^2 = Me, Z = O, R^3 = 4\text{-OMe}$
628	$R^2 = Me, Z = O, R^3 = 4\text{-OCF}_3$
629	$R^2 = Me, Z = O, R^3 = 4\text{-CF}_3$
630	$R^2 = Me, Z = O, R^3 = 4\text{-CHF}_2$
631	$R^2 = Me, Z = O, R^3 = 4\text{-CH}_2\text{F}$
632	$R^2 = Me, Z = O, R^3 = 4\text{-CHO}$
633	$R^2 = Me, Z = O, R^3 = 4\text{-Me}$
634	$R^2 = Me, Z = O, R^3 = 4\text{-Et}$
635	$R^2 = Me, Z = O, R^3 = 4\text{-Ethynyl}$

Table	Header Row
636	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4\text{-Ethenyl}$
637	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4\text{-SO}_2\text{Me}$
638	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4\text{-OAc}$
639	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4\text{-}i\text{-Pr}$
640	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4\text{-}i\text{-Pr}$
641	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4\text{-Ph}$
642	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-F}$
643	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-Cl}$
644	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-Br}$
645	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-I}$
646	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-CN}$
647	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-NO}_2$
648	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-OMe}$
649	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-OCF}_3$
650	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-CF}_3$
651	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-CHF}_2$
652	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-CH}_2\text{F}$
653	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-CHO}$
654	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-Me}$
655	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-Et}$
656	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-Ethinyl}$
657	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-Ethenyl}$
658	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-SO}_2\text{Me}$
659	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-OAc}$
660	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$
661	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$
662	$R^2 = \text{Me}, Z = \text{O}, R^3 = 5\text{-Ph}$
663	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-F}$
664	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-Cl}$
665	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-Br}$
666	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-I}$
667	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-CN}$
668	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-NO}_2$
669	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-OMe}$
670	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-OCF}_3$
671	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-CF}_3$
672	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-CHF}_2$

Table	Header Row
673	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-CH}_2\text{F}$
674	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-CHO}$
675	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-Me}$
676	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-Et}$
677	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-Ethinyl}$
678	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-Ethenyl}$
679	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-SO}_2\text{Me}$
680	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-OAc}$
681	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
682	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
683	$R^2 = \text{Me}, Z = \text{O}, R^3 = 6\text{-Ph}$
684	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-F}$
685	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-F}$
686	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-F}$
687	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-F}$
688	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-Cl}$
689	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-Cl}$
690	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-Cl}$
691	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-Cl}$
692	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-Br}$
693	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-Br}$
694	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-Br}$
695	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-Br}$
696	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-CN}$
697	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-CN}$
698	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-CN}$
699	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-CN}$
700	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-Me}$
701	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-Me}$
702	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-Me}$
703	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-Me}$
704	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-OMe}$
705	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-OMe}$
706	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-OMe}$
707	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-OMe}$
708	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-CF}_3$
709	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-CF}_3$

Table	Header Row
710	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-CF}_3$
711	$R^2 = \text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-CF}_3$
712	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Me}$
713	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-F}$
714	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Br}$
715	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-OMe}$
716	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-CF}_3$
717	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Me}$
718	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-F}$
719	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Br}$
720	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-OMe}$
721	$R^2 = \text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-CF}_3$
722	$R^2 = \text{CN}, Z = \text{O}, R^3 = \text{H} (m = 0)$
723	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-F}$
724	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-Cl}$
725	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-Br}$
726	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-I}$
727	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}$
728	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-NO}_2$
729	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-OMe}$
730	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-OCF}_3$
731	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CF}_3$
732	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CHF}_2$
733	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CH}_2\text{F}$
734	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CHO}$
735	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-Me}$
736	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-Et}$
737	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-Ethyne}$
738	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-Ethenyl}$
739	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-SO}_2\text{Me}$
740	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-OAc}$
741	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-c-Pr}$
742	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-i-Pr}$
743	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-Ph}$
744	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-F}$
745	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-Cl}$
746	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-Br}$

Table	Header Row
747	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-I}$
748	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-CN}$
749	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-NO}_2$
750	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-OMe}$
751	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-OCF}_3$
752	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-CF}_3$
753	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-CHF}_2$
754	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-CH}_2\text{F}$
755	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-CHO}$
756	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-Me}$
757	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-Et}$
758	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-Ethyne}$
759	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-Ethenyl}$
760	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-SO}_2\text{Me}$
761	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-OAc}$
762	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-c-Pr}$
763	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-i-Pr}$
764	$R^2 = \text{CN}, Z = \text{S}, R^3 = 3\text{-Ph}$
765	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-F}$
766	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-Cl}$
767	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-Br}$
768	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-I}$
769	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-CN}$
770	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-NO}_2$
771	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-OMe}$
772	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-OCF}_3$
773	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-CF}_3$
774	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-CHF}_2$
775	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-CH}_2\text{F}$
776	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-CHO}$
777	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-Me}$
778	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-Et}$
779	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-Ethyne}$
780	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-Ethenyl}$
781	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-SO}_2\text{Me}$
782	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-OAc}$
783	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-c-Pr}$

Table	Header Row
784	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-}i\text{-Pr}$
785	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4\text{-Ph}$
786	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-F}$
787	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-Cl}$
788	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-Br}$
789	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-I}$
790	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-CN}$
791	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-NO}_2$
792	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-OMe}$
793	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-OCF}_3$
794	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-CF}_3$
795	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-CHF}_2$
796	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-CH}_2\text{F}$
797	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-CHO}$
798	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-Me}$
799	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-Et}$
800	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-Ethyne}$
801	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-Ethenyl}$
802	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-SO}_2\text{Me}$
803	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-OAc}$
804	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$
805	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$
806	$R^2 = \text{CN}, Z = \text{O}, R^3 = 5\text{-Ph}$
807	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-F}$
808	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-Cl}$
809	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-Br}$
810	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-I}$
811	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-CN}$
812	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-NO}_2$
813	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-OMe}$
814	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-OCF}_3$
815	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-CF}_3$
816	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-CHF}_2$
817	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-CH}_2\text{F}$
818	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-CHO}$
819	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-Me}$
820	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-Et}$

Table	Header Row
821	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-Ethyne}$
822	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-Ethenyl}$
823	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-SO}_2\text{Me}$
824	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-OAc}$
825	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
826	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
827	$R^2 = \text{CN}, Z = \text{O}, R^3 = 6\text{-Ph}$
828	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,4\text{-di-F}$
829	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,5\text{-di-F}$
830	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,6\text{-di-F}$
831	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4,5\text{-di-F}$
832	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,4\text{-di-Cl}$
833	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,5\text{-di-Cl}$
834	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,6\text{-di-Cl}$
835	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4,5\text{-di-Cl}$
836	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,4\text{-di-Br}$
837	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,5\text{-di-Br}$
838	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,6\text{-di-Br}$
839	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4,5\text{-di-Br}$
840	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,4\text{-di-CN}$
841	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,5\text{-di-CN}$
842	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,6\text{-di-CN}$
843	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4,5\text{-di-CN}$
844	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,4\text{-di-Me}$
845	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,5\text{-di-Me}$
846	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,6\text{-di-Me}$
847	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4,5\text{-di-Me}$
848	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,4\text{-di-OMe}$
849	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,5\text{-di-OMe}$
850	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,6\text{-di-OMe}$
851	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4,5\text{-di-OMe}$
852	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,4\text{-di-CF}_3$
853	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,5\text{-di-CF}_3$
854	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3,6\text{-di-CF}_3$
855	$R^2 = \text{CN}, Z = \text{O}, R^3 = 4,5\text{-di-CF}_3$
856	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Me}$
857	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-F}$

Table	Header Row
858	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Br}$
859	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-OMe}$
860	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-CF}_3$
861	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Me}$
862	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-F}$
863	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Br}$
864	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-OMe}$
865	$R^2 = \text{CN}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-CF}_3$
866	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = \text{H} (m = 0)$
867	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-F}$
868	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-Cl}$
869	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-Br}$
870	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-I}$
871	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}$
872	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-NO}_2$
873	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-OMe}$
874	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-OCF}_3$
875	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CF}_3$
876	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CHF}_2$
877	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CH}_2\text{F}$
878	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CHO}$
879	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-Me}$
880	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-Et}$
881	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-Ethyneyl}$
882	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-Ethenyl}$
883	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-SO}_2\text{Me}$
884	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-OAc}$
885	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-c-Pr}$
886	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-i-Pr}$
887	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-Ph}$
888	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-F}$
889	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-Cl}$
890	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-Br}$
891	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-I}$
892	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-CN}$
893	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-NO}_2$
894	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-OMe}$

Table	Header Row
895	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-OCF}_3$
896	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-CF}_3$
897	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-CHF}_2$
898	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-CH}_2\text{F}$
899	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-CHO}$
900	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-Me}$
901	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-Et}$
902	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-Ethyneyl}$
903	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-Ethenyl}$
904	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-SO}_2\text{Me}$
905	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-OAc}$
906	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-c-Pr}$
907	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-i-Pr}$
908	$R^2 = \text{NO}_2, Z = \text{S}, R^3 = 3\text{-Ph}$
909	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-F}$
910	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-Cl}$
911	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-Br}$
912	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-I}$
913	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-CN}$
914	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-NO}_2$
915	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-OMe}$
916	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-OCF}_3$
917	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-CF}_3$
918	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-CHF}_2$
919	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-CH}_2\text{F}$
920	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-CHO}$
921	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-Me}$
922	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-Et}$
923	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-Ethyneyl}$
924	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-Ethenyl}$
925	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-SO}_2\text{Me}$
926	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-OAc}$
927	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-c-Pr}$
928	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-i-Pr}$
929	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4\text{-Ph}$
930	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-F}$
931	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-Cl}$

Table	Header Row
932	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-Br}$
933	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-I}$
934	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-CN}$
935	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-NO}_2$
936	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-OMe}$
937	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-OCF}_3$
938	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-CF}_3$
939	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-CHF}_2$
940	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-CH}_2\text{F}$
941	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-CHO}$
942	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-Me}$
943	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-Et}$
944	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-Ethynyl}$
945	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-Ethenyl}$
946	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-SO}_2\text{Me}$
947	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-OAc}$
948	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-c-Pr}$
949	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-i-Pr}$
950	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 5\text{-Ph}$
951	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-F}$
952	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-Cl}$
953	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-Br}$
954	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-I}$
955	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-CN}$
956	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-NO}_2$
957	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-OMe}$
958	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-OCF}_3$
959	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-CF}_3$
960	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-CHF}_2$
961	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-CH}_2\text{F}$
962	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-CHO}$
963	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-Me}$
964	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-Et}$
965	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-Ethynyl}$
966	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-Ethenyl}$
967	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-SO}_2\text{Me}$
968	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-OAc}$

Table	Header Row
969	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-c-Pr}$
970	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-i-Pr}$
971	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 6\text{-Ph}$
972	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,4\text{-di-F}$
973	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,5\text{-di-F}$
974	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,6\text{-di-F}$
975	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4,5\text{-di-F}$
976	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,4\text{-di-Cl}$
977	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,5\text{-di-Cl}$
978	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,6\text{-di-Cl}$
979	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4,5\text{-di-Cl}$
980	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,4\text{-di-Br}$
981	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,5\text{-di-Br}$
982	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,6\text{-di-Br}$
983	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4,5\text{-di-Br}$
984	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,4\text{-di-CN}$
985	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,5\text{-di-CN}$
986	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,6\text{-di-CN}$
987	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4,5\text{-di-CN}$
988	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,4\text{-di-Me}$
989	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,5\text{-di-Me}$
990	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,6\text{-di-Me}$
991	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4,5\text{-di-Me}$
992	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,4\text{-di-OMe}$
993	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,5\text{-di-OMe}$
994	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,6\text{-di-OMe}$
995	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4,5\text{-di-OMe}$
996	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,4\text{-di-CF}_3$
997	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,5\text{-di-CF}_3$
998	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3,6\text{-di-CF}_3$
999	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 4,5\text{-di-CF}_3$
1000	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Me}$
1001	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-F}$
1002	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Br}$
1003	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-OMe}$
1004	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-CF}_3$
1005	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Me}$

Table	Header Row
1006	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-F}$
1007	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Br}$
1008	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-OMe}$
1009	$R^2 = \text{NO}_2, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-CF}_3$
1010	$R^2 = \text{OMe}, Z = \text{O}, R^3 = \text{H} (m = 0)$
1011	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-F}$
1012	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-Cl}$
1013	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-Br}$
1014	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-I}$
1015	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}$
1016	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-NO}_2$
1017	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-OMe}$
1018	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-OCF}_3$
1019	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CF}_3$
1020	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CHF}_2$
1021	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CH}_2\text{F}$
1022	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CHO}$
1023	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-Me}$
1024	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-Et}$
1025	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-Ethyneyl}$
1026	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-Ethenyl}$
1027	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-SO}_2\text{Me}$
1028	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-OAc}$
1029	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-c-Pr}$
1030	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-i-Pr}$
1031	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-Ph}$
1032	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-F}$
1033	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-Cl}$
1034	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-Br}$
1035	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-I}$
1036	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-CN}$
1037	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-NO}_2$
1038	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-OMe}$
1039	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-OCF}_3$
1040	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-CF}_3$
1041	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-CHF}_2$
1042	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-CH}_2\text{F}$

Table	Header Row
1043	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-CHO}$
1044	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-Me}$
1045	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-Et}$
1046	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-Ethyneyl}$
1047	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-Ethenyl}$
1048	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-SO}_2\text{Me}$
1049	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-OAc}$
1050	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-c-Pr}$
1051	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-i-Pr}$
1052	$R^2 = \text{OMe}, Z = \text{S}, R^3 = 3\text{-Ph}$
1053	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-F}$
1054	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-Cl}$
1055	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-Br}$
1056	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-I}$
1057	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-CN}$
1058	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-NO}_2$
1059	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-OMe}$
1060	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-OCF}_3$
1061	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-CF}_3$
1062	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-CHF}_2$
1063	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-CH}_2\text{F}$
1064	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-CHO}$
1065	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-Me}$
1066	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-Et}$
1067	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-Ethyneyl}$
1068	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-Ethenyl}$
1069	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-SO}_2\text{Me}$
1070	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-OAc}$
1071	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-c-Pr}$
1072	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-i-Pr}$
1073	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4\text{-Ph}$
1074	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-F}$
1075	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-Cl}$
1076	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-Br}$
1077	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-I}$
1078	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-CN}$
1079	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-NO}_2$

Table	Header Row
1080	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-OMe}$
1081	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-OCF}_3$
1082	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-CF}_3$
1083	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-CHF}_2$
1084	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-CH}_2\text{F}$
1085	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-CHO}$
1086	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-Me}$
1087	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-Et}$
1088	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-Ethyneyl}$
1089	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-Ethenyl}$
1090	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-SO}_2\text{Me}$
1091	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-OAc}$
1092	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$
1093	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$
1094	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 5\text{-Ph}$
1095	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-F}$
1096	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-Cl}$
1097	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-Br}$
1098	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-I}$
1099	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-CN}$
1100	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-NO}_2$
1101	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-OMe}$
1102	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-OCF}_3$
1103	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-CF}_3$
1104	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-CHF}_2$
1105	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-CH}_2\text{F}$
1106	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-CHO}$
1107	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-Me}$
1108	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-Et}$
1109	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-Ethyneyl}$
1110	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-Ethenyl}$
1111	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-SO}_2\text{Me}$
1112	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-OAc}$
1113	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
1114	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
1115	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 6\text{-Ph}$
1116	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,4\text{-di-F}$

Table	Header Row
1117	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,5\text{-di-F}$
1118	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,6\text{-di-F}$
1119	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4,5\text{-di-F}$
1120	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,4\text{-di-Cl}$
1121	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,5\text{-di-Cl}$
1122	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,6\text{-di-Cl}$
1123	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4,5\text{-di-Cl}$
1124	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,4\text{-di-Br}$
1125	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,5\text{-di-Br}$
1126	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,6\text{-di-Br}$
1127	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4,5\text{-di-Br}$
1128	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,4\text{-di-CN}$
1129	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,5\text{-di-CN}$
1130	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,6\text{-di-CN}$
1131	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4,5\text{-di-CN}$
1132	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,4\text{-di-Me}$
1133	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,5\text{-di-Me}$
1134	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,6\text{-di-Me}$
1135	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4,5\text{-di-Me}$
1136	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,4\text{-di-OMe}$
1137	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,5\text{-di-OMe}$
1138	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,6\text{-di-OMe}$
1139	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4,5\text{-di-OMe}$
1140	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,4\text{-di-CF}_3$
1141	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,5\text{-di-CF}_3$
1142	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3,6\text{-di-CF}_3$
1143	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 4,5\text{-di-CF}_3$
1144	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Me}$
1145	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-F}$
1146	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Br}$
1147	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-OMe}$
1148	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-CF}_3$
1149	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Me}$
1150	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-F}$
1151	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Br}$
1152	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-OMe}$
1153	$R^2 = \text{OMe}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-CF}_3$

Table	Header Row
1154	$R^2 = CF_3, Z = O, R^3 = H (m = 0)$
1155	$R^2 = CF_3, Z = O, R^3 = 3-F$
1156	$R^2 = CF_3, Z = O, R^3 = 3-Cl$
1157	$R^2 = CF_3, Z = O, R^3 = 3-Br$
1158	$R^2 = CF_3, Z = O, R^3 = 3-I$
1159	$R^2 = CF_3, Z = O, R^3 = 3-CN$
1160	$R^2 = CF_3, Z = O, R^3 = 3-NO_2$
1161	$R^2 = CF_3, Z = O, R^3 = 3-OMe$
1162	$R^2 = CF_3, Z = O, R^3 = 3-OCF_3$
1163	$R^2 = CF_3, Z = O, R^3 = 3-CF_3$
1164	$R^2 = CF_3, Z = O, R^3 = 3-CHF_2$
1165	$R^2 = CF_3, Z = O, R^3 = 3-CH_2F$
1166	$R^2 = CF_3, Z = O, R^3 = 3-CHO$
1167	$R^2 = CF_3, Z = O, R^3 = 3-Me$
1168	$R^2 = CF_3, Z = O, R^3 = 3-Et$
1169	$R^2 = CF_3, Z = O, R^3 = 3-Ethynyl$
1170	$R^2 = CF_3, Z = O, R^3 = 3-Ethenyl$
1171	$R^2 = CF_3, Z = O, R^3 = 3-SO_2Me$
1172	$R^2 = CF_3, Z = O, R^3 = 3-OAc$
1173	$R^2 = CF_3, Z = O, R^3 = 3-c-Pr$
1174	$R^2 = CF_3, Z = O, R^3 = 3-i-Pr$
1175	$R^2 = CF_3, Z = O, R^3 = 3-Ph$
1176	$R^2 = CF_3, Z = S, R^3 = 3-F$
1177	$R^2 = CF_3, Z = S, R^3 = 3-Cl$
1178	$R^2 = CF_3, Z = S, R^3 = 3-Br$
1179	$R^2 = CF_3, Z = S, R^3 = 3-I$
1180	$R^2 = CF_3, Z = S, R^3 = 3-CN$
1181	$R^2 = CF_3, Z = S, R^3 = 3-NO_2$
1182	$R^2 = CF_3, Z = S, R^3 = 3-OMe$
1183	$R^2 = CF_3, Z = S, R^3 = 3-OCF_3$
1184	$R^2 = CF_3, Z = S, R^3 = 3-CF_3$
1185	$R^2 = CF_3, Z = S, R^3 = 3-CHF_2$
1186	$R^2 = CF_3, Z = S, R^3 = 3-CH_2F$
1187	$R^2 = CF_3, Z = S, R^3 = 3-CHO$
1188	$R^2 = CF_3, Z = S, R^3 = 3-Me$
1189	$R^2 = CF_3, Z = S, R^3 = 3-Et$
1190	$R^2 = CF_3, Z = S, R^3 = 3-Ethynyl$

Table	Header Row
1191	$R^2 = CF_3, Z = S, R^3 = 3-Ethenyl$
1192	$R^2 = CF_3, Z = S, R^3 = 3-SO_2Me$
1193	$R^2 = CF_3, Z = S, R^3 = 3-OAc$
1194	$R^2 = CF_3, Z = S, R^3 = 3-c-Pr$
1195	$R^2 = CF_3, Z = S, R^3 = 3-i-Pr$
1196	$R^2 = CF_3, Z = S, R^3 = 3-Ph$
1197	$R^2 = CF_3, Z = O, R^3 = 4-F$
1198	$R^2 = CF_3, Z = O, R^3 = 4-Cl$
1199	$R^2 = CF_3, Z = O, R^3 = 4-Br$
1200	$R^2 = CF_3, Z = O, R^3 = 4-I$
1201	$R^2 = CF_3, Z = O, R^3 = 4-CN$
1202	$R^2 = CF_3, Z = O, R^3 = 4-NO_2$
1203	$R^2 = CF_3, Z = O, R^3 = 4-OMe$
1204	$R^2 = CF_3, Z = O, R^3 = 4-OCF_3$
1205	$R^2 = CF_3, Z = O, R^3 = 4-CF_3$
1206	$R^2 = CF_3, Z = O, R^3 = 4-CHF_2$
1207	$R^2 = CF_3, Z = O, R^3 = 4-CH_2F$
1208	$R^2 = CF_3, Z = O, R^3 = 4-CHO$
1209	$R^2 = CF_3, Z = O, R^3 = 4-Me$
1210	$R^2 = CF_3, Z = O, R^3 = 4-Et$
1211	$R^2 = CF_3, Z = O, R^3 = 4-Ethynyl$
1212	$R^2 = CF_3, Z = O, R^3 = 4-Ethenyl$
1213	$R^2 = CF_3, Z = O, R^3 = 4-SO_2Me$
1214	$R^2 = CF_3, Z = O, R^3 = 4-OAc$
1215	$R^2 = CF_3, Z = O, R^3 = 4-c-Pr$
1216	$R^2 = CF_3, Z = O, R^3 = 4-i-Pr$
1217	$R^2 = CF_3, Z = O, R^3 = 4-Ph$
1218	$R^2 = CF_3, Z = O, R^3 = 5-F$
1219	$R^2 = CF_3, Z = O, R^3 = 5-Cl$
1220	$R^2 = CF_3, Z = O, R^3 = 5-Br$
1221	$R^2 = CF_3, Z = O, R^3 = 5-I$
1222	$R^2 = CF_3, Z = O, R^3 = 5-CN$
1223	$R^2 = CF_3, Z = O, R^3 = 5-NO_2$
1224	$R^2 = CF_3, Z = O, R^3 = 5-OMe$
1225	$R^2 = CF_3, Z = O, R^3 = 5-OCF_3$
1226	$R^2 = CF_3, Z = O, R^3 = 5-CF_3$
1227	$R^2 = CF_3, Z = O, R^3 = 5-CHF_2$

Table	Header Row
1228	$R^2 = CF_3, Z = O, R^3 = 5-CH_2F$
1229	$R^2 = CF_3, Z = O, R^3 = 5-CHO$
1230	$R^2 = CF_3, Z = O, R^3 = 5-Me$
1231	$R^2 = CF_3, Z = O, R^3 = 5-Et$
1232	$R^2 = CF_3, Z = O, R^3 = 5-Ethynyl$
1233	$R^2 = CF_3, Z = O, R^3 = 5-Ethenyl$
1234	$R^2 = CF_3, Z = O, R^3 = 5-SO_2Me$
1235	$R^2 = CF_3, Z = O, R^3 = 5-OAc$
1236	$R^2 = CF_3, Z = O, R^3 = 5-c-Pr$
1237	$R^2 = CF_3, Z = O, R^3 = 5-i-Pr$
1238	$R^2 = CF_3, Z = O, R^3 = 5-Ph$
1239	$R^2 = CF_3, Z = O, R^3 = 6-F$
1240	$R^2 = CF_3, Z = O, R^3 = 6-Cl$
1241	$R^2 = CF_3, Z = O, R^3 = 6-Br$
1242	$R^2 = CF_3, Z = O, R^3 = 6-I$
1243	$R^2 = CF_3, Z = O, R^3 = 6-CN$
1244	$R^2 = CF_3, Z = O, R^3 = 6-NO_2$
1245	$R^2 = CF_3, Z = O, R^3 = 6-OMe$
1246	$R^2 = CF_3, Z = O, R^3 = 6-OCF_3$
1247	$R^2 = CF_3, Z = O, R^3 = 6-CF_3$
1248	$R^2 = CF_3, Z = O, R^3 = 6-CHF_2$
1249	$R^2 = CF_3, Z = O, R^3 = 6-CH_2F$
1250	$R^2 = CF_3, Z = O, R^3 = 6-CHO$
1251	$R^2 = CF_3, Z = O, R^3 = 6-Me$
1252	$R^2 = CF_3, Z = O, R^3 = 6-Et$
1253	$R^2 = CF_3, Z = O, R^3 = 6-Ethynyl$
1254	$R^2 = CF_3, Z = O, R^3 = 6-Ethenyl$
1255	$R^2 = CF_3, Z = O, R^3 = 6-SO_2Me$
1256	$R^2 = CF_3, Z = O, R^3 = 6-OAc$
1257	$R^2 = CF_3, Z = O, R^3 = 6-c-Pr$
1258	$R^2 = CF_3, Z = O, R^3 = 6-i-Pr$
1259	$R^2 = CF_3, Z = O, R^3 = 6-Ph$
1260	$R^2 = CF_3, Z = O, R^3 = 3,4-di-F$
1261	$R^2 = CF_3, Z = O, R^3 = 3,5-di-F$
1262	$R^2 = CF_3, Z = O, R^3 = 3,6-di-F$
1263	$R^2 = CF_3, Z = O, R^3 = 4,5-di-F$
1264	$R^2 = CF_3, Z = O, R^3 = 3,4-di-Cl$

Table	Header Row
1265	$R^2 = CF_3, Z = O, R^3 = 3,5-di-Cl$
1266	$R^2 = CF_3, Z = O, R^3 = 3,6-di-Cl$
1267	$R^2 = CF_3, Z = O, R^3 = 4,5-di-Cl$
1268	$R^2 = CF_3, Z = O, R^3 = 3,4-di-Br$
1269	$R^2 = CF_3, Z = O, R^3 = 3,5-di-Br$
1270	$R^2 = CF_3, Z = O, R^3 = 3,6-di-Br$
1271	$R^2 = CF_3, Z = O, R^3 = 4,5-di-Br$
1272	$R^2 = CF_3, Z = O, R^3 = 3,4-di-CN$
1273	$R^2 = CF_3, Z = O, R^3 = 3,5-di-CN$
1274	$R^2 = CF_3, Z = O, R^3 = 3,6-di-CN$
1275	$R^2 = CF_3, Z = O, R^3 = 4,5-di-CN$
1276	$R^2 = CF_3, Z = O, R^3 = 3,4-di-Me$
1277	$R^2 = CF_3, Z = O, R^3 = 3,5-di-Me$
1278	$R^2 = CF_3, Z = O, R^3 = 3,6-di-Me$
1279	$R^2 = CF_3, Z = O, R^3 = 4,5-di-Me$
1280	$R^2 = CF_3, Z = O, R^3 = 3,4-di-OMe$
1281	$R^2 = CF_3, Z = O, R^3 = 3,5-di-OMe$
1282	$R^2 = CF_3, Z = O, R^3 = 3,6-di-OMe$
1283	$R^2 = CF_3, Z = O, R^3 = 4,5-di-OMe$
1284	$R^2 = CF_3, Z = O, R^3 = 3,4-di-CF_3$
1285	$R^2 = CF_3, Z = O, R^3 = 3,5-di-CF_3$
1286	$R^2 = CF_3, Z = O, R^3 = 3,6-di-CF_3$
1287	$R^2 = CF_3, Z = O, R^3 = 4,5-di-CF_3$
1288	$R^2 = CF_3, Z = O, R^3 = 3-CN, 4-Me$
1289	$R^2 = CF_3, Z = O, R^3 = 3-CN, 4-F$
1290	$R^2 = CF_3, Z = O, R^3 = 3-CN, 4-Br$
1291	$R^2 = CF_3, Z = O, R^3 = 3-CN, 4-OMe$
1292	$R^2 = CF_3, Z = O, R^3 = 3-CN, 4-CF_3$
1293	$R^2 = CF_3, Z = O, R^3 = 3-CN, 6-Me$
1294	$R^2 = CF_3, Z = O, R^3 = 3-CN, 6-F$
1295	$R^2 = CF_3, Z = O, R^3 = 3-CN, 6-Br$
1296	$R^2 = CF_3, Z = O, R^3 = 3-CN, 6-OMe$
1297	$R^2 = CF_3, Z = O, R^3 = 3-CN, 6-CF_3$
1298	$R^2 = CHF_2, Z = O, R^3 = H (m = 0)$
1299	$R^2 = CHF_2, Z = O, R^3 = 3-F$
1300	$R^2 = CHF_2, Z = O, R^3 = 3-Cl$
1301	$R^2 = CHF_2, Z = O, R^3 = 3-Br$

Table	Header Row
1302	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-I}$
1303	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-CN}$
1304	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-NO}_2$
1305	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-OMe}$
1306	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-OCF}_3$
1307	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-CF}_3$
1308	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-CHF}_2$
1309	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-CH}_2\text{F}$
1310	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-CHO}$
1311	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-Me}$
1312	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-Et}$
1313	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-Ethyneyl}$
1314	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-Ethenyl}$
1315	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-SO}_2\text{Me}$
1316	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-OAc}$
1317	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-}i\text{-Pr}$
1318	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-}i\text{-Pr}$
1319	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 3\text{-Ph}$
1320	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-F}$
1321	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-Cl}$
1322	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-Br}$
1323	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-I}$
1324	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-CN}$
1325	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-NO}_2$
1326	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-OMe}$
1327	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-OCF}_3$
1328	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-CHF}_2$
1329	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-CH}_2\text{F}$
1330	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-CHO}$
1331	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-Me}$
1332	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-Et}$
1333	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-Ethyneyl}$
1334	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-Ethenyl}$
1335	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-SO}_2\text{Me}$
1336	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-OAc}$
1337	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-}i\text{-Pr}$
1338	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-}i\text{-Pr}$

Table	Header Row
1339	$R^2 = \text{CHF}_2, Z = \text{S}, R^3 = 3\text{-Ph}$
1340	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-F}$
1341	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-Cl}$
1342	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-Br}$
1343	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-I}$
1344	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-CN}$
1345	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-NO}_2$
1346	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-OMe}$
1347	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-OCF}_3$
1348	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-CF}_3$
1349	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-CHF}_2$
1350	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-CH}_2\text{F}$
1351	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-CHO}$
1352	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-Me}$
1353	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-Et}$
1354	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-Ethyneyl}$
1355	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-Ethenyl}$
1356	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-SO}_2\text{Me}$
1357	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-OAc}$
1358	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-}i\text{-Pr}$
1359	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-}i\text{-Pr}$
1360	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 4\text{-Ph}$
1361	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-F}$
1362	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-Cl}$
1363	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-Br}$
1364	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-I}$
1365	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-CN}$
1366	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-NO}_2$
1367	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-OMe}$
1368	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-OCF}_3$
1369	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-CF}_3$
1370	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-CHF}_2$
1371	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-CH}_2\text{F}$
1372	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-CHO}$
1373	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-Me}$
1374	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-Et}$
1375	$R^2 = \text{CHF}_2, Z = \text{O}, R^3 = 5\text{-Ethyneyl}$

Table	Header Row
1376	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 5\text{-Ethenyl}$
1377	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 5\text{-SO}_2\text{Me}$
1378	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 5\text{-OAc}$
1379	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 5\text{-}i\text{-Pr}$
1380	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 5\text{-}i\text{-Pr}$
1381	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 5\text{-Ph}$
1382	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-F}$
1383	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-Cl}$
1384	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-Br}$
1385	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-I}$
1386	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-CN}$
1387	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-NO}_2$
1388	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-OMe}$
1389	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-OCF}_3$
1390	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-CF}_3$
1391	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-CHF}_2$
1392	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-CH}_2\text{F}$
1393	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-CHO}$
1394	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-Me}$
1395	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-Et}$
1396	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-Ethinyl}$
1397	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-Ethenyl}$
1398	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-SO}_2\text{Me}$
1399	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-OAc}$
1400	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-}i\text{-Pr}$
1401	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-}i\text{-Pr}$
1402	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 6\text{-Ph}$
1403	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,4\text{-di-F}$
1404	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,5\text{-di-F}$
1405	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,6\text{-di-F}$
1406	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 4,5\text{-di-F}$
1407	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,4\text{-di-Cl}$
1408	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,5\text{-di-Cl}$
1409	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,6\text{-di-Cl}$
1410	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 4,5\text{-di-Cl}$
1411	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,4\text{-di-Br}$
1412	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,5\text{-di-Br}$

Table	Header Row
1413	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,6\text{-di-Br}$
1414	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 4,5\text{-di-Br}$
1415	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,4\text{-di-CN}$
1416	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,5\text{-di-CN}$
1417	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,6\text{-di-CN}$
1418	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 4,5\text{-di-CN}$
1419	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,4\text{-di-Me}$
1420	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,5\text{-di-Me}$
1421	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,6\text{-di-Me}$
1422	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 4,5\text{-di-Me}$
1423	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,4\text{-di-OMe}$
1424	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,5\text{-di-OMe}$
1425	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,6\text{-di-OMe}$
1426	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 4,5\text{-di-OMe}$
1427	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,4\text{-di-CF}_3$
1428	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,5\text{-di-CF}_3$
1429	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3,6\text{-di-CF}_3$
1430	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 4,5\text{-di-CF}_3$
1431	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 4\text{-Me}$
1432	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 4\text{-F}$
1433	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 4\text{-Br}$
1434	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 4\text{-OMe}$
1435	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 4\text{-CF}_3$
1436	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 6\text{-Me}$
1437	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 6\text{-F}$
1438	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 6\text{-Br}$
1439	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 6\text{-OMe}$
1440	$R^2 = \text{CHF}_2$, $Z = \text{O}$, $R^3 = 3\text{-CN, } 6\text{-CF}_3$
1441	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = \text{H (m = 0)}$
1442	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = 3\text{-F}$
1443	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = 3\text{-Cl}$
1444	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = 3\text{-Br}$
1445	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = 3\text{-I}$
1446	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = 3\text{-CN}$
1447	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = 3\text{-NO}_2$
1448	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = 3\text{-OMe}$
1449	$R^2 = \text{SO}_2\text{Me}$, $Z = \text{O}$, $R^3 = 3\text{-OCF}_3$

Table	Header Row
1450	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CF}_3$
1451	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CHF}_2$
1452	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CH}_2\text{F}$
1453	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CHO}$
1454	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-Me}$
1455	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-Et}$
1456	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-Ethyneyl}$
1457	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-Ethenyl}$
1458	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-SO}_2\text{Me}$
1459	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-OAc}$
1460	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-c-Pr}$
1461	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-i-Pr}$
1462	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-Ph}$
1463	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-F}$
1464	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-Cl}$
1465	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-Br}$
1466	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-I}$
1467	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-CN}$
1468	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-NO}_2$
1469	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-OMe}$
1470	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-OCF}_3$
1471	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-CF}_3$
1472	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-CHF}_2$
1473	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-CH}_2\text{F}$
1474	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-CHO}$
1475	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-Me}$
1476	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-Et}$
1477	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-Ethyneyl}$
1478	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-Ethenyl}$
1479	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-SO}_2\text{Me}$
1480	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-OAc}$
1481	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-c-Pr}$
1482	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-i-Pr}$
1483	$R^2 = \text{SO}_2\text{Me}, Z = \text{S}, R^3 = 3\text{-Ph}$
1484	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-F}$
1485	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-Cl}$
1486	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-Br}$

Table	Header Row
1487	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-I}$
1488	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-CN}$
1489	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-NO}_2$
1490	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-OMe}$
1491	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-OCF}_3$
1492	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-CF}_3$
1493	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-CHF}_2$
1494	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-CH}_2\text{F}$
1495	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-CHO}$
1496	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-Me}$
1497	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-Et}$
1498	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-Ethyneyl}$
1499	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-Ethenyl}$
1500	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-SO}_2\text{Me}$
1501	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-OAc}$
1502	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-c-Pr}$
1503	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-i-Pr}$
1504	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4\text{-Ph}$
1505	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-F}$
1506	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-Cl}$
1507	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-Br}$
1508	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-I}$
1509	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-CN}$
1510	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-NO}_2$
1511	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-OMe}$
1512	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-OCF}_3$
1513	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-CF}_3$
1514	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-CHF}_2$
1515	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-CH}_2\text{F}$
1516	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-CHO}$
1517	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-Me}$
1518	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-Et}$
1519	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-Ethyneyl}$
1520	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-Ethenyl}$
1521	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-SO}_2\text{Me}$
1522	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-OAc}$
1523	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-c-Pr}$

Table	Header Row
1524	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-}i\text{-Pr}$
1525	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 5\text{-Ph}$
1526	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-F}$
1527	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-Cl}$
1528	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-Br}$
1529	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-I}$
1530	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-CN}$
1531	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-NO}_2$
1532	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-OMe}$
1533	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-OCF}_3$
1534	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-CF}_3$
1535	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-CHF}_2$
1536	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-CH}_2\text{F}$
1537	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-CHO}$
1538	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-Me}$
1539	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-Et}$
1540	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-Ethynyl}$
1541	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-Ethenyl}$
1542	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-SO}_2\text{Me}$
1543	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-OAc}$
1544	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-}c\text{-Pr}$
1545	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-}i\text{-Pr}$
1546	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 6\text{-Ph}$
1547	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-F}$
1548	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-F}$
1549	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-F}$
1550	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-F}$
1551	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-Cl}$
1552	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-Cl}$
1553	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-Cl}$
1554	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-Cl}$

Table	Header Row
1555	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-Br}$
1556	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-Br}$
1557	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-Br}$
1558	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-Br}$
1559	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-CN}$
1560	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-CN}$
1561	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-CN}$
1562	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-CN}$
1563	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-Me}$
1564	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-Me}$
1565	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-Me}$
1566	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-Me}$
1567	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-OMe}$
1568	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-OMe}$
1569	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-OMe}$
1570	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-OMe}$
1571	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,4\text{-di-CF}_3$
1572	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,5\text{-di-CF}_3$
1573	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3,6\text{-di-CF}_3$
1574	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 4,5\text{-di-CF}_3$
1575	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Me}$
1576	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-F}$
1577	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-Br}$
1578	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-OMe}$
1579	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 4\text{-CF}_3$
1580	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Me}$
1581	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-F}$
1582	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-Br}$
1583	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-OMe}$
1584	$R^2 = \text{SO}_2\text{Me}, Z = \text{O}, R^3 = 3\text{-CN}, 6\text{-CF}_3$

A compound of this invention will generally be used as a herbicidal active ingredient in a composition, i.e. formulation, with at least one additional component selected from the group consisting of surfactants, solid diluents and liquid diluents, which serves as a carrier. The formulation or composition ingredients are selected to be consistent with the physical properties of the active ingredient, mode of application and environmental factors such as soil type, moisture and temperature.

Useful formulations include both liquid and solid compositions. Liquid compositions include solutions (including emulsifiable concentrates), suspensions, emulsions (including microemulsions, oil-in-water emulsions, flowable concentrates and/or suspoemulsions) and the like, which optionally can be thickened into gels. The general types of aqueous liquid compositions are soluble concentrate, suspension concentrate, capsule suspension, concentrated emulsion, microemulsion, oil-in-water emulsion, flowable concentrate and suspo-emulsion. The general types of nonaqueous liquid compositions are emulsifiable concentrate, microemulsifiable concentrate, dispersible concentrate and oil dispersion.

The general types of solid compositions are dusts, powders, granules, pellets, prills, pastilles, tablets, filled films (including seed coatings) and the like, which can be water-dispersible ("wetttable") or water-soluble. Films and coatings formed from film-forming solutions or flowable suspensions are particularly useful for seed treatment. Active ingredient can be (micro)encapsulated and further formed into a suspension or solid formulation; alternatively the entire formulation of active ingredient can be encapsulated (or "overcoated"). Encapsulation can control or delay release of the active ingredient. An emulsifiable granule combines the advantages of both an emulsifiable concentrate formulation and a dry granular formulation. High-strength compositions are primarily used as intermediates for further formulation.

Sprayable formulations are typically extended in a suitable medium before spraying. Such liquid and solid formulations are formulated to be readily diluted in the spray medium, usually water, but occasionally another suitable medium like an aromatic or paraffinic hydrocarbon or vegetable oil. Spray volumes can range from about one to several thousand liters per hectare, but more typically are in the range from about ten to several hundred liters per hectare. Sprayable formulations can be tank mixed with water or another suitable medium for foliar treatment by aerial or ground application, or for application to the growing medium of the plant. Liquid and dry formulations can be metered directly into drip irrigation systems or metered into the furrow during planting.

The formulations will typically contain effective amounts of active ingredient, diluent and surfactant within the following approximate ranges which add up to 100 percent by weight.

	Weight Percent		
	<u>Active Ingredient</u>	<u>Diluent</u>	<u>Surfactant</u>
Water-Dispersible and Water-soluble Granules, Tablets and Powders	0.001–90	0–99.999	0–15
Oil Dispersions, Suspensions, Emulsions, Solutions (including Emulsifiable Concentrates)	1–50	40–99	0–50
Dusts	1–25	70–99	0–5
Granules and Pellets	0.001–99	5–99.999	0–15
High Strength Compositions	90–99	0–10	0–2

Solid diluents include, for example, clays such as bentonite, montmorillonite, attapulgite and kaolin, gypsum, cellulose, titanium dioxide, zinc oxide, starch, dextrin, sugars (e.g., lactose, sucrose), silica, talc, mica, diatomaceous earth, urea, calcium carbonate, sodium carbonate and bicarbonate, and sodium sulfate. Typical solid diluents are described in Watkins et al., *Handbook of Insecticide Dust Diluents and Carriers*, 2nd Ed., Dorland Books, Caldwell, New Jersey.

Liquid diluents include, for example, water, *N,N*-dimethylalkanamides (e.g., *N,N*-dimethylformamide), limonene, dimethyl sulfoxide, *N*-alkylpyrrolidones (e.g., *N*-methylpyrrolidinone), alkyl phosphates (e.g., triethyl phosphate), ethylene glycol, triethylene glycol, propylene glycol, dipropylene glycol, polypropylene glycol, propylene carbonate, butylene carbonate, paraffins (e.g., white mineral oils, normal paraffins, isoparaffins), alkylbenzenes, alkylnaphthalenes, glycerine, glycerol triacetate, sorbitol, aromatic hydrocarbons, dearomatized aliphatics, alkylbenzenes, alkylnaphthalenes, ketones such as cyclohexanone, 2-heptanone, isophorone and 4-hydroxy-4-methyl-2-pentanone, acetates such as isoamyl acetate, hexyl acetate, heptyl acetate, octyl acetate, nonyl acetate, tridecyl acetate and isobornyl acetate, other esters such as alkylated lactate esters, dibasic esters, alkyl and aryl benzoates and γ -butyrolactone, and alcohols, which can be linear, branched, saturated or unsaturated, such as methanol, ethanol, *n*-propanol, isopropyl alcohol, *n*-butanol, isobutyl alcohol, *n*-hexanol, 2-ethylhexanol, *n*-octanol, decanol, isodecyl alcohol, isooctadecanol, cetyl alcohol, lauryl alcohol, tridecyl alcohol, oleyl alcohol, cyclohexanol, tetrahydrofurfuryl alcohol, diacetone alcohol, cresol and benzyl alcohol. Liquid diluents also include glycerol esters of saturated and unsaturated fatty acids (typically C₆–C₂₂), such as plant seed and fruit oils (e.g., oils of olive, castor, linseed, sesame, corn (maize), peanut, sunflower, grapeseed, safflower, cottonseed, soybean, rapeseed, coconut and palm kernel), animal-sourced fats (e.g., beef tallow, pork tallow, lard, cod liver oil, fish oil), and mixtures thereof. Liquid diluents also include alkylated fatty acids (e.g.,

methyated, ethyated, butyated) wherein the fatty acids may be obtained by hydrolysis of glycerol esters from plant and animal sources, and can be purified by distillation. Typical liquid diluents are described in Marsden, *Solvents Guide*, 2nd Ed., Interscience, New York, 1950.

5 The solid and liquid compositions of the present invention often include one or more surfactants. When added to a liquid, surfactants (also known as “surface-active agents”) generally modify, most often reduce, the surface tension of the liquid. Depending on the nature of the hydrophilic and lipophilic groups in a surfactant molecule, surfactants can be useful as wetting agents, dispersants, emulsifiers or defoaming agents.

10 Surfactants can be classified as nonionic, anionic or cationic. Nonionic surfactants useful for the present compositions include, but are not limited to: alcohol alkoxylates such as alcohol alkoxylates based on natural and synthetic alcohols (which may be branched or linear) and prepared from the alcohols and ethylene oxide, propylene oxide, butylene oxide or mixtures thereof; amine ethoxylates, alkanolamides and ethoxylated alkanolamides;
15 alkoxylated triglycerides such as ethoxylated soybean, castor and rapeseed oils; alkylphenol alkoxylates such as octylphenol ethoxylates, nonylphenol ethoxylates, dinonyl phenol ethoxylates and dodecyl phenol ethoxylates (prepared from the phenols and ethylene oxide, propylene oxide, butylene oxide or mixtures thereof); block polymers prepared from ethylene oxide or propylene oxide and reverse block polymers where the terminal blocks are
20 prepared from propylene oxide; ethoxylated fatty acids; ethoxylated fatty esters and oils; ethoxylated methyl esters; ethoxylated tristyrylphenol (including those prepared from ethylene oxide, propylene oxide, butylene oxide or mixtures thereof); fatty acid esters, glycerol esters, lanolin-based derivatives, polyethoxylate esters such as polyethoxylated sorbitan fatty acid esters, polyethoxylated sorbitol fatty acid esters and polyethoxylated
25 glycerol fatty acid esters; other sorbitan derivatives such as sorbitan esters; polymeric surfactants such as random copolymers, block copolymers, alkyd peg (polyethylene glycol) resins, graft or comb polymers and star polymers; polyethylene glycols (pegs); polyethylene glycol fatty acid esters; silicone-based surfactants; and sugar-derivatives such as sucrose esters, alkyl polyglycosides and alkyl polysaccharides.

30 Useful anionic surfactants include, but are not limited to: alkylaryl sulfonic acids and their salts; carboxylated alcohol or alkylphenol ethoxylates; diphenyl sulfonate derivatives; lignin and lignin derivatives such as lignosulfonates; maleic or succinic acids or their anhydrides; olefin sulfonates; phosphate esters such as phosphate esters of alcohol alkoxylates, phosphate esters of alkylphenol alkoxylates and phosphate esters of styryl
35 phenol ethoxylates; protein-based surfactants; sarcosine derivatives; styryl phenol ether sulfate; sulfates and sulfonates of oils and fatty acids; sulfates and sulfonates of ethoxylated alkylphenols; sulfates of alcohols; sulfates of ethoxylated alcohols; sulfonates of amines and amides such as *N,N*-alkyltaurates; sulfonates of benzene, cumene, toluene, xylene, and

dodecyl and tridecylbenzenes; sulfonates of condensed naphthalenes; sulfonates of naphthalene and alkyl naphthalene; sulfonates of fractionated petroleum; sulfosuccinamates; and sulfosuccinates and their derivatives such as dialkyl sulfosuccinate salts.

Useful cationic surfactants include, but are not limited to: amides and ethoxylated
5 amides; amines such as *N*-alkyl propanediamines, tripropylenetriamines and dipropylenetetramines, and ethoxylated amines, ethoxylated diamines and propoxylated amines (prepared from the amines and ethylene oxide, propylene oxide, butylene oxide or mixtures thereof); amine salts such as amine acetates and diamine salts; quaternary ammonium salts such as quaternary salts, ethoxylated quaternary salts and diquaternary salts;
10 and amine oxides such as alkyldimethylamine oxides and bis-(2-hydroxyethyl)-alkylamine oxides.

Also useful for the present compositions are mixtures of nonionic and anionic surfactants or mixtures of nonionic and cationic surfactants. Nonionic, anionic and cationic surfactants and their recommended uses are disclosed in a variety of published references
15 including *McCutcheon's Emulsifiers and Detergents*, annual American and International Editions published by McCutcheon's Division, The Manufacturing Confectioner Publishing Co.; Sisely and Wood, *Encyclopedia of Surface Active Agents*, Chemical Publ. Co., Inc., New York, 1964; and A. S. Davidson and B. Milwidsky, *Synthetic Detergents*, Seventh Edition, John Wiley and Sons, New York, 1987.

20 Compositions of this invention may also contain formulation auxiliaries and additives, known to those skilled in the art as formulation aids (some of which may be considered to also function as solid diluents, liquid diluents or surfactants). Such formulation auxiliaries and additives may control: pH (buffers), foaming during processing (antifoams such polyorganosiloxanes), sedimentation of active ingredients (suspending agents), viscosity
25 (thixotropic thickeners), in-container microbial growth (antimicrobials), product freezing (antifreezes), color (dyes/pigment dispersions), wash-off (film formers or stickers), evaporation (evaporation retardants), and other formulation attributes. Film formers include, for example, polyvinyl acetates, polyvinyl acetate copolymers, polyvinylpyrrolidone-vinyl acetate copolymer, polyvinyl alcohols, polyvinyl alcohol copolymers and waxes. Examples
30 of formulation auxiliaries and additives include those listed in *McCutcheon's Volume 2: Functional Materials*, annual International and North American editions published by McCutcheon's Division, The Manufacturing Confectioner Publishing Co.; and PCT Publication WO 03/024222.

The compound of Formula 1 and any other active ingredients are typically
35 incorporated into the present compositions by dissolving the active ingredient in a solvent or by grinding in a liquid or dry diluent. Solutions, including emulsifiable concentrates, can be prepared by simply mixing the ingredients. If the solvent of a liquid composition intended for use as an emulsifiable concentrate is water-immiscible, an emulsifier is typically added to

emulsify the active-containing solvent upon dilution with water. Active ingredient slurries, with particle diameters of up to 2,000 μm can be wet milled using media mills to obtain particles with average diameters below 3 μm . Aqueous slurries can be made into finished suspension concentrates (see, for example, U.S. 3,060,084) or further processed by spray
5 drying to form water-dispersible granules. Dry formulations usually require dry milling processes, which produce average particle diameters in the 2 to 10 μm range. Dusts and powders can be prepared by blending and usually grinding (such as with a hammer mill or fluid-energy mill). Granules and pellets can be prepared by spraying the active material upon preformed granular carriers or by agglomeration techniques. See Browning,
10 "Agglomeration", *Chemical Engineering*, December 4, 1967, pp 147–48, *Perry's Chemical Engineer's Handbook*, 4th Ed., McGraw-Hill, New York, 1963, pages 8–57 and following, and WO 91/13546. Pellets can be prepared as described in U.S. 4,172,714. Water-dispersible and water-soluble granules can be prepared as taught in U.S. 4,144,050, U.S. 3,920,442 and DE 3,246,493. Tablets can be prepared as taught in U.S. 5,180,587, U.S.
15 5,232,701 and U.S. 5,208,030. Films can be prepared as taught in GB 2,095,558 and U.S. 3,299,566.

For further information regarding the art of formulation, see T. S. Woods, "The Formulator's Toolbox – Product Forms for Modern Agriculture" in *Pesticide Chemistry and Bioscience, The Food–Environment Challenge*, T. Brooks and T. R. Roberts, Eds.,
20 Proceedings of the 9th International Congress on Pesticide Chemistry, The Royal Society of Chemistry, Cambridge, 1999, pp. 120–133. See also U.S. 3,235,361, Col. 6, line 16 through Col. 7, line 19 and Examples 10–41; U.S. 3,309,192, Col. 5, line 43 through Col. 7, line 62 and Examples 8, 12, 15, 39, 41, 52, 53, 58, 132, 138–140, 162–164, 166, 167 and 169–182; U.S. 2,891,855, Col. 3, line 66 through Col. 5, line 17 and Examples 1–4; Klingman, *Weed*
25 *Control as a Science*, John Wiley and Sons, Inc., New York, 1961, pp 81–96; Hance et al., *Weed Control Handbook*, 8th Ed., Blackwell Scientific Publications, Oxford, 1989; and *Developments in formulation technology*, PJB Publications, Richmond, UK, 2000.

In the following Examples, all percentages are by weight and all formulations are prepared in conventional ways. Compound numbers refer to compounds in Index Table A.
30 Without further elaboration, it is believed that one skilled in the art using the preceding description can utilize the present invention to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative, and not limiting of the disclosure in any way whatsoever. Percentages are by weight except where otherwise indicated.

Example AHigh Strength Concentrate

Compound 1	98.5%
silica aerogel	0.5%
synthetic amorphous fine silica	1.0%

Example BWettable Powder

Compound 12	65.0%
dodecylphenol polyethylene glycol ether	2.0%
sodium ligninsulfonate	4.0%
sodium silicoaluminate	6.0%
montmorillonite (calcined)	23.0%

Example CGranule

Compound 15	10.0%
attapulgit granules (low volatile matter, 0.71/0.30 mm; U.S.S. No. 25–50 sieves)	90.0%

Example DExtruded Pellet

Compound 21	25.0%
anhydrous sodium sulfate	10.0%
crude calcium ligninsulfonate	5.0%
sodium alkyl naphthalenesulfonate	1.0%
calcium/magnesium bentonite	59.0%

5

Example EEmulsifiable Concentrate

Compound 23	10.0%
polyoxyethylene sorbitol hexoleate	20.0%
C ₆ –C ₁₀ fatty acid methyl ester	70.0%

Example FMicroemulsion

Compound 24	5.0%
polyvinylpyrrolidone-vinyl acetate copolymer	30.0%
alkylpolyglycoside	30.0%
glyceryl monooleate	15.0%
water	20.0%

Example GSuspension Concentrate

Compound 27	35%
butyl polyoxyethylene/polypropylene block copolymer	4.0%
stearic acid/polyethylene glycol copolymer	1.0%
styrene acrylic polymer	1.0%
xanthan gum	0.1%
propylene glycol	5.0%
silicone based defoamer	0.1%
1,2-benzisothiazolin-3-one	0.1%
water	53.7%

Example HEmulsion in Water

Compound 32	10.0%
butyl polyoxyethylene/polypropylene block copolymer	4.0%
stearic acid/polyethylene glycol copolymer	1.0%
styrene acrylic polymer	1.0%
xanthan gum	0.1%
propylene glycol	5.0%
silicone based defoamer	0.1%
1,2-benzisothiazolin-3-one	0.1%
aromatic petroleum based hydrocarbon	20.0
water	58.7%

Example IOil Dispersion

Compound 42	25%
polyoxyethylene sorbitol hexaoleate	15%
organically modified bentonite clay	2.5%
fatty acid methyl ester	57.5%

Example JSuspoemulsion

Compound 1	10.0%
imidacloprid	5.0%
butyl polyoxyethylene/polypropylene block copolymer	4.0%
stearic acid/polyethylene glycol copolymer	1.0%
styrene acrylic polymer	1.0%
xanthan gum	0.1%
propylene glycol	5.0%

silicone based defoamer	0.1%
1,2-benzisothiazolin-3-one	0.1%
aromatic petroleum based hydrocarbon	20.0%
water	53.7%

Test results indicate that the compounds of the present invention are highly active preemergent and/or postemergent herbicides and/or plant growth regulants. The compounds of the invention generally show highest activity for postemergence weed control (i.e. applied after weed seedlings emerge from the soil) and preemergence weed control (i.e. applied before weed seedlings emerge from the soil). Many of them have utility for broad-spectrum pre- and/or postemergence weed control in areas where complete control of all vegetation is desired such as around fuel storage tanks, industrial storage areas, parking lots, drive-in theaters, air fields, river banks, irrigation and other waterways, around billboards and highway and railroad structures. Many of the compounds of this invention, by virtue of selective metabolism in crops versus weeds, or by selective activity at the locus of physiological inhibition in crops and weeds, or by selective placement on or within the environment of a mixture of crops and weeds, are useful for the selective control of grass and broadleaf weeds within a crop/weed mixture. One skilled in the art will recognize that the preferred combination of these selectivity factors within a compound or group of compounds can readily be determined by performing routine biological and/or biochemical assays. Compounds of this invention may show tolerance to important agronomic crops including, but is not limited to, alfalfa, barley, cotton, wheat, rape, sugar beets, corn (maize), sorghum, soybeans, rice, oats, peanuts, vegetables, tomato, potato, perennial plantation crops including coffee, cocoa, oil palm, rubber, sugarcane, citrus, grapes, fruit trees, nut trees, banana, plantain, pineapple, hops, tea and forests such as eucalyptus and conifers (e.g., loblolly pine), and turf species (e.g., Kentucky bluegrass, St. Augustine grass, Kentucky fescue and Bermuda grass). Compounds of this invention can be used in crops genetically transformed or bred to incorporate resistance to herbicides, express proteins toxic to invertebrate pests (such as *Bacillus thuringiensis* toxin), and/or express other useful traits. Those skilled in the art will appreciate that not all compounds are equally effective against all weeds. Alternatively, the subject compounds are useful to modify plant growth.

As the compounds of the invention have both preemergent and postemergent herbicidal activity, to control undesired vegetation by killing or injuring the vegetation or reducing its growth, the compounds can be usefully applied by a variety of methods involving contacting a herbicidally effective amount of a compound of the invention, or a composition comprising said compound and at least one of a surfactant, a solid diluent or a liquid diluent, to the foliage or other part of the undesired vegetation or to the environment

of the undesired vegetation such as the soil or water in which the undesired vegetation is growing or which surrounds the seed or other propagule of the undesired vegetation.

A herbicidally effective amount of the compounds of this invention is determined by a number of factors. These factors include: formulation selected, method of application, amount and type of vegetation present, growing conditions, etc. In general, a herbicidally effective amount of compounds of this invention is about 0.001 to 20 kg/ha with a preferred range of about 0.004 to 1 kg/ha. One skilled in the art can easily determine the herbicidally effective amount necessary for the desired level of weed control.

Compounds of the invention are useful in treating all plants and plant parts. Plant varieties and cultivars can be obtained by conventional propagation and breeding methods or by genetic engineering methods. Genetically modified plants (transgenic plants) are those in which a heterologous gene (transgene) has been stably integrated into the plant's genome. A transgene that is defined by its particular location in the plant genome is called a transformation or transgenic event.

Genetically modified plant cultivars which can be treated according to the invention include those that are resistant against one or more biotic stresses (pests such as nematodes, insects, mites, fungi, etc.) or abiotic stresses (drought, cold temperature, soil salinity, etc.), or that contain other desirable characteristics. Plants can be genetically modified to exhibit traits of, for example, herbicide tolerance, insect-resistance, modified oil profiles or drought tolerance. Useful genetically modified plants containing single gene transformation events or combinations of transformation events are listed in Exhibit C. Additional information for the genetic modifications listed in Exhibit C can be obtained from publicly available databases maintained, for example, by the U.S. Department of Agriculture.

The following abbreviations, 1 through 37, are used in Exhibit C for traits. A “-” means the entry is not available.

Trait	Description	Trait	Description	Trait	Description
1	Glyphosate tolerance	15	Cold tolerance	27	High tryptophan
2	High lauric acid oil	16	Imidazolinone herb. tol.	28	Erect leaves semidwarf
3	Glufosinate tolerance	17	Modified alpha-amylase	29	Semidwarf
4	Phytate breakdown	18	Pollination control	30	Low iron tolerance
5	Oxynil tolerance	19	2,4-D tolerance	31	Modified oil/fatty acid
6	Disease resistance	20	Increased lysine	32	HPPD tolerance
7	Insect resistance	21	Drought tolerance	33	High oil
9	Modified flower color	22	Delayed ripening/senescence	34	Aryloxyalkanoate tol.
11	ALS Herbicide Tol.	23	Modified product quality	35	Mesotrione tolerance
12	Dicamba Tolerance	24	High cellulose	36	Reduced nicotine
13	Anti-allergy	25	Modified starch/carbohydrate	37	Modified product
14	Salt tolerance	26	Insect & disease resist.		

Exhibit C

Crop	Event Name	Event Code	Trait(s)	Gene(s)
Alfalfa	J101	MON-00101-8	1	cp4 epsps (aroA:CP4)
Alfalfa	J163	MON-ØØ163-7	1	cp4 epsps (aroA:CP4)
Canola*	23-18-17 (Event 18)	CGN-89465-2	2	te
Canola*	23-198 (Event 23)	CGN-89465-2	2	te
Canola*	61061	DP-Ø61Ø61-7	1	gat4621
Canola*	73496	DP-Ø73496-4	1	gat4621
Canola*	GT200 (RT200)	MON-89249-2	1	cp4 epsps (aroA:CP4); goxv247
Canola*	GT73 (RT73)	MON-ØØØ73-7	1	cp4 epsps (aroA:CP4); goxv247
Canola*	HCN10 (Topas 19/2)	-	3	bar
Canola*	HCN28 (T45)	ACS-BNØØ8-2	3	pat (syn)
Canola*	HCN92 (Topas 19/2)	ACS-BNØØ7-1	3	bar
Canola*	MON88302	MON-883Ø2-9	1	cp4 epsps (aroA:CP4)
Canola*	MPS961	-	4	phyA
Canola*	MPS962	-	4	phyA
Canola*	MPS963	-	4	phyA
Canola*	MPS964	-	4	phyA
Canola*	MPS965	-	4	phyA
Canola*	MS1 (B91-4)	ACS-BNØØ4-7	3	bar
Canola*	MS8	ACS-BNØØ5-8	3	bar
Canola*	OXY-235	ACS-BNØ11-5	5	bxn
Canola*	PHY14	-	3	bar
Canola*	PHY23	-	3	bar
Canola*	PHY35	-	3	bar
Canola*	PHY36	-	3	bar
Canola*	RF1 (B93-101)	ACS-BNØØ1-4	3	bar
Canola*	RF2 (B94-2)	ACS-BNØØ2-5	3	bar
Canola*	RF3	ACS-BNØØ3-6	3	bar
Bean	EMBRAPA 5.1	EMB-PV051-1	6	ac1 (sense and antisense)
Brinjal #	EE-1	-	7	cry1Ac
Cotton	19-51a	DD-Ø1951A-7	11	S4-HrA
Cotton	281-24-236	DAS-24236-5	3,7	pat (syn); cry1F
Cotton	3006-210-23	DAS-21Ø23-5	3,7	pat (syn); cry1Ac
Cotton	31707	-	5,7	bxn; cry1Ac
Cotton	31803	-	5,7	bxn; cry1Ac
Cotton	31807	-	5,7	bxn; cry1Ac
Cotton	31808	-	5,7	bxn; cry1Ac
Cotton	42317	-	5,7	bxn; cry1Ac
Cotton	BNLA-601	-	7	cry1Ac
Cotton	BXN10211	BXN10211-9	5	bxn; cry1Ac
Cotton	BXN10215	BXN10215-4	5	bxn; cry1Ac

Cotton	BXN10222	BXN10222-2	5	bxn; cry1Ac
Cotton	BXN10224	BXN10224-4	5	bxn; cry1Ac
Cotton	COT102	SYN-IR102-7	7	vip3A(a)
Cotton	COT67B	SYN-IR67B-1	7	cry1Ab
Cotton	COT202	-	7	vip3A
Cotton	Event 1	-	7	cry1Ac
Cotton	GMF Cry1A	GTL-GMF311-7	7	cry1Ab-Ac
Cotton	GHB119	BCS-GH005-8	7	cry2Ac
Cotton	GHB614	BCS-GH002-5	1	2mepsps
Cotton	GK12	-	7	cry1Ab-Ac
Cotton	LLCotton25	ACS-GH001-3	3	bar
Cotton	MLS 9124	-	7	cry1C
Cotton	MON1076	MON-89924-2	7	cry1Ac
Cotton	MON1445	MON-01445-2	1	cp4 epsps (aroA:CP4)
Cotton	MON15985	MON-15985-7	7	cry1Ac; cry2Ab2
Cotton	MON1698	MON-89383-1	7	cp4 epsps (aroA:CP4)
Cotton	MON531	MON-00531-6	7	cry1Ac
Cotton	MON757	MON-00757-7	7	cry1Ac
Cotton	MON88913	MON-88913-8	1	cp4 epsps (aroA:CP4)
Cotton	Nqwe Chi 6 Bt	-	7	-
Cotton	SKG321	-	7	cry1A; CpTI
Cotton	T303-3	BCS-GH003-6	3,7	cry1Ab; bar
Cotton	T304-40	BCS-GH004-7	3,7	cry1Ab; bar
Cotton	CE43-67B	-	7	cry1Ab
Cotton	CE46-02A	-	7	cry1Ab
Cotton	CE44-69D	-	7	cry1Ab
Cotton	1143-14A	-	7	cry1Ab
Cotton	1143-51B	-	7	cry1Ab
Cotton	T342-142	-	7	cry1Ab
Cotton	PV-GHGT07 (1445)	-	1	cp4 epsps (aroA:CP4)
Cotton	EE-GH3	-	1	mepsps
Cotton	EE-GH5	-	7	cry1Ab
Cotton	MON88701	MON-88701-3	3,12	Modified dmo; bar
Cotton	OsCr11	-	13	Modified Cry j
Flax	FP967	CDC-FL001-2	11	als
Lentil	RH44	-	16	als
Maize	3272	SYN-E3272-5	17	amy797E
Maize	5307	SYN-05307-1	7	ecry3.1Ab
Maize	59122	DAS-59122-7	3,7	cry34Ab1; cry35Ab1; pat
Maize	676	PH-000676-7	3,18	pat; dam
Maize	678	PH-000678-9	3,18	pat; dam

Maize	680	PH-000680-2	3,18	pat; dam
Maize	98140	DP-098140-6	1,11	gat4621; zm-hra
Maize	Bt10	-	3,7	cry1Ab; pat
Maize	Bt176 (176)	SYN-EV176-9	3,7	cry1Ab; bar
Maize	BVLA430101	-	4	phyA2
Maize	CBH-351	ACS-ZM004-3	3,7	cry9C; bar
Maize	DAS40278-9	DAS40278-9	19	aad-1
Maize	DBT418	DKB-89614-9	3,7	cry1Ac; pinII; bar
Maize	DLL25 (B16)	DKB-89790-5	3	bar
Maize	GA21	MON-00021-9	1	mepsps
Maize	GG25	-	1	mepsps
Maize	GJ11	-	1	mepsps
Maize	Fl117	-	1	mepsps
Maize	GAT-ZM1	-	3	pat
Maize	LY038	REN-00038-3	20	cordapA
Maize	MIR162	SYN-IR162-4	7	vip3Aa20
Maize	MIR604	SYN-IR604-5	7	mcry3A
Maize	MON801 (MON80100)	MON801	1,7	cry1Ab; cp4 epsps (aroA:CP4); goxv247
Maize	MON802	MON-80200-7	1,7	cry1Ab; cp4 epsps (aroA:CP4); goxv247
Maize	MON809	PH-MON-809-2	1,7	cry1Ab; cp4 epsps (aroA:CP4); goxv247
Maize	MON810	MON-00810-6	1,7	cry1Ab; cp4 epsps (aroA:CP4); goxv247
Maize	MON832	-	1	cp4 epsps (aroA:CP4); goxv247
Maize	MON863	MON-00863-5	7	cry3Bb1
Maize	MON87427	MON-87427-7	1	cp4 epsps (aroA:CP4)
Maize	MON87460	MON-87460-4	21	cspB
Maize	MON88017	MON-88017-3	1,7	cry3Bb1; cp4 epsps (aroA:CP4)
Maize	MON89034	MON-89034-3	7	cry2Ab2; cry1A.105
Maize	MS3	ACS-ZM001-9	3,18	bar; barnase
Maize	MS6	ACS-ZM005-4	3,18	bar; barnase
Maize	NK603	MON-00603-6	1	cp4 epsps (aroA:CP4)
Maize	T14	ACS-ZM002-1	3	pat (syn)
Maize	T25	ACS-ZM003-2	3	pat (syn)
Maize	TC1507	DAS-01507-1	3,7	cry1Fa2; pat
Maize	TC6275	DAS-06275-8	3,7	mocry1F; bar
Maize	VIP1034	-	3,7	vip3A; pat
Maize	43A47	DP-043A47-3	3,7	cry1F; cry34Ab1; cry35Ab1; pat
Maize	40416	DP-040416-8	3,7	cry1F; cry34Ab1; cry35Ab1; pat
Maize	32316	DP-032316-8	3,7	cry1F; cry34Ab1; cry35Ab1; pat
Maize	4114	DP-004114-3	3,7	cry1F; cry34Ab1; cry35Ab1; pat
Melon	Melon A	-	22	sam-k

Melon	Melon B	-	22	sam-k
Papaya	55-1	CUH-CP551-8	6	prsv cp
Papaya	63-1	CUH-CP631-7	6	prsv cp
Papaya	Huanong No. 1	-	6	prsv rep
Papaya	X17-2	UFL-X17CP-6	6	prsv cp
Plum	C-5	ARS-PLMC5-6	6	ppv cp
Canola**	ZSR500	-	1	cp4 epsps (aroA:CP4); goxv247
Canola**	ZSR502	-	1	cp4 epsps (aroA:CP4); goxv247
Canola**	ZSR503	-	1	cp4 epsps (aroA:CP4); goxv247
Rice	7Crp#242-95-7	-	13	7crp
Rice	7Crp#10	-	13	7crp
Rice	GM Shanyou 63	-	7	cry1Ab; cry1Ac
Rice	Huahui-1/TT51-1	-	7	cry1Ab; cry1Ac
Rice	LLRICE06	ACS-OS001-4	3	bar
Rice	LLRICE601	BCS-OS003-7	3	bar
Rice	LLRICE62	ACS-OS002-5	3	bar
Rice	Tarom molaii + cry1Ab	-	7	cry1Ab (truncated)
Rice	GAT-OS2	-	3	bar
Rice	GAT-OS3	-	3	bar
Rice	PE-7	-	7	Cry1Ac
Rice	7Crp#10	-	13	7crp
Rice	KPD627-8	-	27	OASA1D
Rice	KPD722-4	-	27	OASA1D
Rice	KA317	-	27	OASA1D
Rice	HW5	-	27	OASA1D
Rice	HW1	-	27	OASA1D
Rice	B-4-1-18	-	28	Δ OsBRI1
Rice	G-3-3-22	-	29	OSGA2ox1
Rice	AD77	-	6	DEF
Rice	AD51	-	6	DEF
Rice	AD48	-	6	DEF
Rice	AD41	-	6	DEF
Rice	13pNasNa800725atAprt1	-	30	HvNAS1; HvNAAT-A; APRT
Rice	13pAprt1	-	30	APRT
Rice	gHvNAS1-gHvNAAT-1	-	30	HvNAS1; HvNAAT-A; HvNAAT-B
Rice	gHvIDS3-1	-	30	HvIDS3
Rice	gHvNAAT1	-	30	HvNAAT-A; HvNAAT-B
Rice	gHvNAS1-1	-	30	HvNAS1
Rice	NIA-OS006-4	-	6	WRKY45
Rice	NIA-OS005-3	-	6	WRKY45
Rice	NIA-OS004-2	-	6	WRKY45

Rice	NIA-OS003-1	-	6	WRKY45
Rice	NIA-OS002-9	-	6	WRKY45
Rice	NIA-OS001-8	-	6	WRKY45
Rice	OsCr11	-	13	Modified Cry j
Rice	17053	-	1	cp4 epsps (aroA:CP4)
Rice	17314	-	1	cp4 epsps (aroA:CP4)
Rose	WKS82 / 130-4-1	IFD-52401-4	9	5AT; bp40 (f3'5'h)
Rose	WKS92 / 130-9-1	IFD-52901-9	9	5AT; bp40 (f3'5'h)
Soybean	260-05 (G94-1, G94-19, G168)	-	9	gm-fad2-1 (silencing locus)
Soybean	A2704-12	ACS-GM005-3	3	pat
Soybean	A2704-21	ACS-GM004-2	3	pat
Soybean	A5547-127	ACS-GM006-4	3	pat
Soybean	A5547-35	ACS-GM008-6	3	pat
Soybean	CV127	BPS-CV127-9	16	csr1-2
Soybean	DAS68416-4	DAS68416-4	3	pat
Soybean	DP305423	DP-305423-1	11,31	gm-fad2-1 (silencing locus); gm-hra
Soybean	DP356043	DP-356043-5	1,31	gm-fad2-1 (silencing locus); gat4601
Soybean	FG72	MST-FG072-3	32,1	2mepsps; hppdPF W336
Soybean	GTS 40-3-2 (40-3-2)	MON-04032-6	1	cp4 epsps (aroA:CP4)
Soybean	GU262	ACS-GM003-1	3	pat
Soybean	MON87701	MON-87701-2	7	cry1Ac
Soybean	MON87705	MON-87705-6	1,31	fatb1-A (sense & antisense); fad2-1A (sense & antisense); cp4 epsps (aroA:CP4)
Soybean	MON87708	MON-87708-9	1,12	dmo; cp4 epsps (aroA:CP4)
Soybean	MON87769	MON-87769-7	1,31	Pj.D6D; Nc.Fad3; cp4 epsps (aroA:CP4)
Soybean	MON89788	MON-89788-1	1	cp4 epsps (aroA:CP4)
Soybean	W62	ACS-GM002-9	3	bar
Soybean	W98	ACS-GM001-8	3	bar
Soybean	MON87754	MON-87754-1	33	dga2A
Soybean	DAS21606	DAS-21606	34,3	Modified aad-12; pat
Soybean	DAS44406	DAS-44406-6	1,3,34	Modified aad-12; 2mepsps; pat
Soybean	SYHT04R	SYN-0004R-8	35	Modified avhppd
Soybean	9582.814.19.1	-	3,7	cry1Ac; cry1F; pat
Squash	CZW3	SEM-ØCZW3-2	6	cmv cp; zymv cp; wmv cp
Squash	ZW20	SEM-ØZW20-7	6	zymv cp; wmv cp
Sugar Beet	GTSB77 (T9100152)	SY-GTSB77-8	1	cp4 epsps (aroA:CP4); goxv247
Sugar Beet	H7-1	KM-000H71-4	1	cp4 epsps (aroA:CP4)
Sugar Beet	T120-7	ACS-BV001-3	3	pat
Sugar Beet	T227-1	-	1	cp4 epsps (aroA:CP4)
Sugarcane	NXI-1T	-	21	EcbetA

Sunflower	X81359	-	16	als
Pepper	PK-SP01	-	6	cmv cp
Tobacco	C/F/93/08-02	-	5	bxn
Tobacco	Vector 21-41	-	36	NtQPT1 (antisense)
Wheat	MON71800	MON-71800-3	1	cp4 epsps (aroA:CP4)

* Argentine (*Brassica napus*), ** Polish (*B. rapa*), # Eggplant

Treatment of genetically modified plants with compounds of the invention may result in super-additive or synergistic effects. For example, reduction in application rates, broadening of the activity spectrum, increased tolerance to biotic/abiotic stresses or enhanced storage stability may be greater than expected from just simple additive effects of the application of compounds of the invention on genetically modified plants.

Compounds of this invention can also be mixed with one or more other biologically active compounds or agents including herbicides, herbicide safeners, fungicides, insecticides, nematocides, bactericides, acaricides, growth regulators such as insect molting inhibitors and rooting stimulants, chemosterilants, semiochemicals, repellents, attractants, pheromones, feeding stimulants, plant nutrients, other biologically active compounds or entomopathogenic bacteria, virus or fungi to form a multi-component pesticide giving an even broader spectrum of agricultural protection. Mixtures of the compounds of the invention with other herbicides can broaden the spectrum of activity against additional weed species, and suppress the proliferation of any resistant biotypes. Thus the present invention also pertains to a composition comprising a compound of Formula 1 (in a herbicidally effective amount) and at least one additional biologically active compound or agent (in a biologically effective amount) and can further comprise at least one of a surfactant, a solid diluent or a liquid diluent. The other biologically active compounds or agents can be formulated in compositions comprising at least one of a surfactant, solid or liquid diluent. For mixtures of the present invention, one or more other biologically active compounds or agents can be formulated together with a compound of Formula 1, to form a premix, or one or more other biologically active compounds or agents can be formulated separately from the compound of Formula 1, and the formulations combined together before application (e.g., in a spray tank) or, alternatively, applied in succession.

A mixture of one or more of the following herbicides with a compound of this invention may be particularly useful for weed control: acetochlor, acifluorfen and its sodium salt, aclonifen, acrolein (2-propenal), alachlor, alloxymid, ametryn, amicarbazone, amidosulfuron, aminocyclopyrachlor and its esters (e.g., methyl, ethyl) and salts (e.g., sodium, potassium), aminopyralid, amitrole, ammonium sulfamate, anilofos, asulam, atrazine, azimsulfuron, beflubutamid, benazolin, benazolin-ethyl, bencarbazone, benfluralin, benfuresate, bensulfuron-methyl, bensulide, bentazone, benzobicyclon, benzofenap, bicyclopiron, bifenox, bilanafos, bispyribac and its sodium salt, bromacil, bromobutide,

bromofenoxim, bromoxynil, bromoxynil octanoate, butachlor, butafenacil, butamifos, butralin, butoxydim, butylate, cafenstrole, carbetamide, carfentrazone-ethyl, catechin, chlomethoxyfen, chloramben, chlorbromuron, chlorflurenol-methyl, chloridazon, chlorimuron-ethyl, chlorotoluron, chlorpropham, chlorsulfuron, chlorthal-dimethyl,

5 chlorthiamid, cinidon-ethyl, cinmethylin, cinosulfuron, clacyfos, clefoxydim, clethodim, clodinafop-propargyl, clomazone, clomeprop, clopyralid, clopyralid-olamine, cloransulam-methyl, cumyluron, cyanazine, cycloate, cyclopyrimorate, cyclosulfamuron, cycloxydim, cyhalofop-butyl, 2,4-D and its butotyl, butyl, isooctyl and isopropyl esters and its dimethylammonium, diolamine and trolamine salts, daimuron, dalapon, dalapon-sodium,

10 dazomet, 2,4-DB and its dimethylammonium, potassium and sodium salts, desmedipham, desmetryn, dicamba and its diglycolammonium, dimethylammonium, potassium and sodium salts, dichlobenil, dichlorprop, diclofop-methyl, diclosulam, difenzoquat metilsulfate, diflufenican, diflufenzopyr, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, dimethipin, dimethylarsinic acid and its sodium salt,

15 dinitramine, dinoterb, diphenamid, diquat dibromide, dithiopyr, diuron, DNOC, endothal, EPTC, esprocarb, ethalfluralin, ethametsulfuron-methyl, ethiozin, ethofumesate, ethoxyfen, ethoxysulfuron, etobenzanid, fenoxaprop-ethyl, fenoxaprop-P-ethyl, fenoxasulfone, fenquinotriene, fentrazamide, fenuron, fenuron-TCA, flamprop-methyl, flamprop-M-isopropyl, flamprop-M-methyl, flazasulfuron, florasulam, fluazifop-butyl,

20 fluazifop-P-butyl, fluazolate, flucarbazone, flucetosulfuron, fluchloralin, flufenacet, flufenpyr, flufenpyr-ethyl, flumetsulam, flumiclorac-pentyl, flumioxazin, fluometuron, fluoroglycofen-ethyl, flupoxam, flupyrsulfuron-methyl and its sodium salt, flurenol, flurenol-butyl, fluridone, flurochloridone, fluroxypyr, flurtamone, fluthiacet-methyl, fomesafen, foramsulfuron, fosamine-ammonium, glufosinate, glufosinate-ammonium,

25 glufosinate-P, glyphosate and its salts such as ammonium, isopropylammonium, potassium, sodium (including sesquisodium) and trimesium (alternatively named sulfosate), halauxifen, halauxifen-methyl, halosulfuron-methyl, haloxyfop-etotyl, haloxyfop-methyl, hexazinone, imazamethabenz-methyl, imazamox, imazapic, imazapyr, imazaquin, imazaquin-ammonium, imazethapyr, imazethapyr-ammonium, imazosulfuron, indanofan, indaziflam, iofensulfuron,

30 iodosulfuron-methyl, ioxynil, ioxynil octanoate, ioxynil-sodium, ipfencarbazone, isoproturon, isouron, isoxaben, isoxaflutole, isoxachlortole, lactofen, lenacil, linuron, maleic hydrazide, MCPA and its salts (e.g., MCPA-dimethylammonium, MCPA-potassium and MCPA-sodium, esters (e.g., MCPA-2-ethylhexyl, MCPA-butotyl) and thioesters (e.g., MCPA-thioethyl), MCPB and its salts (e.g., MCPB-sodium) and esters (e.g., MCPB-ethyl),

35 mecoprop, mecoprop-P, mefenacet, mefluidide, mesosulfuron-methyl, mesotrione, metam-sodium, metamifop, metamitron, metazachlor, metazosulfuron, methabenzthiazuron, methylarsonic acid and its calcium, monoammonium, monosodium and disodium salts, methyldymron, metobenzuron, metobromuron, metolachlor, S-metolachlor, metosulam,

metoxuron, metribuzin, metsulfuron-methyl, molinate, monolinuron, naproanilide, napropamide, napropamide-M, naptalam, neburon, nicosulfuron, norflurazon, orbencarb, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxasulfuron, oxaziclomefone, oxyfluorfen, paraquat dichloride, pebulate, pelargonic acid, pendimethalin, penoxsulam, pentanochlor, pentoxazone, perfluidone, pethoxamid, pethoxyamid, phenmedipham, picloram, picloram-potassium, picolinafen, pinoxaden, piperophos, pretilachlor, primisulfuron-methyl, prodiamine, profoxydim, prometon, prometryn, propachlor, propanil, propaquizafop, propazine, propham, propisochlor, propoxycarbazone, propyrisulfuron, propyzamide, prosulfocarb, prosulfuron, pyraclonil, pyraflufen-ethyl, pyrasulfotole, pyrazogyl, pyrazolynate, pyrazoxyfen, pyrazosulfuron-ethyl, pyribenzoxim, pyributicarb, pyridate, pyriftalid, pyriminobac-methyl, pyrimisulfan, pyrithiobac, pyrithiobac-sodium, pyroxasulfone, pyroxsulam, quinclorac, quinmerac, quinochloramine, quizalofop-ethyl, quizalofop-P-ethyl, quizalofop-P-tefuryl, rimsulfuron, saflufenacil, sethoxydim, siduron, simazine, simetryn, sulcotrione, sulfentrazone, sulfometuron-methyl, sulfosulfuron, 2,3,6-TBA, TCA, TCA-sodium, tebutam, tebuthiuron, tefuryltrione, tembotrione, tepraloxydim, terbacil, terbutometon, terbuthylazine, terbutryn, thenylchlor, thiazopyr, thiencarbazone, thifensulfuron-methyl, thiobencarb, tiafenacil, tiocarbamil, topramezone, tralkoxydim, tri-allate, triafamone, triasulfuron, triaziflam, tribenuron-methyl, triclopyr, triclopyr-butyl, triclopyr-triethylammonium, tridiphane, trietazine, trifloxysulfuron, trifluralin, triflurosulfuron-methyl, tritosulfuron, vernolate, 3-(2-chloro-3,6-difluorophenyl)-4-hydroxy-1-methyl-1,5-naphthyridin-2(1H)-one, 5-chloro-3-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-1-(4-methoxyphenyl)-2(1H)-quinoxalinone, 2-chloro-N-(1-methyl-1H-tetrazol-5-yl)-6-(trifluoromethyl)-3-pyridinecarboxamide, 7-(3,5-dichloro-4-pyridinyl)-5-(2,2-difluoroethyl)-8-hydroxypyrido[2,3-b]pyrazin-6(5H)-one, 4-(2,6-diethyl-4-methylphenyl)-5-hydroxy-2,6-dimethyl-3(2H)-pyridazinone, 5-[[[(2,6-difluorophenyl)methoxy]methyl]-4,5-dihydro-5-methyl-3-(3-methyl-2-thienyl)isoxazole (previously methioxolin), 3-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propyn-1-yl)-2H-1,4-benzoxazin-6-yl]dihydro-1,5-dimethyl-6-thioxo-1,3,5-triazine-2,4(1H,3H)-dione, 4-(4-fluorophenyl)-6-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-2-methyl-1,2,4-triazine-3,5(2H,4H)-dione, methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoro-2-pyridinecarboxylate, 2-methyl-3-(methylsulfonyl)-N-(1-methyl-1H-tetrazol-5-yl)-4-(trifluoromethyl)benzamide and 2-methyl-N-(4-methyl-1,2,5-oxadiazol-3-yl)-3-(methylsulfinyl)-4-(trifluoromethyl)benzamide. Other herbicides also include bioherbicides such as *Alternaria destruens* Simmons, *Colletotrichum gloeosporioides* (Penz.) Penz. & Sacc., *Drechslera monoceras* (MTB-951), *Myrothecium verrucaria* (Albertini & Schweinitz) Ditmar: Fries, *Phytophthora palmivora* (Butl.) Butl. and *Puccinia thlaspeos* Schub.

Compounds of this invention can also be used in combination with plant growth regulators such as aviglycine, N-(phenylmethyl)-1H-purin-6-amine, epocholeone, gibberellic

acid, gibberellin A₄ and A₇, harpin protein, mepiquat chloride, prohexadione calcium, prohydrojasmon, sodium nitrophenolate and trinexapac-methyl, and plant growth modifying organisms such as *Bacillus cereus* strain BP01.

General references for agricultural protectants (i.e. herbicides, herbicide safeners, insecticides, fungicides, nematocides, acaricides and biological agents) include *The Pesticide Manual, 13th Edition*, C. D. S. Tomlin, Ed., British Crop Protection Council, Farnham, Surrey, U.K., 2003 and *The BioPesticide Manual, 2nd Edition*, L. G. Copping, Ed., British Crop Protection Council, Farnham, Surrey, U.K., 2001.

For embodiments where one or more of these various mixing partners are used, the weight ratio of these various mixing partners (in total) to the compound of Formula 1 is typically between about 1:3000 and about 3000:1. Of note are weight ratios between about 1:300 and about 300:1 (for example ratios between about 1:30 and about 30:1). One skilled in the art can easily determine through simple experimentation the biologically effective amounts of active ingredients necessary for the desired spectrum of biological activity. It will be evident that including these additional components may expand the spectrum of weeds controlled beyond the spectrum controlled by the compound of Formula 1 alone.

In certain instances, combinations of a compound of this invention with other biologically active (particularly herbicidal) compounds or agents (i.e. active ingredients) can result in a greater-than-additive (i.e. synergistic) effect on weeds and/or a less-than-additive effect (i.e. safening) on crops or other desirable plants. Reducing the quantity of active ingredients released in the environment while ensuring effective pest control is always desirable. Ability to use greater amounts of active ingredients to provide more effective weed control without excessive crop injury is also desirable. When synergism of herbicidal active ingredients occurs on weeds at application rates giving agronomically satisfactory levels of weed control, such combinations can be advantageous for reducing crop production cost and decreasing environmental load. When safening of herbicidal active ingredients occurs on crops, such combinations can be advantageous for increasing crop protection by reducing weed competition.

Of note is a combination of a compound of the invention with at least one other herbicidal active ingredient. Of particular note is such a combination where the other herbicidal active ingredient has different site of action from the compound of the invention. In certain instances, a combination with at least one other herbicidal active ingredient having a similar spectrum of control but a different site of action will be particularly advantageous for resistance management. Thus, a composition of the present invention can further comprise (in a herbicidally effective amount) at least one additional herbicidal active ingredient having a similar spectrum of control but a different site of action.

Compounds of this invention can also be used in combination with herbicide safeners such as allidochlor, benoxacor, cloquintocet-mexyl, cumyluron, cyometrinil,

cyprosulfonamide, daimuron, dichlormid, dicyclonon, dietholate, dimepiperate, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, isoxadifen-ethyl, mefenpyr-diethyl, mephenate, methoxyphenone naphthalic anhydride (1,8-naphthalic anhydride), oxabetrinil, *N*-(aminocarbonyl)-2-methylbenzenesulfonamide, *N*-(aminocarbonyl)-2-fluorobenzenesulfonamide, 1-bromo-4-[(chloromethyl)sulfonyl]benzene (BCS), 4-(dichloroacetyl)-1-oxa-4-azospiro[4.5]decane (MON 4660), 2-(dichloromethyl)-2-methyl-1,3-dioxolane (MG 191), ethyl 1,6-dihydro-1-(2-methoxyphenyl)-6-oxo-2-phenyl-5-pyrimidinecarboxylate, 2-hydroxy-*N,N*-dimethyl-6-(trifluoromethyl)pyridine-3-carboxamide, and 3-oxo-1-cyclohexen-1-yl 1-(3,4-dimethylphenyl)-1,6-dihydro-6-oxo-2-phenyl-5-pyrimidinecarboxylate to increase safety to certain crops. Antidotally effective amounts of the herbicide safeners can be applied at the same time as the compounds of this invention, or applied as seed treatments. Therefore an aspect of the present invention relates to a herbicidal mixture comprising a compound of this invention and an antidotally effective amount of a herbicide safener. Seed treatment is particularly useful for selective weed control, because it physically restricts antidoting to the crop plants. Therefore a particularly useful embodiment of the present invention is a method for selectively controlling the growth of undesired vegetation in a crop comprising contacting the locus of the crop with a herbicidally effective amount of a compound of this invention wherein seed from which the crop is grown is treated with an antidotally effective amount of safener. Antidotally effective amounts of safeners can be easily determined by one skilled in the art through simple experimentation.

Of note is a composition comprising a compound of the invention (in a herbicidally effective amount), at least one additional active ingredient selected from the group consisting of other herbicides and herbicide safeners (in an effective amount), and at least one component selected from the group consisting of surfactants, solid diluents and liquid diluents.

Table A1 lists specific combinations of a Component (a) with Component (b) illustrative of the mixtures, compositions and methods of the present invention. Compound 1 in the Component (a) column is identified in Index Table A. The second column of Table A1 lists the specific Component (b) compound (e.g., "2,4-D" in the first line). The third, fourth and fifth columns of Table A1 lists ranges of weight ratios for rates at which the Component (a) compound is typically applied to a field-grown crop relative to Component (b) (i.e. (a):(b)). Thus, for example, the first line of Table A1 specifically discloses the combination of Component (a) (i.e. Compound 1 in Index Table A) with 2,4-D is typically applied in a weight ratio between 1:168 – 6:1. The remaining lines of Table A1 are to be construed similarly.

TABLE A1

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
1	2,4-D	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Acetochlor	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Acifluorfen	1:84 to 11:1	1:28 to 4:1	1:8 to 2:1
1	Aclonifen	1:750 to 2:1	1:250 to 1:3	1:75 to 1:9
1	Alachlor	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Ametryn	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Amicarbazone	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Amidosulfuron	1:6 to 150:1	1:2 to 50:1	1:1 to 15:1
1	Aminocyclopyrachlor	1:42 to 22:1	1:14 to 8:1	1:4 to 3:1
1	Aminopyralid	1:18 to 50:1	1:6 to 17:1	1:1 to 5:1
1	Amitrole	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Anilofos	1:84 to 11:1	1:28 to 4:1	1:8 to 2:1
1	Asulam	1:840 to 2:1	1:280 to 1:3	1:84 to 1:10
1	Atrazine	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Azimsulfuron	1:6 to 150:1	1:2 to 50:1	1:1 to 15:1
1	Beflubutamid	1:300 to 3:1	1:100 to 1:1	1:30 to 1:4
1	Benfuresate	1:540 to 2:1	1:180 to 1:2	1:54 to 1:6
1	Bensulfuron-methyl	1:22 to 40:1	1:7 to 14:1	1:2 to 4:1
1	Bentazon	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Benzobicyclon	1:75 to 12:1	1:25 to 4:1	1:7 to 2:1
1	Benzofenap	1:225 to 4:1	1:75 to 2:1	1:22 to 1:3
1	Bicyclopyrone	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Bifenox	1:225 to 4:1	1:75 to 2:1	1:22 to 1:3
1	Bispyribac-sodium	1:9 to 100:1	1:3 to 34:1	1:1 to 10:1
1	Bromacil	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Bromobutide	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Bromoxynil	1:84 to 11:1	1:28 to 4:1	1:8 to 2:1
1	Butachlor	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Butafenacil	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Butylate	1:1350 to 1:2	1:450 to 1:5	1:135 to 1:15
1	Carfenstrole	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Carfentrazone-ethyl	1:112 to 8:1	1:37 to 3:1	1:11 to 1:2
1	Chlorimuron-ethyl	1:7 to 120:1	1:2 to 40:1	1:1 to 12:1
1	Chlorotoluron	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
1	Chlorsulfuron	1:6 to 150:1	1:2 to 50:1	1:1 to 15:1
1	Cincosulfuron	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Cinidon-ethyl	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Cinmethylin	1:30 to 30:1	1:10 to 10:1	1:3 to 3:1
1	Clacyfos	1:84 to 6:1	1:28 to 2:1	1:16 to 1:2
1	Clethodim	1:42 to 22:1	1:14 to 8:1	1:4 to 3:1
1	Clodinafop-propargyl	1:18 to 50:1	1:6 to 17:1	1:1 to 5:1
1	Clomazone	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Clomeprop	1:150 to 6:1	1:50 to 2:1	1:15 to 1:2
1	Clopyralid	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Cloransulam-methyl	1:10 to 86:1	1:3 to 29:1	1:1 to 9:1
1	Cumyluron	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Cyanazine	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Cyclopyrimorate	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Cyclosulfamuron	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Cycloxydim	1:84 to 11:1	1:28 to 4:1	1:8 to 2:1
1	Cyhalofop	1:22 to 40:1	1:7 to 14:1	1:2 to 4:1
1	Daimuron	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Desmedipham	1:282 to 4:1	1:94 to 2:1	1:28 to 1:4
1	Dicamba	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Dichlobenil	1:1200 to 1:2	1:400 to 1:4	1:120 to 1:14
1	Dichlorprop	1:810 to 2:1	1:270 to 1:3	1:81 to 1:9
1	Diclofop-methyl	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Diclosulam	1:9 to 100:1	1:3 to 34:1	1:1 to 10:1
1	Difenzoquat	1:252 to 4:1	1:84 to 2:1	1:25 to 1:3
1	Diflufenican	1:750 to 2:1	1:250 to 1:3	1:75 to 1:9
1	Diflufenzopyr	1:10 to 86:1	1:3 to 29:1	1:1 to 9:1
1	Dimethachlor	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Dimethametryn	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Dimethenamid-p	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Dithiopyr	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Diuron	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	EPTC	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Esprocarb	1:1200 to 1:2	1:400 to 1:4	1:120 to 1:14
1	Ethalfuralin	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
1	Ethametsulfuron-methyl	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Ethoxyfen	1:7 to 120:1	1:2 to 40:1	1:1 to 12:1
1	Ethoxysulfuron	1:18 to 50:1	1:6 to 17:1	1:1 to 5:1
1	Etobenzanid	1:225 to 4:1	1:75 to 2:1	1:22 to 1:3
1	Fenoxaprop-ethyl	1:105 to 9:1	1:35 to 3:1	1:10 to 1:2
1	Fenoxasulfone	1:75 to 12:1	1:25 to 4:1	1:7 to 2:1
1	Fenquinotrione	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Fentrazamide	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Flazasulfuron	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Florasulam	1:2 to 375:1	1:1 to 125:1	4:1 to 38:1
1	Fluazifop-butyl	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Flucarbazone	1:7 to 120:1	1:2 to 40:1	1:1 to 12:1
1	Flucetosulfuron	1:7 to 120:1	1:2 to 40:1	1:1 to 12:1
1	Flufenacet	1:225 to 4:1	1:75 to 2:1	1:22 to 1:3
1	Flumetsulam	1:21 to 43:1	1:7 to 15:1	1:2 to 5:1
1	Flumiclorac-pentyl	1:9 to 100:1	1:3 to 34:1	1:1 to 10:1
1	Flumioxazin	1:22 to 40:1	1:7 to 14:1	1:2 to 4:1
1	Fluometuron	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Flupyr-sulfuron-methyl	1:3 to 300:1	1:1 to 100:1	3:1 to 30:1
1	Fluridone	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Fluroxypyr-meptyl	1:84 to 11:1	1:28 to 4:1	1:8 to 2:1
1	Flurtamone	1:750 to 2:1	1:250 to 1:3	1:75 to 1:9
1	Fluthiacet-methyl	1:42 to 38:1	1:14 to 13:1	1:2 to 5:1
1	Fomesafen	1:84 to 11:1	1:28 to 4:1	1:8 to 2:1
1	Foramsulfuron	1:12 to 75:1	1:4 to 25:1	1:1 to 8:1
1	Glufosinate	1:252 to 4:1	1:84 to 2:1	1:25 to 1:3
1	Glyphosate	1:252 to 4:1	1:84 to 2:1	1:25 to 1:3
1	Halauxifen	1:18 to 50:1	1:6 to 17:1	1:1 to 5:1
1	Halauxifen-methyl	1:18 to 50:1	1:6 to 17:1	1:1 to 5:1
1	Halosulfuron-methyl	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Haloxypyr-methyl	1:30 to 30:1	1:10 to 10:1	1:3 to 3:1
1	Hexazinone	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Imazamox	1:12 to 75:1	1:4 to 25:1	1:1 to 8:1
1	Imazapic	1:18 to 50:1	1:6 to 17:1	1:1 to 5:1
1	Imazapyr	1:75 to 12:1	1:25 to 4:1	1:7 to 2:1

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
1	Imazaquin	1:30 to 30:1	1:10 to 10:1	1:3 to 3:1
1	Imazethabenz-methyl	1:150 to 6:1	1:50 to 2:1	1:15 to 1:2
1	Imazethapyr	1:21 to 43:1	1:7 to 15:1	1:2 to 5:1
1	Imazosulfuron	1:24 to 38:1	1:8 to 13:1	1:2 to 4:1
1	Indanofan	1:300 to 3:1	1:100 to 1:1	1:30 to 1:4
1	Indaziflam	1:22 to 40:1	1:7 to 14:1	1:2 to 4:1
1	Iodosulfuron-methyl	1:3 to 300:1	1:1 to 100:1	3:1 to 30:1
1	Ioxynil	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Ipfencarbazone	1:75 to 12:1	1:25 to 4:1	1:7 to 2:1
1	Isoproturon	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Isoxaben	1:252 to 4:1	1:84 to 2:1	1:25 to 1:3
1	Isoxaflutole	1:52 to 18:1	1:17 to 6:1	1:5 to 2:1
1	Lactofen	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Lenacil	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Linuron	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	MCPA	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	MCPB	1:252 to 4:1	1:84 to 2:1	1:25 to 1:3
1	Mecoprop	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Mefenacet	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Mefluidide	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Mesosulfuron-methyl	1:4 to 200:1	1:1 to 67:1	2:1 to 20:1
1	Mesotrione	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Metamifop	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Metazachlor	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Metazosulfuron	1:22 to 40:1	1:7 to 14:1	1:2 to 4:1
1	Methabenzthiazuron	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Metolachlor	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Metosulam	1:7 to 120:1	1:2 to 40:1	1:1 to 12:1
1	Metribuzin	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Metsulfuron-methyl	1:1 to 500:1	1:1 to 167:1	5:1 to 50:1
1	Molinate	1:900 to 1:1	1:300 to 1:3	1:90 to 1:10
1	Napropamide	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Napropamide-M	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Naptalam	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Nicosulfuron	1:10 to 86:1	1:3 to 29:1	1:1 to 9:1

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
1	Norflurazon	1:1008 to 1:2	1:336 to 1:4	1:100 to 1:12
1	Orbencarb	1:1200 to 1:2	1:400 to 1:4	1:120 to 1:14
1	Orthosulfamuron	1:18 to 50:1	1:6 to 17:1	1:1 to 5:1
1	Oryzalin	1:450 to 2:1	1:150 to 1:2	1:45 to 1:5
1	Oxadiargyl	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Oxadiazon	1:480 to 2:1	1:160 to 1:2	1:48 to 1:6
1	Oxasulfuron	1:24 to 38:1	1:8 to 13:1	1:2 to 4:1
1	Oxaziclomefone	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Oxyfluorfen	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Paraquat	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Pendimethalin	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Penoxsulam	1:9 to 100:1	1:3 to 34:1	1:1 to 10:1
1	Pentoxamid	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Pentoxazone	1:90 to 10:1	1:30 to 4:1	1:9 to 1:1
1	Phenmedipham	1:90 to 10:1	1:30 to 4:1	1:9 to 1:1
1	Picloram	1:84 to 11:1	1:28 to 4:1	1:8 to 2:1
1	Picolinafen	1:30 to 30:1	1:10 to 10:1	1:3 to 3:1
1	Pinoxaden	1:22 to 40:1	1:7 to 14:1	1:2 to 4:1
1	Pretilachlor	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Primisulfuron-methyl	1:7 to 120:1	1:2 to 40:1	1:1 to 12:1
1	Prodiamine	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Profoxydim	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Prometryn	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Propachlor	1:1008 to 1:2	1:336 to 1:4	1:100 to 1:12
1	Propanil	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Propaquizafop	1:42 to 22:1	1:14 to 8:1	1:4 to 3:1
1	Propoxycarbazon	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Propyrisulfuron	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Propyzamide	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Prosulfocarb	1:1050 to 1:2	1:350 to 1:4	1:105 to 1:12
1	Prosulfuron	1:6 to 150:1	1:2 to 50:1	1:1 to 15:1
1	Pyraclonil	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Pyraflufen-ethyl	1:4 to 200:1	1:1 to 67:1	2:1 to 20:1
1	Pyrasulfotole	1:12 to 75:1	1:4 to 25:1	1:1 to 8:1
1	Pyrazolynate	1:750 to 2:1	1:250 to 1:3	1:75 to 1:9

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
1	Pyrazosulfuron-ethyl	1:9 to 100:1	1:3 to 34:1	1:1 to 10:1
1	Pyrazoxyfen	1:4 to 200:1	1:1 to 67:1	2:1 to 20:1
1	Pyribenzoxim	1:9 to 100:1	1:3 to 34:1	1:1 to 10:1
1	Pyributicarb	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Pyridate	1:252 to 4:1	1:84 to 2:1	1:25 to 1:3
1	Pyriftalid	1:9 to 100:1	1:3 to 34:1	1:1 to 10:1
1	Pyriminobac-methyl	1:18 to 50:1	1:6 to 17:1	1:1 to 5:1
1	Pyrimisulfan	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Pyrihiobac	1:21 to 43:1	1:7 to 15:1	1:2 to 5:1
1	Pyroxasulfone	1:75 to 12:1	1:25 to 4:1	1:7 to 2:1
1	Pyroxulam	1:4 to 200:1	1:1 to 67:1	2:1 to 20:1
1	Quinclorac	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Quizalofop-ethyl	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Rimsulfuron	1:12 to 75:1	1:4 to 25:1	1:1 to 8:1
1	Saflufenacil	1:22 to 40:1	1:7 to 14:1	1:2 to 4:1
1	Sethoxydim	1:84 to 11:1	1:28 to 4:1	1:8 to 2:1
1	Simazine	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Sulcotrione	1:105 to 9:1	1:35 to 3:1	1:10 to 1:2
1	Sulfentrazone	1:129 to 7:1	1:43 to 3:1	1:12 to 1:2
1	Sulfometuron-methyl	1:30 to 30:1	1:10 to 10:1	1:3 to 3:1
1	Sulfosulfuron	1:7 to 120:1	1:2 to 40:1	1:1 to 12:1
1	Tebuthiuron	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Tefuryltrione	1:37 to 24:1	1:12 to 8:1	1:3 to 3:1
1	Tembotrione	1:27 to 33:1	1:9 to 11:1	1:2 to 4:1
1	Tepraloxym	1:22 to 40:1	1:7 to 14:1	1:2 to 4:1
1	Terbacil	1:252 to 4:1	1:84 to 2:1	1:25 to 1:3
1	Terbuthylatrazine	1:750 to 2:1	1:250 to 1:3	1:75 to 1:9
1	Terbutryn	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Thenylchlor	1:75 to 12:1	1:25 to 4:1	1:7 to 2:1
1	Thiazopyr	1:336 to 3:1	1:112 to 1:2	1:33 to 1:4
1	Thiencarbazone	1:3 to 300:1	1:1 to 100:1	3:1 to 30:1
1	Thifensulfuron-methyl	1:4 to 200:1	1:1 to 67:1	2:1 to 20:1
1	Thiobencarb	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Topramazone	1:6 to 150:1	1:2 to 50:1	1:1 to 15:1
1	Tralkoxydim	1:60 to 15:1	1:20 to 5:1	1:6 to 2:1

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
1	Triafamone	1:3 to 38:1	1:1 to 13:1	1:1 to 8:1
1	Triallate	1:672 to 2:1	1:224 to 1:3	1:67 to 1:8
1	Triasulfuron	1:4 to 200:1	1:1 to 67:1	2:1 to 20:1
1	Triaziflam	1:150 to 6:1	1:50 to 2:1	1:15 to 1:2
1	Tribenuron-methyl	1:3 to 300:1	1:1 to 100:1	3:1 to 30:1
1	Triclopyr	1:168 to 6:1	1:56 to 2:1	1:16 to 1:2
1	Trifloxysulfuron	1:2 to 375:1	1:1 to 125:1	4:1 to 38:1
1	Trifluralin	1:252 to 4:1	1:84 to 2:1	1:25 to 1:3
1	Triflusulfuron-methyl	1:15 to 60:1	1:5 to 20:1	1:1 to 6:1
1	Tritosulfuron	1:12 to 75:1	1:4 to 25:1	1:1 to 8:1

Table A2 is constructed the same as Table A1 above except that entries below the “Component (a)” column heading are replaced with the respective Component (a) Column Entry shown below. Compound 1 in the Component (a) column is identified in Index Table A. Thus, for example, in Table A2 the entries below the “Component (a)” column heading all recite “Compound 12” (i.e. Compound 12 identified in Index Table A), and the first line below the column headings in Table A2 specifically discloses a mixture of Compound 12 with 2,4-D. Tables A3 through A9 are constructed similarly.

<u>Table Number</u>	<u>Component (a) Column Entries</u>	<u>Table Number</u>	<u>Component (a) Column Entries</u>
A2	Compound 12	A11	Compound 53
A3	Compound 15	A12	Compound 55
A4	Compound 21	A13	Compound 62
A5	Compound 23	A14	Compound 63
A6	Compound 24	A15	Compound 144
A7	Compound 27	A16	Compound 145
A8	Compound 32	A17	Compound 168
A9	Compound 42	A18	Compound 200
A10	Compound 35		

Preferred for better control of undesired vegetation (e.g., lower use rate such as from synergism, broader spectrum of weeds controlled, or enhanced crop safety) or for preventing the development of resistant weeds are mixtures of a compound of this invention with a herbicide selected from the group consisting of chlorimuron-ethyl, nicosulfuron, diuron, hexazinone, thifensulfuron-methyl and S-metolachlor.


The compounds of the present invention are useful for the control of weed species that are resistant to herbicides with the AHAS-inhibitor or (b2) [chemical compound that inhibits

acetoxy acid synthase (AHAS), also known as acetolactate synthase (ALS)] mode of action.

The following Tests demonstrate the control efficacy of the compounds of this invention against specific weeds. The weed control afforded by the compounds is not limited, however, to these species. See Index Table A for compound descriptions. Mass spectra are reported as the molecular weight of the highest isotopic abundance parent ion (M+1) formed by addition of H⁺ (molecular weight of 1) to the molecule, observed by mass spectrometry using atmospheric pressure chemical ionization (AP⁺) or electrospray ionization (ESI). The following abbreviations are used in the Index Table A which follow:

Ph is phenyl, pyridyl is pyridinyl, OEt is ethoxy, CN is cyano, , CHO is formyl, *t*-Bu is *tertiary*-butyl, *i*-Pr is *iso*-propyl, *c*-Pr is cyclopropyl, Me is methyl, Et is ethyl and C(=O)CH₃ is acyl. The abbreviation "Ex." stands for "Example" and is followed by a number indicating in which example the compound is prepared.

INDEX TABLE A



No.	Q	R ²	(R ³) _m	M.S.(AP ⁺) or m.p.
1	5-Cl-2-pyridyl	Cl	m = 0	*
2	6-N(CH ₃) ₂ -3-pyridyl	Cl	m = 0	327 ^a
3	5-Cl-2-pyrimidinyl	Cl	m = 0	320 ^a
4	3-pyridyl	Cl	m = 0	285 ^a
5	1,3,4-oxadiazol-2-yl	Br	m = 0	319
6	1,3,4-oxadiazol-2-yl	Cl	m = 0	275
7	5-oxazolyl	Cl	m = 0	274 ^a
8	4-thiazolyl	Cl	m = 0	290
9	5-thiazolyl	Cl	m = 0	290
10	1-CH ₃ -1 <i>H</i> -pyrazol-3-yl	Cl	m = 0	*
11	1-CH ₃ -1 <i>H</i> -pyrazol-4-yl	Cl	m = 0	*
12	3-Br-5-isoxazolyl	Cl	m = 0	352
13	5-thiazolyl	CF ₃	m = 0	324
14	4-thiazolyl	CF ₃	m = 0	324
15	2-Br-5-thiazolyl	CF ₃	m = 0	402

No.	Q	R ²	(R ³) _m	M.S.(AP+)
				or m.p.
16	6-Cl-2-pyridyl	Cl	m = 0	319 ^a *
17	6-Cl-2-pyridyl	Br	m = 0	363 ^a *
18	4-Cl-2-pyrimidinyl	Cl	m = 0	320 ^a *
19	6-CF ₃ -2-pyridyl	Cl	m = 0	352 ^a *
20	2-CF ₃ -4-pyrimidinyl	Cl	m = 0	353 ^a *
21	2-CF ₃ -4-pyrimidinyl	Br	m = 0	398 ^a *
22	6-CF ₃ -3-pyridyl	Cl	4-CH ₃	366 ^a
23	5-Cl-2-pyridyl	Cl	3-Cl	*
24	2-CF ₃ -4-pyridyl	Cl	4-CH ₃	366
25	4-CF ₃ -2-pyridyl	Cl	m = 0	352
26	4-CF ₃ -2-pyridyl	Br	m = 0	397
27	5-CF ₃ -2-pyridyl	Cl	m = 0	352
28	5-CF ₃ -2-pyridyl	Br	m = 0	397
29	1-CH ₃ -3-CF ₃ -1 <i>H</i> -pyrazol-5-yl	Cl	m = 0	355
30	1-CH ₃ -3-CF ₃ -1 <i>H</i> -pyrazol-5-yl	Br	m = 0	399
31	5-CH ₂ OH-3-isoxazolyl	Cl	m = 0	304
32	5-CH ₂ F-3-isoxazolyl	Cl	m = 0	306
33	5-CHO-3-isoxazolyl	Cl	m = 0	302
34	5-CH ₂ Cl-3-isoxazolyl	Cl	m = 0	322
35	5-CF ₂ H-3-isoxazolyl	Cl	m = 0	324
36	5-CH ₂ CN-3-isoxazolyl	Cl	m = 0	311 ^b
37	5-CH=NOH-3-isoxazolyl	Cl	m = 0	317
38	5-CN-3-isoxazolyl	Cl	m = 0	299 ^a
39	3-CF ₃ -Ph	Cl	4-CH ₃	365
40	3-OCF ₃ -Ph	Cl	m = 0	367 ^a *
41	3,5-di-Cl-Ph	Cl	m = 0	352 ^a
42	4-OCF ₃ -Ph	Cl	m = 0	367 ^a
43	4-CF ₃ -Ph	Cl	m = 0	351 ^a
44	3-OCF ₃ -Ph	Cl	4-CH ₃	381 ^a *
45	3-OCF ₃ -Ph	Cl	6-OCF ₃	397 ^a
46	3-OCF ₃ -Ph	Cl	3-F	385 ^a *
47	1-methyl-1 <i>H</i> -tetrazol-5-yl	Cl	m = 0	289
48	2-methyl-2 <i>H</i> -tetrazol-5-yl	Cl	m = 0	289
49	1-methyl-1 <i>H</i> -tetrazol-5-yl	Br	m = 0	334
50	2-methyl-2 <i>H</i> -tetrazol-5-yl	Br	m = 0	334
51	1-methyl-1 <i>H</i> -1,2,3-triazol-4-yl	Cl	m = 0	288

No.	Q	R ²	(R ³) _m	M.S.(AP+)
				or m.p.
52	1-methyl-1 <i>H</i> -1,2,3-triazol-5-yl	Cl	m = 0	288
53	3-CHF ₂ -5-isoxazolyl	Cl	m = 0	324
54	6-CF ₃ -4-pyrimidinyl	Cl	m = 0	*
55	5-CHF ₂ -3-isoxazolyl	Cl	3-F	96-98
56	5-CHO-3-isoxazolyl	Cl	3-F	137-139
57	5-CH ₂ F-3-isoxazolyl	Cl	3-F	324
58	3-CH ₃ -5-isoxazolyl	Cl	m = 0	288
59	5-(<i>t</i> -Bu)-3-isoxazolyl	Cl	m = 0	330
60	5-CH ₃ -3-isoxazolyl	Cl	m = 0	288
61	2-oxazolyl	Cl	m = 0	274
62	5-CHF ₂ -3-isoxazolyl	Br	m = 0	89-93
63	3-CF ₃ -5-isoxazolyl	Cl	m = 0	342
64	3-CF ₃ -5-isoxazolyl	Br	m = 0	388
65	3-CHF ₂ -5-isoxazolyl	Cl	5-F	342
66	3-CHF ₂ -5-isoxazolyl	Br	5-F	387
67	3-CClF ₂ -5-isoxazolyl	Cl	m = 0	359
68	3-CHF ₂ -5-isoxazolyl	Cl	6-F	342
69	3-CHF ₂ -5-isoxazolyl	Br	6-F	387
70	2-CH ₃ -4-oxazolyl	Cl	m = 0	288
71	2-CF ₃ -4-pyridinyl	Cl	m = 0	352
72	2-CF ₃ -4-pyridinyl	Br	m = 0	396
73	1-(<i>i</i> -Pr)-1 <i>H</i> -1,2,4-triazol-3-yl	Cl	m = 0	316
74	3-(<i>c</i> -Pr)-5-isoxazolyl	Cl	m = 0	314
75	3-CHF ₂ -5-isoxazolyl	Cl	4-F	342
76	3-CHF ₂ -5-isoxazolyl	Br	4-F	387
77	3,5-di-Me-4-isoxazolyl	Cl	3-F	320
78	3,5-di-Me-4-isoxazolyl	Cl	m = 0	302
79	2-CH ₂ CF ₃ -2 <i>H</i> -1,2,4-triazol-3-yl	Cl	m = 0	102-106
80	2-CF ₃ -4-pyridinyl	F	m = 0	336
81	2-CF ₃ -4-pyridinyl	Cl	3-F	370
82	2-CF ₃ -4-pyridinyl	Br	3-F	414
83	2-CF ₃ -4-pyridinyl	CH ₃	3-F	350
84	3-CF ₃ -5-isoxazolyl	Cl	4-F	360
85	3-CF ₃ -5-isoxazolyl	Br	4-F	405
86	5-(C≡CH)-3-isoxazolyl	Cl	m = 0	156-160
87	2-CF ₃ -4-pyridinyl	F	3-F	354

No.	Q	R ²	(R ³) _m	M.S.(AP+)
				or m.p.
88	5-(OCH ₂ CF ₂ H)-3-isoxazolyl	Cl	m = 0	352
89	1-Et-3-CF ₃ -1 <i>H</i> -pyrazol-5-yl	Cl	3-CN	394
90	1-(<i>i</i> -Pr)-3-CF ₃ -1 <i>H</i> -pyrazol-5-yl	Cl	3-CN	408
91	5-(CH=CF ₂)-3-isoxazolyl	Cl	3-F	354
92	3-(<i>c</i> -Pr)-5-isoxazolyl	Br	m = 0	359
93	1-CH ₂ CF ₃ -1 <i>H</i> -1,2,4-triazol-3-yl	Cl	m = 0	155-158
94	5-(OCH ₂ CF ₃)-3-isoxazolyl	Cl	m = 0	85-89
95	3-CHF ₂ -5-isoxazolyl	CF ₃	3-F	376
96	5-CHF ₂ -3-isoxazolyl	Cl	3-Cl, 4-F	376
97	5-CHCl ₂ -3-isoxazolyl	Cl	m = 0	88-91
98	3-CHF ₂ -5-isoxazolyl	CF ₃	m = 0	358
99	5-(C≡CCF ₃)-3-isoxazolyl	Cl	m = 0	63-65
100	3-CHF ₂ -1,2,4-oxadiazol-5-yl	Cl	m = 0	107-109
101	3-CHF ₂ -5-isoxazolyl	Cl	3-CH ₃	338
102	3-CHF ₂ -5-isoxazolyl	Br	3-CH ₃	383
103	3-CHF ₂ -5-isoxazolyl	Cl	3-OMe	354
104	3-CF ₃ -5-isoxazolyl	Cl	3-OMe	372
105	5-CF ₃ -3-isoxazolyl	Cl	3-OMe	372
106	5-CH ₃ -1,3,4-oxadiazol-2-yl	Cl	m = 0	289
107	3-CHF ₂ -5-isoxazolyl	Cl	3,5-di-F	358
108	3-CH(OEt) ₂ -5-isoxazolyl	Cl	m = 0	398 ^c
109	5-CHF ₂ -3-isoxazolyl	Cl	3-OMe	354
110	3-CH ₃ -5-isoxazolyl	Cl	3-OMe	318
111	3-CH ₃ -5-isoxazolyl	F	3-OMe	302
112	3-thienyl	Cl	3-CN	103-105
113	5-CHF ₂ -3-isoxazolyl	Cl	3,4-di-F	102-105
114	5-CHF ₂ -3-isoxazolyl	Cl	3-Br, 4-F	420
115	1-CH ₃ -1 <i>H</i> -1,2,4-triazol-3-yl	Cl	m = 0	119-122
116	5-CHClF-3-isoxazolyl	Cl	m = 0	108-112
117	1-CH ₃ -1 <i>H</i> -1,2,4-triazol-5-yl	Cl	m = 0	134-138
118	5-CHF ₂ -3-isoxazolyl	Br	3-F	386
119	3-Br-5-isoxazolyl	Br	m = 0	398
120	3-CHF ₂ -5-isoxazolyl	Cl	3-Cl	359
121	3-CHF ₂ -5-isoxazolyl	Br	3-Cl	403
122	5-(<i>c</i> -Pr)-1,3,4-oxadiazol-2-yl	Cl	m = 0	315
123	1-(<i>i</i> -Pr)-1 <i>H</i> -1,2,4-triazol-5-yl	Cl	m = 0	316

No.	Q	R ²	(R ³) _m	M.S.(AP+)
				or m.p.
124	3-CF ₃ -5-isoxazolyl	Cl	5-F	360
125	3-CF ₃ -5-isoxazolyl	Br	5-F	405
126	3-CF ₃ -5-isoxazolyl	Cl	3-Cl	377
127	3-CF ₃ -5-isoxazolyl	Br	3-Cl	421
128	3-(CH ₂ OCH ₂ CF ₃)-5-isoxazolyl	Cl	m = 0	386
129	3-(CH ₂ OCH ₂ CF ₃)-5-isoxazolyl	Br	m = 0	430 ^a
130	5-(<i>c</i> -Pr)-3-isoxazolyl	Cl	3-F	332
131	5-CHF ₂ -3-isoxazolyl	Cl	3-OCHF ₂	390
132	3-CHF ₂ -5-isoxazolyl	Cl	3-OCHF ₂	390
133	5-CHFCF ₃ -3-isoxazolyl	Cl	m = 0	374
134	3,5-di-Me-4-isoxazolyl	Cl	3-CN	*
135	5-Cl-2-pyridinyl	Cl	3-Br	*
136	2-Me-5-CF ₃ -2 <i>H</i> -pyrazol-3-yl	Cl	3-CN	380
137	4-CF ₃ -2-thiazolyl	Cl	m = 0	358
138	4-CF ₃ -2-thiazolyl	I	m = 0	450
139	4-CF ₃ -2-thiazolyl	Br	m = 0	403
140	5-CHO-3-furanyl	Cl	m = 0	*
141	5-CHF ₂ -3-furanyl	Cl	m = 0	*
142	5-CF ₂ CF ₃ -3-isoxazolyl	Cl	m = 0	392
143	5-CF ₂ Cl-3-isoxazolyl	Cl	m = 0	358
144	3-CHF ₂ -5-isoxazolyl	Cl	3-F	342
145	3-CHF ₂ -5-isoxazolyl	Br	3-F	387
146	1-Me-5-CF ₃ -1 <i>H</i> -pyrazol-3-yl	Cl	3-CN	380
147	1-Me-5-CF ₃ -1 <i>H</i> -pyrazol-3-yl	Cl	3-Br	434
148	5-CO ₂ Et-3-isoxazolyl	Cl	m = 0	346
149	5-CF ₂ CH ₃ -3-isoxazolyl	Br	m = 0	*
150	5-C(=O)CH ₃ -3-isoxazolyl	Br	m = 0	360
151	1-Me-1 <i>H</i> -imidazol-2-yl	Cl	m = 0	287
152	1-Me-1 <i>H</i> -imidazol-2-yl	Br	m = 0	332
153	5-CH ₃ -3-isoxazolyl	Cl	m = 0	288
154	5-isoxazolyl	Cl	m = 0	274
155	5-isoxazolyl	Br	m = 0	319
156	5-CF ₃ -3-isoxazolyl	Cl	3-I	*
157	5-CF ₃ -3-isoxazolyl	Cl	3-CN	367
158	4-CF ₃ -2-pyridinyl	Cl	3-CN	377
159	4-CF ₃ -2-pyridinyl	Cl	3-Cl	386

No.	Q	R ²	(R ³) _m	M.S.(AP+)
				or m.p.
160	5-CF ₃ -2-pyridinyl	Cl	3-Cl	386
161	1-CH ₂ CF ₃ -1 <i>H</i> -imidazol-4-yl	Cl	m = 0	*
162	5-CBrF ₂ -3-isoxazolyl	Cl	m = 0	402
163	2-pyrazinyl	Cl	m = 0	79-82
164	2-pyrazinyl	F	m = 0	95-97
165	2-pyrazinyl	Cl	3-F	303
166	2-pyrazinyl	F	3-F	287
167	4-Me-5-CF ₃ -3-isoxazolyl	Cl	m = 0	356
168	5-CF ₃ -3-isoxazolyl	Cl	3-F	360
169	5-CF ₃ -3-isoxazolyl	Cl	3-Cl	376
170	5-CF ₃ -3-isoxazolyl	Cl	3-Br	420
171	3-CH ₃ -5-isothiazolyl	Cl	m = 0	304
172	1,2,4-thiadiazol-5-yl	Cl	m = 0	132-135
173	5-CF ₂ CH ₃ -3-isoxazolyl	CH ₃	m = 0	*
174	4-Cl-2-pyridinyl	Cl	m = 0	319
175	4-F-2-pyridinyl	Cl	m = 0	302
176	3-(OCH ₂ CF ₃)-5-isoxazolyl	Cl	m = 0	372
177	3-Et-5-isoxazolyl	Cl	m = 0	318
178	3-CF ₂ CH ₃ -5-isoxazolyl	Cl	m = 0	338
179	5-CHF ₂ -3-isoxazolyl	Cl	3-Cl	358
180	5-CHF ₂ -3-isoxazolyl	Cl	3-CN	349
181	5-Br-2-thienyl	Cl	m = 0	368
182	2-thienyl	Cl	m = 0	290
183	5-(<i>c</i> -Pr)-3-isoxazolyl	Cl	m = 0	314
184	6-CHF ₂ -4-pyrimidinyl	Cl	m = 0	334
185	3-isoxazolyl	Cl	m = 0	274
186	6-Cl-3-pyridazinyl	Cl	m = 0	319
187	4-Me-2-pyridinyl	Cl	m = 0	298
188	4-CN-2-pyridinyl	Cl	m = 0	309
189	5-Cl-3-pyridazinyl	Cl	m = 0	320
190	6-Cl-4-pyrimidinyl	Cl	m = 0	320
191	5-CH ₂ F-3-isoxazolyl	Cl	3-Br	384
192	5-Cl-3-isothiazolyl	Cl	m = 0	86-88
193	5-Cl-3-isothiazolyl	F	m = 0	100-102
194	5-Cl-3-isothiazolyl	CH ₃	m = 0	78-82
195	5-CHO-3-isoxazolyl	Cl	3-Br	379

No.	Q	R ²	(R ³) _m	M.S.(AP+)
				or m.p.
196	3-C(CH ₃)=CH ₂ -5-isoxazolyl	Cl	m = 0	314
197	3-C(CH ₃)=CH ₂ -5-isoxazolyl	Br	m = 0	359
198	5-CFCl ₂ -3-isoxazolyl	Cl	m = 0	374
199	2-thiazolyl	Cl	m = 0	290
200	5-CF ₃ -3-isoxazolyl	Cl	m = 0	342
201	3-NO ₂ ,5-Cl-2-pyridinyl	Cl	m = 0	363 ^a
202	5-Cl-2-pyridinyl	Cl	3-CN	344
203	5-Cl-2-pyridinyl	Cl	5-Br	398
204	5-Cl-2-pyridinyl	Br	3-CN	388
205	5-Cl-2-pyridinyl	Cl	3-I	445
206	5-CF ₃ -3-isoxazolyl	F	m = 0	326
207	5-Cl-2-pyridinyl	F	3-CN	327
208	5-Cl-2-pyridinyl	Cl	3-&	464
209	4-Br-1 <i>H</i> -pyrazol-1-yl	Cl	6-CN	377
210	3-CHF ₂ -5-isoxazolyl	F	m = 0	308
211	3-CHF ₂ -5-isoxazolyl	CH ₃	m = 0	304
212	3-CHF ₂ -5-isoxazolyl	Br	m = 0	369
213	3-CHF ₂ -5-isoxazolyl	OMe	m = 0	320
214	4-oxazolyl	Cl	m = 0	274
215	5-C(=O)CH ₃ -3-isoxazolyl	Cl	m = 0	316
216	4-CH ₃ -2-thienyl	Cl	m = 0	303
217	1-CH ₂ CF ₃ -1 <i>H</i> -imidazol-4-yl	Cl	3-F	373
218	3-C(=O)CH ₃ -5-isoxazolyl	Cl	m = 0	316
219	4-OMe-2-pyridinyl	Cl	m = 0	314
220	5-CF ₂ CH ₃ -3-isoxazolyl	Cl	m = 0	338
221	5-CFCl ₂ -3-isoxazolyl	Cl	4-F	129-132
222	5-CN-3-isoxazolyl	Cl	3-F	317
223	3-CN-5-isoxazolyl	Cl	m = 0	297
224	3-CH ₂ F-5-isoxazolyl	Cl	m = 0	306
225	3-CO ₂ Et-5-isoxazolyl	Cl	m = 0	346
226	5-CFH ₂ -3-isoxazolyl	Cl	m = 0	320
227	6-OCH ₂ CF ₃ -4-pyrimidinyl	Cl	m = 0	383
228	3-CF ₃ -5-isoxazolyl	Cl	3-F	360
229	3-CH ₃ -5-isoxazolyl	Cl	3-F	306
230	3-CH ₃ -5-isoxazolyl	Br	3-F	351
231	3-thienyl	Cl	m = 0	289

No.	Q	R ²	(R ³) _m	M.S.(AP+)
				or m.p.
232	3-thienyl	Br	m = 0	334
233	5-isothiazolyl	Cl	m = 0	290
234	5-isothiazolyl	Br	m = 0	335
235	5-CO ₂ Me-3-isoxazolyl	Cl	m = 0	332
236	5-CF(CH ₃) ₂ -3-isoxazolyl	Cl	m = 0	334
237	1-Me-5-CF ₃ -1 <i>H</i> -pyrazol-3-yl	Cl	m = 0	355
238	4-CH ₃ -2-thienyl	Cl	3-CN	328
239	5-CHF ₂ -2-furanyl	Cl	m = 0	323

^a ES⁺, ^b AP⁻, ^cM+Na.

* See Index Table B for ¹H NMR data.

& 4-Br-1*H*-pyrazol-1-yl

INDEX TABLE B

Cmpd	¹ H NMR (CDCl ₃ solution unless indicated otherwise) ^Z
1	8.54 (d, 1H), 8.39 (s, 2H), 7.87 (d, 1H), 7.69 (d, 1H), 7.60 (m, 1H), 7.50 (m, 1H), 7.42 (m, 1H), 7.24 (d, 1H)
10	8.43 (s, 2H), 8.03 (m, 2H), 7.36 (m, 2H), 7.26 (m, 1H), 7.19 (m, 1H), 6.56 (s, 1H), 3.85 (s, 3H)
11	8.45 (s, 2H), 7.77 (s, 1H), 7.72 (s, 1H), 7.62 (m, 1H), 7.32 (m, 2H), 7.18 (m, 1H), 3.86 (s, 3H)
16	8.41 (s, 2H), 7.88 (m, 1H), 7.64 (m, 1H), 7.59 (m, 1H), 7.50 (m, 1H), 7.40 (m, 1H), 7.24 (m, 1H), 7.18 (m, 1H)
17	8.49 (s, 2H), 7.88 (m, 1H), 7.64 (m, 1H), 7.58 (m, 1H), 7.50 (m, 1H), 7.41 (m, 1H), 7.25 (m, 1H), 7.17 (m, 1H)
18	8.55 (m, 1H), 8.43 (s, 2H), 8.05 (m, 1H), 7.75 (m, 1H), 7.58 (m, 1H), 7.45 (m, 1H), 7.26 (m, 1H)
19	8.41 (s, 2H), 7.89 (m, 2H), 7.82 (m, 1H), 7.54 (m, 2H), 7.43 (m, 1H), 7.26 (m, 1H)
20	8.82 (m, 1H), 8.52 (s, 2H), 8.06 (m, 1H), 7.96 (m, 1H), 7.61 (m, 1H), 7.48 (m, 1H), 7.30 (m, 1H)
21	8.83 (d, 1H), 8.52 (s, 2H), 8.08 (m, 1H), 7.97 (d, 1H), 7.62 (m, 1H), 7.47 (m, 1H), 7.30 (m, 1H)
23	8.52 (m, 1H), 8.40 (s, 2H), 7.66 (m, 1H), 7.43 (m, 2H), 7.33 (m, 1H), 7.18 (m, 1H)
40	8.26 (s, 2H), 7.39 (m, 2H), 7.32 (m, 2H), 7.27 (m, 1H), 7.24 (s, 1H), 7.17 (m, 1H), 7.02 (d, 1H)
44	8.33 (s, 2H), 7.39 (s, 1H), 7.28–7.34 (m, 2H), 7.25 (m, 2H), 7.12 (d, 1H), 7.07 (m, 1H), 2.43 (s, 3H)
46	(300 MHz) 8.33 (s, 2H), 7.43 (m, 1H), 7.34 (m, 2H), 7.22 (s, 1H), 7.10 (m, 3H)
54	9.42 (s, 1H), 8.43 (s, 2H), 8.12 (s, 1H), 8.05 (d, 1H), 7.72 (m, 1H), 7.53 (m, 1H), 7.32 (m, 1H)
134	8.36 (s, 2 H), 7.74 (m, 1 H), 7.62 (m, 1 H), 7.48 - 7.56 (m, 1 H), 2.33 (s, 3 H), 2.22 (s, 3 H)
135	7.22 (m, 1H), 7.31 (m, 1H), 7.36 (m, 1H), 7.65 (m, 2H), 8.40 (s, 2H), 8.52 (m, 1H)
140	9.60 (s, 1H), 8.44 (s, 2H), 8.05 (s, 1H), 7.59 (d, 1H), 7.54 (s, 1H), 7.43 (t, 1H), 7.37 (t, 1H), 7.23 (d, 1H)
141	8.43 (s, 2H), 7.86 (s, 1H), 7.57 (d, 1H), 7.37 (t, 1H), 7.34 (t, 1H), 7.22 (d, 1H), 6.98 (s, 1H), 6.57 (t, 1H)

- 149 8.54 (2, 2H), 7.96 (dd, 1H), 7.49-7.63 (m, 1H), 7.42 (t, 1H), 7.26-7.29 (m, 1H), 6.86 (t, 1H), 2.00 (t, 3H)
- 156 7.06 - 7.08 (m, 1 H) 7.18 (s, 1 H) 7.93 (s, 1 H) 8.01 - 8.06 (m, 1 H) 8.46 (s, 2 H)
- 161 8.34 (s,2H), 6.59 (s,1H), 6.56 (t,1H), 6.40-6.48 (m,2H), 7.28 (d,1H), 6.92 (s,1H), 6.55 (q,2H)
- 173 8.34 (s, 2H), 7.99 (dd, 1H), 7.51-7.59 (m, 1H), 7.38 (dt, 1H), 7.26-7.31 (m, 1H), 6.90 (t, 1H), 2.25 (s, 3H), 1.99 (t, 3H)

^z ¹H NMR data are in ppm downfield from tetramethylsilane at 500 MHz unless otherwise indicated.

Couplings are designated by (s)-singlet, (d)-doublet and (m)-multiplet.

BIOLOGICAL EXAMPLES OF THE INVENTION

TEST A

- 5 Seeds of plant species selected from downy brome grass (*Bromus tectorum*), cocklebur (common cocklebur, *Xanthium strumarium*), wild oat (*Avena fatua*), barnyardgrass (*Echinochloa crus-galli*), large (Lg) crabgrass (*Digitaria sanguinalis*), giant foxtail (*Setaria faberii*), morningglory (*Ipomoea* spp.), velvetleaf (*Abutilon theophrasti*), and sorghum (*Sorghum vulgare*) were planted into a sandy loam soil and treated preemergence by soil
- 10 drench using test a chemical formulated in a non-phytotoxic solvent mixture which included a surfactant. At the same time these species were also treated postemergence sprayed to runoff using a test chemical formulated in the same manner.

- Plants ranged in height from 2 to 18 cm and were in the one- to two-leaf stage for the postemergence treatment. Treated plants and untreated controls were maintained in a
- 15 greenhouse for approximately 11 days, after which time all treated plants were compared to untreated controls and visually evaluated for injury. Plant response ratings, summarized in Table A, are based on a 0 to 100 scale where 0 is no effect and 100 is complete control. A dash (–) response means no test result.

Table A	Compound	Table A	Compound
1000 g ai/ha	39	2000 g ai/ha	39
Post Sprayed to Runoff		Pre Soil Drench	
Barnyardgrass	50	Barnyardgrass	90
Brome grass, Downy	20	Brome grass, Downy	70
Cocklebur	40	Cocklebur	0
Crabgrass, Large	50	Crabgrass, Large	90
Foxtail, Giant	30	Foxtail, Giant	100
Morningglory	30	Morningglory	0
Oat, Wild	20	Oat, Wild	90
Sorghum	30	Sorghum	50
Velvetleaf	60	Velvetleaf	70

TEST B

Seeds of plant species selected from barnyardgrass (*Echinochloa crus-galli*), kochia (*Kochia scoparia*), ragweed (common ragweed, *Ambrosia elatior*), Italian ryegrass (*Lolium multiflorum*), large (Lg) crabgrass (*Digitaria sanguinalis*), giant foxtail (*Setaria faberii*), morningglory (*Ipomoea* spp.), pigweed (*Amaranthus retroflexus*), velvetleaf (*Abutilon theophrasti*), wheat (*Triticum aestivum*), and corn (*Zea mays*) were planted into a blend of loam soil and sand and treated preemergence with a directed soil spray using test chemicals formulated in a non-phytotoxic solvent mixture which included a surfactant.

At the same time, plants selected from these crop and weed species and also blackgrass (*Alopecurus myosuroides*), and galium (catchweed bedstraw, *Galium aparine*) were planted in pots containing the same blend of loam soil and sand and treated with postemergence applications of test chemicals formulated in the same manner. Plants ranged in height from 2 to 10 cm and were in the one- to two-leaf stage for the postemergence treatment. Treated plants and untreated controls were maintained in a greenhouse for approximately 10 days, after which time all treated plants were compared to untreated controls and visually evaluated for injury. Plant response ratings, summarized in Table B, are based on a 0 to 100 scale where 0 is no effect and 100 is complete control. A dash (–) response means no test result.

Table B	Compounds													
1000 g ai/ha	1	2	3	5	6	7	8	9	10	12	23	31	33	34
Postemergence														
Barnyardgrass	70	0	50	20	0	10	0	10	10	40	50	10	0	0
Blackgrass	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Corn	10	0	10	0	0	10	0	10	0	0	20	0	0	0
Crabgrass, Large	70	0	10	0	20	20	10	50	10	30	60	50	10	10
Foxtail, Giant	80	0	20	0	0	20	0	20	10	40	70	30	10	0
Galium	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Kochia	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Morningglory	60	40	60	20	20	40	10	50	10	30	70	30	10	0
Pigweed	100	70	70	70	70	20	70	60	50	100	100	30	10	40
Ragweed	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ryegrass, Italian	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Velvetleaf	100	50	100	30	30	10	30	40	20	90	70	30	0	20
Wheat	0	0	10	0	0	10	0	10	0	10	0	0	0	0

5	Barnyardgrass	100	90	30	90	100	100	10	10	30	10	0	100	100	90
	Blackgrass	-	-	-	-	100	90	-	10	80	0	20	100	90	90
	Corn	90	50	20	20	100	90	0	0	10	10	0	60	30	10
	Crabgrass, Large	100	80	50	100	-	-	0	-	-	-	-	-	-	-
	Foxtail, Giant	90	90	50	100	100	100	0	10	100	10	30	100	90	90
10	Galium	-	-	-	-	100	90	-	0	100	10	20	100	100	100
	Kochia	-	-	-	-	100	90	-	0	100	30	30	100	100	100
	Morningglory	100	20	100	70	-	-	10	-	-	-	-	-	-	-
	Pigweed	100	90	100	100	100	100	60	0	100	80	20	100	100	100
	Ragweed	-	-	-	-	100	100	-	0	30	10	50	90	60	90
10	Ryegrass, Italian	-	-	-	-	100	80	-	0	10	0	0	100	50	30
	Velvetleaf	100	100	100	100	-	-	70	-	-	-	-	-	-	-
	Wheat	80	50	10	20	100	90	0	0	30	0	0	30	30	20

Table B	Compounds			Table B	Compounds		
1000 g ai/ha	194	195	235	1000 g ai/ha	194	195	235
Postemergence				Postemergence			
Barnyardgrass	100	10	0	Morningglory	-	-	-
Blackgrass	90	0	0	Pigweed	90	50	0
Corn	60	10	0	Ragweed	80	10	0
Crabgrass, Large	-	-	-	Ryegrass, Italian	50	10	0
Foxtail, Giant	90	10	0	Velvetleaf	-	-	-
Galium	100	10	0	Wheat	20	0	0
Kochia	100	10	0				

[illegible]

[illegible]

5	Barnyardgrass	90	40	90	10	0	0	0	0	70	20	60	100	20	80	
	Blackgrass	-	-	-	0	0	0	0	0	90	40	90	90	30	90	
	Corn	80	30	70	20	0	0	0	0	40	40	40	90	30	90	
	Crabgrass, Large	100	50	90	-	-	-	-	-	-	-	-	-	-	-	
	Foxtail, Giant	90	60	80	20	0	0	0	0	90	90	80	100	70	100	
	Galium	-	-	-	20	0	0	0	0	100	100	100	100	100	100	
	Kochia	-	-	-	60	0	0	0	0	90	90	100	90	80	90	
	Morningglory	90	50	90	-	-	-	-	-	-	-	-	-	-	-	
	Pigweed	100	100	100	90	0	0	0	0	100	100	100	100	100	100	
	10	Ragweed	-	-	-	0	0	0	0	0	100	40	50	30	0	30
Ryegrass, Italian	-	-	-	0	0	0	0	0	60	70	100	70	0	90		
Velvetleaf	90	80	100	-	-	-	-	-	-	-	-	-	-	-		
Wheat	30	10	40	0	0	0	0	0	50	30	70	70	20	80		
15	Table B	Compounds														
	500 g ai/ha	161	167	168	169	170	179	180	181	182	183	184	185	190	199	
	Postemergence															
	Barnyardgrass	0	10	100	30	20	80	90	20	10	70	50	40	0	0	
	Blackgrass	0	10	90	90	60	100	90	10	10	90	60	50	0	-	
	Corn	0	20	100	20	20	40	70	10	20	30	20	10	0	0	
	20	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	10	
	Foxtail, Giant	0	10	100	60	50	80	100	20	50	80	70	70	0	0	
	Galium	0	60	100	100	100	100	100	50	50	100	100	70	10	-	
	Kochia	0	60	90	90	100	100	100	50	80	100	100	100	0	-	
25	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	10	
	Pigweed	0	70	100	100	100	100	100	90	80	100	100	100	20	30	
	Ragweed	0	0	90	30	50	30	70	10	10	30	40	100	10	-	
	Ryegrass, Italian	0	0	90	50	20	100	100	0	0	70	20	0	0	-	
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	10	
	Wheat	0	20	100	0	30	90	90	0	0	60	10	0	0	0	
	30	Table B	Compounds													
		500 g ai/ha	200	201	202	203	204	205	206	207	208	209	215	217		
		Postemergence														
		Barnyardgrass	100	0	40	0	10	10	10	70	20	0	90	0		
Blackgrass		-	-	-	-	-	-	-	-	-	-	40	0			
35		Corn	90	0	30	0	0	10	0	0	0	0	40	0		
Crabgrass, Large		100	0	70	0	40	20	20	30	0	30	-	-			
Foxtail, Giant		100	0	80	0	30	20	20	30	0	30	80	0			

126

	Galium	-	-	-	-	-	-	-	-	-	-	70	30	
	Kochia	-	-	-	-	-	-	-	-	-	-	10	30	
	Morningglory	100	0	40	0	10	10	30	30	0	30	-	-	
	Pigweed	100	0	100	0	90	90	70	100	60	100	100	60	
5	Ragweed	-	-	-	-	-	-	-	-	-	-	20	30	
	Ryegrass, Italian	-	-	-	-	-	-	-	-	-	-	50	0	
	Velvetleaf	100	0	30	0	10	10	60	60	30	30	-	-	
	Wheat	100	0	0	0	0	0	0	0	0	0	20	0	
	Table B	Compounds												
10	125 g ai/ha	4	10	11	13	14	15	16	17	18	19	20	21	22 23
	Postemergence													
	Barnyardgrass	0	0	0	0	0	0	0	0	0	0	0	0	0 0
	Blackgrass	-	-	-	-	-	-	-	-	-	-	-	-	-
	Corn	-	0	0	0	0	0	0	0	0	0	0	0	0 10
15	Crabgrass, Large	20	0	10	0	0	10	0	0	0	0	0	0	0 10
	Foxtail, Giant	0	0	0	0	0	0	0	0	0	0	0	0	0 10
	Galium	-	-	-	-	-	-	-	-	-	-	-	-	-
	Kochia	-	-	-	-	-	-	-	-	-	-	-	-	-
	Morningglory	0	20	10	0	0	10	-	-	0	0	0	20	0 20
20	Pigweed	30	0	10	0	0	70	0	0	0	0	40	80	0 70
	Ragweed	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ryegrass, Italian	-	-	-	-	-	-	-	-	-	-	-	-	-
	Velvetleaf	30	0	10	30	20	0	0	0	0	0	0	20	50 40
	Wheat	0	0	0	0	0	0	0	0	0	0	0	0	0 0
25	Table B	Compounds												
	125 g ai/ha	24	25	26	27	28	29	30	32	38	41	42	43	44 45
	Postemergence													
	Barnyardgrass	0	0	0	20	0	0	0	70	0	0	20	0	10 0
	Blackgrass	-	-	-	-	-	-	-	-	0	-	-	-	-
30	Corn	0	10	10	0	0	0	0	10	0	0	20	0	0 0
	Crabgrass, Large	80	10	10	20	20	0	0	60	-	10	30	0	10 10
	Foxtail, Giant	30	0	0	20	20	0	0	70	0	10	10	0	0 10
	Galium	-	-	-	-	-	-	-	-	30	-	-	-	-
	Kochia	-	-	-	-	-	-	-	-	30	-	-	-	-
35	Morningglory	50	10	10	10	0	0	0	70	-	0	20	0	20 30
	Pigweed	20	30	30	50	30	0	0	100	20	40	60	30	20 10
	Ragweed	-	-	-	-	-	-	-	-	0	-	-	-	-

	Ryegrass, Italian	-	-	-	-	-	-	-	-	0	-	-	-	-	-
	Velvetleaf	40	20	10	70	40	0	10	90	-	10	100	20	30	50
	Wheat	0	0	0	0	0	0	0	30	0	0	0	0	0	0
	Table B	Compounds													
5	125 g ai/ha	47	48	49	50	51	52	53	54	58	59	60	62	63	64
	Postemergence														
	Barnyardgrass	0	0	0	0	0	0	100	20	70	10	50	90	80	90
	Blackgrass	0	40	0	40	0	0	100	70	70	0	50	-	-	-
	Corn	0	0	0	0	0	0	10	0	0	0	30	10	20	20
10	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	70	80	60
	Foxtail, Giant	0	0	0	0	10	0	100	80	70	10	50	80	70	60
	Galium	30	70	20	90	60	40	100	100	100	40	100	-	-	-
	Kochia	60	50	30	60	70	20	100	100	100	0	90	-	-	-
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	90	100	70
15	Pigweed	50	70	10	70	70	30	100	100	100	40	100	100	100	100
	Ragweed	20	50	30	20	20	20	60	50	100	0	60	-	-	-
	Ryegrass, Italian	0	0	0	0	0	0	80	30	20	0	0	-	-	-
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	100	100	100
	Wheat	0	0	10	0	0	0	40	0	0	0	30	0	0	0
20	Table B	Compounds													
	125 g ai/ha	70	73	89	90	91	96	97	103	104	105	106	107	108	109
	Postemergence														
	Barnyardgrass	0	0	30	0	0	30	20	40	60	70	0	0	0	90
	Blackgrass	0	0	20	0	0	70	0	90	100	100	0	0	0	90
25	Corn	0	0	30	0	0	20	20	30	30	20	0	0	10	60
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	0	0	40	0	0	50	30	70	80	80	0	0	0	90
	Galium	0	10	50	30	0	100	50	100	100	100	0	0	50	100
	Kochia	0	0	70	10	0	100	90	30	80	70	0	0	0	30
30	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	20	10	90	10	0	100	90	80	90	100	0	0	0	100
	Ragweed	0	0	10	0	0	60	50	60	10	30	0	0	0	90
	Ryegrass, Italian	0	0	0	0	0	60	0	10	30	10	0	0	0	50
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35	Wheat	0	0	0	0	0	0	0	0	10	10	0	0	0	10
	Table B	Compounds													
	125 g ai/ha	110	111	112	113	114	115	116	117	118	119	120	121	122	123

[illegible]

	Foxtail, Giant	0	0	0	0	80	100	90	20	0	60	10	0	0	20
	Galium	0	0	0	10	100	100	100	50	0	90	10	0	0	90
	Kochia	0	0	0	20	100	100	100	60	0	100	0	0	0	90
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5	Pigweed	0	0	20	20	100	100	100	60	0	100	20	0	0	100
	Ragweed	0	0	0	0	20	60	50	0	0	20	0	0	0	20
	Ryegrass, Italian	0	0	0	0	50	100	80	0	0	20	0	0	0	30
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	0	0	0	0	30	90	70	0	0	10	0	0	0	30
10	Table B	Compounds													
	125 g ai/ha	154	155	156	157	158	159	160	161	162	163	164	165	166	167
	Postemergence														
	Barnyardgrass	0	0	0	20	60	0	0	0	90	0	0	20	0	0
	Blackgrass	0	0	20	70	30	0	50	0	30	20	0	30	0	0
15	Corn	0	0	20	0	30	20	30	0	40	10	0	0	10	0
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	0	0	20	70	50	0	30	0	90	10	0	20	0	0
	Galium	0	0	90	100	30	70	70	0	90	30	30	40	30	10
	Kochia	0	0	90	90	70	20	40	0	100	50	0	40	0	10
20	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	0	0	90	100	80	90	90	0	100	30	30	50	30	10
	Ragweed	0	0	20	10	0	0	0	0	40	30	30	60	30	0
	Ryegrass, Italian	0	0	30	90	0	0	0	0	50	0	0	0	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25	Wheat	0	0	20	50	20	20	20	0	30	0	0	0	0	10
	Table B	Compounds													
	125 g ai/ha	168	169	170	173	174	175	176	177	178	179	180	181	182	183
	Postemergence														
	Barnyardgrass	100	10	0	70	10	10	30	30	40	10	50	10	0	40
30	Blackgrass	90	50	50	70	50	0	30	30	40	80	90	0	0	80
	Corn	100	10	20	40	0	0	30	10	20	10	20	10	10	10
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	100	30	30	90	0	0	50	40	60	40	80	0	0	60
	Galium	100	100	80	90	70	40	90	100	100	100	100	10	30	100
35	Kochia	90	90	70	100	70	30	100	70	100	90	100	30	30	100
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	100	100	100	80	50	60	100	100	100	100	100	50	40	100

130

Ragweed	90	20	10	60	30	30	20	20	10	10	50	10	0	20
Ryegrass, Italian	90	20	0	20	0	0	10	0	30	90	100	0	0	40
Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wheat	100	0	0	0	0	0	0	0	10	40	60	0	0	10

5 Table B

Compounds

125 g ai/ha 184 185 187 188 189 190 196 197 198 199 200 201 202 203

Postemergence

Barnyardgrass	0	10	0	20	0	0	10	0	50	0	90	0	10	0
Blackgrass	10	0	0	20	0	0	0	0	40	-	-	-	-	-
10 Corn	10	0	0	30	0	0	0	0	20	0	50	0	0	0
Crabgrass, Large	-	-	-	-	-	-	-	-	-	0	100	0	10	0
Foxtail, Giant	10	10	0	20	0	0	10	0	90	0	100	0	30	0
Galium	70	40	0	50	0	0	100	50	100	-	-	-	-	-
Kochia	70	90	0	90	0	0	40	0	80	-	-	-	-	-
15 Morningglory	-	-	-	-	-	-	-	-	-	0	90	0	10	0
Pigweed	90	90	0	60	0	0	60	10	100	10	100	0	90	0
Ragweed	20	70	0	20	0	10	20	0	20	-	-	-	-	-
Ryegrass, Italian	0	0	0	0	0	0	0	0	50	-	-	-	-	-
Velvetleaf	-	-	-	-	-	-	-	-	-	0	100	0	10	0
20 Wheat	0	0	0	20	0	0	0	0	10	0	60	0	0	0

Table B

Compounds

125 g ai/ha 204 205 206 207 208 209 210 211 212 213 214 215 216 217

Postemergence

Barnyardgrass	0	0	0	0	0	0	0	10	30	0	0	20	10	0
25 Blackgrass	-	-	-	-	-	-	30	50	50	10	0	0	10	0
Corn	0	0	0	0	0	0	10	10	20	10	0	20	10	0
Crabgrass, Large	10	10	10	0	0	10	-	-	-	-	-	-	-	-
Foxtail, Giant	0	0	0	0	0	0	40	50	70	20	0	20	0	0
Galium	-	-	-	-	-	-	90	100	100	80	10	20	50	10
30 Kochia	-	-	-	-	-	-	80	90	100	100	0	0	20	10
Morningglory	0	0	20	10	0	30	-	-	-	-	-	-	-	-
Pigweed	50	60	40	70	20	90	80	50	100	80	10	20	30	10
Ragweed	-	-	-	-	-	-	30	30	60	60	0	0	0	10
Ryegrass, Italian	-	-	-	-	-	-	0	0	20	0	0	0	0	0
35 Velvetleaf	0	0	30	10	20	30	-	-	-	-	-	-	-	-
Wheat	0	0	0	0	0	0	20	0	10	10	0	0	0	0

Table B		Compounds													
125 g ai/ha		218	219	220	221	222	223	224	225	226	227	228	229	230	231
Postemergence															
5	Barnyardgrass	0	0	40	90	0	10	50	0	70	60	70	70	60	10
	Blackgrass	0	0	90	60	10	0	70	0	70	70	100	80	70	0
	Corn	0	0	20	50	30	0	20	0	30	20	30	30	10	0
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	0	0	80	90	0	10	50	0	70	60	90	80	70	10
10	Galium	10	30	100	100	90	50	100	0	100	80	100	100	100	20
	Kochia	0	0	100	100	30	10	100	0	100	80	100	100	100	40
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	40	10	100	90	100	60	100	0	100	90	100	100	90	40
	Ragweed	0	30	10	50	20	10	100	0	40	30	40	100	100	0
15	Ryegrass, Italian	0	0	40	40	0	0	0	0	50	30	90	20	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	0	0	50	30	20	0	0	0	50	20	50	40	10	0

Table B		Compounds						
125 g ai/ha		232	233	234	236	237	238	239
Postemergence								
20	Barnyardgrass	0	10	0	30	0	30	0
	Blackgrass	0	10	10	40	0	0	0
	Corn	0	10	10	30	0	20	20
	Crabgrass, Large	-	-	-	-	-	-	-
	Foxtail, Giant	0	0	0	30	0	10	0
25	Galium	0	0	0	100	0	40	0
	Kochia	0	0	0	90	0	80	0
	Morningglory	-	-	-	-	-	-	-
	Pigweed	40	20	20	70	0	20	20
	Ragweed	0	0	10	20	0	0	0
30	Ryegrass, Italian	0	0	0	20	0	0	0
	Velvetleaf	-	-	-	-	-	-	-
	Wheat	0	0	0	20	0	0	0

Table B		Compounds													
31 g ai/ha		15	38	70	73	89	90	91	96	97	103	104	105	106	107
Postemergence															
35	Barnyardgrass	0	0	0	0	10	0	0	10	0	0	10	10	0	0
	Blackgrass	-	0	0	0	10	0	0	10	0	30	70	70	0	0

132

	Corn	0	0	0	0	10	0	0	0	0	0	10	10	0	0
	Crabgrass, Large	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	0	0	0	0	10	0	0	0	0	40	20	40	0	0
	Galium	-	20	0	0	10	0	0	100	10	90	90	70	0	0
5	Kochia	-	0	0	0	50	0	0	70	60	0	50	30	0	0
	Morningglory	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	60	0	10	0	60	0	0	100	60	30	90	90	0	0
	Ragweed	-	0	0	0	10	0	0	50	20	20	10	10	0	0
	Ryegrass, Italian	-	0	0	0	0	0	0	30	0	0	0	0	0	0
10	Velvetleaf	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table B

Compounds

	31 g ai/ha	108	109	110	111	112	113	114	115	116	117	118	119	120	121
	Postemergence														
15	Barnyardgrass	0	60	0	0	10	60	10	0	10	0	70	0	50	50
	Blackgrass	0	50	0	0	0	60	10	0	20	0	60	0	70	50
	Corn	0	10	0	0	10	10	30	0	20	0	50	0	30	20
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	0	60	0	0	10	60	20	0	20	0	90	0	40	70
20	Galium	30	50	20	10	30	90	70	0	60	0	100	30	100	100
	Kochia	0	0	10	20	60	100	70	0	90	0	100	0	90	90
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	0	80	10	0	30	100	100	0	100	0	100	40	100	100
	Ragweed	0	60	30	10	20	50	0	0	20	0	30	10	40	50
25	Ryegrass, Italian	0	10	0	0	0	20	10	0	0	0	20	0	30	50
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	0	0	0	0	0	0	10	0	10	0	20	0	0	0

Table B

Compounds

	31 g ai/ha	122	124	125	126	127	128	129	130	131	132	133	140	141	142
30	Postemergence														
	Barnyardgrass	0	0	10	80	30	0	0	40	50	20	20	0	0	0
	Blackgrass	0	0	10	60	30	0	0	50	50	20	30	0	0	0
	Corn	0	0	0	20	20	0	0	20	20	10	10	0	0	0
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35	Foxtail, Giant	0	0	0	60	50	0	0	50	50	30	20	0	0	0
	Galium	10	60	50	90	80	10	10	80	80	80	70	0	0	0
	Kochia	10	40	60	90	90	20	0	90	60	30	90	0	0	0

Table B	Compounds														
31 g ai/ha	143	144	145	146	147	149	150	154	155	162	163	164	165	166	
Postemergence															
Barnyardgrass	10	30	40	0	0	10	0	0	0	10	0	0	0	0	
Blackgrass	30	80	70	0	0	10	0	0	0	30	20	0	20	0	
Corn	40	30	30	0	0	10	0	0	0	20	0	0	0	0	
Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Foxtail, Giant	10	80	70	0	0	10	0	0	0	10	0	0	10	0	
Galium	50	100	100	30	0	90	0	0	0	90	20	10	20	20	
Kochia	100	100	100	50	0	90	0	0	0	90	10	0	20	0	
Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Pigweed	90	100	100	30	0	90	10	0	0	100	20	0	30	20	
Ragweed	0	20	20	0	0	10	0	0	0	0	30	20	30	10	
Ryegrass, Italian	0	20	20	0	0	0	0	0	0	0	0	0	0	0	
Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Wheat	20	30	20	0	0	0	0	0	0	20	0	0	0	0	

[illegible]

	Wheat	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Table B	Compounds													
	31 g ai/ha	212	213	214	216	218	219	220	221	222	223	224	225	226	227
	Postemergence														
5	Barnyardgrass	0	0	0	10	0	0	10	10	0	0	30	0	10	10
	Blackgrass	10	0	0	0	0	0	60	30	0	0	40	0	10	10
	Corn	10	0	0	0	0	0	10	20	0	0	0	0	20	10
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	10	0	0	0	0	0	30	10	0	0	20	0	10	10
10	Galium	60	50	0	20	0	20	90	90	30	30	80	0	60	70
	Kochia	90	60	0	10	0	0	90	80	20	0	70	0	100	40
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	90	30	0	10	0	10	100	90	30	20	70	0	50	60
	Ragweed	20	10	0	0	0	20	0	10	0	0	50	0	10	10
15	Ryegrass, Italian	0	0	0	0	0	0	0	0	0	0	0	0	10	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	0	0	0	0	0	0	10	20	0	0	0	0	0	0
	Table B	Compounds													
	31 g ai/ha	228	229	230	231	232	233	234	236	237	238	239			
20	Postemergence														
	Barnyardgrass	30	20	0	0	0	0	0	0	0	0	0			
	Blackgrass	60	30	20	0	0	0	0	20	0	0	0			
	Corn	20	0	0	0	0	0	0	0	0	0	0			
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-			
25	Foxtail, Giant	40	50	0	0	0	0	0	0	0	0	0			
	Galium	100	80	80	0	0	0	0	30	0	0	0			
	Kochia	100	90	80	20	0	0	0	20	0	20	0			
	Morningglory	-	-	-	-	-	-	-	-	-	-	-			
	Pigweed	100	70	60	20	10	0	0	30	0	10	0			
30	Ragweed	30	100	70	0	0	0	0	0	0	0	0			
	Ryegrass, Italian	20	0	0	0	0	0	0	0	0	0	0			
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-			
	Wheat	0	0	0	0	0	0	0	0	0	0	0			
	Table B	Compounds													
35	1000 g ai/ha	1	2	3	5	6	7	8	9	10	12	23	31	33	34
	Preemergence														
	Barnyardgrass	80	0	60	0	0	10	30	20	0	20	40	0	0	0

135

	Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Crabgrass, Large	100	10	40	0	0	40	80	70	10	80	100	80	80	10
	Foxtail, Giant	100	10	70	0	0	20	60	60	0	90	100	30	10	10
	Kochia	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5	Morningglory	20	0	0	0	0	0	0	0	0	0	0	0	0	0
	Pigweed	100	100	50	90	70	40	100	90	10	90	100	0	40	100
	Ragweed	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ryegrass, Italian	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Velvetleaf	80	20	90	20	20	10	30	10	0	70	80	0	0	10
10	Wheat	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table B

Compounds

	1000 g ai/ha	35	36	40	46	55	57	61	148	171	172	186	191	192	193
	Preemergence														
	Barnyardgrass	100	30	50	80	100	100	20	0	70	0	20	100	100	100
15	Corn	30	0	0	0	-	-	0	-	-	-	-	-	-	-
	Crabgrass, Large	100	100	70	100	-	-	50	-	-	-	-	-	-	-
	Foxtail, Giant	100	90	100	100	100	100	40	0	100	30	100	100	100	100
	Kochia	-	-	-	-	100	100	-	0	90	20	100	100	100	100
	Morningglory	90	10	0	10	-	-	0	-	-	-	-	-	-	-
20	Pigweed	10	100	100	100	100	90	90	0	100	90	100	100	100	100
	Ragweed	-	-	-	-	100	90	-	-	0	0	70	90	20	80
	Ryegrass, Italian	-	-	-	-	100	80	-	0	10	0	0	90	50	0
	Velvetleaf	100	-	90	20	-	-	50	-	-	-	-	-	-	-
	Wheat	60	0	0	10	-	-	0	-	-	-	-	-	-	-

Table B

Compounds

	1000 g ai/ha	194	195	235
	Preemergence			
	Barnyardgrass	100	40	0
	Corn	-	-	-
	Crabgrass, Large	-	-	-
	Foxtail, Giant	100	30	0
	Kochia	100	0	0
	Morningglory	-	-	-

Table B

Compounds

	1000 g ai/ha	194	195	235
	Preemergence			
	Pigweed	100	60	0
	Ragweed	30	10	0
	Ryegrass, Italian	50	0	0
	Velvetleaf	-	-	-
	Wheat	-	-	-

25

Table B

Compounds

	500 g ai/ha	4	10	11	13	14	16	17	18	19	20	21	22	23	24
	Preemergence														
	Barnyardgrass	0	0	0	0	0	0	0	0	0	0	50	0	30	80

136

	Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	40
	Crabgrass, Large	80	0	0	10	20	10	10	20	50	70	100	0	80	100
	Foxtail, Giant	50	0	0	10	30	10	0	10	70	60	100	0	70	90
	Kochia	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5	Morningglory	0	0	-	0	0	-	-	0	0	0	0	0	0	0
	Pigweed	70	30	30	80	100	20	0	30	20	20	90	0	90	40
	Ragweed	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Ryegrass, Italian	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Velvetleaf	30	0	10	10	20	20	10	0	0	20	20	0	50	0
10	Wheat	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Table B	Compounds													
	500 g ai/ha	25	26	27	28	29	30	32	37	41	42	43	44	45	47
	Preemergence														
	Barnyardgrass	70	30	80	60	0	0	90	0	0	70	20	10	0	0
15	Corn	0	0	0	0	0	0	0	-	0	0	0	0	0	-
	Crabgrass, Large	100	100	100	100	30	20	100	-	10	90	30	70	20	-
	Foxtail, Giant	90	90	90	90	10	10	100	0	20	100	70	10	20	30
	Kochia	-	-	-	-	-	-	-	0	-	-	-	-	-	70
	Morningglory	10	0	0	0	0	0	30	-	0	0	10	0	0	-
20	Pigweed	100	100	100	80	50	30	100	40	0	90	0	10	0	60
	Ragweed	-	-	-	-	-	-	-	0	-	-	-	-	-	60
	Ryegrass, Italian	-	-	-	-	-	-	-	0	-	-	-	-	-	0
	Velvetleaf	100	80	60	60	0	0	100	-	0	70	10	20	0	-
	Wheat	0	0	0	0	0	0	10	-	0	0	0	0	0	-
25	Table B	Compounds													
	500 g ai/ha	48	49	50	51	52	53	54	58	59	60	62	63	64	123
	Preemergence														
	Barnyardgrass	40	0	10	10	0	100	100	100	10	100	100	100	100	0
	Corn	-	-	-	-	-	-	-	-	-	-	20	30	20	-
30	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	100	100	100	-
	Foxtail, Giant	50	20	50	60	0	100	100	100	60	100	100	100	100	0
	Kochia	80	70	60	70	30	100	100	100	0	100	-	-	-	0
	Morningglory	-	-	-	-	-	-	-	-	-	-	90	60	60	-
	Pigweed	100	30	100	100	50	100	100	100	30	100	100	100	100	0
35	Ragweed	60	0	50	50	0	50	70	80	0	50	-	-	-	0
	Ryegrass, Italian	50	0	50	0	0	40	60	20	0	30	-	-	-	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	100	90	90	-

	Wheat	-	-	-	-	-	-	-	-	-	-	30	40	30	-
	Table B	Compounds													
	500 g ai/ha	134	135	136	137	138	139	151	152	153	156	157	158	159	160
	Preemergence														
5	Barnyardgrass	90	60	100	10	0	0	0	0	100	20	80	100	60	80
	Corn	60	0	40	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	100	100	100	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	100	100	100	0	0	0	0	0	100	80	100	100	90	100
	Kochia	-	-	-	10	0	0	0	0	100	40	100	100	30	90
10	Morningglory	80	10	90	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	100	100	100	20	0	0	0	0	100	100	100	100	100	100
	Ragweed	-	-	-	0	0	0	0	0	40	0	70	20	0	0
	Ryegrass, Italian	-	-	-	0	0	0	0	0	20	0	90	20	0	0
	Velvetleaf	100	60	100	-	-	-	-	-	-	-	-	-	-	-
15	Wheat	20	0	30	-	-	-	-	-	-	-	-	-	-	-
	Table B	Compounds													
	500 g ai/ha	161	167	168	169	170	179	180	181	182	183	184	185	190	199
	Preemergence														
	Barnyardgrass	0	10	100	30	20	90	80	10	20	80	90	70	0	0
20	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	0
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	10
	Foxtail, Giant	0	30	100	90	90	100	100	40	40	90	90	90	10	0
	Kochia	0	30	100	90	80	100	100	0	30	100	100	100	0	-
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	0
25	Pigweed	0	20	100	100	100	100	100	80	40	100	100	100	10	30
	Ragweed	0	10	80	20	60	40	70	0	10	60	50	90	0	-
	Ryegrass, Italian	0	0	100	20	20	40	70	0	10	40	10	0	0	-
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	10
	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	0
30	Table B	Compounds													
	500 g ai/ha	200	201	202	203	204	205	206	207	208	209	215	217		
	Preemergence														
	Barnyardgrass	100	0	80	0	30	10	30	90	0	10	90	10		
	Corn	100	0	0	0	0	0	0	0	0	0	-	-		
35	Crabgrass, Large	100	0	100	0	100	70	60	100	0	100	-	-		
	Foxtail, Giant	100	0	100	0	100	70	60	100	0	100	90	30		
	Kochia	-	-	-	-	-	-	-	-	-	-	0	0		

[illegible]

	Barnyardgrass	0	0	0	0	0	0	90	60	60	0	80	80	90	60
	Corn	-	-	-	-	-	-	-	-	-	-	-	0	10	0
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	100	100	100
	Foxtail, Giant	10	10	0	0	10	0	90	100	90	10	80	100	100	90
5	Kochia	60	60	0	10	10	0	100	80	80	0	90	-	-	-
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	60	30	0
	Pigweed	30	80	20	60	50	40	100	100	100	0	100	100	100	100
	Ragweed	10	50	0	0	50	0	10	60	70	0	20	-	-	-
	Ryegrass, Italian	0	40	0	20	0	0	10	30	0	0	0	-	-	-
10	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	90	70	80
	Wheat	-	-	-	-	-	-	-	-	-	-	-	20	20	0

Table B

Compounds

	125 g ai/ha	70	73	89	90	91	96	97	103	104	105	106	107	108	109
	Preemergence														
15	Barnyardgrass	0	0	70	0	0	20	0	50	80	90	0	0	0	100
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	0	0	90	0	0	40	30	90	100	100	0	0	0	100
	Kochia	0	0	0	0	0	70	30	10	10	10	0	0	0	30
20	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	0	0	100	20	0	100	50	100	100	100	0	0	0	100
	Ragweed	0	0	0	0	0	0	0	10	0	0	0	0	0	70
	Ryegrass, Italian	0	0	0	0	0	0	0	0	10	20	0	0	0	10
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Table B

Compounds

	125 g ai/ha	110	111	112	113	114	115	116	117	118	119	120	121	122	123
	Preemergence														
30	Barnyardgrass	40	20	30	100	20	0	30	0	100	0	100	90	0	0
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	70	50	70	100	30	0	90	0	100	50	100	100	0	0
	Kochia	0	10	50	100	20	0	60	0	100	10	90	100	10	0
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35	Pigweed	70	80	70	100	80	0	100	0	100	50	100	100	50	0
	Ragweed	0	10	0	70	20	0	20	0	30	0	30	20	0	0
	Ryegrass, Italian	0	0	10	60	20	0	20	0	80	0	90	80	0	0

Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Table B	Compounds													
125 g ai/ha	124	125	126	127	128	129	130	131	132	133	134	135	137	138

5	Preemergence														
	Barnyardgrass	30	50	100	90	0	0	90	100	60	70	70	10	0	0
	Corn	-	-	-	-	-	-	-	-	-	-	20	0	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	100	80	-	-
	Foxtail, Giant	100	90	100	100	10	0	100	100	90	90	90	60	0	0
10	Kochia	60	40	90	90	0	0	70	10	0	90	-	-	0	0
	Morningglory	-	-	-	-	-	-	-	-	-	-	10	0	-	-
	Pigweed	100	80	100	100	40	0	100	100	100	100	100	90	0	0
	Ragweed	10	0	10	30	0	0	20	60	40	10	-	-	0	0
	Ryegrass, Italian	10	10	100	90	0	0	50	90	30	50	-	-	0	0
15	Velvetleaf	-	-	-	-	-	-	-	-	-	-	40	20	-	-
	Wheat	-	-	-	-	-	-	-	-	-	-	20	0	-	-

Table B	Compounds													
125 g ai/ha	139	140	141	142	143	144	145	146	147	149	150	151	152	153

20	Preemergence														
	Barnyardgrass	0	0	0	0	90	100	100	30	0	70	20	0	0	0
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	0	0	0	20	100	100	100	20	0	100	20	0	0	30
	Kochia	0	0	0	0	90	100	100	0	0	80	0	0	0	20
25	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	0	0	0	0	100	100	100	70	0	100	20	0	0	100
	Ragweed	0	0	0	0	30	50	20	0	0	0	0	0	0	0
	Ryegrass, Italian	0	0	0	0	60	90	40	0	0	10	0	0	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Table B	Compounds													
125 g ai/ha	154	155	156	157	158	159	160	161	162	163	164	165	166	167

35	Preemergence														
	Barnyardgrass	0	0	0	20	90	0	0	0	100	0	0	10	0	0
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	0	0	20	80	90	20	50	0	100	0	0	10	0	0

141

	Kochia	0	0	20	100	100	0	30	0	70	0	0	60	0	0
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	0	0	80	100	90	40	90	0	100	30	30	80	60	0
	Ragweed	0	0	0	20	0	0	0	0	20	40	0	50	20	0
5	Ryegrass, Italian	0	0	0	20	0	0	0	0	40	0	0	0	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Table B

Compounds

	125 g ai/ha	168	169	170	173	174	175	176	177	178	179	180	181	182	183
10	Preemergence														
	Barnyardgrass	100	20	0	90	0	0	50	50	70	10	20	0	0	30
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	100	50	60	100	40	0	90	70	90	80	90	0	0	80
15	Kochia	100	70	30	100	0	0	20	20	20	100	100	0	0	80
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	100	100	100	100	80	70	90	90	100	100	100	10	0	90
	Ragweed	70	10	20	70	0	0	0	0	0	10	50	0	0	0
	Ryegrass, Italian	80	0	0	20	0	0	10	0	10	20	20	0	0	20
20	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Table B

Compounds

	125 g ai/ha	184	185	187	188	189	190	196	197	198	199	200	201	202	203
	Preemergence														
25	Barnyardgrass	10	20	0	20	0	0	10	0	40	0	100	0	10	0
	Corn	-	-	-	-	-	-	-	-	-	0	20	0	0	0
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	0	100	0	90	0
	Foxtail, Giant	20	30	0	40	0	0	20	0	90	0	100	0	70	0
	Kochia	20	40	0	30	0	0	0	0	30	-	-	-	-	-
30	Morningglory	-	-	-	-	-	-	-	-	-	0	20	0	0	0
	Pigweed	80	90	0	100	0	0	10	0	100	0	100	0	100	0
	Ragweed	0	10	0	0	0	0	0	0	10	-	-	-	-	-
	Ryegrass, Italian	0	0	0	0	0	0	0	0	20	-	-	-	-	-
	Velvetleaf	-	-	-	-	-	-	-	-	-	0	80	0	0	0
35	Wheat	-	-	-	-	-	-	-	-	-	0	20	0	0	0

Table B

Compounds

	125 g ai/ha	204	205	206	207	208	209	210	211	212	213	214	215	216	217
--	-------------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Preemergence															
5	Barnyardgrass	0	0	0	10	0	0	80	50	60	40	0	40	10	0
	Corn	0	0	0	0	0	0	-	-	-	-	-	-	-	-
	Crabgrass, Large	90	30	10	100	0	70	-	-	-	-	-	-	-	-
	Foxtail, Giant	60	20	10	70	0	30	90	100	100	80	0	20	0	0
	Kochia	-	-	-	-	-	-	90	80	90	50	0	0	0	0
10	Morningglory	0	0	0	0	0	0	-	-	-	-	-	-	-	-
	Pigweed	100	10	50	90	0	90	100	100	100	100	0	20	0	10
	Ragweed	-	-	-	-	-	-	20	30	40	0	0	0	0	0
	Ryegrass, Italian	-	-	-	-	-	-	10	0	40	0	0	0	0	0
	Velvetleaf	0	0	30	0	0	0	-	-	-	-	-	-	-	-
	Wheat	0	0	0	0	0	0	-	-	-	-	-	-	-	-

Table B		Compounds													
125 g ai/ha		218	219	220	221	222	223	224	225	226	227	228	229	230	231
15	Preemergence														
	Barnyardgrass	50	0	100	90	0	0	50	0	100	50	100	90	70	10
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	60	0	100	90	0	0	90	0	100	90	100	90	90	10
20	Kochia	0	0	90	70	0	0	90	0	90	50	100	100	100	0
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	90	0	100	100	90	10	100	0	100	80	100	100	100	20
	Ragweed	0	0	10	0	0	0	-	0	20	0	30	90	80	0
	Ryegrass, Italian	0	0	30	40	0	0	0	0	20	10	70	30	0	0
25	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Table B		Compounds						
125 g ai/ha		232	233	234	236	237	238	239
Preemergence								
30	Barnyardgrass	20	0	0	80	0	40	0
	Corn	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-
	Foxtail, Giant	10	0	0	90	0	20	0
	Kochia	0	0	0	100	0	0	0
35	Morningglory	-	-	-	-	-	-	-
	Pigweed	30	0	0	100	0	10	0
	Ragweed	0	0	-	0	0	0	0

5	Foxtail, Giant	0	20	10	100	100	0	0	90	100	60	70	0	0	0
	Kochia	0	10	0	80	80	0	0	50	0	0	50	0	0	0
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	10	40	0	100	100	0	0	100	100	90	90	0	0	0
	Ragweed	0	0	0	10	10	0	0	10	20	0	0	0	0	0
	Ryegrass, Italian	0	0	0	40	30	0	0	20	20	0	0	0	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10	Table B	Compounds													
	31 g ai/ha	143	144	145	146	147	149	150	154	155	162	163	164	165	166
	Preemergence														
	Barnyardgrass	20	80	70	0	0	10	0	0	0	20	0	0	0	0
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	70	100	90	0	0	70	0	0	0	90	0	0	0	0
	Kochia	30	90	90	0	0	20	0	0	0	70	0	0	0	0
15	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	90	100	100	0	0	80	0	0	0	80	0	0	30	40
	Ragweed	0	0	0	0	0	0	0	0	0	0	0	0	20	0
	Ryegrass, Italian	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Table B	Compounds													
	31 g ai/ha	173	174	175	176	177	178	187	188	189	196	197	198	210	211
25	Preemergence														
	Barnyardgrass	60	0	0	10	0	20	0	0	0	0	0	10	10	10
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	50	0	0	60	10	60	0	0	0	0	0	70	40	10
	Kochia	70	-	0	0	0	10	0	0	0	0	0	0	50	20
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	80	60	0	30	50	90	0	0	0	0	0	30	100	100
30	Ragweed	10	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ryegrass, Italian	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Table B	Compounds													
	31 g ai/ha	173	174	175	176	177	178	187	188	189	196	197	198	210	211
	Preemergence														
	Barnyardgrass	60	0	0	10	0	20	0	0	0	0	0	0	10	10
35	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	50	0	0	60	10	60	0	0	0	0	0	70	40	10
	Kochia	70	-	0	0	0	10	0	0	0	0	0	0	50	20
	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	80	60	0	30	50	90	0	0	0	0	0	30	100	100
	Ragweed	10	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ryegrass, Italian	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table B		Compounds													
31 g ai/ha		212	213	214	216	218	219	220	221	222	223	224	225	226	227
Preemergence															
5	Barnyardgrass	10	0	0	0	0	0	10	20	0	0	10	0	30	10
	Corn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	60	0	0	0	0	0	70	50	0	0	40	0	60	40
	Kochia	40	10	0	0	0	0	50	50	0	0	60	0	70	0
10	Morningglory	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	100	80	0	0	0	0	100	100	20	0	90	0	100	40
	Ragweed	0	0	0	0	0	0	0	0	0	0	-	0	0	0
	Ryegrass, Italian	0	0	0	0	0	0	0	20	0	0	0	0	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wheat		-	-	-	-	-	-	-	-	-	-	-	-	-	-

Table B		Compounds										
31 g ai/ha		228	229	230	231	232	233	234	236	237	238	239
Preemergence												
20	Barnyardgrass	50	50	10	0	0	0	0	10	0	0	0
	Corn	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass, Large	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	90	90	60	0	0	0	0	10	0	0	0
	Kochia	70	80	60	0	0	0	0	30	0	0	0
25	Morningglory	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	100	100	90	0	20	0	0	60	0	0	0
	Ragweed	0	70	20	0	0	0	0	0	0	0	0
	Ryegrass, Italian	30	0	0	0	0	0	0	0	0	0	0
	Velvetleaf	-	-	-	-	-	-	-	-	-	-	-
Wheat		-	-	-	-	-	-	-	-	-	-	-

30 TEST C

Plant species in the flooded paddy test selected from rice (*Oryza sativa*), small-flower umbrella sedge (*Cyperus difformis*), ducksalad (*Heteranthera limosa*), and barnyardgrass (*Echinochloa crus-galli*) were grown to the 2-leaf stage for testing. At time of treatment, test pots were flooded to 3 cm above the soil surface, treated by application of test compounds directly to the paddy water, and then maintained at that water depth for the duration of the test.

Treated plants and controls were maintained in a greenhouse for 13 to 15 days, after which time all species were compared to controls and visually evaluated. Plant response ratings, summarized in Table C, are based on a scale of 0 to 100 where 0 is no effect and 100 is complete control. A dash (–) response means no test result.

5	Table C					Compounds										
	250 g ai/ha	4	10	11	12	13	14	16	17	18	19	20	21	22	23	
	Flood															
	Barnyardgrass	0	0	0	10	0	0	0	0	0	0	0	0	0	0	
	Ducksalad	0	0	0	40	40	40	0	0	0	0	0	0	0	60	
10	Rice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	Sedge, Umbrella	0	0	0	0	0	0	0	0	0	0	0	0	0	40	
	Table C					Compounds										
	250 g ai/ha	24	25	26	27	28	29	30	32	35	38	41	42	43	44	
	Flood															
15	Barnyardgrass	0	0	0	0	0	0	0	0	65	0	0	0	0	0	
	Ducksalad	0	30	0	30	0	0	0	90	100	60	0	0	0	0	
	Rice	0	0	0	0	0	0	0	20	80	0	0	0	0	0	
	Sedge, Umbrella	40	20	0	40	0	0	0	85	100	30	0	0	0	0	
	Table C					Compounds										
20	250 g ai/ha	45	46	47	48	49	50	51	52	53	54	55	57	58	59	
	Flood															
	Barnyardgrass	0	20	0	0	0	0	0	0	20	0	25	0	0	0	
	Ducksalad	0	60	0	0	0	30	30	0	80	60	100	90	80	0	
	Rice	0	35	0	0	0	0	0	0	20	0	25	0	0	0	
25	Sedge, Umbrella	0	50	0	0	0	0	0	0	75	0	90	75	40	0	
	Table C					Compounds										
	250 g ai/ha	60	62	63	64	70	73	89	90	96	97	103	104	105	106	
	Flood															
	Barnyardgrass	20	35	60	35	0	0	0	0	15	0	60	20	45	0	
30	Ducksalad	80	100	100	95	0	0	30	0	90	0	40	75	40	0	
	Rice	30	40	25	0	0	0	0	0	30	0	30	0	35	0	
	Sedge, Umbrella	50	90	85	80	0	0	80	0	100	50	80	80	70	0	
	Table C					Compounds										
	250 g ai/ha	107	108	109	110	111	112	113	114	115	116	117	118	119	120	
35	Flood															
	Barnyardgrass	0	0	30	0	0	0	0	0	0	0	0	50	0	90	

147

Ducksalad	0	0	85	30	0	30	95	75	0	95	0	100	40	100
Rice	0	0	35	0	0	0	15	15	0	30	0	45	0	90
Sedge, Umbrella	0	0	80	60	0	95	95	75	0	90	0	100	40	95

Table C

Compounds

5	250 g ai/ha	121	122	123	124	125	126	127	128	129	130	131	132	133	134
	Flood														
	Barneyardgrass	60	0	0	0	0	95	50	0	0	65	75	30	40	0
	Ducksalad	90	0	0	80	80	100	100	0	0	90	95	40	80	40
	Rice	45	0	0	0	0	75	60	0	0	30	60	20	40	0
10	Sedge, Umbrella	90	0	30	85	90	100	100	0	0	90	95	75	80	75

Table C

Compounds

	250 g ai/ha	135	136	137	138	139	140	141	142	143	144	145	146	147	149
	Flood														
	Barneyardgrass	0	0	0	0	0	0	0	0	30	98	45	0	0	0
15	Ducksalad	0	40	20	20	30	0	30	0	85	90	90	0	0	80
	Rice	0	0	0	0	20	0	0	0	35	40	35	0	0	0
	Sedge, Umbrella	0	40	0	0	0	0	0	0	100	85	95	0	0	65

Table C

Compounds

	250 g ai/ha	150	151	152	153	154	155	156	157	158	159	160	161	162	163
20	Flood														
	Barneyardgrass	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ducksalad	0	0	0	0	0	0	0	40	50	0	0	0	0	0
	Rice	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Sedge, Umbrella	0	0	0	0	0	0	0	40	0	0	0	0	0	0

25 Table C

Compounds

	250 g ai/ha	164	165	166	167	168	169	170	171	173	174	175	176	177	178
	Flood														
	Barneyardgrass	0	0	0	0	90	0	0	0	0	0	0	0	0	40
	Ducksalad	0	0	0	0	90	0	0	0	80	0	0	75	60	90
30	Rice	0	0	0	0	40	0	0	0	0	0	0	0	20	0
	Sedge, Umbrella	0	0	0	0	90	0	0	0	60	0	0	30	40	85

Table C

Compounds

	250 g ai/ha	179	180	181	182	183	184	185	186	187	188	189	190	191	192
	Flood														
35	Barneyardgrass	0	0	0	0	0	0	0	0	0	0	0	0	20	0
	Ducksalad	80	90	0	0	85	0	0	0	50	30	0	0	80	60

148

Rice	40	0	0	0	0	0	0	0	0	0	0	0	0	0
Sedge, Umbrella	80	80	0	0	75	0	0	0	0	0	0	0	85	40

Table C

Compounds

250 g ai/ha	193	194	196	197	198	199	200	201	202	203	204	205	206	207
-------------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

5 Flood

Barnyardgrass	0	0	30	20	40	0	90	0	0	0	0	0	0	0
Ducksalad	65	80	55	0	85	0	100	0	80	0	0	0	70	30
Rice	0	0	0	0	25	0	35	0	0	0	0	0	0	0
Sedge, Umbrella	0	0	70	0	100	80	100	0	80	0	0	0	70	30

10 Table C

Compounds

250 g ai/ha	208	209	210	211	212	213	214	215	216	217	218	219	220	221
-------------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Flood

Barnyardgrass	0	0	0	0	0	0	0	0	0	20	0	0	20	0
Ducksalad	0	0	85	70	75	75	0	0	0	40	0	0	80	100
15 Rice	0	0	0	0	20	0	0	0	0	0	0	0	0	0
Sedge, Umbrella	0	0	40	20	75	0	0	0	0	0	0	0	75	95

Table C

Compounds

250 g ai/ha	222	223	224	225	226	227	228	229	230	231	232	233	234	236
-------------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Flood

20 Barnyardgrass	0	0	0	0	20	0	40	55	40	0	0	0	0	0
Ducksalad	30	65	70	0	85	98	85	90	85	65	0	0	0	75
Rice	0	0	0	0	20	0	0	15	20	0	0	0	0	0
Sedge, Umbrella	0	70	65	0	65	85	90	90	85	70	0	0	0	40

Table C

Compounds

25 250 g ai/ha	237	238	239
----------------	-----	-----	-----

Flood

Barnyardgrass	0	20	0
Ducksalad	0	45	0
Rice	0	15	0

30 Sedge, Umbrella	0	65	0
--------------------	---	----	---

TEST D

Seeds of plant species selected from blackgrass (*Alopecurus myosuroides*), Italian ryegrass (*Lolium multiflorum*), winter wheat (*Triticum aestivum*), galium (catchweed bedstraw, *Galium aparine*), corn (*Zea mays*), large (Lg) crabgrass (*Digitaria sanguinalis*),

35

giant foxtail (*Setaria faberii*), johnsongrass (*Sorghum halepense*), lambsquarters (*Chenopodium album*), morningglory (*Ipomoea coccinea*), yellow nutsedge (*Cyperus esculentus*), pigweed (*Amaranthus retroflexus*), ragweed (common ragweed, *Ambrosia elatior*), soybean (*Glycine max*), barnyardgrass (*Echinochloa crus-galli*), oilseed rape (*Brassica napus*), waterhemp (common waterhemp, *Amaranthus rudis*), and velvetleaf (*Abutilon theophrasti*) were planted into a blend of loam soil and sand and treated preemergence with test chemicals formulated in a non-phytotoxic solvent mixture which included a surfactant.

At the same time, plants selected from these crop and weed species and also kochia (*Kochia scoparia*), wild oat (*Avena fatua*), and chickweed (common chickweed, *Stellaria media*) were planted in pots containing Redi-Earth[®] planting medium (Scotts Company, 14111 Scottslawn Road, Marysville, Ohio 43041) comprising sphagnum peat moss, vermiculite, wetting agent and starter nutrients and treated with postemergence applications of test chemicals formulated in the same manner. Plants ranged in height from 2 to 18 cm (1- to 4-leaf stage) for postemergence treatments.

Plant species in the flooded paddy test consisted of rice (*Oryza sativa*), small-flower umbrella sedge (*Cyperus difformis*), ducksalad (*Heteranthera limosa*), and barnyardgrass (*Echinochloa crus-galli*) grown to the 2-leaf stage for testing. At time of treatment, test pots were flooded to 3 cm above the soil surface, treated by application of test compounds directly to the paddy water, and then maintained at that water depth for the duration of the test.

Treated plants and controls were maintained in a greenhouse for 13 to 15 days, after which time all species were compared to controls and visually evaluated. Plant response ratings, summarized in Table D, are based on a scale of 0 to 100 where 0 is no effect and 100 is complete control. A dash (–) response means no test result.

Table D	Compound	Table D	Compound
250 g ai/ha	62	250 g ai/ha	62
Postemergence		Postemergence	
Barnyardgrass	95	Nutsedge, Yellow	10
Blackgrass	60	Oat, Wild	60
Chickweed	100	Oilseed Rape	100
Corn	10	Pigweed	100
Crabgrass, Large	65	Ragweed	85
Foxtail, Giant	55	Ryegrass, Italian	40
Galium	100	Soybean	98
Johnsongrass	85	Velvetleaf	85
Kochia	100	Waterhemp	100

150

Lambsquarters 100
Morningglory 98

Wheat 40

Table D

Compounds

125 g ai/ha 50 53 54 55 57 58 60 62 63 96 113 116 118 120

Postemergence

5	Barnyardgrass	0	35	10	15	10	5	10	30	10	5	25	30	30	40
	Blackgrass	0	30	5	50	15	5	20	40	25	20	40	35	35	55
	Chickweed	5	90	5	100	75	30	75	98	100	98	95	90	98	100
	Corn	5	0	5	20	10	5	5	15	10	5	25	10	15	15
	Crabgrass, Large	5	30	5	15	10	5	5	50	35	10	25	40	20	65
10	Foxtail, Giant	35	30	50	75	20	30	10	20	20	10	10	60	20	25
	Galium	60	90	75	90	95	55	65	100	70	98	90	90	98	100
	Johnsongrass	0	5	5	15	5	5	5	75	45	10	10	-	45	85
	Kochia	15	100	80	100	100	80	90	100	100	100	100	100	100	100
	Lambsquarters	5	100	70	98	80	90	75	100	90	95	90	100	95	100
15	Morningglory	20	55	85	85	65	65	40	95	70	30	80	30	80	100
	Nutsedge, Yellow	-	-	-	5	5	0	5	5	5	5	15	25	10	15
	Oat, Wild	0	20	5	40	15	0	5	50	35	20	25	15	50	60
	Oilseed Rape	30	85	60	100	90	75	90	98	100	65	90	70	95	95
	Pigweed	30	100	60	100	75	75	70	100	95	98	100	90	98	100
20	Ragweed	15	55	35	90	98	60	70	65	55	40	90	75	85	85
	Ryegrass, Italian	0	5	0	15	0	0	0	30	30	10	5	10	5	50
	Soybean	40	98	80	95	90	75	95	98	98	35	75	95	90	95
	Velvetleaf	40	100	40	85	80	85	80	75	40	35	70	75	85	100
	Waterhemp	40	100	75	100	98	75	70	98	98	100	90	100	98	100
	Wheat	5	0	5	35	20	10	20	10	5	10	5	5	5	0

25 Table D

Compounds

125 g ai/ha 121 130 131 143 144 145 149 153 162 168 169 170 178 179

Postemergence

30	Barnyardgrass	45	25	20	65	15	30	10	5	15	75	5	5	20	5
	Blackgrass	50	30	30	30	75	80	30	0	45	20	25	5	15	20
	Chickweed	100	95	100	80	95	100	60	25	85	100	80	100	80	95
	Corn	25	10	25	10	5	10	5	5	5	20	10	5	10	15
	Crabgrass, Large	55	25	20	35	20	20	15	10	20	40	10	10	15	5
35	Foxtail, Giant	40	25	15	50	25	20	30	5	20	60	55	40	25	30
	Galium	95	100	95	100	95	90	95	80	80	100	75	80	85	75
	Johnsongrass	60	10	45	30	50	40	10	10	20	20	5	5	10	5

5	Kochia	100	100	100	100	100	100	100	60	100	100	95	100	100	100
	Lambsquarters	100	100	100	100	95	95	85	50	95	90	80	80	90	100
	Morningglory	98	85	95	100	-	80	80	25	75	85	65	98	90	90
	Nutsedge, Yellow	20	25	10	5	5	10	5	5	10	15	5	10	5	5
	Oat, Wild	70	60	35	5	10	15	5	0	10	30	10	10	5	20
	Oilseed Rape	95	95	85	65	-	-	85	85	-	100	60	25	95	100
	Pigweed	100	98	100	100	100	98	90	70	100	98	98	98	90	98
	Ragweed	98	40	75	30	75	65	45	55	50	55	35	10	35	60
	Ryegrass, Italian	55	15	25	15	10	10	0	0	5	15	15	5	5	20
	Soybean	95	95	65	60	-	75	65	90	50	90	25	35	45	70
10	Velvetleaf	85	100	85	65	98	80	65	50	-	90	30	25	75	70
	Waterhemp	100	98	98	100	100	100	100	65	100	98	90	85	90	98
	Wheat	0	35	0	5	10	10	10	10	5	35	10	15	10	35
Table D		Compounds													
15	125 g ai/ha	180	183	185	191	192	198	200	212	221	224	228	229	230	
	Postemergence														
	Barnyardgrass	5	5	5	10	5	35	15	25	10	15	35	10	5	
	Blackgrass	5	20	0	5	5	15	90	10	40	0	35	10	10	
20	Chickweed	95	70	5	90	55	95	100	80	100	80	100	85	80	
	Corn	15	20	5	10	10	15	15	15	10	5	10	5	15	
	Crabgrass, Large	5	10	10	20	10	30	35	15	30	10	45	5	10	
	Foxtail, Giant	45	5	5	5	15	25	15	10	35	40	40	15	10	
25	Galium	100	95	5	85	50	98	100	80	100	60	100	80	95	
	Johnsongrass	5	10	5	5	5	35	40	20	15	5	60	25	5	
	Kochia	100	100	30	90	95	100	100	100	100	100	100	98	100	
	Lambsquarters	98	75	20	85	75	75	98	98	98	85	95	85	70	
30	Morningglory	90	90	65	85	45	85	98	98	85	80	75	35	70	
	Nutsedge, Yellow	15	5	30	10	5	10	15	5	10	5	10	10	5	
	Oat, Wild	30	25	5	5	5	10	55	10	35	0	55	10	10	
	Oilseed Rape	90	100	0	90	50	90	100	35	90	98	98	80	75	
35	Pigweed	98	95	50	60	55	100	98	100	100	90	98	80	85	
	Ragweed	65	65	25	65	45	60	40	60	75	70	60	60	55	
	Ryegrass, Italian	20	10	0	5	0	10	35	5	10	0	25	5	5	
	Soybean	85	90	45	95	85	60	98	90	90	75	80	70	85	
35	Velvetleaf	55	55	25	50	55	55	75	90	100	60	80	70	65	
	Waterhemp	98	98	30	65	70	98	100	100	100	85	100	85	80	
	Wheat	30	30	0	0	5	5	20	15	0	15	10	10	15	

	Table D	Compounds														
	62 g ai/ha	35	50	53	54	55	57	58	60	62	63	96	113	116	118	
5	Postemergence															
	Barnyardgrass	20	0	25	5	15	5	0	10	35	10	5	20	20	20	
	Blackgrass	60	0	20	5	30	0	0	5	25	20	10	10	5	15	
	Chickweed	100	5	80	5	98	55	15	60	95	98	90	90	30	90	
	Corn	20	0	0	5	10	5	5	5	10	5	5	20	5	15	
10	Crabgrass, Large	10	0	15	5	10	10	5	5	35	10	5	25	25	15	
	Foxtail, Giant	15	5	25	10	40	20	15	5	25	10	10	10	10	10	
	Galium	85	5	95	55	90	65	60	50	85	70	80	90	70	90	
	Johnsongrass	35	0	5	5	10	5	0	0	25	25	5	5	20	10	
	Kochia	100	5	100	80	100	95	60	55	100	98	85	100	100	95	
15	Lambsquarters	95	50	100	5	98	70	60	55	98	85	90	95	80	90	
	Morningglory	98	10	85	75	80	45	45	25	95	98	40	85	10	85	
	Nutsedge, Yellow	5	-	-	-	5	5	0	0	5	5	5	5	10	10	
	Oat, Wild	45	0	30	5	40	5	0	0	40	25	20	10	10	15	
	Oilseed Rape	98	10	95	50	100	75	60	65	90	98	65	90	55	60	
20	Pigweed	95	5	90	80	98	55	40	55	100	90	98	95	90	95	
	Ragweed	75	5	55	5	65	60	65	40	60	55	10	90	60	55	
	Ryegrass, Italian	20	0	5	0	10	0	0	0	5	5	10	5	5	5	
	Soybean	95	20	95	50	90	85	60	95	95	55	55	65	30	60	
	Velvetleaf	65	0	70	40	75	70	60	65	70	40	35	70	70	70	
25	Waterhemp	98	0	90	80	98	75	55	50	98	98	98	85	90	95	
	Wheat	10	5	0	0	30	15	10	5	20	5	0	5	5	0	
	Table D															
	Compounds															
	62 g ai/ha	120	121	130	131	143	144	145	149	153	156	162	168	169	170	
30	Postemergence															
	Barnyardgrass	25	25	10	10	10	10	10	5	5	5	15	10	5	5	
	Blackgrass	60	30	20	5	20	15	25	0	0	5	30	15	0	5	
	Chickweed	100	100	80	98	60	95	95	30	20	70	60	100	70	90	
	Corn	10	20	10	10	5	10	5	5	5	5	5	10	10	5	
35	Crabgrass, Large	35	40	20	10	15	15	10	10	5	10	10	15	10	5	
	Foxtail, Giant	15	15	10	15	10	25	20	10	5	35	10	40	25	40	
	Galium	100	100	95	95	70	90	80	70	70	70	60	95	70	65	
	Johnsongrass	40	60	5	25	30	10	5	15	5	10	10	5	0	0	
	Kochia	100	100	100	95	100	100	100	100	40	45	100	100	90	85	
	Lambsquarters	100	100	100	98	98	90	85	80	30	40	75	85	35	40	

153

	Morningglory	80	90	85	75	90	85	85	70	15	65	75	85	55	70
	Nutsedge, Yellow	10	5	20	5	5	5	5	5	5	5	5	10	0	5
	Oat, Wild	55	70	35	20	5	5	15	0	0	0	5	20	5	5
	Oilseed Rape	90	95	75	80	60	95	-	60	60	45	40	100	10	40
5	Pigweed	100	100	98	98	100	100	98	95	40	75	100	95	80	98
	Ragweed	70	75	75	65	10	70	50	30	25	15	40	55	25	10
	Ryegrass, Italian	10	40	10	5	0	10	10	0	0	0	5	5	5	5
	Soybean	70	90	98	75	60	95	75	65	70	30	40	40	15	10
	Velvetleaf	100	80	75	80	50	75	75	40	60	25	75	45	20	25
10	Waterhemp	100	100	98	90	100	100	100	95	35	90	100	98	90	100
	Wheat	0	0	30	0	5	10	5	0	5	10	5	30	10	10

Table D

Compounds

	62 g ai/ha	178	179	180	183	185	191	192	198	200	212	221	224	228	229
	Postemergence														
15	Barnyardgrass	10	5	5	5	0	5	5	20	10	15	5	10	20	5
	Blackgrass	10	15	5	5	0	0	0	10	40	0	5	5	25	15
	Chickweed	60	95	85	65	5	70	50	60	100	30	80	40	98	80
	Corn	5	10	10	15	5	10	5	15	10	10	5	5	10	5
	Crabgrass, Large	20	5	5	5	10	10	10	20	25	10	25	10	35	5
20	Foxtail, Giant	5	40	30	5	5	5	25	15	10	15	5	5	20	5
	Galium	98	70	100	95	5	85	50	98	100	65	100	60	98	65
	Johnsongrass	15	5	5	5	5	5	5	35	45	20	15	5	25	0
	Kochia	100	100	100	100	5	85	90	100	100	100	100	95	100	98
	Lambsquarters	75	75	98	65	15	70	40	70	85	98	95	80	90	70
25	Morningglory	100	85	90	90	10	60	10	80	85	75	80	35	80	25
	Nutsedge, Yellow	10	5	10	5	0	5	0	15	5	10	5	5	5	10
	Oat, Wild	5	15	5	10	0	5	0	10	30	5	10	0	40	10
	Oilseed Rape	80	85	80	100	0	60	5	60	100	20	90	50	80	60
	Pigweed	85	98	75	85	35	30	55	90	85	100	98	85	95	70
30	Ragweed	20	50	35	60	15	60	15	55	60	55	45	55	60	70
	Ryegrass, Italian	5	5	5	5	0	0	0	10	35	0	20	0	15	0
	Soybean	35	70	70	75	35	75	55	30	90	85	85	65	80	45
	Velvetleaf	60	30	35	55	10	30	30	55	70	60	80	30	65	45
	Waterhemp	85	95	90	75	25	70	55	95	98	100	100	85	95	80
35	Wheat	5	30	15	30	0	0	5	5	25	5	5	0	10	5

Table D

Compound

62 g ai/ha

230

Table D

Compound

62 g ai/ha

230

Postemergence

Barnyardgrass	10
Blackgrass	10
Chickweed	75
Corn	0
Crabgrass, Large	5
Foxtail, Giant	10
Galium	85
Johnsongrass	5
Kochia	100
Lambsquarters	80
Morningglory	5

Postemergence

Nutsedge, Yellow	5
Oat, Wild	5
Oilseed Rape	75
Pigweed	70
Ragweed	55
Ryegrass, Italian	5
Soybean	65
Velvetleaf	55
Waterhemp	75
Wheat	5

Table D

Compounds

	31 g ai/ha	35	50	53	54	55	57	58	60	62	63	96	113	116	118
Postemergence															
5	Barnyardgrass	20	0	10	5	10	5	0	0	50	10	0	15	5	20
	Blackgrass	40	0	40	5	10	0	0	0	20	10	10	15	5	10
	Chickweed	85	0	80	5	75	50	10	50	70	65	85	85	15	70
	Corn	10	0	0	0	10	5	0	0	10	5	15	10	10	10
	Crabgrass, Large	10	0	20	5	10	5	5	5	55	5	5	20	5	10
10	Foxtail, Giant	10	0	20	30	50	10	0	5	25	5	5	5	5	15
	Galium	75	15	85	55	90	50	45	40	85	70	90	80	60	90
	Johnsongrass	5	0	0	0	5	0	0	0	25	5	0	5	20	5
	Kochia	100	0	100	55	100	90	50	50	98	95	60	100	100	95
	Lambsquarters	85	5	80	5	85	40	75	20	98	70	65	85	60	85
15	Morningglory	85	0	80	75	70	45	45	40	85	50	40	75	10	85
	Nutsedge, Yellow	5	-	-	-	5	5	0	0	5	0	5	5	5	5
	Oat, Wild	40	0	10	0	15	0	0	0	30	20	20	15	5	10
	Oilseed Rape	80	5	80	20	95	55	10	50	80	90	45	50	10	30
	Pigweed	90	5	90	80	85	55	25	35	98	85	95	90	75	95
20	Ragweed	85	5	50	0	40	50	45	30	65	40	10	60	35	45
	Ryegrass, Italian	5	0	5	0	5	0	0	0	5	0	5	5	0	0
	Soybean	80	5	90	10	75	70	40	65	80	45	50	55	55	60
	Velvetleaf	55	0	60	5	50	50	35	50	60	25	35	60	65	75
	Waterhemp	95	0	90	60	90	55	40	30	98	85	95	80	90	90
	Wheat	5	0	0	0	25	5	5	5	30	0	0	0	0	0

Table D

Compounds

	31 g ai/ha	120	121	130	131	143	144	145	149	153	156	157	162	168	169
	Postemergence														
	Barnyardgrass	15	25	10	10	5	5	5	5	5	0	5	10	5	5
5	Blackgrass	35	25	5	5	10	15	20	0	0	5	5	30	15	0
	Chickweed	100	100	80	80	50	80	80	30	20	45	75	60	98	40
	Corn	15	20	25	5	5	5	5	5	0	5	5	5	5	5
	Crabgrass, Large	25	10	20	5	10	10	10	5	5	5	5	5	5	5
	Foxtail, Giant	10	10	5	10	10	10	10	10	0	10	20	10	35	5
10	Galium	98	90	90	85	70	85	80	60	70	60	70	60	95	70
	Johnsongrass	30	35	5	5	5	10	5	10	0	5	0	5	5	0
	Kochia	100	100	95	90	100	100	100	90	50	30	90	100	100	60
	Lambsquarters	100	100	100	98	80	80	85	70	35	35	55	65	85	40
	Morningglory	98	75	80	75	-	85	85	60	5	15	35	65	85	15
15	Nutsedge, Yellow	10	10	5	0	5	5	5	0	0	5	10	0	5	0
	Oat, Wild	40	45	15	15	0	0	10	0	0	0	10	5	10	5
	Oilseed Rape	85	80	80	45	50	80	70	30	40	40	25	5	100	5
	Pigweed	100	98	95	98	90	98	95	85	20	60	65	90	85	80
	Ragweed	55	65	60	45	5	60	50	30	10	10	25	30	50	5
20	Ryegrass, Italian	5	10	5	5	0	5	0	0	0	0	5	0	5	0
	Soybean	70	60	55	15	30	80	60	40	40	25	35	10	75	5
	Velvetleaf	85	80	80	65	30	55	70	30	20	15	40	40	40	10
	Waterhemp	100	95	85	90	100	85	90	80	10	70	75	98	90	90
	Wheat	0	0	5	0	5	10	5	0	0	5	0	0	10	5
25	Table D														
	Compounds														
	31 g ai/ha	170	178	179	180	183	185	191	192	198	200	212	221	224	228
	Postemergence														
	Barnyardgrass	0	15	0	0	5	0	5	5	10	10	5	5	5	20
	Blackgrass	0	5	5	0	5	0	0	0	5	30	0	10	0	15
30	Chickweed	65	50	80	60	60	0	55	50	60	95	30	85	15	85
	Corn	5	5	10	5	15	0	5	0	10	5	5	0	5	10
	Crabgrass, Large	5	10	5	5	5	0	10	5	15	10	10	10	5	20
	Foxtail, Giant	30	5	5	50	5	5	5	25	15	20	10	5	5	10
	Galium	50	80	80	75	65	0	70	50	75	100	60	80	50	98
35	Johnsongrass	0	10	5	0	5	0	5	5	20	20	5	5	5	10
	Kochia	60	95	100	90	95	0	80	80	95	100	100	100	70	100
	Lambsquarters	40	70	75	80	65	10	70	50	55	80	80	95	75	85

156

5	Morningglory	30	100	65	55	75	25	65	0	55	50	98	65	35	65
	Nutsedge, Yellow	5	5	5	0	5	0	5	5	5	5	5	5	0	5
	Oat, Wild	5	0	5	5	5	0	0	0	5	30	0	10	0	20
	Oilseed Rape	5	80	75	60	85	0	40	0	50	85	-	70	10	85
	Pigweed	90	85	90	75	85	35	30	20	98	85	100	95	85	95
10	Ragweed	5	5	25	40	35	20	40	5	55	55	40	50	55	35
	Ryegrass, Italian	0	0	5	0	0	0	0	0	0	25	0	5	0	5
	Soybean	10	35	45	60	55	30	65	30	25	85	70	25	40	55
	Velvetleaf	5	40	20	40	55	10	30	20	45	60	50	65	25	40
	Waterhemp	80	80	90	75	80	25	60	40	85	90	98	85	75	95
	Wheat	20	5	10	5	20	0	0	0	0	10	5	0	0	5

Table D Compounds

31 g ai/ha 229 230

Postemergence

Barnyardgrass 5 5

Blackgrass 10 10

Chickweed 60 65

Corn 20 0

Crabgrass, Large 0 5

Foxtail, Giant 5 5

Galium 70 85

Johnsongrass 0 0

Kochia 90 85

Lambsquarters 70 70

Morningglory 25 5

Table D Compounds

31 g ai/ha 229 230

Postemergence

Nutsedge, Yellow 0 0

Oat, Wild 10 0

Oilseed Rape 70 60

Pigweed 60 65

Ragweed 50 40

Ryegrass, Italian 0 0

Soybean 40 55

Velvetleaf 60 40

Waterhemp 70 70

Wheat 10 5

Table D Compounds

16 g ai/ha 35 50 53 54 55 57 58 60 63 96 113 116 118 120

Postemergence

15 Barnyardgrass 15 0 5 0 5 0 0 0 5 0 10 5 10 10

Blackgrass 35 0 5 0 5 0 0 0 5 0 5 0 5 20

Chickweed 85 0 50 5 70 30 5 45 60 60 70 5 75 95

Corn 5 0 0 0 10 0 0 0 5 5 5 5 5 15

Crabgrass, Large 10 0 10 5 5 5 5 5 5 5 10 5 10 30

20 Foxtail, Giant 5 0 40 0 40 0 0 0 0 0 5 5 5 10

Galium 70 30 60 50 80 40 40 30 60 70 80 60 85 98

Johnsongrass 10 0 0 0 5 0 0 0 5 0 5 10 5 5

Kochia 100 0 90 50 98 75 35 50 90 25 90 100 95 98

157

	Lambsquarters	80	0	70	0	75	60	10	20	10	65	70	40	75	100
	Morningglory	60	0	50	5	70	5	20	0	50	85	85	15	85	85
	Nutsedge, Yellow	5	-	-	-	0	0	0	0	0	0	5	5	5	5
	Oat, Wild	10	0	5	0	5	0	0	5	0	15	10	5	5	35
5	Oilseed Rape	45	5	70	5	95	30	5	60	60	40	80	50	5	80
	Pigweed	80	0	85	60	85	30	20	20	85	98	90	75	80	98
	Ragweed	45	5	50	0	25	20	20	15	10	0	80	40	80	40
	Ryegrass, Italian	0	0	0	0	0	0	0	0	0	0	5	0	0	5
	Soybean	65	5	60	15	65	40	55	65	35	15	80	40	75	90
10	Velvetleaf	50	0	55	0	35	10	60	40	10	20	60	55	65	70
	Waterhemp	75	0	80	45	85	60	35	15	85	85	85	70	85	90
	Wheat	0	0	0	0	5	0	0	0	0	0	0	70	0	0

Table D

Compounds

	16 g ai/ha	121	130	131	143	144	145	149	153	156	157	162	168	169	170
15	Postemergence														
	Barnyardgrass	20	10	10	5	5	5	5	5	0	5	5	5	5	0
	Blackgrass	5	0	5	10	5	15	0	0	0	5	0	10	0	0
	Chickweed	95	80	65	45	70	70	5	20	5	50	50	65	5	30
	Corn	10	10	5	5	5	5	0	0	5	5	5	5	0	5
20	Crabgrass, Large	15	10	5	10	5	5	5	0	5	5	5	5	5	5
	Foxtail, Giant	10	5	10	5	5	5	5	0	5	40	10	35	0	5
	Galium	80	65	80	50	80	80	60	70	20	60	60	70	55	50
	Johnsongrass	35	0	5	0	5	5	5	0	0	0	5	5	0	0
	Kochia	98	90	85	100	100	100	90	50	10	80	100	100	40	5
25	Lambsquarters	98	85	95	80	85	70	55	15	25	25	50	55	5	10
	Morningglory	65	55	70	95	85	85	25	5	5	20	45	40	15	30
	Nutsedge, Yellow	5	0	0	0	5	5	0	0	0	5	0	5	0	0
	Oat, Wild	25	5	10	0	0	5	0	0	0	0	5	10	0	0
	Oilseed Rape	70	60	60	50	25	60	20	50	5	5	5	85	0	5
30	Pigweed	98	85	85	75	85	90	85	20	65	65	75	60	75	60
	Ragweed	20	70	30	0	25	10	5	10	5	20	20	20	5	5
	Ryegrass, Italian	0	0	0	0	0	0	0	0	0	0	0	5	0	0
	Soybean	55	50	30	30	60	50	30	35	15	10	25	45	5	10
	Velvetleaf	75	80	45	30	35	60	10	5	10	10	20	40	10	5
35	Waterhemp	95	75	85	85	85	85	80	5	35	75	95	80	55	65
	Wheat	0	5	0	0	5	5	0	0	0	0	0	10	0	0

Table D

Compounds

16 g ai/ha	178	179	180	183	185	191	192	198	200	212	220	221	224	228
Postemergence														
Barnyardgrass	10	0	0	5	0	5	5	10	5	5	10	0	5	15
5 Blackgrass	0	5	0	0	0	0	0	5	30	0	5	5	0	15
Chickweed	50	70	60	30	0	50	0	5	60	10	40	75	10	85
Corn	0	5	5	10	0	5	0	5	5	5	5	0	0	5
Crabgrass, Large	5	5	5	5	0	5	5	10	10	5	15	10	5	15
Foxtail, Giant	0	5	60	5	35	5	10	10	5	10	5	35	0	10
10 Galium	75	70	65	65	0	40	35	70	75	60	60	80	40	85
Johnsongrass	10	0	0	0	0	5	0	5	15	5	5	0	5	10
Kochia	90	90	85	90	0	50	50	60	95	90	100	98	35	100
Lambsquarters	25	70	70	35	10	50	35	10	70	40	70	85	55	85
Morningglory	70	30	40	65	0	65	0	55	55	75	75	65	15	65
15 Nutsedge, Yellow	5	5	0	5	0	0	0	5	5	10	5	5	0	5
Oat, Wild	0	0	0	0	0	0	0	5	20	0	5	5	0	10
Oilseed Rape	35	40	5	5	0	0	0	30	70	10	10	70	0	65
Pigweed	80	80	70	55	10	35	15	85	90	100	85	95	85	85
Ragweed	5	30	30	15	10	30	0	30	30	35	25	35	30	20
20 Ryegrass, Italian	0	0	0	0	0	0	0	0	20	0	0	5	0	0
Soybean	15	25	45	65	15	45	10	10	60	60	10	35	35	60
Velvetleaf	60	15	45	30	5	15	20	50	30	20	25	70	10	35
Waterhemp	70	60	40	80	5	25	20	85	80	80	80	70	70	90
Wheat	0	5	5	5	0	0	0	0	5	0	0	0	0	0

Table D

Compounds

16 g ai/ha

229 230

Postemergence

Barnyardgrass

5 5

Blackgrass

0 0

Chickweed

60 65

Corn

5 0

Crabgrass, Large

0 0

Foxtail, Giant

0 5

Galium

70 80

Johnsongrass

0 0

Kochia

85 65

Lambsquarters

60 55

Table D

Compounds

8 g ai/ha

35 156 157 220

Postemergence

Barnyardgrass

10 0 0 10

Blackgrass

30 0 0 0

Chickweed

50 5 10 40

Corn

5 0 0 5

Crabgrass, Large

5 0 5 5

Foxtail, Giant

5 0 20 5

Galium

80 20 50 55

Johnsongrass

0 0 0 0

Kochia

95 0 20 100

Lambsquarters

75 10 25 35

Morningglory	40	5
Nutsedge, Yellow	0	0
Oat, Wild	0	0
Oilseed Rape	45	50
Pigweed	55	60
Ragweed	35	30
Ryegrass, Italian	0	0
Soybean	20	50
Velvetleaf	15	25
Waterhemp	60	65
Wheat	0	5

Morningglory	25	5	5	25
Nutsedge, Yellow	5	0	5	0
Oat, Wild	5	0	0	0
Oilseed Rape	5	5	5	5
Pigweed	80	15	70	80
Ragweed	40	0	15	15
Ryegrass, Italian	0	0	0	0
Soybean	40	5	10	10
Velvetleaf	30	0	5	10
Waterhemp	75	5	55	70
Wheat	0	0	0	0

Table D	Compounds	
4 g ai/ha	157	220
Postemergence		
Barnyardgrass	0	5
Blackgrass	0	0
Chickweed	5	10
Corn	0	5
Crabgrass, Large	0	5
Foxtail, Giant	0	5
Galium	10	50
Johnsongrass	0	0
Kochia	0	50
Lambsquarters	20	45
Morningglory	0	15
Nutsedge, Yellow	0	0
Oat, Wild	0	0
Oilseed Rape	0	5
Pigweed	35	75
Ragweed	50	15
Ryegrass, Italian	0	0
Soybean	5	10
Velvetleaf	5	15
Waterhemp	20	60
Wheat	0	0

Table D	Compound
2 g ai/ha	220
Postemergence	
Barnyardgrass	5
Blackgrass	0
Chickweed	5
Corn	0
Crabgrass, Large	5
Foxtail, Giant	0
Galium	50
Johnsongrass	0
Kochia	20
Lambsquarters	55
Morningglory	35
Nutsedge, Yellow	0
Oat, Wild	0
Oilseed Rape	0
Pigweed	70
Ragweed	5
Ryegrass, Italian	0
Soybean	15
Velvetleaf	5
Waterhemp	15
Wheat	0

Table D	Compound
250 g ai/ha	62
Preemergence	
Barnyardgrass	100
Blackgrass	90
Corn	70
Crabgrass, Large	100
Foxtail, Giant	100
Galium	100
Johnsongrass	98
Lambsquarters	100
Morningglory	100

Table D	Compound
250 g ai/ha	62
Preemergence	
Nutsedge, Yellow	60
Oilseed Rape	100
Pigweed	100
Ragweed	85
Ryegrass, Italian	95
Soybean	95
Velvetleaf	100
Waterhemp	100
Wheat	50

	Table D	Compounds													
	125 g ai/ha	53	54	55	57	58	60	62	63	104	113	118	120	131	144
	Preemergence														
5	Barnyardgrass	95	95	100	85	80	75	100	100	70	100	100	100	100	100
	Blackgrass	95	70	90	60	95	70	90	90	30	80	90	90	95	95
	Corn	5	0	80	30	20	5	60	15	0	40	30	75	65	65
	Crabgrass, Large	100	100	100	100	100	100	100	100	100	100	100	100	100	100
	Foxtail, Giant	100	95	100	100	98	95	100	100	100	100	100	100	100	100
10	Galium	98	85	100	90	100	98	100	98	70	90	95	90	90	98
	Johnsongrass	55	40	100	80	50	45	90	85	65	100	100	98	85	100
	Lambsquarters	100	85	98	100	100	100	98	95	90	98	98	100	98	100
	Morningglory	40	15	100	45	45	20	98	60	25	95	90	75	85	100
	Nutsedge, Yellow	5	15	70	20	15	5	10	5	0	60	65	45	70	85
15	Oilseed Rape	100	100	100	100	100	100	100	98	50	100	100	100	95	100
	Pigweed	100	100	100	100	100	100	100	100	100	100	100	100	100	100
	Ragweed	85	35	95	100	100	90	75	70	60	90	85	100	25	100
	Ryegrass, Italian	60	30	95	50	5	30	90	90	5	70	60	90	90	98
	Soybean	80	20	98	95	85	95	85	60	25	80	95	50	85	95
20	Velvetleaf	100	100	100	100	100	100	100	100	65	100	100	100	98	100
	Waterhemp	100	100	100	100	100	100	100	100	100	100	100	100	100	100
	Wheat	5	0	60	5	0	0	40	45	5	20	5	5	5	25

	Table D	Compounds						
	125 g ai/ha	145	158	168	179	180	183	200 229
25	Preemergence							

5	Barnyardgrass	100	20	100	35	80	90	100	100						
	Blackgrass	95	90	90	80	70	50	90	85						
	Corn	45	15	55	5	30	5	20	40						
	Crabgrass, Large	100	100	100	100	100	100	100	100						
	Foxtail, Giant	100	100	100	85	100	100	100	98						
	Galium	100	100	100	98	98	95	95	98						
	Johnsongrass	95	40	100	65	90	45	100	70						
	Lambsquarters	100	80	98	100	90	100	100	95						
	Morningglory	100	70	100	15	100	40	90	45						
10	Nutsedge, Yellow	45	5	45	25	30	5	15	45						
	Oilseed Rape	100	98	100	100	100	100	100	100						
	Pigweed	100	100	100	100	100	100	100	100						
	Ragweed	100	50	100	60	55	10	80	95						
	Ryegrass, Italian	85	20	95	80	65	50	90	60						
	Soybean	90	95	85	45	80	80	85	95						
	Velvetleaf	100	70	100	100	100	100	100	100						
	Waterhemp	100	100	100	100	100	100	100	98						
	Wheat	10	0	85	35	30	0	40	15						
20	Table D														
	62 g ai/ha	32	35	53	54	55	57	58	60	62	63	104	113	118	120
	Preemergence														
	Barnyardgrass	98	100	40	10	100	75	65	30	98	98	25	98	100	90
	Blackgrass	35	90	90	40	90	50	80	10	90	90	0	70	60	90
	Corn	0	15	100	0	35	5	10	0	30	15	0	25	30	25
	Crabgrass, Large	100	100	98	90	100	100	100	100	100	98	98	100	100	100
	Foxtail, Giant	95	100	98	95	100	100	85	40	100	98	90	100	100	98
	Galium	75	100	100	60	100	90	100	100	100	98	75	98	95	90
25	Johnsongrass	85	98	35	10	85	40	20	15	75	70	15	90	85	95
	Lambsquarters	100	100	100	85	100	100	100	85	95	90	90	98	95	90
	Morningglory	0	90	45	5	100	35	0	10	55	25	25	85	85	60
	Nutsedge, Yellow	25	5	0	0	55	20	0	0	5	0	0	15	15	35
	Oilseed Rape	100	100	100	100	100	100	100	90	98	98	30	100	100	95
	Pigweed	100	100	100	100	100	100	100	100	100	100	100	100	100	100
	Ragweed	70	80	55	10	90	100	75	65	65	55	55	85	70	25
	Ryegrass, Italian	10	80	15	5	85	5	0	0	60	80	0	60	65	90
	Soybean	65	85	55	20	95	95	80	65	70	45	25	90	80	60
35	Velvetleaf	85	100	100	85	100	100	100	100	100	100	70	100	100	100

Waterhemp	98	100	100	100	100	100	100	100	100	100	100	100	100	100
Wheat	5	15	0	0	25	0	0	0	10	10	5	10	0	5

Table D

Compounds

62 g ai/ha 131 143 144 145 158 168 179 180 183 200 229

5	Preemergence													
	Barnyardgrass	85	98	100	85	35	100	15	35	30	100	65		
	Blackgrass	90	90	95	95	50	90	45	60	0	90	70		
	Corn	60	100	15	10	5	25	0	5	0	10	5		
	Crabgrass, Large	100	100	100	100	85	100	75	100	98	100	100		
10	Foxtail, Giant	100	100	100	100	85	100	85	85	80	100	90		
	Galium	90	80	98	100	100	100	90	98	95	100	98		
	Johnsongrass	80	80	100	90	10	85	20	60	20	85	45		
	Lambsquarters	85	85	100	100	100	100	100	100	90	95	80		
	Morningglory	35	55	100	90	15	80	0	5	5	70	35		
15	Nutsedge, Yellow	5	5	50	30	5	15	0	0	0	0	10		
	Oilseed Rape	85	95	100	100	90	100	-	100	100	100	100		
	Pigweed	100	100	100	100	100	100	100	100	100	100	100		
	Ragweed	0	40	100	75	70	100	40	15	0	70	55		
	Ryegrass, Italian	35	70	80	70	0	95	15	50	15	85	20		
20	Soybean	50	25	90	75	45	55	30	60	35	40	60		
	Velvetleaf	80	85	100	100	40	100	85	90	90	100	100		
	Waterhemp	100	100	100	100	100	100	90	95	95	100	98		
	Wheat	0	0	25	5	0	45	0	0	0	20	0		

Table D

Compounds

25 31 g ai/ha 32 35 53 54 55 57 58 60 62 63 104 113 118 120

	Preemergence														
	Barnyardgrass	55	75	5	5	85	20	5	5	50	65	10	95	60	75
	Blackgrass	30	90	25	5	90	5	5	5	5	85	0	40	10	60
	Corn	0	0	10	0	20	0	20	0	35	5	0	0	5	35
30	Crabgrass, Large	98	100	95	75	100	85	90	55	95	98	85	100	100	100
	Foxtail, Giant	85	98	85	10	100	90	15	5	95	80	80	100	100	98
	Galium	65	100	95	85	100	70	98	85	100	98	80	90	98	98
	Johnsongrass	75	60	0	0	75	10	10	0	30	35	20	70	70	55
	Lambsquarters	98	98	100	60	100	100	100	80	100	90	80	98	90	95
35	Morningglory	5	55	0	0	50	20	25	0	25	0	5	35	5	30
	Nutsedge, Yellow	0	5	0	0	10	10	0	0	0	0	0	10	0	5
	Oilseed Rape	100	100	100	85	100	100	98	98	100	98	0	100	98	90

163

5	Pigweed	98	100	100	100	100	85	100	100	100	100	100	100	100	100
	Ragweed	75	75	40	40	75	35	35	55	0	20	25	55	20	0
	Ryegrass, Italian	0	30	10	5	35	0	0	0	40	40	0	10	5	35
	Soybean	25	60	25	10	35	65	35	55	60	15	20	40	75	10
	Velvetleaf	55	85	100	70	100	100	100	100	90	100	50	85	70	75
	Waterhemp	100	100	100	85	100	90	100	100	100	100	100	100	100	100
	Wheat	0	5	0	0	0	0	0	0	0	0	0	5	0	0
Table D		Compounds													
10	31 g ai/ha	131	143	144	145	157	158	168	179	180	183	200	229		
	Preemergence														
	Barnyardgrass	75	70	80	85	5	5	75	5	0	5	95	55		
	Blackgrass	80	15	95	90	5	15	90	30	0	0	90	15		
	Corn	10	5	5	5	0	0	15	0	0	0	0	5		
15	Crabgrass, Large	100	100	98	100	40	90	100	65	75	70	100	85		
	Foxtail, Giant	100	100	100	100	20	55	100	30	60	35	100	70		
	Galium	30	80	98	98	85	50	100	90	80	50	98	80		
	Johnsongrass	30	60	75	75	5	0	95	0	10	5	60	10		
	Lambsquarters	80	85	100	100	100	100	100	100	100	85	98	80		
20	Morningglory	40	35	75	55	10	15	70	0	0	0	45	0		
	Nutsedge, Yellow	5	0	30	0	0	0	5	0	0	0	0	0		
	Oilseed Rape	50	85	100	100	100	50	100	95	95	100	100	100		
	Pigweed	100	100	100	100	100	75	100	100	98	60	100	100		
	Ragweed	0	30	85	60	0	60	45	30	5	0	5	50		
25	Ryegrass, Italian	20	5	60	35	5	0	95	5	0	0	80	20		
	Soybean	10	10	75	60	50	25	30	0	25	10	10	35		
	Velvetleaf	65	75	100	100	75	25	100	65	85	80	100	90		
	Waterhemp	100	98	100	100	100	100	100	90	95	100	100	100		
	Wheat	0	0	5	0	5	0	5	0	0	0	5	0		
Table D		Compounds													
30	16 g ai/ha	32	35	53	54	55	57	58	60	63	104	113	118	120	131
	Preemergence														
	Barnyardgrass	5	65	0	0	25	5	5	0	30	0	30	10	25	30
	Blackgrass	0	70	5	5	15	5	0	0	30	0	10	0	70	5
	Corn	0	0	5	0	5	0	5	0	0	0	0	0	0	0
35	Crabgrass, Large	90	100	65	10	100	75	25	5	65	65	98	100	98	98
	Foxtail, Giant	35	90	10	0	100	50	0	0	30	30	85	98	95	80
	Galium	25	100	85	50	100	80	0	98	75	90	90	60	90	95

164

	Johnsongrass	45	10	0	0	30	0	10	0	5	5	10	10	50	0
	Lambsquarters	95	100	100	5	100	80	25	10	95	65	85	90	70	75
	Morningglory	0	0	0	0	10	10	10	0	0	5	20	30	0	0
	Nutsedge, Yellow	65	0	20	0	0	0	0	0	0	0	0	0	0	0
5	Oilseed Rape	80	98	80	60	100	100	60	80	70	0	98	85	50	10
	Pigweed	100	100	95	5	100	75	50	75	95	100	100	100	100	100
	Ragweed	60	40	0	10	70	25	100	15	75	40	40	85	0	0
	Ryegrass, Italian	0	0	0	0	15	0	0	0	0	0	5	0	5	0
	Soybean	15	30	0	0	25	-	30	35	5	10	35	80	75	5
10	Velvetleaf	40	85	100	15	90	100	85	60	40	25	70	80	50	20
	Waterhemp	100	100	100	25	100	100	100	80	90	100	100	100	100	100
	Wheat	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table D

Compounds

	16 g ai/ha	143	144	145	157	158	168	179	180	183	200	229
15	Preemergence											
	Barnyardgrass	20	70	65	0	0	35	0	0	5	35	15
	Blackgrass	10	55	15	0	0	85	0	0	0	85	15
	Corn	0	0	0	5	0	0	0	0	0	0	0
	Crabgrass, Large	98	90	85	5	25	100	10	30	5	98	75
20	Foxtail, Giant	85	100	85	0	5	100	5	5	5	95	5
	Galium	10	95	90	80	0	100	90	75	50	85	40
	Johnsongrass	15	20	10	0	0	35	0	0	0	25	10
	Lambsquarters	70	100	100	100	75	90	75	95	80	75	35
	Morningglory	10	20	15	0	5	5	0	0	0	5	0
25	Nutsedge, Yellow	0	15	0	0	0	5	0	0	0	0	0
	Oilseed Rape	0	98	40	40	40	95	30	70	50	85	90
	Pigweed	100	100	100	100	70	85	30	80	50	100	100
	Ragweed	0	75	25	0	80	35	25	0	0	0	35
	Ryegrass, Italian	5	5	0	0	0	70	0	0	0	30	0
30	Soybean	5	35	10	0	0	5	0	5	0	100	5
	Velvetleaf	30	100	90	45	35	85	15	75	40	100	70
	Waterhemp	100	100	100	100	100	100	75	40	10	98	95
	Wheat	0	0	0	0	0	0	0	0	0	0	0

Table D

Compounds

Table D

Compound

8 g ai/ha

32 35 143 157

4 g ai/ha

157

Preemergence

Preemergence

Barnyardgrass

0 5 5 0

Barnyardgrass

0

165

Blackgrass	0	70	0	0	Blackgrass	0
Corn	0	0	0	0	Corn	0
Crabgrass, Large	100	80	60	0	Crabgrass, Large	0
Foxtail, Giant	0	70	40	0	Foxtail, Giant	0
Galium	0	85	10	80	Galium	70
Johnsongrass	0	0	35	0	Johnsongrass	0
Lambsquarters	95	100	25	25	Lambsquarters	0
Morningglory	0	0	0	0	Morningglory	0
Nutsedge, Yellow	60	0	0	0	Nutsedge, Yellow	0
Oilseed Rape	50	85	0	0	Oilseed Rape	0
Pigweed	100	100	90	100	Pigweed	20
Ragweed	75	55	0	0	Ragweed	0
Ryegrass, Italian	0	0	0	0	Ryegrass, Italian	0
Soybean	10	10	0	0	Soybean	0
Velvetleaf	5	65	35	5	Velvetleaf	0
Waterhemp	100	100	95	70	Waterhemp	0
Wheat	0	0	0	0	Wheat	0

Table D	Compounds									
250 g ai/ha	1	32	40	53	55	144	145	178	180	221
Flood										
Barnyardgrass	0	20	0	0	30	35	35	0	10	60
Ducksalad	0	95	40	80	90	100	95	70	75	90
Rice	0	15	0	0	0	40	35	0	0	40
Sedge, Umbrella	0	90	50	75	100	95	90	75	80	95

Table D	Compounds									
125 g ai/ha	1	32	40	53	55	144	145	178	180	
Flood										
Barnyardgrass	0	10	0	0	10	15	0	0	10	
Ducksalad	0	65	20	80	85	90	95	70	75	
Rice	0	0	0	0	0	25	15	0	0	
Sedge, Umbrella	0	65	30	70	85	95	90	60	75	

Table D	Compounds									
62 g ai/ha	1	32	40	53	55	144	145	178	180	221
Flood										
Barnyardgrass	0	0	0	0	0	0	0	0	0	40
Ducksalad	0	60	0	50	80	85	85	65	0	80
Rice	0	0	0	0	0	15	15	0	0	15

Sedge, Umbrella	0	40	0	0	75	80	75	60	50	80
-----------------	---	----	---	---	----	----	----	----	----	----

Table D

Compounds

31 g ai/ha	1	32	40	53	55	144	145	178	180
------------	---	----	----	----	----	-----	-----	-----	-----

Flood

5	Barnyardgrass	0	0	0	0	0	0	0	0	0
	Ducksalad	0	0	0	40	75	75	80	40	0
	Rice	0	0	0	0	0	0	0	0	0
	Sedge, Umbrella	0	0	0	0	75	65	60	30	0

TEST E

10 Seeds of plant species selected from bluegrass (annual bluegrass, *Poa annua*), blackgrass (*Alopecurus myosuroides*), canarygrass (*Phalaris minor*), chickweed (common chickweed, *Stellaria media*), galium (catchweed bedstraw, *Galium aparine*), downy brome grass (*Bromus tectorum*), field poppy (*Papaver rhoeas*), field violet (*Viola arvensis*), green foxtail (*Setaria viridis*), deadnettle (henbit deadnettle, *Lamium amplexicaule*), Italian ryegrass (*Lolium multiflorum*), kochia (*Kochia scoparia*), lambsquarters (*Chenopodium album*), oilseed rape (*Brassica napus*), pigweed (*Amaranthus retroflexus*), Russian thistle (*Salsola iberica*), chamomile (scentless chamomile, *Matricaria inodora*), speedwell (bird's-eye speedwell, *Veronica persica*), spring barley (*Hordeum vulgare*), spring wheat (*Triticum aestivum*), wild buckwheat (*Polygonum convolvulus*), wild mustard (*Sinapis arvensis*), wild

15 oat (*Avena fatua*), wild radish (*Raphanus raphanistrum*), windgrass (*Apera spica-venti*), winter barley (*Hordeum vulgare*), and winter wheat (*Triticum aestivum*) were planted into a silt loam soil and treated preemergence with test chemicals formulated in a non-phytotoxic solvent mixture which included a surfactant. At the same time, these species were planted in pots containing Redi-Earth[®] planting medium (Scotts Company, 14111 Scottslawn Road,

20 Marysville, Ohio 43041) comprising sphagnum peat moss, vermiculite, wetting agent and starter nutrients and treated with postemergence applications of the test chemicals formulated in the same manner. Plants ranged in height from 2 to 18 cm (1- to 4-leaf stage).

25 Treated plants and controls were maintained in a controlled growth environment for 7 to 21 days after which time all species were compared to controls and visually evaluated.

30 Plant response ratings, summarized in Table E, are based on a scale of 0 to 100 where 0 is no effect and 100 is complete control. A dash (–) response means no test result.

Table E

Compounds

125 g ai/ha	35	53	55	62	144	145	168	200
-------------	----	----	----	----	-----	-----	-----	-----

Postemergence

35	Barley, Spring	25	10	20	25	45	30	25	35
	Barley, Winter	30	10	10	35	45	40	15	25
	Blackgrass	60	25	40	75	85	80	60	75

	Bluegrass	60	5	25	60	65	70	15	70
	Bromegrass, Downy	30	5	10	40	75	55	35	35
	Buckwheat, Wild	95	95	95	100	100	100	100	100
	Canarygrass	60	40	30	55	85	65	55	60
5	Chamomile	10	0	5	10	95	30	10	15
	Chickweed	100	65	70	90	100	95	90	100
	Deadnettle	80	65	80	70	-	100	80	90
	Field Poppy	100	80	95	75	100	100	100	100
	Field Violet	75	70	90	85	100	100	95	95
10	Foxtail, Green	55	25	60	65	100	98	75	55
	Galium	85	75	85	90	100	100	85	90
	Kochia	90	80	85	95	100	100	85	90
	Lambsquarters	90	75	90	95	100	95	95	95
	Mustard, Wild	100	95	95	100	100	100	95	100
15	Oat, Wild	60	25	50	60	90	80	65	35
	Oilseed Rape	100	80	95	100	100	100	100	100
	Pigweed	90	90	95	95	100	100	95	90
	Radish, Wild	95	90	80	100	100	100	85	95
	Russian Thistle	-	-	-	-	98	100	-	-
20	Ryegrass, Italian	60	5	20	35	65	40	25	40
	Speedwell	100	100	100	100	100	100	100	100
	Wheat, Spring	40	35	45	40	75	55	40	35
	Wheat, Winter	40	35	40	35	70	55	35	25
	Windgrass	40	20	40	65	55	60	40	45
25	Table E	Compounds							
	62 g ai/ha	35	53	55	62	144	145	168	200
	Postemergence								
	Barley, Spring	10	10	15	20	35	20	15	25
	Barley, Winter	5	5	5	20	35	30	10	20
30	Blackgrass	55	10	20	40	60	65	35	55
	Bluegrass	40	5	15	55	55	65	10	55
	Bromegrass, Downy	15	5	10	20	40	55	25	25
	Buckwheat, Wild	90	90	95	95	100	100	95	95
	Canarygrass	55	25	30	45	65	40	25	45
35	Chamomile	5	5	5	5	80	30	10	10
	Chickweed	75	60	70	80	100	100	70	100
	Deadnettle	70	60	70	65	-	100	75	80

	Field Poppy	80	80	100	70	100	100	100	100
	Field Violet	75	70	85	70	100	90	90	75
	Foxtail, Green	55	10	50	25	80	40	60	45
	Galium	75	75	80	75	100	98	80	75
5	Kochia	80	70	85	90	100	100	85	85
	Lambsquarters	85	65	85	95	98	95	90	90
	Mustard, Wild	85	95	95	100	100	100	95	100
	Oat, Wild	35	10	40	35	65	40	40	30
	Oilseed Rape	95	80	90	80	100	100	95	100
10	Pigweed	90	90	95	95	100	100	95	90
	Radish, Wild	85	90	80	90	100	100	80	85
	Russian Thistle	-	-	-	-	95	90	-	-
	Ryegrass, Italian	25	5	15	30	40	25	20	25
	Speedwell	100	95	100	100	100	100	90	100
15	Wheat, Spring	25	20	25	25	45	45	25	20
	Wheat, Winter	25	15	30	30	35	35	25	15
	Windgrass	30	10	25	30	25	20	25	35
	Table E					Compounds			
	31 g ai/ha	35	53	55	62	144	145	168	200
20	Postemergence								
	Barley, Spring	5	5	10	15	25	15	15	15
	Barley, Winter	0	5	0	10	20	15	10	10
	Blackgrass	35	5	10	30	50	35	25	35
	Bluegrass	20	0	0	35	45	60	5	35
25	Bromegrass, Downy	5	0	0	15	35	35	15	10
	Buckwheat, Wild	85	65	95	80	95	100	95	70
	Canarygrass	25	10	20	35	60	25	15	35
	Chamomile	5	5	5	5	75	30	5	10
	Chickweed	75	50	55	70	100	85	45	75
30	Deadnettle	60	30	50	35	-	100	55	60
	Field Poppy	75	65	90	55	100	100	85	80
	Field Violet	65	50	70	65	90	100	75	80
	Foxtail, Green	35	10	45	20	50	30	60	35
	Galium	80	75	85	70	95	95	75	75
35	Kochia	75	55	70	75	100	100	80	85
	Lambsquarters	80	70	80	95	95	90	85	85
	Mustard, Wild	85	75	85	100	100	100	95	95

	Oat, Wild	15	5	15	25	25	30	10	20
	Oilseed Rape	80	75	75	80	100	100	85	85
	Pigweed	90	85	90	90	100	100	80	90
	Radish, Wild	75	60	70	70	100	100	75	80
5	Russian Thistle	-	-	-	-	90	85	-	-
	Ryegrass, Italian	5	0	20	20	15	10	10	10
	Speedwell	100	60	80	95	100	100	75	100
	Wheat, Spring	10	10	15	20	35	35	20	10
	Wheat, Winter	15	5	15	20	30	20	10	5
10	Windgrass	25	5	10	20	20	10	15	20
Table E Compounds									
	16 g ai/ha	35	53	55	62	144	145	168	200
	Postemergence								
	Barley, Spring	5	5	10	10	15	15	10	10
15	Barley, Winter	0	5	0	5	30	15	5	5
	Blackgrass	20	0	5	15	25	25	15	15
	Bluegrass	15	0	0	10	25	15	10	20
	Bromegrass, Downy	5	0	5	10	25	25	5	5
	Buckwheat, Wild	75	70	65	75	100	100	65	65
20	Canarygrass	10	5	10	25	35	15	15	10
	Chamomile	5	5	5	0	80	20	5	5
	Chickweed	65	30	55	65	85	80	50	65
	Deadnettle	40	20	20	35	-	100	20	50
	Field Poppy	70	60	85	40	100	100	55	75
25	Field Violet	60	25	65	60	95	65	75	70
	Foxtail, Green	20	10	30	15	40	15	20	25
	Galium	70	75	70	70	80	95	75	70
	Kochia	75	50	65	65	100	98	80	70
	Lambsquarters	80	65	75	75	80	90	75	75
30	Mustard, Wild	90	65	70	75	100	100	75	75
	Oat, Wild	10	5	10	15	20	15	5	10
	Oilseed Rape	70	70	75	65	100	70	70	75
	Pigweed	85	85	85	90	98	100	90	85
	Radish, Wild	65	60	75	65	100	95	70	65
35	Russian Thistle	-	-	-	-	85	80	-	-
	Ryegrass, Italian	0	0	20	10	5	5	5	5
	Speedwell	75	60	75	70	100	100	80	70

Wheat, Spring	0	10	5	5	25	30	10	5
Wheat, Winter	5	5	0	10	20	15	5	0
Windgrass	10	5	5	10	10	5	10	10

Table E Compounds

5	125 g ai/ha	35	53	55	62	144	145	168	200
	Preemergence								
	Barley, Spring	-	35	50	0	65	80	35	-
	Barley, Winter	-	25	35	5	55	85	40	-
	Blackgrass	95	30	100	65	100	75	100	100
10	Bluegrass	70	25	100	75	85	90	100	100
	Bromegrass, Downy	25	15	55	20	55	60	40	80
	Buckwheat, Wild	100	100	100	100	100	100	100	100
	Canarygrass	100	95	100	90	100	100	100	100
	Chamomile	70	-	-	65	100	100	-	75
15	Chickweed	100	100	100	100	100	100	100	100
	Deadnettle	95	65	0	100	100	100	100	100
	Field Poppy	90	100	100	100	100	100	100	95
	Field Violet	100	100	100	100	100	100	100	100
	Foxtail, Green	100	55	100	85	100	100	100	100
20	Galium	100	100	100	100	100	100	100	100
	Kochia	100	100	100	100	100	100	100	100
	Lambsquarters	100	100	100	100	100	100	100	100
	Mustard, Wild	95	100	100	100	100	100	100	100
	Oat, Wild	35	25	45	20	75	85	50	75
25	Oilseed Rape	100	100	100	100	100	100	100	100
	Pigweed	100	100	100	100	100	100	100	100
	Radish, Wild	100	90	100	100	100	100	100	95
	Russian Thistle	-	-	-	-	100	100	-	-
	Ryegrass, Italian	75	25	50	25	100	70	75	85
30	Speedwell	100	100	100	100	100	100	100	100
	Wheat, Spring	-	10	35	10	55	70	30	-
	Wheat, Winter	-	30	20	5	35	75	30	-
	Windgrass	100	50	100	50	100	98	100	100
	Table E Compounds								
35	62 g ai/ha	35	53	55	62	144	145	168	200
	Preemergence								
	Barley, Spring	-	35	45	0	45	45	30	-

	Barley, Winter	-	30	10	5	55	30	25	-
	Blackgrass	65	25	60	15	98	60	100	90
	Bluegrass	35	20	10	10	40	10	80	95
	Bromegrass, Downy	35	15	55	10	35	35	10	45
5	Buckwheat, Wild	90	100	100	80	100	85	100	100
	Canarygrass	95	65	100	50	100	100	100	100
	Chamomile	70	-	-	60	100	100	-	60
	Chickweed	95	100	100	100	100	100	100	100
	Deadnettle	95	50	0	40	100	100	80	100
10	Field Poppy	90	100	100	100	100	100	100	95
	Field Violet	90	60	100	35	100	100	100	-
	Foxtail, Green	90	20	100	65	95	55	100	100
	Galium	100	60	70	65	100	100	100	55
	Kochia	100	55	100	85	100	100	100	100
15	Lambsquarters	90	100	100	95	100	100	100	100
	Mustard, Wild	95	100	100	95	100	100	95	85
	Oat, Wild	45	10	35	15	65	30	35	30
	Oilseed Rape	100	100	100	65	100	100	65	100
	Pigweed	100	100	100	100	100	100	100	100
20	Radish, Wild	100	100	100	95	100	85	100	95
	Russian Thistle	-	-	-	-	100	60	-	-
	Ryegrass, Italian	35	25	50	15	70	50	30	30
	Speedwell	100	100	100	95	100	100	100	100
	Wheat, Spring	-	10	15	0	40	20	20	-
25	Wheat, Winter	-	25	10	0	25	25	10	-
	Windgrass	85	35	100	50	100	75	100	100
Table E									
Compounds									
	31 g ai/ha	35	53	55	62	144	145	168	200
Preemergence									
30	Barley, Spring	-	35	10	0	25	25	15	-
	Barley, Winter	-	10	10	0	35	15	5	-
	Blackgrass	60	10	15	15	75	50	35	75
	Bluegrass	35	10	10	5	15	0	10	70
	Bromegrass, Downy	35	15	20	0	10	20	10	35
35	Buckwheat, Wild	75	100	100	35	100	100	65	75
	Canarygrass	55	20	20	0	80	75	15	55
	Chamomile	10	-	-	65	20	100	-	25

	Chickweed	100	100	100	75	100	100	80	90
	Deadnettle	80	10	0	20	100	100	50	90
	Field Poppy	80	100	100	75	100	100	100	90
	Field Violet	70	25	80	10	100	100	100	50
5	Foxtail, Green	90	10	25	40	35	45	100	25
	Galium	55	30	100	60	70	100	60	25
	Kochia	65	15	100	5	100	100	100	100
	Lambsquarters	85	75	100	85	100	100	100	85
	Mustard, Wild	60	100	100	80	100	100	80	95
10	Oat, Wild	0	0	15	10	30	30	10	0
	Oilseed Rape	95	75	100	25	80	100	40	100
	Pigweed	95	20	100	100	100	100	100	100
	Radish, Wild	95	70	100	40	80	15	45	70
	Russian Thistle	-	-	-	-	65	10	-	-
15	Ryegrass, Italian	25	10	0	0	25	20	0	25
	Speedwell	100	70	100	95	100	100	100	100
	Wheat, Spring	-	5	15	0	15	15	15	-
	Wheat, Winter	-	15	10	0	15	20	10	-
	Windgrass	80	15	25	15	75	25	15	55
20	Table E	Compounds							
	16 g ai/ha	35	53	55	62	144	145	168	200
	Preemergence								
	Barley, Spring	-	35	10	0	10	15	0	-
	Barley, Winter	-	15	5	0	30	10	0	-
25	Blackgrass	30	10	0	5	15	10	10	25
	Bluegrass	0	10	0	0	10	0	0	20
	Bromegrass, Downy	0	0	20	0	10	10	0	10
	Buckwheat, Wild	60	65	65	25	85	100	0	65
	Canarygrass	40	10	10	0	35	40	15	10
30	Chamomile	5	-	-	0	15	5	-	25
	Chickweed	55	100	65	65	100	100	100	70
	Deadnettle	35	10	0	15	90	0	10	60
	Field Poppy	75	100	100	65	100	100	80	80
	Field Violet	15	50	70	0	95	100	85	-
35	Foxtail, Green	5	10	10	20	20	0	10	-
	Galium	25	25	10	5	20	25	60	15
	Kochia	25	10	55	0	98	60	35	70

	Lambsquarters	65	60	95	35	100	100	45	10
	Mustard, Wild	20	90	80	25	90	95	80	85
	Oat, Wild	0	0	0	0	35	25	10	0
	Oilseed Rape	60	50	20	10	35	50	10	60
5	Pigweed	90	30	75	95	100	100	55	100
	Radish, Wild	100	50	70	25	80	15	0	80
	Russian Thistle	-	-	-	-	15	0	-	-
	Ryegrass, Italian	0	10	0	0	10	10	0	0
	Speedwell	100	100	100	95	100	95	100	100
10	Wheat, Spring	-	5	15	0	5	15	15	-
	Wheat, Winter	-	15	5	0	0	0	5	-
	Windgrass	35	0	5	0	10	10	0	15

TEST F

Seeds of plant species selected from corn (*Zea mays*), soybean (*Glycine max*),
 15 velvetleaf (*Abutilon theophrasti*), lambsquarters (*Chenopodium album*), wild poinsettia
 (*Euphorbia heterophylla*), palmer pigweed (*Amaranthus palmeri*), waterhemp (common
 waterhemp, *Amaranthus rudis*), surinam grass (*Brachiaria decumbens*), large (Lg) crabgrass
 (*Digitaria sanguinalis*), Brazilian crabgrass (*Digitaria horizontalis*), fall panicum (*Panicum*
dichotomiflorum), giant foxtail (*Setaria faberii*), green foxtail (*Setaria viridis*), goosegrass
 20 (*Eleusine indica*), johnsongrass (*Sorghum halepense*), ragweed (common ragweed,
Ambrosia elatior), barnyardgrass (*Echinochloa crus-galli*), sandbur (southern sandbur,
Cenchrus echinatus), arrowleaf sida (*Sida rhombifolia*), Italian ryegrass (*Lolium*
multiflorum), dayflower (Virginia (VA) dayflower, *Commelina virginica*), field bindweed
 (*Convolvulus arvensis*), cocklebur (common cocklebur, *Xanthium strumarium*),
 25 morningglory (*Ipomoea coccinea*), nightshade (eastern black nightshade, *Solanum*
ptycanthum), kochia (*Kochia scoparia*), yellow nutsedge (*Cyperus esculentus*), and hairy
 beggarticks (*Bidens pilosa*), were planted into a silt loam soil and treated preemergence with
 test chemicals formulated in a non-phytotoxic solvent mixture which included a surfactant.

At the same time, plants from these crop and weed species and also waterhemp_RES1,
 30 (ALS & Triazine resistant common waterhemp, *Amaranthus rudis*), and waterhemp_RES2,
 (ALS & HPPD resistant common waterhemp, *Amaranthus rudis*) were treated with
 postemergence applications of test chemicals formulated in the same manner. Plants ranged
 in height from 2 to 18 cm for postemergence treatments (1- to 4-leaf stage).

Treated plants and controls were maintained in a greenhouse for 14 to 21 days, after
 35 which time all species were compared to controls and visually evaluated. Plant response
 ratings, summarized in Table F, are based on a scale of 0 to 100 where 0 is no effect and 100
 is complete control. A dash (–) response means no test result.

Table F

Compounds

	125 g ai/ha	35	53	58	62	96	144	145	149	179	200	212	221
	Postemergence												
	Arrowleaf Sida	98	98	90	98	95	100	95	98	100	95	100	98
5	Barnyardgrass	20	10	5	20	20	75	70	10	0	25	10	70
	Beggarticks	70	60	65	80	40	75	70	50	50	10	40	70
	Corn	20	10	5	10	10	10	20	15	15	25	0	20
	Crabgrass, Brazil	40	20	10	10	25	55	50	25	30	20	35	75
	Dayflower, VA	80	60	10	65	70	85	85	50	-	35	50	80
10	Field Bindweed	70	60	50	75	50	90	90	75	50	35	70	80
	Horseweed	-	-	-	-	10	25	40	20	10	-	10	30
	Kochia	-	98	75	-	95	95	100	95	95	-	100	100
	Panicum, Fall	50	60	15	60	25	60	80	40	10	15	10	60
	Pigweed, Palmer	100	100	60	100	90	100	100	100	85	100	98	100
15	Poinsettia, Wild	80	70	60	50	50	65	80	70	35	80	60	80
	Ragweed	-	-	-	-	40	90	70	35	50	-	50	80
	Ryegrass, Italian	0	0	0	10	10	40	15	5	0	20	0	5
	Sandbur	0	10	10	0	0	20	15	10	10	5	10	0
	Soybean	98	95	95	85	80	95	95	90	90	90	95	95
20	Waterhemp	100	100	90	100	95	100	100	98	100	100	100	100
	Waterhemp_RES1	100	100	75	100	95	100	100	100	100	98	98	98
	Waterhemp_RES2	100	100	80	100	95	100	100	100	100	98	100	100

Table F

Compounds

	62 g ai/ha	35	53	55	58	62	96	144	145	149	179	200	212	221
25	Postemergence													
	Arrowleaf Sida	98	100	100	75	98	90	100	98	95	85	95	85	98
	Barnyardgrass	10	0	35	0	20	20	50	50	10	0	15	0	30
	Beggarticks	65	50	70	50	50	40	70	65	40	40	10	40	70
	Corn	15	0	20	5	5	10	10	20	10	10	20	0	10
30	Crabgrass, Brazil	35	20	30	10	15	20	40	40	20	20	20	30	45
	Dayflower, VA	50	50	-	20	25	70	60	75	40	-	15	40	80
	Field Bindweed	50	50	70	30	50	20	80	80	50	35	35	60	-
	Horseweed	-	-	20	-	-	0	10	20	10	10	-	15	10
	Kochia	-	95	98	60	-	75	95	95	90	90	-	100	100
35	Panicum, Fall	50	15	30	10	10	10	35	50	20	15	5	10	40
	Pigweed, Palmer	100	100	98	70	100	95	100	100	98	85	100	100	100
	Poinsettia, Wild	50	70	45	50	50	60	70	80	60	20	60	50	70

175

	Ragweed	-	-	55	-	-	20	85	60	30	40	-	30	70
	Ryegrass, Italian	0	5	10	0	10	0	20	10	10	0	15	0	0
	Sandbur	0	5	15	0	0	0	20	5	15	5	0	0	0
	Soybean	95	95	95	90	65	40	98	95	90	75	65	75	98
5	Waterhemp	100	100	100	80	100	100	100	100	95	95	98	95	95
	Waterhemp_RES1	100	95	100	70	100	95	100	100	90	100	95	95	100
	Waterhemp_RES2	100	100	100	75	100	90	100	100	100	100	98	98	95
	Table F	Compounds												
	31 g ai/ha	35	53	55	58	62	96	144	145	149	179	200	212	221
10	Postemergence													
	Arrowleaf Sida	95	98	95	60	95	80	100	95	85	80	85	70	90
	Barnyardgrass	0	0	50	0	20	10	30	25	5	0	0	0	25
	Beggarticks	60	40	55	50	10	30	60	50	30	50	5	40	60
	Corn	15	0	15	0	0	5	5	10	10	0	10	0	10
15	Crabgrass, Brazil	30	15	40	5	10	5	50	20	20	20	15	30	25
	Dayflower, VA	40	30	70	10	5	40	50	60	20	-	25	20	70
	Field Bindweed	35	50	25	20	35	10	75	75	50	25	35	40	75
	Horseweed	-	-	15	-	-	0	0	20	20	0	-	0	5
	Kochia	-	80	95	30	-	75	95	100	80	75	-	90	98
20	Panicum, Fall	40	20	40	0	5	15	30	10	20	10	5	0	15
	Pigweed, Palmer	95	90	98	50	95	80	100	100	95	90	98	75	90
	Poinsettia, Wild	50	60	35	60	40	30	50	75	40	15	40	40	75
	Ragweed	-	-	55	-	-	20	50	50	40	40	-	20	60
	Ryegrass, Italian	0	0	0	0	0	0	20	5	10	0	0	0	0
25	Sandbur	0	0	25	0	0	0	15	5	5	0	0	0	0
	Soybean	85	95	70	70	40	-	95	95	85	50	40	50	95
	Waterhemp	90	98	90	65	100	90	100	98	95	95	95	90	98
	Waterhemp_RES1	90	90	100	65	100	90	95	98	85	90	90	80	95
	Waterhemp_RES2	95	90	100	70	100	90	100	90	95	100	95	95	98
30	Table F	Compounds												
	16 g ai/ha	35	53	58	62	96	144	145	149	179	200	212	221	
	Postemergence													
	Arrowleaf Sida	80	90	70	80	70	100	95	75	70	75	75	80	
	Barnyardgrass	0	0	0	15	0	10	10	0	0	0	0	15	
35	Beggarticks	50	30	40	5	0	70	55	5	20	0	25	50	
	Corn	10	0	0	0	0	10	5	0	5	0	0	5	
	Crabgrass, Brazil	20	10	5	0	0	40	20	20	10	10	20	20	

5	Dayflower, VA	40	40	5	0	20	50	50	10	-	0	30	60
	Field Bindweed	30	55	10	40	5	70	70	60	15	20	30	70
	Horseweed	-	-	-	-	0	0	0	0	0	-	0	0
	Kochia	-	60	40	-	65	95	98	60	65	-	50	95
	Panicum, Fall	40	5	0	0	0	25	10	10	15	5	0	10
	Pigweed, Palmer	75	65	50	70	70	90	95	65	70	95	30	85
	Poinsettia, Wild	40	50	30	35	0	50	70	50	15	50	35	60
	Ragweed	-	-	-	-	15	50	30	10	45	-	20	50
	Ryegrass, Italian	0	0	0	0	0	10	10	0	0	0	0	0
10	Sandbur	0	0	0	0	0	10	5	0	0	0	0	0
	Soybean	80	80	50	20	40	65	95	50	60	20	50	60
	Waterhemp	85	95	60	90	75	100	95	85	70	80	90	90
	Waterhemp_RES1	80	85	20	85	85	90	95	80	90	75	70	85
	Waterhemp_RES2	90	80	60	80	85	98	95	95	95	95	70	90
15	Table F	Compounds											
	8 g ai/ha	35	53	58	62	200							
	Postemergence												
	Arrowleaf Sida	60	70	50	65	65							
	Barnyardgrass	0	0	0	10	0							
20	Beggarticks	50	15	40	5	0							
	Corn	10	0	0	0	0							
	Crabgrass, Brazil	10	0	5	0	10							
	Dayflower, VA	10	30	5	0	0							
	Field Bindweed	20	30	10	40	20							
25	Kochia	-	50	30	-	-							
	Panicum, Fall	30	0	15	0	5							
	Pigweed, Palmer	60	70	50	90	85							
	Poinsettia, Wild	30	50	30	25	35							
	Ryegrass, Italian	0	0	0	0	0							
30	Sandbur	0	0	0	0	0							
	Soybean	70	40	50	10	15							
	Waterhemp	70	80	65	80	75							
	Waterhemp_RES1	90	70	10	80	70							
	Waterhemp_RES2	90	70	65	80	75							
35	Table F	Compounds											
	125 g ai/ha	35	53	58	62	96	144	145	149	168	179	200	
	Preemergence												

	Arrowleaf Sida	100	100	98	100	90	100	98	100	100	90	100	
	Barnyardgrass	65	50	15	25	0	100	75	35	100	5	60	
	Beggarticks	50	20	20	0	20	0	25	0	0	0	0	
	Cocklebur	0	-	-	-	-	-	-	-	-	-	-	
5	Corn	20	15	0	0	0	35	60	0	35	40	0	
	Crabgrass, Brazil	100	90	75	100	5	100	100	85	100	25	100	
	Crabgrass, Large	100	98	80	98	95	100	100	85	100	60	100	
	Dayflower, VA	95	95	80	80	20	95	70	40	90	85	90	
	Field Bindweed	50	0	0	0	40	70	95	25	70	0	30	
10	Foxtail, Giant	100	95	40	75	20	100	100	85	100	20	100	
	Foxtail, Green	98	90	25	90	25	100	100	50	100	15	100	
	Goosegrass	95	60	75	75	60	100	95	60	100	0	98	
	Johnsongrass	65	5	20	40	98	70	85	15	100	10	90	
	Kochia	100	100	30	95	90	100	100	90	100	98	100	
15	Lambsquarters	100	98	100	100	100	100	100	100	100	100	100	
	Morningglory	65	0	10	50	25	95	35	10	75	25	15	
	Nightshade	100	98	98	100	90	100	98	100	98	100	98	
	Nutsedge, Yellow	35	20	0	0	0	0	0	15	20	0	35	
	Panicum, Fall	100	98	100	100	70	100	100	100	100	75	100	
20	Pigweed, Palmer	-	100	35	100	100	100	100	100	100	65	100	
	Poinsettia, Wild	40	0	20	35	25	35	65	35	90	0	70	
	Ragweed	70	40	75	35	35	85	80	20	98	0	40	
	Ryegrass, Italian	70	0	0	40	30	100	90	40	70	10	40	
	Sandbur	65	0	0	35	30	95	95	10	90	0	75	
25	Soybean	90	35	75	40	0	75	40	30	40	35	50	
	Surinam Grass	35	0	0	10	0	80	25	15	100	0	35	
	Velvetleaf	100	100	100	100	80	100	100	100	100	100	100	
	Waterhemp	100	100	100	100	100	100	100	100	100	100	100	
	Table F	Compounds											
30	62 g ai/ha	35	53	55	58	62	96	144	145	149	168	179	200
	Preemergence												
	Arrowleaf Sida	100	100	100	75	95	80	100	98	65	100	80	100
	Barnyardgrass	10	20	85	0	0	0	70	30	15	80	0	30
	Beggarticks	20	30	0	10	0	0	0	30	0	0	0	0
35	Corn	0	0	40	0	0	0	40	40	0	40	0	0
	Crabgrass, Brazil	100	85	100	35	100	0	98	95	35	100	15	100
	Crabgrass, Large	75	80	100	65	70	65	100	98	70	98	35	95

	Dayflower, VA	90	50	40	-	30	0	85	35	5	70	35	70
	Field Bindweed	50	0	10	0	0	60	65	65	35	50	0	30
	Foxtail, Giant	70	70	98	0	65	0	95	85	40	100	30	100
	Foxtail, Green	75	20	70	0	40	35	98	100	50	100	0	100
5	Goosegrass	75	10	50	5	5	40	95	80	40	98	0	95
	Johnsongrass	35	0	90	0	35	50	30	100	0	35	10	35
	Kochia	100	50	100	0	35	65	100	100	75	100	70	80
	Lambsquarters	100	95	100	90	95	100	100	100	100	100	100	98
	Morningglory	30	0	98	10	0	35	65	15	0	15	35	0
10	Nightshade	98	98	98	90	98	65	100	98	98	98	100	98
	Nutsedge, Yellow	10	0	0	0	0	0	0	0	0	0	0	10
	Panicum, Fall	100	95	100	90	95	50	100	95	98	100	50	98
	Pigweed, Palmer	-	100	100	35	100	100	100	100	100	100	35	100
	Poinsettia, Wild	70	0	30	20	25	0	20	65	0	50	0	60
15	Ragweed	40	30	50	25	40	50	95	65	35	65	0	35
	Ryegrass, Italian	65	0	30	0	0	30	40	40	20	65	0	20
	Sandbur	40	0	0	0	0	40	50	25	0	65	0	40
	Soybean	70	0	50	35	10	0	40	15	0	20	0	30
	Surinam Grass	20	0	10	0	15	0	75	0	15	70	0	10
20	Velvetleaf	100	100	100	100	100	80	100	100	85	100	90	100
	Waterhemp	100	100	100	90	100	100	100	100	100	100	100	100
Table F		Compounds											
	31 g ai/ha	35	53	55	58	62	96	144	145	149	168	179	200
	Preemergence												
25	Arrowleaf Sida	100	65	98	35	70	70	100	90	35	100	80	100
	Barnyardgrass	20	0	10	0	0	0	20	5	0	25	0	0
	Beggarticks	20	0	0	0	0	0	0	35	0	0	0	0
	Cocklebur	0	-	-	-	-	-	-	-	-	-	-	-
	Corn	30	0	0	0	0	0	0	0	0	0	0	0
30	Crabgrass, Brazil	98	5	75	0	75	0	95	80	15	100	0	98
	Crabgrass, Large	75	15	98	50	25	30	100	95	0	95	0	85
	Dayflower, VA	65	0	60	0	0	0	30	0	0	10	0	50
	Field Bindweed	50	0	0	0	0	0	60	50	50	60	0	30
	Foxtail, Giant	70	20	90	0	35	0	90	65	20	98	0	65
35	Foxtail, Green	30	0	20	0	10	30	95	40	0	100	0	98
	Goosegrass	60	0	5	5	0	40	60	70	0	95	0	80
	Johnsongrass	0	0	20	0	20	60	0	20	0	30	0	5

	Kochia	85	20	100	0	20	0	100	98	0	100	0	50
	Lambsquarters	100	95	100	65	95	100	100	100	0	100	100	98
	Morningglory	0	0	30	10	0	35	0	10	0	0	0	0
	Nightshade	98	90	98	80	70	0	98	80	80	98	65	60
5	Nutsedge, Yellow	0	0	0	0	0	0	0	0	0	0	0	0
	Panicum, Fall	100	0	90	65	0	40	100	80	80	95	0	35
	Pigweed, Palmer	-	90	75	0	75	98	100	100	90	100	0	100
	Poinsettia, Wild	35	0	0	20	30	0	0	0	0	0	0	20
	Ragweed	0	0	40	0	25	20	0	70	25	35	0	10
10	Ryegrass, Italian	15	0	0	0	0	0	0	20	0	30	0	0
	Sandbur	10	0	0	0	0	10	10	35	0	30	0	5
	Soybean	40	0	40	20	0	0	10	0	0	0	0	20
	Surinam Grass	0	0	0	0	15	0	75	0	0	65	0	0
	Velvetleaf	70	90	100	100	75	60	100	98	70	75	65	80
15	Waterhemp	100	95	100	75	98	100	100	100	100	100	95	100

Table F

Compounds

	16 g ai/ha	35	53	58	62	96	144	145	149	168	179	200
	Preemergence											
	Arrowleaf Sida	80	20	0	20	50	95	65	0	95	0	65
20	Barnyardgrass	0	0	0	0	0	0	0	0	0	0	0
	Beggarticks	0	0	0	0	0	0	0	0	0	0	0
	Corn	0	0	0	0	0	0	0	0	0	0	0
	Crabgrass, Brazil	95	0	0	0	0	75	5	0	65	0	90
	Crabgrass, Large	60	0	20	0	0	80	10	0	65	0	50
25	Dayflower, VA	25	0	0	0	0	0	0	0	5	0	15
	Field Bindweed	35	0	0	0	0	40	0	0	40	0	0
	Foxtail, Giant	10	0	0	15	0	15	0	0	65	0	35
	Foxtail, Green	0	0	0	0	25	25	5	0	70	0	10
	Goosegrass	0	0	0	0	50	75	0	0	80	0	70
30	Johnsongrass	0	0	0	0	20	0	0	0	0	0	0
	Kochia	40	0	0	-	0	98	75	0	95	0	25
	Lambsquarters	98	65	0	90	5	100	100	-	100	98	0
	Morningglory	0	0	0	0	35	0	0	0	0	0	0
	Nightshade	95	35	65	0	0	98	90	0	5	-	35
35	Nutsedge, Yellow	0	0	0	0	0	0	0	0	0	0	0
	Panicum, Fall	35	0	35	0	-	95	65	0	35	0	70
	Pigweed, Palmer	-	0	0	80	80	100	100	25	98	0	85

180

	Poinsettia, Wild	30	0	0	0	0	0	0	0	0	0	0
	Ragweed	0	0	0	20	0	0	0	20	10	0	0
	Ryegrass, Italian	0	0	0	0	0	0	25	0	15	0	0
	Sandbur	10	0	0	0	0	0	0	0	10	0	5
5	Soybean	35	0	10	0	0	-	0	0	0	0	0
	Surinam Grass	0	0	0	0	0	0	0	0	50	0	0
	Velvetleaf	70	65	70	35	0	80	35	10	35	15	25
	Waterhemp	100	0	75	85	40	98	100	35	100	95	98
	Table F	Compounds										
10	8 g ai/ha	35	53	58	62	200						
	Preemergence											
	Arrowleaf Sida	50	0	20	0	0						
	Barnyardgrass	0	0	0	0	0						
	Beggarticks	0	0	0	0	0						
15	Corn	0	0	0	0	0						
	Crabgrass, Brazil	0	0	0	0	0						
	Crabgrass, Large	0	0	0	0	0						
	Dayflower, VA	0	0	0	0	0						
	Field Bindweed	0	0	0	0	0						
20	Foxtail, Giant	0	0	0	0	0						
	Foxtail, Green	0	0	0	0	0						
	Goosegrass	0	0	0	0	10						
	Johnsongrass	0	0	0	0	0						
	Kochia	0	0	0	10	20						
25	Lambsquarters	90	65	-	70	0						
	Morningglory	0	0	0	0	0						
	Nightshade	75	-	0	0	0						
	Nutsedge, Yellow	0	0	0	0	0						
	Panicum, Fall	0	0	0	0	0						
30	Pigweed, Palmer	-	0	0	10	0						
	Poinsettia, Wild	20	0	0	0	0						
	Ragweed	0	0	0	0	0						
	Ryegrass, Italian	0	0	0	0	0						
	Sandbur	0	0	0	0	0						
35	Soybean	0	0	0	0	0						
	Surinam Grass	0	0	0	0	0						
	Velvetleaf	40	0	20	0	0						

Waterhemp 100 0 20 80 70

TEST G

Three plastic pots (ca. 16-cm diameter) per rate were partially filled with sterilized Tama silt loam soil comprising a 35:50:15 ratio of sand, silt and clay and 2.6% organic matter. Separate plantings for each of the three pots were as follows. Seeds from the U.S. of monochoria (*Monochoria vaginalis*), small-flower umbrella sedge (*Cyperus difformis*), hardstem bulrush (*Scirpus juncooides*), and redstem (purple redstem, *Ammannia coccinea*), were planted into one 16-cm pot for each rate. Seeds from the U.S. of rice flatsedge (*Cyperus iria*), bearded sprangletop (*Leptochloa fascicularis*), one stand of 9 or 10 water seeded rice seedlings (Indica rice, *Oryza sativa*), and two stands of 3 or 4 transplanted rice seedlings (*Oryza sativa* cv. 'Japonica – M202') were planted into one 16-cm pot for each rate. Seeds from the U.S. of barnyardgrass (*Echinochloa crus-galli*), and late watergrass (*Echinochloa oryzicola*) were planted into one 16-cm pot for each rate. Plantings were sequential so that crop and weed species were at the 2.0 to 2.5-leaf stage at time of treatment.

Potted plants were grown in a greenhouse with day/night temperature settings of 30/27 °C, and supplemental balanced lighting was provided to maintain a 16-hour photoperiod. Test pots were maintained in the greenhouse until test completion.

At time of treatment, test pots were flooded to 3 cm above the soil surface, treated by application of test compounds directly to the paddy water, and then maintained at that water depth for the duration of the test. Effects of treatments on rice and weeds were visually evaluated by comparison to untreated controls after 21 days. Plant response ratings, summarized in Table G, are based on a scale of 0 to 100 where 0 is no effect and 100 is complete control. A dash (–) response means no test result.


Table G	Compounds	Table G	Compounds
250 g ai/ha	35 55	125 g ai/ha	35 55
Flood		Flood	
Barnyardgrass	100 100	Barnyardgrass	100 100
Bulrush, Hardstem	100 100	Bulrush, Hardstem	100 100
Flatsedge, Rice	100 100	Flatsedge, Rice	100 100
Monochoria	100 100	Monochoria	100 100
Redstem	100 100	Redstem	90 100
Rice, Transplanted	100 100	Rice, Transplanted	45 80
Rice, Water Seeded	100 100	Rice, Water Seeded	100 100
Sedge, Umbrella	100 100	Sedge, Umbrella	100 100
Sprangletop, Brdd.	100 100	Sprangletop, Brdd.	100 100
Watergrass, Late	100 100	Watergrass, Late	40 75

Table G	Compounds	
64 g ai/ha	35	55
Flood		
Barnyardgrass	45	75
Bulrush, Hardstem	95	100
Flatsedge, Rice	100	100
Monochoria	100	100
Redstem	85	95
Rice, Transplanted	30	60
Rice, Water Seeded	70	100
Sedge, Umbrella	98	100
Sprangletop, Brdd.	100	80
Watergrass, Late	30	0

Table G	Compounds	
32 g ai/ha	35	55
Flood		
Barnyardgrass	30	0
Bulrush, Hardstem	0	80
Flatsedge, Rice	100	85
Monochoria	98	100
Redstem	0	85
Rice, Transplanted	15	20
Rice, Water Seeded	60	45
Sedge, Umbrella	0	98
Sprangletop, Brdd.	60	45
Watergrass, Late	0	0

CLAIMS


1. A compound selected from Formula 1, *N*-oxides and salts thereof,





- Q is a 5- or 6-membered aromatic heterocyclic ring, bound to the remainder of Formula 1 through a carbon atom, and optionally substituted with 1 to 4 R¹;
Z is O or S;
each R¹ is independently halogen, cyano, nitro, SF₅, CHO, C(=O)NH₂, C(=S)NH₂, SO₂NH₂, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₄-C₈ alkylcycloalkyl, C₄-C₈ cycloalkylalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₃-C₇ cycloalkylcarbonyl, C₂-C₈ alkylaminocarbonyl, C₃-C₁₀ dialkylaminocarbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₃-C₆ cycloalkoxy, C₃-C₆ halocycloalkoxy, C₄-C₈ cycloalkylalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ alkoxyhaloalkyl, C₂-C₆ alkoxyalkoxy, C₂-C₄ alkylcarbonyloxy, C₂-C₆ cyanoalkyl, C₂-C₆ cyanoalkoxy, C₁-C₄ hydroxyalkyl, C₂-C₄ alkylthioalkyl, SO_nR^{1A}, Si(CH₃)₃ or B(-OC(R^{1B})₂C(R^{1B})₂O-); or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{1C}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{1C} on carbon atom ring members and R^{1D} on nitrogen atom ring members;
R² is halogen, cyano, nitro, C₁-C₄ alkoxy, C₁-C₄ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, SO_nR^{2A}, C₁-C₄ haloalkyl or C₃-C₆ cycloalkyl;
each R³ is independently halogen, cyano, hydroxy, nitro, amino, CHO, C(=O)NH₂, C(=S)NH₂, SO₂NH₂, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₄-C₈ alkylcycloalkyl, C₄-C₈ cycloalkylalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₃-C₇

- cycloalkylcarbonyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₃-C₆ cycloalkoxy, C₃-C₆ halocycloalkoxy, C₄-C₈ cycloalkylalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ alkoxyhaloalkyl, C₂-C₆ alkoxyalkoxy, C₂-C₄ alkylcarbonyloxy, C₂-C₆ cyanoalkyl, C₂-C₆ cyanoalkoxy, C₂-C₄ alkylthioalkyl, Si(CH₃)₃, C≡CSi(CH₃)₃, C(=O)N(R^{3A})(R^{3B}), C(=NOR^{3C})H, C(=NR^{3D})H, SO_nR^{3E}; or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{3F}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members; or pyrimidinyl;
- m is 0, 1, 2 or 3;
- each n is independently 0, 1 or 2;
- each R^{1A}, R^{2A} and R^{3E} is independently C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkylamino or C₂-C₆ dialkylamino;
- each R^{1B} is independently H or C₁-C₄ alkyl;
- each R^{1C} is independently hydroxy, halogen, cyano, nitro, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₁-C₆ haloalkoxy;
- each R^{1D} is independently cyano, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₂-C₆ alkylcarbonyl;
- each R^{3A} is independently C₁-C₄ alkyl or C₁-C₄ haloalkyl;
- each R^{3B} is independently H, C₁-C₄ alkyl or C₁-C₄ haloalkyl;
- each R^{3C} is independently H or C₁-C₄ alkyl;
- each R^{3D} is independently H, amino, C₁-C₄ alkyl or C₁-C₄ alkylamino;
- each R^{3F} is independently hydroxy, halogen, cyano, nitro, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₁-C₆ haloalkoxy; and
- each R^{3G} is independently cyano, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy or C₂-C₆ alkylcarbonyl;

2. A compound of Claim 1 wherein
Q is selected from







wherein r is 0, 1, 2 or 3; and s is 0 or 1;

each R^1 is independently halogen, cyano, SF_5 , CHO, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_2\text{-C}_4$ alkenyl,

$\text{C}_2\text{-C}_4$ alkynyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_2\text{-C}_4$ haloalkenyl, $\text{C}_2\text{-C}_4$ haloalkynyl, $\text{C}_2\text{-C}_6$ alkylcarbonyl, $\text{C}_2\text{-C}_6$ haloalkylcarbonyl, $\text{C}_2\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_3\text{-C}_4$ alkenyloxy, $\text{C}_3\text{-C}_4$ alkynyloxy, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_3\text{-C}_4$ haloalkenyloxy, $\text{C}_3\text{-C}_4$ haloalkynyloxy, $\text{C}_2\text{-C}_6$ alkoxyalkyl, $\text{C}_2\text{-C}_6$ haloalkoxyalkyl, $\text{C}_2\text{-C}_6$ cyanoalkyl, $\text{C}_1\text{-C}_4$ hydroxyalkyl, $\text{C}_2\text{-C}_4$ alkylthioalkyl or SO_nR^{1A} ;

R^3 is independently halogen, cyano, CHO, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_2\text{-C}_4$ alkenyl, $\text{C}_2\text{-C}_4$ alkynyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_2\text{-C}_4$ haloalkenyl, $\text{C}_2\text{-C}_4$ haloalkynyl, $\text{C}_3\text{-C}_6$ cycloalkyl, $\text{C}_3\text{-C}_6$ halocycloalkyl, $\text{C}_4\text{-C}_8$ alkylcycloalkyl, $\text{C}_2\text{-C}_6$ alkylcarbonyl, $\text{C}_2\text{-C}_6$ haloalkylcarbonyl, $\text{C}_2\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_3\text{-C}_4$ alkenyloxy, $\text{C}_3\text{-C}_4$ alkynyloxy, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_3\text{-C}_4$ haloalkenyloxy, $\text{C}_3\text{-C}_4$ haloalkynyloxy, $\text{C}_3\text{-C}_6$ cycloalkoxy, $\text{C}_3\text{-C}_6$ halocycloalkoxy, $\text{C}_2\text{-C}_6$ alkoxyalkyl, $\text{C}_2\text{-C}_6$ haloalkoxyalkyl, $\text{C}_2\text{-C}_4$ alkylcarbonyloxy, $\text{C}_2\text{-C}_6$ cyanoalkyl, $\text{C}(=\text{O})\text{N}(\text{R}^{3A})(\text{R}^{3B})$, $\text{C}(=\text{NOR}^{3C})\text{H}$, SO_nR^{3E} ; or a phenyl ring optionally substituted with up to 5 substituents independently selected from R^{3F} ; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members;

Z is O; and
m is 0, 1 or 2.

3. A compound of Claim 2 wherein

each R¹ is independently halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₁-C₄ alkoxy, C₃-C₄ alkenyloxy, C₃-C₄ alkynyloxy, C₁-C₄ haloalkoxy, C₃-C₄ haloalkenyloxy, C₃-C₄ haloalkynyloxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₁-C₄ hydroxyalkyl, C₂-C₄ alkylthioalkyl or SO_nR^{1A};

R² is halogen, C₁-C₄ alkyl or C₁-C₄ haloalkyl;

each R³ is independently halogen, cyano, CHO, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₂-C₄ haloalkynyl, C₃-C₆ cycloalkyl, C₃-C₆ halocycloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxycarbonyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₂-C₆ alkoxyalkyl, C₂-C₆ haloalkoxyalkyl, C₂-C₆ cyanoalkyl, SO_nR^{3E}; or a 5- or 6-membered heteroaromatic ring containing ring members selected from carbon atoms and up to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, each ring optionally substituted with up to 3 substituents independently selected from R^{3F} on carbon atom ring members and R^{3G} on nitrogen atom ring members; and

m is 0 or 1.

4. A compound of Claim 3 wherein

Q is selected from Q-7 through Q-24;

each R¹ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy or SO_nR^{1A};

R² is halogen or C₁-C₄ alkyl;

each R³ is independently halogen, cyano, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxycarbonyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₂-C₆ alkoxyalkyl or C₂-C₆ haloalkoxyalkyl; and

each R^{1A} is independently C₁-C₄ alkyl or C₁-C₄ haloalkyl.

5. A compound of Claim 4 wherein

Q is selected from Q-16 and Q-18;

each R¹ is independently halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy; R² is halogen or CH₃; and

each R³ is independently halogen, cyano, C₁-C₄ alkyl or C₁-C₄ haloalkyl.

6. A compound of Claim 3 wherein

Q is selected from Q-43, Q-44, Q-45, Q-48, Q-49 and Q-50;

each R¹ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy or SO_nR^{1A};

R² is halogen or C₁-C₄ alkyl;

5 each R³ is independently halogen, cyano, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ haloalkyl, C₂-C₆ alkylcarbonyl, C₂-C₆ haloalkylcarbonyl, C₂-C₆ alkoxy carbonyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₂-C₆ alkoxyalkyl or C₂-C₆ haloalkoxyalkyl; and

each R^{1A} is independently C₁-C₄ alkyl or C₁-C₄ haloalkyl.

10 7. A compound of Claim 6 wherein

Q is selected from Q-43, Q-44 and Q-45;

each R¹ is independently halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy; R² is halogen or CH₃; and

each R³ is independently halogen, cyano, C₁-C₄ alkyl or C₁-C₄ haloalkyl.

15 8. A compound of Claim 1 selected from the group consisting of

5-chloro-2-[2-(5-chloro-2-pyridinyl)phenoxy]pyrimidine,
 5-chloro-2-[2-[5-(fluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine,
 2-[2-(3-bromo-5-isoxazolyl)phenoxy]-5-chloropyrimidine,
 5-chloro-2-[2-[5-(trifluoromethyl)-2-pyridinyl]phenoxy]pyrimidine,
 5-chloro-2-[3-chloro-2-(5-chloro-2-pyridinyl)phenoxy]pyrimidine,
 4-[2-[(5-bromo-2-pyrimidinyl)oxy]phenyl]-2-(trifluoromethyl)pyrimidine,
 2-[2-(2-bromo-5-thiazolyl)phenoxy]-5-(trifluoromethyl)pyrimidine,
 5-chloro-2-[4-methyl-2-[2-(trifluoromethyl)-4-pyridinyl]phenoxy]pyrimidine,
 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine,
 5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]phenoxy]pyrimidine,
 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine,
 5-bromo-2-[2-[5-(difluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine,
 5-chloro-2-[2-[3-(trifluoromethyl)-5-isoxazolyl]phenoxy]pyrimidine,
 5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine,
 5-bromo-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine,
 5-chloro-2-[2-[5-(trifluoromethyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine,
 5-chloro-2-[2-[5-(trifluoromethyl)-3-isoxazolyl]phenoxy]pyrimidine,
 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]-3-difluoromethoxyphenoxy]pyrimidine,
 5-chloro-2-[2-[5-(cyclopropyl)-3-isoxazolyl]-3-fluorophenoxy]pyrimidine,
 5-chloro-2-[2-[5-(trifluoromethyl)-3-isoxazolyl]-3-chlorophenoxy]pyrimidine,
 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]-3-chlorophenoxy]pyrimidine,

5-chloro-2-[2-[3-(trifluoromethyl)-5-isoxazolyl]-3-chlorophenoxy]pyrimidine,
 5-chloro-2-[2-[3-(trifluoromethyl)-5-isoxazolyl]-3-fluorophenoxy]pyrimidine,
 5-chloro-2-[2-[5-(trifluoromethyl)-3-isoxazolyl]-3-bromophenoxy]pyrimidine,
 5-chloro-2-[2-[3-(difluoromethyl)-5-isoxazolyl]-3-chlorophenoxy]pyrimidine,
 5-bromo-2-[2-[3-(trifluoromethyl)-5-isoxazolyl]-3-chlorophenoxy]pyrimidine,
 5-bromo-2-[2-[3-(trifluoromethyl)-5-isoxazolyl]-3-chlorophenoxy]pyrimidine,
 5-chloro-2-[2-[3-(difluoromethyl)-3-isoxazolyl]-3-bromophenoxy]pyrimidine,
 and
 5-chloro-2-[2-[5-(difluoromethyl)-3-isoxazolyl]-3-bromophenoxy]pyrimidine.

9. A herbicidal composition comprising a compound of any of Claims 1-8 and at least one component selected from the group consisting of surfactants, solid diluents and liquid diluents.

10. A herbicidal composition comprising a compound of any of Claims 1-8, at least one additional active ingredient selected from the group consisting of other herbicides and herbicide safeners, and at least one component selected from the group consisting of surfactants, solid diluents and liquid diluents.

11. A herbicidal mixture comprising (a) a compound of any of Claims 1-8, and (b) at least one additional active ingredient selected from (b1) photosystem II inhibitors, (b2) acetohydroxy acid synthase (AHAS) inhibitors, (b3) acetyl-CoA carboxylase (ACCase) inhibitors, (b4) auxin mimics and (b5) 5-enol-pyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, (b6) photosystem I electron diverters, (b7) protoporphyrinogen oxidase (PPO) inhibitors, (b8) glutamine synthetase (GS) inhibitors, (b9) very long chain fatty acid (VLCFA) elongase inhibitors, (b10) auxin transport inhibitors, (b11) phytoene desaturase (PDS) inhibitors, (b12) 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors, (b13) homogentisate solanesyltransferase (HST) inhibitors, (b14) cellulose biosynthesis inhibitors, (b15) other herbicides selected from mitotic disruptors, organic arsenicals, asulam, bromobutide, cinmethylin, cumyluron, dazomet, difenzoquat, dymron, etobenzanid, flurenol, fosamine, fosamine-ammonium, metam, methyldymron, oleic acid, oxaziclomefone, pelargonic acid and pyributicarb, and (b16) herbicide safeners and salts of compounds of (b1) through (b16).

12. A method for controlling the growth of undesired vegetation comprising contacting the vegetation or its environment with a herbicidally effective amount of a compound of any of Claims 1-8.

13. A method for controlling the growth of undesired vegetation in a genetically modified plants that exhibit traits of glyphosate tolerance, glufosinate tolerance, ALS herbicide tolerance, dicamba tolerance, imidazolinone herbicide tolerance, 2,4-D tolerance,

HPPD tolerance and mesotrione tolerance, comprising contacting the vegetation or its environment with a herbicidally effective amount of a compound of any of Claims 1-8.

14. A herbicidal mixture comprising (a) a compound of any of Claims 1-8, and (b) at least one additional active ingredient selected from (b1) photosystem II inhibitors, (b2) acetohydroxy acid synthase (AHAS) inhibitors, (b4) auxin mimics, (b5) 5-enol-pyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, (b7) protoporphyrinogen oxidase (PPO) inhibitors, (b9) very long chain fatty acid (VLCFA) elongase inhibitors and (b12) 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors.


15. A herbicidal mixture comprising (a) a compound of any of Claims 1-8, and (b) at least one additional active ingredient selected from the group consisting of 2,4-D, acetochlor, alachlor, atrazine, bromoxynil, bentazon, bicyclopyrone, carfentrazone-ethyl, cloransulam-methyl, dicamba, dimethenamid-p, florasulam, flufenacet, flumioxazin, flupyr-sulfuron-methyl, fluroxypyr-meptyl, glyphosate, halauxifen-methyl, isoxaflutole, MCPA, mesotrione, metolachlor, metsulfuron-methyl, nicosulfuron, pyrasulfotole, pyroxasulfone, pyroxsulam, rimsulfuron, saflufenacil, tembotrione, thifensulfuron-methyl, topramazine and tribenuron.

TITLE

PYRIMIDINYLOXY BENZENE DERIVATIVES AS HERBICIDES

ABSTRACT

Disclosed are compounds of Formula **1**, including all stereoisomers, *N*-oxides, and salts thereof,



wherein

Q, Z, R², R³ and m are as defined in the disclosure.

Also disclosed are compositions containing the compounds of Formula **1** and methods for controlling undesired vegetation comprising contacting the undesired vegetation or its environment with an effective amount of a compound or a composition of the invention.

[NO FIGURE]