




7. A compound according to claim 1, of the formula




(A),




(D),




(G),




(B),




(E),




(H),




(C),




(F),




(I),




(J),




(K),




(M),




(V),




(L),




(P),




(W),




(M),




(Q),




(X),




(N),




(T),




(R),




(U),




(S),




(Y),




(Zc),




(Zg),




(Z),




(Zd),




(Zh),




(Za),




(Ze),




(Zi).




(Zb).




(Zf).




(Zl).




(Zp).




(Zx),




(Zm),




(Zy),




(Zn),




(Zr),




(Zz),




(Zo),




(Zs),




(Zt),




(Zu),




(Zw),




(Zzb),



(Zzc),




(Zv),



(Zzd),

or




(Zza),

wherein said compound is selected from the group consisting of:

alkyl-, phenyl and benzyl, wherein phenyl and benzyl optionally may be substituted with one or more substituents independently selected from each other from the group of fluoro, chloro, bromo, -CN, C₁₋₄ alkyl-O-, C₁₋₄ alkyl-; and


- 5 PG is a protecting group for an amino function selected from the group consisting of tert-butoxycarbonyl-, 9-fluorenylmethoxycarbonyl-, benzyl-, and 2,4-dimethoxybenzyl-.

15. A compound according to claim 1, of the formula



- 10 wherein said compound is selected from the group consisting of:

20. A compound according to claim 1, of the formula



wherein said compound is selected from the group consisting of:

5 the stereoisomer with R-configuration at the chiral carbon atom designated by the numeral 1 and R-configuration at the chiral carbon atom designated by the numeral 3;

10 the stereoisomer with S-configuration at the chiral carbon atom designated by the numeral 1 and S-configuration at the chiral carbon atom designated by the numeral 3;


15 the stereoisomer with R-configuration at the chiral carbon atom designated by the numeral 1 and S-configuration at the chiral carbon atom designated by the numeral 3; and

20 the stereoisomer with S-configuration at the chiral carbon atom designated by the numeral 1 and R-configuration at the chiral carbon atom designated by the numeral 3;

wherein, in each such stereoisomer, the chiral carbon atom designated by the numeral 2 is always in the syn configuration with respect to the chiral carbon atom designated by the numeral 1;

wherein the chiral carbon atom designated by the numeral 2 is in the syn configuration with respect to the chiral carbon atom designated by the numeral 1.

25. A compound according to claim 1, of the formula



(I).

wherein said compound is selected from the group consisting of:

the stereoisomer with R-configuration at the chiral carbon atom designated by the numeral 1 and R-configuration at the chiral carbon atom designated by the numeral 3;

10

the stereoisomer with S-configuration at the chiral carbon atom designated by the numeral 1 and S-configuration at the chiral carbon atom designated by the numeral 3;

15


the diastereomer with R-configuration at the chiral carbon atom designated by the numeral 1 and S-configuration at the chiral carbon atom designated by the numeral 3; and

the stereoisomer with S-configuration at the chiral carbon atom designated by the numeral 1 and R-configuration at the chiral carbon atom designated by the numeral 3;

20

wherein the chiral carbon atom designated by the numeral 2 is in the syn configuration with respect to the chiral carbon atom designated by the numeral 1.

30. A compound according to claim 1, of the formula



wherein said compound is selected from the group consisting of:


10 the stereoisomer with R-configuration at the chiral carbon atom designated by the numeral 1 and R-configuration at the chiral carbon atom designated by the numeral 3;

the stereoisomer with S-configuration at the chiral carbon atom designated by the numeral 1 and S-configuration at the chiral carbon atom designated by the numeral 3;

15 the stereoisomer with R-configuration at the chiral carbon atom designated by the numeral 1 and S-configuration at the chiral carbon atom designated by the numeral 3; and

20 the stereoisomer with S-configuration at the chiral carbon atom designated by the numeral 1 and R-configuration at the chiral carbon atom designated by the numeral 3;


34. A compound according to claim 1, of the formula



with S-configuration at the chiral carbon atom designated by the numeral 1;

- 5 wherein the chiral carbon atom designated by the numeral 2 is in the syn configuration with respect to the chiral carbon atom designated by the numeral 1.


35. A compound according to claim 1, of the formula



- 10 wherein said compound is selected from the group consisting of:

the stereoisomer with R-configuration at the chiral carbon atom designated by the numeral 1 and R-configuration at the chiral carbon atom designated by the numeral 3;


39. A compound according to claim 1, of the formula



with S-configuration at the chiral carbon atom designated by the numeral 1;

- 5 wherein the chiral carbon atom designated by the numeral 2 is in the syn configuration with respect to the chiral carbon atom designated by the numeral 1.

40. A compound according to claim 1, of the formula




- 10 wherein said compound is selected from the group consisting of:

the stereoisomer with R-configuration at the chiral carbon atom designated by the numeral 1 and R-configuration at the chiral carbon atom designated by the numeral 3;

wherein the chiral carbon atom designated by the numeral 2 is in the syn configuration with respect to the chiral carbon atom designated by the numeral 1.


44. A compound according to claim 1, of the formula



with S-configuration at the chiral carbon atom designated by the numeral 1;

wherein the chiral carbon atom designated by the numeral 2 is in the syn configuration with respect to the chiral carbon atom designated by the numeral 1.

45. A compound according to claim 1, of the formula



wherein said compound is selected from the group consisting of: