


5

10 **TRICYCLIC AMIDE AND UREA COMPOUNDS USEFUL FOR**
INHIBITION OF G-PROTEIN FUNCTION AND FOR TREATMENT
OF PROLIFERATIVE DISEASES

BACKGROUND

15 International Publication Number WO92/11034, published July 9,
1992, discloses a method of increasing the sensitivity of a tumor to an
antineoplastic agent, which tumor is resistant to the antineoplastic agent,
by the concurrent administration of the antineoplastic agent and a
potentiating agent of the formula:



20 wherein the dotted line represents an optional double bond, X' is
hydrogen or halo, and Y' is hydrogen, substituted carboxylate or
substituted sulfonyl. For example, Y' can be, amongst others, -COOR'
wherein R' is C1 to C6 alkyl or substituted alkyl, phenyl, substituted
phenyl, C7 to C12 aralkyl or substituted aralkyl or -2, -3, or -4 piperidyl or
25 N-substituted piperidyl. Y' can also be, amongst others, SO₂R' wherein R'
is C1 to C6 alkyl, phenyl, substituted phenyl, C7 to C12 aralkyl or
substituted aralkyl. Examples of such potentiating agents include 11-(4-
piperidylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridines such as
Loratadine.

Oncogenes frequently encode protein components of signal transduction pathways which lead to stimulation of cell growth and mitogenesis. Oncogene expression in cultured cells leads to cellular transformation, characterized by the ability of cells to grow in soft agar and the growth of cells as dense foci lacking the contact inhibition exhibited by non-transformed cells. Mutation and/or overexpression of certain
5 oncogenes is frequently associated with human cancer.

To acquire transforming potential, the precursor of the Ras oncoprotein must undergo farnesylation of the cysteine residue located in a carboxyl-terminal tetrapeptide. Inhibitors of the enzyme that catalyzes this modification, farnesyl protein transferase, have therefore been suggested as anticancer agents for tumors in which Ras contributes to transformation. Mutated, oncogenic forms of ras are frequently found in many human cancers, most notably in more than 50% of colon and
10 pancreatic carcinomas (Kohl et al., Science, Vol. 260, 1834 to 1837, 1993).

In view of the current interest in inhibitors of farnesyl protein transferase, a welcome contribution to the art would be compounds useful for the inhibition of farnesyl protein transferase. Such a contribution is
15 provided by this invention.


SUMMARY OF THE INVENTION

Inhibition of farnesyl protein transferase by tricyclic compounds of this invention has not been reported previously. Thus, this invention
25 provides a method for inhibiting farnesyl protein transferase using tricyclic compounds of this invention which: (i) potently inhibit farnesyl protein transferase, but not geranylgeranyl protein transferase I, *in vitro*; (ii) block the phenotypic change induced by a form of transforming Ras which is a farnesyl acceptor but not by a form of transforming Ras engineered to be a
30 geranylgeranyl acceptor; (iii) block intracellular processing of Ras which is a farnesyl acceptor but not of Ras engineered to be a geranylgeranyl acceptor; and (iv) block abnormal cell growth in culture induced by transforming Ras. Several compounds of this invention have been demonstrated to have anti-tumor activity in animal models.

35 This invention provides a method for inhibiting the abnormal growth of cells, including transformed cells, by administering an effective amount of a compound of this invention. Abnormal growth of cells refers to cell growth independent of normal regulatory mechanisms (e.g., loss of

contact inhibition). This includes the abnormal growth of: (1) tumor cells (tumors) expressing an activated Ras oncogene; (2) tumor cells in which the Ras protein is activated as a result of oncogenic mutation in another gene; and (3) benign and malignant cells of other proliferative diseases in which aberrant Ras activation occurs.

Compounds useful in the claimed methods are represented by Formula 1.0:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

- 10 one of a, b, c and d represents N or NR⁹ wherein R⁹ is O⁻, -CH₃ or -(CH₂)_nCO₂H wherein n is 1 to 3, and the remaining a, b, c and d groups represent CR¹ or CR²; or
each of a, b, c, and d are independently selected from CR¹ or CR²;
each R¹ and each R² is independently selected from H, halo, -CF₃,
15 -OR¹⁰ (e.g., -OCH₃), -COR¹⁰, -SR¹⁰ (e.g., -SCH₃ and -SCH₂C₆H₅),
-S(O)_tR¹¹ (wherein t is 0, 1 or 2, e.g., -SOCH₃ and -SO₂CH₃), -N(R¹⁰)₂,
-NO₂, -OC(O)R¹⁰, -CO₂R¹⁰, -OCO₂R¹¹, -CN, -NR¹⁰COOR¹¹,
-SR¹¹C(O)OR¹¹ (e.g., -SCH₂CO₂CH₃), -SR¹¹N(R⁷⁵)₂ wherein each R⁷⁵
is independently selected from H and -C(O)OR¹¹ (e.g.,
20 -S(CH₂)₂NHC(O)O-t-butyl and -S(CH₂)₂NH₂), benzotriazol-1-yloxy,
tetrazol-5-ylthio, or substituted tetrazol-5-ylthio (e.g., alkyl substituted
tetrazol-5-ylthio such as 1-methyl-tetrazol-5-ylthio), alkynyl, alkenyl or
alkyl, said alkyl or alkenyl group optionally being substituted with halo,
-OR¹⁰ or -CO₂R¹⁰;
25 R³ and R⁴ are the same or different and each independently
represents H, any of the substituents of R¹ and R², or R³ and R⁴ taken

together represent a saturated or unsaturated C₅-C₇ fused ring to the benzene ring (Ring III);

R⁵, R⁶, R⁷ and R⁸ each independently represents H, -CF₃, -COR¹⁰, alkyl or aryl, said alkyl or aryl optionally being substituted with -OR¹⁰,
 5 -SR¹⁰, -S(O)_iR¹¹, -NR¹⁰COOR¹¹, -N(R¹⁰)₂, -NO₂, -COR¹⁰, -OCOR¹⁰,
 -OCO₂R¹¹, -CO₂R¹⁰, OPO₃R¹⁰ or one of R⁵, R⁶, R⁷ and R⁸ can be taken
 in combination with R⁴⁰ as defined below to represent -(CH₂)_r wherein r
 is 1 to 4 which can be substituted with lower alkyl, lower alkoxy, -CF₃ or
 aryl, or R⁵ is combined with R⁶ to represent =O or =S and/or R⁷ is
 10 combined with R⁸ to represent =O or =S;

R¹⁰ represents H, alkyl, aryl, or aralkyl (e.g., benzyl);


R¹¹ represents alkyl or aryl;

X represents N, CH or C, which C may contain an optional double
 bond (represented by the dotted line) to carbon atom 11;

15 the dotted line between carbon atoms 5 and 6 represents an
 optional double bond, such that when a double bond is present, A and B
 independently represent -R¹⁰, halo, -OR¹¹, -OCO₂R¹¹ or -OC(O)R¹⁰, and
 when no double bond is present between carbon atoms 5 and 6, A and B
 each independently represent H₂, -(OR¹¹)₂; H and halo, dihalo, alkyl and
 20 H, (alkyl)₂, -H and -OC(O)R¹⁰, H and -OR¹⁰, =O, aryl and H, =NOR¹⁰ or
 -O-(CH₂)_p-O- wherein p is 2, 3 or 4;

R represents R⁴⁰, R⁴², R⁴⁴, or R⁵⁴, as defined below;

R⁴⁰ represents H, aryl, alkyl, cycloalkyl, alkenyl, alkynyl or -D
 wherein -D represents



25

wherein R³ and R⁴ are as previously defined and W is O, S or NR¹⁰

wherein R¹⁰ is as defined above; said R⁴⁰ cycloalkyl, alkenyl and alkynyl
 groups being optionally substituted with from 1-3 groups selected from
 halo, -CON(R¹⁰)₂, aryl, -CO₂R¹⁰, -OR¹², -SR¹², -N(R¹⁰)₂,

30


-N(R¹⁰)CO₂R¹¹, -COR¹², -NO₂ or D, wherein -D, R¹⁰ and R¹¹ are as
 defined above and R¹² represents R¹⁰, -(CH₂)_mOR¹⁰ or -(CH₂)_qCO₂R¹⁰

wherein R¹⁰ is as previously defined, m is 1 to 4 and q is 0 to 4; said alkenyl and alkynyl R⁴⁰ groups not containing -OH, -SH or -N(R¹⁰)₂ on a carbon containing a double or triple bond respectively; or



R⁴⁰ represents phenyl substituted with a group selected from

5 -SO₂NH₂, -NH₂SO₂CH₃, -SO₂NHCH₃, -SO₂CH₃, -SOCH₃, -SCH₃, or -NH₂SO₂CF₃, preferably, said group is located in the para (p-) position of the phenyl ring; or


R⁴⁰ represents a group selected from




10



15



R42 represents



wherein R²⁰, R²¹ and R⁴⁶ are each independently selected from the group consisting of:

20

- (1) H;
- (2) $-(CH_2)_qSC(O)CH_3$ wherein q is 1 to 3 (e.g., $-CH_2SC(O)CH_3$);
- (3) $-(CH_2)_qOSO_2CH_3$ wherein q is 1 to 3 (e.g., $-CH_2OSO_2CH_3$);
- (4) $-OH$;
- (5) $-CS(CH_2)_w(\text{substituted phenyl})$ wherein w is 1 to 3 and the

25

substitutents on said substituted phenyl group are the same substitutents

as described below for said substituted phenyl (e.g., -C-S-CH₂-4-methoxyphenyl);

(6) -NH₂;

(7) -NHCBZ (wherein CBZ stands for carbonylbenzyloxy--i.e.,

5 CBZ represents -C(O)OCH₂C₆H₅);

(8) -NHC(O)OR²² wherein R²² is an alkyl group having from 1 to 5 carbon atoms (e.g., R²² is t-butyl thus forming -NHBOC wherein BOC stands for tert-butyloxycarbonyl--i.e., BOC represents -C(O)OC(CH₃)₃), or R²² represents phenyl substituted with 1 to 3 alkyl groups (e.g., 4-methylphenyl);

(9) alkyl (e.g., ethyl);

(10) -(CH₂)_kphenyl wherein k is 1 to 6, usually 1 to 4 and preferably 1 (e.g., benzyl);

(11) phenyl;

15 (12) substituted phenyl (i.e., phenyl substituted with from 1 to 3 substituents, preferably one) wherein the substituents are selected from the group consisting of: halo (e.g., Br, Cl, or I, with Br being preferred); NO₂; -OH; -OCH₃; -NH₂; -NHR²²; -N(R²²)₂; alkyl (e.g., alkyl having from 1 to 3 carbons with methyl being preferred); -O(CH₂)_tphenyl (wherein t is from 1 to 3 with 1 being preferred); and -O(CH₂)_tsubstituted phenyl (wherein t is from 1 to 3 with 1 being preferred); examples of substituted phenyls include, but are not limited to, p-bromophenyl, m-nitrophenyl, o-nitrophenyl, m-hydroxy-phenyl, o-hydroxyphenyl, methoxyphenyl, p-methylphenyl, m-methyl-phenyl, and -OCH₂C₆H₅;

25 (13) naphthyl;

(14) substituted naphthyl, wherein the substituents are as defined for substituted phenyl above;


(15) bridged polycyclic hydrocarbons having from 5 to 10 carbon atoms (e.g., adamantyl and norbornyl);

30 (16) cycloalkyl having from 5 to 7 carbon atoms (e.g., cyclopentyl, and cyclohexyl);

(17) heteroaryl (e.g., pyridyl, and pyridyl N-oxide);


(18) hydroxyalkyl (e.g., -(CH₂)_vOH wherein v is 1 to 3, such as, for example, -CH₂OH);

35 (19) substituted pyridyl or substituted pyridyl N-oxide wherein the substituents are selected from the substituents given above for said substituted phenyl, and said substituents are bound to a ring carbon by replacement of the hydrogen bound to said carbon;




(23) $-\text{NHC(O)}-(\text{CH}_2)_k\text{-phenyl}$ or $-\text{NH(O)}-(\text{CH}_2)_k\text{-substitued phenyl}$,
wherein said k is as defined above (i.e., 1-6, usually 1-4 and preferably
5 1);


(24) piperidine Ring V:



wherein R^{50} represents H, alkyl (e.g., methyl), alkylcarbonyl (e.g.,
 $\text{CH}_3\text{C(O)-}$), alkylloxycarbonyl (e.g., $-\text{C(O)O-t-C}_4\text{H}_9$, $-\text{C(O)OC}_2\text{H}_5$, and
10 $-\text{C(O)OCH}_3$), haloalkyl (e.g., trifluoromethyl), or $-\text{C(O)NH(R}^{10})$ wherein
 R^{10} is H or alkyl; Ring V includes




examples of Ring V include:




(25) $-\text{NHC(O)CH}_2\text{C}_6\text{H}_5$ or $-\text{NHC(O)CH}_2\text{-substitued-C}_6\text{H}_5$, for
example $-\text{NHC(O)CH}_2\text{-p-hydroxyphenyl}$, $-\text{NHC(O)CH}_2\text{-m-hydroxyphenyl}$,
and $-\text{NHC(O)CH}_2\text{-o-hydroxyphenyl}$;

20 (26) $-\text{NHC(O)OC}_6\text{H}_5$;



(30) -OC(O)-heteroaryl, for example




5 (31) -O-alkyl (e.g., -OCH₃); and

(32) -CF₃; or


R²⁰ and R²¹ taken together form a =O group and the remaining R⁴⁶ is as defined above; or

Two of R²⁰, R²¹ and R⁴⁶ taken together form piperidine Ring V




10


wherein R⁵⁰ represents H, alkyl (e.g., methyl), alkylcarbonyl (e.g., CH₃C(O)-), alkoxycarbonyl (e.g., -C(O)O-t-C₄H₉, -C(O)OC₂H₅, and -C(O)OCH₃), haloalkyl (e.g., trifluoro-methyl), or -C(O)NH(R¹⁰) wherein R¹⁰ is H or alkyl; Ring V includes




15

examples of Ring V include:






with the proviso that R^{46} , R^{20} , and R^{21} are selected such that the carbon atom to which they are bound does not contain more than one heteroatom (i.e., R^{46} , R^{20} , and R^{21} are selected such that the carbon atom to which they are bound contains 0 or 1 heteroatom);
 R^{44} represents




wherein R^{25} represents heteroaryl (e.g., pyridyl or pyridyl N-oxide) or aryl (e.g., phenyl and substituted phenyl); and R^{48} represents H or alkyl (e.g., methyl);

R^{54} represents an N-oxide heterocyclic group of the formula (i), (ii), (iii) or (iv):



wherein R^{56} , R^{58} , and R^{60} are the same or different and each is independently selected from H, halo, $-CF_3$, $-OR^{10}$, $-C(O)R^{10}$, $-SR^{10}$, $-S(O)_eR^{11}$ (wherein e is 1 or 2), $-N(R^{10})_2$, $-NO_2$, $-CO_2R^{10}$, $-OCO_2R^{11}$, $-OCOR^{10}$, alkyl, aryl, alkenyl or alkynyl, which alkyl may be substituted with $-OR^{10}$, $-SR^{10}$ or $-N(R^{10})_2$ and which alkenyl may be substituted with OR^{11} or SR^{11} ; or

R^{54} represents an N-oxide heterocyclic group of the formula (ia), (iia), (iiia) or (iva):






wherein Y represents N⁺-O⁻ and E represents N; or


R⁵⁴ represents an alkyl group substituted with one of said N-oxide heterocyclic groups (i), (ii), (iii), (iv), (ia), (iia), (iiaa) or (iva);

5 Z represents O or S such that R can be taken in combination with R⁵, R⁶, R⁷ or R⁸ as defined above, or R represents R⁴⁰, R⁴², R⁴⁴ or R⁵⁴.

Examples of R²⁰, R²¹, and R⁴⁶ for the above formulas include:




Examples of R²⁵ groups include:



wherein Y represents N or NO, R²⁸ is selected from the group consisting of: C₁ to C₄ alkyl, halo, hydroxy, NO₂, amino (-NH₂), -NHR³⁰, and -N(R³⁰)₂ wherein R³⁰ represents C₁ to C₆ alkyl.


- 5 Tricyclic compounds useful in the methods of this invention are described in: (1) U.S. 5,151,423; (2) U.S. 4,826,853; (3) U.S. 5,089,496; (4) WO 88/03138 published on May 5, 1988 (PCT/US87/02777); and (5) U.S. 5,104,876; the disclosures of each being incorporated herein by reference thereto. Those compounds within the scope of this invention
- 10 which are not described in these documents are described herein.

This invention also provides novel compounds of Formula 1.0 having the formula:




wherein all substituents are as defined for Formula 1.0

- 15 This invention further provides novel compounds of Formula 1.0 having the formula:



wherein all substituents are as defined for Formula 1.0

Additionally, this invention provides novel compounds of Formula 1.0 having the formula:



5

wherein all substituents are as defined for Formula 1.0.


Compounds of Formula 5.2 include compounds wherein the substituents R^{20} , R^{21} , and R^{46} are selected such that when one of said substituents R^{20} , R^{21} , and R^{46} (e.g., R^{46}) is selected from the group consisting of: (1) H, (2) -OH, (3) -NH₂, (4) -NHC(O)OR²², (5) alkyl, (6) phenyl, (7) heteroaryl, (8) hydroxyalkyl, (9) substituted pyridyl, (10) substituted phenyl and (11) -O-alkyl, then the remaining two of said substituents R^{20} , R^{21} and R^{46} (e.g., R^{20} and R^{21}) cannot both be H when:

10 (a) R^1 and R^2 are both H, and (b) the double bond between C-5 and C-6 is absent, and (c) both A and B are H₂, and (d) R^4 is H, and (e) R^3 is H or Cl at C-8. Compounds of Formula 5.2 also include compounds wherein

15 when R^{46} is a group (1) to (11) defined above then R^{20} and R^{21} cannot


- both be H when: R¹ and R² are both H, and both A and B are H or H₂. Compounds of Formula 5.2 further include compounds wherein when R⁴⁶ is a group (1) to (11) defined above then R²⁰ and R²¹ cannot both be H when R¹ and R² are both H. Compounds of Formula 5.2 also include
- 5 compounds wherein two of R²⁰, R²¹ and R⁴⁶ are not H when R¹ and R² are both H.

This invention further provides novel compounds of Formula 1.0 having the formula:



or

10



wherein all the substituents are as defined for Formula 1.0. Preferably R²⁵ represents heteroaryl.

This invention also provides a method for inhibiting tumor growth by administering an effective amount of the tricyclic compounds, described herein, to a mammal (e.g., a human) in need of such treatment. In particular, this invention provides a method for inhibiting the growth of
5 tumors expressing an activated Ras oncogene by the administration of an effective amount of the above described compounds. Examples of tumors which may be inhibited include, but are not limited to, lung cancer (e.g., lung adenocarcinoma), pancreatic cancers (e.g., pancreatic carcinoma such as, for example, exocrine pancreatic carcinoma), colon cancers (e.g.,
10 colorectal carcinomas, such as, for example, colon adenocarcinoma and colon adenoma), myeloid leukemias (for example, acute myelogenous leukemia (AML)), thyroid follicular cancer, myelodysplastic syndrome (MDS), bladder carcinoma and epidermal carcinoma.


It is believed that this invention also provides a method for
15 inhibiting proliferative diseases, both benign and malignant, wherein Ras proteins are aberrantly activated as a result of oncogenic mutation in other genes--i.e., the Ras gene itself is not activated by mutation to an oncogenic form--with said inhibition being accomplished by the administration of an effective amount of the tricyclic compounds described
20 herein, to a mammal (e.g., a human) in need of such treatment. For example, the benign proliferative disorder neurofibromatosis, or tumors in which Ras is activated due to mutation or overexpression of tyrosine kinase oncogenes (e.g., neu, src, abl, lck, and fyn), may be inhibited by the tricyclic compounds described herein.

25 The compounds of this invention inhibit farnesyl protein transferase and the farnesylation of the oncogene protein Ras. This invention further provides a method of inhibiting ras farnesyl protein transferase, in mammals, especially humans, by the administration of an effective amount of the tricyclic compounds described above. The administration of
30 the compounds of this invention to patients, to inhibit farnesyl protein transferase, is useful in the treatment of the cancers described above.

The tricyclic compounds useful in the methods of this invention inhibit the abnormal growth of cells. Without wishing to be bound by theory, it is believed that these compounds may function through the
35 inhibition of G-protein function, such as ras p21, by blocking G-protein isoprenylation, thus making them useful in the treatment of proliferative diseases such as tumor growth and cancer. Without wishing to be bound by theory, it is believed that these compounds inhibit ras farnesyl protein


transferase, and thus show antiproliferative activity against ras transformed cells.

This invention also provides a process for producing 3-nitro substituted compounds. The process comprises reacting one molar equivalent of a compound:




wherein R^1 , R^2 , R^3 , R^4 , A, B, a, b, d, and the dotted lines are as defined for Formula 1.0; and R^{65} represents H or $-OR^{66}$ wherein R^{66} represents alkyl (e.g., C_1 to C_4 alkyl, preferably ethyl); with one molar equivalent of a
 10 nitrating reagent, said nitrating reagent being preformed (i.e., prepared first) by mixing, at cold temperature (e.g., at 0°C) equimolar amounts of tetrabutyl ammonium nitrate with trifluoroacetic anhydride; the reaction of the nitrating reagent with the compound of Formula 1.0g taking place in a suitable aprotic solvent (e.g., methylene chloride, chloroform, toluene or tetrahydrofuran); said reaction with said nitrating reagent being conducted
 15 at a temperature and for a period of time sufficient to allow the reaction to proceed at a reasonable rate to produce the desired final 3-nitro compound of Formula 1.0h (described below)--i.e., the reaction of the compound of Formula 1.0g with said nitrating reagent is conducted at an
 20 initial temperature of 0°C , and said reaction temperature is thereafter allowed to rise to about 25°C during the reaction time period. The reaction usually proceeds overnight to completion, i.e., the reaction usually proceeds for about 16 hours. The reaction can be conducted within a temperature of 0°C to about 25°C during a time period of about
 25 10 to about 24 hours. Preferably the reaction is initially conducted at 0°C

and the temperature is allowed to warm up to 25°C. The reaction produces the 3-nitro compound:



The compound of Formula 1.0h can then be converted to other 3-substituted products by methods well known to those skilled in the art. For example, the 3-nitro compounds can be converted to 3-amino, 3-halo, 3-cyano, 3-alkyl, 3-aryl, 3-thio, 3-arylalkyl, 3-hydroxyl, and 3-OR⁶⁷ wherein R⁶⁷ is alkyl or aryl. The 3-substituted compounds can then be converted to final products (wherein R⁶⁵ is R⁴² or R⁴⁴) by the procedures described herein.

This invention also provides a process for producing 3-nitro compounds of the formula:



by producing a compound of Formula 1.0h from 1.0g as described above; and then hydrolyzing the compound of Formula 1.0h by dissolving the


compound of Formula 1.0h in a sufficient amount of concentrated acid (e.g., concentrated HCl or aqueous sulfuric acid), and heating the resulting mixture to a temperature sufficient to remove (hydrolyze) the -C(O)R⁶⁵ substituent, for example, heating to reflux or to a temperature of about 100°C. This hydrolysis process is exemplified in Preparative

5 Example 28.


The compound of Formula 1.0i can then be converted to other 3-substituted compounds as discussed above for the compounds of Formula 1.0h. The compounds of Formula 1.0i can then be converted to

10 compounds of this invention by the methods described herein.

This invention also provides a process for producing compounds of the formula:



by reacting one molar equivalent a compound of formula:



15

with one molar equivalent of a nitrating reagent, said nitrating reagent being preformed (i.e., prepared first) by mixing, at cold temperature (e.g., at 0°C) equimolar amounts of tetrabutyl ammonium nitrate with trifluoroacetic anhydride; the reaction of the nitrating reagent with the


20 compound of Formula 1.0k taking place in a suitable aprotic solvent (e.g., methylene chloride, chloroform, toluene or tetrahydrofuran); said reaction with said nitrating reagent being conducted at a temperature and for a period of time sufficient to allow the reaction to proceed at a reasonable rate to produce the desired final 3-nitro compound of Formula 1.0j--i.e.,

25 the reaction of the compound of Formula 1.0k with said nitrating reagent is conducted at an initial temperature of 0°C, and said reaction temperature


is thereafter allowed to rise to about 25°C during the reaction time period. The reaction usually proceeds overnight to completion, i.e., the reaction usually proceeds for about 16 hours. The reaction can be conducted within a temperature of 0°C to about 25°C during a time period of about 10 to about 24 hours. Preferably the reaction is initially conducted at 0°C and the temperature is allowed to warm up to 25°C. In Formulas 1.0j and 1.0k, R¹, R², R³, R⁴, A, B, a, b, d, and the dotted lines are as defined for Formula 1.0

The compounds of Formula 1.0j can be converted to compounds of Formula 1.0h by methods described below. Also, as discussed above for the compounds of Formula 1.0h, the compounds of Formula 1.0j can be converted to other 3-substituted compounds wherein the substituents are those discussed above for Formula 1.0h.


The compounds of Formula 1.0j can be converted to compounds of Formula 1.0m:






wherein R⁶⁸ is H or -COOR^a wherein R^a is a C₁ to C₃ alkyl group (preferably R⁶⁸ is H), by reducing a compound of Formula 1.0j with a suitable reducing agent (such as sodium borohydride) in a suitable solvent (such as ethanol or methanol) at a suitable temperature to allow the reaction to proceed at a reasonable rate (e.g., 0 to about 25°C); reacting the resulting product (Formula 1.0j wherein the =O has been reduced to a -OH) with a chlorinating agent (e.g., thionyl chloride) in an suitable organic solvent (e.g., benzene, toluene or pyridine) at a suitable temperature to allow the reaction to proceed at a reasonable rate (e.g., about -20 to about 20°C, preferably at -15°C, see, for example Preparative Example 7) to produce a compound of Formula 1.0n:






and reacting a compound of Formula 1.0n with a compound of the formula:



- 5 wherein R^{68} is as previously defined, and is preferably H, in a suitable organic solvent (such as tetrahydrofuran or toluene) containing a suitable base (such as triethylamine or N-methylmorpholine) at a suitable temperature to allow the reaction to proceed at a reasonable rate (e.g., 25 to about 120°C).
- 10 Compounds of Formula 1.0m can be converted to compounds of this invention by the methods disclosed herein. Also, as discussed above for the compounds of Formula 1.0h, the compounds of Formula 1.0m can be converted to other 3-substituted compounds wherein the substituents are those discussed above for Formula 1.0h.
- 15 This invention also provides novel compounds (produced in the above described processes as intermediates to the compounds of this invention) having the formulas:







5

wherein all substituents are as defined herein.

10 Preferably, for the intermediate compounds of the processes of this invention, R¹ and R² are H; R³ is halo, most preferably Cl, in the C-8 position; R⁴ is H; and A and B are H when the double between C-5 and C-6 is present, and A and B are H₂ when the bond between C-5 and C-6 is a single bond (most preferably the bond between C-5 and C-6 is a single bond). Those skilled in the art will appreciate that Rings I, II, and/or III can be further substituted, as described herein, to produce the desired compounds of the invention.

15

Examples of such novel intermediate compounds include:



5 DETAILED DESCRIPTION OF THE INVENTION

As used herein, the following terms are used as defined below unless otherwise indicated:

M⁺-represents the molecular ion of the molecule in the mass spectrum;

10 MH⁺-represents the molecular ion plus hydrogen of the molecule in the mass spectrum;


Bu-represents butyl;

Et-represents ethyl;


Me-represents methyl;

15 Ph-represents phenyl;

benzotriazol-1-yloxy represents



1-methyl-tetrazol-5-ylthio represents



alkyl-(including the alkyl portions of alkoxy, alkylamino and dialkylamino)-represents straight and branched carbon chains and contains from one to twenty carbon atoms, preferably one to six carbon atoms;

5 alkanediyl-represents a divalent, straight or branched hydrocarbon chain having from 1 to 20 carbon atoms, preferably 1 to 6 carbon atoms, the two available bonds being from the same or different carbon atoms thereof, e.g., methylene, ethylene, ethylidene,
10 -CH₂CH₂CH₂-, -CH₂CHCH₃-, -CHCH₂CH₃-, etc.

cycloalkyl-represents saturated carbocyclic rings branched or unbranched of from 3 to 20 carbon atoms, preferably 3 to 7 carbon atoms;

heterocycloalkyl-represents a saturated, branched or unbranched carbocyclic ring containing from 3 to 15 carbon atoms, preferably from 4 to 6 carbon atoms, which carbocyclic ring is interrupted by 1 to 3 hetero groups selected from -O-, -S- or -NR¹⁰-(suitable heterocycloalkyl groups including 2- or 3-tetrahydrofuranyl, 2- or 3- tetrahydrothienyl, 2-, 3- or 4-
15 piperidinyl, 2- or 3-pyrrolidinyl, 2- or 3-piperizinyl, 2- or 4-dioxanyl, etc.);


alkenyl-represents straight and branched carbon chains having at least one carbon to carbon double bond and containing from 2 to 12 carbon atoms, preferably from 2 to 6 carbon atoms and most preferably from 3 to 6 carbon atoms;

alkynyl-represents straight and branched carbon chains having at least one carbon to carbon triple bond and containing from 2 to 12 carbon atoms, preferably from 2 to 6 carbon atoms;


25 aryl (including the aryl portion of aryloxy and aralkyl)-represents a carbocyclic group containing from 6 to 15 carbon atoms and having at least one aromatic ring (e.g., aryl is a phenyl ring), with all available substitutable carbon atoms of the carbocyclic group being intended as possible points of attachment, said carbocyclic group being optionally substituted (e.g., 1 to 3) with one or more of halo, alkyl, hydroxy, alkoxy, phenoxy, CF₃, amino, alkylamino, dialkylamino, -COOR¹⁰ or -NO₂; and
30

halo-represents fluoro, chloro, bromo and iodo; and

heteroaryl-represents cyclic groups, optionally substituted with R^3 and R^4 , having at least one heteroatom selected from O, S or N, said heteroatom interrupting a carbocyclic ring structure and having a sufficient number of delocalized pi electrons to provide aromatic character, with the aromatic heterocyclic groups preferably containing from 2 to 14 carbon atoms, e.g., 2-, 3- or 4-pyridyl or pyridyl N-oxide (optionally substituted with R^3 and R^4), wherein pyridyl N-oxide can be represented as:






Reference to the position of the substituents R^1 , R^2 , R^3 , and R^4 is based on the numbered ring structure:



For example, R^1 can be at the C-4 position and R^2 can be at the C-2 or C-3 position. Also, for example, R^3 can be at the C-8 position and R^4 can be at the C-9 position.

Representative structures of Formula 1.0 include but are not limited to:





Preferably, for the compounds of Formula 1.0 (including 1.0a to 1.0d):

each of a, b, c, and d are C (carbon); or

- 5 one of a, b, c and d (most preferably a) represents N or NO, most preferably N, and the remaining a, b, c and d groups represent CR¹ or CR²;

- each R¹ and each R² is independently selected from H, halo (e.g., Cl, Br and F), -CF₃, -OR¹⁰ (e.g., hydroxy and alkoxy (e.g., -OCH₃)), alkyl (e.g., methyl and t-butyl, said alkyl group being optionally substituted with halo), benzotriazol-1-yloxy, -S(O)_tR¹¹ (e.g., -SCH₂CH₃), -SR¹¹C(O)OR¹¹ (e.g., -SCH₂CO₂CH₃), -SR¹⁰ (e.g., R¹⁰ represents -CH₂C₆H₅) and 1-methyl-tetrazol-5-ylthio; most preferably R¹ and R² are independently H, halo, -CF₃, lower alkyl (e.g., C₁ to C₄, more preferably methyl) or
- 10 benzotriazol-1-yloxy; more preferably R¹ is Cl or H, and R² is H, Cl or Br; still more preferably R¹ is at the C-4 position, and R² is at the C-3 position; even more preferably R² is Br or H;

- R³ and R⁴ are the same or different and each independently represents H, halo, -CF₃, -OR¹⁰, -COR¹⁰, -SR¹⁰, -S(O)_tR¹¹ (wherein t is
- 20 0, 1 or 2), -N(R¹⁰)₂, -NO₂, -OC(O)R¹⁰, -CO₂R¹⁰, -OCO₂R¹¹, -CN, -NR¹⁰COOR¹¹, alkynyl, alkenyl or alkyl, said alkyl or alkenyl group optionally being substituted with halo, -OR¹⁰ or -CO₂R¹⁰; most preferably R³ and R⁴ independently represent H, halo, -CF₃, -OR¹⁰ or alkyl (said alkyl group being optionally substituted with halo); more preferably R³ and R⁴ independently represent H or halo (e.g., Cl, Br, or F); even more
- 25


preferably R^3 is at the C-8 position and R^4 is at the C-9 position; still more preferably R^3 is Cl at the C-8 position and R^4 is H at the C-9 position;






R^5 , R^6 , R^7 and R^8 each independently represents H, $-CF_3$ or alkyl (said alkyl optionally being substituted with $-OR^{10}$); most preferably R^5 , R^6 , R^7 and R^8 independently represent H and alkyl, and more preferably H;

when the optional double bond between carbon atoms 5 and 6 is present, A and B independently represent H, $-R^{10}$ or $-OR^{10}$, most preferably H, lower alkyl (C_1 to C_4) and alkyloxy (i.e., R^{10} represents alkyl), more preferably H and $-OH$, and still more preferably H; and when no double bond is present between carbon atoms 5 and 6, A and B each independently represent H_2 , $-(OR^{10})_2$, alkyl and H, $(alkyl)_2$, - H and $-OR^{10}$ or $=O$, most preferably H_2 , -H and $-OH$, or $=O$, and more preferably A represents H_2 and B represents H_2 or $=O$;


R represents R^{42} or R^{44} ; and
Z represents O or S, and most preferably O.

Compounds of Formula 5.0 include:












and




Compounds of Formula 5.1 include:











Compounds of Formula 5.2 additionally include:




Compounds of Formula 5.3 include:












and



Compounds of formula 5.3A include:







For the compounds of Formulas 5.0, 5.0a-5.0g, 5.1, 5.1a-5.1g, 5.2, 5.2a-5.2b, 5.3, 5.3a-5.3g, 5.3A, 5.3Aa-5.3Ag, and 5.3B, the definitions of the substituents are as defined for Formula 1.0.

- 5 Preferably, for compounds of Formulas 5.0, 5.0a-5.0g, 5.1, 5.1a-5.1g, 5.2, and 5.2a-5.2b, R^{46} is selected from piperidine Ring V, heteroaryl, phenyl, substituted phenyl, substituted pyridyl or substituted pyridyl N-oxide, and R^{20} and R^{21} are independently selected from H or alkyl. Most preferably, R^{46} is pyridyl, pyridyl N-oxide or piperidine Ring V.
- 10 More preferably, R^{46} is pyridyl, pyridyl N-oxide or piperidine Ring V and both R^{20} and R^{21} are hydrogen or both R^{20} and R^{21} are alkyl (still more preferably methyl).

- 15 Even more preferably, R^{46} is selected from 3-pyridyl, 4-pyridyl, 3-pyridyl N-oxide, 4-pyridyl N-oxide, 4-N-methylpiperidinyl, 3-N-methylpiperidinyl, 4-N-acetylpiperidinyl or 3-N-acetylpiperidinyl, and both R^{20} and R^{21} are hydrogen or both R^{20} and R^{21} are alkyl (still even more preferably methyl). Even still more preferably, R^{46} is selected from 3-pyridyl, 3-pyridyl N-oxide, 4-pyridyl, and 4-pyridyl N-oxide, and both R^{20} and R^{21} are hydrogen or both R^{20} and R^{21} are methyl.







- 20 Examples of the R^{42} groups include:























Preferably for the compounds of Formulas 5.3, 5.3a-5.3g, 5.3A, 5.3Aa-5.3Ag, and 5.3B, R²⁵ represents phenyl, 2-pyridyl, 3-pyridyl or 4-pyridyl, and most preferably 3-pyridyl. More preferably, R⁴⁸ represents H or methyl and still more preferably H.







Representative compounds of the invention include:

































and



Preferred compounds of this invention are selected from the group consisting of compounds of Examples: 1, 2, 3, 4, 5, 6, 19, 42, 43, 44, 45, 46, 47, 48, 49, 75, 76, 78, 82, 83, 84, 85, 89, 121, 180, 182, 183, 184, 187
5 (6.7 and 6.8), 192, 196, 197, 198, 200, 201, 206, 222, 223, 224, 225, 226, 227, 233, 234, 236, 239, 246, 247, 248, 249, 250, 251, 261, 262, 266, 267, 269, 273, 276, 283, 285, 286, 287, 288, 289, 291, 292, 293, 299, 300, 301, 303, 307, 309, 311, 312, 313, 314, 316, 350, 351, 352, 354 and 356.

10 More preferred compounds of this invention are selected from the group consisting of compounds of Examples: 1, 2, 42, 43, 75, 78, 82, 180, 183, 187 (6.7 and 6.8), 196, 197, 198, 200, 222, 223, 224, 227, 233, 234, 246, 247, 248, 249, 250, 251, 266, 269, 273, 283, 285, 286, 291, 292, 300, 301, 303, 307, 311, 312, 313, 314, 350, 351, 352, 354 and 356.

15 Even more preferred compounds of this invention are selected from the group consisting of compounds of Examples: 82, 197, 233, 246, 266, 312, 351, 352, 354 and 356.

Lines drawn into the ring systems indicate that the indicated bond may be attached to any of the substitutable ring carbon atoms.

20 Certain compounds of the invention may exist in different isomeric (e.g., enantiomers and diastereoisomers) forms. The invention contemplates all such isomers both in pure form and in admixture, including racemic mixtures. Enol forms are also included.

25 Certain tricyclic compounds will be acidic in nature, e.g. those compounds which possess a carboxyl or phenolic hydroxyl group. These compounds may form pharmaceutically acceptable salts. Examples of


such salts may include sodium, potassium, calcium, aluminum, gold and silver salts. Also contemplated are salts formed with pharmaceutically acceptable amines such as ammonia, alkyl amines, hydroxyalkylamines, N-methylglucamine and the like.

5 Certain basic tricyclic compounds also form pharmaceutically acceptable salts, e.g., acid addition salts. For example, the pyrido-
nitrogen atoms may form salts with strong acid, while compounds having
basic substituents such as amino groups also form salts with weaker
acids. Examples of suitable acids for salt formation are hydrochloric,
10 sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric,
succinic, ascorbic, maleic, methanesulfonic and other mineral and
carboxylic acids well known to those in the art. The salts are prepared by
contacting the free base form with a sufficient amount of the desired acid
to produce a salt in the conventional manner. The free base forms may be
15 regenerated by treating the salt with a suitable dilute aqueous base
solution such as dilute aqueous sodium hydroxide, potassium carbonate,
ammonia and sodium bicarbonate. The free base forms differ from their
respective salt forms somewhat in certain physical properties, such as
solubility in polar solvents, but the acid and base salts are otherwise
20 equivalent to their respective free base forms for purposes of the
invention.

 All such acid and base salts are intended to be pharmaceutically
acceptable salts within the scope of the invention and all acid and base
salts are considered equivalent to the free forms of the corresponding
25 compounds for purposes of the invention.


 Compounds of Formula 1.0 wherein R is $-N(R^{10})_2$, and compounds
of Formulas 5.3, 5.3A and 5.3B can be prepared by reacting compound
405.00 (described below) with an isocyanate ($R^{10}-N=C=O$) in a solvent
such as DMF, dichloromethane or THF in accordance with methods
30 known in the art.

 The following processes may be employed to produce compounds
of the invention--i.e., compounds of Formula 1.0 represented by
compounds of Formulas 5.0, 5.1, 5.2 and 5.3. For purposes of describing
the processes, the compounds are represented by Formula 400.00:




wherein R represents R⁴² or R⁴⁴, and all other substituents are as described herein.

- 5 A. A compound of Formula 405.00 may be coupled with a compound of the formula RCOOH in the presence of coupling agent such as 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (DEC), N,N'-dicyclohexylcarbodiimide (DCC) or N,N'-carbonyl-diimidazole (CDI) to produce compounds of Formula 400.00:




- 10 The reaction is usually conducted in an inert solvent such as tetrahydrofuran, DMF or methylene chloride at a temperature between about 0°C and reflux, preferably at about room temperature. When the coupling agent is DCC or DEC, the reaction is preferably run in the presence of 1-hydroxybenzotriazole (HOBT). Method A is the method of
15 choice for preparing compounds of this invention.

B. A compound of Formula 405.00 may also be reacted with a compound of Formula 410.00 in the presence of base to produce compounds of Formula 400.00:



- 5 Representative examples of appropriate bases are pyridine and triethylamine. L designates a suitable leaving group. For example, a compound of compound 410.00 may be an acyl halide (e.g., L represents halo) or an acyl anhydride, (e.g., L is -O-C(O)-R). The leaving group may also be alkoxy, in which case the compounds of Formula 400.00 may be produced by refluxing a compound of Formula 405.00 with an excess of a compound of Formula 410.00.


Compounds of Formula 405.00 may be prepared by cleaving the group COOR^a from the corresponding carbamates 415.00, for example, via acid hydrolysis (e.g., HCl) or base hydrolysis (e.g., KOH):




- 15 wherein R^a is a group which does not prevent the cleavage reaction, e.g., R^a is an optionally substituted alkyl such as ethyl.

- Alternatively, depending upon the nature of R^a, as determined by one skilled in the art, Compound 415.00 may be treated with an organometallic reagent (e.g., CH₃Li), a reductive reagent (e.g., Zn in acid), etc., to form compounds of Formula 405.00.


Compound 415.00 may be prepared from the N-alkyl compound shown as Formula 420.00 below, in the manner disclosed in U.S. Patents 4,282,233 and 4,335,036.



- It also will be apparent to one skilled in the art that there are other methods for converting Compound 420.00 to Compound 405.00. For example, treatment of Compound 420.00 with BrCN via von Braun
- 5 reaction conditions would provide nitrile 420.00a. Subsequent hydrolysis of the nitrile under either aqueous basic or acidic conditions would produce Compound 405.00. This method is preferable when there is substitution on the piperidine or piperazine ring.



157



Compound 420.00 is prepared as described in part B above.

PREPARATION OF SINGLE BOND COMPOUNDS


- 5 Compounds of Formula 400.00, wherein X is carbon and the bond to carbon 11 (C-11) is a single bond, can be prepared by reducing compounds of Formula 405.00, wherein X is carbon and the bond to C-11 is a double bond, with lithium aluminum hydride in tetrahydrofuran. Conversion to final products can be done following the process described
- 10 above for conversion of compounds of Formula 405.00 to compounds of Formula 400.00.

PREPARATION OF DOUBLE BOND COMPOUNDS


- 15 Compounds of Formula 400.00, wherein X is a carbon atom having an exocyclic double bond to carbon 11, may be prepared from compound 420.00 as described above. Compounds of Formula 420.00 may be produced by the methods disclosed generally in U.S. Patent 3,326,924 or alternatively may be prepared by a ring closure reaction, wherein the desired cycloheptene ring is formed by treating compound 425.00 with a
- 20 super acid. Suitable super acids for this purpose include, for example, HF/BF₃, CF₃SO₃H (triflic acid), CH₃SO₃H/BF₃, etc. The reaction can be performed in the absence of, or with, an inert co-solvent such as CH₂Cl₂. The temperature and time of the reaction vary with the acid employed. For example, with HF/BF₃ as the super acid system the temperature may be
- 25 controlled so as to minimize side reactions, such as HF addition to the exocyclic double bond. For this purpose, the temperature is generally in the range of from about +5°C to -50°C. With CF₃SO₃H as the super acid

system, the reaction may be run at elevated temperatures, e.g., from about 25°C to about 150°C and at lower temperatures but the reaction then takes longer to complete.


Generally the super acid is employed in excess, preferably in amounts of from about 1.5 to about 30 equivalents.



A ketone compound of Formula 425.00 may be formed by hydrolysis of 430.00, e.g., such as by reacting a Grignard intermediate of Formula 430.00 with an aqueous acid (e.g., aqueous HCl). I^a in Formula 430.00 represents chloro, bromo or iodo.




- The Grignard intermediate 430.00 is formed by the reaction of the cyano compound 435.00 with an appropriate Grignard reagent 440.00 prepared from 1-alkyl-4-halopiperidine. The reaction is generally performed in an inert solvent, such as ether, toluene, or tetrahydrofuran, under general Grignard conditions e.g., temperature of from about 0°C to about 75°C. Alternatively, other organometallic derivatives of the 1alkyl-4-halo piperidine can be employed.




- The cyano compound of Formula 435.00 is produced by converting the tertiary butyl amide of Formula 445.00 with a suitable dehydrating agent, such as POCl₃, SOCl₂, P₂O₅, toluene sulfonyl chloride in pyridine, oxalyl chloride in pyridine, etc. This reaction can be performed in the absence of or with a co-solvent, such as xylene.
- The dehydrating agent such as POCl₃ is employed in equivalent amounts or greater and preferably in amounts of from about 2 to about 15 equivalents. Any suitable temperature and time can be employed for

performing the reaction, but generally heat is added to accelerate the reaction. Preferably the reaction is performed at or near reflux.




- 5 The tert-butylamide of Formula 445.00 may be produced by reaction of a compound of Formula 450.00a and 450.00b, in the presence of base, wherein G is chloro, bromo or iodo.




- 10 The compound of Formula 450.00a may be formed by hydrolysis of the corresponding nitrile wherein the appropriate cyanomethyl pyridine, such as 2-cyano-3-pyridine, is reacted with a tertiary butyl compound in acid, such as concentrated sulfuric acid or concentrated sulfuric acid in glacial acetic acid. Suitable tertiary butyl compounds include, but are not limited to, t-butyl alcohol, t-butyl chloride, t-butyl bromide, t-butyl iodide, isobutylene or any other compound which under hydrolytic conditions
- 15 forms t-butyl carboxamides with cyano compounds. The temperature of the reaction will vary depending upon the reactants, but generally the reaction is conducted in the range of from about 50°C to about 100°C with t-butyl alcohol. The reaction may be performed with inert solvents, but is usually run neat.

- 20 An alternative process for the formation of compounds of Formula 400.00a may involve direct cyclization of Compound 455.00 as shown below.




Cyclization to form the cycloheptene ring may be accomplished with a strong acid (e.g., triflic, polyphosphoric, HF/BF₃), and may be performed in an inert solvent, such as ether, toluene or THF. The temperature and time may vary with the acid employed, as described in process A above.

Compounds of Formula 455.00 wherein Z = O or S may be prepared by treating a compound of Formula 425.00 with an appropriate acyl halide or acyl anhydride of formula 410.00. Most preferably this reaction is run in the presence of a good nucleophile, such as LiI, in the appropriate solvent, such as toluene, dioxane or xylene, and at a temperature ranging from 50-150°C, preferably 100-120°C.




A second method of preparing compounds of Formula 455.00 involves reacting an unsubstituted piperidylidene compound of Formula 460.00 with the appropriate acyl halide or acyl anhydride of Formula 410.00 in the presence of base, such as pyridine or triethylamine. Alternatively, if L = OH in compound 410.00, then coupling of compound

460.00 with compound 410.00 may require use of a conventional coupling reagent, such as DCC or CDI.



- Compounds of Formula 460.00 may be produced from the
- 5 corresponding carbamates of Formula 465.00, via acid hydrolysis, using for example, aqueous hydrochloric acid, or base hydrolysis using for example, potassium hydroxide. Alternatively, some compounds can be prepared by treating the carbamate, Formula 465.00, with an
 - 10 organometallic reagent, such as methyl lithium or a reductive reagent, such as zinc in acid, etc., depending upon the nature of the R^a group. For example, if R^a is a simple alkyl group, CO_2R^a may be cleaved by alkaline hydrolysis at $100^\circ C$.



- The carbamate compounds of Formula 465.00 may be prepared
- 15 from the appropriate alkyl compound of Formula 425.00 by treatment with a chloroformate, preferably in an inert solvent, such as toluene, with warming to approximately $80^\circ C$. Other alternative methods are available


for the conversion of 425.00 to 455.00 as previously described (e.g. Von Braun reaction conditions). Compounds of Formula 425.00 may be prepared as described above.

5

SUBSTITUTION ON THE PYRIDINE RING

Various methods can be used as described in WO 88/03138 to provide compounds which are substituted on the pyridine ring, i.e., in positions 2-, 3- and or 4- positions of the tricyclic ring system. For example, the cyclization methods described on pages 20-30 of WO 88/03138 can already have the appropriate substituents on the pyridine ring in place. A variety of substituted pyridines are known in the literature and can be employed in these syntheses. Alternatively, the azaketone of Formula XIX (from page 27 of WO 88/03138)

10




(XIX) p.27 WO88/03138

15 wherein R¹ and R² are both H can be converted to the appropriately substituted azaketone wherein R¹ and R² are non-H substituents. If both R¹ and R² are desired to be non-H substituents the procedure would be repeated.

The azaketone is thus reacted with an oxidizing agent such as meta-chloroperoxybenzoic acid (MCPBA) or hydrogen peroxide to produce the corresponding compound in which the nitrogen of the pyridine ring is as an N-oxide:

20




wherein one of a', b', c' or d' is N→O and the others are CH or CR¹ or CR². This reaction is normally run at temperatures from -15°C to reflux,


25

more typically at about 0°C. The reaction is preferably conducted in an inert solvent such as methylene chloride for MCPBA or acetic acid for hydrogen peroxide.

- 5 The azaketone N-oxide of Formula 470.00a can then be reacted with a chlorinating agent such as SO_2Cl_2 or SOCl_2 to form a compound of Formula 470.00b. Typically, this reaction results in monosubstitution of Cl in the ortho or para-position relative to the N atom of the ring.




- 10 To provide the disubstituted products, steps 1 and 2 above are repeated.




Typically, the resulting disubstituted compounds have Cl ortho and para relative to the N atom of the pyridine ring.

- 15 The mono or disubstituted compounds of Formulas 470.00b and 470.00c above can be reacted with various nucleophiles such as alkoxides, amines, thiols, etc. This will result in compounds where one or both of the Cl substituents are replaced by the nucleophile to provide a compound of Formula 470.00d or a compound easily converted to Formula 470.00d.



The substituted ketone of Formula 470.00 can then be converted to the desired compound by the methods described above and in WO 88/03138 and in U.S. Patent No. 3,326,924.

- 5 Formula 405.00, wherein R¹ or R² are chlorine, can be made by the following alternate process.



- 10 The N-oxide of Formula 415.00 can be treated with POCl₃ to form a compound of Formula 415.01. Typically, this reaction results in mono-substitution of Cl in the ortho or para position relative to the N atom of the ring.

- 15 Alternatively, the Cl substituted azaketones of Formula 470.00b or 470.00c above can be converted to the corresponding derivatives of Formula 405.00 above wherein R¹ and/or R² is Cl by methods analogous to those described above. At this point the Cl substituent(s) can be displaced by an appropriate nucleophile to provide the desired substituent. Suitable nucleophiles include alkoxide, amines, thiols, etc. This reaction usually requires higher temperatures (e.g., from about 100° to about 200°C) than the displacement reaction to produce ketone 470.00d above. It is also usually conducted in a sealed vessel in an inert solvent. The compound of Formula 405.00 is then converted to a compound of Formula 400.00 as described above.
- 20


Various electrophilic species can also be added to the pyridine ring from the corresponding halo-substituted pyridine (Formula 405.00 wherein R¹ is halo, preferably bromo or iodo). Transmetalation of the halo derivative using an alkyl lithium (e.g. n-BuLi) provides the lithio derivative, which can then be quenched with the appropriate electrophile (e.g. R¹L, etc.).

An alternative process for introducing substituents at the C-3 position of pyridine Ring I of Formula 1.0, involves nitrating a compound of Formula 415.00 (except wherein X is nitrogen) or a compound of Formula 470.00d with tetrabutylammonium nitrate - trifluoroacetic anhydride in methylene chloride at a temperature of 0°C to room temperature (about 25°C). The nitro group may then be reduced to the corresponding amine using iron filings in ethanol, or powdered zinc - acetic acid in aqueous THF. By methods known to those skilled in the art, the amine group can be converted to a variety of substituents, such as, halo, cyano, thio, hydroxyl, alkyl, alkenyl, alkynyl and haloalkyl.

Wherein Z represents sulfur, a compound of Formula 400.00 wherein Z is oxygen is reacted with P₂S₅, Lawesson's reagent, or another reagent capable of introducing sulfur in place of oxygen. The reaction may take place at elevated temperature in pyridine, toluene or other suitable solvents. In this and other reactions, numerous conversions of a compound of Formula 400.00 (Z = O) to another compound of Formula 400.00 (Z = S) are possible.


25 PREPARATION OF C5-C6-ENE DERIVATIVES

Compounds of formula 400.00 with a double bond between C-5 and C-6 can be prepared by heating a compound of Formula 470.00h in acetic acid with SeO₂ to produce a compound of Formula 470.00i. Compounds of Formula 470.00i can be converted to final products according to methods already described.




PREPARATION OF PIPERAZINE ANALOGS

- Compounds having a piperazine ring bound to the C-11 of the tricyclic nucleus, i.e., Formula 1.0 wherein X is N, are best prepared via alkylation of the appropriately substituted piperazine compound of
- 5 Formula 700.00 with a compound of Formula 705.00. Compounds of Formula 705.00 contain the appropriately substituted halide (such as Cl, Br, or I) or other similar leaving group (e.g., tosyloxy or mesyloxy). The reaction is usually conducted in an inert solvent, such as THF or toluene, optionally with a base such as triethylamine or potassium carbonate, and
- 10 typically at a temperature range of ambient to reflux to produce a compound of Formula 710.00.



- In this reaction R⁹ is H, CO₂R^a (wherein R^a is a C₁ to C₄ alkyl group) or C(Z)R. The preparation of compound 705.00 wherein L is Cl is analogous
- 15 to the procedure described in U.S. 3, 409,621. One skilled in the art can prepare other derivatives of 705.00 (e.g., L is Br, I, mesyloxy, or tosyloxy). When R⁹ is H, C(Z)R or CO₂R^a, these are converted to compounds of the invention by processes known in the art.

- An alternate route for generating the compound of Formula 710.00
- 20 is by reductive amination of the aza ketone 715.00 with the piperazine 700.00



The reaction is typically carried out in a polar solvent, such as methanol or ethanol, optionally in the presence of a dehydrating agent, such as 3Å molecular sieves. The intermediate Schiff base can be reduced to the compound of Formula 710.00 by employing a variety of reducing agents, such as NaCNBH₃, or catalytic hydrogenation, for example, hydrogen over Pd/C.

When R⁹ is C(Z)R, these are the compounds of the invention. When R⁹ is H or CO₂R^a, these are converted to compounds of the invention as described herein.

Compounds of Formulas 5.3A and 5.3B, wherein R²⁵ represents a pyridyl N-oxide, can be produced by reacting compounds of Formulas 5.3A and 5.3B, wherein R²⁵ is pyridyl, with a one molar equivalent of an oxidizing agent (such as oxone).







Compounds of Formulas 5.3, 5.3A and 5.3B, wherein R²⁵ represents a pyridyl N-oxide, can be produced by reacting the product of Preparative Example 12 with a peroxyacid (such as m-chloroperbenzoic acid) to give the corresponding N-oxide intermediate. The desired N-oxide product may be obtained from the N-oxide intermediate by following the procedure of Example 183.

In the above processes, it is sometimes desirable and/or necessary to protect certain R¹, R², R³ and R⁴ etc., groups during the reactions. Conventional protecting groups are operable as described in Greene, T.W., "Protective Groups In Organic Synthesis," John Wiley & Sons, New

York, 1981. For example, the groups listed in column 1 of Table 1 may be protected as indicated in column 2 of the table:

TABLE 1
PROTECTED GROUPS

5

1. GROUP TO BE PROTECTED	2. PROTECTED GROUP
-COOH	-COOalkyl, -COObenzyl, -COOphenyl, 
	 ,  , 
	 , 
-OH	 , -OCH ₂ phenyl, -OCH ₃ , OSi(CH ₃) ₂ (t-Bu),
-NHR, wherein R is any substituent on an amino group within the scope of the claims	 , -NR-CO-CF ₃ , -NR-COCH ₃ , -NRCH ₂ 
-NH ₂	 , -NH-C(O)-O(t-Bu)


Other protecting groups well known in the art also may be used. After the reaction or reactions, the protecting groups may be removed by standard procedures.

Compounds useful in this invention are exemplified by the following preparative examples, which should not be construed to limit the scope of the disclosure. Alternative mechanistic pathways and analogous structures within the scope of the invention may be apparent to those skilled in the art.

10

PREPARATIVE EXAMPLE 1


A. N-(1,1-DIMETHYLETHYL)-3-METHYL-2-PYRIDINE CARBOXAMIDE



Suspend 2-cyano-3-methyl pyridine (400 g) in t-butanol (800 mL) and heat to 70°C. Add concentrated sulphuric acid (400 mL) dropwise over 45 minutes. Maintain the temperature at 75°C, until the reaction is complete, and for an additional 30 minutes. Dilute the mixture with water (400 mL), charge with toluene (600 mL) and bring to pH 10 with concentrated aqueous ammonia. Maintain the temperature at 50-55°C during the work up. Separate the toluene phase, and reextract the aqueous layer. Combine toluene phases and wash with water. Remove the toluene to yield the title compound N-(1,1-dimethylethyl)-3-methyl-2-pyridine carboxamide, as an oil, from which solid product is crystallized. (Yield 97%, as determined by an internal standard assay with gas chromatography).

25

B. 3-[2-(3-CHLOROPHENYL)ETHYL]-N-(1,1-DIMETHYLETHYL)-2-PYRIDINE CARBOXAMIDE




Dissolve the title compound of Preparative Example 1A, N-(1,1-dimethylethyl)-3-methyl-2-pyridine carboxamide (31.5 g.) in

30


tetrahydrofuran (600 mL) and cool the resulting solution to -40°C. Add n-butyllithium (2 eq.) in hexane while maintaining the temperature at -40°C. The solution turns deep purple-red. Add sodium bromide (1.6 g) and stir the mixture. Add solution of m-chlorobenzylchloride (26.5 g., 0.174 mole) in tetrahydrofuran (125 mL) while maintaining the temperature at -40°C. Stir the reaction mixture until the reaction is complete as determined by thin layer chromatography. Add water to the reaction until the color is dissipated. Extract the reaction mixture with ethyl acetate, wash with water, and concentrate to a residue which is the title compound. (Yield 92% as shown by chromatography).

C. 3-[2-(3-CHLOROPHENYL)ETHYL]-2-PYRIDINE-CARBONITRILE




Heat a solution of the title compound of Preparative Example 1B, 3-[2-(3-chlorophenyl)ethyl]-N-(1,1-dimethylethyl)-2-pyridine carboxamide (175 g, 0.554 mole) in phosphorous oxychloride (525 mL, 863 g, 5.63 mole) and reflux for 3 hours. Determine completion of the reaction by thin layer chromatography. Remove any excess phosphorous oxychloride by distillation at reduced pressure and quench the reaction in a mixture of water and isopropanol. Bring to pH 5-7 by adding 50% aqueous sodium hydroxide solution while maintaining the temperature below 30°C. Filter the crystalline slurry of crude product and wash with water. Purify the crude product by slurrying the wet cake in hot isopropanol, and cool to 0-5°C. Filter the product, wash with hexane and dry at a temperature below 50°C to yield the title compound. (Yield: 118g (HPLC purity 95.7%), m.p. 72°C-73°C, 89.4% of theory).

D. 1-(METHYL-4-PIPERIDINYL)[3-(2-(3-CHLORO-PHENYL)ETHYL)-2-PYRIDINYL]METHANONE HYDROCHLORIDE



- Dissolve the title compound of Preparative Example 1C, (118 g, 0.487 mole) in dry tetrahydrofuran (1.2L) and add N-methyl-piperidyl magnesium chloride (395 mL, 2.48 mole/liter, 0.585 mole, 1.2 eq.) over 15 minutes. Maintain the temperature at 40°C-50°C by cooling with water as necessary, for 30 minutes. Determine completion of the reaction by thin layer chromatography. Quench the reaction by reducing the pH to below 2 with 2N HCl and stir the resulting solution at 25°C for 1 hour. Remove the bulk of the tetrahydrofuran by distillation and adjust the resulting solution to pH 3.5 by addition of aqueous sodium hydroxide. Cool to 0 to 5°C and filter off the crystalline hydrochloride salt product. Wash with ice cold water and dry to constant weight at 60°C to yield the title compound. (Yield: 168.2 g (HPLC purity 94%), m.p. 183°-185°C, 89% of theory).

E. 8-CHLORO-11-(1-METHYL-4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE




- Dissolve the title compound of Preparative Example 1D above (59 g, 0.15 mole) in hydrofluoric acid (120 mL, 120 g, 6.0 mole) at -35°C and add boron trifluoride (44.3 g, 0.66 mole) over 1 hour. Determine completeness of the reaction by thin layer chromatography. Quench the

reaction using ice, water and potassium hydroxide bringing the solution to a final pH of 10. Extract the product with toluene and wash with water and brine. Concentrate the toluene solution to a residue, and dissolve in hot hexane. Remove the insolubles by filtration and concentrate the filtrate to yield the title compound as an off-white powder. (Yield: 45.7 g (HPLC purity: 95%), 92% of theory).

Alternative Step E: 8-CHLORO-11-(1-METHYL-4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE

React the title compound of Preparative Example 1D above (177 g, 0.49 mole) in trifluoromethanesulfonic acid (480 ml, 814.1 g, 5.31 mole) at 90-95°C for 18 hours under nitrogen. Determine the completeness of the reaction by thin layer chromatography. Cool the reaction and quench the reaction with ice-water and adjust the pH to 6 with barium carbonate. Extract the product with methylene chloride, and concentrate under reduced pressure to about 1 liter. Wash with water, and extract the product into 1 N HCl which is treated with 30 g of activated charcoal, and filter through celite. Adjust the pH of the filtrate to 10 with aqueous sodium hydroxide (50%), extract the product into methylene chloride, and remove under reduced pressure to form a residue. Dissolve the residue in hot hexane, and filter to remove insolubles. Concentrate the filtrate to yield the title compound as a beige powder. (Yield: 126 g (HPLC purity 80%), 65% of theory).


F. 8-CHLORO-11-(1-ETHOXYCARBONYL-4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE



Dissolve the title compound of Preparative Example 1E above (45.6 g, 0.141 mole) in toluene (320 mL) at 80°C and to it gradually add ethyl chloroformate (40.4 mL, 45.9 g, 0.423 mole). Following complete

- addition, maintain the temperature at 80°C for 1 hour, then add diisopropylethylamine (2.7 mL, 2.00 g, 0.016 mole) and additional ethyl chloroformate (4.1 mL, 4.65 g, 0.0429 mole). Monitor completeness of the reaction by thin layer chromatography. Upon completion, cool the
- 5 reaction mixture to ambient temperature, and wash the toluene solution with water. Concentrate the organic layer to a residue and dissolve in hot acetonitrile (320 mL). Decolorize the solution with 14 g of activated charcoal. Remove the activated charcoal by filtration and concentrate the filtrate to a crystalline slurry. Cool the mixture to 0-5°C, and isolate the
- 10 product by filtration. Wash with cold acetonitrile and dry the product at below 70°C to yield the title compound. (Yield: 42.4 g (HPLC purity 97.4%), 80% of theory).


G. 8-CHLORO-11-(4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE




15

- Hydrolyze the title compound of Preparative Example 1F, 8-chloro-11-(1-ethoxycarbonyl-4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (39 g, 0.101 mole) with KOH (50 g) in ethanol (305 mL) and water (270 mL) at reflux under an argon
- 20 atmosphere for 64 hours. Partially distill off the ethanol and dilute the residue with brine, and extract with ethyl acetate (3x). Wash the combined organic phases with water and dry with Na₂SO₄. Remove the solvent to give a solid which can be recrystallized from toluene to give the title
- 25 compound as a white solid. (Yield: 24.5 g, 77%, melting point 154-155°C).

H. By substituting in step 1B above, the benzylic halide:




for meta-chlorobenzylchloride, and employing basically the same methods as steps C through G, the compound



is prepared. Dichloro compound (I) is recrystallized from toluene and has a melting point of 150-152°C. Reaction times are determined by TLC or HPLC. In some instances purification of the product by chromatography is necessary.


PREPARATIVE EXAMPLE 2

10 A. N-(1,1-DIMETHYLETHYL)-3-[2-(4-FLUOROPHENYL)-ETHYL]-2-PYRIDINE CARBOXAMIDE




Cool a solution of N-(1,1-dimethylethyl)-3-methyl-2-pyridine-carboxamide (38.4 g, 0.2 mole) in dry THF (250 mL) to -40°C and add n-butyl lithium (185 mL, 0.44 mole). Add sodium bromide (1.9 g, 18 mmol.) and stir for 15 minutes. Add 4-fluorobenzylchloride (31.8 g, 0.22 mole) and stir for 2.5 hours while warming to -5°C. Quench the reaction with water and extract the product twice with ethyl acetate, then wash with brine (2X). Dry the organic phase over Na₂SO₄, filter and remove the solvent to give the title compound. (60.0 g; Yield 99%, m.p. 59-61°C.)

20 B. 3-[2-(4-FLUOROPHENYL)ETHYL]-2-PYRIDINE CARBONITRILE




- Heat the title compound of Preparative Example 2A above (60.0 g, 0.2 mole) in POCl_3 (200 mL) to 110°C under an argon atmosphere for 3.5 hours. Pour the reaction mixture onto ice and basify with NaOH (50%) solution. Extract the mixture with ethyl acetate (3x) and wash with water.
- 5 Wash with brine and dry over Na_2SO_4 . Remove the solvent and pass the residue through a coarse SiO_2 (60-200 mesh) column to give the title compound as a white solid (40 g, Yield 88%, m.p. $48-49^\circ\text{C}$).

C. 9-FLUORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA-[1,2-b]PYRIDIN-11-ONE



- Cyclize the title compound of Preparative Example 2B above (31.5 g, 139 mmol) in polyphosphoric acid (1.24 kg) at 200°C for 5.5 hours. Pour onto ice and basify with NaOH solution (50%). Extract the product with chloroform (3x) and wash with brine. Dry the organic phase with Na_2SO_4 , filter and remove the solvent to give the title compound (20.4 g, yield 64%, m.p. $78-81^\circ\text{C}$ after recrystallization from diisopropyl ether).
- 15


D. 9-FLUORO-11-(1-METHYL-4-PIPERIDINYL)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOPHEPTA[1,2-b]PYRIDIN-11-OL



- 20 Dissolve the title compound of Preparative Example 2C above (10.0 g, 44 mmol) in THF (100 mL) and add slowly to a cooled (-40°C) solution of the Grignard reagent prepared from N-methyl-4-chloropiperidine (57.9 mL, 88 mmol) and magnesium in THF (70 mL). Stir the mixture for about 1 hour while warming up to 0°C . Quench the reaction with NH_4Cl solution and extract with ethyl acetate (2X). Wash the organic phase with brine and dry over Na_2SO_4 , filter and remove the solvent.
- 25

Purify the residue with flash chromatography and elute with methanol (5%) in CHCl_3 to give the title compound as white granular crystals. (10.1 g, Yield 70%, m.p. 126-127°C after recrystallization from diisopropyl ether.)


5 E. 9-FLUORO-11-(1-METHYL-4-PIPERIDYLENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE



Add the title compound of Preparative Example 2D above (7.3 g, 22.3 mmol) to a mixture of cooled H_2SO_4 and $\text{CF}_3\text{SO}_3\text{H}$ (1:1), (146 mL).

- 10 Stir the reaction mixture for 0.5 hours at ice bath temperature and then at room temperature for 1.5 hours. Pour the reaction mixture onto ice and basify with NaOH (50%) solution. Extract the product with ethyl acetate (3X) and wash with brine. Dry the organic phase over Na_2SO_4 , filter and remove the solvent to give a crude oil. Charcoal the oil and recrystallize from ethyl acetate and isopropyl ether to give the title compound. (5.6 g, Yield 82%, m.p. 134.5-135.5°C.).

F. 9-FLUORO-11-(1-ETHOXYCARBONYL-4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE




20

Stir a solution of the title compound of Preparative Example 2E above (5.0 g, 16.2 mmol) and triethylamine (2.6 g, 26 mmol) in dry toluene


(60 mL) at 80°C under an argon atmosphere, and add ethyl chloroformate (9.8 g, 90 mmol) via a syringe. Stir the reaction at this temperature for 30 minutes and at room temperature for one hour. Filter the reaction and remove the solvent. Pass the residue through a coarse SiO₂ column (60-
5 200 mesh), and elute with CHCl₃ to yield the title compound as a white solid. (4.5 g, Yield 76%, m.p. 112-114°C after trituration with pentane).

G. 9-FLUORO-11-(4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE




10 Reflux the title compound of Preparative Example 2F above (3.83 g, 10.4 mmol) with KOH (4.6 g) in 50 mL of ethanol/H₂O (1:1) for 4 hours under an argon atmosphere. Pour the reaction mixture into a brine solution and extract with ethyl acetate (2X), dry over Na₂SO₄ and filter. Remove the solvent to give the title compound (2.86 g, Yield 90%, m.p.
15 138-140°C.).

H. By employing the benzyl halide



in place of 4-fluorobenzyl chloride in step 2A above, the product




20 is prepared (m.p. 138-140°C, triturated with pentane) by employing basically the same process as described in steps 2A-2G. Workup time is

determined by either TLC or HPLC. In some instances purification of the product by chromatography is necessary.

PREPARATIVE EXAMPLE 3

5

A. 3,5-DIMETHYLPYRIDINIUM N-OXIDE




10

A solution of 285 mL (1.31 mol) of 35% peracetic acid was slowly added to a stirred solution of 149 g (1.39 mol) of 3,5-dimethylpyridine during which the temperature rose to 85°C and was maintained at this temperature during addition. After the temperature of the mixture dropped to about 35°C the reaction was stored at 5°C overnight.

15

After partial removal of 185 ml of acetic acid via distillation under vacuum, the reaction was washed with NaHSO₄ solution and then neutralized with 10% NaOH solution to pH of about 7. The product was extracted with CH₂Cl₂ to give the title compound as a white solid (yield 142 g, 83%).


B. 1-METHOXY-3,5-DIMETHYLPYRIDINIUM METHYL SULFATE



20

Dimethylsulfate (42.0 g, 0.33 mol) was slowly added to 41.0 g (0.33 mol) of 3,5-dimethylpyridinium N-oxide with mechanical stirring. The mixture was then heated on a steam bath for 1 hr. Then vacuum was applied while cooling to give a brownish solid of the title compound in quantitative yield.


C. 2-CYANO-3,5-DIMETHYLPYRIDINE



25

To a cooled (0°C) solution of sodium cyanide (49.0 g, 0.999 mol, 3.0 eq.) in 135 mL of water (air free) was dripped 1-methoxy-3,5-dimethyl pyridinium methyl sulfate (83.0g, 0.33 mol) in 100 mL water (air free) in 1.25 hr., keeping the temperature below 3°C. The reaction mixture was stored at about 3°C overnight. The mixture was filtered and washed with water to give 40g of the title compound. An analytical sample was recrystallized from isopropyl ether and pentane (4:1) (m.p.: 61-62°C).

D. N-(1,1-DIMETHYLETHYL)-3,5-DIMETHYL-2-PYRIDINE CARBOXAMIDE




10

To a stirred solution of 20.3 g (0.153 mol) of 2-cyano-3,5-dimethylpyridine in 100 mL of 20 mL of conc. sulfuric acid within 10 minutes, followed by 20 mL of t-butanol over an additional 15 minutes. The solution was warmed at 75°C for 30 minutes after which it was cooled to room temperature and basified with 25% NaOH. The product was extracted 3X with EtOAc (600 mL), which was combined and washed 1X with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo* to give the title compound (31.26 g) as a yellowish oil.

E. 8-CHLORO-3-METHYL-11-(4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE

20




By substituting in step 1B above N-(1,1-dimethylethyl)-3,5-dimethyl-2-pyridine carboxamide for N-(1,1-dimethylethyl)-3-methyl-2-pyridine carboxamide and employing basically the same methods as steps B through G of Preparative Example 1, one obtains 8-chloro-3-methyl-11-(4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine. Reaction times are determined by TLC or HPLC.


25

PREPARATIVE EXAMPLE 4

By substituting




for 3,5-dimethylpyridine in Preparative Example 3 above and following
5 basically the same procedure (steps A-E), the compounds



respectively, can be prepared. Note that the addition of the nitrile group
to the pyridine in Step C of Preparative Example 3 can result in the
10 formation of other undesirable isomers which can be removed via flash
chromatography.


PREPARATIVE EXAMPLE 5

A. 8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA-[1,2-b]PYRIDIN-11-ONE N-OXIDE



- 5 To a mixture of 25.1 grams (0.103 mole) of 8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-one in 175 ml of dry methylene chloride at 0°C under an argon atmosphere was added dropwise over 70 minutes a solution of 24.12 grams of 3-chloroperoxybenzoic acid in 150 ml of methylene chloride. After the addition the
- 10 solution was stirred for 1/2 hour after which the ice bath was removed. After two days the reaction was poured into 1.0 N aqueous sodium hydroxide and extracted with methylene chloride. The organic portions were combined, washed once with water, dried over magnesium sulfate, filtered and concentrated in vacuo. The resultant product was triturated
- 15 with isopropyl ether and filtered to provide 25.8 grams (96%) yield of the title compound.

B. 2,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-ONE AND 4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-ONE




20

To a mixture of 29.13 grams (112.2 mmol) of the title compound from Preparative Example 5A above, in 40 ml of dry methylene chloride at 0°C and under argon atmosphere was added 500 ml of 1.0 M SO₂Cl₂ dropwise over 1 hour. The ice bath was then removed and the reaction

stirred at room temperature for 1 hr and then refluxed for seven hours. The mixture was poured into 1.0 N aqueous NaOH and extracted three times with CH₂Cl₂. The organic portions were combined, dried over MgSO₄, filtered and concentrated in vacuo to yield a product which was purified and separated via flash chromatography to yield the two title compounds.


5 C. 4-(2,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)PIPERIDINE AND 4-(4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA-[1,2-b]-
10 PYRIDIN-11-YLIDENE)PIPERIDINE



By following essentially the same procedure as that described in parts D-G of Preparative Example 2 above, the 2,8-dichloro and 4,8-dichloro products of Preparative Example 5B above were converted to the corresponding title compounds.

PREPARATIVE EXAMPLE 6


A. 3-(1,1-DIMETHYL-1-ETHYL)-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-ONE



To a mixture of 20.05 grams (82.28 mmol) of 8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-one in 400 ml of dry

tetrahydrofuran at -72°C and under an atmosphere of nitrogen was added dropwise over 40 minutes 66.0 ml of 2.7 M t-butyl magnesium chloride in tetrahydrofuran. The reaction mixture was slowly warmed to room temperature and stirred overnight. The mixture was then poured into 10% aqueous ammonium chloride and extracted four times with methylene chloride. The combined organic portions were dried over magnesium sulfate, filtered, and concentrated in vacuo to give the title compound, along with 8-chloro-11-(1,1-dimethyl-1-ethyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ol. These compounds were separated via flash chromatography to give the title compound, which was recrystallized from isopropyl ether to give 4.37 grams (18%) of the title compound as a white solid.

B. 4-[3-(1,1-DIMETHYL-1-ETHYL)-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]PIPERIDINE




By using the title compound of Part A above and applying essentially the same procedure described in parts D-G of Preparative Example 2 above, one can obtain the title compound.

20

PREPARATIVE EXAMPLE 7


A. 8-CHLORO-6,11-DIHYDRO-11-HYDROXY-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE



To a mixture of 25.03 g (103 mmol) of 8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-one in 200 mL of methanol at room temperature and under a nitrogen atmosphere was added portionwise over a period of about 1 hour 4.82 g (124 mmol) of sodium borohydride. Occasional cooling with an ice bath was necessary at times during the


addition in order to avoid excessive reflux. After 1.6 hours the mixture was poured into ice cold water and then extracted with ethyl acetate (3X). The combined organic portions were washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. The residue was recrystallized from hot isopropyl ether. The remaining filtrate was purified via flash chromatography (20% ethyl acetate in hexanes) to yield more product which solidified on standing. Both batches were combined to yield 20.41 g of the title compound as a white solid.

10 B. 8,11-DICHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDINE



To a mixture of 13.3 g (54 mmol) of 8-chloro-6,11-dihydro-11-hydroxy-5H-benzo[5,6]cyclohepta[1,2-b]pyridine in 290 mL of toluene at -15°C and under an atmosphere of nitrogen was added via syringe pump over a period of 1 hour 6.20 mL (85.7 mmol) of thionyl chloride. The extent of reaction was monitored by TLC (50% ethyl acetate in hexanes). When completed the mixture was poured into 300 mL of 1.0 N aqueous sodium hydroxide and extracted with ethyl acetate (5X). The combined organic portions were washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. The residue was taken up in ethyl acetate, quickly filtered through basic alumina, and concentrated again to yield a product which was triturated with pentane to yield 10.22 g of the title compound as a tan solid.

25 C. 8-CHLORO-11-(1-PIPERAZINYL)-6,11-DIHYDRO-5H-
BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE




To a mixture of 10.0 g (37.9 mmol) of 8,11-dichloro-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine and 1.0 mL of triethylamine in 200

mL of dry tetrahydrofuran at room temperature and under a nitrogen atmosphere was added 33.0 g of piperazine. The mixture was stirred at room temperature for 22.5 hours and then refluxed for 5.5 hours. It was then cooled to room temperature, poured into 250 mL of 5% aqueous sodium hydroxide, and extracted with methylene chloride (3X). The combined organic portions were washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo. The residue was purified via flash chromatography (2→5% methanol saturated with ammonia in methylene chloride) to yield the title compound as a glass.

10

PREPARATIVE EXAMPLE 8


A. ETHYL 3-PYRIDYLACETIC ACID 1-N-OXIDE



Ethyl 3-pyridylacetic acid (10grams) (60.6 mmoles) was dissolved in dry dichloromethane (120ml) and the solution was stirred at -18°C for 30 minutes. 3-Chloroperbenzoic acid (31.34 grams) (181.6 mmoles) was added and the mixture was stirred at -18°C for 1 hour and then at 25°C for 87 hours. The reaction mixture was diluted with dichloromethane and washed with saturated aqueous sodium bicarbonate and then water. The dichloromethane was then dried (magnesium sulphate), filtered and evaporated to dryness. The residue was chromatographed on silica gel using 3% (10% concentrated ammonium hydroxide in methanol)-dichloromethane as the eluant to give the title compound (Yield: 8.45 grams, 77%, MH⁺ 182).

25


B. 3-PYRIDYLACETIC ACID 1-N-OXIDE



3-Pyridylacetic acid (0.2747 grams) (1.5 mmoles) was dissolved in ethanol (200 proof) (1.22 ml.) and a 1M solution of lithium hydroxide in water (3.64 ml.) (3.0 mmoles) was added and the mixture was stirred at 25°C for 4 hours. 1N Hydrochloric acid (4.28 ml.) was added and the mixture was pumped down to dryness on a rotary evaporator to give the title compound (Yield: 0.2931 grams, 100%).

PREPARATIVE EXAMPLE 9

A. ETHYL α -METHYL-3-PYRIDYLACETIC ACID.




10

To ethyl 3-pyridylacetic acid (10.86 grams) (65.7 mmoles) was added a 2.0M solution of lithium diisopropylamide in THF / heptane / ethyl benzene (32.87 ml.) (65.8 mmoles) at -30°C. The semi-solid mixture was agitated and sonicated for 1 hour. The mixture was allowed to remain at 25°C for 1 hour, whereupon methyl iodide (4.09 ml.) (65.7 mmoles) was added. After 1 hour at 25°C the mixture was taken up in dichloromethane and washed with saturated aqueous sodium bicarbonate and water. The dichloromethane was dried (magnesium sulphate), filtered and evaporated to dryness. The residue was chromatographed on silica gel using 10% ethyl acetate in hexane as the eluant to give the title compound (Yield: 3.48 grams, 30%, MH⁺ 180).

15

20

B. α -METHYL-3-PYRIDYLACETIC ACID.




The title compound from Preparative Example 9A above (2.16 grams) (12.05 mmoles) was dissolved in ethanol (10 ml.) and 1.0M lithium hydroxide in water (29.15 ml.) (29.2 mmoles) was added. The mixture was stirred at 25°C for 4 hours, whereupon 1N hydrochloric acid (34.27

25

ml.) (34.2 mmoles) was added and the solution was evaporated to dryness to give the title compound (Yield 2.33 grams, 100%).

PREPARATIVE EXAMPLE 10

5 α,α -DIMETHYL-3-PYRIDYLACETIC ACID.




Ethyl α,α -dimethyl -3-pyridylacetate (disclosed in EP Application 0 288 279, published October 26, 1988) (2.67 grams, 13.8 mmoles) was dissolved in ethanol (11.1 ml.) and a 1.0M lithium hydroxide in water (33.3 ml.) (33.4 mmoles) was added. The mixture was stirred at 25°C for 4 hours. 1N Hydrochloric acid (38.73 ml.) was added and after 5 minutes the mixture was evaporated to dryness to give the title compound (Yield: 100%).

15


PREPARATIVE EXAMPLE 11

A. 8-CHLORO-6,11-DIHYDRO-11-(1-PIPERAZINYL)-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE 1-N-OXIDE




To a mixture of 8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta-
20 [1,2-b]pyridin-11-one (5 grams) (20.6 mmoles) in dry dichloromethane (35 ml) was added dropwise 3-chloroperbenzoic acid (4.7 grams) (27.3 mmoles) in dry dichloromethane (75 ml) at 0-25°C over 1 hour. The mixture was diluted with dichloromethane and washed with saturated aqueous sodium bicarbonate and water. The dichloromethane was dried
25 (magnesium sulphate), filtered and evaporated to dryness. The residue was chromatographed on silica gel using 1% (10% saturated ammonium hydroxide in methanol)-dichloromethane as the eluant to give the title compound (Yield: 2.81 grams, 53%, MH⁺ 260).

B. 8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA-[1,2-b]PYRIDIN-11-OL 1-N-OXIDE



By using the title compound (8.6 grams) from Preparative Example 11A and reducing it by the procedure described in Preparative Example 7A above the title alcohol was obtained (Yield: 7.03 grams, 81%, MH⁺ 262).

C. 8,11-DICHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE 1-N-OXIDE




10

The title compound from Preparative Example 11B (6.2 grams) (23.7 mmoles) was reacted with thionyl chloride as described in Preparative Example 7B to give the title compound.

D. 8-CHLORO-6,11-DIHYDRO-11-(1-PIPERAZINYL)-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE 1-N-OXIDE


15



The title compound from Preparative Example 11C above was reacted with piperazine (9.9 grams) (115.0 mmoles) as described in Preparative Example 7C to give the title compound (Yield: 6.78 grams, 87%, MH⁺ 330).


20

PREPARATIVE EXAMPLE 12
4-ETHOXYCARBONYLAMINOPYRIDINE



- 4-Aminopyridine (17.34 grams) (184.3) was dissolved in dry
5 pyridine (217 ml.) and cooled to 0°C over 30 minutes. Ethyl chloroformate
(17.2 ml.) (180.7 mmoles) was added and the solution was stirred at 0°C
for 1 hour and then at 25°C for 40 hours. The mixture was diluted with
dichloromethane and washed with saturated aqueous sodium
bicarbonate and water. The dichloromethane was dried (magnesium
10 sulphate), filtered and evaporated to dryness. The residue was
chromatographed on silica gel using 2%(10% saturated ammonium
hydroxide in methanol)-dichloromethane to give the title compound (Yield:
10 grams, 33%, M⁺ 166).


By using essentially the same procedure, with the exception that



was obtained, respectively.


PREPARATIVE EXAMPLE 13

A. N-ACETYLISONIPECOTIC ACID



Isonipecotic acid (10 grams) (77.5 mmoles) and acetic anhydride (23.7 grams) (232.5 mmoles) were dissolved in methanol (100 ml.) and the mixture was stirred at 25°C for 24 hours. The mixture was evaporated to dryness and the residue was azeotroped with toluene to give the title compound (Yield: 12.8 grams, 97%, MH^+ 172).

B. 1-N-tert-BUTOXYCARBONYLISONIPECOTIC ACID




10

Isonipecotic acid (20 grams) (155.0 mmoles) was dissolved in THF-water (1:1) (400 ml) and sodium hydroxide (6.2 grams) (155.0 mmoles) and di-tert-butylidicarbonate (37.2 grams) (170.5 mmoles) were added. The mixture was stirred at 25°C for 72 hours. The solution was then eluted through a bed of washed BioRad 50WX4 (RSO3H resin) (150 ml bed) and the resin was eluted with a 1:1 mixture of THF and water. The eluate was evaporated to dryness to give the title compound (Yield: 33.78 grams, 90%).

20

PREPARATIVE EXAMPLE 14

1-N-ACETYLNIEPECOTIC ACID




Nipecotic acid (3.87 grams) (30.0 mmoles) was reacted with acetic anhydride (9.17 grams) (90 mmoles) as described in Preparative Example 13A to give the title compound (Yield: 5.0 grams, 97%, MH^+ 172).

5

PREPARATIVE EXAMPLE 15

1-N-METHYLNIPECOTIC ACID




Arecaidine hydrochloride (4 grams) (22.6 mmoles) was hydrogenated in water (100 ml) using 10% Pd-C at 40 psi at 25°C for 24 hours. The catalyst was filtered off and washed with water. The aqueous solution was shaken with BioRad AG1X8 resin (OH^- form) (23 ml bed) and after 5 minutes the resin was filtered off and washed with water. The aqueous solution was evaporated to give the title compound (Yield: 2.95 grams, 92%).

15

PREPARATIVE EXAMPLE 16

1-N-ACETYL D,L-PIPECOLINIC ACID




D,L-Pipecolic acid (10 grams) (77.5 mmoles) and acetic anhydride (23.7 grams) (232.5 mmoles) were reacted as described in Preparative Example 13A above to give the title compound (Yield: 12.94 grams, 98%, MH^+ 172).

20


PREPARATIVE EXAMPLE 17

A. PIPERIDINE-4-ACETIC ACID




- 5 4-Pyridylacetic acid (7 grams) (40.4 mmoles) was hydrogenated as described in Preparative Example 15 to give the title compound (Yield: 5.2 grams, 90%, MH⁺ 144).

B. 1-N-ACETYL-4-PIPERIDINYLACETIC ACID



- 10 4-Piperidylacetic acid (5 grams) (35.0 mmoles) was reacted with acetic anhydride (10.7 grams) (105.0 mmoles) as described in Preparative Example 13A to give the title compound (Yield: 6.4 grams, 99%, MH⁺ 185).


C. 1-N-METHYL-4-PIPERIDINYLACETIC ACID



- 15 4-Piperidylacetic acid (4 grams) (28.0 mmoles) from Preparative Example 17A was dissolved in water (50 ml) and 37% formalin (2.72 ml) (33.6 mmoles) was added. The mixture was hydrogenated over 10% Pd-C at 55psi at 25°C for 68 hours. The catalyst was filtered off and washed

with water. The combined filtrates were evaporated to dryness to give the title compound (MH^+158).

D. 1-N-tert-BUTOXYCARBONYLPIPERIDINYL-4-ACETIC ACID




- 5 4-Piperidylacetic acid (41.24 grams) (288.4 mmoles) from Preparative Example 17A was reacted with di-tert-butylidicarbonate (69.14 grams) (317.3 mmoles) and sodium hydroxide (11.52 grams) (288.4 mmoles) as described in Preparative Example 13B above to give the title compound (Yield: 53.0 grams, 76%).

10


PREPARATIVE EXAMPLE 18

A. 3-PIPERIDINYLACETIC ACID



- 15 3-Pyridylacetic acid hydrochloride (13 grams) (74.9 mmoles) was hydrogenated as described in Preparative Example 15 to give a mixture of unreacted 3-pyridylacetic acid and the title compound (76:24) (8.63 grams, $MH^+ 144$).


B. 1-N-ACETYL-3-PIPERIDINYLACETIC ACID



- 20 The mixture of compounds from Preparative Example 18A (8.56 grams) were reacted with acetic anhydride (8.56 grams) as described in

Preparative Example 13A and the crude mixture of products was taken up in methanol (60 ml) and passed over a bed of BioRad AG50WX4 resin (RSO₃H) and the latter was eluted with methanol. The eluates were evaporated to dryness to give the title compound (Yield: 1.23 grams, MH⁺ 186).

C. 1-N-METHYL-3-PIPERIDINYLACETIC ACID




The mixture of compounds from Preparative Example 18A (4 grams) and 37% formalin (2.72 ml.) were hydrogenated as described in Preparative Example 17C to give the title compound (MH⁺ 158).

PREPARATIVE EXAMPLE 19


PREPARATION OF THE R(+) AND S(-) DIASTEREOISOMERS

The racemic 8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo-[5,6]cyclohepta[1,2-b]pyridine prepared in Preparative Example 7C above was resolved by the method described in Preparative Example 15 A-C, pages 116-118, of WO 92/00293, published January 9, 1992, to give the R(+) and S(-) diastereoisomers:




PREPARATIVE EXAMPLE 20

A. 3-BROMO-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11-ONE




5 Cyclize 3-[2-(3-chlorophenyl)ethyl]-4-bromo-2-pyridine carbonitrile (10.7g, 32.8 mmol) in triflic acid (82 mL) at 60°C for 2 hours and then at room temperature for 2 hours. Add 80 mL of 5N HCl carefully, then reflux in an oil bath (120°C) for 30 minutes. Cool the solution and pour into ice and basify with 25% NaOH solution. Extract the product with CH₂Cl₂ and
10 wash with brine. Dry the organic layer with Na₂SO₄, filter and remove the solvent to give crude product (10.4g). Purify the crude product with flash chromatography on silica gel and elute with 15% ethyl acetate-hexane to give the title compound as a white solid (9g, 27.95 mmol, Yield 85.2% MH⁺ 322).

15 B. 8-CHLORO-3-METHOXY-5,6-DIHYDRO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11-ONE



Dissolve the title compound of Preparative Example 20A (2.37g, 7.4 mmol) in dry methanol and add sodium metal (3.37g, 180 mmol). the
20 reaction is stirred overnight at room temperature. Reflux the reaction for 3 hours, cool to room temperature and extract with dichloromethane-water. Dry the CH₂Cl₂ fraction and chromatograph on silica gel eluting with 50% EtOAc-hexanes to give the title compound as a light yellow solid (1.5g, Yield 72% MH⁺ 274).


C. 8-CHLORO-3-METHOXY- 11-(-4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDINE



By substituting in Preparative Example 2 step D, 8-chloro-3-methoxy-5,6-dihydro-11H-benzo[5,6]-cyclohepta[1,2-b]pyridin-11-one for 9-fluoro-5,6-dihydro-11H-benzo[5,6]-cyclohepta[1,2-b]pyridin-11-one and employing basically the same methods as steps D through H of Preparative Example 2, one obtains 8-chloro-3-methoxy-11-(4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]-cyclohepta[1,2-b]pyridine as a white solid (MH^+ 340).

PREPARATIVE EXAMPLE 25


A. ETHYL α -METHYL-4-PYRIDYL ACETIC ACID



To dry THF at -78°C was added diisopropylamine (5.05g 48 mmol, 7mL) and then n-butyl lithium. The reaction mixture was stirred for 0.5 h and then ethyl 4-pyridyl acetic acid (7.85g, 46 mmol) was added, and after stirring for 0.5 h at that -78°C the reaction temperature was raised to room temperature. DMF (20 mL) was added and the reaction mixture cooled to -78°C again. Methyl iodide (7.07g, 50.2 mmol, 3.15 mL) was added and the reaction mixture stirred at that temperature for 1h and then at room temperature overnight. All the volatiles were then stripped off and the reaction mixture was partitioned between water- CH_2Cl_2 . The aqueous phase was washed twice with CH_2Cl_2 . The combined CH_2Cl_2 phases were dried and evaporated. The crude product was chromatographed on

silica gel eluting with 80% ethyl acetate hexane to give the title compound (7.88g, MH^+ 179).


B. α -METHYL-4-PYRIDYL ACETIC ACID



- 5 The compound from Preparative Example 25A was hydrolysed in a similar manner to Preparative Example 9B to give the title compound (MH^+ 152).

PREPARATIVE EXAMPLE 26


10 A. α,α -DIMETHYL-4-PYRIDYL ACETIC ACID



- 15 By essentially the same procedure as set forth in Preparative Example 10A-B, but using ethyl α -methyl-4-pyridylacetic acid (from Preparative Example 25) instead of ethyl pyridyl acetic acid the title compound was obtained as an oil (MH^+ 166).

PREPARATIVE EXAMPLE 27


ETHYL 4-[4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-PIPERIDINECARBOXYLATE and
ETHYL 4-[2,8-DICHLORO-5,6-DIHYDRO-11H-
5 BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-PIPERIDINE-
CARBOXYLATE



To phosphorous oxychloride (256 mL) stirring at reflux was added dropwise a solution of the title compound (109 grams) from Example 231A dissolved in chloroform (850 mL). After stirring the resulting solution for an additional 20 minutes at reflux, the reaction mixture was cooled to room temperature and the chloroform removed *in vacuo*. The resulting solution was cooled in an ice-water bath and to it was slowly added 1N aqueous sodium hydroxide (850 mL) followed by 50% aqueous sodium hydroxide until the resulting mixture was slightly basic. Extraction with ethyl acetate, drying of the organic phase over anhydrous magnesium sulfate, concentration *in vacuo*, and purification by flash column chromatography provided the 4,8-dichloro product (27 grams, 23% yield, mp 141.6-145.6 °C) and the 2,8-dichloro product (9 grams, 8% yield, 176.5-177.9 °C).

PREPARATIVE EXAMPLE 28


4,8-DICHLORO-11-(4-PIPERIDYLIDENE)-6,11-5H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDINE



- 5 A solution of the 4,8-dichloro compound from Preparative Example 27 (2.6 grams) dissolved in absolute ethanol (50 mL) and concentrated hydrochloric acid (100 mL) was stirred at reflux for 48 hours. The reaction mixture was cooled in an ice-water bath and was made basic by addition of solid potassium hydroxide. Concentration *in vacuo* afforded a solid
- 10 which was diluted with dichloromethane and water. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo* to provide the title compound (2.0 grams, 93% yield, mp = 181.1-183.2°C).

PREPARATIVE EXAMPLE 29

- 15 ETHYL 4-[4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-PIPERIDINE CARBOXYLATE, N-OXIDE




- To a cooled (0°C) solution of the 4,8-dichloro compound from
- 20 Preparative Example 27 (9.5 grams) dissolved in dichloromethane (300

mL) under N₂ was added dropwise a solution of *meta*-chloroperoxybenzoic acid (6.8 grams) dissolved in ethyl acetate (200 mL). The resulting mixture was washed with 1*N* aqueous sodium hydroxide, dried over anhydrous magnesium sulfate and concentrated *in vacuo*. The residue was purified by flash column chromatography (silica gel) using 100% ethyl acetate then 10% methanol-dichloromethane to afford the title compound (4.9 grams, 50%, MH⁺ 433).

PREPARATIVE EXAMPLE 30

10 ETHYL 4-[4-(2-AMINOETHYLTHIO)-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-PIPERIDINE CARBOXYLATE




15 A mixture of the title compound from Preparative Example 29 (0.53 grams), 2-aminoethanethiol hydrochloride (0.74 grams) and absolute ethanol (15 mL) was stirred at reflux for 48 hours. The mixture was cooled to 25°C, diluted with dichloromethane and washed with 1*N* aqueous sodium hydroxide. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo* to provide the title compound (0.5 grams, 88%, MH⁺ 458).

20

PREPARATIVE EXAMPLE 31

1,1-DIMETHYLETHYL [2-[[8-CHLORO-6,11-DIHYDRO-11-(1-ETHOXYCARBONYL)-4-PIPERIDINYLIDENE]-5H-BENZO[5,6]CYCLO-HEPTA[1,2-b]PYRIDIN-4-YL]THIO]ETHYL]CARBAMATE




5

To the title compound from Preparative Example 30 (0.33 grams) dissolved in dichloromethane (60 mL) was added di-*tert*-butyldicarbonate (0.17 grams). The solution was stirred at 25°C under N₂ overnight. An additional 0.1 grams of di-*tert*-butyldicarbonate was added and after 4
10 hours the reaction mixture was diluted with dichloromethane, washed with 1*N* aqueous sodium hydroxide and concentrated *in vacuo* to afford the title compound (0.5 grams, 100%, MH⁺ 558).

PREPARATIVE EXAMPLE 32


1,1-DIMETHYLETHYL [2-[[8-CHLORO-6,11-DIHYDRO-11-[4-PIPERIDINYLIDENE]-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-4-YL]THIO]ETHYL]CARBAMATE



5 To the title compound from Preparative Example 31 (0.22 grams) dissolved in absolute ethanol (5 mL) was added water (5 mL) and solid potassium hydroxide (0.33 grams). The solution was stirred at reflux for 4 days, then cooled to 25°C, diluted with dichloromethane and washed with
10 water. The organic phase was concentrated *in vacuo* and the residue purified by flash column chromatography (silica gel) using 5% methanol-dichloromethane saturated with ammonium hydroxide to afford the title compound (0.04 grams, 19%, MH⁺ 486).

PREPARATIVE EXAMPLE 33

3-PYRIDYLISOCYANATE, HYDROCHLORIDE



20 A 1.93 solution of phosgene in toluene (20%) (584 mL) was diluted with dry dichloromethane (1 L) and the mixture was stirred at 0°C under nitrogen atmosphere. A solution of 3-aminopyridine (21.1 grams) and dry pyridine (19 mL) dissolved in dry dichloromethane (600 mL) was added dropwise to the stirred solution at 0°C over a period of 5.5 hours. The mixture was stirred at 0-25°C for an additional 48 hours. A stream of nitrogen was passed through the solution to remove most of the phosgene


and the solution was then evaporated until almost all of the solvent was removed to give the title compound which was then taken up in dry pyridine (850 mL) to give a stock solution of the title compound.

5

PREPARATIVE EXAMPLE 34

A. 8-CHLORO-11-(1-ETHOXYCARBONYL-4-PIPERIDINYL)-
11H-BENZO[5,6]CYCLOHEPTA(1,2-b)PYRIDINE

B. 8-CHLORO-11-(1-ETHOXYCARBONYL-4-PIPERIDINYL)-9-
ETHYL-11H-BENZO[5,6]CYCLOHEPTA(1,2-b)PYRIDINE




10

The title compound of Preparative Example 1F above (51.15 grams, 0.1336 mole) was dissolved in trifluoromethanesulfonic acid (170 mL). The dark mixture was heated to reflux for 70h. The solution was cooled to room temperature and was then poured into 800 mL of an ice/water slurry and the resulting mixture stirred. Concentrated ammonium hydroxide solution (175 mL) was added to the mixture in small portions so that the temperature of the mixture was below 20°C. The resulting basic mixture was extracted with dichloromethane. The dichloromethane extract was washed with brine and was then evaporated to give a brown residue. This residue was dissolved in dichloromethane

20

(750 mL) and the solution cooled to 0°C. Ethyl chloroformate (14.8 grams, 0.136 mole) was added over 5 minutes and the resulting mixture stirred at 0°C for 15 minutes. Saturated sodium bicarbonate solution (150 mL) was added and the cooling bath was removed. The resulting biphasic mixture was stirred rapidly for 3h. The layers were separated and the dichloromethane layer was filtered through silica gel. The filtrate was evaporated to dryness and the residue chromatographed on silica gel using a gradient of hexane-dichloromethane-acetone 16:2.5:1.5 to hexane-dichloromethane-acetone 28:7.5:4.5 as eluent to give title compound A (25.02g 49% MH⁺ 383) and title compound B (4.85g, 9%, MH⁺ 411).


C. 8-CHLORO-11-(4-PIPERIDINYL)-11H-BENZO[5.6]CYCLO-HEPTA(1.2-b)PYRIDINE



Hydrolyze the title compound of Preparative Example 34A by dissolving in 50% aqueous sulfuric acid (v/v) and heating to 90° to 100°C for 16 h. The cooled acidic mixture was neutralized with 25% sodium hydroxide solution (w/v). The resulting mixture was extracted with ethyl acetate and the ethyl acetate extract was dried with sodium sulfate. Filtration and evaporation of the ethyl acetate afforded the title compound (MH⁺ 311).

PREPARATIVE EXAMPLE 35

8-CHLORO-9-ETHYL-11-(4-PIPERIDINYL)-11H-BENZO-[5.6]CYCLOHEPTA[1.2-b]PYRIDINE




- 5 Hydrolyze the title compound of Preparative Example 34B following the procedure described in Preparative Example 34C to provide the title compound. 9Decomposes between 205.7-215.4°C, heating 2-3°C per minute.

10


PREPARATIVE EXAMPLE 36

A. 8-CHLORO-11-(1-ETHOXYCARBONYL-4-PIPERIDINYL)-11H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDINE-1-OXIDE



- 15 The title compound from Preparative Example 34A above (20.23 grams, 52.84 mmoles) was dissolved in dichloromethane (250 mL). 3-Chloroperoxybenzoic acid (1.25 equivalents) was added in one portion and this solution was stirred for 45 minutes. Sodium bisulfite solution (20% w/v) was added and the biphasic mixture rapidly stirred for 30 minutes. The layers were separated and the organic layer was washed with saturated sodium carbonate solution and dried with sodium sulfate.
- 20 Filtration and evaporation afforded the title compound (21g, 99%, mp 78.6-89.4°C, MH+ 399).


B. 4,8-DICHLORO-11-(1-ETHOXYCARBONYL-4-PIPERIDIN-
YL)-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE (A) and
2,8-DICHLORO-11-(1-ETHOXYCARBONYL-4-PIPERIDIN-
YL)-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE (B)



The title compound from Preparative Example 36A (21 grams, 53 mmoles) above was dissolved in anhydrous dichloroethane (250 mL) and the solution cooled to 0°C. POCl₃ (49.4 grams, 0.322 mole) was added dropwise to the dichloroethane solution over 15 minutes. After the POCl₃ was added the reaction mixture was warmed to 45 - 50°C and stirred for 18h. Additional POCl₃ (8.2 grams) was added and the mixture heated to reflux for 9h. The mixture was cooled and added to an ice cooled, stirred solution of sodium hydroxide (15% w/v). The resulting biphasic mixture was stirred rapidly for 18h. The layers were separated and the aqueous layer was extracted with dichloromethane. The combined organic layers were washed with water followed by brine and dried (sodium sulfate). The mixture was filtered and evaporated, and the residue chromatographed on silica gel using a gradient of 25% ethyl acetate in hexane to 45% ethyl acetate in hexane as eluent. The title compound A

was obtained as a yellow solid (5.98 g M^+ 417), and title compound B was obtained as a yellow solid (1.0 g, mp 84.4-90.6°C).

C. 4,8-DICHLORO-11-(4-PIPERIDINYL)-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDINE




The title compound A from Preparative Example 36B was hydrolyzed under the conditions described in Preparative Example 34C to give the title compound (M^+ 345).

10

PREPARATIVE EXAMPLE 37


A. 4-(8-CHLORO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-(ETHOXYCARBONYL)-PIPERIDINE



15 The preparation of the starting material for this reaction was described in *The Journal of Organic Chemistry*, 1990, 55, pp. 3341-3350 by Piwinski, J.J.; Wong, J.K.; Chan, T.-M.; Green, M.J.; and Ganguly, A.K. By substituting in Preparative Example 2, 8-chloro-11H-benzo[5,6]-cyclohepta[1,2-b]pyridin-11-one for 9-fluoro-5,6-dihydro-11H-benzo[5,6]-cyclohepta[1,2-b]pyridin-11-one and employing basically the same methods as steps D through F of Preparative Example 2, one obtains the title compound (mp 154.7 - 155.5°C).

20


B. 8-CHLORO-11-(4-PIPERIDINYL)-BENZO[5.6]CYCLO-
HEPTA[1.2-b]PYRIDINE



Hydrolyze the title compound of Preparative Example 37A following
the procedure described in Preparative Example 334C (mp 168.5 -
171.2°C, decomposition).

PREPARATIVE EXAMPLE 38

10 8-CHLORO-11-(1-PIPERAZINYL)-11H-BENZO[5.6]CYCLO-
HEPTA[1.2b]PYRIDINE




The preparation of the starting material for this reaction was
described in *The Journal of Organic Chemistry*, 1990, 55 , pp. 3341-3350
by Piwinski, J.J.; Wong, J.K.; Chan, T.-M.; Green, M.J.; and Ganguly, A.K.
By substituting in Preparative Example 7A, 8-chloro-11H-benzo[5,6]cyclo-
hepta[1,2-b]pyridin-11-one (11.53g) (47.71mmoles) for 8-chloro-5,6-
dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-one and employing

basically the same methods as steps A through C of Preparative Example 7, one obtains 11.53g (36%) of the title compound (MH^+ 312).


PREPARATIVE EXAMPLE 39

5 A. **ETHYL α,α -DIMETHYL-3-PYRIDYLACETIC ACID N-OXIDE**



By substituting in Preparative Example 8A, ethyl α,α -dimethyl-3-pyridylacetic acid (4.0g, 20.7mmoles) for ethyl 3-pyridylacetic acid and using the same method as described in Preparative Example 8A, one
10 obtains the title compound (3.2g, 74%, MH^+ 210).


B. **α,α -DIMETHYL-3-PYRIDYLACETIC ACID N-OXIDE**



By substituting in Preparative Example 8B, ethyl α,α -dimethyl-3-pyridylacetic acid N-oxide (0.142g, 0.68mmoles) (Preparative Example
15 39A) for ethyl 3-pyridylacetic acid N-oxide and using the same method as described in Preparative Example 8B, one obtains the title compound.

PREPARATIVE EXAMPLE 40


4-BROMO-8-CHLORO-11-(1-PIPERAZINYL)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE



- 5 By substituting in Preparative Example 7A, 4-bromo-8-chloro-11-(1-piperazinyl)-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-one (1.5g, 4.65mmoles) (Preparative Example 20A) for 8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-one and using the same methods as described in steps A through C of Preparative Example 7, one
- 10 obtains the title compound (1.31g, 72%, MH^+ 392).

PREPARATIVE EXAMPLE 41


4,8-DICHLORO-11-(1-PIPERAZINYL)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE



- 5 By substituting in Preparative Example 7A 4,8-Dichloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-one (6.64g, 28.37mmoles) (Preparative Example 5B) for 8-chloro-5,6-dihydro-11H-benzo[5,6] cyclohepta [1,2-b]pyridin-11-one and using the same methods as described in steps A through C of Preparative Example 7, one obtains
- 10 the title compound (2.59g, 26%, MH^+ 348).

PREPARATIVE EXAMPLE 42


ETHYL 4-[4-[(1H-BENZOTRIAZOL-1-YL)OXY]-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-PIPERIDINE CARBOXYLATE



To a solution of the 4,8-dichloro compound from Preparative Example 27 (1.5 grams) in dry dimethylformamide (20 mL) was added 1-hydroxybenzotriazole (1.5 grams). After stirring for 14 days at 25°C, sodium hydride (0.84 grams, 60% in mineral oil) was added and after an additional 24 hours, the mixture was poured into water. Filtration provided the title compound (Yield: 1.7 grams, 89%, mp = 181.5 - 183.9 °C, MH⁺ 516).

PREPARATIVE EXAMPLE 43

ETHYL 4-[4-HYDROXY-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-PIPERIDINE CARBOXYLATE




5

To a solution of the title compound from Preparative Example 42 (0.15 grams) and glacial acetic acid (5 mL) was added zinc dust (0.2 grams). After stirring at 25°C for 1 hour, the mixture was filtered through celite and the filtrate concentrated *in vacuo*. The residue was diluted with ethyl acetate, washed with saturated aqueous sodium bicarbonate and brine. The organic layer was separated, dried over magnesium sulfate and concentrated *in vacuo* to give the title compound (Yield: 0.11 grams, 95%, MH+ 399).

10

PREPARATIVE EXAMPLE 44


ETHYL 4-[3-BROMO-4-HYDROXY-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-PIPERIDINE CARBOXYLATE



To a solution of the title compound from Preparative Example 43 (1.3 grams) and glacial acetic acid (5 mL) was added a 0.7 M bromine-acetic acid solution (4 mL) at 25°C under N₂. The solution was poured into 200 mL of water and the resulting solid was filtered, then washed with water. The solid was dried under vacuum overnight to provide the title compound (Yield: 1.2 grams, 81%, MH⁺ 477).

PREPARATIVE EXAMPLE 45

15 ETHYL 4-[3-BROMO-4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-PIPERIDINE CARBOXYLATE




A mixture of the title compound from Preparative Example 44 (5.1 grams), phosphorous oxychloride (20 mL) and chloroform (40 mL) was stirred at reflux over night. The reaction mixture was made basic by the

slow addition of 1N aqueous sodium hydroxide, and the resultant mixture was diluted with dichloromethane. The mixture was shaken well and after separation of the phases, the organic phase was washed with 1 N aqueous sodium hydroxide. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo* to provide a solid which was mixed with methanol and filtered. Concentration of the filtrate provided the title compound as a solid (Yield: 5.7 grams, MH⁺ 497).

10

PREPARATIVE EXAMPLE 46


3-BROMO-4,8-DICHLORO-11-(4-PIPERIDYLIDENE)-6,11-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE



A solution of the title compound from Preparative Example 45 (5.7 grams) dissolved in absolute ethanol (100 mL) and concentrated hydrochloric acid (200 mL) was stirred at reflux for 24 hours. The reaction mixture was cooled in an ice-water bath and was made basic by the addition of solid potassium hydroxide. Extraction with dichloromethane and concentration of the organic phase *in vacuo* afforded the title compound as a solid (1.7 grams, 35% yield, MH⁺ 425).

PREPARATIVE EXAMPLE 47

A. 4-(8-CHLORO-3-NITRO-5,6-DIHYDRO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-PIPERIDINE-1-CARBOXYLIC ACID ETHYL ESTER




5

Tetrabutyl ammonium nitrate(4.98g, 16.3 mmol) was dissolved in dichloromethane(20 mL) and trifluoroacetic anhydride(3.12g,14.9 mmol, 2.1 mL) was then added. The solution was cooled to 0°C and then added (by cannulation) to a solution of 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]-cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidine-1-carboxylic acid ethyl ester (5.69g, 14.9 mmol) in methylene chloride (35 mL) also cooled to 0°C. The reaction mixture was stirred at 0°C for 3h and then allowed to go to room temperature (25°C) overnight. The reaction mixture was then extracted with saturated sodium bicarbonate (60 mL) dried over magnesium sulfate and concentrated to give a semi-solid material that was chromatographed on silica gel eluting first with 10% and then 20% ethyl acetate -hexane. Removal of the organic solvents gave the title compound in 44% yield as a light yellow solid. MP = 90.4-91.0°C, MH⁺ 428.

10


15

B. 4-(8-CHLORO-3-AMINO-5,6-DIHYDRO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-PIPERIDINE-1-CARBOXYLIC ACID ETHYL ESTER



5 The title compound from Preparative Example 47A (5.99g, 14 mmol) was dissolved in 85% aqueous ethanol. To this solution was added iron filings (7.01g, 125.57 mmol) and calcium chloride (0.69g, 6.29 mmol) and the reaction mixture was refluxed for 16h. The reaction mixture was filtered through a bed of celite while hot and the celite was washed with
10 hot ethanol (700 mL). The ethanol solution was then decolorized with activated charcoal (2.4g) and then filtered through celite. Ethanol was then rotary evaporated to give the title compound in 100% yield as an off-white solid. MP= 102.4-103.1°C, MH + 398.

15 C. 4-(8-CHLORO-3-BROMO-5,6-DIHYDRO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-PIPERIDINE-1-CARBOXYLIC ACID ETHYL ESTER




20 The title compound from Preparative Example 47B (3.00g, 7.60 mmol) was dissolved in hydrobromic acid (48%, 30 mL). The reaction mixture was cooled to -5°C (ice-ethylene glycol bath) and bromine(2 mL) was added dropwise. The reaction mixture was stirred at -5°C for 15 minutes. Sodium nitrite (1.57g, 22.8 mmol) dissolved in water (15 mL)

was slowly added to the reaction mixture. The reaction mixture was then stirred for 45 minutes and then quenched with 40% NaOH to pH ~10. The aqueous phase was then extracted with ethyl acetate(3x100mL). Combined ethyl acetate fractions were dried over sodium sulfate and then concentrated to give the title compound in 83% yield as a light brown solid. Mp = 146-148°C, MH+ 463.

EXAMPLE 1


10 1-(4-PYRIDYLACETYL)-4-(8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)PIPERIDINE



To a mixture of 528 mg (1.7 mmol) of 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)piperidine (product from Preparative Example 1, Step G), 274 mg (1.7 mmol) of 4-pyridylacetic acid hydrochloride, and 242 mg (1.8 mmol) of 1-hydroxybenzotriazole hydrate in 5 mL of dry methylene chloride at -15 °C and under a nitrogen atmosphere was added dropwise 0.17 mL (1.5 mmol) of triethylamine followed by a solution of 363 mg (1.9 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (DEC) in 5 mL of dry methylene chloride. The reaction mixture was slowly allowed to warm to room temperature. After 4 hours the mixture was poured into water and extracted several times with methylene chloride. The combined organic portions were dried over MgSO₄, filtered, and concentrated *in vacuo* to give a product which was purified via flash chromatography (3% methanol saturated with ammonia in methylene chloride) 155 mg of 1-(4-pyridylacetyl)-4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)piperidine as a solid: mp 152 - 155 °C.

By essentially the same procedure as set forth in Example 1, but using the carboxylic acids set forth in column 1, of Table 2 below, in place of 4-pyridylacetic acid, one can obtain the compounds listed in column 2

of Table 2. The compounds listed in Table 2 refer to compounds of Formula 500.00:



wherein R is the substituent in Column 2.

5

TABLE 2

EXAMPLE	COLUMN 1	COLUMN 2	COMPOUND
2			glass
3			white powder
4			white solid
5			white crystals mp 200°C
6			mp 122 - 125°C
7			----

TABLE 2 - continued





















EXAMPLE	COLUMN 1	COLUMN 2	COMPOUND
8			off white powder
9			glass
10			white solid
11			white solid
12			glass
13			white solid
14			glass
15			mp 176 - 178°C
16			glass
17			mp 200 - 204°C

TABLE 2 - continued





















EXAMPLE	COLUMN 1	COLUMN 2	COMPOUND
18			glass
19			glass
20			yellow solid
21			off white solid
22			white solid mp 228°C (dec)
23			white solid mp 205-207°C
24			white powder
25			white powder
26			glass
27			glass

TABLE 2 continued



















EXAMPLE	COLUMN 1	COLUMN 2	COMPOUND
28			glass
29			glass
30			mp 211 - 215°C
31			yellow solid
32			white solid
33			white solid
34			glass
35			solid mp 190 - 193°C
36			solid

TABLE 2 - continued




































EXAMPLE	COLUMN 1	COLUMN 2	COMPOUND
37			glass
38			white solid
39			glass
40			mp 218 - 220°C
41			light brown solid mp = 92.7 - 93°C MS M+ = 459
42			white solid mp = 114.2 - 115.8°C MS M+ = 506
43			white solid mp = 93.3 - 94.6°C MS M+ = 506
44			white solid mp = 112 - 114.6°C MS M+ = 428
45			white solid mp = 94.3 - 95.5°C MS M+ = 474

TABLE 2 - continued

EXAMPLE	COLUMN 1	COLUMN 2	COMPOUND
46			white solid mp = 126.5 - 127.5°C MS M+ = 607
47			white solid mp = 83.6 - 85.0°C
48			white solid mp = 82.7 - 83.8°C MS M+ = 456
49			white solid MS M+ = 534
49a			white solid mp = 73.5 - 73.8°C
288	 Preparative Ex. 25		white solid MH+ 452
299	 Preparative Ex. 26		off white solid MH+ 459
300	 Preparative Ex. 10		white solid MH+ 459

EXAMPLE 50

1-(2-THIOPHENEACETYL)-4-(8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)PIPERIDINE




5 To a solution of 1.0 gm (3.22 mmole) of 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta [1,2-b] pyridin-11-ylidene) piperidine and 0.29 mL of pyridine in 20 mL of dry methylene chloride at 0°C and under an argon atmosphere was added dropwise 0.438 mL (3.55 mmol) of 2-thiopheneacetyl chloride. After 30 minutes the mixture was washed with
10 1.0 N aqueous sodium hydroxide and then brine. The organic portion was dried over sodium sulfate, filtered and converted in vacuo to provide a residue which was purified via flash chromatography (3% methanol in methylene chloride) and treated with activated carbon to provide the title compound as a glass.

15

















EXAMPLE 51

By essentially the same procedure as set forth in Example 50, but using the acid chlorides set forth in Column 1, in Table 3 below, in place of 2-thiopheneacetyl chloride, one can obtain the compounds listed in
20 Column 2 of Table 3. The compounds listed in Table 3 refer to compounds of Formula 500.00:




wherein R is the substituent in Column 2

TABLE 3


EXAMPLE	COLUMN 1	COLUMN 2	COMPOUND
52			solid
53			solid mp 158 - 160°C
54			glass
55			white powder
56			solid mp 126 - 128°C
57			solid mp 137 - 139°C
58			solid mp 104 - 106°C
59			white solid mp 155 - 157°C

EXAMPLE 65

- 5 By essentially the same procedures as set forth in Example 50 above, or Example 4 of US 5,089,496, but using




in place of 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]-pyridin-11ylidene)piperidine, one can obtain the compound



5 as a white solid.

EXAMPLE 75

1-(8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLO-HEPTA[1,2-b]PYRIDIN-11-YL-4-(4-PYRIDYLACETYL)-PIPERAZINE




10

To a mixture of 8.5 g (27.2 m mole) of 8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (Preparative Example 7) in 256 mL of anhydrous dimethylformamide at room temperature and under an argon atmosphere was added 2.98 g (27.2 m

mole of 4-methylmorpholine, 7.81 g (27.2 m mole) of 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride, 3.68 g (27.2 m mole) of 1-hydroxybenzotriazole, and 3.72 g (27.2 m mole) of 4-pyridylacetic acid. The mixture was stirred at room temperature for 22 hours. The mixture was poured into 3300 mL of methylene chloride and washed with 500 mL of water. The aqueous layer was extracted with 500 mL of methylene chloride. The combined organic portions were dried over magnesium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel column chromatography using a solution of 1.5% (10% ammonium hydroxide in methanol) in methylene chloride. The product was obtained as a white amorphous solid, M.S. (Mass Spec) $M^+ = 433$.

By essentially the same procedures as set forth in Example 75 above but using the compounds set forth in Column 1, Table 4 below, in place of 4-pyridylacetic acid, one can obtain compounds of the formula











wherein R is as listed in Column 2 of Table 4.

TABLE 4


EX.	COLUMN 1	COLUMN 2	CMPD
76			white amorphous solid Mass Spec $M^+ = 501$
77			white amorphous solid Mass Spec $M^+ = 512$

TABLE 4 - continued

EX.	COLUMN 1	COLUMN 2	CMPD
78			white amorphous solid Mass Spec M+ = 433
79			white amorphous solid Mass Spec M+ = 508
80			white amorphous solid Mass Spec M+ = 432
81			white amorphous solid Mass Spec M+ = 459

EXAMPLE 82


8-CHLORO-11-[1-(2-(4-PYRIDYL)ACETYL)-4-PIPERIDYL]-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]-PYRIDINE



- 5 Dissolve 0.1 g (0.32 m mole) of 8-chloro-11-4-piperidyl]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]-pyridine (from Example 233), 0.06 g (0.32 m mole) of 4-pyridylacetic acid, 0.092 g (0.48 m mole) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 0.065 g (0.48 m mole) of N-hydroxy benzotriazole and 0.048 g (0.50 m mole) of N-methyl morpholine in 5 mL of dimethylformamide and stir at room temperature for
- 10 18 hours under nitrogen. Concentrate under vacuo and partition between 100 mL each of ethyl acetate and water. Dry the organic layer over sodium sulfate and concentrate under vacuo. The resulting residue is chromatographed on silica gel using 98% dichloro methane and 2%
- 15 methanol, saturated with ammonia as the solvent, giving the product as a white waxy solid, mass spec $M^+ = 431$.

EXAMPLE 82A

20 8-CHLORO-11-[1-(2-(PYRIDYL)ACETYL)-4-PIPERIDYL]-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE




By essentially the same procedure as set forth in Example 82, but using 3-pyridylacetic acid instead of 4-pyridylacetic acid, the title compound is obtained as a white solid ($M^+ = 431$, $mp = 81.7-82^\circ C$).

5

EXAMPLE 83

8-CHLORO-11-[1-(2-METHYLSULFONYLOXY-1-PHENYLETHYL-CARBONYL)-4-PIPERIDYLIDENE]-6,11-DIHYDRO-5H-BENZO[5,6]-CYCLOHEPTA[1,2-b]-PYRIDINE



10


Dissolve 0.40 g (0.9 m mole) of 8-chloro-11-[1-(2-hydroxy-1-phenylethylcarbonyl)-4-piperidylidene]-6,11-dihydro-5H-benzo[5,6]-cyclohepta-[1,2-b]pyridine (Example 41 of Table 2) in 10 mL of pyridine and stir under nitrogen. Add 0.15 g (1.3 m mole) of methanesulfonyl chloride and stir for 20 hours. Concentrate under vacuo and triturate the residue with ether. Purify the resulting solid by silica gel chromatography using 2% methanol saturated with ammonia, and 98% dichloromethane as the solvent. The product is obtained as a white solid, $mp = 110.7-111.6^\circ C$.

15

C.

EXAMPLE 84

8-CHLORO-11-[1-(2-ACETYL MERCAPTO-1-PHENYLETHYL-CARBONYL)-4-PIPERIDYLIDENE]-6,11-DIHYDRO-5H-BENZO[5,6]-CYCLOHEPTA[1,2-b]-PYRIDINE




5

Dissolve 0.3 g (0.56 m mole) of 8-chloro-11-[1-(2-methanesulfonyloxy-1-phenylethylcarbonyl)-4-piperidylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (Formula 5.6 of Example 83) in 5 mL of dimethylformamide and add 0.2 g (0.6 m mole) of cesium thioacetate (preparation described in Synthetic Communications, 13, 553, 1983). Stir the reaction at 80°C for twenty hours then concentrate under vacuo. Purify the residue by silica gel chromatography using 70% ethyl acetate and 30% hexane as the solvent. The product is obtained as a light brown solid, mp = 92.7-93°C.

15

EXAMPLE 85

8-CHLORO-11-[1-(1-(2,3-DIHYDRO-3-OXO-1,2-BENZOISOTHIAZOL-S,S-DIOXIDE-2-YL)METHYLCARBONYL)-4-PIPERIDYLIDENE]-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]-PYRIDINE




20

Dissolve 0.46g (1.7 m mole) of 8-chloro-11-[1-(2-hydroxyethyl-carbonyl)-4-piperidylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2,b]pyridine (Example 49a of Table 2) in 30 mL of dimethylformamide and stir at 0°C under nitrogen. Add 0.55 g (2.1 m mole) of triphenyl phosphine and 0.36 g (2.1 m mole) of diethyl azodicarboxylate. Stir reaction mixture at 70°C for 3 days, then concentrate under vacuo. The residue was dissolved in 50 mL of 1 N hydrochloric acid and washed with 100 mL of ethyl acetate. The water layer was neutralized with 1 N sodium hydroxide and the mixture was extracted with ethyl acetate. The organic layer was dried over magnesium sulfate and concentrated under vacuo. The residue was purified by silica gel chromatography using 90% ethyl acetate and 10% hexane as the solvent, giving the product as a white solid, mass spec. M+ = 534.

15

EXAMPLE 86

8-CHLORO-11-[1-(1-(3-PYRIDYL)METHYLTHIOCARBONYL)-4-PIPERIDYLIDENE]-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]-PYRIDINE




20 Dissolve 0.50 g (0.12 m mole) of 8-chloro-11-[1-(1-(3-pyridyl)-methylcarbonyl)-4-piperidylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]-pyridine (Example 2 of Table 2) and 0.5 g (0.12 m mole) of 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (Lawesson's Reagent) in 15 mL of toluene and stir for 18 hours at room temperature and 18 hours at 80°C, under nitrogen. Filter the reaction mixture and concentrate under vacuo. Dissolve the residue in 50 mL of 1N hydrochloric acid and extract with 200 mL of dichloromethane. The water layer was neutralized with sodium carbonate and extracted with dichloromethane. The organic layer was dried over magnesium sulfate

25

and concentrated under vacuo giving the product as a white solid, mp = 92-93°C.

EXAMPLE 87


5 10,11-DIHYDRO-5-[1-(1-(4-PYRIDYL)METHYLCARBONYL)-4-PIPERIDYLIDENE]-5H-DIBENZO[a,d]CYCLOHEPTENE



Dissolve 0.15 g (0.6m mole) of 10,11-dihydro-5-(4-piperidylidene)-5H-dibenzo[a,d]cycloheptene, 0.096 g (0.55 m mole) of 4-pyridylacetic acid hydrochloride, 0.16 g (0.83 m mole) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and 0.075 g (0.55 m mole) of N-hydroxy benzotriazole in 5 mL of dimethylformamide and stir at room temperature for 18 hours under nitrogen. Concentrate under vacuo and partition between 100 mL each of ethyl acetate and 10% aqueous sodium hydrogenphosphate. Dry the organic layer over magnesium sulfate and concentrate under vacuo. The resulting residue is chromatographed on silica gel using 98% dichloro methane and 2% methanol, saturated with ammonia as the solvent, giving the product as a white waxy solid, mp = 162.8-163.4 °C.


EXAMPLE 180

1- 1-(4-PYRIDINYLACETYL) -4-[3,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE] - PIPERIDINE



- 5 Dissolve 0.18 g (0.51 mmole) of 3,8-dichloro 11-(1-acetyl-4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine, 0.088g (0.51 mmole) 4-pyridylacetic acid, 0.117g (0.61 mmole) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, 0.082g (0.61
- 10 mmole) N-hydroxybenzotriazole and 0.071g (0.71 mmole) N-methyl morpholine in 5 mL of dimethylformamide and stir for 18 hours under nitrogen. Concentrate under vacuo and partition between ethyl acetate and water. Dry organic layer over sodium sulfate and concentrate in vacuo. The resulting residue is chromatographed on silica gel using 95%
- 15 dichloromethane and 5% methanol, saturated with ammonia as the solvent. The product is obtained as white solid, mp = 113-114 °C.


EXAMPLE 181



- 20 By essentially the same procedure as set forth in Exampe 180, but using 8-bromo-11-(1-acetyl-4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine instead of 3,8-dichloro-11-(1-acetyl-4-

piperidylidene)-6,11-dihydro-5H-benzo[5,6]-cyclohepta[1,2-b]pyridine, compound 5.48 was obtained as an off-white solid, mp= 94.3-94.7 °C.

EXAMPLE 182




5

To a stirred solution of phenyl isocyanate (1.27 mmole) in 15 ml of anhydrous methylene chloride at room temperature and under an argon atmosphere was added dropwise over 20 minutes, a solution of 8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]-cyclohepta[1,2-b]pyridine (1.27 mmole) in 5 ml of anhydrous methylene chloride. The mixture was stirred at room temperature for 20 hours. The mixture was poured into 700 ml of methylene chloride and washed with 100 ml of saturated sodium bicarbonate. The organic portion was dried over magnesium sulfate, filtered, and concentrated in vacuo. The residue was purified by silica gel chromatography using a solution of 1.0% (10% ammonium hydroxide in methanol) in methylene chloride. The product was obtained as a white amorphous solid, M.S. (Mass Spec) $M^+ = 433$.

10

15

EXAMPLE 183




To a 5.0 ml reaction vial was added 8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (1.0 mmole) and N-ethoxycarbonyl-4-aminopyridine (0.99 mmole). The vial was capped and placed in an oil bath at 170 °C and stirred for 5 hours. The residue was purified by silica gel chromatography using a solution of 3.0% (10% ammonium hydroxide in methanol) in methylene chloride. The product was obtained as a white amorphous solid, M.S. (Mass Spec) $M^+ = 434$.

10

EXAMPLE 184

1-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA [1,2-b]PYRIDIN-11-YL)-4-(3-PYRIDINYLACETYL)piperazine 1-N-OXIDE




The title compound from Preparative Example 11D (0.5 grams) (1.5 mmoles) was reacted with 3-pyridylacetic acid (0.208 grams) (1.5 mmoles) under the conditions described in Example 75 to give the title compound (Yield: 0.439 grams, 95%, $MH^+ 449$).

15

EXAMPLE 185

**1-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA
[1,2-b]PYRIDIN-11-YL)-4-(3-PYRIDINYLACETYL 1-N-OXIDE)PIPERAZINE
1-N-OXIDE**




5

The title compound from Preparative Example 11D (0.5 grams) (1.5 mmoles) was reacted with the title compound from Preparative Example 8 (0.232 grams) (1.5 mmoles) under the conditions described in Example 75 to give the title compound (Yield: 0.6454 grams, 92%, MH⁺ 465.2).

10

EXAMPLE 186

**N-BENZYL 4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11-YL)-1-PIPERAZINECARBOXAMIDE**




The title compound from Example 75 was reacted with benzyl isocyanate under the conditions described in Example 182 above to give the title compound (Yield: 79%, MH^+ 447).

5


EXAMPLE 187

By essentially the same procedure as Example 183, with the exception that 3-ethoxycarbonylaminopyridine or 2-ethoxycarbonylaminopyridine (Preparative Example 12) is used instead of using 4-ethoxycarbonylaminopyridine, the compound



White amorphous
solid, MH^+ 434.3

or




White amorphous
solid, MH^+ 434.3

10

was obtained, respectively.

EXAMPLE 188

4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YL)-N-METHYL-N-(3-PYRIDINYL)-1-PIPERAZINE-CARBOXAMIDE



5


Compound 6.7 from Example 187 (10 grams) (23.1 mmoles) in DMSO (37.6 ml.) was added to a solution of powdered potassium hydroxide (2.62 grams) (23.1 mmoles) in DMSO (25 ml.) and the mixture was stirred at 25°C for 3 minutes. Iodomethane (1.4518 ml.) (23.1 mmoles) was added and the mixture was stirred at 25°C for 15 minutes. The mixture was poured into water and extracted with dichloromethane. The latter was dried (magnesium sulphate), filtered and evaporated to dryness. The product was purified by chromatography on silica gel using 3-5% (10% concentrated ammonium hydroxide in methanol)-dichloromethane as the eluant to give the title compound (Yield: 6.28 grams, 61%, MH⁺ 448).

15

EXAMPLES 189 - 218

By essentially the same procedures as set forth in Example 75 above but using the compounds set forth in Column 1, Table 5 below, in place of 4-pyridylacetic acid, one can obtain compounds of the formula

20



wherein R is as listed in Column 2 of Table 5.

TABLE 5







EX.	COLUMN 1	COLUMN 2	CMPD
189		 (5.62)	white amorphous solid MH ⁺ 475
190		 (5.63)	white amorphous solid MH ⁺ 460
191		 (5.64)	white amorphous solid MH ⁺ 447

TABLE 5 - continued







EX.	COLUMN 1	COLUMN 2	CMPD
192			white amorphous solid MH ⁺ 467
193			white amorphous solid MH ⁺ 539
194			white amorphous solid MH ⁺ 467

TABLE 5 - continued







EX.	COLUMN 1	COLUMN 2	CMPD
195			white amorphous solid MH ⁺ 439
196			white amorphous solid MH ⁺ 433
197			white amorphous solid MH ⁺ 449

TABLE 5 - continued







EX.	COLUMN 1	COLUMN 2	CMPD
198			white amorphous solid MH ⁺ 461
199		 5.72A = Isomer A 5.72B = Isomer B	white amorphous solid MH ⁺ 467
200			white amorphous solid MH ⁺ 467

TABLE 5 - continued








EX.	COLUMN 1	COLUMN 2	CMPD
201			white amorphous solid MH ⁺ 453
202			white amorphous solid MH ⁺ 525
203	Trifluoroacetic Acid deprotection of Compound 5.75 of Example 202		white amorphous solid MH ⁺ 525
204			white amorphous solid MH ⁺ 497

TABLE 5 - continued











EX.	COLUMN 1	COLUMN 2	CMPD
205		(5.78) 	white amorphous solid MH ⁺ 481
206		(5.79) 	white amorphous solid MH ⁺ 453
207	 (Ph=phenyl)	(5.80) 	white amorphous solid MH ⁺ 505
208	 (tBu=t-butyl)	(5.81) 	white amorphous solid MH ⁺ 471
209		(5.82) 	white amorphous solid MH ⁺ 489

TABLE 5 - continued




















EX.	COLUMN 1	COLUMN 2	CMPD
210		 (5.83)	white amorphous solid MH+ 505
211		 (5.84)	white amorphous solid MH+ 505
212		 (5.85)	white amorphous solid MH+ 505
213		 (5.86)	white amorphous solid MH+ 595
214		 (5.87)	white amorphous solid MH+ 561

TABLE 5 - continued

EX.	COLUMN 1	COLUMN 2	CMPD
215		 (5.88)	white amorphous solid MH+ 461
216		 (5.89)	white amorphous solid MH+ 591
217		 (5.90)	white amorphous solid MH+ 503
218		 (5.91)	white amorphous solid MH+ 519

EXAMPLE 219 - 222

- 5 By essentially the same procedure as set forth in Example 1, but using the acids set forth in Column 1 of Table 6 below in place of 4-pyridylacetic acid, the compounds listed in Column 2 of Table 6 are obtained. The compounds listed in Table 6 refer to compounds of Formula 500.00:




wherein R is the substituent in Column 2.

TABLE 6

EX.	COLUMN 1	COLUMN 2	CMPD
219		(5.92)	white amorphous solid MH ⁺ 482 m.p. = 192-193°C
220		(5.93)	white amorphous solid MH ⁺ 502 m.p. = 282-285°C
221		(5.94)	white amorphous solid MH ⁺ 485
222		(5.95)	white amorphous solid MH ⁺ 514


EXAMPLE 223

**A. (+)-1-(8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11(R)-YL)-4-(3-PYRIDINYLA-
CETYL)PIPERAZINE**



5 The title R(+) diastereoisomer from Preparative Example 19 was reacted with 3-pyridylacetic acid under the same conditions as described in Example 75 to give the title compound (Yield: 88%, MH⁺ 433).

**B. (-)-1-(8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11(S)-YL)-4-(3-PYRIDINYLA-
CETYL)PIPERAZINE**




10 The title S(-) diastereoisomer from Preparative Example 19 above was reacted with 3-pyridylacetic acid under the same conditions as described in Example 75 to give the title compound (Yield: 96%, MH⁺ 433).

15

EXAMPLE 224

A. (+)-4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5.6]CYCLO-
HEPTA[1.2-b]PYRIDIN-11(R)-YL)-N-(3-PYRIDINYL)-1-PIPERAZINE-
CARBOXYLATE




5

The title R(+) diastereoisomer from Preparative Example 19 was reacted with 3-ethoxycarbonylaminopyridine under the same conditions as described in Example 75 to give the title compound (Yield: 81%, MH⁺ 434).

10

B. (-)-4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5.6]CYCLO-
HEPTA[1.2-b]PYRIDIN-11(S)-YL)-N-(3-PYRIDINYL)-1-PIPERAZINE-
CARBOXAMIDE




15

The title S(-) diastereoisomer from Preparative Example 19 was reacted with 3-ethoxycarbonylaminopyridine under the same conditions

as described in Example 75 to give the title compound (Yield: 80%, MH^+ 434).


EXAMPLE 225

- 5 A. (+)-1-(8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11(R)-YL)-4-[(1-ACETYL-4-PIPERIDINYL)-
ACETYL]PIPERAZINE



- 10 The title R(+) diastereoisomer from Preparative Example 19 above was reacted with 1-N-acetylpiperidinyl-3-acetic acid under the same conditions as described in Example 75 to give the title compound (Yield: 52%, MH^+ 481).

B. (-)-1-(8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11(S)-YL)-4-[(1-ACETYL-4-PIPERIDINYL)-
ACETYL]PIPERAZINE




- 5 The title S(-) diastereoisomer from Preparative Example 19 above was reacted with 1-N-acetylpiperidinyl-3-acetic acid under the same conditions as described in Example 75 to give the title compound (Yield: 53%, MH^+ 481).

10


EXAMPLE 226

A. (+)-1-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11(R)-YL)-4-[(1-ACETYL-4-PIPERIDINYL)-
CARBONYL]PIPERAZINE



The title R(+) diastereoisomer from Preparative Example 19 was reacted with 1-N-acetylisonipecotic acid under the same conditions as described in Example 75 to give the title compound (Yield: 90%, MH⁺ 467).


5 B. (-)-1-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11(S)-YL)-4-[(1-ACETYL-4-PIPERIDINYL)-CARBONYL]PIPERAZINE



10 The title S(-) diastereoisomer from Preparative Example 19 was reacted with 1-N-acetylisonipecotic acid under the same conditions as described in Example 75 to give the title compound (Yield: 93%, MH⁺ 467).

EXAMPLE 227


15 4-(8-CHLORO-5,6-DIHYDRO-11H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-[(4-PYRIDINYL)ACETYL]-PIPERIDINE N1 OXIDE



To a mixture of 0.933g(3 mmol) of 4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta(1,2-b)pyridin-11-ylidene)-piperidine (product from Preparative Example 1, step G), 0.46g(3 mmol) of 4-pyridyl acetic acid N-oxide (prepared as described in Preparative Example 8) 1-hydroxybenzo-
5 triazole (0.40g, 3 mmol) in 20 mL of DMF at ~ 4°C and under nitrogen atmosphere was added N- methyl morpholine(1.65 mL, 15 mmol) followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride salt (DEC) an reaction stirred overnight at room temperature. The
10 volatiles were stripped off and the resulting semi-solid was partitioned between water and ethyl acetate. The aqueous phase was washed twice with ethyl acetate. Combined ethyl acetate fractions were dried over Na₂SO₄ and concentrated. The crude product was purified via flash chromatography on silica gel (first eluting with 3% and then 5% methanol saturated with ammonia in methylene chloride) to give the title compound
15 as a light brown solid(0.2g mp=128-130 °C MH⁺446).

EXAMPLE 228


4-(8-CHLORO-5,6-DIHYDRO-11H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-[(3-PYRIDINYL)ACETYL]-PIPERIDINE N1
20 OXIDE



By essentially the same procedure as set forth in Example 227, but using 3-pyridyl acetic acid N-oxide (Preparative Example 9) instead of 4-pyridyl acetic acid N-oxide the title compound was obtained as a white
25 solid (mp = 120-121°C, MH⁺=466).

EXAMPLE 229

4-(8-CHLORO-5,6-DIHYDRO-11H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-[(3-PYRIDINYL)ACETYL]-PIPERIDINE N4 OXIDE




4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-[(3-pyridinyl)acetyl]piperidine (1.0g, 2.33mmol) was dissolved in dry methylene chloride(50mL) at -10°C. 3-Chloroperbenzoic acid (80-85% purity 1.1g, 5.13 mmol) was added and the reaction stirred at that temperature for 95 minutes. The reaction mixture was washed with sodium bisulfite and then with 10% NaOH. The organic phase was dried over magnesium sulfate and then concentrated. Purification on silica gel eluting, first with 4%, 6% and then 10% methanol in methylene chloride gave rise to the title compound as a white solid (0.2g, 0.77 mmol

10

15 $MH^+=446$).

EXAMPLE 230

4-(8-CHLORO-5,6-DIHYDRO-11H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-[(4-PYRIDINYL)ACETYL]-PIPERIDINE N4 OXIDE




By essentially the same procedures as set forth in Example 229 above, but using 4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta[1,2-

b]pyridin-11-ylidene)-1-[(4-pyridinyl)acetyl]-piperidine instead of 4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-[(3-pyridinyl)acetyl]-piperidine the title compound was obtained as an off-white solid ($MH^+=446$).

5

EXAMPLE 231

A. 8-CHLORO-11-(1-ETHOXYCARBONYL-4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE N-OXIDE




10

8-chloro-11-(1-ethoxycarbonyl-4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (5g, 13.06 mmol) was dissolved in methylene chloride at -10°C . 3-Chlorobenzoic acid (4.9g, 15.67 mmol) was then added and the reaction mixture stirred for 95 minutes. The reaction mixture was taken up in methylene chloride and extracted with sodium bisulfite, 10% sodium hydroxide. The crude reaction product was purified on silica gel eluting first with 1% and then with 2% methanol in methylene chloride to give the title compound (2.7g, $MH^+ 399$).

15

20


B. 8-CHLORO-11-(4-PIPERIDYLIDENE)-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE N-OXIDE



By essentially the same procedures as set forth in Preparative Example 1 step G, but using 8-chloro-11-(1-ethoxycarbonyl-4-

piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine N-oxide instead of 8-chloro-11-(1-ethoxycarbonyl-4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine, the title compound was obtained and used for the next reaction without further purification (MH+ 327).


C. 4-(8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA(1,2-b)PYRIDIN-11-YLIDENE)-1-[(3-PYRIDINYL)ACETYL]-PIPERIDINE N1,N4 DIOXIDE



By essentially the same procedure as set forth in Example 227, but using 3-pyridyl acetic acid N-oxide (Preparative Example 9) instead of 4-pyridyl acetic acid N-oxide and 8-chloro-11-(4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine N-oxide instead of 8-chloro-11-(4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine, the title compound was obtained as a white solid (mp=105-107°C, MH+=462).

EXAMPLE 232


4-(8-CHLORO-5,6-DIHYDRO-11H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-[(4-PYRIDINYL)ACETYL]-PIPERIDINE N1,N4 DIOXIDE



By essentially the same procedure as set forth in Example 227, but using 8-chloro-11-(4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine N-oxide instead of 8-chloro-11-(4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine, the title compound was obtained as a light brown solid (MH⁺=462).


EXAMPLE 233

A. 8-CHLORO-6,11-DIHYDRO-11-(4-PIPERIDINYL)-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE (Product A) and
10 6,11-DIHYDRO-11-(4-PIPERIDINYL)-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE (Product B)



To a solution 66.27g (0.21mole) of 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta(1,2-b)pyridin-11-ylidene)-piperidine (product from Preparative Example 1 Example, step G), in THF (1L) was added lithium aluminum hydride (24.32g, 0.64 mole) and the reaction mixture was heated to reflux overnight. The reaction mixture was then cooled to room temperature and ~ 3L of diethyl ether is added followed by dropwise addition of saturated sodium sulfate until a white gray precipitate forms.
20 Magnesium sulfate was then added to the separated organic layer and stirred for 30 minutes. All the volatiles were then removed and the resulting crude mixture was chromatographed on a silica gel column eluting with 10% methanol saturated with ammonia in methylene chloride. The material obtained contained both the desired compound and the des-chloro compound. Separation on HPLC using reverse phase column and eluting with 40% methanol-water afforded the desired compounds as white solids (Product A's mp = 95.2-96.1°C, Product B's mp = 145.1-145.7°C).


B. 4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YL)-1-[(3-PYRIDINYL)ACETYL]-PIPERIDINE N1 OXIDE



- 5 By essentially the same procedure as set forth in Example 227, but using 3-pyridyl acetic acid N-oxide (Preparative Example 9) instead of 4-pyridyl acetic acid N-oxide and 8-chloro-6,11-dihydro-11-(4-piperidinyl)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (product from Example 233A) instead of 4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta(1,2-b)pyridin-11-ylidene)-piperidine, the title compound was obtained as a white solid (mp=117-118°C, MH⁺=414).

EXAMPLE 234


15 4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA(1,2-b)PYRIDIN-11-YL)-1-[(4-PYRIDINYL)ACETYL]-PIPERIDINE N1 OXIDE



- 20 By essentially the same procedure as set forth in Example 227, but using 8-chloro-6,11-dihydro-11-(4-piperidinyl)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (product from Example 233A) instead of 4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta(1,2-b)pyridin-11-ylidene)-piperidine (product from Preparative Example 1, step G), the title compound was obtained as a white solid (mp=125-126°C, MH⁺=414).

EXAMPLE 235

A. ETHYL 4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO-[5,6]CYCLOHEPTA(1,2-b)PYRIDIN-11-YL)-1-PIPERIDINE-CARBOXYLATE




5

8-Chloro-6,11-dihydro-11-(4-piperidyl)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine (product from Example 233A) (4.18g, 13mmol) was dissolved in toluene (175mL). Ethyl chloroformate(11.6g,110 mmol, 10.2 mL) was then added and the reaction mixture was heated to ~120°C overnight. All volatiles were stripped off and the crude product was purified on silica gel column eluting with 50% ethyl acetate- hexanes to give the title compound as a white solid(MH⁺ 385).

10

B. ETHYL 4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO-[5,6]CYCLOHEPTA(1,2-b)PYRIDIN-11-YL)-1-PIPERIDINECARBOXYLATE N OXIDE


15



20


By essentially the same procedure as set forth in Example 231, but using ethyl 4-(8-chloro-6,11-dihydro-5H-benzo-[5,6]cyclohepta(1,2-b)pyridin-11-yl)-1-piperidinecarboxylate (product from Example 235A) instead of 8-chloro-11-(1-ethoxycarbonyl-4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine, the title compound was obtained as a white solid (mp= 81.7-82.5°C, MH⁺=400).

C. 4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO-[5,6]CYCLOHEPTA(1,2-b)PYRIDIN-11-YL)-1- PIPERIDINE N OXIDE



By essentially the same procedure as set forth in Preparative Example 1 step G, but using ethyl 4-(8-chloro-6,11-dihydro-5H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-yl)-1- piperidinecarboxylate N1 oxide (product from Example 235B) instead of 8-chloro-11-(1-ethoxycarbonyl-4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine, the title compound was obtained as a solid ($MH^+=329$).


10 D. 4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO[5,6]CYCLOHEPTA(1,2-b)PYRIDIN-11-YL)-1-[(3-PYRIDINYL)ACETYL]- PIPERIDINE N4 OXIDE



By essentially the same procedure as set forth in Example 227, but using 3-pyridyl acetic acid instead of 4-pyridyl acetic acid N-oxide and 4-(8-chloro-6,11-dihydro-5H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-yl)-1-piperidine N oxide (product from Example 235C) instead of 4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-piperidine, the title compound was obtained as a white solid (mp=61.8-62.3°C, $MH^+=448$).

EXAMPLE 236


4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO-[5,6]CYCLOHEPTA(1,2-b)PYRIDIN-11-YL)-1-[(4-PYRIDINYL)ACETYL]- PIPERIDINE N4 OXIDE



- 5 By essentially the same procedure as set forth in Example 227, but using 4-pyridyl acetic acid instead of 4-pyridyl acetic acid N-oxide and 4-(8-chloro-6,11-dihydro-5H-benzo-[5,6]cyclohepta(1,2-b)pyridin-11-yl)-1-piperidine N oxide (product from Example 235C) instead of 4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-piperidine, the title compound was obtained as a white solid (mp=116.7-117.6°C, MH⁺=448).

EXAMPLE 237

15 4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO-[5,6]CYCLOHEPTA(1,2-b)PYRIDIN-11-YL)-1-[(3-PYRIDINYL)ACETYL]- PIPERIDINE N1, N4 OXIDE




- 20 4-(8-Chloro-6,11-dihydro-5H-benzo-[5,6]cyclohepta(1,2-b)pyridin-11-yl)-1-[(3-pyridinyl)acetyl]piperidine, from Example 82A, (0.5g, 1.2mmol) was dissolved in methylene chloride at about -18°C. 3-Chloroperbenzoic acid (0.62g, 3.6 mmol) was added and the reaction stirred for 1.5 hours. The reaction mixture was extracted with 10% sodium bisulfite, 10% sodium hydroxide and then dried with magnesium sulfate, filtered

and concentrated. The crude product was purified on silica gel eluting with 7% methanol saturated with ammonia in methylene chloride to give the title compound as a white solid (0.51g, 91% yield MH^+ 464).

5

EXAMPLE 238

4-(8-CHLORO-6,11-DIHYDRO-5H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YL)-1-[(4-PYRIDINYL)ACETYL]-PIPERIDINE N1, N4 OXIDE



10

By essentially the same procedure as set forth in Example 237, but using 4-(8-chloro-6,11-dihydro-5H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-yl)-1-[(4-pyridinyl)acetyl]piperidine (product from Example 82) instead of 4-(8-chloro-6,11-dihydro-5H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-yl)-1-[(3-pyridinyl)acetyl]piperidine, the title compound was obtained as a


15

white solid (mp=85-85.6 °C, MH^+ =464).

EXAMPLE 239

4-(8-CHLORO-3-METHOXY-5,6-DIHYDRO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-[(3-PYRIDINYL)ACETYL]-PIPERIDINE


20



By essentially the same procedure as set forth in Example 180, but using 8-chloro-3-methoxy-11-(4-piperidylidene)-6,11-dihydro-5H-benzo-[5,6]-cyclohepta[1,2-b]pyridine (Preparative Example 20) instead of 3,8-dichloro-11-(1-acetyl-4-piperidylidene)-6,11-dihydro-5H-benzo[5,6]-cyclohepta[1,2-b]pyridine the title compound was obtained as a white solid (MH^+ 460).

EXAMPLE 240


4-(8-CHLORO-3-HYDROXY-5,6-DIHYDRO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE)-1-[(3-PYRIDINYL)-ACETYL]-PIPERIDINE



4-(8-Chloro-3-methoxy-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-[(3-pyridinyl)acetyl]-piperidine (0.24g, 0.54 mmol) (Example 239) was dissolved in methylene chloride at 0°C under nitrogen atmosphere. Bromine tribromide (0.9g, 3.6 mmol, 3.6 mL) was added and the reaction was run at room temperature for two days. The reaction mixture was concentrated and chromatographed on a silica gel column eluting with 3% methanol saturated with ammonia in methylene chloride to give an off white borate salt solid (0.14g, 61% yield, MH^+ 446).

EXAMPLE 246

1-1(4-PYRIDINYLACETYL)-4-[3-BROMO8-CHLORO5-6-DIHYDRO-11H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11YLIDENE]-PIPERIDINE




5 By essentially the same procedure as set forth in Example 180 but using 4-(3-bromo-8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta(1,2-b]pyridin-11-ylidene)-piperidine instead of 4-(3,8-dichloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta(1,2-b]pyridin-11-ylidene)-piperidine, the title compound was obtained as a glassy solid (MH^+ 508).

10

EXAMPLE 247

1-1(3-PYRIDINYLACETYL)-4-[3-BROMO8-CHLORO5-6-DIHYDRO-11H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11YLIDENE]-PIPERIDINE




15 By essentially the same procedure as set forth in Example 180, but using 4-(3-bromo-8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta(1,2-b]pyridin-11-ylidene)-piperidine instead of 4-(3,8-dichloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-piperidine and 3-pyridyl acetic acid instead of 4-pyridyl acetic acid, the title compound was
20 obtained as a white solid (mp=92-93°C MH^+ 508).

EXAMPLE 248

4-[4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA-[1,2-b]PYRIDIN-11-YLIDENE]-1-(4-PYRIDINYLACETYL)-PIPERIDINE and


5 • 4-[2,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA-[1,2-b]PYRIDIN-11-YLIDENE]-1-(4-PYRIDINYLACETYL)-PIPERIDINE



A solution of the title compound from Example 230 (1.7 grams) and phosphorous oxychloride (21 mL) dissolved in chloroform (12 mL) was stirred at reflux for 1 hour. Concentration *in vacuo* provided a residue
 10 which was diluted with dichloromethane and washed with saturated aqueous sodium bicarbonate and brine. The organic phase was dried over anhydrous magnesium sulfate, concentrated *in vacuo*, and purified by flash column chromatography (silica gel) using 2% methanol-dichloromethane to afford the title 4,8-dichloro compound (0.34 grams,
 15 20% yield, mp 84-91 °C, MH⁺ 464) and the title 2,8-dichloro compound (0.18 grams, 11% yield, mp 163.8-164.6 °C, MH⁺ 464).

EXAMPLE 249

4-[4-[(1H-BENZOTRIAZOL-1-YL)OXY]-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(4-PYRIDINYLACETYL)-PIPERIDINE




5

A mixture of the 4,8-dichloro compound from Example 248 (0.5 grams), 1-hydroxybenzotriazole hydrate (0.4 grams) and anhydrous dimethylformamide (20 mL) was stirred at 25°C under N₂ for 5 days. The mixture was concentrated *in vacuo*, diluted with dichloromethane, and washed with 1N aqueous sodium hydroxide. The organic phase was dried over anhydrous magnesium sulfate, concentrated *in vacuo* and purified by flash column chromatography (silica gel) using 3-5% methanol-dichloromethane to give the title compound (0.58 grams, 96%, mp 98.6-101.6 °C, MH⁺ 563).

15

EXAMPLE 250

4-[4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(3-PYRIDINYLACETYL)-
PIPERIDINE





5

A mixture of the 4,8-dichloro product from Preparative Example 28 (1.91 grams), 3-pyridylacetic acid hydrochloride (2.1 grams), 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (1.6 grams), 4-methylmorpholine (1.4 mL) and anhydrous dimethylformamide (100 mL) was stirred at 25 °C overnight. Concentration *in vacuo* provided a residue which was diluted with dichloromethane and water. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo* to provide the title compound (2.2 grams, 87%, mp 59.8-63.5 °C, MH⁺ 464).

15

EXAMPLE 251

4-[4-[(1H-BENZOTRIAZOL-1-YL)OXY]-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(3-PYRIDINYLA CETYL)-PIPERIDINE



5


The 4,8-dichloro compound from Example 250 (0.8 grams) was added to a solution of 1-hydroxybenzotriazole hydrate (1.2 grams) and sodium hydride (0.14 grams, 60% in mineral oil) in anhydrous dimethylformamide (60 mL). The resulting solution was irradiated with a 200 W lamp while stirring at 25°C for 60 hours. The solution was poured into 1N aqueous sodium hydroxide while stirring and an additional 400 mL of water was added to the resulting mixture. Filtration provided a solid which was washed with water several times. The solid was dissolved in dichloromethane-acetone, dried over anhydrous magnesium sulfate, and concentrated in vacuo to provide the title compound (0.87 grams, 90%, mp = 120-122°C, MH⁺ 563).

10

15

EXAMPLE 252

4-[4-HYDROXY-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]-
CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(3-PYRIDINYLA
CETYL)-
PIPERIDINE




5

To a solution of the title compound from Example 251 (0.8 grams) and glacial acetic acid (30 mL) was added zinc dust (0.4 grams). After stirring at 25°C for 18 hour, the mixture was filtered through celite and the filtrate concentrated *in vacuo*. The residue was diluted with ethyl acetate, washed with saturated aqueous sodium bicarbonate and brine. The organic layer was separated, dried over magnesium sulfate and concentrated *in vacuo* to give the title compound (Yield 0.346 grams, 58%, MH+ 446).

10

EXAMPLE 253

4-[3-BROMO-4-HYDROXY-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(3-PYRIDINYLAACETYL)-PIPERIDINE



5


To a solution of the title compound from Example 252 (0.19 grams) and glacial acetic acid (4 mL) was added a 0.7 M bromine-acetic acid solution (0.7 mL) at 25°C under N₂. After 10 minutes, water was added and the resulting solid was filtered and washed with water several times and dried to give the title compound (0.18 grams, 71%, MH⁺ 526).

10

EXAMPLE 255

4-[8-CHLORO-4-(METHYLTHIO)-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(3-PYRIDINYLAACETYL)-PIPERIDINE

15




A mixture of the title compound from Example 250 (0.26 grams), sodium methylthiolate (0.06 grams) and anhydrous dimethylformamide (10 mL) was stirred while being irradiated with a 200 W lamp for 1 hour. After stirring an additional 12 hours at room temperature without

20

irradiation, the mixture was concentrated *in vacuo*, diluted with dichloromethane, and washed with 1N aqueous sodium hydroxide and brine. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo* to afford the title compound as a white foam
5 (0.3 grams, 100%, MH⁺ 476).

EXAMPLE 256


4-[8-CHLORO-4-(METHYLSULFINYL)-5,6-DIHYDRO-11H-
BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(3-
10 PYRIDINYLACETYL)-PIPERIDINE



To the title compound from Example 255 (0.18 grams) dissolved in anhydrous tetrahydrofuran (10 mL) was added 30% aqueous hydrogen peroxide (3 mL) and the resulting solution was stirred for 12 hours at
15 73°C. The solution was concentrated *in vacuo*, diluted with dichloromethane, and washed with water. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo* to afford the title compound after preparative plate chromatography (silica gel) using 3% methanol-dichloromethane (0.04 grams, 26%, MH⁺ 492).

EXAMPLE 257

METHYL [(8-CHLORO-6,11-DIHYDRO-11-[1-[1-OXO-2-(3-PYRIDINYL)ETHYL]-4-PIPERIDINYLIDENE]-5H-BENZO[5,6]CYCLO-HEPTA[1,2-b]PYRIDIN-4-YL)THIO]ACETATE



5


A mixture of the title compound from Example 250 (0.26 grams), sodium hydride (0.08 grams, 60% in mineral oil), methyl thioglycolate (0.19 mL) and anhydrous dimethylformamide (15 mL) was stirred while being irradiated with a 200 W lamp for 16 hours. The mixture was diluted with methanol, concentrated *in vacuo*, diluted with dichloromethane and water, and washed with 1N aqueous sodium hydroxide and brine. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo* and the residue purified by preparative plate chromatography (silica gel) using 3% methanol-dichloromethane to afford the title compound (0.05 grams, 15%, MH⁺ 534).

10

15

EXAMPLE 258

4-[8-CHLORO-5,6-DIHYDRO-4-(PHENYLMETHYLTHIO)-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(3-PYRIDINYLACETYL)-PIPERIDINE



5


A mixture of the title compound from Example 250 (0.25 grams), sodium hydride (0.11 grams, 60% in mineral oil), benzyl mercaptan (0.13 mL) and anhydrous dimethylformamide (15 mL) was stirred while being irradiated with a 200 W lamp for 10 days. Isolation and purification as in Example 257 provided the title compound (0.02 grams, 8%, MH⁺ 552).

10

EXAMPLE 259

4-[8-CHLORO-5,6-DIHYDRO-4-[(2-METHYL-2H-TETRAZOL-5-YL)THIO]-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(3-PYRIDINYLACETYL)-PIPERIDINE


15



A mixture of the title compound from Example 250 (0.24 grams), 5-mercapto-1-methyltetrazole sodium salt (0.6 grams) and anhydrous dimethylformamide (10 mL) was stirred while being irradiated with a 200 W lamp for 10 days. Isolation and purification as in Example 257 provided the title compound (0.2 grams, 68%, MH⁺ 544).

EXAMPLE 260


1,1-DIMETHYLETHYL [2-[[[8-CHLORO-6,11-DIHYDRO-11-[1-[1-OXO-2-(3-PYRIDINYL)ETHYL]-4-PIPERIDINYLDENE]-5H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-4-YL]THIO]ETHYL] CARBAMATE



A mixture of the title compound from Preparative Example 32 (0.032 grams), 3-pyridylacetic acid hydrochloride (0.05 grams), 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (0.03 grams), triethylamine (0.08 mL) and anhydrous dimethylformamide (4 mL) was stirred at 25 °C for 48 hours. Concentration *in vacuo* provided a residue which was diluted with dichloromethane and washed with 1N aqueous sodium hydroxide. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo* to provide the title compound (0.02 grams, 50%, mp 59.8-63.5 °C, MH⁺ 605).

EXAMPLE 261

4-(4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLO-
HEPTA[1,2-b]PYRIDIN-11-YLIDENE)-N-(3-PYRIDYL)-1-PIPERIDINE-
CARBOXAMIDE




5 A portion of the stock solution of 3-pyridylisocyanate (32 mL) prepared as described in Preparative Example 33 was added to the 4,8-dichloro product from Preparative Example 28 (1.37 grams) and the mixture was stirred at 25°C for 4 days. The mixture was evaporated to dryness and the residue was taken up in dichloromethane and washed with saturated aqueous sodium bicarbonate and then water. The organic solution was dried over magnesium sulfate, filtered and evaporated to dryness. The residue was purified by flash column chromatography silica gel) using 2% methanol-dichloromethane as eluent to give the title compound (Yield 1.25 grams, 70%, MH+ 465).

10

15

EXAMPLE 262


4-[4-[(1H-BENZOTRIAZOL-1-YL)OXY]-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-N-(3-PYRIDYL)-1-PIPERIDINECARBOXAMIDE



To a solution of the title compound from Example 261 (1.0 grams) in dry dimethylformamide (60 mL) was added 1-hydroxybenzotriazole (1.4 grams), sodium hydride (0.2 grams, 60% in mineral oil) and distilled water (0.5 mL). The solution was stirred at 25°C under nitrogen while being irradiated with a 200 Watt lamp for 20 hours. The reaction mixture was concentrated in vacuo, diluted with dichloromethane and saturated aqueous sodium bicarbonate and after two hours, the organic phase was separated, dried over magnesium sulfate and concentrated. Purification by flash column chromatography (silica gel) using 3-5% methanol-dichloromethane afforded the title compound (Yield 1.1 grams, 87%, MH+ 564).

EXAMPLE 263

4-[4-HYDROXY-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]-
CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-N-(3-PYRIDYL)-1-
PIPERIDINECARBOXAMIDE



5


To a solution of the title compound from Example 262 (0.86 grams) and glacial acetic acid (20 mL) was added zinc dust (0.5 grams). After stirring at 25°C for 1.5 hours, the mixture was filtered through celite and the filtrate concentrated *in vacuo*. The residue was purified by flash column chromatography (silica gel) using 5-10% methanol-dichloromethane saturated with ammonium hydroxide to give the title compound (Yield 0.47 grams, 69%, MH+448).

10

EXAMPLE 264

4-[3-BROMO-4-HYDROXY-8-CHLORO-5,6-DIHYDRO-11H-
BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-N-(3-PYRIDYL)-
1-PIPERIDINECARBOXAMIDE


15



To a solution of the title compound from Example 263 (0.34 grams) and glacial acetic acid (10 mL) was added a 0.7 M bromine-acetic acid solution (4 mL) at 25°C under N₂. After 10 minutes, water was added and the resulting solid was filtered and washed with water several times and dried to give the title compound (Yield 0.31 grams, 67%, MH⁺ 527).

EXAMPLE 266

4-(8-CHLORO-5H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11-YL)-1-(3-PYRIDINYLACETYL)-PIPERIDINE



10

The title compound from Preparative Example 34C (2.0g, 6.4 mmole) was dissolved in anhydrous dimethylformamide (70 mL) and the solution was cooled with an ice bath for 30 minutes. 4-Methylmorpholine (3.3 g, 32 mmole), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.8 g, 9.7 mmole) and 1-hydroxybenzotriazole (0.87g 6.4 mmole) were added to the cold solution. 3-Pyridylacetic acid (0.88 g, 6.4 mmole) was added and the cooling bath removed. Stir the mixture at room temperature for 18 hours. The reaction mixture was evaporated to dryness and the residue was diluted with water (50 mL). The aqueous mixture was extracted with ethyl acetate and the combined extracts dried (MgSO₄), filtered and evaporated. The resulting residue was purified by silica gel chromatography using a gradient of 97% dichloromethane/ 3% methanol saturated with ammonia to 93% dichloromethane/7% methanol saturated with ammonia as eluent to give the title compound (0.87g MH⁺ 430).


15

20

25

EXAMPLE 267


E. 4-(8-CHLORO-11H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11-YL)-N-(3-PYRIDINYL)-1-PIPERIDINECARBOXAMIDE



- 5 The title compound from Preparative Example 34C was treated with 3-pyridylisocyanate, similar to the procedure in Example 261, to afford the title compound (MH^+ 431).

EXAMPLE 268


10 **4-(8-CHLORO-5H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11-YL)-1-[2-METHYL-2-(3-PYRIDINYL)-1-OXOPROPYL]-PIPERIDINE**



- 15 The title compound from Preparative Example 34C was treated as described in Example 266, using α,α -dimethyl-3-pyridylacetic acid (described in Preparative Example 10B) in place of 3-pyridylacetic acid, to afford the title compound (M^+ 458).

EXAMPLE 269


4-(8-CHLORO-5H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11-YL)-1-(4-PYRIDINYLACETYL)-PIPERIDINE



- 5 The title compound from Preparative Example 34C above was treated as described in Example 266, using 4-pyridylacetic acid in place of 3-pyridylacetic acid, to give the title compound (M^+ 430).

EXAMPLE 270


10 4-(8-CHLORO-9-ETHYL-5H-BENZO[5.6]CYCLOHEPTA-[1.2-b]PYRIDIN-11-YL)-1-(3-PYRIDINYLACETYL)-PIPERIDINE



- 15 The title compound from Preparative Example 2A was treated as described in Example 266 to give the title compound (M^+ = 458, mp = 67.2-69.8°C).

EXAMPLE 273


4-(4,8-DICHLORO-5H-BENZO[5,6]CYCLOHEPTA[1,2-B]PYRIDIN-11-YL)-1-(3-PYRIDINYLALETYL)-PIPERIDINE



- 5 The title compound from Preparative Example 36C was treated as described in Example 266 to give the title compound (mp 100.1 - 103.4°C).

EXAMPLE 274

10 4-(4,8-DICHLORO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YL)-N-(3-PYRIDINYL)-1-PIPERIDINECARBOXAMIDE




- 15 The title compound from Preparative Example 36C (0.75g, 2.17mmol) was treated with a pyridine solution of 3-pyridylisocyanate (from Preparative Example 33). The reaction mixture was evaporated to dryness and the residue dissolved in dichloromethane. This solution was washed with saturated sodium bicarbonate solution and brine, dried (MgSO₄), filtered and evaporated to give a dark syrup. The syrup was purified by silica gel chromatography using a gradient of 97% dichloromethane/3% methanol saturated with ammonia to 93%

dichloromethane/7% methanol saturated with ammonia. The title compound was obtained as a yellow solid (0.13g, 13%, M+ 465)

EXAMPLE 276

5 4-(8-CHLORO-11H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11YLIDENE)-1-(3-PYRIDINYLACETYL)-PIPERIDINE




The title compound from Preparative Example 37B was treated as described in Example 266 to give the title compound (MH⁺ 428).

10

EXAMPLE 277

4-(8CHLORO-11H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11YLIDENE)-N-(3-PYRIDINYL)-1-PIPERIDINECARBOXAMIDE




15

The title compound from Preparative Example 37B above was treated as described in Example 261 above to give the title compound (mp 95.9 - 97.6°C).

EXAMPLE 278

4-(8-CHLORO-11H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11YLIDENE)-1-[2-METHYL-2-(3-PYRIDINYL)-1-OXO-PROPYL]-PIPERIDINE




5

The title compound from Preparative Example 37B was treated as described in Example 266 using α,α -dimethyl-3-pyridylacetic acid (described in Preparative Example 10B) in place of 3-pyridylacetic acid, to give the title compound (M+ 456).

10

EXAMPLE 279

4-(8-CHLORO-5,6-DIHYDRO-5-OXO-11H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11YLIDENE)-1-(3-PYRIDINYLALETYL)-PIPERIDINE




15

The preparation of the starting material for this reaction was described in *The Journal of Organic Chemistry*, 1990, 55, pp. 3341-3350 by Piwinski, J.J.; Wong, J.K.; Chan, T.-M.; Green, M.J.; and Ganguly, A.K. The procedure described in Example 266 was followed using 8-chloro-

6,11-dihydro-11-(4-piperidinylidene)-5H-benzo[5,6]cyclohepta[1,2-b]-pyridin-5-one to give the title compound (M^+ 443).

EXAMPLE 280

5 **4-(8-CHLORO-5,6-DIHYDRO-5-HYDROXY-11H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11YLIDENE)-1-(3-PYRIDINYL-ACETYL)-PIPERIDINE**




The preparation of the starting material for this reaction was described in *The Journal of Organic Chemistry*, 1990, 55, pp. 3341-3350 by Piwinski, J.J.; Wong, J.K.; Chan, T.-M.; Green, M.J.; and Ganguly, A.K. The procedure described in Example 266 was followed using 8-chloro-6,11-dihydro-5-hydroxy-11-(4-piperidinylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine to give the title compound (MH^+ 446).

15

EXAMPLE 281

4-(8-CHLORO-5,6-DIHYDRO-5-OXO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11YLIDENE)-1-(4-PYRIDINYLACETYL)-PIPERIDINE




20

The procedure of Example 279 was followed with the exception that 4-pyridylacetic acid was used in place of 3-pyridylacetic acid to give the title compound (MH^+ 444).

5

EXAMPLE 282

4-(8-CHLORO-5,6-DIHYDRO-5-HYDROXY-11H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11YLIDENE)-1-(4-PYRIDINYL-ACETYL)-PIPERIDINE




10

The procedure of Example 280 was followed with the exception that 4-pyridylacetic acid was used in place of 3-pyridylacetic acid to give the title compound (MH^+ 446).

EXAMPLE 283

15

4-(8-CHLORO-5,6-DIHYDRO-6-OXO-11H-BENZO[5,6]-CYCLOHEPTA[1,2-b]PYRIDIN-11YLIDENE)-1-(3-PYRIDINYLACETYL)-PIPERIDINE



The preparation of the starting material for this reaction was described in *The Journal of Organic Chemistry*, 1990, 55, pp. 3341-3350


20

by Piwinski, J.J.; Wong, J.K.; Chan, T.-M.; Green, M.J.; and Ganguly, A.K. The procedure described in Example 266 was followed using 8-chloro-6,11-dihydro-11-(4-piperidinylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridin-6-one to give the title compound (M^+ 444).

5

EXAMPLE 284

4-(8-CHLORO-5,6-DIHYDRO-6-HYDROXY-11H-BENZO-[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11YLIDENE)-1-(3-PYRIDINYL-ACETYL)-PIPERIDINE




10

The preparation of the starting material for this reaction was described in *The Journal of Organic Chemistry*, 1990, 55, pp. 3341-3350 by Piwinski, J.J.; Wong, J.K.; Chan, T.-M.; Green, M.J.; and Ganguly, A.K. The procedure described in Example 266 above was followed using 8-chloro-6,11-dihydro-6-hydroxy-11-(4-piperidinylidene)-5H-benzo[5,6]-cyclohepta[1,2-b]pyridine to give the title compound (MH^+ 446).

15

EXAMPLE 285

4-(8-CHLORO-5,6-DIHYDRO-6-OXO-11H-BENZO[5,6]-
CYCLOHEPTA[1,2-b]PYRIDIN-11YLIDENE)-1-(4-PYRIDINYLA
CETYL)-PIPERIDINE




5

The procedure of Example 283 was followed with the exception that 4-pyridylacetic acid was used in place of 3-pyridylacetic acid to give the title compound (M^+ 444).

10

EXAMPLE 286

4-(8-CHLORO-5,6-DIHYDRO-6-HYDROXY-11H-BENZO-
[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11YLIDENE)-1-(4-PYRIDINYLA
CETYL)-PIPERIDINE



15

The procedure of Example 284 was followed with the exception that 4-pyridylacetic acid was used in place of 3-pyridylacetic acid to give the title compound (MH^+ 446).

EXAMPLES 287, 289 AND 290


By essentially the same procedure as in Example 1, but using either (R)-(+)- α -methoxy- α -(trifluoromethyl)-phenylacetic acid (Example 290), (S)-(-)- α -methoxy- α -(trifluoromethyl)-phenylacetic acid (Example 287), or α,α -dimethylphenylacetic acid (Example 289), the compounds of Example 290, 287 and 289 were obtained. The structures for these compounds are in Table 7. Data for these compounds are: compound of Example 290, white solid MH+ 527; compound of Example 287, white solid MH+ 527; and compound of Example 289, white solid M+ 457.

EXAMPLES 291, 292, 294, 313 AND 314

By essentially the same procedure as in Example 183, and using either 4-, 3-, or 2-ethoxycarbonylaminopyridine and either 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)piperidine or 8-chloro-6,11-dihydro-11-(4-piperidiny)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (product of Example 233A), the compounds of Examples 291, 292, 294, 313 and 314 were obtained. The structures for the compounds of Examples 291, 292, and 294 are given in Table 7. The structures for the compounds of Examples 313 and 314 are given in Table 10. Data are: the compound of Example 291 was a yellow solid (MH+ 431), the compound of Example 292 was an off white solid (MH+ 431), the compound of Example 294 was an off white solid (MH+ 431), the compound of Example 313 was a white solid (MH+ 433), and the compound of Example 314 was a white solid (MH+ 433).

EXAMPLE 301

1-1-(4-PYRIDINYLACETYL)-4-[3-METHYL-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-PIPERIDINE




By essentially the same procedure as set forth in Example 180, but using 4-(8-chloro-3-methyl-5,6-dihydro-11-(4-piperidylidene)-11H-benzo[5,6]cyclohepta[1,2-b]pyridine (from Preparative Example 3E)

instead of 4-(3,8-dichloro-5,6-dihydro-11-(4-piperidylidene)-11H-benzo[5,6]cyclohepta[1,2-b]pyridine, the title compound was obtained as an off-white solid MH⁺ 444

5

EXAMPLE 303

1-1-(3-PYRIDINYLACETYL)-4-[3-METHYL-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-PIPERIDINE




- 10 By essentially the same procedure as set forth in Example 180, but using 4-(8-chloro-3-methyl-5,6-dihydro-11-(4-piperidylidene)-11H-benzo[5,6]cyclohepta[1,2-b]pyridine (from Preparative Example 3E) instead of 4-(3,8-dichloro-5,6-dihydro-11-(4-piperidylidene)-11H-benzo[5,6]cyclohepta[1,2-b]pyridine, and 3-pyridylacetic acid instead of 4-pyridylacetic acid, the title compound was obtained as white solid MH⁺ 444.
- 15

EXAMPLE 307


- 20 By essentially the same procedure as in Example 1, using the title compound from Preparative Example 37B, and 4-pyridylacetic acid the compound of Example 307, identified in Table 8, was obtained, MH⁺ 428.

EXAMPLE 309

- 25 1-1-(4-PYRIDINYLACETYL)-4-[2-METHYL-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-PIPERIDINE



10 1-1-(4-PYRIDINYLACETYL)-4-[8.9 DICHLORO-5.6-DIHYDRO-11H-
BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11-YLIDENE] - PIPERIDINE




By essentially the same procedure as set forth in Example 180, but using 4-(8,9-dichloro-5,6-dihydro-11-(4-piperidylidene)-11H-benzo[5,6]-cyclohepta[1,2-b]pyridine (from Preparative Example 1H) instead of 4-(3,8-dichloro-5,6-dihydro-11-(4-piperidylidene)-11H-benzo[5,6]cyclohepta[1,2-b]pyridine, and 3-pyridylacetic acid instead of 4-pyridylacetic acid, the title compound was obtained as white solid MH⁺ 464.

By essentially the same procedure as in Example 182, with the exception that 8-chloro-6,11-dihydro-11-(4-piperidiny)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine is used instead of 8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine, the compound of Example 312 was obtained as a white solid (MH⁺ 432). The structure for this compound is given in Table 10.

• EXAMPLE 350


8-CHLORO-11H-BENZO[5.5]CYCLOHEPTA[1.2-b]PYRIDIN-11-
YLIDENE)-4-(3-PYRIDINYLACETYL)PIPERAZINE



- 5 By substituting in Example 75, 0.4g (1.28mmoles) of 8-chloro-11-(1-piperazinyl)-11H-benzo[5,6]cyclohepta[1,2-b]pyridine (Preparative Example 38) for 8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine and 0.1765g (1.28mmoles) of 3-pyridylacetic acid for 4-pyridylacetic acid and using the same method as described in
- 10 Example 75, one obtains the title compound (0.513g, 93%, MH^+ 431).

EXAMPLE 351


1-(3-BROMO-8-CHLORO-6,11-DIHYDRO-5H-BENZO[5.6]CYCLO-
HEPTA[1.2-b]PYRIDIN-11-YL)-4-(3-PYRIDINYLACETYL)PIPERAZINE




- By substituting in Example 75, 3-bromo-8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (0.32g, 0.81mmoles) (Preparative Example 40) for 8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine and 3-pyridylacetic acid (0.117g, 0.86mmoles) for 4-pyridylacetic acid and using the method described in Example 50, one obtains the title compound (0.3942g, 95%, MH^+ 511).

EXAMPLES 352-353

- By essentially the same procedures as set forth in Example 351, but using




in place of 3-pyridylacetic acid, one obtains compounds of the formulas



- respectively. The compound of Example 352 is a white amorphous solid, yield 65%, Mass Spec MH^+ 555. The compound of Example 353 is a white amorphous solid, yield 59%, Mass Spec MH^+ 539.

EXAMPLE 354

4-(3-BROMO 8-CHLORO-6,11-DIHYDRO-5H-BENZO[5.6]CYCLOHEPTA[1.2-b]PYRIDIN-11-YL)-N-(3-PYRIDINYL)-1-PIPERAZINE-CARBOXAMIDE



5


The title compound from Preparative Example 40 (0.37g, 0.94mmoles) was reacted with 3-ethoxycarbonylaminopyridine (Preparative Example 12) (0.123g, 0.94mmoles) under essentially the same conditions as described in Example 183, to give the title compound

10

(0.3164g, 66%, MH^+ 512).

EXAMPLE 355

1-(4,8-DICHLORO-6,11-DIHYDRO-5H-BENZO[5.6]CYCLOHEPTA-[1.2-b]PYRIDIN-11-YL)-4-(3-PYRIDYLACETYL) PIPERAZINE




15

By substituting in Example 75, 4,8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine (0.3g, 0.86mmoles) (Preparative Example 41) for 8-chloro-11-(1-piperazinyl)-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine and 3-pyridylacetic acid (0.1181g, 0.86mmoles) for 4-pyridylacetic acid and using the method described in Example 50, one obtains the title compound (0.357g, 88%, MH^+ 467).

EXAMPLE 356


10 4-[3-BROMO-4,8-DICHLORO-5,6-DIHYDRO-11H-BENZO[5,6]-
CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-1-(4-PYRIDINYLACETYL)-
PIPERIDINE



A mixture of the title compound from Preparative Example 46 (0.68 grams), 4-pyridylacetic acid hydrochloride (0.60 grams), 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (0.65 grams), 4-methylmorpholine (0.6 mL) and anhydrous dimethylformamide (20 mL) was stirred at 25°C for 48 hours. Concentration *in vacuo* provided a residue which was diluted with dichloromethane and washed with 1 *N* aqueous sodium hydroxide and brine. The organic phase was dried over anhydrous magnesium sulfate and concentrated *in vacuo* to provide a residue which was purified by flash column chromatography (silica gel) using 2-5% methanol-dichloromethane saturated with ammonium hydroxide to afford the title compound (0.06 grams, 7%, MH^+ 544).


EXAMPLE 358

A. 4-(8-CHLORO-3-NITRO-5,6-DIHYDRO-11-(4-PIPERIDYLIDENE)-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDINE.



- 5 Hydrolyze the title compound of Preparative Example 47A (10.0g, mmol) by dissolving in conc. HCl (250mL) and heating to 100°C for 16h. The cooled acidic mixture was neutralized with 1M NaOH (950 mL). The mixture was extracted with methylene chloride. The latter was dried over magnesium sulfate. Filtration and concentration afforded the title
- 10 compound in 99% yield as a solid. MH+ 358.

B. 1-1-(4-PYRIDINYLACETYL)-4-[3-BROMO-8-CHLORO-5,6-DIHYDRO-11H-BENZO[5,6]CYCLOHEPTA[1,2-b]PYRIDIN-11-YLIDENE]-PIPERIDINE



- 15 By essentially the same procedure as set forth in Example 180, but using 4-(8-chloro-3-nitro-5,6-dihydro-11-(4-piperidylidene)-11H-benzo[5,6]cyclohepta[1,2-b]pyridine instead of 4-(3,8-dichloro-5,6-dihydro-11-(4-piperidylidene)-11H-benzo[5,6]cyclohepta[1,2-b]pyridine, the title compound was obtained as an off-white solid. Mp = 111.3-
- 20 112.2°C, MH+ 475.

ASSAYS

1. In vitro enzyme assays: Inhibition of farnesyl protein transferase and geranylgeranyl protein transferase.

Both farnesyl protein transferase (FPT) and geranylgeranyl protein transferase (GGPT) I were partially purified from rat brain by ammonium sulfate fractionation followed by Q-Sepharose (Pharmacia, Inc.) anion exchange chromatography essentially as described by Yokoyama et al (Yokoyama, K., et al., (1991), A protein geranylgeranyltransferase from bovine brain: Implications for protein prenylation specificity, **Proc. Natl. Acad. Sci USA** **88**: 5302-5306, the disclosure of which is incorporated herein by reference thereto). Human farnesyl protein transferase was also expressed in *E. coli*, using cDNA clones encoding both the α and β subunits. The methods used were similar to those published (Omer, C. et al., (1993), Characterization of recombinant human farnesyl protein transferase: Cloning, expression, farnesyl diphosphate binding, and functional homology with yeast prenyl-protein transferases, *Biochemistry* **32**:5167-5176). Human farnesyl protein transferase was partially-purified from the soluble protein fraction of *E. coli* as described above. The tricyclic farnesyl protein transferase inhibitors disclosed herein inhibited both human and rat enzyme with similar potencies. Two forms of val¹²-Ha-Ras protein were prepared as substrates for these enzymes, differing in their carboxy terminal sequence. One form terminated in cysteine-valine-leucine-serine (Ras-CVLS) the other in cysteine-valine-leucine-leucine (Ras-CVLL). Ras-CVLS is a substrate for the farnesyl protein transferase while Ras-CVLL is a substrate for geranylgeranyl protein transferase I. The cDNAs encoding these proteins were constructed so that the proteins contain an amino-terminal extension of 6 histidine residues. Both proteins were expressed in Escherichia coli and purified using metal chelate affinity chromatography. The radiolabelled isoprenyl pyrophosphate substrates, [³H]farnesyl pyrophosphate and [³H]geranylgeranyl pyrophosphate, were purchased from DuPont/New England Nuclear.

Several methods for measuring farnesyl protein transferase activity have been described (Reiss et al 1990, *Cell* **62**: 81; Schaber et al 1990, *J. Biol. Chem.* **265**: 14701; Manne et al 1990, *PNAS* **87**: 7541; and Barbacid & Manne 1993, U.S. Patent No. 5,185,248). The activity was assayed by measuring the transfer of [³H]farnesyl from [³H]farnesyl pyrophosphate to Ras-CVLS using conditions similar to those described

by Reiss et al. 1990 (*Cell* **62**: 81) The reaction mixture contained 40 mM Hepes, pH 7.5; 20 mM magnesium chloride; 5 mM dithiothreitol; 0.25 μ M [3 H]farnesyl pyrophosphate; 10 μ l Q-Sepharose-purified farnesyl protein transferase; the indicated concentration of tricyclic compound or dimethylsulfoxide (DMSO) vehicle control (5% DMSO final); and 5 μ M Ras-CVLS in a total volume of 100 μ l. The reaction was allowed to proceed for 30 minutes at room temperature and then stopped with 0.5 ml of 4% sodium dodecyl sulfate (SDS) followed by 0.5 ml of cold 30% trichloroacetic acid (TCA). Samples were allowed to sit on ice for 45 minutes and precipitated Ras protein was then collected on GF/C filter paper mats using a Brandel cell harvester. Filter mats were washed once with 6% TCA, 2% SDS and radioactivity was measured in a Wallac 1204 Betaplate BS liquid scintillation counter. Percent inhibition was calculated relative to the DMSO vehicle control.

The geranylgeranyl protein transferase I assay was essentially identical to the farnesyl protein transferase assay described above, with two exceptions: [3 H]geranylgeranylpyrophosphate replaced farnesyl pyrophosphate as the isoprenoid donor and Ras-CVLL was the protein acceptor. This is similar to the assay reported by Casey et al (Casey, P.J., et al., (1991), Enzymatic modification of proteins with a geranylgeranyl isoprenoid, *Proc. Natl. Acad. Sci, USA* **88**: 8631-8635, the disclosure of which is incorporated herein by reference thereto).

2. Cell-Based Assay: Transient expression of val¹²-Ha-Ras-CVLS and val¹²-Ha-Ras-CVLL in COS monkey kidney cells: Effect of farnesyl protein transferase inhibitors on Ras processing and on disordered cell growth induced by transforming Ras.

COS monkey kidney cells were transfected by electroporation with the plasmid pSV-SPORT (Gibco/BRL) containing a cDNA insert encoding either Ras-CVLS or Ras-CVLL, leading to transient overexpression of a Ras substrate for either farnesyl protein transferase or geranylgeranyl protein transferase I, respectively (see above).

Following electroporation, cells were plated into 6-well tissue culture dishes containing 1.5 ml of Dulbecco's-modified Eagle's media (GIBCO, Inc.) supplemented with 10% fetal calf serum and the appropriate farnesyl protein transferase inhibitors. After 24 hours, media was removed and fresh media containing the appropriate drugs was re-added.

48 hours after electroporation cells were examined under the microscope to monitor disordered cell growth induced by transforming

Ras. Cells expressing transforming Ras become more rounded and refractile and overgrow the monolayer, reminiscent of the transformed phenotype. Cells were then photographed, washed twice with 1 ml of cold phosphate-buffered saline (PBS) and removed from the dish by
5 scraping with a rubber policeman into 1 ml of a buffer containing 25 mM Tris, pH 8.0; 1 mM ethylenediamine tetraacetic acid; 1 mM phenylmethylsulfonyl fluoride; 50 μ M leupeptin; and 0.1 μ M pepstatin. Cells were lysed by homogenization and cell debris was removed by centrifugation at 2000 x g for 10 min.

10 Cellular protein was precipitated by addition of ice-cold trichloroacetic acid and redissolved in 100 μ l of SDS-electrophoresis sample buffer. Samples (5-10 μ l) were loaded onto 14% polyacrylamide minigels (Novex, Inc.) and electrophoresed until the tracking dye neared the bottom of the gel. Proteins resolved on the gels were electroblotted
15 onto nitrocellulose membranes for immunodetection.

Membranes were blocked by incubation overnight at 4°C in PBS containing 2.5% dried milk and 0.5% Tween-20 and then incubated with a Ras-specific monoclonal antibody, Y13-259 (Furth, M.E., et al., (1982), Monoclonal antibodies to the p21 products of the transforming gene of
20 Harvey murine sarcoma virus and of the cellular ras gene family, J. Virol. 43: 294-304), in PBS containing 1% fetal calf serum for one hour at room temperature. After washing, membranes were incubated for one hour at room temperature with a 1:5000 dilution of secondary antibody, rabbit anti-rat IgG conjugated to horseradish peroxidase, in PBS
25 containing 1% fetal calf serum. The presence of processed and unprocessed Ras-CVLS or Ras-CVLL was detected using a colorimetric peroxidase reagent (4-chloro-1-naphthol) as described by the manufacturer (Bio-Rad).

3. Cell Mat Assay:

30 Normal human HEPM fibroblasts were planted in 3.5 cm dishes at a density of 5×10^4 cells/dish in 2 ml growth medium, and incubated for 3-5d to achieve confluence. Medium was aspirated from each dish and the indicator tumor cells, T24-BAG4 human bladder carcinoma cells expressing an activated H-ras gene, were planted on top of the fibroblast
35 monolayer at a density of 2×10^3 cells/dish in 2 ml growth medium, and allowed to attach overnight. Compound-induced colony inhibition was assayed by addition of serial dilutions of compound directly to the growth medium 24 h after tumor cell planting, and incubating cells for an

additional 14 d to allow colony formation. Assays were terminated by rinsing monolayers twice with phosphate-buffered saline (PBS), fixing the monolayers with a 1% glutaraldehyde solution in PBS, then visualizing tumor cells by staining with X-Gal (Price, J., et al., Lineage analysis in the vertebrate nervous system by retrovirus-mediated gene transfer, Proc. Natl. Acad. Sci. 84, 156-160(1987)). In the colony inhibition assay, compounds were evaluated on the basis of two IC_{50} values: the concentration of drug required to prevent the increase in tumor cell number by 50% (tIC_{50}) and the concentration of drug required to reduce the density of cells comprising the cell mat by 50% (mIC_{50}). Both IC_{50} values were obtained by determining the density of tumor cells and mat cells by visual inspection and enumeration of cells per colony and the number of colonies under the microscope. The therapeutic index of the compound was quantitatively expressed as the ratio of mIC_{50}/tIC_{50} , with values greater than one indicative of tumor target specificity.

RESULTS OF ASSAYS - TABLES 7 TO 19

The compounds listed in Table 7 refer to compounds of Formula 500.00:




TABLE 7











EXAMPLE	R	FPT IC ₅₀ (μM)
1		0.01-10
2		0.01-10
3		0.01-10
88		0.01-10
4		0.01-10
53		0.01-10
5		0.01-10
89		0.01-10
6		0.01-10
7		0.01-10

TABLE 7 - continued












EXAMPLE	R	FPT IC ₅₀ (μM)
8		0.01-10
90		0.01-10
91		0.01-10
92		0.01-10
93		0.01-10
94		0.01-10
9		0.01-10
95		0.01-10
10		0.01-10
11		0.01-10
12		0.01-10

TABLE 7 - continued












EXAMPLE	R	FPT IC ₅₀ (μM)
96		0.01-10
97		0.01-10
98		0.01-10
99		0.01-10
13		0.01-10
100		0.01-10
101		0.01-10
102		0.01-10
14		0.01-10
103		0.01-10
104		0.01-10

TABLE 7 - continued












EXAMPLE	R	FPT IC ₅₀ (μM)
15		0.01-10
16		0.01-10
17		0.01-10
52		0.01-10
18		0.01-10
105		0.01-10
19A		0.01-10
20		0.01-10
54		0.01-10
106		0.01-10
21		0.01-10

TABLE 7 - continued











EXAMPLE	R	FPT IC ₅₀ (μM)
55		0.01-10
22		0.01-10
23		0.01-10
24		0.01-10
25		0.01-10
26		0.01-10
108		0.01-10
109		0.01-10
27		0.01-10
28		0.01-10

TABLE 7 - continued











EXAMPLE	R	FPT IC ₅₀ (μM)
56		0.01-10
29		0.01-10
30		0.01-10
31		0.01-10
110		0.01-10
111		0.01-10
32		0.01-10
33		0.01-10
112		0.01-10
34		0.01-10

TABLE 7 - continued











EXAMPLE	R	FPT IC ₅₀ (μM)
57		0.01-10
35		0.01-10
113		10-100
114		10-100
36		10-100
37		10-100
38		10-100
39		10-100
115		10-100
58		10-100

TABLE 7 - continued











EXAMPLE	R	FPT IC ₅₀ (μM)
116		10-100
117		10-100
118		10-100
59		10-100
119		10-100
120		10-100
121		0.01-10
40		10-100
122		10-100
123		10-100

TABLE 7 - continued











EXAMPLE	R	FPT IC ₅₀ (μM)
124		10-100
125		10-100
126		0.01-10
127		0.01-10
128		0.01-10
129		10-100
41	 (5.4)	0.01-10
42	 (5.7)	0.01-10
43	 (5.8)	0.01-10
44	 (5.9)	0.01-10

TABLE 7 - continued










EXAMPLE	R	FPT IC ₅₀ (μM)
45	 (5.11)	0.01-10
46	 (5.12)	0.01-10
47	 (5.13)	0.01-10
48	 (5.14)	0.01-10
49	 (5.15)	0.01-10
84	 (5.5)	0.01-10
83	 (5.6)	0.01-10
19	 (5.10)	0.01-10
85	 (5.16)	0.01-10

TABLE 7 - continued








EXAMPLE	R	FPT IC ₅₀ (μM)
219	 (5.92)	0.01-10
220	 (5.93)	0.01-10
221	 (5.94)	0.01-10
222	 (5.95)	0.01-10
227	 (5.102)	0.01-10
228	 (5.103)	0.01-10
287	 (5.104)	0.01-10

TABLE 7 - continued














EXAMPLE	R	FPT IC ₅₀ (μM)
288		0.01-10
289		0.01-10
290		0.01-10
291		0.01-10
292		0.01-10
293		0.01-10
294		0.01-10
295		0.01-10

TABLE 7 - continued

EXAMPLE	R	FPT IC ₅₀ (μM)
296		0.01-10
297		0.01-10
298		0.01-10
299		0.01-10
300		0.01-10

The compounds listed in Table 8 refer to compounds of Formula 505.00:




TABLE 8









EXAMPLE	A	B	FPT IC ₅₀ (μM)
74B		CH ₃	0.01-10
74C		CH ₃	10-100
74A		CH ₃	10-100
130		CH ₃	10-100
131			10-100
73		CH ₃	10-100
132		CH ₃	10-100

TABLE 8 - continued










EXAMPLE	A	B	FPT IC ₅₀ (μ M)
133		CH ₃	10-100
134			>100
135		CH ₃	0.01-10
62		CH ₃	0.01-10
64		CH ₃	10-100
136		CH ₃	10-100
66		CH ₃	10-100
137		CH ₃	10-100

TABLE 8 - continued












EXAMPLE	A	B	FPT IC ₅₀ (μM)
63		CH ₃	100
139		CH ₃	10-100
140		CH ₃	10-100
71A		CH ₃	10-100
141		CH ₃	10-100
143		CH ₃	10-100
71B		CH ₃	10-100
144		CH ₃	10-100
144a			10-100
71C		CH ₃	10-100

TABLE 8 - continued

















EXAMPLE	A	B	FPT IC ₅₀ (μM)
145			>100
146			10-100
147		CH ₃	>100
301			0.01-10
180			0.01-10
303			0.01-10
304			0.01-10
305			10-100
230			0.01-10

TABLE 8 - continued

















EXAMPLE	A	B	FPT IC ₅₀ (μM)
307			0.01-10
235			0.01-10
309			0.01-10
310			0.01-10
311			0.01-10
323			0.01-10
5.39			0.01-10
358			0.01-10

TABLE 8 - continued













EXAMPLE	A	B	FPT IC ₅₀ (μ M)
360			0.01-10
361			0.01-10
362			0.01-10
365			0.01-10
366			0.01-10
367			0.01-10

Table 9 lists FPT IC₅₀ results for additional compounds.

TABLE 9

EXAMPLE	FPT IC ₅₀ (μM)	EXAMPLE	FPT IC ₅₀ (μM)	EXAMPLE	FPT IC ₅₀ (μM)
229 (5.104)	0.01-10	231 (5.106)	10-100	232 (5.107)	0.01-10
236 (5.111)	0.01-10	237 (5.112)	10-100	238 (5.113)	10-100
239 (5.114)	0.01-10	240 (5.115)	0.01-10	246 (5.121)	0.01-10
247 (5.122)	0.01-10	248 (5.124)	0.01-10	248 (5.123)	0.01-10
249 (5.125)	0.01-10	250 (5.126)	0.01-10	256 (5.132)	0.01-10
257 (5.133)	0.01-10	258 (5.134)	10-100	259 (5.135)	0.01-10
260 (5.136)	0.01-10	266 (5.138)	0.01-10	269 (5.140)	0.01-10
276 (5.145)	0.01-10	279 (5.147)	0.01-10	280 (5.148)	10-100
281 (5.149)	0.01-10	282 (5.150)	0.01-10	283 (5.151)	0.01-10
284 (5.152)	0.01-10	285 (5.153)	0.01-10	286 (5.154)	0.01-10

The compounds listed in Table 10 refer to compounds of Formula 510.00:




TABLE 10








EXAMPLE	R	FPT IC ₅₀ (μM)
149		10-100
150		10-100
75	 (5.17)	0.01-10
76	 (5.18)	0.01-10
77	 (5.19)	0.01-10
78	 (5.20)	0.01-10
79	 (5.21)	0.01-10

TABLE 10 - continued








EXAMPLE	R	FPT IC ₅₀ (μM)
80	 (5.22)	0.01-10
81	 (5.23)	0.01-10
82	 (5.25)	0.01-10
312		0.01-10
313		0.01-10
314		0.01-10
234		0.01-10

TABLE 10 - continued







EXAMPLE	R	FPT IC ₅₀ (μM)
316		0.01-10
317		10-100
318		0.01-10
182		0.01-10
183		0.01-10
321		0.01-10

TABLE 10 - continued


EXAMPLE	R	FPT IC ₅₀ (μM)
368		>100

Table 11 lists FPT IC₅₀ results for additional compounds.

5

TABLE 11

EXAMPLE	FPT IC ₅₀ (μM)	EXAMPLE	FPT IC ₅₀ (μM)	EXAMPLE	FPT IC ₅₀ (μM)
187 (6.7)	0.01-10	187 (6.8)	0.01-10	188 (6.9)	0.01-10
189 (5.62)	0.01-10	190 (5.63)	0.01-10	191 (5.64)	0.01-10
192 (5.65)	0.01-10	194 (5.67)	0.01-10	195 (5.68)	0.01-10
196 (5.69)	0.01-10	197 (5.70)	0.01-10	198 (5.71)	0.01-10
199 (5.72A)	10-100	199 (5.72B)	10-100	200 (5.73)	0.01-10
201 (5.74)	0.01-10	202 (5.75)	10-100	203 (5.76)	10-100
205 (5.78)	0.01-10	206 (5.79)	0.01-10	207 (5.80)	10-100
208 (5.81)	0.01-10	209 (5.82)	0.01-10	210 (5.83)	0.01-10
211 (5.84)	0.01-10	212 (5.85)	0.01-10	213 (5.86)	10-100
214 (5.87)	>100	215 (5.88)	10-100	216 (5.89)	10-100

TABLE 11 - continued

EXAMPLE	FPT IC ₅₀ (μ M)	EXAMPLE	FPT IC ₅₀ (μ M)	EXAMPLE	FPT IC ₅₀ (μ M)
217 (5.90)	0.01-10	218 (5.91)	0.01-10	233 (5.108)	0.01-10
251 (5.127)	0.01-10	261 (6.12)	0.01-10	351	0.01-10
352	0.01-10	353	0.01-10	354	0.01-10
355	0.01-10	273 (5.143)	0.01-10	267 (6.17)	0.01-10
356	0.01-10	264 (6.15)	>100	262 (6.13)	0.01-10
263 (6.14)	0.01-10	253 (5.129)	0.01-10	350	0.01-10
252 (5.128)	0.01-10	182 (6.4)	0.01-10	268 (5.139)	0.01-10
277 (6.20)	0.01-10	----	----	----	----

The compounds listed in Table 12 refer to compounds of Formula 525.00:




TABLE 12
















EXAMPLE	A	B	FPT IC ₅₀ (μ M)
184 (5.60)			0.01-10
185 (5.61)			0.01-10
223 (5.96)			0.01-10
223 (5.97)			0.01-10
224 (6.10)			0.01-10
224 (6.11)			0.01-10
225 (5.98)			0.01-10

TABLE 12 - continued

EXAMPLE	A	B	FPT IC ₅₀ (μM)
225 (5.99)			0.01-10
226 (5.100)			0.01-10
226 (5.101)			0.01-10
351			0.01-10
354			0.01-10

The compounds listed in Table 13 refer to compounds of Formula 515.00:



circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

The amount and frequency of administration of the compounds of the invention and the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended dosage regimen is oral administration of from 10 mg to 2000 mg/day preferably 10 to 1000 mg/day, in two to four divided doses to block tumor growth. The compounds are non-toxic when administered within this dosage range.

The following are examples of pharmaceutical dosage forms which contain a compound of the invention. The scope of the invention in its pharmaceutical composition aspect is not to be limited by the examples provided.

Pharmaceutical Dosage Form Examples

EXAMPLE A

Tablets

No.	Ingredients	mg/tablet	mg/tablet
1.	Active compound	100	500
2.	Lactose USP	122	113
3.	Corn Starch, Food Grade, as a 10% paste in Purified Water	30	40
4.	Corn Starch, Food Grade	45	40
5.	Magnesium Stearate	<u>3</u>	<u>7</u>
Total		300	700

Method of Manufacture

Mix Item Nos. 1 and 2 in a suitable mixer for 10–15 minutes. Granulate the mixture with Item No. 3. Mill the damp granules through a coarse screen (e.g., 1/4", 0.63 cm) if necessary. Dry the damp granules. Screen the dried granules if necessary and mix with Item No. 4 and mix for 10–15 minutes. Add Item No. 5 and mix for 1–3 minutes. Compress the mixture to appropriate size and weigh on a suitable tablet machine.

EXAMPLE B

Capsules


No.	Ingredient	mg/capsule	mg/capsule
1.	Active compound	100	500
2.	Lactose USP	106	123
3.	Corn Starch, Food Grade	40	70
4.	Magnesium Stearate NF	<u>7</u>	<u>7</u>
Total		253	700

5 Method of Manufacture

Mix Item Nos. 1, 2 and 3 in a suitable blender for 10-15 minutes. Add Item No. 4 and mix for 1-3 minutes. Fill the mixture into suitable two-piece hard gelatin capsules on a suitable encapsulating machine.


10 While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

15

WHAT IS CLAIMED IS:

1. A method for inhibiting the abnormal growth of cells comprising administering an effective amount of a compound of Formula

5 1.0:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

one of a, b, c and d represents N or NR⁹ wherein R⁹ is O⁻, -CH₃ or -(CH₂)_nCO₂H wherein n is 1 to 3, and the remaining a, b, c and d groups represent CR¹ or CR²; or

10 each of a, b, c, and d are independently selected from CR¹ or CR²;
 each R¹ and each R² is independently selected from H, halo, -CF₃, -OR¹⁰, -COR¹⁰, -SR¹⁰, -S(O)_tR¹¹ (wherein t is 0, 1 or 2), -N(R¹⁰)₂, -NO₂, -OC(O)R¹⁰, -CO₂R¹⁰, -OCO₂R¹¹, -CN, -NR¹⁰COOR¹¹, -SR¹¹C(O)OR¹¹,
 15 -SR¹¹N(R⁷⁵)₂ (wherein each R⁷⁵ is independently selected from H and -C(O)OR¹¹), benzotriazol-1-yloxy, tetrazol-5-ylthio, or substituted tetrazol-5-ylthio, alkynyl, alkenyl or alkyl, said alkyl or alkenyl group optionally being substituted with halo, -OR¹⁰ or -CO₂R¹⁰;

20 R³ and R⁴ are the same or different and each independently represents H, any of the substituents of R¹ and R², or R³ and R⁴ taken together represent a saturated or unsaturated C₅-C₇ fused ring to the benzene ring;

25 R⁵, R⁶, R⁷ and R⁸ each independently represents H, -CF₃, -COR¹⁰, alkyl or aryl, said alkyl or aryl optionally being substituted with -OR¹⁰, -SR¹⁰, -S(O)_tR¹¹, -NR¹⁰COOR¹¹, -N(R¹⁰)₂, -NO₂, -COR¹⁰, -OCOR¹⁰, -OCO₂R¹¹, -CO₂R¹⁰, OPO₃R¹⁰ or one of R⁵, R⁶, R⁷ and R⁸ can be taken



in combination with R^{40} as defined below to represent $-(CH_2)_r$ wherein r is 1 to 4 which can be substituted with lower alkyl, lower alkoxy, $-CF_3$ or aryl, or R^5 is combined with R^6 to represent $=O$ or $=S$ and/or R^7 is combined with R^8 to represent $=O$ or $=S$;

5 R^{10} represents H, alkyl, aryl, or aralkyl;


R^{11} represents alkyl or aryl;

X represents N, CH or C, which C may contain an optional double bond, represented by the dotted line, to carbon atom 11;

the dotted line between carbon atoms 5 and 6 represents an
10 optional double bond, such that when a double bond is present, A and B independently represent $-R^{10}$, halo, $-OR^{11}$, $-OCO_2R^{11}$ or $-OC(O)R^{10}$, and when no double bond is present between carbon atoms 5 and 6, A and B each independently represent H_2 , $-(OR^{11})_2$, H and halo, dihalo, alkyl and H, $(alkyl)_2$, $-H$ and $-OC(O)R^{10}$, H and $-OR^{10}$, $=O$, aryl and H, $=NOR^{10}$ or
15 $-O-(CH_2)_p-O-$ wherein p is 2, 3 or 4;

R represents R^{40} , R^{42} , R^{44} , or R^{54} , as defined below;

R^{40} represents H, aryl, alkyl, cycloalkyl, alkenyl, alkynyl or -D wherein -D represents







20 wherein R^3 and R^4 are as previously defined and W is O, S or NR^{10} wherein R^{10} is as defined above; said R^{40} cycloalkyl, alkenyl and alkynyl groups being optionally substituted with from 1-3 groups selected from halo, $-CON(R^{10})_2$, aryl, $-CO_2R^{10}$, $-OR^{12}$, $-SR^{12}$, $-N(R^{10})_2$, $-N(R^{10})CO_2R^{11}$, $-COR^{12}$, $-NO_2$ or D, wherein -D, R^{10} and R^{11} are as
25 defined above and R^{12} represents R^{10} , $-(CH_2)_mOR^{10}$ or $-(CH_2)_qCO_2R^{10}$ wherein R^{10} is as previously defined, m is 1 to 4 and q is 0 to 4; said alkenyl and alkynyl R^{40} groups not containing $-OH$, $-SH$ or $-N(R^{10})_2$ on a carbon containing a double or triple bond respectively; or
30 R^{40} represents phenyl substituted with a group selected from $-SO_2NH_2$, $-NH SO_2CH_3$, $-SO_2NHCH_3$, $-SO_2CH_3$, $-SOCH_3$, $-SCH_3$, or






-NHSO₂CF₃, preferably, said group is located in the para position of the phenyl ring; or

R⁴⁰ represents a group selected from



R⁴² represents



wherein R²⁰, R²¹ and R⁴⁶ are each independently selected from the group consisting of:

- 15
- (1) H;
 - (2) -(CH₂)_qSC(O)CH₃ wherein q is 1 to 3;
 - (3) -(CH₂)_qOSO₂CH₃ wherein q is 1 to 3;
 - (4) -OH;
 - (5) -CS(CH₂)_w(substituted phenyl) wherein w is 1 to 3 and the
 - 20 substituents on said substituted phenyl group are the same substituents as described below for said substituted phenyl;
 - (6) -NH₂;
 - (7) -NHCBZ;





- 240 -

(8) -NHC(O)OR^{22} wherein R^{22} is an alkyl group having from 1 to 5 carbon atoms, or R^{22} represents phenyl substituted with 1 to 3 alkyl groups;

(9) alkyl;

5 (10) $\text{-(CH}_2\text{)}_k\text{phenyl}$ wherein k is 1 to 6;

(11) phenyl;

(12) substituted phenyl wherein the substituents are selected from the group consisting of: halo, NO_2 , -OH , -OCH_3 , -NH_2 , -NHR^{22} , $\text{-N(R}^{22}\text{)}_2$, alkyl, $\text{-O(CH}_2\text{)}_t\text{phenyl}$ (wherein t is from 1 to 3), and $\text{-O(CH}_2\text{)}_t\text{substituted}$
10 phenyl (wherein t is from 1 to 3);

(13) naphthyl;

(14) substituted naphthyl, wherein the substituents are as defined for substituted phenyl above;

(15) bridged polycyclic hydrocarbons having from 5 to 10 carbon
15 atoms;


(16) cycloalkyl having from 5 to 7 carbon atoms;

(17) heteroaryl;


(18) hydroxyalkyl;

(19) substituted pyridyl or substituted pyridyl N-oxide wherein the
20 substituents are selected from the substituents given above for said substituted phenyl, and said substituents are bound to a ring carbon by replacement of the hydrogen bound to said carbon;


(20)



(21)




(22)



25 (23) $\text{-NHC(O)-(CH}_2\text{)}_k\text{phenyl}$ or $\text{-NH(O)-(CH}_2\text{)}_k\text{-substituted phenyl}$, wherein said k is as defined above;

(24) piperidine Ring V:



wherein R^{50} represents H, alkyl, alkylcarbonyl, alkylloxycarbonyl,
30 haloalkyl, or $\text{-C(O)NH(R}^{10}\text{)}$ wherein R^{10} is H or alkyl;

(25) $\text{-NHC(O)CH}_2\text{C}_6\text{H}_5$ or $\text{-NHC(O)CH}_2\text{-substituted-C}_6\text{H}_5$;

THIS SHEET HAS BEEN
REPLACED







- 241 -

(26) $-\text{NHC(O)OC}_6\text{H}_5$;

(27)

(28)

(29)


(30) $-\text{OC(O)}-\text{heteroaryl}$, for example

5

(31) $-\text{O-alkyl}$ (e.g., $-\text{OCH}_3$); and(32) $-\text{CF}_3$; or

R^{20} and R^{21} taken together form a $=\text{O}$ group and the remaining R^{46} is as defined above; or

10 Two of R^{20} , R^{21} , and R^{46} taken together form piperidine Ring V




wherein R^{50} is as defined above;

with the proviso that R^{46} , R^{20} , and R^{21} are selected such that the carbon atom to which they are bound does not contain more than one

15 heteroatom;




R^{44} represents




wherein R^{25} represents heteroaryl or aryl; and R^{48} represents H or alkyl;

R^{54} represents an N-oxide heterocyclic group of the formula (i), (ii),

20 (iii) or (iv):



or




THIS SHEET HAS BEEN
REPLACED



wherein R^{56} , R^{58} , and R^{60} are the same or different and each is independently selected from H, halo, $-CF_3$, $-OR^{10}$, $-C(O)R^{10}$, $-SR^{10}$, $-S(O)_eR^{11}$ (wherein e is 1 or 2), $-N(R^{10})_2$, $-NO_2$, $-CO_2R^{10}$, $-OCO_2R^{11}$, $-OCOR^{10}$, alkyl, aryl, alkenyl or alkynyl, which alkyl may be substituted with $-OR^{10}$, $-SR^{10}$ or $-N(R^{10})_2$ and which alkenyl may be substituted with OR^{11} or SR^{11} ; or

R^{54} represents an N-oxide heterocyclic group of the formula (ia), (iia), (iiia) or (iva):



wherein Y represents N^+-O^- and E represents N; or

R^{54} represents an alkyl group substituted with one of said N-oxide heterocyclic groups (i), (ii), (iii), (iv), (ia), (iia), (iiia) or (iva);

Z represents O or S such that R can be taken in combination with R^5 , R^6 , R^7 or R^8 as defined above, or R represents R^{40} , R^{42} , R^{44} or R^{54} .

15

2. The method of Claim 1 wherein a is N and b, c, and d are carbon; R^1 and R^2 are the same or different and each is independently selected from H, halo, $-CF_3$, lower alkyl, or benzotriazol-1-yloxy, and R^1 is at the C-4 position and R^2 is at the C-3 position; R^3 and R^4 are the same or different and each is independently selected from H or halo, and R^3 is at the C-8 position and R^4 is at the C-9 position; when the double bond between carbon atoms 5 and 6 is present, A and B independently represent H, lower alkyl or alkyloxy; and when the double bond between carbon atoms 5 and 6 is absent, A and B independently represent H_2 , $(-H$ and $-OH)$ or $=O$; R^5 , R^6 , R^7 , and R^8 are H; Z is O; and R represents R^{42} and the R^{46} is selected from phenyl, substituted phenyl, heteroaryl or piperidine Ring V.

3. The method of Claim 2 wherein R^{20} and R^{21} are each independently selected from H and alkyl; R^3 is Cl; R^4 is H; R^1 and R^2 are individually selected from H, benzotriazol-1-yloxy, C_1 to C_4 alkyl or halo; and R^{46} represents 3-pyridyl, 3-pyridyl N-oxide, 4-pyridyl, 4-pyridyl N-oxide, 3-N-methylpiperidinyl, 4-N-methylpiperidinyl, 3-N-acetylpiperidinyl or 4-N-acetylpiperidinyl.





4. The method of Claim 3 wherein both R²⁰ and R²¹ are H, or both R²⁰ and R²¹ are methyl; R¹ and R² are individually selected from H, Br, Cl, methyl or benzotriazol-1-yloxy; and R⁴⁶ represents 3-pyridyl, 3-pyridyl N-oxide, 4-pyridyl, 4-pyridyl N-oxide, 3-N-methylpiperidiny or 4-N-methylpiperidiny.

5. The method of Claim 1 wherein a is N and b, c, and d are carbon; R¹ and R² are the same or different and each is independently selected from H, halo, -CF₃, lower alkyl, or benzotriazol-1-yloxy, and R¹ is at the C-4 position and R² is at the C-3 position; R³ and R⁴ are the same or different and each is independently selected from H or halo, and R³ is at the C-8 position and R⁴ is at the C-9 position; when the double bond between carbon atoms 5 and 6 is present, A and B independently represent H, lower alkyl or alkyloxy; and when the double bond between carbon atoms 5 and 6 is absent, A and B independently represent H₂, (-H and -OH) or =O; R⁵, R⁶, R⁷, and R⁸ are H; Z is O; and R represents R⁴⁴ and the R²⁵ represents pyridyl or phenyl.

6. The method of Claim 5 wherein R²⁵ represents 3-pyridyl or phenyl; R³ is Cl; R⁴ is H; R⁴⁸ represents are H or methyl; and R¹ and R² are individually selected from H, benzotriazol-1-yloxy, C₁ to C₄ alkyl or halo.

7. The method of Claim 6 wherein R¹ and R² are individually selected from H, Br, Cl, methyl or benzotriazol-1-yloxy.

8. The method of Claim 1 wherein the the cells inhibited are tumor cells expressing an activated ras oncogene.

9. The method of Claim 8 wherein the cells inhibited are pancreatic tumor cells, lung cancer cells, myeloid leukemia tumor cells, thyroid follicular tumor cells, myelodysplastic tumor cells, epidermal carcinoma tumor cells, bladder carcinoma tumor cells or colon tumors cells.

10. The method of Claim 1 wherein the inhibition of the abnormal growth of cells occurs by the inhibition of ras farnesyl protein transferase.









11. The method of Claim 1 wherein the inhibition is of tumor cells wherein the Ras protein is activated as a result of oncogenic mutation in genes other than the Ras gene.



5 12. The method of Claim 1 wherein the compound is selected from the compounds of Examples: 1, 2, 3, 4, 5, 6, 19, 42, 43, 44, 45, 46, 47, 48, 49, 75, 76, 78, 82, 83, 84, 85, 89, 121, 180, 182, 183, 184, 187
10 structure 6.7, 187 structure 6.8, 192, 196, 197, 198, 200, 201, 206, 222, 223, 224, 225, 226, 227, 233, 234, 236, 239, 246, 247, 248, 249, 250, 251, 261, 262, 266, 267, 269, 273, 276, 283, 285, 286, 287, 288, 289, 291, 292, 293, 299, 300, 301, 303, 307, 309, 311, 312, 313, 314, 316, 350, 351, 352, 354 or 356.

13. A compound selected from a compound of the formula:





- 245 -



or

- or a pharmaceutically acceptable salt or solvate thereof, wherein all the substituents are as defined in Claim 1, and wherein for the compounds of Formula 5.2 the substituents R^{20} , R^{21} , and R^{46} are selected such that
- 5 when one of said substituents R^{20} , R^{21} , and R^{46} is selected from the group consisting of: (1) H, (2) -OH, (3) -NH₂, (4) -NHC(O)OR²², (5) alkyl, (6) phenyl, (7) heteroaryl, (8) hydroxyalkyl, (9) substituted pyridyl, (10) substituted phenyl and (11) -O-alkyl, then the remaining two of said substituents R^{20} , R^{21} and R^{46} cannot both be H when: (a) R^1 and R^2 are both H, and (b) the double bond between C-5 and C-6 is absent, and (c)
- 10 both A and B are H₂, and (d) R^4 is H, and (e) R^3 is H or Cl at C-8.

14. The compound of Claim 13 wherein a is N and b, c, and d are carbon; R^1 and R^2 are the same or different and each is
- 15 independently selected from H, halo, -CF₃, lower alkyl, or benzotriazol-1-yloxy, and R^1 is at the C-4 position and R^2 is at the C-3 position; R^3 and R^4 are the same or different and each is independently selected from H or halo, and R^3 is at the C-8 position and R^4 is at the C-9 position; when the double bond between carbon atoms 5 and 6 is present, A and B
- 20 independently represent H, lower alkyl or alkyloxy; and when the double bond between carbon atoms 5 and 6 is absent, A and B independently represent H₂, (-H and -OH) or =O; R^5 , R^6 , R^7 , and R^8 are H; Z is O; and R^{46} is selected from phenyl, substituted phenyl, heteroaryl or piperidine Ring V.

25

15. The compound of Claim 14 wherein R^{20} and R^{21} are each independently selected from H and alkyl; R^3 is Cl; R^4 is H; R^1 and R^2 are

THIS SHEET HAS BEEN
REPLACED



individually selected from H, benzotriazol-1-yloxy, C₁ to C₄ alkyl or halo; and R⁴⁶ represents 3-pyridyl, 3-pyridyl N-oxide, 4-pyridyl, 4-pyridyl N-oxide, 3-N-methylpiperidinyl, 4-N-methylpiperidinyl, 3-N-acetylpiperidinyl or 4-N-acetylpiperidinyl.

5

16. The compound of Claim 15 wherein both R²⁰ and R²¹ are H, or both R²⁰ and R²¹ are methyl; R¹ and R² are individually selected from H, Br, Cl, methyl or benzotriazol-1-yloxy; and R⁴⁶ represents 3-pyridyl, 3-pyridyl N-oxide, 4-pyridyl, 4-pyridyl N-oxide, 3-N-methylpiperidinyl or 4-N-methylpiperidinyl.

10

17. The compound of Claim 13 wherein a is N and b, c, and d are carbon; R¹ and R² are the same or different and each is independently selected from H, halo, -CF₃, lower alkyl, or benzotriazol-1-yloxy, and R¹ is at the C-4 position and R² is at the C-3 position; R³ and R⁴ are the same or different and each is independently selected from H or halo, and R³ is at the C-8 position and R⁴ is at the C-9 position; when the double bond between carbon atoms 5 and 6 is present, A and B independently represent H, lower alkyl or alkyloxy; and when the double bond between carbon atoms 5 and 6 is absent, A and B independently represent H₂, (-H and -OH) or =O; R⁵, R⁶, R⁷, and R⁸ are H; Z is O; and R²⁵ represents phenyl, 3-pyridyl, 3-pyridyl N-oxide, 4-pyridyl, 4-pyridyl N-oxide, 3-N-methylpiperidinyl, 4-N-methylpiperidinyl, 3-N-acetylpiperidinyl or 4-N-acetylpiperidinyl.

25


18. The compound of Claim 17 wherein R²⁵ represents phenyl, 3-pyridyl, 3-pyridyl N-oxide, 4-pyridyl, 4-pyridyl N-oxide, 3-N-methylpiperidinyl or 4-N-methylpiperidinyl; R³ is Cl; R⁴ is H; R⁴⁸ represents are H or metnly; and R¹ and R² are individually selected from H, benzotriazol-1-yloxy, C₁ to C₄ alkyl or halo.

30

19. The compound of Claim 18 wherein R¹ and R² are individually selected from H, Br, Cl, methyl or benzotriazol-1-yloxy.

35

20. The compound of Claim 13 selected from a compound having the structure number: 5.17, 5.18, 5.19, 5.20, 5.21, 5.22, 5.23, 5.60, 5.61, 5.62, 5.63, 5.64, 5.65, 5.66, 5.67, 5.68, 5.69, 5.70, 5.71, 5.72, 5.73, 5.74, 5.75, 5.76, 5.77, 5.78, 5.79, 5.81, 5.82, 5.83, 5.84, 5.85, 5.90, 5.91,



5

15

Formula 1.0h:




reacting one molar equivalent of a compound of Formula 1.0g:






- 248 -



wherein R^1 , R^2 , R^3 , R^4 , A , B , a , b , d , and the dotted lines are as defined for Formula 1.0 in Claim 1, and R^{65} represents H or $-OR^{66}$ wherein R^{66} represents alkyl;

- 5 with one molar equivalent of a nitrating reagent, said nitrating reagent being preformed by mixing, at cold temperature, equimolar amounts of tetrabutyl ammonium nitrate with trifluoroacetic anhydride; the reaction of said nitrating reagent with said compound of Formula 1.0g taking place in a suitable aprotic solvent; and
- 10 said reaction with said nitrating reagent being conducted at a temperature and for a period of time sufficient to allow the reaction to proceed at a reasonable rate to produce the 3-nitro compound of Formula 1.0h.

- 15 23. A process for producing 3-nitro compounds of the formula:




* C L M S *

THIS SHEET HAS BEEN
REPLACED


wherein R^1 , R^2 , R^3 , R^4 , A, B, a, b, d, and the dotted lines are as defined for Formula 1.0 in Claim 1, comprising:

reacting one molar equivalent of a compound of Formula 1.0g:



- 5 wherein R^1 , R^2 , R^3 , R^4 , A, B, a, b, d, and the dotted lines are as defined for Formula 1.0 in Claim 1, and R^{65} represents H or $-OR^{66}$ wherein R^{66} represents alkyl;
- with one molar equivalent of a nitrating reagent, said nitrating reagent being preformed by mixing, at cold temperature, equimolar
- 10 amounts of tetrabutyl ammonium nitrate with trifluoroacetic anhydride;
- the reaction of said nitrating reagent with said compound of Formula 1.0g taking place in a suitable aprotic solvent; and
- said reaction with said nitrating reagent being conducted at a temperature and for a period of time sufficient to allow the reaction to
- 15 proceed at a reasonable rate to produce the 3-nitro compound of Formula 1.0h:






hydrolyzing the compound of Formula 1.0h by dissolving the compound of Formula 1.0h in a sufficient amount of concentrated acid, and heating the resulting mixture to a temperature sufficient to remove the

5 -C(O)R⁶⁵ substituent to produce the compound of Formula 1.0i.


24. A process for producing compounds of the formula:



wherein R¹, R², R³, R⁴, A, B, a, b, d, and the dotted lines are as defined for

10 Formula 1.0 in Claim 1, comprising:

reacting one molar equivalent a compound of formula:



with one molar equivalent of a nitrating reagent;





- 251 -


said nitrating reagent being preformed, by mixing at a cold temperature, equimolar amounts of tetrabutyl ammonium nitrate with trifluoroacetic anhydride;

- the reaction of said nitrating reagent with the compound of Formula 1.0k taking place in a suitable aprotic solvent;

said reaction with said nitrating reagent being conducted at a temperature and for a period of time sufficient to allow the reaction to proceed at a reasonable rate to produce the 3-nitro compound of Formula 1.0j.

10


25. A process for producing a compound of Formula 1.0m:



wherein R^1 , R^2 , R^3 , R^4 , A, B, a, b, d, and the dotted lines are as defined for Formula 1.0 in Claim 1, and wherein R^{68} is H or $-COOR^a$ wherein R^a is a

- 15 C_1 to C_3 alkyl group, comprising:

reacting one molar equivalent a compound of formula:



with one molar equivalent of a nitrating reagent;

- said nitrating reagent being preformed, by mixing at a cold temperature, equimolar amounts of tetrabutyl ammonium nitrate with trifluoroacetic anhydride;

20




THIS SHEET HAS BEEN REPLACED



- 252 -


the reaction of said nitrating reagent with the compound of Formula 1.0k taking place in a suitable aprotic solvent;

- 5 said reaction with said nitrating reagent being conducted at a temperature and for a period of time sufficient to allow the reaction to proceed at a reasonable rate to produce the 3-nitro compound of Formula 1.0j:




- 10 reducing said compound of Formula 1.0j with a suitable reducing agent in a suitable solvent at a suitable temperature to allow the reaction to proceed at a reasonable rate;

reacting the resulting hydroxy product with a chlorinating agent in a suitable organic solvent at a suitable temperature to allow the reaction to proceed at a reasonable rate to produce a compound of Formula 1.0n:



; and

- 15 reacting said compound of Formula 1.0n with a compound of the formula:



wherein R⁶⁸ is as previously defined, in a suitable organic solvent containing a suitable base at a suitable temperature to allow the reaction




* C L M S *

THIS SHEET HAS BEEN
REPLACED




to proceed at a reasonable rate to produce the compounds of Formula 1.0m.

26. A compound selected from a compound of the formula:



5

wherein R^1 , R^2 , R^3 , R^4 , A , B , a , b , d , and R^{65} are as defined for Formula 1.0h in Claim 22;




10 wherein R^1 , R^2 , R^3 , R^4 , A , B , a , b , and d are as defined for Formula 1.0i in Claim 23;






- 254 -




wherein R^1 , R^2 , R^3 , R^4 , A, B, a, b, and d are as defined for Formula 1.0j in Claim 24;



5

wherein R^1 , R^2 , R^3 , R^4 , A, B, a, b, d and R^{68} are as defined for Formula 1.0m in Claim 25;




10 wherein R^1 , R^2 , R^3 , R^4 , A, B, a, b, and d are as defined for Formula 1.0j in Claim 24; or



* C L N S *


THIS SHEET HAS BEEN
REPLACED

- 255 -



wherein R^1 , R^2 , R^3 , R^4 , A, B, a, b, and d are as defined for Formula 1.0j in Claim 24.

- 5 27. A compound selected from a compound of the formula:



28. The use of a compound of Claim 13 for the manufacture of a medicament for use in inhibiting the abnormal growth of cells.

10

29. The use of a compound of Claim 13 for inhibiting the abnormal growth of cells.

30. A process for producing a compound of Formula 5.0, 5.1 or 5.3, comprising:

- 15 (a) reacting a compound of Formula 405.00 with $RCOOH$ at a temperature between $0^\circ C$ and reflux in an inert solvent;
- (b) reacting a compound of Formula 405.00 with $RC(O)L$, wherein L represents a suitable leaving group, in the presence of a base;
- 20 (c) reacting a compound of Formula 420.00 with an appropriate acyl halide or an acyl anhydride;
- (d) reducing compounds of Formula 405.00, wherein X is carbon and the bond to C-11 is a double bond, with lithium aluminum hydride in tetrahydrofuran, and then reacting reduced Formula 405.00



* C L H S *

THIS SHEET HAS BEEN
REPLACED



with RCOOH or RC(O)L to produce compounds of Formula 1.0 wherein X is carbon and the bond to C-11 is a single bond;

5 (e) reducing compounds of Formula 470.00d with sodium borohydride in methanol or ethanol, converting the resulting alcohol (470.00e) to a chloride (470.00f) with thionyl chloride, displacing with piperazine (405.00c), and reacting the resulting compound with RCOOH or RC(O)L as described above for the conversion of Formula 405.00 to Formula 1.0 wherein the bond to C-11 is a single bond;

10 (f) cyclizing a compound of Formula 455.00 to produce a compound of Formula 1.0 wherein the bond to C-11 is a double bond;

(g) reacting the azaketone of Formula 470.00 with an oxidizing agent to produce the corresponding compound wherein the nitrogen of the pyridine ring is an N-oxide, reacting the resulting product with a chlorinating agent to produce a compound of Formula 470.00b, optionally
15 repeating the chlorination to produce the disubstituted compound of Formula 470.00c, optionally reacting the compound of Formula 470.00b or 470.00c with nucleophiles to obtain the substituted compound of Formula 470.00d, reacting the compound of 470.00d as described in (f) above to produce compounds of Formula 1.0;

20 (h) reacting the N-oxide of the compound of Formula 405.00 with POCl₃ to form a compound of Formula 415.01 wherein R¹ or R² is Cl;

(i) reacting a halo-substituted compound of formula 405.00, wherein R¹ is halo, an alkyl lithium to provide the lithio derivative and then quenching the lithio derivative with the appropriate electrophile;

25 (j) reacting a compound of Formula 1.0, wherein Z is O, with P₂S₅ to produce a compound of Formula 1.0 wherein Z is S;

(k) reacting a compound of Formula 470.00h in acetic acid with SeO₂ to produce a compound of Formula 470.00i and then converting the compound of 470.00i to a compound of Formula 1.0.

30 (l) alkylating the appropriately substituted piperazine compound of Formula 700.00 with a compound of Formula 705.00 to produce a compound of formula 1.0 wherein X is N; or

(m) reductive amination of the aza ketone of Formula 715.00 with the piperazine of Formula 700.00.

35

