

PROTEIN S ANTIBODIES, METHODS OF MAKING AND USES THEREOF**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims priority to U.S. Provisional Patent Application No. 63/020,505, filed on May 5, 2020 and to U.S. Provisional Patent Application No. 63/169,755 filed April 1, 2021. The contents of each of the preceding applications are incorporated by reference herein in their entirety.

BACKGROUND

[0002] Protein S (also known as ProS, ProS1) is a vitamin K-dependent plasma protein involved in the anti-coagulation cascade. The protein is multi-modular, comprising a γ -carboxy-glutamic acid domain (Gla domain), an epidermal growth factor-like domain (EGF domain), a thrombin-sensitive region (TSR), and a sex hormone binding globulin-like domain (SHBG-like domain). The protein is found in both a free form and as part of a complex with proteins such as C4 binding protein (C4BP) and tissue factor pathway inhibitor (TFPI). Among other functions, in its free form, Protein S is a cofactor in at least two pathways of the anti-coagulation cascade: (1) Protein S is a cofactor for plasma activated Protein C (APC), involved in the inactivation and degradation of coagulation factors Factor Va and Factor VIIIa; and (2) Protein S is also a cofactor for TFPI, also present in plasma, involved in the inactivation of coagulation factors Factor Xa and Factor VIIa.

[0003] Protein S is a potential therapeutic target for bleeding disorders, thus there is a need for agents that bind and modulate its activities within the coagulation pathway.

SUMMARY

[0004] Provided here are antibodies that bind Protein S, and methods of making and using such antibodies. In some embodiments, the Protein S antibodies provided herein are useful for treating a bleeding disorder, or a condition characterized by reduced or impaired blood coagulation and/or clotting.

[0005] Accordingly, in one aspect, provided herein are antibodies that bind Protein S, wherein the antibodies are inhibitors of the cofactor activity of Protein S for activated Protein C (APC), inhibitors of the cofactor activity of Protein S for tissue factor pathway inhibitor (TFPI), or

inhibitors of the cofactor activity of Protein S for both APC and TFPI (dual inhibitor of cofactor activity), and wherein the antibody is human, humanized, or chimeric. In some preferred embodiments, the antibodies provided herein specifically bind Protein S.

[0006] In another aspect, provided herein are antibodies that bind Protein S, wherein the antibodies are capable of promoting coagulation and/or modulating a component in the coagulation cascade.

[0007] In another aspect, provided herein are exemplary Protein S antibodies comprising any one or more of the amino acid sequences of the complementarity determining region (CDR) sequences provided in Tables 1A, 1B, 1C, 2A, 2B, and 2C. In another aspect, the exemplary Protein S antibodies comprise any one of the CDR-L1 amino acid sequences of Table 1A; any one of the CDR-L2 amino acid sequences of Table 1B; any one of the CDR-L3 amino acid sequences of Table 1C; any one of the CDR-H1 amino acid sequences of Table 2A; any one of the CDR-H2 amino acid sequences of Table 2B; any one of the CDR-H3 amino acid sequences of Table 2B. In another aspect, provided herein are exemplary Protein S antibodies comprising the combinations of variable light chains and variable heavy chains presented in Table 4C. In another aspect, provided herein are the sequences of different antibodies as presented in Table 6. In another aspect, provided herein are nucleic acids encoding for any of the Protein S antibodies provided herein.

[0008] In another aspect, provided herein are pharmaceutical compositions comprising any one of the Protein S antibodies provided herein, and optionally a pharmaceutically acceptable excipient.

[0009] Provided herein are methods of using the exemplary Protein S antibodies described herein. Accordingly, in one aspect, provided herein is an *in vitro* method of promoting the coagulation of blood, comprising contacting any one of the Protein S antibodies provided herein with a blood sample.

[0010] In another aspect, provided herein is a method of promoting the coagulation of blood in a subject in need thereof, comprising administering to the subject any of the exemplary Protein S antibodies provided herein, or pharmaceutical compositions provided herein. In exemplary embodiments, the route of administration is subcutaneous.

[0011] In another aspect, provided herein is a method of promoting the generation of thrombin in a subject in need thereof, comprising administering to the subject any of the Protein S antibodies provided herein, or pharmaceutical compositions provided herein.

[0012] In another aspect, provided herein is a method of treating a condition in a subject in need thereof, comprising administering to the subject any of the Protein S antibodies provided herein, or pharmaceutical compositions provided herein, wherein the condition is selected from the group consisting of: bleeding disorders, platelet disorders, trauma, bleeding resulting from a surgery or a medical procedure, and combinations thereof.

[0013] In another aspect, provided herein is the use of any one of the Protein S antibodies or pharmaceutical compositions provided herein, for the treatment of a condition in a subject in need thereof. The condition may be selected from the group consisting of: bleeding disorders, platelet disorders, trauma, bleeding resulting from a surgery or a medical procedure, and combinations thereof.

[0014] In another aspect, any one of the Protein S antibodies or pharmaceutical compositions provided herein, may be used for the manufacture of a medicament for the treatment of a condition in a subject in need thereof. The condition may be selected from the group consisting of: bleeding disorders, platelet disorders, trauma, bleeding resulting from a surgery or a medical procedure, and combinations thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] **FIG. 1** depicts a schematic diagram of Protein S showing the modular domains of Protein S.

[0016] **FIGS. 2A-2B** depict the results of screening assays performed to assess APC and TFPI cofactor activity, respectively, using control Protein S antibodies. In this and subsequent figures, ProS = Protein S; mAb = monoclonal antibody; pAb = polyclonal antibody; PRP = platelet rich plasma.

[0017] **FIGS. 3-4** depict peak thrombin levels generated by Protein S monoclonal antibodies, identified from hybridoma libraries, in the APC cofactor and TFPI cofactor screening assays, respectively.

[0018] **FIGS. 5-10** depict the prototypic thrombin generation profiles of a dual inhibitor of APC and TFPI (Antibody 13; FIGS 5, 6), an APC cofactor inhibitor (Antibody 21; FIGS 7, 8), and a TFPI cofactor inhibitor (Antibody 23; FIGS 9, 10) when using a TFPI cofactor assay and an APC cofactor assay.

[0019] **FIGS. 11A-11H** depict the characterization of Antibody 19 and Antibody 7, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0020] **FIGS. 12A-12H** depict the characterization of Antibody 13 and Antibody 1, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0021] **FIGS. 13A-13H** depict the characterization of Antibody 14 and Antibody 2, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0022] **FIGS. 14A-14H** depict the characterization of Antibody 15 and Antibody 3, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0023] **FIGS. 15A-15H** depict the characterization of Antibody 16 and Antibody 4, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0024] **FIGS. 16A-16H** depict the characterization of Antibody 20 and Antibody 8, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0025] **FIGS. 17A-17H** depict the characterization of Antibody 21 and Antibody 9, antibodies sharing the same human variable region, and are characterized as APC cofactor inhibitors.

[0026] **FIGS. 18A-18G** depict the characterization of Antibody 23 and Antibody 11, antibodies sharing the same human variable region, and are characterized as TFPI cofactor inhibitors.

[0027] **FIGS. 19A-19H** depict the characterization of Antibody 24 and Antibody 12, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0028] **FIGS. 20A-20H** depict the characterization of Antibody 18 and Antibody 6, antibodies sharing the same human variable region, and are characterized as APC cofactor inhibitors.

[0029] **FIGS. 21A-21G** depict enhanced thrombin generation when Antibody 15 is added to various samples of congenital factor deficient plasma containing soluble thrombomodulin.

[0030] **FIGS 22A-22B** depict the levels of D-dimer, as a marker of coagulation activity, observed over time in cynomolgus monkeys injected with either 1 mg/kg or 3 mg/kg of Antibody 1.

[0031] **FIGS. 23A-23C** depict the level of fibrin deposited onto collagen coated spots in Factor VIII (FVIII) neutralized blood treated with various Protein S antibodies, showing a restoration of fibrin deposition activity by the Protein S antibodies in the FVIII neutralized blood.

[0032] **FIGS. 24A-24B** depict exemplary dose response curves exhibited by antibodies 19 and 15 in an APC cofactor screening assay. The antibodies exhibit graded inhibition and switch-like inhibition, respectively, in the assay.

[0033] **FIGS. 25A-25B** depict the effect of selected Protein S antibodies on fibrin deposition.

[0034] **FIGS. 26A-26B** depict the effects of Protein S antibodies in an *in vitro* microfluidic Hemophilia A bleeding model.

[0035] **FIGS. 27A-27F** depict the effect of selected Protein S antibodies on enhanced dose-dependent thrombin generation, wherein the Protein S antibodies were added to plasma obtained from patients with various types of von Willebrand disease.

[0036] **FIGS. 28A-28N** depict the results of thrombin generation assays (*ex vivo* pharmacodynamic assay) as well as pharmacokinetic assays used to measure free antibody, in cynomolgus monkeys injected subcutaneously or intravenously with Protein S antibodies. These figures also depict levels of D-dimer, used as a marker of coagulation activity, observed in the monkeys.

[0037] **FIG. 29** depicts the effect of Antibody 2 on the ability of Protein S to bind TFPI.

[0038] **FIGS. 30A-30B** depict similar dose-titration curves resulting from a full-length antibody or a Fab fragment of the same antibody in an APC cofactor assay.

DETAILED DESCRIPTION

[0039] Provided herein are antibodies that bind Protein S, and methods of making and using such antibodies. In some embodiments, the Protein S antibodies provided herein specifically bind Protein S. For example, antibodies that bind to Protein S and inhibit its cofactor activity for TFPI and/or APC, may be useful in the treatment of bleeding disorders and other related diseases by promoting clot formation

[0040] Where elements are presented in a list format (e.g., in a Markush group), it should be understood that each possible subgroup of the elements is also disclosed, and that any one or more elements can be removed from the list or group.

[0041] It should be understood that, unless clearly indicated, in any method described or disclosed herein that includes more than one act, the order of the acts is not necessarily limited to the order in which the acts of the method are recited, but the disclosure encompasses exemplary embodiments in which the order of the acts is so limited.

[0042] The terms used throughout the specification are defined as follows unless otherwise limited in specific instances. As used in the specification and the claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. All technical and scientific terms, acronyms, and abbreviates used in the specification and claims have the same meaning as commonly understood by one of ordinary skill in the art to which the disclosure pertains, unless defined or stated otherwise. All numerical ranges are inclusive of the values defining the range as well as all integer values in between, unless indicated or defined otherwise.

[0043] The term “antibody” as used herein throughout is used in the broadest sense and may include a monoclonal antibody, polyclonal antibody, human antibody, humanized antibody, non-human antibody, chimeric antibody, a monovalent antibody, an antigen-binding fragment (e.g., a Fab fragment, a Fab’2 fragment, an scFv), and other antibody fragments that retain specificity for and binding of Protein S. In some embodiments, the antibodies are monoclonal antibodies. In some embodiments, the antibodies are monoclonal antibodies. In some embodiments, the antibodies are monoclonal human antibodies. In some embodiments, the antibodies are

monoclonal humanized antibodies. In some embodiments, the antibodies are monoclonal chimeric antibodies.

[0044] Also provided herein are antibody-drug conjugates, bispecific antibodies, and multispecific antibodies that exhibit specificity for and binding of Protein S.

[0045] The terms “polynucleotide” and “nucleic acid” are used interchangeably herein, and refer to a polymeric form of nucleotides of any length, which may be ribonucleotides or deoxyribonucleotides. The terms include, but are not limited to, single-, double-, or multi-stranded DNA or RNA, genomic DNA, cDNA, DNA-RNA hybrids, or a polymer comprising purine and pyrimidine bases or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases. The terms encompass nucleic acids containing known analogues of natural nucleotides and having similar binding properties, and are metabolized in a manner similar to naturally-occurring nucleotides, unless specifically limited or stated otherwise.

[0046] When a nucleic acid or amino acid sequence is said to have a certain percent “sequence identity” or “identity” or is a certain percent “identical” to another nucleic acid or amino acid sequence, that percentage of bases or amino acids are the same, and in the same relative position, when the sequences are aligned, when comparing the two sequences.

[0047] The term “subject,” as used herein refers to any subject for whom treatment or therapy is provided. The subject may be a mammalian subject. Mammalian subjects include, e. g., humans, non-human primates (e.g., cynomolgus monkey), rodents, (e.g., rats, mice), lagomorphs (e.g., rabbits), ungulates (e.g., cows, sheep, pigs, horses, goats, and the like), etc. In some embodiments, the subject is a human. In some embodiments, the subject is a non-human primate, e.g. a cynomolgus monkey. In some embodiments, the subject is a companion animal (e.g. cats, dogs).

[0048] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

I. Antibodies That Bind and Modulate Protein S Activity

A. Protein S Antibodies

[0049] Provided here are antibodies that bind to Protein S, and in some embodiments are specific for Protein S. The Protein S can be of any species, e.g. any mammalian species. In some embodiments, the Protein S antibody binds to human Protein S. In some embodiments, the Protein S antibody binds to the Protein S of non-human primates. In some embodiments, the non-human primate is cynomolgus monkey.

[0050] The amino acid sequence of human Protein S, targeted by antibodies of the disclosure, is shown below as SEQ ID. NO: 216.

10	20	30	40	50
MRVLGGRCGA	LLACLLLVLP	VSEANFLSKQ	QASQVLVRKR	RANSLLEETK
60	70	80	90	100
QGNLERECIE	ELCNKEEARE	VFENDPETDY	FYPKYLVLCL	SFQTGLFTAA
110	120	130	140	150
RQSTNAYPDL	RSCVNAIPDQ	CSPLPCNEDG	YMSCKDGKAS	FTCTCKPGWQ
160	170	180	190	200
GEKCEFDINE	CKDPSNINGG	CSQICDNTPG	SYHCSCKNKF	VMLSNKKDCK
210	220	230	240	250
DVDECSLKPS	ICGTAVCKNI	PGDFECECPE	GYRYNLKSKS	CEDIDECSEN
260	270	280	290	300
MCAQLCVNYP	GGYTCYCDGK	KGFKLAQDQK	SCEVVSVCCLP	LNLDTKYELL
310	320	330	340	350
YLAEQFAGVV	LYLKFRLEPI	SRFSAEFDFR	TYDSEGVILY	AESIDHSAWL
360	370	380	390	400
LIALRGGKIE	VQLKNEHTSK	ITTGGDVINN	GLWNMVSVEE	LEHSISIKIA
410	420	430	440	450
KEAVMDINKP	GPLFKPENGL	LETKVYFAGF	PRKVESELIK	PINPRLDGCI
460	470	480	490	500
RSWNLMKQGA	SGIKEIIQEK	QNKHCLVTVE	KGSYYPGSGI	AQFHIDYNNV
510	520	530	540	550
SSAEGWHVNV	TLNIRPSTGT	GVMLALVSGN	NTVPFAVSLV	DSTSEKSQDI
560	570	580	590	600
LLSVENTVIY	RIQALSLCSD	QQSHLEFRVN	RNNLELSTPL	KIETISHEDL
610	620	630	640	650
QRQLAVLDKA	MKAKVATYLG	GLPDVPFSAT	PVNAFYNGCM	EVNINGVQLD
660	670			

LDEAISKHND IRAHSCPSVW KKTKNS (SEQ ID. NO: 216)

[0051] In some embodiments, provided herein are Protein S antibodies comprising a binding affinity (KD) to Protein S of about 0.0005 nM or lower, 0.001 nM or lower, 0.005 nM or lower, 0.01 nM or lower, 0.05 nM or lower, about 0.1 nM or lower, about 0.5 nM or lower, about 1nM

or lower, about 5 nM or lower, about 10 nM or lower, about 50 nM or lower, about 100 nM or lower, about 500 nM or lower, or about 1 μ M or lower.

[0052] The Protein S antibodies provided herein are capable of modulating one or more activities of Protein S, including, for example, modulating Protein S cofactor activity, as well as promoting coagulation and altering levels of markers associated with coagulation, and modulating a component in the coagulation cascade.

[0053] In some embodiments, the Protein S antibodies provided herein are capable of modulating the Protein S ability to act as a cofactor within pathways of the coagulation cascade.

[0054] Accordingly, in some embodiments, the Protein S antibodies provided herein are useful for reducing or inhibiting the cofactor activity of Protein S for activated Protein C (“APC”).

[0055] In some embodiments, the Protein S antibodies provided herein are useful for reducing or inhibiting the cofactor activity of Protein S for tissue factor pathway inhibitor (“TFPI”). TFPI is an inhibitor of procoagulant activity and is produced as at least two alternatively spliced isoforms in humans, TFPI α , and TFPI β , which differ in domain structure and mechanism for cell surface association. TFPI α , but not TFPI β contains Kunitz domain 3, the domain which is believed to be involved in binding to Protein S. Without being held to any theory or mechanism, it is believed that the Protein S antibodies of the disclosure inhibit the cofactor activity of Protein S for at least TFPI α , as it contains Kunitz domain 3.

[0056] In some embodiments, the Protein S antibodies provided herein are useful for reducing or inhibiting the cofactor activity of Protein S for APC, but show negligible or no effect on cofactor activity of Protein S for TFPI (such antibodies are referred to interchangeably herein as “APC cofactor inhibitors”, “APC cofactor specific inhibitors”, or “APC pathway inhibitors”).

[0057] In some embodiments, the Protein S antibodies provided herein are useful for reducing or inhibiting the cofactor activity of Protein S for TFPI, but show negligible or no effect on cofactor activity of Protein S for APC (such antibodies are referred to interchangeably herein as “TFPI cofactor inhibitors”, “TFPI cofactor specific inhibitors”, or “TFPI pathway inhibitors”).

[0058] In some embodiments, the Protein S antibodies provided herein are useful for reducing or inhibiting the cofactor activity of Protein S for both APC and TFPI (such antibodies are

referred to herein as “dual inhibitors”). The dual inhibitors of the disclosure may reduce the activities of APC and TFPI to different degrees.

[0059] In some embodiments, the Protein S antibodies provided herein are capable of causing a reduction in the activity of APC. For example, APC activity may be reduced by about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or about 100% as compared to in the absence of the Protein S antibodies.

[0060] In some embodiments, the Protein S antibodies provided herein are capable of causing a reduction in the activity of TFPI. For example, TFPI activity may be reduced by about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or about 100% as compared to in the absence of the Protein S antibodies.

[0061] In some embodiments, the Protein S antibodies provided herein are capable of causing a reduction in the activity of both APC and TFPI. For example, APC activity may be reduced by about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or about 100%, and TFPI activity may also be reduced by about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or about 100% as compared to in the absence of the Protein S antibodies. The reduction in APC activity and TFPI activity may be affected to different degrees by the same Protein S antibody.

[0062] In some embodiments, the Protein S antibodies provided herein are APC cofactor inhibitors, and the capability of the antibody for affecting the cofactor activity of Protein S for TFPI is negligible. In other embodiments, the Protein S antibodies provided herein are TFPI cofactor inhibitors, and the capability of the antibody for affecting the cofactor activity of Protein S for APC is negligible.

[0063] In some embodiments, the Protein S antibodies provided herein are capable of promoting coagulation and/or modulating a component in the coagulation cascade, e.g., in a subject in need thereof, or in a sample. In some embodiments, the Protein S antibodies provided herein are capable of promoting the clotting of blood. In some embodiments, the antibodies are capable of promoting clotting of blood by reducing the ability of Protein S to act as a cofactor for APC. For example, clotting of blood may be promoted by reducing APC activity by about

10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or about 100%, as compared to in the absence of the Protein S antibodies.

[0064] In some embodiments, the antibodies are capable of promoting clotting of blood by reducing an ability of Protein S to act as a cofactor for TFPI. For example, clotting of blood may be promoted by reducing TFPI activity by about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or about 100%, as compared to in the absence of the Protein S antibodies.

[0065] In some embodiments, the antibodies are capable of promoting clotting of blood by reducing an ability of Protein S to act as a cofactor for both APC and TFPI. For example, clotting of blood may be promoted by reducing both APC activity and TFPI activity by about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or about 100%, as compared to in the absence of the Protein S antibodies.

[0066] In some embodiments, the Protein S antibodies provided herein are capable of promoting the generation of a marker associated with coagulation activity, and this can be exhibited *in vitro* (e.g. in a sample) and/or *in vivo* (e.g. upon administration to a subject). Such markers include, but are not limited to, thrombin, fibrin, D-dimer, clot formation, thrombin-thrombin complex, fibrin degradation products, and prothrombin fragment F1.2.

[0067] In some embodiments, the Protein S antibodies provided herein are capable of promoting thrombin generation (e.g. includes restoring thrombin generation or restoring the levels of thrombin, e.g., in a subject in need thereof, or in a sample). In some embodiments, the antibodies are capable of promoting generation of thrombin in a subject in need thereof. In some embodiments, the generation of thrombin does not exceed a predetermined threshold level. In some embodiments, the generation of thrombin is partially restored. In some embodiments, the generation of thrombin does not exceed a predetermined percentage of a maximum thrombin generation. In some embodiments, the generation of thrombin does not exceed a predetermined percentage of an area under the curve of the maximum thrombin generation. In some embodiments, thrombin generation may be increased by about 5-fold to 50-fold, e.g. by about 5-fold, by about 10-fold, by about 15-fold, by about 20-fold, by about 25-fold, by about 30-fold, by about 35-fold, by about 40-fold, by about 45-fold, or even by about 50-fold, as compared to in the absence of the Protein S antibodies. Exemplary antibodies that promote thrombin

generation are described in greater detail below, and include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0068] In some embodiments, the Protein S antibodies provided herein can restore or promote thrombin generation in a subject who is deficient in coagulation factors. In some exemplary embodiments, the coagulation factor deficiency is congenital. In some exemplary embodiments, the coagulation factor deficiency is acquired. In some embodiments, the Protein S antibodies provided herein can promote thrombin generation in a subject who is deficient in Factor VII, Factor VIII, Factor IX, Factor XI. For example, thrombin generation can be increased by about 5-fold to 50-fold, e.g. by about 5-fold, by about 10-fold, by about 15-fold, by about 20-fold, by about 25-fold, by about 30-fold, by about 35-fold, by about 40-fold, by about 45-fold, or even by about 50-fold, as compared to in the absence of the Protein S antibodies, when used to promote thrombin generation in a sample from a subject with a factor deficiency.

[0069] In some embodiments, the Protein S antibodies provided herein can promote thrombin generation in a subject who suffers from von Willebrand Disease (vWD) disease. In some embodiments, the vWD is a subtype selected from: vWD Type 1, vWD Type 2A, vWD Type 2B, vWD Type 2N, vWD Type 2M, vWD Type 3, and acquired vWD. In some exemplary embodiments, the vWD is a subtype selected from Type 1, Type 2, or Type 3. For example,

thrombin generation can be increased by about 5-fold to 50-fold, e.g. by about 5-fold, by about 10-fold, by about 15-fold, by about 20-fold, by about 25-fold, by about 30-fold, by about 35-fold, by about 40-fold, by about 45-fold, or even by about 50-fold, as compared to in the absence of the Protein S antibodies, when used to promote thrombin generation in a sample from a subject with vWD disease.

[0070] Exemplary antibodies that may promote thrombin generation in a subject who is deficient in a coagulation factor (such as Factor VII, Factor VIII, Factor IX, Factor XI) or who has von Willebrand disease (such as Type 1, Type 2A, Type 2B, Type 2M, Type 2N or Type 3) include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0071] In some embodiments, the Protein S antibodies provided herein are capable of promoting fibrin generation, e.g., in a subject in need thereof, or in a sample. In some embodiments, the Protein S antibodies provided herein are capable of promoting fibrin deposition e.g., in a sample. In some embodiments, the Protein S antibodies provided herein are capable of promoting coagulation activity, wherein the coagulation activity is marked by a promotion of fibrin generation. For example, fibrin generation may be increased by about 5-fold to 50-fold, e.g. by about 5-fold, by about 10-fold, by about 15-fold, by about 20-fold, by about 25-fold, by about 30-fold, by about 35-fold, by about 40-fold, by about 45-fold, or even by about

50-fold, as compared to in the absence of the Protein S antibodies. as compared to in the absence of the Protein S antibodies. Exemplary antibodies that may promote fibrin generation include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0072] In some embodiments, the Protein S antibodies provided herein are capable of promoting increasing D-dimer levels, e.g., in a subject in need thereof, or in a sample. In some embodiments, the Protein S antibodies provided herein are capable of promoting coagulation activity, wherein the coagulation activity is marked by an increase in D-dimer levels. For example, D- dimer levels may be increased by about two-fold to about 10,000-fold. Exemplary antibodies that may promote an increase in D-dimer levels include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f)

comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0073] In some embodiments, the Protein S antibodies provided herein are capable of promoting coagulation in a sample or in a subject. In some embodiments, the Protein S antibodies provided herein alter the levels of markers associated with coagulation activity in a sample or in a subject. For example, in some embodiments, the antibodies are capable of restoring or promoting thrombin generation in a sample, or in a subject. In some embodiments, the antibodies are capable of restoring fibrin deposition in a sample, or in a subject. In some embodiments, the antibodies provided herein are capable of promoting a restoration of fibrin deposition. In some embodiments, the antibodies provided herein are capable of increasing the levels of D-dimer in a sample, or in a subject. In some embodiments, the antibodies provided herein are capable of promoting an increase of D-dimer. In some embodiments, activity of the antibodies provided herein is dose-dependent. In some embodiments, activity of the antibodies provided herein is measured *in vitro*. In some embodiments, activity of the antibodies provided herein is measured *in vivo*.

[0074] In some embodiments, the Protein S antibodies provided herein are administered to a subject, wherein the antibody remains active in the subject for a period of time.

[0075] In some embodiments, the Protein S antibodies provided herein are administered to a subject, wherein the effect of the antibody is dose-dependent, such antibodies exhibit graded inhibition. FIG. 24A (Antibody 19) depicts an exemplary dose response curve of thrombin generation resulting from an antibody exhibiting graded inhibition. Antibodies exhibiting graded inhibition show concentration-dependent inhibition *in vitro*, where increasing concentrations of antibody result in incremental increases in thrombin generation over a wide concentration range. Exemplary antibodies that exhibit such graded inhibition include, but are not limited to, antibodies that comprise (a) the CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57; or (b) the CDR amino

acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, SEQ ID NO: 67.

[0076] In such embodiments, the graded inhibition can allow for increasing the dose of administration of the Protein S antibodies to a subject for added efficacy, or decreasing the dose to prevent excess thrombin generation, and/or potential thromboembolic complications, for example. Effectively, the dose can be adjusted to achieve the desired level of inhibition.

[0077] In some embodiments, the Protein S antibodies provided herein exhibit switch-like inhibition, wherein inhibition can be switched on or off. FIG. 24B (Antibody 15) depicts an exemplary dose response curve resulting from an antibody exhibiting switch-like inhibition. Antibodies exhibiting switch-like inhibition show abrupt concentration-dependent increases in thrombin generation in vitro, where upon reaching a concentration capable of promoting thrombin generation, achieves maximal thrombin generated within a narrow concentration range. Exemplary antibodies that exhibit such graded inhibition include, but are not limited to, antibodies that comprise (a) the CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58; or (b) the CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60.

[0078] In some embodiments, the Protein S antibodies provided herein are antibody fragments. In some embodiments, the antibody fragments are antigen-binding fragments (Fab), variable fragments (Fv) containing VH and VL, single chain variable fragments (scFv) containing VH and VL linked together in one chain, or other antibody variable region fragments, such as Fab', F(ab)2, F(ab')2, dsFv diabody, Fc, and Fd polypeptide fragments. The antibody fragments contain a Fc domain.

[0079] In some embodiments, the Protein S antibodies provided herein are monoclonal antibodies (mAbs). In some embodiments, the Protein S antibodies provided herein are human antibodies. In some embodiments, the Protein S antibodies provided herein are monoclonal human antibodies. In some embodiments, the Protein S antibodies provided herein are humanized antibodies. In some embodiments, the Protein S antibodies provided herein are monoclonal humanized antibodies. In some embodiments, the Protein S antibodies provided

herein are chimeric antibodies. In some embodiments, the Protein S antibodies provided herein are monoclonal chimeric antibodies.

[0080] In some embodiments, the Protein S antibodies provided herein are full-length antibodies.

[0081] In some embodiments, the Protein S antibodies provided herein contain an Fc domain (either are full-length or for example, a single chain antibody linked to a Fc domain).

[0082] In some embodiments, the constant region (herein referred to also as a Fc domain, a Fc sequence or simply as a Fc) of a Protein S antibody is a human Fc domain. In some embodiments, the Fc domain of a full-length Protein S antibody is human IgG1, human IgG2, human IgG3, or human IgG4. In some embodiments, the Fc domain of a full-length Protein S antibody is that of a rat. In some embodiments, the Fc domain of a full-length Protein S antibody is rat IgG1 or rat IgG2b. In some embodiments, the Fc domain of a full-length Protein S antibody is that of a non-human primate, e.g. it is a cynomolgus monkey Fc domain.

[0083] In some embodiments, the Protein S antibodies provided herein are chimeric and comprise a variable region from one species, and a constant region from another species, e.g. comprise a human variable region and a rat constant region. In some embodiments, the rat constant region is rat IgG1 or IgG2b. In some embodiments, the antibodies comprise a human variable region and a human constant region. In exemplary embodiments, the human constant region is human IgG1, or human IgG4.

[0084] In some embodiments, the Protein S antibody contains an Fc domain, and the Fc domain of the antibody is a human IgG1 Fc. Exemplary, but non-limiting, human IgG1 Fc domain sequences are provided as SEQ ID NOS: 217.

```

1   ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS
61  GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDKKVEP KSCDKHTTCP PCPAPELLGG
121 PSVFLFPPKP KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN
181 STYRVVSVLT VLNQDNLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE
241 LTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDSDGSFFLY SKLTVDKSRW
301 QQGNVFSCSV MHEALHNHYT QKSLSLSPGK (SEQ ID NO: 217)

```

[0085] In some embodiments, the Protein S antibody contains an Fc domain, and the Fc domain of the antibody is a human IgG4 Fc. An exemplary human IgG4 heavy chain Fc domain sequence is provided as SEQ ID NO: 218.

```
ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLG
TKTYTCNVDPKPSNTKVDKRVESKYGPPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQ
FNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTL
PPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVM
HEALHNNHYTQKSLSLSLGK (SEQ ID NO: 218)
```

[0086] The EU numbering scheme is one of many available antibody numbering schemes based on the residue numbers assigned to a canonical antibody sequence. Accordingly, a skilled artisan would understand that reference to a particular residue using the EU numbering scheme may or may not be exactly the residue in one of the antibodies of the disclosure. For example, if a Protein S antibody of the disclosure comprises a V215A substitution in the Fc, wherein the position number of the amino acid residue is of the EU numbering scheme, the residue may not be the actual residue 215 in that particular Protein S antibody. It may be actual residue number 213, or 214, or 215, or 216. Accordingly a skilled artisan will understand how to correspond the recited residue using the EU numbering scheme, to the actual residue in a Protein S antibody of the disclosure. The EU numbering system for antibodies is known in the art and is described, for example, at imgt.org/IMGTScientificChart/Numbering/Hu_IGHGnber.html.

[0087] In some embodiments, the Fc domain of a Protein S antibody is an IgG1 or IgG4 human Fc domain, and Fc variants comprise at least one amino acid substitution at a position selected from the group consisting of: 215, 221, 222, 228, 234, 235, 236, 239, 240, 241, 243, 244, 245, 247, 250, 252, 254, 256, 262, 263, 264, 265, 266, 267, 268, 269, 270, 292, 296, 297, 298, 299, 300, 305, 313, 324, 325, 326, 327, 328, 329, 330, 332, 333, 334, 345, 396, 428, 430, 433, 434, and 440, wherein the position numbers of the amino acid residues are of the EU numbering scheme.

[0088] In some embodiments, the Fc domain of a Protein S antibody is a human IgG1, and substitutions are introduced to reduce effector function, including N297A, N297Q, N297G, L235E, L234A, and L235A, wherein the position numbers of the amino acid residues are of the EU numbering scheme. In some embodiments, the Fc domain of a full-length Protein S antibody is human IgG4, and substitutions are introduced to reduce effector function, including

L235E, and F234A/L235A, wherein the position numbers of the amino acid residues are of the EU numbering scheme. In some embodiments, the Fc domain of a full-length Protein S antibody is human IgG2, and substitutions are introduced to reduce effector function, including H268Q/V309L/A330S/P331S and V234A/G237A/P238S/H268A/V309L/A330S/P331S, wherein the position numbers of the amino acid residues are of the EU numbering scheme.

[0089] In some embodiments, the Fc domain of a Protein S antibody is a human IgG1, and substitutions are introduced to increase effector function, including G236A/S239D/I332E, K326W/E333S, S267E/H268F/S324T, and E345R/E430G/S440Y, wherein the position numbers of the amino acid residues are of the EU numbering scheme.

[0090] In some embodiments, the Fc domain of a Protein S antibody is an IgG4 human Fc domain, and the antibody is prone to the dynamic process of Fab-arm exchange. Accordingly, in some embodiments the IgG4 Fc domain comprises a S228P substitution, resulting in the reduction of this process, wherein the position number of the amino acid residues are of the EU numbering scheme.

[0091] In other embodiments, the Fc domain of a Protein S antibody is altered to increase its serum half-life. Such alterations include substitutions of a human IgG1 (e.g. SEQ ID NO: 217) such as T250Q/M428L, M252Y/S254T/T256E, M428L/N434S, S267E/L328F, N325S/L328F, and H433K/N434F, wherein the position number of the amino acid residues are of the EU numbering scheme.

[0092] In other embodiments, the Fc domain of a Protein S antibody is altered to increase its serum half-life. Such alterations include substitutions of a human IgG4 (e.g. SEQ ID NO: 218) such as T250Q/M428L, M252Y/S254T/T256E, M428L/N434S, S267E/L328F, N325S/L328F, and H433K/N434F, wherein the position number of the amino acid residues are of the EU numbering scheme.

[0093] In some embodiments, the Fc domain of a Protein S antibody is an IgG1 human Fc domain, and substitutions are introduced enhance effector function, including F243L/R292P/Y300L/V305I/P396L, S239D/I332E, S298A/E333A/K334A, L234Y/L235Q/G236W/S239M/H268D/D270E/S298A, D270E/K326D/A330M/K334E, wherein the position number of the amino acid residues are of the EU numbering scheme.

Exemplary Protein S Antibodies – CDR Sequences

[0094] Provided herein are exemplary CDR sequences of the Protein S antibodies disclosed herein.

[0095] Exemplary CDR sequences presented in Tables 1A-1C and 2A-2C below. As referred below, a light chain variable (VL) domain CDR1 region is referred to as CDR-L1; a VL CDR2 region is referred to as CDR-L2; a VL CDR3 region is referred to as CDR-L3; a heavy chain variable (VH) domain CDR1 region is referred to as CDR-H1; a VH CDR2 region is referred to as CDR-H2; and a VH CDR3 region is referred to as CDR-H3.

Table 1A: Exemplary Protein S Antibody CDR-L1 Sequences
KLGDKY (SEQ ID NO: 1)
SLRNYY (SEQ ID NO: 2)
SSDVGGYEF (SEQ ID NO: 3)
QSVSIY (SEQ ID NO: 4)
QRINSN (SEQ ID NO: 5)
QSLHNSGYNY (SEQ ID NO: 6)
TGAVTASNY (SEQ ID NO: 9)
QSVTSN (SEQ ID NO: 10)
QSLVHSDGNTY (SEQ ID NO: 11)
QGINNY (SEQ ID NO: 117)
QSISTF (SEQ ID NO: 127)
QSVGSSY (SEQ ID NO: 136)
QNIHMW (SEQ ID NO: 141)
QSISSY (SEQ ID NO: 174)
NIGGKS (SEQ ID NO: 184)
KLGDKY (SEQ ID NO: 194)

KLGDKY (SEQ ID NO: 204)

Table 1B: Exemplary Protein S Antibody CDR-L2 Sequences
--

QDT (SEQ ID NO: 12)

GKN (SEQ ID NO: 13)

DVS (SEQ ID NO: 14)

QNS (SEQ ID NO: 15)

DAS (SEQ ID NO: 16)

GAS (SEQ ID NO: 17)

LGS (SEQ ID NO: 18)

STN (SEQ ID NO: 19)

KIS (SEQ ID NO: 20)

AAS (SEQ ID NO: 118)

ATS (SEQ ID NO: 128)

KTS (SEQ ID NO: 142)

AAS (SEQ ID NO: 175)

DDS (SEQ ID NO: 185)

QDS (SEQ ID NO: 195)

QDN (SEQ ID NO: 205)

Table 1C: Exemplary Protein S Antibody CDR-L3 Sequences
--

QAWDSNTVV (SEQ ID NO: 21)

NSRDSSGNHVV (SEQ ID NO: 22)

SSYTRSSTVV (SEQ ID NO: 23)

QAWDSSTWV (SEQ ID NO: 24)
QQRSNWPLT (SEQ ID NO: 25)
QQYDNWPLT (SEQ ID NO: 26)
MQALQTFT (SEQ ID NO: 27)
ALWYSDHFV (SEQ ID NO: 30)
QQYNNWPT (SEQ ID NO: 31)
MQATQFPHLT (SEQ ID NO: 32)
QQYNSYPRT (SEQ ID NO: 119)
QQYNSYPIT (SEQ ID NO: 123)
QQSYSTPRT (SEQ ID NO: 129)
QQYGSSPYT (SEQ ID NO: 137)
LQGQSYPT (SEQ ID NO: 143)
QQSYSSLT (SEQ ID NO: 176)
QVWEITSDHPA (SEQ ID NO: 186)
QAWDSSTVG (SEQ ID NO: 196)
QAWDSSTAV (SEQ ID NO: 206)

Table 2A: Exemplary Protein S Antibody CDR-H1 Sequences
GGSISSSSYY (SEQ ID NO: 33)
GGTFSSYS (SEQ ID NO: 34)
GGSITSDGYH (SEQ ID NO: 35)
GFTFDDYA (SEQ ID NO: 36)
GFTFSTYG (SEQ ID NO: 37)
GYSISSGYY (SEQ ID NO: 38)
GDTFNHA (SEQ ID NO: 39)

GHTFTGYY (SEQ ID NO: 42)
GGISSTNW (SEQ ID NO: 43)
GGISNYY (SEQ ID NO: 44)
GGITNSNYY (SEQ ID NO: 120)
GFTFSSYN (SEQ ID NO: 124)
GGISGNY (SEQ ID NO: 130)
GDSVSNNAA (SEQ ID NO: 138)
GYFTNHW (SEQ ID NO: 144)
GISFSAW (SEQ ID NO: 179)
GFTFSSYS (SEQ ID NO: 189)
GYFTNYY (SEQ ID NO: 199)
GYFTSYY (SEQ ID NO: 209)

Table 2B: Exemplary Protein S Antibody CDR-H2 Sequences
IYYSGNT (SEQ ID NO: 45)
IPIFGTT (SEQ ID NO: 46)
IYYTGNT (SEQ ID NO: 47)
ITWNSGNI (SEQ ID NO: 48)
IYYDGINK (SEQ ID NO: 49)
IYYSGST (SEQ ID NO: 50)
YIPIFGTT (SEQ ID NO: 51)
INPNSGDT (SEQ ID NO: 54)
IYQTGST (SEQ ID NO: 55)
IYYIGIT (SEQ ID NO: 56)
VYYSGTT (SEQ ID NO: 121)

ISSSSSYI (SEQ ID NO: 125)
TYYRSKWYN (SEQ ID NO: 139)
IYPGGGYT (SEQ ID NO: 145)
IKANPDGGTT (SEQ ID NO: 180)
ISSSTRTI (SEQ ID NO: 190)
ITPSGGTT (SEQ ID NO: 200)
TSPSGRST (SEQ ID NO: 210)

Table 2C: Exemplary Protein S Antibody CDR-H3 Sequences
ARCSGYGYSSGRSYFDY (SEQ ID NO: 57)
EGGRVGADFDY (SEQ ID NO: 58)
ARRLSTGPYFDY (SEQ ID NO: 59)
AKGRAVSDTFDI (SEQ ID NO: 60)
AESDL DY (SEQ ID NO: 61)
ATTYS DIVTGYYNDAFDI (SEQ ID NO: 62)
ARGGLAGSHYKNYYYDGMDV (SEQ ID NO: 63)
ARDSQILWFGELGY (SEQ ID NO: 66)
ARRFGELDY (SEQ ID NO: 67)
AALSGDHAFDI (SEQ ID NO: 68)
VRESESYYYYGSDV (SEQ ID NO: 122)
ARDEEWELLTGFDY (SEQ ID NO: 126)
ARDLDYFTWGAYSDWYFDL (SEQ ID NO: 131)
ARGSSWYRFFDY (SEQ ID NO: 140)
SRFGDQNWAWFAY (SEQ ID NO: 146)
TTELDILLWFTSFDY (SEQ ID NO: 181)

ARERSAFDY (SEQ ID NO: 191)
ARAGVQLDRRGWFDP (SEQ ID NO: 201)
ARGGVTIHLERRGYFDY (SEQ ID NO: 211)

[0096] In some embodiments, the Protein S antibodies provided herein include any one or more of the amino acid sequences of the CDR sequences provided in Tables 1A, 1B, 1C, 2A, 2B, and 2C.

[0097] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise:

- (a) any one of the CDR-L1 amino acid sequences of Table 1A;
- (b) any one of the CDR-L2 amino acid sequences of Table 1B;
- (c) any one of the CDR-L3 amino acid sequences of Table 1C;
- (d) any one of the CDR-H1 amino acid sequences of Table 2A;
- (e) any one of the CDR-H2 amino acid sequences of Table 2B; and
- (f) any one of the CDR-H3 amino acid sequences of Table 2C.

[0098] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of any one of SEQ ID NOs: 1-6, 9-11, 117, 127, 136, 141, 174, 184, 194, and 204;
- (b) a CDR-L2 comprising the amino acid sequence of any one of SEQ ID NOs: 12-20, 118, 128, 142, 175, 185, 195, and 205; and
- (c) a CDR-L3 comprising the amino acid sequence of any one of SEQ ID NOs: 21-27, 30-32, 119, 123, 129, 137, 143, 176, 186, 196, and 206.

[0099] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of any one of SEQ ID NOs: 33-39, 42-44, 120, 124, 130, 138, 144, 179, 189, 199, and 209;
- (b) a CDR-H2 comprising the amino acid sequence of any one of SEQ ID NOs: 45-51, 54-56, 121, 125, 139, 145, 180, 190, 200, and 210;
- (c) a CDR-H3 comprising the amino acid sequence of any one of SEQ ID NOs: 57-63, 66-68, 122, 126, 131, 140, 146, 181, 191, 201, and 211.

[0100] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 1;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 12; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 21.

[0101] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 2;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 13; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 22.

[0102] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 3;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 14; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 23.

[0103] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 1;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 15; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 24.

[0104] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 4;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 25.

[0105] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 5;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 17; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 26.

[0106] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 6;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 18; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 27.

[0107] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 9;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 30.

[0108] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 10;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 31.

[0109] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 11;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 20; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[0110] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 117;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 118; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 119.

[0111] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 117;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 118; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 123.

(d)

[0112] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 127;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 128; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 129.

[0113] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 136;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 17; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 137.

[0114] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 141;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 142; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 143.

[0115] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 174;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 175; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 176.

[0116] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 184;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 185; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 186.

[0117] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 194;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 195; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 196.

[0118] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 204;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 205; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 206.

[0119] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 33;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 45; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 57.

[0120] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 34;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 46; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 58.

[0121] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 35;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 47; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 59.

[0122] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 36;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 48; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 60.

[0123] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 37;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 49; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 61.

[0124] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 38;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 50; and

- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 62.

[0125] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 39;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 51; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 63.

[0126] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 42;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 54; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 66.

[0127] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 43;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 55; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 67.

[0128] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 44;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 56; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 68.

[0129] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 120;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 121; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 122.

[0130] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 124;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 125; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 126.

[0131] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 130;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 50; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 131.

[0132] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 138;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 139; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 140.

[0133] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 144;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 145; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 146.

[0134] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 179;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 180; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 181.

[0135] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 189;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 190; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 191.

[0136] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 199;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 200; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 201.

[0137] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 209;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 210; and

- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 211.

[0138] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 33;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 45; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 57.

[0139] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 34;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 46; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 58.

[0140] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 35;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 47; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 59.

[0141] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from

Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 36;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 48; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 60.

[0142] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 37;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 49; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 61.

[0143] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 38;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 50; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 62.

[0144] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 39;

- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 51; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 63.

[0145] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 42;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 54; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 66.

[0146] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises any one or more of the sequences provided in Tables 1A, 1B, and 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 43;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 55; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 67.

[0147] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 44;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 56; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 68.

[0148] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 120;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 121; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 122.

[0149] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 124;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 125; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 126.

[0150] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 130;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 50; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 131.

[0151] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 138;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 139; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 140.

[0152] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 144;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 145; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 146.

In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (d) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 179;
- (e) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 180; and
- (f) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 181.

[0153] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 189;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 190; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 191.

[0154] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 199;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 200; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 201.

[0155] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprises a CDR-L1 from Table 1A, a CDR-L2 from Table 1B, and a CDR-L3 from Table 1C, and wherein the heavy chain variable domain of the antibodies comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 209;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 210; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 211.

[0156] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 1;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 12; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 21.

[0157] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 2;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 13; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 22.

[0158] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 3;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 14; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 23.

[0159] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 1;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 15; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 24.

[0160] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 4;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 25.

[0161] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 5;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 17; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 26.

[0162] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 6;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 18; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 27.

[0163] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 9;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 30.

[0164] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 10;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 31.

[0165] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 11;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 20; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[0166] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 117;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 118; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 119.

[0167] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 117;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 118; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 123.

[0168] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 127;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 128; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 129.

[0169] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 136;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 17; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 137.

[0170] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 141;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 142; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 143.

[0171] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 174;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 175; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 176.

[0172] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 184;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 185; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 186.

[0173] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 194;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 195; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 196.

[0174] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprises a CDR-H1 from Table 2A, a CDR-H2 from Table 2B, and a CDR-H3 from Table 2C, and wherein the light chain variable domain of the antibodies comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 204;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 205; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 206.

[0175] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57.

[0176] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58.

[0177] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 3, SEQ ID NO: 14, SEQ ID NO: 23, SEQ ID NO: 35, SEQ ID NO: 47, and SEQ ID NO: 59.

[0178] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60.

[0179] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 4, SEQ ID NO: 16, SEQ ID NO: 25, SEQ ID NO: 37, SEQ ID NO: 49, and SEQ ID NO: 61.

[0180] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 5, SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 38, SEQ ID NO: 50, and SEQ ID NO: 62.

[0181] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 6, SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 39, SEQ ID NO: 51, and SEQ ID NO: 63.

[0182] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 9, SEQ ID NO: 19, SEQ ID NO: 30, SEQ ID NO: 42, SEQ ID NO: 54, and SEQ ID NO: 66.

[0183] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0184] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 11, SEQ ID NO: 20, SEQ ID NO: 32, SEQ ID NO: 44, SEQ ID NO: 56, and SEQ ID NO: 68.

[0185] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 117, SEQ ID NO: 118, SEQ ID NO: 119, SEQ ID NO: 120, SEQ ID NO: 121, and SEQ ID NO: 122.

[0186] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 117, SEQ ID NO: 118, SEQ ID NO: 123, SEQ ID NO: 124, SEQ ID NO: 125, and SEQ ID NO: 126.

[0187] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 127, SEQ ID NO: 128, SEQ ID NO: 129, SEQ ID NO: 130, SEQ ID NO: 50, and SEQ ID NO: 131.

[0188] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 136, SEQ ID NO: 17, SEQ ID NO: 137, SEQ ID NO: 138, SEQ ID NO: 139, and SEQ ID NO: 140.

[0189] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 141, SEQ ID NO: 142, SEQ ID NO: 143, SEQ ID NO: 144, SEQ ID NO: 145, and SEQ ID NO: 146.

[0190] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 174, SEQ ID NO: 175, SEQ ID NO: 176, SEQ ID NO: 179, SEQ ID NO: 180, and SEQ ID NO: 181.

[0191] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 184, SEQ ID NO: 185, SEQ ID NO: 186, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191.

[0192] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 194, SEQ ID NO: 195, SEQ ID NO: 196, SEQ ID NO: 199, SEQ ID NO: 200, and SEQ ID NO: 201.

[0193] In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise the CDR amino acid sequences of SEQ ID NO: 204, SEQ ID NO: 205, SEQ ID NO: 206, SEQ ID NO: 209, SEQ ID NO: 210, and SEQ ID NO: 211.

Exemplary Protein S Antibodies – Kappa and Lambda Light Chains

[0194] Provided herein are amino acid sequences for the kappa and lambda light chain constant regions of exemplary Protein S antibodies of the disclosure. Any of the Protein S antibodies provided herein (provided in the preceding section) may have a kappa light chain constant region or a lambda light chain constant region. The sequences of the kappa and lambda light chain constant regions are provided in Table 3.

Table 3

Kappa light chain sequence	RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWK VDNALQSGNSQESVTEQDSKDSTYLSSTLTLSKADYEKHK VYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 214)
Lambda light chain sequence	GQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAW KADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSHR SYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO: 215)

[0195] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 214 comprises CDRs having the amino acid sequences as set forth in SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58.

[0196] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 215 comprises CDRs having the amino acid sequences as set forth in SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58.

[0197] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 214 comprises CDRs having the amino acid sequences as set forth in SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0198] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 215 comprises CDRs having the amino acid sequences as set forth in SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0199] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 214 comprises CDRs having the amino acid sequences as set forth in SEQ ID NO: 174, SEQ ID NO: 175, SEQ ID NO: 176, SEQ ID NO: 179, SEQ ID NO: 180, and SEQ ID NO: 181.

[0200] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 215 comprises CDRs having the amino acid sequences as set forth in SEQ ID NO: 174, SEQ ID NO: 175, SEQ ID NO: 176, SEQ ID NO: 179, SEQ ID NO: 180, and SEQ ID NO: 181.

[0201] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 214 comprises the variable heavy and variable light chains having the amino acid sequences as set forth in SEQ ID NO: 71 and SEQ ID NO: 72.

[0202] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 215 comprises the variable heavy and variable light chains having the amino acid sequences as set forth in SEQ ID NO: 71 and SEQ ID NO: 72.

[0203] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 214 comprises the variable heavy and variable light chains having the amino acid sequences as set forth in SEQ ID NO: 89 and SEQ ID NO: 90.

[0204] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 215 comprises the variable heavy and variable light chains having the amino acid sequences as set forth in SEQ ID NO: 89 and SEQ ID NO: 90.

[0205] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 214 comprises the variable heavy and variable light chains having the amino acid sequences as set forth in SEQ ID NO: 177 and SEQ ID NO: 182.

[0206] In exemplary embodiments, the Protein S antibody having the light chain sequence of SEQ ID NO: 215 comprises the variable heavy and variable light chains having the amino acid sequences as set forth in SEQ ID NO: 177 and SEQ ID NO: 182.

Exemplary Protein S Antibodies – Variable Region Sequences

[0207] Provided herein are amino acid sequences for the variable domains of exemplary Protein S antibodies of the disclosure. The exemplary variable light chain amino acid sequences and exemplary variable heavy chain amino acid sequences are presented in Tables 4A and 4B below. Table 4C presents exemplary combinations of variable heavy and variable light chains.

[0208] Accordingly, in some embodiments, the Protein S antibodies of the disclosure comprise the variable chain amino acid sequence of any one of the combinations provided in Table 4C. In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise a variable light chain comprising the amino acid sequence selected from SEQ ID NO: 69, 71, 73, 75, 77, 79, 81, 87, 89, 91, 148, 150, 152, 156 158, 177, 187, 197, and 207. In some embodiments, provided herein are Protein S antibodies, wherein the antibodies comprise a variable heavy chain comprising the amino acid sequence selected from SEQ ID NO: 70, 72, 74, 76, 78, 80, 82, 88, 90, 92, 149, 151, 153, 157 159, 182, 192, 202, and 212.

Table 4A: Exemplary Variable Light Chain Amino Acid Sequences of Protein S Monoclonal Antibodies
SYELTQPPSVSVSPGQTASITCSGDKLGDKYACWYQQKPGQSPVLVIYQDTRK PSGIPERFSGSNSGNTATLTISGTQAMDEADYYCQAWDSNTVVFGGGTKLTV L (SEQ ID NO: 69)
SSDLTQGPAVSVALGQTVRITCQGDSLNRNYASWYQQKPGQAPVPVIYGKND RPSGIPDRFSGSISGNTASLTITGAQAEDEAHYYCNSRDSSGNHVVFGGGTKLT VL (SEQ ID NO: 71)
QSALTQPASVSGSPGQSITISCTGTSSDVGGYEFVSWYQHHPGKAPKLMIYDV SSRPSGVSNRFGSGKSGNTASLTISGLQAEDEADYYCSSYTRSSTVVFGGGARL TVL (SEQ ID NO: 73)
SYELNQPPSVSVSPGQTASITCSGDKLGDKYASWYQQKPGQSPVVAIYQNSK RPSGIPERFSASNSGNTATLTISGTQALDEADYYCQAWDSSTWVFGGGTKLTV L (SEQ ID NO: 75)

Table 4A: Exemplary Variable Light Chain Amino Acid Sequences of Protein S Monoclonal Antibodies
EIVLTQSPATLSLSPGERATLSCRASQSVSIYLAWYQQKPGQAPRLLIYDASNR ATGIPARFSGSGSGTDFTLTISLSEPEDFAVYYCQQRSNWPLTFGPGTKVDIK (SEQ ID NO: 77)
EIVMTQSPATLSVSPGERATLSCRASQRINSNLAWYQQKPGQAPRLLIYGAST RATGIPARFSGSGSGTEFTLTISLQSEDFAAYYCQQYDNWPLTFGGGTKVEIK (SEQ ID NO: 79)
DIVMTQSPLSLPVTGPGEASISCRSSQSLLSNGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQALQTFTFGPGT KVDIK (SEQ ID NO: 81)
QAVVTQESALTTSPGETVTLTCRSSTGAVTASNYANWVQEKPDHLFTGLIGST NNRAPGVPARFSGSLIGDKAALTITGAQTEDEAIYFCALWYSDHFVFGGGTKL TVL (SEQ ID NO: 87)
EIVMTQSPATLSVSPGERATLSCRASQSVTSNLAWYQQKPGQAPRLLIYDAST RATGIPARFSGSGSGTEFTLTISLQSEDFAIYYCQQYNNWPTFGQGTRLEIK (SEQ ID NO: 89)
DIVMTQTPLSPVTLGQPASISCRSSQSLVHSDGNTYLSWLQQRPGQPPRLLIY KISNRFSGVPDRFSGSGAGTDFTLKISRVEAEDVGVYYCMQATQFPHLTFGGG TKVEIK (SEQ ID NO: 91)
DIQMTQSPSSLSASVGDRVTITCRASQGINNYLAWFQQKPGKAPKSLIYAASS LQSGVPSKFSGSGSGTDFTLTISLQPEDFATYYCQQYNSYPRTFGQGTKVEIK (SEQ ID NO: 148)
DIQMTQSPSSLSASVGDRVTITCRASQGINNYLAWFQQKPGKAPKSLIYAASN LQSGVPLKFSGSGSGTDFTLTISLQPEDFATYYCQQYNSYPITFGQGTRLEIK (SEQ ID NO: 150)
DIQMTQSPSSLSASVGDRVTITCRASQSISTFLNWIYQQKPGKAPKLLIYATSSL RSGVPSRFSGSGSGTDFTLTISLQPEDFAIYYCQQSYSTPRTFGQGTQVEIK (SEQ ID NO: 152)
EIVLTQSPGTLSPGERATLSCRASQSVGSSYLAWYQQKPGQAPRLLISGASG RATGIPDRFSGSGSGTDFTLTISRLEPEDFTVYYCQQYGSSPYTFGQGTKLEIK (SEQ ID NO: 156)
DIQMNQSPSSLSASLGDTITITCRASQNIHMWLSWYQQKPGNIPKLLIFKTSNL HTGVPSRFSGSGSGTDFTLTISLQPEDATYYCLQGQSYPTFGGGTKLEIK (SEQ ID NO: 158)

Table 4A: Exemplary Variable Light Chain Amino Acid Sequences of Protein S Monoclonal Antibodies
DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSL QSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQSYSSLTFGQGTRLEIK (SEQ ID NO: 177)
SYVLTPPPSVSVAPGQTARITCGGDNIGGKSVHWYQQKPGQAPVMVYDDSD DRPSGIPERFAGSNSGNTATLAISRVEAGDEADYYCQVWEITSDHPAFGGGTR LTVL (SEQ ID NO: 187)
SYELTQPPSVSVSPGQTASITCSGDKLGDKYVFWYQQKPGQSPVLVIYQDSKR PSGIPERFSGSNSGNTATLTISGTQTMDEADYYCQAWDSSTVFGGGTKLAVL (SEQ ID NO: 197)
SYELTQPPSVSVSPGQTASITCSGDKLGDKYAFWYQQKPGQSPVLVIYQDNKR PSGIPERFSGSNSGNTATLTISGTQAVDEADYYCQAWDSSTAVFGGGTKLTVL (SEQ ID NO: 207)

Table 4B: Exemplary Variable Heavy Chain Amino Acid Sequences of Protein S Monoclonal Antibodies
QLQLQESGPGLVKPSETLSLTCTVSGGSISSSSYYWGWIRQPPGKGLEWIGNIY YSGNTYYNPSLKSRTISVDTSKNQFSLKLSSMTAADTAVYYCARCSGYGYS SGRSYFDYWQGQETLTVSS (SEQ ID NO: 70)
QVQLVQSGAEVKKPGSSVKVSCKVSGGTFSSYSISWVRQAPGQGLEWMGGII PIFGTTNYAQKFQGRVTITADESTSTAYMDLSSLKSEDTAMYYCEGGRVGAD FDYWGGQGLTVTVSS (SEQ ID NO: 72)
QVQLQESGPGLVKPSQTLSTCTVSGGSITSDGYHWSWIRQYPGKGLDWIGYI YYTGNTYYNPSLKSRTISVGTSQNQFSLKLISVTAADTAVYYCARRLSTGPY FDYWGGQGLTVTVSS (SEQ ID NO: 74)
EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSGI TWNSGNIGYADSVKGRFTISRDNKNSLYLHMNSLRIEDTAFYYCAKGRAVS DTFDIWGGQGMVTVSS (SEQ ID NO: 76)
QVQLVESGGGVVQPGRSLRLSCAASGFTFSTYGFHWVRQPPGKGLEWVAVIY YDGINKYYADSVKGRFTISRDNKNTLFLQMNSLRAEDTAVYYCAESDLDY WGQGLTVTVSS (SEQ ID NO: 78)
QVQLQESGPGLVKPSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLDWIGSIY YSGSTYYNPSLKSRTISVDTSKNQISLKLSSVTAADTAVYYCATTYSIVTGY YNDAFDIWGGQGMVTVSS (SEQ ID NO: 80)

Table 4B: Exemplary Variable Heavy Chain Amino Acid Sequences of Protein S Monoclonal Antibodies
QVQLVQSGAEVKKPGSSVKVSCASGDTFSNHAINWVRQAPGQGLEWMGG YIPFGTTNSAQKFRGRVTITADKSTNTAYMALSSLRSEDVAVYYCARGGLAG SHYKNYYYDGMVWGQGTTVTVSS (SEQ ID NO: 82)
QVQLVQSGAEVKKPGASVKVSCKSSGHTFTGYMHVVRQAPGQGLEWMG WINPNSGDTNYAQKFQGRVTMTRDTSISTAYMEMSRLRSDDTAVYYCARDS QILWFGELGYWGQGLTVTVSS (SEQ ID NO: 88)
QVQLQESGPGLVKPSETLSLTCTVSGGSISSTNWWVVRQPPGKGLEWIGEII QTGSTDYDPSLKSRTISIDKSKNQFSLKLYSVTAADTAVYYCARRFGELDY WGQGLTVTVSS (SEQ ID NO: 90)
QVQLQESGPGLVKPSETLSLTCTVSGGSISNYWNWVRQPPGKGLEWIGYIYY IGITDYNPSLKSRTISVDTSKNQFSLKVTSTVTAADTAVYYCAALSGDHAFDI WGQGLTVTVSS (SEQ ID NO: 92)
QLQLQESGPGLVKPSETLSLTCTVSGGSITNSNYWGWVRQPPGKGLEWIGSV YYSGTYYNPSLKSRTISVDPSKNQFSLKLSSVTAADTAVYYCVRESESYYY YGSDVWGQGTTVTVSS (SEQ ID NO: 149)
EVQLVESGGGLVKPGGSLRLSCAASGFTFSSYNMNWVRQAPGRGLDWVSSIS SSSSYIYYADSVKGRFTISRDNANKNSLYLQMNTLRAEDTAVYYCARDEEWEL LTGFDYWGQGLTVTVSS (SEQ ID NO: 151)
QVQLQESGPGLVKPSETLSLTCTVSGGSISGNYWSWVRQPPGKGLEWIGYIYY SGSTNYPNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARDLDYFTWG AYSDDYFDLWGRGTLTVTVSS (SEQ ID NO: 153)
QVQLQQSGPGLVKPSQTLSTCAISGDSVSNNAAWNVRQSPSRGLEWLGG TYYRSKWYNDYAVSVKSRIINPVTSKNQFSLQLNSVTPEDTAVYYCARGSS WYRFFDYWGQGLTVTVSS (SEQ ID NO: 157)
QVQLQQSGTELVRPGTSVKMSCKAAGYTFTNHWIGWVKQRPBGHGLEWIGDI YPGGGYTNYNEKFKGKASLTADTSSTTAYMQLSSLTSEDSAIYYCSRFGDQN WAWFAYWGQGLTVTVSA (SEQ ID NO: 159)
EVQLVESGGGLVKPGGSLRLSCAASGISFSNAWMWVRQAPGKGLEWVGRI KANPDGGTDDYAAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTTELD ILLWFTSFDYWGQGLTVTVSS (SEQ ID NO: 182)
EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVAYIS SSTRIFYADSVKGRFTISRDNANKNSLYLQMNSLRDEDTAFYYCARERSAFDY WGQGLTVTVSS (SEQ ID NO: 192)

Table 4B: Exemplary Variable Heavy Chain Amino Acid Sequences of Protein S Monoclonal Antibodies
QVQLVQSGSEVKKPGASVKVSCKASGYTFTNYYIHWVRQAPGQGLEWMGII TPSGGTTSYAQKFQGRVTMTRDTSTNTVYMGLSSLRSEDAMYYCARAGVQ LDRRGWFDWPWGQGTLLTVSS (SEQ ID NO: 202)
QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYIHWVRQAPGQGLEWMGV TSPSGRSTSAQKFQGRVTMTRDTSTSAVYMDLDSLRS EDTAVYYCARGGV IHLERRGYFDYWGGTLLVIVSS (SEQ ID NO: 212)

Table 4C: Exemplary Variable Light Chain and Variable Heavy Chain Amino Acid Sequence Combinations of Protein S Monoclonal Antibodies		
Combination Number	Variable Light Chain Amino Acid Sequence	Variable Heavy Chain Amino Acid Sequence
Combination 1	SEQ ID NO: 69	SEQ ID NO: 70
Combination 2	SEQ ID NO: 71	SEQ ID NO: 72
Combination 3	SEQ ID NO: 73	SEQ ID NO: 74
Combination 4	SEQ ID NO: 75	SEQ ID NO: 76
Combination 5	SEQ ID NO: 77	SEQ ID NO: 78
Combination 6	SEQ ID NO: 79	SEQ ID NO: 80
Combination 7	SEQ ID NO: 81	SEQ ID NO: 82
Combination 10	SEQ ID NO: 87	SEQ ID NO: 88
Combination 11	SEQ ID NO: 89	SEQ ID NO: 90
Combination 12	SEQ ID NO: 91	SEQ ID NO: 92
Combination 13	SEQ ID NO: 148	SEQ ID NO: 149
Combination 14	SEQ ID NO: 150	SEQ ID NO: 151
Combination 15	SEQ ID NO: 152	SEQ ID NO: 153
Combination 17	SEQ ID NO: 156	SEQ ID NO: 157
Combination 18	SEQ ID NO: 158	SEQ ID NO: 159

Table 4C: Exemplary Variable Light Chain and Variable Heavy Chain Amino Acid Sequence Combinations of Protein S Monoclonal Antibodies		
Combination 19	SEQ ID NO: 177	SEQ ID NO: 182
Combination 20	SEQ ID NO: 187	SEQ ID NO: 192
Combination 21	SEQ ID NO: 197	SEQ ID NO: 202
Combination 22	SEQ ID NO: 207	SEQ ID NO: 212

[0209] In some embodiments, the heavy and light chain variable domains of the Protein S antibodies provided herein include the amino acid sequence of any one of the numbered combinations presented in Table 4C.

[0210] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 69 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0211] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 70, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0212] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 69 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 70, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57.

[0213] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 71 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0214] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 72, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0215] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 71 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 72, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58.

[0216] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 73 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0217] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 74, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0218] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 73 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%,

90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 74, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 3, SEQ ID NO: 14, SEQ ID NO: 23, SEQ ID NO: 35, SEQ ID NO: 47, and SEQ ID NO: 59.

[0219] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 75 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0220] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 76, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0221] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 75 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 76, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60.

[0222] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 77 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0223] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 78, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0224] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 77 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 78, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 4, SEQ ID NO: 16, SEQ ID NO: 25, SEQ ID NO: 37, SEQ ID NO: 49, and SEQ ID NO: 61.

[0225] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 79 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0226] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 80, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0227] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 79 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 80, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino

acid sequences of SEQ ID NO: 5, SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 38, SEQ ID NO: 50, and SEQ ID NO: 62.

[0228] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 81 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0229] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 82, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0230] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 81 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 82, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 6, SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 39, SEQ ID NO: 51, and SEQ ID NO: 63.

[0231] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 87 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0232] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 88, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0233] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 87 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 88, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 9, SEQ ID NO: 19, SEQ ID NO: 30, SEQ ID NO: 42, SEQ ID NO: 54, and SEQ ID NO: 66.

[0234] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 89 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0235] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 90, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0236] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 89 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 90, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0237] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 91 or

an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0238] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 92, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0239] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 91 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 92, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 11, SEQ ID NO: 20, SEQ ID NO: 32, SEQ ID NO: 44, SEQ ID NO: 56, and SEQ ID NO: 68.

[0240] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 148 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0241] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 149, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0242] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 148 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid

sequence of SEQ ID NO: 149, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 117, SEQ ID NO: 118, SEQ ID NO: 119, SEQ ID NO: 120, SEQ ID NO: 121, and SEQ ID NO: 122.

[0243] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 150 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0244] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 151, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0245] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 150 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 151, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 117, SEQ ID NO: 118, SEQ ID NO: 123, SEQ ID NO: 124, SEQ ID NO: 125, and SEQ ID NO: 126.

[0246] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 152 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0247] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 153,

or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0248] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 152 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 153, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 127, SEQ ID NO: 128, SEQ ID NO: 129, SEQ ID NO: 130, SEQ ID NO: 50, and SEQ ID NO: 131.

[0249] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 156 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0250] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 157, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0251] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 156 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 157, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 136, SEQ ID NO: 17, SEQ ID NO: 137, SEQ ID NO: 138, SEQ ID NO: 139, and SEQ ID NO: 140.

[0252] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 158 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0253] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 159, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0254] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 158 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 159, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 141, SEQ ID NO: 142, SEQ ID NO: 143, SEQ ID NO: 144, SEQ ID NO: 145, and SEQ ID NO: 146.

[0255] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 177 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0256] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 182, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0257] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 177 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%,

89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 182, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 174, SEQ ID NO: 175, SEQ ID NO: 176, SEQ ID NO: 179, SEQ ID NO: 180, and SEQ ID NO: 181.

[0258] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 187 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0259] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 192, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0260] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 187 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 192, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 184, SEQ ID NO: 185, SEQ ID NO: 186, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191.

[0261] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 197, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0262] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 202, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0263] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 197 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 202, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR amino acid sequences of SEQ ID NO: 194, SEQ ID NO: 195, SEQ ID NO: 196, SEQ ID NO: 199, SEQ ID NO: 200, and SEQ ID NO: 201.

[0264] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 207, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0265] In some embodiments, provided herein are Protein S antibodies, wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 212, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0266] In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 207, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 212, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto. In such embodiments, the Protein S antibodies may comprise the CDR

amino acid sequences of SEQ ID NO: 204, SEQ ID NO: 205, SEQ ID NO: 206, SEQ ID NO: 209, SEQ ID NO: 210, and SEQ ID NO: 211.

[0267] As noted above, Protein S comprises four domains: the γ -carboxy-glutamic acid domain (Gla-domain), the thrombin-sensitive region (TSR), the epidermal growth factor-like domain (EGF domain), and the sex hormone binding globulin-like domain (SHBG domain). FIG. 1 depicts the schematic diagram of Protein S showing these modular domains of Protein S. The TSR is within the heavy chain of Protein S. The heavy chain of Protein S represents amino acids 42-296 of Protein S, the TSR represents amino acids 88-116, the signal peptide represents amino acids 1-24 and the propeptide represents amino acids 25-41.

[0268] In some embodiments, the Protein S antibodies provided herein bind to the Gla domain of Protein S. In some embodiments, the Protein S antibodies provided herein bind to the Gla domain and inhibit the cofactor activity of Protein S for both APC and TFPI. Exemplary antibodies that bind to the Gla domain and inhibit the cofactor activity of Protein S for both APC and TFPI may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 77 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 78; or such antibodies may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 4, SEQ ID NO: 16, SEQ ID NO: 25, SEQ ID NO: 37, SEQ ID NO: 49, and SEQ ID NO: 61.

[0269] In some embodiments, the Protein S antibodies provided herein bind to EGF domain of Protein S. In some embodiments, the Protein S antibodies provided herein bind to the SHBG-like domain of Protein S. In some embodiments, the Protein S antibodies provided herein bind to the C-terminal region of Protein S. In some embodiments, the Protein S antibodies provided herein bind the C-terminal region of Protein S, and inhibit the cofactor activity of Protein S for TFPI.

[0270] In some embodiments, the Protein S antibodies provided herein bind to Protein S fragments. The Protein S fragments are referred to herein as the Protein S heavy chain when they are expressed recombinantly in a cell line, such as HEK293 cells, for example. The Protein S heavy chain comprises amino acids 42-296. In some embodiments, the Protein S antibodies provided herein bind the Protein S heavy chain. Exemplary antibodies that bind to the Protein S heavy chain may comprise the light chain variable domain comprising the amino acid sequence

of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57. As another example, antibodies that bind to the Protein S heavy chain may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 79 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 80, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 5, SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 38, SEQ ID NO: 50, and SEQ ID NO: 62.

[0271] In some embodiments, the Protein S antibodies provided herein bind to the Protein S heavy chain comprising the TSR. In some embodiments, the Protein S antibodies provided herein bind to the Protein S heavy chain not comprising the TSR.

[0272] In some embodiments, the Protein S antibodies provided herein bind to the TSR of Protein S. In some embodiments, the Protein S antibodies provided herein do not bind to the TSR. In some embodiments, the Protein S antibodies provided herein bind to the heavy chain of Protein S, but do not bind the TSR region of the heavy chain. In some embodiments, the Protein S antibodies provided are dual inhibitors of APC and TFPI, and bind to the TSR of Protein S. In some embodiments, the Protein S antibodies provided are dual inhibitors of APC and TFPI, and do not bind to the TSR of Protein S. In some embodiments, the Protein S antibodies provided are inhibitors of APC, and do not bind to the TSR of Protein S. In some embodiments, the Protein S antibodies provided are inhibitors of TFPI, and do not bind to the TSR of Protein S.

[0273] Exemplary antibodies that bind to the TSR and are dual inhibitors of APC and TFPI may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67. As another example, antibodies that bind to the TSR and are dual inhibitors of APC and TFPI may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60.

[0274] In some embodiments, the Protein S antibodies provided herein do not bind to the TSR of Protein S, and cause a dual inhibition of the activity of APC and TFPI. In some embodiments, the Protein S antibodies provided herein do not bind to the TSR of Protein S, and cause an inhibition of the activity of APC. In some embodiments, the Protein S antibodies provided herein do not bind to the TSR of Protein S, and cause an inhibition of the activity of TFPI.

[0275] Exemplary antibodies that do not bind to the TSR and are dual inhibitors of APC and TFPI may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 73 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 74, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 3, SEQ ID NO: 14, SEQ ID NO: 23, SEQ ID NO: 35, SEQ ID NO: 47, and SEQ ID NO: 59.

[0276] In some embodiments, the Protein S antibodies provided herein bind to a linear epitope of Protein S. Exemplary antibodies that bind to a linear epitope of Protein S may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 77 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 78, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 4, SEQ ID NO: 16, SEQ ID NO: 25, SEQ ID NO: 37, SEQ ID NO: 49, and SEQ ID NO: 61.

[0277] In some embodiments, the Protein S antibodies provided herein bind to a conformational epitope, i.e., an epitope that is non-linear. Exemplary antibodies that bind to a non-linear epitope and inhibit the cofactor activity of Protein S for APC may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 91 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 92, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 11, SEQ ID NO: 20, SEQ ID NO: 32, SEQ ID NO: 44, SEQ ID NO: 56, and SEQ ID NO: 68. Exemplary antibodies that bind to a non-linear epitope and inhibit the cofactor activity of Protein S for APC and TFPI may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of 90, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0278] As discussed above, Protein S can be found in a free form, or in a complexed form. Protein S can form a complex with at least TFPI, or with C4b-binding protein (C4BP). In some

embodiments, the Protein S antibodies provided herein are capable of binding to Protein S in a free form, but not when in complexed form. In some embodiments, the Protein S antibodies provided herein are capable of binding to Protein S when Protein S is complexed, but not in free form. In some embodiments, the Protein S antibodies provided herein are capable of binding to Protein S that is either in free or complexed form. In some embodiments, the Protein S antibodies provided herein bind to a complexed Protein S, wherein the Protein S is bound to TFPI. In some embodiments, the Protein S antibodies provided herein bind to a complexed Protein S, wherein the Protein S is bound to C4BP.

[0279] In some embodiments, the Protein S antibodies provided herein bind Protein S, and the binding affinity of the antibodies to Protein S is calcium-dependent. In some embodiments, the Protein S antibodies provided herein bind Protein S, and the binding affinity of the antibodies to Protein S is not calcium-dependent. In some embodiments, the Protein S antibodies provided herein are dual inhibitors of APC and TFPI, and the binding affinity of the antibodies to Protein S is calcium-dependent. In some embodiments, the Protein S antibodies provided herein are dual inhibitors of APC and TFPI, and the binding affinity of the antibodies to Protein S is calcium-independent. In some embodiments, the Protein S antibodies provided herein are inhibitors of APC, and the binding affinity of the antibodies to Protein S is calcium-dependent. In some embodiments, the Protein S antibodies provided herein are inhibitors of APC, and the binding affinity of the antibodies to Protein S is calcium-independent. In some embodiments, the Protein S antibodies provided herein are inhibitors of TFPI, and the binding affinity of the antibodies to Protein S is calcium-independent. In some embodiments, the Protein S antibodies provided herein are inhibitors of TFPI, and the binding affinity of the antibodies to Protein S is calcium-dependent.

[0280] Exemplary antibodies that are dual inhibitors of the Protein S cofactor activity of APC and TFPI, wherein the binding affinity of the antibodies to Protein S is calcium-independent, may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of 72, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58.

[0281] Exemplary antibodies that are inhibitors of the Protein S cofactor activity of TFPI, wherein the wherein the binding affinity of the antibodies to Protein S is calcium-independent,

may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 87 and the heavy chain variable domain comprising the amino acid sequence of 88, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 9, SEQ ID NO: 19, SEQ ID NO: 30, SEQ ID NO: 42, SEQ ID NO: 54, and SEQ ID NO: 66.

[0282] Exemplary antibodies that are inhibitors of the Protein S cofactor activity of APC, wherein the wherein the binding affinity of the antibodies to Protein S is calcium-dependent, may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 91 and the heavy chain variable domain comprising the amino acid sequence of 92, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 11, SEQ ID NO: 20, SEQ ID NO: 32, SEQ ID NO: 44, SEQ ID NO: 56, and SEQ ID NO: 68.

[0283] Exemplary antibodies that are dual inhibitors of the Protein S cofactor activity of APC and TFPI, wherein the wherein the binding affinity of the antibodies to Protein S is calcium-dependent, may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of 90, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

B. Generation of Protein S Antibodies

[0284] Production of the Protein S antibodies provided herein may be by use of any method known to those of ordinary skill in the art. In some embodiments, the antibodies are produced by hybridomas. In some embodiments, the antibodies are encoded by a nucleic acid and are expressed, purified, and isolated.

[0285] Accordingly, provided herein are nucleic acids encoding any of the antibodies disclosed herein, vectors comprising any of the nucleic acids encoding such antibodies, and host cells comprising such vectors. Also provided herein are exemplary nucleic acid sequences encoding for the variable heavy chains and variable light chains of the Protein S antibodies disclosed herein.

[0286] Tables 5A and 5B provide exemplary variable light chain nucleic acid sequences and exemplary variable heavy chain nucleic acid sequences. Exemplary combinations of nucleic acid sequences encoding for the variable heavy and light chain domains of the Protein S antibodies

disclosed herein are presented in Table 5C. The exemplary amino acid sequences of Tables 4A-4C correspond to the nucleic acid sequences of Tables 5A-5C. The exemplary combinations of Table 5C correspond to the numbered combinations presented in Table 4C.

[0287] The person of ordinary skill in the art will appreciate that, because of redundancy in the triplet code, multiple nucleic acids may encode the same amino acid sequence. Thus, nucleic acid sequences that are not identical to those set forth in the tables below may still encode the Protein S antibodies of the disclosure.

Table 5A: Exemplary Variable Light Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
TCCTATGAGCTGACTCAGCCACCCTCAGTGTCCGTGTCCCCAGGACAGACA GCCAGCATCACCTGCTCTGGAGATAAATTGGGGGATAAATATGCTTGCTG GTATCAGCAGAAGCCAGGCCAGTCCCCTGTACTGGTCATCTATCAAGATA CTAAGCGGCCCTCAGGGATCCCTGAGCGATTCTCTGGCTCCAACTCTGGGA ACACAGCCACTCTGACCATCAGCGGGACCCAGGCTATGGATGAGGCTGAC TATTACTGTCAGGCGTGGGACAGCAACACTGTGGTCTTCGGCGGAGGGAC CAAGCTGACCGTCCTA (SEQ ID NO: 93)
TCCTCTGACCTGACTCAGGGCCCTGCTGTGTCTGTGGCCCTGGGACAGACA GTCAGGATCACATGCCAAGGAGACAGCCTCAGAACTATTATGCAAGCTG GTACCAGCAGAAGCCAGGACAGGCCCTGTACCTGTCATCTATGGTAAAA ACGACCGGCCCTCAGGGATCCCAGACCGATTCTCTGGCTCCATCTCAGGA AACACAGCTTCCTTGACCATCACTGGGGCTCAGGCGGAAGATGAGGCTCA CTATTACTGTAACCTCCCGGGACAGCAGTGGTAACCATGTGGTATTCGGCG GAGGGACCAAGCTGACCGTCCTG (SEQ ID NO: 95)
CAGTCTGCCCTGACTCAGCCTGCCTCCGTGTCTGGGTCTCCTGGACAGTCG ATCACCATCTCCTGCACTGGAACCAGCAGTGACGTTGGTGGTTATGAATTT GTCTCCTGGTACCAACATACCCAGGCAAAGCCCCAAACTCATGATTTAT GATGTCAGTAGTCGGGCCCTCAGGGGTTTCTAATCGCTTCTCTGGCTCCAAG TCTGGCAACACGGCCTCCCTGACCATCTCTGGGCTCCAGGCTGAGGACGA GGCTGATTACTGTCAGCTCATATACGCGCAGCAGCACTGTGGTGTTCGG CGGCGGGGCCAGGCTGACCGTCCTA (SEQ ID NO: 97)
TCCTATGAGCTGAATCAGCCACCCTCAGTGTCCGTGTCCCCAGGACAGAC AGCCAGCATCACCTGCTCTGGAGATAAATTGGGGGATAAATATGCTTCCT GGTATCAGCAGAAGCCAGGCCAGTCCCCTGTGGTGGCCATCTATCAAAAT AGCAAGCGGCCCTCAGGGATCCCTGAGCGATTCTCTGCCTCCAACTCTGG GAACACAGCCACTCTGACCATCAGCGGGACCCAGGCTTTGGATGAGGCTG ACTATTACTGTCAGGCGTGGGACAGCAGCACTTGGGTGTTCGGCGGAGGG ACCAAGCTGACCGTCCTA (SEQ ID NO: 99)

Table 5A: Exemplary Variable Light Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
GAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAA AGAGCCACCCTCTCCTGCAGGGGCCAGTCAGAGTGTTAGTATCTACTTAGCC TGGTACCAACAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGATGC ATCCAACAGGGCCACTGGCATCCCAGCCAGGTTCAAGTGGCAGTGGGTCTG GGACAGACTTCACTCTCACCATCAGCAGCCTAGAGCCTGAAGATTTTGA GTTTATTACTGTCAGCAGCGTAGCAACTGGCCCCCTCACTTTCGGCCCTGGG ACCAAAGTGGATATCAAA (SEQ ID NO: 101)
GAAATAGTGATGACGCAGTCTCCAGCCACCCTGTCTGTGTCTCCAGGGGA AAGAGCCACCCTCTCCTGCAGGGGCCAGTCAGAGGATTAACAGCAACTTAG CCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGGT GCATCCACCAGGGCCACTGGTATCCCCGCCAGGTTCAAGTGGCAGTGGGTCT TGGGACAGAGTTCACTCTCACCATCAGCAGCCTGCAGTCTGAAGATTTTGC AGCTTATTACTGTCAGCAGTATGATAACTGGCCGCTCACTTTCGGCGGAGG GACCAAGGTGGAGATCAAA (SEQ ID NO: 103)
GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAG CCGGCCTCCATCTCCTGCAGGTCTAGTCAGAGCCTCCTGCATAGTAATGGA TACAATATTTGGATTGGTACCTGCAGAAGCCAGGGCAGTCTCCACAGCT CCTGATCTATTTGGGTTCTAATCGGGCCTCCGGGGTCCCTGACAGGTTTCA TGGCAGTGGATCAGGCACAGATTTTACACTGAAAATCAGCAGAGTGGAGG CTGAGGATGTTGGGGTTTATTATTGTATGCAAGCTCTACAACTTTCACTT TCGGCCCTGGGACCAAAGTGGATATCAAA (SEQ ID NO: 105)
CAGGCTGTTGTGACTCAGGAATCTGCACTCACCACATCACCTGGTGAAAC AGTCACACTCACTTGTGCTCAAGTACTGGGGCTGTTACAGCTAGTAACTA TGCCAACTGGGTCCAAGAAAAACCAGATCATTGTTCACTGGTCTAATAG GTAGTACCAATAACCGAGCTCCAGGTGTTCTGCCAGATTCTCAGGCTCCC TGATTGGAGACAAGGCTGCCCTCACCATCACAGGGGCACAGACTGAGGAT GAGGCAATATATTTCTGTGCTCTATGGTACAGCGACCATTTCGTGTTCCGT GGAGGAACCAAAGTACTGTCTA (SEQ ID NO: 111)
GAAATAGTGATGACGCAGTCTCCAGCCACCCTGTCTGTGTCTCCAGGGGA AAGAGCCACCCTCTCCTGCAGGGGCCAGTCAGAGTGTTACCAGCAACTTAG CCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGAT GCATCCACCAGGGCCACTGGTATCCCAGCCAGGTTCAAGTGGCAGTGGGTCT TGGGACAGAGTTCACTCTCACCATCAGCAGCCTGCAGTCTGAAGATTTTGC AATTTATTACTGTCAGCAGTATAATAACTGGCCACCTTCGGCCAAGGGAC ACGACTGGAGATTAAA (SEQ ID NO: 113)
GATATTGTGATGACCCAGACTCCACTCTCCTCACCTGTCACCCCTGGACAG CCGGCCTCCATCTCCTGCAGGTCTAGTCAAAGCCTCGTACACAGTGATGGA AACACCTACTTGAGTTGGCTTCAGCAGAGGCCAGGCCAGCCTCCAAGACT CCTAATTTATAAGATTTCTAACCAGTTCTCTGGGGTCCCAGACAGATTCA TGGCAGTGGGGCAGGGACAGATTTCACTGAAAATCAGCAGGGTGGAA

Table 5A: Exemplary Variable Light Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
GCTGAGGATGTCGGGGTTTATTACTGCATGCAAGCTACACAATTTCCCATCTCACTTTCGGCGGAGGGACCAAGGTGGAGATCAAA (SEQ ID NO: 115)
GACATCCAGATGACCCAGTCTCCATCCTCACTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGTCGGGCGAGTCAGGGCATTAAACAATTATTTAGCTGGTTTCAGCAGAAACCAGGGAAAGCCCCTAAGTCCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAAGTTCAGCGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGCCTGAAGATTTTGCAACTTATTACTGCCAACAGTATAATAGTTACCCCTCGGACGTTTCGGCCAAGGGACCAAGGTGGAAATCAAA (SEQ ID NO: 162)
GACATCCAGATGACCCAGTCTCCATCCTCACTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGTCGGGCGAGTCAGGGCATTAAACAATTATTTAGCTGGTTTCAGCAGAAACCAGGGAAAGCCCCTAAGTCCCTGATCTATGCTGCATCCAATTTGCAAAGTGGGGTCCCATTAAGTTCAGCGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGCCTGAAGATTTTGCAACTTATTACTGCCAACAGTATAATAGTTACCCGATCACCTTCGGCCAAGGGACACGACTGGAGATTAAA (SEQ ID NO: 164)
GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGGCAAGTCAGAGCATTAGCACCTTTTTAAATTTGGTATCAGCAGAAACCAGGGAAAGCCCCTAAACTCCTGATCTATGCTACATCCAGTTTGCGAAGTGGGGTCCCATCAAAGTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCAATTTATTATTGTCAACAGAGTTACAGTACCCCTCGGACGTTTCGGCCAAGGGACCCAGGTGGAAATCAAA (SEQ ID NO: 166)
GAAATTGTGTTGACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCCTGCAGGGCCAGTCAGAGTGTTGGCAGCAGCTACTTAGCCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTCTGTGTCATCCGGCAGGGCCACTGGCATCCCAGACAGGTTCAAGTGGCAGTGGGTCTGGGACAGACTTCACTCTCACCATCAGCAGACTGGAGCCTGAAGATTTTACAGTGTATTACTGTCAGCAGTATGGTAGCTACCGTACACTTTTGGCCAGGGGACCAAGCTGGAGATCAAA (SEQ ID NO: 170)
GACATCCAGATGAACCAGTCTCCATCCAGTCTGTCTGCATCCCTCGGAGACACAATTACCATCACTTGCCGTGCCAGTCAGAACATTCATATGTGGTTAAGCTGGTACCAGCAGAAACCAGGAAATATTCCTAAACTATTGATCTTTAAGACTTCCAATTTGCACACAGGCGTCCCATCAAAGTTTAGTGGCAGTGGATCTGGAACAGATTTACATTAACCATCAGCAGTCTGCAGCCTGAAGACATTGCCACTTACTACTGTCTACAGGGTCAAAGTTATCCGTTCACGTTTCGGAGGGGGGACCAAGCTGGAAATAAAG (SEQ ID NO: 172)
GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGGCAAGTCAGAGCATTAGCAGCTATTTAAATTTGGTATCAGCAGAAACCAGGGAAAGCCCCTAAGTCCCTGATCTATGCTG

Table 5A: Exemplary Variable Light Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
CATCCAGTTTGC AAAGTGGGGTCCCATCAAGGTTCAAGTGGCAGTGGATCT GGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCTGAAGATTTTGCA ACTTACTACTGTCAACAGAGTTACAGTTCCCTCACCTTCGGCCAAGGGACA CGACTGGAGATTAAA (SEQ ID NO: 178)
TCCTATGTGCTGACTCAGCCACCCTCGGTGTCAGTGGCCCCAGGACAGAC GGCCAGGATTACCTGTGGGGGAGACAACATTGGAGGTAAAAGTGTGCACT GGTACCAGCAGAAGCCAGGCCAGGCCCTGTGATGGTCGTCTATGATGAT AGCGACCGGCCCTCAGGGATCCCTGAGCGATTCTGCTGGCTCCAATTCTGG GAACACGGCCACCCTGGCCATCAGCAGGGTCGAAGCCGGGGATGAGGCC GACTATTACTGTGTCAGGTGTGGGAGATAACTAGTGATCATCCGGCATTCTGG CGGAGGGACCAGGCTGACCGTCCTA (SEQ ID NO: 188)
TCCTATGAGCTGACTCAGCCACCCTCAGTGTCCGTGTCCCCAGGACAGACA GCCAGCATCACCTGCTCTGGAGATAAATTGGGGGATAAATATGTTTTCTGG TATCAGCAGAAGCCAGGCCAGTCCCCTGTGTTGGTCATCTATCAAGATAG CAAGCGGCCCTCAGGGATCCCTGAGCGATTCTCTGGCTCCAACTCTGGGA ACACAGCCACTCTGACCATCAGCGGGACCCAGACTATGGATGAGGCTGAC TATTACTGTGTCAGGCGTGGGACAGCAGCACTGTGGGATTCGGCGGAGGGAC CAAGCTGGCCGTCTCTG (SEQ ID NO: 198)
TCCTATGAGCTGACTCAGCCACCCTCAGTGTCCGTGTCCCCGGGACAGACA GCCAGCATCACCTGCTCTGGAGATAAATTGGGGGATAAATATGCTTTCTG GTATCAGCAGAAGCCAGGCCAGTCCCCTGTGCTGGTCATCTATCAAGATA ACAAGCGGCCCTCAGGGATCCCTGAGCGATTCTCTGGCTCCAACTCTGGG AACACAGCCACTCTAACCATCAGCGGGACCCAGGCTGTGGATGAGGCTGA CTATTACTGTGTCAGGCGTGGGACAGCAGCACTGCGGTATTCGGCGGAGGGGA CCAAGCTGACCGTCCTA (SEQ ID NO: 208)

Table 5B: Exemplary Variable Heavy Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
CAGCTGCAGCTGCAGGAGTCGGGCCAGGACTGGTGAAGCCTTCGGAGAC CCTGTCCCTCACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTAGTAGTTA CTACTGGGGCTGGATCCGCCAGCCCCGGGGAAGGGACTGGAGTGGATTG GGAATATCTATTATAGTGGGAACACCTACTACAACCCGTCCCTCAAGAGT CGAGTCACCATATCCGTAGACACGTCCAAGAACCAGTTCTCCCTGAAGCT GAGCTCTATGACCGCCGCAGACACGGCTGTGTATTACTGTGCGAGATGTA GTGGCTACGGGTATAGCAGTGGCCGGTCTACTTTGACTACTGGGGCCAG GAAACCCTGGTCACCGTCTCCTCA (SEQ ID NO: 94)
CAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGTCTC GGTGAAGGTCTCCTGCAAGGTTTCTGGAGGCACCTTCAGCAGCTATTCTAT

Table 5B: Exemplary Variable Heavy Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
CAGCTGGGTGCGACAGGCCCTGGACAAGGGCTTGAGTGGATGGGAGGG ATCATCCCTATATTTGGTACAACAACTACGCACAGAAGTTCCAGGGCAG AGTCACGATCACCGCGGACGAATCCACGAGCACAGCCTACATGGATCTGA GCAGCCTGAAATCTGAGGACACGGCCATGTATTACTGTGAGGGGGGTAGA GTGGGAGCGGACTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTC CTCA (SEQ ID NO: 96)
CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGAC CCTGTCCCTCACCTGCACTGTCTCTGGTGGCTCCATCACCAGTGATGGTTA CCACTGGAGCTGGATCCGCCAGTACCCAGGGAAGGGCCTGGACTGGATTG GATACATCTATTACACTGGGAACACCTACTACAACCCGTCCCTCAAGAGTC GAGTGACCATATCAGTAGGCACGTCTCAGAACCAGTTCTCCCTGAAGCTG ATCTCTGTGACTGCCGCGGACACGGCCGTTTATTACTGTGCGAGAAGGCTG TCGACTGGGCCCTACTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTC TCCTCC (SEQ ID NO: 98)
GAAGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGCAGGTC CCTGAGACTCTCCTGTGCAGCCTCTGGATTACCTTTGATGATTATGCCAT GCACTGGGTCCGGCAAGCTCCAGGGAAGGGCCTGGAATGGGTCTCAGGTA TTACTTGGAATAGTGGTAACATAGGCTATGCGGACTCTGTGAAGGGCCGA TTCACCATCTCCAGAGACAACGCCAAGAAGTCCCTGTATCTGCACATGAA CAGTCTGAGAATTGAGGACACGGCCTTCTATTACTGTGCAAAGGCCGAG CAGTGTCTGATACTTTTGATATCTGGGGCCAAGGGACAATGGTCACCGTCT CTTCA (SEQ ID NO: 100)
CAGGTGCAGTTGGTGGAAATCTGGGGGAGGCGTGGTCCAGCCTGGGAGGTC CCTGAGACTCTCCTGTGCAGCGTCTGGATTACCTTCAGTACCTATGGCTT TCACTGGGTCCGCCAGCCTCCAGGCAAGGGACTGGAGTGGGTGGCAGTTA TATATTATGATGGAATTAATAAATATTATGCAGACTCCGTGAAGGGCCGA TTCACCATCTCCAGAGACAATTCCAAGAACACGCTGTTTCTTCAAATGAAC AGCCTGAGAGCCGAGGACACGGCTGTGTATTACTGTGCGGAGTCCGACTT GGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCA (SEQ ID NO: 102)
CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGAC CCTGTCCCTCACCTGCGCTGTCTCTGGTTATTCCATCAGCAGTGGTTACTAC TGGGGCTGGATCCGGCAGCCCCAGGGAAGGGGCTGGACTGGATTGGGA GTATCTATTATAGTGGGAGTACCTACTACAACCCGTCCCTCAAGAGTCGAG TCACCATATCAGTTGACACGTCCAAGAACCAGATCTCCCTGAAGCTGAGC TCTGTGACCGCCGACACACGGCCGTGTATTACTGTGCGGACACGTATTCC GATATTGTGACTGGTTATTATAATGATGCTTTTGATATCTGGGGCCAAGGG ACAATGGTCACCGTGTCTTCA (SEQ ID NO: 104)
CAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGTCTC GGTGAAGGTCTCCTGTAAGGCTTCTGGAGACACCTTCAGCAACCATGCTAT CAACTGGGTGCGACAGGCCCTGGACAAGGGCTTGAGTGGATGGGAGGGT

Table 5B: Exemplary Variable Heavy Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
ACATCCCTATCTTTGGTACAACAACTCCGCACAGAAGTTCCGGGGCAGAGTCACGATTACCGCGGACAAATCCACGAACACAGCCTACATGGCGCTGAGCAGCCTGAGATCTGAGGACACGGCCGTTTATTACTGTGCGAGAGGGGGGCTCGCGGGGAGTCATTATAAGAACTACTACTATGACGGTATGGACGTCTGGGCCAGGGGACCACGGTCACCGTCTCCTCA (SEQ ID NO: 106)
CAGGTGCAACTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGGCCTCAGTGAAGGTCTCCTGCAAGTCTTCTGGCCACACCTTCACCGGCTACTATATGCACTGGGTGCGACAGGCCCCTGGACAAGGGCTTGAGTGGATGGGATGGATCAACCCTAACAGTGGTGCACAACTACGCACAGAAGTTTCAGGGCAGGTGTCACCATGACCAGGGACACGTCCATCAGCACAGCCTACATGGAGATGAGCAGGCTGAGATCTGACGACACGGCCGTGTATTACTGTGCGAGAGACTCCCAAATACTATGGTTTCGGGGAGTTAGGCTACTGGGGCCAGGGAACCTGGTCAACCGTCTCCTCC (SEQ ID NO: 112)
CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCACCTGCGGTGTCTCTGGTGGCTCCATCAGCAGTACTAAGTGTGGAGTTGGGTCCGCCAGCCCCCAGGGAAGGGGCTGGAGTGGATTGGGAAATCTATCAAAGTGGGAGTACCGACTACGACCCGTCCCTCAAGAGTCGAGTCAACCATATCAATAGACAAGTCCAAGAACCAGTTCTCCCTGAAGCTGTACTCTGTGACCGCCGCGGACACGGCCGTGTATTACTGTGCGAGAAGGTTCTGGGAGTTAGACTACTGGGGCCAGGGAACCTGGTCAACCGTCTCCTCA (SEQ ID NO: 114)
CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCACCTGCACTGTCTCTGGTGGCTCCATCAGTAATTACTACTGGAAGTGGATCCGGCAGCCCCCAGGGAAGGGACTGGAGTGGATTGGGTATATCTATTACATTGGGATCACCAGTACAACCCCTCCCTCAAGAGTCGAGTCAACCATATCAGTAGACACGTCCAAGAACCAGTTCTCCCTGAAGGTGACCTCTGTGACCGCTGCGGACACGGCCGTGTATTACTGTGCGGCTCTAAGTGGGATCATGCTTTTGACATCTGGGGCCAAGGGACACTGGTCAACCGTCTCTTCA (SEQ ID NO: 116)
CAGCTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTCACCTGCACTGTCTCTGGTGGCTCCATCACCAATAGTAATTACTACTGGGGCTGGATCCGCCAGCCCCCAGGGAAGGGACTGGAGTGGATTGGAGTGTCTATTATAGTGGGACCACCTACTACAACCCGTCCCTCAAGAGTCGAGTCACCATATCCGTAGACCCGTCCAAGAACCAGTTCTCCCTGAAGCTGAGCTCTGTGACCGCCGCGAGACACGGCTGTGTATTACTGTGTGAGAGAGAGTGAGAGCTACTACTACGGTTTCGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCA (SEQ ID NO: 163)
GAGGTGCAGCTGGTTGAGTCTGGGGGAGGCCTGGTCAAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAGCCTCTGGATTACCTTCAGTAGCTATAACATGAAGTGGGTCCGCCAGGCTCCAGGGAGGGGGCTGGACTGGGTCTCATCCATTAGTAGTAGTAGTAGTTACATATACTACGCAGACTCAGTGAAGGGCCGA

Table 5B: Exemplary Variable Heavy Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
TTCACCATCTCCAGAGACAACGCCAAGAACTCACTGTATCTGCAAATGAA TACCCTGAGAGCCGAGGACACGGCTGTTTATTACTGTGCGAGAGATGAGG AGTGGGAGCTACTGACGGGCTTTGACTACTGGGGCCAGGGAACCCTGGTC ACCGTCTCCTCA (SEQ ID NO: 165)
CAGGTGCAGCTGCAGGAGTCGGGCCAGGACTGGTGAAGCCTTCGGAGAC CCTGTCCCTCACCTGCACTGTCTCTGGTGGCTCCATCAGTGGTAACTACTG GAGCTGGATCCGGCAGCCCCAGGGAAGGGACTGGAGTGGATTGGGTATA TCTATTACAGTGGGAGCACCAACTACAATCCCTCCCTCAAGAGTCGAGTC ACCATATCAGTTGACACGTCCAAGAACCAGTTCTCCCTGAAGCTGAGCTCT GTGACCGCTGCGGATACGGCCGTGTATTACTGTGCGAGAGATCTTGATTAC TTTACTTGGGGGGCTTATTCTGACTGGTACTTCGATCTCTGGGGCCGTGGC ACCCTGGTCACTGTCTCCTCA (SEQ ID NO: 167)
CAGGTACAGCTGCAGCAGTCAGGTCCAGGACTGGTGAAGCCCTCGCAGAC CCTCTCACTCACCTGTGCCATCTCCGGGGACAGTGTCTCTAACAACAATGC TGCTTGGAACTGGATCAGGCAGTCCCCATCGAGAGGCCTTGAGTGGCTGG GAGGGACATACTACAGGTCCAAGTGGTATAATGATTATGCAGTATCTGTG AAAAGTCGAATAATCATCAACCCAGTCACATCCAAGAACCAGTTCTCCCT ACAGCTGAACTCTGTGACTCCCGAGGACACGGCTGTGTATTACTGTGCAA GAGGCAGCAGCTGGTACAGGTTTTTTGACTACTGGGGCCAGGGAACCCTG GTCACCGTCTCCTCA (SEQ ID NO: 171)
CAGGTCCAGCTGCAGCAGTCTGGAAGTGAAGCTGGTAAGGCCTGGGACTTC AGTGAAGATGTCCTGTAAGGCTGCTGGATACACCTTCACTAACCCTGGA TAGGTTGGGTAAAGCAGAGGCCTGGACATGGCCTTGAGTGGATTGGAGAT ATTTACCCTGGAGGTGGTTATACTAACTACAATGAGAAGTTCAAGGGCAA GGCCTCACTGACTGCAGACACATCCTCCACCACAGCCTACATGCAGCTCA GCAGCCTGACATCTGAGGACTCTGCCATCTATTACTGTTCAAGATTCGGGG ATCAAAACTGGGCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCACT GTCTCTGCA (SEQ ID NO: 173)
GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTAAAGCCTGGGGGGTC CCTTAGACTCTCCTGTGCAGCCTCTGGAATCAGTTTCAGTAACGCCTGGAT GAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAATGGGTTGGCCGTA TTAAAGCCAATCCTGATGGTGGGACAACAGACTACGCTGCACCCGTGAAA GGCAGATTACCATCTCAAGAGATGATTCAAAAAACACGCTATATCTGCA AATGAACAGCCTGAAAACCGAGGACACAGCCGTGTATTACTGTACCACAG AGTTGGACATTTTACTATGGTTACCTCCTTTGACTACTGGGGCCAGGGAA CCCTGGTCACCGTCTCCTCA (SEQ ID NO: 183)
GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTC CCTGAGACTCTCCTGTGCAGCCTCTGGATTACCTTCAGTAGCTATAGCAT GAACTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTTGCATACA TTAGTAGTAGTACTCGTACCATATTCTACGCAGACTCTGTGAAGGGCCGAT TCACCATCTCCAGAGACAATGCCAAGAACTCACTGTATCTGCAAATGAAC

Table 5B: Exemplary Variable Heavy Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies
AGCCTGAGAGACGAGGACACGGCTTTTATTATTGTGCGAGAGAACGTTCCGGCCTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCA (SEQ ID NO: 193)
CAGGTGCAGCTGGTGCAGTCTGGGTCTGAGGTGAAGAAGCCTGGGGCCTCAGTGAAGGTTTCCTGCAAGGCATCTGGATACACCTTCACCAACTACTATATACACTGGGTGCGGCAGGCCCCTGGACAAGGGCTTGAGTGGATGGGAATAATCACCCTAGTGGTGGTACCACAAGCTACGCACAGAAGTTCCAGGGCAGAGTCACTATGACCAGGGACACGTCCACGAACACAGTCTACATGGGGCTGAGCAGCCTGAGATCTGAGGACACGGCCATGTATTACTGTGCGAGAGCCGGGGTACAACCTGGATCGACGAGGGTGGTTCGACCCCTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCA (SEQ ID NO: 203)
CAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGGCCTCAGTGAAGGTTTCCTGCAAGGCATCTGGATACACCTTCACCAGCTACTATATACACTGGGTACGACAGGCCCCTGGACAAGGGCTTGAGTGGATGGGAGTAACAGCCCTAGTGGTCGTAGCACAAGCTTCGCACAGAAGTTCCAGGGCAGAGTCAACCATGACCAGGGACACGTCCACGAGCGCAGTCTATATGGACCTGGA CAGCCTGAGATCTGAGGACACGGCCGTGTATTACTGTGCGAGAGGGGGAGTGACGATACACCTGGAACGACGGGGCTACTTTGACTACTGGGGCCAGGGAACCCTGGTCATTGTCTCCTCA (SEQ ID NO: 213)

Table 5C: Exemplary Variable Light Chain and Variable Heavy Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies		
Combination Number	Variable Light Chain Nucleic Acid Sequence	Variable Heavy Chain Nucleic Acid Sequence
Combination 1	SEQ ID NO: 93	SEQ ID NO: 94
Combination 2	SEQ ID NO: 95	SEQ ID NO: 96
Combination 3	SEQ ID NO: 97	SEQ ID NO: 98
Combination 4	SEQ ID NO: 99	SEQ ID NO: 100
Combination 5	SEQ ID NO: 101	SEQ ID NO: 102
Combination 6	SEQ ID NO: 103	SEQ ID NO: 104
Combination 7	SEQ ID NO: 105	SEQ ID NO: 106
Combination 10	SEQ ID NO: 111	SEQ ID NO: 112

Table 5C: Exemplary Variable Light Chain and Variable Heavy Chain Nucleic Acid Sequences of Anti-Protein S Monoclonal Antibodies		
Combination 11	SEQ ID NO: 113	SEQ ID NO: 114
Combination 12	SEQ ID NO: 115	SEQ ID NO: 116
Combination 13	SEQ ID NO: 162	SEQ ID NO: 163
Combination 14	SEQ ID NO: 164	SEQ ID NO: 165
Combination 15	SEQ ID NO: 166	SEQ ID NO: 167
Combination 17	SEQ ID NO: 170	SEQ ID NO: 171
Combination 18	SEQ ID NO: 172	SEQ ID NO: 173
Combination 19	SEQ ID NO: 178	SEQ ID NO: 183
Combination 20	SEQ ID NO: 188	SEQ ID NO: 193
Combination 21	SEQ ID NO: 198	SEQ ID NO: 203
Combination 22	SEQ ID NO: 208	SEQ ID NO: 213

[0288] In some embodiments, provided herein are nucleic acids encoding any of the Protein S antibodies disclosed herein. In some embodiments, provided herein are nucleic acids comprising any one or more of the nucleic acid sequences of Tables 5A-5B. In some embodiments, the heavy chain and light chain variable domains of the Protein S antibodies disclosed herein are encoded by a nucleic acid comprising any one or more of the nucleic acid sequences of Tables 5A-5B.

[0289] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 93 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0290] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 94, or a nucleic acid sequence with at least

80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0291] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 93 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 94, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0292] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 95 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0293] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 96, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0294] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 95 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 96, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0295] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 97 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0296] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 98, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0297] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 97 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 98, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0298] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 99 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0299] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 100, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0300] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 99 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 100, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0301] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 101 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0302] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 102, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0303] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 101 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 102, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0304] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 103 or a nucleic acid sequence with at

least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0305] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 104, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0306] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 103 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 104, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0307] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 105 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0308] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 106, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, or 99% sequence identity thereto.

[0309] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 105 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%,

95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 106, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0310] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 111 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0311] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 112, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0312] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 111 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 112, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0313] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 113 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0314] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is

encoded by the nucleic acid sequence of SEQ ID NO: 114, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0315] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 113 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 114, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0316] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 115 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0317] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 116, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0318] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 115 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 116, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0319] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 162 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0320] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 163, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0321] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 162 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 163, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0322] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 164 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0323] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 165, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0324] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 164 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 165, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0325] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 166 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0326] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 167, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0327] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 166 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 167, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0328] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 170 or a nucleic acid sequence with at

least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0329] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 171, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0330] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 170 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 171, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0331] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 172 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0332] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 173, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0333] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 172 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%,

95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 173, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0334] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 178 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0335] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 183, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0336] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 178 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 183, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0337] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 188 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0338] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is

encoded by the nucleic acid sequence of SEQ ID NO: 193, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0339] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 188 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 193, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0340] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 198 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0341] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 203, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0342] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 198 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 203, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0343] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 208 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0344] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 213, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0345] In some embodiments, the variable domain of the Protein S antibodies of the disclosure are encoded by a nucleic acid, wherein the light chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 208 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 213, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0346] The disclosure also provides vectors comprising any nucleic acid of the disclosure. In some embodiments, the nucleic acid of the vector comprises any one or more of the nucleic acid sequences provided in Tables 5A-5B. In some embodiments, the vector is an expression vector or an expression construct. In some embodiments, the vector is a mammalian vector. In some embodiments, the vector is a viral vector.

[0347] In some embodiments, the Protein S antibodies provided herein are produced by culturing a cell under suitable conditions for leading to the expression of the Protein S antibody, wherein the cell comprises a vector.

II. Uses of Protein S Antibodies

A. Therapeutic Protein S Antibodies

[0348] Provided herein are antibodies that recognize and selectively and/or specifically bind to Protein S, including Protein S fragments. The antibodies disclosed herein may be used for therapeutics in a subject. In some embodiments, the subject is a mammalian subject. In some embodiments, the mammalian subject is a human subject. In some embodiments, the mammalian subject is a non-human primate, e.g. a cynomolgus monkey.

[0349] In some embodiments, the Protein S antibodies provided herein are useful for treating a condition in a subject, wherein the condition is associated with the coagulation cascade. In some embodiments, the Protein S antibodies provided herein are useful for reducing an ability of Protein S to act as a cofactor within the coagulation cascade for the treatment of a condition in a subject.

[0350] In some embodiments, the Protein S antibodies provided herein are useful for reducing an ability of Protein S to act as a cofactor for APC, TFPI, or APC and TFPI for the treatment of a bleeding disorder or other diseases, e.g., a platelet disorder.

[0351] In some embodiments, provided herein is a method of promoting the coagulation of blood, the method comprising contacting any one of the Protein S antibodies disclosed herein with Protein S. In some embodiments, the contacting takes place in plasma. In some embodiments, the method is *in vitro*. In some embodiments, the method is *in vivo*. In some embodiments, the method is *in vivo*, and the method further comprises administering any one of the Protein S antibodies disclosed herein to a subject in need thereof.

[0352] In some embodiments, provided herein is a method of promoting the coagulation of blood, the method comprising contacting any one of the Protein S antibodies disclosed herein with a blood sample. Exemplary antibodies that may be used in a method for promoting the coagulation of blood include, but are not limited to, antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ

ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0353] In some embodiments, the coagulation of blood is marked by an increase in thrombin generation. Exemplary antibodies wherein use of the antibodies in the method for promoting the coagulation of blood, and wherein the coagulation of blood is marked by an increase in thrombin generation, include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0354] In some embodiments, the coagulation of blood is marked by an increase in fibrin generation. Exemplary antibodies wherein use of the antibodies in the method of promoting the coagulation of blood, and wherein the coagulation of blood is marked by an increase in fibrin

generation include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0355] In some embodiments, the coagulation of blood is marked by an increase in D-dimer.

[0356] In some embodiments, the blood sample is obtained from a subject having a coagulation factor deficiency or von Willebrand disease (vWD), or a platelet disorder. In some embodiments, the vWD is a subtype selected from: vWD Type 1, vWD Type 2A, vWD Type 2B, vWD Type 2N, vWD Type 2M, vWD Type 3, and acquired vWD. Exemplary antibodies wherein use of the antibodies for a method for promoting coagulation of blood, and wherein the blood is a blood sample obtained from a subject having a coagulation factor deficiency (e.g. such as Factor VII deficiency, Factor VIII deficiency, Factor IX deficiency, Factor XI deficiency) or von Willebrand disease may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60.

[0357] In some embodiments, provided herein is a method of promoting the coagulation of blood in a subject in need thereof, the method comprising administering to the subject any one of the Protein S antibodies disclosed herein, or any one of the pharmaceutical compositions

disclosed herein. In some embodiments, the antibody remains active in the subject for a period of time, wherein the period of time is antibody dose-dependent. In some embodiments, the period of time is about 50 hours to about 170 hours. Exemplary antibodies wherein use of the antibodies for a method of promoting the coagulation of blood in a subject in need thereof, wherein the activity of the antibody is dose-dependent, may comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58.

[0358] In some embodiments, provided herein is a method of promoting generation of thrombin in a subject in need thereof, the method comprising administering to the subject any one of the Protein S antibodies disclosed herein, or any one of the pharmaceutical compositions disclosed herein. In some embodiments, the subject suffers from a disease or condition selected from the group consisting of bleeding disorders, and platelet disorders. In some embodiments, the subject suffers from trauma and/or bleeding resulting from a surgery or a medical procedure. For example, the medical procedure may be a procedure in which bleeding may occur, but not necessarily so. In some embodiments, the medical procedure is a dental procedure. Exemplary antibodies that may be used in a method of promoting generation of thrombin in a subject in need thereof may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, or may comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, or may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, or may comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, and may comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57.

[0359] In some embodiments, provided herein is a method of treating a condition in a subject in need thereof, wherein the disease is selected from the group consisting of bleeding disorders, and platelet disorders, and the method comprises administering to the subject any one of the Protein S antibodies disclosed herein, or any one of the pharmaceutical compositions disclosed

herein. In some embodiments, the condition is a bleeding disorder. In some embodiments, the bleeding disorder is selected from the group consisting of hemophilia A, hemophilia B, von Willebrand disease (vWD, which may be a subtype selected from: vWD Type 1, vWD Type 2A, vWD Type 2B, vWD Type 2N, vWD Type 2M, vWD Type 3, and acquired vWD), menorrhagia including menorrhagia due to a congenital or acquired factor deficiency, Factor I deficiency, Factor II deficiency, Factor V deficiency, Factor VII deficiency, Factor X deficiency, Factor XI deficiency (hemophilia C), Factor VIII deficiency (hemophilia A), Factor IX deficiency (hemophilia B), trauma, and hereditary hemorrhagic telangiectasia. In some embodiments, the bleeding is associated with surgery, e.g. in a subject with a type of hemophilia. In some embodiments, the bleeding is associated with a medical procedure, e.g., a dental procedure. In some embodiments, the bleeding disorder is vWD, and the subject also suffers from menorrhagia. In some embodiments, the bleeding disorder is vWD, and the subject is undergoing a prophylactic treatment. In some embodiments, the subject suffers from menorrhagia associated with any one or more bleeding disorders and/or platelet disorders. In some embodiments, the subject is a hemophilia carrier. In some embodiments, the subject is a hemophilia carrier, and suffers from menorrhagia. In some embodiments, the condition is a platelet disorder. In some embodiments, the platelet disorder includes but is not limited to Bernard-Soulier syndrome, Glanzmann's thrombasthenia, and platelet storage pool deficiencies. In some embodiments, the platelet disorder is a platelet storage pool deficiency. In some embodiments, the platelet storage pool deficiency includes but is not limited to: Gray platelet syndrome, Quebec platelet disorder, and MYH9-related thrombocytopenia (MYH9RD). In some embodiments, the subject has a bleeding disorder, and has inhibitors. In some embodiments, the bleeding disorder is hemophilia A or hemophilia B, wherein the subject has inhibitors. In some embodiments, the bleeding disorder is vWD. For example, the inhibitors may be developed in the subject as a response to factor replacement therapy. Exemplary antibodies that may be used for a method of treating a condition in a subject in need thereof, wherein the disease is selected from the group consisting of bleeding disorders, and platelet disorders, include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain

variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0360] In some embodiments, a subject in need thereof may be treated with any of the Protein S antibodies provided herein, wherein the treatment is a routine prophylaxis to prevent or reduce the frequency of bleeding episodes. In some embodiments, a subject in need thereof may be treated with any of the Protein S antibodies provided herein, wherein the treatment is an on-demand treatment used for the control of bleeding episodes. In some embodiments, a subject in need thereof may be treated with any of the Protein S antibodies provided herein, wherein the treatment is a perioperative management of bleeding. For example, a perioperative management treatment may be used for treating a subject prior to, during, and/or after surgery or other medical procedure, or prior to, during, and/or after trauma.

[0361] In some embodiments, the treatment with any of the Protein S antibodies provided herein is provided as a chronic therapy, with dosing occurring continuously over time. In some embodiments, the treatment with any of the Protein S antibodies provided herein is provided as an intermittent therapy, with dosing occurring at irregular intervals. As an example, such an intermittent therapy can be used for a subject having menorrhagia. In some embodiments, the treatment with any of the Protein S antibodies provided herein is provided as an acute therapy, with dosing occurring for a short finite period of time. For example, the acute therapy may be administered for spontaneous bleeding episodes, or in conjunction with a surgery or other medical procedure, or after experiencing a trauma.

[0362] In some embodiments, the method of treatment of a subject may be a combination one of the above, e.g., the method of treatment may be prophylactic, and on-demand. In some embodiments, a prophylactic method of treatment may be a chronic therapy. In some

embodiments, a prophylactic method of treatment may be an acute therapy. In some embodiments, a prophylactic method of treatment may be an intermittent therapy. In some embodiments, an on-demand treatment may be an acute therapy. In some embodiments, an on-demand treatment may be an intermittent treatment.

[0363] In some embodiments, treatment of a subject in need thereof comprises administering to the subject any of the Protein S antibodies provided herein, wherein the Protein S antibodies provided herein are Fab fragments. Without being bound to any theory, in some embodiments a shorter half-life of a Fab fragment, in relation to a full-length antibody with the same VH/VL may be beneficial for an acute treatment or on-demand. In some embodiments, the Fab fragment Protein S antibodies are administered to a subject in need thereof to reduce risk of bleeding.

B. Combination Therapies

[0364] The administration of any one of the therapeutic Protein S antibodies provided herein may be a monotherapy, or may be in combination with any other known drugs or treatments for diseases or conditions. In some embodiments, the other known drugs or treatments are useful for treating disorders, diseases, or conditions associated with reduced or impaired clotting. In some embodiments, the disorder, condition is a bleeding disorder. In some embodiments, the disorder, disease, or condition is a bleeding disorder or a platelet disorder.

[0365] In some embodiments, the administration of any of the therapeutic Protein S antibodies provided herein may be with a factor replacement therapy. In some embodiments, the administration of any of the therapeutic Protein S antibodies provided herein may be with the administration of a recombinant Factor VII.

C. Administration of Therapeutic Protein S Antibodies

[0366] The *in vivo* administration of the therapeutic Protein S antibodies described herein may be carried out intravenously, intramuscularly, subcutaneously, topically, orally, transdermally, intraperitoneally, intraorbitally, intrathecally, intraventricularly, intranasally, transmucosally, through implantation, or through inhalation. Administration of the therapeutic Protein S antibodies may be performed with any suitable excipients, carriers, or other agents to provide suitable or improved tolerance, transfer, delivery, and the like.

[0367] In some embodiments, the *in vivo* administration of any of the therapeutic Protein S antibodies provided herein may be an intravenous administration. In some embodiments, the intravenous administration may be provided as a prophylactic treatment. In some embodiments, the prophylactic treatment may be a routine prophylaxis. In some embodiments, the routine prophylaxis may have a regular dosing schedule. In some exemplary embodiments, the regular dosing schedule may be once weekly, twice weekly, once monthly, twice monthly, or three times monthly. In some embodiments, the intravenous administration may be provided as an on-demand treatment. In some embodiments, the intravenous administration may be provided as a chronic therapy. In some embodiments, the intravenous administration may be provided as an intermittent therapy. In some embodiments, the intravenous administration may be provided as an acute therapy. In some embodiments, an intermittent therapy may have a regular dosing schedule for the duration of the intermittent therapy. In some embodiments, an acute therapy may have a regular dosing schedule for the duration of the acute therapy. For example, administration of any of the therapeutic Protein S antibodies provided herein for an acute therapy by intravenous administration may occur on a regular dosing schedule for a predetermined duration of days, e.g., 7 days, 14 days, or more. Exemplary antibodies that may be used for an *in vivo* intravenous administration include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0368] In some embodiments, the *in vivo* administration of any of the therapeutic Protein S antibodies provided herein may be a subcutaneous administration. In some embodiments, the subcutaneous administration may be provided as a prophylactic treatment. In some embodiments, the prophylactic treatment may be a routine prophylaxis. In some embodiments, the routine prophylaxis may have a regular dosing schedule. In some exemplary embodiments, the regular dosing schedule may be once weekly, twice weekly, once monthly, twice monthly, or three times monthly. In some embodiments, the subcutaneous administration may be provided as an on-demand treatment. In some embodiments, the subcutaneous administration may be provided as a chronic therapy. In some embodiments, the subcutaneous administration may be provided as an intermittent therapy. In some embodiments, the subcutaneous administration may be provided as an acute therapy. In some embodiments, an intermittent therapy may have a regular dosing schedule for the duration of the intermittent therapy. In some embodiments, an acute therapy may have a regular dosing schedule for the duration of the acute therapy. For example, administration of any of the therapeutic Protein S antibodies provided herein for an acute therapy by subcutaneous administration may occur on a regular dosing schedule for a predetermined duration of days, e.g., 7 days, 14 days, or more. Exemplary antibodies that may be used for an *in vivo* subcutaneous administration include, but are not limited to antibodies that (a) comprise the light chain variable domain comprising the amino acid sequence of SEQ ID NO: 71 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 72, (b) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58, (c) comprise the amino acid sequence of SEQ ID NO: 75 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 76, (d) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60, (e) comprise the amino acid sequence of SEQ ID NO: 69 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 70, (f) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57, (g) comprise the amino acid sequence of SEQ ID NO: 89 and the heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 90, and (h) comprise the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0369] In some embodiments, the Protein S antibodies provided herein may be provided in a small volume amenable for injection, such as for subcutaneous administration. In some embodiments, the Protein S antibodies provided herein may be provided in a small volume amenable for injection by use of a pen-like auto-injector device. In some embodiments, the device is a syringe, for example a pre-filled syringe.

[0370] In some embodiments provided herein are single-dose vials useful for either subcutaneous or intravenous administration.

[0371] Accordingly, a therapeutically effective amount of the Protein S antibodies provided herein may be provided in a small volume for subcutaneous administration to a subject in need thereof. In some embodiments, the Protein S antibodies provided herein may be provided in a large volume amenable for administration by a subcutaneous infusion device, for subcutaneous infusion to a subject in need thereof.

D. Pharmaceutical Compositions

[0372] The disclosure also provides pharmaceutical compositions comprising any one of the Protein S antibodies disclosed herein, and optionally a pharmaceutical acceptable excipient or carrier. In some embodiments, the pharmaceutical composition is sterile. The pharmaceutical compositions may be formulated to be compatible with their intended routes of administration. In some embodiments, the pharmaceutical compositions of the disclosure are suitable for administration to a human subject.

E. Diagnostic Antibodies

[0373] The antibodies provided herein may also be used for diagnostic purposes. For example, diagnostic antibodies could be used for detecting protein S deficiencies, or for detecting protein S levels in plasma prior to dosing (e.g. as a companion diagnostic).

[0374] Accordingly, in some embodiments, a Protein S antibody of the disclosure is conjugated to a label, for example a detectable label, a spin label, a colorimetric label, a radioactive label, an enzymatic label, a fluorescent label, or a magnetic label. The label may be detectable by spectroscopic, photochemical, biochemical, immunochemical, fluorescent, electrical, optical or chemical methods. Useful labels include, but are not limited to, magnetic

beads (e.g. DYNABEADS®), fluorescent dyes (e.g., fluorescein isothiocyanate, red, rhodamine, and the like), radiolabels (e.g., ³H, ¹²⁵I, ³⁵S, ¹⁴C, or ³²P), enzymes (e.g., LacZ, CAT, horseradish peroxidase, alkaline phosphatase and others, commonly used as detectable enzymes, either as marker gene products or in an ELISA), biotin, avidin, or streptavidin and colorimetric labels such as colloidal gold colored glass or plastic (e.g. polystyrene, polypropylene, latex, etc.) beads, and nanoparticles. In some embodiments, provided herein are substrates to which one or more Protein S antibodies of the disclosure is attached.

[0375] Detection may be carried out on any biological sample obtained from a subject. Biological samples include, but are not limited to whole blood, plasma, serum, saliva, urine, feces, synovial fluid, cerebrospinal fluid, bronchial lavage, ascites fluid, bone marrow aspirate, pleural effusion, tissue, cells, a biopsy, interstitial fluid, lymphatic fluid, or fractions thereof derived from a subject. In some embodiments, the biological sample comprises cells and the cells are in culture, in a suspension, on a slide, in intact tissue, or in preparation ready for a FACs analysis.

III. Kits and Articles of Manufacture

[0376] The disclosure also provides a kit or article of manufacture comprising any one of the antibodies disclosed herein, or any pharmaceutical composition disclosed herein. In some embodiments, the kits may further include instructional materials for carrying out any of the methods disclosed herein. In some embodiments, the kits may further include sterile containers or vials for holding the antibodies and/or pharmaceutical compositions disclosed herein. In some embodiments, the kits may further include sterile delivery devices for administering the antibodies and/or pharmaceutical compositions disclosed herein. In some embodiments, an article of manufacture comprises any pharmaceutical composition of the disclosure.

IV. Exemplary Enumerated Embodiments

[0377] Exemplary enumerated embodiments of the disclosure are as follows.

[0378] 1. An antibody that binds Protein S, wherein the antibody is an inhibitor of the cofactor activity of Protein S for activated Protein C (APC), an inhibitor of the cofactor activity of Protein S for tissue factor pathway inhibitor (TFPI), or an inhibitor of the cofactor activity of Protein S for both APC and TFPI, and wherein the antibody is human, humanized, or chimeric.

- [0379]** 2. An antibody that binds Protein S, wherein the antibody is capable of promoting coagulation and/or modulating a component in the coagulation cascade.
- [0380]** 3. The antibody of any one of Enumerated Embodiments 1-2, wherein the antibody is an inhibitor for the cofactor activity of Protein S for APC.
- [0381]** 4. The antibody of any one of Enumerated Embodiments 1-2, wherein the antibody is an inhibitor for the cofactor activity of Protein S for TFPI.
- [0382]** 5. The antibody of any one of Enumerated Embodiments 1-2, wherein the antibody is an inhibitor for the cofactor activity of Protein S for both APC and TFPI.
- [0383]** 6. The antibody of Enumerated Embodiment 3, wherein the capability of the antibody for inhibiting the cofactor activity of Protein S for TFPI is negligible.
- [0384]** 7. The antibody of Enumerated Embodiment 4, wherein the capability of the antibody for inhibiting the cofactor activity of Protein S for APC is negligible.
- [0385]** 8. The antibody of any one of Enumerated Embodiments 1-7, wherein the antibody binds to the C-terminus of Protein S.
- [0386]** 9. The antibody of any one of Enumerated Embodiments 1-7, wherein the antibody binds to the N-terminus of Protein S.
- [0387]** 10. The antibody of any one of Enumerated Embodiments 1-9, wherein the antibody binds to a thrombin-sensitive region of the Protein S.
- [0388]** 11. The antibody of any one of Enumerated Embodiments 1-9, wherein the antibody binds to an EGF region of the Protein S.
- [0389]** 12. The antibody of any one of Enumerated Embodiments 1-9, wherein the antibody binds to an SHBG region of Protein S.
- [0390]** 13. The antibody of any one of Enumerated Embodiments 1-12, wherein the antibody is capable of promoting generation of a marker associated with coagulation activity.

[0391] 14. The antibody of any one of Enumerated Embodiments 1-13, wherein the antibody is capable of promoting thrombin generation.

[0392] 15. The antibody of any one of Enumerated Embodiments 1-14, wherein the antibody is capable of promoting D-dimer levels.

[0393] 16. The antibody of any one of Enumerated Embodiments 1-15, wherein the antibody is capable of promoting fibrin generation.

[0394] 17. The antibody of any one of Enumerated Embodiments 1-16, wherein activity of the antibody is dose-dependent.

[0395] 18. The antibody of any one of Enumerated Embodiments 1-17, wherein activity of the antibody is measured in vitro.

[0396] 19. The antibody of any one of Enumerated Embodiments 1-17, wherein activity of the antibody is measured in vivo.

[0397] 20. The antibody of any one of Enumerated Embodiments 1-19, wherein the binding affinity of the antibody to Protein S is calcium-dependent.

[0398] 21. The antibody of any one of Enumerated Embodiments 1-19, wherein the binding affinity of the antibody to Protein S is calcium-independent.

[0399] 22. The antibody of any one of Enumerated Embodiments 1-21, wherein the antibody binds to free Protein S.

[0400] 23. The antibody of any one of Enumerated Embodiments 1-21, wherein the antibody binds to complexed Protein S.

[0401] 24. The antibody of Enumerated Embodiment 23, wherein the Protein S is complexed with C4BP.

[0402] 25. The antibody of Enumerated Embodiment 23, wherein the Protein S is complexed with TFPI.

[0403] 26. The antibody of any one of Enumerated Embodiments 1-25, wherein the antibody is a monoclonal antibody.

[0404] 27. The antibody of any one of Enumerated Embodiments 1-26, wherein the antibody is a full-length antibody.

[0405] 28. The antibody of any one of Enumerated Embodiments 1-26, wherein the antibody is an antibody fragment.

[0406] 29. The antibody of any one of Enumerated Embodiments 1-28, wherein the antibody is a humanized antibody.

[0407] 30. The antibody of any of Enumerated Embodiments 1-29, wherein the antibody comprises a Fc domain.

[0408] 31. The antibody of Enumerated Embodiment 30, wherein the Fc domain is human.

[0409] 32. The antibody of Enumerated Embodiment 31, wherein the human Fc domain is IgG1, IgG2, IgG3, or IgG4.

[0410] 33. The antibody of Enumerated Embodiment 32, wherein the Fc domain of the antibody is human IgG4, optionally SEQ ID NO: 218, and comprises at least one amino acid substitution at a position selected from the group consisting of: 215, 221, 222, 228, 234, 235, 236, 239, 240, 241, 243, 244, 245, 247, 250, 252, 254, 256, 262, 263, 264, 265, 266, 267, 268, 269, 270, 292, 296, 297, 298, 299, 300, 305, 313, 324, 325, 326, 327, 328, 329, 330, 332, 333, 334, 345, 396, 428, 430, 433, 434, and 440, or comprises one or more of the substitutions selected from the group consisting of T250Q/M428L, M252Y/S254T/T256E, M428L/N434S, S267E/L328F, N325S/L328F, and H433K/N434F, wherein the position numbers of the amino acid residues are of the EU numbering scheme.

[0411] 34. The antibody of Enumerated Embodiment 27, wherein the antibody is a human antibody.

[0412] 35. The antibody of Enumerated Embodiment 27, wherein the antibody is a chimeric antibody.

[0413] 36. The antibody of any one of Enumerated Embodiments 1-35, wherein the antibody is conjugated.

[0414] 37. The antibody of Enumerated Embodiment 36, wherein the antibody is conjugated to a label.

[0415] 38. The antibody of any one of Enumerated Embodiments 1-37, wherein the antibody comprises any one or more of the amino acid sequences of the CDR sequences provided in Tables 1A, 1B, 1C, 2A, 2B, and 2C.

[0416] 39. The Protein S antibody of any one of Enumerated Embodiments 1-38, wherein the antibody comprises:

- (a) any one of the CDR-L1 amino acid sequences of Table 1A;
- (b) any one of the CDR-L2 amino acid sequences of Table 1B;
- (c) any one of the CDR-L3 amino acid sequences of Table 1C;
- (d) any one of the CDR-H1 amino acid sequences of Table 2A;
- (e) any one of the CDR-H2 amino acid sequences of Table 2B; and
- (f) any one of the CDR-H3 amino acid sequences of Table 2C.

[0417] 40. The Protein S antibody of any one of Enumerated Embodiments 1-39, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of any one of SEQ ID NOs: 1-6, 9-11, 117, 127, 136, 141, 174, 184, 194, and 204;
- (b) a CDR-L2 comprising the amino acid sequence of any one of SEQ ID NOs: 12-20, 118, 128, 142, 175, 185, 195, and 205; and
- (c) a CDR-L3 comprising the amino acid sequence of any one of SEQ ID NOs: 21-27, 30-32, 119, 123, 129, 137, 143, 176, 186, 196, and 206.

[0418] 41. The Protein S antibody of any one of Enumerated Embodiments 1-40, wherein the heavy chain variable domain of antibody comprises:

(a) a CDR-H1 comprising the amino acid sequence of any one of SEQ ID NOs: 33-39, 42-44, 120, 124, 130, 138, 144, 179, 189, 199, and 209;

(b) a CDR-H2 comprising the amino acid sequence of any one of SEQ ID NOs: 45-51, 54-56, 121, 125, 139, 145, 180, 190, 200, and 210; and

(c) a CDR-H3 comprising the amino acid sequence of any one of SEQ ID NOs: 57-63, 66-68, 122, 126, 131, 140, 146, 181, 191, 201, and 211.

[0419] 42. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 1;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 12; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 21.

[0420] 43. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 2;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 13; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 22.

[0421] 44. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 3;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 14; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 23.

[0422] 45. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 1;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 15; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 24.

[0423] 46. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 4;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 25.

[0424] 47. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 5;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 17; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 26.

[0425] 48. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 6;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 18; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 27.

[0426] 49. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 9;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 30.

[0427] 50. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 10;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 31.

[0428] 51. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 11;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 20; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[0429] 52. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 117;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 118; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 119.

[0430] 53. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 117;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 118; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 123.

[0431] 54. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 127;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 128; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 129.

[0432] 55. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 136;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 137; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 137.

[0433] 56. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 141;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 142; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 143.

[0434] 57. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

(a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 174;

(b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 175; and

(c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 176.

[0435] 58. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 184;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 185; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 186.

[0436] 59. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 194;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 195; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 196.

[0437] 60. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises:

- (a) a CDR-L1 comprising the amino acid sequence of SEQ ID NO: 204;
- (b) a CDR-L2 comprising the amino acid sequence of SEQ ID NO: 205; and
- (c) a CDR-L3 comprising the amino acid sequence of SEQ ID NO: 206.

[0438] 61. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 33;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 45; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 57.

[0439] 62. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 34;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 46; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 58.

[0440] 63. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 35;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 47; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 59.

[0441] 64. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 36;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 48; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 60.

[0442] 65. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 37;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 49; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 61.

[0443] 66. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 38;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 50; and

(c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 62.

[0444] 67. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

(a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 39;

(b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 51; and

(c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 63.

[0445] 68. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

(a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 42;

(b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 54; and

(c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 66.

[0446] 69. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

(a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 43;

(b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 55; and

(c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 67.

[0447] 70. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

(a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 44;

(b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 56; and

(c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 68.

[0448] 71. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 120;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 121; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 122.

[0449] 72. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 124;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 125; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 126.

[0450] 73. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 130;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 50; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 131.

[0451] 74. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 138;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 139; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 140.

[0452] 75. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 144;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 145; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 146.

[0453] 76. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 179;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 180; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 181.

[0454] 77. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 189;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 190; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 191.

[0455] 78. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 199;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 200; and
- (c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 201.

[0456] 79. The Protein S antibody of any one of Enumerated Embodiments 1-60, wherein the heavy chain variable domain of the antibody comprises:

- (a) a CDR-H1 comprising the amino acid sequence of SEQ ID NO: 209;
- (b) a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 210; and

(c) a CDR-H3 comprising the amino acid sequence of SEQ ID NO: 211.

[0457] 80. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, and SEQ ID NO: 57.

[0458] 81. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, and SEQ ID NO: 58.

[0459] 82. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 3, SEQ ID NO: 14, SEQ ID NO: 23, SEQ ID NO: 35, SEQ ID NO: 47, and SEQ ID NO: 59.

[0460] 83. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, and SEQ ID NO: 60.

[0461] 84. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy amino acid sequences of SEQ ID NO: 4, SEQ ID NO: 16, SEQ ID NO: 25, SEQ ID NO: 37, SEQ ID NO: 49, and SEQ ID NO: 61.

[0462] 85. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 5, SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 38, SEQ ID NO: 50, and SEQ ID NO: 62.

[0463] 86. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 6, SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 39, SEQ ID NO: 51, and SEQ ID NO: 63.

[0464] 87. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 9, SEQ ID NO: 19, SEQ ID NO: 30, SEQ ID NO: 42, SEQ ID NO: 54, and SEQ ID NO: 66.

[0465] 88. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, and SEQ ID NO: 67.

[0466] 89. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 11, SEQ ID NO: 20, SEQ ID NO: 32, SEQ ID NO: 44, SEQ ID NO: 56, and SEQ ID NO: 68.

[0467] 90. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 117, SEQ ID NO: 118, SEQ ID NO: 119, SEQ ID NO: 120, SEQ ID NO: 121, and SEQ ID NO: 122.

[0468] 91. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 117, SEQ ID NO: 118, SEQ ID NO: 123, SEQ ID NO: 124, SEQ ID NO: 125, and SEQ ID NO: 126.

[0469] 92. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 127, SEQ ID NO: 128, SEQ ID NO: 129, SEQ ID NO: 130, SEQ ID NO: 50, and SEQ ID NO: 131.

[0470] 93. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 136, SEQ ID NO: 17, SEQ ID NO: 137, SEQ ID NO: 138, SEQ ID NO: 139, and SEQ ID NO: 140.

[0471] 94. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 141, SEQ ID NO: 142, SEQ ID NO: 143, SEQ ID NO: 144, SEQ ID NO: 145, and SEQ ID NO: 146.

[0472] 95. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 174, SEQ ID NO: 175, SEQ ID NO: 176, SEQ ID NO: 179, SEQ ID NO: 180, and SEQ ID NO: 181.

[0473] 96. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 184, SEQ ID NO: 185, SEQ ID NO: 186, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191.

[0474] 97. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 194, SEQ ID NO: 195, SEQ ID NO: 196, SEQ ID NO: 199, SEQ ID NO: 200, and SEQ ID NO: 201.

[0475] 98. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the antibody comprises the light and heavy CDR amino acid sequences of SEQ ID NO: 204, SEQ ID NO: 205, SEQ ID NO: 206, SEQ ID NO: 209, SEQ ID NO: 210, and SEQ ID NO: 211.

[0476] 99. The Protein S antibody of any one of Enumerated Embodiments 1-98, wherein the antibody comprises the variable chain amino acid sequence of any one of the amino acid sequences provided in Table 4A, and/or the variable chain amino acid sequence of any one of the amino acid sequences provided in Table 4B.

[0477] 100. The Protein S antibody of any one of Enumerated Embodiments 1-99, wherein the antibody comprises the light and heavy variable chain amino acid sequence of any one of the amino acid sequence combinations provided in Table 4C.

[0478] 101. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 69 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 70, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0479] 102. The Protein S antibody of one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 71 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 72, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0480] 103. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 73 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 74, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0481] 104. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 75 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 76, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0482] 105. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 77 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 78, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0483] 106. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 79 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 80, or an amino acid sequence with at least 80%, 81%, 82%, 83%,

84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0484] 107. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 81 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 82, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0485] 108. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 87 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 88, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0486] 109. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises In some embodiments, provided herein are Protein S antibodies, wherein the light chain variable domain of the antibodies comprise the amino acid sequence of SEQ ID NO: 89 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 90, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0487] 110. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 91 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%,

88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 92, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0488] 111. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 148 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 149, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0489] 112. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 150 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 151, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0490] 113. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 152 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 153, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0491] 114. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 156 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 157, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0492] 115. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 158 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 159, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0493] 116. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 177 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 182, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0494] 117. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 187 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 192, or an amino acid sequence with at least 80%, 81%, 82%,

83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0495] 118. The Protein S antibody of any one of Enumerated Embodiments 1-41, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 197 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 202, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0496] 119. The Protein S antibody of any one of Enumerated Embodiments 1-100, wherein the light chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 207 or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody comprises the amino acid sequence of SEQ ID NO: 212, or an amino acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0497] 120. The Protein S antibody of any one of Enumerated Embodiments 1-119, wherein the light chain comprises the amino acid sequence of SEQ ID NO: 214.

[0498] 121. The Protein S antibody of any one of Enumerated Embodiments 1-119, wherein the light chain comprises the amino acid sequence of SEQ ID NO: 215.

[0499] 122. A pharmaceutical composition comprising any one of the antibodies of Enumerated Embodiments 1-121, and optionally a pharmaceutically acceptable excipient.

[0500] 123. A complex comprising Protein S and the antibody of any one of Enumerated Embodiments 1-119, wherein the antibody is bound to the Protein S, and the Protein S is free.

[0501] 124. A complex comprising Protein S and the antibody of any one of Enumerated Embodiments 1-119, wherein the antibody is bound to the Protein S, and the Protein S is complexed.

[0502] 125. The complex of Enumerated Embodiment 124, wherein the Protein S is bound to C4BP.

[0503] 126. The complex of Enumerated Embodiment 124, wherein the Protein S is bound to TFPI.

[0504] 127. A nucleic acid encoding for any one of the antibodies of Enumerated Embodiments 1-121.

[0505] 128. The nucleic acid of Enumerated Embodiment 127, comprising any one of the nucleic acid sequences selected from Table 5A.

[0506] 129. The nucleic acid of any one of Enumerated Embodiments 127-128, comprising any one of the nucleic acid sequences selected from Table 5B.

[0507] 130. The nucleic acid of any one of Enumerated Embodiments 127-129, wherein the nucleic acid comprises the nucleic acid sequence of any one of the nucleic acid sequence combinations provided in Table 5C.

[0508] 131. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 93 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 94, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0509] 132. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 95 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity

thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 96, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0510] 133. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 97 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 98, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0511] 134. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 99 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 100, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0512] 135. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 101 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 102, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0513] 136. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of

SEQ ID NO: 103 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 104, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0514] 137. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 105 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 106, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0515] 138. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 111 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 112, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0516] 139. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 113 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 114, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0517] 140. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 115 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 116, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0518] 141. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 162 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 163, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0519] 142. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 164 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 165, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0520] 143. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 166 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 167, or a nucleic acid sequence with at least 80%,

81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0521] 144. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 170 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 171, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0522] 145. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 172 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 173, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0523] 146. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 178 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 183, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0524] 147. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 188 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence

identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 193, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0525] 148. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 198 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 203, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0526] 149. The nucleic acid of any one of Enumerated Embodiments 127-130, wherein the light chain variable domain of the Protein S antibody is encoded by the nucleic acid sequence of SEQ ID NO: 208 or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto; and/or wherein the heavy chain variable domain of the antibody is encoded by the nucleic acid sequence of SEQ ID NO: 213, or a nucleic acid sequence with at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity thereto.

[0527] 150. A vector comprising the nucleic acid of any one of Enumerated Embodiments 127-149.

[0528] 151. An in vitro method of promoting the coagulation of blood, comprising contacting the antibody of any one of Enumerated Embodiments 1-121 with a blood sample.

[0529] 152. The method of Enumerated Embodiment 151, wherein the blood sample comprises plasma.

[0530] 153. The method of any one of Enumerated Embodiments 151-152, wherein a marker associated with coagulation activity is increased.

[0531] 154. The method of any one of Enumerated Embodiments 151-153, wherein thrombin generation is promoted.

[0532] 155. The method of any one of Enumerated Embodiments 151-154, wherein fibrin generation is promoted.

[0533] 156. The method of any one of Enumerated Embodiments 151-155, wherein D-dimer levels are promoted.

[0534] 157. The method of any one of Enumerated Embodiments 151-156, wherein the blood sample is obtained from a subject having a coagulation factor deficiency, von Willebrand disease, or a platelet disorder.

[0535] 158. A method of promoting the coagulation of blood in a subject in need thereof, comprising administering to the subject the antibody of any one of Enumerated Embodiments 1-121 or the pharmaceutical composition of Enumerated Embodiment 122.

[0536] 159. The method of Enumerated Embodiment 158, wherein a marker associated with coagulation activity is increased.

[0537] 160. The method of any of Enumerated Embodiments 158-159, wherein thrombin generation is promoted in the subject.

[0538] 161. The method of any of Enumerated Embodiments 158-160, wherein fibrin generation is promoted in the subject.

[0539] 162. The method of any of Enumerated Embodiments 153-161, wherein D-dimer levels are increased in the subject.

[0540] 163. The method of any of Enumerated Embodiments 153-162, wherein the antibody remains active in the subject for a period of time.

[0541] 164. method of any of Enumerated Embodiments 151-163, wherein activity of the antibody is dose-dependent.

[0542] 165. A method of promoting the generation of thrombin in a subject in need thereof, comprising administering to the subject the antibody of any one of Enumerated Embodiments 1-121 or the pharmaceutical composition of Enumerated Embodiment 122.

[0543] 166. The method of Enumerated Embodiment 165, wherein the subject has a coagulation factor deficiency, von Willebrand disease, or a platelet disorder, and the antibody restores or promotes the generation of thrombin.

[0544] 167. A method of treating a condition in a subject in need thereof, comprising administering to the subject the antibody of Enumerated Embodiments 1-121 or the pharmaceutical composition of Enumerated Embodiment 122, wherein the condition is selected from the group consisting of: bleeding disorders, platelet disorders, trauma, and bleeding resulting from a surgery or a medical procedure.

[0545] 168. The method of Enumerated Embodiment 167, wherein the method of treating is prophylactic.

[0546] 169. The method of Enumerated Embodiment 167, wherein the method of treating is on-demand.

[0547] 170. The method of any one of Enumerated Embodiments 167-169, wherein the method is prophylactic and on-demand.

[0548] 171. The method of any one of Enumerated Embodiments 168 or 170, wherein the prophylactic method of treating is a routine prophylaxis.

[0549] 172. The method of any one of Enumerated Embodiments 167-171, wherein the administration of the antibody of any one of Enumerated Embodiments 1-119 is a subcutaneous administration.

[0550] 173. The method of any one of Enumerated Embodiments 158-172, wherein the method of treating is acute.

[0551] 174. The method of any one of Enumerated Embodiments 158-172, wherein the method of treating is chronic.

[0552] 175. The method of any one of Enumerated Embodiments 158-172, wherein the method of treating is perioperative.

[0553] 176. The method of any one of Enumerated Embodiments 158-172, wherein the method of treating is intermittent.

[0554] 177. The method of any one of Enumerated Embodiments 158-176, wherein the antibody exhibits graded inhibition.

[0555] 178. The method of any one of Enumerated Embodiments 158-176, wherein the antibody exhibits switch-like inhibition.

[0556] 179. The method of any one of Enumerated Embodiments 167-178, wherein the subject suffers from two or more conditions selected from the group consisting of: bleeding disorders, platelet disorders, trauma, and bleeding resulting from a surgery or a medical procedure.

[0557] 180. The method of any one of Enumerated Embodiments 165-179, wherein the subject suffers from a bleeding disorder selected from the group consisting of: hemophilia A, hemophilia B, von Willebrand disease (vWD) disease, menorrhagia, Factor I deficiency, Factor II deficiency, Factor V deficiency, Factor VII deficiency, Factor X deficiency, Factor XI deficiency, Factor VIII deficiency (hemophilia A), Factor IX deficiency (hemophilia B), trauma, and hereditary hemorrhagic telangiectasia.

[0558] 181. The method of any one of Enumerated Embodiments 158-180, wherein the subject is a hemophilia carrier.

[0559] 182. The method of any one of Enumerated Embodiments 165-181, wherein the subject suffers from menorrhagia.

[0560] 183. The method of any one of Enumerated Embodiments 158-182, wherein the subject suffers from menorrhagia associated with one or more of the bleeding disorders or the platelet disorders.

[0561] 184. The method of Enumerated Embodiment 180, wherein the bleeding disorder is vWD, and wherein the subject is undergoing a prophylactic treatment.

[0562] 185. The method of Enumerated Embodiment 180, wherein the bleeding disorder is vWD, and the vWD is a subtype selected from: vWD Type 1, vWD Type 2A, vWD Type 2B, vWD Type 2N, vWD Type 2M, vWD Type 3, and acquired vWD.

[0563] 186. The method of any one of Enumerated Embodiments 167-185, wherein the condition is a platelet disorder selected from the group consisting of: Bernard-Soulier syndrome, Glanzmann's thrombasthenia, and platelet storage pool deficiency.

[0564] 187. The method of Enumerated Embodiment 186, wherein the platelet disorder is a platelet storage pool deficiency selected from the group consisting of: Gray platelet syndrome, Quebec platelet disorder, and MYH9-related thrombocytopenia (MYH9RD).

[0565] 188. The method of Enumerated Embodiment 180, wherein the bleeding disorder is selected from Factor I deficiency, Factor II deficiency, Factor V deficiency, Factor VII deficiency, Factor X deficiency, Factor XI deficiency, Factor VIII deficiency (hemophilia A), Factor IX deficiency (hemophilia B), and vWD disease, and wherein the subject has inhibitors.

[0566] 189. The method of any one of Enumerated Embodiments 167-188, wherein the antibody or the pharmaceutical composition is capable of promoting thrombin generation in the subject.

[0567] 190. The method of any one of Enumerated Embodiments 165-166, wherein the thrombin generation does not exceed a predetermined threshold level.

[0568] 191. The method of any one of Enumerated Embodiments 165-166 and 190, wherein the thrombin generation is antibody concentration-dependent.

[0569] 193. A kit or article of manufacture comprising an antibody of any one of Enumerated Embodiments 1-121 or the pharmaceutical composition of Enumerated Embodiment 122.

[0570] 193. Use of the antibody of any one of Enumerated Embodiments 1-121 or the pharmaceutical composition of Enumerated Embodiment 122 for the treatment of a condition in a subject in need thereof.

[0571] 194. Use of the antibody of any one of Enumerated Embodiments 1-121 or the pharmaceutical composition of Enumerated Embodiment 122 for the manufacture of a medicament for the treatment of a condition in a subject in need thereof.

[0572] The present invention is not limited in scope by the specific embodiments described herein, which are intended as illustrations of individual aspects or embodiments of the invention. Functionally equivalent methods and components are within the scope of the invention. Various modifications of the invention, in addition to those described here, are apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications fall within the scope of the invention.

[0573] The invention is now described with reference to the following Examples. These Examples are provided for the purpose of illustration only and the invention should in no way be construed as being limited to these Examples, but rather should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

[0574] Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the antibodies of the present invention and practice the claimed methods. The following examples therefore, specifically point out the preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

EXAMPLES

Example 1: Generation of Protein S Antibodies

[0575] The Protein S antibodies of the disclosure were generated as follows. Animals (e.g., mice, rats) were immunized with a full-length purified human plasma Protein S. Three immunization campaigns were carried out and standard techniques were used to generate hybridoma libraries from the animals. Flow cytometry and single cell sorting was used to generate single cell clones. Supernatants from these single clones were then screened for binding to both human and cynomolgus monkey Protein S. Clones that exhibited binding to both human and cynomolgus monkey Protein S were selected for expanded growth. The expanded cultures were then purified over Protein G or Protein A Sepharose using standard techniques. These purified antibody preparations were used in subsequent functional assays.

The selected Protein S antibodies had a human or mouse variable region and a rat Fc domain, or a mouse Fc domain.

[0576] Antibodies having a human variable region and a rat or mouse Fc domain were made into fully human antibodies maintaining the human variable domain as the parent antibody, but with a human IgG4 Fc domain. Table 6 below lists the Antibody number used to designate the human antibodies in the first column and the corresponding parental rodent antibodies (with the human variable region) in the second column. The second column also includes a single antibody with a mouse variable domain and a mouse Fc domain. The subsequent columns provide the variable light chain/variable heavy chain amino acid sequences and nucleic acid sequences, and the last column provides the amino acid sequences of the set of six CDRs that map to each Antibody.

Table 6

Human Antibody: Protein S Antibody with Human Fc	Rat Antibody: Protein S Antibody with Rat Fc	Variable Light Chain, Variable Heavy Chain (Amino Acid Sequences)	Variable Light Chain, Variable Heavy Chain (Nucleic Acid Sequences)	CDR Combination (Amino Acid Sequences)
Antibody 1	Antibody 13	SEQ ID NO: 71, SEQ ID NO: 72	SEQ ID NO: 95, SEQ ID NO: 96	SEQ ID NO: 2, SEQ ID NO: 13, SEQ ID NO: 22, SEQ ID NO: 34, SEQ ID NO: 46, SEQ ID NO: 58
Antibody 2	Antibody 14	SEQ ID NO: 73, SEQ ID NO: 74	SEQ ID NO: 97, SEQ ID NO: 98	SEQ ID NO: 3, SEQ ID NO: 14, SEQ ID NO: 23, SEQ ID NO: 35, SEQ ID NO: 47, SEQ ID NO: 59
Antibody 3	Antibody 15	SEQ ID NO: 75, SEQ ID NO: 76	SEQ ID NO: 99, SEQ ID NO: 100	SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 36, SEQ ID NO: 48, SEQ ID NO: 60

Human Antibody: Protein S Antibody with Human Fc	Rat Antibody: Protein S Antibody with Rat Fc	Variable Light Chain, Variable Heavy Chain (Amino Acid Sequences)	Variable Light Chain, Variable Heavy Chain (Nucleic Acid Sequences)	CDR Combination (Amino Acid Sequences)
Antibody 4	Antibody 16	SEQ ID NO: 77, SEQ ID NO: 78	SEQ ID NO: 101, SEQ ID NO: 102	SEQ ID NO: 4, SEQ ID NO: 16, SEQ ID NO: 25, SEQ ID NO: 37, SEQ ID NO: 49, SEQ ID NO: 61
Antibody 6	Antibody 18	SEQ ID NO: 91, SEQ ID NO: 92	SEQ ID NO: 115, SEQ ID NO: 116	SEQ ID NO: 11, SEQ ID NO: 20, SEQ ID NO: 32, SEQ ID NO: 44, SEQ ID NO: 56, SEQ ID NO: 68
Antibody 7	Antibody 19	SEQ ID NO: 69, SEQ ID NO: 70	SEQ ID NO: 93, SEQ ID NO: 94	SEQ ID NO: 1, SEQ ID NO: 12, SEQ ID NO: 21, SEQ ID NO: 33, SEQ ID NO: 45, SEQ ID NO: 57
Antibody 8	Antibody 20	SEQ ID NO: 79, SEQ ID NO: 80	SEQ ID NO: 103, SEQ ID NO: 104	SEQ ID NO: 5, SEQ ID NO: 17, SEQ ID NO: 26, SEQ ID NO: 38, SEQ ID NO: 50, SEQ ID NO: 62
Antibody 9	Antibody 21	SEQ ID NO: 81, SEQ ID NO: 82	SEQ ID NO: 105, SEQ ID NO: 106	SEQ ID NO: 6, SEQ ID NO: 18, SEQ ID NO: 27, SEQ ID NO: 39, SEQ ID NO: 51, SEQ ID NO: 63
Antibody 11	Antibody 23	SEQ ID NO: 87, SEQ ID NO: 88	SEQ ID NO: 111, SEQ ID NO: 112	SEQ ID NO: 9, SEQ ID NO: 19, SEQ ID NO: 30, SEQ ID NO: 42, SEQ ID NO: 54, SEQ ID NO: 66

Human Antibody: Protein S Antibody with Human Fc	Rat Antibody: Protein S Antibody with Rat Fc	Variable Light Chain, Variable Heavy Chain (Amino Acid Sequences)	Variable Light Chain, Variable Heavy Chain (Nucleic Acid Sequences)	CDR Combination (Amino Acid Sequences)
Antibody 12	Antibody 24	SEQ ID NO: 89, SEQ ID NO: 90	SEQ ID NO: 113, SEQ ID NO: 114	SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 31, SEQ ID NO: 43, SEQ ID NO: 55, SEQ ID NO: 67
Antibody 29	Antibody 35	SEQ ID NO: 148, SEQ ID NO: 149	SEQ ID NO: 162, SEQ ID NO: 163	SEQ ID NO: 117, SEQ ID NO: 118, SEQ ID NO: 119, SEQ ID NO: 120, SEQ ID NO: 121, SEQ ID NO: 122
Antibody 30	Antibody 36	SEQ ID NO: 150, SEQ ID NO: 151	SEQ ID NO: 164, SEQ ID NO: 165	SEQ ID NO: 117, SEQ ID NO: 118, SEQ ID NO: 123, SEQ ID NO: 124, SEQ ID NO: 125, SEQ ID NO: 126
Antibody 25	Antibody 31	SEQ ID NO: 152, SEQ ID NO: 153	SEQ ID NO: 166, SEQ ID NO: 167	SEQ ID NO: 127, SEQ ID NO: 128, SEQ ID NO: 129, SEQ ID NO: 130, SEQ ID NO: 50, SEQ ID NO: 131
Antibody 27	Antibody 33	SEQ ID NO: 156, SEQ ID NO: 157	SEQ ID NO: 170, SEQ ID NO: 171	SEQ ID NO: 136, SEQ ID NO: 17, SEQ ID NO: 137, SEQ ID NO: 138, SEQ ID NO: 139, SEQ ID NO: 140
Antibody 28	Antibody 34* (*mouse variable region/mouse Fc)	SEQ ID NO: 158, SEQ ID NO: 159	SEQ ID NO: 172, SEQ ID NO: 173	SEQ ID NO: 141, SEQ ID NO: 142, SEQ ID NO: 143, SEQ ID NO: 144, SEQ ID NO: 145, SEQ ID NO: 146

Human Antibody: Protein S Antibody with Human Fc	Rat Antibody: Protein S Antibody with Rat Fc	Variable Light Chain, Variable Heavy Chain (Amino Acid Sequences)	Variable Light Chain, Variable Heavy Chain (Nucleic Acid Sequences)	CDR Combination (Amino Acid Sequences)
Antibody 37	Antibody 38	SEQ ID NO: 177, SEQ ID NO: 182	SEQ ID NO: 178, SEQ ID NO: 183	SEQ ID NO: 174, SEQ ID NO: 175, SEQ ID NO: 176, SEQ ID NO: 179, SEQ ID NO: 180, SEQ ID NO: 181
Antibody 39	Antibody 40	SEQ ID NO: 187, SEQ ID NO: 192	SEQ ID NO: 188, SEQ ID NO: 193	SEQ ID NO: 184, SEQ ID NO: 185, SEQ ID NO: 186, SEQ ID NO: 189, SEQ ID NO: 190, SEQ ID NO: 191
Antibody 41	Antibody 42	SEQ ID NO: 197, SEQ ID NO: 202	SEQ ID NO: 198, SEQ ID NO: 203	SEQ ID NO: 194, SEQ ID NO: 195, SEQ ID NO: 196, SEQ ID NO: 199, SEQ ID NO: 200, SEQ ID NO: 201
Antibody 43	Antibody 44	SEQ ID NO: 207, SEQ ID NO: 212	SEQ ID NO: 208, SEQ ID NO: 213	SEQ ID NO: 204, SEQ ID NO: 205, SEQ ID NO: 206, SEQ ID NO: 209, SEQ ID NO: 210, SEQ ID NO: 211

Example 2: Assessing Thrombin Generation in APC and TFPI Cofactor Activity Screening Assays

Assessing APC and TFPI Cofactor Activity with Screening Assays

[0577] FIGS. 2A-2B depict exemplary assays performed to assess APC and TFPI cofactor activity, respectively and demonstrate how a APC cofactor inhibitor and a TFPI cofactor inhibitor would behave in the assay.

[0578] FIGS. 2A-2B depict assays performed to assess APC and TFPI cofactor activity, respectively. For the APC cofactor activity screening assay depicted in FIG. 2A, the following were tested: Protein S depleted platelet poor plasma, Protein S depleted plasma reconstituted with a Protein S and APC mixture, and Protein S depleted plasma reconstituted with a Protein S and APC mixture in the presence of varying concentrations of an anti-Protein S monoclonal antibody (mAb). The Protein S and APC mixture was made by pre-mixing Protein S and APC and adding to the Protein S depleted plasma to create a Protein S-dependent assay, due to APC not showing inhibition of thrombin generation in the absence of Protein S. As shown, the addition of the pre-mixed Protein S and APC mixture to Protein S depleted plasma shows significantly less tissue factor-induced thrombin generation than the Protein S depleted plasma alone. Addition of anti-Protein S mAbs to the Protein S depleted plasma + Protein S and APC mixture restored tissue factor-induced thrombin generation in a concentration-dependent manner.

[0579] For the TFPI cofactor activity screening assay depicted in FIG. 2B, the following were tested: platelet rich plasma with no treatment (also referred to herein as platelet-rich plasma, or PRP), a commercial Factor VIII (FVIII) neutralizing antibody, and a FVIII antibody and Protein S polyclonal antibody (pAB) mixture. The FVIII antibody acts as a neutralizing antibody and inhibits tissue factor induced thrombin generation. Addition of the neutralizing Protein S antibodies enhanced thrombin generation in the presence of the FVIII antibody. Neutralizing Protein C antibodies did not show an effect on thrombin generation in the TFPI cofactor activity assay, suggesting that endogenous Protein C or activated Protein C does not inhibit thrombin generation in this assay.

[0580] Clones M1-M40 were generated from a rat hybridoma library described in Example 1. The cofactor activity assays performed as described when referring to FIGS. 2A-2B were used to measure the amount of thrombin generation in the presence of clones M1-M40 generated from a rat hybridoma library. FIGS. 3-4 depict the peak thrombin levels generated from an APC cofactor screening assay (APC assay) and a TFPI cofactor screening assay, respectively, of the clones M1-M40. Controls used for the APC assay depicted in FIG. 3 were platelet-poor plasma ("PPP") alone and PPP with APC. Controls used for the TFPI assay depicted in FIG. 4 were platelet-rich plasma ("PRP") alone and PRP with a commercial neutralizing FVIII antibody.

Example 3: Characterization of Selected Protein S Antibodies

Characterization of Inhibitor Type by Cofactor Activity Assays

[0581] As a general matter, Protein S antibodies can be identified as either a dual inhibitor of Protein S cofactor activity for both APC and TFPI, for APC only (APC cofactor inhibitor), or for TFPI (TFPI cofactor inhibitor) only. The antibodies can be categorized by assessment of the thrombin generation profiles shown when assaying each antibody using the cofactor activity assays described in FIGS. 2A-2B.

[0582] FIGS. 5-6 depict thrombin generation in the presence of Antibody 13 in the TFPI cofactor assay (FIG. 5) and the APC cofactor assay (FIG. 6). As shown in the TFPI cofactor activity assay using PRP, robust thrombin generation occurred in the absence of FVIII antibody and was reduced by addition of the FVIII antibody. Addition of 267 nM Antibody 13 promoted thrombin generation in the presence of the FVIII antibody. As shown in the APC cofactor activity assay, thrombin generation was reduced by addition of both APC and Protein S to Protein S depleted plasma. Addition of 267 nM Antibody 13 to Protein S depleted plasma containing APC and Protein S rescued thrombin generation. Since Antibody 13 induced thrombin generation in both the TFPI and APC cofactor assays, Antibody 13 was characterized as a dual inhibitor.

[0583] FIGS. 7-8 depict the thrombin generation of Antibody 21 when using a TFPI cofactor assay (FIG. 7) and an APC cofactor assay (FIG. 8). As shown in the TFPI cofactor activity assay depicted in FIG. 7, in PRP, robust thrombin generation occurred in the absence of FVIII antibody. Thrombin generation was reduced by addition of the FVIII antibody only. Addition of 267 nM Antibody 21 showed no rescue of the thrombin generation. As shown in the APC cofactor activity assay depicted in FIG. 8, thrombin generation was reduced by addition of a mixture of APC and Protein S, which was rescued by the addition of 267 nM Antibody 21. Because Antibody 21 did not induce thrombin generation in the TFPI cofactor assay but did rescue thrombin generation in the APC cofactor assay, Antibody 21 was characterized as an APC cofactor inhibitor.

[0584] FIGS. 9-10 depict the thrombin generation of Antibody 23 when using a TFPI cofactor assay (FIG. 9) and an APC cofactor assay (FIG. 10). As shown in the TFPI cofactor activity assay depicted in FIG. 9, in PRP, robust thrombin generation occurred in the absence of FVIII antibody. Thrombin generation was reduced by addition of the FVIII antibody only. Addition of

267 nM Antibody 23 promoted thrombin generation. As shown in the APC cofactor activity assay depicted in FIG. 10, in Protein S depleted plasma, thrombin generation was reduced by addition of a mixture of APC and Protein S. Addition of 267 nM Antibody 23 to Protein S depleted plasma with APC + Protein S showed limited rescue of the thrombin generation. Since Antibody 23 induced thrombin generation in the TFPI cofactor assay but not in the APC cofactor assay, Antibody 23 was characterized as a TFPI cofactor inhibitor.

Characterization of Protein S Antibodies

[0585] The assay results depicted in FIGS. 11A-20H were used to characterize the exemplary Protein S antibodies provided herein. The antibodies were characterized as an inhibitor of APC cofactor activity (APC cofactor inhibitor), an inhibitor of TFPI cofactor activity (TFPI cofactor inhibitor), or a dual inhibitor. Binding regions and calcium dependence were also determined. The results of the characterization assays are summarized in Tables 7 and 8 below and described in further detail in the below sections.

Table 7

Protein S Antibody	Fc	Pathway Inhibitor (APC, TFPI, or Dual)	IC50 APC (nM)	Hill Coefficient	IC50 TFPI (nM)	Hill Coefficient
Antibody 13	Rat	Dual	5.30E-09	2.56	5.11E-08	5.62
Antibody 14	Rat	Dual	8.76E-09	2.26	7.33E-08	2.46
Antibody 15	Rat	Dual	1.35E-08	2.80	7.73E-08	3.79
Antibody 16	Rat	Dual	1.40E-08	2.69	4.63E-08	3.9
Antibody 18	Rat	APC only	1.28E-08	2.7	N/A	N/A
Antibody 19	Rat	Dual	8.12E-08	1.06	1.05E-07	1.39
Antibody 20	Rat	Dual	1.28E-08	2.70	N/A	N/A
Antibody 21	Rat	APC only	6.23E-09	2.10	N/A	N/A
Antibody 23	Rat	TFPI only	N/A	N/A	<4.4E-09	N/A

Protein S Antibody	Fc	Pathway Inhibitor (APC, TFPI, or Dual)	IC50 APC (nM)	Hill Coefficient	IC50 TFPI (nM)	Hill Coefficient
Antibody 24	Rat	Dual	1.79E-08	1.6	7.36E-08	1.59
Antibody 31	Rat	APC only	9.90E-08	1.12	N/A	N/A
Antibody 33	Rat	Dual	5.81E-08	2.48	3.4E-08	2.21
Antibody 34	Mouse	APC only	9.58E-08	2.28	N/A	N/A
Antibody 35	Rat	APC only	3.71E-07	0.68	N/A	N/A
Antibody 36	Rat	Dual	1.53E-08	1.55	6.05E-09	3.53
Antibody 38	Rat	APC only	8.40E-08	0.70	N/A	N/A
Antibody 40	Rat	APC only	1.19E-08	2.82	N/A	N/A
Antibody 42	Rat	APC only	2.15E-07	1.37	N/A	N/A
Antibody 44	Rat	APC only	4.92E-07	0.87	N/A	N/A

Table 8

Protein S Antibody with Rat Fc	Pathway Inhibitor	Binds Heavy Chain	Calcium-Dependent Binding	Binds Thrombin-Sensitive Region of Protein S
Antibody 13	Dual	Yes	No	No
Antibody 14	Dual	Yes	No	No
Antibody 15	Dual	Yes	Yes	Yes
Antibody 16	Dual	Yes	Yes	No
Antibody 19	Dual	Yes	Yes	Yes
Antibody 20	Dual	Yes	Yes	No
Antibody 24	Dual	Yes	Yes	Yes
Antibody 18	APC	Yes	Yes	No
Antibody 21	APC	Yes	No	No
Antibody 23	TFPI	No	No	No
Antibody 31	APC	No	No	No
Antibody 33	Dual	No	Yes	No
Antibody 34	APC	No	Yes	No
Antibody 35	APC	No	Yes	No

Antibody 36	Dual	No	Yes	No
Antibody 38	APC	No	Yes	No
Antibody 40	APC	No	Yes	No
Antibody 42	APC	No	No	No
Antibody 44	APC	Yes	No	No

[0586] Binding characteristics of the Protein S antibodies were also determined. Binding interaction analysis was obtained by Surface Plasmon Resonance in a Biacore X100 instrument. The human anti-IgG (Fc) antibody was immobilized to the carboxymethylated dextran membrane on the sensor chip surface (CM5) via the free amine method for a contact time of 420 seconds. Human anti-IgG antibody at 25 µg/mL in 0.15 M NaCl was immobilized to 9,785 RU in 10 mM sodium acetate pH 5.0. Each tested monoclonal antibody (Antibody 1-12) was captured at a fixed concentration (0.25 µg/mL) with immobilized human anti-IgG antibody. Experimentally, the capture ligand level for the antibodies tested was determined to be 66-99 RL, corresponding to a R_{max} of 61-91 RU. No signs of mass transport limitation were observed.

[0587] The following experimental conditions were used: purified Protein S was injected at concentrations: 2-128 nM, 1:2-fold dilution. Dilution and running buffer were as follows: Hbs-EP + 5 mM CaCl₂. Regeneration was obtained by 3 M MgCl₂. Binding constants (k_a , k_d , and KD) were determined using the Biacore X100 evaluation software, assuming a 1:1 interaction of Protein S and the tested monoclonal antibody under investigation. The resulting data are presented in Table 9 below.

Table 9

	K_a (1/MS)	K_d (1/S)	KD nM
Antibody 1	2.90E+05	3.90E-04	1.33
Antibody 2	3.46E+05	5.60E-04	1.63
Antibody 3	2.53E+05	5.69E-04	2.24
Antibody 4	3.99E+05	1.65E-03	4.13
Antibody 6	2.80E+05	1.60E-03	5.79
Antibody 7	2.52E+05	5.00E-03	20.00
Antibody 8	1.32E+05	8.23E-04	6.25
Antibody 9	2.12E+05	3.36E-03	15.90
Antibody 11	1.16E+05	1.13E-03	9.60

Antibody 12	6.57E+05	9.47E-03	14.40
-------------	----------	----------	-------

[0588] Binding characteristics of the Protein S antibodies were also further characterized using Octet. Using the Octet System (Sartorius), the binding of each antibody to both human and cynomolgus monkey (“cyno” in Table 10) Protein S was determined. The human Fc antibodies were immobilized onto anti-human Fc capture probes by placing the probes into 10 µg/ml antibody solution in 10 mg/ml bovine serum albumin, 20 mM Tris pH 7.0, 150 mM NaCl, and 4 mM calcium chloride. Next, the bound antibodies were placed into solutions containing 500 nM, 250 nM, 125 nM, 62.5 nM, 31.25 nM, 15.625 nM and 7.81 nM Protein S and the association rates were measured. Next, the probes were placed into buffer and the dissociation rates were measured. A summary of the resulting data is provided in Table 10 below.

Table 10

Antibody	k_{on} (1/Ms)	k_{dis} (1/s)	KD (human) (M)	k_{on} (1/Ms)	k_{dis} (1/s)	KD (cyno) (M)
Antibody 1	1.69E+05	<1.0E-07	<1.0E-12	1.39E+05	<1.0E-07	<1.0E-12
Antibody 2	2.07E+05	<1.0E-07	<1.0E-12	1.54E+05	<1.0E-07	<1.0E-12
Antibody 3	1.64E+05	8.86E-06	5.42E-11	1.27E+05	<1.0E-07	<1.0E-12
Antibody 4	2.71E+05	7.08E-04	2.62E-09	2.01E+05	3.08E-03	1.53E-08
Antibody 6	2.26E+05	3.32E-04	1.47E-09	1.40E+05	3.94E-04	2.81E-09
Antibody 7	1.72E+05	1.41E-03	8.18E-09	1.37E+05	2.32E-03	1.69E-08
Antibody 8	1.31E+05	6.70E-05	5.14E-10	8.58E+04	1.33E-04	1.55E-09
Antibody 9	3.04E+05	1.62E-04	5.33E-10	1.66E+05	4.42E-05	2.66E-10
Antibody 11	8.52E+04	3.40E-05	3.99E-10	8.21E+04	<1.0E-07	<1.0E-12
Antibody 12	3.09E+05	2.32E-03	7.49E-09	2.58E+05	2.91E-03	1.13E-08
Antibody 25	1.43E+05	<1.0E-07	<1.0E-12	1.24E+05	<1.0E-07	<1.0E-12
Antibody 27	2.77E+05	1.59E-04	5.74E-10	6.79E+04	2.33E-03	3.43E-08
Antibody 28	1.56E+05	1.06E-03	6.78E-09	1.09E+05	1.06E-03	9.81E-09
Antibody 29	8.19E+04	1.73E-03	2.11E-08	8.82E+04	1.77E-03	2.00E-08

Antibody	k_{on} (1/Ms)	k_{dis} (1/s)	KD (human) (M)	k_{on} (1/Ms)	k_{dis} (1/s)	KD (cyno) (M)
Antibody 30	2.42E+04	2.79E-04	1.15E-08	3.50E+04	7.30E-04	2.09E-08
Antibody 37	2.18E+05	8.32E-05	3.82E-10	1.77E+05	1.19E-04	6.72E-10
Antibody 39	2.34E+05	5.84E-05	2.50E-10	1.91E+05	8.02E-05	4.19E-10
Antibody 41	8.00E+04	1.10E-03	1.38E-08	9.64E+04	1.88E-03	1.95E-08
Antibody 43	3.20E+05	6.36E-04	1.99E-09	2.68E+05	1.04E-03	3.88E-09

Antibody 19 and Antibody 7

[0589] FIGS. 11A-11H depict the characterization of Antibody 19 and Antibody 7. Antibody 19 and Antibody 7 both comprise the same human variable regions, while Antibody 19 comprises a rat Fc domain and Antibody 7 comprises a human IgG4 Fc domain.

[0590] FIG. 11A depicts a Western blot showing that Antibody 19 binds Protein S in the Thrombin Sensitive Region (TSR) of Protein S. A full-length human Protein S protein (Reduced (Red) and Non-reduced (NR)) and a thrombin-cleaved Protein S fragment (cleaved in the Thrombin Sensitive Region (TSR), NR and Red) were used. By Western blot, Antibody 19 bound the full Protein S, but not the thrombin-cleaved Protein S, indicating that Antibody 19 binds at the TSR. Because no signal was observed with the reduced Protein S, the epitope for this antibody is not a linear epitope.

[0591] Binding to the heavy chain of Protein S was determined using an ELISA based method. Briefly, high binding ELISA plates were coated with recombinant heavy chain Protein S expressed and purified from HEK293 cells. The “heavy chain” of Protein S represents a fragment of Protein S spanning amino acids 42-296. The coated plate was blocked with 1% casein solution. Then, buffer containing 1 μ g/ml of antibody was applied to the well with buffer containing 1 mM calcium chloride. For antibodies that bound this fragment of Protein S, it was concluded that the epitope on Protein S for that antibody is between amino acids 42-296 of Protein S. Table 11 below summarizes the results of the heavy chain binding assay for all Antibodies. (antibodies that do bind this region have ODs greater than about 0.48 OD).

Table 11

Antibody	Binding to heavy chain (OD)
Antibody 13	1.63
Antibody 14	2.03
Antibody 15	1.78
Antibody 16	1.67
Antibody 18	0.60
Antibody 19	0.61
Antibody 20	1.54
Antibody 21	1.58
Antibody 23	0.10
Antibody 24	0.71
Antibody 31	0.13
Antibody 33	0.11
Antibody 34	0.16
Antibody 35	0.08
Antibody 36	0.08
Antibody 38	0.10
Antibody 40	0.11
Antibody 42	0.11
Antibody 44	0.48

[0592] Calcium dependence of binding was determined using an ELISA based method. Briefly, high binding ELISA plates were coated with human Protein S. Then, buffer containing 1 µg/ml of antibody was applied to the well with either buffer containing 1 mM EDTA or 1 mM calcium chloride. If the level of binding of Protein S was reduced dramatically (greater than 85% of the absorbance observed in the presence of calcium) when EDTA was added, it was concluded that the binding was calcium-dependent. Table 12 summarizes the results of the calcium dependence assay for all Antibodies.

Table 12

	Binding to Protein S in EDTA (OD)	Binding to Protein S in CaCl ₂ (OD)
Antibody 13	0.48	1.77
Antibody 14	0.75	2.03
Antibody 15	0.07	1.86
Antibody 16	0.17	1.95
Antibody 18	0.10	1.96
Antibody 19	0.06	1.68
Antibody 20	0.26	1.93
Antibody 21	1.40	1.92
Antibody 23	0.26	0.57
Antibody 24	0.06	1.72
Antibody 31	0.63	1.03
Antibody 33	0.20	1.23
Antibody 34	0.06	0.51
Antibody 35	0.11	0.96
Antibody 36	0.03	0.60
Antibody 38	0.09	1.38
Antibody 40	0.11	0.51
Antibody 42	0.53	0.67
Antibody 44	0.89	1.17

[0593] FIGS. 11B-11D depict the results of an APC cofactor assay for Antibody 19 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0594] FIGS. 11E and 11F depict the results of a TFPI cofactor assay in human PRP using Antibody 7 and controls, respectively. A rabbit polyclonal human Protein S antibody labeled Dako was used as a positive control.

[0595] FIGS. 11G-11H depict the results of a PPP with APC assay performed with Antibody 7, and controls, respectively.

[0596] These results indicate that Antibody 19 and Antibody 7 are dual inhibitors.

Antibody 13 and Antibody 1

[0597] FIGS. 12A-12H depict the characterization of Antibody 13 and Antibody 1, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0598] FIG. 12A depicts a Western blot showing that Antibody 13 does not bind to Protein S in the TSR of Protein S. The Western blot was carried out as described for FIG. 11A. By Western blot, Antibody 13 bound the full Protein S, the thrombin-cleaved Protein S, the reduced Protein S, and the reduced thrombin-cleaved Protein S, indicating that Antibody 13 does not bind Protein S at the TSR. Binding to reduced Protein S showed that the epitope for Antibody 13 is a linear epitope.

[0599] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described herein. Antibody 13 was determined to bind to the heavy chain of Protein S. Calcium dependence was determined, also as described above, and Antibody 13 showed calcium-independent binding.

[0600] FIGS. 12B-12D depict the results of an APC cofactor assay for Antibody 13 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0601] FIGS. 12E and 12F depict the results of a TFPI cofactor assay in human PRP using Antibody 1 and controls, respectively. A rabbit polyclonal human Protein S antibody labeled Dako was used as a positive control.

[0602] FIGS. 12G-12H depict the results of the APC cofactor assay performed with Antibody 1, and controls, respectively.

[0603] These results indicate that Antibody 13 and Antibody 1 are dual inhibitors.

Antibody 14 and Antibody 2

[0604] FIGS. 13A-13H depict the characterization of Antibody 14 and Antibody 2, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0605] FIG. 13A depicts a Western blot showing that Antibody 14 does not bind Protein S in the TSR of Protein S. The Western blot was carried out as described for FIG. 11A. By Western blot, Antibody 14 bound the full Protein S and the thrombin-cleaved Protein S, but did not bind the reduced Protein S, and the reduced thrombin-cleaved Protein S. These results indicate that Antibody 14 does not bind Protein S at the TSR, and the lack of signal observed with the reduced Protein S showed that the epitope for this antibody is not a linear epitope.

[0606] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described herein. Antibody 14 was determined to bind to the heavy chain of Protein S. Calcium dependence was determined, also as described above, and Antibody 14 showed binding that was calcium-independent.

[0607] FIGS. 13B-13D depict the results of an APC cofactor assay for Antibody 14 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0608] FIGS. 13E and 13F depict the results of a TFPI cofactor assay in human PRP using Antibody 2 and controls, respectively. A rabbit polyclonal human Protein S antibody labeled Dako was used as a positive control.

[0609] FIGS. 13G-13H depict the results of the APC cofactor assay performed with Antibody 2 and controls, respectively.

[0610] These results indicate that Antibody 14 and Antibody 2 are dual inhibitors.

Antibody 15 and Antibody 3

[0611] FIGS. 14A-14H depict the characterization of Antibody 15 and Antibody 3, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0612] FIG. 14A depicts a Western blot showing that Antibody 15 binds Protein S in the TSR region of Protein S. The Western blot was carried out as described for FIG. 11A. By Western blot, Antibody 15 bound the full Protein S and showed a small band for the thrombin-cleaved Protein S, but did not bind the reduced Protein S, and the reduced thrombin-cleaved Protein S. These results indicate that Antibody 15 binds Protein S at the TSR, and the lack of signal observed with the reduced Protein S showed that the epitope for this antibody is not a linear epitope.

[0613] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described above. Antibody 15 was determined to bind to the heavy chain of Protein S. Calcium dependence was determined, also as described above, and Antibody 15 showed binding that was calcium-dependent.

[0614] FIGS. 14B-14D depict the results of an APC cofactor assay for Antibody 15 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0615] FIGS. 14E and 14F depict the results of a TFPI cofactor assay in human PRP using Antibody 3 and controls, respectively. A rabbit polyclonal human Protein S antibody labeled Dako was used as a positive control.

[0616] FIGS. 14G-14H depict the results of the APC cofactor assay performed with Antibody 3 and controls, respectively.

[0617] These results indicate that Antibody 15 and Antibody 3 are dual inhibitors.

Antibody 16 and Antibody 4

[0618] FIGS. 15A-15H depict the characterization of Antibody 16 and Antibody 4, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0619] FIG. 15A depicts a Western blot showing that Antibody 16 binds Protein S in the Gla-domain of Protein S. The Western blot was carried out as described for FIG. 11A. By Western blot, Antibody 16 bound the full Protein S, the thrombin-cleaved Protein S, the reduced Protein S, but did not bind the reduced thrombin-cleaved Protein S. These results indicate that Antibody 16 binds Protein S at the Gla-domain, and the signal observed with the reduced Protein S showed that the epitope for this antibody is a linear epitope.

[0620] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described herein. Antibody 16 was determined to bind to the heavy chain of Protein S. Calcium dependence was determined, also as described herein, and Antibody 16 showed binding that was calcium-dependent.

[0621] FIGS. 15B-15D depict the results of an APC cofactor assay for Antibody 16 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0622] FIGS. 15E and 15F depict the results of a TFPI cofactor assay in human PRP using Antibody 4 and controls, respectively. A rabbit polyclonal human Protein S antibody labeled Dako was used as a positive control.

[0623] FIGS. 15G-15H depict the results of the APC cofactor assay performed with Antibody 4, and controls, respectively.

[0624] These results indicate that Antibody 16 and Antibody 4 are dual inhibitors.

Antibody 20 and Antibody 8

[0625] FIGS. 16A-16H depict the characterization of Antibody 20 and Antibody 8, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0626] FIG. 16A depicts a Western blot showing that Antibody 20 does not bind Protein S in the TSR of Protein S. The Western blot was carried out as described for FIG. 11A. By Western blot, Antibody 20 bound the full Protein S and the thrombin-cleaved Protein S, but did not bind the reduced Protein S or the reduced thrombin-cleaved Protein S. These results indicate that Antibody 20 does not bind Protein S at the TSR, and the lack of signal observed with the reduced Protein S showed that the epitope for this antibody is not a linear epitope.

[0627] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described herein. Antibody 20 was determined to bind to the heavy chain of Protein S. Calcium dependence was determined, also as described herein, and Antibody 20 showed binding that was calcium-dependent.

[0628] FIGS. 16B-16D depict the results of an APC cofactor assay for Antibody 20 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0629] FIGS. 16E and 16F depict the results of a TFPI cofactor assay in human PRP using Antibody 8 and controls, respectively. A rabbit polyclonal human Protein S antibody labeled Dako was used as a positive control.

[0630] FIGS. 16G-16H depict the results of the APC cofactor assay performed with Antibody 8 and controls, respectively.

[0631] These results indicate that Antibody 20 and Antibody 8 are dual inhibitors.

Antibody 21 and Antibody 9

[0632] FIGS. 17A-17H depict the characterization of Antibody 21 and Antibody 9, antibodies sharing the same human variable region, and are characterized as APC cofactor inhibitors.

[0633] FIG. 17A depicts a Western blot showing that Antibody 21 does not bind Protein S in the TSR of Protein S. The Western blot was carried out as described for FIG. 11A. By Western blot, Antibody 21 bound the full Protein S and the thrombin-cleaved Protein S, but did not bind the reduced Protein S or the reduced thrombin-cleaved Protein S. These results indicate that Antibody 21 does not bind Protein S at the TSR, and the lack of signal observed with the reduced Protein S showed that the epitope for this antibody is not a linear epitope.

[0634] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described herein. Antibody 21 was determined to bind to the heavy chain of Protein S. Calcium dependence was determined, also as described herein, and Antibody 21 showed binding that was calcium-independent.

[0635] FIGS. 17B-17D depict the results of an APC cofactor assay for Antibody 21 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0636] FIGS. 17E and 17F depict the results of a TFPI cofactor assay in human PRP using Antibody 9 and controls, respectively. A rabbit polyclonal human Protein S antibody labeled Dako was used as a positive control.

[0637] FIGS. 17G-17H depict the results of the APC cofactor assay performed with Antibody 9 and controls, respectively.

[0638] These results indicate that Antibody 21 and Antibody 9 are APC cofactor inhibitors.

Antibody 23 and Antibody 11

[0639] FIGS. 18A-18G depict the characterization of Antibody 23 and Antibody 11, antibodies sharing the same human variable region, and are characterized as TFPI cofactor inhibitors.

[0640] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described herein. Antibody 23 was determined to not bind to the heavy chain of Protein S. Calcium dependence was determined, also as described herein, and Antibody 23 showed binding that was calcium-independent.

[0641] FIGS. 18A-18C depict the results of a APC cofactor assay for Antibody 23 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0642] FIGS. 18D and 18E depict the results of a TFPI cofactor assay in human PRP using Antibody 11 and controls, respectively.

[0643] FIGS. 18F and 18G depict the results of a APC cofactor assay with Antibody 11 and controls, respectively.

[0644] These results indicate that Antibody 23 and Antibody 11 are TFPI cofactor inhibitors.

Antibody 24 and Antibody 12

[0645] FIGS. 19A-19H depict the characterization of Antibody 24 and Antibody 12, antibodies sharing the same human variable region, and are characterized as dual inhibitors.

[0646] FIG. 19A depicts a Western blot showing that Antibody 24 binds Protein S in the TSR of Protein S. The Western blot was carried out as described for FIG. 11A. By Western blot, Antibody 24 bound the full Protein S but did not bind the thrombin-cleaved Protein S. Antibody 24 did not bind the reduced Protein S or the reduced thrombin-cleaved Protein S. These results indicate that Antibody 24 binds Protein S at the TSR, and the lack of signal observed with the reduced Protein S showed that the epitope for this antibody is not a linear epitope.

[0647] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described herein. Antibody 24 was determined to bind to the heavy chain of Protein S. Calcium dependence was determined, also as described herein, and Antibody 24 showed binding that was calcium-dependent.

[0648] FIGS. 19B-19D depict the results of a APC cofactor assay for Antibody 24 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0649] FIGS. 19E and 19F depict the results of a TFPI cofactor assay in human PRP using Antibody 12 and controls, respectively.

[0650] FIGS. 19G and 19H depict the results of a APC cofactor assay with Antibody 12 and controls, respectively.

[0651] These results indicate that Antibody 24 and Antibody 12 are dual inhibitors.

Antibody 18 and Antibody 6

[0652] FIGS. 20A-20H depict the characterization of Antibody 18 and Antibody 6, antibodies sharing the same human variable region, and are characterized as APC cofactor inhibitors.

[0653] FIG. 20A depicts a Western blot showing that Antibody 18 does not bind Protein S in the TSR of Protein S. The Western blot was carried out as described for FIG. 20A. By Western blot, Antibody 18 bound the full Protein S and the thrombin-cleaved Protein S, but did not bind the reduced Protein S or the reduced thrombin-cleaved Protein S. These results indicate that Antibody 18 does not bind Protein S at the TSR, and the lack of signal observed with the reduced Protein S showed that the epitope for this antibody is not a linear epitope.

[0654] Binding to the heavy chain of Protein S was determined using an ELISA based method, as described herein. Antibody 18 was determined to bind to the heavy chain of Protein S. Calcium dependence was determined, also as described herein, and Antibody 18 showed binding that was calcium-dependent.

[0655] FIGS. 20B-20D depict the results of a APC cofactor assay for Antibody 18 using Protein S deficient human PPP, cynomolgus monkey PPP, and a TFPI cofactor assay using human PRP, respectively.

[0656] FIGS. 20E and 20F depict the results of a TFPI cofactor assay in human PRP using Antibody 6 and controls, respectively.

[0657] FIGS. 20G and 20H depict the results of a APC cofactor assay with Antibody 6 and controls, respectively.

[0658] These results indicate that Antibody 18 and Antibody 6 are APC cofactor inhibitors.

[0659] A summary of the characterizations exemplified above is presented in Table 13 below.

Table 13

Human Antibody: Corresponding Protein S Antibody with Human Fc	Rat Antibody: Protein S Antibody with Rat Fc	Inhibitor Type
Antibody 1	Antibody 13	Dual
Antibody 2	Antibody 14	Dual
Antibody 3	Antibody 15	Dual
Antibody 4	Antibody 16	Dual
Antibody 6	Antibody 18	APC cofactor
Antibody 7	Antibody 19	Dual
Antibody 8	Antibody 20	Dual
Antibody 9	Antibody 21	APC cofactor
Antibody 11	Antibody 23	TFPI cofactor
Antibody 12	Antibody 24	Dual

Example 4: Thrombin Generation Improvement in Plasma Samples from Patients with Various Factor Deficiencies and von Willebrand Disease

[0660] FIGS. 21A-21G depict enhanced thrombin generation when Antibody 15 is added to various samples of congenital factor deficient plasma containing soluble thrombomodulin, a cofactor of thrombin that is involved in conversion of thrombin to an anti-coagulant enzyme. Plasma samples were taken from patients having various factor deficiencies and conditions: Factor VII, Factor VIII, Factor IX, or Factor XI deficiencies, and von Willebrand disease (vWD) type 1, 2, or 3.

[0661] FIG. 21A depicts soluble tissue factor added to congenital Factor VII deficient plasma in the presence or absence of 40 nM soluble thrombomodulin. The level of thrombin generated is measured over time. When Antibody 15 is added to the deficient plasma containing thrombomodulin, increased thrombin was observed. FIGS. 21B-21G depict data generated from experiments performed in a similar manner, but using congenital Factor VIII deficient plasma, congenital Factor IX deficient plasma, congenital Factor XI deficient plasma, vWD type 1 plasma, vWD type 2B plasma or vWD type 3 plasma, respectively. These results show that the tested antibodies were able to promote thrombin generation in samples having a factor deficiency or vWD disease.

Example 5: Activity of Protein S Antibodies Post-Treatment in Cynomolgus Monkeys Using D-Dimer as a Marker

[0662] FIGS. 22A and 22B depict D-dimer levels observed in cynomolgus monkeys treated with 1 mg/kg or 3 mg/kg, respectively, of Antibody 1. At varying times post-injection of the antibody, blood was collected in sodium citrate tubes. Plasma samples were analyzed for the presence of D-dimer, a fibrin degradation product which acts as a marker of coagulation. Increases in the levels of D-dimer indicated that the tested antibodies activate the coagulation cascade and remain active *in vivo* for a period of time after treatment. FIG. 22A shows that treatment with 1 mg/kg of Antibody 1 showed activity for approximately 70 hours after treatment, and FIG. 22B shows that treatment with 3 mg/kg of Antibody 1 showed activity for approximately 105-110 hours after treatment.

Example 6: Restoration of Fibrin Deposition by Protein S Antibodies

[0663] FIGS 23A-23C depict the level of fibrin deposited onto collagen spots in a microfluidic system. Certain coagulation factor deficiencies can cause fibrin deposition to be at levels lower than that of subjects not having a factor deficiency. The microfluidic system can allow measurement of coagulation occurring during flow of blood.

[0664] Normal human blood, from a patient without any factor deficiencies, was untreated or treated with Factor VIII (FVIII) neutralizing antibody. This was then passed over collagen spots in a microfluidic system. Strong fibrin deposition was observed in untreated blood, but not in treated blood. When Antibodies 1, 3 and 7 were added to the treated blood, fibrin deposition was

restored in a dose dependent manner. These results show that in blood neutralized with FVIII antibody, the Protein S antibodies disclosed herein may be used for restoration of fibrin formation or deposition, which may serve as a desired marker for coagulation activity.

Example 7: Binding of Protein S Antibodies to the Protein S-C4BP Complex

[0665] The Octet assay was used to determine the capability of the Protein S antibodies provided herein to bind to Protein S either alone, or when complexed to C4BP. A summary of the binding to Protein S or the Protein S-C4BP complex is provided in Table 14. Table 14 presents an X where a tested antibody was able to bind to either the Protein S-C4BP Complex, and/or the Protein S alone. A dash indicates that the binding did not occur.

[0666] Using the Octet System (Sartorius), the binding of each antibody to both Protein S and Protein S in complex with C4BP was determined. It had previously been determined that the C4BP preparation from Complement Technologies contains Protein S precomplexed to human C4BP. Therefore, this preparation represented the Protein S-C4BP complex and plasma purified Protein S from Haematologic Technologies was used as the source of free Protein S. The human Fc antibodies were immobilized onto anti-human Fc capture probes by placing the probes into 10 µg/ml antibody solution in 10 mg/ml bovine serum albumin, 20 mM Tris pH 7.0, 150 mM NaCl, and 4 mM calcium chloride. Then the bound antibodies were placed into solutions containing either 75 µg/ml Protein S or 100 µg/ml C4BP (C4BP-Protein S complex). With antibodies that do not bind Protein S in complex with C4BP, no binding is observed. With antibodies that bind Protein S in free form or in complex with C4BP, an association is observed under both conditions. All antibodies tested bound free Protein S, as depicted in Table 14.

Table 14

Antibody with Human Fc	Binding to ProS-C4BP Complex	Binding to ProS
Antibody 1	X	X
Antibody 2	X	X
Antibody 3	X	X
Antibody 4	X	X
Antibody 6	X	X
Antibody 7	X	X

Antibody with Human Fc	Binding to ProS-C4BP Complex	Binding to ProS
Antibody 8	X	X
Antibody 9	X	X
Antibody 11	-	X
Antibody 12	X	X
Antibody 25	-	X
Antibody 27	X	X
Antibody 28	X	X
Antibody 29	-	X
Antibody 30	-	X
Antibody 37	-	X
Antibody 39	-	X
Antibody 41	-	X
Antibody 43	X	X

Example 8: Effects of Protein S Antibodies on Fibrin Deposition

[0667] These experiments were carried out as follows. Bioflux 1000z 48-well high shear microfluidics plates (0-200 dynes/cm²) employed in the experiments were purchased from Bioflux. The device has the #1.5 borosilicate glass coverslip which forms the floor of the microfluidic channel engineer to facilitate the coated of collagen. Collagen type 1 was purchase from Chrono-Par collagen (chrono-Log Corp, Havertown, PA). Collagen was perfused from the wells at room temperature and incubated for 1 hour. Precoated plates with collagen type 1 were rinsed with PBS and channels were blocked with 0.5% (v/v) BSA for 10 min in PBS prior to the addition of the labeled blood to the wells. Sodium citrate anti-coagulated whole human blood (ALLCells, Oakland, CA) was used within 4-8h of collection. Whole blood was incubated for 1 hour with 100 µg/mL Sheep anti-Human Factor VIII (Haematologic Technologies, Essex, VT). The antibodies were added and incubated with whole blood. Fibrinogen from human plasma, Alexa Fluor 546 (Invitrogen, Carlsbad, CA) was added at a final concentration of 50 µg/mL prior to biological experiments. Whole blood was added to the input wells and perfused at 30 dyn/cm² using the BioFlux Controller and software. The samples were illuminated for no more than 30 ms for each capture. The BioFlux software imaging module was used to control the image acquisition settings and to process the fluorescence intensity measurements. Fluorescent micrographs were captured with the blood under flow using an inverted microscope (ZEISS

Axio Observer 7) and sCMOS Camera. Images were timelapse recorded using the BioFlux 1000 imaging system (Fluxion Biosciences). Data was processed using BioFlux Montage Software.

[0668] FIGS. 25A-25B depict the level of fibrin deposition resulting from the Protein S antibodies added to whole blood treated with Factor VIII neutralizing antibodies. This experiment tests the ability of the Protein S antibodies to restore or increase fibrin deposition in an *in vitro* microfluidic Hemophilia A model. Whole blood was treated with Factor VIII antibody to decrease fibrin deposition. FIG. 25A depicts the effects of Antibodies 12, and 3, and a control on fibrin deposition in anti-FVIII treated human plasma, and FIG. 25B depicts the effects of Antibodies 1 and 6 and a control on fibrin deposition in anti-FVIII treated human plasma. These results demonstrate that the Protein S antibodies 12, 3, 1 and 6 could restore or increase fibrin deposition in an *in vitro* microfluidic Hemophilia A model.

Example 9: Effect of Protein S Antibodies in an *in vitro* Microfluidics Hemophilia A Bleeding Model

[0669] These experiments were carried out using a fully endothelialized microfluidic system that was coupled to a microengineered pneumatic valve that mimics vascular damage (Sakurai, et al. Nature Communications 2018). Briefly, whole blood collected from healthy volunteers were treated with a sheep anti-human FVIII antibody to mimic whole blood from a hemophilia A patient. After treatment, the whole blood was perfused into the microfluidics system at which time an “injury” was introduced using the pneumatic valve. The localization of both platelets and fibrin at the site of “injury” was monitored and the time to cessation of “bleeding” was measured.

[0670] FIGS. 26A-26B depict the effects of Protein S antibodies in an *in vitro* microfluidics Hemophilia A bleeding model, measured by microfluidics experiments. These experiments were performed to measure the time of various Protein S antibodies to stop bleeding, and to determine the extent to which these antibodies induced fibrin deposition, respectively, in a microengineered, vascularized Hemophilia A bleeding model. The tested antibodies were Antibodies 6, 3, and 1. These results demonstrate that the antibodies were able to restore or increase fibrin deposition in a Hemophilia A bleeding model and therefore stop bleeding.

Example 10: Effect of Protein S Antibodies on Thrombin Generation

[0671] These experiments were carried out as follows. The Thrombin Generation Assay (TGA) was performed using a Thermo Fluoroskan Ascent Microplate Fluorometer and Thrombinoscope software. The PPP low reagent (Diagnostica Stago) was used in this experiment. Briefly, plasma from von Willebrand diseased patients was mixed with increasing levels of Protein S antibodies and incubated at room temperature. Then soluble human thrombomodulin was added prior to initiation of the reaction. To start the reaction, PPP low reagent was added along with calcium and the thrombin substrate. The levels of thrombin were then monitored.

[0672] FIGS. 27A-27F depict various results of enhanced dose-dependent thrombin generation when Protein S antibodies were added to plasma obtained from patients with various types of von Willebrand disease, with thrombomodulin added. These results demonstrate that the Protein S antibodies could effectively increase thrombin generation in a dose-dependent manner, and that different Protein S antibodies were more efficacious for different types of von Willebrand disease. Generally, the tested antibodies were more effective at increasing thrombin generation for Type 1 than Types 2A, 2B, and 3.

[0673] FIGS. 27A-27B depict enhanced dose-dependent thrombin generation when Antibodies 1 and 2 were added to plasma obtained from patients with various types of von Willebrand disease. FIGS. 27C-27D depict enhanced dose-dependent thrombin generation when Antibodies 6 and 7 were added to plasma obtained from patients with various types of von Willebrand disease, with thrombomodulin added. FIGS. 27E-27F depict enhanced dose-dependent thrombin generation when Antibodies 12 and 29 were added to plasma obtained from patients with various types of von Willebrand disease, with thrombomodulin added.

Example 11: APC Cofactor Assay in Cynomolgus Monkeys with Subcutaneous and Intravenous Injection

[0674] The APC cofactor assay was performed using a Thermo Fluoroskan Ascent Microplate Fluorometer and Thrombinoscope software. The PPP reagent (Diagnostica Stago) was used in this experiment. Briefly, cynomolgus monkey plasma was mixed with PPP reagent and 5 µg/ml of activated Protein C along with calcium and substrate and the levels of thrombin generation

was monitored over 1 hour. The PK assay was performed by incubating diluted cynomolgus monkey plasma onto plates immobilized with human Protein S. The plates used were MSD 96-well plates. 30 μ l of 2 μ g/ml plasma purified Protein S in Tris buffer containing calcium was used to coat the plate overnight. After blocking, the wells were incubated with samples, standards and QCs. 25 μ l per well of 2 μ g/ml sulfo-tagged goat anti-human IgG, monkey ads and incubated at room temperature for 1 hr. After washing, 150 μ l of 1X MSD Read Buffer T in water was added to each well and the plate was read on a MSD plate reader. The levels of D-dimer were measured using the D-dimer assay kit from Diagnostica Stago following manufacturer's recommended protocol.

[0675] FIGS. 28A-28N depict the results of APC cofactor assays, PK assays, and D-dimer assays for plasma samples collected after administration of Protein S antibodies into cynomolgus monkeys at 3 mg/kg subcutaneously and intravenously. The APC cofactor assay was used to measure the pharmacodynamic activity of the administered Protein S antibodies at the various times. In addition, FIGS. 28A-28N also show the levels of free antibody which was measured using an ELISA assay. And finally, elevated D-dimer, which was used as a biomarker of Protein S antibody induced coagulation activity, was observed in the monkeys and the data shown.

[0676] Specifically, FIGS. 28A-28B depict the levels of free antibody (left axis, filled circles) and percent inhibition in an APC cofactor assay (right axis, filled triangles) when Antibody 1 was administered to cynomolgus monkeys at 3 mg/kg subcutaneously and intravenously, respectively.

[0677] FIG. 28C depicts levels of D-dimer, as a marker of coagulation activity, measured over time in cynomolgus monkeys injected with Antibody 1, subcutaneously (SC) and intravenously (IV).

[0678] FIG. 28D depicts the levels of free antibody (left axis, filled squares) and percent inhibition in an APC cofactor assay (right axis, filled circles) when Antibody 3 was administered to cynomolgus monkeys at 3 mg/kg subcutaneously.

[0679] FIG. 28E depicts levels of D-dimer, as a marker of coagulation activity, measured over time in cynomolgus monkeys injected with Antibody 3.

[0680] FIGS. 28F-28G depict the levels of free antibody (left axis, filled squares) and percent inhibition in an APC cofactor assay (right axis, filled triangles) when Antibody 6 was administered to cynomolgus monkeys at 3 mg/kg subcutaneously and intravenously, respectively.

[0681] FIG. 28H depicts levels of D-dimer, as a marker of coagulation activity, measured over time in cynomolgus monkeys injected with Antibody 6.

[0682] FIGS. 28I-28J depict the levels of free antibody (left axis, filled circles) and percent inhibition in an APC cofactor assay (right axis, filled triangles) when Antibody 12 was administered to cynomolgus monkeys at 3 mg/kg subcutaneously and intravenously, respectively.

[0683] FIG. 28K depicts levels of D-dimer, as a marker of coagulation activity, measured over time in cynomolgus monkeys injected with Antibody 12.

[0684] FIGS. 28L-28M depict the levels of free antibody (left axis, filled squares) and percent inhibition in an APC cofactor assay (right axis, filled triangles) when Antibody 29 was administered to cynomolgus monkeys at 3 mg/kg subcutaneously and intravenously, respectively.

[0685] FIG. 28N depicts levels of D-dimer, as a marker of coagulation activity, measured over time in cynomolgus monkeys injected with Antibody 29.

Example 12: Effects of Protein S Antibodies on the Binding of Protein S to TFPI

[0686] The effect of the antibodies of the disclosure on the binding of Protein S to TFPI was measured.

[0687] Using the Octet System (Sartorius), an assay to measure the binding of TFPI to Protein S was developed. Briefly, the human Fc antibodies were immobilized onto anti-human Fc capture probes by placing the probes into 10 µg/ml antibody solution in 10 mg/ml bovine serum albumin, 20 mM Tris pH 7.0, 150 mM NaCl, and 4 mM calcium chloride. Then the bound antibodies were placed into a solution containing 10 µg/ml human Protein S followed by a 10 µg/ml solution containing human TFPI. Finally, the probe was placed into a buffer solution

(wash) to observe the dissociation of TFPI from Protein S. The kinetics of association and dissociation of TFPI to Protein S were measured.

[0688] FIG. 31 depicts the binding of human TFPI to Protein S after the binding of Protein S to Antibody 2. These results demonstrate that Antibody 2 does not block the binding of TFPI to Protein S.

Example 13: Full-length and Fab Fragments of Protein S Antibodies in APC Cofactor Assays

[0689] FIGS. 30A-30B depict similar dose-titration curves resulting from a full-length antibody or a Fab fragment of the same antibody in an APC cofactor assay. FIG. 30A shows the results using the full-length and Fab fragment of Antibody 15, and FIG. 30B shows the results using the full-length and Fab fragment of Antibody 16. These results demonstrate that both the full-length and the Fab fragments of the antibodies tested gave similar results in the APC cofactor assay.