

Exatecan derivatives, linker-payloads, and conjugates and thereof

Technical Field

[0001] The present disclosure relates to the biopharmaceutical field, in particular, Exatecan derivatives, linker-payloads, and conjugates and thereof antibody-drug conjugates, and the corresponding preparing process and use thereof.

Background

[0002] In traditional tumor treatment, chemotherapy is one of the main treatment strategies. However, the off-target toxicity from non-specific accumulation in normal tissues, narrow therapeutic window and low tolerance limit the chemotherapy drug development. In recent decades, targeted therapies using monoclonal antibodies and polypeptides which bind to specific markers on the tumor cell's surface has demonstrated less toxicity than chemotherapy. However, they both lack potency in killing tumor cells.

[0003] The treatment strategy of antibodies armed with toxins to selectively kill target cells was first proposed in 1970. The tumor-targeting drug conjugates mainly compose of ADCs that generally couple antibodies which specifically recognize the surface antigens of tumor cells with chemical toxins which effectively kill tumor tissues through linkers.

[0004] Until 2021, more than 120 ADCs have been in the process of clinical development.

[0005] Most of cytotoxic payloads developed belong to two major families: tubulin inhibitors (maytansinoids or auristatins) and DNA-damaging agents (mainly calicheamicins). All of them are extremely potent cytotoxic drugs characterized by an IC_{50} (the inhibitory concentrations that inhibited 50% of cells) in the nanomolar and picomolar range, and an unfavorable toxicity profile if administered systemically. Conjugation into ADC hides the cytotoxic drug in the bloodstream to convey it directly into tumor cells, thus significantly reducing toxicity of these potent agents. In general, drugs with a more favorable therapeutic index are considered unsuitable for development as warheads, with the IC_{50} too low to be effective when released in the amount delivered into tumor cells.

[0006] Researches on ADCs with an increased DAR or exhibiting the bystander effect propose a promising new approach to exploit the less cytotoxic drugs as warheads.

[0007] So far, anti-microtubules agents such as Auristatins (e.g., monomethyl auristatin E (MMAE) and F (MMAF)) represent the large majority of warheads developed. Brentuximab vedotin, polatuzumab vedotin and enfortumab vedotin are all approved ADCs carrying the MMAE payload.

[0008] Topoisomerase I (TOP1) inhibitors are used as warheads in several ADCs. Sacituzumab govitecan bears SN-38, the active metabolite of irinotecan, as the warhead. Enhertu (fam-


trastuzumab deruxtecan (DS8201a), also known as T-Dxd) bears an Exatecan derivative deruxtecan (Dxd) as the warhead (WO2014057687A), and was recently approved for treating breast cancer. On this basis, novel small molecule topoisomerase I inhibitors, either used alone or as a component of ADCs, with greater activity, stability, physiochemical properties are required in the field.

[0001] However, on one hand, Enhertu, as well as the other commercially available ADCs and most of the ADCs in clinical trials, are prepared by chemical conjugation, using a thiosuccinimide structure (thiosuccinimide linkage) to conjugate the small molecule drug with the targeting antibody or protein. The thiosuccinimide structure is formed by the reaction of a thiol group and a maleimide. The thiosuccinimide linkage is not stable. In organisms, reverse Michael addition or exchange with other thiol groups leads to the fall-off of the cytotoxin from the ADC and off-target toxicity, which reduces the safety and limits the clinical application. For example, albeit the great efficacy, Enhertu caused more than 10% of interstitial lung disease, which limited its usage in part of patients. On another hand, chemical coupling reaction is not site specific, and the resulting ADCs show high heterogeneity.


[0009] New ADCs with higher efficacy and less toxicity are still in urgent need.

Summary


[0010] In a first aspect, provided is a compound of formula (I):



wherein,



P1 and P2 are independently a payload having the structure of formula (i'):



R¹ and R² are each independently selected from hydrogen, C₁₋₆ alkyl, halogen and C₁₋₆ alkoxy.

[illegible]

3

wherein,

A is an antibody or an antigen binding fragment thereof; the antibody or antigen binding fragment is preferably modified to connect with the (Gly)_n moiety in the compound of formula (I);


z is an integer of 1 to 20; preferably 1 to 4; particularly 2;

P1, P2, R⁰, *op*Su, n, k1, k2, i and j are as defined in formula (I).

[0012] In a third aspect, provided is a pharmaceutical composition comprising the conjugate of the present disclosure, and at least one pharmaceutically acceptable carrier.

[0013] In a fourth aspect, provided is use of the conjugate of the present disclosure or the pharmaceutical composition of the present disclosure in the manufacture of a medicament for treating a disease; wherein the disease is a tumor or an autoimmune disease; preferably a tumor comprising HER2-positive tumor cells.


[0014] In a fifth aspect, provided is a compound, having the structure of formula (i):



[0015] wherein,

[0016] a, the carbon atoms marked with p1* and p2*, L¹, M, L², R¹ and R² are as defined in formula (I).

[0017] In a sixth aspect, provided is a compound of formula (1):



[0018] wherein,

[0019] k is an integer of 1 to 7; preferably 1, or 3 or 5, particularly 5

[0020] P is a payload having the structure of formula (i'), wherein the structure of formula (i') is as defined in formula (I).

Brief Description of the Drawings

- [0021] Figure 1 shows the efficacy of conjugates in the SK-BR-3 HER2-high cell line.
- [0022] Figure 2 shows the efficacy of conjugates in the NCI-N87 HER2-high cell line.
- [0023] Figure 3 shows the efficacy of conjugates in the MDA-MB-468 HER2-negative cell line.
- [0024] Figure 4a and Figure 4b show the efficacy of conjugates in the co-culture system of SK-BR-3 and MDA-MB-468. Figure 4a shows result detected by GFP fluorescence. Figure 4b shows result detected by luciferase substrate chemiluminescence. In Figure 4a and Figure 4b, from left to right are CH-2-589, CH-2-593, CH-2-518, LC1184(8) and LC302-2-4(4) respectively.
- [0025] Figure 5 shows the efficacy of conjugates in the co-culture system of SK-BR-3 and MDA-MB-468 detected by GFP fluorescence.
- [0026] Figure 6 shows the mean tumor volume change over time in SCID Beige mice with JIMT1 CDX model dosed with 5 mg/kg of conjugates.
- [0027] Figure 7 shows the mean tumor volume change over time in BALB/c nude mice with Capan-1 CDX model dosed with 5 mg/kg of conjugates.
- [0028] Figure 8a and Figure 8b show the mean tumor volume change and body weight change over time in BALB/c nude mice with Capan-1 CDX model dosed with 5 mg/kg of conjugates.
- [0029] Figure 9a and Figure 9b shows the mean tumor volume change and body weight change over time in BALB/c Nude mice with NCI-N87 CDX model dosed with 5 mg/kg of conjugates.
- [0030] Figure 10 shows the ex-vivo serum stability of the conjugates after 0, 24, 48 and 96 h of incubation in pooled human serum.

Detailed Description

[0031] The specific embodiments are provided below to illustrate technical contents of the present disclosure. Those skilled in the art can easily understand other advantages and effects of the present disclosure through the contents disclosed in the specification. The present disclosure can also be implemented or applied through other different specific embodiments. Various modifications and variations can be made by those skilled in the art without departing from the spirit of the present disclosure.

Definitions

[0032] Unless otherwise defined hereinafter, all technical and scientific terms used herein have the same meaning as commonly understood by those skilled in the art. The techniques used herein refer to those that are generally understood in the art, including the variants and equivalent substitutions that are obvious to those skilled in the art. While the following terms

are believed to be readily comprehensible by those skilled in the art, the following definitions are set forth to better illustrate the present disclosure. When a trade name is present herein, it refers to the corresponding commodity or the active ingredient thereof. All patents, published patent applications and publications cited herein are hereby incorporated by reference.

[0033] When a certain amount, concentration, or other value or parameter is set forth in the form of a range, a preferred range, or a preferred upper limit or a preferred lower limit, it should be understood that it is equivalent to specifically revealing any range formed by combining any upper limit or preferred value with any lower limit or preferred value, regardless of whether the said range is explicitly recited. Unless otherwise stated, the numerical ranges listed herein are intended to include the endpoints of the range and all integers and fractions (decimals) within the range. For example, the expression “i is an integer of 1 to 20” means that i is any integer of 1 to 20, for example, i can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20. Other similar expressions such as j, k and g should also be understood in a similar manner.

[0034] Unless the context clearly dictates otherwise, singular forms like “a” and “the” include the plural forms. The expression “one or more” or “at least one” may mean 1, 2, 3, 4, 5, 6, 7, 8, 9 or more.

[0035] The terms “about” and “approximately”, when used in connection with a numerical variable, generally mean that the value of the variable and all values of the variable are within experimental error (for example, within a 95% confidence interval for the mean) or within $\pm 10\%$ of a specified value, or a wider range.

[0036] The term “optional” or “optionally” means the event described subsequent thereto may, but not necessarily happen, and the description includes the cases wherein said event or circumstance happens or does not happen.

[0037] The expressions “comprising”, “including”, “containing” and “having” are open-ended, and do not exclude additional unrecited elements, steps, or ingredients. The expression “consisting of” excludes any element, step, or ingredient not designated. The expression “consisting essentially of” means that the scope is limited to the designated elements, steps or ingredients, plus elements, steps or ingredients that are optionally present that do not substantially affect the essential and novel characteristics of the claimed subject matter. It should be understood that the expression “comprising” encompasses the expressions “consisting essentially of” and “consisting of”.

[0038] As used herein, the term “antibody” is used in a broad way and particularly includes intact monoclonal antibodies, polyclonal antibodies, monospecific antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments, as long as they have the desired biological activity. The antibody may be of any subtype (such as IgG, IgE, IgM, IgD, and IgA) or subclass, and may be derived from any suitable species. In some embodiments, the antibody

is of human or murine origin. The antibody may also be a fully human antibody, humanized antibody or chimeric antibody prepared by recombinant methods.

[0039] Monoclonal antibodies are used herein to refer to antibodies obtained from a substantially homogeneous population of antibodies, i.e., the individual antibodies constituting the population are identical except for a small number of possible natural mutations. Monoclonal antibodies are highly specific for a single antigenic site. The word “monoclonal” refers to that the characteristics of the antibody are derived from a substantially homogeneous population of antibodies and are not to be construed as requiring some particular methods to produce the antibody.

[0040] An intact antibody or full-length antibody essentially comprises the antigen-binding variable region(s) as well as the light chain constant region(s) (CL) and heavy chain constant region(s) (CH), which could include CH1, CH2, CH3 and CH4, depending on the subtype of the antibody. An antigen-binding variable region (also known as a fragment variable region, Fv fragment) typically comprises a light chain variable region (V_L) and a heavy chain variable region (V_H). A constant region can be a constant region with a native sequence (such as a constant region with a human native sequence) or an amino acid sequence variant thereof. The variable region recognizes and interacts with the target antigen. The constant region can be recognized by and interacts with the immune system.

[0041] An antibody fragment may comprise a portion of an intact antibody, preferably its antigen binding region or variable region. Examples of antibody fragments include Fab, Fab', F(ab')₂, Fd fragment consisting of V_H and CH1 domains, Fv fragment, single-domain antibody (dAb) fragment, and isolated complementarity determining region (CDR). The Fab fragment is an antibody fragment obtained by papain digestion of a full-length immunoglobulin, or a fragment having the same structure produced by, for example, recombinant expression. A Fab fragment comprises a light chain (comprising a V_L and a CL) and another chain, wherein the said other chain comprises a variable domain of the heavy chain (V_H) and a constant region domain of the heavy chain (CH1). The F(ab')₂ fragment is an antibody fragment obtained by pepsin digestion of an immunoglobulin at pH 4.0-4.5, or a fragment having the same structure produced by, for example, recombinant expression. The F(ab')₂ fragment essentially comprises two Fab fragments, wherein each heavy chain portion comprises a few additional amino acids, including the cysteines that form disulfide bonds connecting the two fragments. A Fab' fragment is a fragment comprising one half of a F(ab')₂ fragment (one heavy chain and one light chain). The antibody fragment may comprise a plurality of chains joined together, for example, via a disulfide bond and/or via a peptide linker. Examples of antibody fragments also include single-chain Fv (scFv), Fv, dsFv, diabody, Fd and Fd' fragments, and other fragments, including modified fragments. An antibody fragment typically comprises at least or about 50 amino acids,

and typically at least or about 200 amino acids. An antigen-binding fragment can include any antibody fragment that, when inserted into an antibody framework (e.g., by substitution of the corresponding region), can result in an antibody that immunospecifically binds to the antigen.

[0042] Antibodies according to the present disclosure can be prepared using techniques well known in the art, such as the following techniques or a combination thereof: recombinant techniques, phage display techniques, synthetic techniques, or other techniques known in the art. For example, a genetically engineered recombinant antibody (or antibody mimic) can be expressed by a suitable culture system (e.g., *E. coli* or mammalian cells). The engineering can refer to, for example, the introduction of a ligase-specific recognition sequence at its terminals.

[0043] HER2 refers to human epidermal growth factor receptor-2, which belongs to the epidermal growth factor (EGFR) receptor tyrosine kinase family. In the present application, the terms ErbB2 and HER2 have the same meaning and can be used interchangeably.

[0044] As used herein, the term “antibody-drug conjugate” is referred to as “conjugate”.

[0045] A small molecule compound refers to a molecule with a size comparable to that of an organic molecule commonly used in medicine. The term does not encompass biological macromolecules (e.g., proteins, nucleic acids, etc.), but encompasses low molecular weight peptides or derivatives thereof, such as dipeptides, tripeptides, tetrapeptides, pentapeptides, and the like. Typically, the molecular weight of the small molecule compound can be, for example, about 100 to about 2000 Da, about 200 to about 1000 Da, about 200 to about 900 Da, about 200 to about 800 Da, about 200 to about 700 Da, about 200 to about 600 Da, about 200 to about 500 Da.

[0046] A spacer is a structure that is located between different structural modules and can spatially separate the structural modules. The definition of spacer is not limited by whether it has a certain function or whether it can be cleaved or degraded *in vivo*. Examples of spacers include but are not limited to amino acids and non-amino acid structures, wherein non-amino acid structures can be, but are not limited to, amino acid derivatives or analogues. “Spacer sequence” refers to an amino acid sequence serving as a spacer, and examples thereof include but are not limited to a single amino acid, a sequence containing a plurality of amino acids, for example, a sequence containing two amino acids such as GA, etc., or, for example, GGGGS, GGGGSGGGGS, GGGGSGGGGSGGGGS, etc. Self-immolative spacers are covalent assemblies tailored to correlate the cleavage of two chemical bonds after activation of a protective part in a precursor: Upon stimulation, the protective moiety (such as a cleavable sequence) is removed, which generates a cascade of disassembling reactions leading to the temporally sequential release of smaller molecules. Examples of self-immolative spacers include but not limited to PABC (p-benzyloxycarbonyl), acetal, heteroacetal and the combination thereof.

[0047] The term “alkyl” refers to a saturated aliphatic hydrocarbon group consisting of carbon atoms and hydrogen atoms, which is connected to the rest of the molecule through a single bond. Alkyl groups comprise straight alkyl, branched alkyl or cyclic alkyl (cycloalkyl) or partially cyclic alkyl (e.g., cycloalkyl–linear alkyl and cycloalkyl–branched alkyl). The alkyl group may contain 1 to 10 carbon atoms, referring to C₁₋₁₀ alkyl group, for example, C₁₋₆ alkyl group, C₁₋₄ alkyl group, C₁₋₃ alkyl group, C₁₋₂ alkyl, C₃ alkyl, C₄ alkyl, C₃₋₆ alkyl. Non-limiting examples of linear alkyl groups include but are not limited to methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, etc. Non-limiting examples of branched alkyl groups include but are not limited to isopropyl, isobutyl, sec-butyl, tert-butyl, isopentyl, 2-methylbutyl, 1-methylbutyl, 1-ethylpropyl, 1,2-dimethylpropyl, neopentyl, 1,1-dimethylpropyl, 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl, 2-ethylbutyl, 1-ethylbutyl, 3,3-dimethylbutyl, 2,2-dimethyl butyl, 1,1-dimethylbutyl, 2,3-dimethylbutyl, 1,3-dimethylbutyl or 1,2-dimethylbutyl, etc.

[0048] The terms “cyclic alkyl” and “cycloalkyl” have the same meaning and are used herein interchangeably. Cyclic alkyl groups can include mono- or polycyclic (e.g., having 2 or more than 2 fused rings) groups. In a multicyclic cycloalkyl, two or more rings can be fused or bridged or spiro together. Ring-forming carbon atoms of a cyclic alkyl group can be optionally substituted by oxo (i.e., C(O)). Cyclic alkyl groups can have 3, 4, 5, 6, 7, 8, 9, or 10 ring-forming carbon atoms (C₃₋₁₀). Examples of cyclic alkyl groups include but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, bicyclo[1.1.1]pentyl and bicyclo[2.1.1]hexyl. In some embodiments, the cyclic alkyl is a C₃₋₆ monocyclic or bicyclic cyclic alkyl, preferably C₃₋₆ monocyclic cyclic alkyl, especially cyclopropyl.

[0049] The term “partially cyclic” refers to that the group contains one or more cyclic moieties and one or more acyclic (i.e. linear or branched) moieties. Partially cyclic alkyl groups can include cyclic alkyl–linear alkyl groups and cyclic alkyl–branched alkyl groups. Partially cyclic alkyl groups can have 4, 5, 6, 7, 8, 9, or 10 carbon atoms (C₃₋₁₀), including ring-forming carbon atoms and non-ring-forming carbon atoms. Examples of partially cyclic alkyl groups include but are not limited to C₃₋₉ cyclic alkyl–C₁ alkyl groups, C₃₋₈ cyclic alkyl–C₂ alkyl groups, C₃₋₇ cyclic alkyl–C₃ linear alkyl groups, C₃₋₆ cyclic alkyl–C₄ linear alkyl groups, C₃₋₅ cyclic alkyl–C₅ linear alkyl groups, C₃₋₄ cyclic alkyl–C₆ linear alkyl groups, C₃ cyclic alkyl–C₇ linear alkyl groups, C₃₋₇ cyclic alkyl–C₃ branched alkyl groups, C₃₋₆ cyclic alkyl–C₄ branched alkyl groups, C₃₋₅ cyclic alkyl–C₅ branched alkyl groups, C₃₋₄ cyclic alkyl–C₆ branched alkyl groups, and C₃ cyclic alkyl–C₇ branched alkyl groups. In some embodiments, the partially cyclic alkyl is a C₃₋₉ cyclic alkyl–C₁ alkyl group, preferably a C₃₋₆ cyclic alkyl–C₁₋₂ alkyl group, C₃₋₆ cyclic alkyl–C₁ alkyl group, more preferably C₃₋₄ cyclic alkyl–C₁₋₂ alkyl group, particularly C₃₋₄ cyclic alkyl–C₁ alkyl group, especially cyclopropyl-methyl.

[0050] A bivalent radical refers to a group obtained from the corresponding monovalent radical by removing one hydrogen atom from a carbon atom with free valence electron(s). A bivalent radical have two connecting sites which are connected to the rest of the molecule, wherein the two connecting sites may be on the same atom or on two different atoms of the bivalent radical.

[0051] An “alkylene” or an “alkylidene” refers to a saturated divalent hydrocarbon group. Alkylene groups comprise linear, branched, cyclic or partially cyclic groups. Examples of linear alkylene groups include but are not limited to methylene (-CH₂-), -(CH₂)₂-, -(CH₂)₃-, -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, etc. Examples of branched alkylene groups include but are not limited to -CH(CH₃)-, -CH(C₂H₅)-, -CH(CH₃)-CH₂-, -CH(C₃H₇)-, -CH(C₂H₅)-CH₂-, -C(CH₃)₂-CH₂-, -(CH(CH₃))₂-, -CH(CH₃)-(CH₂)₂-, -CH₂-CH(CH₃)-CH₂-, -CH(C₄H₉)-, -C(CH₃)(C₃H₇)-, -C(C₂H₅)₂-, -CH(C₃H₇)-CH₂-, -CH(C₂H₅)-CH(CH₃)-, -CH(C₂H₅)-(CH₂)₂-, -CH₂-CH(C₂H₅)-CH₂-, -C(CH₃)₂-(CH₂)₂-, -CH₂-C(CH₃)₂-CH₂-, -CH(CH₃)-(CH₂)₃-, -CH₂-CH(CH₃)-(CH₂)₂-, -CH(C₅H₁₁)-, -C(C₂H₅)(C₃H₇)-, -C(CH₃)(C₄H₉)-, -CH(C₄H₉)-CH₂-, -C(C₂H₅)₂-CH₂-, -C(CH₃)(C₃H₇)-CH₂-, -CH(C₂H₅)-CH(C₂H₅)-, -CH(CH₃)-CH(C₃H₇)-, -C(CH₃)₂-C(CH₃)₂-, -CH(C₃H₇)-(CH₂)₂-, -CH₂-CH(C₃H₇)-CH₂-, -CH(C₂H₅)-C(CH₃)₂-, -C(CH₃)₂-CH(CH₃)-CH₂-, -CH(CH₃)-C(CH₃)₂-CH₂-, -CH(C₂H₅)-CH(CH₃)-CH₂-, -CH(CH₃)-CH(C₂H₅)-CH₂-, -CH(CH₃)-CH₂-CH(C₂H₅)-, -CH(CH₃)-C(CH₃)₂-CH₂-, -(CH(CH₃))₃-, -C(CH₃)₂-(CH₂)₃-, -CH(C₂H₅)-(CH₂)₃-, -CH₂-CH(C₂H₅)-(CH₂)₂-, -CH₂-CH(CH₃)-CH(CH₃)-CH₂-, -(CH(CH₃))₂-(CH₂)₂-, -CH(CH₃)-(CH₂)₂-CH(CH₃)-, -(CH₂)₂-CH(CH₃)-(CH₂)₂-, -CH₂-CH(CH₃)-(CH₂)₃-, -CH(CH₃)-(CH₂)₄-, etc.

[0052] The terms “cyclic alkylene” and “cycloalkylene” have the same meaning and are used herein interchangeably. Examples of cyclic alkylene groups include but are not limited to cyclopropylene, cyclobutylene, cyclopentylene, cyclohexylene, cycloheptylene and cyclooctylene, and divalent multicyclic alkyl groups containing fused, spiro or bridged rings. In some embodiments, the cyclic alkylene is a C₃₋₆ cyclic alkylene group, particularly C₃₋₄ cyclic alkylene group, especially cyclopropyl.

[0053] Partially cyclic alkylene groups can include bivalent radicals wherein the two connecting sites which are connected to the rest of the molecule can be both on the one or more linear or branched alkyl moieties, or both on the one or more cyclic alkyl moieties, or respectively on a cyclic alkyl moiety and a linear or branched alkyl moiety. Examples of partially cyclic alkylene groups include but are not limited to: (1) cyclopropylene, cyclobutylene, cyclopentylene, cyclohexylene, cycloheptylene and cyclooctylene, and divalent multicyclic alkyl groups containing fused, spiro or bridged rings, which are each independently substituted by one or more linear or branched alkyl groups; (2) linear or branched alkylene groups, which are each independently substituted by one or more cyclic alkyl groups; and (3) a group formed by combining one or more cyclic alkylene groups and one or more linear or


branched alkylene groups, provided that a chemically stable structure is formed. In some embodiments, the partially cyclic alkylene is a C₃₋₉ cyclic alkyl-C₁ alkylene group, preferably a C₃₋₆ cyclic alkyl-C₁₋₂ alkylene group, C₃₋₆ cyclic alkyl-C₁ alkylene group, more preferably C₃₋₄ cyclic alkyl-C₁₋₂ alkylene group, particularly C₃₋₄ cyclic alkyl-C₁ alkylene group, especially cyclopropyl-methylene.

[0054] Compounds of the invention can exist in isotope-labeled or -enriched form containing one or more atoms having an atomic mass or mass number different from the atomic mass or mass number most abundantly found in nature. Isotopes can be radioactive or non-radioactive isotopes. Isotopes of atoms such as hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine, chlorine and iodine include, but are not limited to, ²H, ³H, ¹³C, ¹⁴C, ¹⁵N, ¹⁸O, ³²P, ³⁵S, ¹⁸F, ³⁶Cl and ¹²⁵I. In one embodiment, the payload (for example the compound of formula (i)) can exist in isotope-labeled or -enriched form, especially deuterated form.

[0055] As used herein, the expressions "antibody-conjugated drug" and "antibody-drug conjugate" has the same meaning.

Compound of formula (i)

[0056] In an aspect, provided is a compound or a pharmaceutically acceptable salt, stereoisomer, solvate, polymorph, tautomer, isotopic compound, metabolite or prodrug thereof, wherein the compound has the structure of formula (i):



[0057] wherein,

[0058] a is 0 or 1;

[0059] the carbon atoms marked with p1* and p2* each is asymmetric center, and the asymmetric center is *S* configured, *R* configured or racemic;





[0060] L¹ is selected from C₁₋₆ alkylene, which is unsubstituted or substituted with one substituent selected from halogen, -OH and -NH₂;

[0061] M is -CH₂-, -NH- or -O-;



[0062] L² is C₁₋₃ alkylene;


[0063] R^1 and R^2 are each independently selected from hydrogen, C_{1-6} alkyl, halogen and C_{1-6} alkoxy.

[0064] In an embodiment, L^1 is selected from C_{1-6} linear alkylene, C_{1-6} branched alkylene, C_{3-6} cyclic alkylene and C_{3-4} cyclic alkyl- C_{1-2} linear alkylene group, which are each independently unsubstituted or substituted with one substituent selected from halogen, -OH and -NH₂. In an embodiment, L^1 is selected from C_{1-4} alkylene, which is unsubstituted or substituted with one substituent selected from halogen, -OH and -NH₂. In a preferred embodiment, L^1 is selected


from -CH₂-, -C₂H₄-, , , , , , which are each independently unsubstituted or substituted with at least one substituent selected from halogen,

-OH and -NH₂. In a preferred embodiment, L^1 is selected from -CH₂-, , , ,

, , , , , wherein “#” marks the position attached to

carbonyl. In a more preferred embodiment, L^1 is selected from -CH₂-, , ,


, , wherein “#” marks the position attached to carbonyl. In a particular


embodiment, L^1 is selected from -CH₂-, , and , wherein “#” marks the position attached to carbonyl. In a preferred embodiment, the halogen is selected from F, Cl and Br, especially F.

[0065] In an embodiment, a is 1, M is -CH₂-, -NH- or -O-; and L^2 is -C₂H₄-. In another embodiment, a is 1, M is -CH₂-, and L^2 is -CH₂-.

[0066] In an embodiment, the carbon atom marked with $p1^*$ is *S* configured or racemic, preferably *S* configured. In another embodiment, the carbon atom marked with $p2^*$ is *S* configured or racemic, preferably *S* configured.


[0067] In an embodiment, R^1 and R^2 are each independently selected from hydrogen, C_{1-3} alkyl, halogen and C_{1-3} alkoxy. In a preferred embodiment, R^1 and R^2 are each independently selected from CH₃-, F, Cl, Br and CH₃O-. In an embodiment, R^1 is selected from CH₃- and Cl. In another embodiment, R^2 is F.


[0068] In an embodiment, a is 0, L¹ is selected from -CH₂-, , ,  and , wherein “#” marks the position attached to carbonyl. In an embodiment, a is 1, L¹ is , M is O, and L² is -C₂H₄-.





[0069] In an embodiment, a is 0, R¹ is Cl, R² is F, and L¹ is selected from -CH₂-, ,  and . In an embodiment, a is 0, R¹ is CH₃-, R² is F, and L¹ is selected from  and , wherein “#” marks the position attached to carbonyl.

[0070] In an embodiment, a is 1, R¹ is CH₃-, R² is F, L¹ is , M is O, and L² is -C₂H₄-.


[0071] In an embodiment, the compound is selected from:










[0072] In an embodiment, the compound is selected from:









[0073] In a preferred embodiment, the compound is selected from:







[0074] In a preferred embodiment the compound is selected from:




[0075] In a more preferred embodiment, the compound is selected from:






[0076] In a particular embodiment the compound is selected from:




[0077] In an aspect, provided is a compound selected from:




Compound of Formula (I)

[0078] In an aspect, provided is a compound having the structure of formula (I):



[0079] wherein,



[0080] *opSu* is

or ;

[0081] R^0 is C_{1-10} alkyl;


[0082] n is any integer of 2 to 20;

[0083] k_1 and k_2 are independently an integer of 1 to 7;

[0084] i is an integer of 1-100;

[0085] j is an integer of 1-100;


[0086] P_1 and P_2 are independently a payload having the structure of formula (i'):








[0087] wherein,

[0088] a , the carbon atoms marked with p_1^* and p_2^* , L^1 , M , L^2 , R^1 and R^2 are as defined in formula (i).


[0089] In an embodiment, the payload is selected from:










[0090] In an embodiment, the payload is selected from:









[0091] In a preferred embodiment, the payload is selected from:







[0092] In a preferred embodiment, the payload is selected from:




[0093] In a more preferred embodiment, the payload is selected from:





[0094] In a particular embodiment, the payload is selected from:



[0095] In an embodiment, R^0 is C_{1-6} alkyl. In a preferred embodiment, R^0 is C_{1-3} alkyl. In a particular embodiment, R^0 is methyl.


[0096] In an embodiment, n is an integer of 2 to 5. In a particular embodiment, n is 3.

[0097] In an embodiment, k_1 and k_2 are independently 1, or 3 or 5. In a particular embodiment, k_1 and k_2 are independently 5.

[0098] In an embodiment, i is independently an integer of 1 to 20, preferably 1 to 12, more preferably 2 to 8. In a particular embodiment, i is 4.

[0099] In an embodiment, j is independently an integer of 1 to 20, preferably 1 to 12, more preferably 8 to 12, especially 8 or 12. In a particular embodiment, j is 12.

[0100] In an embodiment, the compound of formula (I) has the structure of formula (I-1)




(I-1)


[0101] wherein,

[0102] P_1 , P_2 , R^0 , $opSu$, n , i and j are as defined in formula (I).

[0103] In an embodiment, the compound of formula (I) is selected from the group consisting of:



BH-1 and



BH-2

Specific Embodiments of Compound of Formula (I)

[0104] In an embodiment, the structure of the compound of formula (I) is as shown in the following table.

Compound of formula (I)	Formula	Payload	Compound of formula (I)	Formula	Payload
BH-1-730	BH-1	P730	BH-1-731	BH-1	P731
BH-1-518	BH-1	P518	BH-1-519	BH-1	P519
BH-1-586	BH-1	P586	BH-1-587	BH-1	P587
BH-1-593	BH-1	P593	BH-1-594	BH-1	P594
BH-1-622	BH-1	P622	BH-1-624	BH-1	P624
BH-1-625	BH-1	P625	BH-1-677	BH-1	P677
BH-1-678	BH-1	P678	BH-1-679	BH-1	P679
BH-1-680	BH-1	P680	BH-1-703	BH-1	P703
BH-1-704	BH-1	P704	BH-1-627	BH-1	P627
BH-1-626	BH-1	P626	BH-1-636	BH-1	P636
BH-1-637	BH-1	P637	BH-1-595	BH-1	P595
BH-1-596	BH-1	P596	BH-1-597	BH-1	P597


BH-1-600	BH-1	P600	BH-1-601	BH-1	P601
BH-1-602	BH-1	P602	BH-1-603	BH-1	P603
BH-1-619	BH-1	P619	BH-1-620	BH-1	P620
BH-1-664	BH-1	P664	BH-1-665	BH-1	P665
BH-1-667	BH-1	P667	BH-1-668	BH-1	P668
BH-1-671	BH-1	P671	BH-1-589	BH-1	P589
BH-1-588	BH-1	P588	BH-1-504	BH-1	P504
BH-1-562	BH-1	P562	BH-1-563	BH-1	P563
BH-1-565	BH-1	P565	BH-1-687	BH-1	P687
BH-1-688	BH-1	P688	LB302-2-1	BH-1	Dxd
BH-2-730	BH-2	P730	BH-2-731	BH-2	P731
BH-2-518	BH-2	P518	BH-2-519	BH-2	P519
BH-2-586	BH-2	P586	BH-2-587	BH-2	P587
BH-2-593	BH-2	P593	BH-2-594	BH-2	P594
BH-2-622	BH-2	P622	BH-2-624	BH-2	P624
BH-2-625	BH-2	P625	BH-2-677	BH-2	P677
BH-2-678	BH-2	P678	BH-2-679	BH-2	P679
BH-2-680	BH-2	P680	BH-2-703	BH-2	P703
BH-2-704	BH-2	P704	BH-2-627	BH-2	P627
BH-2-626	BH-2	P626	BH-2-636	BH-2	P636
BH-2-637	BH-2	P637	BH-2-595	BH-2	P595
BH-2-596	BH-2	P596	BH-2-597	BH-2	P597
BH-2-600	BH-2	P600	BH-2-601	BH-2	P601
BH-2-602	BH-2	P602	BH-2-603	BH-2	P603
BH-2-619	BH-2	P619	BH-2-620	BH-2	P620
BH-2-664	BH-2	P664	BH-2-665	BH-2	P665
BH-2-667	BH-2	P667	BH-2-668	BH-2	P668
BH-2-671	BH-2	P671	BH-2-589	BH-2	P589
BH-2-588	BH-2	P588	BH-2-504	BH-2	P504
BH-2-562	BH-2	P562	BH-2-563	BH-2	P563
BH-2-565	BH-2	P565	BH-2-687	BH-2	P687
BH-2-688	BH-2	P688	LB302-2-4	BH-2	Dxd

Note: For all the compounds listed, P1 and P2 are the same, and are described as Payload.

Preparation of the Compound of Formula (I)

[0105] In an embodiment, compound of formula (I) can be synthesized by connecting a linker with a payload or by connecting a serial of suitable building blocks. Such building blocks can be easily designed by retrosynthetic analysis, and any reaction known in the art can be used.

[0106] In an aspect, provided is a compound having the structure of formula (1):



formula (1)



[0107] wherein,

[0108] k is an integer of 1 to 7;

[0109] P is a payload having the structure of formula (i'), wherein the structure of formula (i') is as defined above.

[0110] In a preferred embodiment, k is 1, or 3 or 5. In a particular embodiment, k is 5.

[0111] The compound of formula (1) can be synthesized using the compound of formula (i) and other necessary building blocks, using a method similar to the synthetic method as disclosed in EP2907824A (e.g., synthetic method for formula (2) or (2b) of EP2907824A). Suitable building blocks include but not limited to linking-unit HX20113 and linking-unit HX20111.



[0112] . Then the maleimide group therein can be reacted to a thiol group on another building block for the compound of formula (I). The resulting thiosuccinimide is unstable under physiological conditions and is liable to reverse Michael addition which leads to cleavage at the connection site. Moreover, when another thiol compound is present in the system, thiosuccinimide may also undergo thiol exchange with the other thiol compound. Both of these reactions cause the fall-off of the payload and result in toxic side effects. The thiosuccinimide is then subjected to ring opening reaction. The compound of formula (I) can then be obtained.

[0113] Method of ring opening reaction can be found in WO2015165413A1. The compound comprising ring-opened succinimide moiety can be purified by semi-preparative/preparative HPLC or other suitable separation means to obtain with high purity and defined composition, regardless of the efficiency of the succinimide ring opening reaction.

[0114] In the present disclosure, when applied in the linker-payload (linker–small molecule intermediate), the ring-opened succinimide structure no longer undergoes reverse Michael addition or thiol exchange, and thus the product is more stable.

Moiety Comprising Recognition Sequence of the Ligase Acceptor or Donor Substrate

[0115] In an embodiment, the (Gly)_n moiety of the compound of formula (I) is a recognition sequence of a ligase acceptor or donor substrate, which facilitates enzyme-catalyzed coupling of compound of formula (I) with the an antibody or an antigen binding fragment thereof under the catalysis of the ligase. The antibody or the antigen binding fragment thereof is optionally modified and comprises the corresponding recognition sequence of a ligase acceptor or donor substrate.

[0116] In an embodiment, the ligase is a transpeptidase. In an embodiment, the ligase is selected from the group consisting of a natural transpeptidase, an unnatural transpeptidase, variants thereof, and the combination thereof. Unnatural transpeptidase enzymes can be, but are not limited to, those obtained by engineering of natural transpeptidase. In a preferred embodiment, the ligase is selected from the group consisting of a natural Sortase, an unnatural Sortase, and the combination thereof. The species of natural Sortase include Sortase A, Sortase B, Sortase C, Sortase D, Sortase *L. plantarum*, etc. (US20110321183A1). The type of ligase corresponds to the ligase recognition sequence and is thereby used to achieve specific conjugation between different molecules or structural fragments.

[0117] In an embodiment, the (Gly)_n moiety of the compound of formula (I) is a recognition sequence of a ligase acceptor substrate; and the antibody or the antigen binding fragment thereof is optionally modified and comprises the corresponding recognition sequence of a ligase donor substrate.

[0118] In some embodiments, the ligase is a Sortase selected from Sortase A, Sortase B, Sortase C, Sortase D and Sortase *L. plantarum*.

[0119] In a particular embodiment, the ligase is Sortase A from *Staphylococcus aureus*. Accordingly, the ligase recognition sequence of the ligase donor substrate may be the typical recognition sequence LPXTG of the enzyme. In yet another particular embodiment, the recognition sequence of the ligase donor substrate is LPXTGJ, wherein X can be any single amino acid that is natural or unnatural; J is absent, or is an amino acid fragment comprising 1-10 amino acids, optionally labeled. In an embodiment, J is absent. In yet another embodiment, J is an amino acid fragment comprising 1-10 amino acids, wherein each amino acid is independently any natural or unnatural amino acid. In another embodiment, J is (Gly)_m, wherein m is an integer of 1 to 10. In yet another particular embodiment, the recognition sequence of

[0123] The ligase recognition sequence can also be other totally new recognition sequence for transpeptidase optimized by manual screening.

[0124] Furthermore, the payload-bearing compound (compound of formula (I)) which has the moiety comprising ligase recognition sequence can be conjugated with other molecules comprising a ligase recognition sequence, and can be thereby used in for example, the preparation of a antibody–small molecule Conjugate, such as an antibody-drug conjugate. Accordingly, in yet another aspect, provided is a conjugate which comprises a compound of formula (I), and an antibody or an antigen binding fragment.

$$\left[\begin{array}{c} \text{HN} \left[\text{CH}_2 \text{O} \right]_i \text{CH}_2 \text{CH}_2 \text{C}(=\text{O}) \text{NH} \text{CH}(\text{NH}_2) \text{CH}_2 \text{S} \text{opSu} \left[\text{CH}_2 \right]_{k1} \text{C}(=\text{O}) \text{NH} \text{CH}_2 \text{C}(=\text{O}) \text{NH} \text{CH}_2 \text{C}(=\text{O}) \text{NH} \text{CH}(\text{CH}_2 \text{C}_6\text{H}_5) \text{C}(=\text{O}) \text{NH} \text{CH}_2 \text{C}(=\text{O}) \text{NH} \text{CH}_2 \text{OP1} \\ | \\ \text{A} - \left[\text{NH} \text{CH}_2 \text{C}(=\text{O}) \right]_n \text{NH} \text{CH}(\text{CH}_2 \text{CH}_2 \text{CH}_2 \text{CH}_2 \text{NH} \text{C}(=\text{O}) \text{CH}_2 \text{CH}_2 \text{CH}_2 \text{CH}_2 \text{OP2}) \text{C}(=\text{O}) \text{NH} \text{CH}(\text{CH}_2 \text{S} \text{opSu} \left[\text{CH}_2 \right]_{k2} \text{C}(=\text{O}) \text{NH} \text{CH}_2 \text{C}(=\text{O}) \text{NH} \text{CH}_2 \text{C}(=\text{O}) \text{NH} \text{CH}(\text{CH}_2 \text{C}_6\text{H}_5) \text{C}(=\text{O}) \text{NH} \text{CH}_2 \text{C}(=\text{O}) \text{NH} \text{CH}_2 \text{OP2} \\ | \\ \text{HN} \text{C}(=\text{O}) \text{CH}_2 \text{CH}_2 \text{CH}_2 \text{CH}_2 \text{OP0} \end{array} \right]$$

z is an integer of 1 to 20;

P1, P2, R⁰, *opSu*, n, k1, k2, i and j are as defined in formula (I).

[0126] In a preferred embodiment, the antibody or antigen binding fragment is modified to connect with the (Gly)_n moiety in the compound of formula (I).

[0127] In a preferred embodiment, z is 1 to 4. In a particular embodiment, z is 2.

Antibody or the antigen binding fragment thereof

[0128] In an embodiment, the antibody is an anti-CD19 antibody, anti-CD20 antibody, anti-CD22 antibody, anti-CD25 antibody, anti-CD30/TNFRSF8 antibody, anti-CD33 antibody, anti-CD37 antibody, anti-CD44v6 antibody, anti-CD56 antibody, anti-CD70 antibody, anti-CD71 antibody, anti-CD74 antibody, anti-CD79b antibody, anti-CD117/KIT antibody, anti-CD123 antibody, anti-CD138 antibody, anti-CD142 antibody, anti-CD174 antibody, anti-CD227/MUC1 antibody, anti-CD352 antibody, anti-CLDN18.2 antibody, anti-DLL3 antibody, anti-ErbB2/HER2 antibody, anti-CN33 antibody, anti-GPNMB antibody, anti-ENPP3 antibody, anti-Nectin-4 antibody, anti-EGFRvIII antibody, anti-SLC44A4/AGS-5 antibody, anti-CEACAM5 antibody, anti-PSMA antibody, anti-TIM1 antibody, anti-LY6E antibody, anti-LIV1 antibody, anti-Nectin4 antibody, anti-SLITRK6 antibody, anti-HGFR/cMet antibody, anti-SLAMF7/CS1 antibody, anti-EGFR antibody, anti-BCMA antibody, anti-AXL antibody, anti-NaPi2B antibody, anti-GCC antibody, anti-STEAP1 antibody, anti-MUC16 antibody, anti-Mesothelin antibody, anti-ETBR antibody, anti-EphA2 antibody, anti-5T4 antibody, anti-FOLR1 antibody, anti-LAMP1 antibody, anti-Cadherin 6 antibody, anti-FGFR2 antibody, anti-FGFR3 antibody, anti-CA6 antibody, anti-CanAg antibody, anti-integrin α V antibody, anti-TDGF1 antibody, anti-Ephrin A4 antibody, anti-TROP2 antibody, anti-PTK7 antibody, anti-NOTCH3 antibody, anti-C4.4A antibody, anti-FLT3 antibody, anti-B7H3/4 antibody, anti-Tissue Factor antibody, or anti-ROR1/2 antibody; preferably anti-HER2 antibody, anti-FGFR3 antibody, anti-Trop2 antibody, anti-HER3 antibody or anti-FR α antibody.

[0129] In a preferred embodiment, the antibody is anti-HER2 antibody. In an embodiment, the antibody is an anti-human HER2 antibody. Examples of anti-human HER2 antibodies include but are not limited to Pertuzumab and Trastuzumab. Pertuzumab binds to the second extracellular domain (ECD2) of HER2 and is approved for the treatment of HER2-positive breast cancer. Trastuzumab binds to the fourth extracellular domain (ECD4) of HER2 and is approved for the treatment of Her2-positive breast cancer and gastric cancer.

[0130] In one embodiment, the antibody is an anti-HER2 antibody or antigen-binding fragment thereof, comprising a light chain variable region (V_L) and a heavy chain variable region (V_H), wherein the V_L comprises LCDR1 comprising the amino acid sequence of SEQ ID NO: 1, LCDR2 comprising the amino acid sequence of SEQ ID NO: 2, LCDR3 comprising the amino acid sequence of SEQ ID NO: 3, and wherein the V_H comprises HCDR1 comprising the amino acid sequence of SEQ ID NO: 4, HCDR2 comprising the amino acid sequence of SEQ ID NO:

5, HCDR3 comprising the amino acid sequence of SEQ ID NO: 6. In one embodiment, the CDR regions are defined by KABAT. In one embodiment, the antibody is an anti-HER2 antibody or antigen-binding fragment thereof, comprising a light chain variable region (V_L) and a heavy chain variable region (V_H), wherein the V_L has an amino acid sequence having at least about 85%, at least about 90% or at least about 95% sequence identity with SEQ ID NO: 7, and the V_H has an amino acid sequence having at least about 85%, at least about 90% or at least about 95% sequence identity with SEQ ID NO: 8. In a preferred embodiment, the anti-HER2 antibody or antigen-binding fragment thereof comprises a light chain variable region (V_L) and a heavy chain variable region (V_H), wherein the V_L has the amino acid sequence of SEQ ID NO: 7, and the V_H has the amino acid sequence of SEQ ID NO: 8. In a particular embodiment, the anti-HER2 antibody comprises a light chain and a heavy chain, wherein the light chain has an amino acid sequence having at least about 85%, at least about 90% or at least about 95% sequence identity with the amino acid sequence of amino acids 1 to 214 of SEQ ID NO: 9, and the heavy chain has an amino acid sequence having at least about 85%, at least about 90% or at least about 95% sequence identity with SEQ ID NO: 10. In a preferred embodiment, the anti-human HER2 antibody is one or more selected from engineered anti-HER2 antibodies based on Trastuzumab. In one embodiment, the antibody is a modified Trastuzumab, preferably Ab0001-LCCT_L-HC (having light chain SEQ ID NO: 9, heavy chain: SEQ ID NO: 10). The sequence of Ab0001-LCCT_L-HC is based on the amino acid sequence of Trastuzumab, and GALPETGG was introduced at the C-terminal of the light chain, wherein LPETGG is the recognition sequence of the ligase donor substrate, and GA is a spacer sequence.


[0131] In a preferred embodiment, the anti-human HER2 antibody is a recombinant antibody selected from monoclonal antibody, chimeric antibody, humanized antibody, antibody fragment, and antibody mimic. In an embodiment, the antibody mimic is selected from scFv, minibody, diabody, nanobody. For the conjugation with the compound of formula (I), the antibody or the antigen binding fragment thereof of the present disclosure may comprise a modified moiety to conjugate with the (Gly)_n moiety in the compound of formula (I). The introduction position of such modified moiety is not limited, for example, the introduction position on the antibody or the antigen binding fragment thereof can be, but not limited to, located at the C-terminal or the N-terminal of the heavy chain or light chain of the antibody.

[0132] In an alternative embodiment, a modified moiety for the conjugation with the (Gly)_n moiety in the compound of formula (I) can be introduced at a non-terminal position of the heavy chain or light chain of the antibody using, for example, chemical modification methods.

[0133] In an embodiment, the antibody or the antigen-binding fragment thereof may comprise terminal modification. A terminal modification refers to a modification at the C-terminal or N-terminal of the heavy chain or light chain of the antibody, which for example comprises a ligase

[0134] In a preferred embodiment, the light chain of the antibody or antigen-binding fragment thereof includes 3 types: wild-type (LC); the C-terminus modified light chain (LCCT), which is modified by direct introduction of a ligase recognition sequence LPXTG and C-terminus modified light chain (LCCT_L), which is modified by introduction of short peptide spacers plus the ligase donor substrate recognition sequence LPXTG. The heavy chain of the antibody or antigen-binding fragment thereof includes 3 types: wild-type (HC); the C-terminus modified heavy chain (HCCT), which is modified by direct introduction of a ligase recognition sequence LPXTG; and C-terminus modified heavy chain (HCCT_L), which is modified by introduction of short peptide spacers plus the ligase donor substrate recognition sequence LPXTG. X can be any natural or non-natural single amino acid. When z in the compound of formula (II) is 1 or 2, the combination of the above heavy and light chains can form 8 preferred antibody molecules.

[0135] In an embodiment, the compound of formula (II) is selected from the group consisting:



Specific Embodiments of Conjugates

[0136] In an embodiment, the payload is a cytotoxin or a fragment thereof. In an embodiment, the antibody-drug conjugate is as shown in the following table.

[0137] Nomenclature of the ADCs: the number in the parenthesis indicates the number of payload (drug) molecules that is intended to be connected to the antibody.

ADCs	Compound of formula (I)	ADCs	Compound of formula (I)	ADCs	Compound of formula (I)
CH-1-730	BH-1-730	CH-1-731	BH-1-731	CH-1-518	BH-1-518
CH-1-519	BH-1-519	CH-1-586	BH-1-586	CH-1-587	BH-1-587
CH-1-593	BH-1-593	CH-1-594	BH-1-594	CH-1-622	BH-1-622
CH-1-624	BH-1-624	CH-1-625	BH-1-625	CH-1-677	BH-1-677
CH-1-678	BH-1-678	CH-1-679	BH-1-679	CH-1-680	BH-1-680
CH-1-703	BH-1-703	CH-1-704	BH-1-704	CH-1-627	BH-1-627
CH-1-626	BH-1-626	CH-1-636	BH-1-636	CH-1-637	BH-1-637
CH-1-595	BH-1-595	CH-1-596	BH-1-596	CH-1-597	BH-1-597
CH-1-600	BH-1-600	CH-1-601	BH-1-601	CH-1-602	BH-1-602
CH-1-603	BH-1-603	CH-1-619	BH-1-619	CH-1-620	BH-1-620
CH-1-664	BH-1-664	CH-1-665	BH-1-665	CH-1-667	BH-1-667
CH-1-668	BH-1-668	CH-1-671	BH-1-671	CH-1-589	BH-1-589
CH-1-588	BH-1-588	CH-1-504	BH-1-504	CH-1-562	BH-1-562
CH-1-563	BH-1-563	CH-1-565	BH-1-565	CH-1-687	BH-1-687
CH-1-688	BH-1-688	LC302-2-1(4)	LB302-2-1	-	-
CH-2-730	BH-2-730	CH-2-731	BH-2-731	CH-2-518	BH-2-518
CH-2-519	BH-2-519	CH-2-586	BH-2-586	CH-2-587	BH-2-587
CH-2-593	BH-2-593	CH-2-594	BH-2-594	CH-2-622	BH-2-622
CH-2-624	BH-2-624	CH-2-625	BH-2-625	CH-2-677	BH-2-677
CH-2-678	BH-2-678	CH-2-679	BH-2-679	CH-2-680	BH-2-680
CH-2-703	BH-2-703	CH-2-704	BH-2-704	CH-2-627	BH-2-627
CH-2-626	BH-2-626	CH-2-636	BH-2-636	CH-2-637	BH-2-637
CH-2-595	BH-2-595	CH-2-596	BH-2-596	CH-2-597	BH-2-597
CH-2-600	BH-2-600	CH-2-601	BH-2-601	CH-2-602	BH-2-602
CH-2-603	BH-2-603	CH-2-619	BH-2-619	CH-2-620	BH-2-620
CH-2-664	BH-2-664	CH-2-665	BH-2-665	CH-2-667	BH-2-667
CH-2-668	BH-2-668	CH-2-671	BH-2-671	CH-2-589	BH-2-589
CH-2-588	BH-2-588	CH-2-504	BH-2-504	CH-2-562	BH-2-562
CH-2-563	BH-2-563	CH-2-565	BH-2-565	CH-2-687	BH-2-687
CH-2-688	BH-2-688	LC302-2-4(4)	LB302-2-4	-	-

Note: 1. For all the ADCs listed, P1 and P2 are the same, and are described as Payload; z is 2; the antibody is Ab0001-LCCT₁-HC (light chain SEQ ID NO: 9, heavy chain: SEQ ID NO: 10).

2. When the above-listed ADC is formed by the corresponding above-listed linker-payload and the corresponding above-listed antibody (targeting molecule), the upstream peptide bond of GG in the LPETGG


sequence in the antibody is cleaved by Sortase A, and the resulting intermediate is linked to the free N-terminal of G₃ in linker-payload to generate a new peptide bond, thereby forming the resulting amino acid sequence LPXTG₃.

Preparation of the Conjugate

[0138] The conjugates (i.e., the compound of formula (II)) of the present disclosure can be prepared by any method known in the art. In some embodiments, the conjugate is prepared by the ligase-catalyzed site-specific conjugation of an antibody or an antigen binding fragment and a compound of formula (I), wherein the antibody or the antigen binding fragment thereof is modified by a ligase recognition sequence.

[0139] The antibody or the antigen binding fragment thereof and the compound of formula (I) are linked to each other via the ligase-specific recognition sequences of the substrates. The recognition sequence depends on the particular ligase employed. In an embodiment, the antibody or the antigen binding fragment thereof is an antibody with recognition sequence-based terminal modifications introduced at the C-terminal of the light chain and/or the heavy chain, and the antibody or the antigen binding fragment thereof is conjugated with the compound of formula (I), under the catalysis of the wild type or optimized engineered ligase or any combination thereof, and under suitable catalytic reaction conditions.

[0140] In a specific embodiment, the ligase is Sortase A and the conjugation reaction can be represented by the following scheme:



[0141] The triangle represents a portion of an antibody; the pentagon represents a portion of a compound of formula (II); and G_n represents the (Gly)_n moiety. n, X and J are respectively as defined above. When conjugated with G_n, which is the corresponding recognition sequence of the acceptor substrate, the upstream peptide bond of the glycine in the LPXTGJ sequence is cleaved by Sortase A, and the resulting intermediate is linked to the free N-terminal of G_n to generate a new peptide bond. The resulting amino acid sequence is LPXTG_n. The sequences G_n and LPXTGJ are as defined above.

[0142] Therefore, in one embodiment, the G_n moiety in the compound of formula (I) is the recognition sequence of the ligase acceptor substrate of a ligase; and the antibody is modified to comprises the recognition sequence of the ligase donor substrate in order to connect with the G_n moiety in the compound of formula (I);

preferably,

- (a) the ligase is a Sortase selected from Sortase A, Sortase B, Sortase C, Sortase D and Sortase *L. plantarum*; preferably Sortase A from *Staphylococcus aureus*; and/or
- (b) the recognition sequence of the ligase donor substrate is LPXTGJ, J is absent or is G_m,

wherein G is glycine, m is an integer of 1 to 10, X is any single amino acid that is natural or unnatural; preferably, the recognition sequence of the ligase donor substrate is LPXTG or LPETGG; and/or

the connection of LPXTGJ with G_n results in LPXTG_n.

[0143] The compound of formula (I) of the present disclosure has defined structure, defined composition and high purity, so that when the conjugation reaction with an antibody is conducted, fewer impurities are introduced or no other impurities are introduced. When such an intermediate is used for the ligase-catalyzed site-specific conjugation with a modified antibody containing a ligase recognition sequence, a homogeneous ADC with highly controllable quality is obtained.

Metabolism of the Conjugate in a physiological environment

[0144] When a part or whole linker is cleaved in tumor cells, the payload is released. As the linker is cleaved at a connecting position to the antitumor compound, the antitumor compound is released in its intrinsic structure to exhibit its intrinsic antitumor effect.

[0145] In an embodiment, the GGFG (Gly-Gly-Phe-Gly) moiety comprised by the compound of formula (I) can be cleaved by lysosomal enzymes (such as cathepsin B and/or cathepsin L).

[0146] In an embodiment, the compound of formula (I) comprises a self-immolative spacer. In an embodiment, the self-immolative spacer is an acetal or a heteroacetal. In an embodiment, the -GGFG-NH-CH₂-O- moiety comprised by the compound of formula (I) represents a combination of a restriction enzyme site and a self-immolative spacer, which would cleave in the cell and release the aimed molecule (such as the antitumor compound).

Pharmaceutical Composition and Pharmaceutical Preparation

[0147] Another object of the disclosure is to provide a pharmaceutical composition comprising the conjugate of the present disclosure, and at least one pharmaceutically acceptable carrier.

[0148] The pharmaceutical composition of the present disclosure may be administered in any manner as long as it achieves the effect of preventing, alleviating, preventing or curing the symptoms of a human or animal. For example, various suitable dosage forms can be prepared according to the administration route, especially injections such as lyophilized powder for injection, injection, or sterile powder for injection.

[0149] The term “pharmaceutically acceptable” means that when contacted with tissues of the patient within the scope of normal medical judgment, no undue toxicity, irritation or allergic reaction, etc. shall arise, having reasonable advantage-disadvantage ratios and effective for the intended use.

[0150] The term pharmaceutically acceptable carrier refers to those carrier materials which are pharmaceutically acceptable and which do not interfere with the bioactivities and properties of

the conjugate. Examples of aqueous carriers include but are not limited to buffered saline, and the like. The pharmaceutically acceptable carrier also includes carrier materials which brings the composition close to physiological conditions, such as pH adjusting agents, buffering agents, toxicity adjusting agents and the like, and sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate, and the like.

[0151] In an embodiment, the pharmaceutical composition of the present disclosure has a drug to antibody ratio (DAR) of an integer or non-integer of about 1 to about 20, such as about 1 to about 10, about 1 to about 8, about 1 to about 6, about 1 to about 4, about 2 to about 5, about 2 to about 4, or about 3 to about 4. In a particular embodiment, the conjugate of the present disclosure has a DAR of about 4. In a particular embodiment, the conjugate of the present disclosure has a DAR of about 3.3 to about 3.75, especially about 3.4 to about 3.65.

Treatment Method and Use

[0152] The conjugates of the present disclosure are useful for the treatment of tumors and/or autoimmune diseases. Tumors susceptible to conjugate treatment include those characterized by specific tumor-associated antigens or cell surface receptors, and those will be recognized by the targeting molecule (antibody) in the conjugate and can be killed by the payload/cytotoxin in the conjugate.

[0153] Accordingly, in yet another aspect, also provided is use of a conjugate of the present disclosure or a pharmaceutical composition of the present disclosure in the manufacture of a medicament for treating a disease, disorder or condition selected from a tumor or an autoimmune disease.

[0154] In another aspect, provided is a conjugate of the present disclosure or a pharmaceutical composition of the present disclosure for use in the treatment of a tumor or an autoimmune disease.

[0155] In a further aspect, provided is a method of treating a tumor or an autoimmune disease, the method comprising administering to an individual in need thereof an effective amount of a conjugate of the present disclosure or a pharmaceutical composition of the present disclosure


[0156] In a preferred embodiment, the conjugate of the present disclosure formed by conjugation of the anti-HER2 antibody and the small molecule cytotoxin can specifically bind to HER2 on the surface of the tumor cell and selectively kill the HER2-expressing tumor cells. In another preferred embodiment, the anti-HER2 antibody is an anti-human HER2 antibody. In another preferred embodiment, provided is use of a conjugate of the present disclosure or a pharmaceutical composition of the present disclosure in the manufacture of a medicament for treating a disease, disorder or condition selected from HER2-positive tumors. In a more

preferred embodiment, the disease, disorder or condition is selected from the group consisting of breast cancer, gastric cancer, lung cancer, ovarian cancer, urothelial cancer, and the like.


[0157] The dosage of the conjugate administered to the subject can be adjusted to a considerable extent. The dosage can vary according to the particular route of administration and the needs of the subject, and can be subjected to the judgment of the health care professional.

[0158] As defined above, the embodiments of the present disclosure can also be described as the following [1] to [20]:

[1] A compound of formula (I):



wherein,



R^0 is C_{1-10} alkyl;


n is any integer of 2 to 20;

k_1 and k_2 are independently an integer of 1 to 7;

i is an integer of 1-100;

j is an integer of 1-100;

P_1 and P_2 are independently a payload having the structure of formula (i'):



wherein,

a is 0 or 1;

the carbon atoms marked with p1* and p2* each is asymmetric center, and the asymmetric center is *S* configured, *R* configured or racemic;


L¹ is selected from C₁₋₆ alkylene, which is unsubstituted or substituted with one substituent selected from halogen, -OH and -NH₂;

M is -CH₂-, -NH- or -O-;

L² is C₁₋₃ alkylene;

R¹ and R² are each independently selected from hydrogen, C₁₋₆ alkyl, halogen and C₁₋₆ alkoxy.

[2] The compound of [1], having the structure of formula (I-1)



wherein,

P1, P2, R⁰, *opSu*, n, i and j are as defined in [1].



[3] The compound of [1] or [2], wherein in formula (i'):







L¹ is selected from C₁₋₆ linear alkylene, C₁₋₆ branched alkylene, C₃₋₆ cyclic alkylene and





C₃₋₄ cyclic alkyl-C₁₋₂ linear alkylene group, which are each independently unsubstituted or substituted with one substituent selected from halogen, -OH and -NH₂;



preferably,

L¹ is selected from C₁₋₄ linear alkylene, C₁₋₄ branched alkylene, C₃₋₄ cyclic alkylene and cyclopropyl-methylene, which are each independently unsubstituted or substituted with one substituent selected from halogen, -OH and -NH₂;

preferably, L¹ is selected from -CH₂-, -C₂H₄-, , , , , , which are each independently unsubstituted or substituted with at least one substituent selected from halogen, -OH and -NH₂;

more preferably, L¹ is selected from -CH₂-, , , , , , , , , wherein “#” marks the position attached to carbonyl;

further preferably, L¹ is selected from -CH₂-, , , , , wherein “#” marks the position attached to carbonyl;

particularly, L¹ is selected from -CH₂-, , and , wherein “#” marks the position attached to carbonyl;

and/or

the halogen is selected from F, Cl and Br.

[4] The compound of any one of [1] to [3], wherein in formula (i'):

M is -CH₂-, -NH- or -O-; and L² is -C₂H₄-; or

M is -CH₂-, and L² is -CH₂-.

[5] The compound of any one of [1] to [4], wherein in formula (i'):

the carbon atom marked with p1* is *S* configured or racemic, preferably *S* configured;
and/or

the carbon atom marked with p2* is *S* configured or racemic, preferably *S* configured.

[6] The compound of any one of [1] to [5], wherein in formula (i'):

R^1 and R^2 are each independently selected from hydrogen, C_{1-3} alkyl, halogen and C_{1-3} alkoxy;

preferably, R^1 and R^2 are each independently selected from CH_3- , F, Cl, Br and CH_3O- ;
more preferably,


R^1 is selected from CH_3- and Cl; and/or

R^2 is F;

further preferably,


a is 0, R^1 is Cl, R^2 is F, and L^1 is selected from $-CH_2-$, ,  and ;


or

a is 0, R^1 is CH_3- , R^2 is F, and L^1 is selected from  and , wherein
“#” marks the position attached to carbonyl; or


a is 1, R^1 is CH_3- , R^2 is F, L^1 is , M is O, and L^2 is $-C_2H_4-$.


[7] The compound of any one of [1] to [6], wherein the payload is selected from:






preferably selected from:






more preferably selected from:







particularly selected from:




- [8] The compound of any one of [1] to [7], wherein
 R^0 is C_{1-6} alkyl, preferably C_{1-3} alkyl; particularly methyl; and/or
 n is an integer of 2 to 5, particularly 3; and/or
 k_1 and k_2 are independently 1, or 3 or 5, particularly 5;
 i is independently an integer of 1 to 20, preferably 1 to 12, more preferably 2 to 8,
 particularly 4; and/or
 j is independently an integer of 1 to 20, preferably 1 to 12, more preferably 8 to 12,


especially 8 or 12, particularly 12.

[9] The compound of any one of [1] to [8], which is selected from





[10] A conjugate having the structure of formula (II):






wherein,

A is an antibody or an antigen binding fragment thereof; the antibody or antigen binding fragment is preferably modified to connect with the (Gly)_n moiety in the compound of formula (I);

z is an integer of 1 to 20; preferably 1 to 4; particularly 2;

P1, P2, R⁰, *opSu*, n, k1, k2, i and j are as defined in any one of [1] to [8].

[11] The conjugate of any one of [10], which is selected from:



[12] The conjugate of any one of [10] to [11], wherein the antibody is an anti-human HER2 antibody, preferably Trastuzumab.

[13] A pharmaceutical composition comprising a prophylactically or therapeutically effective amount of a conjugate of any one of [10] to [12], and at least one pharmaceutically acceptable carrier.


[14] The pharmaceutical composition of [13], wherein the conjugate has a drug to antibody ratio (DAR) of an integer or non-integer of 1 to 8, particularly 3 to 4.

[15] Use of the conjugate of any one of [10] to [12] or the pharmaceutical composition of claim [13] or [14] in the manufacture of a medicament for treating a disease; wherein the disease is a tumor or an autoimmune disease; preferably a tumor comprising HER2-positive tumor cells.

[16] Use of [15], wherein the tumor comprising HER2-positive tumor cells further comprises HER2 low-expressing tumor cells;

wherein the HER2 low-expressing tumor cells express lower levels of HER2 than the HER2-positive tumor cells.


[17] A compound, or a pharmaceutically acceptable salt, stereoisomer, solvate, polymorph, tautomer, isotopic compound, metabolite or prodrug thereof, wherein the compound has the structure of formula (i):




wherein,


a, the carbon atoms marked with p1* and p2*, L¹, M, L², R¹ and R² are as defined in any one of [1] to [7].

[18] The compound of [17], which is selected from:







preferably selected from:




more preferably selected from:



particularly selected from:



[19] A compound of formula (1):




wherein,

k is an integer of 1 to 7; preferably 1, or 3 or 5, particularly 5

P is a payload having the structure of formula (i'), wherein the structure of formula (i') is as defined in any one of [1] to [7].

[20] The compound of [19], which is selected from:



Beneficial effects

[0159] The novel small molecule topoisomerase I inhibitors provided in the present disclosure, either used alone or as a component of ADCs, could show greater activity, stability, physiochemical properties over the prior art.

[0160] The antibody-drug conjugate of the present invention uses specially designed linker-payload, and can achieve great efficacy and bystander killing effects. At the same time, it has a lower DAR, and therefore can reduce side effects and increase the therapeutic index, which is

of special importance for bystander killing. The antibody-drug conjugate of the present invention is more stable in its structure such as the ring-opened succinimide structure.

[0161] The present disclosure utilizes a linker with unique structure and uses a ligase to catalyze the conjugation of the targeting molecule (antibody) and the payload. The conjugate of the present disclosure has good homogeneity and high activity. Furthermore, the toxicity of the linker-payload intermediate is much lower than that of the free payload, and thus the manufacture process of the drug is less detrimental, which is advantageous for industrial production.

[0162] The conjugate of the present disclosure achieves at least one of the following technical effects:

- (1) High inhibitory activity against target cells, and strong bystander killing effect.
- (2) Good physicochemical properties (e.g., solubility, physical and/or chemical stability).
- (3) Good pharmacokinetic properties (e.g., good stability in plasma, appropriate half-life and duration of action).
- (4) Good safety (low toxicity on non-target normal cells or tissues, and/or fewer side effects, wider treatment window), etc.
- (5) Highly modular design, simple assembly of multiple drugs.

Examples

[0163] Preparation example

[0164] In order to more clearly illustrate the objects and technical solutions, the present disclosure is further described below with reference to specific examples. It is to be understood that the examples are not intended to limit the scope of the disclosure. The specific experimental methods which were not mentioned in the following examples were carried out according to conventional experimental method.

Instruments, Materials and Reagents

[0165] Unless otherwise stated, the instruments and reagents used in the examples are commercially available. The reagents can be used directly without further purification.

MS: Thermo Fisher Q Exactive Plus, Water2795-Quattro micro triple quadrupole mass spectrometer

HPLC : Waters 2695, Agilent 1100, Agilent 1200

Semi-preparative HPLC: Lisure HP plus 50D

Flow Cytometry: CyttoFLEX S

HIC-HPLC: Butyl-HIC; mobile phase A: 25 mM PB, 2M (NH₄)₂SO₄, pH 7.0; mobile phase B: 25 mM PB, pH 7.0; flow rate: 0.8 ml/min; acquisition time: 25 min; injection amount: 20 µg; column temperature: 25 °C; detection wavelength: 280 nm; sample chamber temperature:

8 °C.


SEC-HPLC: column: TSK-gel G3000 SWXL, TOSOH 7.8 mm ID × 300 mm, 5 μm; mobile phase: 0.2 M KH₂PO₄, 0.25 M KCl, pH 6.2; flow rate : 0.5 ml/min; acquisition time: 30 min; injection volume: 50 μl; column temperature: 25 °C; detection wavelength; 280 nm; sample tray temperature: 8 °C.

CHO was obtained from Thermo Fisher Scientific; pcDNA 3.3 was obtained from Life Technology; HEK293F was obtained from Prejin; PEIMAX transfection reagent was obtained from Polyscience; MabSelect Sure ProA was obtained from GE; Capto S ImpAct was obtained from GE; Rink-amide-MBHA-resin and dichloro resin were obtained from Nankai synthesis; HCC1954 was obtained from ATCC CAT# CRL-2338; SK-BR-3 was obtained from ATCC CAT# HTB-30; BT-474 was obtained from ATCC CAT# HTB-20; NCI-N87 cells was obtained from ATCC CAT# CRL-5822; MCF7 was obtained from ATCC CAT# HTB-22; MDA-MB-231 was obtained from ATCC CAT# HTB-26; MDA-MB-468 was obtained from ATCC CAT# HTB-132; CFPAC-1 was obtained from ATCC CAT# CRL-1918; NCI-H2110 was obtained from ATCC CAT# CRL-5924; JIMT-1 was obtained from Wuxi Apptech; Capan-1 was obtained from ATCC CAT# CRL-1573; antibody Trastuzumab is prepared according to the known sequence; optimized recombinant enzyme Sortase A derived from *Staphylococcus aureus* is prepared in *E.coli*.

[0166] In some cases, the order of carrying out the foregoing reaction schemes may be varied to facilitate the reaction or to avoid unwanted reaction products. The following examples are provided so that the invention might be more fully understood. These examples are illustrative only and should not be construed as limiting the invention in any way.

Example 1 Preparation of payload small molecules

Intermediate 11



Step A: N-(2-bromo-5-fluorophenyl)acetamide: To a stirred solution of acetic anhydride (214 g, 2.10 mol) in acetic acid (500 mL) was added con. H₂SO₄ (3 mL), followed with 2-bromo-5-fluoroaniline (100 g, 526.27 mmol) in portions at room temperature. The mixture was stirred for 3 h, then poured into 2000 mL ice-water. A precipitate was formed, which was collected by filtration and dried in vacuo at room temperature to afford N-(2-bromo-5-fluorophenyl)acetamide (105 g) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.68 (dd, *J* = 8.9, 6.0 Hz, 1H), 7.61 (ddd, *J* = 10.7, 5.3, 3.1 Hz, 1H), 7.02 (ddd, *J* = 8.9, 8.0, 3.1 Hz, 1H), 2.11 (s, 3H). LCMS *m/z* 232.0(M+H).

Step B: N-(5-fluoro-2-(1-hydroxycyclobutyl)phenyl)acetamide: To a stirred solution of N-(2-bromo-5-fluorophenyl)acetamide (105 g, 452.48 mmol) in THF (1000 mL) was added *n*-BuLi (594 mL, 1.6 M in *n*-hexane, 950.22 mmol) dropwise over 1 h at -78 °C. After completion, the mixture was stirred for 0.5 h under N₂. Then a solution of cyclobutanone (38.06 g, 542.98 mmol) in THF (50 mL) was added dropwise at -78 °C over 0.5 h, the mixture was stirred at -78 °C to room temperature for 6 h. The mixture was poured into 500 mL saturated NH₄Cl aq at 0 °C. Extracted with ethyl acetate (500 mL x 3), washed with brine (250 mL x 2), dried over Na₂SO₄ and concentrated. The mixture was triturated with (PE/EA=1:1, 100 mL) for 10 mins, filtered and the cake was collected and dried in vacuo to afford N-(5-fluoro-2-(1-hydroxycyclobutyl)phenyl)acetamide (24 g) as a yellow solid. LCMS *m/z* 206.1(M-18+H), 246.1(M+Na).

Step C: N-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)acetamide: To a stirred mixture of N-(5-fluoro-2-(1-hydroxycyclobutyl)phenyl)acetamide (24 g, 107.50 mmol) in CH₂Cl₂ (170 mL) and water (170 mL) was added silver nitrate (AgNO₃) (5.48 g, 32.25 mmol) and potassium persulfate (K₂S₂O₈) (58.12 g, 215.01 mmol), the mixture was stirred at 30 °C for 6 h. The mixture was filtered on Celite and washed with CH₂Cl₂ (100 mL), the filtrate was concentrated and purified by FCC(EA/PE=0-40%) to afford N-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)acetamide (14 g) as a light yellow solid. LCMS *m/z* 222.1(M+H).

Step D: N-(3-fluoro-7-(hydroxyimino)-8-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)acetamide: To a stirring mixture of N-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)acetamide (14 g, 63.28 mmol) in THF (500 mL) at 0 °C was added 1-butyl nitrite (8.48 g, 63.28 mmol), followed with *t*-BuOK (8.52 g, 75.94 mmol). The mixture was stirred at 0 °C for 2 h. After completion, the mixture was acidified by HCl (2 N) to adjust pH=3. The mixture was extracted by ethyl acetate (200 mL x 3), washed by brine (100 mL x 2), dried over Na₂SO₄ and concentrated under reduced pressure. The crude mixture was triturated with tert-butyl methyl ether (200 mL) for 10 mins, filtered and the cake was collected and dried in vacuo to afford N-(3-fluoro-7-(hydroxyimino)-8-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)acetamide (12 g) as a yellow solid. LCMS *m/z* 251.1(M+H).

Step E: N,N'-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalene-1,7-diyl)diacetamide: To a solution of N-(3-fluoro-7-(hydroxyimino)-8-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)acetamide (12 g, 47.96 mmol) in acetic anhydride (90 mL) and THF (90 mL) was added 10% Pd/C (1 g), the mixture was stirred at 25 °C under H₂ atmosphere for 16 h. After cooling to 0 °C, Et₃N (20 mL) was added dropwise, the mixture was stirred at 0 °C for 1 h. Filtered on Celite, the filtrate was poured into ice-water (500 mL). Extracted with ethyl acetate (500 mL x 3), washed with brine (250 mL x 2), dried over Na₂SO₄ and concentrated. The residue was triturated with tert-butyl methyl ether (120 mL) for 10 mins, filtered and the cake was collected and dried in

vacuo to give N,N'-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalene-1,7-diyl)diacetamide (7.9 g) as a yellow solid. LCMS m/z 279.1(M+H).


Step F: N,N'-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalene-1,7-diyl)diacetamide: To a solution of N,N'-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalene-1,7-diyl)diacetamide (7.9 g, 28.39 mmol) in MeOH (150 mL) was added HCl aq (2 N, 150 mL), the mixture was stirred at 50 °C for 7 h. After cooling to 0 °C, Sat. NaHCO₃ aq was added dropwise to adjust pH = 8. Extracted with ethyl acetate (200 mL x 3), washed with brine (200 mL x 2), dried over Na₂SO₄ and concentrated under reduced pressure to give N,N'-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalene-1,7-diyl)diacetamide (6.0 g) as a yellow solid. ¹H NMR (400 MHz, Chloroform-d) δ 6.57 (s, 3H), 6.18 (td, J = 11.1, 2.4 Hz, 2H), 4.52 (dt, J = 13.3, 5.0 Hz, 1H), 3.13 (ddd, J = 17.5, 13.0, 4.6 Hz, 1H), 3.00 – 2.81 (m, 1H), 2.69 (dtd, J = 9.4, 4.6, 2.5 Hz, 1H), 2.09 (s, 3H), 1.79 (qd, J = 13.0, 4.3 Hz, 1H). LCMS m/z 237.1(M+H).

Step G: N-(8-amino-5-chloro-6-fluoro-1-oxo-1,2,3,4-tetrahydronaphthalen-2-yl)acetamide: To a solution of N,N'-(3-fluoro-8-oxo-5,6,7,8-tetrahydronaphthalene-1,7-diyl)diacetamide (4.0 g, 16.93 mmol) in DMF (80 mL) was added NCS (2.26 g, 16.93 mmol) in portions at 0 °C, the mixture was stirred at room temperature for 16 h. The mixture was poured into 200 mL ice-water. A precipitate was formed, which was collected by filtration and dried in vacuo at room temperature to afford N-(8-amino-5-chloro-6-fluoro-1-oxo-1,2,3,4-tetrahydronaphthalen-2-yl)acetamide (4.0 g) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.11 (d, J = 8.0 Hz, 1H), 7.71 (s, 2H), 6.62 (d, J = 11.9 Hz, 1H), 4.53 (ddd, J = 13.0, 8.0, 4.7 Hz, 1H), 3.18 – 3.04 (m, 1H), 2.91 (ddd, J = 17.5, 12.4, 4.8 Hz, 1H), 2.21 – 2.08 (m, 1H), 1.99 – 1.83 (m, 4H). LCMS m/z 271.0(M+H).

Step H: N-((9S)-4-chloro-9-ethyl-5-fluoro-9-hydroxy-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)acetamide: To a mixture of N-(8-amino-5-chloro-6-fluoro-1-oxo-1,2,3,4-tetrahydronaphthalen-2-yl)acetamide (4.0 g, 14.78 mmol) in toluene (400 mL) was added (S)-4-ethyl-4-hydroxy-7,8-dihydro-1H-pyrano[3,4-f]indolizine-3,6,10(4H)-trione (4.28 g, 16.25 mmol), pyridinium p-Toluenesulfonate (1.11 g, 4.43 mmol) and o-cresol (10 mL), the mixture was heated to reflux under N₂ for 24 h. The solvent was removed by reduced pressure and the mixture was purified by FCC(THF/CH₂Cl₂=0-60%) to afford N-((9S)-4-chloro-9-ethyl-5-fluoro-9-hydroxy-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)acetamide (4.1 g) as a brown solid. LCMS m/z 498.1(M+H).


Step I: (9S)-1-amino-4-chloro-9-ethyl-5-fluoro-9-hydroxy-1,2,3,9,12,15-hexahydro-10H,13H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione: A mixture of N-((9S)-4-chloro-9-ethyl-5-fluoro-9-hydroxy-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)acetamide (2.0 g, 4.02 mmol) in 20 mL con. HCl aq was stirred at 70 °C under N₂ for 36 h. The mixture was concentrated under reduced pressure to give crude (9S)-1-amino-4-chloro-9-ethyl-5-fluoro-9-hydroxy-1,2,3,9,12,15-hexahydro-10H,13H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione hydrochloride (2 g) as a brown solid. LCMS (ESI) m/z 456.1 [M+H]⁺.

Intermediate 12




Intermediate 12 : (9S)-1-amino-4-bromo-9-ethyl-5-fluoro-9-hydroxy-1,2,3,9,12,15-hexahydro-10H,13H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione hydrobromide was synthesized from 2-bromo-5-fluoroaniline using a similar procedure as above for intermediate **11**. LCMS (ESI) m/z 500.0 [M+H]⁺.

Intermediate 13





Intermediate 13 : (9S)-1-amino-9-ethyl-5-fluoro-9-hydroxy-4-methoxy-1,2,3,9,12,15-hexahydro-10H,13H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione hydrochloride was synthesized from 2-bromo-5-fluoro-4-methoxyaniline using a similar procedure as above for intermediate **11**. LCMS (ESI) m/z 452.1 [M+H]⁺.

Procedure of C730 and C731




C730 and **C731** was prepared by prep-HPLC from (9S)-1-amino-4-chloro-9-ethyl-5-fluoro-9-hydroxy-1,2,3,9,12,15-hexahydro-10H,13H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione hydrochloride (intermediate **11**) as TFA salt.

Table I.

Number	Structure	¹ HNMR	MS [M+H] ⁺	Retention time on HPLC (min)
C730		¹ H NMR (400 MHz, DMSO-d ₆) δ 8.51 (d, <i>J</i> = 4.8 Hz, 3H), 8.17 (d, <i>J</i> = 10.2 Hz, 1H), 7.38 (s, 1H), 6.56 (s, 1H), 5.74 (d, <i>J</i> = 19.4 Hz, 1H), 5.52 – 5.40 (m, 3H), 5.16 (s, 1H), 3.44 (dd, <i>J</i> = 16.3, 4.1 Hz, 1H), 3.19 (t, <i>J</i> = 13.9 Hz, 1H), 2.57 (d, <i>J</i> = 14.0 Hz, 1H), 2.26 (t, <i>J</i> = 14.3 Hz, 1H), 1.89 (hept, <i>J</i> = 7.0 Hz, 2H), 0.89 (t, <i>J</i> = 7.3 Hz, 3H).	456.0	1.395
C731		¹ H NMR (400 MHz, DMSO-d ₆) δ 8.56 (s, 2H), 8.17 (d, <i>J</i> = 10.1 Hz, 1H), 7.38 (s, 1H), 6.57 (s, 1H), 5.74 (d, <i>J</i> = 19.4 Hz, 1H), 5.53 – 5.28 (m, 3H), 5.16 (s, 1H), 3.59 – 3.01 (m, 2H), 2.35 – 2.19 (m, 2H), 2.00 – 1.78 (m, 2H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H).	456.0	1.541

Conditions of HPLC above: Equipment: Agilent 1200; Chromatographic column: Waters XBridge C18 4.6*50mm,3.5μm; Flow:2.0mL/min; Gradient elute: 5.0%-95.0%-95.0%-5.0%-5.0%, 0.00min-1.50min-2.50min-2.52min-3.00min; Temperature : 40°C; Mobile phase : A: Acetonitrile, B: H₂O (0.05% TFA); Wavelength: 214 nm/254 nm.

Procedure of C518












Step J: N-((1S,9S)-4-chloro-9-ethyl-5-fluoro-9-hydroxy-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)-2-hydroxyacetamide: To a solution of crude (9S)-1-amino-4-chloro-9-ethyl-5-fluoro-9-hydroxy-1,2,3,9,12,15-hexahydro-10H,13H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione hydrochloride (50 mg, 0.10 mmol) in 2 mL DMF was added 2-hydroxyacetic acid (11.6 mg, 0.15mmol), HATU (57.9 mg, 0.15 mmol) and followed with Et₃N (20.5 mg, 0.20mmol) dropwise at 0 °C, the mixture was stirred at 0 °C to room temperature for 2 h. The mixture was poured into 10 mL ice-water, extracted with ethyl acetate (20 mL x 3), washed with brine (20 mL x 2), dried over Na₂SO₄ and concentrated. The residue was purified by prep-HPLC to give N-((1S,9S)-4-chloro-9-ethyl-5-fluoro-9-hydroxy-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)-






2-hydroxyacetamide (6 mg) as a yellow solid and N-((1R,9S)-4-chloro-9-ethyl-5-fluoro-9-hydroxy-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)-2-hydroxyacetamide (5 mg) as a yellow solid.




The compounds prepared in the table II below was reacted and post-treated in the same manner as in Example C518.

Table II.

Number	Structure	¹ H NMR	MS [M+H] ⁺	Retention time on HPLC (min)
C518		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.50 (d, <i>J</i> = 8.9 Hz, 1H), 8.04 (d, <i>J</i> = 10.3 Hz, 1H), 7.33 (s, 1H), 6.54 (s, 1H), 5.80 – 5.48 (m, 2H), 5.42 (s, 2H), 5.21 (s, 2H), 3.96 (d, <i>J</i> = 5.8 Hz, 2H), 3.26 (d, <i>J</i> = 20.3 Hz, 2H), 2.22 (dd, <i>J</i> = 18.8, 9.7 Hz, 2H), 1.86 (p, <i>J</i> = 7.5 Hz, 2H), 0.88 (d, <i>J</i> = 7.3 Hz, 3H).	514.0	1.548
C519		¹ H NMR (400 MHz, Methanol- <i>d</i> ₄) δ 7.86 (d, <i>J</i> = 9.9 Hz, 1H), 7.66 (s, 1H), 5.74 (s, 1H), 5.61 (d, <i>J</i> = 16.4 Hz, 1H), 5.42 (dd, <i>J</i> = 17.4, 8.4 Hz, 2H), 5.22 (d, <i>J</i> = 18.9 Hz, 1H), 4.34 – 4.12 (m, 2H), 3.53 (d, <i>J</i> = 18.8 Hz, 2H), 2.35 (d, <i>J</i> = 30.7 Hz, 2H), 1.98 (s, 2H), 1.02 (t, <i>J</i> = 7.4 Hz, 3H).	514.0	1.602
C586		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.56 (d, <i>J</i> = 8.9 Hz, 1H), 8.07 (d, <i>J</i> = 10.2 Hz, 1H), 7.34 (s, 1H), 6.55 (s, 1H), 5.74 – 5.61 (m, 1H), 5.44 (s, 2H), 5.24 (s, 2H), 4.69 (s, 1H), 4.05 (s, 2H), 3.62 – 3.42 (m, 4H), 3.41 – 3.21 (m, 2H), 2.33 – 2.15 (m, 2H), 1.97 – 1.78 (m, 2H), 0.88 (t, <i>J</i> = 7.2 Hz, 3H).	558.0	1.621
C587		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.56 (d, <i>J</i> = 8.9 Hz, 1H), 8.07 (d, <i>J</i> = 10.2 Hz, 1H), 7.35 (s, 1H), 6.54 (s, 1H), 5.76 – 5.61 (m, 1H), 5.44 (s, 2H), 5.23 (s, 2H), 4.68 (s, 1H), 4.05 (s, 2H), 3.60 – 3.45 (m, 4H), 3.34 – 3.23 (m, 2H), 2.26 – 2.16 (m, 2H), 2.04 – 1.81 (m, 2H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H).	558.0	1.675

C593		^1H NMR (400 MHz, DMSO- d_6) δ 8.66 (d, J = 9.2 Hz, 1H), 8.05 (d, J = 10.7 Hz, 1H), 7.34 (s, 1H), 6.54 (s, 1H), 6.29 (s, 1H), 5.71 – 5.48 (m, 1H), 5.44 (s, 2H), 5.32 – 5.04 (m, 2H), 3.33 – 3.06 (m, 2H), 2.37 – 2.13 (m, 2H), 2.04 – 1.69 (m, 2H), 1.30 – 1.09 (m, 2H), 1.00 – 0.64 (m, 5H).	540.0	1.746
C594		^1H NMR (400 MHz, DMSO- d_6) δ 8.69 (d, J = 9.4 Hz, 1H), 8.05 (d, J = 10.2 Hz, 1H), 7.34 (s, 1H), 6.54 (s, 1H), 6.31 (s, 1H), 5.67 – 5.55 (m, 1H), 5.44 (s, 2H), 5.36 – 5.06 (m, 2H), 3.33 – 3.07 (m, 2H), 2.38 – 2.15 (m, 2H), 1.96 – 1.79 (m, 2H), 1.28 – 1.13 (m, 2H), 1.05 – 0.92 (m, 2H), 0.88 (t, J = 7.2 Hz, 3H).	540.0	1.783
C622		^1H NMR (400 MHz, DMSO- d_6) δ 8.43 (d, J = 8.9 Hz, 1H), 8.06 (d, J = 10.2 Hz, 1H), 7.34 (s, 1H), 6.53 (s, 1H), 5.62 (d, J = 8.0 Hz, 1H), 5.51 (d, J = 5.2 Hz, 1H), 5.43 (s, 2H), 5.37 – 5.15 (m, 2H), 3.61 (t, J = 5.8 Hz, 1H), 3.34 – 3.12 (m, 2H), 2.25 – 2.15 (m, 2H), 1.96 – 1.80 (m, 2H), 1.29 – 1.15 (m, 1H), 0.88 (t, J = 7.3 Hz, 3H), 0.58 – 0.27 (m, 4H).	554.0	1.694
C624	 (mixture)	^1H NMR (400 MHz, DMSO- d_6) δ 8.43 (dd, J = 8.7, 5.3 Hz, 1H), 8.05 (d, J = 10.2 Hz, 1H), 7.34 (d, J = 1.6 Hz, 1H), 6.54 (d, J = 2.0 Hz, 1H), 5.68 – 5.52 (m, 1H), 5.49 – 5.35 (m, 2H), 5.34 – 5.11 (m, 2H), 3.81 – 3.57 (m, 1H), 3.35 – 3.22 (m, 2H), 2.29 – 2.07 (m, 2H), 1.96 – 1.71 (m, 2H), 1.19 – 1.06 (m, 1H), 0.88 (t, J = 7.3 Hz, 3H), 0.46 – 0.24 (m, 4H).	554.0	1.753
C625	 (mixture)	^1H NMR (400 MHz, DMSO- d_6) δ 8.44 (t, J = 7.7 Hz, 1H), 8.08 (d, J = 10.2 Hz, 1H), 7.34 (s, 1H), 6.54 (s, 1H), 5.70 – 5.54 (m, 1H), 5.44 (s, 2H), 5.37 – 5.16 (m, 2H), 4.16 – 3.93 (m, 1H), 3.38 – 2.84 (m, 2H), 2.35 – 2.21 (m, 2H), 2.22 – 2.09 (m, 2H), 1.96 – 1.74 (m, 2H), 1.09 (d, J = 6.2 Hz, 3H), 0.88 (t, J = 7.4 Hz, 3H).	542.0	1.646


C677		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.50 (d, <i>J</i> = 9.2 Hz, 1H), 8.04 (d, <i>J</i> = 10.2 Hz, 1H), 7.33 (s, 1H), 6.53 (s, 1H), 5.87 – 5.53 (m, 1H), 5.42 (s, 2H), 5.33 – 5.04 (m, 2H), 4.26 – 4.06 (m, 1H), 3.34 – 3.15 (m, 2H), 2.27 – 2.12 (m, 2H), 2.02 – 1.75 (m, 2H), 1.40 (d, <i>J</i> = 6.8 Hz, 3H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	528.0	1.658
C678		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.46 (d, <i>J</i> = 8.9 Hz, 1H), 8.05 (d, <i>J</i> = 10.2 Hz, 1H), 7.34 (s, 1H), 6.53 (s, 1H), 5.71 – 5.52 (m, 1H), 5.43 (s, 2H), 5.34 – 5.02 (m, 2H), 4.47 – 3.80 (m, 1H), 3.34 – 3.13 (m, 2H), 2.31 – 2.10 (m, 2H), 2.02 – 1.76 (m, 2H), 1.30 (d, <i>J</i> = 6.7 Hz, 3H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	528.0	1.717
C679		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.47 (d, <i>J</i> = 8.8 Hz, 1H), 8.06 (d, <i>J</i> = 10.3 Hz, 1H), 7.34 (s, 1H), 6.54 (s, 1H), 5.77 – 5.51 (m, 1H), 5.44 (s, 2H), 5.34 – 5.10 (m, 2H), 3.50 – 3.35 (m, 2H), 3.33 – 3.14 (m, 2H), 2.28 – 2.19 (m, 2H), 2.19 – 2.09 (m, 2H), 1.92 – 1.80 (m, 2H), 1.79 – 1.62 (m, 2H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	542.0	1.603
C680		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.45 (d, <i>J</i> = 8.8 Hz, 1H), 8.06 (d, <i>J</i> = 10.2 Hz, 1H), 7.33 (s, 1H), 6.54 (s, 1H), 5.69 – 5.54 (m, 1H), 5.43 (s, 2H), 5.36 – 5.10 (m, 2H), 3.42 – 3.37 (m, 2H), 3.33 – 3.26 (m, 2H), 2.27 – 2.07 (m, 4H), 1.99 – 1.79 (m, 2H), 1.77 – 1.67 (m, 2H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	542.0	1.645
C703		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.96 (d, <i>J</i> = 8.7 Hz, 1H), 8.07 (d, <i>J</i> = 10.2 Hz, 1H), 7.34 (s, 1H), 7.13 (s, 1H), 6.55 (s, 1H), 5.63 – 5.53 (m, 1H), 5.43 (s, 2H), 5.31 – 5.03 (m, 2H), 3.31 – 3.24 (m, 2H), 2.31 – 2.11 (m, 2H), 1.95 – 1.75 (m, 2H), 1.55 (s, 3H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H).	596.0	1.792







C704		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.06 (d, <i>J</i> = 9.2 Hz, 1H), 8.06 (d, <i>J</i> = 10.3 Hz, 1H), 7.35 (s, 1H), 7.33 (s, 1H), 6.56 (s, 1H), 5.67 – 5.59 (m, 1H), 5.44 (s, 2H), 5.31 – 5.02 (m, 2H), 3.32 – 3.19 (m, 2H), 2.30 – 2.10 (m, 2H), 1.96 – 1.80 (m, 2H), 1.71 (s, 3H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H).	596.0	1.896
C627		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.00 (d, <i>J</i> = 8.5 Hz, 1H), 8.07 (d, <i>J</i> = 10.3 Hz, 1H), 7.34 (s, 1H), 6.55 (s, 1H), 5.70 – 5.47 (m, 1H), 5.43 (s, 2H), 5.34 – 5.06 (m, 2H), 4.72 – 4.52 (m, 1H), 3.34 – 3.16 (m, 2H), 2.29 – 2.09 (m, 2H), 1.99 – 1.75 (m, 2H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	582.0	1.835
C626		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.02 (d, <i>J</i> = 8.6 Hz, 1H), 8.07 (d, <i>J</i> = 10.2 Hz, 1H), 7.34 (s, 1H), 7.21 (d, <i>J</i> = 6.9 Hz, 1H), 6.55 (s, 1H), 5.73 – 5.58 (m, 1H), 5.43 (s, 2H), 5.29 (d, <i>J</i> = 19.2 Hz, 1H), 5.13 (d, <i>J</i> = 19.2 Hz, 1H), 4.63 (d, <i>J</i> = 8.2 Hz, 1H), 3.26 (d, <i>J</i> = 7.7 Hz, 2H), 2.20 (p, <i>J</i> = 6.8 Hz, 2H), 1.86 (dt, <i>J</i> = 15.5, 7.2 Hz, 2H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	582.0	1.820






Conditions of HPLC above: Equipment: Agilent 1200; Chromatographic column: Waters XBridge C18 4.6*50mm,3.5μm; Flow:2.0mL/min; Gradient elute: 5.0%-95.0%-95.0%-5.0%-5.0%, 0.00min-1.50min-2.50min-2.52min-3.00min; Temperature : 40°C; Mobile phase : A: Acetonitrile, B: H₂O (0.05% TFA); Wavelength: 214 nm/254 nm.





The compounds prepared in the table **III** below was synthesized from intermediate **12** or intermediate **13** using a similar procedure as above for **C518**.

Table III.

Number	Structure	¹ H NMR	MS [M+H] ⁺	Retention time on HPLC (min)
C636		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.49 (d, <i>J</i> = 9.1 Hz, 1H), 8.01 (d, <i>J</i> = 9.8 Hz, 1H), 7.34 (s, 1H), 6.54 (s, 1H), 5.71 – 5.56 (m, 1H), 5.58 – 5.45 (m, 1H), 5.43 (s, 2H), 5.22 (s, 2H), 3.96 (d, <i>J</i> = 5.7 Hz, 2H), 3.35 – 3.23 (m, 2H), 2.30 – 2.10 (m, 2H), 1.96 – 1.79 (m, 2H), 0.87 (t, <i>J</i> = 8.3, 5.9 Hz, 3H).	558.0	1.612

C637		^1H NMR (400 MHz, DMSO- d_6) δ 8.51 (d, J = 8.9 Hz, 1H), 8.01 (d, J = 9.8 Hz, 1H), 7.35 (s, 1H), 6.54 (s, 1H), 5.74 – 5.58 (m, 1H), 5.52 (t, J = 5.8 Hz, 1H), 5.43 (s, 2H), 5.29 – 5.16 (m, 2H), 3.97 (d, J = 5.5 Hz, 2H), 3.36 – 3.14 (m, 2H), 2.26 – 2.14 (m, 2H), 1.96 – 1.78 (m, 2H), 0.88 (t, J = 7.3 Hz, 3H).	558.0	1.642
C595	 (mixture)	^1H NMR (400 MHz, DMSO- d_6) δ 8.62 – 8.36 (m, 1H), 8.01 (d, J = 10.2 Hz, 1H), 7.34 (s, 1H), 6.68 – 6.45 (m, 2H), 5.73 – 5.49 (m, 1H), 5.43 (s, 2H), 5.36 – 5.04 (m, 2H), 4.13 (s, 1H), 3.22 – 2.99 (m, 2H), 2.36 – 2.03 (m, 2H), 1.97 – 1.69 (m, 2H), 1.48 – 1.10 (m, 3H), 0.87 (t, 3H).	572.0	1.747
C596		^1H NMR (400 MHz, DMSO- d_6) δ 8.52 (d, 1H), 8.02 (d, J = 14.2 Hz, 1H), 7.33 (s, 1H), 6.56 (d, J = 17.9 Hz, 2H), 5.71 – 5.54 (m, 1H), 5.42 (s, 2H), 5.34 – 5.06 (m, 2H), 4.13 (s, 1H), 3.25 – 2.94 (m, 2H), 2.27 – 2.07 (m, 2H), 2.03 – 1.72 (m, 2H), 1.53 – 1.10 (m, 3H), 0.86 (t, 3H).	572.0	1.720
C597		^1H NMR (400 MHz, DMSO- d_6) δ 8.47 (d, J = 8.9 Hz, 1H), 8.02 (d, J = 9.7 Hz, 1H), 7.35 (s, 1H), 6.54 (s, 2H), 5.64 – 5.50 (m, 1H), 5.44 (s, 2H), 5.31 – 5.08 (m, 2H), 4.17 – 4.08 (m, 1H), 3.33 – 3.00 (m, 2H), 2.26 – 2.12 (m, 2H), 1.96 – 1.74 (m, 2H), 1.30 (d, J = 6.7 Hz, 3H), 0.88 (t, J = 7.3 Hz, 3H).	572.0	1.773
C600			586.0	1.792
C601			586.0	1.712


C602			586.0	1.741
C603			586.0	1.682
C619	 (mixture)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.06 (d, <i>J</i> = 8.1 Hz, 1H), 8.03 (d, <i>J</i> = 9.7 Hz, 1H), 7.38 (s, 1H), 6.55 (s, 1H), 5.74 – 5.54 (m, 1H), 5.43 (s, 2H), 5.36 – 5.05 (m, 2H), 4.77 – 4.56 (m, 1H), 3.13 – 3.00 (m, 2H), 2.27 – 2.12 (m, 2H), 1.95 – 1.78 (m, 2H), 0.87 (t, <i>J</i> = 7.2 Hz, 3H).	625.9	1.785
C620	 (mixture)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.06 (d, <i>J</i> = 8.1 Hz, 1H), 8.03 (d, <i>J</i> = 9.7 Hz, 1H), 7.38 (s, 1H), 6.55 (s, 1H), 5.74 – 5.54 (m, 1H), 5.43 (s, 2H), 5.36 – 5.05 (m, 2H), 4.77 – 4.56 (m, 1H), 3.13 – 3.00 (m, 2H), 2.27 – 2.12 (m, 2H), 1.95 – 1.78 (m, 2H), 0.87 (t, <i>J</i> = 7.2 Hz, 3H).	626.0	1.847
C664	 (mixture)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.46 (d, <i>J</i> = 8.9 Hz, 1H), 7.91 (d, <i>J</i> = 12.4 Hz, 1H), 7.31 (s, 1H), 6.53 (s, 1H), 5.59 (s, 1H), 5.42 (s, 2H), 5.19 (d, <i>J</i> = 2.8 Hz, 2H), 3.97 (s, 3H), 3.20 (dt, <i>J</i> = 13.7, 6.8 Hz, 3H), 2.23 – 2.07 (m, 2H), 1.86 (s, 2H), 1.24 (s, 2H), 0.87 (t, <i>J</i> = 7.2 Hz, 3H).	510.1	1.510

C665	 (mixture)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.42 (d, <i>J</i> = 8.7 Hz, 1H), 7.93 (d, <i>J</i> = 12.4 Hz, 1H), 7.30 (s, 1H), 6.52 (s, 1H), 5.57 (s, 1H), 5.43 (s, 2H), 5.24 (d, <i>J</i> = 6.7 Hz, 2H), 4.04 (d, <i>J</i> = 6.5 Hz, 1H), 3.97 (s, 3H), 3.18 (d, <i>J</i> = 25.1 Hz, 2H), 2.26 (d, <i>J</i> = 7.5 Hz, 2H), 2.20 (d, <i>J</i> = 5.8 Hz, 1H), 2.09 (s, 2H), 1.93 – 1.79 (m, 2H), 1.09 (d, <i>J</i> = 6.2 Hz, 3H), 0.87 (s, 3H).	538.2	1.521
C667	 (mixture)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.60 (d, <i>J</i> = 9.0 Hz, 1H), 7.88 (d, <i>J</i> = 12.4 Hz, 1H), 7.30 (s, 1H), 6.52 (s, 1H), 6.30 (s, 1H), 5.56 (d, <i>J</i> = 7.0 Hz, 1H), 5.42 (s, 2H), 5.34 – 5.01 (m, 2H), 3.96 (s, 3H), 3.13 (dd, <i>J</i> = 16.9, 7.7 Hz, 2H), 2.18 (t, <i>J</i> = 6.2 Hz, 2H), 1.96 – 1.77 (m, 2H), 1.26 – 1.14 (m, 2H), 0.94 (d, <i>J</i> = 3.5 Hz, 2H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	536.1	1.578
C668	 (mixture)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.48 (d, <i>J</i> = 9.2 Hz, 1H), 7.92 (s, 1H), 7.30 (s, 1H), 6.53 (s, 1H), 5.61 (dd, <i>J</i> = 21.7, 6.1 Hz, 2H), 5.42 (s, 2H), 5.23 (d, <i>J</i> = 19.1 Hz, 2H), 5.10 (d, <i>J</i> = 19.0 Hz, 1H), 4.12 (d, <i>J</i> = 6.8 Hz, 1H), 3.96 (d, <i>J</i> = 1.1 Hz, 3H), 3.25 – 3.04 (m, 2H), 2.13 (d, <i>J</i> = 6.4 Hz, 2H), 1.86 (s, 2H), 1.40 (d, <i>J</i> = 6.8 Hz, 3H), 0.87 (s, 3H).	524.2	1.740
C671	 (mixture)	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.01 (d, <i>J</i> = 8.5 Hz, 1H), 7.94 (d, <i>J</i> = 12.4 Hz, 1H), 7.31 (s, 1H), 7.24 (d, <i>J</i> = 6.8 Hz, 1H), 6.55 (s, 1H), 5.59 (d, <i>J</i> = 6.3 Hz, 1H), 5.43 (s, 2H), 5.32 (dd, <i>J</i> = 5.5, 4.2 Hz, 1H), 4.64 (t, <i>J</i> = 7.3 Hz, 1H), 3.97 (d, <i>J</i> = 1.1 Hz, 3H), 3.19 (d, <i>J</i> = 6.2 Hz, 3H), 2.05 – 1.96 (m, 2H), 1.94 – 1.78 (m, 2H), 1.46 (d, <i>J</i> = 7.7 Hz, 1H), 0.86 (d, <i>J</i> = 8.2 Hz, 3H).	578.1	1.736

Conditions of HPLC above: Equipment: Agilent 1200; Chromatographic column: Waters XBridge C18 4.6*50mm,3.5um; Flow:2.0mL/min; Gradient elute: 5.0%-95.0%-95.0%-5.0%-5.0%, 0.00min-1.50min-

2.50min-2.52min-3.00min; Temperature : 40°C; Mobile phase : A: Acetonitrile, B: H₂O (0.05% TFA); Wavelength: 214 nm/254 nm.

Procedure of C589







Step K: (S)-N-((1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)-3-hydroxybutanamide: To a solution of (1S,9S)-1-amino-9-ethyl-5-fluoro-9-hydroxy-4-methyl-1,2,3,9,12,15-hexahydro-10H,13H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione mesylate (50 mg, 0.10 mmol, CAS:169869-90-3) in 2 mL DMF was added (S)-3-hydroxybutanoic acid (15.6 mg, 0.15 mmol), HATU (57.9 mg, 0.15 mmol) and followed with Et₃N (20.5 mg, 0.20 mmol) dropwise at 0 °C, the mixture was stirred at 0 °C to room temperature for 2 h. The mixture was poured into 10 mL ice-water, extracted with ethyl acetate (20 mL x 3), washed with brine (20 mL x 2), dried over Na₂SO₄ and concentrated. The residue was purified by prep-HPLC to give (S)-N-((1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)-3-hydroxybutanamide (9.4 mg) as a yellow solid.

The compounds prepared in the table **IV** below was reacted and post-treated in the same manner as in Example **C589**.

Table IV.

Number	Structure	¹ H NMR	MS [M+H] ⁺	Retention time on HPLC (min)
C589		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.44 (d, <i>J</i> = 8.8 Hz, 1H), 7.80 (d, <i>J</i> = 11.0 Hz, 1H), 7.31 (s, 1H), 6.53 (s, 1H), 5.71 – 5.51 (m, 1H), 5.43 (s, 2H), 5.31 – 5.11 (m, 2H), 4.67 (s, 1H), 4.05 (dd, <i>J</i> = 12.5, 6.3 Hz, 1H), 3.23 – 3.09 (m, 2H), 2.41 (s, 3H), 2.36 – 2.26 (m, 1H), 2.26 – 2.18 (m, 1H), 2.20 – 2.08 (m, 2H), 1.96 – 1.77 (m, 2H), 1.09 (d, <i>J</i> = 6.2 Hz, 3H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H).	522.1	1.643


C588		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.43 (d, <i>J</i> = 8.7 Hz, 1H), 7.80 (d, <i>J</i> = 11.0 Hz, 1H), 7.31 (s, 1H), 6.53 (s, 1H), 5.83 – 5.49 (m, 1H), 5.43 (s, 2H), 5.32 – 5.12 (m, 2H), 4.64 (s, 1H), 4.05 (dd, <i>J</i> = 12.6, 6.1 Hz, 1H), 3.18 (t, <i>J</i> = 6.2 Hz, 2H), 2.41 (s, 3H), 2.33 – 2.24 (m, 1H), 2.23 – 2.05 (m, 3H), 1.96 – 1.75 (m, 2H), 1.09 (d, <i>J</i> = 6.2 Hz, 3H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H).	522.1	1.629
C504		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.51 (d, <i>J</i> = 8.9 Hz, 1H), 7.79 (d, <i>J</i> = 10.9 Hz, 1H), 7.31 (s, 1H), 6.52 (s, 1H), 5.68 – 5.58 (m, 1H), 5.43 (s, 2H), 5.27 – 5.15 (m, 2H), 4.05 (s, 2H), 3.64 – 3.52 (m, 2H), 3.49 – 3.47 (m, 2H), 3.28 – 3.10 (m, 2H), 2.40 (s, 3H), 2.22 – 2.11 (m, 2H), 1.94 – 1.80 (m, 2H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H).	538.5	1.429
C562		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.49 (d, <i>J</i> = 9.0 Hz, 1H), 7.79 (d, <i>J</i> = 10.9 Hz, 1H), 7.31 (s, 1H), 6.52 (s, 1H), 5.82 – 5.54 (m, 1H), 5.42 (s, 2H), 5.31 – 5.09 (m, 2H), 4.67 (s, 1H), 3.59 – 3.45 (m, 2H), 3.43 – 3.33 (m, 2H), 3.25 – 3.09 (m, 2H), 2.40 (s, 3H), 2.23 – 2.07 (m, 2H), 2.03 – 1.62 (m, 2H), 1.39 – 1.07 (m, 1H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H), 0.72 – 0.31 (m, 4H).	578.2	1.581
C563		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.50 (d, <i>J</i> = 8.5 Hz, 1H), 7.80 (d, <i>J</i> = 10.9 Hz, 1H), 7.32 (s, 1H), 6.52 (s, 1H), 5.56 (d, <i>J</i> = 6.7 Hz, 1H), 5.43 (s, 2H), 5.35 – 5.12 (m, 2H), 4.55 (s, 1H), 3.56 – 3.42 (m, 2H), 3.42 – 3.35 (m, 2H), 3.24 – 3.11 (m, 2H), 2.40 (s, 3H), 2.27 – 2.08 (m, 2H), 1.98 – 1.82 (m, 2H), 1.20 – 1.05 (m, 1H), 0.89 (t, <i>J</i> = 7.3 Hz, 3H), 0.58 – 0.46 (m, 2H), 0.47 – 0.35 (m, 2H).	578.2	1.628

C565		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.82 (d, <i>J</i> = 8.8 Hz, 1H), 7.78 (d, <i>J</i> = 10.9 Hz, 1H), 7.31 (s, 1H), 6.52 (s, 1H), 5.67 – 5.52 (m, 1H), 5.43 (s, 2H), 5.28 – 5.10 (m, 2H), 4.95 (s, 1H), 3.59 – 3.48 (m, 2H), 3.50 – 3.41 (m, 2H), 3.29 – 3.03 (m, 2H), 2.40 (s, 3H), 2.28 – 2.17 (m, 2H), 2.02 – 1.73 (m, 2H), 1.30 – 1.19 (m, 2H), 1.19 – 1.02 (m, 2H), 0.88 (t, <i>J</i> = 7.3 Hz, 3H).	564.1	1.624
C687		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.85 (d, <i>J</i> = 8.7 Hz, 1H), 7.78 (d, <i>J</i> = 10.9 Hz, 1H), 7.30 (s, 1H), 5.66 – 5.49 (m, 1H), 5.41 (s, 2H), 5.31 – 4.91 (m, 2H), 3.23 – 3.06 (m, 2H), 2.39 (s, 3H), 2.26 – 2.07 (m, 2H), 1.95 – 1.75 (m, 2H), 1.54 (s, 3H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	576.1	1.719
C688		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.95 (d, <i>J</i> = 9.1 Hz, 1H), 7.77 (d, <i>J</i> = 10.9 Hz, 1H), 7.31 (s, 1H), 6.52 (s, 1H), 5.68 – 5.51 (m, 1H), 5.42 (s, 2H), 5.29 – 4.93 (m, 2H), 3.28 – 3.06 (m, 2H), 2.38 (s, 3H), 2.24 – 2.08 (m, 2H), 1.97 – 1.79 (m, 2H), 1.71 (s, 3H), 0.87 (t, <i>J</i> = 7.3 Hz, 3H).	576.1	1.806

Conditions of HPLC above: Equipment: Agilent 1200; Chromatographic column: Waters XBridge C18 4.6*50mm,3.5um; Flow:2.0mL/min; Gradient elute: 5.0%-95.0%-95.0%-5.0%-5.0%, 0.00min-1.50min-2.50min-2.52min-3.00min; Temperature : 40°C; Mobile phase : A: Acetonitrile, B: H₂O (0.05% TFA); Wavelength: 214 nm/254 nm.


Example 2 Preparation of linkers

[0167] 2.1 Preparation of linker HX20113



[0168] HX20113 synthesized by a conventional solid phase polypeptide synthesis using Rink-amide-MBHA-resin. Fmoc was used to protect the amino acid in the linking unit. The conjugation reagent was selected from HOBt, HOAt/DIC, DCC, EDCI or HATU. After synthesis, the resin was cleaved using trifluoroacetic acid. The product was purified by HPLC, lyophilized and stored for use. Theoretical Mass: 1207.59, measured: [M-H]⁻ = 1206.7.

[0169] Preparation of linker HX20111




[0170] HX20111 synthesized by a conventional solid phase polypeptide synthesis using Rink-amide-MBHA-resin. Fmoc was used to protect the amino acid in the linking unit. The conjugation reagent was selected from HOBt, HOAt/DIC, DCC, EDCI or HATU. After synthesis, the resin was cleaved using trifluoroacetic acid. The product was purified by HPLC, lyophilized and stored for use. Theoretical Mass: 1383.70, measured: $[M-H]^+ = 1382.6$.

Example 3 Preparation of linker-payload intermediates

[0171] 3.1 Preparation of intermediate Mc-GGFG-Dxd

[0172] The intermediate Mc-GGFG-Dxd is commercial available or prepared following the procedures as described in EP2907824. This compound is used to prepare the Linker-Payload intermediate (formula (I) compound), and is also used to directly connect to the (optionally modified) antibody to prepare reference ADCs LC1184(8) and LC1184(4).

[0173] 3.2 Preparation of H0019



[0174] Synthesis of HX21008-a

[0175] 4.33 g Fmoc-Gly-Gly-OH and 6.84 g Pb(OAc)₄ were weighed and added into a 500 ml single-neck round bottom flask. Anhydrous THF/Toluene (120/40 ml) was added under nitrogen atmosphere and stirred for dissolving. Then 1.16 mL of pyridine was added to the reaction system. The reaction system was heated to 80°C and refluxed for 5hr under nitrogen atmosphere. Samples were taken and detected by HPLC to monitor the reaction.

[0176] The reaction system was cooled to room temperature, filtered, and the filter cake was washed with EA for 3 times. The filtrates were combined and concentrated to dryness. Column chromatography was performed (PE: EA = 100: 0 ~ 50: 100) to give about 2000 mg of the target product in white solid with a yield of 44%.

[0177] Synthesis of H0005-a

[0178] 200 mg HX21008-a was weighed and added into a 100 ml single-neck round bottom flask. Then 15 ml THF was added and stirred for dissolving. Then H0005 (316 mg, 3.0 eq) and TsOH·H₂O (15 mg, 0.15 eq) were added to the reaction system. The reaction system was reacted overnight at room temperature. Samples were taken and detected by TLC (PE/EA=1: 1) to monitor the reaction. The raw material basically disappeared, and a new point was detected.

[0179] Saturated sodium bicarbonate solution was added to quench reaction. Extraction was conducted with EA for 3 times. The organic phase was combined and washed with saline, dried

with anhydrous magnesium sulfate and concentrated. The crude product was purified by column chromatography (PE: EA = 5: 1 ~ 1: 1) to give about 165 mg of the target product in colorless oil with a yield of 60%. MS: $[M+H]^+ = 503.4$.

[0180] Synthesis of H0005-b

[0181] 200 mg of H0005-a was weighed and added into a 100 ml single-neck round bottom flask. Then 10 ml of EtOH and 5 ml of EA were added with complete dissolution. Then 40 mg of palladium carbon was added to the reaction system under nitrogen atmosphere, and the reaction system was purged with hydrogen gas for three times. The reaction system was kept under hydrogen atmosphere and stirred for 0.5 hr at room temperature. Samples were taken and detected by TLC (DCM/MeOH=10: 1) to monitor the reaction. The raw material basically disappeared, and a new point was detected.

[0182] The reaction system was filtered, and the filter cake was washed with EA for 3 times. The filtrates were combined and concentrated to dryness to give 200 mg product in white solid with 100% yield. The product can be directly used in the next reaction without purification. $[M+H]^+ = 413.3$.

[0183] Synthesis of H0005

[0184] 2.0 g of dichlororesin was weighed and placed in a polypeptide synthesis tube. DCM (10 ml) was added and swelled at room temperature for 30 minutes. The solvent was removed by vacuum suction. The resin was washed twice with DCM, with a volume of 7 mL and a time length of 1 minute for each wash. The solvent was removed by vacuum suction. Then H0005-b (200 mg) was weighed and added into a 50 ml centrifuge tube. DCM (about 10 ml) was added. the solid was dissolved by shaking. Added to the above resin. Stirring was conducted to soak all the resin in the solution (if there was resin attached to the tube wall, a small amount of DCM was used to wash the tube wall). Stirring was conducted for 4-5 hours. After the reaction was complete, an appropriate amount of methanol was added. Stirring was conducted for 30 min. The solvent was removed by vacuum suction. The resin was washed with DMF once, methanol once, DMF once, methanol once and DMF twice in sequence, with a volume of 10 mL and a time length of 1 minute for each wash. The solvent was removed by vacuum suction. A small amount of dry resin was taken for ninhydrin detection. The resin was colorless and transparent, and the solution was yellowish, indicating qualified for the next coupling step.

[0185] The deprotection was conducted twice by adding 10 mL readymade 20% piperidine/DMF solution and reacting for 10 minutes for each time. After the reaction was complete, the solution was removed by vacuum suction. The resin was washed with DMF twice, methanol once, DMF once, methanol once and DMF twice in sequence, with a volume of 10 mL and a time length of 1 minute for each wash. The solvent was removed by vacuum suction.

A small amount of dry resin was taken for ninhydrin detection. Both the resin and solution were dark blue.

[0186] To a 50 mL centrifuge tube was added 563 mg Fmoc-Phe-OH, 197 mg HOBt. Then about 7 mL DMF was added. The solid was dissolved by shaking. Then 0.24 mL DIC was added. Activated for 10-30 minutes to give the activated reaction solution.

[0187] 3 molar equivalent of activated reaction solution added to the resin. Stirring was conducted to soak the resin completely in the solution (if there was resin attached to the tube wall, a small amount of DCM was used to wash the tube wall). Stirring was conducted for 2-3 hours. After the reaction was complete, the solvent was removed by vacuum suction. The resin was washed with DMF twice, methanol once, DMF once, methanol once and DMF twice in sequence, with a volume of 10 mL and a time length of 1 minute for each wash. The solvent was removed by vacuum suction. A small amount of dry resin was taken for ninhydrin detection. The resin was colorless and transparent, and the solution was yellowish, indicating qualified for the next coupling step.

[0188] The deprotection was conducted twice by adding 10 mL readymade 20% piperidine/DMF solution and reacting for 10 minutes for each time. After the reaction was complete, the solution was removed by vacuum suction. The resin was washed with DMF twice, methanol once, DMF once, methanol once and DMF twice in sequence, with a volume of 10 mL and a time length of 1 minute for each wash. The solvent was removed by vacuum suction. A small amount of dry resin was taken for ninhydrin detection. Both the resin and solution were dark blue.

[0189] To a 50 mL centrifuge tube was added 531 mg Fmoc-GG-OH, 197mg HOBt. Then about 10 mL DMF was added. The solid was dissolved by shaking. Then 0.24 mL DIC was added. Activated for 10-30 minutes to give the activated reaction solution.

[0190] 3 molar equivalent of activated reaction solution was added to the resin. Stirring was conducted to soak the resin completely in the solution (if there was resin attached to the tube wall, a small amount of DCM was used to wash the tube wall). Stirring was conducted for 2-3 hours. After the reaction was complete, the reaction solution was removed by vacuum suction. The resin was washed with DMF twice, methanol once, DMF once, methanol once and DMF twice in sequence, with a volume of 10 mL and a time length of 1 minute for each wash. The solvent was removed by vacuum suction. A small amount of dry resin was taken for ninhydrin detection. The resin was colorless and transparent, and the solution was yellowish, indicating qualified for the next coupling step.

[0191] The deprotection was conducted twice by adding 10 mL readymade 20% piperidine/DMF solution and reacting for 10 minutes for each time. After the reaction was complete, the solution was removed by vacuum suction. The resin was washed with DMF twice,

methanol once, DMF once, methanol once and DMF twice in sequence, with a volume of 10 mL and a time length of 1 minute for each wash. The solvent was removed by vacuum suction. A small amount of dry resin was taken for ninhydrin detection. Both the resin and solution were dark blue. Then, 462 mg MC-OSu was placed in a 50 mL centrifuge tube, about 10 mL DMF was added. The solid was dissolved by shaking. Then 0.24 mL DIEA was added to the resin. Stirring was conducted to soak the resin completely in the solution (if there was resin attached to the tube wall, a small amount of DCM was used to wash the tube wall). Stirring was conducted for 2-3 hours. After the reaction was complete, the reaction solution was removed by vacuum suction. The resin was washed with DMF twice, methanol once, DMF once, methanol once and DMF twice in sequence, with a volume of 10 mL and a time length of 1 minute for each wash. The solvent was removed by vacuum suction. A small amount of dry resin was taken for ninhydrin detection. The resin was colorless and transparent, and the solution was yellowish, indicating qualified for the next coupling step.

[0192] The resin was washed twice with 10 mL of methanol. Then the solvent was removed thoroughly by vacuum suction. The resin was poured out and weighed. The lysis buffer was prepared in a 250 mL conical flask, wherein: the ratio of TFE/DCM was 80%/20%, and the volume was 7-8 times of the weight of peptide resin. The lysis buffer was added into the peptide resin, shaken well. The resin was fully soaked in the lysis buffer, and lysis was carried out at room temperature for 2-3 hours. The lysis buffer was then filtered out using a simple filter made of a syringe, and the resin was washed with 1-2 mL DCM and discarded. Then 150 mL precooled anhydrous ether was added to the lysis buffer, shaken well and then stood for 20-30 minutes. Using a 50 mL centrifuge tube, the above system was centrifuged in a centrifuge at 3500 rpm for 3 minutes, and the supernatant was poured out and discarded. The solid was shaken with precooled anhydrous ether, washed once under ultrasound, centrifuged at 3500rpm for 3 minutes, and the supernatant was poured out and discarded. The solid was placed in a centrifuge tube and allowed to air dry overnight, and then subjected to preparative purification to give 125 mg of product in white solid with a yield of 40%. $[M+H]^+ = 645.4$.

[0193] Synthesis of H0013

[0194] 150 mg of raw material H0005 and 55 mg of TSTU were weighed and added into a 10 mL single-neck round bottom flask, and anhydrous DMF (3 mL) was added under nitrogen atmosphere and stirred for 20 min. Then 18 mg TSN00643 and 20 μ L DIEA were added in sequence to the reaction system. Stirring was conducted at room temperature for 2 hr under nitrogen atmosphere. Samples were taken and detected by HPLC to monitor the reaction. The raw material peak completely disappeared, and new peaks were detected.

[0195] The reaction system was subjected to preparative purification, and the target product was collected and lyophilized to give about 22mg of product in yellowish solid. $[M+H]^+ = 1062.7$.


[0196] Synthesis of H0019




[0197] H0013 (30mg) was weighed and added into a 10 ml single-neck round bottom flask, purified water (2ml) was added. Stirring was conducted for dissolving. DMF solution (2 ml) containing HX20111 (19.5 mg) was added to the reaction system and stirred. After reacting overnight, HPLC was used to monitor the reaction until all of the raw material had converted into intermediates. Then the reaction system was cooled to $0 \sim 10^\circ\text{C}$, and 0.5% LiOH solution was added to the reaction system under controlled temperature until the pH of the reaction system was about 12. The reaction was stirred for 20 min, and the reaction was monitored by HPLC until all the intermediates were consumed and then 30% acetic acid (1.5 mL) was added to the reaction system to quench the reaction.


[0198] The reaction system was subjected to preparative purification, and the target product was collected and lyophilized to give about 25mg of product in yellowish solid. $[(M+3H)/3]^+ = 1182.3$.

[0199] 3.3 H0020, H0021, H0034 and H0035

[0200] The following linker-payload intermediates H0020, H0021, H0034 and H0035 can be prepared using similar synthetic routes and reagents. The intermediates H0014, H0015, H0026 and H0032 are synthesized using similar method for H0013, and then used for synthesizing H0020, H0021, H0034 and H0035, respectively.








[0201] The LC-MS data is shown in the following table:

	MS data
H0020	$[(M+3H)/3]^+ = 1193.9$
H0021	$[(M+3H)/3]^+ = 1177.9$


[0202] 3.4 Preparation of LB302-2-4



[0203] HX20111 and intermediate MC-GGFG-Dxd (molar ratio ~1:2) were weighed and dissolved in water and DMF, respectively, and then thoroughly mixed to give a mixture, which was reacted at 0-40°C for 0.5-30h. Once the reaction was completed, the reaction mixture was directly added with an appropriate amount of Tris Base solution or other solution that promotes the ring-opening reaction, and the reaction was performed at 0-40°C for another 0.2-20h. After the reaction was completed, the product was purified by semi-preparative/preparative HPLC and lyophilized to obtain linker-payload intermediate LB302-2-4. Theoretical Mass: 3486.52, measured: $[(M+3H)/3]^+ = 1163.3$.

[0204] 3.5 Preparation of LB302-2-1

[0205] LB302-2-1 can be prepared using similar synthetic routes and reagents.



Example 4 Construction of antibody expression vector, antibody expression, purification and identification

[0206] 4.1 Production of the modified anti-human HER2 antibody Ab0001-LCCT_L-HC

[0207] The expression plasmids for antibody Ab0001-LCCT_L-HC (light chain SEQ ID NO: 9, heavy chain: SEQ ID NO: 10) were constructed as follows. The sequence of the antibody Ab0001-LCCT_L-HC: based on the amino acid sequence of Trastuzumab, and GALPETGG was introduced at the C-terminal of the light chain, wherein LPETGG is the recognition sequence of the ligase donor substrate, and GA is a spacer sequence. The plasmids were transfected into CHO cells and the cell population was established and screened for a highly expressed cell population, which was cultured with reference to the culture process of Trastuzumab in a 5-10L reactor, and supernatant was collected.

[0208] 4.2 The purification of antibody Ab0001-LCCT_L-HC

[0209] The purification of Ab0001-LCCT_L-HC was carried out in a standard process using the combination of MabSelect affinity chromatography and Sepharose S cation exchange chromatography, the purified products were dissolved in the original Trastuzumab drug buffer (5mM histidine-HCl, 2% Trehalose, 0.009% Polysorbate 20, PH 6.0), and frozen in small aliquots.

[0210] 4.3 The quality control of antibody Ab0001-LCCT_L-HC

[0211] The purity of the above purified antibody Ab0001-LCCT_L-HC is 98.5% by SDS-PAGE; the content of high molecular weight polymer of the sample is less than 0.4% by SEC-HPLC; endotoxin content is less than 0.098 EU/mg.

[0212] 4.4 Preparation of other modified anti-human antibodies

[0213] According to a similar method, a terminal modification based on the ligase recognition sequence was introduced at the C-terminal of the light and/or heavy chain of the Trastuzumab, respectively, giving a modified antibody.

[0214] The modified anti-human HER2 antibodies based on Ab0001 (Trastuzumab) are listed in the following table. LPETGG in the terminal modification sequence is a recognition sequence of the ligase donor substrate, and GA is a spacer sequence.

Modified anti-human HER2 antibodies

	Sequence	Sequence introduced at the terminal
Ab0001-LCCT _L -HC light chain	SEQ ID NO: 9	GALPETGG
Ab0001-LCCT _L -HC heavy chain	SEQ ID NO: 10	_*

*: “-” indicates no terminal modification

Example 5 Preparation of antibody-small molecule Conjugates

[0215] 5.1 The Linker-payload intermediates were respectively conjugated to an antibody in a site-specific manner by a ligase to form an ADC. The method for conjugation reaction can be found in WO2015165413A1. The resulting ADCs are as listed in the following table:

Name of ADC	Linker-Payload	Antibody	DAR
CH-2-589	BH-2-589	Ab0001-LCCT _L -HC	3.44
CH-2-593	BH-2-593	Ab0001-LCCT _L -HC	3.49
CH-2-518	BH-2-518	Ab0001-LCCT _L -HC	3.61
LC302-2-1	LB302-2-1	Ab0001-LCCT _L -HC	3.42
LC302-2-4(4)	LB302-2-4	Ab0001-LCCT _L -HC	3.41

[0216] 5.2 The reference ADCs LC1184(8) and LC1184(4) are prepared respectively by directly connecting the intermediate Mc-GGFG-Dxd to Ab0001-LCCT_L-HC (Cys conjugation, i.e. conjugation through connections formed by maleimide structure(s) with thiol group(s) of Cys). The method for conjugation reaction is known in the art. LC1184(8) has eight Mc-GGFG-Dxd introduced to the reduced inter-chain cysteines. LC1184(4) has four Mc-GGFG-Dxd introduced to the reduced inter-chain cysteines.

Effect Example 1 Effect of small molecules on cell proliferation

[0217] **Study Purpose:** In this experiment, cell viability was detected using CellTiter-Glo to evaluate the effects of the small molecule compounds on the proliferation of a high HER2 tumor cell line SK-BR-3, and to analyze the effects of the small molecule compounds on the proliferation of tumor cells.

[0218] **Materials and Equipment**

[0219] PBS: Gibco, Cat# 10010-023); Trypsin: Gibco, Cat# 25200056); FBS: Gibco, Cat# 10270-106); CellTiter-Glo® Luminescent Cell Viability Assay: Promega, Cat# G7573); McCoy's 5A: Gibco, Cat# 16600-082); L15: Hyclone, Cat# SH30525.01); DMEM: Gibco, Cat# 11995-065). 96-well plate: Corning/3603; 96-well deep-well Plate: Thermo/278743; 15 ml EP tube: Thermo/339650; 1.5 ml EP tube: Beaver.

[0220] **Cell plating -Day 1**

[0221] (1) Microscopic examination of cells

[0222] (2) Digestion: digested using 2mL 0.25% trypsin for 3 minutes;

[0223] (3) Centrifuge: 1000rpm for 5min;

[0224] (4) Cell counting

[0225] A. Cell dilution:

[0226] B. Cell plating: in 100 µl/well

[0227] C. Incubation: 37 °C, 5% CO₂, overnight

[0228] **Cell test**

[0229] (1) Microscopic examination

[0230] (2) Drug preparation:

[0231] A. Preparation of Buffer for preparation of samples: the desired amount of Buffer for preparation of samples was prepared using the culture medium (10% FBS) for the testing cells;

[0232] B. Sample preparation: the sample was diluted from the first concentration to the desired concentration in a 96-well deep-well Plate.

[0233] Test compounds: Ten final concentrations are prepared from dilution with the cell medium: 1000, 200, 40, 8, 1.6, 0.32, 0.064, 0.0128, 0.00256 and 0.000512 nM.

[0234] Positive control Puromycin: A stock solution of 2.5 mg/ml was used, and diluted with the cell medium. A dilution of 250-fold was preformed, and then a dilution of 2-fold was preformed when applying the compounds to reach the testing concentration 5 µg/ml for puromycin.

[0235] (3) Application of the compounds: for each concentration of the diluted series, the compound was applied in triplicate. In 100µl/well.

[0236] (4) After the drug was added, the cells were incubated in the incubator for 120h.

[0237] **Cell viability test**

[0238] (1) Preparation of detection reagents: CellTiter-Glo Luminescent Cell Viability Assay detection reagent was allowed to warm to room temperature with protection from light

[0239] (2) Cell preparation: The cells to be tested were taken out from the incubator and equilibrated at room temperature (25 °C) 30min.

[0240] (3) ATP detection: Discard the culture medium and 100 μ l/well DMEM was added, 50 μ l/hole CTG was added into 96-well plate, protected from light using aluminum foil and oscillated using a vortex oscillator at room temperature of 200 rpm for 10 min.

[0241] (4) Detection procedure: After setting up the procedure, the black-wall clear-bottom plate was placed in the instrument without its cover.

[0242] (5) Data acquisition.

[0243] 5. Data processing and analysis

[0244] GraphPad software was used to process the data. The cell proliferation inhibition rate was plotted against the concentration of the control sample or the test sample using four-parameter curve model for data-fitting, and the R^2 (rounded to 4 significant digits) and IC_{50} (rounded to 4 significant digits), or EC_{50} of the control sample and the test sample were obtained respectively. If necessary, the relative biological activity of sample A, i.e., ratio activity was calculated. Dxd was used as the control sample for the ratio activity calculation.

[0245] Cell proliferation inhibition rate (%) = (luminescence value of drug-treated group - luminescence value of puro group)/(luminescence value of blank control group - luminescence value of puro group) $\times$ 100; wherein the puro group refers to the group treated with puromycin

[0246] Ratio activity of sample (%) = (IC_{50} value of the control sample/ IC_{50} value of the test sample) $\times$ 100

[0247] Ratio activity of control sample (%) = (IC_{50} value of the control sample/ IC_{50} value of the previous control sample) $\times$ 100

[0248] 6. Result verification

[0249] R^2 Value $\geq$ 0.950; CV% between parallel pores $\leq$ 30%.

[0250] **Results:**

[0251] The results are listed in the following table.

Test compound	Ratio activity (%)	Test compound	Ratio activity (%)	Test compound	Ratio activity (%)
C730	1.35	C703	0.846	C667	0.658
C731	0.82	C704	2.271	C668	0.231
C518	2.272	C636	0.682	C671	0.365
C519	1.059	C637	1.585	C588	1.019
C586	0.259	C595	0.832	C589	1.325
C587	0.285	C596	0.911	C504	0.703
C593	2.514	C597	0.684	C562	1.029
C594	0.568	C600	0.62	C563	1.284
C622	0.669	C601	1.826	C565	1.381
C624	0.916	C602	0.29	C687	1.835
C625	0.783	C603	1.439	C688	0.744
C677	1.008	C619	0.434	C716	2.305
C678	0.277	C620	0.384	C626	0.54
C679	0.49	C664	0.269	C627	5.054
C680	0.546	C665	0.495	Dxd	1

Effect Example 2 Effect of conjugates targeting HER2 on cell proliferation

[0252] **Study Purpose:** Cell viability was detected by CellTiter-Glo assay to evaluate the effects of the ADCs on the proliferation of two high HER2 tumor cell lines SK-BR-3, NCI-N87 and a HER2-negative tumor cell line MDA-MB-468, and to analyze the effects of the ADCs on the proliferation of tumor cells.

[0253] Materials and Equipment

[0254] PBS: Gibco, Cat# 10010-023); Trypsin: Gibco, Cat# 25200056); FBS: Gibco, Cat# 10270-106); CellTiter-Glo® Luminescent Cell Viability Assay: Promega, Cat# G7573); McCoy's 5A: Gibco, Cat# 16600-082); L15: Hyclone, Cat# SH30525.01); DMEM: Gibco, Cat# 11995-065). 96-well plate: Corning/3603; 96-well deep-well Plate: Thermo/278743; 15 ml EP tube: Thermo/339650; 1.5 ml EP tube: Beaver.

[0255] Cell plating -Day 1

[0256] (1) Microscopic examination of cells

[0257] (2) Digestion: digested using 2mL 0.25% trypsin for 3 minutes;

[0258] (3) Centrifuge: 1000rpm for 5min;

[0259] (4) Cell counting

[0260] A. Cell dilution:

[0261] B. Cell plating: in 100 µl/well

[0262] C. Incubation: 37 °C, 5% CO₂, overnight

[0263] Cell test

[0264] (1) Microscopic examination

[0265] (2) Drug preparation:

[0266] A. Preparation of Buffer for preparation of samples: the desired amount of Buffer for preparation of samples was prepared using the culture medium (10% FBS) for the testing cells;

[0267] B. Sample preparation: the sample was diluted from the first concentration to the desired concentration in a 96-well deep-well Plate.

[0268] Test samples: Ten final concentrations are prepared from dilution with the cell medium: 200, 40, 8, 1.6, 0.32, 0.064, 0.0128, 0.00256, 0.000512 and 0.0001024 nM.

[0269] Positive control Puromycin: A stock solution of 10 mg/ml was used, and diluted with the cell medium. A dilution of 100-fold was preformed, and then a dilution of 10-fold was preformed when applying the compounds to reach the testing concentration 5 µg/ml for puromycin.

[0270] (4) Application of the compounds: for each concentration of the diluted series, the compound was applied in triplicate. In 100µl/well.

[0271] (5) After the drug was added, the cells were incubated in the incubator for 120h.

[0272] **Cell viability detection**

[0273] (1) Preparation of detection reagents: CellTiter-Glo Luminescent Cell Viability Assay detection reagent was allowed to warm to room temperature with protection from light

[0274] (2) Cell preparation: The cells to be tested were taken out from the incubator and equilibrated at room temperature (25 °C) 30min.

[0275] (3) ATP detection: Discard the culture medium and 100 µl/well DMEM was added, 50 µl/hole CTG was added into 96-well plate, protected from light using aluminum foil and oscillated using a vortex oscillator at room temperature of 200 rpm for 10 min.

[0276] (4) Detection procedure: After setting up the procedure, the black-wall clear-bottom plate was placed in the instrument without its cover.

[0277] (5) Data acquisition.

[0278] 5. Data processing and analysis

[0279] GraphPad software was used to process the data. The cell proliferation inhibition rate was plotted against the concentration of the control sample or the test sample using four-parameter curve model for data-fitting, and the R^2 (rounded to 4 significant digits) and IC50 (rounded to 4 significant digits), or EC50 of the control sample and the test sample were obtained respectively.

[0280] Cell proliferation inhibition rate (%) = (luminescence value of drug-treated group - luminescence value of puro group)/(luminescence value of blank control group - luminescence value of puro group) × 100; wherein the puro group refers to the group treated with puromycin

[0281] 6. Result verification

[0282] R^2 Value ≥ 0.950 ; CV% between parallel pores $\leq 30\%$.

[0283] **Results:**

[0284] The results are shown in the following table and Figure 1, Figure 2 and Figure 3.

Samples	IC ₅₀ (nM)		
	SK-BR-3	NCI-N87	MDA-MB-468
CH-2-589	0.1737	0.4363	N/A
CH-2-593	~0.2679	0.3431	N/A
CH-2-518	0.1621	0.2888	N/A
LC302-2-1(4)	0.1926	0.6353	N/A
LC302-2-4(4)	0.2208	0.7226	N/A
Dxd	1.472	4.682	N/A

[0285] In HER2 high-expressing SK-BR-3 and NCI-N87, the conjugates (CH-2-589, CH-2-593, CH-2-518, LC302-2-1(4) and LC302-2-4(4)) exhibited outstanding efficacy. The IC₅₀ values of the conjugates are lower than the small molecule payload Dxd. In HER2 negative cells, the conjugates exhibited minimal efficacy. No cytotoxicity was observed in HER2 negative cell MDA-MB-468, demonstrating the stability of the conjugates, with reduced off-target release of the payload and thereby good safety. These indicate that the conjugates exerts HER2-dependent cytotoxic activity in a dose dependent manner, exhibiting good targeting ability, outstanding efficacy and good safety.

Effect Example 3 Assessment of bystander killing effect

[0286] **Purpose of experiment** The Bystander killing activities of ADCs (CH-2-589, CH-2-593, CH-2-518 and LC302-2-4(4)) on MDA-MB-468 cells were tested in a co-culture system of a HER2-positive tumor cell line SK-BR-3 and a HER2-negative tumor cell line MDA-MB-468, and compared the bystander killing effect with a reference ADC (LC1184 (8)) with a structure similar to DS8201a.

[0287] **Materials and Equipment**

[0288] L15: Hyclone, SH30525.01; McCoy's 5A: Gibco, 16600-082; FBS: Gibco, 10099-141; Firefly luciferase reporter gene assay kit: Beyotime, RG006; 50ml Centrifugal tube: corning, 430829; 15ml Centrifugal tube: corning, 430791; 96-well plate: corning, 3599; CO₂ Cell incubator: Thermo; Centrifuge: Thermos, L550; Microscope: Nikon, ISZ; Cell counter: Countstar, IC1000; Cytation3 Plate reader: BioTek, Cytation3; HTX Multi-mode microplate reader: BioTek, Synergy HTX; SK-BR-3: ATCC, HTB-30; MDA-MB-468-Luc-GFP: prepared in-house using lentivirus infection.

[0289] **Experimental method**

[0290] 1) Cell collection and counting.

[0291] 2) The cell density was adjusted to 1×10^5 cells/ml

[0292] 3) The cell mixture was prepared, wherein SKBR3: MDA-MB-468-Luc-GFP=4: 1, 5ml.

[0293] Cell plating: 1×10^4 cells/ well, 100 μ l/well

[0294] Test samples: Three final concentrations are prepared from dilution with the cell medium: 50, 10 and 1 nM.

[0295] 4) The drug was administered the next day, 100 μ l/well.

[0296] 5) Incubate in a 5% CO₂ incubator at 37 °C for 120h.

[0297] ① washed by PBS to remove dead cells, and detected by Cytation3 plate reader;

[0298] ② Biyuntian luciferase reporter gene assay kit: lysed in 37 °C incubator for 10min using 100 μ l lysis buffer. Centrifuged at 2000 rpm for 5 minutes. 100 μ l supernatant was taken. 100 μ l substrate was added. The chemiluminescence was detected by microplate reader (gain: 200).

[0299] **Result:**

[0300] Two methods (GFP fluorescence counting and luciferase substrate luminescence method) were used for determination of bystander killing effects (Figure 4a, 4b and Figure 5).

[0301] The conjugates (CH-2-593, CH-2-589, CH-2-518 and LC302-2-4(4)) all show outstanding bystander killing effects in the detections using both of the two methods. CH-2-593 shows a bystander killing effect comparable to or stronger than LC1184(8).

Effect Example 4 Assessment of conjugates *in vivo*

[0302] 4.1 On JIMT-1 human breast cancer xenograft model

[0303] 5×10^6 JIMT-1 human breast cancer cells (HER2 medium) were inoculated subcutaneously in the right flank in SCID Beige mice. After 7 days, when tumor volume reached 142mm³ on average, the tumor bearing mice were assigned and administrated intravenously of LC1184(8) and other five different conjugates at 5mg/kg. The tumor volume was measured twice weekly with calipers. LC1184(8) showed better efficacy than LC1184(4), suggesting that with same payload, higher DAR would result in better efficacy. LC302-2-1(4) and LC302-2-4(4) showed better efficacy than LC1184(8) (Figure 6). The conjugate of the present invention achieves great efficacy in lower DAR.

[0304] 4.2 On Capan-1 human pancreatic cancer xenograft model (1)

[0305] 5×10^6 Capan-1 human pancreatic cancer cells (HER2 low) were inoculated subcutaneously in the right flank in BALB/c nude mice to generate xenograft model. After 8 days, when tumor volume reached 178mm³ on average, the tumor bearing mice were administrated intravenously of LC1184(8) and other five different conjugates at 5mg/kg. The

tumor volume was measured twice weekly with calipers. LC1184(8) showed better efficacy than LC1184(4). LC302-2-1(4) and LC302-2-4(4) showed comparable efficacy to LC1184(8) (Figure 7).

[0306] 4.3 On Capan-1 human pancreatic cancer xenograft model (2)

[0307] **Study Purpose:** The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of CH-2-589, CH-2-593, CH-2-518 and LC302-2-4(4) in the treatment of the subcutaneous Capan-1 human pancreatic cancer xenograft model in female BALB/c Nude mice.

[0308] **Study Design:**

[0309] **Cell Culture:** The Capan-1 tumor cells (ATCC-HTB-79) will be maintained *in vitro* as a monolayer culture in IMDM supplemented with 20% fetal bovine serum, 1% Antibiotic-Antimycotic at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells will be routinely subcultured twice weekly by trypsin-EDTA treatment. The cells growing in an exponential growth phase will be harvested and counted for tumor inoculation.

[0310] **Animals:** BALB/c Nude, female, 6~8 weeks, weighing approximately 18~20g. A total of 48 (30 plus 60%) will be needed for the study, which will be purchased from Shanghai Lingchang Laboratory Animal Co., LTD. or other certified vendors.

[0311] **Tumor Inoculation:** Each mouse will be inoculated subcutaneously at the right flank with Capan-1 tumor cells (5×10^6) in 0.2 mL of PBS with Matrigel (1:1) for tumor development. The animals will be randomized and treatment will be started when the average tumor volume reaches approximately 150~200mm³ for the efficacy study. The test article administration and the animal numbers in each group are shown in the following experimental design table.

Groups and Treatments

Group	N ¹	Treatment	Dose (mg/kg)	Dosing Volume (mL/kg) ²	Dosing Route	Schedule ³
1	6	Vehicle	--	10	IV	Single dose
2	6	CH-2-589	5	10	IV	Single dose
3	6	CH-2-593	5	10	IV	Single dose
4	6	CH-2-518	5	10	IV	Single dose
5	6	LC302-2-4(4)	5	10	IV	Single dose

N: animal number;

Dosing volume: adjust dosing volume based on body weight 10 mL/kg.

The experiment duration will be adjusted according to tumor volume.

[0312] **Animal Housing:** An acclimation period of approximately one week will be allowed between animal arrival and tumor inoculation in order to accustom the animals to the laboratory environment. The mice will be maintained in a special pathogen-free environment and in individual ventilation cages (3 mice per cage). All cages, bedding, and water will be sterilized before use. When working in the mouse room, the investigators will wear lab coat and latex or

vinyl gloves. Each cage will be clearly labeled with a cage card indicating number of animals, sex, strain, date received, treatment, study number, group number, and the starting date of the treatment. The cages with food and water will be changed twice a week. The targeted conditions for animal room environment and photoperiod will be as follows:

[0313] **Temperature** 20~26 °C

[0314] **Humidity** 40~70 %

[0315] **Light cycle** 12 hours light and 12 hours dark

[0316] **Dietary Materials:** All animals will have free access to a standard certified commercial laboratory diet. Maximum allowable concentrations of contaminants in the diet are controlled and routinely analyzed by the manufacturers. Autoclaved municipal tap water, suitable for human consumption will be available to the animals *ad libitum*. It is considered that there are no known contaminants in the dietary materials that could influence the tumor growth.

[0317] **Assignment to Groups:** Before commencement of treatment, all animals will be weighed and the tumor volumes will be measured. Since the tumor volume can affect the effectiveness of any given treatment, mice will be assigned into groups using an Excel-based randomization software performing stratified randomization based upon their tumor volumes. This ensures that all the groups are comparable at the baseline.

[0318] **Observations:** The protocol and any amendment(s) or procedures involving the care and use of animals in this study will be reviewed and approved by the Institutional Animal Care and Use Committee (IACUC) of WuXi AppTec prior to conduct. During the study, the care and use of animals will be conducted in accordance with the regulations of the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC). After inoculation, the animals will be checked daily for morbidity and mortality. At the time of routine monitoring, the animals will be checked for any effects of tumor growth and treatments on normal behavior such as mobility, food and water consumption, body weight gain/loss (body weights will be measured twice weekly), eye/hair matting and any other abnormal effect. Death and observed clinical signs will be recorded on the basis of the numbers of animals within each subset.

[0319] **Endpoints:** The major endpoint is to see if the tumor growth can be delayed or mice can be cured. Tumor sizes will be measured twice weekly in two dimensions using a caliper, and the volume will be expressed in mm³ using the formula: $V = 0.5 a \times b^2$ where *a* and *b* are the long and short diameters of the tumor, respectively. The tumor sizes are then used for the calculations of both TGI(%) and the relative tumor proliferation rate T/C(%) values. TGI is calculated for each group using the formula: $TGI (\%) = [1 - (T_i - T_0) / (V_i - V_0)] \times 100$; *T_i* is the average tumor volume of a treatment group on a given day, *T₀* is the average tumor volume of the treatment group on the first day of treatment, *V_i* is the average tumor volume of the vehicle

control group on the same day with V_0 , and V_0 is the average tumor volume of the vehicle group on the first day of treatment.

[0320] The T/C(%) value is calculated for each group using the formula: $T/C \% = T_{RTV} / C_{RTV} \times 100 \%$ (T_{RTV} : relative average tumor volume (RTV) of the treatment group; C_{RTV} : relative average tumor volume (RTV) of the vehicle control group on the same day with T_{RTV}). The relative tumor volume (RTV) is calculated for each group using the formula: $RTV = V_t / V_0$; V_0 is the tumor volume on the first day of treatment, V_t is the tumor volume on a given day.

[0321] **Termination:**

[0322] **1) Bodyweight loss:** Any animal exhibiting 20% bodyweight loss at any one day will be humanely killed or the veterinary staff will be contacted.

[0323] **2) Tumor burden:** Tumor burden should not exceed 3,000 mm³. Individual animal will be sacrificed immediately when its tumor volume reaches 3,000 mm³.

Ulceration: If tumor ulceration occurs, the following procedures will apply:

- Animals with ulcerated tumors will be monitored at least 3 times per week with increasing frequency, up to daily, depending upon clinical signs.
- Ulcerated tumors, which have not scabbed over, should be cleaned with an appropriate wound cleansing solution (e.g., Novalsan). Antibiotic cream is to be applied to the ulceration/lesion only if directed by the Veterinary staff.

Criteria for euthanasia include if the lesion (meet one or more):

- Does not heal or form a scab within 1 week.
- Is greater than 5 mm diameter.
- Becomes cavitated.

Develops signs of infection (such as presence of pus) or bleeding, or if the animal shows signs of discomfort (e.g. excessive licking and biting directed at the site) or systemic signs of illness (lethargy, decreased activity, decreased food consumption, decreased body condition or weight loss). Contact the veterinary staff to discuss any possible exceptions.

[0324] **3) Clinical signs:** Animals must be euthanized if they found to be moribund (unless special permission is granted by the IACUC based on adequate justification, which must be included in the protocol and increased supportive care provided such as warm SQ fluids, Diet Gel food cup next to animal so they can reach food, cage on a warming pad for supplemental heat, etc. Note: a moribund condition indicates an animal is unlikely to survive.) For questions regarding these endpoints, please contact the Veterinary Staff.

[0325] Clinical examples of morbidity may include:

- Hunched.
- Persistent recumbency and lack of response to handling or other stimuli.
- Signs of severe organ or system failure.

- Emaciation.
- Hypothermia.
- CNS deficits: convulsions.
- Respiratory: rapid respiratory rate, labored breathing, coughing, rales.
- GI: diarrhea lasting > 2 days, jaundice.

[0326] Any animal that exhibits the above clinical issues will be humanely sacrificed by CO₂.

[0327] Necropsy will not be performed in the event of an unexpected death.

[0328] **Statistical Analysis:** For comparison between two groups, an independent sample *t*-test will be used. For comparison among three or more groups, a one-way ANOVA will be performed. If a significant F -statistics (a ratio of treatment variance to the error variance) is obtained, *multiple comparison* procedures will be applied after *ANOVA*. All data will be analyzed using SPSS 17.0. *p* < 0.05 is considered to be statistically significant.

[0329] **Results:**

[0330] All the conjugates show outstanding tumor inhibitory effects ((Figure 8a and Figure 8b).

[0331] 4.4 On NCI-N87 human gastric cancer xenograft model

[0332] **Study Purpose:** The objective of the research is to evaluate the *in vivo* anti-tumor efficacy of CH-2-589, CH-2-593, CH-2-518 and LC302-2-4(4) in the treatment of the subcutaneous NCI-N87 human gastric cancer xenograft model in female BALB/c Nude mice.

[0333] **Study Design:**

[0334] **Cell Culture:** The NCI-N87 tumor cells (ATCC, Manassas, VA, cat # CRL-5822) will be maintained *in vitro* as a monolayer culture in RPMI 1640 medium supplemented with 10% fetal bovine serum, 1% Antibiotic-Antimycotic, at 37°C in an atmosphere of 5% CO₂ in air. The tumor cells will be routinely subcultured twice weekly by trypsin-EDTA treatment. The cells growing in an exponential growth phase will be harvested and counted for tumor inoculation.

[0335] **Animals:** BALB/c Nude, female, 6~8 weeks, weighing approximately 18~22g. A total of 36 (30 plus 20%) will be needed for the study, which will be purchased from Shanghai Lingchang Laboratory Animal Co., LTD. or other certified vendors.

[0336] **Tumor Inoculation:** Each mouse will be inoculated subcutaneously at the right flank with NCI-N87 tumor cells (10×10^6) in 0.2 mL of PBS with Matrigel (1:1) for tumor development. The animals will be randomized and treatment will be started when the average tumor volume reaches approximately 150-200 mm³ for the efficacy study. The test article administration and the animal numbers in each group are shown in the following experimental design table.

Groups and Treatments

Group	N ¹	Treatment	Dose (mg/kg)	Dosing Volume (mL/kg) ²	Dosing Route	Schedule ³
1	6	Vehicle	--	10	IV	Single dose
2	6	CH-2-589	5	10	IV	Single dose
3	6	CH-2-593	5	10	IV	Single dose
4	6	CH-2-518	5	10	IV	Single dose
5	6	LC302-2-4(4)	5	10	IV	Single dose

N: animal number;

Dosing volume: adjust dosing volume based on body weight 10 mL/kg.

The experiment duration will be adjusted according to tumor volume.

[0337] **Results:**

[0338] All the conjugates show outstanding tumor inhibitory effects (Figure 9a and Figure 9b).

Effect Example 5 Ex-vivo serum stability of the conjugates

[0339] For ex-vivo serum stability study, the conjugates LC302-2-1(4), LC302-2-4(4) and LC1184(8) were inoculated in pooled human serum at 37°C, respectively. At 0, 24, 48 and 96 h, the conjugates were captured by antigen, then deglycosylated by glycosidase and dissociated by acid. Supernatant were collected and centrifuged, and then detected by high resolution LC-MS to determine DAR.

[0340] LC302-2-1(4) and LC302-2-4(4) are quite stable after incubated in serum for 96 hours. There is no observable decrease in the DARs of LC302-2-1(4) and LC302-2-4(4) after incubation for 96 h. However, the DAR of LC1184(8) dropped from 7.8 to 2.6 after incubation for 96 h (Figure 10). Stability of the linker indicates the delivery of more payloads to target tumour and decrease of off-target free payload release, and thus increases therapeutic index.

Sequence Listing

SEQ ID NO: 1: Ab0001-LCCT_L-HC LCDR1
RASQDVNTAVA

SEQ ID NO: 2: Ab0001-LCCT_L-HC LCDR2
SASFLYS

SEQ ID NO: 3: Ab0001-LCCT_L-HC LCDR3
QQHYTTPPT

SEQ ID NO: 4: Ab0001-LCCT_L-HC HCDR1
DTYIH

SEQ ID NO: 5: Ab0001-LCCT_L-HC HCDR2
RIYPTNGYTRYADSVKG

SEQ ID NO: 6: Ab0001-LCCT_L-HC HCDR3
WGGDGFYAMDY

SEQ ID No. 7: Ab0001-LCCT_L-HC Light chain variable region:
DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAPKLLIYSASFLYSGV
PSRFGSGRSGTDFLTISLQPEDFATYYCQQHYTTPPTFGQGTKVEIKRTV


SEQ ID NO: 8: Ab0001-LCCT_L-HC Heavy chain variable region:
EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGLEWVARIYPTNGY
TRYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCSRWGGDGFYAMDYWGQ
GTLVTVSS

SEQ ID No. 9: Ab0001-LCCT_L-HC Light chain full length:
DIQMTQSPSSLSASVGDRVTITCRASQDVNTAVAWYQQKPGKAPKLLIYSASFLYSGV
PSRFGSGRSGTDFLTISLQPEDFATYYCQQHYTTPPTFGQGTKVEIKRTVAAPSVFIFP
PSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSTYLS
STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECGALPETGG


SEQ ID No. 10: Ab0001-LCCT_L-HC Heavy chain full length:
EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGLEWVARIYPTNGY
TRYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCSRWGGDGFYAMDYWGQ
GTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGV
HTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHT
CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE
VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAK
GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVL
DSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPGK

Claim

1. A compound of formula (I):



wherein,



R^0 is C_{1-3} alkyl;


n is any integer of 2 to 5;

k_1 and k_2 are independently an integer of 1 to 7;

i is an integer of 1-20;

j is an integer of 1-20;

P_1 and P_2 are independently a payload having the structure of formula (i'):



wherein,

a is 0 or 1;

the carbon atoms marked with p_1^* and p_2^* each is asymmetric center, and the

asymmetric center is *S* configured, *R* configured or racemic;

L^1 is selected from C_{1-6} alkylene, which is unsubstituted or substituted with at least one substituent selected from halogen, -OH and -NH₂;

M is -CH₂-, -NH- or -O-;

L^2 is C_{1-3} alkylene;


R^1 is selected from Cl and Br;

R^2 is F.


2. The compound of claim 1, wherein in formula (i'):

L^1 is selected from C_{1-6} linear alkylene, C_{1-6} branched alkylene, C_{3-6} cyclic alkylene and C_{3-4} cyclic alkyl- C_{1-2} linear alkylene group, which are each independently unsubstituted or substituted with at least one substituent selected from halogen, -OH and -NH₂.

3. The compound of claim 2, wherein in formula (i'):

L^1 is selected from -CH₂-, -C₂H₄-, , which are each independently unsubstituted or substituted with at least one substituent selected from halogen, -OH and -NH₂.

4. The compound of claim 3, wherein in formula (i'):

L^1 is selected from -CH₂-, , wherein “#” marks the position attached to carbonyl.

5. The compound of any one of the claims 1 to 4, wherein a is 0.

6. The compound of any one of the claims 1 to 4, wherein in formula (i'):
the carbon atom marked with p1* is *S* configured.

7. The compound of any one of the claims 1 to 4, wherein in formula (i'):
the carbon atom marked with p2* is *S* configured.

8. The compound of any one of the claims 1 to 4, wherein in formula (i'), R^1 is Cl.


9. The compound of any one of the claims 1 to 4, wherein a is 0, R^1 is Cl, R^2 is F, and L^1

is selected from $-\text{CH}_2-$, ,  and .


10. The compound of any one of the claims 1 to 4, wherein a is 0, R^1 is Cl, R^2 is F, and L^1

is .

11. The compound of any one of the claims 1 to 4, wherein the payload is selected from:



12. The compound of any one of the claims 1 to 4, wherein the payload is selected from:




13. The compound of any one of the claims 1 to 4, wherein n is 3.

14. The compound of any one of the claims 1 to 4, wherein k1 and k2 are 5.

15. The compound of any one of the claims 1 to 4, wherein i is 2 to 8.

16. The compound of any one of the claims 1 to 4, wherein j is 8 to 12.

17. A conjugate having the structure of formula (II):



wherein,

A is an antibody or an antigen binding fragment thereof;

z is an integer of 1 to 4;

P1, P2, R^0 , $opSu$, n , $k1$, $k2$, i and j are as defined in any one of claims 1 to 16.

18. The conjugate of claim 17, wherein the antibody or antigen binding fragment is modified to connect with the (Gly)_n moiety in the compound of formula (I) as defined in any one of claims 1 to 16.

19. The conjugate of claim 18, wherein z is 2.

20. The conjugate of any one of the claims 17 to 19, wherein the antibody is an anti-HER2 antibody.


21. A pharmaceutical composition comprising a prophylactically or therapeutically effective amount of a conjugate of any one of the claims 17 to 20, and at least one pharmaceutically acceptable carrier.

22. Use of the conjugate of any one of the claims 17 to 20 or the pharmaceutical composition of claim 21 in the manufacture of a medicament for treating a disease; wherein the

disease is a tumor.

23. The use of claim 22, wherein the disease is a tumor comprising HER2-positive tumor cells.

24. A compound, or a pharmaceutically acceptable salt thereof, wherein the compound has the structure of formula (i):



wherein,

a is 0 or 1;

the carbon atoms marked with p1* and p2* each is asymmetric center, and the asymmetric center is *S* configured, *R* configured or racemic;

L¹ is selected from C₁₋₆ alkylene, which is unsubstituted or substituted with at least one substituent selected from halogen, -OH and -NH₂;

M is -CH₂-, -NH- or -O-;

L² is C₁₋₃ alkylene;

R¹ is selected from Cl and Br;

R² is F.

25. The compound of claim 24, wherein in formula (i'):


L¹ is selected from C₁₋₆ linear alkylene, C₁₋₆ branched alkylene, C₃₋₆ cyclic alkylene and C₃₋₄ cyclic alkyl-C₁₋₂ linear alkylene group, which are each independently unsubstituted or substituted with at least one substituent selected from halogen, -OH and -NH₂.

26. The compound of claim 25, wherein in formula (i'):


L¹ is selected from -CH₂-, -C₂H₄-, , , , , , which are each independently unsubstituted or substituted with at least one substituent selected from

halogen, -OH and -NH₂.


27. The compound of claim 25, wherein in formula (i'):



28. The compound of claim 27, wherein in formula (i'):



29. The compound of claim 28, wherein in formula (i'):



30. The compound of any one of the claims 24 to 29, wherein a is 0.


31. The compound of any one of the claims 24 to 29, wherein in formula (i'):
the carbon atom marked with p1* is *S* configured or racemic.

32. The compound of claim 31, wherein in formula (i'):
the carbon atom marked with p1* is *S* configured.


33. The compound of any one of the claims 24 to 29, wherein in formula (i'):
the carbon atom marked with p2* is *S* configured.

34. The compound of any one of the claims 24 to 29, wherein in formula (i'), R¹ is Cl.


35. The compound of any one of the claims 24 to 28, wherein a is 0, R¹ is Cl, R² is F, and



36. The compound of any one of the claims 24 to 29, wherein a is 0, R¹ is Cl, R² is F, and



37. A compound of formula (1):




wherein,

k is an integer of 1 to 7;


P is a payload having the structure of formula (i'), wherein the structure of formula (i') is as defined in any one of claims 1 to 16.

38. The compound of claim 37, wherein k is 1, or 3 or 5.

39. A compound of formula (I):



wherein,



R⁰ is C₁₋₃ alkyl;


n is any integer of 2 to 5;

k1 and k2 are independently 1, or 3 or 5;


i is independently an integer of 1 to 20;

j is independently an integer of 1 to 20;

P1 and P2 are independently a payload having the following structure:



40. The compound of claim 39, having the structure of formula (I-1)



wherein,

P1, P2, R⁰, opSu, n, i and j are as defined in claim 39.

41. The compound of claim 39 or 40, wherein

R⁰ is methyl; and/or

n is 3; and/or


i is 4; and/or

j is 8 or 12.


42. The compound of claim 39 or 40, wherein

R⁰ is methyl; n is 3; i is 4; j is 12.

43. The compound of claim 39 or 40, which is selected from



44. A conjugate having the structure of formula (II):




46. The conjugate of claim 44 or 45, wherein the antibody is Trastuzumab.

47. A pharmaceutical composition comprising a prophylactically or therapeutically effective amount of a conjugate of any one of the claims 44 to 46, and at least one pharmaceutically acceptable carrier.


48. Use of the conjugate of any one of the claims 44 to 46 or the pharmaceutical composition of claim 47 in the manufacture of a medicament for treating a disease; wherein the disease is a tumor.

49. A compound, or a pharmaceutically acceptable salt thereof, wherein the compound has a structure selected from the following formulae:



and


50. A compound of formula (1):



wherein,

k is an integer of 1 to 7;

P has a structure of the following:



51. The compound of claim 50, wherein k is 1, or 3 or 5.

52. The compound of claim 50, which is selected from:

