


Description

The present invention refers to complexes between bilobalide, a sesquiterpene extracted from the leaves of *Ginkgo biloba*, and natural or synthetic phospholipids, to the preparation thereof and to the therapeutical use thereof. The complexes prepared according to the procedure described in the present invention should be considered as new compounds. In fact, they possess physico-chemical and spectroscopic characteristics which are markedly different from those of the original components and as such they can be incorporated as active principles into pharmaceutical formulations.

The production of complexes of natural products with phospholipids has already been investigated by the Applicant as a method to enhance the bioavailability of highly polar substances such as flavonoids (US-4,764,508; EP-0 275 005 A2) or saponins (EP-0 283 713 A2), in order to improve their absorption and duration of action. Also in the case of the agents described in the present invention, formation of complexes with phospholipids enabled the preparation of new biologically active compounds.

Bilobalide forms with phospholipids new compounds having the following general formula



wherein R and R₁ which can be the same or different, are each an acyl residue of a C₁₆-C₂₂ saturated or unsaturated fatty acid; R₂ is one of the following residues: -CH₂-CH₂-N-(CH₃)₃, -CH₂-CH₂-NH₃, -CH₂-CH₂-(COOH)-NH₃. The phospholipids used for the preparation of these complexes may be either natural or synthetic.

The complexes described in the present invention are normally prepared by reacting in aprotic solvents, such as aromatic hydrocarbons, halogen derivatives or some cyclic ethers, one mole of phospholipid with one mole of bilobalide. The final products are then isolated by evaporation of the reaction solvent or by precipitation of the complex with appropriate solvents. These complexes may also be prepared by lyophilizing the product obtained from the reaction of the components in dioxane.

The formation of the complex is confirmed by nuclear magnetic resonance analysis, which enables the identification of the interaction between the two reacting molecular species and the portion of the molecule which is involved in the reaction. As mentioned above, the phospholipids selected for the preparation of the complexes are derivatives from choline or ethanolamine or inositol, in which the glycerol moiety is bound to saturated or unsaturated C₁₆-C₂₂ acyl chains, the natural phospholipids are normally those extracted from soy-bean or from animal organs, whereas the synthetic phospholipids have homogeneous acyl chains and are prepared according to methods described in the literature. Bilobalide shows a strong affinity for phospholipids, resulting in the generation of bonds which markedly modify the physico-chemical and spectroscopic characteristics of the new molecules. In the ¹H-NMR spectrum of the complex, in fact, some of the signals from the protons of the terpenic group are absent, while others are shifted and markedly broadened. The signals from the protons of the lipid are visible but partially modified. The N-CH₃ signal of the phospholipid, which appears as a sharp singlet in the spectrum of the free phospholipid, takes in the complex the appearance of a broader band, indicating that this portion of the molecule is involved in the generation of the complex. In the proton spectrum the signals from the lipid chains are clearly visible, indicating that these chains retain their mobility in the solvent. The comparison of the ¹³C-NMR spectrum of the complex with that of its individual components evidences the occurrence of marked changes, similar to those reported for the protons. The spectrum of the complex lacks many of the signals from the sesquiterpene. The signals from the carbons belonging to the choline and glyceric components of the phospholipid are also virtually absent. The signals from the lipid chains can be recorded (in a somewhat

broadened form) in both the proton and the carbon spectra, indicating that these chains retain their mobility in the solvent. In contrast with the situation encountered with polar compounds such as flavonoids and saponins, the formation of these complexes involves a clear participation of the acyl portion of the phospholipid. The signals from the acyl chains are broadened, probably due to partial immobilization caused by the interaction with the substance bound in the complex. The latter is probably partially enwrapped by the acyl chains, but some of its groups remain exposed to the surrounding medium, as in the case of the *terbutyl* moiety, whose signal at 1.1 ppm can be recorded in broadened form in the proton spectrum and is also visible in the carbon spectrum. In this respect, the complexes described in the present invention differ from those described previously for other compounds.

If exception is made for particular applications, the complexes are preferentially prepared by using natural phosphatidylcholine of vegetal origin with a standardized content in fatty acids (linoleic acid 63%, palmitic acid 16%, oleic acid 11%, stearic acid 3,5%, expressed as percentage of the total fatty acid content).

As described in the present invention, production of the complex was found to provide a useful mode of administration for bilobalide, which is a lipophilic molecule *per se*. The possibility of producing new derivatives of bilobalide which retain unaltered the structure of the basic compound but exhibit a several-fold increase in specific activity represents a significant advantage which certainly could not be predicted beforehand. Due to the enhancement in specific activity, the dosage can be reduced with considerable economic savings. The advantage of using derivatives with higher specific activity is immediately clear when it is considered that bilobalide is found in the leaves of *Ginkgo biloba* at concentrations below 0.01%.

From the therapeutic point of view, bilobalide has been previously reported to possess a useful activity (DE-3338995) for the treatment of neuropathies of various origin. In the same report, an antiinflammatory activity was excluded or was not determined. On the contrary, the antiinflammatory activity represents part of the present invention, in addition to the neurotrophic activity which is markedly increased as a result of the formation of complexes with phospholipid substances. For the latter specific application, it was found that not only pure phospholipids, but also phospholipid mixtures extracted from brain, skin and peripheral nerves are particularly useful. Natural or purified mixtures of vegetal phospholipids may advantageously be used for some applications such as cosmetic products and mild dermatological remedies.

From the biological point of view, the complexes synthesized according to the procedure outlined above have been evaluated for antiinflammatory activity by using classical models such as the carrageenin-induced oedema in the rat and the croton oil-induced oedema in the mouse after systemic and topical administration respectively.

When the topical route is used, formation of complexes with phospholipids results in a clear-cut and significant increase in bioavailability as assessed by an increase in anti-oedema activity.

Table 1 illustrates the results concerning the antiinflammatory activity obtained with free bilobalide and with its equimolar complex with distearoyl-phosphatidylcholine.

Table 1

Anti-oedema activity of the complex of bilobalide with distearoyl-phosphatidylcholine in the croton oil test.				
Substances	Number of animals	Dose μ mole/ear	Oadema	Reduction
Controls	35	-	5.8 +/-0.3	-
Bilobalide	30	0.4	2.1 +/-0.3	63.8% *
	30	0.2	3.3 +/-0.3	43.1% *
	30	0.1	4.8 +/-0.4	19.5%
Bilobalide/distearoyl-phosphatidylcholine complex	30	0.4	1.0 +/-0.1	82.8% *
	30	0.2	2.9 +/-0.3	50.0% *
	30	0.1	3.4 +/-0.3	41.1% *
	30	0.05	4.6 +/-0.7	20.7% *
Distearoyl-phosphatidylcholine	30	0.4	4.9 +/-0.6	18.8%
	30	0.2	4.6 +/-0.7	20.7%

* statistically significant as compared to controls
 $P < 0.05$

The complexes with other saturated phospholipids exhibit a similar behaviour and activity.

The complexes with unsaturated phospholipids such as lineoyl-phosphatidylcholine, or with a soy-bean phosphatidylcholine product containing the average percent proportions of fatty acids reported above for natural phosphatidylcholine, or with phosphatidylcholine extracted from bovine or swine brain, exhibit greater activity in this test, due to a significant contribution from the phospholipids themselves. As observed for other substances, formation of a complex of bilobalide with phospholipids leads to a prolongation of the duration of action of the active principle as compared to free bilobalide administered at the same dosage by the same route.

After oral administration, the complexes of bilobalide with phospholipids exhibit a greater bioavailability which results in an enhanced pharmacodynamic response, as shown in Table II.

Table II

Anti-inflammatory activity on carrageenin paw edema in rats of complex bilobalide/distearylphosphatidylcoline

Substances	Dose (mg/kg)	No. of animals	Baseline paw volume (ml)	Mean increments ml over baseline after carrageenin (a ± s.e.)	1h	2h	3h	4h	5h
Controls	-	8	1.74 ± 0.03	0.25 ± 0.04	0.60 ± 0.09	0.75 ± 0.11	0.67 ± 0.10	0.52 ± 0.08	
Bilobalide	200	8	1.72 ± 0.05	0.26 ± 0.03	0.55 ± 0.07	0.66 ± 0.05	0.66 ± 0.07	0.50 ± 0.05	
			-1.1X	4.0X	-8.3X	-12.0X	-1.5X	-3.8X	
Bilobalide complex (as bilobalide)	70	8	1.73 ± 0.03	0.25 ± 0.03	0.48 ± 0.06	0.76 ± 0.11	0.77 ± 0.09	0.62 ± 0.09	
			-0.6X	0.0X	-20.0X	1.3X	14.9X	19.2X	
Bilobalide complex (as bilobalide)	200	8	1.72 ± 0.04	0.20 ± 0.03	0.35 ± 0.06	0.51 ± 0.08	0.52 ± 0.06	0.39 ± 0.07	
			-1.1X	-20.0X	-41.7X	-32.0X	-22.4X	-25.0X	
Indomethacin	10	8	1.70 ± 0.02	0.12 ± 0.02	0.30 ± 0.02	0.35 ± 0.05	0.49 ± 0.05	0.44 ± 0.06	
			-2.3X	-52.0X	-50.0X	-53.3X	-26.9X	-15.4X	

Substances were given by gastric gavage one hour before carrageenin injection into the foot.

* p < 0.05 Dunnett's t test

As far as the more complex neurotrophic activity and anti-cerebral oedema activity are concerned, investigations were carried out by using indirect tests which provide an indication of the prospected effect. The nerve protecting activities of bilobalide and its complexes were evaluated by using the inflammatory pain test of Randall and Selitto or the hot plate test in animals intoxicated with neurotoxic agents such as nerve gases or organic derivatives of tin and lead. Electrophysiological investigations on the rate of

recovery of damaged nerves showed an increased velocity of conduction of the impulse in motor nerves and a reduced conduction in sensory nerves. Surprisingly, in all these experiments the complex of bilobalide with phospholipids was found to exert greater activity as compared to bilobalide in free form. The electrophysiological investigations on Bilobalide and its complexes with distearoylphosphatidylcholine were extended also to the study of their ability to interfere with the reinnervation process of the muscle cells of the extensor digitorum longus muscle of the rat, consequent on the lesion and the regeneration of the sciatic nerve. The products were administered to the animals at a dosage of 10 mg/kg of Bilobalide for 6 days the week either by intraperitoneal or by oral route and their effect was evaluated at different times after denervation through an estimate of the muscle cells receiving more than one motor nerve terminal. Considering that the polyinnervation is a negative condition from the physiological point of view, since it leads to a destabilization of the innervated cell, any product favouring the reinstatement of physiological conditions can be regarded as highly valuable. We found in our test that Bilobalide in its free form exhibits a significant inhibitory activity on the polyinnervation only at 30 days after denervation, while its complex with distearoylphosphatidylcholine is endowed with a more precocious action, antagonizing the polyinnervation significantly already 20 days after the lesion. At 25 days after the lesion Bilobalide in its free form even produces an increase, albeit of modest degree and statistically not significant with respect to controls, of the percentage of polyinnervated muscle cells. The results of the investigation are shown in Table III.

Table III

Percentage of polyinnervated muscle cells following the lesion and the regeneration of the sciatic nerve at different times after denervation.					
	days				
	18	20	25	30	60
controls					
mean	33.4	43.8	21.8	24.4	9.4
S.E.	2.8	7.4	3.9	4.4	4.7
bilobalide					
mean	31.1	29.1	26.9	10.4*	6.2
S.E.	3.3	5.2	4.7	0.7	2.4
bilobalide complex					
mean	35.8	22.4*	19.6	20.3	5.3
S.E.	2.3	4.8	3.7	3.8	3.0

* $p < 0.05$ analysis of variance

The anti-oedema activity may be partly related to interference with prostaglandin synthesis at vascular level, a possibility which does not affect the validity of the current invention. By contrast, the improvement in motor nerve conduction may be ascribed to a more rapid functional recovery of the axon. Bilobalide, which is structurally related to the better known ginkgolides, both in free form and as a complex with phospholipids possesses a very weak anti-PAF activity which cannot account in any way for the antagonistic effect exerted in different models of inflammation. Conversely, the compound exhibits a non-specific immunomodulating activity which may influence favourably its main action.

On the basis of the evidence discussed above, the complexes of bilobalide with appropriate phospholipids may be usefully employed in the treatment of inflammatory states of different aetiology, including many neuritic disorders of various origin such as those associated with diabetes and autoimmune diseases. Because of their high extent of percutaneous absorption, the products described in the present invention may be particularly useful for the topical treatment of superficial or deep inflammatory processes, including those affecting the joints, and for the management of neuropathies secondary to inflammatory processes of infectious, traumatic or metabolic origin.

The reaction of bilobalide with phospholipids under appropriate conditions results in the generation of new products whose physico-chemical characteristics are completely different from those of the individual constituents. These products exhibit a greater activity as compared to bilobalide in free form and are

suitable for incorporation into the most common pharmaceutical formulations.

In human therapeutics, the new products can be used at dosages ranging from 1 to 500 mg/day in single or divided daily administrations. These products may be administered also as topical formulations, at variable dosages depending on the severity of the pathological condition being treated.

- 5 The following examples are designed to better illustrate the invention, without representing a limitation in any case.

EXAMPLE 1

- 10 Preparation of the equimolar complex of bilobalide with distearoyl-phosphatidylcholine.

50 ml of methylene chloride, 3.26 g of bilobalide and 7.95 g of distearoyl-phosphatidylcholine are suspended in a 100 ml-flask, then heated under light reflux until the contents are completely dissolved. The solution is cooled and evaporated to dryness, the residue is re-dissolved in 25 ml of methylene chloride.

- 15 The clear solution is concentrated to a small volume and the concentrate is poured into 50 ml n-hexane: this results in the formation of an abundant precipitate which is filtered and dried. This procedure yields 11 g of bilobalide/distearoyl-phosphatidylcholine complex, which exhibits the following characteristics: melting point 244-6 °C, ¹H-NMR (CDCl₃): -CH₃ at 0.8 ppm, C-(CH₃)₃ at 1.14 ppm, -CH₂- between 1.4 and 2.1 ppm, N-(CH₃)₃ at 3.3 ppm, 12-H at 6.28 ppm.

- 20 The superimposed spectra of bilobalide, distearoyl-phosphatidylcholine and their equimolar complex are shown in Figure 1 (attached).

EXAMPLE 2

- 25 Preparation of the complex of bilobalide with pure phosphatidylcholine extracted from soy-bean

3.26 g of bilobalide are added under continuous stirring to a solution containing 7.7 g of soy-bean phosphatidylcholine (Lipoid S-100) in 50 ml methylene chloride: the mixture is heated under reflux until the contents are completely dissolved, then evaporated to dryness under vacuum. The residue is re-dissolved

- 30 in methylene chloride and concentrated again to dryness. This procedure yields 10.4 g of a yellowish product which exhibits the following spectroscopic characteristics:

¹H-NMR (CDCl₃): -CH₃ at 0.8 ppm, C-(CH₃)₃ at 1.18 ppm, -CH₂- between 1.4 and 2.4 ppm, N-(CH₃)₃ at 3.3 ppm, olefinic protons at 5.1-5.4 ppm, 12-H at 6.28 ppm.

- 35 **EXAMPLE 3**

Preparation of the complex of bilobalide with distearoyl-phosphatidylethanolamine

- 40 3.26 g of bilobalide together with 7.56 g of distearoyl-phosphatidylethanolamine are dissolved in 40 ml of anhydrous dioxane. The resulting solution is lyophilized.

This procedure yields a soft solid white substance soluble in methylene chloride and in benzene, which exhibits the following spectroscopic characteristics:

¹H-NMR (CDCl₃): -CH₃ at 0.8 ppm, C-(CH₃)₃ at 1.14 ppm, -CH₂- between 1.4 and 2.1 ppm, 12-H at 6.28 ppm.

- 45

EXAMPLE 4

Preparation of the complex of bilobalide with a natural mixture of phospholipids extracted from bovine brain

- 50 3.26 g of bilobalide together with 7.5 g of brain phospholipids are suspended in 40 ml ethyl acetate. This suspension is heated to 40 °C until the contents are completely dissolved, then concentrated to dryness and the waxy residue is re-dissolved in 25 ml methylene chloride. 0.1 g of decolourizing vegetal charcoal are added to this solution, which is then filtered. The clear filtrate is concentrated to dryness under vacuum. This procedure yields 9.8 g of a straw-coloured product with a dense oily appearance, which
- 55 exhibits the spectroscopic characteristics typical of a complex.

EXAMPLE 5**Preparation of an aqueous microdispersion of the complex of bilobalide with brain phospholipids**

- 5 3.26 g of bilobalide are dissolved in 50 ml methylene chloride together with 7.5 g of a mixture of phospholipids extracted from bovine cerebral cortex and characterized by the following composition: 31% phosphatidylcholine, 14% phosphatidylethanolamine, 11% phosphatidylinositol and the remaining fraction consisting of minor phospholipids. When all contents have been dissolved, the solvent is evaporated under vacuum. The residue is re-suspended in 750 ml of distilled water using as an aid initially an Ultraturax and
 10 subsequently ultrasonication. The microdispersion prepared according to this procedure may be applied directly for the epicutaneous treatment of the pathological conditions outlined in the general section.

EXAMPLE 6

- 15 **Preparation of capsules containing the bilobalide/phosphatidylcholine complex**

Each 250 mg capsule contains:

20	Bilobalide/phosphatidylcholine complex	230 mg
	Micronized silica gel	15 mg
	Polivinylpyrrolidone	2.5 mg
	Magnesium stearate	2.5 mg

25 **EXAMPLE 7**

Preparation of a cream containing the bilobalide/phosphatidylcholine complex

- 30 100 g of cream contain:

35	Bilobalide/soy-bean phosphatidylcholine complex	1 g
	Carboxyvinylpolymer (Carbomer 934)	13 g
	Sodium laurylsarcosinate (30%)	0.5 g
	Polysorbate	3 g
	Hydrogenated lanoline	5 g
	Spermaceti	5 g
	Polyisoprene	5 g
40	Silicon oil	0.5 g
	Sodium hydroxide (1N)	4 g
	Water	to 100 g

45 **Claims**

Claims for the following Contracting States : BE, CH, DE, FR, GB, IT, LI, LU, NL, SE, AT, DK

1. Complexes of bilobalide with natural or synthetic, saturated or unsaturated phospholipids, having the following general formula:

50

55

EXAMPLE 5Preparation of an aqueous microdispersion of the complex of bilobalide with brain phospholipids

- 5 3.26 g of bilobalide are dissolved in 50 ml methylene chloride together with 7.5 g of a mixture of phospholipids extracted from bovine cerebral cortex and characterized by the following composition: 31% phosphatidylcholine, 14% phosphatidylethanolamine, 11% phosphatidylinositol and the remaining fraction consisting of minor phospholipids. When all contents have been dissolved, the solvent is evaporated under vacuum. The residue is re-suspended in 750 ml of distilled water using as an aid initially an Ultraturax and
- 10 subsequently ultrasonication. The microdispersion prepared according to this procedure may be applied directly for the epicutaneous treatment of the pathological conditions outlined in the general section.

EXAMPLE 6

- 15 Preparation of capsules containing the bilobalide/phosphatidylcholine complex

Each 250 mg capsule contains:

Bilobalide/phosphatidylcholine complex	230 mg
Micronized silica gel	15 mg
Polivinylpyrrolidone	2.5 mg
Magnesium stearate	2.5 mg

25

EXAMPLE 7Preparation of a cream containing the bilobalide/phosphatidylcholine complex

- 30 100 g of cream contain:

Bilobalide/soy-bean phosphatidylcholine complex	1 g
Carboxyvinylpolymer (Carbomer 934)	13 g
Sodium laurylsarcosinate (30%)	0.5 g
Polysorbate	3 g
Hydrogenated lanoline	5 g
Spermaceti	5 g
Polyisoprene	5 g
Silicon oil	0.5 g
Sodium hydroxide (1N)	4 g
Water	to 100 g

45 **Claims**


Claims for the following Contracting States : BE, CH, DE, FR, GB, IT, LI, LU, NL, SE, AT, DK

1. Complexes of bilobalide with natural or synthetic, saturated or unsaturated phospholipids, having the following general formula:

50

55






wherein R and R₁, which can be the same or different, are each an acyl residue of a C₁₆-C₂₂ saturated or unsaturated fatty acid; R₂ is one of the following residues: -CH₂-CH₂-N-(CH₃)₃, -CH₂-CH₂-NH₃, -CH₂-CH(COOH)-NH₃.

2. Complexes as claimed in claim 1, in which the phospholipids are derived from choline, ethanolamine and inositol.
3. A process for the preparation of the complexes as claimed in claim 1, in which process bilobalide and the phospholipid are reacted in aprotic solvents and the resulting complex is precipitated with a non-solvent or isolated by lyophilization.
4. A process as claimed in claim 3, in which process the aprotic solvents used are aromatic hydrocarbons, halogenated derivatives or cyclic ethers.
5. A process as claimed in claims 3 and 4, in which the reagents are reacted in stoichiometric ratios.
6. Pharmaceutical formulations containing as active principle the complexes claimed in claims 1 and 2, mixed with an appropriate vehicle.
7. Cosmetic or dermatological formulations containing as active principle the complexes claimed in claims 1 and 2.
8. Use of the complexes claimed in claim 1 for the preparation of a drug with antiinflammatory and neurotrophic properties.

Claims for the following Contracting States : GR, ES

1. A process for the preparation of complexes of bilobalide with natural or synthetic, saturated or unsaturated phospholipids, having the following general formula:






wherein R and R₁, which can be the same or different, are each an acyl residue of a C₁₆-C₂₂ saturated or unsaturated fatty acid; R₂ is one of the following residues: -CH₂-CH₂-N-(CH₃)₃, -CH₂-CH₂-NH₃, -CH₂-CH(COOH)-NH₃, in which process bilobalide and the phospholipid are reacted in aprotic solvents and the resulting complex is precipitated with a non-solvent or isolated by lyophilization.

2. A process as claimed in claim 1, in which process the aprotic solvents used are aromatic hydrocarbons, halogenated derivatives or cyclic ethers.
3. A process as claimed in claims 1 and 2, in which the reagents are reacted in stoichiometric ratios.
4. Cosmetic or dermatological formulations containing as active principle the complexes obtained according to the process claimed in claims 1 to 3.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : BE, CH, DE, FR, GB, IT, LI, LU, NL, SE, AT, DK

1. Komplexe von Bilobalid mit natürlichen oder synthetischen, gesättigten oder ungesättigten Phospholipiden der allgemeinen Formel:



wobei R und R₁ gleich oder unterschiedlich sind und beide ein Acylrest einer gesättigten oder ungesättigten C₁₆-C₂₂-Fettsäure sind; R₂ einer der folgenden Reste ist: -CH₂-CH₂-N-(CH₃)₃, -CH₂-CH₂-NH₃, -CH₂-CH(COOH)-NH₃.


2. Komplexe nach Anspruch 1, wobei sich die Phospholipide von Cholin, Ethanolamin und Inositol ableiten.
3. Verfahren zur Herstellung der Komplexe nach Anspruch 1, wobei Bilobalid und das Phospholipid in aprotischen Lösungsmitteln umgesetzt werden und der erhaltene Komplex mit einem Nichtlöser ausgefällt wird oder durch Lyophilisierung isoliert wird.



4. Verfahren nach Anspruch 3, in welchem die verwendeten aprotischen Lösungsmittel aromatische Kohlenwasserstoffe, halogenierte Derivate oder cyclische Ether sind.
5. Verfahren nach den Ansprüchen 3 und 4, in welchem die Reagenzien in stöchiometrischen Verhältnissen umgesetzt werden.
6. Pharmazeutische Zubereitungen, welche als Wirkstoff die Komplexe nach Anspruch 1 und 2 im Gemisch mit einem geeigneten Träger enthalten.
7. Kosmetische oder dermatologische Zubereitungen, welche als Wirkstoff die Komplexe nach Anspruch 1 und 2 enthalten.
8. Verwendung der Komplexe nach Anspruch 1 für die Herstellung eines Arzneimittels mit entzündungshemmenden und neurotrophen Eigenschaften.

Patentansprüche für folgende Vertragsstaaten : GR, ES

1. Verfahren zur Herstellung von Komplexen von Bilobalid mit natürlichen oder synthetischen, gesättigten oder ungesättigten Phospholipiden der allgemeinen Formel:



wobei R und R₁ gleich oder unterschiedlich sind und beide ein Acylrest einer gesättigten oder ungesättigten C₁₆-C₂₂-Fettsäure sind; R₂ einer der folgenden Reste ist: -CH₂-CH₂-N-(CH₃)₃, -CH₂-CH₂-NH₃, -CH₂-CH(COOH)-NH₃, in welchem Verfahren Bilobalid und das Phospholipid in aprotischen Lösungsmitteln umgesetzt werden und der erhaltene Komplex mit einem Nichtlöser ausgefällt wird oder durch Lyophilisierung isoliert wird.


2. Verfahren nach Anspruch 1, in welchem die verwendeten aprotischen Lösungsmittel aromatische Kohlenwasserstoffe, halogenierte Derivate oder cyclische Ether sind.
3. Verfahren nach den Ansprüchen 1 und 2, in welchem die Reagenzien in stöchiometrischen Verhältnissen umgesetzt werden.
4. Kosmetische oder dermatologische Zubereitungen, welche als Wirkstoff die aus dem Verfahren nach den Ansprüchen 1 bis 3 erhaltenen Komplexe enthalten.

Revendications

Revendications pour les Etats contractants suivants : BE, CH, DE, FR, GB, IT, LI, LU, NL, SE, AT, DK

1. Complexes de bilobalide et de phospholipides saturés ou insaturés, d'origine naturelle ou synthétique, répondant à la formule générale suivante:






dans laquelle R et R₁, qui peuvent être identiques ou différents, sont chacun un résidu acyle d'un acide gras saturé ou insaturé en C₁₆-C₂₂; R₂ est l'un des résidus suivants: -CH₂-CH₂-N-(CH₃)₃, -CH₂-CH₂-NH₃, -CH₂-CH(COOH)-NH₃.

2. Complexes tels que revendiqués dans la revendication 1, dans lesquels les phospholipides sont dérivés de la choline, de l'éthanolamine et de l'inositol.
3. Procédé pour la préparation de complexes tels que revendiqués dans la revendication 1, dans lequel le bilobalide et le phospholipide sont mis à réagir dans des solvants aprotiques et le complexe obtenu est précipité dans un non solvant ou isolé par lyophilisation.
4. Procédé tel que revendiqué dans la revendication 3, dans lequel les solvants aprotiques employés sont des hydrocarbures aromatiques, des dérivés halogénés ou des éthers cycliques.
5. Procédé tel que revendiqué dans les revendications 3 et 4, dans lequel les réactifs sont mis à réagir en proportions stoechiométriques.
6. Formulations pharmaceutiques contenant à titre de principe actif les complexes revendiqués dans les revendications 1 et 2, en mélange avec un véhicule approprié.
7. Formulations cosmétiques ou dermatologiques contenant à titre de principe actif les complexes revendiqués dans les revendications 1 et 2.
8. Utilisation des complexes revendiqués dans la revendication 1 pour la préparation d'un médicament doué de propriétés anti-inflammatoires et trophoneurotiques.

Revendications pour les Etats contractants suivants : GR, ES

1. Procédé pour la préparation de complexes de bilobalide et de phospholipides saturés ou insaturés, d'origine naturelle ou synthétique, répondant à la formule générale suivante:





dans laquelle R et R₁, qui peuvent être identiques ou différents, sont chacun un résidu acyle d'un acide gras saturé ou insaturé en C₁₆-C₂₂; R₂ est l'un des résidus suivants: -CH₂-CH₂-N-(CH₃)₃, -CH₂-CH₂-NH₃, -CH₂-CH(COOH)-NH₃, procédé dans lequel le bilobalide et le phospholipide sont mis à réagir dans des solvants aprotiques et le complexe obtenu est précipité dans un non solvant ou isolé par lyophilisation.

2. Procédé tel que revendiqué dans la revendication 1, dans lequel les solvants aprotiques employés sont des hydrocarbures aromatiques, des dérivés halogénés ou des éthers cycliques.
3. Procédé tel que revendiqué dans les revendications 1 et 2, dans lequel les réactifs sont mis à réagir en proportions stoechiométriques.
4. Formulations cosmétiques ou dermatologiques contenant à titre de principe actif les complexes obtenus conformément au procédé revendiqué dans les revendications 1 à 3.

