

DESCRIPTION

TEST AGENT FOR DIAGNOSING DYSPEPSIA

TECHNICAL FIELD

5 The present invention relates to a composition for measuring the gastric-emptying function, which can be used to effectively diagnose dyspepsia, and a method for measuring the gastric-emptying function. Specifically, the present invention relates to a composition that can be used to measure reductions
10 of the gastric-emptying function in a noninvasive manner using expired air, and a method for measuring the gastric-emptying function using the composition. The present invention also relates to a dyspepsia diagnostic agent and a method for diagnosing dyspepsia. Furthermore, the present invention relates
15 to a method for measuring the drug efficiency and treatment effects of a gastrointestinal medicine, in particular, a medicine affecting the alimentary tract motor function on a patient suffering from gastrointestinal disorders caused by dyspepsia and like insufficiencies in the gastric-emptying function.

20

BACKGROUND ART

Nowadays, we live in a society with much stress due to rapid and drastic changes in various matters. Accordingly, there has been an increase in patients who complain of gastroenteric
25 pain and discomfort (such as epigastralgia/epigastric discomfort, heavy stomach, heartburn, epigastric fullness, early satiety, nausea, vomiting, etc.). Even if a subject suffers from such symptoms for a long period of time, if no disorders are detected by magendurchleuchtung, endoscopy or abdominal ultrasonography,
30 such symptoms would be diagnosed as unidentified complaints in the abdominal region and would not be subjected to medical treatment, or diagnosed and treated as chronic gastritis.

However, it has recently been considered that such symptoms may be attributable to dysfunctions of the alimentary
35 tract. In the United States, when unidentified complaints last at

least four weeks in spite of the absence of structural abnormalities when checked with an endoscope, such conditions are referred to as "symptoms of non-ulcer dyspepsia in the alimentary tract" (Non-ulcer Dyspepsia: NUD, hereunder referred to as
5 "dyspepsia") and distinguished from chronic gastritis (1987, the American Gastroenterological Association).

Dyspepsia is roughly categorized into the following four groups according to subjective symptoms:

(1) a gastroesophageal reflux type (heartburn, acid
10 reflux, reflux feeling, belching), (2) a dysmotility type (early satiety, bloating, anorexia, nausea, vomiting), (3) an ulcer type (night pain, hunger pain, periodic discomfort, abdominal pain), and (4) a non-specific type (symptoms not applicable to (1) to (3) above). These symptoms excluding (1), i.e., the symptoms of
15 (2) to (4), are collectively called Functional Dyspepsia: FD (see, ROME II scale: Non-Patent Document 1). Among these dyspepsias, in particular, the dysmotility type (2) is the most common, and it accounts for 30 to 40% of the total dyspepsias (Non-Patent Document 2).

20 Treatment of dyspepsia is often conducted as a symptomatic therapy using medicines. Since the gastroesophageal reflux type (1) is significantly related to gastric acid, a medicine that can control the secretion of gastric acid is used. The non-specific type (4) is assumed to be caused by
25 psychological factors, and therefore antianxiety medicines and antidepressants are often used. However, as shown below, medicines for improving the alimentary tract motor function serve a key role in treating these symptoms:

(1) gastroesophageal reflux type: acid antisecretory,
30 antacid, and medicines for improving the alimentary tract motor function;

(2) dysmotility type and (3) ulcer type: medicines for improving alimentary tract motor function; and

(4) non-specific type: medicines for improving the
35 alimentary tract motor function, antianxiety medicines, and

antidepressants.

Dyspepsia is diagnosed by exclusion, i.e., when a subject complains of gastroenteric pain or discomfort (epigastralgia or epigastric discomfort, heavy stomach, heartburn, epigastric fullness, extreme sensation of fullness after a small amount of food, nausea, vomiting, etc.) for a long time, but no structural abnormalities were found even after being examined with endoscopy, abdominal ultrasonography, etc. Dyspepsia is generally diagnosed on the basis of the symptoms reported by the subject, and this makes the diagnosis of dyspepsia difficult. Furthermore, the difficulty of such diagnosis imposes mentally and monetary burdens on the patient and delays appropriate treatment, causing deterioration in the patient's quality of life (QOL).

Therefore, development of an easy and highly accurate method of diagnosing dyspepsia will reduce the mentally and monetary burdens on dyspepsia patients and greatly contribute to its treatment.

One of the primary causes of dyspepsia is a lowering of the gastric-emptying function, and therefore a technique for measuring the gastric-emptying function can be utilized in its diagnosis. However, conventional methods for measuring the gastric-emptying function have the following drawbacks. They are expensive or invasive, the subject is constrained for a long time so that they impose a psychological and physical burden on the subject, and the measurement accuracy is insufficient. For example, among conventional methods for measuring the gastric-emptying function, isotope methods (e.g., scintigraphy or the like) use a radioactive isotope; as a result, the administration of such a method is complicated. Furthermore, since such a method requires an expensive γ -ray camera, its use is restricted to special facilities. In the case of an X-ray impermeable marker method, the marker is not excreted from the stomach simultaneously with the food contents, but is instead excreted from the stomach after all of the food contents have been

excreted; as a result, the actual gastric-emptying function cannot be accurately examined. In the case of an acetaminophen method, there is a danger of drug allergy and liver damage due to the side effects of acetaminophen; furthermore, since this drug is subject to other effects in the body, such as absorption in the small intestine, metabolism by the liver, excretion from the kidneys and the like, the gastric-emptying function cannot be accurately examined. Since the concentration of acetaminophen in the blood is measured after administering acetaminophen, the invasive procedure of blood collection is required.

Furthermore, methods such as measuring the endogastric volume and saburra by means of ultrasonic waves (an ultrasound method), measuring the gastric-emptying function by MRI (a magnetic resonance imaging method), evaluating the gastric motor function by measuring with an electrogastrogram (elctrogastrography) and the like have also been proposed for measuring the gastric-emptying function; however, such methods pose the following problems: (i) There are problems in the accuracy of the diagnostic method, (ii) there are no fixed criteria, so that evaluations vary according to the evaluator, and (iii) the subject must be constrained for a long period of time while the diagnosis is being made. (Non-Patent Documents 3 and 4).

Non-Patent Document 1: Talley Nj et al., Gut 45 (Suppl 2): II 37-42, 1999

Non-Patent Document 2: Quarteo AO et al., Dig Dis Sci 43: 2028-2033, 1998

Non-Patent Document 3: J. Smooth Muscle Res. (Jpn. Sec.) 6: J-75-J-91, 2002

Non-Patent Document 4: J. Smooth Muscle Res. (Jpn. Sec.) 6: J-129-J-138, 2002

DISCLOSURE OF THE INVENTION

An object of the present invention is to provide a

composition for measuring the gastric-emptying function used to effectively diagnose dyspepsia. Specifically, the present invention aims to provide a composition that allows an easy and noninvasive measurement of the gastric-emptying function using
5 expired air. The present invention also aims to provide a method for measuring the gastric-emptying function that can be used to effectively diagnose dyspepsia. Specifically, the main objects of the present invention are to provide a dyspepsia diagnostic agent, and a method for diagnosing dyspepsia.

10 Furthermore, the present invention aims to provide a method for measuring therapeutic effects (including pharmacotherapy) on patients, such as dyspepsia patients, suffering from gastrointestinal disorders caused by an insufficient gastric-emptying function. In particular, the
15 present invention aims to provide a method for measuring the pharmaceutical effects on the patients of medicines affecting the function of the alimentary tract and/or therapeutic effects thereof.

20 Means for Solving the Problem

Shown below are the four characteristics desirable for inspection probes for use in diagnosing and evaluating gastric-emptying functions using expired air.

25 (1) The inspection probe is not absorbed from the stomach, but from the alimentary tracts including and below the duodenum (such as the duodenum, jejunum, ileum, etc.);

(2) Its absorption is not affected by the change in pH (the pH in the alimentary tract);

30 (3) The probe exhibits high absorption and metabolic rates, and a high excretion rate in expired air as isotope-labeled CO₂ (a high recovery rate); and

(4) The probe is promptly metabolized after absorption.

35 In order to solve the above problems, the present

inventors conducted an intensive search to develop a gastric-emptying function inspection probe having the above characteristics, and found that uracil, thymine and like pyrimidine compounds labeled with an isotope of C or O have the
5 four characteristics. The present inventors also found that by measuring the amount or behavior of the isotope-labeled CO₂ excreted in expired air after orally administering these compounds to a subject, the gastric-emptying function of the subject could be easily measured. They confirmed that the
10 measuring method is effective for diagnosing dyspepsia in a noninvasive manner. They also confirmed that by using the method for measuring the gastric-emptying function using a pyrimidine compound, the effects of the medicine or like treatment on the subject can be measured, allowing to select a treatment method
15 most desirable and effective for the subject (the patient). The present invention has been accomplished based on such findings.

The present invention includes the following embodiments:

20 (I) Composition for measuring gastric-emptying function

(I-1) A composition for measuring the gastric-emptying function containing as an active ingredient a pyrimidine compound that is labeled with either or both isotopes of C and O, and that is converted into labeled CO₂ in the body and excreted in expired
25 air.

(I-2) The composition for measuring the gastric-emptying function according to Item (I-1), wherein the isotope is at least one member selected from a group consisting of ¹³C, ¹⁴C and ¹⁸O.

30 (I-3) The composition for measuring the gastric-emptying function according to Item (I-1) or (I-2), wherein the labeled pyrimidine compound is uracil or thymine.

(II) Dyspepsia diagnostic agent

35 (II-1) A dyspepsia diagnostic agent consisting of a

composition of any one of Items (I-1) to (I-3).

(II-2) The dyspepsia diagnostic agent according to Item (II-1), wherein the dyspepsia is caused by an insufficient gastric-emptying function.

5 (II-3) The dyspepsia diagnostic agent according to Item (II-2), wherein the dyspepsia is dysmotility-type dyspepsia.

(III) Method for measuring gastric-emptying function

(III-1) A method for measuring the gastric-emptying
10 function comprising: orally administering a composition of any one of Items (I-1) to (I-3) to a subject, and measuring the amount or behavior of the labeled CO₂ excreted in expired air.

(III-2) The method for measuring the gastric-emptying function according to Item (III-1), wherein a composition of any
15 one of Items (I-1) to (I-3) is orally administered to a subject in whom a drop or acceleration in the gastric-emptying function is suspected, and the amount or behavior of the labeled CO₂ excreted in expired air is measured and compared with the amount or behavior of the labeled CO₂ excreted in expired air from a healthy
20 subject, measured using the same composition.

More specifically, the methods of (III-1) and (III-2) are as below:

(III-1') A method for measuring the gastric-emptying function comprising: using expired air from a subject who had
25 orally taken a composition of any one of Items (I-1) to (I-3) as a test sample, and measuring *in vitro* the amount of the labeled CO₂ in the test sample.

(III-2') The method for measuring the gastric-emptying function according to Item (III-1'), which comprises:
30 using expired air from a subject in whom a drop or acceleration in the gastric-emptying rate is suspected and who had orally taken a composition of any one of Items (I-1) to (I-3), as a test sample;

measuring *in vitro* the amount of the labeled CO₂ in the
35 test sample; and

comparing the measured value with the amount of labeled CO₂ in expired air from a healthy subject who had taken the same composition.

5 (IV) Method for diagnosing dyspepsia

 (IV-1) A method for diagnosing dyspepsia comprising:
orally administering to a subject a dyspepsia diagnostic agent of
any one of Items (II-1) to (II-3), and measuring the gastric-
emptying function based on the amount or behavior of the labeled
10 CO₂ excreted in expired air.

 (IV-2) The method for diagnosing dyspepsia according to
Item (IV-1), which comprises orally administering a dyspepsia
diagnostic agent of any one of Items (II-1) to (II-3) to a subject
in whom dyspepsia is suspected;
15 comparing the amount or behavior of the labeled CO₂
excreted in expired air from the subject with that from a healthy
subject measured using the same diagnostic agent; and

 measuring the gastric-emptying function of the subject.

 (IV-3) The diagnosis method according to Item (IV-1) or
20 (IV-2), which is for diagnosing dyspepsia caused by an
insufficient gastric-emptying function.

 (IV-4) The diagnosis method according to Item (IV-3),
which is for diagnosing dysmotility-type dyspepsia.

 The methods of Items (IV-1) to (IV-4) can also be
25 expressed as below.

 (IV-1') A method for diagnosing dyspepsia comprising:
using expired air from a subject who had orally taken a dyspepsia
diagnostic agent of any one of Items (II-1) to (II-3) as a test
sample, measuring the amount of the labeled CO₂ in the test sample
30 *in vitro*, and measuring the gastric-emptying function.

 (IV-2') The method for diagnosing dyspepsia according
to Item (IV-1'), wherein expired air from a subject in whom
dyspepsia is suspected and who had orally taken a dyspepsia
diagnostic agent of any one of Items (II-1) to (II-3) is used as a
35 test sample, the amount of the labeled CO₂ in the test sample is

measured *in vitro*, the amount is compared with that of the labeled CO₂ in expired air from a healthy subject who had taken the same diagnostic agent, and the gastric-emptying function of the subject is then measured.

5 (IV-3') The diagnosis method of Item (IV-1') or (IV-2'), which is for diagnosing dyspepsia caused by an insufficient gastric-emptying function.

(IV-4') The diagnosis method according to Item (IV-3'), which is for diagnosing dysmotility-type dyspepsia.

10

(V) Method for measuring pharmaceutical or therapeutic effect

(V-1) A method for measuring the effect of the gastrointestinal treatment on a subject comprising:

before and after conducting a gastrointestinal treatment
15 to a subject, orally administering a composition of any one of Items (I-1) to (I-3) or a diagnostic agent of any one of Items (II-1) to (II-3) to the subject; and

comparing the amounts or behaviors of the labeled CO₂ excreted in expired air before and after the gastrointestinal
20 treatment.

(V-2) A method for measuring the pharmaceutical effect of the gastrointestinal medicine or the therapeutic effect on a subject comprising:

before and after the administration of a
25 gastrointestinal medicine to a subject, orally administering a composition of any one of Items (I-1) to (I-3) or a diagnostic agent of any one of Items (II-1) to (II-3) to the subject; and

comparing the amounts or behaviors of the labeled CO₂ excreted in expired air before and after the administration of the
30 gastrointestinal medicine.

(V-3) The measuring method of Item (V-2), wherein the gastrointestinal medicine affects the alimentary tract motor function.

(V-4) The measuring method according to any one of
35 Items (V-1) to (V-3) that measures the pharmaceutical effect of

the gastrointestinal medicine or the therapeutic effect on a patient suffering from dyspepsia caused by an insufficient gastric-emptying function.

(V-5) The measuring method according to any one of
5 Items (V-1) to (V-3) that measures the pharmaceutical effect of the gastrointestinal medicine or the therapeutic effect on a patient suffering from dysmotility-type dyspepsia.

The methods of Items (V-1) to (V-5) can also be expressed as below:

10 (V-1') A method for measuring the effect of a gastrointestinal treatment on a subject comprising:

before and after the gastrointestinal treatment, collecting expired air from a subject who had orally taken a composition of any one of Items (I-1) to (I-3) or a diagnostic
15 agent of any one of Items (II-1) to (II-3) as a test sample; and comparing the amounts of the labeled CO₂ in the test samples collected before and after the gastrointestinal treatment.

(V-2') A method for measuring the pharmaceutical effect of the gastrointestinal medicine or the therapeutic effect on a
20 subject comprising:

before and after administering a gastrointestinal medicine, collecting expired air from a subject who had orally taken a composition of any one of Items (I-1) to (I-3) or a diagnostic agent of any one of Items (II-1) to (II-3) as a test
25 sample; and

comparing the amounts of the labeled CO₂ in the test samples collected before and after the administration of the gastrointestinal medicine.

(V-3') The measuring method according to Item (V-2'),
30 wherein the gastrointestinal medicine affects the alimentary tract motor function.

(V-4') The measuring method according to any one of Items (V-1') to (V-3') that measures the pharmaceutical effect of the gastrointestinal medicine or the therapeutic effect on a
35 patient suffering from dyspepsia caused by an insufficient

gastric-emptying function.

(V-5') The measuring method according to any one of Items (V-1') to (V-3') that measures the pharmaceutical effect of the gastrointestinal medicine or the therapeutic effect on a patient suffering from dysmotility-type dyspepsia.

(VI) Use

(VI-1) A use of a pyrimidine compound for producing a dyspepsia diagnostic agent, wherein the pyrimidine compound is labeled with either or both isotopes of C and O, converted into an isotope-labeled CO₂ gas in the body, and then excreted in expired air.

(VI-2) The use of the pyrimidine compound according to Item (VI-1), wherein the isotope is at least one member selected from the group consisting of ¹³C, ¹⁴C and ¹⁸O.

(VI-3) The use of the pyrimidine compound according to Item (VI-1) or (VI-2), wherein the pyrimidine compound is uracil or thymine.

In the present invention, the term "dyspepsia" means "a condition in which there is an unidentified complaint in the upper abdomen for at least four weeks, regardless of the absence of structural abnormalities verified with an endoscope", i.e., non-ulcer dyspepsia (NUD). The dyspepsia targeted by the present invention is preferably caused mainly by an insufficient gastric-emptying function. Such dyspepsia includes functional dyspepsia (FD), in particular, dysmotility-type dyspepsia. Here, "an insufficient gastric-emptying function" means that the gastric-emptying function is abnormal, including a decrease in the gastric-emptying function as well as the loss of the gastric-emptying function.

Effect of the Invention

The composition of the present invention makes it possible to measure the gastric-emptying function in humans or animals in

a simple and accurate manner. Specifically, the composition of the present invention is usable in objectively measuring the gastric motor function, diagnosing the diseases caused by the gastric motor dysfunction, and measuring the pharmaceutical effect of medicines affecting the alimentary tract motor function and/or therapeutic effect on a patient.

The dyspepsia diagnostic agent of the present invention makes it possible to diagnose dyspepsia by readily and accurately measuring the reduction of a subject's gastric-emptying function. In particular, the diagnostic agent of the present invention uses, as an active ingredient, a labeled piperidine compound excreted in expired air in a form of a labeled carbon dioxide gas. This allows using the dyspepsia diagnostic agent by merely testing the expired air, without imposing a psychological or physical burden on the subject. The diagnostic agent is particularly effective in measuring the effects of treating a dyspeptic patient (including pharmacotherapy), in particular measuring the pharmaceutical effects of medicines affecting the motor function of the alimentary tract or the therapeutic effects thereof.

20

BEST MODE FOR CARRYING OUT THE INVENTION

(I) Composition for measuring gastric-emptying function, and (II) Dyspepsia diagnostic agent

25 The composition for measuring the gastric-emptying function of the present invention comprises as an active ingredient a pyrimidine compound that is labeled with at least one of isotopes of C and O, and that is excreted in expired air after being converted into a labeled CO₂ gas in the body.

30 The pyrimidine compound used in the composition has a pyrimidine skeleton labeled with at least one of isotopes of C and O, so that it is excreted in expired air after being converted into a labeled CO₂ gas in the body when orally administered. Specific examples thereof include uracil, thymine, 35 cytosine, 5-methylcytosine and like pyrimidine bases. Preferable

properties of the pyrimidine compound used in the present invention are as follows:

- (1) All or almost all thereof is absorbed from the alimentary tracts including and below the duodenum (such as the duodenum, jejunum, ileum) without completely or barely being absorbed in the stomach, and then excreted in expired air as an isotope-labeled CO₂ gas after being decomposed or metabolized.
- (2) Its absorption is not readily affected by the change in pH in the alimentary tract. More preferable properties are:
- (3) Having a high absorption rate, a high metabolic rate, and a high excretion rate (recovery rate) as a labeled CO₂ gas in expired air. A particularly preferable property, in addition to the above properties, is the ability to
- (4) be promptly metabolized after absorption. Examples of pyrimidine compounds having such properties include uracil and thymine.

There is no limitation to the isotopes used in labeling the carbon or oxygen atom in a pyrimidine compound, and usable examples include ¹³C, ¹⁴C and ¹⁸O. Such isotopes may be radioactive or non-radioactive; however, from the standpoint of safety, non-radioactive isotopes are preferable. In particular, ¹³C is especially desirable for use as such an isotope.

Specifically, the pyrimidine compound used in the present invention is isotope labeled in such a manner that at least a portion of the CO₂ formed through the pyrimidine metabolic pathway is isotope labeled. Examples of such pyrimidine compounds include those having the carbon atom at the 2-position in the pyrimidine skeleton labeled with an isotope. Specific examples include 2-¹³C labeled uracil, 2-¹³C labeled thymine, 2-¹³C labeled cytosine, etc. Among these, 2-¹³C labeled uracil and 2-¹³C labeled thymine are preferable.

There are no particular restrictions on the method for labeling a pyrimidine compound with these isotopes, and wide various commonly used methods may be employed. (see Sasaki, "5.1 Use of Stable Isotopes in Clinical Diagnoses"; Kagaku no Ryoiki 107, "Use of Stable Isotopes in Medicine, Pharmacy and Biology",

pp. 149-163 (1975) Nankodo; Kajiwara, RADIOISOTOPES, 41, 45-48 (1992); and the like). Furthermore, some isotope-labeled pyrimidine compounds, such as 2-¹³C labeled uracil, are commercially available and these commercial products are
5 conveniently usable.

There is no particular limitation to the composition of the present invention in terms of its form, components other than the isotope-labeled pyrimidine compound, proportion of each component, and preparation method of the composition, as long as
10 the pyrimidine compound contained therein is absorbed in and below the duodenum after oral administration, and excreted in expired air as a labeled CO₂ gas after being metabolized.

There is no limitation to the form of the composition, as long as it can be taken orally. Examples of the forms of the
15 composition include any orally administrable forms, such as solutions (including syrup), suspensions, emulsions and like liquids; tablets (with and without coating), chewable tablets, capsules, pills, pulvis (powders), fine particles, granules and like solids.

20 The application of the composition of the present invention is not limited to formulation, such as pharmaceutical preparation, as long as it contains the labeled pyrimidine compound and does not adversely affect the effects of the present invention. The labeled pyrimidine compound may be combined with
25 any foodstuff and formed into solid food, fluid food or liquid food.

The composition of the present invention may substantially consist of the isotope-labeled pyrimidine compound, which is an active ingredient; however, as long as the effects of
30 the present invention are not adversely affected, any pharmaceutically acceptable carriers and/or additives that are generally used in this field may be added.

In this case, there is no limitation to the amount of the isotope-labeled pyrimidine compound added as an active
35 ingredient. For example, the amount of the isotope-labeled

pyrimidine compound falls in the range from 1 to 95 wt.% per 100 wt.% of the composition, and is preferably controlled within this range.

When the composition of the present invention is formed
5 into, for example, tablets, chewable tablets, capsules, pills, pulvis (powders), fine particles, granules and like solid forms, various carriers and/or additives suitable to such forms may be added.

Specific examples of usable carriers or additives
10 include lactose, sucrose, dextrin, mannitol, xylitol, sorbitol, erythritol, calcium dihydrogen phosphate, sodium chloride, glucose, urine, starch, calcium carbonate, kaolin, crystalline cellulose, silicic acid and like excipients; water, ethanol, simple syrup, glucose liquid, starch liquid, gelatin liquid,
15 carboxymethyl cellulose, sodium carboxymethyl cellulose, shellac, methyl cellulose, hydroxypropylmethyl cellulose, hydroxypropyl cellulose, potassium phosphate, polyvinyl alcohol, polyvinyl pyrrolidone, dextrin, pullulan and like binders; dry starch, sodium alginate, agar powder, laminaran powder, sodium
20 bicarbonate, calcium carbonate, polyoxyethylenesorbitan fatty acid esters, sodium lauryl sulfate, monoglyceride stearate, starch, lactose, carmellose calcium, low substituted hydroxypropyl cellulose, carmellose, croscarmellose sodium, sodium carboxymethyl starch, crospovidone and like
25 disintegrators; saccharose, stearic acid, cacao butter, hydrogenated oil and like disintegration inhibitors; polysorbate 80, quaternary-ammonium salt, laurylsodium sulfate and like absorbefacients; glycerin, starch and like humectants; starch, lactose, kaolin, bentonite, colloidal silicic acid and like
30 adsorbents; purified talc, stearate, boric acid powder, polyethylene glycol, colloidal silicic acid, sucrose fatty acids, hardened oil and like lubricants; citric acid, anhydrous citric acid, sodium citrate, sodium citrate dihydrate, anhydrous sodium monohydrogenphosphate, anhydrous sodium dihydrogenphosphate,
35 sodium hydrogenphosphate and like pH adjustors; iron oxide, β

carotene, titanium oxide, edible pigment, copper chlorophyll, riboflavin and like coloring agents; and ascorbic acid, sodium chloride, various sweeteners and like corrigents.

Tablets may be provided with an ordinary coating, if
5 necessary. Specific examples thereof include sugar-coated tablets, gelatin-coated tablets, film-coated tablets, double-coated tablets, multi-coated tablets, etc. Capsules are prepared in a commonly employed method, i.e., mixing the isotope-labeled pyrimidine compound, which is an active ingredient, with various
10 carriers mentioned above and placing it in a hard gelatin capsule, a soft capsule, etc.

A preferable composition of the present invention for measuring the gastric-emptying function in an accurate manner with little variation attributable to individual subject
15 differences can be obtained by mixing (a) an isotope-labeled pyrimidine compound and (b) sugar and/or sugar alcohol, and then pulverizing the resulting mixture. The thus-obtained powder is formed into a preparation.

There is no limitation to the sugar and sugar alcohol
20 used here as far as it is pharmaceutically acceptable. Usable sugars include glucose, galactose, fructose, xylose, arabinose, mannose and like monosaccharides; maltose, isomaltose, cellobiose, lactose, sucrose, trehalose and like disaccharides. Among these, glucose and sucrose are preferable. Examples of sugar alcohols
25 include erythritol, mannitol, xylitol, sorbitol, maltitol, hydrogenated palatinose, lactitol, etc. Among these, mannitol, xylitol and erythritol are preferable, and mannitol is particularly preferable. Sugar alcohol is more preferably than sugar used as component (b).

30 The content of component (a) in the composition is preferably 5 to 20 wt.%, more preferably 6 to 18 wt.%, and particularly preferably 8 to 15 wt.%, per total weight of the composition. The content of component (b) is generally 80 to 95 wt.%, preferably 82 to 94 wt.%, and more preferably 85 to 92 wt.%,
35 per total weight of the composition.

There is no limitation to the ratio of the component (a) to the component (b), and, for example, component (b) may generally be added in 400 to 1,900 parts by weight, preferably 450 to 1,550 parts by weight, and more preferably 550 to 1,150 parts by weight, per 100 parts by weight of component (a).

Such a composition is preferably prepared using a powder material containing component (a) and component (b), and forming the material into preparations. The powder material can preferably be prepared by mixing component (a) with component (b) in such a manner so as to achieve the above-mentioned proportions, and then pulverizing the mixture. There is no limitation to the particle diameter of the powder material, but in order to minimize the variations attributable to individual subject differences and increase the accuracy of the measurement in the gastric-emptying function, the particle diameter at 50% is generally not greater than 40 μm , preferably not greater than 30 μm , and particularly preferably from 5 to 20 μm . The distribution of the particle size is preferably such that the particle diameter at 50% is not greater than 40 μm , and the particle diameter at 90% is not greater than 200 μm ; more preferably, the particle diameter at 50% is not greater than 30 μm , and the particle diameter at 90% is not greater than 100 μm ; and particularly preferably, the particle diameter at 50% is from 5 to 20 μm , and the particle diameter at 90% is from 10 to 70 μm . Such particle size distribution can be measured by a dry laser method (measurement conditions: focal distance: 100 mm; a number of averaging processes: 10 times; an averaging interval: 5 milliseconds; air pressure: 0.4 MPa).

There is no limitation to the pulverization treatment employed in the preparation of the powder material, but a pulverization treatment using a dry pulverizer is preferable. Examples of usable dry pulverizers include hammer mills, pin mills, jet mills, etc.

The composition may consist of components (a) and (b), and may contain other components as long as the above-mentioned

proportions of components (a) and (b) are included. In this case, usable components other than components (a) and (b) include the above-described pharmaceutically acceptable carriers and/or additives (for example, excipients, binders, pH adjustors, 5 disintegrators, absorbefacients, lubricants, coloring agents, corrigents, flavor, etc.). Preferably, those carriers and/or additives had been subjected to the same pulverization treatment as components (a) and (b). There is no limitation to the form of the preparation as long as it can be taken orally, and it may be 10 in the form of a fine particle, granule, pulvis (powder), tablet (with and without coating), capsule, pill, etc. Among these, fine particle, granule and like particles, in particular particles obtained by extrusion granulation, are preferable.

When formed into particles, the average particle 15 diameter of the preparation is, for example, generally not greater than 1,400 μm , preferably within the range of 50 to 1,200 μm , and more preferably within the range of 100 to 1,000 μm . By forming the preparation into particles having such a particle diameter, the gastric-emptying function can be measured in a more 20 accurate manner. The measurement of particle diameters can be conducted by a vibration sieving method (specifically, using a measurement apparatus: Robot Sifter RPS-95, Seishin Enterprise Co., Ltd.; vibration level: 5; shift time: 5 minutes; pulse interval: 1 second).

25 The amount of the isotope-labeled pyrimidine compound (active ingredient) contained in a unit dosage of the composition of the present invention cannot be generalized because it varies depending on the types of the measurement sample, the active ingredients etc., and is therefore suitably selected according to 30 each case. For example, when 2-¹³C labeled uracil and like isotope-labeled uracils are used as an isotope-labeled pyrimidine compound, i.e., the active ingredient, generally 1 to 1,000 mg/body and preferably 10 to 100 mg/body of isotope-labeled uracil is contained in one unit dosage of the composition. When 35 other isotope-labeled pyrimidine compounds are used as an active

ingredient, the content may be suitably selected based on the above amounts.

The composition for measuring the gastric-emptying function of the present invention allows measuring the gastric-emptying function of a subject when taken orally, by measuring
5 the amount or behavior of the labeled CO₂ gas excreted in expired air.

Specifically, the composition for measuring the gastric-emptying function of the present invention, after being
10 taken orally, enters into the stomach, and is then excreted from the pylorus of the stomach by systolic and diastolic motion and/or peristaltic movement. After being excreted from the pylorus of the stomach, the isotope-labeled pyrimidine compound, which is an active ingredient, is promptly absorbed and
15 metabolized in the alimentary tract including and below the duodenum (the duodenum, jejunum, ileum, etc.), and then excreted in expired air as a labeled CO₂ gas. One of the major features of the isotope-labeled pyrimidine compound used in the present invention is that it is not absorbed, or barely absorbed, in the
20 stomach, and that it is promptly absorbed and metabolized after being discharged from the stomach, and finally excreted in expired air as a labeled CO₂ gas. Therefore, the behavior of the labeled CO₂ gas excreted in expired air (specifically, for example, the behavior is expressed as the proportion of the isotope-labeled CO₂ gas in the ¹²CO₂ excreted in expired air, (isotope-labeled CO₂/¹²CO₂)), depends on the gastric emptying rate (gastric emptying time) of the composition of the present invention, i.e.,
25 the isotope-labeled pyrimidine compound.

The following factors can be used as the indices
30 expressing the gastric-emptying function. Specifically, the amount of the ¹³CO₂ gas in expired air after a predetermined time from the administration of the composition, the carbon dioxide gas value Δ(%) (the difference of the ¹³CO₂/¹²CO₂ concentration ratios (δ¹³C values) before and after the administration of the
35 composition in expired air collected), or the initial speed of

the $^{13}\text{CO}_2$ gas. For example, when the carbon dioxide gas value $\Delta(\%)$ or the initial speed of the $^{13}\text{CO}_2$ gas of a healthy subject is determined as a criterion value, and if the subject exhibited a carbon dioxide gas value $\Delta(\%)$ or the initial speed lower than
5 that of a healthy subject, the subject is diagnosed as having a reduced gastric-emptying function.

The composition for measuring the gastric-emptying function of the present invention may be taken singly, together with an experimental diet, or immediately before or after taking
10 the experimental diet. One preferable example of the method is that the composition for measuring the gastric-emptying function of the present invention is taken immediately after taking an experimental diet. There is no limitation to the experimental diet taken, as far as it does not adversely affect the
15 measurement of the gastric-emptying function using the composition of the present invention, and it may be a solid food, fluid food or liquid food.

As described above, one of the main causes of dyspepsia (non-ulcer dyspepsia symptoms in the alimentary tract) is
20 dysfunction of the alimentary tract motor, especially lowering of the gastric-emptying function. Accordingly, the composition for measuring the gastric-emptying function of the present invention is effectively usable as an agent for diagnosing dyspepsia, especially dyspepsia mainly caused by insufficiencies in the
25 gastric-emptying function (for example, dysmotility-type dyspepsia). Therefore, all explanations relating to the composition for measuring the gastric-emptying function of the present invention can also be applied to the dyspepsia diagnostic agent.

30

(III) Method for measuring gastric-emptying function and (iv) Method for diagnosing dyspepsia

The present invention also relates to a method for measuring the gastric-emptying function conducted using the
35 composition for measuring the gastric-emptying function described

above. The measurement of the gastric-emptying function can be conducted by orally administering the composition for measuring the gastric-emptying function of the present invention, which contains an isotope-labeled pyrimidine compound as an active
5 ingredient, to animals or humans; collecting expired air; and measuring the amount or behavior of the labeled CO₂ gas excreted in expired air.

When ¹³C is used as an isotope, the gastric-emptying function can be measured according to an ordinary ¹³C breath test,
10 i.e., after orally administering the composition for measuring the gastric-emptying function of the present invention to a subject, expired air is collected over the lapse of time, and, with the amount of ¹³CO₂ excreted in expired air being expressed as ¹³CO₂/¹²CO₂ ratio (δ¹³C value), the behavior is observed over
15 time.

The composition for measuring the gastric-emptying function of the present invention contains, as an active ingredient, an isotope-labeled pyrimidine compound that is not absorbed or is barely absorbed in the stomach. After being
20 excreted from the stomach, the isotope-labeled pyrimidine compound is hardly affected by the pH in the alimentary tract, and is promptly absorbed and metabolized in and below the duodenum (the duodenum, jejunum, ileum, etc.). Thereafter, a large percentage thereof is excreted into expired air as an
25 isotope-labeled CO₂ gas. This achieves the measurement directly and accurately reflecting the gastric excretion motor function. By preparing the composition for measuring the gastric-emptying function using a powder material containing an isotope-labeled pyrimidine compound and a sugar and/or sugar alcohol, the
30 variance between subjects can be reduced and the gastric excretion motor function can be accurately measured.

The measurement of the gastric-emptying function is conducted with higher accuracy by administering the composition for measuring the gastric-emptying function of the present
35 invention not only once but several times in a repeated manner

and under various conditions including after fasting, after eating, etc. The measurement and analysis methods of the isotope-labeled CO₂ in expired air vary depending on whether the isotope used is radioactive or non-radioactive, but generally used methods include the liquid scintillation counter method, mass spectrometry, infrared spectroscopy, emission spectrometry, and the magnetic resonance spectrum method. From the viewpoint of measurement accuracy, infrared spectroscopy and mass spectrometry are preferable.

The method for administering the composition for measuring the gastric-emptying function of the present invention may be the same as that described above, but is not limited to this.

The amount of the isotope-labeled pyrimidine compound per unit dose of the composition of the present invention cannot be generalized because it varies depending on the types of the isotope-labeled compound, etc., and may be suitably selected in each case. For example, when measurement is conducted by an expired air test using 2-¹³C uracil as an isotope-labeled pyrimidine compound, preferably 1 to 2,000 mg, and more preferably 10 to 300 mg of 2-¹³C uracil is contained in a preparation per unit dose. When other isotope-labeled pyrimidine compounds are used as an active ingredient, the amounts may be suitably selected based on the above.

By employing the method for measuring the gastric-emptying function of the present invention, the drop or acceleration of a subject's excretion function can be diagnosed and measured. Specifically, the diagnosis can be conducted by the comparison between the amount or behavior of the isotope-labeled CO₂ gas excreted in expired air from a subject measured by the above method with the standard control (the amount or behavior of the isotope-labeled CO₂ gas excreted in expired air from a healthy subject).

The gastric-emptying function of a subject can be diagnosed and measured, for example, in the manner as describe

below.

After administering the composition for measuring the gastric-emptying function of the present invention to a subject, the amounts of the isotope-labeled carbon dioxide ($^{13}\text{CO}_2$) excreted in expired air or carbon dioxide gas value $\Delta(\%)$ (the difference between the $\delta^{13}\text{C}$ values before and after the administration of the composition) were measured with lapse of time, and compared its excretion pattern with that of the standard control (from a healthy subject).

Lowering of the gastric-emptying function can also be detected from the initial speed of the $^{13}\text{CO}_2$ excreted in expired air. This method further shortens the total hours the subject has to be constrained. In this case, if the initial speed of the $^{13}\text{CO}_2$ excreted into expired air is slower than that of the standard control (the healthy subject), the gastric-emptying function is judged to be reduced.

As described above, one of major causes of dyspepsia (non-ulcerative upper-gastrointestinal tract syndrome) is alimentary tract motor dysfunction, in particular the lowering of the gastric-emptying function. Therefore, the above-explained method for measuring the gastric-emptying function can also be effectively employed for diagnosing dyspepsia, in particular dyspepsia mainly attributable to insufficiencies in the gastric-emptying function (for example, dysmotility-type dyspepsia). Accordingly, all explanations of the method for measuring the gastric-emptying function of the present invention can be applied to a dyspepsia diagnosis method. In this case, the above-described dyspepsia diagnostic agent is used instead of the composition for measuring the gastric-emptying function.

(V) Method for measuring pharmaceutical or therapeutic effect

By employing the method for measuring the gastric-emptying function described above, the pharmaceutical effect of the gastrointestinal medicine, in particular, a medicine that affects the alimentary tract motor function, or therapeutic

effect on each subject can be measured. Specifically, the measurement can be conducted by comparing the gastric-emptying functions measured using the composition for measuring the gastric-emptying function of the present invention before and
5 after administering to a subject a gastrointestinal medicine, in particular, a medicine relating to the alimentary tract motor function. This makes it possible to evaluate the pharmaceutical effect of the medicine itself. Furthermore, the therapeutic effect of the medicine on each subject can also be measured. As a
10 result, this method can be used as a means to suitably select a medicine applicable to each subject.

Examples of the medicines affecting the alimentary tract motor function include alimentary tract motor function improving agents, alimentary tract motor function accelerators,
15 and alimentary tract motor function activation agents (specifically, acetylcholine agonists, dopamine receptor antagonists, dopamine D₂ receptor antagonists, serotonin receptor agonists, opioid receptor agonist, Chinese medicines (Rikkunshi-to, Hange-shashin-to, and Annaka powder); alimentary tract motor
20 function inhibitors (anticholinergic agents, muscarinic receptor antagonists, etc.) and the like, which control the peristaltic movement of the stomach in an accelerating or inhibitory manner.

In this method, the subject may be a patient with dyspepsia, especially a patient whose dyspepsia is mainly caused
25 by a gastric motor dysfunction (a dysmotility-type dyspepsia patient). In this case, the pharmacotherapy effect on each dyspepsia patient can be measured, so that a suitable medicine, e.g., a medicine affecting the alimentary tract motor functions (alimentary tract motor function improving agents, alimentary
30 tract motor function accelerators, or alimentary tract motor function activation agents) can be selected. In other words, the method is effectively usable in measuring the pharmaceutical effect of a medicine for a dyspepsia patient, in particular, a medicine affecting the alimentary tract motor function, or
35 measuring the therapeutic effect of a medicine on a dyspepsia

patient. The above explanation of the method for measuring the gastric-emptying function can also be applied to the specific measuring method of the present invention.

5

EXAMPLES

The present invention is further explained with reference to Examples and Experimental Examples. However, the scope of the present invention is not limited to these Examples,
10 etc.

Example 1 Solution

2-¹³C uracil (molecular weight: 113.08; Cambridge Isotope Laboratory; 100 mg) was dissolved in 50 ml of 0.1N-NaOH/saline solution (adjusted), preparing a composition taking a
15 form of an aqueous solution (containing 2-¹³C uracil at a concentration of 20 μmol/ml).

Example 2 Granules

(1) Preparation of granules

2-¹³C uracil (Cambridge Isotope Laboratory, 20 g) and D-mannitol (Mannite; Kyowa Hakko Kogyo Co., Ltd., 380 g) were mixed,
20 introduced into a SampleMill (KIIWG-1F; Fuji Paudal Co., Ltd.) and then subjected to pulverization while mixing (pulverization conditions; the number of revolutions of the pulverization rotor: 12,800 rpm; the number of revolutions of the sample supply motor:
25 about 10 rpm; screen: 1 mm punch screen), thereby preparing a powder material. The thus-obtained powder material (200 g) was weighed, placed in a speed kneader (NSK-150; Okada Seiko Co., Ltd.), and then kneaded after adding 20 g of purified water. Subsequently, the thus-obtained wet powder was extruded using an
30 extrusion granulator equipped with a dome-shaped die having openings of φ1 mm (Dome Gran DG-1L; Fuji Paudal Co., Ltd.), and then dried at 60°C using a ventilation drier (SPHH-200; ESPEC Corp.). From the dried preparation, after passing through a sieve having openings of 1,400 μm but not through openings of 355 μm, 5
35 wt.% of 2-¹³C uracil-containing granules were obtained.

The particle diameters of the thus-obtained 5 wt.% of 2-¹³C uracil-containing granules were measured by vibration sieving (specifically, measurement apparatus: Robot Sifter RPS-95, Seishin Enterprise Co., Ltd.; vibration level: 5; shift time: 5 minutes; interpulse interval: 1 second). Table 1 shows the results.

Table 1

Particle Diameter	Content (wt.%)
Not less than 1400 μm	2.09
1000 μm to less than 1400 μm	7.29
850 μm to less than 1000 μm	22.07
710 μm to less than 850 μm	59.04
500 μm to less than 710 μm	8.99
355 μm to less than 500 μm	0.09
250 μm to less than 355 μm	0.00
150 μm to less than 250 μm	0.09
Less than 150 μm	0.34
Total	100.0

(2) Evaluation of solubility

Tap water (100 mL) was placed in a 200 mL beaker at room temperature, 2,000 mg of the above-obtained granules were added thereto while stirring at 200 rpm using a magnetic stirrer (RCN-7D; EYELA), and then the time until dissolution was measured by visual observation. When three minutes had passed after the addition of the granules, the remaining, undissolved preparation was observed by visual observation. The results show that the time before completion of the dissolution was 1 minute and 10 seconds, and that very little preparation remained after 3 minutes.

Example 3 Granules

2-¹³C uracil (Cambridge Isotope Laboratory, 20 g) and D-mannitol (Mannite; Kyowa Hakko Kogyo Co., Ltd.; 180 g) were mixed, introduced into a SampleMill (KIIWG-1F; Fuji Paudal Co., Ltd.) and then subjected to pulverization while mixing (pulverization conditions: the number of revolutions of the pulverization rotor: 12,800 rpm; the number of revolutions of the sample supply motor:

about 10 rpm; screen: 1 mm punch screen), thereby preparing a powder material. The thus-obtained powder material (144 g) was weighed, placed in a speed kneader (NSK-150; Okada Seiko Co., Ltd.), and then kneaded after adding 14.4 g of purified water. Subsequently, the thus-obtained wet powder was extruded using an extrusion granulator equipped with a dome-shaped die having openings of $\phi 1$ mm (Dome Gran DG-1L; Fuji Paudal Co., Ltd.), and then dried at 60°C using a ventilation drier (SPHH-201, ESPEC Corp.). From the dried preparation, after passing through a sieve having openings of 1,400 μm but not through openings of 355 μm , 10 wt.% of 2- ^{13}C uracil-containing granules were obtained.

Example 4 Tablets

2- ^{13}C uracil (Cambridge Isotope Laboratory, 100 g) and lactose (H.M.S, 60 g), corn starch (Cornstarch; Nihon Shokuhin Kako Co., Ltd.; 25 g), crystalline cellulose (CEOLUS PH301; Asahi Kasei Corporation; 10 g) and hydroxypropyl cellulose (HPC-L fine powder; Nippon Soda Co., Ltd.; 4 g) were placed in a speed kneader (NSK-150; Okada Seiko Co.; Ltd.) and mixed, and then kneaded after adding 40 g of purified water. Subsequently, the thus-obtained kneaded powder was granulated using a speed mill equipped with a 3 mm punch screen (ND-02; Okada Seiko Co., Ltd.), and then dried by a ventilation drier (SPHH-200; ESPEC Corp.) having a temperature of 70°C. The dried granules were subjected to sizing by passing through a No. 16 sieve. Magnesium stearate (Taihei Chemicals Limited, 1 g) was added to 199 g of the granules after sizing, giving granules for tablets. The resulting granules for tablets were formed into tablets each having a weight of 200 mg using a single-shot tableting machine equipped with a punch and die having a corner angle R of $\phi 8$ mm (No.2B; Kikusui Seisakusho Ltd.).

Example 5 Powder preparation

(1) Formation of powder preparation

2- ^{13}C uracil (Cambridge Isotope Laboratory, 20 g) and D-

mannitol (Mannite; Kyowa Hakko Kogyo Co., Ltd.; 180 g) were well mixed, placed in a SampleMill (SAM; Nara Machinery Co., Ltd. and subjected to pulverization while mixing (shape of the grinding vane: pin-type; number of revolutions of rotor: 4,000 rpm; 5 screen: 3 mm punch screen), producing powder preparations.

(2) Measurement of distribution of particle size

The particle size distribution of the powder preparations was measured using a dry particle size distribution measurement apparatus (LDSA-1500A, Tohnichi Computer Applications 10 Co., Ltd.; focal distance: 100 mm; number of averaging processes: 10 times; averaging interval: 5 milliseconds; and air pressure: 0.4 MPa). Based on the obtained particle size distribution, particle diameter at 10% (10% D (μm)), particle diameter at 50% (50% D (μm)), and particle diameter at 90% (90% D (μm)) were 15 calculated. Table 2 shows the results.

Table 2

10% D (μm)	50% D (μm)	90% D (μm)
5.74	14.95	56.58

As shown in Table 2, the resulting particles had very small diameters, and a satisfactory pulverization effect was 20 obtained.

Examples 6-10 and 11-15

A solution (Example 6), granules (Examples 7 and 8), tablets (Example 9) and powder preparations (Example 10) were 25 prepared in the same manner as in Examples 1 to 5, except that 2-¹³C thymine was used instead of 2-¹³C uracil. A solution (Example 11), granules (Examples 12 and 13), tablets (Example 14) and powder preparations (Example 15) were prepared in the same manner as in Examples 1 to 5, except that 2-¹³C cytosine was used instead 30 of 2-¹³C uracil.

Experimental Example 1

Rats (female, Wister rat, 8 weeks old) were

anesthetized with pentobarbital and subjected to laparotomy. Thereafter, each of the stomach, duodenum, jejunum and ileum was tied to form a loop in each of the digestive organs (e.g., the stomach, duodenum, jejunum, and ileum). Specifically, the stomach
5 was tied at the pylorus portion, the duodenum was tied at the pylorus portion and 20 cm below the pylorus portion, the jejunum was tied 10 cm below the Treitz' ligament and further 20 cm therebelow, and the ileum was tied 20 cm above the cecum and ileocecal portion.

10 Subsequently, an aqueous solution of 20 $\mu\text{mol/mL}$ 6- ^{14}C uracil (Moravek Biochemicals, Inc.) dissolved in distillation water was injected at 1 mL/kg to each alimentary tract loop using an injection needle, and the injection portion was sealed using an adhesive. Blood was collected 5, 10, 20, 30, 45 and 60 minutes
15 after the injection of the aqueous 6- ^{14}C uracil solution from the jugular vein using an injection syringe, and then transferred into a tube containing a serum separation agent. The thus-collected blood was centrifuged for 15 minutes at 3,000 rpm, obtaining serum. Subsequently, the radioactivity in the serum was
20 measured using a liquid scintillation counter. Table 1 shows the results.

As is clear from the results of Table 1, uracil is not absorbed or is barely absorbed in the stomach, but is absorbed in alimentary tracts including and below the duodenum (e.g., the
25 duodenum, jejunum, and ileum).

Experimental Example 2

2- ^{13}C uracil (Cambridge Isotope Laboratory) was dissolved in Britton-Robinson buffer solutions of differing pH
30 values (pH 2, 4, 6 and 8) to prepare a saturated solution. Each solution was subjected to high performance liquid chromatography, the amount of the 2- ^{13}C uracil dissolved in the solution was measured, and then the solubility (w/v%) was calculated. Table 3 shows the results.

Table 3

pH of buffer solution	Solubility (w/v%)
2.0	0.28
4.0	0.28
6.0	0.28
8.0	0.29

As is clear from the results, the solubility of uracil is almost unchanged depending on the pH of the solution (i.e., the alimentary tract pH).

Experimental Example 3

An aqueous solution containing 6-¹⁴C uracil (Moravek Biochemicals, Inc.) was orally administered to rats (female, Wister rats, 8 weeks old: n=3) and dogs (female, beagle, 11 kg: n=3) once in an amount of 20 µmol/kg under fasting. Up to 168 hours after the administration, the amount of the metabolite excreted in expired air, urine and feces were measured with a liquid scintillation counter using the radioactivity as its index, so that the accumulated excretion rate was obtained.

The granules containing 2-¹³C uracil (Cambridge Isotope Laboratory) (100 mg calculated as the amount of 2-¹³C uracil) prepared in Example 2 were orally administered to humans (12 cases), and then the rates of the metabolite excreted into urine and expired air were measured. The excretion rate to urine was expressed as the metabolite excretion rate (the accumulated excretion rate) in the urine pooled for 12 hours from the administration of 2-¹³C uracil, calculated based on the metabolite concentration in urine measured using an LC/MS/MS and the amount of urine. The excretion rate in expired air was measured based on the ¹³CO₂ concentration excreted in expired air (before administration, and 10, 20, 30, 40, 50, 60, 90 minutes and 2, 4, 6, 8, 12 hours after administration) using a GC/MS and the conversion equation of Ghooos, et al. (Ghooos YF, Maes BD, Geypens BJ, et al., Measurement of gastric-emptying function of solids by

means of a carbon-labeled octanoic acid breath test. Gastroenterology 1993; 104: 1640-7). FIG. 2 shows the results.

As shown in FIG. 2, the uracil administered to rats, dogs and humans was metabolized within 12 hours from administration, and a large amount, i.e., over 80%, thereof was excreted into expired air as carbon dioxide gas.

Experimental Example 4

"Propantheline", an anticholinergic agent, was intravenously administered to rats (female, Wister rat, 8 weeks old: n=3) in such an amount of 1 mg/kg to obtain model animals whose gastric-emptying function was decreased (a delayed gastric-emptying model). Five minutes after the administration of propantheline, 10 mg/kg of aqueous 2-¹³C uracil solution was orally administered to each rat, and expired air was collected by suction 5, 10, 20, 30, 40, 50 and 60 minutes after administration at the rate of 100 mL/60 sec, using an expired air collection device for rats. The ¹³CO₂ concentrations in the thus-collected expired air samples and expired air samples (pre) collected in the same manner before administration of 2-¹³C uracil were measured using a GC-MS (ABCA-G; Europe Scientific Ltd.). As a control experiment, 2-¹³C uracil (10 mg/kg) was administered in the same manner as described above to rats to which propantheline was not intravenously injected, and therefore having normal gastric-emptying function. The expired air samples were collected over time, and the ¹³CO₂ concentration in each expired air sample was measured.

FIG. 3 shows the change in the ¹³CO₂ concentration in expired air after administration of 2-¹³C uracil. In FIG. 3, the vertical axis indicates the $\Delta^{13}\text{C}$ value (%), which is the difference between the $\delta^{13}\text{C}$ values (%) (¹³CO₂/¹²CO₂ concentration ratio in expired air) in the expired air samples collected before and after administration of 2-¹³C uracil. The horizontal axis indicates the time (in minutes) collected expired air from the administration of 2-¹³C uracil. As is clear from FIG. 3, the ¹³CO₂

concentration ($\Delta^{13}\text{C}$ value (%)) excreted in expired air from the rats having reduced gastric-emptying function by administering propantheline was significantly lower than that of normal rats. From this result, it became clear that the gastric-emptying function can be measured based on the change of the $^{13}\text{CO}_2$ concentration in expired air after the administration of 2- ^{13}C uracil; and that by comparing the change of the $^{13}\text{CO}_2$ concentration in expired air after the administration of 2- ^{13}C uracil with that of controls having normal gastric-emptying function, drop or acceleration in the gastric-emptying function can be detected.

Experimental Example 5

The 2- ^{13}C uracil granules prepared in Example 2 were orally administered to human patients (on or before 20 days from the extraction of the stomach; a total of 20 cases, with 7 cases undergoing total gastric resection) who were suspected of having post-operative gastroparesis, in a 100 mg amount calculated as the 2- ^{13}C uracil amount. The expired air was collected 10, 20, 30, 40, 50 and 60 minutes after the administration, and the $^{13}\text{CO}_2$ concentration in each of the expired air samples including expired air samples (pre-) collected before administering 2- ^{13}C uracil was measured using a GC/MS. Subsequently, the change of the $^{13}\text{CO}_2$ concentration ($\Delta^{13}\text{C}$ (%)) in expired air was calculated. FIG. 4 shows the results.

As shown in FIG. 4, based on the expired air test of the present invention using the 2- ^{13}C uracil, the human patients (20 cases) can be divided into three groups: patients with normal gastric-emptying function (normal type: solid line), patients with decreased gastric-emptying function (delayed gastric-emptying type: dashed line), and patients with insufficient gastric-emptying function (dysfunction type: dotted line). Regarding these patients, 2- ^{13}C uracil concentrations in plasma were measured 20 minutes after the administration of 2- ^{13}C uracil. As shown in FIG. 5, patients with decreased gastric-emptying

function (delayed gastric emptying), and patients with insufficient gastric-emptying function exhibited lowered 2-¹³C uracil concentrations in plasma corresponding to the reduction of the gastric-emptying function. This indicates that expired air
5 test of the present invention using 2-¹³C uracil accurately reflects the gastric-emptying function.

Experimental Example 6 Accuracy of diagnosis evaluation

Granules of Example 2 (1 g each) were orally
10 administered to three healthy subjects (subjects A, B and C); their expired air samples were collected over time, and then the ¹³CO₂ concentration in the expired air was measured using a GC-MS analysis apparatus (ABCA-G; Europe Scientific Ltd.).

FIG. 6 shows the change in the ¹³CO₂ concentration in
15 expired air after the administration of the preparation. In FIG. 6, the vertical axis indicates the $\Delta^{13}\text{C}$ value (%), which is the difference between the $\delta^{13}\text{C}$ values (%) (¹³CO₂/¹²CO₂ concentration ratio in expired air) in the expired air samples collected before and after administration of 2-¹³C uracil. The horizontal axis
20 indicates the time (in minutes) collected expired air from the administration of the granules. As is clear from FIG. 6, when the granules prepared by mixing and pulverizing a powder material containing an isotope-labeled pyrimidine compound and a sugar and/or sugar alcohol as those prepared in Example 2 were used in
25 an expired air test, the individual subjects showed similar patterns of change in ¹³CO₂ concentration and the variance attributable to the differences in the individual subjects can be reduced. This result indicates that by measuring the gastric-emptying function using, as the index, the ¹³CO₂ concentration in
30 expired air 20 to 30 minutes after administering the above-mentioned preparation, dyspepsia can be diagnosed in a quick and accurate manner with little variance between subjects.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 shows the absorption of uracil ($6\text{-}^{14}\text{C}$ uracil) in the stomach, duodenum, jejunum, and ileum.

FIG. 2 shows in the body dynamics of uracil.

FIG. 3 shows the behavior with lapse of time of $^{13}\text{CO}_2$ excreted in expired air after orally administering $2\text{-}^{13}\text{C}$ uracil to delayed gastric-emptying model rats (pretreatment with propantheline) and rats with normal gastric-emptying function (control) in Experimental Example 3.

FIG. 4 shows the behavior with lapse of time of $^{13}\text{CO}_2$ excreted in expired air when $2\text{-}^{13}\text{C}$ uracil was administered to 20 patients suspected of having gastroparesis in Experimental Example 4.

FIG. 5 shows the $2\text{-}^{13}\text{C}$ uracil concentration in plasma 20 minutes after the administration of $2\text{-}^{13}\text{C}$ uracil in the patients divided into three groups based on the results shown in FIG. 4: i.e., patients with normal gastric-emptying function, patients with decreased gastric-emptying function, and patients with insufficient gastric-emptying function.

FIG. 6 shows the behavior with lapse of time of $^{13}\text{CO}_2$ excreted in expired air when the granules of Example 2 were orally administered to three healthy subjects (subjects A, B and C).

CLAIMS

1. A composition for measuring the gastric-emptying function comprising a pyrimidine compound as an active ingredient,
5 the pyrimidine compound being labeled with at least one isotope of C or O, converted into labeled CO₂ in the body, and excreted in expired air.

2. The composition for measuring the gastric-emptying
10 function according to claim 1, wherein the isotope is at least one member selected from the group consisting of ¹³C, ¹⁴C and ¹⁸O.

3. The composition for measuring the gastric-emptying function according to claim 1, wherein the pyrimidine compound is
15 uracil or thymine.

4. A dyspepsia diagnostic agent consisting of a composition of any one of claims 1 to 3.

20 5. The dyspepsia diagnostic agent according to claim 4, wherein the dyspepsia is caused by an insufficient gastric-emptying function.

6. A method for measuring the gastric-emptying function
25 comprising:

orally administering a composition of any one of claims 1 to 3 to a subject; and

measuring the amount or behavior of the labeled CO₂ excreted in expired air.

30

7. The method for measuring the gastric-emptying function according to claim 6, which comprises:

orally administering a composition of any one of claims 1 to 3 to a subject in whom a drop or acceleration in the
35 gastric-emptying function is suspected;

measuring the amount or behavior of the labeled CO₂ excreted in expired air; and

comparing the resulting amount or behavior of the labeled CO₂ with that of the labeled CO₂ excreted in expired air
5 collected from a healthy subject measured using the same composition.

8. A method for measuring the gastric-emptying function comprising:

10 using expired air collected from a subject who had orally taken a composition of any one of claims 1 to 3, as a test sample; and

measuring *in vitro* the amount of the labeled CO₂ in the test sample.

15

9. The method for measuring the gastric-emptying function according to claim 8, which comprises:

using expired air from a subject in whom a drop or acceleration in the gastric-emptying function is suspected and who
20 had orally taken a composition of any one of claims 1 to 3, as a test sample;

measuring *in vitro* the amount of the labeled CO₂ in the test sample; and

comparing the measured value with the amount of the
25 labeled CO₂ in expired air collected from a healthy subject who had taken the same composition.

10. A method for diagnosing dyspepsia comprising:

orally administering to a subject a dyspepsia
30 diagnostic agent of claim 4 or 5; and

measuring the gastric-emptying function based on the amount or behavior of the labeled CO₂ excreted in expired air.

11. The method for diagnosing dyspepsia according to
35 claim 10, which comprises:

orally administering a dyspepsia diagnostic agent of claim 4 or 5 to a subject in whom dyspepsia is suspected;

comparing the amount or behavior of the labeled CO₂ excreted in expired air collected from the subject with the amount
5 or behavior of the labeled CO₂ collected from a healthy subject measured using the same diagnostic agent; and

measuring the gastric-emptying function of the subject.

12. A method for diagnosing dyspepsia comprising:

10 using expired air collected from a subject who had orally taken a dyspepsia diagnostic agent of claim 4 or 5, as a test sample;

measuring *in vitro* the amount of the labeled CO₂ in the test sample; and

15 measuring the gastric-emptying function.

13. The method for diagnosing dyspepsia according to claim 12, which comprises:

using expired air collected from a subject in whom
20 dyspepsia is suspected and who had orally taken a dyspepsia diagnostic agent of claim 4 or 5, as a test sample;

measuring *in vitro* the amount of the labeled CO₂ in the test sample;

comparing the measured value with the amount of labeled
25 CO₂ in expired air collected from a healthy subject who had taken the same composition; and

measuring the gastric-emptying function.

14. The method for diagnosing dyspepsia according to
30 claim 10 or 12, wherein the dyspepsia is caused by an insufficient gastric-emptying function.

15. A method for measuring a gastrointestinal treatment effect on a subject comprising:

35 before and after conducting the gastrointestinal

treatment to a subject, orally administering a composition of any one of claims 1 to 3 or a diagnostic agent of claim 4 or 5 to the subject; and

5 comparing the amounts or behaviors of the labeled CO₂ excreted in expired air before and after the gastrointestinal treatment.

16. A method for measuring a pharmaceutical effect of a gastrointestinal medicine or a therapeutic effect on a subject
10 comprising:

before and after administering the gastrointestinal medicine to the subject, orally administering to the subject a composition of any one of claims 1 to 3 or a diagnostic agent of claim 4 or 5; and

15 comparing the amounts or behaviors of the labeled CO₂ excreted in expired air before and after the administration of the gastrointestinal medicine.

17. A method for measuring a gastrointestinal treatment
20 effect on a subject comprising:

using, as a test sample, expired air collected from a subject who had orally taken a composition of any one of claims 1 to 3 or a diagnostic agent of claim 4 or 5 before and after conducting the gastrointestinal treatment; and

25 comparing the amounts of the labeled CO₂ excreted in expired air before and after the gastrointestinal treatment.

18. A method for evaluating a pharmaceutical effect of a gastrointestinal medicine or a therapeutic effect on a subject
30 comprising:

using, as a test sample, expired air collected from a subject who had orally taken a composition of any one of claims 1 to 3 or a diagnostic agent of claim 4 or 5 before and after administering the gastrointestinal medicine to the subject; and

35 comparing the amounts of the labeled CO₂ excreted in

expired air before and after the administration of the gastrointestinal medicine.

19. The measuring method according to claim 16 or 18,
5 wherein the gastrointestinal medicine affects the alimentary tract motor function.

20. The measuring method according to claim 16 or 18,
which is used to measure the pharmaceutical effect of a
10 gastrointestinal medicine or the therapeutic effect on a patient suffering from dyspepsia caused by an insufficient gastric-emptying function.

21. A use of a pyrimidine compound for producing a
15 dyspepsia diagnostic agent, the pyrimidine compound being labeled with either or both isotopes of C and O, converted into an isotope-labeled CO₂ gas in the body, and then excreted in expired air.