

[DESCRIPTION]

[TITLE OF INVENTION]

EDIBLE OIL/FAT IN WHICH GLYCIDOL CONTENT AND GLYCIDOL FATTY
ACID ESTER CONTENT ARE REDUCED, AND METHOD FOR

5 MANUFACTURING SAID EDIBLE OIL/FAT

[Technical Field]

[0001]

The present invention relates to an edible oil/fat in which the content of glycidol
and glycidol fatty acid esters is reduced, and a method for manufacturing the same.

10 [Background Art]

[0002]

Glycidol is a compound in which an epoxy ring is formed at the 2- and
3-positions of a glycerol framework. In recent years, glycidol fatty acid esters in which
fatty acids are ester-bonded to glycidol have been detected in a plurality of types of
15 edible oils/fats. It has been reported that, particularly, the content of palm oils/fats and
refined edible oils/fats made from palm oils/fats as a raw material is relatively high
among oils/fats in which the presence of glycidol fatty acid esters is confirmed
(Non-Patent Literature 1).

[0003]

20 When a glycidol fatty acid ester is ingested in the body, there is a concern that it
will be hydrolyzed by the action of lipase present in the body and produce glycidol.
Glycidol is classified as “Group 2A” by the International Agency for Research on Cancer
(IARC). “Group 2A” is the group into which compounds that are probably
carcinogenic to humans are classified (Non-Patent Literature 2).

25 [0004]

Based on the above circumstances, it is desired to develop an oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced.

[0005]

Patent Literature 1 to 3 disclose methods for reducing the content of glycidol
5 and glycidol fatty acid esters. Glycidol and glycidol fatty acid esters tend to be produced during a deodorizing treatment, and Patent Literature 1 discloses a glyceride composition that is subjected to a low-temperature deodorizing treatment at 190 to 230°C.

[Citation List]

10 [Patent Literature]

[0006]

PCT International Publication No. WO 2016/121604

[Patent Literature 1]

Japanese Patent Laid-Open No. 2011-147436

15 [Patent Literature 2]

PCT International Publication No. WO 2019/038320

[Patent Literature 3]

PCT International Publication No. WO 2019/027315

[Non-Patent Literature]

20 [0007]

[Non-Patent Literature 1]

December 17, 2014, The results of examination of the content of 3-MCPD fatty acid ester and glycidol fatty acid ester in foods, Food Safety and Consumer Affairs Bureau, Ministry of Agriculture, Forestry and Fisheries (2012, 2013).

25 [Non-Patent Literature 2]

[Summary of Invention]

[Technical Problem]

[0008]

5 Patent Literature 2 and 3 disclose a process of refining a physically refined palm oil (RBDPO) from a crude palm oil (CPO), and do not describe a step of refining a physically refined palm oil (RBDPO).

[0009]

 It has been found that the method in Patent Literature 1 has a problem that a
10 decrease in the acid value of refined oil after deodorization is insufficient, and the refined oil flavor becomes oily.

[0010]

 In order to address the above problem, an objective of the present invention is to provide an edible oil/fat which has favorable flavor and color, and in which the content
15 of glycidol and glycidol fatty acid esters is reduced to a satisfactory level, and a method for manufacturing the same.

[Solution to Problem]

[0011]

 The inventors conducted extensive studies, and as a result, found that, when a
20 raw oil/fat having a 3-chloro-1,2-propanediol (3-MCPD) concentration of a certain level or less is subjected to a step of performing an acid-induced contacting process during a decolorization work step, the flavor and color become favorable and the content of glycidol and glycidol fatty acid esters is reduced, and completed the present invention.

[0012]

25 Specifically, the present invention is as follows.

(1) A method for manufacturing an edible oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced, and in which, in a step of decolorizing and deodorizing an oil/fat containing a palm-based oil/fat using a physically refined palm oil (RBDPO) as a raw material, the following raw oil/fat is used, a decolorization work step
5 includes a step of adding an acid to an aqueous solution, and the following edible oil/fat is obtained after the step of decolorizing and deodorizing:

a raw oil/fat: an oil/fat containing a palm-based oil/fat and having a 3-MCPD concentration of 1.5 mg/kg or less

an edible oil/fat after the step of decolorizing and deodorizing: the content of glycidol
10 and glycidol fatty acid esters is 1.0 mg/kg or less in an equal amount of glycidol.

(2) The method for manufacturing an edible oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced according to (1), wherein 0.01 to 10 weight% of the following aqueous solution as an acid is added.

· a citric acid and/or phosphoric acid aqueous solution having a concentration of 1 to 85
15 weight%.

(3) The method for manufacturing an edible oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced according to (1) or (2), wherein the deodorizing step is performed under the following conditions:

· a degree of vacuum of 100 to 800 Pa
20 · an amount of water vapor used of 0.5 to 5 weight%
· a deodorizing temperature of 230 to 260°C

(4) A method for reducing the content of glycidol and glycidol fatty acid esters in an edible oil/fat by the method for manufacturing an edible oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced according to any one of (1) to (3).

25 [Advantageous Effects of Invention]

[0013]

According to the present invention, it is possible to provide an edible oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced.

The content of glycidol and glycidol fatty acid esters in the obtained deodorized oil/fat can be reduced to 1.0 mg/kg or less, and preferably 0.5 mg/kg or less in an equal amount of glycidol.

[Description of Embodiments]

[0014]

Hereinafter, the present invention will be described in detail.

10 [0015]

As the raw oil/fat that can be used in the present invention, an oil/fat containing a palm-based oil/fat having a 3-MCPD concentration of 1.5 mg/kg or less can be used. When a raw oil/fat having a 3-MCPD concentration exceeding 1.5 mg/kg is used, the effect of the present invention may not be obtained.

15 [0016]

A palm oil manufacturing step includes a step of refining a crude palm oil (CPO) in which a physically refined palm oil (RBDPO) is obtained from a crude palm oil (CPO) obtained by squeezing the fruit pulp of oil palm through a refining step such as degumming, deoxidizing, decolorizing, deodorizing, and dewaxing, and a step of refining from the physically refined palm oil (RBDPO) in which the physically refined palm oil (RBDPO) is additionally subjected to refining such as decolorizing and deodorizing.

[0017]

The manufacturing method of the present invention is applied to a step of manufacturing from a physically refined palm oil (RBDPO).

25 [0018]

In the present invention, as the palm-based oil/fat, a physically refined palm oil, and an oil fraction obtained from a physically refined palm oil as a raw material can be used. Examples of oil fractions made from a physically refined palm oil as a raw material include palm olein, palm superolein, palm stearin, and palm mid fractions.

5 [0019]

In the present invention, as the palm-based oil/fat, an oil/fat (hereinafter referred to as (B)) processed using a palm-based oil/fat (hereinafter referred to as (A)) can be used. Examples of processing methods include hardening, separation using a hardened oil as a raw material, hardening using a fractionated oil as a raw material, and transesterification.

10 As the raw material of (B), in addition to (A), oils/fats of (B) can also be used, and (A) and/or (B) is contained as an essential component, but otherwise, other edible oils/fats may be contained. Examples of oils/fats that can be used include plant oils/fats such as soybean oil, rapeseed oil, corn oil, cottonseed oil, peanut oil, sunflower oil, rice bran oil, safflower oil, olive oil, sesame oil, medium-chain fatty acid-bound oil/fat
15 (MCT), coconut oil, and palm kernel oil, animal fats such as milk fat, beef tallow, and lard, and their hardened oils, fractionated oils, hardened fractionated oils, and separately hardened oils, processed oils/fats subjected to transesterification or the like, and mixed oils/fats thereof.

[0020]

20 In the present invention, the raw oil/fat contains the above palm-based oil/fat as an essential component, but it may contain other edible oils/fats in addition to the palm-based oil/fat. Examples of oils/fats that can be used include plant oils/fats such as soybean oil, rapeseed oil, corn oil, cottonseed oil, peanut oil, sunflower oil, rice bran oil, safflower oil, olive oil, sesame oil, medium-chain fatty acid-bound oil/fat (MCT),
25 coconut oil, and palm kernel oil, animal fats such as milk fat, beef tallow, and lard, and

their hardened oils, fractionated oil, hardened fractionated oil, and separately hardened oil, processed oils/fats subjected to transesterification or the like, and mixed oils/fats thereof.

[0021]

5 In the present invention, the step of decolorizing and deodorizing includes all a series of steps in the manufacturing step, and is not limited to the manufacturing step in a reaction device such as a decolorization can or a deodorizing tower. The step of decolorizing and deodorizing is composed of a decolorization work step and a deodorizing work step.

10 In the present invention, the decolorization work step includes a decolorization preparation step in which a raw oil/fat is transferred to a reaction device (decolorization can) that performs a decolorization step, a decolorization step that is performed under a reduced pressure by adding a processing aid such as decolorized white clay to the decolorization can and heating the decolorization can, and a decolorization filtration step
15 in which cooling is performed as necessary after decolorization and white clay is filtered off, and is composed of a series of steps: the above decolorization preparation step, decolorization step, and decolorization filtration step.

 In the present invention, the deodorizing work step includes a deodorization preparation step in which a decolorizing oil after decolorization filtration is transferred to
20 a deodorizing container (deodorizing tower), a deodorizing step in the deodorizing tower, and a deodorizing post-treatment step including cooling, addition of an antioxidant, and the like, and is composed of a series of steps: the above deodorization preparation step, deodorizing step, and deodorizing post-treatment step.

[0022]

In the present invention, in the step of decolorizing and deodorizing an oil/fat containing a palm-based oil/fat, the decolorization work step includes a step of adding an acid to an aqueous solution. Preferably, a step of adding an acid to an aqueous solution is included immediately before the decolorization step in the decolorization work step.

- 5 Inclusion of the step of adding an acid to an aqueous solution immediately before the decolorization step in the decolorization work step means that no other step is provided between the step of adding an acid to an aqueous solution and the decolorization step. Examples of other steps include an alkaline deoxidizing step, a washing step, a dehydration step, a transesterification reaction step, and a deodorizing step.

10 [0023]

Examples of acids used in the present invention include organic acids and/or inorganic acids. Examples of organic acids include at least one organic acid selected from the group consisting of ascorbic acid, erythorbic acid, lactic acid, tartaric acid, citric acid, gallic acid, phosphoric acid, and malic acid, and salts thereof, and derivatives thereof. As the inorganic acid, it is preferable to use phosphoric acids, and examples thereof include orthophosphoric acid, diphosphoric acid (pyrophosphoric acid), and metaphosphoric acid.

[0024]

- As a preferable form of the present invention, an aqueous solution containing citric acid and/or phosphoric acid as an acid is used. Preferably, the concentration of the citric acid and/or phosphoric acid aqueous solution to be added is 1 to 85 weight%, and the amount of the citric acid and/or phosphoric acid aqueous solution added is 0.01 to 10 weight%. Within the above range, no particular conditions are designated, except that the amount of the citric acid and/or phosphoric acid aqueous solution added is more preferably 0.01 to 5 weight%, still more preferably 0.05 to 1 weight%, and yet more
- 20
- 25

preferably 0.05 to 0.5 weight% within the above range. If the amount of the citric acid and/or phosphoric acid aqueous solution added is less than 0.01 weight%, the amount of glycidol and glycidol fatty acid esters reduced may be insufficient. On the other hand, if the amount thereof exceeds 10 weight%, the load on the facility may increase.

5 [0025]

The deodorizing step is preferably performed at a deodorizing temperature of 230 to 260°C, a degree of vacuum 100 to 800 Pa, and an amount of water vapor used of 0.5 to 5 weight%. If the deodorizing temperature is lower than 230°C, this is not preferable because the flavor and color of the edible oil/fat after deodorization may
10 deteriorate. If the deodorizing temperature exceeds 260°C, this is not preferable because the content of glycidol and glycidol fatty acid esters may increase. The degree of vacuum is more preferably 200 to 800 Pa, and still more preferably 200 to 700 Pa. If the degree of vacuum exceeds 800 Pa, this is not preferable because the flavor of the edible oil/fat after deodorization may deteriorate. In addition, if the amount of water
15 vapor used is 0.5 weight% or less, similarly, this is not preferable because the flavor of the edible oil/fat is worse after deodorization. On the other hand, if the amount of water vapor used exceeds 5 weight%, this is not preferable because the load on the facility may increase. The deodorizing time is preferably 30 to 120 minutes, and particularly preferably 60 minutes to 90 minutes. If the deodorizing time is shorter than 30 minutes,
20 this is not preferable because the flavor of the edible oil/fat after deodorization may deteriorate. If the deodorizing time exceeds 120 minutes, this is not preferable because the content of glycidol and glycidol fatty acid esters may increase.

[0026]

Examples of deodorizing devices used in the present invention include vacuum
25 water vapor distillation devices, which may be of a batch type, a semi-continuous type, a

continuous type or the like. If the amount of the oil/fat to be treated is small, it is preferable to use a batch type device, and if the amount thereof is large, it is preferable to use a semi-continuous type or continuous type device.

As the semi-continuous type device, for example, a Gardler type deodorizing
5 device composed of a deodorizing tower including several trays can be used. This device performs processing by supplying an oil/fat from the upper part, making contact between the oil/fat on the tray and water vapor for an appropriate time, and then lowering the oil/fat to the lower tray, and moving it while intermittently lowering it one batch at a time. As the continuous type device, a thin film deodorizing device or the like that can
10 bring a thin film-like oil/fat into contact with water vapor can be used. It is preferable to use a semi-continuous type device because the temperature and the amount of water vapor used can be managed.

[0027]

In the present invention, the content of glycidol and glycidol fatty acid esters is
15 measured by the following method.

[0028]

0.3 mL of isooctane is added to 100 mg of an oil/fat sample, and the oil/fat sample is dissolved. 3 mL of a sodium bromide solution containing 90 U/mL lipase is put into a test tube, and the test tube is shaken with a shaker for 30 minutes. After the
20 test tube is heated at 80°C for 10 minutes, the test tube is allowed to cool or water-cooled until it reaches room temperature. 50 µL of a 2.0 µg/mL internal standard mixed solution and 3 mL of hexane are put into the test tube, and the lid is closed. The test tube is stirred with a Vortex mixer and the centrifuge, and an aqueous layer is then transferred to a new test tube with a Pasteur pipette. 3 mL of hexane is added to the test
25 tube to which the aqueous layer has been transferred, the lid is closed, stirring is

performed with a Vortex mixer, the centrifuge is performed and an organic layer is then removed. 20 μ L of a phenylboronic acid solution and 3 mL of hexane are added to the test tube, and the lid is closed. The test tube is shaken with a shaker for 10 minutes. The organic layer is transferred to a new test tube with a Pasteur pipette. The organic
5 layer in the new test tube is concentrated to 0.5 to 0.8 mL with nitrogen gas, and analyzed through GC-MS.

Here, for GC-MS, quantification is performed using molecular ion peaks m/z 147 and m/z 150.

Examples

10 [0029]

Hereinafter, the present invention will be described in more detail with reference to examples. Here, in the examples, % is based on weight.

[0030]

As the physically refined palm oil, a physically refined palm oil (RBDPO)
15 obtained by degumming, decolorizing, and deodorizing a crude palm oil (CPO) with a phosphoric acid aqueous solution was used.

[0031]

(Example 1)

A physically refined palm oil having a 3-MCPD concentration of 0.31 mg/kg
20 was subjected to a decolorizing-deodorizing step. 0.1 weight% of a 50% citric acid aqueous solution based on the weight of the oil/fat was added to a physically refined palm oil as a raw material. 1.5 weight% of activated clay based on the weight of the oil/fat was added, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was
25 completed, the oil/fat from which the activated clay was removed was deodorized under

conditions of 230°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes.

After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0032]

(Example 2)

5 A physically refined palm oil having a 3-MCPD concentration of 0.45 mg/kg was subjected to a decolorizing-deodorizing step. 0.1 weight% of a 85% phosphoric acid aqueous solution based on the weight of the oil/fat was added to a physically refined palm oil as a raw material. The activated clay is set to be 1.5 weight% based on the weight of the oil/fat and a decolorization treatment was performed under conditions of
10 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from which the activated clay was removed was deodorized under conditions of 230°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0033]

15 (Example 3)

 An oil/fat containing a physically refined palm oil having an iodine value of 53, a physically refined palm fractionated oil having an iodine value of 31, and a high erucic acid rapeseed extremely hardened oil mixed at a ratio of 44:57:3 was subjected to a transesterification reaction using 0.2 weight% sodium methoxide as a catalyst. The
20 reaction temperature was set to 80°C, and the reaction time was set to 30 minutes. After the reaction was completed, 5 weight% of a 50% citric acid aqueous solution was added to the oil/fat stirred at a rotational speed of 5,000 rpm using a homo mixer, and the mixture was brought into contact for 1 minute. Then, 20 weight% of warm water at 80°C was added, the mixture was treated for 5 minutes, and the oil/fat was washed with
25 water. The oil/fat washed with water was centrifuged, the aqueous phase was removed,

and a dehydration treatment was then performed under conditions of 110°C and 1,330 Pa for 10 minutes. The 3-MCPD concentration in the transesterified oil obtained in this manner was 0.14 mg/kg. This transesterified oil was subjected to a decolorizing-deodorizing step. 0.1 weight% of a 50% citric acid aqueous solution based on the weight of the oil/fat was added to a physically refined palm oil as a raw material, and additionally, 1.5 weight% of activated clay based on the weight of the oil/fat was added, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from which the activated clay was removed was deodorized under conditions of 250°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0034]

(Example 4)

A physically refined palm oil having a 3-MCPD concentration of 1.01 mg/kg was subjected to a decolorizing-deodorizing step. 0.1 weight% of a 50% citric acid aqueous solution based on the weight of the oil/fat was added to a physically refined palm oil as a raw material. 1.5 weight% of activated clay based on the weight of the oil/fat was added, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from which the activated clay was removed was deodorized under conditions of 250°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0035]

(Comparative Example 1)

A physically refined palm oil having a 3-MCPD concentration of 2.12 mg/kg was subjected to a decolorizing-deodorizing step. 0.1 weight% of a 50% citric acid aqueous solution based on the weight of the oil/fat was added to a physically refined palm oil as a raw material. 1.5 weight% of activated clay based on the weight of the oil/fat was added, and a decolorization treatment was performed under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from which the activated clay was removed was deodorized under conditions of 230°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

10 [0036]

(Comparative Example 2)

A physically refined palm oil having a 3-MCPD concentration of 0.39 mg/kg was subjected to a decolorizing-deodorizing step. 1.5 weight% of activated clay based on the weight of the oil/fat was added to a physically refined palm oil as a raw material, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from which the activated clay was removed was deodorized under conditions of 230°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

20 [0037]

(Comparative Example 3)

A physically refined palm oil having a 3-MCPD concentration of 0.22 mg/kg was subjected to a decolorizing-deodorizing step. 0.1 weight% of a 50% citric acid aqueous solution based on the weight of the oil/fat was added to a physically refined palm oil as a raw material. 1.5 weight% of activated clay based on the weight of the

oil/fat was added, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from which the activated clay was removed, and a deodorizing treatment was performed under conditions of 200°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0038]

(Comparative Example 4)

A physically refined palm oil having a 3-MCPD concentration of 1.01 mg/kg was subjected to a decolorizing-deodorizing step. 1.5 weight% of activated clay based on the weight of the oil/fat was added, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, 0.1 weight% of a 50% citric acid aqueous solution based on the weight of the oil/fat was added to the oil/fat from which the activated clay was removed, and immediately, a deodorizing treatment was performed under conditions of 250°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0039]

(Comparative Example 5)

A physically refined palm oil having a 3-MCPD concentration of 1.01 mg/kg was subjected to a decolorizing-deodorizing step. 0.05 weight% of citric acid powder based on the weight of the oil/fat was added to a physically refined palm oil as a raw material. 1.5 weight% of activated clay based on the weight of the oil/fat was added, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from

which the activated clay was removed was deodorized under conditions of 250°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0040]

5 (Comparative Example 6)

A physically refined palm oil having a 3-MCPD concentration of 1.01 mg/kg was subjected to a decolorizing-deodorizing step. 1.5 weight% of activated clay based on the weight of the oil/fat was added, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the
10 decolorization was completed, 0.05 weight% of citric acid powder based on the weight of the oil/fat was added to the oil/fat from which the activated clay was removed, and immediately, a deodorizing treatment was performed under conditions of 250°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

15 [0041]

(Comparative Example 7)

A physically refined palm oil having a 3-MCPD concentration of 1.17 mg/kg was subjected to a decolorizing-deodorizing step. 0.1 weight% of a 50% citric acid aqueous solution was added to the oil/fat stirred at a rotational speed of 5,000 rpm using
20 a homo mixer, and the mixture was brought into contact for 1 minute. Then, 20 weight% of warm water at 80°C was added, the mixture was treated for 5 minutes, and the oil/fat was washed with water. The oil/fat washed with water was centrifuged, the aqueous phase was removed, and a dehydration treatment was then performed under conditions of 110°C and 1,330 Pa for 10 minutes. 1.5 weight% of activated clay based
25 on the weight of the oil/fat was added, and a decolorization treatment was performed

with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from which the activated clay was removed was deodorized under conditions of 250°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes. After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0042]

(Comparative Example 8)

An oil/fat containing a physically refined palm oil having an iodine value of 53, a physically refined palm fractionated oil having an iodine value of 31, and a high erucic acid rapeseed extremely hardened oil mixed at a ratio of 44:57:3 was subjected to a transesterification reaction using a 0.2 weight% of sodium methoxide as a catalyst. The reaction temperature was set to 80°C, and the reaction time was set to 30 minutes. After the reaction was completed, 5 weight% of a 50% citric acid aqueous solution was added to the oil/fat stirred at a rotational speed of 5,000 rpm using a homo mixer, and the mixture was brought into contact for 1 minute. Then, 20 weight% of warm water at 80°C was added, the mixture was treated for 5 minutes, and the oil/fat was washed with water. The oil/fat washed with water was centrifuged, the aqueous phase was removed, and a dehydration treatment was then performed under conditions of 110°C and 1,330 Pa for 10 minutes. The 3-MCPD concentration in the transesterified oil obtained in this manner was 0.14 mg/kg. This transesterified oil was subjected to a decolorizing-deodorizing step. 1.5 weight% of activated clay based on the weight of the oil/fat was added, and a decolorization treatment was performed with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the decolorization was completed, the oil/fat from which the activated clay was removed was deodorized under

conditions of 250°C, 266 Pa, and an amount of vapor used of 3 weight% for 90 minutes.
After the deodorization, glycidol fatty acid ester contained in the oil/fat was analyzed.

[0043]

(Comparative Example 9)

5 A physically refined palm oil having a 3-MCPD concentration of 1.01 mg/kg
was subjected to a decolorizing-deodorizing step. 1.5 weight% of activated clay based
on the weight of the oil/fat was added, and a decolorization treatment was performed
with stirring under conditions of 110°C and 1,330 Pa for 10 minutes. After the
decolorization was completed, the oil/fat from which the activated clay was removed was
10 deodorized under conditions of 250°C, 266 Pa, and an amount of vapor used of 3
weight% for 90 minutes. After the deodorization, 0.1 weight% of a 50% citric acid
aqueous solution was added while the vacuum was maintained at 1,330 Pa. Finally,
glycidol fatty acid ester contained in the oil/fat was analyzed.

[0044]

15 Table 1 shows the analysis results of the oils/fats obtained in Examples 1 to 4
and Comparative Examples 1 to 9 and the evaluation results of the flavor. The flavor
was evaluated by a plurality of people, the score was 5 for the best flavor (tasteless and
odorless), and the score decreased as the flavor worsened in the evaluation. The score
of the flavor in the table indicates an average value of the scores scored by each evaluator.
20 3 points or more was satisfactory, and less than 3 points was unsatisfactory.

[0045]

[Table 1]

	3-MCPD concentration in raw oil/fat (mg/kg)	Equal amount of glycidol after refining (mg/kg)	Flavor after refining
Example 1	0.31	0.23	4.6
Example 2	0.45	0.35	4.4
Example 3	0.14	0.78	4.6
Example 4	1.01	0.38	4.8
Comparative Example 1	2.12	1.45	4.6
Comparative Example 2	0.39	1.39	3.8
Comparative Example 3	0.22	0.10	2.8
Comparative Example 4	1.01	1.02	4.4
Comparative Example 5	1.01	1.31	4.4
Comparative Example 6	1.01	1.01	4.6
Comparative Example 7	1.17	1.67	4.5
Comparative Example 8	0.16	1.27	4.5
Comparative Example 9	1.01	1.25	4.6

[0046]

As shown in Table 1, in examples in which a palm oil having a 3-MCPD

- 5 concentration in the raw material of 1.5 mg/kg or less was used as a raw material, and a citric acid and phosphoric acid aqueous solution was added, the flavor was favorable, and the amount of glycidol and glycidol fatty acid esters could be reduced to 1.0 mg/kg or less in an equal amount of glycidol. The deodorizing temperature in this case was 230 to 260°C.

10 [0047]

In Comparative Example 1, the 3-MCPD concentration in the palm oil as a raw material was 1.5 mg/kg or more, and in this case, an equal amount of glycidol after deodorization was 1.45 mg/kg.

In Comparative Example 2, no citric acid aqueous solution and/or phosphoric acid aqueous solution was added during decolorization, but an equal amount of glycidol in this case was 1.39 mg/kg.

In Comparative Example 3, the deodorizing temperature was set to 200°C, but in this case, the flavor was 2.8 points.

In Comparative Example 4 in which a citric acid aqueous solution was added immediately before deodorization, Comparative Example 5 and Comparative Example 6 in which citric acid powder was used, and Comparative Examples 4 to 9 in which the timing at which the citric acid aqueous solution was added was different from that of the examples, an equal amount of glycidol after refining could not be reduced to 1.0 mg/kg or less.

[0048]

As the physically refined palm oil used in the present invention, a physically refined palm oil degummed with phosphoric acid was used as in Patent Literature 2 and 3, but in comparative examples that did not satisfy the formulation of the present invention, the content of glycidol and glycidol fatty acid esters could not be reduced.

[0049]

The above results are summarized in Table 2. An edible oil/fat having a favorable flavor and in which the content of glycidol and glycidol fatty acid esters was reduced to 1.0 mg/kg or less in an equal amount of glycidol could be obtained only when a citric acid aqueous solution and/or phosphoric acid aqueous solution was added during the decolorization work step, a raw material having a 3-MCPD concentration in a raw oil

of 1.5 mg/kg or less was used, and deodorizing was performed at a deodorizing temperature of 230 to 260°C.

[0050]

[Table 2]

	Refining conditions			Quality of refined oil	
	3-MCPD concentration in raw oil/fat was 1.5 mg/kg or less	Add acid to aqueous solution during decolorization work step	Deodorizing temperature was 230 to 260°C	Glycidol concentration	Flavor
Example	O	O	O	O	O
Comparative Example 1	x	O	O	x	O
Comparative Example 2	O	x	O	x	O
Comparative Example 3	O	O	x	O	x
Comparative Examples 4 to 9	O	x	O	x	O

5 [Industrial Applicability]

[0051]

According to the present invention, it is possible to manufacture an oil/fat in which the content of glycidol and glycidol fatty acid esters, which may adversely affect human health, is reduced without impairing the flavor.

[CLAIMS]

[Claim 1]

A method for manufacturing an edible oil/fat in which a content of glycidol and glycidol fatty acid esters is reduced, and in which, in a step of decolorizing and

5 deodorizing an oil/fat containing a palm-based oil/fat using a physically refined palm oil (RBDPO) as a raw material, the following raw oil/fat is used, a decolorization work step includes a step of adding an acid to an aqueous solution, and the following edible oil/fat is obtained after the step of decolorizing and deodorizing:

a raw oil/fat:

10 an oil/fat containing a palm-based oil/fat and having a 3-MCPD concentration of 1.5 mg/kg or less,

an edible oil/fat after the step of decolorizing and deodorizing:

the content of glycidol and glycidol fatty acid esters is 1.0 mg/kg or less in an equal amount of glycidol.

15 [Claim 2]

The method for manufacturing an edible oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced according to claim 1,

wherein 0.01 to 10 weight% of the following aqueous solution as an acid is added:

20 a citric acid and/or phosphoric acid aqueous solution having a concentration of 1 to 85 weight%.

[Claim 3]

The method for manufacturing an edible oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced according to claim 1 or 2,

25 wherein the deodorizing step is performed under the following conditions:

a degree of vacuum of 100 to 800 Pa,
an amount of water vapor used of 0.5 to 5 weight%,
a deodorizing temperature of 230 to 260°C.

[Claim 4]

- 5 A method for reducing a content of glycidol and glycidol fatty acid esters in an edible oil/fat by the method for manufacturing an edible oil/fat in which the content of glycidol and glycidol fatty acid esters is reduced according to any one of claims 1 to 3.