

DESCRIPTION

DIHYDROPYRIDAZINE-3,5-DIONE DERIVATIVE

TECHNICAL FIELD

[0001] The present invention relates to a dihydropyridazine-3,5-dione derivative or a salt thereof, or a solvate of the compound or the salt. The present invention also relates to a pharmaceutical drug, a pharmaceutical composition, a sodium-dependent phosphate transporter inhibitor, and a preventive and/or therapeutic agent for hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, containing the dihydropyridazine-3,5-dione derivative or the salt thereof, or the solvate of the compound or the salt as an active ingredient, and a method for prevention and/or treatment.

BACKGROUND ART

[0002] Phosphorus is found in every cell and makes up 1% of a person's body weight. This element plays an essential role in sustaining life, such as the energy metabolism of cells. The concentration of phosphorus in blood is determined by absorption from the gastrointestinal tract and excretion from the kidney as well as bone formation and bone resorption and adjusted to a constant level. The phosphorus absorption in the gastrointestinal tract is performed mainly by a sodium-dependent phosphate transporter NaPi-IIb (SLC34A2) (Non-Patent Documents 1 and 2). Phosphorus in blood is filtered by renal glomerulus and reabsorbed in necessary amounts mainly by NaPi-IIa (SLC34A1) and NaPi-IIc (SLC34A3) in the renal tubule (Non-Patent Documents 1 and 3). The kidney plays a very important role in regulating phosphorus *in vivo*. In end-stage renal failure patients and dialysis patients with impaired renal functions, phosphorus accumulates in the body, resulting in a rise in phosphorus concentration in blood, i.e., hyperphosphatemia.

[0003] Hyperphosphatemia brings about the calcification of soft tissues. Particularly, vascular calcification is considered responsible for the dysfunction of the heart, leading to the death of the patient. Hyperphosphatemia also brings about the hypersecretion of parathyroid hormones, i.e., secondary hyperparathyroidism, and causes bone lesions. Thus,

hyperphosphatemia is viewed as a problematic factor that deteriorates the prognosis and QOL of end-stage renal failure patients and dialysis patients.

[0004] In the current treatment of hyperphosphatemia, phosphorus adsorbents are used for the purpose of suppressing phosphorus absorption in the gastrointestinal tract. Nonmetallic polymer adsorbents typified by sevelamer hydrochloride, calcium salt preparations typified by precipitated calcium carbonate, or metallic adsorbents typified by lanthanum carbonate are used as the phosphorus adsorbents. These adsorbents, however, have each been reported to have adverse reactions such as gastrointestinal disorders including constipation and diarrhea, hypercalcemia caused by a rise in serum calcium concentration, and *in vivo* metal accumulation. In addition, these adsorbents require a daily intake of a few grams and therefore present noncompliance problems. Accordingly, there is a strong demand for the development of novel hyperphosphatemia therapy improved in terms of these problems of the phosphorus adsorbents.

[0005] The inhibition of NaPi-IIb, which plays a major role in phosphorus absorption in the gastrointestinal tract, may suppress phosphorus absorption in the gastrointestinal tract, as with the phosphorus adsorbents, to decrease phosphorus concentration in blood (Non-Patent Documents 2 and 4). Also, PiT-1 (SLC20A1) and PiT-2 (SLC20A2), which are sodium-dependent phosphate transporters like NaPi-IIb, are partially responsible for phosphorus absorption in the gastrointestinal tract (Non-Patent Documents 1 and 6). Thus, a compound that inhibits NaPi-IIb, PiT-1, and PiT-2 can be expected to produce a stronger phosphorus absorption inhibitory effect and decrease phosphorus concentration in blood, compared with an inhibitor for only NaPi-IIb. Meanwhile, the suppression of phosphorus absorption by the inhibition of these sodium-dependent phosphate transporters is based on the mechanism of action different from that of the phosphorus adsorbents currently used. Thus, the sodium-dependent phosphate transporter inhibitor can be expected to serve as a novel preventive and/or therapeutic agent for hyperphosphatemia in place of the conventional phosphorus adsorbents. The sodium-dependent phosphate transporter inhibitor is further expected to exert preventive and/or therapeutic effects on secondary hyperparathyroidism and chronic

renal failure by decreasing phosphorus concentration in blood (Non-Patent Document 5). NTX1942 (Patent Document 1) and compounds described in Patent Documents 2 and 4 have been reported so far as NaPi-IIb inhibitors. Also, a compound having a pyridazine skeleton has been reported in Patent Document 3 which makes mention about the treatment of anemia, ischemia, and hypoxia by the HIF hydroxylase inhibitory activity of the compound.

CITATION LIST

PATENT DOCUMENTS

[0006] Patent Document 1: WO2012/006475

Patent Document 2: WO2011/136269

Patent Document 3: WO2011/048611

Patent Document 4: WO2013/062065

NON-PATENT DOCUMENTS

[0007] Non-Patent Document 1: J Pharm Sci. 2011 Sep; 100 (9): 3719-30;

Non-Patent Document 2: J Am Soc Nephrol. 2009 Nov; 20 (11): 2348-58;

Non-Patent Document 3: Pflugers Arch. 2004 Feb; 447(5): 763-7;

Non-Patent Document 4: Am J Physiol Renal Physiol. 2011 Nov; 301 (5): F1105-13;

Non-Patent Document 5: Kidney Int. 2003 Nov; 64 (5): 1653-61; and

Non-Patent Document 6: Mol Aspects Med. 2013 Apr-Jun; 34 (2-3): 386-95.

SUMMARY OF INVENTION

TECHNICAL PROBLEM

[0008] There is a strong demand for the development of novel hyperphosphatemia therapy improved in terms of these problems of the existing phosphorus adsorbents.

SOLUTION TO PROBLEM


[0009] The present inventors have conducted diligent studies in light of the problems described above and consequently completed the present invention by finding for the first time that a compound represented by the formula (I) shown below, which largely differ in chemical structure from NaPi-IIb inhibitors known in the art, has an excellent NaPi-IIb

inhibitory effect, PiT-1 inhibitory effect, and/or PiT-2 inhibitory effect, is useful in the prevention and/or treatment of hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, and has excellent drug efficacy on these diseases. Specifically, one aspect of the present invention provides compounds shown below or salts thereof, or solvates of the compounds or the salts.

[0010] One embodiment of the present invention provides the following compounds (1-1) to (1-55):

[0011] (1-1) A compound represented by the formula (I) or a salt thereof, or a solvate of the compound or the salt:

[0012] [Formula 1]



[0013] wherein R^1 , R^4 , and R^5 are as defined in any one of the following (1) to (3):

(1) R^1 is a hydrogen atom or C_{1-10} alkyl;

R^4 is a hydrogen atom, C_{1-4} alkyl optionally substituted with one or more substituents R_f , C_{6-10} aryl optionally substituted with one or more substituents R_g , (C_{1-6} alkyl)carbonyl, (C_{6-10} aryl)carbonyl, a group $-C(O)NR^{37}R^{38}$, C_{3-7} cycloalkyl, or 5- to 8-membered heterocycloalkyl; and

R^5 is a hydrogen atom or C_{1-4} alkyl;

(2) R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-6} saturated carbocyclic ring; and

R^4 is as defined above; and

(3) R^1 is a hydrogen atom or linear C_{1-10} alkyl;

R^4 and R^5 together with the carbon atom and the nitrogen atom to which they are

attached form a 5- to 8-membered saturated heterocyclic ring, wherein the saturated heterocyclic ring is optionally substituted with one or more substituents R^2 ;

R^3 is C_{1-10} alkyl optionally substituted with one or more substituents R_h , or R^3 is C_{1-4} alkyl substituted with R_e ;

R^{37} and R^{38} are each independently selected from a hydrogen atom and C_{1-3} alkyl; each R^2 is independently selected from C_{1-5} alkyl and a halogen atom; and/or two or more substituents R^2 on the 5- to 8-membered saturated heterocyclic ring may together form C_{1-5} alkylene that links the ring atoms to which they are attached;

each R_h is independently selected from a halogen atom, $(C_{1-4}$ alkoxy)carbonyl, and a group $-(O(CH_2)_a)_b-C_{1-4}$ alkoxy, wherein a is an integer selected from 2 to 4, and b is an integer selected from 1 to 4;

R_e is C_{6-10} aryl optionally substituted with one or more substituents R_a , or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R_a ;

each R_f is independently selected from a halogen atom, hydroxy, cyano, carboxy, $(C_{1-6}$ alkoxy)carbonyl, C_{1-6} alkoxy, and C_{6-10} aryl optionally substituted with one or more substituents R_g ;

each R_g is independently selected from a halogen atom, C_{1-6} alkyl, C_{2-6} alkynyl, and C_{1-6} alkoxy, wherein the alkyl, alkynyl, and alkoxy groups are each optionally substituted with one or more substituents selected from hydroxy and cyano;

each R_a is independently selected from a halogen atom, hydroxy, nitro, cyano, $(C_{1-4}$ alkoxy)carbonyl, 3- to 10-membered heterocycloalkyloxy (the heterocycloalkyloxy group is optionally substituted with optionally C_{1-4} alkoxy-substituted C_{1-4} alkyl), C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkenyl optionally substituted with one or more substituents R^{15} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} , a group $-(O(CH_2)_{q1})_{q2}-NR^{41}R^{42}$ (wherein $q1$ is an integer selected from 1 to 4, and $q2$ is an integer selected from 2 to 6), a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43}R^{44}$ (wherein $r1$ is an integer selected from 1 to 4, and $r2$

is an integer selected from 1 to 4), a group $-(O(CH_2)_{s1})_{s2}-NR^{45}-C(O)R^{46}$ (wherein $s1$ and $s2$ are each independently an integer selected from 2 to 4), a group $-C(O)NR^{47}R^{48}$, pyridinyl, pyrrolyl, a group $-NR^{49}R^{50}$, and a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-C(O)NR^{51}R^{52}$ (wherein $y1$ is an integer selected from 1 to 4, and $y2$ is an integer selected from 1 to 4);

R^{10} , R^{11} , R^{12} , R^{13} , and R^{15} are each independently selected from a halogen atom, hydroxy, carboxy, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-4} \text{ alkoxy})C_{1-6}$ alkoxy, $(C_{1-4} \text{ alkoxy})$ carbonyl, a group $-(O(CH_2)_o)_p-OH$ (wherein o and p are each independently an integer selected from 2 to 4), C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups, 3- to 10-membered heterocycloalkyl, 5- to 10-membered heteroaryl, and a group $-NR^{39}R^{40}$, wherein the 3- to 10-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, C_{1-4} alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups), $(C_{1-4} \text{ alkoxy})C_{1-4}$ alkyl (wherein the alkoxy moiety is optionally substituted with one or more hydroxy groups), C_{1-4} alkoxy, $(C_{1-4} \text{ alkoxy})$ carbonyl, C_{1-4} alkylthio, morpholino, $(C_{1-3} \text{ alkyl})$ sulfonyl, and $-C(O)NR^{53}R^{54}$;

Ar^1 is C_{6-10} aryl or 5- to 10-membered heteroaryl, wherein the aryl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from optionally C_{1-4} alkoxy-substituted C_{1-5} alkoxy, a halogen atom, C_{1-10} alkyl optionally substituted with one or more halogen atoms, a group $-SF_5$, cyano, hydroxy, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R^{14} , C_{6-10} aryl optionally substituted with one or more substituents R^{14} , and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14} ;

each R^{14} is independently selected from a halogen atom, oxo, cyano, nitro, C_{1-4} alkyl optionally substituted with one or more halogen atoms, C_{1-4} alkoxy optionally substituted with one or more halogen atoms, $(C_{1-6} \text{ alkoxy})$ carbonyl, a group $-NR^{27}R^{28}$, a group $-SO_2NR^{35}R^{36}$, C_{1-4} alkylthio, and 5- to 10-membered heterocycloalkyl;

R^{27} and R^{28} are each independently selected from a hydrogen atom and C_{1-4} alkyl optionally substituted with (C_{1-4} alkoxy)carbonyl;

R^{35} and R^{36} are each independently selected from a hydrogen atom and C_{1-4} alkyl;

R^{39} is a hydrogen atom, or optionally C_{1-6} alkoxy-substituted C_{1-6} alkyl;

R^{40} is a hydrogen atom, optionally C_{1-6} alkoxy-substituted C_{1-6} alkyl, ((C_{1-4} alkoxy)carbonyl) C_{1-6} alkyl, hydroxy-substituted C_{1-6} alkyl, a group $-(CH_2)_u-NR^{55}R^{56}$ (wherein u is an integer selected from 1 to 4), a group $-CH((CH_2)_{v1}COOR^{57})-(CH_2)_{v2}-COOR^{57}$ (wherein $v1$ is an integer selected from 0 to 2, and $v2$ is an integer selected from 1 to 3), a group $-(CH_2)_w-SO_3H$ (wherein w is an integer selected from 1 to 4), a group $-(CH_2)_{x1}-CH(COOH)-(CH_2)_{x2}-SO_3H$ (wherein $x1$ is an integer selected from 0 to 2, and $x2$ is an integer selected from 1 to 3), 3- to 6-membered oxacycloalkyl, or a group $-(CH_2)_{t1}-O-(CH_2)_{t2}-C(O)NR^{58}R^{59}$ (wherein $t1$ and $t2$ are each independently an integer selected from 1 to 3);

R^{41} is a hydrogen atom or C_{1-3} alkyl;

R^{42} is C_{1-8} alkyl substituted with one or more hydroxy groups, or (C_{1-3} alkoxy) C_{1-4} alkyl;

R^{43} is a hydrogen atom or C_{1-3} alkyl;

R^{44} is C_{1-8} alkyl substituted with one or more hydroxy groups, or

R^{43} and R^{44} together with the nitrogen atom to which they are attached may form morpholino;

R^{45} is a hydrogen atom or C_{1-3} alkyl;

R^{46} is C_{1-6} alkyl substituted with one or more hydroxy groups;

R^{47} is C_{1-3} alkyl;

R^{48} is (C_{1-3} alkoxy) C_{1-4} alkyl;

R^{49} is a hydrogen atom and C_{1-4} alkyl;

R^{50} is $-(CH_2)_z-NR^{60}R^{61}$ (z is an integer selected from 1 to 4, R^{60} is a hydrogen atom or C_{1-4} alkyl, and R^{61} is (C_{1-3} alkoxy) C_{1-4} alkyl, or R^{60} and R^{61} together with the nitrogen atom to which they are attached may form morpholino);

R^{51} is a hydrogen atom or C_{1-4} alkyl;

R^{52} is C_{1-8} alkyl substituted with one or more hydroxy groups;

R^{53} and R^{54} are each independently selected from a hydrogen atom and C_{1-4} alkyl;

R^{55} is a hydrogen atom or C_{1-4} alkyl;

R^{56} is $(C_{1-4}$ alkyl)carbonyl;

R^{57} is a hydrogen atom or C_{1-4} alkyl;

R^{58} is a hydrogen atom or C_{1-3} alkyl; and

R^{59} is C_{1-8} alkyl substituted with one or more hydroxy groups.

[0014] (1-2) The compound according to (1-1) or a salt thereof, or a solvate of the compound or the salt, wherein

each R_a is independently selected from a halogen atom, hydroxy, nitro, cyano, $(C_{1-4}$ alkoxy)carbonyl, 3- to 10-membered heterocycloalkyloxy, C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkenyl optionally substituted with one or more substituents R^{15} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , and C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} ;

R^{10} , R^{11} , R^{12} , R^{13} , and R^{15} are each independently selected from a halogen atom, hydroxy, carboxy, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-4}$ alkoxy)carbonyl, a group $-(O(CH_2)_o)_p-OH$ (wherein o and p are each independently an integer selected from 2 to 4), C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups, 3- to 10-membered heterocycloalkyl, 5- to 10-membered heteroaryl, and a group $-NR^{39}R^{40}$, wherein the 3- to 10-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, and C_{1-4} alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups); and

R^{39} and R^{40} are each independently selected from a hydrogen atom and optionally C_{1-6} alkoxy-substituted C_{1-6} alkyl.

[0015] (1-3) The compound according to (1-1) or (1-2) or a salt thereof, or a solvate of the compound or the salt, wherein

3- to 10-membered heterocycloalkyloxy in the definition of R_a is 3- to 6- membered

heterocycloalkyloxy;

3- to 10-membered heterocycloalkyl in the definition of R^{10} , R^{11} , R^{12} , R^{13} and R^{15} is 3- to 6- membered heterocycloalkyl; and

5- to 10-membered heteroaryl in the definition of R^{10} , R^{11} , R^{12} , R^{13} and R^{15} is 5- to 6-membered heteroaryl.

[0016] (1-4) The compound according to any of (1-1) to (1-3) or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is optionally C_{1-4} alkoxy-substituted C_{1-5} alkoxy or a halogen atom;

Rc is a halogen atom, C_{1-10} alkyl optionally substituted with one or more halogen atoms, or a group $-SF_5$; and

Rd is cyano, hydroxy, a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R^{14} , C_{6-10} aryl optionally substituted with one or more substituents R^{14} , or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14} .

[0017] (1-5) The compound according to any of (1-1) to (1-4) or a salt thereof, or a solvate of the compound or the salt, wherein

each Ra is independently selected from a halogen atom, nitro, cyano, (C_{1-4} alkoxy)carbonyl, 3- to 6-membered heterocycloalkyloxy (the heterocycloalkyloxy group is optionally substituted with optionally C_{1-4} alkoxy-substituted C_{1-4} alkyl), C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} , a group $-(O(CH_2)_{q1})_{q2}-NR^{41}R^{42}$, a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43}R^{44}$, a group $-(O(CH_2)_{s1})_{s2}-NR^{45}-C(O)R^{46}$, a group $-C(O)NR^{47}R^{48}$, pyridinyl, pyrrolyl, a group $-NR^{49}R^{50}$, and a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-NR^{51}R^{52}$;

R^{10} is carboxy, 3- to 6-membered heterocycloalkyl, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, or a group $-(O(CH_2)_6)_p-OH$;

R^{11} is hydroxy, carboxy, 5- or 6-membered heterocycloalkyl optionally substituted

with one or more oxo groups, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, a group -(O(CH₂)_o)_p-OH, C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups or a group -NR³⁹R⁴⁰;

R¹² is a halogen atom, hydroxy, carboxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₄ alkoxy)C₁₋₆ alkoxy, (C₁₋₃ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyl, 5- or 6-membered heteroaryl, or a group -NR³⁹R⁴⁰, wherein the 3- to 6-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, C₁₋₄ alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups), (C₁₋₄ alkoxy)C₁₋₄ alkyl (wherein the alkoxy moiety is optionally substituted with one or more hydroxy groups), C₁₋₄ alkoxy, (C₁₋₄ alkoxy)carbonyl, C₁₋₄ alkylthio, morpholinyl, (C₁₋₃ alkyl)sulfonyl, and -C(O)NR⁵³R⁵⁴;

R¹³ is 5- or 6-membered heterocycloalkyl; and

o and p are each independently an integer selected from 2 to 4.

[0018] (1-6) The compound according to any of (1-1) to (1-5) or a salt thereof, or a solvate of the compound or the salt, wherein

each R_a is independently selected from a halogen atom, nitro, cyano, (C₁₋₄ alkoxy)carbonyl, 3- to 6-membered heterocycloalkoxy, C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², and C₁₋₄ alkylthio optionally substituted with one or more substituents R¹³;

each R¹⁰ is independently selected from carboxy, 3- to 6-membered heterocycloalkyl, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, and a group -(O(CH₂)_o)_p-OH;

each R¹¹ is independently selected from hydroxy, carboxy, 5- or 6-membered heterocycloalkyl optionally substituted with one or more oxo groups, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, a group -(O(CH₂)_o)_p-OH, and C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups;

each R¹² is independently selected from a halogen atom, hydroxy, C₁₋₆ alkoxy

optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyl, 5- or 6-membered heteroaryl, and a group -NR³⁹R⁴⁰;

R¹³ is 5- or 6-membered heterocycloalkyl; and

o and p are each independently an integer selected from 2 to 4.

[0019] (1-7) The compound according to any of (1-1) to (1-5) or a salt thereof, or a solvate of the compound or the salt, wherein each R_a is independently selected from a halogen atom, hydroxy, nitro, cyano, C₁₋₆ alkyl, (carboxy)C₁₋₈ alkyl, (morpholino)C₁₋₄ alkyl, [HO-((CH₂)_oO)]C₁₋₆ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl optionally substituted with one or more hydroxy groups, C₁₋₆ alkoxy optionally substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, C₁₋₆ alkoxy substituted with one or more hydroxy groups, ((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkoxy, (C₁₋₆ alkoxy)C₁₋₈ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy(C₁₋₄ alkoxy))C₁₋₄ alkoxy, (carboxy)C₁₋₈ alkoxy, (3- to 6-membered heterocycloalkyl)C₁₋₆ alkoxy (the heterocycloalkyl moiety is optionally substituted with one or more substituents selected from oxo, a halogen atom, C₁₋₄ alkyl optionally substituted with one or more hydroxy groups, (C₁₋₄ alkoxy)C₁₋₄ alkyl (the alkoxy moiety is optionally substituted with one or more hydroxy groups), C₁₋₄ alkoxy, (C₁₋₄ alkoxy)carbonyl, C₁₋₄ alkylthio, morpholino, (C₁₋₃ alkyl)sulfonyl, and (di(C₁₋₃ alkyl)amino)carbonyl), C₁₋₄ alkoxy substituted with a group -NH-CH((CH₂)_{v1}COOR⁵⁷)-(CH₂)_{v2}-COOR⁵⁷, [N-(3- to 6-membered oxacycloalkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di((hydroxy)C₁₋₄ alkyl)-amino]C₁₋₄ alkoxy, [N-((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (pyridinyl)C₁₋₄ alkoxy, (pyrimidinyl)C₁₋₄ alkoxy, (1,2,4-triazolyl)C₁₋₄alkoxy, [N-(hydroxy)C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di((C₁₋₃ alkoxy)C₁₋₄ alkyl)amino]C₁₋₆ alkoxy, [N,N-di(C₁₋₃alkyl)amino]C₁₋₆ alkoxy, [N-[N-(C₁₋₄alkyl)carbonyl-N-(C₁₋₃alkyl)amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino]C₁₋₄ alkoxy, [N-[N-(C₁₋₄alkyl)carbonyl-amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino]C₁₋₄ alkoxy, a group -(O(CH₂)_{t1})_{t2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_w-SO₃H, C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_{x1}-CH(COOH)-(CH₂)_{x2}-SO₃H, a group -(O(CH₂)_{s1})_{s2}-NR⁴⁵-C(O)R⁴⁶, a group -C(O)NR⁴⁷R⁴⁸, pyridinyl, a group -NR⁴⁹R⁵⁰, a

group $-(O(CH_2)_{y1})_{y2}-O-CH_2-C(O)NR^{51}R^{52}$, (carboxy) C_{2-6} alkynyl, (3- to 6-membered heterocycloalkyl) C_{2-6} alkynyl optionally substituted with one or more oxo groups, C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, $[HO-((CH_2)_oO)_p]C_{2-8}$ alkynyl, $(C_{1-6}$ alkoxy) C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, $(C_{3-6}$ cycloalkyl) C_{2-6} alkynyl optionally substituted with one or more hydroxy groups, $[N-((C_{1-3}$ alkoxy) C_{1-4} alkyl)-N-(C_{1-3} alkyl)amino] C_{2-6} alkynyl, $(C_{1-3}$ alkoxy)carbonyl, (morpholino) C_{1-4} alkylthio optionally substituted with one or more oxo groups, 3- to 6-membered oxacycloalkyloxy, or 4- to 6-membered nitrogen containing heterocycloalkyloxy (the nitrogen containing heterocycloalkyl moiety is optionally substituted with one substituent selected from $(C_{1-3}$ alkoxy) C_{1-4} alkyl and C_{1-3} alkyl).

[0020] (1-8) The compound according to any of (1-1) to (1-7) or a salt thereof, or a solvate of the compound or the salt, wherein each R_a is independently selected from a halogen atom, hydroxy, cyano, C_{1-6} alkyl, (carboxy) C_{1-8} alkyl, (morpholino) C_{1-4} alkyl, $(C_{1-6}$ alkoxy) C_{1-8} alkyl substituted with one or more hydroxy groups, C_{1-6} alkoxy optionally substituted with one or more halogen atoms, $[N-((C_{1-3}$ alkoxy) C_{1-4} alkyl)-N-(C_{1-3} alkyl)amino] C_{1-4} alkoxy, (morpholino) C_{1-6} alkoxy (the morpholino moiety is optionally substituted with one or two substituent(s) selected from oxo and C_{1-3} alkyl), $(C_{1-6}$ alkoxy) C_{1-8} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-3}$ alkoxy(C_{1-4} alkoxy)) C_{1-4} alkoxy, (carboxy) C_{1-8} alkoxy, (pyrrolidinyl) C_{1-4} alkoxy (the pyrrolidinyl moiety is optionally substituted with $(C_{1-3}$ alkyl) C_{1-4} alkoxy), C_{1-4} alkoxy substituted with a group $-NH-CH((CH_2)_{v1}COOR^{57})-(CH_2)_{v2}-COOR^{57}$, $[N-(oxetanyl)-N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, $[N,N-di(C_{1-3}$ alkyl)amino] C_{1-6} alkoxy, a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43}R^{44}$, a group $-(O(CH_2)_{q1})_{q2}-NR^{41}R^{42}$, C_{1-4} alkoxy substituted with a group $-NH-(CH_2)_w-SO_3H$, C_{1-4} alkoxy substituted with a group $-NH-(CH_2)_{x1}-CH(COOH)-(CH_2)_{x2}-SO_3H$, (carboxy) C_{2-6} alkynyl, (morpholino) C_{2-6} alkynyl, C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, $[HO-((CH_2)_oO)_p]C_{2-8}$ alkynyl, $(C_{1-6}$ alkoxy) C_{2-8} alkynyl substituted with one or more hydroxy groups, or $(C_{1-3}$ alkoxy)carbonyl.

[0021] (1-9) The compound according to any of (1-1) to (1-8) or a salt thereof, or a solvate

of the compound or the salt, wherein each Ra is independently selected from a halogen atom, hydroxy, cyano, C₁₋₃ alkyl, (carboxy)C₁₋₈ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl substituted with one or more hydroxy groups, C₁₋₃ alkoxy optionally substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (morpholino)C₁₋₆ alkoxy, (C₁₋₆ alkoxy)C₁₋₈ alkoxy substituted with one or more hydroxy groups, (carboxy)C₂₋₆ alkynyl, (morpholino)C₂₋₆ alkynyl, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)₆O)_p]C₂₋₈ alkynyl, (C₁₋₆ alkoxy)C₂₋₈ alkynyl substituted with one or more hydroxy groups, and (C₁₋₃ alkoxy)carbonyl.

[0022] (1-10) The compound according to any of (1-1) to (1-9) or a salt thereof, or a solvate of the compound or the salt, wherein R³ is methyl substituted with Re.

[0023] (1-11) The compound according to any of (1-1) to (1-10) or a salt thereof, or a solvate of the compound or the salt, wherein R³ is benzyl optionally substituted with one or more substituents Ra on the benzene ring.

[0024] (1-12) The compound according to any of (1-1) to (1-11) or a salt thereof, or a solvate of the compound or the salt, wherein R³ is benzyl optionally substituted with one to three substituents Ra on the benzene ring.

[0025] (1-13) The compound according to any of (1-1) to (1-6), and (1-10) to (1-12) or a salt thereof, or a solvate of the compound or the salt, wherein Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri is a halogen atom or C₁₋₃ alkoxy;

Rj is a halogen atom, nitro, or cyano; and

Rk is hydroxy, a halogen atom, (C₁₋₄ alkoxy)carbonyl, 5- to 10-membered heterocycloalkyloxy (the heterocycloalkyloxy group is optionally substituted with optionally C₁₋₄ alkoxy-substituted C₁₋₄ alkyl), C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkenyl optionally substituted with one or more substituents R¹⁵, C₂₋₁₀

alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} , a group $-(O(CH_2)_{q1})_{q2}-NR^{41}R^{42}$, a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43}R^{44}$, a group $-(O(CH_2)_{s1})_{s2}-NR^{45}-C(O)R^{46}$, a group $-C(O)NR^{47}R^{48}$, pyridinyl, pyrrolyl, a group $-NR^{49}R^{50}$, or a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-C(O)NR^{51}R^{52}$.

[0026] (1-14) The compound according to (1-13) or a salt thereof, or a solvate of the compound or the salt, wherein

Rk is hydroxy, a halogen atom, (C_{1-4} alkoxy)carbonyl, 3- to 6-membered heterocycloalkyloxy (the heterocycloalkyloxy group is optionally substituted with optionally C_{1-4} alkoxy-substituted C_{1-4} alkyl), C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} , a group $-(O(CH_2)_{q1})_{q2}-NR^{41}R^{42}$, a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43}R^{44}$, a group $-(O(CH_2)_{s1})_{s2}-NR^{45}-C(O)R^{46}$, a group $-C(O)NR^{47}R^{48}$, pyridinyl, pyrrolyl, a group $-NR^{49}R^{50}$, or a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-C(O)NR^{51}R^{52}$;

R^{10} is a carboxy, hydroxy, morpholinyl, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, or a group $-(O(CH_2)_o)_p-OH$;

R^{11} is a hydroxy, carboxy, morpholinyl optionally substituted with one or more oxo groups, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, a group $-(O(CH_2)_o)_p-OH$, C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups or a group $-NR^{39}R^{40}$;

R^{12} is a halogen atom, hydroxy, carboxy, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, (C_{1-4} alkoxy) C_{1-6} alkoxy, (C_{1-3} alkoxy)carbonyl, 3- to 6-membered heterocycloalkyl, 5- or 6-membered heteroaryl containing one nitrogen atom, or a group $-NR^{39}R^{40}$;

wherein the 3- to 6-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, C_{1-4} alkyl (wherein the alkyl

group is optionally substituted with one or more hydroxy groups), (C₁₋₄ alkoxy)C₁₋₄ alkyl (wherein the alkoxy moiety is optionally substituted with one or more hydroxy groups), C₁₋₄ alkoxy, (C₁₋₄ alkoxy)carbonyl, C₁₋₄ alkylthio, morpholinyl, (C₁₋₃ alkyl)sulfonyl, and -C(O)NR⁵³R⁵⁴;

R¹³ is morpholinyl; and

o and p are each independently an integer selected from 2 to 4.

[0027] (1-15) The compound according to any of (1-1) to (1-6) and (1-10) to (1-13) or a salt thereof, or a solvate of the compound or the salt,

wherein Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri is a halogen atom or C₁₋₃ alkoxy;

Rj is a halogen atom, nitro, or cyano; and

Rk is hydroxy, a halogen atom, (C₁₋₄ alkoxy)carbonyl, 5- to 10-membered heterocycloalkyloxy, C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkenyl optionally substituted with one or more substituents R¹⁵, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², or C₁₋₄ alkylthio optionally substituted with one or more substituents R¹³.

[0028] (1-16) The compound according to (1-14) or (1-15) or a salt thereof, or a solvate of the compound or the salt,

wherein Rk is hydroxy, a halogen atom, (C₁₋₄ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyloxy, C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², or C₁₋₄ alkylthio optionally substituted with one or more substituents R¹³;

each R¹⁰ is independently selected from hydroxy, carboxy, morpholinyl, C₁₋₆ alkoxy

optionally substituted with one or more hydroxy groups, and a group $-(O(CH_2)_o)_p-OH$;

each R^{11} is independently selected from hydroxy, carboxy, morpholinyl optionally substituted with one or more oxo groups, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, a group $-(O(CH_2)_o)_p-OH$, and C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups;

each R^{12} is independently selected from a halogen atom, hydroxy, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-3}$ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyl, pyridinyl, pyrrolidyl, and a group $-NR^{39}R^{40}$;

R^{13} is morpholinyl; and

o and p are each independently an integer selected from 2 to 4.

[0029] (1-17) The compound according to (1-13) or (1-14) or a salt thereof, or a solvate of the compound or the salt, wherein R_k is a halogen atom, hydroxy, C_{1-6} alkyl, (carboxy) C_{1-8} alkyl, (morpholino) C_{1-4} alkyl, $[HO-((CH_2)_oO)]_pC_{1-6}$ alkyl, $(C_{1-6}$ alkoxy) C_{1-8} alkyl optionally substituted with one or more hydroxy groups, C_{1-6} alkoxy optionally substituted with one or more halogen atoms, $[N-((C_{1-3}$ alkoxy) C_{1-4} alkyl)- $N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, C_{1-6} alkoxy substituted with one or more hydroxy groups, $((C_{1-3}$ alkoxy)carbonyl) C_{1-3} alkoxy, $(C_{1-6}$ alkoxy) C_{1-8} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-3}$ alkoxy(C_{1-4} alkoxy)) C_{1-4} alkoxy, (carboxy) C_{1-8} alkoxy, (3- to 6-membered heterocycloalkyl) C_{1-6} alkoxy (the heterocycloalkyl moiety contains one to three heteroatoms selected from O or N, and is optionally substituted with one or more substituents selected from oxo, a halogen atom, C_{1-4} alkyl (wherein the alkyl moiety is optionally substituted with one or more hydroxy groups), $(C_{1-4}$ alkoxy) C_{1-4} alkyl (the alkoxy moiety is optionally substituted with one or more hydroxy groups), C_{1-4} alkoxy, $(C_{1-4}$ alkoxy)carbonyl, C_{1-4} alkylthio, morpholino, $(C_{1-3}$ alkyl)sulfonyl, and $(di(C_{1-3}$ alkyl)amino)carbonyl), C_{1-4} alkoxy substituted with a group $-NH-CH((CH_2)_{v1}COOR^{57})-(CH_2)_{v2}-COOR^{57}$, $[N-(3- to 6-membered oxacycloalkyl)-N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, $[N,N-di((hydroxy)C_{1-4}$ alkyl)-amino] C_{1-4} alkoxy, $[N-((C_{1-3}$ alkoxy)carbonyl) C_{1-3} alkyl- $N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, (pyridinyl) C_{1-4} alkoxy, (pyrimidinyl) C_{1-4} alkoxy, (1,2,4-triazolyl) C_{1-4} alkoxy, $[N-(hydroxy)C_{1-4}$ alkyl- $N-(C_{1-3}$

alkyl)amino]C₁₋₄ alkoxy, [N,N-di(C₁₋₃ alkoxy(C₁₋₃ alkyl))amino]C₁₋₆ alkoxy, [N,N-di(C₁₋₃alkyl)amino]C₁₋₆ alkoxy, [N-[N-(C₁₋₄ alkyl)carbonyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N-[N-(C₁₋₄ alkyl)carbonyl-amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, a group -(O(CH₂)_{r1})_{r2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_w-SO₃H, C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_{x1}-CH(COOH)-(CH₂)_{x2}-SO₃H, a group -(O(CH₂)_{s1})_{s2}-NR⁴⁵-C(O)R⁴⁶, a group -C(O)NR⁴⁷R⁴⁸, pyridinyl, a group -NR⁴⁹R⁵⁰, a group -(O(CH₂)_{y1})_{y2}-O-CH₂-C(O)NR⁵¹R⁵², (carboxy)C₂₋₈ alkynyl, (3- to 6-membered heterocycloalkyl)C₂₋₆ alkynyl optionally substituted with one or more oxo groups, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)_oO)_p]C₂₋₈ alkynyl, (C₁₋₆ alkoxy)C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, (C₃₋₆ cycloalkyl)C₂₋₆ alkynyl optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy)carbonyl, (morpholino)C₁₋₄ alkylthio, 3- to 6-membered oxacycloalkyloxy, or 3- to 6- membered nitrogen containing heterocycloalkyloxy (the nitrogen containing heterocycloalkyl moiety is optionally substituted with one substituent selected from (C₁₋₃ alkoxy)C₁₋₄ alkyl and C₁₋₃ alkyl).

[0030] (1-18) The compound according to any of (1-13), (1-14) and (1-17) or a salt thereof, or a solvate of the compound or the salt, wherein Rk is hydroxy, a halogen atom, C₁₋₆ alkyl, (carboxy)C₁₋₈ alkyl, (morpholino)C₁₋₄ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl substituted with one or more hydroxy groups, C₁₋₆ alkoxy optionally substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (morpholino)C₁₋₆ alkoxy, (C₁₋₆ alkoxy)C₁₋₈ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy(C₁₋₄ alkoxy))C₁₋₄ alkoxy, (carboxy)C₁₋₈ alkoxy, (pyrrolidinyl)C₁₋₄ alkoxy (the pyrrolidinyl moiety is substituted with (C₁₋₃ alkyl)C₁₋₄ alkoxy), C₁₋₄ alkoxy substituted with a group -NH-CH((CH₂)_{v1}COOR⁵⁷)-(CH₂)_{v2}-COOR⁵⁷, [N-(oxetanyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di(C₁₋₃ alkyl)amino]C₁₋₆ alkoxy, a group -(O(CH₂)_{r1})_{r2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_w-SO₃H, C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_{x1}-CH(COOH)-(CH₂)_{x2}-SO₃H, (carboxy)C₂₋₈

alkynyl, (morpholino) C_{2-6} alkynyl, C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, $[HO-((CH_2)_6O)_p]C_{2-8}$ alkynyl, $(C_{1-6}$ alkoxy) C_{2-8} alkynyl substituted with one or more hydroxy groups, or $(C_{1-3}$ alkoxy)carbonyl.

[0031] (1-19) The compound according to any of (1-13) to (1-18) or a salt thereof, or a solvate of the compound or the salt, wherein R_k is hydroxy, a halogen atom, C_{1-6} alkyl, (carboxy) C_{1-8} alkyl, $(C_{1-6}$ alkoxy) C_{1-8} alkyl substituted with one or more hydroxy groups, C_{1-3} alkoxy optionally substituted with one or more halogen atoms, $[N-((C_{1-3}$ alkoxy) C_{1-4} alkyl)- $N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, (morpholino) C_{1-6} alkoxy, $(C_{1-6}$ alkoxy) C_{1-8} alkoxy substituted with one or more hydroxy groups, (carboxy) C_{2-8} alkynyl, (morpholino) C_{2-6} alkynyl, C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, $[HO-((CH_2)_6O)_p]C_{2-8}$ alkynyl, $(C_{1-6}$ alkoxy) C_{2-8} alkynyl substituted with one or more hydroxy groups, or $(C_{1-3}$ alkoxy)carbonyl.

(1-20) The compound according to any of (1-1) to (1-19) or a salt thereof, or a solvate of the compound or the salt,

wherein R^1 , R^4 , and R^5 are as defined in any one of the following (1) to (3):

(1) R^1 is a hydrogen atom or C_{1-6} alkyl;

R^4 is C_{1-4} alkyl optionally substituted with one or more halogen atom(s), or phenyl;

and

R^5 is a hydrogen atom or C_{1-4} alkyl;

(2) R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-6} saturated carbocyclic ring; and

R^4 is as defined above; and

(3) R^1 is a hydrogen atom or linear C_{1-6} alkyl;

R^4 and R^5 together with the carbon atom and the nitrogen atom to which they are attached form a 5- to 8-membered saturated heterocyclic ring.

[0032] (1-21) The compound according to any of (1-1) to (1-3), (1-10) to (1-12), and (1-20) or a salt thereof, or a solvate of the compound or the salt, wherein

R^1 , R^4 , and R^5 are as defined in the following (1) or (2):

(1) R¹ is selected from C₁₋₆ alkyl;

R⁴ is optionally halogen atom-substituted C₁₋₄ alkyl or phenyl; and

R⁵ is a hydrogen atom or C₁₋₄ alkyl; or

(2) R¹ and R⁵ together with the carbon atom to which they are attached form a C₃₋₆ saturated carbocyclic ring; and

R⁴ is as defined above;

R³ is C₁₋₄ alkyl substituted with Re;

Re is phenyl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom, C₂₋₆ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₆ alkoxy optionally substituted with one or more substituents R¹², 3- to 6-membered heterocycloalkyloxy (the heterocycloalkyloxy group is optionally substituted with optionally C₁₋₄ alkoxy-substituted C₁₋₄ alkyl), a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², a group -(O(CH₂)_{r1})_{r2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{s1})_{s2}-NR⁴⁵-C(O)R⁴⁶, a group -C(O)NR⁴⁷R⁴⁸, pyridinyl, pyrrolyl, a group -NR⁴⁹R⁵⁰, and a group -(O(CH₂)_{y1})_{y2}-O-CH₂-C(O)NR⁵¹R⁵²;

R¹¹ and R¹² are each independently selected from halogen atom, hydroxy, carboxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy, (C₁₋₄ alkoxy)C₁₋₆ alkoxy, 3- to 6-membered heterocycloalkyl, 5- to 6-membered heteroaryl, and a group -NR³⁹R⁴⁰, wherein the 3- to 6-membered heterocycloalkyl group is optionally substituted with one or two substituents selected from oxo, a halogen atom, C₁₋₄ alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups), (C₁₋₄ alkoxy)C₁₋₄ alkyl (wherein the alkoxy moiety is optionally substituted with one or more hydroxy groups), C₁₋₄ alkoxy, (C₁₋₄ alkoxy)carbonyl, C₁₋₄ alkylthio, morpholino, (C₁₋₃ alkyl)sulfonyl, and -C(O)NR⁵³R⁵⁴;

Ar¹ is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, phenyl optionally substituted with

one or more substituents R^{14} , and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from a halogen atom, cyano, C_{1-4} alkyl optionally substituted with one or more halogen atoms, C_{1-4} alkoxy, a group $-SO_2NR^{35}R^{36}$ (wherein R^{35} and R^{36} are each independently selected from C_{1-4} alkyl), and C_{1-4} alkylthio.

[0033] (1-22) The compound according to any of (1-1) to (1-3), (1-5) to (1-7), (1-10) to (1-12), (1-20), and (1-21) or a salt thereof, or a solvate of the compound or the salt, wherein each R_a is independently selected from a halogen atom, $[N-((C_{1-3}$ alkoxy) C_{1-4} alkyl)- $N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, $(C_{1-4}$ alkoxy) C_{1-6} alkoxy (the C_{1-4} alkoxy moiety is optionally substituted with one or more hydroxy groups), $(C_{1-3}$ alkoxy(C_{1-4} alkoxy)) C_{1-4} alkoxy, (carboxy) C_{1-6} alkoxy, (3- to 6-membered heterocycloalkyl) C_{1-6} alkoxy (the heterocycloalkyl moiety is optionally substituted with one or two substituents selected from oxo, a halogen atom, C_{1-4} alkyl optionally substituted with one or more hydroxy groups, $(C_{1-4}$ alkoxy) C_{1-4} alkyl (the alkoxy moiety is optionally substituted with one or more hydroxy groups), C_{1-4} alkoxy, $(C_{1-4}$ alkoxy)carbonyl, C_{1-4} alkylthio, morpholino, $(C_{1-3}$ alkyl)sulfonyl, and $(di(C_{1-3}$ alkyl)amino)carbonyl), C_{1-4} alkoxy substituted with a group $-NH-CH((CH_2)_{v1}COOR^{57})-(CH_2)_{v2}-COOR^{57}$, $[N-(3- \text{ to } 6\text{-membered oxacycloalkyl})-N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, $[N,N-di((hydroxy)C_{1-4}$ alkyl)-amino] C_{1-4} alkoxy, $[N-((C_{1-3}$ alkoxy)carbonyl) C_{1-3} alkyl- $N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, (1,2,4-triazolyl) C_{1-4} alkoxy, $[N-(hydroxy)C_{1-4}$ alkyl- $N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, $[N,N-di((C_{1-3}$ alkoxy) C_{1-4} alkyl)amino] C_{1-6} alkoxy, $[N,N-di(C_{1-3}alkyl)amino]C_{1-6}$ alkoxy, $[N-[N-(C_{1-4}alkyl)carbonyl-N-(C_{1-3}alkyl)amino]C_{1-4}$ alkyl- $N-(C_{1-3}$ alkyl) amino] C_{1-4} alkoxy, $[N-[N-(C_{1-4}alkyl)carbonyl-amino]C_{1-4}$ alkyl- $N-(C_{1-3}$ alkyl) amino] C_{1-4} alkoxy, C_{1-6} alkoxy optionally substituted with one or more halogen atoms, a group $-(O(CH_2)_{t1})_{t2}-C(O)NR^{43}R^{44}$, a group $-(O(CH_2)_{q1})_{q2}-NR^{41}R^{42}$, C_{1-4} alkoxy substituted with a group $-NH-(CH_2)_w-SO_3H$, C_{1-4} alkoxy substituted with a group $-NH-(CH_2)_{x1}-CH(COOH)-(CH_2)_{x2}-SO_3H$, a group $-(O(CH_2)_{s1})_{s2}-NR^{45}-C(O)R^{46}$, a group $-C(O)NR^{47}R^{48}$, pyridinyl, a group $-NR^{49}R^{50}$, a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-C(O)NR^{51}R^{52}$, (3- to 6-membered heterocycloalkyl) C_{2-6} alkynyl (the heterocycloalkyl moiety

is optionally substituted with one or more oxo groups), [N-((C₁₋₃alkoxy)C₁₋₄alkyl)-N-(C₁₋₃alkyl)amino]C₂₋₆alkynyl, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, 3- to 6-membered oxacycloalkyloxy, or 4- to 6- membered nitrogen containing heterocycloalkyloxy (the heterocycloalkyl moiety is optionally substituted with one substituent selected from (C₁₋₃ alkoxy)C₁₋₄ alkyl and C₁₋₃ alkyl).

[0034] (1-23) The compound according to any of (1-1) to (1-3), (1-10) to (1-13), (1-20), and (1-21) or a salt thereof, or a solvate of the compound or the salt, wherein Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri is a halogen atom;

Rj is a halogen atom; and

Rk is a halogen atom, 3- to 6-membered heterocycloalkyloxy (the heterocycloalkyl moiety comprises one heteroatom selected from O and N and is optionally substituted with optionally C₁₋₄ alkoxy-substituted C₁₋₄ alkyl), optionally R¹¹-substituted C₂₋₆ alkynyl, optionally R¹²-substituted C₁₋₆ alkoxy, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², a group -(O(CH₂)_{r1})_{r2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{s1})_{s2}-NR⁴⁵-C(O)R⁴⁶, a group -C(O)NR⁴⁷R⁴⁸, a group -NR⁴⁹R⁵⁰, or a group -(O(CH₂)_{y1})_{y2}-O-CH₂-C(O)NR⁵¹R⁵².

[0035] (1-24) The compound according to (1-21) or (1-23) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹¹ is morpholino;

R¹² is selected from 5- to 6- membered heterocycloalkyl which contains one or two hetero atoms selected from O and N, (C₁₋₄alkoxy)C₁₋₆alkoxy or a group -NR³⁹R⁴⁰.

[0036] (1-25) The compound according to any of (1-13), (1-14), (1-17) and (1-23) or a salt thereof, or a solvate of the compound or the salt,

wherein Rk is [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (C₁₋₄ alkoxy)C₁₋₆ alkoxy (the C₁₋₄ alkoxy moiety is optionally substituted with one or more hydroxy groups),

(C₁₋₃ alkoxy(C₁₋₄ alkoxy))C₁₋₄ alkoxy, (carboxy)C₁₋₆ alkoxy, (3- to 6-membered heterocycloalkyl)C₁₋₆ alkoxy (the heterocycloalkyl moiety comprises one to three hetero atom(s) which is selected from O or N and is optionally substituted with one or two substituents selected from oxo, a halogen atom, C₁₋₄ alkyl (wherein the alkyl moiety is optionally substituted with one or more hydroxy groups), (C₁₋₄ alkoxy)C₁₋₄ alkyl (the alkoxy moiety is optionally substituted with one or more hydroxy groups), C₁₋₄ alkoxy, (C₁₋₄ alkoxy)carbonyl, C₁₋₄ alkylthio, morpholino, (C₁₋₃ alkyl)sulfonyl, and (di(C₁₋₃ alkyl)amino)carbonyl), C₁₋₄ alkoxy substituted with a group -NH-CH((CH₂)_{v1}COOR⁵⁷)-(CH₂)_{v2}-COOR⁵⁷, [N-(3- to 6-membered oxacycloalkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di((hydroxy)C₁₋₄ alkyl)-amino]C₁₋₄ alkoxy, [N-((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (1,2,4-triazolyl)C₁₋₄ alkoxy, [N-(hydroxy)C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di(C₁₋₃ alkoxy(C₁₋₄ alkyl))amino]C₁₋₆ alkoxy, [N,N-di(C₁₋₃ alkyl)amino]C₁₋₆ alkoxy, [N-[N-(C₁₋₄ alkyl)carbonyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N-[N-(C₁₋₄ alkyl)carbonyl-amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, a group -(O(CH₂)_{t1})_{t2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_w-SO₃H, C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_{x1}-CH(COOH)-(CH₂)_{x2}-SO₃H, a group -(O(CH₂)_{s1})_{s2}-NR⁴⁵-C(O)R⁴⁶, a group -C(O)NR⁴⁷R⁴⁸, pyridinyl, a group -NR⁴⁹R⁵⁰, a group -(O(CH₂)_{y1})_{y2}-O-CH₂-C(O)NR⁵¹R⁵², (3- to 6-membered heterocycloalkyl)C₂₋₆ alkynyl optionally substituted with one or more oxo groups (the heterocycloalkyl moiety comprises one to three hetero atoms selected from O and N), [N-((C₁₋₃alkoxy)C₁₋₄alkyl-N-(C₁₋₃alkyl)amino]C₂₋₆alkynyl, 3- to 6-membered oxacycloalkyloxy, or 4- to 6-membered nitrogen containing heterocycloalkyloxy (the nitrogen containing heterocycloalkyl moiety is optionally substituted with one substituent selected from (C₁₋₃ alkoxy)C₁₋₄ alkyl and C₁₋₃ alkyl).

[0037] (1-26) The compound according to any of (1-1) to (1-3), (1-5), (1-7), (1-8), (1-10) to (1-20), and (1-21) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹, R⁴, and R⁵ are as defined in the following (1) or (2):

(1) R¹ is C₁₋₆ alkyl;

R^4 is optionally halogen atom-substituted C_{1-4} alkyl or phenyl; and

R^5 is a hydrogen atom or C_{1-4} alkyl; or

(2) R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-6} saturated carbocyclic ring; and

R^4 is as defined above;

R^3 is C_{1-4} alkyl substituted with R_e ;

R_e is phenyl optionally substituted with one or more substituents R_a ;

each R_a is independently selected from a halogen atom, optionally R^{11} -substituted C_{2-6} alkynyl, and optionally R^{12} -substituted C_{1-6} alkoxy;

R^{11} and R^{12} are each independently selected from 5- or 6-membered heterocycloalkyl and $-NR^{39}R^{40}$;

R^{39} and R^{40} are each independently selected from hydrogen atom and optionally C_{1-6} alkoxy-substituted C_{1-6} alkyl;

Ar^1 is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from R_b , R_c , and R_d ;

R_b , R_c , and R_d are each independently selected from a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, phenyl optionally substituted with one or more substituents R^{14} , and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from a halogen atom, cyano, C_{1-4} alkyl optionally substituted with one or more halogen atoms, C_{1-4} alkoxy, a group $-SO_2NR^{35}R^{36}$ (wherein R^{35} and R^{36} are each independently selected from C_{1-4} alkyl), and C_{1-4} alkylthio.

[0038] (1-27) The compound according to any of (1-1) to (1-26) or a salt thereof, or a solvate of the compound or the salt, wherein

R_b is a halogen atom;

R_c is a halogen atom or C_{1-4} alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, phenyl optionally substituted with one or more substituents R¹⁴, or 5- or 6-membered heteroaryl optionally substituted with one or more substituents R¹⁴.

[0039] (1-28) The compound according to any of (1-1) to (1-12), (1-20), (1-21), (1-23), (1-26) and (1-27) or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one or more substituents Ra;

R¹¹ and R¹² are each independently selected from morpholinyl and a group -NR³⁹R⁴⁰;

R⁴ is optionally halogen atom-substituted C₁₋₄ alkyl or phenyl;

Ar¹ is phenyl, pyridinyl, or pyrimidinyl, wherein the phenyl, pyridinyl, and pyrimidinyl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb is a halogen atom;

Rc is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, phenyl, pyridinyl, or pyrimidinyl, wherein the phenyl, pyridinyl, and pyrimidinyl groups are each optionally substituted with one or more substituents R¹⁴.

[0040] (1-29) The compound according to any of (1-1) to (1-4), (1-10) to (1-17), (1-20), (1-21), (1-23), and (1-26) to (1-28) or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri is a halogen atom;

Rj is a halogen atom; and

Rk is hydroxy, C₂₋₆ alkynyl optionally substituted with a substituent R¹¹, or C₁₋₆

alkoxy optionally substituted with a substituent R¹².

[0041] (1-30) The compound according to any of (1-26) to (1-29) or a salt thereof, or a solvate of the compound or the salt, wherein R¹¹ and R¹² are each independently selected from morpholinyl, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino] and [N,N-di(C₁₋₃alkyl)amino].

[0042] (1-31) The compound according to any of (1-26) to (1-30) or a salt thereof, or a solvate of the compound or the salt, wherein R¹¹ and R¹² are each independently selected from morpholinyl, and [N-((methoxy)ethyl)-N-(methyl)amino].

[0043] (1-32) The compound according to any of (1-1) to (1-31) or a salt thereof, or a solvate of the compound or the salt, wherein R¹ and R⁵ together with the carbon atom to which they are attached form a C₃₋₆ saturated carbocyclic ring, and R⁴ is C₁₋₄ alkyl.

[0044] (1-33) The compound according to any of (1-1) to (1-3), (1-5), (1-6), (1-10) to (1-12) and (1-20) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹ and R⁵ together with the carbon atom to which they are attached form a C₃₋₆ saturated carbocyclic ring;

R⁴ is C₁₋₄ alkyl;

R³ is C₁₋₄ alkyl substituted with Re;

Re is phenyl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom and C₁₋₄ alkoxy optionally substituted with one or more substituents R¹²;

each R¹² is independently selected from 5- or 6-membered heterocycloalkyl and -NR³⁹R⁴⁰;

R³⁹ and R⁴⁰ are each independently selected from a hydrogen atom and optionally C₁₋₄ alkoxy-substituted C₁₋₄ alkyl;

Ar¹ is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from a halogen atom, C₁₋₄ alkyl

optionally substituted with one or more halogen atoms, and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from a halogen atom, cyano, C_{1-4} alkyl optionally substituted with one or more halogen atoms, and C_{1-4} alkylthio.

[0045] (1-34) The compound according to any of (1-1) to (1-33) or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is a halogen atom;

Rc is a halogen atom or C_{1-4} alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom or 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14} .

[0046] (1-35) The compound according to any of (1-1) to (1-34) or a salt thereof, or a solvate of the compound or the salt, wherein R^3 is benzyl optionally substituted with one to three substituents Ra on the benzene ring.

[0047] (1-36) The compound according to any of (1-1) to (1-35) or a salt thereof, or a solvate of the compound or the salt, wherein Ra is selected from a halogen atom, (morpholino) C_{1-4} alkoxy, [N-((C_{1-3} alkoxy) C_{1-4} alkyl)-N-(C_{1-3} alkyl)amino] C_{1-4} alkoxy, [N,N-di(C_{1-3} alkyl)amino] C_{1-4} alkoxy.

[0048] (1-37) The compound according to any of (1-1) to (1-5), (1-10) to (1-16), (1-20), and (1-33) to (1-36) or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri and Rj are each independently a halogen atom; and

Rk is C_{1-4} alkoxy optionally substituted with one or more substituents R^{12} .

[0049] (1-38) The compound according to any of (1-13) to (1-18), (1-20), (1-25), (1-29) and (1-37) or a salt thereof, or a solvate of the compound or the salt, wherein

R_k is a halogen atom, (morpholino)C₁₋₄ alkoxy, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃alkyl)amino]C₁₋₄ alkoxy, [N,N-di(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy.

[0050] (1-39) The compound according to any of (1-1) to (1-6), (1-10) to (1-16), (1-20) to (1-23), (1-26), (1-33) to (1-35), (1-37) and (1-38) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar¹ is phenyl or pyridinyl;

R_d is pyridinyl or pyrimidinyl;

R¹² is selected from morpholinyl, [N-(2-(methoxy)ethyl)-N-(methyl)amiono], [N,N-dimethylamino].

[0051] (1-40) The compound according to any of (1-1) to (1-39) or a salt thereof, or a solvate of the compound or the salt, wherein

R_b is a halogen atom;

R_c is a halogen atom, methyl or trifluoromethyl;

R_d is a halogen atom, trifluoromethyl, phenyl optionally substituted with one to three substituents R¹⁴ and 5- to 6-membered heteroaryl optionally substituted with one to three substituents R¹⁴, wherein the heteroaryl comprises one to three hetero atoms selected from O, S, and N.

[0052] (1-41) The compound according to any of (1-1) to (1-40) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹⁴ is each selected from methyl, trifluoromethyl, cyano, nitro, a halogen atom, methoxy, ethoxy, trifluoromethoxy, methylthio, methoxycarbonyl and dimethylaminosulfonyl.

[0053] (1-42) The compound according to any of (1-1) to (1-41) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹⁴ is each selected from methyl, trifluoromethyl, cyano, a chlorine atom and methylthio.


[0054] (1-43) The compound according to any of (1-1) to (1-42) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar¹ is phenyl or 5- to 6-membered heteroaryl which comprises one to three hetero atoms

selected from O, S, and N, wherein the phenyl and the heteroaryl is substituted with one to three substituents selected from Rb, Rc, and Rd.

[0055] (1-44) The compound, according to any of (1-1) to (1-11), (1-20) to (1-22), (1-24), (1-26) to (1-28), (1-30) to (1-36), and (1-39) to (1-43) or a salt thereof, or a solvate of the compound or the salt, wherein the compound is represented by the formula (I-c);

[0056] [Formula 2]




[0057] wherein, n1 is an interger selected from 1 to 4, n2 is an interger selected from 0 or more, R¹, R⁴, R⁵, Ar¹, Ra, Rb, Rc, Rd, Re are as defined in any of (1-1) to (1-11), (1-20) to (1-22), (1-24), (1-26) to (1-28), (1-30) to (1-36), and (1-39) to (1-43).

[0058] (1-45) The compound according to (1-44) or a salt thereof, or a solvate of the compound or the salt, wherein n1 is 1 and n2 is 3.

[0059] (1-46) The compound, according to any of (1-13) to (1-20), (1-23) to (1-25), (1-29) to (1-32) and (1-37) to (1-43) or a salt thereof, or a solvate of the compound or the salt, wherein the compound is represented by the formula (I-d);

[0060] [Formula 3]



[0061] wherein n1 is an interger selected from 1 to 4, n3, n4 and n5 is a interger independently selected from 0 or 1, provided that at least one of n3, n4 and n5 is 1, R¹, R⁴, R⁵,

Ar¹, Ra, Rb, Rc, Rd, Re, Ri, Rj, and Rk are as defined in any of (1-13) to (1-20), (1-23) to (1-25), (1-29) to (1-32), and (1-37) to (1-43).

[0062] (1-47) The compound according to (1-46) or a salt thereof, or a solvate of the compound or the salt, wherein n1 is 1.

[0063] (1-48) The compound according to any of (1-1) to (1-47) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar¹ is 4-(trifluoromethyl)-2-(6-methylthiopyridin-3-yl)phenyl, 4-(trifluoromethyl)-2-(6-trifluoromethylpyridin-3-yl)phenyl, 4-(trifluoromethyl)-2-(4-trifluoromethylpyrimidin-5-yl)phenyl, 4-(trifluoromethyl)-2-(6-trifluoromethylpyrimidin-4-yl)phenyl, 4-(trifluoromethyl)-2-(6-cyano-5-methylpyrimidin-4-yl)phenyl, 4-(trifluoromethyl)-2-(2-cyanopyridin-4-yl)phenyl, 4-chloro-2-(6-methylthiopyridin-3-yl)phenyl, 4-chloro-2-(6-trifluoromethylpyridin-3-yl)phenyl, 4-chloro-2-(4-trifluoromethylpyrimidin-5-yl)phenyl, 4-chloro-2-(6-cyano-5-methylpyrimidin-4-yl)phenyl, 4-chloro-2-(6-trifluoromethylpyrimidin-4-yl)phenyl, or 4-chloro-2-(2-cyanopyridin-4-yl)phenyl.

[0064] (1-49) A compound selected from:

(4aR)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-[2-(6-cyano-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-[2-(2-cyanopyridin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

6-[4-[(4aR)-4-hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]hex-5-ynoic acid;

(4aR)-1-[[2,3-difluoro-4-(3-morpholin-4-ylprop-1-ynyl)phenyl]methyl]-4-hydroxy-

4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[3-[(2R)-2,3-dihydroxypropoxy]propoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[4-[(2R)-2,3-dihydroxypropoxy]butoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[6-[(2R)-2,3-dihydroxypropoxy]hexoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[2,3-difluoro-4-(morpholin-4-ylmethyl)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-(4-bromo-3,5-difluorophenyl)-1-[(3-chloro-2-fluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(3S)-3-tert-butyl-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-(6-methylsulfanylpurin-3-yl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[2-(6-cyano-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

6-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

6-[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

4-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]butanoic acid;

5-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]pentanoic acid;

6-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]hexanoic acid;

7-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-

(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]heptanoic acid;

7-[[2,3-difluoro-4-[2-[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

(2S)-2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]butanedioic acid;

3-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]pentanedioic acid;

6-(2,3-difluoro-4-(2-(methyl(oxetan-3-yl)amino)ethoxy)benzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-[2-[methyl(oxetan-3-yl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

7-(2,3-difluoro-4-(2-(methyl(oxetan-3-yl)amino)ethoxy)benzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

7-(4-(3-(dimethylamino)-2,2-dimethylpropoxy)-2,3-difluorobenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

6-[[2,3-difluoro-4-[1-(2-methoxyethyl)azetidin-3-yl]oxyphenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-[4-[methyl-[(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]-4-oxobutoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-

(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

7-[[2,3-difluoro-4-[2-[2-[methyl-(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

7-[[2,3-difluoro-4-[2-[2-[2-[2-methoxyethyl(methyl)amino]ethoxy]ethoxy]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]ethanesulfonic acid; and

2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]ethanesulfonic acid

or a salt thereof, or a solvate of the compound or the salt.

[0065] (1-50) The compound according to any of (1-1) to (1-49) or the salt thereof, or the solvate of the compound or the salt for use in the prevention and/or treatment of a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure.

[0066] (1-51) A NaPi-IIb inhibitor comprising a compound according to any of (1-1) to (1-49) or a salt thereof, or a solvate of the compound or the salt.

[0067] (1-52) A PiT-1 inhibitor comprising a compound according to any of (1-1) to (1-49) or a salt thereof, or a solvate of the compound or the salt.

[0068] (1-53) A PiT-2 inhibitor comprising a compound according to any of (1-1) to (1-49) or a salt thereof, or a solvate of the compound or the salt.

[0069] (1-54) A preventive and/or therapeutic agent for a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, the agent comprising a compound according to any of (1-1) to (1-49) or a salt thereof, or a solvate of


the compound or the salt as an active ingredient.

[0070] (1-55) A method for preventing and/or treating a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, comprising administering a therapeutically effective amount of a compound according to any of (1-1) to (1-49) or a salt thereof, or a solvate of the compound or the salt to a patient.

[0071] Another embodiment of the present invention provides the following compounds (2-1) to (2-26):

[0072] (2-1) A compound represented by the formula (I) or a salt thereof, or a solvate of the compound or the salt:

[0073] [Formula 4]



[0074] wherein R^1 , R^4 , and R^5 are as defined in any one of the following (1) to (3):

(1) R^1 is a hydrogen atom or C_{1-10} alkyl;

R^4 is a hydrogen atom, C_{1-4} alkyl optionally substituted with one or more substituents R_f , C_{6-10} aryl optionally substituted with one or more substituents R_g , (C_{1-6} alkyl)carbonyl, (C_{6-10} aryl)carbonyl, a group $-C(O)NR^{37}R^{38}$, C_{3-7} cycloalkyl, or 5- to 8-membered heterocycloalkyl; and

R^5 is a hydrogen atom or C_{1-4} alkyl;

(2) R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-6} saturated carbocyclic ring; and

R^4 is as defined above; and

(3) R^1 is a hydrogen atom or linear C_{1-10} alkyl;

R^4 and R^5 together with the carbon atom and the nitrogen atom to which they are

attached form a 5- to 8-membered saturated heterocyclic ring, wherein the saturated heterocyclic ring is optionally substituted with one or more substituents R^2 ;

R^3 is C_{1-10} alkyl optionally substituted with one or more substituents Rh or C_{1-4} alkyl substituted with Re;

R^{37} and R^{38} are each independently selected from a hydrogen atom and C_{1-3} alkyl; each R^2 is independently selected from C_{1-5} alkyl and a halogen atom; and/or two or more substituents R^2 on the 5- to 8-membered saturated heterocyclic ring may together form C_{1-5} alkylene that links the ring atoms to which they are attached;

each Rh is independently selected from a halogen atom, (C_{1-4} alkoxy)carbonyl, and a group $-(O(CH_2)_a)_b-C_{1-4}$ alkoxy (wherein a is an integer selected from 2 to 4, and b is an integer selected from 1 to 4);

Re is C_{6-10} aryl optionally substituted with one or more substituents Ra or 5- to 10-membered heteroaryl optionally substituted with one or more substituents Ra;

each Rf is independently selected from a halogen atom, hydroxy, cyano, carboxy, (C_{1-6} alkoxy)carbonyl, C_{1-6} alkoxy, and C_{6-10} aryl optionally substituted with one or more substituents Rg;

each Rg is independently selected from a halogen atom, C_{1-6} alkyl, C_{2-6} alkynyl, and C_{1-6} alkoxy, wherein the alkyl, alkynyl, and alkoxy groups are each optionally substituted with one or more substituents selected from hydroxy and cyano;

each Ra is independently selected from a halogen atom, hydroxy, nitro, cyano, (C_{1-4} alkoxy)carbonyl, 3- to 10-membered heterocycloalkyloxy, C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkenyl optionally substituted with one or more substituents R^{15} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , and C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} ;

R^{10} , R^{11} , R^{12} , R^{13} , and R^{15} are each independently selected from a halogen atom, hydroxy, carboxy, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, (C_{1-4} alkoxy)carbonyl, a group $-(O(CH_2)_o)_p-OH$ (wherein o and p are each independently an

integer selected from 2 to 4), C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups, 3- to 10-membered heterocycloalkyl, 5- to 10-membered heteroaryl, and a group -NR³⁹R⁴⁰, wherein the 3- to 10-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, and C₁₋₄ alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups);

Ar¹ is C₆₋₁₀ aryl or 5- to 10-membered heteroaryl, wherein the aryl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from optionally C₁₋₄ alkoxy-substituted C₁₋₅ alkoxy, a halogen atom, C₁₋₁₀ alkyl optionally substituted with one or more halogen atoms, a group -SF₅, cyano, hydroxy, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R¹⁴, C₆₋₁₀ aryl optionally substituted with one or more substituents R¹⁴, and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R¹⁴;

each R¹⁴ is independently selected from a halogen atom, oxo, cyano, nitro, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₁₋₄ alkoxy optionally substituted with one or more halogen atoms, (C₁₋₆ alkoxy)carbonyl, a group -NR²⁷R²⁸, a group -SO₂NR³⁵R³⁶, C₁₋₄ alkylthio, and 5- to 10-membered heterocycloalkyl;

R²⁷ and R²⁸ are each independently selected from a hydrogen atom and optionally (C₁₋₄ alkoxy)carbonyl-substituted C₁₋₄ alkyl;

R³⁵ and R³⁶ are each independently selected from a hydrogen atom and C₁₋₄ alkyl; and

R³⁹ and R⁴⁰ are each independently selected from a hydrogen atom and optionally C₁₋₆ alkoxy-substituted C₁₋₆ alkyl.

[0075] (2-2) The compound according to (2-1) or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is optionally C₁₋₄ alkoxy-substituted C₁₋₅ alkoxy or a halogen atom;

Rc is a halogen atom, C₁₋₁₀ alkyl optionally substituted with one or more halogen

atoms, or a group -SF₅; and

R_d is cyano, hydroxy, a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R¹⁴, C₆₋₁₀ aryl optionally substituted with one or more substituents R¹⁴, or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R¹⁴.

[0076] (2-3) The compound according to (2-1) or (2-2) or a salt thereof, or a solvate of the compound or the salt, wherein

each R_a is independently selected from a halogen atom, nitro, cyano, (C₁₋₄ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyloxy, C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², and C₁₋₄ alkylthio optionally substituted with one or more substituents R¹³;

each R¹⁰ is independently selected from carboxy, 3- to 6-membered heterocycloalkyl, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, and a group -(O(CH₂)_o)_p-OH;

each R¹¹ is independently selected from hydroxy, carboxy, 5- or 6-membered heterocycloalkyl optionally substituted with one or more oxo groups, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, a group -(O(CH₂)_o)_p-OH, and C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups;

each R¹² is independently selected from a halogen atom, hydroxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyl, 5- or 6-membered heteroaryl, and a group -NR³⁹R⁴⁰;

R¹³ is 5- or 6-membered heterocycloalkyl; and

o and p are each independently an integer selected from 2 to 4.

[0077] (2-4) The compound according to any of (2-1) to (2-3) or a salt thereof, or a solvate of the compound or the salt, wherein each R_a is independently selected from a halogen atom, hydroxy, nitro, cyano, C₁₋₆ alkyl, (carboxy)C₁₋₈ alkyl, (morpholino)C₁₋₄ alkyl, (oxetanyl)C₁₋₄ alkyl, [HO-((CH₂)_oO)_p]C₁₋₆ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl optionally substituted with one or

more hydroxy groups, C₁₋₃ alkoxy substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, ((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkoxy, (morpholino)C₁₋₆ alkoxy optionally substituted with one or more oxo groups, (3- to 6-membered oxacycloalkyl)C₁₋₄ alkoxy, (pyridinyl)C₁₋₄ alkoxy, (pyrimidinyl)C₁₋₄ alkoxy, (C₁₋₆ alkoxy)C₁₋₈ alkoxy optionally substituted with one or more hydroxy groups, (carboxy)C₂₋₆ alkynyl, (morpholino)C₂₋₆ alkynyl optionally substituted with one or more oxo groups, (3- to 6-membered oxacycloalkyl)C₂₋₆ alkynyl, (pyrrolidino)C₂₋₆ alkynyl optionally substituted with one or more oxo groups, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)₆O)_p]C₂₋₈ alkynyl, (C₁₋₆ alkoxy)C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, (C₃₋₆ cycloalkyl)C₂₋₆ alkynyl optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy)carbonyl, (morpholino)C₁₋₄ alkylthio optionally substituted with one or more oxo groups, and 3- to 6-membered oxacycloalkyloxy.

[0078] (2-5) The compound according to any of (2-1) to (2-4) or a salt thereof, or a solvate of the compound or the salt, wherein each R_a is independently selected from a halogen atom, hydroxy, cyano, C₁₋₃ alkyl, (carboxy)C₁₋₈ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl substituted with one or more hydroxy groups, C₁₋₃ alkoxy optionally substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (morpholino)C₁₋₆ alkoxy, (C₁₋₆ alkoxy)C₁₋₈ alkoxy substituted with one or more hydroxy groups, (carboxy)C₂₋₆ alkynyl, (morpholino)C₂₋₆ alkynyl, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)₆O)_p]C₂₋₈ alkynyl, (C₁₋₆ alkoxy)C₂₋₈ alkynyl substituted with one or more hydroxy groups, and (C₁₋₃ alkoxy)carbonyl.

[0079] (2-6) The compound according to any of (2-1) to (2-5) or a salt thereof, or a solvate of the compound or the salt, wherein R³ is methyl substituted with Re.

[0080] (2-7) The compound according to any of (2-1) to (2-6) or a salt thereof, or a solvate of the compound or the salt, wherein R³ is benzyl optionally substituted with one or more substituents R_a on the benzene ring.

[0081] (2-8) The compound according to any of (2-1) to (2-7) or a salt thereof, or a solvate

of the compound or the salt, wherein R^3 is benzyl optionally substituted with one to three substituents R_a on the benzene ring.

[0082] (2-9) The compound according to any of (2-1), (2-4) and (2-6) to (2-8) or a salt thereof, or a solvate of the compound or the salt, wherein R_e is phenyl optionally substituted with one to three substituents R_a ;

the one to three substituents R_a are one substituent selected from R_i , R_j , and R_k , two substituents selected from combinations of R_i and R_j , R_i and R_k , and R_j and R_k , or three substituents R_i , R_j , and R_k ;

R_i is a halogen atom or C_{1-3} alkoxy;

R_j is a halogen atom, nitro, or cyano; and

R_k is hydroxy, a halogen atom, (C_{1-4} alkoxy)carbonyl, 5- to 10-membered heterocycloalkyloxy, C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkenyl optionally substituted with one or more substituents R^{15} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , or C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} .

[0083] (2-10) The compound according to (2-9) or a salt thereof, or a solvate of the compound or the salt, wherein R_k is hydroxy, a halogen atom, C_{1-6} alkyl, (carboxy) C_{1-8} alkyl, (C_{1-6} alkoxy) C_{1-8} alkyl substituted with one or more hydroxy groups, C_{1-3} alkoxy optionally substituted with one or more halogen atoms, [N-((C_{1-3} alkoxy) C_{1-4} alkyl)-N-(C_{1-3} alkyl)amino] C_{1-4} alkoxy, (morpholino) C_{1-6} alkoxy, (C_{1-6} alkoxy) C_{1-8} alkoxy substituted with one or more hydroxy groups, (carboxy) C_{2-8} alkynyl, (morpholino) C_{2-6} alkynyl, C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)₆O)_p] C_{2-8} alkynyl, (C_{1-6} alkoxy) C_{2-8} alkynyl substituted with one or more hydroxy groups, or (C_{1-3} alkoxy)carbonyl.

[0084] (2-11) The compound according to (2-1) or a salt thereof, or a solvate of the compound or the salt, wherein

R^1 , R^4 , and R^5 are as defined in the following (1) or (2):

(1) R^1 is a hydrogen atom or C_{1-6} alkyl;

R^4 is a hydrogen atom, C_{1-4} alkyl optionally substituted with one or more substituents R_f , C_{6-10} aryl optionally substituted with one or more substituents R_g , (C_{1-6} alkyl)carbonyl, (C_{6-10} aryl)carbonyl, $-C(O)NR^{37}R^{38}$, C_{3-7} cycloalkyl, or 5- to 8-membered heterocycloalkyl; and

R^5 is a hydrogen atom or C_{1-4} alkyl; or

(2) R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-6} saturated carbocyclic ring; and

R^4 is as defined above;

R^3 is C_{1-4} alkyl substituted with R_e ;

R_e is C_{6-10} aryl optionally substituted with one or more substituents R_a ;

each R_a is independently selected from a halogen atom, cyano, hydroxy, C_{1-6} alkyl optionally substituted with one or more substituents R^{10} , C_{2-6} alkynyl optionally substituted with one or more substituents R^{11} , and C_{1-6} alkoxy optionally substituted with one or more substituents R^{12} ;

R^{10} , R^{11} , and R^{12} are each independently selected from 4- to 10-membered heterocycloalkyl optionally substituted with one or more substituents and a group $-NR^{39}R^{40}$, wherein the substituents are each independently selected from oxo, a halogen atom, and C_{1-4} alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups);

Ar^1 is C_{6-10} aryl or 5- to 10-membered heteroaryl, wherein the aryl and heteroaryl groups are each optionally substituted with one to three substituents selected from R_b , R_c , and R_d ;

R_b , R_c , and R_d are each independently selected from a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, optionally C_{1-4} alkyl-substituted 5- to 10-membered heterocycloalkyl, C_{6-10} aryl optionally substituted with one or more substituents R^{14} , and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from a halogen atom, cyano, C_{1-4} alkyl optionally substituted with one or more halogen atoms, C_{1-4} alkoxy, a group $-SO_2NR^{35}R^{36}$, and C_{1-4}

alkylthio.

[0085] (2-12) The compound according to (2-11) or a salt thereof, or a solvate of the compound or the salt, wherein R^{10} , R^{11} , and R^{12} are each independently selected from 5- or 6-membered heterocycloalkyl optionally substituted with one or more substituents.

[0086] (2-13) The compound according to any of (2-1) and (2-11) or a salt thereof, or a solvate of the compound or the salt, wherein

R^1 , R^4 , and R^5 are as defined in the following (1) or (2):

(1) R^1 is C_{1-6} alkyl;

R^4 is optionally halogen atom-substituted C_{1-4} alkyl or phenyl; and

R^5 is a hydrogen atom or C_{1-4} alkyl; or

(2) R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-6} saturated carbocyclic ring; and

R^4 is as defined above;

R^3 is C_{1-4} alkyl substituted with Re ;

Re is phenyl optionally substituted with one or more substituents Ra ;

each Ra is independently selected from a halogen atom, optionally R^{11} -substituted C_{2-6} alkynyl, and optionally R^{12} -substituted C_{1-6} alkoxy;

R^{11} and R^{12} are each independently selected from 5- or 6-membered heterocycloalkyl and $-NR^{39}R^{40}$;

Ar^1 is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb , Rc , and Rd ;

Rb , Rc , and Rd are each independently selected from a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, phenyl optionally substituted with one or more substituents R^{14} , and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from a halogen atom, cyano, C_{1-4} alkyl optionally substituted with one or more halogen atoms, C_{1-4} alkoxy, a group $-SO_2NR^{35}R^{36}$ (wherein R^{35}

and R^{36} are each independently selected from C_{1-4} alkyl), and C_{1-4} alkylthio.

[0087] (2-14) The compound according to any of (2-11) to (2-13) or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is a halogen atom;

Rc is a halogen atom or C_{1-4} alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, phenyl optionally substituted with one or more substituents R^{14} , or 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14} .

[0088] (2-15) The compound according to (2-13) or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one or more substituents Ra;

R^{11} and R^{12} are each independently selected from morpholinyl and a group $-NR^{39}R^{40}$;

R^4 is optionally halogen atom-substituted C_{1-4} alkyl or phenyl;

Ar^1 is phenyl, pyridinyl, or pyrimidinyl, wherein the phenyl, pyridinyl, and pyrimidinyl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb is a halogen atom;

Rc is a halogen atom or C_{1-4} alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, phenyl, pyridinyl, or pyrimidinyl, wherein the phenyl, pyridinyl, and pyrimidinyl groups are each optionally substituted with one or more substituents R^{14} .

[0089] (2-16) The compound according to (2-15) or a salt thereof, or a solvate of the compound or the salt, wherein R^{11} and R^{12} are each independently selected from morpholinyl and [N-((methoxy)ethyl)-N-(methyl)amino].

[0090] (2-17) The compound according to (2-11) or (2-12) or a salt thereof, or a solvate of

the compound or the salt, wherein R^{10} , R^{11} , and R^{12} are each independently selected from morpholinyl, thiomorpholinyl, azetidiny, pyrrolidinyl, 2-oxa-6-azaspiro[3.3]heptanyl, 3,7-dioxa-9-azabicyclo[3.3.1]nonanyl, piperazinyl, piperidyl, and [N-((methoxy)ethyl)-N-(methyl)amino].

[0091] (2-18) The compound according to any of (2-1) to (2-17) or a salt thereof, or a solvate of the compound or the salt, wherein

R^1 , R^4 , and R^5 are as defined in any one of the following (1) to (3):

(1) R^1 is a hydrogen atom, and R^4 and R^5 are each independently selected from C_{1-6} alkyl;

(2) R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-5} saturated carbocyclic ring, and R^4 is C_{1-6} alkyl; and

(3) R^1 is linear C_{1-6} alkyl, and

R^4 and R^5 together with the carbon atom to which they are attached form a pyrrolidine ring;

R^3 is benzyl substituted with one or more substituents R_a on the benzene ring;

each R_a is independently selected from a halogen atom, (morpholino) C_{2-6} alkoxy, (morpholino) C_{1-6} alkyl, (carboxy) C_{2-6} alkynyl, (C_{1-6} alkoxy) C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, and [N-((methoxy)ethyl)-N-(methyl)amino]ethoxy;

Ar^1 is phenyl or pyrimidyl, wherein these groups are each substituted with one to three groups each independently selected from a halogen atom, trifluoromethyl, pyridinyl substituted with one or more substituents R^{14} , and pyrimidyl substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from methyl, cyano, trifluoromethyl, and methylthio.

[0092] (2-19) The compound according to any of (2-1) to (2-18) or a salt thereof, or a solvate of the compound or the salt, wherein Ar^1 is 4-(trifluoromethyl)-2-(6-methylthiopyridin-3-yl)phenyl, 4-(trifluoromethyl)-2-(6-trifluoromethylpyridin-3-yl)phenyl, 4-(trifluoromethyl)-2-(4-trifluoromethylpyrimidin-5-yl)phenyl, 4-(trifluoromethyl)-2-(6-

trifluoromethylpyrimidin-4-yl)phenyl, 4-(trifluoromethyl)-2-(6-cyano-5-methylpyrimidin-4-yl)phenyl, 4-(trifluoromethyl)-2-(2-cyanopyridin-4-yl)phenyl, 4-chloro-2-(6-methylthiopyridin-3-yl)phenyl, 4-chloro-2-(6-trifluoromethylpyridin-3-yl)phenyl, 4-chloro-2-(4-trifluoromethylpyrimidin-5-yl)phenyl, 4-chloro-2-(6-cyano-5-methylpyrimidin-4-yl)phenyl, 4-chloro-2-(6-trifluoromethylpyrimidin-4-yl)phenyl, or 4-chloro-2-(2-cyanopyridin-4-yl)phenyl.

[0093] (2-20) The compound according to any of (2-1) to (2-19) or a salt thereof, or a solvate of the compound or the salt, wherein

R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-6} saturated carbocyclic ring;

R^4 is C_{1-4} alkyl;

R^3 is C_{1-4} alkyl substituted with Re;

Re is phenyl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom and C_{1-4} alkoxy optionally substituted with one or more substituents R^{12} ;

each R^{12} is independently selected from 5- or 6-membered heterocycloalkyl and a group $-NR^{39}R^{40}$;

R^{39} and R^{40} are each independently selected from a hydrogen atom and optionally C_{1-4} alkoxy-substituted C_{1-4} alkyl;

Ar^1 is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from a halogen atom, cyano, C_{1-4} alkyl optionally substituted with one or more halogen atoms, and C_{1-4} alkylthio.

[0094] (2-21) The compound according to (2-20) or a salt thereof, or a solvate of the

compound or the salt, wherein

Rb is a halogen atom;

Rc is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen atoms; and

Rd is selected from a halogen atom and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R¹⁴.

[0095] (2-22) The compound according to any of (2-1), (2-20) and (2-21) or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri and Rj are each independently a halogen atom; and

Rk is C₁₋₄ alkoxy optionally substituted with one or more substituents R¹².

[0096] (2-23) The compound according to any of (2-20) to (2-22) or a salt thereof, or a solvate of the compound or the salt, wherein R³ is benzyl, wherein the benzyl group is optionally substituted with one to three substituents Ra.

[0097] (2-24) The compound according to any of (2-20) to (2-23) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar¹ is phenyl or pyridinyl;


Rd is pyridinyl or pyrimidinyl; and

R¹² is morpholinyl or [N-((methoxy)ethyl)-N-(methyl)amino].

[0098] (2-25) The compound according to any of (2-1) to (2-24) or a salt thereof, or a solvate of the compound or the salt, wherein the compound is represented by the formula I-a:

[0099]

[Formula 5]



[0100] wherein q is an integer selected from 0 to 3, and R³, R⁴, and Ar¹ are as defined above in any of (2-1) to (2-24).

[0101] (2-26) A compound selected from:

(4aR)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-[2-(6-cyano-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-[2-(2-cyanopyridin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

6-[4-[[[(4aR)-4-hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]hex-5-ynoic acid;

(4aR)-1-[[2,3-difluoro-4-(3-morpholin-4-ylprop-1-ynyl)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[3-[(2R)-2,3-dihydroxypropoxy]propoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[4-[(2R)-2,3-dihydroxypropoxy]butoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[6-[(2R)-2,3-dihydroxypropoxy]hexoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[2,3-difluoro-4-(morpholin-4-ylmethyl)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-(4-bromo-3,5-difluorophenyl)-1-[(3-chloro-2-fluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(3S)-3-tert-butyl-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-(6-methylsulfanylpyridin-3-yl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[2-(6-cyano-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-

hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

6-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;


6-[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide; and

7-[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide
or a salt thereof, or a solvate of the compound or the salt.

[0102] Another embodiment of the present invention provides the following compounds (3-1) to (3-13):

[0103] (3-1) A compound represented by the formula (I) or a salt thereof or a solvate of the compound or the salt:

[0104] [Formula 6]



[0105] wherein

R¹ is a hydrogen atom or C₁₋₁₀ alkyl;

R⁴ is a hydrogen atom, C₁₋₄ alkyl optionally substituted with one or more

substituents Rf, C₆₋₁₀ aryl optionally substituted with one or more substituents Rg, (C₁₋₆ alkyl)carbonyl, (C₆₋₁₀ aryl)carbonyl, -C(O)NR³⁷R³⁸, C₃₋₇ cycloalkyl, or 5- to 8-membered heterocycloalkyl;

R⁵ is a hydrogen atom or C₁₋₄ alkyl; or

R¹ and R⁵ together with the carbon atom to which they are attached may form a C₃₋₆ saturated carbocyclic ring; or

R⁴ and R⁵ together with the carbon atom and the nitrogen atom to which they are attached may form a 5- to 8-membered saturated heterocyclic ring, wherein the 5- to 8-membered saturated heterocyclic ring is optionally substituted with one or more substituents R²;

R³ is C₁₋₁₀ alkyl optionally substituted with one or more substituents Rh, or C₁₋₄ alkyl substituted with Re;

R³⁷ and R³⁸ are each independently selected from a hydrogen atom and C₁₋₃ alkyl;

each R² is independently selected from C₁₋₅ alkyl and a halogen atom; and/or

two or more substituents R² on the 5- to 8-membered saturated heterocyclic ring may together form C₁₋₅ alkylene that links the ring atoms to which they are attached;

each Rh is independently selected from a halogen atom, (C₁₋₄ alkoxy)carbonyl, and -(O(CH₂)_a)_b-C₁₋₄ alkoxy (wherein a is an integer selected from 2 to 4, and b is an integer selected from 1 to 4);

Re is C₆₋₁₀ aryl optionally substituted with one or more substituents Ra or 5- to 10-membered heteroaryl optionally substituted with one or more substituents Ra;

each Rf is independently selected from a halogen atom, hydroxy, cyano, carboxy, (C₁₋₆ alkoxy)carbonyl, C₁₋₆ alkoxy, and C₆₋₁₀ aryl optionally substituted with one or more substituents Rg;

each Rg is independently selected from a halogen atom, C₁₋₆ alkyl, C₂₋₆ alkynyl, and C₁₋₆ alkoxy, wherein the alkyl, alkynyl, and alkoxy groups are each optionally substituted with one or more substituents selected from hydroxy and cyano;

each Ra is independently selected from a halogen atom, hydroxy, nitro, cyano, (C₁₋₄

alkoxy)carbonyl, 5- to 10-membered heterocycloalkoxy, C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkenyl optionally substituted with one or more substituents R¹⁵, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², and C₁₋₄ alkylthio optionally substituted with one or more substituents R¹³;

the substituents R¹⁰, R¹¹, R¹², R¹³, and R¹⁵ are each independently selected from a halogen atom, hydroxy, carboxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₄ alkoxy)carbonyl, a group -(O(CH₂)_o)_p-OH (wherein o and p are each independently an integer selected from 2 to 4), C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups, 4- to 10-membered heterocycloalkyl, and 5- to 10-membered heteroaryl, wherein the 4- to 10-membered heterocycloalkyl is optionally substituted with one or more substituents selected from oxo, a halogen atom, and C₁₋₄ alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups);

Ar¹ is C₆₋₁₀ aryl or 5- to 10-membered heteroaryl, wherein the aryl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from optionally C₁₋₄ alkoxy-substituted C₁₋₅ alkoxy, a halogen atom, C₁₋₁₀ alkyl optionally substituted with one or more halogen atoms, a group -SF₅, cyano, hydroxy, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R¹⁴, C₆₋₁₀ aryl optionally substituted with one or more substituents R¹⁴, and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R¹⁴;

each R¹⁴ is independently selected from a halogen atom, oxo, cyano, nitro, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₁₋₄ alkoxy optionally substituted with one or more halogen atoms, (C₁₋₆ alkoxy)carbonyl, -NR²⁷R²⁸, -SO₂NR³⁵R³⁶, C₁₋₄ alkylthio, and 5- to 10-membered heterocycloalkyl;

R²⁷ and R²⁸ are each independently selected from a hydrogen atom and optionally (C₁₋₄ alkoxy)carbonyl-substituted C₁₋₄ alkyl; and

R^{35} and R^{36} are each independently selected from a hydrogen atom and C_{1-4} alkyl.

[0106] (3-2) The compound according to (3-1) or a salt thereof, or a solvate of the compound or the salt, wherein R^1 is C_{1-10} alkyl.

[0107] (3-3) The compound according to (3-1) or (3-2) or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is optionally C_{1-4} alkoxy-substituted C_{1-5} alkoxy or a halogen atom;

Rc is a halogen atom, C_{1-10} alkyl optionally substituted with one or more halogen atoms, or a group $-SF_5$; and

Rd is selected from cyano, hydroxy, a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R^{14} , C_{6-10} aryl optionally substituted with one or more substituents R^{14} , or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14} .

[0108] (3-4) The compound according to any of (3-1) to (3-3) or a salt thereof, or a solvate of the compound or the salt, wherein

each Ra is independently selected from a halogen atom, nitro, cyano, (C_{1-4} alkoxy)carbonyl, 5- or 6-membered heterocycloalkyloxy, C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , and C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} ;

each R^{10} is independently selected from carboxy, 5- or 6-membered heterocycloalkyl, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, and a group $-(O(CH_2)_o)_p-OH$;

each R^{11} is independently selected from hydroxy, carboxy, 5- or 6-membered heterocycloalkyl optionally substituted with one or more oxo groups, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, a group $-(O(CH_2)_o)_p-OH$, and C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups;

each R^{12} is independently selected from a halogen atom, hydroxy, C_{1-6} alkoxy

optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy)carbonyl, 5- or 6-membered heterocycloalkyl, and 5- or 6-membered heteroaryl;

each R¹³ is independently selected from 5- or 6-membered heterocycloalkyl; and

o and p are each independently an integer selected from 2 to 4.

[0109] (3-5) The compound according to any of (3-1) to (3-4) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹ is selected from C₁₋₄ alkyl;

R³ is selected from C₁₋₄ alkyl substituted with Re;

Re is selected from C₆₋₁₀ aryl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom and optionally R¹²-substituted C₁₋₆ alkoxy;

R⁴ is selected from C₁₋₄ alkyl;

R⁵ is selected from C₁₋₄ alkyl; or

of R¹, R⁴, and R⁵, R¹ and R⁵ together with the carbon atom to which they are attached form a C₃₋₆ saturated carbocyclic ring, or R⁴ and R⁵ together with the nitrogen atom and the carbon atom to which they are attached form a 5- to 8-membered saturated heterocyclic ring;

Ar¹ is C₆₋₁₀ aryl, wherein the aryl group is substituted with two substituents selected from Rc and Rd;

Rc is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen atoms;

Rd is 5- to 10-membered heterocycloalkyl optionally substituted with one or more C₁₋₄ alkyl groups and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R¹⁴;

R¹² is 5- to 10-membered heterocycloalkyl; and

R¹⁴ is C₁₋₄ alkyl optionally substituted with one or more halogen atoms.

[0110] (3-6) The compound according to (3-1) or (3-2) or a salt thereof, or a solvate of the

compound or the salt, wherein

R^1 is a hydrogen atom or C_{1-6} alkyl;

R^3 is C_{1-4} alkyl substituted with Re;

Re is C_{6-10} aryl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom, cyano, hydroxy, C_{1-6} alkyl optionally substituted with one or more substituents R^{10} , C_{2-6} alkynyl optionally substituted with one or more substituents R^{11} , and C_{1-6} alkoxy optionally substituted with one or more substituents R^{12} ;

R^{10} , R^{11} , and R^{12} are each independently selected from 4- to 10-membered heterocycloalkyl optionally substituted with one or more substituents, wherein the substituents are each independently selected from oxo, a halogen atom, and C_{1-4} alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups);

R^4 is a hydrogen atom, C_{1-4} alkyl optionally substituted with one or more substituents Rf, C_{6-10} aryl optionally substituted with one or more substituents Rg, (C_{1-6} alkyl)carbonyl, (C_{6-10} aryl)carbonyl, $-C(O)NR^{37}R^{38}$, C_{3-7} cycloalkyl, or 5- to 8-membered heterocycloalkyl;

R^5 is a hydrogen atom or C_{1-4} alkyl; or

R^1 and R^5 together with the carbon atom to which they are attached may form a C_{3-6} saturated carbocyclic ring;

Ar^1 is C_{6-10} aryl or 5- to 10-membered heteroaryl, wherein the aryl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, optionally C_{1-4} alkyl-substituted 5- to 10-membered heterocycloalkyl, C_{6-10} aryl optionally substituted with one or more substituents R^{14} , and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from a halogen atom, cyano, C_{1-4} alkyl optionally

substituted with one or more halogen atoms, C₁₋₄ alkoxy, -SO₂NR³⁵R³⁶, and C₁₋₄ alkylthio.

[0111] (3-7) The compound according to any of (3-1), (3-2), and (3-6) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹ is C₁₋₆ alkyl;

R³ is C₁₋₄ alkyl substituted with Re;

Re is C₆₋₁₀ aryl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom, optionally R¹¹-substituted C₂₋₆ alkynyl, and optionally R¹²-substituted C₁₋₆ alkoxy;

R¹¹ and R¹² are each independently selected from 5- to 10-membered heterocycloalkyl;

R⁴ is optionally halogen atom-substituted C₁₋₄ alkyl or C₆₋₁₀ aryl;

R⁵ is a hydrogen atom or C₁₋₄ alkyl; or

R¹ and R⁵ together with the carbon atom to which they are attached may form a C₃₋₆ saturated carbocyclic ring;

Ar¹ is C₆₋₁₀ aryl or 5- to 10-membered heteroaryl, wherein the aryl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₆₋₁₀ aryl optionally substituted with one or more substituents R¹⁴, and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R¹⁴; and

each R¹⁴ is independently selected from a halogen atom, cyano, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₁₋₄ alkoxy, -SO₂NR³⁵R³⁶ (wherein R³⁵ and R³⁶ are each independently selected from C₁₋₄ alkyl), and C₁₋₄ alkylthio.

[0112] (3-8) The compound according to (3-7) or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is a halogen atom;

Rc is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen

atoms; and

Rd is a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₆₋₁₀ aryl optionally substituted with one or more substituents R¹⁴, or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R¹⁴.

[0113] (3-9) The compound according to (3-7) or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one or more substituents Ra;

R¹¹ and R¹² are each independently morpholinyl;

R⁴ is optionally halogen atom-substituted C₁₋₄ alkyl or phenyl;

Ar¹ is phenyl, pyridinyl, or pyrimidinyl, wherein the phenyl, pyridinyl, and pyrimidinyl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb is a halogen atom;

Rc is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, phenyl, pyridinyl, or pyrimidinyl, wherein the phenyl, pyridinyl, and pyrimidinyl groups are each optionally substituted with one or more substituents R¹⁴.

[0114] (3-10) The compound according to (3-6) or (3-7) or a salt thereof, or a solvate of the compound or the salt, wherein R¹⁰, R¹¹, and R¹² are each independently selected from morpholinyl, thiomorpholinyl, azetidiny, pyrrolidinyl, 2-oxa-6-azaspiro[3.3]heptanyl, 3,7-dioxa-9-azabicyclo[3.3.1]nonanyl, piperazinyl, and piperidyl.

[0115] (3-11) The compound according to any of (3-1) to (3-10) or a salt thereof, or a solvate of the compound or the salt, wherein R³ is benzyl optionally substituted with one or more substituents Ra on the benzene ring.

[0116] (3-12) The compound according to (3-1) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹ is C₁₋₆ alkyl, and R⁴ and R⁵ together with the carbon atom to which they are

attached form a pyrrolidine ring; or

R^1 is a hydrogen atom, and R^4 and R^5 are each independently selected from C_{1-6} alkyl; or

R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-5} saturated carbocyclic ring, and R^4 is C_{1-6} alkyl;

R^3 is benzyl substituted with one or more substituents R_a on the benzene ring;

each R_a is independently selected from a halogen atom, (morpholino) C_{2-6} alkoxy, (morpholino) C_{1-6} alkyl, (carboxy) C_{2-6} alkynyl, and (C_{1-6} alkoxy) C_{1-6} alkoxy optionally substituted with one or more hydroxy groups;

Ar^1 is phenyl or pyrimidyl, wherein these groups are each substituted with one to three groups each independently selected from a halogen atom, trifluoromethyl, pyridinyl substituted with one or more substituents R^{14} , and pyrimidyl substituted with one or more substituents R^{14} ; and

each R^{14} is independently selected from methyl, cyano, trifluoromethyl, and methylthio.

[0117] (3-13) The compound according to any of (3-1) to (3-11) or a salt thereof, or a solvate of the compound or the salt, wherein Ar^1 is 4-(trifluoromethyl)-2-(6-methylthiopyridin-3-yl)phenyl, 4-(trifluoromethyl)-2-(6-trifluoromethylpyridin-3-yl)phenyl, 4-(trifluoromethyl)-2-(4-trifluoromethylpyrimidin-5-yl)phenyl, 4-(trifluoromethyl)-2-(6-trifluoromethylpyrimidin-4-yl)phenyl, 4-(trifluoromethyl)-2-(6-cyano-5-methylpyrimidin-4-yl)phenyl, 4-(trifluoromethyl)-2-(2-cyanopyridin-4-yl)phenyl, 4-chloro-2-(6-methylthiopyridin-3-yl)phenyl, 4-chloro-2-(6-trifluoromethylpyridin-3-yl)phenyl, 4-chloro-2-(4-trifluoromethylpyrimidin-5-yl)phenyl, 4-chloro-2-(6-cyano-5-methylpyrimidin-4-yl)phenyl, 4-chloro-2-(6-trifluoromethylpyrimidin-4-yl)phenyl, or 4-chloro-2-(2-cyanopyridin-4-yl)phenyl.

[0118] An alternative aspect of the present invention provides a pharmaceutical drug comprising a compound according to any of (2-1) to (2-26) and (3-1) to (3-13) or a salt thereof, or a solvate of the compound or the salt.

[0119] A further alternative aspect of the present invention provides a pharmaceutical composition comprising a compound according to any of (2-1) to (2-26) and (3-1) to (3-13) or a salt thereof, or a solvate of the compound or the salt.

[0120] A further alternative aspect of the present invention provides the compound according to any of (2-1) to (2-26) and (3-1) to (3-13) or the salt thereof, or the solvate of the compound or the salt for use in the prevention and/or treatment of a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure.

[0121] A further alternative aspect of the present invention provides a NaPi-IIb inhibitor comprising a compound according to any of (2-1) to (2-26) and (3-1) to (3-13) or a salt thereof, or a solvate of the compound or the salt.

[0122] A further alternative aspect of the present invention provides a preventive and/or therapeutic agent for a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, the agent comprising a compound according to any of (2-1) to (2-26) and (3-1) to (3-13) or a salt thereof, or a solvate of the compound or the salt as an active ingredient.


[0123] A further alternative aspect of the present invention provides a method for preventing and/or treating a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, comprising administering a therapeutically effective amount of a compound according to any of (2-1) to (2-26) and (3-1) to (3-13) or a salt thereof, or a solvate of the compound or the salt to a patient.

[0124] According to one embodiment of the compound represented by the formula (I) wherein R⁴ and R⁵ together with the carbon atom and the nitrogen atom to which they are attached form a 5- to 8-membered saturated heterocyclic ring, the following compound (4-1) is provided:

[0125] (4-1) A compound represented by the formula (II) or a salt thereof, or a solvate of the compound or the salt:

[0126]

[Formula 7]



[0127] wherein R^{101} is selected from a hydrogen atom and linear C_{1-10} alkyl;

R^{102} may be the same or different and are each independently selected from a hydrogen atom, C_{1-5} alkyl, and a halogen atom; or

two R^{102} moieties bonded to different carbon atoms may together form C_{1-5} alkylene that links the carbon atoms;

$R^{102'}$ may be the same or different and are each independently selected from a hydrogen atom, C_{1-5} alkyl, and a halogen atom;

n is an integer selected from 1 to 4;

R^{103} is C_{1-10} alkyl optionally substituted with one or more substituents R_w or C_{1-4} alkyl substituted with one or more substituents R_x ;

each R_w is independently selected from a halogen atom, $(C_{1-4} \text{ alkoxy})\text{carbonyl}$, and a group $-(O(CH_2)_a)_b-C_{1-4} \text{ alkoxy}$ (wherein a is an integer selected from 2 to 4, and b is an integer selected from 1 to 4);

each R_x is independently selected from C_{6-10} aryl optionally substituted with one or more substituents R_y and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R_y ;

each R_y is independently selected from a halogen atom, hydroxy, nitro, cyano, $(C_{1-4} \text{ alkoxy})\text{carbonyl}$, 5- to 10-membered heterocycloalkoxy, C_{1-10} alkyl optionally substituted with one or more substituents R^{110} , C_{2-10} alkenyl optionally substituted with one or more substituents R^{115} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{111} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{112} , and C_{1-4} alkylthio optionally

substituted with one or more substituents R^{113} ;

R^{110} , R^{111} , R^{112} , R^{113} , and R^{115} are each independently selected from a halogen atom, hydroxy, carboxy, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, (C_{1-4} alkoxy)carbonyl, a group $-(O(CH_2)_o)_p-OH$ (wherein o and p are each independently an integer selected from 2 to 4), C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups, 5- to 10-membered heterocycloalkyl optionally substituted with one or more oxo groups, and 5- to 10-membered heteroaryl;

Ar^{101} is C_{6-10} aryl or 5- to 10-membered heteroaryl, wherein the C_{6-10} aryl and 5- to 10-membered heteroaryl groups are each optionally substituted with one to three substituents selected from Rz^1 , Rz^2 , and Rz^3 ;

Rz^1 , Rz^2 , and Rz^3 are each independently selected from optionally C_{1-4} alkoxy-substituted C_{1-5} alkoxy, a halogen atom, C_{1-10} alkyl optionally substituted with one or more halogen atoms, a group $-SF_5$, cyano, hydroxy, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R^{114} , C_{6-10} aryl optionally substituted with one or more substituents R^{114} , and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{114} ; and

each R^{114} is independently selected from a halogen atom, cyano, nitro, C_{1-4} alkyl optionally substituted with one or more halogen atoms, C_{1-4} alkoxy optionally substituted with one or more halogen atoms, (C_{1-6} alkoxy)carbonyl, a group $-NR^{127}R^{128}$ (wherein R^{127} and R^{128} are each independently selected from a hydrogen atom and optionally (C_{1-4} alkoxy)carbonyl-substituted C_{1-4} alkyl), a group $-SO_2NR^{135}R^{136}$ (wherein R^{135} and R^{136} are each independently selected from a hydrogen atom and C_{1-4} alkyl), C_{1-4} alkylthio, and 5- to 10-membered heterocycloalkyl.

[0128] According to another embodiment, the present invention provides the following compounds (4-2) to (4-12) represented by the formula (II).

[0129] (4-2) The compound according to (4-1) or a salt thereof, or a solvate of the compound or the salt, wherein

Rz^1 is optionally C_{1-4} alkoxy-substituted C_{1-5} alkoxy or a halogen atom;

Rz^2 is a halogen atom, C_{1-10} alkyl optionally substituted with one or more halogen atoms, and a group $-SF_5$; and

Rz^3 is cyano, hydroxy, a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R^{114} , C_{6-10} aryl optionally substituted with one or more substituents R^{114} , or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{114} .

[0130] (4-3) The compound according to (4-1) or (4-2) or a salt thereof, or a solvate of the compound or the salt, wherein Rz^3 is cyano, hydroxy, a halogen atom, C_{1-4} alkyl optionally substituted with one or more halogen atoms, phenyl, and 5- to 6-membered heteroaryl, wherein the phenyl and 5- to 6-membered heteroaryl may be substituted with one or two substituents R^{114} .

[0131] (4-4) The compound according to any of (4-1) to (4-3) or a salt thereof, or a solvate of the compound or the salt, wherein

R^{101} is a hydrogen atom or linear C_{1-6} alkyl;

R^{102} may be the same or different and are each independently selected from a hydrogen atom, C_{1-3} alkyl, and a halogen atom;

$R^{102'}$ may be the same or different and are each independently selected from a hydrogen atom, C_{1-3} alkyl, and a halogen atom;

n is an integer selected from 1 to 3;

Rx is independently selected from C_{6-10} aryl optionally substituted with one or more substituents Ry and 5- to 10-membered heteroaryl optionally substituted with one or more substituents Ry ;

Ry is independently selected from a halogen atom, hydroxy, nitro, cyano, (C_{1-3} alkoxy)carbonyl, 5- to 10-membered heterocycloalkyloxy, C_{1-10} alkyl optionally substituted with one or more substituents R^{110} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{111} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{112} , and C_{1-3} alkylthio optionally substituted with one or more substituents R^{113} ;

R^{110} is independently selected from carboxyl, 5- to 10-membered heterocycloalkyl,

C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, and a group -(O(CH₂)_o)_p-OH (wherein o and p are each independently an integer selected from 2 to 4);

R¹¹¹ is independently selected from hydroxy, carboxy, 5- to 10-membered heterocycloalkyl optionally substituted with one or more oxo groups, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, a group -(O(CH₂)_o)_p-OH (wherein o and p are each independently an integer selected from 2 to 4), and C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups;

R¹¹² is independently selected from a halogen atom, hydroxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy)carbonyl, 5- to 10-membered heterocycloalkyl, and 5- to 10-membered heteroaryl;

R¹¹³ is independently selected from 5- to 10-membered heterocycloalkyl;

R^{z1} is selected from optionally C₁₋₃ alkoxy-substituted C₁₋₄ alkoxy and a halogen atom; and

R¹¹⁴ is independently selected from a halogen atom, cyano, nitro, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₁₋₄ alkoxy optionally substituted with one or more halogen atoms, (C₁₋₆ alkoxy)carbonyl, a group -NR¹²⁷R¹²⁸ (wherein R¹²⁷ and R¹²⁸ are each independently selected from a hydrogen atom and optionally (C₁₋₃ alkoxy)carbonyl-substituted C₁₋₄ alkyl), a group -SO₂NR¹³⁵R¹³⁶ (wherein R¹³⁵ and R¹³⁶ are each independently selected from a hydrogen atom and C₁₋₃ alkyl), C₁₋₃ alkylthio, and 5- to 10-membered heterocycloalkyl.

[0132] (4-5) The compound according to any of (4-1) to (4-4) or a salt thereof, or a solvate of the compound or the salt, wherein Rx is C₆₋₁₀ aryl optionally substituted with one or more substituents Ry.

[0133] (4-6) The compound according to any of (4-1) to (4-5) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar¹⁰¹ is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with R^{z1}, R^{z2}, and/or R^{z3};

Rz^1 is a halogen atom or optionally C_{1-3} alkoxy-substituted C_{1-4} alkoxy;

Rz^2 is a halogen atom or C_{1-10} alkyl optionally substituted with one or more halogen atoms; and

Rz^3 is 5- or 6-membered heterocycloalkyl optionally substituted with one or more substituents R^{114} , phenyl optionally substituted with one or more substituents R^{114} , or 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{114} .

[0134] (4-7) The compound according to any of (4-1) to (4-6) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar^{101} is phenyl or 5- or 6-membered heteroaryl, wherein the aryl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rz^1 , Rz^2 , and Rz^3 ;

Rz^1 is a halogen atom;

Rz^2 is selected from a halogen atom, C_{1-10} alkyl optionally substituted with one or more halogen atoms, and a group $-SF_5$; and

Rz^3 is cyano, hydroxy, or a halogen atom.

[0135] (4-8) The compound according to any of (4-1) to (4-7) or a salt thereof, or a solvate of the compound or the salt, wherein Rz^3 is phenyl, thienyl, pyridinyl, pyrimidinyl, or quinolinyl, wherein these groups are each optionally substituted with one or more substituents R^{114} .

[0136] (4-9) The compound according to any of (4-1) to (4-8) or a salt thereof, or a solvate of the compound or the salt, wherein each R^{114} is independently selected from a halogen atom, cyano, nitro, C_{1-4} alkyl optionally substituted with one or more halogen atoms, C_{1-4} alkoxy optionally substituted with one or more halogen atoms, $(C_{1-6}$ alkoxy)carbonyl, a group $-NR^{127}R^{128}$ (each R^{127} and R^{128} are independently selected from a hydrogen atom, and C_{1-4} alkyl optionally substituted with $(C_{1-4}$ alkoxy)carbonyl), a group $-SO_2NR^{135}R^{136}$ (each R^{135} and R^{136} are independently selected from a hydrogen atom, and C_{1-4} alkyl), and C_{1-4} alkylthio.

[0137] (4-10) The compound according to any of (4-1) to (4-9) or a salt thereof, or a solvate of the compound or the salt, wherein R^{101} is linear C_{1-6} alkyl.

[0138] (4-11) The compound according to any of (4-1) to (4-10) or a salt thereof, or a solvate of the compound or the salt, wherein R^{102} and $R^{102'}$ are a hydrogen atom.

[0139] (4-12) The compound according to any of (4-1) to (4-11) or a salt thereof, or a solvate of the compound or the salt, wherein

R^{101} is linear C_{1-6} alkyl;

R^{102} and $R^{102'}$ are a hydrogen atom;

R^{103} is C_{1-10} alkyl optionally substituted with R_x ;

R_x is phenyl optionally substituted with one or more substituents R_y ;

each R_y is independently selected from a halogen atom and C_{1-8} alkoxy optionally substituted with R^{112} ;

R^{112} is 5- or 6-membered heterocycloalkyl;

Ar^{101} is phenyl optionally substituted with one to three substituents selected from Rz^1 , Rz^2 , and Rz^3 ;

Rz^1 is optionally C_{1-3} alkoxy-substituted C_{1-4} alkoxy or a halogen atom;

Rz^2 is a halogen atom or C_{1-10} alkyl optionally substituted with one or more halogen atoms;

Rz^3 is 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{114} ; and

each R^{114} is selected independently from C_{1-4} alkyl optionally substituted with one or more halogen atoms.

[0140] (4-13) The compound according to any of (4-1) to (4-12) or a salt thereof, or a solvate of the compound or the salt, wherein Rz^3 is substituted with one or two substituents R^{114} .

[0141] (4-14) The compound according to any of (4-1) to (4-13) or a salt thereof, or a solvate of the compound or the salt, wherein R^{110} , R^{111} , R^{112} , R^{113} , and R^{115} are each independently selected from tetrahydrofuranyl, pyrrolidinyl, morpholinyl, cyclopentyl, and pyridinyl.

[0142] (4-15) The compound according to any of (4-1) to (4-14) or a salt thereof, or a

solvate of the compound or the salt, wherein R^{103} is benzyl optionally substituted with one or more substituents R_y on the benzene ring.

[0143] (4-16) The compound according to any of (4-1) to (4-15) or a salt thereof, or a solvate of the compound or the salt, wherein R^{103} is benzyl optionally substituted with one to three substituents R_y on the benzene ring.

[0144] (4-17) The compound according to any of (4-1) to (4-16) or a salt thereof, or a solvate of the compound or the salt, wherein the one to three substituents R_y are one substituent selected from R_i' , R_j' , and R_k' , two substituents selected from combinations of R_i' and R_j' , R_i' and R_k' , and R_j' and R_k' , or three substituents R_i' , R_j' , and R_k' ;

R_i' is a halogen atom or C_{1-3} alkoxy;

R_j' is a halogen atom, nitro, or cyano; and

R_k' is hydroxy, a halogen atom, C_{1-6} alkyl, (carboxy) C_{1-8} alkyl, (morpholino) C_{1-4} alkyl, (C_{1-6} alkoxy) C_{1-8} alkyl, (the alkoxy is optionally substituted with one or more hydroxy groups), (5- to 6-membered heterocycloalkyl) C_{1-6} alkoxy, $[HO-((CH_2)_6O)_p]C_{1-6}$ alkyl, (carboxy) C_{2-6} alkynyl, (3- to 6-membered heterocycloalkyl) C_{2-6} alkynyl (the heterocycloalkyl moiety is optionally substituted with oxo), C_{1-3} alkoxy optionally substituted with one or more halogen atoms, C_{1-6} alkoxy substituted with one or more hydroxy groups, ((C_{1-3} alkoxy)carbonyl) C_{1-3} alkoxy, (pyridinyl) C_{1-4} alkoxy, (C_{1-6} alkoxy) C_{1-8} alkoxy (the C_{1-6} alkoxy is optionally substituted with one or more hydroxy groups), optionally hydroxy-substituted C_{2-8} alkynyl, (C_{3-6} cycloalkyl) C_{2-6} alkynyl (the cycloalkyl is optionally substituted with a hydroxy group), $[HO-((CH_2)_6O)_p]C_{2-8}$ alkynyl, (C_{1-6} alkoxy) C_{2-8} alkynyl (the alkoxy is optionally substituted with one or more hydroxy groups), (morpholino) C_{1-4} alkylthio, 3- to 6-membered oxacycloalkyloxy, or (C_{1-3} alkoxy)carbonyl.

[0145] An alternative embodiment of the present invention provides the following compounds (5-1) to (5-12) represented by the formula (I):

[0146] (5-1) The compound according to (1-1), (2-1) or (3-1) or a salt thereof, or a solvate of the compound or the salt, wherein

R^1 is a hydrogen atom or linear C_{1-10} alkyl; and

R⁴ and R⁵ together with the carbon atom and the nitrogen atom to which they are attached form a 5- to 8-membered saturated heterocyclic ring, wherein the saturated heterocyclic ring is optionally substituted with one or more substituents R².

[0147] (5-2) The compound according to (5-1) or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is optionally C₁₋₄ alkoxy-substituted C₁₋₅ alkoxy or a halogen atom; and/or

Rc is a halogen atom, C₁₋₁₀ alkyl optionally substituted with one or more halogen atoms, or a group -SF₅; and/or

Rd is cyano, hydroxy, a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R¹⁴, C₆₋₁₀ aryl optionally substituted with one or more substituents R¹⁴, or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R¹⁴.

[0148] (5-3) The compound according to (5-1) or (5-2) or a salt thereof, or a solvate of the compound or the salt, wherein

R¹ is a hydrogen atom or linear C₁₋₆ alkyl;

R² may be the same or different and are each independently selected from C₁₋₃ alkyl and a halogen atom;

Re is C₆₋₁₀ aryl optionally substituted with one or more substituents Ra or 5- to 10-membered heteroaryl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom, hydroxy, nitro, cyano, (C₁₋₃ alkoxy)carbonyl, 5- to 10-membered heterocycloalkyloxy, C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², and C₁₋₃ alkylthio optionally substituted with one or more substituents R¹³;

each R¹⁰ is independently selected from carboxyl, 5- to 10-membered heterocycloalkyl, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, and a group -(O(CH₂)_o)_p-OH (wherein o and p are each independently an integer selected from 2 to 4);

each R¹¹ is independently selected from hydroxy, carboxy, 5- to 10-membered heterocycloalkyl optionally substituted with one or more oxo groups, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, a group -(O(CH₂)_o)_p-OH (wherein o and p are each independently an integer selected from 2 to 4), and C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups;

each R¹² is selected from a halogen atom, hydroxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy)carbonyl, 5- to 10-membered heterocycloalkyl, and 5- to 10-membered heteroaryl;

each R¹³ is independently selected from 5- to 10-membered heterocycloalkyl;

Rb is optionally C₁₋₃ alkoxy-substituted C₁₋₄ alkoxy or a halogen atom; and

each R¹⁴ is independently selected from a halogen atom, cyano, nitro, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₁₋₄ alkoxy optionally substituted with one or more halogen atoms, (C₁₋₆ alkoxy)carbonyl, a group -NR²⁷R²⁸ (wherein R²⁷ and R²⁸ are each independently selected from a hydrogen atom and optionally (C₁₋₃ alkoxy)carbonyl-substituted C₁₋₄ alkyl), a group -SO₂NR³⁵R³⁶ (wherein R³⁵ and R³⁶ are each independently selected from a hydrogen atom and C₁₋₃ alkyl), C₁₋₃ alkylthio, and 5- to 10-membered heterocycloalkyl.

[0149] (5-4) The compound according to any of (5-1) to (5-3) or a salt thereof, or a solvate of the compound or the salt, wherein Re is selected from C₆₋₁₀ aryl optionally substituted with one or more substituents Ra.

[0150] (5-5) The compound according to any of (5-1) to (5-4) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar¹ is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with Rb, Rc, and/or Rd;

Rb is a halogen atom or optionally C₁₋₃ alkoxy-substituted C₁₋₄ alkoxy; and/or

Rc is a halogen atom or C₁₋₁₀ alkyl optionally substituted with one or more halogen atoms; and/or

Rd is 5- or 6-membered heterocycloalkyl optionally substituted with one or more

substituents R^{14} , phenyl optionally substituted with one or more substituents R^{14} , or 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14} .

[0151] (5-6) The compound according to any of (5-1) to (5-5) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar^1 is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from R_b , R_c , and R_d ;

R_b is a halogen atom;

R_c is a halogen atom, C_{1-10} alkyl optionally substituted with one or more halogen atoms, or a group $-SF_5$; and

R_d is cyano, hydroxy, or a halogen atom.

[0152] (5-7) The compound according to any of (5-1) to (5-6) or a salt thereof, or a solvate of the compound or the salt, wherein R_d is phenyl, thienyl, pyridinyl, pyrimidinyl, or quinolinyl, wherein these groups are each optionally substituted with one or more substituents R^{14} .

[0153] (5-8) The compound according to any of (5-1) to (5-7) or a salt thereof, or a solvate of the compound or the salt, wherein R^1 is linear C_{1-6} alkyl.

[0154] (5-9) The compound according to any of (5-1) to (5-8) or a salt thereof, or a solvate of the compound or the salt, wherein R^2 is a hydrogen atom.

[0155] (5-10) The compound according to any of (5-1) to (5-9) or a salt thereof, or a solvate of the compound or the salt, wherein

R^1 is linear C_{1-6} alkyl;

R^2 is a hydrogen atom;

R^3 is C_{1-4} alkyl substituted with R_e ;

R_e is phenyl optionally substituted with one or more substituents R_a ;

each R_a is independently selected from a halogen atom and optionally R^{12} -

substituted C_{1-8} alkoxy;

each R^{12} is independently selected from 5- or 6-membered heterocycloalkyl;

Ar¹ is phenyl optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb is optionally C₁₋₃ alkoxy-substituted C₁₋₄ alkoxy and a halogen atom; and/or

Rc is a halogen atom or C₁₋₁₀ alkyl optionally substituted with one or more halogen atoms; and/or

Rd is 5- or 6-membered heteroaryl optionally substituted with one or more substituents R¹⁴; and

each R¹⁴ is independently selected from C₁₋₄ alkyl optionally substituted with one or more halogen atoms.


[0156] (5-11) The compound according to any of (5-1) to (5-10) or a salt thereof, or a solvate of the compound or the salt, wherein R¹⁰, R¹¹, R¹², R¹³, and R¹⁵ are each independently selected from tetrahydrofuranyl, pyrrolidinyl, morpholinyl, cyclopentyl, and pyridinyl.

[0157] (5-12) The compound according to any of (5-1) to (5-11) or a salt thereof, or a solvate of the compound or the salt, wherein R³ is benzyl optionally substituted with one or more substituents Ra on the benzene ring.

[0158] An alternative embodiment of the present invention provides the following compounds (6-1) to (6-10) represented by the formula (I):

[0159] (6-1) A compound represented by the formula I-b or a salt thereof, or a solvate of the compound or the salt:

[0160] [Formula 8]



[0161] wherein

$R^{4'}$ is selected from C_{1-4} alkyl;

q is an integer selected from 0 to 3;

$R^{3'}$ is selected from C_{1-4} alkyl substituted with Re' ;

Re' is C_{6-10} aryl, wherein the aryl group is optionally substituted with one to three substituents each independently selected from a halogen atom, C_{1-4} alkoxy optionally substituted with one or more substituents $R^{12'}$, 4- to 6-membered heterocycloalkyloxy (the heterocycloalkyloxy group is optionally substituted with optionally C_{1-4} alkoxy-substituted C_{1-4} alkyl), a group $-(O(CH_2)_{q1})_{q2}-NR^{41'}R^{42'}$ (wherein $q1$ is an integer selected from 1 to 4, and $q2$ is an integer selected from 2 to 6), a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43'}R^{44'}$ (wherein $r1$ is an integer selected from 1 to 4, and $r2$ is an integer selected from 1 to 4), a group $-(O(CH_2)_{s1})_{s2}-NR^{45'}-C(O)R^{46'}$ (wherein $s1$ and $s2$ are each independently an integer selected from 2 to 4), a group $-C(O)NR^{47'}R^{48'}$, pyridinyl, pyrrolyl, a group $-NR^{49'}R^{50'}$, and a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-C(O)NR^{51'}R^{52'}$ (wherein $y1$ is an integer selected from 1 to 4, and $y2$ is an integer selected from 1 to 4);

each $R^{12'}$ is independently selected from $(C_{1-4} \text{ alkoxy})C_{1-6}$ alkoxy, 5- or 6-membered heterocycloalkyl, and a group $-NR^{39'}R^{40'}$, wherein the 5- or 6-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, C_{1-4} alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups), $(C_{1-4} \text{ alkoxy})C_{1-4}$ alkyl (wherein the alkoxy moiety is optionally substituted with one or more hydroxy groups), C_{1-4} alkoxy, C_{1-4} alkoxycarbonyl, C_{1-4} alkylthio, morpholino, $(C_{1-3} \text{ alkyl})$ sulfonyl, and $-C(O)NR^{53'}R^{54'}$;

$R^{39'}$ is selected from a hydrogen and optionally C_{1-6} alkoxy-substituted C_{1-6} alkyl;

$R^{40'}$ is independently selected from a hydrogen atom, optionally C_{1-6} alkoxy-substituted C_{1-6} alkyl, $((C_{1-4} \text{ alkoxy})\text{carbonyl})C_{1-6}$ alkyl, hydroxy-substituted C_{1-6} alkyl, a group $-(CH_2)_u-NR^{55'}R^{56'}$ (wherein u is an integer selected from 1 to 4), a group $-CH((CH_2)_{v1}COOR^{57'})-(CH_2)_{v2}-COOR^{57'}$ (wherein $v1$ is an integer selected from 0 to 2, and $v2$ is an integer selected from 1 to 3), a group $-(CH_2)_w-SO_3H$ (wherein w is an integer selected from 1 to 4), a group $-(CH_2)_{x1}-CH(COOH)-(CH_2)_{x2}-SO_3H$ (wherein $x1$ is an integer

selected from 0 to 2, and x2 is an integer selected from 1 to 3), 3- to 6-membered oxacycloalkyl, and a group $-(CH_2)_{t1}-O-(CH_2)_{t2}-C(O)NR^{58'}R^{59'}$ (wherein t1 and t2 are each independently an integer selected from 1 to 3);

$R^{41'}$ is a hydrogen atom or C_{1-3} alkyl;

$R^{42'}$ is C_{1-8} alkyl substituted with one or more hydroxy groups or $(C_{1-3} \text{ alkoxy})C_{1-4}$ alkyl;

$R^{43'}$ is a hydrogen atom or C_{1-3} alkyl;

$R^{44'}$ is C_{1-8} alkyl substituted with one or more hydroxy groups, wherein

$R^{43'}$ and $R^{44'}$ together with the nitrogen atom to which they are attached may form morpholino;

$R^{45'}$ is a hydrogen atom and C_{1-3} alkyl;

$R^{46'}$ is C_{1-6} alkyl substituted with one or more hydroxy groups;

$R^{47'}$ is C_{1-3} alkyl;

$R^{48'}$ is $(C_{1-3} \text{ alkoxy})C_{1-4}$ alkyl;

$R^{49'}$ is a hydrogen atom or C_{1-4} alkyl;

$R^{50'}$ is $-(CH_2)_z-NR^{60'}R^{61'}$ (z is an integer selected from 1 to 4, $R^{60'}$ is a hydrogen atom and C_{1-4} alkyl, and $R^{61'}$ is $(C_{1-3} \text{ alkoxy})C_{1-4}$ alkyl, or $R^{60'}$ and $R^{61'}$ together with the nitrogen atom to which they are attached may form morpholino);

$R^{51'}$ is a hydrogen atom and C_{1-4} alkyl;

$R^{52'}$ is C_{1-8} alkyl substituted with one or more hydroxy groups;

$R^{53'}$ and $R^{54'}$ are each independently selected from a hydrogen atom and C_{1-4} alkyl;

$R^{55'}$ is a hydrogen atom or C_{1-4} alkyl;

$R^{56'}$ is $(C_{1-4} \text{ alkyl})\text{carbonyl}$;

$R^{57'}$ is a hydrogen atom or C_{1-4} alkyl;

$R^{58'}$ is a hydrogen atom or C_{1-3} alkyl;

$R^{59'}$ is C_{1-8} alkyl substituted with one or more hydroxy groups;

Ar^2 is phenyl or 5- or 6-membered heteroaryl, wherein the aryl and the heteroaryl groups are each optionally substituted with one to three substituents selected from Rb' , Rc' ,

and Rd';

Rb', Rc', and Rd' are each independently selected from optionally C₁₋₄ alkoxy-substituted C₁₋₅ alkoxy, a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14'}; and

each R^{14'} is independently selected from a halogen atom, cyano, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, and C₁₋₄ alkylthio.

[0162] (6-2) The compound according to (6-1) or a salt thereof, or a solvate of the compound or the salt, wherein

R^{4'} is C₁₋₄ alkyl;

q is an integer selected from 0 to 3;

R^{3'} is C₁₋₄ alkyl substituted with Re';

Re' is C₆₋₁₀ aryl, wherein the aryl group is optionally substituted with one to three substituents each independently selected from a halogen atom and C₁₋₄ alkoxy optionally substituted with one or more substituents R^{12'};

each R^{12'} is independently selected from 5- or 6-membered heterocycloalkyl and a group -NR^{39'}R^{40'};

R^{39'} and R^{40'} are each independently selected from a hydrogen atom and optionally C₁₋₄ alkoxy-substituted C₁₋₄ alkyl;

Ar² is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb', Rc', and Rd';

Rb', Rc', and Rd' are each independently selected from a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14'}; and

each R^{14'} is independently selected from a halogen atom, cyano, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, and C₁₋₄ alkylthio.

[0163] (6-3) The compound according to (6-1) or (6-2) or a salt thereof, or a solvate of the

compound or the salt, wherein

Rb' is a halogen atom;

Rc' is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen atoms; and

Rd' is a halogen atom or 5- or 6-membered heteroaryl optionally substituted with one or more substituents R^{14'}.

[0164] (6-4) The compound according to any of (6-1) to (6-3) or a salt thereof, or a solvate of the compound or the salt, wherein R^{3'} is benzyl, wherein phenyl in the benzyl group is optionally substituted with one to three substituents each independently selected from a halogen atom and C₁₋₄ alkoxy optionally substituted with one or more substituents R^{12'}.

[0165] (6-5) The compound according to any of (6-1) to (6-4) or a salt thereof, or a solvate of the compound or the salt, wherein Re' is phenyl, wherein the phenyl group is optionally substituted with one substituent selected from Ri', Rj', and Rk', two substituents selected from combinations of Ri' and Rj', Ri' and Rk', and Rj' and Rk', or three substituents Ri', Rj', and Rk';

Ri' and Rj' are each independently a halogen atom; and

Rk' is C₁₋₄ alkoxy optionally substituted with one or more substituents R^{12'}.

[0166] (6-6) The compound according to any of (6-1) to (6-5) or a salt thereof, or a solvate of the compound or the salt, wherein

Ar² is phenyl or pyridinyl;

Rd' is pyridinyl or pyrimidinyl; and

R^{12'} is morpholinyl or -NR^{39'}R^{40'}.

[0167] (6-7) The compound according to any of (6-1) to (6-6) or a salt thereof, or a solvate of the compound or the salt, wherein R^{12'} is selected from morpholinyl, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃alkyl)amino], and [N,N-di(C₁₋₃ alkyl)amino].

[0168] (6-8) The compound according to any of (6-1) to (6-3) or a salt thereof, or a solvate of the compound or the salt, wherein Ra is selected from a halogen atom, (morpholino)C₁₋₄alkoxy, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃alkyl)amino]C₁₋₄ alkoxy, and

[N,N-di(C₁₋₃ alkyl)amino]C₁₋₄alkoxy.

[0169] (6-9) The compound according to (6-5) or (6-6) or a salt thereof, or a solvate of the compound or the salt, wherein R_k is selected from (morpholino)C₁₋₄alkoxy, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃alkyl)amino]C₁₋₄ alkoxy, and [N,N-di(C₁₋₃ alkyl)amino]C₁₋₄alkoxy.

[0170] (6-10) The compound according to any of (6-1) to (6-6) or a salt thereof, or a solvate of the compound or the salt, wherein R^{12'} is a group -NR^{39'}R^{40'}.

[0171] An alternative aspect of the present invention provides a pharmaceutical drug comprising a compound according to any of (4-1) to (4-12), (5-1) to (5-12), and (6-1) to (6-10) or a salt thereof, or a solvate of the compound or the salt.

[0172] A further alternative aspect of the present invention provides a pharmaceutical composition comprising a compound according to any of (4-1) to (4-12), (5-1) to (5-12), and (6-1) to (6-10) or a salt thereof, or a solvate of the compound or the salt.

[0173] A further alternative aspect of the present invention provides the compound according to any of (4-1) to (4-12), (5-1) to (5-12), and (6-1) to (6-10) or the salt thereof, or the solvate of the compound or the salt for use in the prevention and/or treatment of a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure.

[0174] A further alternative aspect of the present invention provides a NaPi-IIb inhibitor comprising a compound according to any of (4-1) to (4-12), (5-1) to (5-12), and (6-1) to (6-10) or a salt thereof, or a solvate of the compound or the salt.

[0175] A further alternative aspect of the present invention provides a preventive and/or therapeutic agent for a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, the agent comprising a compound according to any of (4-1) to (4-12), (5-1) to (5-12), and (6-1) to (6-10) or a salt thereof, or a solvate of the compound or the salt as an active ingredient.

[0176] A further alternative aspect of the present invention provides a method for preventing and/or treating a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, comprising administering a therapeutically effective amount of a compound according to any of (4-1) to (4-12), (5-1) to (5-12), and (6-1)

to (6-10) or a salt thereof, or a solvate of the compound or the salt to a patient.

ADVANTAGIOUS EFFECTS OF INVENTION

[0177] The compound, the salt thereof or the solvate of the compound or the salt has an excellent NaPi-IIb inhibitory effect, PiT-1 inhibitory effect, or PiT-2 inhibitory effect, and is useful as a prevention and/or treatment agent for hyperphosphatemia. Further, the compound, the salt thereof or the solvate of the compound or the salt is useful as a prevention and/or treatment agent for secondary hyperparathyroidism, or chronic renal failure.

DESCRIPTION OF EMBODIMENTS

[0178] Hereinafter, a compound of the present invention or a salt thereof, or a solvate of the compound or the salt, a method for producing the same, and a pharmaceutical drug containing this compound will be described.

[0179] Definition

In the present invention, a "halogen atom" means a fluorine atom, a chlorine atom, a bromine atom, an iodine atom, and the like. In the present invention, preferred examples of the halogen atom used as a substituent for aryl, heteroaryl, etc. include a fluorine atom, a chlorine atom, and a bromine atom. In the present invention, preferred examples of the halogen atom used as a substituent for alkyl or a group partially containing alkyl (alkoxy, alkenyl, alkylthio, etc.) include a fluorine atom. Specific examples of groups having the halogen atom as a substituent include trifluoromethyl, pentafluoroethyl, trifluoromethoxy, pentafluoroethoxy, trifluoromethylthio, and pentafluoroethylthio.

[0180] In the present invention, "C₁₋₃ alkyl" refers to a monovalent group derived from a linear and branched saturated aliphatic hydrocarbon having 1 to 3 carbon atoms by the loss of one arbitrary hydrogen atom. Specific examples thereof include methyl, ethyl, n-propyl, and isopropyl.

[0181] In the present invention, "C₁₋₄ alkyl" refers to a monovalent group derived from a linear and branched saturated aliphatic hydrocarbon having 1 to 4 carbon atoms by the loss of one arbitrary hydrogen atom. Specific examples thereof include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, and 1-methylpropyl.

[0182] In the present invention, "C₁₋₅ alkyl" refers to a monovalent group derived from a linear and branched saturated aliphatic hydrocarbon having 1 to 5 carbon atoms by the loss of one arbitrary hydrogen atom. Specific examples thereof include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, 1-methylpropyl, n-pentyl, isopentyl, 2-methylbutyl, 1,1-dimethylpropyl, and 1-ethylpropyl.

[0183] In the present invention, "C₁₋₆ alkyl" refers to a monovalent group derived from a linear and branched saturated aliphatic hydrocarbon having 1 to 6 carbon atoms by the loss of one arbitrary hydrogen atom. Specific examples thereof include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, 1-methylpropyl, n-pentyl, isopentyl, 2-methylbutyl, 1,1-dimethylpropyl, 1-ethylpropyl, hexyl, 4-methylpentyl, and 2-ethylbutyl.

[0184] In the present invention, "C₁₋₈ alkyl" refers to a monovalent group derived from a linear and branched saturated aliphatic hydrocarbon having 1 to 8 carbon atoms by the loss of one arbitrary hydrogen atom. Specific examples thereof include the groups listed above in the definition of "C₁₋₆ alkyl" as well as n-heptyl, 5-methylhexyl, 1-propylbutyl, 2-ethyl-2-methylbutyl, n-octyl, 5-methylheptyl, 2,3-dimethylhexyl, 1-methyl-1-propylbutyl, and 2,2-diethylbutyl.

[0185] In the present invention, "C₁₋₁₀ alkyl" refers to a monovalent group derived from a linear and branched saturated aliphatic hydrocarbon having 1 to 10 carbon atoms by the loss of one arbitrary hydrogen atom. Specific examples thereof include the groups listed above in the definition of "C₁₋₈ alkyl" as well as 7-methyloctyl, 5-ethylheptyl, n-decyl, 8-methylnonyl, 5,5-dimethyloctyl, and 4-ethyl-6-methylheptyl.

[0186] In the present invention, "linear C₁₋₆ alkyl" refers to a monovalent group derived from a linear saturated aliphatic hydrocarbon having 1 to 6 carbon atoms by the loss of one arbitrary hydrogen atom. "Linear C₁₋₆ alkyl" specifically means methyl, ethyl, n-propyl, n-butyl, n-pentyl, and n-hexyl.

[0187] In the present invention, "linear C₁₋₁₀ alkyl" refers to a monovalent group derived from a linear saturated aliphatic hydrocarbon having 1 to 10 carbon atoms by the loss of one arbitrary hydrogen atom. "Linear C₁₋₁₀ alkyl" specifically means methyl, ethyl, n-propyl, n-

butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, and n-decyl.

[0188] In the present invention, "morpholino" means morpholin-4-yl.

[0189] In the present invention, "(morpholino) C_{1-6} alkyl" means a C_{1-6} alkyl group substituted with morpholino. " C_{1-6} alkyl" is as defined above. Specific examples thereof include morpholinomethyl, morpholinoethyl, morpholino-n-propyl, morpholinoisopropyl, morpholino-n-butyl, morpholinoisobutyl, morpholino-sec-butyl, morpholino-t-butyl, morpholino-1-methylpropyl, morpholino-n-pentyl, morpholinoisopentyl, morpholino-2-methylbutyl, morpholino-1,1-dimethylpropyl, morpholino-1-ethylpropyl, morpholinohexyl, morpholino-4-methylpentyl, and morpholino-2-ethylbutyl. Preferred examples thereof include morpholinomethyl.

[0190] In the present invention, "(morpholino) C_{1-4} alkyl" means a C_{1-4} alkyl group substituted with morpholino. " C_{1-4} alkyl" is as defined above. Specific examples thereof include morpholinomethyl, morpholinoethyl, morpholino-n-propyl, morpholinoisopropyl, morpholino-n-butyl, morpholinoisobutyl, morpholino-sec-butyl, morpholino-t-butyl, and morpholino-1-methylpropyl. Preferred examples thereof include morpholinomethyl.

[0191] In the present invention, "(oxetanyl) C_{1-4} alkyl" means a C_{1-4} alkyl group substituted with oxetanyl. " C_{1-4} alkyl" is as defined above. Specific examples thereof include oxetanylmethyl, oxetanylethyl, oxetanyl-n-propyl, oxetanylisopropyl, oxetanyl-n-butyl, oxetanylisobutyl, oxetanyl-sec-butyl, oxetanyl-t-butyl, and oxetanyl-1-methylpropyl. Preferred examples thereof include oxetanylmethyl.

[0192] In the present invention, "(4- to 6-membered heterocycloalkyl) C_{1-4} alkyl" means C_{1-4} alkyl substituted with 4- to 6-membered heterocycloalkyl. " C_{1-4} alkyl" is as defined above. "4- to 6-membered heterocycloalkyl" means a saturated heterocyclic group composed of 4 to 6 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. "(4- to 6-membered heterocycloalkyl) C_{1-4} alkyl" includes "(morpholino) C_{1-4} alkyl" and "(oxetanyl) C_{1-4} alkyl".

[0193] In the present invention, "(carboxy) C_{1-8} alkyl" means C_{1-8} alkyl substituted with carboxy. " C_{1-8} alkyl" is as defined above. Specific examples thereof include carboxy-n-

hexyl, carboxy-n-heptyl, carboxy-5-methylhexyl, carboxy-1-propylbutyl, carboxy-2-ethyl-2-methylbutyl, carboxy-n-octyl, carboxy-5-methylheptyl, carboxy-2,3-dimethylhexyl, carboxy-1-methyl-1-propylbutyl, and carboxy-2,2-diethylbutyl. Preferred examples thereof include carboxy-n-hexyl, carboxy-n-heptyl, and carboxy-n-octyl.

In the present invention, "(C₁₋₆ alkoxy)C₁₋₈ alkyl" means C₁₋₈ alkyl substituted with C₁₋₆ alkoxy. "C₁₋₆ alkoxy" and "C₁₋₈ alkyl" are as defined above and below. Specific examples thereof include ethoxy-n-propyl, n-propoxy-n-propyl, n-butoxy-n-propyl, n-heptoxypropyl, n-propoxy-n-hexyl, n-butoxy-n-hexyl, n-propoxy-n-heptyl, and n-butoxy-n-heptyl.

Preferred examples thereof include n-propoxy-n-propyl and n-butoxy-n-heptyl.

[0194] In the present invention, "(C₁₋₃ alkoxy)C₁₋₄ alkyl" means C₁₋₄ alkyl substituted with C₁₋₃ alkoxy. "C₁₋₃ alkoxy" and "C₁₋₄ alkyl" are as defined above and below. Specific examples thereof include methoxymethyl, methoxyethyl, ethoxymethyl, methoxy-n-propyl, ethoxy-n-propyl, methoxy-n-butyl, and ethoxy-n-butyl.

[0195] In the present invention, "(C₁₋₄ alkoxy)C₁₋₄ alkyl" means C₁₋₄ alkyl substituted with C₁₋₄ alkoxy. "C₁₋₄ alkoxy" and "C₁₋₄ alkyl" are as defined above and below. Specific examples thereof include the groups listed above in the definition of "(C₁₋₃ alkoxy)C₁₋₄ alkyl" as well as n-butoxymethyl, n-butoxyethyl, n-butoxy-n-propyl, and n-butoxy-n-butyl.

[0196] In the present invention, "(C₁₋₃ alkoxy)C₁₋₆ alkyl" means C₁₋₆ alkyl substituted with C₁₋₃ alkoxy. "C₁₋₃ alkoxy" and "C₁₋₆ alkyl" are as defined above and below. Specific examples thereof include the groups listed above in the definitions of "(C₁₋₃ alkoxy)C₁₋₄ alkyl" and "(C₁₋₄ alkoxy)C₁₋₄ alkyl". Preferred examples thereof include methoxy-n-propyl.

[0197] In the present invention, "[HO-((CH₂)_oO)_p]C₁₋₆ alkyl" refers to C₁₋₆ alkyl substituted with a group -(O(CH₂)_o)_p-OH. C₁₋₆ alkyl is as defined above. Specific examples thereof include [HO-((CH₂)₂O)₃]propyl.

[0198] In the present invention, "C₁₋₅ alkylene" refers to a divalent group derived from a linear or branched saturated aliphatic hydrocarbon having 1 to 5 carbon atoms by the loss of two arbitrary hydrogen atoms. "C₁₋₅ alkylene" includes "C₁₋₃ alkylene". Specific examples thereof include -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH₂CH₂-, -CH₂CH(CH₃)-, -CH₂C(CH₃)₂-,

-CH(CH₂CH₃)-, -C(CH₂CH₃)₂-, and -CH(CH₂CH₂CH₃)-. -CH₂- is preferred.

[0199] In the present invention, two or more substituents R² on a 5- to 8-membered saturated heterocyclic ring may together form C₁₋₅ alkylene that links the ring atoms to which they are attached. In this case, the saturated heterocyclic ring forms a bicyclo ring or tricycle ring and so on.

[0200] In the present invention, "C₂₋₁₀ alkynyl" refers to a monovalent group derived from a linear or branched aliphatic hydrocarbon having 2 to 10 carbon atoms and having at least one triple bond (two adjacent SP carbon atoms) by the loss of one arbitrary hydrogen atom. Specific examples thereof include ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, 1-methyl-2-propynyl, 1-pentynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 1-methyl-2-butynyl, 1-methyl-3-butynyl, 2-methyl-3-butynyl, 3-methyl-1-butynyl, 1,1-dimethyl-2-propynyl, 1-hexynyl, heptynyl, heptadiynyl, octynyl, and octadiynyl.

[0201] In the present invention, "optionally substituted C₂₋₁₀ alkynyl" means unsubstituted C₂₋₁₀ alkynyl described above or C₂₋₁₀ alkynyl on which one or more hydrogen atoms are replaced with predetermined substituent(s). Two or more substituents on this group may be the same as or different from each other. One carbon atom may be substituted with a plurality of substituents.

[0202] In the present invention, "C₂₋₆ alkynyl" refers to a monovalent group derived from a linear or branched aliphatic hydrocarbon having 2 to 6 carbon atoms and having at least one triple bond (two adjacent SP carbon atoms) by the loss of one arbitrary hydrogen atom. Specific examples thereof include ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, 1-methyl-2-propynyl, 1-pentynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 1-methyl-2-butynyl, 1-methyl-3-butynyl, 2-methyl-3-butynyl, 3-methyl-1-butynyl, 1,1-dimethyl-2-propynyl, and 1-hexynyl.

[0203] In the present invention, "optionally substituted C₂₋₆ alkynyl" means unsubstituted C₂₋₆ alkynyl described above or C₂₋₆ alkynyl on which one or more hydrogen atoms are replaced with predetermined substituent(s). Two or more substituents on this group may be the same as or different from each other. One carbon atom may be substituted with a

plurality of substituents.

[0204] In the present invention, "(carboxy) C_{2-6} alkynyl" means C_{2-6} alkynyl substituted with carboxy. C_{2-6} alkynyl is as defined above. Specific examples thereof include carboxyethynyl, carboxy-1-propynyl, carboxy-2-propynyl, carboxy-1-butynyl, carboxy-2-butynyl, carboxy-3-butynyl, carboxy-1-methyl-2-propynyl, carboxy-1-pentynyl, carboxy-2-pentynyl, carboxy-3-pentynyl, carboxy-4-pentynyl, carboxy-1-methyl-2-butynyl, carboxy-1-methyl-3-butynyl, carboxy-2-methyl-3-butynyl, carboxy-3-methyl-1-butynyl, carboxy-1,1-dimethyl-2-propynyl, and carboxy-1-hexynyl. Preferred examples thereof include carboxy-1-pentynyl and carboxy-1-hexynyl.

[0205] In the present invention, "(C_{1-6} alkoxy) C_{2-8} alkynyl" means C_{2-8} alkynyl substituted with C_{1-6} alkoxy. C_{1-6} alkoxy and C_{2-8} alkynyl are as defined above. Specific examples thereof include ethoxy-1-propynyl, propoxy-1-propynyl, butoxy-1-propynyl, propoxy-1-butynyl, propoxy-1-hexynyl, butoxy-1-hexynyl, propoxy-1-heptynyl, and butoxy-1-heptynyl. Preferred examples thereof include ethoxy-1-propynyl, propoxy-1-propynyl, butoxy-1-propynyl, propoxy-1-butynyl, propoxy-1-hexynyl, and butoxy-1-heptynyl.

[0206] In the present invention, "(morpholino) C_{2-6} alkynyl" means C_{2-6} alkynyl substituted with morpholino. C_{2-6} alkynyl is as defined above. Specific examples thereof include morpholinoethynyl, morpholino-1-propynyl, morpholino-2-propynyl, morpholino-1-butynyl, morpholino-2-butynyl, morpholino-3-butynyl, morpholino-1-pentynyl, morpholino-2-pentynyl, morpholino-3-pentynyl, morpholino-4-pentynyl, and morpholino-1-hexynyl. Preferred examples thereof include morpholino-1-propynyl, morpholino-1-butynyl, and morpholino-1-heptynyl.

[0207] In the present invention, examples of "(morpholino) C_{2-6} alkynyl substituted with oxo group(s)" include (morpholino) C_{2-6} alkynyl substituted with one or two oxo groups. Preferred examples thereof include 3-(3-oxomorpholin-4-yl)-1-propynyl and (1,1-dioxothiomorpholin-4-yl)-1-propynyl.

[0208] In the present invention, "(3- to 6-membered oxacycloalkyl) C_{2-6} alkynyl" means C_{2-6} alkynyl substituted with 3- to 6-membered oxacycloalkyl. 3- to 6-Membered oxacycloalkyl

and C₂₋₆ alkynyl are as defined in the specification.

[0209] In the present invention, "(oxetanyl)C₂₋₆ alkynyl" means C₂₋₆ alkynyl substituted with oxetanyl. C₂₋₆ alkynyl is as defined above. Specific examples thereof include oxetanylethynyl, oxetanyl-1-propynyl, oxetanyl-2-propynyl, oxetanyl-1-butynyl, oxetanyl-2-butynyl, oxetanyl-3-butynyl, oxetanyl-1-pentynyl, oxetanyl-2-pentynyl, oxetanyl-3-pentynyl, oxetanyl-4-pentynyl, and oxetanyl-1-hexynyl. Preferred examples thereof include oxetanyl-1-propynyl, oxetanyl-1-butynyl, and oxetanyl-1-pentynyl.

[0210] In the present invention, "(pyrrolidino)C₂₋₆ alkynyl" means C₂₋₆ alkynyl substituted with pyrrolidino. C₂₋₆ alkynyl is as defined above. Specific examples thereof include pyrrolidinoethynyl, pyrrolidino-1-propynyl, pyrrolidino-2-propynyl, pyrrolidino-1-butynyl, pyrrolidino-2-butynyl, pyrrolidino-3-butynyl, pyrrolidino-1-pentynyl, pyrrolidino-2-pentynyl, pyrrolidino-3-pentynyl, pyrrolidino-4-pentynyl, and pyrrolidino-1-hexynyl. Preferred examples thereof include pyrrolidino-1-propynyl.

[0211] In the present invention, "(C₃₋₆ cycloalkyl)C₂₋₆ alkynyl" means C₂₋₆ alkynyl substituted with C₃₋₆ cycloalkyl. C₃₋₆ cycloalkyl and C₂₋₆ alkynyl are as defined above. Specific examples thereof include cyclopropylethynyl, cyclobutylethynyl, cyclopentylethynyl, cyclohexylethynyl, cyclopropyl-1-propynyl, cyclobutyl-1-propynyl, cyclopentyl-1-propynyl, cyclohexyl-1-propynyl, cyclopropyl-1-butynyl, cyclobutyl-1-butynyl, cyclopentyl-1-butynyl, and cyclohexyl-1-butynyl. Preferred examples thereof include cyclopropylethynyl, cyclobutylethynyl, cyclopentylethynyl, cyclopropyl-1-propynyl, cyclobutyl-1-propynyl, and cyclopentyl-1-propynyl.

[0212] In the present invention, "[HO-((CH₂)_oO)_p]C₂₋₈ alkynyl" means C₂₋₈ alkynyl substituted with a group -(O(CH₂)_o)_p-OH. C₂₋₈ alkynyl is as defined above. Specific examples thereof include [HO-((CH₂)₂O)₃]propynyl and [HO-((CH₂)₂O)₂]propynyl.

[0213] In the present invention, "[N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₂₋₆ alkynyl" means C₂₋₆ alkynyl substituted with N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino. In this context, "[N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]" and "C₂₋₆ alkynyl" are as defined in this specification. Specific examples include [N-(2-methoxyethyl)-N-

(methyl)amino]ethynyl, [N-(2-methoxy-1,1-dimethylethyl)-N-(methyl)amino]ethynyl, and [N-(2-methoxy-2-methyl-1-propyl)-N-(methyl)amino]ethynyl.

[0214] In the present invention, "C₁₋₃ alkoxy" means a C₁₋₃ alkyl-O- group. In this context, C₁₋₃ alkyl is as defined above. Specific examples thereof include methoxy, ethoxy, 1-propoxy, and 2-propoxy.

[0215] In the present invention, "C₁₋₄ alkoxy" means a C₁₋₄ alkyl-O- group. In this context, C₁₋₄ alkyl is as defined above. Specific examples thereof include methoxy, ethoxy, 1-propoxy, 2-propoxy, n-butoxy, i-butoxy, sec-butoxy, and t-butoxy.

[0216] In the present invention, "C₁₋₅ alkoxy" means a C₁₋₅ alkyl-O- group. In this context, C₁₋₅ alkyl is as defined above. Specific examples thereof include methoxy, ethoxy, 1-propoxy, 2-propoxy, n-butoxy, i-butoxy, sec-butoxy, t-butoxy, and 1-pentyloxy.

[0217] In the present invention, "C₁₋₆ alkoxy" means a C₁₋₆ alkyl-O- group. In this context, C₁₋₆ alkyl is as defined above. Specific examples thereof include methoxy, ethoxy, 1-propoxy, 2-propoxy, n-butoxy, i-butoxy, sec-butoxy, t-butoxy, 1-pentyloxy, and 1-hexyloxy.

[0218] In the present invention, "C₁₋₈ alkoxy" means a C₁₋₈ alkyl-O- group. Specific examples thereof include the groups listed above in the definition of "C₁₋₆ alkoxy" as well as 1-heptyloxy and 1-octyloxy.

[0219] In the present invention, "(carboxy)C₁₋₈ alkoxy" means C₁₋₈ alkoxy substituted with one carboxy group. "C₁₋₈ alkoxy" is as defined above. Specific examples thereof include carboxy-n-propoxy, carboxy-n-butoxy, carboxy-n-pentoxy, and carboxy-n-hexoxy.

[0220] In the present invention, "(morpholino)C₁₋₆ alkoxy" means C₁₋₆ alkoxy substituted with morpholino. "C₁₋₆ alkoxy" is as defined above. Specific examples thereof include morpholinomethoxy, morpholinoethoxy, morpholino-1-propoxy, morpholino-2-propoxy, morpholino-n-butoxy, morpholino-i-butoxy, morpholino-sec-butoxy, morpholino-t-butoxy, morpholino-1-pentyloxy, and morpholino-1-hexyloxy. Preferred examples thereof include morpholinoethoxy.

[0221] In the present invention, "(3- to 6-membered oxacycloalkyl)C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with 3- to 6-membered oxacycloalkyl. "C₁₋₄ alkoxy" is as defined above.

3- to 6-membered oxacycloalkyl means a saturated heterocyclic group composed of 3 to 6 ring-constituting atoms containing one or two, (preferably one) oxygen atom. "(3- to 6-membered oxacycloalkyl)_{C₁₋₄} alkoxy" includes (oxiranyl)_{C₁₋₄} alkoxy, (oxetanyl)_{C₁₋₄} alkoxy, and (oxolanyl)_{C₁₋₄} alkoxy. Specific examples thereof include oxiranylmethoxy, oxiranylethoxy, oxiranyl-1-propoxy, oxiranyl-n-butoxy, oxetanylmethoxy, oxetanylethoxy, oxetanyl-1-propoxy, oxetanyl-n-butoxy, oxolanylmethoxy, oxolanylethoxy, oxolanyl-1-propoxy, and oxolanyl-n-butoxy. Preferred examples thereof include oxolanylmethoxy, oxiranylmethoxy, and oxetanylmethoxy.

[0222] In the present invention, "3- to 6-membered oxacycloalkyloxy" means 3- to 6-membered oxacycloalkyl-O-. "3- to 6-membered oxacycloalkyl" is as defined above. Specific examples thereof include oxiranyloxy, oxetanyloxy, oxolanyloxy, and oxanyloxy. Oxetanyloxy, oxolanyloxy, or oxanyloxy is preferred.

[0223] In the present invention, "4- to 6-membered nitrogen containing heterocycloalkyloxy" means 4- to 6-membered nitrogen containing heterocycloalkyl-O-. "4- to 6-membered nitrogen containing heterocycloalkyl" means a saturated heterocyclic group composed of 4 to 6 ring-constituting atoms containing one or two nitrogen atoms. Specific examples thereof include pyrrolidinyloxy and azetidinyloxy.

[0224] In the present invention, "(pyridinyl)_{C₁₋₄} alkoxy" means _{C₁₋₄} alkoxy substituted with pyridinyl. "_{C₁₋₄} alkoxy" is as defined above. Specific examples thereof include pyridinylmethoxy, pyridinylethoxy, pyridinyl-1-propoxy, and pyridinyl-n-butoxy. Preferred examples thereof include pyridinylmethoxy.

[0225] In the present invention, "(pyrimidinyl)_{C₁₋₄} alkoxy" means _{C₁₋₄} alkoxy substituted with pyrimidinyl. "_{C₁₋₄} alkoxy" is as defined above. Specific examples thereof include pyrimidinylmethoxy, pyrimidinylethoxy, pyrimidinyl-1-propoxy, and pyrimidinyl-n-butoxy. Preferred examples thereof include pyrimidinylmethoxy.

[0226] In the present invention, "(1,2,4-triazolyl)_{C₁₋₄} alkoxy" means _{C₁₋₄} alkoxy substituted with 1,2,4-triazolyl. "_{C₁₋₄} alkoxy" is as defined above. Specific examples thereof include 1,2,4-triazolylmethoxy, 1,2,4-triazolylethoxy, 1,2,4-triazolyl-1-propoxy, and 1,2,4-triazolyl-

n-butoxy. Preferred examples thereof include 1,2,4-triazolyloethoxy.

[0227] In the present invention, "(3- to 6-membered heterocycloalkyl)C₁₋₆ alkoxy" means C₁₋₆ alkoxy substituted with 3- to 6-membered heterocycloalkyl. 3- to 6-membered heterocycloalkyl means a saturated heterocyclic group composed of 3 to 6 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. C₁₋₆ alkoxy is as defined above. "(3- to 6-membered heterocycloalkyl)C₁₋₆ alkoxy" includes (morpholino)C₁₋₆alkoxy, (3- to 6-membered oxacycloalkyl)C₁₋₄ alkoxy, (pyrrolidinyl)C₁₋₄ alkoxy, (azetidiny)C₁₋₄ alkoxy, and (piperidinyl)C₁₋₄ alkoxy. Specific examples thereof include morpholinomethoxy, morpholinoethoxy, morpholino-1,1-dimethylethoxy(morpholino-t-butoxy), morpholino-1-methyl-ethoxy(morpholino-2-propoxy), tetrahydrofuranylmethoxy, pyrrolidinylmethoxy, pyrrolidinylethoxy, azetidinylmethoxy, azetidinyloethoxy, piperidinylmethoxy, piperidinylethoxy, piperazinylmethoxy, piperazinylethoxy.

[0228] In the present invention, "(5- or 6-membered heteroaryl)C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with 5- or 6-membered heteroaryl. 5- or 6-membered heteroaryl and C₁₋₄ alkoxy are as defined herein. Specific examples thereof include 1,2,4-triazolyloethoxy.

[0229] In the present invention, "(C₁₋₆ alkoxy)C₁₋₆ alkoxy" means C₁₋₆ alkoxy substituted with C₁₋₆ alkoxy. "C₁₋₆ alkoxy" is as defined above. "(C₁₋₆ alkoxy)C₁₋₆ alkoxy" includes "(C₁₋₄ alkoxy)C₁₋₆ alkoxy", "C₁₋₃ alkoxy(C₁₋₄ alkoxy)", and the like. Specific examples thereof include methoxymethoxy, methoxyethoxy, ethoxymethoxy, ethoxyethoxy, methoxypropoxy, ethoxypropoxy, propoxymethoxy, propoxyethoxy, propoxypropoxy, methoxybutoxy, ethoxybutoxy, propoxybutoxy, butoxymethoxy, butoxyethoxy, butoxypropoxy, butoxybutoxy, methoxypentoxy, ethoxypentoxy, propoxypentoxy, butoxypentoxy, pentoxymethoxy, pentoxyethoxy, pentoxypropoxy, pentoxybutoxy, pentoxypentoxy, methoxyhexoxy, ethoxyhexoxy, propoxyhexoxy, butoxyhexoxy, pentoxyhexoxy, hexoxymethoxy, hexoxyethoxy, hexoxypropoxy, hexoxybutoxy, hexoxypentoxy, and hexoxyhexoxy. Preferred examples thereof include propoxybutoxy, propoxypentoxy, and propoxyhexoxy.

[0230] In the present invention, "(C₁₋₆ alkoxy)C₁₋₈ alkoxy" means C₁₋₈ alkoxy substituted with C₁₋₆ alkoxy. "C₁₋₆ alkoxy" and "C₁₋₈ alkoxy" are as defined above. Specific examples thereof include the groups listed above in the definition of "(C₁₋₆ alkoxy)C₁₋₆ alkoxy" as well as hexoxyheptyloxy and hexoxyoctyloxy.

[0231] In the present invention, "(C₁₋₃ alkoxy(C₁₋₄ alkoxy))C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with C₁₋₃ alkoxy(C₁₋₄ alkoxy). "C₁₋₃ alkoxy(C₁₋₄ alkoxy)" means C₁₋₄ alkoxy substituted with C₁₋₃ alkoxy. "C₁₋₄ alkoxy" and "C₁₋₃ alkoxy" are as defined above. Specific examples thereof include methoxyethoxyethoxy, ethoxyethoxyethoxy, and methoxyethoxymethoxy.

[0232] In the present invention, "[N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino. In this context, "[N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]" means amino substituted with (C₁₋₃ alkoxy)C₁₋₄ alkyl and C₁₋₃ alkyl. (C₁₋₃ alkoxy)C₁₋₄ alkyl and C₁₋₃ alkyl are as defined above. Specific examples of "[N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy" include [Nmethoxyethyl)-N-(methyl)amino]ethoxy, [N-(methoxy-1,1-dimethylethyl)-N-(methyl)amino]ethoxy, and [N-(-methoxy-2-methyl-1-propyl)-N-(methyl)amino]ethoxy.

[0233] In the present invention, "[N-(3- to 6-membered heterocycloalkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with N-(3- to 6-membered heterocycloalkyl)-N-(C₁₋₃ alkyl)amino. In this context, "[N-(3- to 6-membered heterocycloalkyl)-N-(C₁₋₃ alkyl)amino]" means amino substituted with 3- to 6-membered heterocycloalkyl and C₁₋₃ alkyl. 3- to 6-membered heterocycloalkyl is as defined herein. Specific examples thereof include [N-tetrahydrofuranyl-N-(methyl)amino]ethoxy, [N-tetrahydropyranyl-N-(methyl)amino]ethoxy, and [N-oxetanyl-N-(methyl)amino]ethoxy.

[0234] In the present invention, [N,N-di((hydroxy)C₁₋₄ alkyl)-amino]C₁₋₄ alkoxy means C₁₋₄ alkoxy substituted with N,N-di((hydroxy)C₁₋₄ alkyl)-amino. "[N,N-di((hydroxy)C₁₋₄ alkyl)-amino]" means amino substituted with two (hydroxy)C₁₋₄ alkyl groups. (Hydroxy)C₁₋₄ alkyl means C₁₋₄ alkyl substituted with one hydroxy group. Specific examples thereof include

[N,N-di(2-hydroxyethyl)amino]ethoxy.

[0235] In the present invention, "[N-((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with N-((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl-N-(C₁₋₃ alkyl)amino. [N-((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl-N-(C₁₋₃ alkyl)amino] means amino substituted with ((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl and C₁₋₃ alkyl. "((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl" means C₁₋₃ alkyl substituted with (C₁₋₃ alkoxy)carbonyl. Specific examples thereof include [N-((methoxy)carbonyl)methyl-N-methylamino]ethoxy.

[0236] In the present invention, "[N-(hydroxy)C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with N-(hydroxy)C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino. "N-(hydroxy)C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino" means amino substituted with (hydroxy)C₁₋₄ alkyl and C₁₋₃ alkyl. "(Hydroxy)C₁₋₄ alkyl" means C₁₋₄ alkyl substituted with one hydroxy group. Specific examples thereof include [N-(2-hydroxyethyl)-N-methylamino]ethoxy.

[0237] In the present invention, "[N,N-di((C₁₋₃ alkoxy)C₁₋₄ alkyl)amino]C₁₋₆ alkoxy" means C₁₋₆ alkoxy substituted with N,N-di((C₁₋₃ alkoxy)C₁₋₄ alkyl)amino. "N,N-di((C₁₋₃ alkoxy)C₁₋₄ alkyl)amino" means amino substituted with two (C₁₋₃ alkoxy)C₁₋₄ alkyl groups. "(C₁₋₃ alkoxy)C₁₋₄ alkyl" is as defined above. Specific examples thereof include [N,N-di((methoxy)ethoxy)amino]ethoxy.

[0238] In the present invention, "[N,N-di(C₁₋₃ alkyl)amino]C₁₋₆ alkoxy" means C₁₋₆ alkoxy substituted with N,N-di(C₁₋₃ alkyl)amino. "N,N-di (C₁₋₃ alkyl) amino" means amino substituted with two C₁₋₃ alkyl groups. "C₁₋₆ alkoxy" and "C₁₋₃ alkyl" are as defined above. Specific examples thereof include 3-[N,N-di(methyl)amino]-2, 2-dimethyl-propoxy.

[0239] In the present invention, "[N-[N-(C₁₋₄ alkyl) carbonyl-N-(C₁₋₃ alkyl) amino] C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino] C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with N-[N-(C₁₋₄ alkyl) carbonyl-N-(C₁₋₃ alkyl) amino] C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino. C₁₋₄ alkoxy is as defined before. "N-[N-(C₁₋₄ alkyl) carbonyl-N-(C₁₋₃ alkyl) amino] C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino" means amino substituted with [N-(C₁₋₄ alkyl) carbonyl-N-(C₁₋₃ alkyl) amino] C₁₋₄ alkyl and C₁₋₃ alkyl. C₁₋₃ alkyl is as defined before. "[N-(C₁₋₄ alkyl) carbonyl-N-(C₁₋₃ alkyl) amino] C₁₋₄ alkyl" means C₁₋₄ alkyl substituted with N-(C₁₋₄ alkyl) carbonyl-N-(C₁₋₃ alkyl) amino.

"N-(C₁₋₄ alkyl) carbonyl-N-(C₁₋₃ alkyl) amino" means amino substituted with (C₁₋₄ alkyl) carbonyl and C₁₋₃ alkyl. Specific examples thereof include 2-[N-[2-(N-acetyl-N-(methyl) amino)ethyl]-N-(methyl) amino] ethoxy.

[0240] In the present invention, "[N-[N-(C₁₋₄ alkyl) carbonyl-amino] C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino] C₁₋₄ alkoxy" means C₁₋₄ alkoxy substituted with N-[N-(C₁₋₄ alkyl) carbonyl-amino] C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino. C₁₋₄ alkoxy is as defined before. "N-[N-(C₁₋₄ alkyl) carbonyl-amino] C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino" means amino substituted with [N-(C₁₋₄ alkyl) carbonyl-amino] C₁₋₄ alkyl and C₁₋₃ alkyl. C₁₋₃ alkyl is as defined before. "[N-(C₁₋₄ alkyl) carbonyl-amino] C₁₋₄ alkyl" means C₁₋₄ alkyl substituted with a N-(C₁₋₄ alkyl) carbonyl-amino. "N-(C₁₋₄ alkyl) carbonyl-amino" means amino substituted with a (C₁₋₄ alkyl) carbonyl. Specific examples thereof include 2-[N-[2-(acetilamino)ethyl]-N-(methyl) amino] ethoxy.

[0241] In the present invention, "C₁₋₃ alkylthio" means a C₁₋₃ alkyl-S- group. In this context, C₁₋₃ alkyl is as defined above. Specific examples thereof include methylthio, ethylthio, n-propylthio, and i-propylthio.

[0242] In the present invention, "C₁₋₄ alkylthio" means a C₁₋₄ alkyl-S- group. In this context, C₁₋₄ alkyl is as defined above. Specific examples thereof include methylthio, ethylthio, n-propylthio, i-propylthio, n-butylthio, i-butylthio, and t-butylthio.

[0243] In the present invention, "(morpholino)C₁₋₄ alkylthio" means C₁₋₄ alkylthio substituted with morpholino. C₁₋₄ alkylthio is as defined above. Specific examples thereof include morpholinomethylthio, morpholinoethylthio, morpholino-n-propylthio, morpholino-i-propylthio, morpholino-n-butylthio, morpholino-i-butylthio, and morpholino-t-butylthio. Preferred examples thereof include morpholinoethylthio.

[0244] In the present invention, "(5- or 6-membered heterocycloalkyl)C₁₋₄ alkylthio" means C₁₋₄ alkylthio substituted with 5- or 6-membered heterocycloalkyl. 5- or 6-membered heterocycloalkyl and C₁₋₄ alkylthio are as defined herein. "(5- or 6-membered heterocycloalkyl)C₁₋₄ alkylthio" includes "(morpholino)C₁₋₄ alkylthio". Preferred examples thereof include 2-morpholinoethylthio.

[0245] In the present invention, "(C₁₋₆ alkyl)carbonyl" means a C₁₋₆ alkyl-C(O)- group. In this context, C₁₋₆ alkyl is as defined above. "(C₁₋₆ alkyl)carbonyl" includes "(C₁₋₄ alkyl)carbonyl". Specific examples thereof include methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, i-propylcarbonyl, n-butylcarbonyl, i-butylcarbonyl, sec-butylcarbonyl, t-butylcarbonyl, 1-methylpropylcarbonyl, n-pentylcarbonyl, isopentylcarbonyl, 2-methylbutylcarbonyl, 1,1-dimethylpropylcarbonyl, 1-ethylpropylcarbonyl, hexylcarbonyl, 4-methylpentylcarbonyl, and 2-ethylbutylcarbonyl.

[0246] In the present invention, "di(C₁₋₃ alkyl)amino)carbonyl" means carbonyl substituted with di(C₁₋₃ alkyl)amino. "Di(C₁₋₃ alkyl)amino" means amino substituted with two C₁₋₃ alkyl groups. Specific examples thereof include dimethylaminocarbonyl and diethylaminocarbonyl.

[0247] In the present invention, "(C₁₋₃ alkyl)sulfonyl" means a C₁₋₃ alkyl-SO₂- group. In this context, C₁₋₃ alkyl is as defined above. Specific examples thereof include methylsulfonyl, ethylsulfonyl, and n-propylsulfonyl. Preferred examples thereof include methylsulfonyl.

[0248] In the present invention, "(C₁₋₆ alkoxy)carbonyl" means a C₁₋₆ alkyl-O-C(O)- group. In this context, C₁₋₆ alkyl is as defined above. Specific examples thereof include groups listed later in the definition of "(C₁₋₄ alkoxy)carbonyl" as well as n-pentyloxycarbonyl, isopentyloxycarbonyl, 2-methylbutoxycarbonyl, 1,1-dimethylpropoxycarbonyl, 1-ethylpropoxycarbonyl, hexyloxycarbonyl, 4-methylpentyloxycarbonyl, and 2-ethylbutoxycarbonyl.

[0249] In the present invention, "(C₁₋₄ alkoxy)carbonyl" means a C₁₋₄ alkyl-O-C(O)- group. In this context, C₁₋₄ alkyl is as defined above. Specific examples thereof include methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, i-propoxycarbonyl, n-butoxycarbonyl, i-butoxycarbonyl, sec-butoxycarbonyl, t-butoxycarbonyl, and 1-methylpropoxycarbonyl.

[0250] In the present invention, "(C₁₋₃ alkoxy)carbonyl" means a C₁₋₃ alkyl-O-C(O)- group. In this context, C₁₋₃ alkyl is as defined above. Specific examples thereof include methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, and i-propoxycarbonyl.

[0251] In the present invention, " $((C_{1-4} \text{ alkoxy})\text{carbonyl})C_{1-6} \text{ alkyl}$ " means $C_{1-6} \text{ alkyl}$ substituted with $(C_{1-4} \text{ alkoxy})\text{carbonyl}$. In this context, $(C_{1-4} \text{ alkoxy})\text{carbonyl}$ and $C_{1-6} \text{ alkyl}$ are as defined above. Specific examples thereof include methoxycarbonylmethyl, methoxycarbonylethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl, n-propoxycarbonylmethyl, and i-propoxycarbonylmethyl. Preferred examples thereof include methoxycarbonylmethyl and methoxycarbonylethyl.

[0252] In the present invention, " $((C_{1-3} \text{ alkoxy})\text{carbonyl})C_{1-3} \text{ alkoxy}$ " means $C_{1-3} \text{ alkoxy}$ substituted with a $(C_{1-3} \text{ alkoxy})\text{carbonyl}$ group. In this context, $(C_{1-3} \text{ alkoxy})\text{carbonyl}$ and $C_{1-3} \text{ alkyl}$ are as defined above. Specific examples thereof include methoxycarbonylmethoxy, methoxycarbonylethoxy, ethoxycarbonylmethoxy, ethoxycarbonylethoxy, n-propoxycarbonylmethoxy, and i-propoxycarbonylmethoxy. Preferred examples thereof include methoxycarbonylmethoxy and methoxycarbonylethoxy.

[0253] In the present invention, " $C_{6-10} \text{ aryl}$ " means a monovalent aromatic hydrocarbon ring group. Examples of $C_{6-10} \text{ aryl}$ include phenyl, 1-naphthyl, and 2-naphthyl.

[0254] In the present invention, " $(C_{6-10} \text{ aryl})\text{carbonyl}$ " means a $C_{6-10} \text{ aryl-C(O)-}$ group. In this context, $C_{6-10} \text{ aryl}$ is as defined above. Specific examples thereof include phenylcarbonyl, 1-naphthylcarbonyl, and 2-naphthylcarbonyl.

[0255] In the present invention, "5- to 10-membered heteroaryl" means an aromatic ring group composed of 5 to 10 ring-constituting atoms containing one or several (e.g., 1 to 5, preferably 1 to 3) heteroatoms. The ring may be a monocyclic or bicyclic ring. Examples of "5- to 10-membered heteroaryl" include "5- or 6-membered heteroaryl". Specific examples of "5- to 10-membered heteroaryl" include thienyl, pyridazinyl, pyrazinyl, thiazolyl, oxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, isoxazolyl, pyrazolyl, quinolinyl, isoquinolyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, pyridinyl, pyrimidinyl, indolyl, imidazolyl, furyl, thioxazolyl, pyrrolyl, tetrazolyl, oxopyrimidinyl, naphthyl, benzodioxinyl, benzisoxazolyl, benzisothiazolyl, indazolyl, benzothienyl, benzofuranyl, benzopyranyl, and triazolyl.

[0256] In the present invention, "5- or 6-membered heteroaryl" means an aromatic ring

group composed of 5 or 6 ring-constituting atoms containing one or several (e.g., 1 to 4, preferably 1 to 3, more preferably 1 or 2) heteroatoms. Specific examples thereof include thienyl, pyridazinyl, pyrazinyl, thiazolyl, oxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, isoxazolyl, pyrazolyl, pyridinyl, pyrimidinyl, furyl, thioxazolyl, and triazolyl.

[0257] In the present invention, a "C₃₋₆ saturated carbocyclic ring" refers to a cycloalkane ring having 3 to 6 ring-constituting carbon atoms and includes, for example, cyclopropane, cyclobutane, cyclopentane, and cyclohexane. In the formula (I), R¹ and R⁵ together with the carbon atom to which they are attached may form a C₃₋₆ saturated carbocyclic ring. In this case, the C₃₋₆ saturated carbocyclic ring forms a spiro ring. In the formula (I), the C₃₋₆ saturated carbocyclic ring formed by R¹ and R⁵ together with the carbon atom to which they are attached is preferably cyclobutane, cyclopentane, or cyclohexane, particularly preferably cyclobutane or cyclopentane.

[0258] In the present invention, a "5- to 8-membered saturated heterocyclic ring" means a saturated heterocyclic group composed of 5 to 8 ring-constituting atoms containing one N as a heteroatom. Specific examples thereof include pyrrolidine, piperidine, azepane, and azocane and particularly include pyrrolidine and piperidine.

[0259] In the present invention, n is preferably an integer selected from 1 to 3, particularly preferably 1 or 2.

[0260] When Re (or Rx) is C₆₋₁₀ aryl optionally substituted with one or more substituents Ra (or Ry) or 5- to 10-membered heteroaryl optionally substituted with one or more substituents Ra (or Ry), the C₆₋₁₀ aryl is preferably phenyl, and the 5- to 10-membered heteroaryl is preferably indolyl. Particularly, when R⁴ and R⁵ together with the carbon atom and nitrogen atom to which they are attached do not form a 5- to 8-membered saturated heterocyclic ring, Re is preferably phenyl.

[0261] In the present invention, Ar¹ (or Ar¹⁰¹) is preferably phenyl, naphthyl, furyl, thienyl, pyridazinyl, pyrazinyl, thiazolyl, oxazolyl, isothiazolyl, thiadiazolyl, oxadiazolyl, isoxazolyl, pyrazolyl, quinolinyl, isoquinolyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, pyridinyl, pyrimidinyl, or indolyl. These groups are each optionally substituted with one to three

substituents selected from Rb (or Rz¹), Rc (or Rz²), and Rd (or Rz³). When R⁴ and R⁵ together with the carbon atom to which they are attached form a 5- to 8-membered saturated heterocyclic ring or in the case of the compound represented by the formula (II), Ar¹ (or Ar¹⁰¹) is more preferably phenyl, furyl, pyridinyl, or pyrimidinyl.

[0262] When R⁴ and R⁵ together with the carbon atom and nitrogen atom to which they are attached do not form a 5- to 8-membered saturated heterocyclic ring, Ar¹ is preferably phenyl, pyridinyl, or pyrimidinyl.

[0263] In the present invention, Rd (or Rz³) is preferably phenyl, thienyl, pyridinyl, pyrimidinyl, quinolinyl, pyrazolyl, imidazolyl, isoxazolyl, oxazolyl, thiazolyl, isothiazolyl, pyridazinyl, pyrazinyl, furyl, thiadiazolyl, thioxazolyl, oxadiazolyl, pyrrolyl, tetrazolyl, oxopyrimidinyl, naphthyl, benzodioxinyl, isoquinolinyl, benzoxazolyl, benzothiazolyl, benzisoxazolyl, benzisothiazolyl, indolyl, indazolyl, benzothienyl, benzofuranyl, benzopyranyl, piperazinyl, piperidyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, dioxopyrrolidinyl, dioxopiperidinyl, dioxotetrahydropyrimidinyl, oxoimidazolidinyl, or dioxoimidazolidinyl. These groups are each optionally substituted with one or more substituents R¹⁴. When R⁴ and R⁵ together with the carbon atom and the nitrogen atom to which they are attached form a 5- to 8-membered saturated heterocyclic ring or in the case of the compound represented by the formula (II), Rd (or Rz³) is preferably phenyl, thienyl, pyridinyl, pyrimidinyl, quinolinyl, piperidinyl, morpholinyl, thiomorpholinyl, or piperazinyl, more preferably phenyl, thienyl, pyridinyl, pyrimidinyl, quinolinyl, or piperidinyl. When R⁴ and R⁵ together with the carbon atom and the nitrogen atom to which they are attached do not form a 5- to 8-membered saturated heterocyclic ring, Rd (or Rz³) is preferably phenyl, pyridinyl, pyrimidinyl, morpholinyl, thiomorpholinyl, piperidinyl, or piperazinyl, more preferably phenyl, pyridinyl, or pyrimidinyl.

[0264] In the present invention, "C₃₋₆ cycloalkyl" means a cyclic saturated aliphatic hydrocarbon group having 3 to 6 carbon atoms. Specific examples thereof include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

[0265] In the present invention, "C₃₋₇ cycloalkyl" means a cyclic saturated aliphatic

hydrocarbon group having 3 to 7 carbon atoms. Specific examples thereof include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

[0266] In the present invention, "3- to 10-membered heterocycloalkyl" means a saturated heterocyclic group composed of 3 to 10 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. The heterocycloalkyl may be monocyclic, bicyclic, or spiro-cyclic heterocycloalkyl. Specific examples of "3- to 10-membered heterocycloalkyl" include groups listed later in the definition of "4- to 10-membered heterocycloalkyl" as well as oxiranyl.

[0267] In the present invention, "4- to 10-membered heterocycloalkyl" means a saturated heterocyclic group composed of 4 to 10 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. The heterocycloalkyl may be monocyclic, bicyclic, or spiro-cyclic heterocycloalkyl. Examples of "4- to 10-membered heterocycloalkyl" include "5- to 10-membered heterocycloalkyl", "5- to 8-membered heterocycloalkyl", "6- to 8-membered heterocycloalkyl", and "5- or 6-membered heterocycloalkyl". Specific examples of "4- to 10-membered heterocycloalkyl" include groups listed later in the definition of "5- to 10-membered heterocycloalkyl" as well as oxetanyl, azetidiny, and 3,7-dioxa-9-azabicyclo[3.3.1]nonanyl.

[0268] In the present invention, "5- to 10-membered heterocycloalkyl" means a saturated heterocyclic group composed of 5 to 10 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. The heterocycloalkyl may be monocyclic, bicyclic, or spiro-cyclic heterocycloalkyl. Examples of "5- to 10-membered heterocycloalkyl" include "6- to 8-membered heterocycloalkyl" and "5- or 6-membered heterocycloalkyl". Specific examples of "5- to 10-membered heterocycloalkyl" include piperaziny, piperidiny, morpholiny, thiomorpholiny, pyrrolidiny, tetrahydropyrany, tetrahydrofurany, 2-oxa-6-azaspiro[3.3]heptyl, 2-azaspiro[3.3]heptyl, 2,6-diazaspiro[3.3]heptyl, and 2-thia-6-azaspiro[3.3]heptyl.

[0269] In the present invention, "4- to 8-membered heterocycloalkyl" means a saturated heterocyclic group composed of 4 to 8 ring-constituting atoms containing one to three

heteroatoms selected from O, S, and N. The heterocycloalkyl may be monocyclic, bicyclic, or spiro-cyclic heterocycloalkyl. Specific examples of "4- to 8-membered heterocycloalkyl" include those described as specific examples of "5- to 8-membered heterocycloalkyl".

[0270] In the present invention, "5- to 8-membered heterocycloalkyl" means a saturated heterocyclic group composed of 5 to 8 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. The heterocycloalkyl may be monocyclic, bicyclic, or spiro-cyclic heterocycloalkyl. Examples of "5- to 8-membered heterocycloalkyl" include "6- to 8-membered heterocycloalkyl" and "5- or 6-membered heterocycloalkyl". Specific examples of "5- to 8-membered heterocycloalkyl" include those described as specific examples of "6- to 8-membered heterocycloalkyl".

[0271] In the present invention, "6- to 8-membered heterocycloalkyl" means a saturated heterocyclic group composed of 6 to 8 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. The heterocycloalkyl may be monocyclic, bicyclic, or spiro-cyclic heterocycloalkyl. Specific examples thereof include piperazinyl, piperidinyl, morpholinyl, thiomorpholinyl, tetrahydrofuranyl, 2-oxa-6-azaspiro[3.3]heptyl, 2-azaspiro[3.3]heptyl, 2,6-diazaspiro[3.3]heptyl, and 2-thia-6-azaspiro[3.3]heptyl and particularly include 2-oxo-6-azaspiro[3.3]heptyl, 2-azaspiro[3.3]heptyl, 2,6-diazaspiro[3.3]heptyl, and 2-thia-6-azaspiro[3.3]heptyl.

[0272] In the present invention, "3- to 6-membered heterocycloalkyl" means a saturated heterocyclic group composed of 3 to 6 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. Specific examples thereof include the groups listed later in the definition of "5- or 6-membered heterocycloalkyl" as well as oxetanyl.

[0273] In the present invention, "4- to 6-membered heterocycloalkyl" means a saturated heterocyclic group composed of 4 to 6 ring-constituting atoms containing one to three heteroatoms selected from O, S, and N. Specific examples thereof include the groups listed later in the definition of "5- or 6-membered heterocycloalkyl".

[0274] In the present invention, "5- or 6-membered heterocycloalkyl" means a saturated heterocyclic group composed of 5 or 6 ring-constituting atoms containing one to three

heteroatoms selected from O, S, and N. Specific examples of the heterocycloalkyl include piperazinyl, piperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, tetrahydropyranyl, and tetrahydrofuranyl.

[0275] In the present invention, "3- to 6-membered oxacycloalkyl" means a saturated heterocyclic group composed of 3 to 6 ring-constituting atoms containing one or two oxygen atoms. Specific examples thereof include oxetanyl, tetrahydrofuranyl and tetrahydropyranyl.

[0276] In the present invention, "3- to 10-membered heterocycloalkyloxy" means a 3- to 10-membered heterocycloalkyl-O- group. In this context, 3- to 10-membered heterocycloalkyl is as defined above. Specific examples thereof include oxiranyloxy, oxetanyloxy, tetrahydrofuranyloxy, and tetrahydropyranyloxy. Tetrahydropyranyloxy is preferred.

[0277] In the present invention, "5- to 10-membered heterocycloalkyloxy" means a 5- to 10-membered heterocycloalkyl-O- group. In this context, 5- to 10-membered heterocycloalkyl is as defined above. "5- to 10-membered heterocycloalkyloxy" is preferably tetrahydropyranyloxy or tetrahydrofuranyloxy, particularly preferably tetrahydropyranyloxy.

[0278] In the present invention, "4- to 6-membered heterocycloalkyloxy" means a 4- to 6-membered heterocycloalkyl-O- group. In this context, 4- to 6-membered heterocycloalkyl is as defined above.

[0279] In the present invention, "5- or 6-membered heterocycloalkyloxy" means a 5- or 6-membered heterocycloalkyl-O- group. In this context, 5- or 6-membered heterocycloalkyl is as defined above.

[0280] In the present invention, "3- to 6-membered heterocycloalkyloxy" means a 3- to 6-membered heterocycloalkyl-O- group. In this context, 3- to 6-membered heterocycloalkyl is as defined above.

[0281] In the present invention, "3- to 10-membered heterocycloalkyl", "4- to 10-membered heterocycloalkyl", "5- to 10-membered heterocycloalkyl", "5- to 8-membered heterocycloalkyl", "6- to 8-membered heterocycloalkyl", "4- to 8-membered

heterocycloalkyl", "3- to 6-membered heterocycloalkyl", "5- or 6-membered heterocycloalkyl", "5- to 10-membered heterocycloalkyloxy", "3- to 6-membered heterocycloalkyloxy", "5- or 6-membered heterocycloalkyloxy", or "4- to 6- nitrogen containing heterocycloalkyloxy" may be substituted with predetermined substituent(s). In this case, the substituent(s) are added to each of these groups via a carbon atom or a heteroatom constituting the ring of the heterocycloalkyl or the heterocycloalkyloxy.

[0282] When Re (or Rx) is 5- to 10-membered heteroaryl optionally substituted with one or more substituents Ra (or Ry), this group contains, for example, one to three nitrogen atoms as ring-constituting groups.

[0283] When Ra (or Ry) is 5- to 10-membered heterocycloalkyloxy, 3- to 6-membered heterocycloalkyloxy, or 4- to 6-membered heterocycloalkyloxy, this group contains, for example, one to three oxygen atoms or one nitrogen atom as ring-constituting groups.

Examples of the heterocycloalkyloxy group include tetrahydropyranloxy, oxetanyloxy, pyrrolidinylloxy, azetidinyloxy, and piperidinylloxy.

[0284] When R^{10} (or R^{110}) is 3- to 10-membered heterocycloalkyl, 4- to 10-membered heterocycloalkyl, 5- to 10-membered heterocycloalkyl, 3- to 6-membered heterocycloalkyl, or 5- or 6-membered heterocycloalkyl, this group contains, for example, one to three heteroatoms selected from a nitrogen atom, an oxygen atom, and a sulfur atom, preferably a nitrogen atom and an oxygen atom, as ring-constituting groups. When R^{10} (or R^{110}) is 3- to 10-membered heterocycloalkyl, 4- to 10-membered heterocycloalkyl, 5- to 10-membered heterocycloalkyl, 3- to 6-membered heterocycloalkyl, or 5- or 6-membered heterocycloalkyl, this group is preferably morpholinyl, homomorpholinyl, thiomorpholinyl, piperidinyl, piperazinyl, homopiperazinyl, pyrrolidinyl, azetidiny, tetrahydropyranyl, or tetrahydrofuranyl, more preferably morpholinyl. These groups are each optionally substituted with the substituent(s) described as substituents for the corresponding heterocycloalkyl group.

[0285] When R^{11} (or R^{111}) is 3- to 10-membered heterocycloalkyl, 4- to 10-membered heterocycloalkyl, 5- to 10-membered heterocycloalkyl, or 5- or 6-membered heterocycloalkyl,

this group contains, for example, one to three heteroatoms selected from a nitrogen atom and an oxygen atom as ring-constituting groups.

[0286] When R^{11} (or R^{111}) is C_{3-6} cycloalkyl, 3- to 10-membered heterocycloalkyl, 4- to 10-membered heterocycloalkyl, 5- to 10-membered heterocycloalkyl, or 5- or 6-membered heterocycloalkyl, this group is preferably cyclopentyl, morpholinyl, pyrrolidinyl, homomorpholinyl, thiomorpholinyl, piperidinyl, piperazinyl, homopiperazinyl, tetrahydropyranyl, or tetrahydrofuranyl, more preferably cyclopentyl, morpholinyl, or pyrrolidinyl. These groups are each optionally substituted with the substituent(s) described as substituents for the corresponding cycloalkyl or heterocycloalkyl group.

[0287] When R^{12} (or R^{112}) is 3- to 10-membered heterocycloalkyl, 4- to 10-membered heterocycloalkyl, 5- to 10-membered heterocycloalkyl, 3- to 6-membered heterocycloalkyl, or 5- or 6-membered heterocycloalkyl, this group contains, for example, one to three heteroatoms selected from a nitrogen atom, an oxygen atom, and a sulfur atom, preferably a nitrogen atom and an oxygen atom as ring-constituting groups.

[0288] When R^{12} (or R^{112}) is 5- to 10-membered heteroaryl or 5- or 6-membered heteroaryl, this group contains, for example, one to three heteroatoms selected from a nitrogen atom, an oxygen atom, and a sulfur atom, preferably a nitrogen atom and an oxygen atom as ring-constituting groups. Further preferably, this group contains one nitrogen atom as a hetero atom.

[0289] When R^{12} (or R^{112}) is 3- to 10-membered heterocycloalkyl, 4- to 10-membered heterocycloalkyl, 5- to 10-membered heterocycloalkyl, 3- to 6- heterocycloalkyl, 5- or 6-membered heterocycloalkyl, 5- to 10-membered heteroaryl, or 5- or 6-membered heteroaryl, this group is preferably morpholinyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrolidinyl, piperidinyl, piperazinyl, azetidiny, thiomorpholinyl, oxetanyl, pyridinyl or 1,2,4-triazolyl, more preferably morpholinyl, tetrahydrofuranyl, pyridinyl, or 1,2,4-triazolyl. These groups are each optionally substituted with the substituent(s) described as substituents for the corresponding heterocycloalkyl or heteroaryl group.

[0290] When R^{13} (or R^{113}) is 3- to 10-membered heterocycloalkyl, 4- to 10-membered

heterocycloalkyl, 5- to 10-membered heterocycloalkyl, or 5- or 6-membered heterocycloalkyl, this group contains, for example, one to three heteroatoms selected from a nitrogen atom, an oxygen atom, and a sulfur atom, preferably a nitrogen atom and an oxygen atom as ring-constituting groups.

[0291] When R^{13} (or R^{113}) is 4- to 10-membered heterocycloalkyl, 5- to 10-membered heterocycloalkyl, or 5- or 6-membered heterocycloalkyl, this group is preferably morpholinyl, homomorpholinyl, thiomorpholinyl, piperidinyl, piperazinyl, homopiperazinyl, pyrrolidinyl, azetidiny, or tetrahydropyranyl, more preferably morpholinyl. These groups are each optionally substituted with the substituent(s) described as substituents for the corresponding heterocycloalkyl group.

[0292] When R^{10} (or R^{110}), R^{11} (or R^{111}), R^{12} (or R^{112}), R^{13} (or R^{113}), or R^{15} (or R^{115}) is C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups, 3- to 10-membered heterocycloalkyl, 3- to 8-membered heterocycloalkyl, 5- or 6-membered heterocycloalkyl (these heterocycloalkyl groups are each optionally substituted with one or more substituents selected from substituents described as substituents for the corresponding heterocycloalkyl group, such as oxo, a halogen atom, C_{1-4} alkyl (wherein the C_{1-4} alkyl group is optionally substituted with one or more hydroxy groups)), or 5- to 10-membered heteroaryl, this cycloalkyl group is preferably cyclopentyl, cyclohexyl, or cyclobutyl (these groups are each optionally substituted with one or more hydroxy groups); this heterocycloalkyl group is preferably tetrahydrofuranyl, pyrrolidinyl, morpholinyl, homomorpholinyl, thiomorpholinyl, piperidinyl, piperazinyl, homopiperazinyl, azetidiny, or tetrahydropyranyl (these groups are each optionally substituted with one or more of the substituents described above); and this heteroaryl group is preferably pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, 1,2,4-triazolyl. The cycloalkyl group is more preferably cyclopentyl (this group is optionally substituted with one or more hydroxy groups); the heterocycloalkyl group is more preferably tetrahydrofuranyl, pyrrolidinyl, or morpholinyl (these groups are each optionally substituted with one or more of the substituents described above); and the heteroaryl group is more preferably pyridinyl.

[0293] When Ar¹ (or Ar¹⁰¹) is 5- to 10-membered heteroaryl or 5- to 6-membered heteroaryl, this group contains, for example, one to three heteroatoms selected from a nitrogen atom, an oxygen atom, and a sulfur atom as ring-constituting groups. Ar¹ (or Ar¹⁰¹) is optionally substituted with one to three substituents selected from Rb (or Rz¹; the same holds true for the description below), Rc (or Rz²; the same holds true for the description below), and Rd (or Rz³; the same holds true for the description below), i.e., optionally substituted with Rb, Rc, or Rd, by Rb and Rc, Rc and Rd, or Rb and Rd, or by Rb, Rc, and Rd.

[0294] When Rd (or Rz³) is 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R¹⁴ (or R¹¹⁴), this group contains, for example, one to three heteroatoms selected from a nitrogen atom, an oxygen atom and a sulfur atom, as ring-constituting groups.

[0295] When Rd (or Rz³) is 5- to 10-membered heteroaryl or 5- to 6-membered heteroaryl optionally substituted with one or more substituents R¹⁴ (or R¹¹⁴), this group contains, for example, one to three heteroatoms selected from a nitrogen atom and a sulfur atom as ring-constituting groups.

[0296] When R¹⁴ (or R¹¹⁴; the same holds true for the description below) is 5- to 10-membered heterocycloalkyl, this group contains, for example, one to three heteroatoms selected from a nitrogen atom and an oxygen atom as ring-constituting groups. When R¹⁴ is 5- to 10-membered heterocycloalkyl, this group is preferably 2-oxa-6-azaspiro[3.3]heptyl, 2-azaspiro[3.3]heptyl, 2,6-diazaspiro[3.3]heptyl, or 2-thia-6-azaspiro[3.3]heptyl, particularly preferably 2-oxa-6-azaspiro[3.3]heptyl.

[0297] When Re (or Rx) is C₆₋₁₀ aryl (for example phenyl) optionally substituted with one or more substituents Ra (or Ry) or 5- to 10-membered heteroaryl optionally substituted with one or more substituents Ra (or Ry), this group is substituted with, for example, 1 to 4 substituents Ra (or Ry), preferably 1 to 3 substituents Ra (or Ry).

[0298] When R¹⁰ (or R¹¹⁰; the same holds true for the description below), R¹¹ (or R¹¹¹; the same holds true for the description below), R¹² (or R¹¹²; the same holds true for the

description below), R^{13} (or R^{13} ; the same holds true for the description below), or R^{15} (or R^{15} ; the same holds true for the description below) is C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, this group is substituted with, for example, 1 to 5 hydroxy groups, preferably 1 to 4 hydroxy groups.

[0299] When R^{10} , R^{11} , R^{12} , R^{13} , or R^{15} is 5- to 10-membered heterocycloalkyl optionally substituted with one or more oxo groups, this group is substituted with, for example, 1 to 3 oxo groups, preferably 1 or 2 oxo groups.

[0300] When R^{10} , R^{11} , R^{12} , R^{13} , or R^{15} is C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups, this group is substituted with, for example, 1 to 3 hydroxy groups, preferably 1 or 2 hydroxy groups.

[0301] When R_b , R_c , or R_d is 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R^{14} , C_{6-10} aryl optionally substituted with one or more substituents R^{14} , or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14} , this group is substituted with, for example, 1 to 4 substituents R^{14} , preferably 1 to 3 substituents R^{14} .

[0302] When R^4 is C_{6-10} aryl optionally substituted with one or more substituents R_g , this C_{6-10} aryl group is preferably phenyl.

[0303] In the group $-NR^{39}R^{40}$, R^{39} is preferably a hydrogen atom or C_{1-3} alkyl, more preferably methyl. R^{40} is preferably C_{1-3} alkyl, C_{1-3} alkoxy-substituted C_{1-4} alkyl, a group $-\text{CH}((\text{CH}_2)_{v1}\text{COOR}^{57})-(\text{CH}_2)_{v2}\text{COOR}^{57}$, 4- to 6-membered heterocycloalkyl, a group $-(\text{CH}_2)_w\text{SO}_3\text{H}$, or a group $-(\text{CH}_2)_{x1}\text{CH}(\text{COOH})-(\text{CH}_2)_{x2}\text{SO}_3\text{H}$, more preferably methoxyethyl, 2-methoxy-1,1-dimethylethyl, 2-methoxy-2-methyl-1-propyl, a group $-\text{C}_2\text{H}_3(\text{COOH})_2$, a group $-\text{C}_3\text{H}_5(\text{COOH})_2$, oxetanyl, methyl, a group $-\text{C}_2\text{H}_4\text{SO}_3\text{H}$, or a group $-\text{CH}(\text{COOH})\text{CH}_2\text{SO}_3\text{H}$. Examples of the group $-NR^{39}R^{40}$ include $\text{N}-((\text{C}_{1-3}\text{ alkoxy})\text{C}_{1-4}\text{ alkyl})\text{-N}-(\text{C}_{1-3}\text{ alkyl})\text{amino}$. In this context, "[$\text{N}-((\text{C}_{1-3}\text{ alkoxy})\text{C}_{1-4}\text{ alkyl})\text{-N}-(\text{C}_{1-3}\text{ alkyl})\text{amino}$]" is as defined above. The group $-NR^{39}R^{40}$ is preferably $\text{N}-(2\text{-methoxyethyl})\text{-N}-(\text{methyl})\text{amino}$, [$\text{N}-(2\text{-methoxy-1,1-dimethylethyl})\text{-N}-(\text{methyl})\text{amino}$]ethoxy, [$\text{N}-(2\text{-methoxy-2,2-dimethylethyl})\text{-N}-(\text{methyl})\text{amino}$]ethoxy, a group


-NH-CH(COOH)-CH₂(COOH), a group -NH-CH(CH₂COOH)-CH₂COOH, N-oxetanyl-N-methylamino, dimethylamino, a group -NH-C₂H₄-SO₃H, or a group -NH-CH(COOH)-CH₂-SO₃H.

[0304] In the group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², q1 is preferably 2, and q2 is preferably an integer selected from 2 to 5. R⁴¹ is preferably a hydrogen atom or methyl, and R⁴² is preferably C₁₋₆ alkyl (preferably n-hexyl) substituted with 1 to 5 hydroxy groups (preferably 5 hydroxy groups) or methoxyethyl.

[0305] In the group -(O(CH₂)_{r1})_{r2}-C(O)NR⁴³R⁴⁴, r1 is preferably 2 or 3, and r2 is preferably 1. R⁴³ is preferably methyl. R⁴⁴ is preferably C₁₋₆ alkyl (preferably n-hexyl) substituted with 1 to 5 hydroxy groups (preferably 5 hydroxy groups).

[0306] The compound of the formula (I) wherein R¹ and R⁵ together with the carbon atom to which they are attached form a C₃₋₆ saturated carbocyclic ring can be represented by the following formula (I-a) or (I-b):

[0307] [Formula 9]




[0308] The group -(CH₂)_q- wherein q is 0 means a single bond.

[0309] In the present invention, the compound of the formula (I) wherein R³ is C₁₋₄ alkyl substituted with Re, which can be represented by the formula (I-c) shown below. In the present invention, the compound of the formula (I) wherein R³ is C₁₋₄ alkyl substituted with Re, Re is substituted with one to three substituents Ra, and the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk can be represented by the formula (I-d) shown below.

[0310]

[Formula 10]



[0311] In the formula (I-c), n_1 is an integer selected from 1 to 4; n_2 is an integer selected from 0 or more; and R^1 , R^4 , R^5 , Ar^1 , Ra , Rb , Rc , Rd , and Re are as defined in any of (1-1) to (1-48), (2-1) to (2-25), (3-1) to (3-12), (4-1) to (4-12), (5-1) to (5-12), and (6-1) to (6-10).

In the formula (I-d), n_3 , n_4 , and n_5 are each independently an integer selected from 0 and 1, provided that at least one of n_3 , n_4 , and n_5 is 1; n_1 is as defined above; R^1 , R^4 , R^5 , Ar^1 , Rb , Rc , Rd , and Re are as defined in any of (1-1) to (1-48), (2-1) to (2-25), (3-1) to (3-12), (4-1) to (4-12), (5-1) to (5-12), and (6-1) to (6-10).

In the present invention, n_2 is preferably an integer selected from 1 to 4, more preferably an integer selected from 1 to 3.

[0312] In the present invention, the phrase “optionally substituted with” or “substituted with” means “optionally substituted with one substituent” or “substituted with one substituent” respectively unless a number of substituents (for example, “one or more”, “one to three”, “one or two”, “two” or “one”) is specified. For example, “B optionally substituted with A” and “B substituted with A” mean “B optionally substituted with one A” and “B substituted with one A” respectively.

[0313] In the present invention, examples of the salt of the compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) include acid-addition salts and base-addition salts. Examples of the acid-addition salts include: hydrochloride, hydrobromide, hydroiodide, phosphate, phosphonate, and sulfate; sulfonates such as methanesulfonate, ethanesulfonate, benzenesulfonate, and p-toluenesulfonate; and carboxylates such as acetate, citrate, malate, tartrate, succinate, salicylate, maleate, fumarate, benzoate, malonate, glycolate,

oxalate, glucuronate adipate, glutarate, ketoglutarate, and hippurate. Examples of the base-addition salts include: alkali metal salts such as sodium salt and potassium salt; alkaline earth metal salts such as magnesium salt and calcium salt; ammonium salts such as ammonium salt, alkylammonium salt, dialkylammonium salt, trialkylammonium salt, and tetraalkylammonium salt; and amino acid salts such as lysine salt, arginine salt, glycine salt, valine salt, threonine, salt, serine salt, proline salt, and alanine salt. These salts are each produced by the contact of the compound with an acid or a base available in pharmaceutical production.

[0314] In the present invention, the compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) or the salt thereof may be an anhydrate or may form a solvate such as a hydrate. In this context, the "solvate" refers to a solid formed by a complex of the compound molecule and a solvent molecule and refers to, for example, a hydrate formed from water as a solvent. The solvate other than the hydrate includes a solid containing an alcohol (e.g., methanol, ethanol, and n-propanol), dimethylformamide, or the like.

[0315] The compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) and the salt thereof may be found as some tautomers, for example, keto and enol forms, imine and enamine forms, and mixtures thereof. Such tautomers are present as a tautomeric mixture in a solution. In a solid form, one of the tautomers is usually predominant. Although one of the tautomers may be described herein, all tautomers of the compound of the present invention are included in the present invention.

[0316] The present invention further includes all stereoisomers (e.g., enantiomers and diastereomers (including cis and trans geometric isomers)) of the compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d), racemates of the isomers, and other mixtures. The compound of the present invention may have, for example, one or more asymmetric atoms. The compound of the present invention includes racemic mixtures, diastereomeric mixtures, and enantiomers of such a compound.

[0317] The compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) may be obtained in a free form. In this case, the free compound can be converted by a

routine method to the salt that may be formed by the compound or to the hydrate or the solvate of the compound or the salt.

[0318] Alternatively, the compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) may be obtained as the salt, hydrate, or solvate of the compound. In this case, this form can be converted to the free form of the compound according to a routine method.

[0319] Each element constituting the compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) may be any isotope. The present invention encompasses a compound of the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) containing such an isotope. The isotope in the compound refers to a variant of the element in which at least one atom is replaced with an atom with the same atomic number (the same number of protons) and a different mass number (total number of protons and neutrons). Examples of the isotope contained in the pharmaceutical drug of the present invention include a hydrogen atom, a carbon atom, a nitrogen atom, an oxygen atom, a phosphorus atom, a sulfur atom, a fluorine atom, and a chlorine atom including ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl . Particularly, a radioisotope, such as ^3H or ^{14}C , which undergoes radioactive decay are useful in, for example, the *in vivo* histological distribution tests of the pharmaceutical drug or the compound. Stable isotopes rarely vary in abundance without decay and are also free from radioactivity. These stable isotopes can therefore be used with safety. The isotope in the compound serving as an active ingredient in the pharmaceutical drug of the present invention can be converted according to a routine method by the replacement of a reagent used in synthesis with a reagent containing the corresponding isotope.

[0320] The compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) may be administered in the form of a prodrug. The present invention also includes such a prodrug of the compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d). In this context, the "prodrug" of the present invention means a derivative of the compound of the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) that is converted after administration to the compound of the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) or the pharmaceutically acceptable salt thereof through enzymatic or nonenzymatic degradation under physiological

conditions. The prodrug may be inactive when administered to a patient. The prodrug is converted *in vivo* to the active compound of the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d).

[0321] The prodrug is converted to, for example, a desired pharmaceutical formulation at a particular pH or by the action of an enzyme. The prodrug is typically a compound that forms a free acid *in vivo* and a compound having a hydrolyzable ester group. Such a hydrolyzable ester group is, for example, but not limited to, a group represented by the formula -COOR_x, wherein R_x is selected from C₁₋₄ alkyl, C₂₋₇ alkanoyloxymethyl, 1-(C₄₋₉ alkanoyloxy)ethyl, 1-methyl-1-(C₅₋₁₀ alkanoyloxy)-ethyl, (C₃₋₆ alkoxy)carbonyloxymethyl, 1-[(C₄₋₇ alkoxy)carbonyloxy]ethyl, 1-methyl-1-[(C₅₋₈ alkoxy)carbonyloxy]ethyl, N-[(C₃₋₉ alkoxy)carbonyl]aminomethyl, 1-(N-[(C₄₋₁₀ alkoxy)carbonyl]amino)ethyl, 3-phthalidyl, 4-crotonolactonyl, γ-butyrolacton-4-yl, [N,N-di(C₁₋₂ alkyl)amino]C₂₋₃ alkyl (e.g., N,N-dimethylaminoethyl), (carbamoyl)C₁₋₂ alkyl, [N,N-di(C₁₋₂ alkyl)carbamoyl]C₁₋₂ alkyl, (piperidino)C₂₋₃ alkyl, (pyrrolidino)C₂₋₃ alkyl, and (morpholino)C₂₋₃ alkyl.

[0322] Pharmaceutical drug of the present invention

One aspect of the present invention provides a pharmaceutical drug and a pharmaceutical composition containing the compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) or the salt thereof, or the solvate of the compound or the salt.

[0323] The pharmaceutical drug of the present invention contains the compound represented by the formula (I), (II), (III), (I-a), (I-b), (I-c), or (I-d) as an active ingredient and further contains a pharmaceutically acceptable carrier. The term "pharmaceutically acceptable carrier" used herein means one or more compatible solid or liquid excipients or encapsulating materials that are suitable for administration to mammals. The term "pharmaceutically acceptable" used herein means pharmaceutically available from the viewpoint of efficacy, safety, etc. The pharmaceutically acceptable carrier used is suitable for administration to an animal, preferably a mammal, to be treated, from the viewpoint of safety and has sufficiently high purity.

[0324] Examples of a material that may be used as the pharmaceutically acceptable carrier

include: sugars such as lactose, glucose, and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as carboxymethylcellulose sodium, ethylcellulose, and methylcellulose; tragacanth gum powder; malt; gelatin; talc; solid lubricants such as stearic acid and magnesium stearate; calcium sulfate; plant oils such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil, and cacao oil; polyhydric alcohols such as propylene glycol, glycerin, sorbitol, mannitol, and polyethylene glycol; alginic acid; emulsifiers such as TWEEN; wetting agents such as lecithin; colorants; flavors; tableting agents; stabilizers; antioxidants; antiseptics; pyrogen-free water; isotonic saline; and phosphate buffer solutions.

[0325] The compound of the present invention may be used as a NaPi-IIb inhibitor or a preventive and/or therapeutic agent for hyperphosphatemia, secondary hyperparathyroidism, or chronic renal failure. In this case, examples of an administration method thereof include oral, rectal, parenteral (intravenous, intramuscular, or subcutaneous), intracisternal, intravaginal, intraperitoneal, intravesical and local (drip, powder, ointment, gel, or cream) routes, and inhalation (into the oral cavity or using nasal sprays). Examples of the dosage form thereof include tablets, capsules, granules, powders, pills, aqueous and nonaqueous oral solutions and suspensions, and parenteral solutions charged in containers adapted to division into individual doses. Alternatively, the dosage form may be adapted to various administration methods encompassing controlled-release formulations as in subcutaneous implantation.

[0326] These preparations are produced by a well known method using additives such as excipients, lubricants (coating agents), binders, disintegrants, stabilizers, corrigents, and diluents.

[0327] Examples of the excipients can include starches such as starch, potato starch, and corn starch, lactose, crystalline cellulose, and calcium hydrogen phosphate.

[0328] Examples of the coating agents can include ethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, shellac, talc, carnauba wax, and paraffin.

[0329] Examples of the binders can include polyvinylpyrrolidone, Macrogol, and the same compounds as those listed as the excipients.

[0330] Examples of the disintegrants can include the same compounds as those listed as the excipients and chemically modified starches and celluloses such as croscarmellose sodium, carboxymethyl starch sodium, and cross-linked polyvinylpyrrolidone.

[0331] Examples of the stabilizers can include: p-hydroxybenzoate esters such as methylparaben and propylparaben; alcohols such as chlorobutanol, benzyl alcohol, and phenylethyl alcohol; benzalkonium chloride; phenols such as phenol and cresol; thimerosal; dehydroacetic acid; and sorbic acid.

[0332] Examples of the corrigents can include sweeteners, acidulants, and flavors usually used.

[0333] The liquid preparations can be produced using a solvent such as ethanol, phenol, chlorocresol, purified water, or distilled water.

[0334] Examples of the surfactants or emulsifiers can include polyoxyl 40 stearate and Lauromacrogol.

[0335] When the compound of the present invention is used as a NaPi-IIb inhibitor, PiT-1 inhibitor, PiT-2 inhibitor or a preventive and/or therapeutic agent for hyperphosphatemia, secondary hyperparathyroidism, or chronic renal failure, the amount of the pharmaceutical drug of the present invention used differs depending on symptoms, age, body weight, relative health conditions, the presence of other medications, administration methods, etc. In the case of oral agents, a general effective amount, for example, for a patient (warm-blooded animal, particularly a human) is preferably 1 to 20 mg/kg body weight, more preferably 1 to 10 mg/kg body weight, per day in terms of the amount of the active ingredient (the compound of the present invention represented by the formula (I)). The daily dose in an adult patient having a normal body weight is in the range of preferably 60 to 1200 mg.

[0336] Hereinafter, general methods for producing the compound of the present invention and Examples will be shown.

[0337] General synthesis method


The compound of the present invention can be synthesized by various methods. Some of the methods will be described with reference to schemes shown below. These

schemes are provided for illustrative purposes, and the present invention is not limited by chemical reactions and conditions described herein. Although some substituents are excluded from the schemes shown below for easy understanding, such exclusion is not intended to limit the disclosure of the schemes. Typical compounds of the present invention can be synthesized using appropriate intermediates, compounds known in the art, and reagents. In the formulas of the general synthesis methods described below, variable groups represented by R^1 , R^2 , etc. and variables represented by n , etc. have the same meanings as those of the variable groups represented by R^1 , R^2 , etc. and the variables represented by n , etc. in the compounds represented by general formulas defined herein.

[0338] The compound of the present invention can be synthesized by production methods shown below.

[0339] Scheme 1 (Method A)

[0340] [Formula 11]



[0341] wherein R^{21} represents C_{1-3} alkyl; R^{22} represents C_{1-5} alkyl; and $R^3=O$ (2) represents aldehyde or ketone formed by the conversion of a carbon atom at the linkage position of alkyl represented by R^3 to carbonyl.

Step 1 involves reacting a compound of the formula (1) with aldehyde or ketone (2) in an appropriate solvent such as methanol or dichloromethane to synthesize a compound (3). The reaction is performed at a temperature of, for example, 0°C to room temperature for a time of, for example, 0.5 hours to 24 hours. The obtained hydrazone derivative (3) is

isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0342] A large number of applicable methods have been reported as to the synthesis method for the hydrazine (1) shown in Scheme 1 (this reaction can be performed with reference to, for example, Synthetic communications, 40, 654-660; 2010, Journal of Heterocyclic Chemistry, 24 (3), 725-731; 1987, and Synthesis, (6), 423-424; 1979).

[0343] Step 2 involves reducing the hydrazone (3) in the presence of a reducing agent such as sodium cyanoborohydride or borane pyridine in an appropriate solvent such as methanol or acetic acid to obtain hydrazine (4). The reaction is performed at a temperature of, for example, 0°C to 50°C for a time of, for example, 0.5 hours to 60 hours. The obtained hydrazine (4) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0344] Step 3 involves reacting the hydrazine (4) with half ester (5) using a condensing agent such as N,N'-dicyclohexylcarbodiimide (DCC), 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methyl morpholinium chloride n-hydrate (DMT-MM), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyl uronium hexafluorophosphate (HATU), bromo-tris-pyrrolidino-phosphonium hexafluorophosphate (PyBrop), or 1-propanephosphonic acid cyclic anhydride (T3P) in an appropriate solvent such as dichloromethane, N,N-dimethylformamide, N,N-dimethylacetamide, acetonitrile, tetrahydrofuran, or ethyl acetate. The reaction is performed at a temperature of, for example, 0°C to 50°C for a time of, for example, 0.5 hours to 24 hours. The obtained ester form (6) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0345] The half ester (5) shown in Scheme 1 can be synthesized from Meldrum's acid and an alcohol (this reaction can be performed with reference to, for example, Organic Letters, 7 (3), 463-465; 2005).

[0346] Step 4 involves cyclizing the ester form (6) using a base such as potassium carbonate, cesium carbonate, sodium methoxide, or sodium hydride in an appropriate solvent such as methanol, N,N-dimethylformamide, N,N-dimethylacetamide, dimethyl sulfoxide,

diethyl ether, or tetrahydrofuran. The reaction is performed at a temperature of, for example, 0°C to 110°C for a time of, for example, 0.5 hours to 24 hours. The obtained cyclized form (7) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.


[0347] Step 5 involves reacting the cyclized form (7) with various amines (8) in an appropriate solvent such as benzene, toluene, xylene, ethyl acetate, tetrahydrofuran, N,N-dimethylformamide, N,N-dimethylacetamide, or dimethyl sulfoxide. The reaction is performed at a temperature of, for example, 50°C to 120°C for a time of, for example, 0.5 hours to 5 hours. The obtained amide form (formula I) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0348] R²¹ is preferably methyl, and R²² is preferably methyl or i-butyl.

[0349] The compound of the formula I can also be synthesized through the reaction between a keto form (9) and isocyanate as shown in Scheme 2 (Method B).

[0350] Scheme 2 (Method B)

[0351] [Formula 12]



[0352] wherein R²² is as defined in Scheme 1.

The keto form (9) as a product in Step 1 can be synthesized by the decarboxylation of an ester form (7) (this reaction can be performed with reference to, for example, Synthesis, (15), 2487-2491; 2009).


[0353] Step 2 involves reacting the keto form (9) with isocyanate in the presence of a base such as sodium hydride, potassium carbonate, or cesium carbonate in an appropriate solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, or tetrahydrofuran. The reaction is performed at a temperature of, for example, 0°C to 50°C for a time of, for example, 0.5

hours to 5 hours. The obtained amide form (formula I) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0354] The compound of the formula I can also be synthesized by the cyclization of an amide form as shown in Scheme 3 (Method C).

[0355] Scheme 3 (Method C)

[0356] [Formula 13]



[0357] wherein R²¹ is as defined in Scheme 1.

[0358] Step 1 involves reacting the hydrazine (4) with half amide (10) in the presence of a condensing agent such as N,N'-dicyclohexylcarbodiimide (DCC), 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methyl morpholinium chloride n-hydrate (DMT-MM), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyl uronium hexafluorophosphate (HATU), bromo-tris-pyrrolidino-phosphonium hexafluorophosphate (PyBrop), or 1-propanephosphonic acid cyclic anhydride (T3P) in an appropriate solvent such as dichloromethane, N,N-dimethylformamide, N,N-dimethylacetamide, acetonitrile, tetrahydrofuran, or ethyl acetate. The reaction is performed at a temperature of, for example, 0°C to 50°C for a time of, for example, 0.5 hours to 24 hours. The obtained amide form (11) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0359] The half amide (10) shown in Scheme 3 can be synthesized from Meldrum's acid and amine (this reaction can be performed with reference to, for example, Bioorganic & Medicinal Chemistry Letters, 19 (13), 3632-3636; 2009).


[0360] Step 2 involves cyclizing the amide form (11) using a base such as sodium methoxide, potassium carbonate, cesium carbonate, or sodium hydride in an appropriate solvent such as methanol, N,N-dimethylformamide, N,N-dimethylacetamide, or tetrahydrofuran. The reaction is performed at a temperature of, for example, 0°C to 110°C

for a time of, for example, 0.5 hours to 24 hours. The obtained compound of the formula I is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0361] The compound of the formula I can also be synthesized by alkylation as shown in Scheme 4 (Method D).

[0362] Scheme 4 (Method D)

[0363] [Formula 14]



[0364] Three exemplary methods of Step 1 will be shown below.

[0365] Method 1 involves debenzylating a compound (12) using a palladium catalyst such as palladium(0)-carbon or palladium(II) hydroxide-carbon or a platinum catalyst such as platinum oxide (PtO₂) in an appropriate solvent such as methanol, ethyl acetate, N,N-dimethylformamide, or N,N-dimethylacetamide in a hydrogen atmosphere. The reaction is performed at a temperature of, for example, room temperature to the boiling point of the solvent for a time of, for example, 1 hour to 24 hours. Method 2 involves reacting the compound (12) in the presence of an oxidizing agent such as 2,3-dichloro-5,6-dicyano-p-benzoquinone or ammonium hexanitratocerate(IV) in an appropriate solvent such as dichloromethane or acetonitrile. The reaction is performed at a temperature of, for example, room temperature to the boiling point of the solvent for a time of, for example, 1 hour to 60 hours. Method 3 involves reacting the compound (12) in the presence of an organic acid such as trifluoroacetic acid or trifluoromethanesulfonic acid in an appropriate solvent such as dichloromethane or methanol. The reaction is performed at a temperature of, for example, 0°C to the boiling point of the solvent for a time of, for example, 0.5 hours to 5 hours. The obtained NH form (13) is isolated by a general technique and may be purified, if necessary,

by crystallization or chromatography.


[0366] Step 2 involves reacting the NH form (13) with an alkylating agent such as alkyl halide or alkyl sulfonate in the presence of an appropriate base such as sodium hydride, potassium t-butoxide, or potassium pentoxide in an appropriate solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, tetrahydrofuran, or dimethyl sulfoxide. The reaction is performed at a temperature of, for example, 0°C to the boiling point of the solvent for a time of, for example, 0.5 hours to 24 hours. The obtained compound of the formula I is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0367] In $-\text{OSO}_2\text{R}^{23}$ shown in this scheme, R^{23} is C_{1-5} alkyl optionally substituted with one or more halogen atoms or aryl, wherein the aryl group is optionally substituted with one or more halogen atoms or alkyl groups. Specific examples of R^{23} include methyl, trifluoromethyl, phenyl, and 4-methylphenyl.

[0368] The compound of the formula I can also be synthesized by alkylation as shown in Scheme 5 (Method E).

[0369] Scheme 5 (Method E)

[0370] [Formula 15]



[0371] wherein R^{22} is as defined in Scheme 1.

The reaction of Step 1 of Scheme 5 (Method E) can be pursued in the same way as in Step 1 of Scheme 4 (Method D).


[0372] The reaction of Step 2 of Scheme 5 (Method E) can be pursued in the same way as in Step 2 of Scheme 4 (Method D). The compound of the formula I can be obtained in the

same way as in Step 5 of Scheme 1 from the obtained ester form (7).

[0373] Side chain introduction method

Scheme 6 (Method F)

[0374] [Formula 16]




[0375] wherein n is an integer selected from 1 to 10, and R^{24} is C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} .

The method of Scheme 6 (Method F) involves reacting a phenol derivative (16) with an alkylating agent such as alkyl halide or alkyl sulfonate in the presence of an appropriate base such as sodium hydride, potassium carbonate, cesium carbonate, potassium t-butoxide, or potassium pentoxide in an appropriate solvent such as tetrahydrofuran, acetonitrile, N,N-dimethylformamide, N,N-acetamide, dimethyl sulfoxide, or acetone. The reaction is performed at a temperature of, for example, 0°C to the boiling point of the solvent for a time of, for example, 0.5 hours to 12 hours. The obtained compound represented by the formula III is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography. The compound obtained in Scheme 6 can be subjected, if necessary, to the deprotection reaction of various protective groups and other procedures to synthesize a derivative.

[0376] Scheme 7 (Method G)

[0377]

[Formula 17]




wherein n and R^{24} are as defined in Scheme 6.

[0378] The method of Scheme 7 (Method G) involves reacting a phenol derivative (16) with various alcohols using an appropriate Mitsunobu reagent such as diethyl azodicarboxylate (DEAD), diisopropyl azodicarboxylate (DIAD), N,N,N',N' -tetraisopropylazodicarboxamide (TIPA), 1,1'-(azodicarbonyl)dipiperidine (ADDP), or N,N,N',N' -tetramethylazodicarboxamide (TMAD) in the presence of an appropriate trivalent organic phosphorus reagent such as triphenylphosphine or tributylphosphine in an appropriate solvent such as dichloromethane, tetrahydrofuran, acetonitrile, toluene, or benzene. The reaction is performed at a temperature of, for example, 0°C to the boiling point of the solvent for a time of, for example, 0.5 hours to 24 hours. The obtained compound represented by the formula III is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography. The compound obtained in Scheme 7 can be subjected, if necessary, to the deprotection reaction of various protective groups and other procedures to synthesize a derivative.

[0379] Scheme 8 (Method H)

[0380]

[Formula 18]




[0381] wherein n is an integer selected from 1 to 10, and R^{25} is $(\text{C}_{1-4}$ alkoxy)carbonyl, C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , or C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} .

The method of Scheme 8 (Method H) involves reacting a bromobenzene or iodobenzene derivative (17) using a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0) ($\text{Pd}(\text{Ph}_3\text{P})_4$), tris(dibenzylideneacetone)dipalladium(0) ($\text{Pd}_2(\text{dba})_3$), 1,1'-bis(diphenylphosphino)ferrocene-palladium(II) dichloride-dichloromethane complex ($\text{PdCl}_2(\text{dppf})\cdot\text{CH}_2\text{Cl}_2$), palladium(II) acetate ($\text{Pd}(\text{OAc})_2$), or dichlorobis(triphenylphosphine)palladium(II) ($\text{PdCl}_2(\text{Ph}_3\text{P})_2$) supplemented, if necessary, with a phosphine ligand such as triphenylphosphine (Ph_3P), tri-*t*-butylphosphine (tBu_3P), or tri-*o*-tolylphosphine ($(\text{o-tol})_3\text{P}$), a copper catalyst such as copper(I) iodide, and an appropriate base such as sodium carbonate, triethylamine, or potassium carbonate in an appropriate solvent such as dichloromethane, tetrahydrofuran, *N,N*-dimethylformamide, *N,N*-dimethylacetamide, toluene, benzene, 1,2-dimethoxyethane, 1,4-dioxane, ethanol, or acetonitrile to introduce thereinto, for example, sulfur, alkane, alkyne, or alkoxy carbonyl. When R^{25} is $(\text{C}_{1-4}$ alkoxy)carbonyl, for example, carbon monoxide and C_{1-4} alcohol can be used as a reagent in the reaction. When R^{25} is C_{1-10} alkyl optionally substituted with one or more substituents R^{10} or C_{2-10} alkynyl, for example, acetylene or $(\text{C}_{1-8}$ alkyl)- $\text{C}\equiv\text{CH}$ can be used as a reagent in the reaction. When R^{25} is C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} , for example, C_{1-4} alkylmercaptan can be used

as a reagent in the reaction. The reaction is performed at a temperature of, for example, 0°C to the boiling point of the solvent for a time of, for example, 0.5 hours to 60 hours. The obtained compound represented by the formula IV is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography. The compound obtained in Scheme 8 can be subjected, if necessary, to the hydrogenation reaction of a double bond and a triple bond, the oxidation reaction or alkylation of a sulfur atom, the deprotection reaction of various protective groups, and other procedures to synthesize a derivative.

[0382] Scheme 9 (Method I)

[0383] [Formula 19]




[0384] wherein R^{22} , R^{23} , and R^{25} are as defined in Schemes 1 and 8.

The reaction of Scheme 9 (Method I) can be pursued in the same way as in Scheme 8 with a compound (18) as a starting material. The compound of the formula IV can be obtained in the same way as in Step 5 of Scheme 1 from the obtained ester form (19).

[0385] Scheme 10 (Method J)

[0386] [Formula 20]




[0387] wherein

Ar^1 is C_{6-10} aryl or 5- to 10-membered heteroaryl, wherein this group is optionally substituted with Rb and/or Rc ; and

Rd is selected from 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R^{14} , C_{6-10} aryl optionally substituted with one or more substituents R^{14} , and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14} .

Examples of -B(pin) in the above formula include the following structure:

[0388] [Formula 21]




[0389] The method of Scheme 10 (Method J) involves reacting an aryl halide derivative (20) with aryl boronic acid, heteroaryl boronic acid, heterocycloalkyl boronic acid, aryl boronic acid ester, heteroaryl boronic acid ester, heterocycloalkyl boronic acid ester, aryltrialkyltin, heteroaryltrialkyltin, heterocycloalkyltrialkyltin, or the like using a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0) ($Pd(Ph_3P)_4$), tris(dibenzylideneacetone)dipalladium(0) ($Pd_2(dba)_3$), 1,1'-bis(diphenylphosphino)ferrocene-palladium(II) dichloride-dichloromethane complex ($PdCl_2(dppf) \cdot CH_2Cl_2$), or palladium(II) acetate ($Pd(OAc)_2$) supplemented, if necessary, with a phosphine ligand such as triphenylphosphine (Ph_3P), tri-*t*-butylphosphine (tBu_3P), or tri-*o*-tolylphosphine ($((o-tol)_3P)$) and an appropriate base such as sodium carbonate, triethylamine, or potassium carbonate in an appropriate solvent such as tetrahydrofuran, acetonitrile, toluene, *N,N*-dimethylformamide, *N,N*-dimethylacetamide, 1,2-dimethoxyethane (DME), or 1,4-dioxane in a nitrogen atmosphere to introduce an aryl group thereinto. The reaction is performed at a temperature of, for example, room temperature to the boiling point of the solvent for a time of, for example, 0.5 hours to 12 hours. The obtained bisaryl form (formula V) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0390] Scheme 11 (Method K)

[0391]

[Formula 22]



[0392] wherein

R²³ is as defined above;


Ar¹ and Rd are as defined in Scheme 10; and

R¹⁴ is selected from cyano, C₁₋₄ alkoxy optionally substituted with one or more halogen atoms, and -NR²⁷R²⁸ (R²⁷ and R²⁸ are each independently selected from optionally (C₁₋₃ alkoxy)carbonyl-substituted C₁₋₄ alkyl).

The method of Scheme 11 (Method K) involves reacting heteroaryl halide or sulfonic acid heteroaryl ester (21) with a nucleophile such as amine, nitrile, or alcohol, if necessary in the presence of a base such as sodium hydride, sodium alkoxide, potassium carbonate, sodium carbonate, triethylamine, diisopropylethylamine, or diazabicycloundecene (DBU) in an appropriate solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, tetrahydrofuran, or dimethyl sulfoxide. The reaction is performed at a temperature of, for example, room temperature to the boiling point of the solvent for a time of, for example, 0.5 hours to 24 hours. The obtained bisaryl form (formula VI) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0393] Scheme 12 (Method L)

[0394] [Formula 23]



[0395] wherein

R^{23} is as defined in Scheme 11;

Rd is 5- to 10-membered heterocycloalkyl, C_{6-10} aryl, or 5- to 10-membered heteroaryl; and


Y is selected from cyano, C_{1-4} alkoxy optionally substituted with one or more halogen atoms, and $-NR^{27}R^{28}$ (R^{27} and R^{28} are each independently selected from optionally (C_{1-4} alkoxy)carbonyl-substituted C_{1-4} alkyl).

The method of Scheme 12 (Method L) involves reacting heteroaryl halide, sulfonic acid heteroaryl ester, or nitroheteroaryl (22) having an electron-withdrawing substituent (EWG) with a nucleophile such as amine, nitrile, or alcohol, if necessary in the presence of a base such as sodium hydride, sodium alkoxide, potassium carbonate, sodium carbonate, triethylamine, diisopropylethylamine, or diazabicycloundecene (DBU) in an appropriate solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, tetrahydrofuran, dimethyl sulfoxide, or an alcoholic solvent. The reaction is performed at a temperature of, for example, room temperature to the boiling point of the solvent for a time of, for example, 0.5 hours to 24 hours. The obtained bisaryl form (the formula VII) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0396] Starting material synthesis

Scheme 13 (Method M)

[0397] [Formula 24]



R^{20} : $B(OH)_2$, B(pin), $SnBu_3$

Y: Cl, Br, I, OSO_2R^7


[0398] wherein Ar^1 and Rd are as defined in Scheme 10.

According to Scheme 13 (Method M), the arylamine (8) (wherein Rd is 5- to 10-

membered heterocycloalkyl optionally substituted with one or more substituents R^{14} , C_{6-10} aryl optionally substituted with one or more substituents R^{14} , or 5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14}) for use in the reaction of Step 5 of Scheme 1 is synthesized. This method involves reacting aryl halide (23) or (26) with aryl boronic acid, aryl boronic acid ester, aryltrialkyltin, or the like in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0) supplemented, if necessary, with a base such as triethylamine, sodium carbonate, potassium carbonate, cesium carbonate, or sodium hydroxide in an appropriate solvent such as water, toluene, ethanol, N,N-dimethylformamide, N,N-dimethylacetamide, 1,4-dioxane, 1,2-dimethoxyethane, or tetrahydrofuran. The reaction is performed at a temperature of, for example, room temperature to the boiling point of the solvent for a time of, for example, 0.5 hours to 24 hours. The obtained bisaryl form (25) or (28) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0399] Scheme 14 (Method N)

[0400] [Formula 25]




[0401] wherein R^1 , R^4 , R^5 , and R^{21} are as defined above.

According to Scheme 14 (Method N), the hydrazine for use in the reaction of Step 1 of Scheme 1 is synthesized. The compound (1) can be synthesized with reference to, for example, Bioorganic & Medicinal Chemistry Letters 13 (2003) 2413-2418.

[0402] Scheme 15 (Method O)

[0403]

[Formula 26]



[0404] wherein R^{26} is benzyl, methyl, or t-butyl, and R^1 , R^4 , and R^5 are as defined above.


According to Scheme 15 (Method O), the amine (34) for synthesis of the hydrazine (1) for use in the reaction of Step 1 of Scheme 1 is synthesized.

[0405] Step 1 involves N-alkylating a carbamate-protected amino acid using an appropriate base reagent such as sodium hydride, potassium carbonate, or cesium carbonate or a silver salt such as silver(I) oxide in the presence of alkyl halide in an appropriate solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, tetrahydrofuran, or dimethyl sulfoxide. The reaction is performed at a temperature of, for example, 0°C to 80°C for a time of, for example, 0.5 hours to 5 hours. The obtained N-alkyl form (33) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography.

[0406] Step 2 involves deprotecting the carbamate (33), for example, in the presence of an acid such as hydrochloric acid or trifluoroacetic acid or by catalytic hydrogen reduction in the presence of a catalyst such as palladium. The reaction is performed at a temperature of, for example, 0°C to 50°C for a time of, for example, 0.5 hours to 24 hours. The reaction solvent used is, for example, an appropriate solvent such as dichloromethane or acetonitrile for the acidic conditions or, for example, a solvent such as ethyl acetate, N,N-dimethylformamide, N,N-dimethylacetamide, dimethyl sulfoxide, or methanol for the catalytic hydrogen reduction. The obtained N-alkyl form (34) is isolated by a general technique and may be purified, if necessary, by crystallization or chromatography. The hydrazine (1) shown in Scheme 1 can be synthesized from the amine (34) with reference to, for example, Synthetic communications, 40, 654-660; 2010, Journal of Heterocyclic Chemistry, 24 (3), 725-731; 1987, and Synthesis, (6), 423-424; 1979.

[0407] Scheme 16 (Method P)

[0408] [Formula 27]



[0409] wherein R¹, R⁴, R⁵, and R²⁶ are as defined above.

According to Scheme 16 (Method P), the carbamate-protected amino acid (36) for synthesis of the compound (33) for use in the reaction of Step 1 of Scheme 15 is synthesized. The compound (36) can be synthesized using Curtius rearrangement for carboxylic acid (35) (this reaction can be performed with reference to, for example, J. Org. Chem., 1994, 59 (26), 8215-8219). The compound (33) can be synthesized in the same way as in Step 1 of Scheme 15.

[0410] Hereinafter, the present invention will be described in more detail with reference to Reference Examples and Examples. However, the present invention is not limited by these examples.

EXAMPLES

[0411] The contents of the present invention will be further described with reference to Examples and Reference Examples below. However, the present invention is not limited by the contents of these examples. All starting materials and reagents were obtained from commercial suppliers or synthesized using methods known in the art. Each compound is purified by HPLC using AutoPurification HPLC/MS System (manufactured by Waters Corp.). ¹H-NMR or ¹³C-NMR spectra were measured using Me₄Si as an internal standard and Agilent 400-MR (manufactured by Agilent Technologies, Inc.) or AVANCE3 Cryo-TCI (Bruker Corp.) (s = singlet, brs = broad singlet, d = doublet, t = triplet, q = quartet, dd = double doublet, ddd = double double doublet, dt = double triplet, td = triple doublet, and m = multiplet). Mass spectrometry was performed using a mass spectrometer SQD (manufactured by Waters Corp.), 2020 (manufactured by Shimadzu Corp.), or 2010EV

(manufactured by Shimadzu Corp.). Retention time measurement and mass spectrometry in LCMS were performed using the following apparatuses and analysis conditions:


[0412] [Table 1]

LCMS analysis condition No.	Apparatus	Column	Mobile phase and gradient
SMD-TFA05	nexera/2020	Kinetex 2.1mmI.D.x50mm	0.05%TFA/0.05%TFA MeCN =95/5 → 0/100(1.5min) → 0/100(0.5min), 1mL/min
SMD-TFA50	nexera/2020	Kinetex 2.1mmI.D.x50mm	0.05%TFA/0.05%TFA MeCN =50/50 → 0/100(1min) → 0/100(1min), 1mL/min
SMD-FA05	nexera/2020	Ascentis Express C18 2.1mmI.D.x50mm	H ₂ O(0.1%FA)/MeCN(0.1%FA) 95/5 → 0/100(1.5min) → 100(0.6min), 1mL/min
SQD-AA05	UPLC/SQD	Ascentis Express C18 2.1mmI.D.x50mm	10mMAcONH ₄ /MeOH =95/5 → 0/100(1min) → 100(0.4min), 1mL/min
SQD-AA50	UPLC/SQD	Ascentis Express C18 2.1mmI.D.x50mm	10mMAcONH ₄ /MeOH =50/50 → 0/100(0.7min) → 100(0.7min), 1mL/min
SQD-FA05	UPLC/SQD	Ascentis Express C18 2.1mmI.D.x50mm	H ₂ O(0.1%FA)/MeCN(0.1%FA) =95/5 → 0/100(1min) → 100(0.4min), 1mL/min
SQD-FA50	UPLC/SQD	Ascentis Express C18 2.1mmI.D.x50mm	H ₂ O(0.1%FA)/MeCN(0.1%FA) =50/50 → 0/100(0.7min) → 100(0.7min), 1mL/min
SMD-AA05	nexera/2020	Ascentis Express C18 2.1mmI.D.x50mm	10mM AcONH ₄ /MeOH =95/5 → 0/100(1.5min) → 100(0.7min), 1mL/min
QC-SMD-TFA05	nexera/2020	Acquity 2.1mmI.D.x50mm,	0.05%TFA/0.05%TFA MeCN =95/5 → 0/100(1.5min) → 0/100(0.5min), 1mL/min
Ph-SMD-TFA05	UFLC-MS 2010EV	shim-pack XR-ODS 3.0mmI.D.x50mm	0.05%TFA/0.05%TFA MeCN =95/5 → 0/100(2min) → 0/100(1.1min), 1mL/min
ZQ-01	2525BGM/29 96PDA/ZQ20 00	Chromolith Flash RP-18e 4.6mmI.D. x 25mm	10mM AcONH ₄ /MeOH =95/5 → 0/100(3min) → 100(2min), 2mL/min
SQD-TFA05	UPLC/SQD	Ascentis Express C18 2.1mmI.D.x50mm	H ₂ O0.1%TFA)/MeCN(0.1%TF A) =95/5 → 0/100(1min) → 100(0.4min), 1mL/min

[0413] A compound having a C=N bond in a hydrazone structural formula represented by

the following formula

[0414] [Formula 28]



[0415] was used in next reaction without particular confirmation of whether the compound was in a cis form or a trans form or a mixture thereof.


<Example 1>

(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-4a,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-3-carboxamide

First Step

(S)-1-[(3-Fluoro-benzylidene)-amino]-pyrrolidine-2-carboxylic acid methyl ester

[0416] [Formula 29]



[0417] L-proline methyl ester hydrochloride (5.00 g, 30.2 mmol) was suspended in dichloromethane (60.4 mL), p-toluenesulfonic acid monohydrate (6.03 g, 31.7 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 10 minutes. After the reaction mixture was concentrated at reduced pressure and toluene was added, for azeotropic removal of water suspended in dichloromethane (60.4 mL), sodium nitrite (2.19 g, 31.7 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 2 hours. After the reaction mixture was filtered, the resultant was concentrated at reduced pressure to obtain (S)-1-nitroso-pyrrolidine-2-carboxylic acid methyl ester as a crude product. The obtained crude product was dissolved in acetic acid (30.2 mL) and water (30.2 mL), then zinc (19.7 g, 302 mmol) was added at 0°C, and the mixture was stirred under nitrogen atmosphere at room temperature for 1 hour. After sodium

bicarbonate was added to the reaction mixture at 0°C, methanol (60.4 mL) and 3-fluorobenzaldehyde (3.20 mL, 30.2 mmol) were added, and the mixture was stirred under nitrogen atmosphere at room temperature for 2 hours. After the reaction mixture was filtered, the resultant was concentrated at reduced pressure and then extracted with ethyl acetate. The organic layer was washed with water and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (5.68 g, 75%).


[0418] LCMS: m/z 251[M+H]⁺

HPLC retention time: 0.97 minutes (analysis condition SQD-AA05)

[0419] Second Step

(4aS)-1-[(3-Fluorophenyl)methyl]-4a,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-2,4-dione

[0420] [Formula 30]



[0421] (S)-1-[(3-Fluoro-benzylidene)-amino]-pyrrolidine-2-carboxylic acid methyl ester (5.68 g, 22.7 mmol) was dissolved in acetic acid (22.7 mL) and methanol (22.7 mL), sodium cyanoborohydride (5.70 g, 90.8 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 14 hours. A saturated aqueous sodium bicarbonate solution was added to the reaction mixture, then the mixture was concentrated at reduced pressure and extracted with ethyl acetate. The organic layer was washed with a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (S)-1-(3-fluorobenzylamino)-pyrrolidine-2-carboxylic acid methyl ester as a crude product. The obtained crude product was dissolved in tetrahydrofuran (22.7 mL), tripotassium phosphate (9.64 g, 45.4 mmol) and chlorocarbonyl-acetic acid methyl ester (2.67 mL, 45.4 mmol) were added at

0°C, and the mixture was stirred under nitrogen atmosphere at room temperature for 3 hours. 1 N hydrochloric acid was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium bicarbonate solution and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (S)-1-[(3-fluoro-benzyl)-(2-methoxycarbonyl-acetyl)-amino]-pyrrolidine-2-carboxylic acid methyl ester as a crude product. The obtained crude product was dissolved in N,N-dimethylformamide (22.7 mL), cesium carbonate (22.2 g, 68.1 mmol) was added, and the mixture was stirred under nitrogen atmosphere at 80°C for 9 hours. 1 N hydrochloric acid was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (S)-1-(3-fluoro-benzyl)-4-hydroxy-2-oxo-1,2,4a,5,6,7-hexahydro-pyrrolo[1,2-b]pyridazine-3-carboxylic acid methyl ester as a crude product. The obtained crude product was dissolved in acetonitrile (227 mL), water (0.41 mL, 22.7 mmol) was added, and the mixture was heated under reflux for 2 hours under nitrogen atmosphere. The reaction mixture was concentrated at reduced pressure and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (4.01 g, 67%).


[0422] LCMS: m/z 263[M+H]⁺

HPLC retention time: 0.81 minutes (analysis condition SQD-AA05)

[0423] Third Step

(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-4a,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-3-carboxamide

[0424] [Formula 31]



[0425] (4aS)-1-[(3-Fluorophenyl)methyl]-4a,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-2,4-

dione (1.50 g, 5.72 mmol) was dissolved in N,N-dimethylformamide (4.00 mL), and then 3-isocyanato-5-methyl-2-trifluoromethyl-furan (1.20 g, 6.29 mmol) and sodium hydride (50% by weight dispersion in mineral oil, 0.33 g, 6.86 mmol) were added, and the mixture was stirred under nitrogen atmosphere at room temperature for 90 minutes. 1 N hydrochloric acid was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (1.62 g, 63%).

[0426] LCMS: m/z 454[M+H]⁺

HPLC retention time: 1.10 minutes (analysis condition SQD-AA05)




[0427] <Example 2>

(4aR)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-4a,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-3-carboxamide

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from D-proline methyl ester hydrochloride and 3-fluoro-benzaldehyde.

[0428]

[Table 2]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(R)-1-[(3-Fluorobenzylidene)-amino]-pyrrolidine-2-carboxylic acid methyl ester	SQD-AA05	0.97	251
	(4aR)-1-[(3-Fluorophenyl)methyl]-4a,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.81	263
	(4aR)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-4a,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.10	454


[0429] <Example 3>

(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-6,6-dimethyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,7-dihydro-4aH-pyrrolo[1,2-b]pyridazine-3-carboxamide

First Step

(S)-4,4-Dimethyl-pyrrolidine-2-carboxylic acid ethyl ester hydrochloride

[0430] [Formula 32]






[0431] A 1,4-dioxane solution (4 M, 5.00 mL) of hydrogen chloride was added to (S)-4,4-dimethyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-ethyl ester (500 mg, 1.84 mmol), and the mixture was stirred under nitrogen atmosphere at room temperature for 1 hour. The reaction mixture was concentrated at reduced pressure, and toluene was added for azeotropic removal to obtain the title compound as a crude product.

[0432] Second Step

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from the obtained (S)-4,4-dimethyl-pyrrolidine-2-carboxylic acid ethyl ester hydrochloride and 3-fluoro-benzaldehyde.

[0433] [Table 3]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(S)-1-[(3-Fluorobenzylidene)amino]-4,4-dimethylpyrrolidine-2-carboxylic acid ethyl ester	SQD-AA05	1.09	293
	(4aS)-1-[(3-Fluorophenyl)methyl]-6,6-dimethyl-5,7-dihydro-4aH-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.94	291
	(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-6,6-dimethyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,7-dihydro-4aH-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.15	482


[0434] <Example 4>

(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-7,7-dimethyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6-dihydro-4aH-pyrrolo[1,2-b]pyridazine-3-carboxamide

First Step

(S)-5,5-Dimethylpyrrolidine-2-carboxylic acid methyl ester hydrochloride

[0435] [Formula 33]






[0436] (S)-5,5-Dimethyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester (500 mg, 2.06 mmol) was dissolved in N,N-dimethylformamide (2.06 mL), and then cesium carbonate (1.00 g, 3.08 mmol) and iodomethane (0.154 mL, 2.47 mmol) were added, and the mixture was stirred under nitrogen atmosphere at room temperature for 2 hours. 1 N hydrochloric acid was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with water, a saturated sodium bicarbonate solution, and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (S)-5,5-dimethyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester as a crude product. A 1,4-Dioxane solution (4 M, 5.00 mL) of hydrogen chloride was added to the obtained crude product, and the mixture was stirred under nitrogen atmosphere at room temperature for 1 hour. The reaction mixture was concentrated at reduced pressure, and toluene was added for azeotropic removal to obtain the title compound as a crude product.

[0437] Second Step

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from the obtained (S)-5,5-dimethyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride and 3-fluoro-benzaldehyde.

[0438]

[Table 4]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(S)-1-[(3-Fluorobenzylidene)amino]-5,5-dimethyl-pyrrolidine-2-carboxylic acid methyl ester	SQD-AA05	1.10	279
	(4aS)-1-[(3-Fluorophenyl)methyl]-7,7-dimethyl-5,6-dihydro-4aH-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.93	291
	(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-7,7-dimethyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6-dihydro-4aH-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.13	482




[0439] <Example 5>

(4aS)-1-[(2,3-Difluorophenyl)methyl]-6,6-difluoro-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,7-dihydro-4aH-pyrrolo[1,2-b]pyridazine-3-carboxamide

(S)-4,4-Difluoropyrrolidine-1,2-dicarboxylic acid 2-methyl 1-tert-butyl diester was used as a starting material, 2,3-difluorobenzaldehyde was used as a reagent, and operations similar to those of Example 3 were carried out to synthesize the compounds described in the following Table.

[0440]

[Table 5]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(S)-1-((2,3-Difluorobenzylidene)amino)-4,4-difluoropyrrolidine-2-carboxylic acid methyl ester	SMD-TFA05	1.24	305
	(4aS)-1-[(2,3-Difluorophenyl)methyl]-6,6-difluoro-5,7-dihydro-4aH-pyrrolo[1,2-b]pyridazine-2,4-dione	SMD-TFA05	1.04	317
	(4aS)-1-[(2,3-Difluorophenyl)methyl]-6,6-difluoro-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,7-dihydro-4aH-pyrrolo[1,2-b]pyridazine-3-carboxamide	SMD-TFA05	1.55	508


[0441] <Example 6>

(4aR)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-4a-methyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

First Step

(R)-2-Methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride

[0442] [Formula 34]






[0443] (R)-2-Methyl-pyrrolidine-2-carboxylic acid (300 mg, 2.32 mmol) was dissolved in dioxane (1.20 mL) and water (0.60 mL), and then sodium hydroxide (139 mg, 3.48 mmol) and di-tert-butyl dicarbonate (608 mg, 2.79 mmol) were added at 0°C, and the mixture was stirred at room temperature for 2 hours. 1 N hydrochloric acid was added to the reaction

mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (R)-2-methyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester as a crude product. In a similar manner to First Step of Example 4, the title compound was obtained from the obtained (R)-2-methyl-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester as a crude product.

[0444] Second Step

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from the obtained (R)-2-methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride and 3-fluoro-benzaldehyde.

[0445] [Table 6]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(R)-1-[(3-Fluorobenzylidene)amino]-2-methyl-pyrrolidine-2-carboxylic acid methyl ester	SQD-AA05	1.04	265
	(4aR)-1-[(3-Fluorophenyl)methyl]-4a-methyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.89	277
	(4aR)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-4a-methyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.15	468

[0446] <Example 7>




(4aR)-1-[(2,3-Difluorophenyl)methyl]-4-hydroxy-4a-methyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

First Step

In a similar manner to First to Third Steps of Example 1, the compounds described

in the following Table were synthesized from (R)-2-methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride obtained in First Step of Example 6 and 2,3-difluorobenzaldehyde.

[0447] [Table 7]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(R)-1-[(2,3-Difluorobenzylidene)-amino]-2-methyl-pyrrolidine-2-carboxylic acid methyl ester	SQD-AA05	1.08	283
	(4aR)-1-[(2,3-Difluorophenyl)methyl]-4a-methyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.89	295
	(4aR)-1-[(2,3-Difluorophenyl)methyl]-4-hydroxy-4a-methyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.16	486


[0448] <Example 8>

(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-4a-methyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

First Step

(S)-2-Methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride

[0449] [Formula 35]






[0450] In a similar manner to First Step of Example 6, the title compound was obtained from (S)-2-methyl-pyrrolidine-2-carboxylic acid as a crude product.

[0451] Second Step

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from (S)-2-methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride obtained in First Step and 3-fluorobenzaldehyde.

[0452] [Table 8]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(S)-1-[(3-Fluorobenzylidene)amino]-2-methylpyrrolidine-2-carboxylic acid methyl ester	SQD-AA05	1.04	265
	(4aS)-1-[(3-Fluorophenyl)methyl]-4a-methyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.92	277
	(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.15	468




[0453] <Example 9>

(4aR)-4a-Ethyl-1-[(3-fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

By carrying out operations similar to those of First to Third Steps of Example 1, the compounds described in the following Table were synthesized from (S)-2-ethyl-pyrrolidine-2-carboxylic acid ethyl ester hydrochloride and 3-fluorobenzaldehyde.

[0454]

[Table 9]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(R)-2-Ethyl-1-[(3-fluorobenzylidene)-amino]-pyrrolidine-2-carboxylic acid ethyl ester	SQD-AA05	1.14	293
	(4aR)-4a-Ethyl-1-[(3-(4-fluorophenyl)methyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.96	291
	(4aR)-4a-Ethyl-1-[(3-(4-fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.22	482




[0455] <Example 10>

(4aR)-1-[(2,3-Difluorophenyl)methyl]-4a-ethyl-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

By carrying out operations similar to those of First to Third Steps of Example 1, the compounds described in the following Table were synthesized from (R)-2-ethyl-pyrrolidine-2-carboxylic acid ethyl ester hydrochloride and 2,3-difluorobenzaldehyde.

[0456]

[Table 10]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(R)-1-[(2,3-Difluorobenzylidene)-amino]-2-ethyl-pyrrolidine-2-carboxylic acid ethyl ester	SQD-AA05	1.08	283
	(4aR)-1-[(2,3-Difluorophenyl)methyl]-4a-ethyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.89	295
	(4aR)-1-[(2,3-Difluorophenyl)methyl]-4a-ethyl-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.20	500




[0457] <Example 11>

(4aR)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-4a-propyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

By carrying out operations similar to those of First to Third Steps of Example 1, the compounds described in the following Table were synthesized from (R)-2-methyl-pyrrolidine-2-carboxylic acid ethyl ester hydrochloride and 3-fluorobenzaldehyde.

[0458]

[Table 11]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(R)-1-[(3-Fluorobenzylidene)-amino]-2-propyl-pyrrolidine-2-carboxylic acid ethyl ester	SQD-AA05	1.16	307
	(4aR)-1-[(3-Fluorophenyl)methyl]-4a-propyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.99	305
	(4aR)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-4a-propyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.22	496




[0459] <Example 12>

(4aR)-1-[(2,3-Difluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-4a-propyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

By carrying out operations similar to those of First to Third Steps of Example 1, the compounds described in the following Table were synthesized from (R)-2-propyl-pyrrolidine-2-carboxylic acid ethyl ester hydrochloride and 2,3-difluorobenzaldehyde.

[0460]

[Table 12]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(R)-1-[(2,3-Difluorobenzylidene)-amino]-2-propyl-pyrrolidine-2-carboxylic acid ethyl ester	SQD-AA05	1.25	325
	(4aR)-1-[(2,3-Difluorophenyl)methyl]-4a-propyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-2,4-dione	SQD-AA05	1.03	323
	(4aR)-1-[(2,3-Difluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-4a-propyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.23	514


[0461] <Example 13>

6-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

First Step

2,3-Difluoro-1-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzene

[0462] [Formula 36]



[0463] 2,3-Difluoro-1-[2-chloroethoxy]benzene (4.20 g, 22.0 mmol) was dissolved in N,N-dimethylformamide (32.9 mL), N-(2-methoxyethyl)-N-methylamine (4.35 g, 48.8 mmol), potassium iodide (6.08 g, 36.6 mmol), and tripotassium phosphate (7.77 g, 36.6 mmol) were added at 25°C, and the mixture was stirred under nitrogen atmosphere at 100°C for 2 hours.


After the reaction mixture was cooled to 25°C, water (47.0 mL) was added, and the resultant was extracted with isopropyl acetate (47.0 mL). 1 N hydrochloric acid (47.0 mL) was added to the organic layer for extraction, a 5 N aqueous sodium hydroxide solution (14.1 mL) was added to the obtained aqueous solution for basification, and the mixture was extracted with isopropyl acetate (47.0 mL). The organic layer was concentrated at reduced pressure to obtain the title compound (5.15 g, 95%).

[0464] ¹H-NMR (DMSO-D₆) δ: 7.16-7.10 (1H, m), 7.06-7.03 (1H, m), 6.99-6.96 (1H, m), 4.15 (2H, t, J = 5.7 Hz), 3.41 (2H, t, J = 6.0 Hz), 3.22 (3H, s), 2.78 (2H, t, J = 6.0 Hz), 2.59 (2H, t, J = 6.0 Hz), 2.28 (3H, s).

[0465] Second Step

2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzaldehyde

[0466] [Formula 37]



[0467] N,N-Diisopropylamine (5.92 mL) was dissolved in tetrahydrofuran (62.6 mL), and a solution of n-butyllithium in hexane (1.6 M, 26.2 mL, 41.9 mmol) was added dropwise at -20°C. After the mixture was stirred at -20°C for 30 minutes, it was cooled to -50°C, and a solution of 2,3-difluoro-1-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzene (5.15 g, 21.0 mmol) in tetrahydrofuran (21.0 mL) was added dropwise. After the mixture was stirred at -40°C for 2 hours, N,N-dimethylformamide (4.94 mL) was added, and the mixture was stirred for 30 minutes. After the reaction mixture was heated to -15°C, a solution of acetic acid (7.31 mL) in water (26.1 mL) and toluene (15.7 mL) were added for liquid-liquid extraction. The organic layer was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (ethyl acetate/hexane/triethylamine) to obtain the title compound (2.90 g, 50%).


[0468] ¹H-NMR (DMSO-D₆) δ: 10.04 (1H, s), 7.69-7.64 (1H, m), 7.27-7.24 (1H, m), 4.28

(2H, t, J = 5.7 Hz), 3.41 (2H, t, J = 6.0 Hz), 3.22 (3H, s), 2.81 (2H, t, J = 5.7 Hz), 2.60 (2H, t, J = 6.0 Hz), 2.28 (3H, s).

[0469] Third Step

Methyl 1-[[[E)-2,3-difluoro-4-[2-[2-(methoxyethyl(methyl)amino)ethoxy]phenyl]methylideneamino]-methylamino]cyclobutane-1-carboxylate

[0470] [Formula 38]



[0471] Methyl 1-(methylamino)cyclobutanecarboxylate paratoluenesulfonate (1.75 g, 5.54 mmol) (see Reference Example 86) was dissolved in acetic acid (1.60 mL, 27.7 mmol) and water (9.0 mL), a solution of sodium nitrite (0.45 g, 6.59 mmol) dissolved in water (1.00 mL) was added at 25°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. Ethyl acetate (3.50 mL) and sodium chloride (1.00 g) were added to the reaction mixture, and the organic layer after liquid-liquid extraction was washed with a 15% aqueous dipotassium hydrogenphosphate solution. Methanol (10.0 mL) was added to the organic layer after the washing, and 12 N hydrochloric acid (3.60 mL) was added at -10°C. Zinc dust (0.73 g, 11.1 mmol) was added to the reaction mixture at -30°C, and the mixture was stirred for 1 hour. A 28% aqueous ammonia (3.66 mL) was added to the reaction mixture, a solution of 2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzaldehyde (1.00 g, 3.66 mmol) in ethyl acetate (2.75 mL) was added at 0°C, and the mixture was stirred for 30 minutes. After the reaction mixture was concentrated, ethyl acetate and a 15% aqueous potassium dihydrogenphosphate solution were added for liquid-liquid extraction, and subsequently, the organic layer was washed with a 15% aqueous sodium hydrogen sulfite

solution and a 15% aqueous dipotassium hydrogenphosphate solution. The organic layer was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (ethyl acetate/methanol) to obtain the title compound (1.07 g, 70%).


[0472] LCMS: m/z 414[M+H]⁺

[0473] HPLC retention time: 0.65 minutes (SQD-TFA05)

[0474] Fourth Step

Methyl 1-[[[2,3-difluoro-4-{2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl-3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoyl]amino]-methylamino]cyclobutane-1-carboxylate

[0475] [Formula 39]



[0476] Methyl 1-[(E)-[2,3-difluoro-4-{2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclobutane-1-carboxylate (1.00 g, 2.42 mmol) was dissolved in ethyl acetate (8.1 mL), methanesulfonic acid (1.18 mL, 18.1 mmol) was added at 0°C, and the mixture was stirred under nitrogen atmosphere for 10 minutes. 5-Ethyl-2-methylpyridine borane (0.72 mL, 4.84 mmol) was added to the reaction mixture at 0°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. After methanol (0.81 mL) was added to the reaction mixture at 0°C, a 5 M aqueous sodium hydroxide solution (2.44 mL) was added, and the mixture was stirred at 0°C for 1 hour. A 25% aqueous tripotassium phosphate solution was added to the reaction mixture for liquid-liquid extraction. The organic layer was washed with a 15% aqueous sodium chloride solution, and the resultant was concentrated at reduced pressure to obtain

methyl 1-[[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylamino]-methylamino]cyclobutane-1-carboxylate as a crude product.

[0477] The obtained crude product and 3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoate (Reference Example 82) (1.05 g, 2.66 mmol) were dissolved in ethyl acetate (8.00 mL) and N,N-dimethylformamide (4.00 mL), pyridine (1.00 mL) and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (a 1.7 M ethyl acetate solution, 2.85 mL, 4.84 mmol) were added at -10°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. A 10% aqueous sodium chloride solution was added to the reaction mixture for liquid-liquid extraction. The organic layer was washed with a 15% aqueous dipotassium hydrogenphosphate solution, and the resultant was concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (ethyl acetate/methanol) to obtain the title compound (1.90 g, 97%).


[0478] LCMS: m/z 791 $[M+H]^+$

[0479] HPLC retention time: 0.81 minutes (SQD-TFA05)

[0480] Fifth Step

6-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[0481] [Formula 40]



[0482] Methyl 1-[[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl-[3-oxo-3[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoyl]amino]-methylamino]cyclobutane-1-

carboxylate (2.06 g, 2.60 mmol) was suspended in isopropanol (28.8 mL), potassium carbonate (718 mg, 5.20 mmol) was added, and the mixture was stirred under nitrogen atmosphere at 50°C for 8 hours. After the reaction mixture was concentrated, 1 M hydrochloric acid (2.60 mL, 2.60 mmol) was added, and the resultant was extracted with ethyl acetate. After the organic layer was washed with a 10% aqueous potassium dihydrogenphosphate solution and a 10% aqueous sodium chloride solution, the organic layer was concentrated. The resultant residue was purified by column chromatography on silica gel (ethyl acetate/methanol) to obtain the title compound (1.72 g, 87%).

[0483] Although tautomers of the title compound exist, isomers may be observed in some cases and not in other cases according to the type of the solvent for measurement. For example, ¹H-NMR and ¹³C-NMR of major tautomers (chloroform-D) are as follows.

[0484] LCMS: m/z 759[M+H]⁺

[0485] HPLC retention time: 0.91 minutes (SQD-TFA05)

[0486] ¹H-NMR (CDCl₃) δ: 16.57 (1H, s), 12.81 (1H, s), 9.61 (1H, s), 8.50 (1H, d, *J* = 8.7 Hz), 7.95 (1H, s), 7.90 (1H, d, *J* = 1.6 Hz), 7.80 (1H, dd, *J* = 8.7, 1.6 Hz), 7.01 (1H, ddd, *J* = 9.1, 7.2, 1.8 Hz), 6.72 (1H, ddd, *J* = 8.7, 7.2, 1.3 Hz), 5.08-5.03 (1H, m), 4.19-4.16 (1H, m), 4.19-4.16 (1H, m), 3.54-3.51 (2H, m), 3.37 (3H, s), 2.93 (2H, brs), 2.73 (2H, brs), 2.56 (1H, m), 2.43 (3H, brs), 2.41 (3H, s), 1.86 (1H, m), 1.84 (1H, m), 1.82 (1H, m), 1.71 (1H, m), 1.58 (1H, m).

[0487] ¹³C-NMR (CDCl₃) δ: 186.1 (qC), 169.8 (qC), 165.7 (qC), 162.8 (qC), 159.3 (CH), 156.6 (qC, q, *J*_{CF} = 36.3 Hz), 150.4 (qC, dd, *J*_{CF} = 248.4, 10.5 Hz), 147.9 (qC), 141.1 (qC, dd, *J*_{CF} = 248.0, 15.0 Hz), 138.8 (qC), 128.3 (CH, q, *J*_{CF} = 3.3 Hz), 127.8 (qC), 127.1 (CH), 126.9 (qC, q, *J*_{CF} = 33.6 Hz), 125.1 (CH), 124.7 (CH), 123.6 (qC, q, *J*_{CF} = 271.8 Hz), 120.5 (qC, q, *J*_{CF} = 275.4 Hz), 117.9 (qC, d, *J*_{CF} = 11.8 Hz), 115.9 (CH), 109.2 (CH, s), 92.3 (qC), 70.5 (CH₂), 68.1 (CH₂), 64.1 (qC), 58.9 (CH₃), 57.4 (CH₂), 56.3 (CH₂), 43.4 (CH₃), 41.5 (CH₂), 35.0 (CH₃), 32.4 (CH₂), 24.4 (CH₂), 13.4 (CH₂).

[0488] <Example 14>


7-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-

hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

First Step

Methyl 1-[(E)-2,3-[difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate

[0489] [Formula 41]



[0490] Methyl 1-(methylamino)cyclopentanecarboxylate paratoluenesulfonate (3.60 g, 10.9 mmol) (see Reference Example 88) was dissolved in acetic acid (3.10 mL, 54.2 mmol) and water (18.0 mL), a solution of sodium nitrite (0.91 g, 13.2 mmol) dissolved in water (2.00 mL) was added at 25°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. Ethyl acetate (7.00 mL) and sodium chloride (2.00 g) were added to the reaction mixture, and the organic layer after liquid-liquid extraction was washed with a 15% aqueous dipotassium hydrogenphosphate solution. Methanol (20.0 mL) was added to the organic layer after the washing, and 12 N hydrochloric acid (7.30 mL) was added at -10°C. Zinc dust (1.44 g, 22.0 mmol) was added to the reaction mixture at -30°C, and the mixture was stirred for 1 hour. 28% aqueous ammonia (7.30 mL) was added to the reaction mixture, a solution of 2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzaldehyde (2.00 g, 7.30 mmol) (Second Step of Example 13) in ethyl acetate (5.40 mL) was added at 0°C, and the mixture was stirred for 30 minutes. After the reaction mixture was concentrated, ethyl acetate and a 15% aqueous potassium dihydrogenphosphate solution were added for liquid-liquid extraction, and subsequently, the organic layer was washed with a 15% aqueous

sodium hydrogen sulfite solution and a 15% aqueous dipotassium hydrogenphosphate solution. The organic layer was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (ethyl acetate/methanol) to obtain the title compound (2.82 g, 90%).


[0491] LCMS: m/z 428[M+H]⁺

[0492] HPLC retention time: 0.61 minutes (SQD-FA05)

[0493] Second Step

Methyl 1-[[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl-3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoyl]amino]-methylamino]cyclopentane-1-carboxylate

[0494] [Formula 42]



[0495] Methyl 1-[[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate (1.00 g, 2.34 mmol) was dissolved in ethyl acetate (10.0 mL), methanesulfonic acid (1.14 mL, 17.6 mmol) was added at 0°C, and the mixture was stirred under nitrogen atmosphere for 10 minutes. 5-Ethyl-2-methylpyridine borane (0.70 mL, 4.70 mmol) was added to the reaction mixture at 0°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. After methanol (1.00 mL) was added to the reaction mixture at 0°C, a 5 M aqueous sodium hydroxide solution (3.00 mL) was added, and the mixture was stirred at 0°C for 1 hour. A 25% aqueous tripotassium phosphate solution was added to the reaction mixture for liquid-liquid extraction. The organic layer was washed with a 15% aqueous sodium chloride solution, and the resultant was concentrated at reduced pressure to obtain

methyl 1-[[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylamino]-methylamino]cyclopentane-1-carboxylate as a crude product.

[0496] The obtained crude product and 3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoate (1.00 g, 2.54 mmol) were dissolved in ethyl acetate (8.00 mL) and N,N-dimethylformamide (4.00 mL), and then pyridine (1.00 mL) and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (a 1.7 M ethyl acetate solution, 2.75 mL, 4.68 mmol) were added at -10°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. A 10% aqueous sodium chloride solution was added to the reaction mixture for liquid-liquid extraction. The organic layer was washed with a 15% aqueous dipotassium hydrogenphosphate solution, and the resultant was concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (ethyl acetate/methanol) to obtain the title compound (1.82 g, 97%).


[0497] LCMS: m/z 805[M+H]⁺

[0498] HPLC retention time: 0.74 minutes (SQD-FA05)

[0499] Third Step

7-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[0500] [Formula 43]



[0501] Methyl 1-[[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-[3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoyl]amino]-methylamino]cyclopentane-1-carboxylate (500 mg, 0.62 mmol) was suspended in isopropanol (7.00 mL), potassium

carbonate (175 mg, 1.27 mmol) was added, and the mixture was stirred under nitrogen atmosphere at 60°C for 8 hours. After the reaction mixture was concentrated, 1 M hydrochloric acid (1.24 mL, 1.24 mmol) was added, and the mixture was extracted with ethyl acetate. After the organic layer was washed with a 10% aqueous potassium dihydrogenphosphate solution and a 10% aqueous sodium chloride solution, the organic layer was concentrated. The resultant residue was purified by column chromatography on silica gel (ethyl acetate/methanol) to obtain the title compound (432 mg, 90%).

[0502] Although tautomers of the title compound exist, isomers may be observed in some cases and not in other cases according to the type of the solvent for measurement. For example, ¹H-NMR and ¹³C-NMR of major tautomers (chloroform-D) are as follows.

[0503] LCMS: m/z 773[M+H]⁺

[0504] HPLC retention time: 0.95 minutes (SQD-FA05)

[0505] ¹H-NMR (CDCl₃) δ: 16.55 (1H, s), 12.83 (1H, s), 9.62 (1H, s), 8.49 (1H, d, *J* = 8.7 Hz), 7.96 (1H, d, *J* = 1.2 Hz), 7.90 (1H, d, *J* = 1.6 Hz), 7.79 (1H, dd, *J* = 8.7, 2.0 Hz), 7.04 (1H, dd, *J* = 7.4, 7.4 Hz), 6.73 (1H, dd, *J* = 7.4, 7.4 Hz), 5.05 (1H, d, *J* = 14.2 Hz), 4.19-4.18 (1H, m), 4.19 (2H, brs), 3.55 (2H, brs), 3.37 (3H, s), 2.96 (2H, s), 2.76 (2H, s), 2.48 (3H, s), 2.45 (3H, s), 2.15 (1H, m), 1.74 (2H, m), 1.57 (1H, m), 1.50 (1H, m), 1.45 (2H, m), 1.30 (1H, m).

[0506] ¹³C-NMR (CDCl₃) δ: 187.8 (qC), 169.9 (qC), 165.7 (qC), 163.1 (qC), 159.3 (CH), 156.6 (qC, q, *J* = 36.3 Hz), 150.4 (qC, dd, *J*_{CF} = 248.6, 10.6 Hz), 147.9 (qC), 141.1 (qC, dd, *J*_{CF} = 248.3, 14.7 Hz), 138.8 (qC), 128.3 (CH), 127.8 (qC), 127.1 (CH), 126.9 (qC, q, *J*_{CF} = 33.4 Hz), 125.5 (CH), 124.7 (CH), 123.6 (qC, q, *J*_{CF} = 272.1 Hz), 120.5 (qC, q, *J*_{CF} = 275.4 Hz), 117.6 (qC, q, *J*_{CF} = 12.7 Hz), 116.0 (CH), 109.3 (CH), 93.1 (qC), 71.4 (qC), 70.4 (CH₂), 68.0 (CH₂), 58.9 (CH₃), 57.3 (CH₂), 56.3 (CH₂), 43.3 (CH₃), 41.6 (CH₂), 37.6 (CH₂), 36.8 (CH₃), 31.4 (CH₂), 24.8 (CH₂), 23.3 (CH₂).


[0507] <Example 15>

(4aR)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-3-carboxamide

First Step

(R)-Piperidine-2-carboxylic acid methyl ester hydrochloride

[0508] [Formula 44]






[0509] (R)-Piperidine-2-carboxylic acid (300 mg, 2.32 mmol) was dissolved in methanol (4.64 mL), thionyl chloride (0.508 mL, 6.97 mmol) was added dropwise at 0°C, and the mixture was stirred at 50°C for 18 hours. The reaction mixture was concentrated at reduced pressure, and toluene was added for azeotropic removal to obtain the title compound as a crude product.

[0510] Second Step

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from the obtained (R)-piperidine-2-carboxylic acid methyl ester hydrochloride and 3-fluoro-benzaldehyde.

[0511]

[Table 13]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(R)-1-[(3-Fluorobenzylidene)-amino]-piperidine-2-carboxylic acid methyl ester	SQD-AA05	1.02	265
	(4aR)-1-[(3-Fluorophenyl)methyl]-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.90	277
	(4aR)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.17	468




[0512] <Example 16>

(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-3-carboxamide

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from (S)-piperidine-2-carboxylic acid methyl ester hydrochloride and 3-fluoro-benzaldehyde.

[0513]

[Table 14]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(S)-1-[(3-Fluorobenzylidene)-amino]piperidine-2-carboxylic acid methyl ester	SQD-AA05	1.03	265
	(4aS)-1-[(3-Fluorophenyl)methyl]-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-2,4-dione	SQD-AA05	0.91	277
	(4aS)-1-[(3-Fluorophenyl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.16	468




[0514] <Example 17>

(4aS)-1-[(5,6-Difluoro-1H-indol-7-yl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-3-carboxamide

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from (S)-piperidine-2-carboxylic acid methyl ester hydrochloride and 5,6-difluoro-1H-indole-7-carbaldehyde.

[0515]

[Table 15]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(S)-1-[(5,6-Difluoro-1H-indol-7-yl)methylene]-amino]-piperidine-2-carboxylic acid methyl ester	SQD-AA05	1.03	323
	(4aS)-1-[(5,6-Difluoro-1H-indol-7-yl)methyl]-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-2,4-dione	SQD-AA05	1.00	334
	(4aS)-1-[(5,6-Difluoro-1H-indol-7-yl)methyl]-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydro-4aH-pyrido[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.16	525


[0516] <Example 18>

1-[(3-Fluorophenyl)methyl]-4-hydroxy-4a-methyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydropyrido[1,2-b]pyridazine-3-carboxamide

First Step

2-Methyl-piperidine-2-carboxylic acid methyl ester hydrochloride

[0517] [Formula 45]






[0518] In a similar manner to First Step of Example 4, the title compound was obtained from 2-methyl-piperidine-1,2-dicarboxylic acid 1-tert-butyl ester as a crude product.

[0519] Second Step

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from the obtained 2-methyl-piperidine-2-carboxylic acid methyl ester hydrochloride and 3-fluoro-benzaldehyde.

[0520] [Table 16]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	1-[(3-Fluoro-benzylidene)-amino]-2-methyl-piperidine-2-carboxylic acid methyl ester	SQD-AA05	1.07	279
	1-[(3-Fluorophenyl)methyl]-4a-methyl-5,6,7,8-tetrahydropyrido[1,2-b]pyridazine-2,4-dione	ZQ-01	2.53	291
	1-[(3-Fluorophenyl)methyl]-4-hydroxy-4a-methyl-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydropyrido[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.18	482


[0521] <Example 19>

1-[2,3-(Difluorophenyl)methyl]-4a-ethyl-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydropyrido[1,2-b]pyridazine-3-carboxamide

First Step

2-Ethyl-piperidine-2-carboxylic acid methyl ester hydrochloride

[0522] [Formula 46]



[0523] In a similar manner to First Step of Example 6, the title compound was obtained from 2-ethyl-piperidine-2-carboxylic acid hydrochloride as a crude product.

[0524] Second Step

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from the obtained 2-ethyl-piperidine-2-carboxylic

acid methyl ester hydrochloride and 2,3-difluoro-benzaldehyde.

[0525] [Table 17]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	1-([1-(2,3-Difluoro-phenyl)-met-(Z)-ylidene]-amino)-2-ethyl-piperidine-2-carboxylic acid methyl ester	SQD-AA05	1.16	311
	1-[(2,3-Difluorophenyl)methyl]-4a-ethyl-5,6,7,8-tetrahydropyrido[1,2-b]pyridazine-2,4-dione	SQD-AA05	1.00	323
	1-[(2,3-Difluorophenyl)methyl]-4a-ethyl-4-hydroxy-N-[5-methyl-2-(trifluoromethyl)furan-3-yl]-2-oxo-5,6,7,8-tetrahydropyrido[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.23	514

[0526] <Example 20>


(4aS,5S,8R)-1-(3-Fluorobenzyl)-4-hydroxy-N-(5-methyl-2-(trifluoromethyl)furan-3-yl)-2-oxo-2,4a,5,6,7,8-hexahydro-1H-5,8-methanopyrido[1,2-b]pyridazine-3-carboxamide

First Step

(1S,3S,6R)-2-Aza-bicyclo[2.2.1]heptane-3-carboxylic acid methyl ester

hydrochloride

[0527] [Formula 47]



[0528] (1R,3S,4S)-2-Aza-bicyclo[2.2.1]heptane-2,3-dicarboxylic acid 2-tert-butyl ester

(500 mg, 2.07 mmol) was dissolved in N,N-dimethylformamide (2.07 mL), cesium carbonate (1.01 g, 3.11 mmol) and iodomethane (0.155 mL, 2.49 mmol) were added, and the mixture




was stirred under nitrogen atmosphere at room temperature for 1 hour. 1 N hydrochloric acid was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with water, a saturated sodium bicarbonate solution, a 10% aqueous sodium thiosulfate solution, and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (1R,3S,4S)-2-aza-bicyclo[2.2.1]heptane-2,3-dicarboxylic acid 2-tert-butyl ester 3-methyl ester as a crude product. A solution of hydrogen chloride in 1,4-dioxane (4 M, 5.00 mL) was added to the obtained crude product, and the mixture was stirred under nitrogen atmosphere at room temperature for 1 hour. The reaction mixture was concentrated at reduced pressure, and toluene was added for azeotropic removal to obtain the title compound as a crude product.

[0529] Second Step

In a similar manner to First to Third Steps of Example 1, the compounds described in the following Table were synthesized from the obtained (1S,3S,6R)-2-aza-bicyclo[2.2.1]heptane-3-carboxylic acid methyl ester hydrochloride and 3-fluorobenzaldehyde.

[0530]


[Table 18]

Structure	Compound name	Analysis condition	HPLC Retention time (min)	m/z
	(1R,3S,4S)-2-[(3-Fluorobenzylidene)-amino]-2-aza-bicyclo[2.2.1]heptane-3-carboxylic acid methyl ester	SQD-AA05	1.02	277
	(4aS,5S,8R)-1-(3-Fluorobenzyl)tetrahydro-1H-5,8-methanopyrido[1,2-b]pyridazine-2,4(3H,4aH)-dione	SQD-AA05	0.90	289
	(4aS,5S,8R)-1-(3-Fluorobenzyl)-4-hydroxy-N-(5-methyl-2-(trifluoromethyl)furan-3-yl)-2-oxo-2,4a,5,6,7,8-hexahydro-1H-5,8-methanopyrido[1,2-b]pyridazine-3-carboxamide	SQD-AA05	1.17	480

[0531] <Reference Example 1-1>


(4aR)-1-[(2,3-Difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester

[0532] [Formula 48]



[0533] First Step


[0534] [Formula 49]



[0535] (R)-1-((2,3-(Difluorobenzylidene)amino)-2-methylpyrrolidine-2-carboxylic acid methyl ester (1.31 g, 4.63 mmol) was dissolved in methanol (4.6 mL) and acetic acid (4.6 mL), cyano sodium borohydride (1.19 g, 18.98 mmol) was added, and the mixture was stirred at room temperature for 3 hours. A saturated aqueous sodium bicarbonate solution was added, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous sodium sulfate, and filtered, and the solvent was distilled away at reduced pressure to obtain a crude product (1.34 g, 102%) of (R)-1-((2,3-(difluorobenzyl)amino)-2-methylpyrrolidine-2-carboxylic acid methyl ester.

[0536] Second Step

[0537] [Formula 50]



[0538] (R)-1-((2,3-(Difluorobenzyl)amino)-2-methylpyrrolidine-2-carboxylic acid methyl ester (3.24 g, 11.39 mmol) and 3-butyloxy-3-propionic acid (2.19 g, 13.67 mmol) were dissolved in dichloromethane (17.7 mL), and then bromotri(pyrrolidin-1-yl)phosphonium hexafluorophosphate (6.9 g, 14.81 mmol) and triethylamine (6.37 mL, 45.6 mmol) were added, and the mixture was stirred at room temperature for 30 minutes. Dichloromethane and 1 N hydrochloric acid were added to the reaction mixture, and the mixture was stirred at room temperature for 10 minutes. The organic layer was separated and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain (R)-1-(N-(2,3-difluorobenzyl)-3-isobutoxy-3-oxopropanamide)-2-methylpyrrolidine-2-carboxylic acid methyl ester (4.58 g, 94%).

[0539] LCMS: m/z 427[M+H]⁺

HPLC retention time: 1.33 minutes (analysis condition SMD-TFA05)

[0540] Third Step

(R)-1-(N-(2,3-Difluorobenzyl)-3-isobutoxy-3-oxopropanamide)-2-methylpyrrolidine-2-carboxylic acid methyl ester (30.4 g, 71.2 mmol) was dissolved in N,N-dimethylformamide (142 mL), cesium carbonate (69.6 g, 214 mmol) was added, and the mixture was stirred at 80°C for 1 hour. After the mixture was left to cool, it was concentrated at reduced pressure, 2 N hydrochloric acid was added, and the mixture was extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous magnesium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (0-80% ethyl acetate/hexane) to obtain the title compound (25.33 g, 90%).


[0541] LCMS: m/z 395[M+H]⁺

HPLC retention time: 1.05 minutes (analysis condition SQD-FA05)

[0542] <Reference Example 1-2>

Methyl (4aR)-4-hydroxy-4a-methyl-2-oxo-1-pentyl-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylate

[0543] [Formula 51]



[0544] Normal-pentyl aldehyde (0.959 g, 11.1 mmol) was used as a reagent, and operations similar to those of First Step of Reference Example 1-1 were carried out to obtain (R)-1-(pentylamino)-2-methylpyrrolidine-2-carboxylic acid methyl ester (1.73 g) as a crude product. The obtained hydrazine derivative was dissolved in tetrahydrofuran (14 mL), tripotassium phosphate (6.41 g, 30.2 mmol) and methyl 3-chloro-3-oxopropanoate (1.62 mL, 15.1 mmol) were added, and the mixture was stirred at room temperature overnight. Ethyl acetate (100 mL) was added to the reaction mixture, and after the organic layer was washed with 1 N hydrochloric acid, water, and a brine, it was dried over magnesium sulfate. After the resultant was filtered, the filtrate was concentrated at reduced pressure, and the residue was

purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain (R)-1-(3-methoxy-3-oxo-N-pentylpropanamide)-2-methylpyrrolidine-2-carboxylic acid methyl ester (1.84 g, 84%) as yellow oil. The obtained amide body (1.84 g, 5.59 mmol) was dissolved in N,N-dimethylformamide (25 mL), cesium carbonate (5.47 g, 16.8 mmol) was added, and the mixture was stirred at 80°C for 2 hours. After the reaction mixture was cooled to room temperature, ethyl acetate (50 mL) was added, and after the organic layer was washed with 1 N hydrochloric acid, water, and a brine, it was dried over magnesium sulfate. After the resultant was filtered, the filtrate was concentrated at reduced pressure to obtain the title compound (1.84 g).


[0545] LCMS: m/z 297[M+H]⁺

HPLC retention time: 1.22 minutes (analysis condition SMD-TFA05)

[0546] <Reference Example 2>


(R)-2,5,5-Trimethylpyrrolidine-2-carboxylic methyl ester

[0547] [Formula 52]



[0548] First Step

[0549] [Formula 53]




[0550] 4-Chlorobenzaldehyde (0.736 g, 5.23 mmol) was added to a suspension of 2-amino propanoate tert-butyl ester (0.80 g, 5.51 mmol) and magnesium sulfate (0.66 g, 5.51 mmol) in dichloromethane (7.3 mL), and the mixture was stirred at room temperature for 2 days. The reaction mixture was filtered, and the obtained filtrate was concentrated at reduced pressure. The residue was dissolved in diethyl ether, washed with water and a brine, and the organic layer was dried over magnesium sulfate and filtered. The filtrate was concentrated at reduced pressure to obtain 2-((4-chlorobenzylidene)amino)propanoate tert-butyl ester

(1.29 g) as a crude product.

[0551] Second Step


[0552] [Formula 54]



[0553] 2-((4-Chlorobenzylidene)amino)propanoate tert-butyl ester (2.18 g, 12.2 mmol) obtained in First Step and (S)-4,4-dibutyl-2,6-bis(3,4,5-trifluorophenyl)-4,5-dihydro-3H-dinaphtho[7,6,1,2-cde]azepinium bromide (91 mg, 0.122 mmol) were dissolved in toluene (85 mL), and 1-bromo-3-methyl-2-butene (1.71 mL, 14.61 mmol) and cesium hydroxide (9.13 g, 60.9 mmol) were added at 0°C. After the reaction mixture was stirred at 0°C for 4 hours, water was added, and the mixture was extracted with dichloromethane. The organic layer was washed with a brine, dried over sodium sulfate, and filtered. The obtained filtrate was concentrated at reduced pressure to obtain (R)-2-((4-chlorobenzylidene)amino)-2,5-dimethylhex-4-enoic acid tert-butyl ester as a crude product.

[0554] Third Step

[0555] [Formula 55]




[0556] 1 N hydrochloric acid (97 mL, 97 mmol) was slowly added to a solution of (R)-2-((4-chlorobenzylidene)amino)-2,5-dimethylhex-4-enoic acid tert-butyl ester (4.09 g, 12.2 mmol) obtained in Second Step in diethyl ether (97 mL). The reaction mixture was intensely stirred at room temperature for 4 hours. The organic layer was separated, and the aqueous layer was washed with diethyl ether. After the aqueous layer was adjusted to pH 9, it was extracted with dichloromethane. The extracts were combined, dried, and filtered, and the filtrate was concentrated at reduced pressure to obtain (R)-2-amino-2,5-dimethylhex-4-

enoic acid tert-butyl ester as a crude product.

[0557] Fourth Step

[0558] [Formula 56]




[0559] (R)-2-Amino-2,5-dimethylhex-4-enoic acid tert-butyl ester (1.03 g, 4.83 mmol) obtained in Third Step was dissolved in dichloromethane (9.7 mL), p-toluenesulfonyl chloride (1.11 g, 5.79 mmol), triethylamine (0.81 mL, 5.79 mmol), and 4-dimethylaminopyridine (5.90 mg, 0.048 mmol) were added, and the mixture was stirred at room temperature overnight. 1 N hydrochloric acid was added to the reaction mixture, and the mixture was extracted with dichloromethane. The extracts were combined, washed with a brine, dried over sodium sulfate, and filtered, and subsequently, the filtrate was concentrated at reduced pressure. The residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain (R)-2,5-dimethyl-2-(4-methylphenylsulfonamide)hex-4-enoic acid tert-butyl ester (1.47 g, 83%) as yellow oil.

[0560] LCMS: m/z 368[M+H]⁺

HPLC retention time: 1.40 minutes (analysis condition SMD-TFA05)

[0561] Fifth Step

[0562] [Formula 57]




[0563] (R)-2,5-Dimethyl-2-(4-methylphenylsulfonamide)hex-4-enoic acid tert-butyl ester (1.47 g, 4.00 mmol) obtained in Fourth Step was dissolved in chloroform (16.0 mL), and the solution was cooled to 0°C. Trifluoromethanesulfonic acid (0.14 mL, 1.60 mmol) was added at 0°C, and the reaction mixture was stirred at 0°C for 1 hour and at room temperature

for 2 hours. A saturated sodium bicarbonate solution was added to the reaction mixture, and the mixture was extracted with dichloromethane. The extracts were combined, washed with a brine, and dried over sodium sulfate. After the resultant was filtered, the filtrate was concentrated at reduced pressure to obtain (R)-2,5,5-trimethyl-1-tosylpyrrolidine-2-carboxylic acid as a crude product.

[0564] Sixth Step

[0565] [Formula 58]



[0566] Iodomethane (2.99 mL, 4.80 mmol) was added to a suspension of (R)-2,5,5-trimethyl-1-tosylpyrrolidine-2-carboxylic acid (1.25 g, 4.00 mmol) obtained in Fifth Step and potassium carbonate (829 mg, 6.00 mmol) in N,N-dimethylformamide (16.0 mL). The reaction mixture was stirred at room temperature for 1 hour. 1 N hydrochloric acid was added, and the mixture was extracted with ethyl acetate. The extracts were combined, washed with a brine, and dried over sodium sulfate. After the resultant was filtered, the filtrate was concentrated at reduced pressure, and the residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain (R)-2,5,5-trimethyl-1-tosylpyrrolidine-2-carboxylic acid methyl ester (1.24 g, 3.81 mmol) as a pale yellow solid.

[0567] LCMS: m/z 326[M+H]⁺

HPLC retention time: 1.20 minutes (analysis condition SMD-TFA05)

[0568] Seventh Step

(R)-2,5,5-Trimethyl-1-tosylpyrrolidine-2-carboxylic acid methyl ester (1.24 g, 3.81 mmol) obtained in Sixth Step was dissolved in methanol (38.1 mL), and magnesium (1.85 g, 76 mmol) was added. The reaction mixture was stirred at room temperature for 28 hours. 1 N hydrochloric acid was added, and the aqueous layer was washed with dichloromethane. The water layer was adjusted to pH 9 and extracted with dichloromethane.


The extracts were combined, dried over sodium sulfate, and filtered. The filtrate was concentrated at reduced pressure to obtain the title compound (611 mg, 3.57 mmol) as colorless oil.

[0569] $^1\text{H-NMR}$ (CDCl_3) δ : 3.73 (3H, s), 2.36-2.30 (1H, m), 1.88-1.80 (1H, m), 1.69-1.53 (2H, m), 1.39 (3H, s), 1.18 (3H, s), 1.17 (3H, s).

[0570] <Reference Example 3>


2,3-Difluoro-4-(2-morpholinoethoxy)benzaldehyde

[0571] [Formula 59]



[0572] First Step

[0573] [Formula 60]



[0574] 4-(2-Chloroethyl)morpholine hydrochloride (4.72 g, 25.4 mmol), cesium carbonate (18.8 g, 57.7 mmol), and tetrabutylammonium iodide (0.511 g, 1.38 mmol) were added to a solution of 2,3-difluorophenol (3.00 g, 23.1 mmol) in acetonitrile (46.1 mL), and the mixture was stirred at 65°C for 4 hours. After cooling to room temperature, the reaction mixture was filtered through a celite pad, and the filtrate was concentrated at reduced pressure. The residue was dissolved in ethyl acetate, washed with water, a 2 N aqueous sodium hydroxide solution, water, and a brine, and the organic layer was dried over sodium sulfate. After the resultant was filtered, the filtrate was concentrated at reduced pressure, and the residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain 4-(2-(2,3-difluorophenoxy)ethyl)morpholine (5.60 g, 99%) as colorless oil.

[0575] LCMS: m/z 244 $[\text{M}+\text{H}]^+$

HPLC retention time: 0.68 minutes (analysis condition SMD-TFA05)

[0576] Second Step

A solution of 4-(2-(2,3-difluorophenoxy)ethyl)morpholine (1.00 g, 4.11 mmol) and N,N,N',N'-tetramethylethylenediamine (0.62 mL, 4.11 mmol) in tetrahydrofuran (13.7 mL) was cooled to -78°C, and n-butyllithium (2.14 mL, 5.34 mmol) was added. After the mixture was stirred at -78°C for 1 hour, N,N-dimethylformamide (0.35 mL, 4.52 mmol) was added at -78°C. The reaction mixture was slowly heated to -10°C, and the reaction was stopped with a saturated aqueous ammonium chloride solution. The aqueous layer was extracted with ethyl acetate, and the extracts were combined and washed with a brine, dried over sodium sulfate, and filtered. The filtrate was concentrated at reduced pressure, and the residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound (758 mg, 68%) as yellow oil.


[0577] LCMS: m/z 272[M+H]⁺

HPLC retention time: 0.62 minutes (analysis condition SMD-TFA05)

[0578] <Reference Example 4>


(4aR)-1-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester

[0579] [Formula 61]



[0580] First Step

[0581] [Formula 62]




[0582] 2,3-Difluoro-4-(2-morpholinoethoxy)benzaldehyde and methyl (R)-2-methylpyrrolidine-2-carboxylic acid hydrochloride were used, and operations similar to those of First to Third Steps of Example 1 were carried out to synthesize methyl (2R)-1-[(E)-[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methylideneamino]-2-methylpyrrolidine-2-carboxylate.

[0583] LCMS: m/z 412[M+H]⁺

HPLC retention time: 0.95 minutes (analysis condition SMD-TFA05)

[0584] Second Step

[0585] [Formula 63]




[0586] (R)-1-((2,3-Difluoro-4-(2-morpholinoethoxy)benzylidene)amino)-2-methylpyrrolidine-2-carboxylic acid methyl ester was used, and operations similar to those of First Step of Reference Example 1-1 were carried out to obtain (R)-1-((2,3-difluoro-4-(2-morpholinoethoxy)benzyl)amino)-2-methylpyrrolidine-2-carboxylic acid methyl ester as a crude product.

[0587] LCMS: m/z 414[M+H]⁺

HPLC retention time: 0.63 minutes (analysis condition SMD-TFA05)

[0588] Third Step

[0589] [Formula 64]



[0590] (R)-1-((2,3-Difluoro-4-(2-morpholinoethoxy)benzyl)amino)-2-methylpyrrolidine-2-

carboxylic acid methyl ester and 3-butyloxy-3-propionic acid were used, and operations similar to those of Reference Example 1-2 were carried out to synthesize (R)-1-(N-(2,3-difluoro-4-(2-morpholinoethoxy)benzyl)-3-isobutoxy-3-oxopropanamide)-2-methylpyrrolidine-2-carboxylic acid methyl ester.

[0591] LCMS: m/z 556[M+H]⁺

HPLC retention time: 1.03 minutes (analysis condition SMD-TFA05)

[0592] Fourth Step

(R)-1-(N-(2,3-Difluoro-4-(2-morpholinoethoxy)benzyl)-3-isobutoxy-3-oxopropanamide)-2-methylpyrrolidine-2-carboxylic acid methyl ester was used, and operations similar to those of Reference Example 1-2 were carried out to synthesize the title compound.







[0593] LCMS: m/z 524[M+H]⁺

HPLC retention time: 1.04 minutes (analysis condition SMD-TFA05)



[0594] The ester intermediates described in the following Table were synthesized using the corresponding aldehyde reagents and proline derivatives to carry out operations similar to those of Reference Example 1-1 or Reference Example 1-2.

[0595]

[Table 19-1]

Reference Example No.	Ester	LCMS analysis condition No.	Retention time (min)	Ester intermediate m/z
5		SQD-FA05	1.12	521 [M+H] ⁺
6		SQD-FA05	0.62	524 [M+H] ⁺
7		SMD-TFA50	0.47	335 [M+H] ⁺
8		SMD-TFA05	1.08	342 [M+H] ⁺
9		SMD-TFA50	0.88	413 [M+H] ⁺
10		SQD-FA05	0.82	362 [M+H] ⁺


[0596] [Table 19-2]

11		SQD-FA05	1.12	423 [M+H] ⁺
12		SMD-TFA05	1.14	377 [M+H] ⁺

[0597] <Reference Example 13>

4-(Trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]aniline

[0598] [Formula 65]



[0599] Toluene (900 mL), ethanol (226 mL), and water (450 mL) were added to 2-(tetramethyl-1,3,2-dioxaborolan-2-yl)-4-(trifluoromethyl)aniline (60.00 g, 209 mmol), 5-bromo-2-(trifluoromethyl)pyridine (50.80 g, 225 mmol), tetrakis(triphenylphosphine)palladium (9.000 g, 7.79 mmol), and potassium carbonate (129.0 g, 930 mmol) under nitrogen atmosphere, and the mixture was stirred at 110°C for 3 hours. The reaction mixture was concentrated at reduced pressure, water (1,000 mL) was added, and the mixture was extracted with ethyl acetate. The organic layer was washed with water, dried over sodium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (ethyl acetate/petroleum ether) to obtain the title compound (63 g, 98%).


[0600] LCMS: m/z 307[M+H]⁺

HPLC retention time: 0.99 minutes (analysis condition SQD-AA05)

[0601] <Reference Example 14>


2-Iodo-5-(2-methoxyethoxy)-4-(trifluoromethyl)aniline

[0602] [Formula 66]



[0603] First Step

[0604] [Formula 67]



[0605] 2-Methoxyethanol (0.097 mL, 1.24 mmol) was dissolved in N-methylpyrrolidone (2 mL) under nitrogen atmosphere, sodium hydride (60 wt. %, a mineral oil dispersion,


30 mg, 1.24 mmol) was added, and after the mixture was stirred at room temperature for 30 minutes, 4-bromo-2-fluoro-1-(trifluoromethyl)benzene (0.118 mL, 0.823 mmol) was added, and the mixture was stirred at room temperature for 7 days. A 0.37 M aqueous potassium dihydrogenphosphate solution was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous magnesium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (0-10% ethyl acetate/hexane) to obtain 4-bromo-2-(2-methoxyethoxy)-1-(trifluoromethyl)benzene (231.2 mg, 94%).

[0606] HPLC retention time: 0.63 minutes (analysis condition SQD-AA50)

[0607] TLC (silica gel plate) R_f value: 0.37 (10% ethyl acetate/hexane)

[0608] Second Step

[0609] [Formula 68]




[0610] 4-Bromo-2-(2-methoxyethoxy)-1-(trifluoromethyl)benzene (231.2 mg, 0.773 mmol), carbamic acid t-butyl (109 mg, 0.928 mmol), palladium acetate (II) (5.21 mg, 0.023 mmol), 2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl (33.2 mg, 0.07 mmol), and cesium carbonate (353 mg, 1.08 mmol) were dissolved in 1,4-dioxane (5.2 mL), and the mixture was stirred under nitrogen atmosphere at 100°C overnight. A 0.37 M aqueous potassium dihydrogenphosphate solution was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous magnesium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (0 - 20% ethyl acetate/hexane) to obtain (3-(2-methoxyethoxy)-4-(trifluoromethyl)phenyl)carbamic acid tert-butyl (192.5 mg, 74%).

[0611] LCMS: m/z 334[M-H]⁺

HPLC retention time: 1.02 minutes (analysis condition SQD-AA05)

[0612] Third Step

[0613] [Formula 69]



[0614] A solution of 4 N hydrogen chloride/1,4-dioxane (3.9 mL) was added to (3-(2-methoxyethoxy)-4-(trifluoromethyl)phenyl)carbamic acid tert-butyl ester (192.5 mg, 0.574 mmol), and the mixture was stirred at room temperature overnight. The reaction mixture was concentrated at reduced pressure, ether/hexane (1/3) was added to the resultant residue, the deposited solid was collected by filtration, and the resultant was dried under vacuum to obtain 3-(2-methoxyethoxy)-4-(trifluoromethyl)aniline hydrochloride (128.1 mg, 82%).

[0615] LCMS: m/z 236[M+H]⁺

HPLC retention time: 0.83 minutes (analysis condition SQD-AA05)

[0616] Fourth Step

3-(2-Methoxyethoxy)-4-(trifluoromethyl)aniline hydrochloride (55.9 mg, 0.206 mmol) was dissolved in acetic acid (1.03 mL), N-iodosuccinimide (50.9 mg, 0.226 mmol) was added, and the mixture was stirred at room temperature for 2 hours and 20 minutes. The reaction mixture was concentrated at reduced pressure, ethyl acetate was added, washed with 0.5 N sodium hydroxide and a brine, dried over anhydrous magnesium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (0 - 40% ethyl acetate/hexane) to obtain the title compound (69.2 mg, 93%).

[0617] LCMS: m/z 362[M+H]⁺


HPLC retention time: 1.19 minutes (analysis condition SMD-TFA05)

[0618] <Reference Example 15>

2',3'-Dimethoxy-4-(2-methoxyethoxy)-5-(trifluoromethyl)-[1,1'-biphenyl]-2-amine

[0619]

[Formula 70]



[0620] Toluene (2.7 mL) and ethanol (1.1 mL) were added to 2-iodo-5-(2-methoxyethoxy)-4-(trifluoromethyl)aniline (69.2 mg, 0.192 mmol), (2,3-dimethoxyphenyl)boronic acid (34.9 mg, 0.192 mmol), and 1,1'-bis(diphenylphosphino)ferrocene dichloropalladium (II) (7.9 mg, 0.0096 mmol), then a 2M aqueous potassium carbonate solution (0.38 mL) was added, and the mixture was stirred under nitrogen atmosphere at 100°C for 1 hour. A 0.37 M aqueous potassium dihydrogenphosphate solution was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous magnesium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (0 - 40% ethyl acetate/hexane), to obtain the title compound (41.1 mg, 58%).






















[0621] LCMS: m/z 372[M+H]⁺

HPLC retention time: 1.21 minutes (analysis condition SMD-TFA05)

[0622] The boronic acid derivatives and halides described in the following Table were used, and operations similar to those of Reference Example 13 or Reference Example 15 were carried out to synthesize aniline intermediates described in the following Table.

[0623]

[Table 20-1]

Reference Example No.	Aniline	Boronic acid derivative	Halide	Aniline intermediate m/z
16				298 [M+H] ⁺
17				281 [M+H] ⁺
18				359 [M+H] ⁺
19				281 [M+H] ⁺
20				326 [M+H] ⁺
21				312 [M+H] ⁺
22				372 [M+H] ⁺

[0624]

























[Table 20-2]

23				351 [M+H] ⁺
24				323 [M+H] ⁺
25				323 [M+H] ⁺
26				346 [M+H] ⁺
27				264 [M+H] ⁺
28				267 [M-H] ⁻
29				285 [M+H] ⁺
30				273 [M+H] ⁺

























[0625] [Table 20-3]

31				274 [M+H] ⁺
32				264 [M+H] ⁺
33				298 [M+H] ⁺
34				283 [M+H] ⁺
35				307 [M+H] ⁺
36				269 [M+H] ⁺
37				385 [M+H] ⁺
38				307 [M-H] ⁻

























[0626] [Table 20-4]

39				264 [M+H] ⁺
40				294 [M+H] ⁺
41				307 [M+H] ⁺
42				307 [M+H] ⁺
43				278 [M+H] ⁺
44				308 [M+H] ⁺
45				286 [M+H] ⁺
46				308 [M+H] ⁺

[0627] [Table 20-5]

47				289 [M+H] ⁺
48				287 [M+H] ⁺
49				264 [M+H] ⁺
50				308 [M+H] ⁺
51				308 [M+H] ⁺
52				288 [M+H] ⁺
53				274 [M+H] ⁺
54				304 [M+H] ⁺


[0628] [Table 20-6]

55				332 [M+H] ⁺
56				286 [M+H] ⁺
57				300 [M+H] ⁺
58				274 [M+H] ⁺
59				263 [M-H] ⁻
60				287 [M+H] ⁺
61				274 [M+H] ⁺
62				319 [M+H] ⁺

[0629] <Reference Example 63>

6-(2-Amino-5-(trifluoromethyl)phenyl)pyrimidine-4-carbonitrile

[0630] [Formula 71]



[0631] 2-(6-Chloropyrimidin-4-yl)-4-(trifluoromethyl)aniline (1.0 g, 3.65 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.246 g, 2.19 mmol) were dissolved in dimethylsulfoxide (3 mL), an aqueous solution (3 mL) of potassium cyanide (0.95 g, 14.6 mmol) was added at 0°C, and the mixture was stirred at room temperature overnight. A 0.37 M aqueous potassium dihydrogenphosphate solution was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous magnesium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by C18 reverse-phase column chromatography (40 - 100% acetonitrile/water) to obtain the title compound (365.6 mg, 38%).


[0632] LCMS: m/z 265[M+H]⁺

HPLC retention time: 1.16 minutes (analysis condition SMD-TFA05)

[0633] <Reference Example 64>

5-(2-Amino-5-(trifluoromethyl)phenyl)-4-methoxypyrimidine-2-carbonitrile

[0634] [Formula 72]



[0635] A mixture of 2-(2-chloro-4-methoxypyrimidin-5-yl)-4-(trifluoromethyl)aniline (89 mg, 0.293 mmol), zinc dicyanide (20.65 mg, 0.176 mmol), 1,1'-bis(diphenylphosphino)ferrocene (18.07 mg, 0.032 mmol), and tris(dibenzylideneacetone)dipalladium(0)-chloroform adduct (30.3 mg, 0.029 mmol) in N,N-

dimethylformamide (1.5 mL) was heated and stirred under nitrogen atmosphere at 80°C for 1 hour. Further, the reaction mixture was heated and stirred at 110°C for 9 hours. The reaction mixture was purified by C18 reverse-phase column chromatography (acetonitrile-water, 0.03% formic acid) to obtain 5-(2-amino-5-(trifluoromethyl)phenyl)-4-methoxypyrimidine-2-carbonitrile (57 mg, 66%).


[0636] LCMS: m/z 295[M+H]⁺

HPLC retention time: 0.82 minutes (analysis condition SQD-FA05)

[0637] <Reference Example 65>

6-(2-Amino-5-(trifluoromethyl)phenyl)-5-chloropyrimidine-4-carbonitrile

[0638] [Formula 73]



[0639] 2-(5,6-Dichloropyrimidin-4-yl)-4-(trifluoromethyl)aniline was used, and operations similar to those of Reference Example 63 were carried out to synthesize the title compound.


[0640] LCMS: m/z 299[M+H]⁺

HPLC retention time: 0.82 minutes (analysis condition SQD-FA05)

[0641] <Reference Example 66>

6-(2-Amino-5-(trifluoromethyl)phenyl)-5-methoxypyrimidine-4-carbonitrile

[0642] [Formula 74]



[0643] 2-(6-Chloro-5-methoxypyrimidin-4-yl)-4-(trifluoromethyl)aniline was used, and operations similar to those of Reference Example 63 were carried out to synthesize the title compound.


[0643] LCMS: m/z 295[M+H]⁺

HPLC retention time: 0.81 minutes (analysis condition SQD-FA05)

[0645] <Reference Example 67>

6-(2-Amino-5-(trifluoromethyl)phenyl)-5-methylpyrimidine-4-carbonitrile

[0646] [Formula 75]



[0647] 2-(6-Chloro-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)aniline was used, and operations similar to those of Reference Example 63 were carried out to synthesize the title compound.


[0648] LCMS: m/z 279[M+H]⁺

HPLC retention time: 0.78 minutes (analysis condition SQD-FA05)

[0649] <Reference Example 68>

5-(2-Amino-5-(trifluoromethyl)phenyl)pyrazine-2-carbonitrile

[0650] [Formula 76]



[0651] A mixture of 2-(5-bromopyrazin-2-yl)-4-(trifluoromethyl)aniline (27.6 mg, 0.087 mmol), zinc dicyanide (6.1 mg, 0.052 mmol), and tetrakis(triphenylphosphine)palladium(0) (5 mg, 0.0043 mmol) in N,N-dimethylformamide (0.4 mL) was heated and stirred under nitrogen atmosphere at 80°C for 18 hours. The reaction mixture was purified by C18 reverse-phase column chromatography (acetonitrile-water, 0.03% formic acid) to obtain 5-(2-amino-5-(trifluoromethyl)phenyl)pyrazine-2-carbonitrile.


[0652] LCMS: m/z 265[M+H]⁺

HPLC retention time: 0.81 minutes (analysis condition SQD-FA05)

[0653] <Reference Example 69>

2-(6-Methylpyrimidin-4-yl)-4-(trifluoromethyl)aniline

[0654] [Formula 77]



[0655] 2-(6-Chloropyrimidin-4-yl)-4-(trifluoromethyl)aniline (51.9 mg, 0.19 mmol), trimethyl boroxine (0.053 mL, 0.379 mmol), bis(triphenylphosphine)palladium (II) chloride (13.3 mg, 0.019 mmol), and tripotassium phosphate (161 mg, 0.759 mmol) were dissolved in 1,4-dioxane (0.95 mL), and the mixture was stirred at 90°C for 1 hour. A 0.37 M aqueous potassium dihydrogenphosphate solution was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous magnesium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (0 - 30% ethyl acetate/hexane) to obtain the title compound (27.4 mg, 57%).


[0656] LCMS: m/z 254[M+H]⁺

HPLC retention time: 0.93 minutes (analysis condition SQD-AA05)

[0657] <Reference Example 70>

2-(Trifluoromethyl)-4-(6-(trifluoromethyl)pyridin-3-yl)pyrimidin-5-amine


[0658] [Formula 78]



[0659] First Step

[0660]

[Formula 79]




[0661] 4-Iodo-2-(trifluoromethyl)pyrimidine-5-carboxylic acid ethyl ester and (6-(trifluoromethyl)pyridin-3-yl)boronic acid were used, and operations similar to those of Reference Example 13 were carried out to synthesize 2-(trifluoromethyl)-4-(6-(trifluoromethyl)pyridin-3-yl)pyrimidine-5-carboxylic acid ethyl ester.

[0662] LCMS: m/z 366[M+H]⁺

HPLC retention time: 1.26 minutes (analysis condition SMD-TFA05)

[0663] Second Step

[0664] [Formula 80]



[0665] 2-(Trifluoromethyl)-4-(6-(trifluoromethyl)pyridin-3-yl)pyrimidine-5-carboxylic acid ethyl ester obtained in First Step (36 mg, 0.10 mmol) was dissolved in a mixture solution of ethanol (1 mL) and water (1 mL), and sodium hydroxide (7.9 mg, 0.20 mmol) was added at room temperature. After the mixture was stirred at room temperature for 1 hour, the solvent was concentrated at reduced pressure, and 1 N hydrochloric acid (5 mL) was added to the residue. After the resultant was extracted with ethyl acetate, the organic layer was dried over magnesium sulfate and filtered. The filtrate was concentrated at reduced pressure to obtain 2-(trifluoromethyl)-4-(6-(trifluoromethyl)pyridin-3-yl)pyrimidine-5-carboxylic acid as a crude product (30 mg).

[0666] LCMS: m/z 338[M+H]⁺

HPLC retention time: 1.05 minutes (analysis condition SMD-TFA05)

[0667] Third Step

N,N-dimethylformamide (1.5 mL) was added to 2-(trifluoromethyl)-4-(6-(trifluoromethyl)pyridin-3-yl)pyrimidine-5-carboxylic acid obtained in Second Step (30 mg, 0.09 mmol), triethylamine (8.9 mg, 0.09 mmol), and diphenylphosphoryl azide (DPPA, 24.3 mg, 0.09 mmol), and the mixture was stirred at 60°C for 3 hours. Water (1.5 mL) was added to the reaction mixture, and the resultant was stirred at 80°C for 3 hours. The reaction mixture was cooled to room temperature, a 1 N aqueous sodium hydroxide solution was added, and after the resultant was extracted with ethyl acetate, it was dried over magnesium sulfate and filtered. The filtrate was concentrated at reduced pressure to obtain the title compound (26 mg) as a crude product.


[0668] LCMS: m/z 309[M+H]⁺

HPLC retention time: 1.08 minutes (analysis condition SMD-TFA05)

[0669] <Reference Example 71>

4-Bromo-2-(6-(trifluoromethyl)pyrimidin-4-yl)aniline

[0670] [Formula 81]



[0671] 2-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)aniline and 4-chloro-6-(trifluoromethyl)pyrimidine were used as a reagent and a starting material, respectively, and operations similar to those of Reference Example 13 were carried out to obtain 2-(6-(trifluoromethyl)pyrimidin-4-yl)aniline. 1-Bromopyrrolidine-2,5-dione (66.2 mg, 0.372 mmol) was added to a solution of 2-(6-(trifluoromethyl)pyrimidin-4-yl)aniline (89 mg, 0.372 mmol) in acetic acid (0.94 mL)/water (0.0067 mL) at room temperature. The reaction mixture was heated to 55°C and stirred for 3.5 hours. The reaction mixture was cooled to room temperature, and azeotropic removal with toluene was carried out three times. The residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound (60 mg, 50%) as a yellow solid.


[0672] LCMS: m/z 318[M+H]⁺

HPLC retention time: 1.24 minutes (analysis condition SMD-TFA05)

[0673] <Example 21>

(4aR)-1-[(2,3-Difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0674] [Formula 82]



[0675] (4aR)-1-[(2,3-Difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (20 mg, 0.0567 mmol) and 4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)aniline (Reference Example 50, 18 mg, 0.0596 mmol) were dissolved in toluene (0.2 mL), and the mixture was stirred at 90°C for 3.5 hours. After the mixture was left to cool, the resultant was concentrated at reduced pressure, and the resultant residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound (36 mg, 99%).

[0676] LCMS: m/z 628[M+H]⁺

HPLC retention time: 1.17 minutes (analysis condition SQD-AA05)

[0677] Major tautomer

¹H-NMR (CDCl₃) δ: 16.47 (1H, s), 12.91 (1H, s), 9.60 (1H, s), 8.47 (1H, d, *J* = 8.7 Hz), 7.95 (1H, s), 7.90 (1H, s), 7.79 (1H, d, *J* = 8.7 Hz), 7.13-7.10 (2H, m), 7.05-7.03 (1H, m), 5.08 (1H, d, *J* = 14.3 Hz), 4.47 (1H, d, *J* = 14.3 Hz), 3.33 (1H, ddd, *J* = 9.0, 9.0, 3.2 Hz), 2.76 (1H, ddd, *J* = 9.0, 8.7, 8.7 Hz), 2.56-2.54 (1H, m), 1.76-1.73 (1H, m), 1.69-1.64 (2H, m), 1.00 (3H, s).

[0678] Minor tautomer









¹H-NMR (CDCl₃) δ: 18.22 (1H, s), 12.98 (1H, s), 9.59 (1H, s), 8.33 (1H, d, *J* = 8.7 Hz), 7.92

(2H, d, $J = 9.6$ Hz), 7.79 (1H, d, $J = 8.7$ Hz), 7.14-7.06 (3H, m), 5.10 (1H, d, $J = 14.0$ Hz), 4.52 (1H, d, $J = 14.6$ Hz), 3.29-3.25 (1H, m), 2.78-2.75 (1H, m), 2.64-2.59 (1H, m), 1.73-1.66 (2H, m), 1.49-1.46 (1H, m), 0.84 (3H, s).

[0679] The appropriate aniline reagents of Reference Example 13 and 15 to 71 and the appropriate ester intermediates of Reference Examples 1-1 and 1-2 and Reference Examples 4 to 12 were used, and operations similar to those of Example 21 were carried out to synthesize the compounds described in the following Table. Note that in Examples 59 and 60, N,N-dimethylformamide was used as a solvent instead of toluene.










[0680]

[Table 21-1]










Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
22		QC-SMD-TFA05	1.26	728
23		QC-SMD-TFA05	1.65	619
24		QC-SMD-TFA05	1.63	599
25		QC-SMD-TFA05	1.61	615
26		QC-SMD-TFA05	1.75	571
27		QC-SMD-TFA05	1.63	519
28		SQD-AA05	1.18	510
29		SQD-AA05	1.16	528

[0681]










[Table 21-2]

30		SQD-AA05	1.17	560
31		SQD-AA05	1.22	526
32		QC-SMD-TFA05	1.64	528
33		SQD-AA05	1.17	588
34		SQD-AA05	1.17	610
35		SQD-AA05	1.19	492
36		SQD-AA05	1.16	510
37		QC-SMD-TFA05	1.74	570
38		QC-SMD-TFA05	1.72	526

[0682] [Table 21-3]

39		QC-SMD-TFA05	1.76	567
40		SQD-AA05	1.11	577
41		QC-SMD-TFA05	1.8	586
42		QC-SMD-TFA05	1.77	620
43		QC-SMD-TFA05	1.71	526
44		SQD-AA05	1.22	618
45		SQD-AA05	1.16	584
46		SQD-AA05	1.17	601
47		QC-SMD-TFA05	1.71	607










[0683] [Table 21-4]

48		QC-SMD-TFA05	1.61	601
49		QC-SMD-TFA05	1.64	624
50		QC-SMD-TFA05	1.68	679
51		QC-SMD-TFA05	1.59	607
52		QC-SMD-TFA05	1.25	713
53		QC-SMD-TFA05	1.29	748
54		QC-SMD-TFA05	1.62	607
55		QC-SMD-TFA05	1.29	744
56		QC-SMD-TFA05	1.64	615










[0684] [Table 21-5]

57		QC-SMD-TFA05	1.62	585
58		QC-SMD-TFA05	1.65	552
59		QC-SMD-TFA05	1.3	772
60		QC-SMD-TFA05	1.35	772
61		QC-SMD-TFA05	1.73	655
62		QC-SMD-TFA05	1.72	634
63		QC-SMD-TFA05	1.73	655
64		QC-SMD-TFA05	1.65	638
65		QC-SMD-TFA05	1.67	646










[0685] [Table 21-6]

66		QC-SMD-TFA05	1.73	632
67		QC-SMD-TFA05	1.61	594
68		SQD-AA05	1.14	618
69		QC-SMD-TFA05	1.67	608
70		QC-SMD-TFA05	1.70	628
71		QC-SMD-TFA05	1.69	671
72		QC-SMD-TFA05	1.65	620
73		QC-SMD-TFA05	1.72	638
74		QC-SMD-TFA05	1.71	594










[0686] [Table 21-7]

75		SQD-AA05	1.20	510
76		SQD-AA05	1.19	466
77		SQD-AA05	1.24	492
78		SQD-AA05	1.11	493
79		SQD-AA05	1.10	449
80		QC-SMD-TFA05	1.67	603
81		QC-SMD-TFA05	1.40	589
82		QC-SMD-TFA05	1.58	584
83		SQD-AA05	1.19	609








[0687] [Table 21-8]

84		SQD-AA05	1.19	627
85		QC-SMD-TFA05	1.70	705
86		QC-SMD-TFA05	1.66	593
87		QC-SMD-TFA05	1.60	666
88		SQD-AA05	1.16	540
89		SQD-AA05	1.19	482
90		SQD-AA05	1.17	627
91		SMD-AA05	1.71	627
92		QC-SMD-TFA05	1.66	692

[0688] [Table 21-9]

93		QC-SMD-TFA05	1.72	589
94		QC-SMD-TFA05	1.59	584
95		QC-SMD-TFA05	1.60	585
96		QC-SMD-TFA05	1.68	606
97		QC-SMD-TFA05	1.71	605
98		QC-SMD-TFA05	1.65	652
99		QC-SMD-TFA05	1.60	629
100		QC-SMD-TFA05	1.64	585
101		QC-SMD-TFA05	1.65	598

[0689] [Table 21-10]

102		QC-SMD-TFA05	1.29	722
103		QC-SMD-TFA05	1.30	734
104		QC-SMD-TFA05	1.28	735
105		QC-SMD-TFA05	1.64	606
106		QC-SMD-TFA05	1.25	757
107		SMD-TFA05	1.27	723
108		QC-SMD-TFA05	1.30	757

[0690] Although tautomers of for the compounds described in this table exist, for example,

¹H-NMR of Examples 22 and 52 is as follows.

<Example 22>

Major tautomer

¹H-NMR (CDCl₃) δ: 16.12 (1H, s), 11.93 (1H, s), 9.35 (1H, s), 8.54 (1H, d, *J* = 8.8 Hz), 7.77

(1H, d, $J = 8.9$ Hz), 7.59 (1H, s), 6.97 (1H, dd, $J = 7.8, 7.8$ Hz), 6.76 (1H, dd, $J = 7.8, 7.8$ Hz), 4.81 (1H, d, $J = 14.2$ Hz), 4.32 (1H, d, $J = 14.3$ Hz), 4.20 (2H, t, $J = 5.7$ Hz), 3.74 (4H, t, $J = 4.5$ Hz), 3.29 (1H, ddd, $J = 8.8, 8.7, 3.0$ Hz), 2.84 (2H, t, $J = 5.6$ Hz), 2.67 (1H, ddd, $J = 8.8, 8.8, 8.8$ Hz), 2.61-2.59 (4H, m), 2.55-2.53 (1H, m), 2.45 (3H, s), 1.74-1.71 (1H, m), 1.67-1.62 (2H, m), 0.99 (3H, s).

[0691] Minor tautomer

$^1\text{H-NMR}$ (CDCl_3) δ : 17.93 (1H, s), 12.03 (1H, s), 9.32 (1H, s), 8.36 (1H, d, $J = 8.9$ Hz), 7.77 (1H, d, $J = 8.9$ Hz), 7.61 (1H, s), 7.06 (1H, dd, $J = 18.5, 10.0$ Hz), 6.73 (1H, t, $J = 10.0$ Hz), 5.00 (1H, d, $J = 14.4$ Hz), 4.42 (1H, d, $J = 14.6$ Hz), 4.19 (2H, q, $J = 6.7$ Hz), 3.74 (4H, t, $J = 4.5$ Hz), 3.24-3.22 (1H, m), 2.84 (2H, t, $J = 5.6$ Hz), 2.70-2.66 (1H, m), 2.60 (4H, s), 2.55-2.49 (1H, m), 2.40 (3H, s), 1.67-1.62 (2H, m), 1.44-1.41 (1H, m), 0.79 (3H, s).

<Example 52>

Major tautomer

$^1\text{H-NMR}$ (CDCl_3) δ : 16.13 (1H, s), 11.97 (1H, s), 8.87 (1H, d, $J = 5.0$ Hz), 8.46 (1H, d, $J = 8.6$ Hz), 7.81 (1H, s), 7.74 (1H, d, $J = 8.6$ Hz), 7.60 (1H, d, $J = 4.9$ Hz), 7.53 (1H, s), 6.99 (1H, dd, $J = 7.8, 7.8$ Hz), 6.78 (1H, dd, $J = 7.8, 7.8$ Hz), 4.83 (1H, d, $J = 14.3$ Hz), 4.31 (1H, d, $J = 14.3$ Hz), 4.20 (2H, t, $J = 5.7$ Hz), 3.74 (4H, t, $J = 4.2$ Hz), 3.31 (1H, ddd, $J = 8.8, 8.8, 3.0$ Hz), 2.85-2.83 (2H, m), 2.69 (1H, ddd, $J = 8.8, 8.8, 8.8$ Hz), 2.61 (4H, brs), 2.56-2.51 (1H, m), 1.75-1.72 (1H, m), 1.67-1.65 (2H, m), 0.99 (3H, s).

[0692] Minor tautomer


$^1\text{H-NMR}$ (CDCl_3) δ : 17.91 (1H, s), 12.05 (1H, s), 8.82 (1H, d, $J = 4.9$ Hz), 8.24 (1H, d, $J = 8.5$ Hz), 7.77 (1H, s), 7.74 (1H, d, $J = 8.6$ Hz), 7.58 (1H, d, $J = 5.0$ Hz), 7.54 (1H, brs), 7.06 (1H, t, $J = 7.8$ Hz), 6.73 (1H, t, $J = 7.7$ Hz), 5.01 (1H, d, $J = 14.4$ Hz), 4.43 (1H, d, $J = 14.5$ Hz), 4.19 (2H, dd, $J = 12.1, 6.2$ Hz), 3.74 (4H, t, $J = 4.2$ Hz), 3.27-3.24 (1H, m), 2.84 (2H, s), 2.74-2.72 (1H, m), 2.61 (3H, brs), 2.50-2.46 (1H, m), 1.78-1.63 (3H, m), 1.45-1.42 (1H, m), 0.81 (3H, s).

[0693] <Example 109>

(4aR)-N-[2-(5-Cyano-6-methylpyridin-3-yl)-4-(trifluoromethyl)phenyl]-1-[(2,3-

difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0694] [Formula 83]



[0695] (4aR)-N-[2-(6-Chloro-5-cyanopyridin-3-yl)-4-(trifluoromethyl)phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 68) was used as a starting material, and operations similar to those of Reference Example 69 were carried out to obtain the title compound.


[0696] LCMS: m/z 598[M+H]⁺

HPLC retention time: 1.63 minutes (analysis condition QC-SMD-TFA05)

[0697] <Example 110>


(4aR)-1-[(2,3-(Difluorophenyl)methyl)-N-[2-[1-(dimethylsulfamoyl)piperidin-4-yl]-4-(trifluoromethyl)phenyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0698] [Formula 84]



[0699] First Step

[0700] [Formula 85]



[0701] 2-Bromo-4-(trifluoromethyl)aniline and (1-(tert-butoxycarbonyl)-1,2,3,6-tetrahydropyridin-4-yl)boronic acid were used as a reagent and a starting material, respectively, and operations similar to those of Reference Example 13 were carried out to


obtain 4-(2-amino-5-(trifluoromethyl)phenyl)-5,6-dihydropyridine-1(2H)-carboxylic acid tert-butyl.

[0702] LCMS: m/z 243 $[M+H-C_4H_9OCO]^+$

HPLC retention time: 1.31 minutes (analysis condition SMD-TFA05)

[0703] Second Step

[0704] [Formula 86]




[0705] The aniline derivative obtained in First Step (0.73 g, 2.13 mmol) was dissolved in methanol (21.2 mL), palladium hydroxide-carbon (20 wt. %, 0.15 g, 0.21 mmol) was added, and the mixture was stirred under hydrogen atmosphere at room temperature overnight.

After the reaction mixture was filtered through a celite, pad the filtrate was concentrated at reduced pressure to obtain 4-(2-amino-5-(trifluoromethyl)phenyl)piperidine-1-carboxylic acid tert-butyl ester (0.70 g, 91%) as a crude product.

[0706] Third Step

[0707] [Formula 87]




[0708] The aniline derivative obtained in Second Step and the compound described in Reference Example 1-1 were used as a reagent and a starting material, respectively, and operations similar to those of Example 21 were carried out to obtain 4-[2-[(4aR)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carbonyl]amino]-5-(trifluoromethyl)phenyl]piperidine-1-carboxylic acid tert-butyl.

[0709] LCMS: m/z 565 $[M+H-C_4H_9OCO]^+$

HPLC retention time: 1.78 minutes (analysis condition SMD-TFA05)

[0710] Fourth Step

[0711] [Formula 88]



[0712] A 4 N hydrogen chloride/dioxane solution (5 mL) was added to the amide derivative obtained in Third Step (0.81 g, 1.23 mmol) at 0°C, and the mixture was stirred at room temperature for 1 hour. The reaction mixture was concentrated at reduced pressure to obtain (4aR)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-piperidin-4-yl-4-(trifluoromethyl)phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide hydrochloride (776 mg) as a yellow amorphous solid.

[0713] LCMS: m/z 565[M+H]⁺

HPLC retention time: 1.24 minutes (analysis condition SMD-TFA05)

[0714] Fifth Step

The hydrochloride obtained in Fourth Step (0.030 g, 0.050 mmol) was dissolved in N,N-dimethylformamide (0.25 mL), and then triethylamine (0.021 mL, 0.15 mmol) and dimethylsulfamoyl chloride (0.0054 mL, 0.050 mmol) were added at room temperature, and the mixture was stirred for 2 hours. The reaction mixture was purified directly by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound (24 mg, 71%) as a white amorphous solid.

[0715] LCMS: m/z 672[M+H]⁺


HPLC retention time: 1.66 minutes (analysis condition QC-SMD-TFA05)

[0716] <Example 111>

(4aR)-N-[2-[5-Cyano-6-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyridin-3-yl]-4-(trifluoromethyl)phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0717]

[Formula 89]



[0718] (4aR)-N-[2-(6-Chloro-5-cyanopyridin-3-yl)-4-(trifluoromethyl)phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 68) (20 mg, 0.032 mmol) was dissolved in 1,2-dichloroethane (0.32 mL) under nitrogen atmosphere, 2-oxa-6-azaspiro[3.3]heptane oxalate (24.5 mg, 0.129 mmol) and N,N-diisopropylethylamine (0.068 mL, 0.388 mmol) were added, and the mixture was stirred at room temperature overnight and at 60°C for 2 days. A 0.37 M aqueous potassium dihydrogenphosphate solution and dichloromethane were added to the reaction mixture, the mixture was intensely stirred, and subsequently, the water layer was separated by a phase separator, and the organic layer was concentrated at reduced pressure. The resultant residue was purified by C18 reverse-phase column chromatography (30 - 95% acetonitrile/water) to obtain the title compound (19.7 mg, 89%).


[0719] LCMS: m/z 681[M+H]⁺

HPLC retention time: 0.99 minutes (analysis condition SMD-TFA50)

[0720] <Example 112>

Methyl 2-[[5-[2-[[[(4aR)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carbonyl]amino]-5-(trifluoromethyl)phenyl]-3-cyanopyridin-2-yl]-methylamino]acetate

[0721] [Formula 90]



[0722] N-Methylglycine methyl ester was used as a reagent, and operations similar to those of Example 111 were carried out to synthesize the title compound.


[0723] LCMS: m/z 685[M+H]⁺

HPLC retention time: 1.65 minutes (analysis condition QC-SMD-TFA05)

[0724] <Example 113>

(4aR)-N-[2-(6-Cyano-5-methylsulfanylpyridin-3-yl)-4-(trifluoromethyl)phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0725] [Formula 91]



[0726] (4aR)-N-[2-(6-Cyano-5-nitropyridin-3-yl)-4-(trifluoromethyl)phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 99) (26.1 mg, 0.042 mmol) was dissolved in N,N-dimethylformamide (0.3 mL), sodium thiomethoxide (2.91 mg, 0.042 mmol) was added, and the mixture was stirred at room temperature for 3 days. A 0.37 M aqueous potassium dihydrogenphosphate solution was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous magnesium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by C18 reverse-phase column chromatography (30 - 90% acetonitrile (0.1% formic acid)/water (0.1% formic acid) to obtain the title compound (17.7 mg, 49%).

[0727] LCMS: m/z 630[M+H]⁺


HPLC retention time: 1.04 minutes (analysis condition SMD-TFA50)

[0728] <Example 114>

(4aR)-N-[2-(6-Cyano-5-methoxypyridin-3-yl)-4-(trifluoromethyl)phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0729]

[Formula 92]



[0730] Sodium methoxide was used as a reagent, and operations similar to those of Example 113 were carried out to synthesize the title compound.


[0731] LCMS: m/z 614[M+H]⁺

HPLC retention time: 1.62 minutes (analysis condition QC-SMD-TFA05)

[0732] <Reference Example 72>

7-[4-[(4aR)-4-Hydroxy-4a-methyl-3-(2-methylpropoxycarbonyl)-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]hept-6-ynoate

[0733] [Formula 93]



[0734] A suspension of (4aR)-1-[(2,3-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (Reference Example 5) (250 mg, 0.480 mmol), hept-6-ynoic acid (0.182 mL, 1.44 mmol), copper (I) iodide (18.3 mg, 0.0960 mmol), triethylamine (0.335 mL, 2.40 mmol), and bis(triphenylphosphine)palladium (II) chloride (33.7 mg, 0.0480 mmol) in tetrahydrofuran (0.961 mL) was stirred under nitrogen atmosphere at room temperature for 2.5 hours. At the same time, the completely same reaction was run in another vessel on the same scale, and both resultants were combined and purified by C18 reverse-phase column chromatography to obtain the title compound (328 mg, 66%) as yellow oil.


[0735] LCMS: m/z 519[M+H]⁺

HPLC retention time: 1.00 minute (analysis condition SQD-FA05)

[0736] <Reference Example 73>

7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-(2-methylpropoxycarbonyl)-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]heptanoate

[0737] [Formula 94]



[0738] A solution of 7-[4-[[[(4aR)-4-hydroxy-4a-methyl-3-(2-methylpropoxycarbonyl)-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]hept-6-ynoic acid (Reference Example 72) (358 mg, 0.69 mmol) and palladium hydroxide-carbon (20 wt. %, about 50% wet with water, 145 mg) in 2,2,2-trifluoroethanol (7 mL) was stirred under hydrogen atmosphere at room temperature for 2 hours. Further, palladium hydroxide-carbon (20 wt. %, about 50% wet with water, 145 mg) was added, and the reaction mixture was stirred at room temperature for 1 hour. After the reaction mixture was filtered through a celite pad, the filtrate was concentrated to obtain the title compound (361 mg, 100%) as light brown oil.


[0739] LCMS: m/z 523[M+H]⁺

HPLC retention time: 1.05 minutes (analysis condition SQD-AA05)

[0740] <Example 115>

7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-[[2-(6-methylsulfonyl)pyridin-3-yl]-4-(trifluoromethyl)phenyl]carbonyl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]heptanoate

[0741] [Formula 95]



[0742] 7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-(2-methylpropoxycarbonyl)-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]heptanoate (Reference

Example 73) and 2-(6-(methylthio)pyridin-3-yl)-4-(trifluoromethyl)aniline were used, and operations similar to those of Example 21 were carried out to synthesize the title compound.


[0743] LCMS: m/z 733[M+H]⁺

HPLC retention time: 1.25 minutes (analysis condition SQD-FA05)

[0744] <Example 116>

7-[4-[[[(4aR)-3-[[4-Chloro-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]heptanoate

[0745] [Formula 96]



[0746] 7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-(2-methylpropoxycarbonyl)-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]heptanoate (Reference Example 73) and 4-chloro-2-(6-(trifluoromethyl)pyridin-3-yl)aniline were used, and operations similar to those of Example 21 were carried out to synthesize the title compound.


[0747] LCMS: m/z 721[M+H]⁺

HPLC retention time: 1.22 minutes (analysis condition SQD-FA05)

[0748] <Example 117>

7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-[[2-(6-methylsulfanylpyridin-3-yl)-4-(trifluoromethyl)phenyl]carbamoyl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]hept-6-ynoate

[0749] [Formula 97]



[0750] 7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-(2-methylpropoxycarbonyl)-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl)methyl]-2,3-difluorophenyl]hept-6-ynoic acid (Reference Example 72) was used as a starting material, 2-(6-(methylthio)pyridin-3-yl)-4-(trifluoromethyl)aniline was used as a reagent, and operations similar to those of Example 21 were carried out to synthesize the title compound.


[0751] LCMS: m/z 729[M+H]⁺

HPLC retention time: 1.61 minutes (analysis condition SMD-TFA05)

[0752] <Example 118>

7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-[[2-(2-methylsulfanylpirimidin-5-yl)-4-(trifluoromethyl)phenyl]carbamoyl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl)methyl]-2,3-difluorophenyl]hept-6-ynoate

[0753] [Formula 98]



[0754] 7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-(2-methylpropoxycarbonyl)-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl)methyl]-2,3-difluorophenyl]hept-6-ynoic acid (Reference Example 72) and 2-(2-(methylthio)pyrimidin-5-yl)-4-(trifluoromethyl)aniline were used, and operations similar to those of Example 21 were carried out to synthesize the title compound.


[0755] LCMS: m/z 730[M+H]⁺

HPLC retention time: 1.58 minutes (analysis condition SMD-TFA05)

[0756] <Example 119>

7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[5-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl)methyl]-2,3-difluorophenyl]hept-6-ynoate

[0757] [Formula 99]



[0758] 7-[4-[[[(4aR)-4-Hydroxy-4a-methyl-3-(2-methylpropoxycarbonyl)-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]hept-6-ynoic acid (Reference Example 72) and 4-(trifluoromethyl)-2-(5-(trifluoromethyl)pyridin-3-yl)aniline were used, and operations similar to those of Example 21 were carried out to synthesize the title compound.


[0759] LCMS: m/z 751[M+H]⁺

HPLC retention time: 1.60 minutes (analysis condition SMD-TFA05)

[0760] <Reference Example 74>


(4aR)-1-[[4-[4-[[[(4S)-2,2-Dimethyl-1,3-dioxolan-4-yl]methoxy]butoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester

[0761] [Formula 100]



[0762] First Step

[0763] [Formula 101]




[0764] 4-(Benzyloxy)-2,3-difluorobenzaldehyde and (R)-2-methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride were used as a reagent and a starting material, respectively, and operations similar to those of First Step of Example 1 and Reference Example 1-1 were

carried out to synthesize (4aR)-1-[(2,3-difluoro-4-phenylmethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester.

[0765] Second Step

[0766] [Formula 102]




[0767] Palladium-carbon (10 wt. %, 120 mg) was added to a solution of (4aR)-1-[(2,3-difluoro-4-phenylmethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (1.56 g, 3.12 mmol) in ethyl acetate (100 mL), and the mixture was stirred under hydrogen atmosphere at room temperature for 30 minutes. The reaction mixture was filtered through celite, the filtrate was concentrated at reduced pressure, and the residue was purified by column chromatography on silica gel (dichloromethane/methanol) to obtain (4aR)-1-[(2,3-difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (1.02 g, 80%).

[0768] LCMS: m/z 411[M+H]⁺

HPLC retention time: 1.24 minutes (analysis condition SMD-TFA05)

[0769] Third Step

[0770] [Formula 103]



[0771] A 60% sodium hydroxide solution (2 mL, 30.3 mmol) was added to a mixture of (S)-(2,2-dimethyl-1,3-dioxolan-4-yl)methanol (2.00 g, 15.1 mmol) and 1,4-dibromobutane (5.42 mL, 45.4 mmol). Tetrabutylammonium hydrogensulfate (257 mg, 0.757 mmol) was added to this mixture, and the resultant was intensely stirred at room temperature overnight. After water was added, the layer was extracted with hexane/diethyl ether (1:4) four times,

and the organic layers were combined and dried over magnesium sulfate. After the resultant was filtered, the filtrate was concentrated at reduced pressure, and the residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain (S)-4-((4-bromobutoxy)methyl)-2,2-dimethyl-1,3-dioxolane (2.54 g, 63%) as colorless oil.

[0772] Fourth Step

Cesium carbonate (1.20 mg, 3.65 mmol) was added to a solution of (4aR)-1-[(2,3-difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (500 mg, 1.22 mmol) and (S)-4-((4-bromobutoxy)methyl)-2,2-dimethyl-1,3-dioxolane (374 mg, 1.40 mmol) in acetonitrile (15.2 mL), and the mixture was stirred at 70°C for 1 hour. After the mixture was cooled to room temperature, a 0.37 M aqueous potassium dihydrogenphosphate solution was added, and the aqueous layer was extracted with ethyl acetate. The organic layers were combined, washed with a brine, and dried over magnesium sulfate. After the resultant was filtered, the filtrate was concentrated at reduced pressure, and the residue was purified by column chromatography on silica gel (dichloromethane/methanol) to obtain the title compound (414 mg, 57%).


[0773] LCMS: m/z 597[M+H]⁺

HPLC retention time: 1.52 minutes (analysis condition SMD-TFA05)

[0774] <Example 120>

(4aR)-1-[[4-[4-[(2R)-2,3-Dihydroxypropoxy]butoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-N-[2-(6-methylsulfonylpyridin-3-yl)-4-(trifluoromethyl)phenyl]-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0775] [Formula 104]



[0776] (4aR)-1-[[4-[4-[(4S)-2,2-Dimethyl-1,3-dioxolan-4-yl]methoxy]butoxy]-2,3-

difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (Reference Example 74) (10.6 mg, 0.018 mmol) and 2-(6-(methylthio)pyridin-3-yl)-4-(trifluoromethyl)aniline (5.1 mg, 0.018 mmol) were dissolved in toluene (0.5 mL), and the mixture was stirred at 100°C for 1 hour. The reaction mixture was concentrated at reduced pressure, the resultant residue was dissolved in methanol (0.5 mL), p-toluenesulfonic acid monohydrate (3.4 mg, 0.018 mmol) was added, and the mixture was stirred at room temperature for 1 hour and a half. Water was added to the reaction mixture, and the resultant white turbid solution was purified by C18 reverse-phase column chromatography (30 - 90% acetonitrile (0.1% trifluoroacetic acid)/water (0.1% trifluoroacetic acid)) to obtain the title compound (6.3 mg, 46%) as a light brown amorphous solid.


[0777] LCMS: m/z 767[M+H]⁺

HPLC retention time: 1.51 minutes (analysis condition QC-SMD-TFA05)

[0778] <Example 121>

(4aR)-N-[4-Chloro-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-1-[[4-[4-[(2R)-2,3-dihydroxypropoxy]butoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0779] [Formula 105]



[0780] (4aR)-1-[[4-[4-[(4S)-2,2-Dimethyl-1,3-dioxolan-4-yl]methoxy]butoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (Reference Example 74) and 4-chloro-2-(6-(trifluoromethyl)pyridin-3-yl)aniline were used, and operations similar to those of Example 120 were carried out to synthesize the title compound.


[0781] LCMS: m/z 755[M+H]⁺

HPLC retention time: 1.50 minutes (analysis condition QC-SMD-TFA05)

[0782] <Example 122>

(4aR)-N-[4-Bromo-2-(6-chloropyridin-3-yl)phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0783] [Formula 106]



[0784] Dimethoxyethane (0.2 mL), ethylene glycol (0.2 mL), and water (0.1 mL) were added to (4aR)-N-(4-bromo-2-iodophenyl)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 331) (14.8 mg, 0.02 mmol), potassium carbonate (9.7 mg, 0.07 mmol), tetrakis(triphenylphosphine)palladium (2.3 mg, 0.002 mmol), and (6-chloropyridin-3-yl)boronic acid (3.1 mg, 0.02 mmol), and the mixture was stirred at 100 °C for 2 hours. The insolubles were filtered, and the filtrate was purified directly by HPLC (YMC-Actus ODS-A 20 × 100 mm 0.005 mm, 0.1% formic acid acetonitrile/0.1% formic acid water) to obtain the title compound (4.8 mg, 40%) as a white amorphous solid.


[0785] LCMS: m/z 604[M+H]⁺

HPLC retention time: 0.89 minutes (analysis condition SQD-FA05)

[0786] <Reference Example 75>

((4S,5S)-5-((Hept-6-yn-1-yloxy)methyl)-2,2-dimethyl-1,3-dioxolan-4-yl)methanol

[0787] [Formula 107]




[0788] Sodium hydride (60 wt. %, a mineral oil dispersion, 36 mg, 0.824 mmol) was added to a solution of ((4S,5S)-2,2-dimethyl-1,3-dioxolane-4,5-diyl)dimethanol (133 mg,

0.824 mmol) in N,N-dimethylformamide (1.1 mL) at room temperature. After the reaction mixture was stirred at room temperature for 30 minutes, hept-6-yn-1-yl methanesulfonate (105 mg, 0.550 mmol) was added, and the mixture was stirred at room temperature overnight. Water was slowly added to the reaction mixture, and the resultant was extracted with ethyl acetate. The organic layers were combined and concentrated at reduced pressure, and the residue was purified by column chromatography on silica gel to obtain ((4S,5S)-5-((hept-6-yn-1-yloxy)methyl)-2,2-dimethyl-1,3-dioxolan-4-yl)methanol (99.6 mg, 71%).

[0789] <Reference Example 76>

(R)-4-((3-Bromopropoxy)methyl)-2,2-dimethyl-1,3-dioxolane

[0790] [Formula 108]



[0791] (R)-(-)-2,2-Dimethyl-1,3-dioxolane-4-methanol (615 mg, 4.66 mmol) 6-bromo-1-hexyne (500 mg, 3.1 mmol) was dissolved in N,N-dimethylformamide (10 mL), sodium hydride (60 wt. %, a mineral oil dispersion, 186 mg, 4.66 mmol) was added, and the mixture was stirred at room temperature for 12 hours. The reaction mixture was added to water and extracted with diethyl ether. After the organic layer was washed with a brine and dried over sodium sulfate, the resultant was concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel to obtain (R)-4-((hex-5-yn-1-yloxy)methyl)-2,2-dimethyl-1,3-dioxolane (390 mg, 59%) as clear liquid.


[0792] ¹H-NMR (CDCl₃) δ: 4.24 (1H, dt, *J* = 12.0, 6.4 Hz), 4.04 (1H, dd, *J* = 8.0, 6.4 Hz), 3.71 (1H, dd, *J* = 8.3, 6.4 Hz), 3.54-3.44 (3H, m), 3.41 (1H, dd, *J* = 11.0, 5.0 Hz), 2.20 (4H, td, *J* = 7.0, 3.0 Hz), 1.92 (4H, t, *J* = 3.0 Hz), 1.72-1.64 (2H, m), 1.62-1.53 (2H, m), 1.40 (3H, s), 1.34 (3H, s).

[0793] <Reference Example 77>

4-(Prop-2-yn-1-yl)morpholin-3-one

[0794]

[Formula 109]




[0795] Sodium hydride (60 wt. %, a mineral oil dispersion, 187 mg, 4.68 mmol) was added to a solution of morpholin-3-one (430 mg, 4.25 mmol) in tetrahydrofuran (5 mL) at 0°C, and the mixture was stirred at room temperature for 10 minutes, and 3-bromoprop-1-yne (607 mg, 5.1 mmol) was added to the reaction mixture, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was diluted with ethyl acetate (50 mL), and after the reaction mixture was washed serially with a saturated aqueous ammonium chloride solution, water, and a brine and dried over magnesium sulfate, the resultant was concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel to obtain the title compound (312 mg, 53%) as clear oil.

[0796] ¹H-NMR (CDCl₃) δ: 4.29 (2H, d, *J* = 2.4 Hz), 4.19 (2H, s), 3.93 (2H, t, *J* = 5.2 Hz), 3.51 (2H, t, *J* = 5.2 Hz), 2.25 (1H, m).

[0797] <Example 123>

(4aR)-1-[[2,3-Difluoro-4-[3-(4-hydroxybutoxy)prop-1-ynyl]phenyl]methyl]-N-[2-(2,3-dimethoxyphenyl)-4-(trifluoromethyl)phenyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0798] [Formula 110]



[0799] 4-(Prop-2-yn-1-yloxy)butan-1-ol (15.5 mg, 0.121 mmol) and triethylamine (0.020 mL, 0.141 mmol) were added to a solution of (4aR)-1-[2,3-(difluoro-4-iodophenyl)methyl]-N-[2-(2,3-dimethoxyphenyl)-4-(trifluoromethyl)phenyl]-4-hydroxy-4a-

methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 321) (30 mg, 0.040 mmol), bis(triphenylphosphine)palladium (II) chloride (2.83 mg, 0.0404 mmol), and copper (I) iodide (1.54 mg, 0.0081 mmol) in N,N-dimethylformamide (0.40 mL) under nitrogen atmosphere at 0°C. The reaction mixture was stirred at room temperature for 14 hours. Formic acid was added to the reaction mixture, and the mixture was purified directly by C18 reverse-phase column chromatography to obtain the title compound (15 mg, 50%) as a white amorphous solid.


[0800] LCMS: m/z 744[M+H]⁺

HPLC retention time: 1.63 minutes (analysis condition QC-SMD-TFA05)

[0801] <Example 124>

(4aR)-1-[[2,6-Difluoro-4-[7-[(2S,3S)-2,3,4-trihydroxybutoxy]hept-1-ynyl]phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0802] [Formula 111]



[0803] ((4S,5S)-2,2-Dimethyl-1,3-dioxolane-4,5-diyl)dimethanol (23.9 mg, 0.093 mmol) and triethylamine (0.019 mL, 0.140 mmol) were added to a solution of (4aR)-1-[(2,6-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 322) (35 mg, 0.047 mmol), bis(triphenylphosphine)palladium (II) chloride (3.27 mg, 0.0465 mmol), and copper (I) iodide (1.77 mg, 0.0093 mmol), in N,N-dimethylformamide (0.47 mL) under nitrogen atmosphere at 0°C. The reaction mixture was stirred at room temperature overnight. Trifluoroacetic acid/water (1:1, 1.4 mL) and dimethyl sulfoxide (0.7 mL) were added to the reaction mixture, and the mixture was stirred

at room temperature for 1 hour. The reaction mixture was purified directly by C18 reverse-phase column chromatography to obtain the title compound (32 mg, 82%) as an orange amorphous solid.


[0804] LCMS: m/z 841[M+H]⁺

HPLC retention time: 0.90 minutes (analysis condition SMD-TFA50)

[0805] <Example 125>

(4aR)-1-[[2,6-Difluoro-4-[7-[(2S,3S)-2,3,4-trihydroxybutoxy]heptyl]phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0806] [Formula 112]



[0807] A suspension of (4aR)-1-[[2,6-difluoro-4-[7-[(2S,3S)-2,3,4-trihydroxybutoxy]hept-1-ynyl]phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)-pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 124) (20 mg, 0.024 mmol) and palladium hydroxide-carbon (20 wt. %, 3.34 mg, 0.00476 mmol) in ethyl acetate was stirred under hydrogen atmosphere at room temperature for 10 hours. The reaction mixture was filtered through a celite pad, and the filtrate was concentrated at reduced pressure. The residue was purified by C18 reverse-phase column chromatography to obtain the title compound (11.5 mg, 57%) as a colorless amorphous solid.

[0808] LCMS: m/z 845[M+H]⁺


HPLC retention time: 1.60 minutes (analysis condition QC-SMD-TFA05)

[0809] <Example 126>

(4aR)-1-[(4-Ethynyl-2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-

b]pyridazine-3-carboxamide

[0810] [Formula 113]



[0811] Triisopropylsilyl acetylene was used as a reagent, and operations similar to those of First Step of Example 123 were carried out to obtain (4aR)-1-[[2,3-difluoro-4-[2-tri(propan-2-yl)silylethynyl]phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide as a crude product. The obtained crude product was dissolved in tetrahydrofuran (0.15 mL), a 1 M tetrahydrofuran solution (0.12 mL) of tetrabutylammonium fluoride was added, and the mixture was stirred at room temperature for 1 hour. Trifluoroacetic acid (0.5 mL), water (0.5 mL), and dimethylsulfoxide (1 mL) were added to the reaction mixture, and the mixture was purified by C18 reverse-phase column chromatography to obtain the title compound (22 mg, 84%) as a white amorphous solid.









[0812] LCMS: m/z 652[M+H]⁺

HPLC retention time: 0.87 minutes (analysis condition SQD-FA50).

[0813] The appropriate iodide derivatives of Examples 319-323 and appropriate terminal alkyne reagents were used, and operations similar to those of Example 123 were carried out to synthesize the compounds described in the following Table.

[0814]

[Table 22-1]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
127		SQD-AA05	1.18	730
128		QC-SMD-TFA05	1.60	753
129		QC-SMD-TFA05	1.62	738
130		QC-SMD-TFA05	1.69	735
131		SQD-AA05	1.14	631
132		QC-SMD-TFA05	1.33	750
133		QC-SMD-TFA05	1.56	648
134		QC-SMD-TFA05	1.60	686

[0815]

[Table 22-2]

135		QC-SMD-TFA05	1.65	714
136		QC-SMD-TFA05	1.57	804
137		QC-SMD-TFA05	1.58	716
138		QC-SMD-TFA05	1.62	700
139		QC-SMD-TFA05	1.57	760
140		QC-SMD-TFA05	1.34	750

[0816] Although tautomers of some of the compounds described in this table exist, isomers may be observed in some cases and not in other cases according to the type of the solvent for measurement. For example, ¹H-NMR of Example 129 (dimethylsulfoxide-D₆) and Example 140 (chloroform-D) is as follows.

<Example 129>

¹H-NMR (DMSO-*D*₆) δ: 16.58 (1H, brs), 12.80 (1H, brs), 12.15 (1H, brs), 9.59 (1H, s), 8.59-8.56 (1H, brm), 8.48-8.38 (1H, m), 8.27 (1H, brs), 7.99 (1H, d, *J* = 9.8 Hz), 7.28 (1H, q, *J* = 6.8 Hz), 7.24 (1H, q, *J* = 7.3 Hz), 5.11 (1H, d, *J* = 14.7 Hz), 4.47-4.38 (1H, m), 3.41-3.30 (2H,

m), 2.73 (1H, q, $J = 8.6$ Hz), 2.54 (2H, t, $J = 6.4$ Hz), 2.49-2.36 (1H, m), 2.40 (2H, t, $J = 7.8$ Hz), 1.84-1.75 (2H, m), 1.63 (2H, brs), 0.92 (3H, brs).

[0817] <Example 140>

Major tautomer



¹H-NMR (CDCl₃) δ: 11.82 (1H, s), 8.78 (1H, d, J = 2.0 Hz), 8.44 (1H, d, J = 8.6 Hz), 7.97 (1H, dd, J = 8.1, 1.9 Hz), 7.86 (1H, d, J = 8.2 Hz), 7.73 (1H, dd, J = 8.7, 1.9 Hz), 7.56 (1H, d, J = 1.8 Hz), 7.15-7.11 (1H, m), 6.98-6.94 (1H, m), 4.84 (1H, d, J = 14.3 Hz), 4.32 (1H, d, J = 13.9 Hz), 3.85-3.79 (4H, m), 3.68 (2H, s), 3.27 (1H, td, J = 8.7, 3.2 Hz), 2.83-2.77 (4H, m), 2.73-2.64 (1H, m), 2.58-2.48 (1H, m), 1.80-1.58 (3H, m), 0.99 (3H, s).

[0818] Minor tautomer

¹H-NMR (CDCl₃) δ: 11.91 (1H, s), 8.72 (1H, d, J = 2.2 Hz), 8.19 (1H, d, J = 8.2 Hz), 7.96-7.92 (1H, m), 7.80 (1H, d, J = 8.2 Hz), 7.75-7.71 (1H, m), 7.58-7.55 (1H, m), 7.17-7.07 (1H, m), 6.99-6.93 (1H, m), 5.07 (1H, d, J = 14.7 Hz), 4.47 (1H, d, J = 14.9 Hz), 3.79-3.85 (4H, m), 3.67 (2H, s), 3.30-3.20 (1H, m), 2.83-2.77 (4H, m), 2.73-2.64 (1H, m), 2.58-2.48 (1H, m), 1.81-1.58 (3H, m), 0.79 (3H, s).

[0819] The appropriate iodide derivatives of Examples 319 to 323 and appropriate terminal alkyne reagents were used, and operations similar to those of Example 123 and Example 125 were carried out to synthesize the compounds described in the following Table.




[0820] [Table 23]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
141		QC-SMD-TFA05	1.61	743
142		QC-SMD-TFA05	1.59	808

[0821] The appropriate iodide derivatives of Examples 319 to 323 and appropriate terminal

alkyne reagents including that in Reference Example 76 were used, and operations similar to those of Example 124 were carried out to synthesize the compounds described in the following Table.


[0822] [Table 24]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
143		QC-SMD-TFA05	1.56	797
144		QC-SMD-TFA05	1.58	797
145		QC-SMD-TFA05	1.51	769

[0823] <Example 146>

(4aR)-1-[[2,6-Dichloro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide hydrochloride

[0824] [Formula 114]



HCl

[0825] (4aR)-1-[(2,6-Dichloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 328) (22 mg, 0.032 mmol), 4-(2-chloroethyl)morpholine hydrochloride (12.1 mg, 0.065 mmol), and tetra n-butylammonium iodide (1.2 mg, 0.0033 mmol) were dissolved in N,N-dimethylformamide (0.2 mL), cesium carbonate (64 mg, 0.20 mmol) was added, and the mixture was stirred under nitrogen

atmosphere at 60°C for 3 hours. Dimethylsulfoxide (0.4 mL) and concentrated hydrochloric acid (0.1 mL) were added to the reaction mixture at 0°C, and after the mixture was stirred for 5 minutes, the resultant was purified directly by C-18 reverse-phase column chromatography on silica gel (water/acetonitrile) to obtain the title compound (23 mg, 88%) as a white amorphous solid.


[0826] LCMS: m/z 790[M+H]⁺

HPLC retention time: 1.26 minutes (analysis condition SMD-TFA05)

[0827] <Example 147>


(4aR)-1-[[4-[3-[(2R)-2,3-Dihydroxypropoxy]propoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0828] [Formula 115]



[0829] First Step

[0830] [Formula 116]



[0831] Water (10 mL) and sodium hydroxide (1.75 g, 43.8 mmol) were added to 1,3-dibromopropane (9.17 g, 45.4 mmol), (S)-(2,2-dimethyl-1,3-dioxolan-4-yl)methanol (1.5 g, 11.4 mmol), and hydrogensulfate tetra n-butylammonium (0.19 g, 0.57 mmol), and the mixture was stirred under nitrogen atmosphere at room temperature for 16 hours. Water (30 mL) was added to the reaction mixture, and after the resultant was extracted with diethyl ether, the organic layer was dried over magnesium sulfate and filtered. The filtrate was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain (S)-4-((3-bromopropoxy)methyl)-2,2-dimethyl-1,3-dioxolane (640 mg, 22%) as clear liquid.

[0832] Second Step

(4aR)-1-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 324) (1.0 g, 1.56 mmol) and (S)-4-((3-bromopropoxy)methyl)-2,2-dimethyl-1,3-dioxolane (0.47 g, 1.87 mmol) were dissolved in N,N-dimethylformamide (3 mL), cesium carbonate (0.86 g, 6.23 mmol) was added, and the mixture was stirred under nitrogen atmosphere at 80°C for 2 hours. Diethyl ether (30 mL) was added to the reaction mixture, and the insolubles were filtered and concentrated at reduced pressure. After methanol (10 mL) was added to the resultant residue, concentrated hydrochloric acid (2 mL) was added dropwise at 0°C, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was blown with nitrogen for 1 hour to remove an excessive amount of hydrogen chloride, and subsequently, dimethylsulfoxide (5 mL) was added, methanol was removed at reduced pressure, and the resultant residue was purified by C-18 reverse-phase column chromatography on silica gel (water/acetonitrile) to obtain the title compound (1.2 g, 95%) as a white amorphous solid.

[0833] LCMS: m/z 775[M+H]⁺

HPLC retention time: 1.49 minutes (analysis condition SMD-TFA05)

[0834] Major tautomer

¹H-NMR (CDCl₃) δ : 16.21 (1H, s), 11.91 (1H, s), 8.78 (1H, s), 8.44 (1H, d, J = 8.7 Hz), 7.98 (1H, d, J = 8.0 Hz), 7.87 (1H, d, J = 8.0 Hz), 7.73 (1H, d, J = 8.7 Hz), 7.55 (1H, s), 6.92 (1H, dd, J = 7.8, 7.5 Hz), 6.71 (1H, dd, J = 7.8, 7.5 Hz), 4.73 (1H, d, J = 14.2 Hz), 4.30 (1H, d, J = 14.2 Hz), 4.14 (2H, t, J = 6.2 Hz), 3.89-3.87 (1H, m), 3.72-3.69 (3H, m), 3.65-3.64 (1H, m), 3.57-3.54 (2H, m), 3.27-3.24 (1H, m), 2.70-2.62 (1H, m), 2.56-2.55 (1H, m), 2.53-2.50 (1H, m), 2.12-2.08 (3H, m), 1.73-1.70 (1H, m), 1.67-1.61 (2H, m), 1.00 (3H, s).

[0835] Minor tautomer


¹H-NMR (CDCl₃) δ : 17.97 (1H, s), 11.89 (1H, s), 8.71 (1H, s), 8.19 (1H, d, J = 8.6 Hz), 7.94 (1H, d, J = 7.9 Hz), 7.79 (1H, d, J = 8.0 Hz), 7.73 (1H, d, J = 8.6 Hz), 7.55 (1H, s), 7.06-7.04 (1H, m), 6.73-6.70 (1H, m), 4.99 (1H, d, J = 14.4 Hz), 4.42 (1H, d, J = 14.4 Hz), 4.14 (2H, t,

$J = 6.2$ Hz), 3.89-3.88 (1H, m), 3.72-3.69 (3H, m), 3.65-3.64 (1H, m), 3.57-3.54 (2H, m), 3.25-3.24 (1H, m), 2.70-2.62 (2H, m), 2.41-2.39 (1H, m), 2.12-2.08 (3H, m), 1.67-1.61 (2H, m), 1.43-1.37 (1H, m), 0.77 (3H, s).

[0836] <Reference Example 78>

(S)-4-(((5-Bromopentyl)oxy)methyl)-2,2-dimethyl-1,3-dioxolane

[0837] [Formula 117]




[0838] 1,5-Dibromopentane and (S)-(2,2-dimethyl-1,3-dioxolan-4-yl)methanol were used, and operations similar to those of First Step of Example 147 were carried out to obtain (S)-4-(((5-bromopentyl)oxy)methyl)-2,2-dimethyl-1,3-dioxolane as clear liquid.

[0839] $^1\text{H-NMR}$ (CDCl_3) δ : 4.25 (1H, dt, $J = 12.0, 6.0$ Hz), 4.05 (1H, dd, $J = 6.4, 8.0$ Hz), 3.72 (1H, dd, $J = 6.4, 8.0$ Hz), 3.53-3.37 (6H, m), 1.87 (2H, dt, $J = 14.2, 7.1$ Hz), 1.64-1.54 (2H, m), 1.54-1.44 (2H, m), 1.41 (3H, s), 1.35 (3H, s).

[0840] <Reference Example 79>

(R)-4-(((5-Bromopentyl)oxy)methyl)-2,2-dimethyl-1,3-dioxolane

[0841] [Formula 118]




[0842] 1,5-Dibromopentane and (R)-(2,2-dimethyl-1,3-dioxolan-4-yl)methanol were used, and operations similar to those of First Step of Example 147 were carried out to obtain (R)-4-(((5-bromopentyl)oxy)methyl)-2,2-dimethyl-1,3-dioxolane.

[0843] <Example 148>

(4aR)-1-[[2,6-Dichloro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[6-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]pyridin-3-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide hydrochloride

[0844]

[Formula 119]



[0845] (4aR)-1-[(2,6-Dichloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[6-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]pyridin-3-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 329) and 4-(2-chloroethyl)morpholine hydrochloride were used, and operations similar to those of Example 146 were carried out to synthesize the title compound.


[0846] LCMS: m/z 789[M+H]⁺

HPLC retention time: 1.29 minutes (analysis condition QC-SMD-TFA05)

[0847] <Example 149>


(4aR)-1-[[3-Chloro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0848] [Formula 120]



[0849] First Step

[0850] [Formula 121]




[0851] 4-(Benzyloxy)-3-chlorobenzaldehyde and (R)-2-methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride were used, and operations similar to those of Reference Example 4 were carried out to synthesize (4aR)-1-[(3-chloro-4-phenylmethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-

b]pyridazine-3-carboxylic acid 2-methylpropyl ester.

[0852] Second Step


[0853] [Formula 122]



[0854] (4aR)-1-[(3-Chloro-4-phenylmethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester was used, and operations similar to those of Second Step of Reference Example 74 were carried out to synthesize (4aR)-1-[(3-chloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester.

[0855] Third Step

[0856] [Formula 123]



[0857] (4aR)-1-[(3-Chloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester and 4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyridin-3-yl)aniline (Reference Example 13) were used, and operations similar to those of Example 21 were carried out to synthesize (4aR)-1-[(3-chloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide.

[0858] Fourth Step

(4aR)-1-[(3-Chloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide and 4-(2-chloroethyl)morpholine hydrochloride were used, and

operations similar to those of Example 146 were carried out to synthesize the title compound.


[0859] LCMS: m/z 754[M+H]⁺

HPLC retention time: 1.32 minutes (analysis condition QC-SMD-TFA05)

[0860] <Example 150>


Methyl 2-[4-[[[(4aR)-4-Hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenoxy] acetate

[0861] [Formula 124]



[0862] First Step

[0863] [Formula 125]




[0864] (4aR)-1-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 324) and bromoacetic acid tert-butyl ester were used, and operations similar to those of Example 146 were carried out to synthesize 2-[4-[[[(4aR)-4-hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenoxy]acetic acid tert-butyl ester.

[0865] LCMS: m/z 757[M+H]⁺

HPLC retention time: 1.23 minutes (analysis condition SQD-FA05)

[0866] Second Step

[0867] [Formula 126]



[0868] Trifluoroacetic acid (0.3 mL, 4 mmol) was added to a solution of 2-[4-[[[(4aR)-4-hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenoxy]acetic acid tert-butyl ester (14.7 mg, 0.019 mmol) in dichloromethane (0.075 mL) at room temperature, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was concentrated at reduced pressure, and the resultant residue was purified by reverse-phase column chromatography (0.1% formic acid, acetonitrile/water) to obtain 2-[4-[[[(4aR)-4-hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenoxy]acetic acid (12.3 mg, 90%).

[0869] LCMS: m/z 701[M+H]⁺

HPLC retention time: 1.08 minutes (analysis condition SQD-FA05)

[0870] Third Step

(Diazomethyl)trimethylsilane (0.0125 mL, 0.025 mmol) was added to a solution of 2-[4-[[[(4aR)-4-hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenoxy]acetic acid (9 mg, 0.013 mmol) in benzene (0.13 mL)-methanol (0.08 mL) at room temperature, and the mixture was stirred at room temperature for 1 hour. After formic acid (0.005 mL) was added to the reaction mixture, the resultant was concentrated at reduced pressure, and the resultant residue was purified by reverse-phase column chromatography (0.1% formic acid, acetonitrile/water) to obtain the title compound

(8.3 mg, 90%) as a white amorphous solid.


[0871] LCMS: m/z 715[M+H]⁺

HPLC retention time: 1.60 minutes (analysis condition QC-SMD-TFA05)

[0872] <Example 151>

(4aR)-1-[[2,3-Difluoro-4-(pyridin-4-ylmethoxy)phenyl]methyl]-N-[2-(2,3-dimethoxyphenyl)-4-(trifluoromethyl)phenyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0873] [Formula 127]



[0874] (4aR)-1-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-N-[2-(2,3-dimethoxyphenyl)-4-(trifluoromethyl)phenyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 325) and 4-(bromomethyl)pyridine hydrobromate were used, and operations similar to those of Example 146 were carried out to synthesize the title compound.






[0875] LCMS: m/z 725[M+H]⁺

HPLC retention time: 1.37 minutes (analysis condition QC-SMD-TFA05)

[0876] The phenolic derivatives of Examples 326, 327, and 330 and alkyl bromide reagents including those in Reference Examples 78 and 79 were used, and operations similar to those of Second Step of Example 147 were carried out to synthesize the compounds of the Examples described in the following Table.

[0877]

[Table 25]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
152		QC-SMD-TFA05	1.53	790
153		QC-SMD-TFA05	1.56	803
154		QC-SMD-TFA05	1.60	817
155		QC-SMD-TFA05	1.60	817
156		QC-SMD-TFA05	1.49	819

[0878] Although tautomers of some of the compounds described in this table exist, isomers may be observed in some cases and not in other cases according to the type of the solvent for measurement. For example, ¹H-NMR of Example 152 (chloroform-D) and Example 156 (dimethylsulfoxide-D6) is as follows.

<Example 152>

Major tautomer

¹H-NMR (CDCl₃) δ: 16.43 (1H, s), 12.92 (1H, s), 9.61 (1H, s), 8.47 (1H, d, *J* = 8.7 Hz), 7.95 (1H, s), 7.90 (1H, s), 7.79 (1H, d, *J* = 8.7 Hz), 7.04 (1H, dd, *J* = 8.1, 8.1 Hz), 6.70 (1H, dd, *J* = 8.1, 8.1 Hz), 4.98 (1H, d, *J* = 14.3 Hz), 4.41 (1H, d, *J* = 14.3 Hz), 4.07 (2H, t, *J* = 6.2 Hz), 3.87 (1H, brs), 3.73-3.71 (1H, m), 3.66-3.64 (1H, m), 3.56 (2H, m), 3.53-3.52 (2H, m), 3.33-3.31 (1H, m), 2.74 (1H, ddd, *J* = 8.5, 8.5, 8.5 Hz), 2.55-2.53 (2H, m), 2.11 (1H, brs), 1.92-

1.88 (2H, m), 1.82-1.77 (2H, m), 1.74-1.73 (1H, m), 1.68-1.62 (2H, m), 1.01 (2H, s).

[0879] Minor tautomer

¹H-NMR (CDCl₃) δ: 18.16 (1H, s), 12.97 (1H, s), 9.59 (1H, s), 8.33 (1H, d, *J* = 8.7 Hz), 7.92 (2H, d, *J* = 10.9 Hz), 7.79 (1H, d, *J* = 8.4 Hz), 7.04 (1H, t, *J* = 8.1 Hz), 6.70 (1H, t, *J* = 7.5 Hz), 5.01 (1H, d, *J* = 17.3 Hz), 4.45 (1H, d, *J* = 14.6 Hz), 4.07 (2H, t, *J* = 6.2 Hz), 3.87 (1H, s), 3.72 (1H, d, *J* = 10.7 Hz), 3.65 (1H, d, *J* = 11.9 Hz), 3.54 (4H, dt, *J* = 19.0, 6.0 Hz), 3.26 (1H, s), 2.74 (1H, dd, *J* = 16.6, 8.5 Hz), 2.63-2.61 (2H, m), 2.11 (1H, s), 1.92-1.88 (2H, m), 1.74-1.73 (2H, m), 1.67-1.65 (2H, m), 1.50-1.47 (1H, m), 0.85 (3H, s).


[0880] <Example 156>

¹H-NMR (DMSO-D₆) δ: 15.77 (1H, brs), 12.17 (1H, brs), 9.71 (1H, s), 9.14 (1H, s), 8.54 (1H, d, *J* = 6.6 Hz), 8.16 (1H, d, *J* = 8.1 Hz), 7.13 (1H, brs), 7.00 (1H, t, *J* = 6.8 Hz), 4.99-4.17 (3H, m), 4.08 (2H, t, *J* = 6.1 Hz), 3.59-3.54 (1H, m), 3.41-3.24 (6H, m), 2.70 (1H, q, *J* = 8.1 Hz), 2.41 (1H, brs), 1.79-1.48 (8H, m), 1.47-1.31 (4H, m), 0.92 (3H, brs).

[0881] <Example 157>

(4aR)-1-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0882] [Formula 128]



[0883] Triphenylphosphine (17.2 mg, 0.066 mmol), N-(2-hydroxy ethyl)morpholine (7.5 mg, 0.057 mmol), and diethyl azodicarboxylate (0.03 mL, 0.065 mmol) were added to a solution of (4aR)-1-[(2,3-difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 327) (10.6 mg, 0.016 mmol) in

tetrahydrofuran (0.2 mL), and the mixture was stirred at room temperature for 1.5 hours.

After the reaction mixture was concentrated, the residue was purified by reverse-phase column chromatography on silica gel to obtain the title compound (8.0 mg, 0.011 mmol) as a white solid. It was confirmed by measuring ^1H -NMR and ^{13}C -NMR that this solid included two tautomers.

[0884] LCMS: m/z 757 $[\text{M}+\text{H}]^+$

HPLC retention time: 1.30 minutes (SMD-TFA05)

[0885] Major tautomer

^1H -NMR (CDCl_3) δ : 16.44 (1H, s), 12.92 (1H, s), 9.61 (1H, d, $J = 1.0$ Hz), 8.47 (1H, d, $J = 8.7$ Hz), 7.95 (1H, d, $J = 1.0$ Hz), 7.90 (1H, d, $J = 2.0$ Hz), 7.79 (1H, dd, $J = 8.7, 2.0$ Hz), 7.05 (1H, ddd, $J = 9.0, 6.8, 2.2$ Hz), 6.71 (1H, ddd, $J = 9.0, 7.5, 1.2$ Hz), 4.99 (1H, d, $J = 14.2$ Hz), 4.40 (1H, d, $J = 14.2$ Hz), 4.19 (2H, t, $J = 5.7$ Hz), 3.74 (4H, t, $J = 4.6$ Hz), 3.32 (1H, ddd, $J = 8.9, 8.9, 3.5$ Hz), 2.84 (2H, t, $J = 5.7$ Hz), 2.74 (1H, ddd, $J = 8.9, 8.9, 8.9$ Hz), 2.61-2.59 (4H, m), 2.56-2.54 (1H, m), 1.76-1.73 (1H, m), 1.68-1.63 (2H, m), 1.00 (3H, s).

[0886] ^{13}C -NMR (CDCl_3) δ : 187.4 (qC), 169.8 (qC), 165.7 (qC), 162.5 (qC), 159.3 (CH), 156.6 (qC, $J_{\text{CF}} = 36.3$ Hz), 150.3 (qC, $J_{\text{CF}} = 248.7, 10.5$ Hz), 147.8 (qC, $J_{\text{CF}} = 5.5$ Hz), 141.3 (qC, $J_{\text{CF}} = 247.9, 14.6$ Hz), 138.7 (qC), 128.3 (CH, $J_{\text{CF}} = 3.3$ Hz), 128.0 (qC), 127.1 (CH, $J_{\text{CF}} = 3.8$ Hz), 127.0 (qC, $J_{\text{CF}} = 33.5$ Hz), 125.1 (CH), 124.8 (CH), 123.6 (qC, $J_{\text{CF}} = 272.0$ Hz), 120.5 (qC, $J_{\text{CF}} = 275.4$ Hz), 118.4 (qC, $J_{\text{CF}} = 12.7$ Hz), 116.0 (CH, $J_{\text{CF}} = 2.5$ Hz), 109.5 (CH, $J_{\text{CF}} = 2.5$ Hz), 93.6 (qC), 68.0 (CH_2), 66.9 ($\text{CH}_2 \times 2$), 64.2 (qC), 57.5 (CH_2), 54.1 ($\text{CH}_2 \times 2$), 51.9 (CH_2), 43.2 (CH_2), 31.9 (CH_2), 22.4 (CH_3), 19.1 (CH_2).

[0887] Minor tautomer

^1H -NMR (CDCl_3) δ : 18.17 (1H, s), 12.97 (1H, s), 9.59 (1H, s), 8.33 (1H, d, $J = 8.7$ Hz), 7.93 (1H, s), 7.91 (1H, d, $J = 2.0$ Hz), 7.79 (1H, dd, $J = 8.7, 2.0$ Hz), 7.05 (1H, ddd, $J = 9.0, 6.8, 2.2$ Hz), 6.71 (1H, ddd, $J = 9.0, 7.5, 1.2$ Hz), 5.02 (1H, d, $J = 14.5$ Hz), 4.45 (1H, d, $J = 14.5$ Hz), 4.19 (2H, t, $J = 5.7$ Hz), 3.74 (4H, t, $J = 4.6$ Hz), 3.25 (1H, ddd, $J = 8.8, 8.8, 3.2$ Hz), 2.84 (2H, t, $J = 5.7$ Hz), 2.74 (1H, ddd, $J = 8.9, 8.9, 8.9$ Hz), 2.61-2.59 (4H, m), 2.56-2.54 (1H, m), 1.76-1.73 (1H, m), 1.68-1.63 (1H, m), 1.49-1.47 (1H, m), 0.84 (3H, s).

[0888] ^{13}C -NMR (CDCl_3) δ : 191.8 (qC), 170.2 (qC), 168.1 (qC), 165.5 (qC), 159.3 (CH), 156.6 (qC, $J_{\text{CF}} = 36.3$ Hz), 150.3 (qC, $J_{\text{CF}} = 248.7, 10.5$ Hz), 147.8 (qC, $J_{\text{CF}} = 5.5$ Hz), 141.3 (qC, $J_{\text{CF}} = 247.9, 14.6$ Hz), 138.7 (qC), 128.2 (CH, $J_{\text{CF}} = 3.3$ Hz), 128.2 (qC), 127.1 (CH, $J_{\text{CF}} = 3.8$ Hz), 127.0 (qC, $J_{\text{CF}} = 33.5$ Hz), 125.1 (CH), 125.4 (CH), 123.6 (qC, $J_{\text{CF}} = 272.0$ Hz), 120.5 (qC, $J_{\text{CF}} = 275.4$ Hz), 118.4 (qC, $J_{\text{CF}} = 12.7$ Hz), 116.0 (CH, $J_{\text{CF}} = 2.5$ Hz), 109.5 (CH, $J_{\text{CF}} = 2.5$ Hz), 84.3 (qC), 68.0 (CH_2), 66.9 ($\text{CH}_2 \times 2$), 64.2 (qC), 57.5 (CH_2), 54.1 ($\text{CH}_2 \times 2$), 50.7 (CH_2), 44.0 (CH_2), 31.1 (CH_2), 21.8 (CH_3), 18.6 (CH_2).

[0889] The appropriate phenol intermediates of Examples 324, 325, and 327 and appropriate alcohol reagents were used, and operations similar to those of Example 156 were carried out to synthesize the compounds of the Examples described in the following Table.


[0890] [Table 26]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z $[\text{M}+\text{H}]^+$
158		QC-SMD-TFA05	1.68	718
159		QC-SMD-TFA05	1.69	718
160		SMD-TFA50	1.17	727
161		QC-SMD-TFA05	1.32	756
162		QC-SMD-TFA05	1.61	730

[0891] <Example 163>

(4aR)-1-[[2,3-Difluoro-4-(2-morpholin-4-ylethylsulfanyl)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide hydrochloride

[0892] [Formula 129]



[0893] Cesium carbonate (67 mg, 0.206 mmol) was added to a solution of (4aR)-1-[(2,3-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 320) (30 mg, 0.04 mmol), palladium acetate (1.79 mg, 0.0080 mmol), triphenylphosphine (8.37 mg, 0.032 mmol), and triisopropylsilanethiol (9.87 mg, 0.052 mmol) in toluene (0.399 mL), and the mixture was stirred at 100°C for 2 hours and 45 minutes. Subsequently, palladium acetate (1.79 mg, 0.0080 mmol), triisopropylsilanethiol (9.87 mg, 0.052 mmol), and 1,4-dioxane (0.4 mL) were further added, and the mixture was stirred at 110°C for 2 hours.

[0894] After the reaction mixture was cooled to room temperature, 4-(2-chloroethyl)morpholine hydrochloric acid (37.1 mg, 0.199 mmol) was added to this reaction mixture, and after the resultant was heated at 100°C for several hours, it was cooled to room temperature. This reaction mixture was purified by C18 reverse-phase column chromatography. A 4 N hydrogen chloride/1,4-dioxane solution was added to the obtained compound, and the solution was concentrated and dried to obtain the title compound (17 mg, 53%) as a white solid.


[0895] LCMS: m/z 772[M+H]⁺

HPLC retention time: 1.35 minutes (analysis condition QC-SMD-TFA05)

[0896] <Example 164>


(4aR)-1-[[2,3-Difluoro-4-(morpholin-4-ylmethyl)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide hydrochloride

[0897] [Formula 130]



[0898] First Step

[0899] [Formula 131]



[0900] A solution of (bromomethyl)potassium trifluoroborate (100 mg, 0.50 mmol), morpholine (45.7 mg, 0.525 mmol), and potassium carbonate (69.1 mg, 0.50 mmol) in tetrahydrofuran (0.50 mL) was heated at 80°C for 1 hour, and then the resultant was cooled to room temperature. After the reaction mixture was filtered and washed with acetone (15 mL), the solution was dried to obtain trifluoro(morpholinomethyl)potassium borate (68 mg, 66%) as a crude product.

[0901] Second Step

Palladium acetate (2.45 mg, 0.0109 mmol), dicyclohexyl(2',4',6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphine (10.39 mg, 0.022 mmol), and cesium carbonate (89 mg, 0.272 mmol) were added twice to a solution of trifluoro(morpholinomethyl)potassium borate (28.2 mg, 0.136 mmol) obtained in First Step and (4aR)-1-[(2,3-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 320) (41 mg, 0.054 mmol) in a mixture of dioxane and water (0.495 mL and 0.0495 mL), and the mixture was heated and stirred at 80°C overnight. After the reaction mixture was cooled to room temperature, this reaction mixture was purified by

reverse-phase column chromatography. The obtained compound was dissolved in ethanol, then a 4 N hydrogen chloride/dioxane solution were added, and the resultant was dried to obtain the title compound (28 mg, 68%) as a white solid.

[0902] LCMS: m/z 726[M+H]⁺

HPLC retention time: 1.28 minutes (analysis condition QC-SMD-TFA05)

[0903] Major tautomer

¹H-NMR (CDCl₃) δ : 13.60 (1H, s), 11.74 (1H, s), 8.74 (1H, d, J = 2.0 Hz), 8.44 (1H, d, J = 8.6 Hz), 7.95 (1H, dd, J = 8.0, 1.8 Hz), 7.86 (1H, d, J = 8.0 Hz), 7.75 (1H, m), 7.72 (1H, dd, J = 8.6, 2.0 Hz), 7.54 (1H, d, J = 1.8 Hz), 7.17 (1H, t, J = 6.7 Hz), 4.86 (1H, d, J = 14.9 Hz), 4.37-4.22 (4H, m), 3.97 (2H, dd, J = 13.2, 2.6 Hz), 3.35 (2H, d, J = 10.4 Hz), 3.27 (1H, dt, J = 8.7, 3.2 Hz), 2.92 (2H, m), 2.71 (1H, dd, J = 16.7, 8.7 Hz), 2.60-2.51 (1H, m), 1.87-1.58 (4H, m), 1.04 (3H, s).


[0904] Minor tautomer

¹H-NMR (CDCl₃) δ : 11.91 (1H, s), 8.70 (1H, d, J = 2.0 Hz), 8.17 (1H, d, J = 8.6 Hz), 7.80 (1H, d, J = 8.0 Hz), 7.57 (1H, d, J = 2.0 Hz), 7.32 (1H, t, J = 6.9 Hz), 5.12 (1H, d, J = 14.9 Hz), 4.48-4.38 (4H, m), 0.80 (3H, s).

[0905] <Example 165>

Methyl 4-[[[(4aR)-4-Hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-3,5-difluorobenzoate

[0906] [Formula 132]



[0907] Molybdenum hexacarbonyl (70.2 mg, 0.266 mmol), 1,1'-bis(diphenylphosphino)ferrocene dichloropalladium (II) (a 1:1 dichloromethane complex) (4.4 mg, 0.0053 mmol), and 1,8-diazabicyclo[5.4.0]undec-7-ene (0.0024 mL, 0.159 mmol)

were added to a solution of (4aR)-1-[(2,6-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 322) (40 mg, 0.053 mmol) in methanol (2 mL), and the mixture was stirred under nitrogen atmosphere at 70°C for 4 hours. The reaction mixture was added to 1 N hydrochloric acid, the resultant was extracted with dichloromethane, and after the organic layer was dried over anhydrous magnesium sulfate, the resultant was filtered and concentrated at reduced pressure. The resultant residue was purified by preparative TLC (hexane:ethyl acetate = 2:1) to obtain the title compound (29.3 mg, 81%) as light brown oil.


[0908] LCMS: m/z 685[M+H]⁺

HPLC retention time: 1.68 minutes (analysis condition QC-SMD-TFA05)

[0909] <Reference Example 80>


(4aR)-N-(4-Bromo-3,5-difluorophenyl)-4-hydroxy-4a-methyl-2-oxo-1,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-3-carboxamide

[0910] [Formula 133]



[0911] First Step

[0912] [Formula 134]




[0913] A solution of 4-bromo-3,5-difluoroaniline (2.00 g, 9.62 mmol) and Meldrum's acid (2.77 g, 19.2 mmol) in toluene (18 mL) was stirred at 90°C for 3 hours. After the reaction mixture was cooled to 0°C, the deposit was collected by filtration. The obtained solid was purified by reverse-phase column chromatography (0.1% formic acid water/acetonitrile) to obtain 3-((4-bromo-3,5-difluorophenyl)amino)-3-oxopropanoate (2.00 g, 71%) as white powder.

[0914] LCMS: m/z 294 $[M+H]^+$

HPLC retention time: 0.66 minutes (analysis condition SQD-FA05)

[0915] Second Step

[0916] [Formula 135]



[0917] 2,4-Dimethoxybenzaldehyde, (R)-2-methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride, and 3-((4-bromo-3,5-difluorophenyl)amino)-3-oxopropanoate were used, and operations similar to those of Reference Example 1-1 were carried out to synthesize (4aR)-N-(4-bromo-3,5-difluorophenyl)-1-[(2,4-(dimethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide.

[0918] LCMS: m/z 554 $[M+H]^+$

HPLC retention time: 1.64 minutes (analysis condition SQD-FA05)

[0919] Third Step

(4aR)-N-(4-Bromo-3,5-difluorophenyl)-1-[(2,4-(dimethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (910 mg, 1.65 mmol) and triisopropylsilane (0.68 mL, 3.29 mmol) were dissolved in trifluoroacetic acid (6 mL), and after the mixture was cooled to 0°C, trifluoromethanesulfonic acid (0.15 mL, 1.65 mmol) was added dropwise. After the mixture was stirred at room temperature for 30 minutes, the reaction mixture was added to ice-cold water (50 mL), neutralized with potassium carbonate (5.5 g, 39.8 mmol), and extracted three times with ethyl acetate (100 mL). After the organic layer was washed with a brine, the resultant was dried over sodium sulfate. The resultant was filtered, the filtrate was concentrated at reduced pressure, and the resultant residue was purified by C-18 reverse-phase column chromatography (0.1% formic acid water/acetonitrile) to obtain the title compound (590 mg, 89%) as a white solid.


[0920] LCMS: m/z 402 $[M+H]^+$

HPLC retention time: 0.70 minutes (analysis condition SQD-FA05)

[0921] <Reference Example 81>


(4aR)-4-Hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-1,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-3-carboxamide

[0922] [Formula 136]



[0923] First Step

[0924] [Formula 137]



[0925] 4-(Trifluoromethyl)-2-(6-(trifluoromethyl)pyridin-3-yl)aniline (Reference Example 42) (2.0 g) was dissolved in tetrahydrofuran (30 mL), tripotassium phosphate (4.16 g) was added, and the mixture was stirred at room temperature for 3 minutes. Methyl 3-chloro-3-oxopropanoate (1.40 mL) was added, and the mixture was stirred at room temperature for 10 minutes. Ethyl acetate (100 mL) and water (100 mL) were added for separation, and the organic layer was washed with a 12% brine and concentrated at reduced pressure to obtain propanedioic acid 3-O-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl] 1-O-methyl as a crude product.


[0926] LCMS: m/z 407[M+H]⁺

HPLC retention time: 0.81 minutes (analysis condition SQD-FA05)

[0927] Second Step

[0928]

[Formula 138]




[0929] Methanol (20 mL) and a 10 N aqueous sodium hydroxide solution (1.96 mL) were added to 3-O-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl] 1-O-methyl obtained in First Step, and the mixture was stirred at room temperature for 10 hours. 6 N hydrochloric acid was added to adjust the pH to 3, ethyl acetate and water were added for separation, and the organic layer was dried over anhydrous magnesium sulfate and the solvent was distilled away at reduced pressure. Hexane (90 mL) and ethyl acetate (10 mL) were added to the resultant residue, the mixture was stirred and then filtered, and the resultant solid was dried to obtain 3-oxo-3-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyridin-3-yl)phenyl)amino)propanoate (2.34 g, two-step yield 91%).

[0930] LCMS: m/z 393[M+H]⁺

HPLC retention time: 0.29 minutes (analysis condition SQD-FA50)

[0931] Third Step

[0932] [Formula 139]



[0933] (R)-2-Methylpyrrolidine-2-carboxylic acid methyl ester hydrochloride (8.4 g, 46.8 mmol) was suspended in dichloromethane (50 mL), p-toluenesulfonic acid-monohydrate (9.34 g, 49.1 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 10 minutes. The reaction mixture was concentrated at reduced pressure, toluene was added for azeotropic removal, then the residue was suspended in dichloromethane (50 mL), sodium nitrite (3.55 g, 69.0 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 2 hours. After the reaction mixture was filtered, the resultant was concentrated at reduced pressure to obtain (S)-1-nitroso-pyrrolidine-2-carboxylic acid methyl ester as a crude product. The obtained crude


product was dissolved in acetic acid (200 mL) and methanol (30 mL), zinc powder (29.4 g, 4501 mmol) was added in divided portions under nitrogen stream, at 0°C, and it was added, and the mixture was stirred for 1 hour. Methanol (100 mL) was added to the reaction mixture, the mixture was filtered through a celite pad, and then the filtrate was concentrated at reduced pressure. After a similar operation was repeated twice, the resultant residue was purified by column chromatography on silica gel (dichloromethane/methanol). After a 4 N hydrogen chloride/1,4-dioxane solution was added to the obtained fraction, the resultant was concentrated at reduced pressure to obtain (R)-1-amino-2-methylpyrrolidine-2-carboxylic acid methyl ester hydrochloride (6.4 g, 70%).

[0934] LCMS: m/z 158.9[M+H]⁺

HPLC retention time: 0.33 minutes mass chromatogram (analysis condition SMD-TFA05)

[0935] Fourth Step

[0936] [Formula 140]



[0937] (R)-2-Methyl-1-(3-oxo-3-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyridin-3-yl)phenyl)amino)propanamide)pyrrolidine-2-carboxylic acid methyl ester and (R)-1-amino-2-methylpyrrolidine-2-carboxylic acid methyl ester hydrochloride were used as a reagent and a starting material, respectively, and operations similar to those of Second Step of Reference Example 1-1 were carried out to synthesize (R)-2-methyl-1-(3-oxo-3-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyridin-3-yl)phenyl)amino)propanamide)pyrrolidine-2-carboxylic acid methyl ester.

[0938] LCMS: m/z 531[M-H]⁻

HPLC retention time: 0.88 minutes (SQD-FA05)

[0939] Fifth Step

(R)-2-Methyl-1-(3-oxo-3-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyridin-3-yl)phenyl)amino)propanamide)pyrrolidine-2-carboxylic acid methyl ester was used, and operations similar to those of Third Step of Reference Example 1-1 were carried out to synthesize the title compound.


[0940] LCMS: m/z 501[M+H]⁺

HPLC retention time: 1.03 minutes (SQD-FA05)

[0941] <Reference Example 82>


(4aR)-4-Hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-1,5,6,7,-tetrahydropyrrolo[1,2-b]pyridazine-3-carboxamide

[0942] [Formula 141]



[0943] First Step

[0944] [Formula 142]



[0945] 4-(Trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)aniline (Reference Example 50) and Meldrum's acid were used, and operations similar to those of Second Step of Reference Example 80 were carried out to synthesize 3-oxo-3-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)amino)propanoate.

[0946] Second Step

(R)-1-amino-2-methylpyrrolidine-2-carboxylic acid methyl ester hydrochloride and 3-oxo-3-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)amino)propanoate were used, and operations similar to those of Fourth and Fifth Steps of Reference Example 81 were carried out to synthesize the title compound.


[0947] LCMS: m/z 502[M+H]⁺

HPLC retention time: 1.02 minutes (SQD-FA05)

[0948] <Example 166>

(4aR)-N-(4-Bromo-3,5-difluorophenyl)-1-[(4-bromo-2-fluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0949] [Formula 143]



[0950] A 1.7 N toluene solution (0.0644 mL, 0.109 mmol) of 4-bromo-1-(bromomethyl)-2-fluorobenzene (14.7 mg, 0.055 mmol) and potassium 2,2-dimethylpropan-1-olate was added to a solution of (4aR)-N-(4-bromo-3,5-difluorophenyl)-4-hydroxy-4a-methyl-2-oxo-1,5,6,7-tetrahydropyrrolo[1,2-b]pyridazine-3-carboxamide (Reference Example 80) (20.0 mg, 0.050 mmol) in N,N-dimethylformamide (0.249 mL), and the mixture was stirred at room temperature for 15 minutes. The reaction mixture was purified directly by C18 reverse-phase column chromatography (0.1% formic acid acetonitrile/water) to obtain the title compound (6.0 mg, 20%) as a grayish white amorphous solid.








[0951] LCMS: m/z 588[M-H]⁻

HPLC retention time: 1.32 minutes (SQD-FA05)

[0952] The appropriate pyridazinone intermediates of Reference Examples 80 to 82 and appropriate benzylhalide reagents were used, and operations similar to those of Example 166 were carried out to synthesize the compounds of the Examples described in the following Table.









[0953]

[Table 27-1]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
167		QC-SMD-TFA05	1.65	657
168		SQD-AA05	1.20	625
169		SQD-AA05	1.19	625
170		QC-SMD-TFA05	1.67	627
171		QC-SMD-TFA05	1.68	627
172		SQD-AA05	1.16	627
173		QC-SMD-TFA05	1.67	627
174		QC-SMD-TFA05	1.74	544










[0954]

[Table 27-2]










175		QC-SMD-TFA05	1.66	651
176		SQD-AA05	1.18	651
177		QC-SMD-TFA05	1.68	658
178		QC-SMD-TFA05	1.65	617
179		QC-SMD-TFA05	1.64	617
180		QC-SMD-TFA05	1.76	626
181		QC-SMD-TFA05	1.76	626
182		QC-SMD-TFA05	1.76	626

[0955]










[Table 27-3]

183		QC-SMD-TFA05	1.68	650
184		QC-SMD-TFA05	1.76	688
185		QC-SMD-TFA05	1.76	606
186		QC-SMD-TFA05	1.76	606
187		QC-SMD-TFA05	1.77	606
188		QC-SMD-TFA05	1.72	592
189		QC-SMD-TFA05	1.76	676
190		SQD-AA05	1.15	528
191		SQD-AA05	1.16	528










[0956] [Table 27-4]

192		QC-SMD-TFA05	1.70	528
193		SQD-AA05	1.15	528
194		SQD-AA05	1.15	546
195		QC-SMD-TFA05	1.76	524
196		QC-SMD-TFA05	1.75	544
197		QC-SMD-TFA05	1.73	546
198		SQD-AA05	1.18	643
199		QC-SMD-TFA05	1.67	609
200		QC-SMD-TFA05	1.67	609










[Table 27-5]

201		SQD-AA05	1.18	643
202		SQD-AA05	1.17	651
203		QC-SMD-TFA05	1.66	627
204		QC-SMD-TFA05	1.71	643
205		SQD-AA05	1.21	623
206		QC-SMD-TFA05	1.67	609
207		QC-SMD-TFA05	1.68	610
208		QC-SMD-TFA05	1.68	610
209		QC-SMD-TFA05	1.72	644

[Table 27-6]







210		QC-SMD-TFA05	1.72	644
211		QC-SMD-TFA05	1.71	644
212		QC-SMD-TFA05	1.70	628
213		QC-SMD-TFA05	1.71	628
214		QC-SMD-TFA05	1.71	628
215		QC-SMD-TFA05	1.76	624
216		QC-SMD-TFA05	1.78	571
217		QC-SMD-TFA05	1.77	589
218		QC-SMD-TFA05	1.62	517

[Table 27-7]

219		QC-SMD-TFA05	1.76	526
220		QC-SMD-TFA05	1.62	517
221		QC-SMD-TFA05	1.76	526
222		QC-SMD-TFA05	1.62	517
223		QC-SMD-TFA05	1.76	526
224		SQD-AA05	1.18	589
225		QC-SMD-TFA05	1.68	590
226		QC-SMD-TFA05	1.61	615
227		SQD-AA05	1.11	615

[0960]

[Table 27-8]

228		SQD-AA05	1.11	615
229		QC-SMD-TFA05	1.67	644
230		QC-SMD-TFA05	1.66	644
231		QC-SMD-TFA05	1.71	627
232		QC-SMD-TFA05	1.70	627
233		QC-SMD-TFA05	1.70	645

[0961] Although the respective compounds described in this table include their tautomers, for example, ¹H-NMR of Example 196 is as follows.

[0962] Major tautomer

¹H-NMR (400 MHz, CDCl₃) δ: 16.40 (1H, s), 12.00 (1H, s), 7.40-7.28 (4H, m), 7.11-7.04 (1H, m), 5.14 (1H, d, *J* = 14.1 Hz), 4.47 (1H, d, *J* = 14.1 Hz), 3.37-3.32 (1H, m), 2.82-2.76 (1H, m), 2.61-2.54 (1H, m), 1.76-1.61 (3H, m), 1.01 (3H, s).


[0963] Minor tautomer

¹H-NMR (400 MHz, CDCl₃) δ: 16.40 (1H, s), 12.18 (1H, s), 7.40-7.28 (4H, m), 7.11-7.04 (1H, m), 5.11 (1H, d, *J* = 14.4 Hz), 4.53 (1H, d, *J* = 14.4 Hz), 3.32-3.26 (1H, m), 2.84-2.77 (1H, m), 2.65-2.61 (1H, m), 1.81-1.70 (3H, m), 0.88 (3H, s).

[0964] <Example 234>

(4aR)-4-Hydroxy-4a-methyl-2-oxo-1-(4,4,4-trifluorobutyl)-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0965] [Formula 144]



[0966] Methyl (4aR)-1-[(2,4-dimethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylate (Reference Example 12) and 1,1,1-trifluoro-4-iodobutane were used, and operations similar to those of Third Step of Reference Example 80, Example 166, and Example 21 were consecutively carried out to obtain the title compound.


[0967] LCMS: m/z 612[M+H]⁺

HPLC retention time: 1.63 minutes (analysis condition SMD-TFA05)

[0968] <Example 235>

Ethyl 7-[(4aR)-4-hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]heptanoate

[0969] [Formula 145]



[0970] Methyl (4aR)-1-[(2,4-dimethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylate (Reference Example 12) and ethyl 7-bromoheptanoate were used, and operations similar to those of Example 234 were carried out to obtain the title compound.


[0971] LCMS: m/z 657[M+H]⁺

HPLC retention time: 1.70 minutes (analysis condition QC-SMD-TFA05)

[0972] <Example 236>


(4aR)-4-Hydroxy-1-[8-[2-(2-methoxyethoxy)ethoxy]octyl]-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[0973] [Formula 146]



[0974] First Step

[0975] [Formula 147]



[0976] Sodium iodide was added to a solution of 1-bromo-8-(2-(2-methoxyethoxy)ethoxy)octane (1.00 g, 3.21 mmol) in acetone, and the mixture was stirred at 60°C for 2 hours. After the reaction mixture was cooled to room temperature, the precipitate was filtered for separation, and after the filtrate was washed with sodium thiosulfate and a brine, the resultant was dried over sodium sulfate. After the resultant was filtered, the filtrate was concentrated at reduced pressure to obtain 1-iodo-8-(2-(2-methoxyethoxy)ethoxy)octane (1.05 g, 91%) as yellow oil.

[0977] Second Step

Methyl (4aR)-1-[(2,4-(dimethoxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylate (Reference Example 12) and the iodine body obtained in First Step was used, and operations similar to those of Example 234 were carried out to obtain the title compound.

[0978] LCMS: m/z 732[M+H]⁺

HPLC retention time: 1.76 minutes (analysis condition QC-SMD-TFA05)


[0979] <Example 237>

(3S)-3-Tert-butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyrimidine-5-carboxamide

First Step

Methyl (2S)-3,3-Dimethyl-2-[methyl-[(2-methylpropan-2-yl)oxycarbonyl]amino]butanoate

[0980] [Formula 148]




[0981] (S)-2-((Tert-butoxycarbonyl)amino)-3,3-dimethyl butanoic acid (11.3 g, 48.9 mmol) was dissolved in N,N-dimethylformamide (114 mL), silver oxide (34.0 g, 147 mmol) and iodomethane (18.3 mL, 294 mmol) were added, and the mixture was stirred under nitrogen atmosphere at 45°C overnight. The reaction mixture was diluted with ethyl acetate (200 mL) and filtered, water was added, and the resultant was extracted with ethyl acetate. The organic layer was washed with a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (12.7 g, 100%).

[0982] ¹H-NMR (CDCl₃) δ: 4.75 (1H, s), 3.71 (3H, s), 2.94 (3H, s), 1.48 (9H, s), 1.08 (9H, s).

[0983] Second Step

Methyl (S)-3,3-Dimethyl-2-(methylamino)butanoate hydrochloride

[0984] [Formula 149]



[0985] Hydrochloric acid (a dioxane solution, 122 mL, 489 mmol) was added to methyl ((2S)-3,3-dimethyl-2-[methyl-[(2-methylpropan-2-yl)oxycarbonyl]amino]butanoate (12.69 g),


and the mixture was stirred at room temperature for 1 hour. The reaction mixture was concentrated at reduced pressure to obtain the title compound (9.58 g, 100%).

[0986] $^1\text{H-NMR}$ (DMSO-D_6) δ : 9.00 (1H, brs), 3.89-3.86 (1H, m), 3.79 (3H, s), 2.54 (3H, s), 1.02 (9H, s).

[0987] Third Step

Methyl (2S)-2-[[[(2,3-difluorophenyl)methylideneamino]-methylamino]-3,3-dimethylbutanoate

[0988] [Formula 150]



[0989] Methyl (S)-3,3-Dimethyl-2-(methylamino)butanoate hydrochloride (70.0 mg, 0.358 mmol) was dissolved in acetic acid (3.58 mmol, 205 μL) and water (716 μL), sodium nitrite (33.4 mg, 0.393 mmol) was added at 0°C , and the mixture was stirred under nitrogen atmosphere for 1 hour. Zinc (19.7 g, 302 mmol) was added to the reaction mixture at 0°C , and the mixture was stirred under nitrogen atmosphere for 4 hours. After the reaction mixture was filtered, methanol (3.58 mL) and 2,3-difluorobenzaldehyde (39.1 μL , 0.358 mmol) were added, and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated at reduced pressure and diluted with ethyl acetate, and the organic layer was washed with a saturated sodium bicarbonate solution and a brine, and then the resultant was dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (44.2 mg, 41%).

[0990] LCMS: m/z 299 $[\text{M}+\text{H}]^+$


HPLC retention time: 1.13 minutes (analysis condition SQD-AA05)

[0991] Fourth Step

(3S)-3-tert-Butyl-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-2-methyl-6-oxo-3H-

pyrimidine-5-carboxylic acid 2-methylpropyl ester

[0992] [Formula 151]



[0993] Methyl ((2S)-2-[[[(2,3-difluorophenyl)methylideneamino]-methylamino]-3,3-dimethylbutanoate (44.2 mg, 0.148 mmol)) was dissolved in hydrochloric acid (a methanol solution, 442 μ L), borane-pyridine (29.9 μ L, 0.296 mmol) was added at 0°C, and the mixture was stirred under nitrogen atmosphere for 2 hours. The reaction mixture was concentrated at reduced pressure and diluted with ethyl acetate, the organic layer was washed with a 1 N aqueous dipotassium hydrogenphosphate solution and a brine, and the aqueous layer was separated by a phase separator and concentrated at reduced pressure to obtain methyl (2S)-2-[[[(2,3-difluorophenyl)methylamino]-methylamino]-3,3-dimethylbutanoate as a crude product. The obtained crude product was dissolved in ethyl acetate (750 μ L), and then 3-isobutoxy-3-oxopropanoate (0.026 g, 0.163 mmol), pyridine (35.8 μ L, 0.444 mmol) and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (an ethyl acetate solution, 174 μ L, 0.296 mmol) were added at 0°C, and the mixture was stirred under nitrogen atmosphere for 1 hour. Water was added to the reaction mixture, and the resultant was extracted with ethyl acetate. The organic layer was washed with a 1 N aqueous dipotassium hydrogenphosphate solution and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain methyl (2S)-2-[[[(2,3-difluorophenyl)methyl-[3-(2-methylpropoxy)-3-oxopropanoyl]amino]-methylamino]-3,3-dimethylbutanoate as a crude product. The obtained crude product was dissolved in N,N-dimethylformamide (750 μ L), cesium carbonate (121 mg, 0.370 mmol) was added, and the mixture was stirred under nitrogen atmosphere at 80°C for 2 hours. 1 N Hydrochloric acid was added to the reaction mixture at 0°C, and the resultant was extracted with ethyl acetate. The organic layer was washed with a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the

resultant residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound (48.1 mg, 79%).


LCMS: m/z 411.5[M+H]⁺

HPLC retention time: 1.03 minutes (analysis condition SQD-AA05)

Fifth Step

(3S)-3-Tert-butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyrimidine-5-carboxamide

[0994] [Formula 152]



[0995] (3S)-3-tert-Butyl-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxylic acid 2-methylpropyl ester (6.3 mg, 0.015 mmol) was dissolved in toluene (200 μ L), 4-chloro-2-(6-(trifluoromethyl)pyrimidin-4-yl)aniline (Reference Example 53) (4.2 mg, 0.015 mmol) was added, and the mixture was heated for 1 hour. The reaction mixture was concentrated and the resultant residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound (8.0 mg, 85%).

[0996] LCMS: m/z 610.3[M+H]⁺

HPLC retention time: 1.72 minutes (analysis condition QC-SMD-TFA05)

[0997] <Reference Example 83>


Methyl (S)-2-(ethylamino)-3,3-dimethylbutanoate hydrochloride

First Step

Methyl (2S)-2-[benzyl(ethyl)amino]-3,3-dimethylbutanoate

[0998]

[Formula 153]



[0999] Methyl (2S)-2-amino-3,3-dimethylbutanoate hydrochloride (5.27 g, 29.0 mmol) was suspended in dichloroethane (250 mL), and then benzaldehyde (2.93 mL, 29.0 mmol) and sodium triacetoxyhydroborate (12.3 g, 58.0 mmol) were added, and the mixture was stirred under nitrogen atmosphere at room temperature for 10 hours. The reaction mixture was diluted with ethyl acetate (200 mL) and filtered, water was added, and the resultant was extracted with ethyl acetate. A saturated sodium bicarbonate solution was added to the reaction mixture at 0°C, and the mixture was extracted with dichloromethane. The organic layer was washed with a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (2S)-2-(benzylamino)-3,3-dimethylbutanoate methyl ester (6.82 g) as a crude product. A part of the obtained crude product (2.00 g, 8.50 mmol) was dissolved in dichloromethane (42.5 mL), acetaldehyde (2.39 mL, 42.5 mmol) and sodium triacetoxyhydroborate (3.60 g, 17.0 mmol) were added, and the mixture was stirred under nitrogen atmosphere at room temperature for 24 hours. A saturated sodium bicarbonate solution was added to the reaction mixture at 0°C, and the mixture was extracted with dichloromethane. The organic layer was washed with a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (1.65 g, 88%).

[1000] LCMS: m/z 264.3[M+H]⁺


HPLC retention time: 1.14 minutes (analysis condition SMD-TFA05)

[1001] Second Step

Methyl (S)-3,3-dimethyl-2-(methylamino)butanoate hydrochloride

[1002]

[Formula 154]




[1003] Methyl (2S)-2-[benzyl(ethyl)amino]-3,3-dimethylbutanoate (1.64 g, 6.23 mmol) was dissolved in ethyl acetate (6.23 mL), palladium hydroxide-carbon (20 wt. %, 328 mg) was added, and the mixture was stirred under hydrogen atmosphere at room temperature for 6 hours. The reaction mixture was filtered, hydrochloric acid (a dioxane solution, 4.67 mL, 18.7 mmol) was added, and then the resultant was concentrated at reduced pressure to obtain the title compound (922 mg, 71%).

[1004] ¹H-NMR (DMSO-D₆) δ: 8.98 (1H, brs), 8.76 (1H, brs), 3.88 (1H, d, *J* = 10.0 Hz), 3.78 (3H, s), 2.94 (2H, q, *J* = 6.0 Hz), 1.22 (3H, t, *J* = 7.3 Hz), 1.04 (9H, s).

[1005] <Reference Example 84>

Methyl (2S)-3,3-dimethyl-2-(2,2,2-trifluoroethylamino)butanoate

[1006] [Formula 155]



[1007] Methyl (2S)-2-amino-3,3-dimethylbutanoate hydrochloride (1.00 g, 5.50 mmol) was suspended in tetrahydrofuran (250 mL), and then trifluoromethanesulfonic acid 2,2,2-trifluoroethyl (1.59 mL, 11.0 mmol) and diisopropylethylamine (4.79 mL, 27.5 mmol) were added, and the mixture was stirred under nitrogen atmosphere at 60°C for 27 hours. After the mixture was cooled to room temperature, the reaction mixture was filtered, concentrated hydrochloric acid was added, the resultant was concentrated at reduced pressure, and the resultant residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound as a solution of acetonitrile/water.

[1008] LCMS: *m/z* 228.3[M+H]⁺

HPLC retention time: 1.19 minutes (analysis condition SMD-TFA05)


[1009] <Reference Example 85>

Methyl 1-(methylamino)cyclobutanecarboxylate hydrochloride

First Step

Methyl 1-((tert-butoxycarbonyl)(methyl)amino)cyclobutanecarboxylate

[1010] [Formula 156]



[1011] 1-((tert-Butoxycarbonyl)amino)cyclobutanecarboxylic acid (1.00 g, 4.65 mmol) was dissolved in N,N-dimethylformamide (9.29 mL), and then sodium hydride (55%, 0.61 g, 13.9 mmol) and methyl iodide (1.45 mL, 23.3 mmol) were added at 0°C, and the mixture was stirred under nitrogen atmosphere at room temperature for 3 hours. A 1 N aqueous hydrochloric acid solution was added to the reaction mixture at 0°C, and the mixture was extracted with diethyl ether. The organic layer was washed with water, a saturated sodium bicarbonate solution, a 10 wt. % aqueous sodium thiosulfate solution, and a brine, the aqueous layer was separated by a phase separator, the resultant was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (1.13 g, 100%).


[1012] LCMS: m/z 266.3[M+Na]⁺

HPLC retention time: 0.92 minutes (analysis condition-SQD-AA05)

Second Step

Methyl 1-(methylamino)cyclobutanecarboxylate hydrochloride

[1013] [Formula 157]



[1014] In a similar manner to Second Step of Example 237, the title compound was synthesized from methyl 1-((tert-butoxycarbonyl)(methyl)amino)cyclobutanecarboxylate.


[1015] ¹H-NMR (DMSO-D₆) δ: 9.97 (2H, brs), 3.86 (3H, s), 2.60-2.57 (2H, m), 2.50-2.46

(5H, m), 2.06-2.03 (2H, m).

[1016] <Reference Example 86>

Methyl 1-(methylamino)cyclobutanecarboxylate 4-toluenesulfonate

[1017] [Formula 158]




[1018] Methyl 1-(methylamino)cyclobutanecarboxylate hydrochloride (1.39 g, 7.75 mmol) was dissolved in ethyl acetate (15 mL), and then 4-toluenesulfonate monohydrate (1.47 g, 7.75 mmol) was added, the mixture was concentrated at reduced pressure, ethyl acetate (15 mL) was added, and the resultant was filtered to obtain the title compound (1.96 g, 80%).

[1019] ¹H-NMR (DMSO-D₆) δ: 9.10 (2H, brs), 7.48 (2H, d, *J* = 7.9 Hz), 7.12 (2H, d, *J* = 7.9 Hz), 3.82 (3H, s), 2.56-2.31 (7H, m), 2.29 (3H, s), 2.05-1.99 (2H, m).

[1020] <Reference Example 87>

Methyl 1-(methylamino)cyclopentanecarboxylate hydrochloride

[1021] [Formula 159]




[1022] In a similar manner to First and Second Steps of Reference Example 85, the title compound was synthesized from 1-((tert-butoxycarbonyl)amino)cyclopentanecarboxylic acid.

[1023] ¹H-NMR (DMSO-D₆) δ: 9.71 (2H, brs), 3.79 (3H, s), 3.34 (3H, s), 2.20-2.13 (2H, m), 2.05-1.98 (2H, m), 1.89-1.86 (2H, m), 1.74-1.71 (2H, m).

[1024] <Reference Example 88>

Methyl 1-(methylamino)cyclopentanecarboxylate 4-toluenesulfonate

[1025] [Formula 160]



[1026] Methyl 1-(methylamino)cyclopentanecarboxylate hydrochloride (1.50 g, 7.75 mmol)


was dissolved in ethyl acetate (15 mL), and then 4-toluenesulfonate monohydrate (1.47 g, 7.75 mmol) was added, the mixture was concentrated at reduced pressure, ethyl acetate (15 mL) was added, and the resultant was filtered to obtain the title compound (2.00 g, 78%).

[1027] $^1\text{H-NMR}$ (DMSO-D_6) δ : 9.18 (2H, brs), 7.48 (2H, d, $J = 7.9$ Hz), 7.11 (2H, d, $J = 7.9$ Hz), 3.79 (3H, s), 2.57 (3H, brs), 2.29 (3H, s), 2.24-2.10 (2H, m), 1.99-1.86 (2H, m), 1.86-1.65 (2H, m).

[1028] <Reference Example 89>

Methyl 1-(methylamino)cyclohexanecarboxylate hydrochloride

[1029] [Formula 161]



[1030] In a similar manner to First and Second Steps of Reference Example 85, the title compound was synthesized from 1-((tert-butoxycarbonyl)amino)cyclohexanecarboxylic acid.

[1031] $^1\text{H-NMR}$ (DMSO-D_6) δ : 9.43 (2H, brs), 3.84 (3H, s), 2.51 (3H, s), 2.15-2.12 (2H, m), 1.77-1.73 (4H, m), 1.58-1.55 (1H, m), 1.47-1.44 (2H, m), 1.35-1.29 (1H, m).

LCMS: m/z 172.5 $[\text{M}+\text{H}]^+$

[1032] LCMS: m/z 172.5 $[\text{M}+\text{H}]^+$

HPLC retention time: 0.52 minutes (analysis condition SMD-TFA05)


[1033] <Reference Example 90>

Methyl 2-ethyl-2-(methylamino)butanoate hydrochloride

First Step

2-(((Benzyloxy)carbonyl)amino)-2-ethylbutanoate methyl ester

[1034] [Formula 162]



[1035] 2-Ethyl-2-(methoxycarbonyl)butanoate (307.0 mg, 1.76 mmol) was dissolved in toluene (11.7 mL), and then triethylamine (295 μL , 2.12 mmol) and diphenylphosphoryl azide (458 μL , 2.12 mmol) were added, and the mixture was stirred at room temperature for

1.5 hours and at 95°C for 0.5 hours. Benzyl alcohol (1.1 mL, 10.6 mmol) was added, and the mixture was stirred at 95°C overnight. The reaction mixture was concentrated at reduced pressure, diluted with diethyl ether, and the resultant was washed with 1 N hydrochloric acid, a saturated sodium bicarbonate solution, and a brine. The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (ethyl acetate/hexane) to obtain the title compound (488.1 mg, 99%).


[1036] LCMS: m/z 280[M+H]⁺

HPLC retention time: 1.11 minutes (analysis condition SMD-TFA05)

[1037] Second Step

Methyl 2-(((benzyloxy)carbonyl)(methyl)amino)-2-ethylbutanoate

[1038] [Formula 163]



[1039] After a solution of methyl 2-(((benzyloxy)carbonyl)amino)-2-ethylbutanoate (481.3 mg, 1.72 mmol) in N,N-dimethylformamide (3.4 mL) was added to sodium hydride (83 mg, 3.45 mmol) at 0°C, methyl iodide (269 μ L, 4.31 mmol) was added, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was diluted with diethyl ether and washed with 1 N hydrochloric acid, water, a saturated sodium bicarbonate solution, an aqueous sodium thiosulfate solution, and a brine. The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (ethyl acetate/hexane) to obtain the title compound (438.5 mg, 87%).

[1040] LCMS: m/z 294[M+H]⁺


HPLC retention time: 1.14 minutes (analysis condition SMD-TFA05)

[1041] Third Step

Methyl 2-ethyl-2-(methylamino)butanoate hydrochloride

[1042]

[Formula 164]



[1043] Palladium-carbon (10 wt. %, 15.8 mg) was added to a solution of methyl 2-(((benzyloxy)carbonyl)(methyl)amino)-2-ethylbutanoate (434.5 mg, 1.48 mmol) in methanol (7.4 mL), and the mixture was stirred under hydrogen atmosphere at room temperature for 1 hour. A hydrochloric acid-methanol solution (0.5 M, 14.8 mL) was added to the reaction mixture, and the resultant was filtered and concentrated at reduced pressure to obtain the title compound (281.9 mg, 97%).

[1044] $^1\text{H-NMR}$ (CD_3OD) δ : 3.92 (3H, s), 3.34 (1H, m), 2.69 (3H, s), 2.04 (4H, m), 0.99 (6H, t).

[1045] LCMS: m/z 160 $[\text{M}+\text{H}]^+$

HPLC retention time: 0.49 minutes (analysis condition SMD-TFA05)


[1046] <Reference Example 91>

4-Chloro-5-methyl-6-(trifluoromethyl)pyrimidine

First Step

5-Methyl-6-(trifluoromethyl)pyrimidin-4(3H)-one

[1047] [Formula 165]



[1048] Ethyl 4,4,4-trifluoro-2-methyl-3-oxobutanoate (200 mg, 1.01 mmol) and formamidine hydrochloride (122 mg, 1.51 mmol) were dissolved in ethanol (2.0 mL), and the mixture was stirred at room temperature for 2 hours. Sodium ethoxide (172 mg, 2.52 mmol) was added, and the mixture was stirred at 80°C for 4 hours. 1 N hydrochloric acid and water were added to the reaction mixture, and the resultant was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous sodium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was washed with diethyl ether to obtain the title compound (158 mg, 88%).

[1049] LCMS: m/z 179[M+H]⁺

HPLC retention time: 0.46 minutes (analysis condition SQD-FA05)

[1050] Second Step

4-Chloro-5-methyl-6-(trifluoromethyl)pyrimidine

[1051] [Formula 166]



[1052] Phosphoryl chloride (1.06 mL, 11.4 mmol) was added to 5-methyl-6-(trifluoromethyl)pyrimidin-4(3H)-one (135 mg, 0.758 mmol), and the mixture was stirred at 100°C for 1 hour. A saturated sodium bicarbonate solution was added to the reaction mixture, and the resultant was extracted with dichloromethane. The organic layer was washed with a brine, dried over anhydrous sodium sulfate, filtered, and concentrated at reduced pressure to obtain the title compound as a crude product.

[1053] LCMS: m/z 197[M+H]⁺

HPLC retention time: 0.95 minutes (analysis condition SMD-TFA05)


[1054] <Reference Example 92>

4,5-Dichloro-6-(trifluoromethyl)pyrimidine

First Step

5-Chloro-6-(trifluoromethyl)pyrimidin-4(3H)-one

[1055] [Formula 167]



[1056] 6-(Trifluoromethyl)pyrimidin-4(3H)-one (1.0 g, 6.1 mmol) and N-chlorosuccinimide (1.1 g, 7.9 mmol) were dissolved in N,N-dimethylformamide (5.0 mL), and the mixture was stirred at 50°C for 9 hours. The reaction mixture was diluted with ethyl acetate and washed with water and a brine. The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by C18 reverse-phase column chromatography (water/acetonitrile, 0.1% formic acid)

to obtain the title compound (368.5 mg, 31%).


[1057] LCMS: m/z 199[M+H]⁺

HPLC retention time: 0.68 minutes (analysis condition SMD-TFA05)

[1058] Second Step

4,5-Dichloro-6-(trifluoromethyl)pyrimidine

[1059] [Formula 168]



[1060] Phosphoryl chloride (1.4 mL, 15.1 mmol) was added to 5-chloro-6-(trifluoromethyl)pyrimidin-4(3H)-one (150.0 mg, 0.756 mmol), and the mixture was stirred at 100°C for 1.5 hours. Ice-water was added to the reaction mixture and extracted with dichloromethane, the organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated at reduced pressure to obtain the title compound (121.5 mg) as a crude product.


LCMS: m/z 217[M+H]⁺

HPLC retention time: 1.03 minutes (analysis condition SMD-TFA05)

[1061] <Reference Example 93>

4-Chloro-2-(6-chloropyrimidin-4-yl)aniline

[1062] [Formula 169]


















[1063] Dioxane (9.6 mL) was added to 2-amino-4-chloro-phenylboronic acid pinacol ester (608 mg, 2.4 mmol), 4,6-dichloropyridine (892 mg, 6.0 mmol), tetrakis(triphenylphosphine)palladium (138 mg, 0.12 mmol), and potassium phosphate (2.03 g, 9.6 mmol), and the mixture was stirred at 90°C for 3 hours. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic layer was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (ethyl acetate/hexane) to obtain the title compound (349.6 mg, 59%).

[1064] LCMS: m/z 240 $[M+H]^+$

HPLC retention time: 1.14 minutes (analysis condition SMD-TFA05)

[1065] The boronic acid derivatives and halides described in the following Table were used to synthesize the aniline intermediates described in the following Table by carrying out operations similar to those of Reference Example 93.


[1066] [Table 28]

Reference Example No.	Aniline	Boronic acid derivative	Halide	Aniline intermediate m/z
94				251 $[M+H]^+$
95				255 $[M+H]^+$
96				322 $[M+H]^+$
97				342 $[M+H]^+$
98				345 $[M+H]^+$

[1067] <Reference Example 99>

6-(2-Amino-5-chlorophenyl)pyrimidine-4-carbonitrile

[1068] [Formula 170]



[1069] 4-Chloro-2-(6-chloropyrimidin-4-yl)aniline was used, and operations similar to those of Reference Example 63 were carried out to synthesize the title compound.


[1070] LCMS: m/z 265[M+H]⁺

HPLC retention time: 1.16 minutes (analysis condition SMD-TFA05)

<Reference Example 100>

6-(2-Amino-5-chlorophenyl)-5-methylpyrimidine-4-carbonitrile

[1071] [Formula 171]



[1072] 4-Chloro-2-(6-chloropyrimidin-4-yl)aniline was used, and operations similar to those of Reference Example 63 were carried out to synthesize the title compound.

[1073] LCMS: m/z 245[M+H]⁺

HPLC retention time: 0.75 minutes (analysis condition SQD-FA05)

[1074] <Examples 238 to 300>

The appropriate aniline reagents of Reference Examples 94 to 100 described in Table 30, the appropriate benzaldehyde derivatives of Reference Example 3 and those known from literature or commercialized, and the appropriate amines of Reference Examples 85 to 90 and those known from literature or commercialized were used to synthesize the compounds described in the following Table by carrying out operations similar to those of Third to Fifth Steps of Example 237.

Purification condition: HPLC

Mobile phase: MeCN/water (no additive), MeCN/water (0.1% formic acid), MeCN/water (0.1% NEt₃), or CHCl₃

Column:

YMC-Actus ODS-A (100 × 20 mmI.D., S - 5 µm, 12 nm)









YMC-Actus Triart C18 (100 × 30 mmI.D., S - 5 µm, 12 nm)

YMC-Actus Triart C18 (50 × 30 mmI.D., S - 5 µm, 12 nm)

JAI JAIGEL-H (600 × 20 mmI.D., GPC)










[1075]

[Table 29-1]










Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
238		QC-SMD-TFA05	1.71	643
239		QC-SMD-TFA05	1.71	609
240		SQD-AA05	1.72	544
241		QC-SMD-TFA05	1.74	523
242		QC-SMD-TFA05	1.72	577
243		QC-SMD-TFA05	1.75	576
244		QC-SMD-TFA05	1.65	499
245		QC-SMD-TFA05	1.75	623

[1076]










[Table 29-2]

246		QC-SMD-TFA05	1.78	558
247		QC-SMD-TFA05	1.71	677
248		SQD-AA05	1.59	612
249		QC-SMD-TFA05	1.69	630
250		QC-SMD-TFA05	1.69	596
251		QC-SMD-TFA05	1.67	595
252		QC-SMD-TFA05	1.69	630
253		QC-SMD-TFA05	1.68	629
254		QC-SMD-TFA05	1.70	530

[Table 29-3]










255		QC-SMD-TFA05	1.69	594
256		QC-SMD-TFA05	1.68	593
257		QC-SMD-TFA05	1.71	528
258		QC-SMD-TFA05	1.69	561
259		QC-SMD-TFA05	1.70	507
260		SQD-AA05	1.20	642
261		SQD-AA05	1.23	608
262		QC-SMD-TFA05	1.74	521
263		QC-SMD-TFA05	1.72	575

[1078] [Table 29-4]










264		QC-SMD-TFA05	1.36	773
265		QC-SMD-TFA05	1.35	739
266		QC-SMD-TFA05	1.36	772
267		QC-SMD-TFA05	1.34	738
268		QC-SMD-TFA05	1.36	750
269		QC-SMD-TFA05	1.33	716
270		QC-SMD-TFA05	1.31	744
271		QC-SMD-TFA05	1.30	729
272		QC-SMD-TFA05	1.28	696

[1079]

[Table 29-5]

273		QC-SMD-TFA05	1.26	774
274		QC-SMD-TFA05	1.30	710
275		QC-SMD-TFA05	1.33	719
276		QC-SMD-TFA05	1.36	753
277		QC-SMD-TFA05	1.34	695
278		QC-SMD-TFA05	1.38	787
279		QC-SMD-TFA05	1.33	759
280		QC-SMD-TFA05	1.32	758
281		QC-SMD-TFA05	1.30	724

[Table 29-6]

282		QC-SMD-TFA05	1.31	756
283		QC-SMD-TFA05	1.27	728
284		QC-SMD-TFA05	1.35	791
285		QC-SMD-TFA05	1.33	771
286		QC-SMD-TFA05	1.33	737
287		QC-SMD-TFA05	1.34	770
288		QC-SMD-TFA05	1.32	736
289		QC-SMD-TFA05	1.34	748
290		QC-SMD-TFA05	1.35	785

[Table 29-7]


291		QC-SMD-TFA05	1.36	773
292		QC-SMD-TFA05	1.34	739
293		QC-SMD-TFA05	1.35	772
294		QC-SMD-TFA05	1.33	738
295		SMD-TFA05	1.35	785
296		SMD-TFA05	1.35	799
297		SMD-TFA05	1.33	751
298		SMD-TFA05	1.33	718

[1082] <Reference Example 101>

Methyl 2-(2,3-difluorobenzyl)-5-hydroxy-6,6-dimethyl-3-oxo-1-phenyl-1,2,3,6-tetrahydropyridazine-4-carboxylate

First Step

[1083] [Formula 172]




[1084] Phenylhydrazine (0.76 g, 7.1 mmol) and ethyl 2-bromo-2-methylpropionate (1.93 g, 9.9 mmol) were dissolved in N,N-diisopropylethylamine (1.00 g, 7.8 mmol), and the mixture was stirred and heated under nitrogen atmosphere at 110°C for 4 hours. The reaction mixture was cooled to room temperature, concentrated at reduced pressure, and extracted with ethyl acetate. The organic layer was washed with water and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain a crude product. Methanol (6 mL) and 2,3-difluoro-benzaldehyde (0.77 mL, 7.1 mmol) were added to the obtained crude product, and the mixture was stirred under nitrogen atmosphere at room temperature for 1 hour. The reaction mixture was concentrated at reduced pressure, and the resultant residue was purified by C-18 reverse-phase column chromatography on silica gel (0.1% formic acid water/0.1% formic acid acetonitrile) to obtain ethyl 2-(2-(2,3-difluorobenzylidene)-1-phenylhydrazinyl)-2-methylpropanoate (0.51 g, 21%) as yellow oil.

[1085] LCMS: m/z 347[M+H]⁺

HPLC retention time: 1.12 minutes (analysis condition SQD-FA05)

[1086] Second Step

[1087] [Formula 173]



[1088] Ethyl 2-(2-(2,3-difluorobenzylidene)-1-phenylhydrazinyl)-2-methylpropanoate obtained in First Step was used, and operations similar to those of Reference Example 1-2 were carried out to obtain the title compound (0.38 g, 65%) as yellow oil.

[1089] LCMS: m/z 403[M+H]⁺


HPLC retention time: 0.97 minutes (analysis condition SQD-FA05)

[1090] <Reference Example 102>

Methyl 6-(2,3-difluorobenzyl)-9-hydroxy-5-methyl-7-oxo-5,6-diazaspiro[3.5]non-8-ene-8-carboxylate

First Step

[1091] [Formula 174]



[1092] Methylhydrazine (0.10 g, 2.17 mmol) and ethyl 1-bromocyclobutanecarboxylate (0.23 g, 1.1 mmol) were dissolved in N,N-diisopropylethylamine (0.30 g, 2.3 mmol), and the mixture was stirred and heated under nitrogen atmosphere at 110°C for 15 hours. The reaction mixture was cooled to room temperature, filtered, concentrated at reduced pressure, and extracted with ethyl acetate. The organic layer was washed with water and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain a crude product. The obtained crude product was added to the reaction mixture, and then methanol (2 mL) and 2,3-difluoro-benzaldehyde (0.12 mL, 1.1 mmol) were added, and the mixture was stirred under nitrogen atmosphere at room temperature for 1 hour. The reaction mixture was concentrated at reduced pressure, and the resultant residue was purified by C-18 reverse-phase column chromatography on silica gel (0.1% formic acid water/0.1% formic acid acetonitrile) to obtain ethyl 1-(2-(2,3-difluorobenzylidene)-1-methylhydrazinyl)cyclobutanecarboxylate (36 mg, 11%) as colorless oil.


[1093] LCMS: m/z 297[M+H]⁺

HPLC retention time: 1.15 minutes (analysis condition SQD-FA05)

[1094] Second Step

[1095]

[Formula 175]



[1096] Ethyl 1-(2-(2,3-difluorobenzylidene)-1-methylhydrazinyl)cyclobutanecarboxylate obtained in First Step was used, and operations similar to those of Second Step of Reference Example 101 were carried out to obtain the title compound (50 mg, 78%) as red oil.

[1097] LCMS: m/z 353[M+H]⁺


HPLC retention time: 0.93 minutes (analysis condition SQD-FA05)

[1098] <Reference Example 103>

Methyl 2-(2,3-difluorobenzyl)-5-hydroxy-1,6,6-trimethyl-3-oxo-1,2,3,6-tetrahydropyridazine-4-carboxylate

First Step

[1099] [Formula 176]




[1100] Methylhydrazine (0.46 g, 10.0 mmol) and ethyl 2-bromo-2-methylpropionate (1.07 g, 5.5 mmol) were dissolved in tetrahydrofuran (5.0 mL), and then N,N-diisopropylethylamine (0.84 g, 6.5 mmol) was added, and the mixture was stirred and heated under nitrogen atmosphere at 60°C for 66 hours. After the reaction mixture was cooled to room temperature, methanol (60 mL) and 2,3-difluoro-benzaldehyde (0.55 mL, 5.0 mmol) were added to the reaction mixture, and the mixture was stirred under nitrogen atmosphere at room temperature for 1 hour. The reaction mixture was concentrated at reduced pressure, and the resultant residue was purified by C-18 reverse-phase column chromatography on silica gel (0.1% formic acid water/0.1% formic acid acetonitrile) to obtain ethyl 2-[2-(2,3-difluorobenzylidene)-1-methylhydrazinyl]-2-methylpropionate (0.8 g, 56%) as colorless oil.

[1101] LCMS: m/z 285[M+H]⁺

HPLC retention time: 1.00 minute (analysis condition SQD-FA05)

[1102] Second Step

[1103] [Formula 177]



[1104] Ethyl 2-[2-(2,3-difluorobenzylidene)-1-methylhydrazinyl]-2-methylpropionate obtained in First Step was used, and operations similar to those of Second Step of Reference Example 101 were carried out to obtain the title compound (650 mg, 68%) as yellow oil.








[1105] LCMS: m/z 341[M+H]⁺

HPLC retention time: 0.81 minutes (analysis condition SQD-FA05)

[1106] The appropriate aniline reagents and appropriate ester intermediates of Reference Examples 101 to 103 were used to synthesize the compounds described in the following Table by carrying out operations similar to those of Fifth Step of Example 237.








[1107]

[Table 30-1]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
299		QC-SMD-TFA05	1.73	678
300		QC-SMD-TFA05	1.68	628
301		QC-SMD-TFA05	1.67	627
302		QC-SMD-TFA05	1.56	602
303		QC-SMD-TFA05	1.58	573
304		QC-SMD-TFA05	1.59	616
305		QC-SMD-TFA05	1.63	615

[1108]

[Table 30-2]

306		QC-SMD-TFA05	1.58	616
307		QC-SMD-TFA05	1.63	615
308		QC-SMD-TFA05	1.62	653
309		QC-SMD-TFA05	1.63	581
310		QC-SMD-TFA05	1.65	593
311		QC-SMD-TFA05	1.59	587
312		QC-SMD-TFA05	1.62	607


[1109] <Example 313>

2-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-5-hydroxy-1,6,6-trimethyl-3-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]pyridazine-4-carboxamide

First Step

[1110]

[Formula 178]




[1111] Methylhydrazine, ethyl 2-bromo-2-methylpropanoate, and the aldehyde reagent synthesized in Reference Example 3 were used, and operations similar to those of First Step of Reference Example 103 were carried out to obtain ethyl 2-(2-(2,3-difluoro-4-(2-morpholinoethoxy)benzylidene)-1-methylhydrazinyl)-2-methylpropanoate.

[1112] LCMS: m/z 414 $[M+H]^+$

HPLC retention time: 0.96 minutes (analysis condition SMD-TFA05)

[1113] Second Step

[1114] [Formula 179]




[1115] Ethyl 2-(2-(2,3-difluoro-4-(2-morpholinoethoxy)benzylidene)-1-methylhydrazinyl)-2-methylpropanoate (50 mg, 0.12 mmol) obtained in First Step was dissolved in a 10% hydrochloric acid-methanol solution (1.8 mL), borane-pyridine (28 mg, 0.30 mmol) was added at 0°C, and the mixture was stirred for 1.5 hours. The reaction mixture was concentrated at reduced pressure, a 1 N aqueous dipotassium hydrogenphosphate solution was added, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, dried over anhydrous sodium sulfate and filtered, and the solvent was distilled away at reduced pressure to obtain ethyl 2-(2-(2,3-difluoro-4-(2-morpholinoethoxy)benzyl)-1-methylhydrazinyl)-2-methylpropanoate as a crude product.

[1116] LCMS: m/z 416 $[M+H]^+$

HPLC retention time: 0.63 minutes (analysis condition SMD-TFA05)

[1117] Third Step

[1118] [Formula 180]




[1119] The crude product of ethyl 2-(2-(2,3-difluoro-4-(2-morpholinoethoxy)benzyl)-1-methylhydrazinyl)-2-methylpropanoate obtained in Second Step was dissolved in ethyl acetate (800 μ L), and then the carboxylic acid reagent synthesized in First Step of Reference Example 82, pyridine (20 μ L, 0.242 mmol), and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (an ethyl acetate solution, 92 μ L, 0.145 mmol) were added at 0°C, and the mixture was stirred under nitrogen atmosphere overnight. Water was added to the reaction mixture, and the resultant was extracted with ethyl acetate. The organic layer was washed with water and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain ethyl 2-(2-(2,3-difluoro-4-(2-morpholinoethoxy)benzyl)-1-methyl-2-(3-oxo-3-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)amino)propanoyl)hydrazinyl)-2-methylpropanoate as a crude product.

[1120] LCMS: m/z 791[M+H]⁺

HPLC retention time: 1.19 minutes (analysis condition SMD-TFA05)

[1121] Fourth Step

[1122] [Formula 181]



[1123] Ethyl 2-(2-(2,3-difluoro-4-(2-morpholinoethoxy)benzyl)-1-methyl-2-(3-oxo-3-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)amino)propanoyl)hydrazinyl)-2-methylpropanoate obtained in Third Step was used, and operations similar to those of Third

Step of Reference Example 1-1 were carried out to synthesize the title compound (35.1 mg, 39% three-step yield).


[1124] LCMS: m/z 745[M+H]⁺

HPLC retention time: 1.27 minutes (analysis condition QC-SMD-TFA05)

[1125] <Example 314>

2-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-5-hydroxy-1,6,6-trimethyl-3-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]pyridazine-4-carboxamide

[1126] [Formula 182]



[1127] Ethyl 2-(2-(2,3-difluoro-4-(2-morpholinoethoxy)benzyl)-1-methylhydrazinyl)-2-methylpropanoate synthesized in Second Step of Example 313 and the carboxylic acid reagent synthesized in First Step of Reference Example 81 were used, and operations similar to those of Example 313 were carried out to synthesize the title compound (36.6 mg, 41% three-step yield).

[1128] LCMS: m/z 744[M+H]⁺

HPLC retention time: 1.28 minutes (analysis condition QC-SMD-TFA05)


[1129] <Example 315>

2-[[2,3-Difluoro-4-(3-morpholin-4-ylprop-1-ynyl)phenyl]methyl]-5-hydroxy-1,6,6-trimethyl-3-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]pyridazine-4-carboxamide

First Step

[1130]

[Formula 183]




[1131] 2,3-Difluoro-4-iodobenzaldehyde and 3-isobutyloxy-3-oxopropionate were used, and operations similar to those of First and Second Steps of Reference Example 103 were carried out to synthesize isobutyl 2-(2,3-difluoro-4-iodobenzyl)-5-hydroxy-1,6,6-trimethyl-3-oxo-1,2,3,6-tetrahydropyridazine-4-carboxylate.

[1132] LCMS: m/z 509[M+H]⁺

HPLC retention time: 1.10 minutes (analysis condition SQD-FA05)

[1133] Second Step

[1134] [Formula 184]



[1135] (2-(2,3-Difluoro-4-iodobenzyl)-5-hydroxy-1,6,6-trimethyl-3-oxo-1,2,3,6-tetrahydropyridazine-4-carboxylic acid isobutyl ester synthesized in First Step and 4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)aniline (Reference Example 50) were used, and operations similar to those of Fifth Step of Example 237 were carried out to synthesize 2-(2,3-difluoro-4-iodobenzyl)-5-hydroxy-1,6,6-trimethyl-3-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-1,2,3,6-tetrahydropyridazine-4-carboxamide.


[1136] LCMS: m/z 742[M+H]⁺

HPLC retention time: 1.25 minutes (analysis condition SQD-FA05)

[1137] Third Step

[1138]

[Formula 185]



[1139] The iodine body synthesized in Second Step and 4-(prop-2-yn-1-yl)morpholine were used, and operations similar to those of Example 123 were carried out to synthesize the title compound.


[1140] LCMS: m/z 739[M+H]⁺

HPLC retention time: 1.31 minutes (analysis condition QC-SMD-TFA05)

[1141] <Example 316>

N-[4-Chloro-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-2-[[2,3-difluoro-4-(3-morpholin-4-ylprop-1-yn-1-yl)phenyl]methyl]-5-hydroxy-1,6,6-trimethyl-3-oxopyridazine-4-carboxamide

[1142] [Formula 186]



[1143] The aniline reagent synthesized in Reference Example 53 was used, and operations similar to those of Example 315 were carried out to synthesize the title compound.

[1144] LCMS: m/z 705[M+H]⁺

HPLC retention time: 1.29 minutes (analysis condition QC-SMD-TFA05)


[1145] <Example 317>

6-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

First Step

Methyl 1-[[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methylideneamino]-methylamino]cyclobutane-1-carboxylate

[1146] [Formula 187]



[1147] Methyl 1-(methylamino)cyclobutanecarboxylate hydrochloride (Reference Example 85) (300 mg, 1.67 mmol) was dissolved in acetic acid (16.7 mmol, 956 μ L) and water (956 μ L), and then sodium nitrite (132 mg, 1.91 mmol) was added at 0°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. Zinc (1.09 g, 16.7 mmol) was added to the reaction mixture at 0°C, and the mixture was stirred under nitrogen atmosphere for 1.5 hours. After the reaction mixture was filtered, methanol (9.56 mL) and 2,3-difluoro-4-(2-morpholinoethoxy)benzaldehyde (324 mg, 1.19 mmol) were added, and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated at reduced pressure and diluted with ethyl acetate, the organic layer was washed with a 1 M aqueous dipotassium hydrogenphosphate solution and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (aminosilica, hexane/ethyl acetate) to obtain the title compound (402 mg, 82%).

[1148] LCMS: m/z 412[M+H]⁺


HPLC retention time: 0.91 minutes (analysis condition SMD-TFA05)

[1149] Second Step

Methyl 1-[[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl-[3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoyl]amino]-methylamino]cyclobutane-1-carboxylate

[1150]

[Formula 188]



[1151] Methyl 1-[[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methylideneamino]-methylamino]cyclobutane-1-carboxylate (27.5 g, 66.9 mmol) was dissolved in ethyl acetate (158 mL), and then hydrochloric acid (a 4 M ethyl acetate solution, 158 mL) was added at 0°C, and the mixture was stirred under nitrogen atmosphere for 10 minutes. 5-Ethyl-2-methylpyridine borane (19.9 mL, 134 mmol) was added to the reaction mixture at 0°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. After an aqueous sodium hydroxide solution (5 M, 115 mL) and an aqueous potassium phosphate solution (1 M, 50.0 mL) were added to the reaction mixture, the mixture was diluted with ethyl acetate, the organic layer was washed with a 1 M aqueous dipotassium hydrogenphosphate solution and a brine, dried over magnesium sulfate, filtered, and concentrated at reduced pressure to obtain methyl 1-[[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methylamino]-methylamino]cyclobutane-1-carboxylate as a crude product. The obtained crude product and 3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoate (First Step of Reference Example 82) (27.6 g, 70.2 mmol) were dissolved in ethyl acetate (279 mL) and N,N-dimethylformamide (139 mL), and then pyridine (5.40 mL, 66.9 mmol) and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (a 1.7 M ethyl acetate solution, 79.0 mL, 134 mmol) were added at 0°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. A brine was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a 1 M aqueous dipotassium hydrogenphosphate solution and a brine, dried over magnesium sulfate, filtered, and concentrated at reduced pressure to obtain the title compound (1.15 g, 100%).


[1152] LCMS: m/z 789.1[M+H]⁺

HPLC retention time: 1.01 minutes (analysis condition SMD-FA05)

[1153] Third Step

6-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1154] [Formula 189]



[1155] Methyl 1-[[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl-[3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoyl]amino]-methylamino]cyclobutane-1-carboxylate (1.14 g, 1.45 mmol) was dissolved in methanol (7.27 mL) and N,N-dimethylformamide (7.27 mL), and then potassium carbonate (602 mg, 4.36 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 1.5 hours. Formic acid and water were added to the reaction mixture, and the mixture was purified by C18 reverse-phase column chromatography (methanol/water) to obtain the title compound (840 mg, 76%).

[1156] LCMS: m/z 757.4[M+H]⁺

HPLC retention time: 1.27 minutes (analysis condition SMD-TFA05)

[1157] Although tautomers of some of the title compounds exist, isomers may be observed in some cases and not in other cases according to the type of the solvent for measurement. For example, ¹H-NMR and ¹³C-NMR of major tautomers in chloroform-D are as follows.

[1158] <Example 317: Major tautomer>

¹H-NMR (CDCl₃) δ : 16.58 (1H, s), 12.81 (1H, s), 9.60 (1H, s), 8.50 (1H, d, J = 8.6 Hz), 7.95 (1H, s), 7.90 (1H, d, J = 1.6 Hz), 7.80 (1H, dd, J = 8.6, 1.6 Hz), 7.02 (1H, ddd, J = 8.6, J_{CF} = 7.2, 1.6 Hz), 6.72 (1H, ddd, J = 8.6, J_{CF} = 7.2, 1.6 Hz), 5.06 (1H, brs), 4.20 (1H, brs), 4.20 (2H, t, J = 5.5 Hz), 3.74 (4H, t, J = 4.1 Hz), 2.85 (2H, brs), 2.61 (4H, brs), 2.52 (1H, m), 2.41 (3H, s), 1.85-1.83 (3H, m), 1.73 (1H, m), 1.60 (1H, m).

[1159] ^{13}C -NMR (CDCl_3) δ : 186.1 (qC), 169.8 (qC), 165.7 (qC), 162.8 (qC), 159.3 (qC), 156.6 (qC, q, $J = 36.4$ Hz), 150.4 (qC, dd, $J = 248.7, 10.5$ Hz), 147.8 (qC, d, $J = 5.5$ Hz), 141.3 (qC, dd, $J = 248.1, 14.6$ Hz), 138.8 (qC), 128.3 (CH, q, $J = 3.3$ Hz), 127.7 (qC), 127.1 (CH), 127.0 (qC, q, $J = 33.4$ Hz), 125.1 (CH), 124.7 (CH), 123.6 (qC, q, $J = 271.8$ Hz), 120.5 (qC, q, $J = 275.4$ Hz), 118.2 (qC, d, $J = 12.4$ Hz), 115.9 (CH, q, $J = 2.5$ Hz), 109.6 (CH, d, $J = 1.9$ Hz), 92.3 (qC), 67.9 (CH_2), 66.9 ($\text{CH}_2 \times 2$), 64.2 (qC), 57.5 (CH_2), 54.1 ($\text{CH}_2 \times 2$), 41.6 (CH_2), 35.1 (CH_3), 32.4 (CH_2), 24.5 (CH_2), 13.4 (CH_2).


[1160] <Example 318>

7-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

First Step

Methyl 1-[[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate

[1161] [Formula 190]



[1162] Methyl 1-(methylamino)cyclopentanecarboxylate hydrochloride (300 mg, 1.55 mmol) was dissolved in acetic acid (15.5 mmol, 887 μL) and water (887 μL), and then sodium nitrite (122 mg, 1.77 mmol) was added at 0°C , and the mixture was stirred under nitrogen atmosphere for 1 hour. Zinc (1.01 g, 15.5 mmol) was added to the reaction mixture at 0°C , and the mixture was stirred under nitrogen atmosphere for 1.5 hours. After the reaction mixture was filtered, methanol (8.87 mL) and 2,3-difluoro-4-(2-morpholinoethoxy)benzaldehyde (300 mg, 1.11 mmol) were added, and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated at reduced pressure and diluted with ethyl acetate, the organic layer was washed with a 1 M

aqueous dipotassium hydrogenphosphate solution and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (aminosilica, hexane/ethyl acetate) to obtain the title compound (375 mg, 80%).


[1163] LCMS: m/z 426.2[M+H]⁺

HPLC retention time: 0.93 minutes (analysis condition SMD-TFA05)

[1164] Second Step

1-[[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-[3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoyl]amino]-methylamino]cyclopentane-1-carboxylic acid

[1165] [Formula 191]



[1166] Methyl 1-[[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate (29.8 g, 70.3 mmol) was dissolved in ethyl acetate (172 mL), and then hydrochloric acid (a 4 M ethyl acetate solution, 172 mL) was added at 0°C, and the mixture was stirred under nitrogen atmosphere for 10 minutes. 5-Ethyl-2-methylpyridine borane (20.9 mL, 141 mmol) was added to the reaction mixture at 0°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. After an aqueous sodium hydroxide solution (5 M, 125 mL) and an aqueous potassium phosphate solution (1 M, 50.0 mL) were added to the reaction mixture, the mixture was diluted with ethyl acetate, the organic layer was washed with a 1 M aqueous potassium phosphate solution and a brine, dried over magnesium sulfate, filtered, and concentrated at reduced pressure to obtain methyl 1-[[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methylamino]-methylamino]cyclopentane-1-carboxylate as a crude product.

[1167] The obtained crude product and 3-oxo-3-[4-(trifluoromethyl)-2-[6-

(trifluoromethyl)pyrimidin-4-yl]anilino]propanoate (First Step of Reference Example 82) (29.0 g, 73.8 mmol) were dissolved in ethyl acetate (293 mL) and N,N-dimethylformamide (146 mL), and then pyridine (5.67 mL, 70.3 mmol) and 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (a 1.7 M ethyl acetate solution, 83.0 mL, 141 mmol) were added at 0°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. A brine was added to the reaction mixture, and the resultant was extracted with ethyl acetate. The organic layer was washed with a 1 M aqueous dipotassium hydrogenphosphate solution and a brine, dried over magnesium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was washed with methanol to obtain the title compound (39.4 g, 70%).


[1168] LCMS: m/z 803.1[M+H]⁺

HPLC retention time: 1.03 minutes (analysis condition SMD-FA05)

Third Step

7-[[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1169] [Formula 192]



[1170] 1-[[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-3-oxo-3-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]anilino]propanoyl]amino]-methylamino]cyclopentane-1-carboxylic acid (15.0 g, 18.7 mmol) was dissolved in methanol (94.0 mL) and N,N-dimethylformamide (94.0 mL), and then potassium carbonate (7.76 g, 56.1 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 1.5 hours. Formic acid and water were added to the reaction mixture, and the mixture was purified by C18 reverse-phase column chromatography (methanol/water) to obtain the title compound (13.3 g, 92%).

[1171] LCMS: m/z 771.3[M+H]⁺

HPLC retention time: 1.29 minutes (analysis condition SMD-TFA05)

Although tautomers of some of the title compounds exist, isomers may be observed in some cases and not in other cases according to the type of the solvent for measurement.

For example, ¹H-NMR and ¹³C-NMR of major tautomers in chloroform-D are as follows.

[1172] <Example 318: Major tautomer>


¹H-NMR (CDCl₃) δ: 16.55 (1H, s), 12.83 (1H, s), 9.62 (1H, s), 8.49 (1H, d, J = 8.8 Hz), 7.96 (1H, s), 7.90 (1H, d, J = 1.5 Hz), 7.79 (1H, dd, J = 8.8, 1.5 Hz), 7.05 (1H, dd, J = 8.1, 7.4 Hz), 6.73 (1H, dd, J = 8.1, 7.4 Hz), 5.05 (1H, d, J = 14.2 Hz), 4.21 (1H, d, J = 14.2 Hz), 4.19 (2H, t, J = 5.7 Hz), 3.74 (4H, t, J = 4.6 Hz), 2.84 (2H, t, J = 5.7 Hz), 2.60 (4H, m), 2.48 (3H, s), 2.16 (1H, m), 1.74 (2H, m), 1.59 (1H, m), 1.52 (1H, m), 1.47 (2H, m), 1.31 (2H, m).

[1173] ¹³C-NMR (CDCl₃) δ: 187.7 (qC), 169.9 (qC), 165.7 (qC), 163.2 (qC), 159.3 (CH), 156.6 (qC, q, J_{CF} = 36.3 Hz), 150.4 (qC, dd, J_{CF} = 248.7, 10.5 Hz), 147.9 (qC, d, J_{CF} = 5.8 Hz), 141.3 (qC, dd, J_{CF} = 248.3, 14.7 Hz), 138.8 (qC), 128.3 (CH, q, J_{CF} = 3.3 Hz), 127.8 (qC), 127.1 (CH, q, J_{CF} = 3.9 Hz), 126.9 (qC, q, J_{CF} = 33.4 Hz), 125.5 (CH), 124.7 (CH), 123.6 (qC, q, J_{CF} = 272.1 Hz), 120.5 (qC, q, J_{CF} = 275.4 Hz), 117.8 (qC, d, J_{CF} = 12.9 Hz), 116.0 (CH, q, J_{CF} = 2.5 Hz), 109.7 (CH), 93.1 (qC), 71.4 (qC), 68.1 (CH₂), 66.9 (CH₂×2), 57.5 (CH₂), 54.2 (CH₂×2), 41.7 (CH₂), 37.6 (CH₂), 36.8 (CH₂), 31.4 (CH₂), 24.8 (CH₂), 23.3 (CH₂).

[1174] <Example 319>

(4aR)-N-(4-Bromo-3-fluorophenyl)-1-[(2,3-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide


[1175] [Formula 193]



[1176] First Step

[1177]


[Formula 194]



[1178] (R)-2-Methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride and 2,3-difluorobenzaldehyde were used, and then (R)-1-((2,3-difluoro-4-iodobenzylidene)amino)-2-methylpyrrolidine-2-carboxylic acid methyl ester (2.24 g, 5.49 mmol) obtained by carrying out operations similar to those of Example 1 was dissolved in acetic acid (5.0 mL) and methanol (5.5 mL), and then sodium cyanoborohydride (1.76 g, 28.0 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 14 hours. A saturated sodium bicarbonate solution was added to the reaction mixture, and the resultant was concentrated at reduced pressure and extracted with ethyl acetate. The organic layer was washed with a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (R)-1-((2,3-difluoro-4-iodobenzyl)amino)-2-methylpyrrolidine-2-carboxylic acid methyl ester as a crude product.

[1179] Second Step

[1180] [Formula 195]




[1181] The crude product obtained in First Step was dissolved in tetrahydrofuran (5.5 mL), and then tripotassium phosphate (2.38 g, 11.2 mmol) and chlorocarbonyl-acetic acid methyl ester (1.00 mL, 9.06 mmol) were added at 0°C, and the mixture was stirred under nitrogen atmosphere at room temperature for 3 hours. 1 N hydrochloric acid was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a saturated sodium bicarbonate solution and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain (S)-1-[(3-fluoro-benzyl)-(2-

methoxycarbonyl-acetyl)-amino]-pyrrolidine-2-carboxylic acid methyl ester as a crude product.

[1182] Third Step

[1183] [Formula 196]



[1184] The obtained crude product was dissolved in N,N-dimethylformamide (1.5 mL), and then cesium carbonate (672 mg, 2.06 mmol) was added, and the mixture was stirred under nitrogen atmosphere at 80°C for 5.5 hours. 1 N hydrochloric acid was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure to obtain methyl (4aR)-1-[(2,3-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylate as a crude product.

[1185] Fourth Step

The crude product (328 mg, 0.686 mmol) obtained in Third Step and 4-bromo-3-fluoroaniline (145 mg, 0.763 mmol) were dissolved in toluene (3.4 mL), and the mixture was stirred at 110°C for 1 hour. After the reaction mixture was left to cool, it was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (402 mg, 92%) as a grayish white solid.

[1186] LCMS: m/z 479[M+H]⁺


HPLC retention time: 0.97 minutes (analysis condition SQD-FA05)

[1187] <Example 320>

(4aR)-1-[(2,3-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1188]

[Formula 197]



[1189] (4aR)-1-[(2,3-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (Reference Example 5) (100 mg, 0.192 mmol) and 4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyridin-3-yl)aniline (Reference Example 42) (73.6 mg, 0.240 mmol) were dissolved in toluene (0.96 mL), and the mixture was stirred at 100°C for 40 minutes. After the reaction mixture was left to cool, it was concentrated at reduced pressure, and the resultant residue was purified by C18 reverse-phase column chromatography to obtain the title compound (120 mg, 83%) as a white amorphous solid.


[1190] LCMS: m/z 753[M+H]⁺

HPLC retention time: 0.93 minutes (analysis condition SQD-FA50)

[1191] <Example 321>

(4aR)-1-[(2,3-Difluoro-4-iodophenyl)methyl]-N-[2-(2,3-dimethoxyphenyl)-4-(trifluoromethyl)phenyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1192] [Formula 198]



[1193] (4aR)-1-[(2,3-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (Reference Example 5) and 2',3'-dimethoxy-5-(trifluoromethyl)-[1,1'-biphenyl]-2-amine (Reference Example 16) were used, and operations similar to those of Fourth Step of Example 319 were

carried out to synthesize the title compound.


[1194] LCMS: m/z 744 $[M+H]^+$

HPLC retention time: 1.19 minutes (SMD-TFA05)

[1195] <Example 322>


(4aR)-1-[(2,6-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1196] [Formula 199]



[1197] First Step

[1198] [Formula 200]



[1199] 2,6-Difluoro-4-iodobenzaldehyde was used, and operations similar to those of First to Third Steps of Example 319 were carried out to synthesize (4aR)-1-[(2,6-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester.

[1200] LCMS: m/z 521 $[M+H]^+$

HPLC retention time: 1.51 minutes (analysis condition SMD-TFA05)

[1201] Second Step

(4aR)-1-[(2,6-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester and 4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyridin-3-yl)aniline (Reference Example 42) were

used, and operations similar to those of Fourth Step of Example 319 were carried out to synthesize the title compound.


[1202] LCMS: m/z 753[M+H]⁺

HPLC retention time: 1.76 minutes (analysis condition QC-SMD-TFA05)

[1203] <Example 323>

(4aR)-1-[(2,3-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1204] [Formula 201]



[1205] (4aR)-1-[(2,3-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (Reference Example 5) and 4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)aniline (Reference Example 50) were used, and operations similar to those of Fourth Step of Example 319 were carried out to synthesize the title compound.

[1206] LCMS: m/z 754[M+H]⁺


HPLC retention time: 0.93 minutes (analysis condition SQD-FA50)

[1207] <Example 324>

(4aR)-1-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1208]

[Formula 202]



[1209] (4aR)-1-[(2,3-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (Reference Example 5) (1.086 g, 1.444 mmol), copper iodide (31.2 mg, 0.164 mmol), and 2-methyl-8-quinolinol (52.7 mg, 0.331 mmol) were dissolved in dimethylsulfoxide (4.1 mL) under nitrogen atmosphere, tetrabutylammonium hydroxide (4.24 g, 6.54 mmol) and water (1.56 mL) were added, and the mixture was stirred under nitrogen atmosphere at 100°C for 5 hours. After the reaction mixture was left to cool, 1 N hydrochloric acid (5.1 mL), water (30 mL), and an appropriate amount of N-acetyl-L-cysteine were added to the reaction mixture, and the resultant was extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel to obtain the title compound (834 mg, 90%) as a pale yellow amorphous solid.

[1210] LCMS: m/z 643[M+H]⁺


HPLC retention time: 1.12 minutes (analysis condition SQD-FA05)

[1211] <Example 325>

(4aR)-1-[2,3-(Difluoro-4-hydroxyphenyl)methyl]-N-[2-(2,3-dimethoxyphenyl)-4-(trifluoromethyl)phenyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1212]

[Formula 203]



[1213] (4aR)-1-[2,3-(Difluoro-4-iodophenyl)methyl]-N-[2-(2,3-dimethoxyphenyl)-4-(trifluoromethyl)phenyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 321) was used, and operations similar to those of Example 324 were carried out to synthesize the title compound.


[1214] LCMS: m/z 634[M+H]⁺

HPLC retention time: 1.14 minutes (analysis condition SQD-FA05)

[1215] <Example 326>

(4aR)-1-[(2,6-Difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1216] [Formula 204]



[1217] (4aR)-1-[(2,6-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 322) was used, and operations similar to those of Example 324 were carried out to synthesize the title compound.


[1218] LCMS: m/z 643[M+H]⁺

HPLC retention time: 1.12 minutes (analysis condition SQD-FA05)

[1219] <Example 327>

(4aR)-1-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1220] [Formula 205]



[1221] (4aR)-1-[(2,3-Difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide (Example 323) was used, and operations similar to those of Example 324 were carried out to synthesize the title compound.


[1222] LCMS: m/z 644[M+H]⁺

HPLC retention time: 1.11 minutes (analysis condition SQD-FA05)

[1223] <Example 328>

(4aR)-1-[(2,6-Dichloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide


[1224] [Formula 206]



[1225] First Step

[1226]

[Formula 207]



[1227] Methyl (R)-2-methylpyrrolidine-2-carboxylate hydrochloride (20 g, 111 mmol) was suspended in dichloromethane (100 mL), and then p-toluenesulfonic acid-monohydrate (25 g, 145 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 10 minutes. The reaction mixture was concentrated at reduced pressure, toluene was added for azeotropic removal, and then the residue was suspended in dichloromethane (250 mL), sodium nitrite (11.4 g, 165 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 2 hours. The reaction mixture was filtered and then concentrated at reduced pressure to obtain (S)-1-nitroso-pyrrolidine-2-carboxylic acid methyl ester as a crude product. The obtained crude product (362 mg, 2.10 mmol) was dissolved in acetic acid (10 mL) and methanol (1.0 mL), and then zinc (725 mg, 11.2 mmol) was added at 9°C, and the mixture was stirred under nitrogen atmosphere at 9°C for 1 hour. After the reaction mixture was filtered through a celite pad, the filtrate was concentrated at reduced pressure. Methanol (1.0 mL) and 2,4-dichloro-4-hydroxybenzaldehyde (200 mg, 1.05 mmol) were added to the resultant residue, and the mixture was stirred under nitrogen atmosphere at room temperature for 2 hours. The reaction mixture was filtered and then concentrated at reduced pressure, and the resultant was extracted with ethyl acetate. The organic layer was washed with water and a brine, dried over sodium sulfate, filtered, and concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain methyl (R)-1-((2,6-dichloro-4-hydroxybenzylidene)amino)-2-methylpyrrolidine-2-carboxylate. The obtained product (5.3 g, 16.0 mmol) was dissolved in acetonitrile (50 mL), and then 2-(chloromethoxy)ethyl trimethylsilane (3.2 g, 19.2 mmol) and cesium carbonate (6.3 g, 19.3 mmol) were added, and the mixture was stirred under nitrogen atmosphere at 25°C for


24 hours. Water (100 mL) was added to the reaction mixture, and the resultant was extracted with ethyl acetate (100 mL) three times. The organic layer was washed with water and a brine, and then dried over sodium sulfate. The sodium sulfate was filtered off and the resultant was concentrated at reduced pressure to obtain a crude product of methyl (R)-1-((2,6-dichloro-4-((2-(trimethylsilyl)ethoxy)methoxy)benzylidene)amino)-2-methylpyrrolidine-2-carboxylate as a white solid. The obtained crude product was used to obtain methyl (R)-1-N-(2,6-dichloro-4-((2-(trimethylsilyl)ethoxy)methoxy)benzyl)-3-isobutoxy-3-oxopropanamide)-2-methylpyrrolidine-2-carboxylate by carrying out First and Second Steps of Reference Example 1-1.

[1228] LCMS: m/z 461 $[M+H]^+$

HPLC retention time: 4.73 minutes (analysis condition Ph-SMD-TFA05)

[1229] Second Step

[1230] [Formula 208]



[1231] Methyl (R)-1-N-(2,6-dichloro-4-((2-(trimethylsilyl)ethoxy)methoxy)benzyl)-3-isobutoxy-3-oxopropanamide)-2-methylpyrrolidine-2-carboxylate (7.02 g, 11.6 mmol) was dissolved in dichloromethane (85 mL), trifluoroacetic acid (35 mL) was added, and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain methyl (R)-1-N-(2,6-dichloro-4-hydroxybenzyl)-3-isobutoxy-3-oxopropanamide)-2-methylpyrrolidine-2-carboxylate (5.0 g, 91%). The obtained methyl (R)-1-N-(2,6-dichloro-4-hydroxybenzyl)-3-isobutoxy-3-oxopropanamide)-2-methylpyrrolidine-2-carboxylate (5.0 g, 10.5 mmol) was used, and operations similar to those of Third Step of Reference Example 1-1 were carried out to obtain (4aR)-1-[(2,6-dichloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester.

[1232] LCMS: m/z 443[M+H]⁺

HPLC retention time: 1.32 minutes (analysis condition SMD-TFA05)

[1233] Third Step

(4aR)-1-[(2,6-Dichloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester and 2-(trifluoromethyl)-4-(6-(trifluoromethyl)pyridin-3-yl)pyrimidin-5-amine (Reference Example 70) were used, and operations similar to those of Example 21 were carried out to obtain the title compound.


[1234] LCMS: m/z 677[M+H]⁺

HPLC retention time: 1.51 minutes (analysis condition SMD-TFA05)

[1235] <Example 329>

(4aR)-1-[2,6-(Dichloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[6-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]pyridin-3-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1236] [Formula 209]



[1237] (4aR)-1-[(2,6-Dichloro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylic acid 2-methylpropyl ester (Second Step of Example 328) and 6,6'-bis(trifluoromethyl)-[2,3'-bipyridine]-3-amine (Reference Example 44) were used, and operations similar to those of Third Step of Example 328 were carried out to obtain the title compound.

[1238] LCMS: m/z 676[M+H]⁺


HPLC retention time: 1.55 minutes (analysis condition SMD-TFA05)

[1239] <Example 330>

(4aR)-1-[2,3-(Difluoro-4-hydroxyphenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-


(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1240] [Formula 210]



[1241] First Step

[1242] [Formula 211]



[1243] Methyl (4aR)-1-[(2,3-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylate (Third Step of Example 319) and 2-(trifluoromethyl)-4-(6-(trifluoromethyl)pyridin-3-yl)pyrimidin-5-amine (Reference Example 70) were used, and operations similar to those of Fourth Step of Example 319 were carried out to synthesize ((4aR)-1-[(2,3-difluoro-4-iodophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide.

[1244] LCMS: m/z 755[M+H]⁺

HPLC retention time: 1.05 minutes (analysis condition SMD-TFA50)

[1245] Second Step

The iodide derivative obtained in First Step was used, and operations similar to those of Example 324 were carried out to synthesize the title compound.


[1246] LCMS: m/z 645[M+H]⁺

HPLC retention time: 1.43 minutes (analysis condition SMD-TFA05)

[1247] <Example 331>


(4aR)-N-(4-Bromo-2-iodophenyl)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide

[1248] [Formula 212]



[1249] First Step

[1250] [Formula 213]



[1251] (R)-2-Methyl-pyrrolidine-2-carboxylic acid methyl ester hydrochloride and 2,3-difluorobenzaldehyde were used, and operations similar to those of First to Third Steps of Example 319 were carried out to synthesize methyl (4aR)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylate.

[1252] Second Step

Methyl (4aR)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxylate and 4-bromo-2-iodoaniline were used as reagents, and operations similar to those of Example 21 were carried out to synthesize the title compound.

[1253] LCMS: m/z 618[M+H]⁺


HPLC retention time: 1.16 minutes (analysis condition SMD-TFA05)

[1254] <Reference Example 104>

Methyl 1-[[*(E)*]-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclobutane-1-carboxylate hydrochloride

[1255]

[Formula 214]



[1256] Methyl 1-[[[E]-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclobutane-1-carboxylate (Third Step of Example 13) (300 mg, 0.726 mmol) was dissolved in 2-butanone (1.50 mL), and pyridine hydrochloride (84.0 mg, 0.727 mmol) was added at 25°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. Heptane (1.50 mL) and a seed crystal (1.00 mg, 0.00222 mmol) were added to the reaction mixture, and then the mixture was stirred for 1 hour. Heptane (3.00 mL) was added to the reaction mixture, the mixture was stirred for 16 hours, and then the deposited crystal was collected by filtration to obtain the title compound (271 mg, 88%).

[1257] Melting point: 108°C

¹H-NMR (DMSO-D₆) δ: 10.32 (1H, brs), 7.44 (1H, m), 7.29 (1H, s), 7.07 (1H, m), 4.49 (2H, t, *J* = 4.4 Hz), 3.72 (2H, t, *J* = 5.0 Hz), 3.63 (3H, s), 3.55-3.36 (4H, m), 3.44 (3H, s), 2.87 (3H, brs), 2.80 (3H, s), 2.48 (4H, m), 1.93 (2H, m).


[1258] The seed crystal used in Reference Example 104 was obtained by the following method.

[1259] Methyl 1-[[[E]-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclobutane-1-carboxylate (75.3 mg, 0.182 mmol) was dissolved in dimethylsulfoxide (0.317 mL), and 2 M hydrochloric acid (93.0 μL, 0.186 mmol) was added. The prepared solution (15.0 μL) was freeze-dried, 2-butanone (15.0 μL) and heptane (15.0 μL) were added to the obtained powder at 25°C, and the mixture was stirred for 4 days to obtain a deposit.

[1260] <Reference Example 105>

Methyl 1-[[[(E)-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate hydrochloride

[1261] [Formula 215]



[1262] Methyl 1-[[[(E)-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate (First Step of Example 14) (300 mg, 0.702 mmol) was dissolved in 2-butanone (3.00 mL), pyridine hydrochloride (81.0 mg, 0.701 mmol) was added at 25°C, and the mixture was stirred under nitrogen atmosphere for 30 minutes. Heptane (1.00 mL) and a seed crystal (1.00 mg, 0.00215 mmol) were added to the reaction mixture, and then the mixture was stirred for 1 hour. Heptane (2.00 mL) was added to the reaction mixture, then the mixture was stirred for 2 hours, and the deposited crystal was collected by filtration to obtain the title compound (225 mg, 69%).

[1263] Melting point: 97°C

¹H-NMR (DMSO-D₆) δ: 10.34 (1H, brs), 7.44 (1H, m), 7.26 (1H, s), 7.08 (1H, m), 4.50 (2H, t, *J* = 4.4 Hz), 3.72 (2H, t, *J* = 5.0 Hz), 3.60 (3H, s), 3.55-3.44 (4H, m), 3.35 (3H, s), 2.87 (3H, brs), 2.85 (3H, s), 2.25 (2H, m), 2.14 (2H, m), 1.70 (4H, m).

[1264] The seed crystal used in Reference Example 105 was obtained by the following method.


[1265] Methyl 1-[[[(E)-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate (50.0 mg, 0.117 mmol) was dissolved in 2-butanone (0.500 mL), pyridine hydrochloride (15.0 mg, 0.130 mmol) was added at 25°C, and the mixture was stirred under

nitrogen atmosphere for 30 minutes. Heptane (0.500 mL) and methyl 1-[[[(E)-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclobutane-1-carboxylate hydrochloride (1.00 mg, 0.00222 mmol) were added to the reaction mixture, and the mixture was stirred for 1 hour. The deposited crystal was collected by filtration to obtain a deposit (36.6 mg).

[1266] <Example 332>

4-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]butanoate


[1267] [Formula 216]



[1268] First Step

7-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxylic acid 2-methylpropyl ester

[1269] [Formula 217]




[1270] 2,3-Difluoro-4-hydroxybenzaldehyde and methyl 1-(methylamino)cyclopentanecarboxylate hydrochloride (Reference Example 87) were used to synthesize the title compound by a method similar to that of Reference Example 1-1.

[1271] LCMS: m/z 411[M+H]⁺

[1272] HPLC retention time: 1.18 minutes (SMD-TFA05)

[1273] Second Step

[1274] [Formula 218]




[1275] 7-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxylic acid 2-methylpropyl ester and 4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]aniline (Reference Example 50) were used to synthesize 7-[(2,3-difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide with reference to Example 21.

[1276] LCMS: m/z 658[M+H]⁺

[1277] HPLC retention time: 1.57 minutes (SMD-TFA05)

[1278] Third Step

[1279] [Formula 219]



[1280] Cesium carbonate (20.4 mg, 0.113 mmol) was added to a solution of 7-[(2,3-difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (24.7 mg, 0.038 mmol) and methyl 4-bromobutyrate (24.48 mg, 0.075 mmol) in acetonitrile (0.4 mL), and the mixture was stirred at 70°C for 1 hour. The reaction mixture was cooled to 0°C, a 1 N-aqueous hydrochloric acid solution (0.063 mL, 0.375 mmol) was added, and the mixture was stirred for 10 minutes. The resultant was diluted with N,N-dimethyl formamide (0.6 mL) and purified by HPLC to obtain methyl 4-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-

yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl)methyl]phenoxy]butyrate (10.6 mg, 37%).

[1281] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

YMC-Actus Triart C18 (50 × 30 mmI.D., S - 5 µm, 12 nm)

LCMS: m/z 758[M+H]⁺

[1282] HPLC retention time: 1.72 minutes (SMD-FA05)

[1283] Fourth Step

Potassium trimethylsilanolate (5.38 mg, 0.042 mmol) was added to a solution of methyl 4-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl)methyl]phenoxy]butyrate (10.6 mg, 0.014 mmol) in tetrahydrofuran (0.4 mL), and the mixture was stirred at room temperature for 2 hours. After the reaction mixture was concentrated, the resultant was diluted with N,N-dimethyl formamide (0.6 mL) and purified by HPLC to obtain the title compound (6.2 mg, 60%).

[1284] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

YMC-Actus Triart C18 (50 × 30 mmI.D., S - 5 µm, 12 nm)




LCMS: m/z 744[M+H]⁺

[1285] HPLC retention time: 1.60 minutes (SMD-FA05)

[1286] In Examples 333 to 335, appropriate alkyl bromides were used, and operations similar to those of Example 332 were carried out to synthesize the compounds described in the following Table.

[1287]


[Table 31]

Example No.	Structural formula	LCMS Analysis condition	Retention time (min)	m/z [M+H] ⁺
333		SMD-FA05	1.62	758
334		SMD-TFA05	1.64	772
335		SMD-TFA05	1.67	786

[1288] <Reference Example 106>

6-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-5,6-diazaspiro[3.5]non-8-ene-8-carboxylic acid 2-methylpropyl ester

[1289] [Formula 220]



[1290] Methyl 1-(methylamino)cyclobutanecarboxylate hydrochloride was used as a starting material (Reference Example 85) to synthesize the title compound by a method similar to that of Example 332.

[1291] LCMS: m/z 411[M+H]⁺


[1292] HPLC retention time: 1.18 minutes (SMD-TFA05)

[1293] <Reference Example 107>

6-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-

8-carboxamide

[1294] [Formula 221]



[1295] Methyl 1-(methylamino)cyclobutane-1-carboxylate hydrochloride was used as a starting material to synthesize the title compound by a method similar to that of Example 332.


[1296] LCMS: m/z 658[M+H]⁺

[1297] HPLC retention time: 1.48 minutes (SMD-TFA05)

[1298] <Example 336>

8-[[2,3-Difluoro-4-(2-morpholin-2-oxo-ethoxy)phenyl]methyl]-5-hydroxy-9-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-8,9-diazaspiro[3.5]non-5-ene-6-carboxamide

[1299] [Formula 222]



[1300] The title compound was synthesized from 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide and 2-chloro-1-morpholin-4-yl-ethanone (Kapanda, Coco N. Journal of Medicinal Chemistry, 52 (22), 7310-7314; 2009) by a method similar to that of Third Step of Example 322.

[1301] LCMS: m/z 771[M+H]⁺


[1302] HPLC retention time: 1.43 minutes (analysis condition SMD-TFA05)

[1303] <Example 337>

8-[[2,3-Difluoro-4-[2-(2-oxomorpholin-4-yl)ethoxy]phenyl]methyl]-5-hydroxy-9-

methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-8,9-diazaspiro[3.5]non-5-ene-6-carboxamide

[1304] [Formula 223]



[1305] The title compound was obtained from 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide and 4-(2-hydroxyethyl)morpholin-2-one (Khromov-Borisov, N. V. and Remizov, A. L., Zhurnal Obshchei Khimii, 23, 598-605; 1953) by carrying out operations similar to those of Example 407.


[1306] LCMS: m/z 771[M+H]⁺

[1307] HPLC retention time: 1.38 minutes (analysis condition SMD-TFA05)

[1308] <Reference Example 108>

6-[[4-(2-Bromoethoxy)-2,3-difluorophenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1309] [Formula 224]



[1310] 6-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107) (3.00 g, 4.66 mmol) was dissolved in acetone (23.3 mL), then 1,2-dibromoethane (6.85 mL, 79.0 mmol) and potassium carbonate (1.29 g, 9.32 mmol) was added, and the mixture was stirred under nitrogen atmosphere at 60°C for 15 hours. 1 N hydrochloric acid was added to the reaction mixture at 0°C, and the mixture was

extracted with ethyl acetate. The organic layer was washed with a brine, the aqueous layer was separated by a phase separator, the resultant was concentrated at reduced pressure, and the resultant residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound (3.19 g, 91%).


[1311] LCMS: m/z 749[M+H]⁺

HPLC retention time: 1.58 minutes (analysis condition SMD-TFA05)

<Reference Example 109>

7-[[4-(2-Bromoethoxy)-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1312] [Formula 225]



[1313] The title compound was synthesized from 7-[(2,3-difluoro-4-hydroxybenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (Example 332) in a manner similar to that of Reference Example 106.

[1314] LCMS: m/z 764[M+H]⁺


HPLC retention time: 1.62 minutes (analysis condition SMD-TFA05)

<Example 338>

6-[[2,3-Difluoro-4-[2-[methyl(oxolan-3-yl)amino]ethoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1315]

[Formula 226]



[1316] 6-[[4-(2-Bromoethoxy)-2,3-difluorophenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (10.0 mg, 13.0 μ mol) and N-methyltetrahydrofuran-3-amine (2.70 mg, 27.0 μ mol) were dissolved in N,N-dimethyl formamide (100 μ L), then N-ethyl-N-isopropylpropan-2-amine (4.6 μ L, 27.0 μ mol) and tetrabutylammonium iodide (0.5 mg, 1.33 μ mol) were added, and the mixture was stirred at 80°C for 1.5 hours. The reaction mixture was diluted with N,N-dimethyl formamide (0.5 mL) and purified by HPLC to obtain the title compound (8.0 mg, 78%).

[1317] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

Column: YMC-Actus Triart C18 (50 $\times$ 30 mmL.D., S - 5 μ m, 12 nm)

LCMS: m/z 771[M+H]⁺

HPLC retention time: 1.31 minutes (analysis condition SMD-TFA05)

[1318] <Examples 339 to 380>









Various amines which were known from literature or commercialized were used, and operations similar to those of Example 338 were carried out to synthesize the compounds described in the following Table. Note that if the amine was hydrochloride, the amine was not directly used for the reaction and used for the reaction after hydriodic acid (57 wt. %, 2 equivalent weight) was allowed to act and the corresponding hydroiodide was formed.

[1319] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)










Column: YMC-Actus Triart C18 (50 $\times$ 30 mmL.D., S - 5 μ m, 12 nm, 100 $\times$ 30 mmL.D., S - 5 μ m, 12 nm).

[1320] [Table 32-1]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
339		SMD-TFA05	1.34	769
340		SMD-TFA05	1.34	783
341		SMD-TFA05	1.34	785
342		SMD-TFA05	1.35	785
343		SMD-TFA05	1.34	785
344		SMD-TFA05	1.35	785
345		SMD-TFA05	1.37	785
346		SMD-TFA05	1.33	773

[1321]

[Table 32-2]










347		SMD-TFA05	1.28	745
348		SMD-TFA05	1.37	803
349		SMD-TFA05	1.38	787
350		SMD-TFA05	1.38	787
351		SMD-TFA05	1.27	775
352		SMD-TFA05	1.51	739
353		SMD-TFA05	1.28	786
354		SMD-TFA05	1.31	800
355		SMD-TFA05	1.35	799

[1322] [Table 32-3]

356		SMD-TFA05	1.35	771
357		SMD-TFA05	1.35	773
358		SMD-TFA05	1.16	840
359		SMD-TFA05	1.29	784
360		SMD-TFA05	1.22	800
361		SMD-TFA05	1.23	844
362		SMD-TFA05	1.23	814
363		SMD-TFA05	1.31	834
364		SMD-TFA05	1.30	827

[1323]

[Table 32-4]

365		SMD-TFA05	1.35	799
366		SMD-TFA05	1.34	785
367		SMD-TFA05	1.36	797
368		SMD-TFA05	1.34	787
369		SMD-TFA05	1.39	777
370		SMD-TFA05	1.37	799
371		SMD-TFA05	1.36	799
372		SMD-TFA05	1.40	801
373		SMD-TFA05	1.31	759

[Table 32-5]


374		SMD-TFA05	1.19	854
375		SMD-TFA05	1.40	799
376		SMD-TFA05	1.25	814
377		SMD-TFA05	1.30	828
378		SMD-TFA05	1.26	858
379		SMD-TFA05	1.34	848
380		SMD-TFA05	1.37	783

[1325] <Example 381>

(2S)-2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]butanedioate

[1326]

[Formula 227]



[1327] L-aspartate dimethyl ester hydrochloride (142 mg, 0.72 mmol) was dissolved in methanol (1.00 mL), hydroiodic acid (57 wt. %, 95.0 μ L) was added, and the mixture was stirred for 1 minute. After the reaction mixture was concentrated at reduced pressure, methanol and toluene were added for azeotropic removal to obtain L-aspartate dimethyl ester hydroiodide as a crude product. The resultant hydroiodide of amine and 7-[[4-(2-bromoethoxy)-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (110 mg, 0.14 mmol) (Reference Example 109) were dissolved in N,N-dimethyl formamide (480 μ L), and then potassium phosphate (92.0 mg, 0.43 mmol) and tetrabutylammonium iodide (5.3 mg, 14.0 μ mol) were added, and the mixture was stirred at 80°C for 2 hours. After the reaction mixture was diluted with tetrahydrofuran (480 μ L), potassium trimethylsilanolate (92.0 mg, 0.72 mmol) was added, and the mixture was stirred at 50°C for 30 minutes and at 70°C for 2 hours. The reaction mixture was diluted with formic acid and water, and the resultant was purified by HPLC to obtain the title compound (71.5 mg, 61%).

[1328] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

Column: YMC-Actus Triart C18 (100 $\times$ 30 mmI.D., S - 5 μ m, 12 nm)

LCMS: m/z 817[M+H]⁺


HPLC retention time: 1.32 minutes (analysis condition SMD-TFA05)

<Example 382>

3-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbomoyl]-6,7-diazaspiro[4.5]dec-9-en-7-


yl)methyl]phenoxy]ethylamino]pentanedioate

[1329] [Formula 228]



[1330] First Step

[1331] [Formula 229]



[1332] 7-[[4-(2-Bromoethoxy)-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (Reference Example 109) and diethyl 3-aminopentanedioate were used as reagents, and operations similar to those of Example 337 were carried out to synthesize diethyl 3-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]pentanedioate.

[1333] Second Step

Diethyl 3-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]pentanedioate (48.0 mg, 54.0 μ mol) was dissolved in tetrahydrofuran (200 μ L), potassium trimethylsilanolate (34.7 mg, 0.27 mmol) was added, and the mixture was stirred at 50°C for 30 minutes. The reaction mixture was diluted with formic acid and water and purified by HPLC to obtain the title compound (32.4 mg, 72%).

[1334] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

Column: YMC-Actus Triart C18 (50 × 30 mmI.D., S - 5 μm, 12 nm)


LCMS: m/z 831[M+H]⁺

HPLC retention time: 1.32 minutes (analysis condition SMD-TFA05)

[1335] <Example 383>


6-[4-[3-[(2R)-2,3-Dihydroxypropoxy]propoxy]-2,3-difluorophenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1336] [Formula 230]



[1337] First Step


[1338] [Formula 231]



[1339] 6-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-5,6-diazaspiro[3.5]non-8-ene-8-carboxylic acid 2-methylpropyl ester (Reference Example 106) and 4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]aniline (Reference Example 13) were used as reagents, and operations similar to those of Third Step of Example 149 were carried out to synthesize 6-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide.

[1340] Second Step

[1341] [Formula 232]



[1342] (S)-3-((2,2-Dimethyl-1,3-dioxolan-4-yl)methoxy)propan-1-ol (300 mg, 1.58 mmol) and triethylamine (0.33 mL, 2.37 mmol) were dissolved in tetrahydrofuran (7.88 mL), methanesulfonyl chloride (0.15 mL, 1.89 mmol) was added at 0°C, and the mixture was stirred at room temperature for 3 hours. After a 1 N aqueous dipotassium hydrogenphosphate solution was added to the reaction mixture and the mixture was extracted with ethyl acetate, the organic layer was washed with a saturated sodium bicarbonate solution and a brine. The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated at reduced pressure. The resultant residue was purified by column chromatography on silica gel (ethyl acetate/hexane) to obtain 3-[[[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methoxy]propyl methanesulfonate (316 mg, 75%).

[1343] LCMS: m/z 269[M+H]⁺

HPLC retention time: 0.77 minutes (analysis condition SMD-TFA05)

[1344] Third Step

6-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (15.0 mg, 23.0 μ mol) and 3-[[[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methoxy]propyl methanesulfonate (8.77 mg, 33.0 μ mol) were dissolved in acetonitrile (117 μ L), cesium carbonate (22.8 mg, 233 μ mol) was added, and the mixture was stirred at 75°C for 1.5 hours. After the reaction mixture was cooled to room temperature, an aqueous hydrochloric acid solution (6N, 38.9 μ L, 233 μ mol) and methanol (38.9 μ L) were added, and the mixture was stirred at room temperature for 1 hour. A 6 N aqueous sodium hydroxide solution was added to the reaction mixture, and after the mixture was filtered and concentrated, the resultant was diluted with N,N-dimethyl formamide (0.5 mL) and purified by HPLC to obtain the title compound (11.9 mg, 66%).

[1345] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

Column: YMC-Actus Triart C18 (100 $\times$ 30 mmI.D., S - 5 μ m, 12 nm)

LCMS: m/z 775[M+H]⁺

[1346] HPLC retention time: 1.40 minutes (analysis condition SMD-TFA05)

[1347] <Examples 384 to 400>

Bromide(2-[2-(3-bromopropoxy)ethoxy]oxane, 2-[2-(4-bromobutoxy)ethoxy]oxane, 2-[2-(5-bromopentoxy)ethoxy]oxane, 2-[3-(3-bromopropoxy)propoxy]oxane, 2-[3-(4-bromobutoxy)propoxy]oxane, 2-[3-(5-bromopentoxy)propoxy]oxane synthesized by a method similar to that of First Step of Example 147 by using alcohol(2-(oxan-2-yloxy)ethanol, 3-(oxan-2-yloxy)propan-1-ol), and dibromide (1,3-dibromopropane, 1,4-dibromobutane, 1,5-dibromopentane) which were known from literature or commercialized, and 3-[[[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methoxy]propyl methanesulfonate (Example 383), (S)-4-(((4-bromobutoxy)methyl)-2,2-dimethyl-1,3-dioxolane (Reference Example 74), and (S)-4-(((5-bromopentyl)oxy)methyl)-2,2-dimethyl-1,3-dioxolane (Reference Example 78), 7-[(2,3-difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide synthesized by a method similar to that of First Step of Example 383 by using 7-[(2,3-difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxylic acid 2-methylpropyl ester (Example 332) and 4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]aniline (Reference Example 13), 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Example 383), 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107), or 7-(2,3-difluoro-4-hydroxybenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (Example 332) were used, and operations similar to those of Example 383 were carried out to synthesize the compounds described in the following Table.

[1348] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

- 325 -









Column: YMC-Actus Triart C18 (50 × 30 mmI.D., S - 5 µm, 12 nm, 100 ×
30 mmI.D., S - 5 µm, 12 nm)

[1349]

[Table 33-1]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
384		SMD-TFA05	1.51	789
385		SMD-TFA05	1.54	803
386		SMD-TFA05	1.43	789
387		SMD-TFA05	1.54	803
388		SMD-TFA05	1.57	817
389		SMD-TFA05	1.50	776
390		SMD-TFA05	1.52	790
391		SMD-TFA05	1.55	804
392		SMD-TFA05	1.52	790


[1350] [Table 33-2]

393		SMD-TFA05	1.55	804
394		SMD-TFA05	1.58	818
395		SMD-TFA05	1.63	760
396		SMD-TFA05	1.65	774
397		SMD-TFA05	1.68	788
398		SMD-TFA05	1.66	774
399		SMD-TFA05	1.68	788
400		SMD-TFA05	1.57	802

[1351] <Example 401>

6-[[2,3-Difluoro-4-[2-(2-methoxyethoxy)ethoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1352] [Formula 233]



[1353] The title compound was synthesized from 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107) and 1-bromo-2-(2-methoxyethoxy)ethane in a manner similar to that of Third Step of Reference Example 383.

[1354] LCMS: m/z 746[M+H]⁺


HPLC retention time: 1.66 minutes (analysis condition SMD-TFA05)

<Example 402>

6-(2,3-Difluoro-4-(2-(methyl(oxetan-3-yl)amino)ethoxy)benzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

First Step

[1355] [Formula 234]



[1356] 6-(4-(2-Bromoethoxy)-2,3-difluorobenzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (10.0 mg, 0.013 mmol) (Reference Example 108) was dissolved in 1,3-dimethyl-2-imidazolidinone (100 μ L), and then a 2.0 M methylamine-tetrahydrofuran solution (400 μ L, 0.800 mmol), N,N-diisopropyl ethylamine (4.6 μ L, 0.027 mmol), then tetrabutyl ammonium iodide (0.5 mg, 0.001 mmol) were added, and the mixture was stirred


at 80°C for 3 hours. The reaction mixture was concentrated at reduced pressure, a 0.5 M hydrochloric acid-methanol solution (80 µL, 0.040 mmol) was added, and normal butanol and toluene were added for azeotropic removal to obtain 6-(2,3-difluoro-4-(2-(methylamino)ethoxy)benzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide hydrochloride as a crude product.

[1357] LCMS: m/z 701[M+H]⁺

[1358] HPLC retention time: 1.27 minutes (analysis condition SMD-TFA05)

[1359] Second Step

[1360] [Formula 235]



[1361] The crude product of 6-(2,3-difluoro-4-(2-(methylamino)ethoxy)benzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide hydrochloride was dissolved in 1,2-dichloroethane (0.5 mL), then oxetan-3-one (6.25 µL, 0.104 mmol), triacetoxy sodium borohydride (16.5 mg, 0.078 mmol), and acetic acid (10 µL, 0.175 mmol) were added, and the mixture was stirred at room temperature for 1 hour. Oxetan-3-one (6.25 µL, 0.104 mmol) and triacetoxy sodium borohydride (16.5 mg, 0.078 mmol) were further added, and the mixture was stirred at room temperature for 1 hour. After the reaction mixture was concentrated and diluted with N,N-dimethyl formamide, the resultant was purified by HPLC to obtain the title compound (3.8 mg, two-step yield of 39%).

[1362] LCMS: m/z 757[M+H]⁺


[1363] HPLC retention time: 1.25 minutes (analysis condition SMD-TFA05)

[1364] <Example 403>

6-(2,3-Difluoro-4-(oxetan-3-yloxy)benzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-

(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1365] [Formula 236]



[1366] 6-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107) was dissolved in acetonitrile (200 μ L), then oxetan-3-yl 4-methylbenzenesulfonate (14.2 mg, 0.062 mmol) and cesium carbonate (30.4 mg, 0.093 mmol) were added, and the mixture was stirred at 80°C overnight. After the reaction mixture was diluted with N,N-dimethyl formamide and water, the resultant was purified by HPLC to obtain the title compound (14.1 mg, 65%).


[1367] LCMS: m/z 700[M+H]⁺

[1368] HPLC retention time: 1.62 minutes (analysis condition SMD-TFA05)

[1369] <Example 404>

6-[[2,3-Difluoro-4-[[[(2S)-1-(2-methoxyethyl)pyrrolidin-2-yl]methoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide


[1370] [Formula 237]



[1371] First Step

[1372]

[Formula 238]




[1373] 6-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107) (100 mg, 0.16 mmol) and (S)-tert-butyl 2-(hydroxymethyl)pyrrolidine-1-carboxylate (62.6 mg, 0.31 mmol) were dissolved in tetrahydrofuran (311 μ L), then N,N,N',N'-tetramethylazodicarboxamide (53.5 mg, 0.31 mmol) was added, and the mixture was stirred at 70°C for 1 hour. The reaction mixture was concentrated at reduced pressure, and the resultant residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain tert-butyl (2S)-2-[[2,3-difluoro-4-[[9-hydroxy-5-methyl-7-oxo-8-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-5,6-diazaspiro[3.5]non-8-en-6-yl]methyl]phenoxy]methyl]pyrrolidine-1-carboxylate (115 mg, 90%).

[1374] LCMS: m/z 827[M+H]⁺

HPLC retention time: 1.68 minutes (analysis condition SMD-TFA05)

[1375] Second Step

[1376] [Formula 239]



[1377] Hydrochloric acid (a dioxane solution, 4 N, 1.00 mL, 4.00 mmol) was added to tert-butyl (2S)-2-[[2,3-difluoro-4-[[9-hydroxy-5-methyl-7-oxo-8-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-5,6-diazaspiro[3.5]non-8-en-6-yl]methyl]phenoxy]methyl]pyrrolidine-1-carboxylate (104 mg, 0.13 mmol), and the mixture

was stirred at room temperature for 1 hour. After the reaction mixture was concentrated at reduced pressure, toluene was added for azeotropic removal to obtain (S)-6-(2,3-difluoro-4-(pyrrolidin-2-ylmethoxy)benzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide hydrochloride (96 mg, 100%).

[1378] LCMS: m/z 727[M+H]⁺

HPLC retention time: 1.27 minutes (analysis condition SMD-TFA05)

Third Step

(S)-6-(2,3-Difluoro-4-(pyrrolidin-2-ylmethoxy)benzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide hydrochloride (10.0 mg, 13.0 μ mol) and 1-bromo-2-methoxyethane (3.64 mg, 26.0 μ mol) were dissolved in N,N-dimethyl formamide (100 μ L), then N-ethyl-N-isopropylpropan-2-amine (6.9 μ L, 39.0 μ mol) and tetrabutylammonium iodide (0.5 mg, 1.31 μ mol) were added, and the mixture was stirred at 80°C for 1.5 hours. The reaction mixture was diluted with N,N-dimethyl formamide (0.5 mL) and purified by HPLC to obtain the title compound (3.2 mg, 31%).

[1379] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

Column: YMC-Actus Triart C18 (50 $\times$ 30 mmI.D., S - 5 μ m, 12 nm)

LCMS: m/z 785[M+H]⁺


HPLC retention time: 1.32 minutes (analysis condition SMD-TFA05)

<Example 405>

6-[[2,3-Difluoro-4-[(3R)-1-(2-methoxyethyl)pyrrolidin-3-yl]oxyphenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1380]

[Formula 240]



[1381] The title compound was synthesized from 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107) and (S)-tert-butyl 3-hydroxy pyrrolidine-1-carboxylate in a manner similar to that of First to Third Steps of Example 404.


[1382] LCMS: m/z 771[M+H]⁺

HPLC retention time: 1.30 minutes (analysis condition SMD-TFA05)

<Example 406>

6-[[2,3-Difluoro-4-[1-(2-methoxyethyl)azetidin-3-yl]oxyphenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1383] [Formula 241]



[1384] The title compound was synthesized from 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107) and tert-butyl 3-hydroxyazetidine-1-carboxylate in a manner similar to that of First to Third Steps of Example 404.


[1385] LCMS: m/z 756[M+H]⁺

HPLC retention time: 1.33 minutes (analysis condition SMD-TFA05)

<Example 407>

6-[[2,3-Difluoro-4-(2-methyl-1-morpholin-4-ylpropan-2-yl)oxyphenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide

[1386] [Formula 242]



[1387] A tetrahydrofuran solution (1M) (0.109 mL, 0.109 mmol) of trimethylphosphine was added for as long as 30 seconds to a solution of 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (10 mg, 0.016 mmol), 2-methyl-2-morpholinopropan-1-ol (17 mg, 0.106 mmol), and N,N,N',N'-tetramethylazodicarboxamide (18.7 mg, 0.106 mmol) in tetrahydrofuran (0.3 mL) in a state in which they were heated at 60°C. The reaction mixture was stirred at 60°C for 16 hours. After the reaction mixture was cooled to room temperature, the resultant was blown with nitrogen to remove the solvent. The residue was dissolved in dimethylformamide and formic acid, and the resultant was purified by HPLC (0.1% formic acid water/0.1% formic acid acetonitrile) to obtain the title compound (4.3 mg, 34%).

[1388] LCMS: m/z 785[M+H]⁺





[1389] HPLC retention time: 1.30 minutes (SMD-TFA05)

[1390] <Examples 408 to 411>

Appropriate alkyl alcohols were used, and operations similar to those of Example 407 were carried out to synthesize the compounds described in the following Table.

[1391]

[Table 34]


Example No.	Structural formula	LCMS Analysis condition	Retention time (min)	m/z [M+H] ⁺
408		SMD-TFA05	1.30	771
409		SMD-TFA05	1.31	742
410		SMD-TFA05	1.33	756
411		SMD-TFA05	1.30	741

[1392] <Example 412>

7-(2,3-Difluoro-4-(2-(methyl(oxetan-3-yl)amino)ethoxy)benzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

First Step

[1393] [Formula 243-1]



[1394] 7-[[4-(2-Bromoethoxy)-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-


ene-9-carboxamide (20.0 mg, 0.026 mmol) (Reference Example 109) was used, and operations similar to those of First Step of Example 402 were carried out to obtain a crude product of 7-(2,3-difluoro-4-(2-(methylamino)ethoxy)benzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide hydrochloride.

[1395] LCMS: m/z 715[M+H]⁺

[1396] HPLC retention time: 1.29 minutes (analysis condition SMD-TFA05)

[1397] Second Step

[1398] [Formula 243-2]



[1399] 7-(2,3-Difluoro-4-(2-(methylamino)ethoxy)benzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide hydrochloride was used, and operations similar to those of Second Step of Example 402 were carried out to obtain the title compound (11.3 mg, two-step yield of 56%).

[1400] LCMS: m/z 771[M+H]⁺

[1401] HPLC retention time: 1.19 minutes (analysis condition SMD-TFA05)


[1402] ¹H-NMR (DMSO-D₆) δ : 12.72 (1H, s), 9.59 (1H, s), 8.56 (1H, s), 8.42 (1H, d, J = 8.8 Hz), 8.24 (1H, s), 7.97 (1H, d, J = 7.3 Hz), 7.16 (1H, t, J = 7.3 Hz), 7.04 (1H, t, J = 7.6 Hz), 5.07 (1H, d, J = 13.7 Hz), 4.52 (2H, t, J = 6.6 Hz), 4.41 (2H, t, J = 6.1 Hz), 4.16 (3H, t, J = 5.4 Hz), 3.66 (1H, q, J = 6.5 Hz), 2.67 (2H, t, J = 5.4 Hz), 2.46 (3H, s), 2.18 (3H, s), 2.01 (1H, m), 1.59 (3H, m), 1.27 (3H, m), 1.07 (1H, m).

[1403] <Example 413>

6-[[2,3-Difluoro-4-[1-(2-methoxyethyl)azetidin-3-yl]oxyphenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-

diazaspiro[3.5]non-8-ene-8-carboxamide

[1404] [Formula 243-3]



[1405] The title compound was synthesized from 6-[(2,3-difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107) and 2-(chloromethyl)tetrahydrofuran in a manner similar to that of Third Step of Example 383.


[1406] LCMS: m/z 742[M+H]⁺

HPLC retention time: 1.61 minutes (analysis condition SMD-TFA05)

<Example 414>

7-(4-(3-(Dimethylamino)-2,2-dimethylpropoxy)-2,3-difluorobenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1407] [Formula 243-4]



[1408] The title compound was synthesized from 7-[(2,3-difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide and 3-(dimethylamino)-2,2-dimethylpropan-1-ol by a method similar to that of Example 407.

[1409] LCMS: m/z 771[M+H]⁺


[1410] HPLC retention time: 1.25 minutes (analysis condition SQD-FA05)

[1411] $^1\text{H-NMR}$ (CDCl_3) δ : 16.47 (1H, s), 12.87 (1H, s), 9.60 (1H, s), 8.47 (1H, d, $J = 8.4$ Hz), 7.95 (1H, s), 7.93 (1H, s), 7.79 (1H, d, $J = 8.4$ Hz), 6.99 (1H, m), 6.72 (1H, m), 5.04 (1H, d, $J = 14.4$ Hz), 4.4.19 (1H, d, $J = 14.4$ Hz), 3.80 (2H, s), 2.48 (3H, s), 2.43-2.36 (8H, m), 1.85-1.32 (10H, m), 1.02 (6H, s).

[1412] <Example 415>


1-(2-(2,3-Difluoro-4-((10-hydroxy-6-methyl-8-oxo-9-((4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)carbamoyl)-6,7-diazaspiro[4.5]dec-9-en-7-yl)methyl)phenoxy)ethyl)-1,4-diazabicyclo[2.2.2]octan-1-ium formate

[1413] [Formula 243-5]



[1414] First Step

[1415] [Formula 244]



[1416] A solution of 7-[(2,3-difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxylic acid 2-methylpropyl ester, 4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]aniline (100 mg, 0.152 mmol), 1,2-dichloroethane (151 mg, 1.52 mmol), and potassium carbonate (42 mg, 0.304 mmol) in N,N-dimethyl formamide (1.01 mL) was heated at 60°C for 18 hours. The residue was dissolved in N,N-dimethyl formamide and formic acid, and the reaction mixture was purified by C-18 reverse-phase column chromatography on silica gel (0.1% formic acid water/0.1% formic acid acetonitrile) to obtain 7-(4-(2-chloroethoxy)-2,3-difluorobenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-

diazaspiro[4.5]dec-9-ene-9-carboxamide (91 mg, 83%).

[1417] LCMS: m/z 720[M+H]⁺

[1418] HPLC retention time: 1.697 minutes (SMD-TFA05)

[1419] Second Step

A solution of 7-(4-(2-chloroethoxy)-2,3-difluorobenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (93 mg, 0.126 mmol), 1,4-diazabicyclo[2.2.2]octane (142 mg, 1.26 mmol), and potassium carbonate (175 mg, 1.26 mmol) in acetone (3.16 mL) were stirred at 60°C for 4 days. After the reaction mixture was cooled to room temperature, the resultant was blown with nitrogen gas to remove the solvent. The residue was dissolved in N,N-dimethyl formamide and formic acid, and the resultant was purified by HPLC (0.1% formic acid water/0.1% formic acid acetonitrile) to obtain the title compound (78 mg, 73%).


[1420] LCMS: m/z 796[M]⁺

[1421] HPLC retention time: 1.05 minutes (analysis condition SQD-FA05)

[1422] <Example 416>

6-[[2,3-Difluoro-4-[1-(2-methoxyethyl)azetidin-3-yl]oxyphenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide


[1423] [Formula 245]



[1424] First Step

[1425]

[Formula 246]



[1426] 7-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxylic acid 2-methylpropyl ester was synthesized from methyl 1-[[[E]-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate hydrochloride (Reference Example 105) in a manner similar to that of Fourth Step of Example 237.

[1427] LCMS: m/z 540[M+H]⁺

HPLC retention time: 1.11 minutes (analysis condition SMD-TFA05)

[1428] Second Step

The title compound was synthesized from 7-[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxylic acid 2-methylpropyl ester and 2-(5-methyl-6-(trifluoromethyl)pyrimidin-4-yl)-4-(trifluoromethyl)aniline (Reference Example 96) in a manner similar to that of Fifth Step of Example 237.

[1429] LCMS: m/z 787[M+H]⁺


HPLC retention time: 1.41 minutes (analysis condition SMD-TFA05)

[1430] <Example 417>

7-(2,3-Difluoro-4-((2-((2-methoxyethyl)(methyl)amino)ethyl)amino)benzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide


[1431]

[Formula 247]



[1432] First Step

[1433] [Formula 248]




[1434] 2,3-Difluoro-4-iodobenzaldehyde and methyl 1-(methylamino)cyclopentane carboxylate hydrochloride were used, and operations similar to those of Third to Fifth Steps of Example 13 were carried out to synthesize the title compound (2.69 g, three-step yield of 69%).

[1435] LCMS: m/z 768 $[M+H]^+$

HPLC retention time: 0.99 minutes (analysis condition SQD-FA50)

[1436] Second Step

[1437] [Formula 249]



[1438] Dimethyl sulfoxide (1.07 mL) was added to a mixture of 7-(2,3-difluoro-4-iodobenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (205 mg, 0.267 mmol), copper (I) iodide (10.2 mg, 0.053 mmol), 2-((2,6-dimethylphenyl)amino)-2-oxoacetate (20.6 mg, 0.107 mmol), potassium phosphate (284 mg,

1.336 mmol), and N1-(2-methoxyethyl)-N1-methylethane-1,2-diamine (70.6 mg, 0.534 mmol), and the mixture was stirred under nitrogen atmosphere at 90°C for 5 hours. Aqueous ammonia was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The residue was purified by C18 reverse-phase column chromatography to obtain the title compound (210 mg, 95%).


[1439] LCMS: m/z 772[M+H]⁺

HPLC retention time: 0.83 minutes (analysis condition SQD-FA05)

[1440] <Example 418>

7-[[2,3-Difluoro-4-[4-[methyl]-[(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]-4-oxobutoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1441] [Formula 250]




[1442] 4-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]butanoate (40 mg, 0.054 mmol) (Example 332) was dissolved in N,N-dimethyl formamide (1 mL), then (2R,3R,4R,5S)-6-(methylamino)hexane-1,2,3,4,5-pentol (31.5 mg, 0.161 mmol), HATU (40.9 mg, 0.108 mmol), and N-ethyl-N-isopropylpropan-2-amine (20.86 mg, 0.161 mmol) were added, and the mixture was stirred at room temperature for 15 hours. Water was added to the reaction mixture (0.2 mL), and the resultant was purified by C18 reverse-phase column chromatography to obtain the title compound (32 mg, 65%) as a white amorphous solid.

[1443] LCMS: m/z 921[M+H]⁺

[1444] HPLC retention time: 1.01 minutes (analysis condition SQD-FA05)

[1445] <Example 419>


7-[[4-[2-[2-(2,3-Dihydroxypropylamino)ethoxy]ethoxy]-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide
[1446] [Formula 251]



[1447] First Step

2-[2-(4-Methylphenyl)sulfonyloxyethoxy]ethyl 4-methylbenzenesulfonate

[1448] [Formula 252]



[1449] 2,2'-Oxydiethanol (200 mg, 1.9 mmol) and triethylamine (572 mg, 5.65 mmol) were dissolved in dichloromethane (10 mL), then tosyl chloride (898 mg, 4.7 mol) was added, and the mixture was stirred at room temperature for 15 hours. The reaction mixture was purified by chromatography on silica gel (hexane:ethyl acetate = 2:1) to obtain the title compound (620 mg, 79%) as a white plate-like crystal.


[1450] LCMS: m/z 415[M+H]⁺

[1451] HPLC retention time: 0.88 minutes (analysis condition SQD-FA05)

[1452] Second Step

2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethoxy]ethyl 4-methylbenzenesulfonate

[1453] [Formula 253]



[1454] 7-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-

(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (100 mg, 0.152 mmol) was dissolved in acetonitrile (5 mL), then oxybis(ethane-2,1-diyl)bis(4-methylbenzenesulfonate) obtained in First Step (252 mg, 0.61 mmol) and cesium carbonate (49.6 mg, 0.152 mmol) were added, and the mixture was stirred at 50°C for 4 hours. The reaction mixture was purified by C18 reverse-phase column chromatography to obtain the title compound (115 mg, 84%) as a white amorphous solid.

[1455] LCMS: m/z 900[M+H]⁺

[1456] HPLC retention time: 1.25 minutes (analysis condition SQD-FA05)

[1457] Third Step

2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethoxy]ethyl 4-methylbenzenesulfonate obtained in Second Step (30 mg, 0.033 mmol) and 3-aminopropane-1,2-diol (9.11 mg, 0.1 mmol) were dissolved in acetonitrile (1 mL), and the mixture was stirred at 50°C for 5 hours. The reaction mixture was purified by C18 reverse-phase column chromatography to obtain the title compound (18 mg, 66%) as a white amorphous solid.


[1458] LCMS: m/z 819[M+H]⁺

[1459] HPLC retention time: 0.82 minutes (analysis condition SQD-FA05)

[1460] <Example 420>

7-[[2,3-Difluoro-4-[2-[2-[methyl-[(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1461] [Formula 254]



[1462] (2R,3R,4R,5S)-6-(Methylamino)hexane-1,2,3,4,5-pentol was used as a raw material, and operations similar to those of Example 419 were carried out to synthesize the title compound.


[1463] LCMS: m/z 923[M+H]⁺

[1464] HPLC retention time: 0.79 minutes (analysis condition SQD-FA05)

[1465] <Example 421>

7-[[[2,3-Difluoro-4-[2-[2-[2-[2-[2-[methyl]-[(2R,3S,4S,5S)-2,3,4,5,6-pentahydroxyhexyl]amino]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1466] [Formula 255]



[1467] 2-[2-[2-[2-(2-Hydroxyethoxy)ethoxy]ethoxy]ethoxy]ethanol was used as a raw material, and operations similar to those of Example 419 were carried out to synthesize the title compound.


[1468] LCMS: m/z 1055[M+H]⁺

[1469] HPLC retention time: 0.81 minutes (analysis condition SQD-FA05)

[1470] <Example 422>

7-[[[2,3-Difluoro-4-[2-[2-[3-methyl]-[(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]-3-oxopropoxy]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide


[1471] [Formula 256]



[1472] First Step

tert-Butyl 3-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]propanoate

[1473] [Formula257]




[1474] 2-[2-(2-Hydroxyethoxy)ethoxy]ethanol (2 g, 13.3 mmol) was dissolved in THF (5 mL), and NaH (8 mg, 0.2 mol) was added. After the mixture was stirred at room temperature for 10 minutes, tert-butyl prop-2-enoate (0.5 g, 3.9 mmol) was slowly added, and the mixture was stirred at room temperature for 15 hours. The reaction mixture was concentrated, and the resultant residue was purified by column chromatography on silica gel (ethyl acetate) to obtain the target title compound (488 mg, 45%) as colorless clear oil. ¹H-NMR (CDCl₃) δ: 3.68-3.65 (4H, m), 3.62-3.55 (10H, m), 2.47 (2H, t, *J* = 6.4 Hz), 1.44 (9H, s).

[1475] Second Step

tert-Butyl 3-[2-[2-[2-(4-methylphenyl)sulfonyloxyethoxy]ethoxy]ethoxy]propanoate

[1476] [Formula 258]



[1477] tert-Butyl 3-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]propanoate (488 mg, 1.75 mmol) and triethylamine (532 mg, 5.26 mmol) were dissolved in dichloromethane (15 mL), and tosyl chloride (435 mg, 2.28 mmol) was added. After the mixture was stirred at room temperature for 17 hours, the reaction mixture was concentrated, and the resultant residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 1:2) to obtain the title compound (550 mg, 73%) as colorless clear oil.

[1478] LCMS: *m/z* 450[M+H₂O]⁺


[1479] HPLC retention time: 0.90 minutes (analysis condition SQD-FA05)

[1480] Third Step

tert-Butyl 3-[2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-

(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl)methyl]phenoxy]ethoxy]ethoxy]propanoate

[1481] [Formula 259]



[1482] After tert-butyl 3-[2-[2-[2-(4-methylphenyl)sulfonyloxyethoxy]ethoxy]ethoxy]propanoate (70 mg, 0.18 mmol) and 7-[(2,3-difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (79 mg, 0.12 mmol) were dissolved in acetonitrile (2 mL), cesium carbonate (117 mg, 1.36 mol) was added, and the mixture was stirred at 70°C for 2 hours. After water (0.1 mL) was added to the reaction mixture, the resultant was purified by C18 reverse-phase column chromatography to obtain the title compound (18 mg, 66%) as a white amorphous solid.


[1483] LCMS: m/z 874[M+H]⁺

[1484] HPLC retention time: 1.29 minutes (analysis condition SQD-FA05)

[1485] Fourth Step

3-[2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl)methyl]phenoxy]ethoxy]ethoxy]propanoate

[1486] [Formula 260]



[1487] tert-Butyl 3-[2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-

diazaspiro[4.5]dec-9-en-7-yl)methyl]phenoxy]ethoxy]ethoxy]propanoate (100 mg, 0.114 mmol) was dissolved in dichloromethane (2 mL), and TFA (0.54 mL) was added at room temperature. After the reaction mixture was stirred at room temperature for 2 hours, the resultant was concentrated to distill TFA away. The resultant residue was directly used for the next reaction.

[1488] LCMS: m/z 818[M+H]⁺

[1489] HPLC retention time: 1.15 minutes (analysis condition SQD-FA05)

[1490] Fifth Step

3-[2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl)methyl]phenoxy]ethoxy]ethoxy]propanoate (40 mg, 0.05 mmol), 3-aminopropane-1,2-diol (9 mg, 0.1 mol), and HATU (37.2 mg, 0.1 mol) were dissolved in DMF (1 mL), then diisopropylethylamine (12.6 mg, 0.1 mmol) was added, and the mixture was stirred at room temperature for 15 hours. Water (0.1 mL) was added to the reaction mixture, and the resultant was purified by C18 reverse-phase column chromatography to obtain the title compound (18 mg, 41%) as a white amorphous solid.

[1491] LCMS: m/z 995[M+H]⁺



[1492] HPLC retention time: 1.01 minutes (analysis condition SQD-FA05)

[1493] <Examples 423 and 424>

The compounds described in the following Table were synthesized by carrying out Steps similar to those of Example 422 by using 2-[2-(2-hydroxyethoxy)ethoxy]ethanol as a starting material.

[1494]



[Table 35]

Example No.	Structural formula	LCMS Analysis condition	Retention time (min)	m/z [M+H] ⁺
423		SQD-FA05	1.02	1039
424		SQD-FA05	1.06	935

[1495] <Examples 425 and 426>

The compounds described in the following Table were synthesized by carrying out Steps similar to those of Example 422 by using 2-[2-hydroxyethyl(methyl)amino]ethanol as a starting material.

[1496] [Table 36]


Example No.	Structural formula	LCMS Analysis condition	Retention time (min)	m/z [M+H] ⁺
425		SQD-FA05	0.77	1008
426		SQD-FA05	0.78	904

[1497] <Example 427>

7-[[2,3-Difluoro-4-[2-[2-[[[(2S,3R,4S,5S)-2,3,4,5,6-pentahydroxyhexanoyl]amino]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1498]


[Formula 261]



[1499] First Step

tert-Butyl N-[2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethoxy]ethyl]carbamate

[1500] [Formula 262]



[1501] After tert-butyl N-[2-(2-hydroxyethoxy)ethyl]carbamate (28 mg, 0.137 mmol), 7-[(2,3-difluoro-4-hydroxyphenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (30 mg, 0.046 mmol), and TMAD (15.7 mg, 0.091 mmol) were dissolved in THF (1 mL), tributylphosphine (18.5 mg, 0.091 mmol) was added. After the mixture was stirred at room temperature for 5 hours, the resultant was concentrated, and the residue was purified by C18 reverse-phase column chromatography to obtain the title compound (28 mg, 74%) as a white amorphous solid.

LCMS: m/z 845[M+H]⁺


[1502] HPLC retention time: 1.25 minutes (analysis condition SQD-FA05)

[1503] Second Step

7-[[4-[2-(2-Aminoethoxy)ethoxy]-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1504]

[Formula 263]



[1505] tert-Butyl N-[2-[2-[2,3-difluoro-4-[[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethoxy]ethyl]carbamate (30 mg, 0.036 mmol) was dissolved in ethyl acetate (0.3 mL), then 4N-HCl ethyl acetate (1 mL) was added, and the mixture was stirred at room temperature for 15 hours. The reaction mixture was concentrated, and the resultant residue was used for the next reaction without purification.

[1506] LCMS: m/z 745 $[M+H]^+$

[1507] HPLC retention time: 0.81 minutes (analysis condition SQD-FA05)

[1508] Third Step

7-[[[4-[2-(2-Aminoethoxy)ethoxy]-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (42 mg, 0.054 mmol) and (3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-one (19.2 mg, 0.108 mmol) were dissolved in methanol (1 mL), and the resultant was stirred in a sealed tube at 70°C for 20 hours. Water (0.1 mL) was added to the reaction mixture, and the resultant was purified by C18 reverse-phase column chromatography to obtain the title compound (30 mg, 61%) as a white amorphous solid.

[1509] LCMS: m/z 923[M+H]⁺


[1510] HPLC retention time: 1.00 minute (analysis condition SQD-FA05)

[1511] <Example 428>

7-[[2,3-Difluoro-4-[2-[2-[2-[2-[[2,3,4,5,6-pentahydroxyhexanoyl]amino]ethoxy]ethoxy]ethoxy]ethoxy]phenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-

diazaspiro[4.5]dec-9-ene-9-carboxamide

[1512] [Formula 264]



[1513] The title compound was synthesized by carrying out Steps similar to those of Example 427 by using *tert*-butyl *N*-[2-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]ethyl]carbamate as a starting material.


[1514] LCMS: m/z 1011[M+H]⁺

[1515] HPLC retention time: 1.01 minutes (analysis condition SQD-FA05)

[1516] <Example 429>


7-[[2,3-Difluoro-4-[2-[2-[2-methoxyethyl(methyl)amino]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-*N*-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1517] [Formula 265]



[1518] First Step

[1519] [Formula 266]



[1520] 2-[2-(4-Methylphenyl)sulfonyloxyethoxy]ethyl 4-methylbenzenesulfonate (79 mg, 0.192 mmol) and cesium carbonate (15.6 mg, 0.048 mmol) were added to a solution of 7-[[2,3-difluoro-4-(2,2,2-trifluoroethoxy)phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-*N*-[4-(trifluoromethyl)-

2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (Second Step of Example 332) (31.5 mg, 0.048 mmol) in acetonitrile (1 mL), and the mixture was stirred at 50°C for 4 hours. After the reaction was completed, the reaction mixture was neutralized with formic acid, and the reaction mixture was purified by C18 reverse-phase column chromatography to obtain 2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethoxy]ethyl 4-methylbenzenesulfonate (35.9 mg, 83%).

[1521] LCMS: m/z 900[M+H]⁺

HPLC retention time: 1.73 minutes (analysis condition SMD-TFA05)

[1522] Second Step

2-Methoxy-N-methyl ethanamine (0.043 mL, 0.399 mmol) was added to a solution of 2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethoxy]ethyl 4-methylbenzenesulfonate (35.9 mg, 0.04 mmol) in acetonitrile (0.6 mL), and the mixture was stirred at 50°C for 6 hours. After the reaction was completed, the reaction mixture was purified by C18 reverse-phase column chromatography to obtain the title compound (22.3 mg, 68%).

[1523] LCMS: m/z 817[M+H]⁺






HPLC retention time: 1.43 minutes (analysis condition SMD-TFA05)

[1524] <Examples 430 to 434>

Appropriate alkylating agents and commercial amines were used to synthesize the compounds described in the following Table by carrying out operations similar to those of Example 429.

[1525]


[Table 37]

Example No.	Structural formula	LCMS Analysis condition	Retention time (min)	m/z
				[M+H] ⁺
430		SMD-TFA05	1.45	861
431		SMD-TFA05	1.46	905
432		SMD-TFA05	1.46	949
433		SQD-FA05	0.79	923
434		SQD-FA05	0.81	1055

[1526] <Example 435>

2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]ethanesulfonic acid

[1527] [Formula 267]



[1528] 7-[[4-(2-Bromoethoxy)-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-

[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (20 mg, 26.0 μ mol) (Reference Example 109) was dissolved in N,N-dimethyl formamide (200 μ L), then potassium phosphate (27.8 mg, 0.13 mmol) and tetrabutylammonium iodide (1.0 mg, 2.6 μ mol) were added, and the mixture was stirred at 80°C for 3 hours. After the reaction mixture was diluted with formic acid and water and filtered, the resultant was purified by HPLC to obtain the title compound (8.0 mg, 38%).

[1529] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

Column: YMC-Actus Triart C18 (50 $\times$ 30 mmI.D., S - 5 μ m, 12 nm)


LCMS: m/z 809.1[M+H]⁺

HPLC retention time: 1.32 minutes (analysis condition SMD-TFA05)

[1530] <Example 436>

2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]ethanesulfonic acid

[1531] [Formula 268]




[1532] First Step

(2R)-2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]-3-methoxy-3-oxopropane-1-sulfonic acid

[1533]

[Formula 269]



[1534] (2R)-2-[2-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]-3-methoxy-3-oxopropane-1-sulfonic acid was synthesized from 7-[[4-(2-bromoethoxy)-2,3-difluorophenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (Reference Example 109) and (R)-2-amino-3-methoxy-3-oxopropane-1-sulfonic acid hydrochloride in a manner similar to that of Example 381.

[1535] LCMS: m/z 867.0[M+H]⁺

HPLC retention time: 1.37 minutes (analysis condition SMD-TFA05)

[1536] Second Step

The title compound was synthesized from (2R)-2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]-3-methoxy-3-oxopropane-1-sulfonic acid in a manner similar to that of Reference Example 381.


[1537] LCMS: m/z 853.0[M+H]⁺

HPLC retention time: 1.31 minutes (analysis condition SMD-TFA05)

[1538] <Reference Example 110>

2-Chloro-N-(2-methoxyethyl)-N-methylethanamine hydrochloride

[1539] [Formula 270]



[1540] First Step

2-Methoxy-N-methylethanamine (17.0 g, 191 mmol) was dissolved in toluene

(381 mL), then 2-bromoethanol (13.6 mL, 191 mmol) and triethylamine (26.6 mL, 191 mmol) were added, and the mixture was stirred at 100°C for 2 hours. The reaction mixture was dried over magnesium sulfate, and the resultant was filtered and concentrated at reduced pressure to obtain 2-((2-methoxyethyl)(methyl)amino)ethanol (16.5g) as a crude product.

[1541] Second Step


2-((2-Methoxyethyl)(methyl)amino)ethanol (16.4 g, 124 mmol) was dissolved in ethyl acetate (124 mL), then thionyl chloride (13.4 mL, 186 mmol) was added at 0°C, and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated at reduced pressure to obtain the title compound (20.5 g, 88%).

[1542] ¹H-NMR (DMSO-D₆) δ: 10.36 (1H, brs), 4.00 (2H, t, J = 6.8 Hz), 3.70 (2H, t, J = 4.9 Hz), 3.51-3.43 (4H, m), 3.30 (3H, s), 2.81 (3H, s).

[1543] <Example 437>


7-[[2,3-Difluoro-5-iodo-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1544] [Formula 271]



[1545] First Step

[1546] [Formula 272]



[1547] 6-[(2,3-Difluoro-4-hydroxyphenyl)methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-

(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide (Reference Example 107) (500 mg, 0.76 mmol) was dissolved in N,N-dimethyl formamide (2.54 mL), then N-iodosuccinimide (222 mg, 0.99 mmol) was added, and the mixture was stirred under nitrogen atmosphere at room temperature for 2 hours. Methanol was added to the reaction mixture, and the mixture was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain 7-[(2,3-difluoro-4-hydroxy-5-iodophenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (569 mg, 96%).

[1548] LCMS: m/z 784.1[M+H]⁺

HPLC retention time: 1.67 minutes (analysis condition SMD-TFA05)

Second Step

7-[(2,3-Difluoro-4-hydroxy-5-iodophenyl)methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (341 mg, 0.43 mmol), 2-chloro-N-(2-methoxyethyl)-N-methylethanamine hydrochloride (82.0 mg, 0.43 mmol), cesium carbonate (425 mg, 1.31 mmol), tetrabutylammonium iodide (16.1 mg, 44.0 μ mol), and methanol (17.7 μ L) were dissolved in acetonitrile (4.35 mL), and the mixture was stirred under nitrogen atmosphere at 65°C for 1.5 hours. An aqueous formic acid solution was added to the reaction mixture, and the resultant was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain the title compound (341 mg, 87%).


[1549] LCMS: m/z 899.2[M+H]⁺

HPLC retention time: 1.46 minutes (analysis condition SMD-TFA05)

<Example 438>


7-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]-5-[3-methoxypropyl(methyl)carbamoyl]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1550] [Formula 273]



[1551] First Step

[1552] [Formula 274]



[1553] 7-[[2,3-Difluoro-5-iodo-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (140 mg, 156 μmol), a tris(dibenzylideneacetone)dipalladium chloroform complex (16.1 mg, 16.0 μmol), xantphos (9.0 mg, 16.0 μmol), lithium chloride (39.6 mg, 0.94 mmol), lithium formate (40.5 mg, 0.78 mmol), and potassium trimethylsilanolate (60.0 mg, 0.47 mmol) were dissolved in N,N-dimethyl formamide (1.04 mL), then acetic anhydride (58.5 μL , 0.62 mmol) was added, and the mixture was stirred under nitrogen atmosphere at 85°C for 30 minutes. N,N-dimethyl formamide and methanol were added to the reaction mixture, and the resultant was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain 3,4-difluoro-5-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]-2-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzoate (98.4 mg, 77%).

[1554] LCMS: m/z 817.3[M+H]⁺

HPLC retention time: 1.41 minutes (analysis condition SMD-TFA05)

Second Step

3,4-Difluoro-5-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-

(trifluoromethyl)pyrimidin-4-yl]phenyl]carbonyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]-2-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzoate (15.0 mg, 18.0 μmol), 3-methoxy-N-methylpropan-1-amine (3.79 mg, 37.0 μmol), 2-(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate (10.5 mg, 28.0 μmol), and N-ethyl-N-isopropylpropan-2-amine (9.6 μL , 55.0 μmol) were dissolved in N,N-dimethyl formamide (100 μL), and the mixture was stirred under nitrogen atmosphere at 40°C for 2 hours. An aqueous formic acid solution was added to the reaction mixture, and the resultant was purified by HPLC to obtain the title compound (8.0 mg, 38%).

[1555] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

Column: YMC-Actus Triart C18 (50 $\times$ 30 mmI.D., S - 5 μm , 12 nm)

LCMS: m/z 902.1[M+H]⁺


HPLC retention time: 1.17 minutes (analysis condition SMD-FA05)

<Reference Example 111>

5-(2-Methoxyethoxy)-2-[5-methyl-6-(trifluoromethyl)pyrimidin-4-yl]-4-


(trifluoromethyl)aniline

[1556] [Formula 275]



[1557] First Step

[1558] [Formula 276]



[1559] 2-Iodo-5-(2-methoxyethoxy)-4-(trifluoromethyl)aniline (1.30 g, 3.60 mmol)

(Reference Example 13), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (2.74 g, 10.8 mmol), dicyclohexyl(2',4',6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphine (172 mg, 0.36 mmol), potassium acetate (1.06 g, 10.8 mmol), and palladium acetate (40.0 mg, 0.18 mmol) were dissolved in dioxane (12.0 mL), and the mixture was stirred under nitrogen atmosphere at 110°C for 4 hours. Ethyl acetate was added to the reaction mixture, the mixture was filtered, the resultant was concentrated at reduced pressure, and the resultant residue was purified by C18 reverse-phase column chromatography (acetonitrile/water) to obtain 5-(2-methoxyethoxy)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-4-(trifluoromethyl)aniline (183 mg, 14%).

[1560] LCMS: m/z 362.3[M+H]⁺

HPLC retention time: 0.99 minutes (analysis condition SQD-FA05)

[1561] Second Step

5-(2-Methoxyethoxy)-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-4-(trifluoromethyl)aniline (183 mg, 0.51 mmol), 4-chloro-5-methyl-6-(trifluoromethyl)pyrimidine (Reference Example 91) (199 mg, 1.01 mmol), a 1,1'-bis(diphenylphosphino)ferrocene-palladium (II) dichloride-dichloromethane complex (41.7 mg, 51.0 μ mol), and potassium carbonate (210 mg, 1.52 mol) were dissolved in dioxane (2.94 mL) and water (0.44 mL), and the mixture was stirred under nitrogen atmosphere at 90°C for 1 hour. Water was added to the reaction mixture, and the resultant was extracted with ethyl acetate. The organic layer was washed with water and a brine, the aqueous layer was separated by a phase separator, the resultant was concentrated at reduced pressure, and the resultant residue was purified by column chromatography on silica gel (hexane/ethyl acetate) to obtain the title compound (180 mg, 90%).

[1562] LCMS: m/z 396.1[M+H]⁺


HPLC retention time: 1.24 minutes (analysis condition SMD-TFA05)

[1563] <Reference Example 112>

7-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxylic acid 2-methylpropyl

ester

[1564] [Formula 277]



[1565] Methyl 1-[[[(E)-[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methylideneamino]-methylamino]cyclopentane-1-carboxylate hydrochloride (Reference Example 105) was used, and operations similar to those of Fourth Step of Example 237 were carried out to synthesize the title compound.


[1566] LCMS: m/z 540.2[M+H]⁺

HPLC retention time: 1.11 minutes (analysis condition SMD-TFA05)

[1567] <Example 439>

7-[[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-N-[5-(2-methoxyethoxy)-2-[5-methyl-6-(trifluoromethyl)pyrimidin-4-yl]-4-(trifluoromethyl)phenyl]-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1568] [Formula 278]



[1569] 7-[[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-6,7-diazaspiro[4.5]dec-9-ene-9-carboxylic acid 2-methylpropyl ester and 5-(2-methoxyethoxy)-2-[5-methyl-6-(trifluoromethyl)pyrimidin-4-yl]-4-(trifluoromethyl)aniline were used, and operations similar to those of Fifth Step of Example 237 were carried out to synthesize the title compound.


[1570] LCMS: m/z 861.4[M+H]⁺

HPLC retention time: 1.40 minutes (analysis condition SMD-TFA05)

<Example 440>

9-[[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]-5-(3-

pyridinyl]phenyl]methyl]-6-hydroxy-10-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-9,10-diazaspiro[4.5]dec-6-ene-7-carboxamide
[1571] [Formula 279]



[1572] 7-[[2,3-Difluoro-5-iodo-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide (10 mg, 11.0 μ mol) (Example 437) was dissolved in 1,2-dimethoxyethane (80 μ L), ethanol (80 μ L), and water (80 μ L), then 3-pyridinyl borate (2.1 mg, 17.0 μ mol), tetrakis(triphenylphosphine)palladium (3.9 mg, 3.3 μ mol), and sodium carbonate (3.9 mg, 33 μ mol) were added, and the mixture was stirred at 80°C for 16 hours. After the reaction mixture was diluted with formic acid and water and filtered, the resultant was purified by HPLC to obtain the title compound (2.3 mg, 24%).

[1573] Purification condition: HPLC

Mobile phase: MeCN/water (0.1% formic acid)

Column: YMC-Actus Triart C18 (50 $\times$ 30 mmI.D., S - 5 μ m, 12 nm)

LCMS: m/z 850[M+H]⁺


HPLC retention time: 1.26 minutes (analysis condition SMD-TFA05)

<Example 441>

7-[[2,3-Difluoro-4-[2-[2-[2-[2-[methyl-[(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]-2-oxoethoxy]ethoxy]ethoxy]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide

[1574]

[Formula 280]



[1575] The title compound was synthesized by carrying out Steps and operations similar to those of Example 422 by using 2-[2-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]ethoxy]ethanol and tert-butyl 2-bromoacetate as a starting material.

[1576] LCMS: m/z 1069[M+H]⁺

[1577] HPLC retention time: 1.00 minute (analysis condition SQD-FA05)





[1578] <Examples 442 to 445>

The aniline compound of Example 417 and appropriate aldehydes or ketones were used, and the following operations were carried out to synthesize the compounds described in the following Table.

[1579] 7-(2,3-Difluoro-4-((2-((2-methoxyethyl)(methyl)amino)ethyl)amino)benzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide and the corresponding aldehyde or ketone were dissolved in tetrahydrofuran, and 18 M sulfuric acid of an amount equivalent to one-fifth of the amount of tetrahydrofuran was added at 0°C. Subsequently, sodium tetrahydroborate was added in three batches, and the mixture was stirred at 0°C or at room temperature for 3 hours and more. Triethylamine, water, and dimethylsulfoxide were added to the reaction mixture, and the resultant solution was purified by C18 reverse-phase column chromatography to synthesize the compounds of the Examples described in the following Table.

[1580]



[Table 38]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
442		SQD-AA05	1.21	786
443		SQD-AA05	1.22	800
444		SQD-AA05	1.24	814
445		SQD-AA05	1.23	814

[1581] <Examples 446 and 447>

The appropriate iodide derivative of Example 417, 3-morpholinopropan-1-amine, and 2-morpholinoethanamine were used, and operations similar to those of Example 417 were carried out to synthesize the compounds described in the following Table.

[1582] [Table 39]

Example No.	Structural formula	LCMS analysis condition No.	Retention time (min)	m/z [M+H] ⁺
446		SQD-AA50	0.88	784
447		SQD-AA50	0.87	770


[1583] <Reference Example 113>

Methyl 1-((methylamino)cyclopentanecarboxylate 4-toluenesulfonate

First Step

Methyl 1-((tert-butoxycarbonyl)amino)cyclopentane carboxylate

[1584] [Formula 281]




[1585] 1-Aminocyclopentanecarboxylic acid hydrochloride (3.00 g, 18.1 mmol) was dissolved in methanol (18.0 mL), then thionyl chloride (1.32 mL, 18.1 mmol) was added at 0°C, and the mixture was stirred under nitrogen atmosphere at 50°C for 3 hours. The reaction mixture was concentrated at reduced pressure, and water (6.00 mL) and triethylamine (7.55 mL, 54.3 mmol) were added to the resultant residue. Di-tert-butyl dicarbonate (3.95 g, 18.1 mmol) and tert-butyl methyl ester (24.0 mL) were added to this solution, and the mixture was stirred at 50°C for 1 hour. The aqueous layer was separated, tetrahydrofuran (9.00 mL) was added to the organic layer, and the resultant was concentrated at reduced pressure. Tetrahydrofuran (15.0 mL) was added to the resultant residue, and the resultant was concentrated at reduced pressure to obtain the title compound as pale yellow oil.

[1586] ¹H-NMR (DMSO-D₆) δ: 7.28 (1H, brs), 3.58 (3H, s), 2.10-1.80 (4H, m), 1.67-1.55 (4H, m), 1.36 (9H, s).

[1587] Second Step

Methyl 1-((tert-butoxycarbonyl)methylamino)cyclopentanecarboxylate

[1588] [Formula 282]



[1589] Methyl 1-((tert-butoxycarbonyl)amino)cyclopentanecarboxylate obtained in First Step was dissolved in 1-methyl 2-pyrrolidinone (24.0 mL), then sodium hydroxide (1.45 g, 36.2 mmol) and methyl iodide (3.38 mL, 54.3 mmol) were added, and the mixture was stirred at 50°C for 7 hours. Water (30.0 mL) was added to this reaction mixture at room temperature, and the resultant was extracted with isopropyl acetate (15.0 mL) twice. The resultant organic layer was washed serially with a 10% aqueous sodium thiosulfate solution


(15.0 mL) and a 10% aqueous sodium chloride solution (15.0 mL). The resultant organic layer was concentrated at reduced pressure to obtain the title compound (5.88 g, 83%).

[1590] $^1\text{H-NMR}$ (DMSO-D_6) δ : 3.60 (3H, s), 2.87 (3H, s), 2.20-2.10 (2H, m), 1.96-1.87 (2H, m), 1.75-1.60 (4H, m), 1.34 (9H, s).

[1591] Third Step

Methyl 1-(methylamino)cyclopentanecarboxylate 4-toluenesulfonate

[1592] [Formula 283]




[1593] The title compound was synthesized from 1-((tert-butoxycarbonyl)amino)cyclopentanecarboxylic acid in a manner similar to that of Reference Examples 87 and 88.

[1594] <Reference Example 114>

2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzaldehyde hydrochloride

[1595] [Formula 284]



[1596] 2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]benzaldehyde (200 mg, 0.73 mmol) was dissolved in 1 mL dimethoxyethane. Pyridine hydrochloride (85 mg, 0.73 mmol) was added, and the mixture was stirred. The generated crystal was filtered and dried at reduced pressure to obtain the title compound (68 mg, 30%).

[1597] $^1\text{H-NMR}$ (DMSO-D_6) δ : 10.99 (1H, brs), 10.07 (1H, s), 7.73 (1H, m), 7.30 (1H, m), 4.67 (2H, t, $J = 4.9$ Hz), 3.75 (2H, t, $J = 5.1$ Hz), 3.30 (3H, s), 3.30-3.74 (4H, m), 2.87 (3H, brd, $J = 3.7$ Hz).

[1598] Pharmacological test

Test Example 1: Inhibitory effect on $^{33}\text{PO}_4$ uptake into rat small-intestinal brush border membrane vesicles

Brush border membrane vesicles (BBMV_s) were prepared using the upper region of the small intestine of each Wistar female rat (4-5 weeks old). The preparation of BBMV_s was performed according to the method of Murer et al. (Murer H, Hopfer U, Kinne R. Sodium/proton antiport in brush-border-membrane vesicles isolated from rat small intestine and kidney. 1976. J Am Soc Nephrol. 1998 Jan; 9 (1): 143-50). The $^{33}\text{PO}_4$ transport activity of the small-intestinal BBMV_s was determined by the rapid filtration method. Each test compound was added at a final concentration of 1 μM to buffer A (110 mM NaCl, 60 mM mannitol, and 10 mM HEPES (pH 7.5)) containing 340 kBq/mL $^{33}\text{PO}_4$, or DMSO was added to buffer B (110 mM KCl, 60 mM mannitol, and 10 mM HEPES (pH 7.5)) containing 340 kBq/mL $^{33}\text{PO}_4$, and each resulting solution was added to the BBMV_s sample and reacted for 60 seconds. Then, ice-cold buffer C (110 mM NaCl, 1 mM KH_2PO_4 , and 10 mM HEPES (pH 7.5)) was added to each reaction mixture, which was immediately suction-filtered through a Millipore filter. The filter was washed with buffer C, and each sample was then dissolved in a liquid scintillator. The amount of $^{33}\text{PO}_4$ uptake into BBMV_s was measured using a liquid scintillation counter. The rate of inhibition was determined according to the following expression:

$$\text{[1599] Rate (\% of inhibition) = } (1 - (\text{Amount of } ^{33}\text{PO}_4 \text{ uptake into the test compound-supplemented and buffer A-treated BBMV}_s - \text{Amount of } ^{33}\text{PO}_4 \text{ uptake into the DMSO-supplemented and buffer B-treated BBMV}_s) / (\text{Amount of } ^{33}\text{PO}_4 \text{ uptake into the DMSO-supplemented and buffer A-treated BBMV}_s - \text{Amount of } ^{33}\text{PO}_4 \text{ uptake into the DMSO-supplemented and buffer B-treated BBMV}_s)) \times 100$$

The rate of inhibition of $^{33}\text{PO}_4$ uptake into the rat small-intestinal brush border membrane vesicles by 1 μM of the test compound (the rate of uptake inhibition) is shown in Tables 40-1 to 40-8.

[1600]

[Table 40-1]

Example No.	Rate of uptake inhibition (%) (1 μ M)
1	28
2	19
3	11
4	32
5	54
6	48
7	69
8	37
9	63
10	37
11	22
12	27
13	58
14	29
15	42
16	38
17	33
18	12
19	41
20	21
21	61
22	50
23	84
24	76
25	82
26	72
27	76
28	72
29	75
30	71
31	76

Example No.	Rate of uptake inhibition (%) (1 μ M)
32	73
33	77
34	99
35	79
36	80
37	89
38	90
39	90
40	87
41	84
42	67
43	79
44	94
45	73
46	71
47	91
48	87
49	85
50	79
51	72
52	57
53	72
54	72
55	85
56	88
57	71
58	46
59	84
60	78
61	71
62	73

[1601]

[Table 40-2]

Example No.	Rate of uptake inhibition (%) (1 μ M)
63	72
64	77
65	98
66	96
67	91
68	84
69	74
70	79
71	97
72	92
73	74
74	82
75	74
76	77
77	80
78	71
79	71
80	75
81	74
82	100
83	90
84	91
85	71
86	83
87	84
88	39
89	67
90	91
91	87
92	81
93	56

Example No.	Rate of uptake inhibition (%) (1 μ M)
94	73
95	80
96	83
97	89
98	53
99	82
100	69
101	57
102	65
103	78
104	73
105	72
106	79
107	54
108	54
109	52
110	57
111	56
112	43
113	84
114	86
115	85
116	72
117	78
118	81
119	71
120	66
121	72
122	82
123	87
124	81

[1602]

[Table 40-3]

Example No.	Rate of uptake inhibition (%) (1 μ M)
125	76
126	71
127	95
128	72
129	63
130	72
131	87
132	71
133	83
134	94
135	90
136	80
137	92
138	93
139	88
140	77
141	83
142	76
143	71
144	72
145	78
146	71
147	57
148	75
149	80
150	87
151	92
152	54
153	73
154	73
155	77

Example No.	Rate of uptake inhibition (%) (1 μ M)
156	59
157	89
158	84
159	89
160	77
161	94
162	73
163	79
164	64
165	75
166	73
167	73
168	72
169	73
170	73
171	78
172	72
173	73
174	87
175	85
176	75
177	85
178	81
179	83
180	75
181	85
182	86
183	82
184	79
185	83
186	88

[1603]

[Table 40-4]

Example No.	Rate of uptake inhibition (%) (1 μ M)
187	80
188	93
189	73
190	84
191	87
192	88
193	82
194	80
195	76
196	90
197	82
198	75
199	79
200	80
201	77
202	75
203	91
204	87
205	86
206	76
207	73
208	90
209	76
210	71
211	72
212	98
213	79
214	91
215	86
216	76
217	83

[1604]

Example No.	Rate of uptake inhibition (%) (1 μ M)
218	82
219	95
220	96
221	82
222	97
223	77
224	74
225	76
226	71
227	82
228	77
229	73
230	75
231	80
232	79
233	75
234	73
235	65
236	60
237	51
238	47
239	48
240	72
241	86
242	69
243	63
244	71
245	40
246	52
247	51
248	45

[Table 40-5]

Example No.	Rate of uptake inhibition (%) (1 μ M)
249	50
250	52
251	43
252	53
253	40
254	58
255	80
256	79
257	71
258	87
259	52
260	70
261	77
262	43
263	66
264	48
265	52
266	62
267	71
268	63
269	80
270	75
271	60
272	43
273	41
274	43
275	42
276	68
277	41
278	69
279	53

Example No.	Rate of uptake inhibition (%) (1 μ M)
280	50
281	41
282	55
283	45
284	46
285	62
286	50
287	47
288	61
289	55
290	59
291	47
292	44
293	47
294	40
295	21
296	17
297	36
298	10
299	45
300	74
301	79
302	79
303	42
304	51
305	66
306	58
307	41
308	63
309	68
310	63

[1605]

[Table 40-6]

Example No.	Rate of uptake inhibition (%) (1 μ M)
311	65
312	74
313	65
314	72
315	63
316	64
317	57
318	43
332	3
333	29
334	22
335	30
336	44
337	47
338	66
339	66
340	63
341	64
342	43
343	68
344	58
345	73
346	42
347	28
348	31
349	55
350	45
351	22
352	27
353	26

[1606]

Example No.	Rate of uptake inhibition (%) (1 μ M)
354	31
355	36
356	58
357	45
358	34
359	37
360	21
361	20
362	17
363	11
364	36
365	34
366	38
367	43
368	26
369	43
370	41
371	38
372	38
373	27
374	42
375	31
376	33
377	29
378	35
379	37
380	43
381	28
382	21
383	26

[Table 40-7]

Example No.	Rate of uptake inhibition (%) (1 μ M)
384	39
385	39
386	36
387	23
388	15
389	29
390	58
391	42
392	47
393	45
394	45
395	39
396	38
397	40
398	34
399	42
400	36
401	39
402	54
403	71
404	43
405	30
406	27
407	30
408	47
409	43
410	49
411	32
412	52
413	34

Example No.	Rate of uptake inhibition (%) (1 μ M)
414	27
415	21
416	50
417	48
418	29
419	22
420	19
421	24
422	29
423	5
424	17
425	5
426	24
427	12
428	14
429	41
430	29
431	26
432	11
433	19
434	24
435	11
436	12
437	45
438	
439	35
440	28
441	22
442	24
443	28

[1607]

[Table 40-8]

Example No.	Rate of uptake inhibition (%) (1 μ M)	Example No.	Rate of uptake inhibition (%) (1 μ M)
444	24	446	31
445	29	447	50

[1608] Test Example 2: Inhibitory effect on $^{33}\text{PO}_4$ uptake into human NaPi-IIb-expressing cell

CHO cells were transfected with human NaPi-IIb expression plasmids, and a stably human NaPi-IIb-expressing cell line was obtained using G418. The human NaPi-IIb-expressing cells were inoculated to a 96-well plate and incubated overnight in a CO_2 incubator. The medium was replaced with buffer A (145 mM choline chloride, 3 mM KCl, 1 mM CaCl_2 , 0.5 mM MgCl_2 , 5 mM glucose, and 5 mM MES (pH 6.5)) and then replaced with buffer B (145 mM NaCl, 3 mM KCl, 1 mM CaCl_2 , 0.5 mM MgCl_2 , 5 mM glucose, and 5 mM MES (pH 6.5)) supplemented with each test compound at a final concentration of 1, 3, 10, or 30 μM or buffer A supplemented with DMSO. After a given time, a 1/20 volume of buffer A containing $^{33}\text{PO}_4$ was added thereto and reacted at room temperature. After washing with ice-cold buffer A, a liquid scintillator was added to each reaction mixture, and the amount of $^{33}\text{PO}_4$ uptake was measured using TopCount. The rate of inhibition was determined according to the following expression:

[1609] Rate (%) of inhibition = $(1 - (\text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the test compound-supplemented and buffer B-treated well} - \text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the DMSO-supplemented and buffer A-treated well}) / (\text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the DMSO-supplemented and buffer B-treated well} - \text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the DMSO-supplemented and buffer A-treated well})) \times 100$.

IC_{50} values (μM) were calculated from a straight line joining two points intersecting the 50% rate of inhibition. The IC_{50} values of some compounds against $^{33}\text{PO}_4$ uptake into the human NaPi-IIb-expressing cells are shown in Tables 41-1 to 41-7.

[1610]

[Table 41-1]

Example No.	IC50 (μM)
13	7.3
14	8.9
21	2.8
22	7.9
23	6.9
24	8.1
25	10.1
26	2.2
27	9.0
28	4.1
29	2.0
30	2.3
31	3.3
32	2.8
33	6.0
34	7.0
35	12.6
36	6.1
37	3.7
38	6.7
39	8.5
40	6.5
41	6.6
42	5.0
43	8.3
44	30.0
45	13.0
46	14.1
47	7.1
48	16.3
49	12.9
50	8.3

Example No.	IC50 (μM)
51	16.0
52	16.8
53	9.5
54	2.3
55	15.3
56	7.2
57	10.3
58	30.0
59	7.4
60	8.8
61	5.6
62	6.4
63	8.7
64	6.5
65	16.2
66	17.3
67	2.0
68	2.6
69	8.9
70	8.4
71	7.8
72	8.3
73	2.3
74	4.1
76	6.6
77	8.9
78	5.2
79	6.7
80	12.5
81	17.5
82	11.1
83	10.6

[1611]

[Table 41-2]

Example No.	IC50 (μM)
84	7.9
85	6.2
86	5.3
87	8.3
89	4.8
90	2.4
91	15.7
92	20.2
93	17.4
94	8.8
95	5.4
96	2.8
97	2.7
98	10.6
99	4.3
100	6.7
101	9.4
102	10.9
103	11.0
104	10.8
105	6.3
106	12.0
107	25.9
108	26.1
109	9.4
110	19.8
111	20.0
112	30.0
113	15.5
114	7.9
115	6.9

Example No.	IC50 (μM)
116	8.1
117	9.5
118	9.1
119	9.2
120	20.5
121	16.0
122	8.6
123	14.8
124	2.5
125	1.3
126	5.5
127	17.6
128	12.0
129	13.7
130	8.9
131	15.3
132	5.0
133	13.1
134	19.7
135	12.3
136	17.6
137	19.7
138	11.9
139	17.0
140	16.2
141	7.8
142	17.6
143	6.7
144	2.1
145	7.1
146	10.2

[1612]

[Table 41-3]

Example No.	IC50 (μM)
147	21.5
148	8.5
149	19.7
150	9.0
151	8.4
152	20.6
153	6.2
154	2.5
155	2.3
156	3.2
157	19.1
158	12.8
159	11.4
160	7.3
161	11.5
162	15.1
163	6.8
164	5.0
165	1.8
166	10.5
167	4.3
168	2.6
169	2.6
170	2.2
171	2.2
172	2.4
173	1.9
174	2.2
175	3.4
176	4.0
177	11.1

Example No.	IC50 (μM)
178	8.2
179	18.9
180	5.7
181	4.8
182	5.1
183	9.1
184	3.1
185	4.9
186	5.5
187	6.7
188	6.1
189	5.5
190	5.1
191	14.5
192	16.5
193	6.4
194	19.3
195	4.6
196	2.8
197	7.0
198	2.4
199	1.8
200	1.3
201	1.3
202	6.1
203	2.0
204	1.0
205	1.6
206	2.0
207	6.5
208	6.6

[1613]

[Table 41-4]

Example No.	IC50 (μM)
209	4.8
210	1.7
211	5.7
212	2.9
213	6.4
214	7.0
215	6.2
216	8.0
217	4.7
218	7.2
219	4.2
220	12.2
221	6.6
222	17.6
223	17.9
224	7.2
225	5.1
226	3.1
227	3.7
228	5.7
229	1.5
230	2.1
231	5.6
232	4.2
233	2.3
234	9.4
235	4.8
236	24.7
237	12.8
238	8.5
239	8.5

Example No.	IC50 (μM)
240	9.1
241	16.5
242	7.8
243	7.4
244	7.3
245	7.7
246	8.6
247	8.2
248	17.6
249	9.8
250	17.9
251	7.6
252	9.5
253	4.4
254	9.7
255	9.8
256	4.4
257	6.0
258	2.7
259	10.0
260	6.4
261	7.7
262	9.9
263	3.0
264	19.4
265	13.3
266	11.6
267	7.5
268	7.4
269	4.5
270	14.6

[1614]

[Table 41-5]

Example No.	IC ₅₀ (μM)
271	17.3
272	19.5
273	13.2
274	17.0
275	16.8
276	15.8
277	17.8
278	6.6
279	18.7
280	11.9
281	18.8
282	11.1
283	17.7
284	7.8
285	8.5
286	15.8
287	8.0
288	11.0
289	16.3
290	7.0
291	14.7
292	18.4
293	10.7
294	13.4
295	17.0
296	10.0
297	18.3
298	>30
299	18.8
300	9.3
301	2.6
302	7.8

Example No.	IC ₅₀ (μM)
303	16.0
304	3.5
305	5.9
306	8.4
307	4.6
308	12.6
309	6.5
310	4.7
311	8.2
312	6.3
313	18.3
314	11.8
315	9.8
316	14.2
317	9.6
318	8.6
332	23.9
333	19.5
334	10.3
335	8.7
336	19.7
337	17.0
338	13.4
339	19.1
340	10.0
341	7.6
342	8.8
343	21.0
344	13.7
345	7.7
346	6.9
347	25.2

[1615]

[Table 41-6]

Example No.	IC ₅₀ (μM)
348	26.1
349	8.3
350	16.9
351	25.5
352	16.7
353	23.9
354	26.0
355	15.0
356	21.9
357	21.8
358	9.2
359	18.4
360	23.1
361	20.2
362	18.8
363	28.0
364	20.9
365	12.2
366	16.3
367	14.0
368	14.7
369	21.8
370	18.2
371	8.5
372	16.3
373	23.5
374	16.4
375	17.8
376	21.4
377	8.7
378	18.0

Example No.	IC ₅₀ (μM)
379	27.3
380	20.2
381	22.0
382	21.0
383	20.9
384	11.0
385	6.2
386	18.9
387	10.7
388	6.9
389	24.6
390	7.8
391	7.0
392	15.8
393	8.1
394	6.7
395	15.2
396	9.5
397	8.6
398	9.9
399	7.8
400	9.2
401	14.5
402	12.8
403	6.3
404	10.1
405	27.7
406	23.1
407	6.9
408	14.6
409	10.5

[1616]

[Table 41-7]

Example No.	IC ₅₀ (μM)
410	8.4
411	12.8
412	19.0
413	11.3
414	4.6
415	18.5
416	5.8
417	2.2
418	10.9
419	24.8
420	18.6
421	19.5
423	20.0
424	29.0
425	28.1
426	26.1
427	17.4
428	23.0
429	7.7
430	4.6
431	6.2
432	6.3
433	18.6
434	19.5

Example No.	IC ₅₀ (μM)
435	17.1
436	14.0
437	19.5
438	21.9
439	18.9
440	22.6
441	21.4
442	6.3
443	8.9
444	7.4
445	9.2
446	11.1
447	15.8

[1617] Test Example 3: Inhibitory effect on ³³PO₄ uptake into human PiT-1-expressing cell

CHO cells were transfected with human PiT-1 expression plasmids or empty vectors to prepare human PiT-1-expressing cells and empty vector-expressing cells. The human PiT-1-expressing cells or the empty vector-expressing cells were inoculated to a 96-well plate and incubated overnight in a CO₂ incubator. The medium was replaced with buffer A (145 mM choline chloride, 3 mM KCl, 1 mM CaCl₂, 0.5 mM MgCl₂, 5 mM glucose, and 5 mM MES (pH 6.5)) and then replaced with buffer B (145 mM NaCl, 3 mM KCl, 1 mM CaCl₂, 0.5 mM MgCl₂, 5 mM glucose, and 5 mM MES (pH 6.5)) supplemented with each compound at a final concentration of 0.24, 1.2, 6, or 30 μM or buffer B supplemented with DMSO. After a given time, a 1/20 volume of buffer A containing ³³PO₄ was added thereto and reacted at room temperature. After washing with ice-cold buffer A, a liquid scintillator

was added to each reaction mixture, and the amount of $^{33}\text{PO}_4$ uptake was measured using TopCount. The rate of inhibition was determined according to the following expression:

[1618] Rate (%) of inhibition = $(1 - (\text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the compound-supplemented well containing the human PiT-1-expressing cells} - \text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the compound-supplemented well containing the empty vector-expressing cells}) / (\text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the DMSO-supplemented well containing the human PiT-1-expressing cells} - \text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the DMSO-supplemented well containing the empty vector-expressing cells})) \times 100$

IC_{50} values (μM) were calculated from a straight line joining two points intersecting the 50% rate of inhibition. The IC_{50} values of some compounds against $^{33}\text{PO}_4$ uptake into the human PiT-1-expressing cells are shown in Table 42.

[1619]

[Table 42]

Example No.	IC ₅₀ (μM)
13	1.9
14	2.2
21	3.3
22	1.8
52	3.1
129	6.5
140	3.8
147	4.4
152	5.3
156	3.9
157	3.8
164	4.1
196	5.6
264	4.6
265	4.6
267	4.1
269	7.9
270	3.4
317	3.3
318	3.3
333	3.6
345	1.2
410	2.9
414	1.9
416	0.9
418	4.1

[1620] Test Example 4: Inhibitory effect on ³³PO₄ uptake into human PiT-2-expressing cell

CHO cells were transfected with human PiT-2 expression plasmids or empty vectors to prepare human PiT-2-expressing cells and empty vector-expressing cells. The human PiT-2-expressing cells or the empty vector-expressing cells were inoculated to a 96-well plate and incubated overnight in a CO₂ incubator. The medium was replaced with buffer A (145 mM choline chloride, 3 mM KCl, 1 mM CaCl₂, 0.5 mM MgCl₂, 5 mM glucose, and 5 mM MES (pH 6.5)) and then replaced with buffer B (145 mM NaCl, 3 mM KCl, 1 mM CaCl₂, 0.5 mM MgCl₂, 5 mM glucose, and 5 mM MES (pH 6.5)) supplemented with each compound at a final concentration of 0.24, 1.2, 6, or 30 μM or buffer B supplemented with

DMSO. After a given time, a 1/20 volume of buffer A containing $^{33}\text{PO}_4$ was added thereto and reacted at room temperature. After washing with ice-cold buffer A, a liquid scintillator was added to each reaction mixture, and the amount of $^{33}\text{PO}_4$ uptake was measured using TopCount. The rate of inhibition was determined according to the following expression:

[1621] Rate (%) of inhibition = $(1 - (\text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the compound-supplemented well containing the human PiT-2-expressing cells} - \text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the compound-supplemented well containing the empty vector-expressing cells}) / (\text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the DMSO-supplemented well containing the human PiT-2-expressing cells} - \text{Amount of } ^{33}\text{PO}_4 \text{ uptake in the DMSO-supplemented well containing the empty vector-expressing cells})) \times 100$

IC_{50} values (μM) were calculated from a straight line joining two points intersecting the 50% rate of inhibition. The IC_{50} values of some compounds against $^{33}\text{PO}_4$ uptake into the human PiT-2-expressing cells are shown in Table 43.

[1622]

[Table 43]

Example No.	IC ₅₀ (μM)
13	2.7
14	2.6
21	3.6
22	3.2
52	4.0
129	8.7
140	3.8
147	8.4
152	3.5
156	7.2
157	3.9
164	4.1
196	6.2
264	3.8
265	4.3
267	4.7
269	10.6
270	3.6
317	2.9
318	3.7
333	10.4
345	1.4
410	3.3
414	1.2
416	1.4
418	4.6

[1623] Test Example 5: Suppressive effect on rise in serum phosphorus concentration of adenine-induced renal failure rat

Adenine was forcibly administered orally to each Wistar male rat (7-8 weeks old) to impair renal functions and thereby prepare a hyperphosphatemia model (Katsumata K, Kusano K, Hirata M, Tsunemi K, Nagano N, Burke SK, Fukushima N. Sevelamer hydrochloride prevents ectopic calcification and renal osteodystrophy in chronic renal failure rats. *Kidney Int.* 2003 Aug; 64 (2): 441-50). Each test compound was mixed at a ratio of 0.1% of mass concentration with feed. The animal was fed with a given amount of the feed for 3 days. A group given feed non-supplemented with a test compound was used as a






reference pathological group, and an adenine-unadministered group given feed non-supplemented with a test compound was used as a normal group. 3 days after the start of administration of the test compound, blood was collected from the jugular vein, and serum was collected. Serum phosphorus concentrations were measured by the Fiske-Subbarow method. The rate of suppression of a rise in serum phosphorus concentration was determined according to the following expression:

[1624] Rate (%) of suppression of a rise in serum phosphorus concentration = $(1 - [(\text{Serum phosphorus concentration of the test compound-treated pathological group}) - (\text{Serum phosphorus concentration of the normal group})] / [(\text{Serum phosphorus concentration of the reference pathological group}) - (\text{Serum phosphorus concentration of the normal group})]) \times 100$

As a result, each test compound was confirmed to have a suppressive effect on a rise in serum phosphorus concentration. Table 44 shows the rate (%) of suppression of a rise in serum phosphorus concentration by each test compound.






[1625]

[Table 44-1]

Example No.	Structural formula	Compound name	Rate of inhibition (%)
13		6-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide	25%
14		7-[[2,3-Difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide	39%
21		(4aR)-1-[(2,3-Difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	24%
22		(4aR)-N-[2-(6-Cyano-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	26%
52		(4aR)-N-[2-(2-Cyanopyridin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	20%

[1626]

[Table 44-2]

129		6-[4-[[[(4aR)-4-Hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl]methyl]-2,3-difluorophenyl]hex-5-ynoic acid	17%
140		(4aR)-1-[[[2,3-Difluoro-4-(3-morpholin-4-ylprop-1-ynyl)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	39%
147		(4aR)-1-[[[4-[3-[(2R)-2,3-Dihydroxypropoxy]propoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	35%
152		(4aR)-1-[[[4-[4-[(2R)-2,3-Dihydroxypropoxy]butoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	26%
156		(4aR)-1-[[[4-[6-[(2R)-2,3-Dihydroxypropoxy]hexoxy]-2,3-difluorophenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	21%

[1627]

[Table 44-3]

157		(4aR)-1-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	30%
164		(4aR)-1-[[2,3-Difluoro-4-(morpholin-4-ylmethyl)phenyl]methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide; hydrochloride	14%
196		(4aR)-N-(4-Bromo-3,5-difluorophenyl)-1-[(3-chloro-2-fluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide	35%
264		(3S)-3-tert-Butyl-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-3H-pyridazine-5-carboxamide	15%
265		(3S)-3-tert-Butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide	11%




[1628]

[Table 44-4]

267		(3S)-3-tert-Butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide	18%
269		(3S)-3-tert-Butyl-N-[4-chloro-2-(6-methylsulfanylpyridin-3-yl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide	15%
270		(3S)-3-tert-Butyl-N-[2-(6-cyano-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide	20%
317		6-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide	22%
318		7-[[2,3-Difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide	32%

[1629]

[Table 44-5]

333		5-[2,3-Difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]pentanoic acid	8%
345		6-[[2,3-Difluoro-4-[2-[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]ethoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide	19%
410		7-[[2,3-Difluoro-4-[2-[methyl(oxetan-3-yl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide	25%




[1630] Test Example 6: Suppressive effect on rise in serum phosphorus concentration of adenine-induced renal failure rat

Each adenine-induced renal failure rat was prepared in the same way as in Test Example 5. Each test compound was mixed at a ratio of 0.05% of mass concentration with feed. The animal was fed with a given amount of the feed for 8 days. A group given feed non-supplemented with a test compound was used as a reference pathological group, and an adenine-unadministered group given feed non-supplemented with a test compound was used as a normal group. 8 days after the start of administration of the test compound, blood was collected from the jugular vein, and serum was collected. Serum phosphorus concentrations were measured by the Fiske-Subbarow method. The rate of suppression of a rise in serum phosphorus concentration was determined according to the following expression:

[1631] Rate (%) of suppression of a rise in serum phosphorus concentration = $(1 - [(\text{Serum phosphorus concentration of the test compound-treated pathological group}) - (\text{Serum phosphorus concentration of the normal group})] / [(\text{Serum phosphorus concentration of the reference pathological group}) - (\text{Serum phosphorus concentration of the normal group})]) \times$

As a result, each test compound was confirmed to have a suppressive effect on a rise in serum phosphorus concentration. Table 45 shows the rate (%) of suppression of a rise in serum phosphorus concentration by each test compound.


[1632] [Table 45]

Example No.	Structural formula	Compound name	Rate of inhibition (%)
414		7-(4-(3-(Dimethylamino)-2,2-dimethylpropoxy)-2,3-difluorobenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide	37%
416		6-[[2,3-Difluoro-4-[1-(2-methoxyethyl)azetidin-3-yl]oxyphenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide	38%
418		7-[[2,3-Difluoro-4-[4-[methyl-[(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]-4-oxobutoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide	38%

CLAIMS

1. A compound represented by the formula (I) or a salt thereof, or a solvate of the compound or the salt:

[Formula 1]



wherein R^1 , R^4 , and R^5 are as defined in any one of the following (1) to (3):

(1) R^1 is a hydrogen atom or C_{1-10} alkyl;

R^4 is a hydrogen atom, C_{1-4} alkyl optionally substituted with one or more substituents R_f , C_{6-10} aryl optionally substituted with one or more substituents R_g , (C_{1-6} alkyl)carbonyl, (C_{6-10} aryl)carbonyl, a group $-C(O)NR^{37}R^{38}$, C_{3-7} cycloalkyl, or 5- to 8-membered heterocycloalkyl; and

R^5 is a hydrogen atom or C_{1-4} alkyl;

(2) R^1 and R^5 together with the carbon atom to which they are attached form a C_{3-6} saturated carbocyclic ring; and

R^4 is as defined above; and

(3) R^1 is a hydrogen atom or linear C_{1-10} alkyl;

R^4 and R^5 together with the carbon atom and the nitrogen atom to which they are attached form a 5- to 8-membered saturated heterocyclic ring, wherein the saturated heterocyclic ring is optionally substituted with one or more substituents R^2 ;

R^3 is C_{1-10} alkyl optionally substituted with one or more substituents R_h , or R^3 is C_{1-4} alkyl substituted with R_e ;

R^{37} and R^{38} are each independently selected from a hydrogen atom and C_{1-3} alkyl;

each R^2 is independently selected from C_{1-5} alkyl and a halogen atom; and/or
two or more substituents R^2 on the 5- to 8-membered saturated heterocyclic ring
may together form C_{1-5} alkylene that links the ring atoms to which they are attached;

each R_h is independently selected from a halogen atom, $(C_{1-4}$ alkoxy)carbonyl, and a
group $-(O(CH_2)_a)_b-C_{1-4}$ alkoxy, wherein a is an integer selected from 2 to 4, and b is an
integer selected from 1 to 4;

R_e is C_{6-10} aryl optionally substituted with one or more substituents R_a , or 5- to 10-
membered heteroaryl optionally substituted with one or more substituents R_a ;

each R_f is independently selected from a halogen atom, hydroxy, cyano, carboxy,
 $(C_{1-6}$ alkoxy)carbonyl, C_{1-6} alkoxy, and C_{6-10} aryl optionally substituted with one or more
substituents R_g ;

each R_g is independently selected from a halogen atom, C_{1-6} alkyl, C_{2-6} alkynyl, and
 C_{1-6} alkoxy, wherein the alkyl, alkynyl, and alkoxy groups are each optionally substituted
with one or more substituents selected from hydroxy and cyano;

each R_a is independently selected from a halogen atom, hydroxy, nitro, cyano, $(C_{1-4}$
alkoxy)carbonyl, 3- to 10-membered heterocycloalkyloxy (the heterocycloalkyloxy group is
optionally substituted with optionally C_{1-4} alkoxy-substituted C_{1-4} alkyl), C_{1-10} alkyl
optionally substituted with one or more substituents R^{10} , C_{2-10} alkenyl optionally substituted
with one or more substituents R^{15} , C_{2-10} alkynyl optionally substituted with one or more
substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , C_{1-4}
alkylthio optionally substituted with one or more substituents R^{13} , a group $-(O(CH_2)_{q1})_{q2}-$
 $NR^{41}R^{42}$ (wherein $q1$ is an integer selected from 1 to 4, and $q2$ is an integer selected from 2 to
6), a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43}R^{44}$ (wherein $r1$ is an integer selected from 1 to 4, and $r2$
is an integer selected from 1 to 4), a group $-(O(CH_2)_{s1})_{s2}-NR^{45}-C(O)R^{46}$ (wherein $s1$ and $s2$
are each independently an integer selected from 2 to 4), a group $-C(O)NR^{47}R^{48}$, pyridinyl,
pyrrolyl, a group $-NR^{49}R^{50}$, and a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-C(O)NR^{51}R^{52}$ (wherein $y1$ is
an integer selected from 1 to 4, and $y2$ is an integer selected from 1 to 4);

R^{10} , R^{11} , R^{12} , R^{13} , and R^{15} are each independently selected from a halogen atom,

hydroxy, carboxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₄ alkoxy)C₁₋₆ alkoxy, (C₁₋₄ alkoxy)carbonyl, a group -(O(CH₂)_o)_p-OH (wherein o and p are each independently an integer selected from 2 to 4), C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups, 3- to 10-membered heterocycloalkyl, 5- to 10-membered heteroaryl, and a group -NR³⁹R⁴⁰, wherein the 3- to 10-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, C₁₋₄ alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups), (C₁₋₄ alkoxy)C₁₋₄ alkyl (wherein the alkoxy moiety is optionally substituted with one or more hydroxy groups), C₁₋₄ alkoxy, (C₁₋₄ alkoxy)carbonyl, C₁₋₄ alkylthio, morpholino, (C₁₋₃ alkyl)sulfonyl, and -C(O)NR⁵³R⁵⁴;

Ar¹ is C₆₋₁₀ aryl or 5- to 10-membered heteroaryl, wherein the aryl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from optionally C₁₋₄ alkoxy-substituted C₁₋₅ alkoxy, a halogen atom, C₁₋₁₀ alkyl optionally substituted with one or more halogen atoms, a group -SF₅, cyano, hydroxy, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R¹⁴, C₆₋₁₀ aryl optionally substituted with one or more substituents R¹⁴, and 5- to 10-membered heteroaryl optionally substituted with one or more substituents R¹⁴;

each R¹⁴ is independently selected from a halogen atom, oxo, cyano, nitro, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₁₋₄ alkoxy optionally substituted with one or more halogen atoms, (C₁₋₆ alkoxy)carbonyl, a group -NR²⁷R²⁸, a group -SO₂NR³⁵R³⁶, C₁₋₄ alkylthio, and 5- to 10-membered heterocycloalkyl;

R²⁷ and R²⁸ are each independently selected from a hydrogen atom and C₁₋₄ alkyl optionally substituted with (C₁₋₄ alkoxy)carbonyl;

R³⁵ and R³⁶ are each independently selected from a hydrogen atom and C₁₋₄ alkyl;

R³⁹ is a hydrogen atom, or optionally C₁₋₆ alkoxy-substituted C₁₋₆ alkyl;

R⁴⁰ is a hydrogen atom, optionally C₁₋₆ alkoxy-substituted C₁₋₆ alkyl, ((C₁₋₄

alkoxy)carbonyl)C₁₋₆ alkyl, hydroxy-substituted C₁₋₆ alkyl, a group -(CH₂)_u-NR⁵⁵R⁵⁶ (wherein u is an integer selected from 1 to 4), a group -CH((CH₂)_{v1}COOR⁵⁷)-(CH₂)_{v2}-COOR⁵⁷ (wherein v1 is an integer selected from 0 to 2, and v2 is an integer selected from 1 to 3), a group -(CH₂)_w-SO₃H (wherein w is an integer selected from 1 to 4), a group -(CH₂)_{x1}-CH(COOH)-(CH₂)_{x2}-SO₃H (wherein x1 is an integer selected from 0 to 2, and x2 is an integer selected from 1 to 3), 3- to 6-membered oxacycloalkyl, or a group -(CH₂)_{t1}-O-(CH₂)_{t2}-C(O)NR⁵⁸R⁵⁹ (wherein t1 and t2 are each independently an integer selected from 1 to 3);

R⁴¹ is a hydrogen atom or C₁₋₃ alkyl;

R⁴² is C₁₋₈ alkyl substituted with one or more hydroxy groups, or (C₁₋₃ alkoxy)C₁₋₄ alkyl;

R⁴³ is a hydrogen atom or C₁₋₃ alkyl;

R⁴⁴ is C₁₋₈ alkyl substituted with one or more hydroxy groups, or

R⁴³ and R⁴⁴ together with the nitrogen atom to which they are attached may form morpholino;

R⁴⁵ is a hydrogen atom or C₁₋₃ alkyl;

R⁴⁶ is C₁₋₆ alkyl substituted with one or more hydroxy groups;

R⁴⁷ is C₁₋₃ alkyl;

R⁴⁸ is (C₁₋₃ alkoxy)C₁₋₄ alkyl;

R⁴⁹ is a hydrogen atom and C₁₋₄ alkyl;

R⁵⁰ is -(CH₂)_z-NR⁶⁰R⁶¹ (z is an integer selected from 1 to 4, R⁶⁰ is a hydrogen atom or C₁₋₄ alkyl, and R⁶¹ is (C₁₋₃ alkoxy)C₁₋₄ alkyl, or R⁶⁰ and R⁶¹ together with the nitrogen atom to which they are attached may form morpholino);

R⁵¹ is a hydrogen atom or C₁₋₄ alkyl;

R⁵² is C₁₋₈ alkyl substituted with one or more hydroxy groups;

R⁵³ and R⁵⁴ are each independently selected from a hydrogen atom and C₁₋₄ alkyl;

R⁵⁵ is a hydrogen atom or C₁₋₄ alkyl;

R⁵⁶ is (C₁₋₄ alkyl)carbonyl;

R⁵⁷ is a hydrogen atom or C₁₋₄ alkyl;

R⁵⁸ is a hydrogen atom or C₁₋₃ alkyl; and

R⁵⁹ is C₁₋₈ alkyl substituted with one or more hydroxy groups.

2. The compound according to claim 1 or a salt thereof, or a solvate of the compound or the salt, wherein

each R_a is independently selected from a halogen atom, hydroxy, nitro, cyano, (C₁₋₄ alkoxy)carbonyl, 3- to 10-membered heterocycloalkyloxy, C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkenyl optionally substituted with one or more substituents R¹⁵, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², and C₁₋₄ alkylthio optionally substituted with one or more substituents R¹³;

R¹⁰, R¹¹, R¹², R¹³, and R¹⁵ are each independently selected from a halogen atom, hydroxy, carboxy, C₁₋₆ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₄ alkoxy)carbonyl, a group -(O(CH₂)_o)_p-OH (wherein o and p are each independently an integer selected from 2 to 4), C₃₋₆ cycloalkyl optionally substituted with one or more hydroxy groups, 3- to 10-membered heterocycloalkyl, 5- to 10-membered heteroaryl, and a group -NR³⁹R⁴⁰, wherein the 3- to 10-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, and C₁₋₄ alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups); and

R³⁹ and R⁴⁰ are each independently selected from a hydrogen atom and optionally C₁₋₆ alkoxy-substituted C₁₋₆ alkyl.

3. The compound according to claim 1 or 2 or a salt thereof, or a solvate of the compound or the salt, wherein

R_b is optionally C₁₋₄ alkoxy-substituted C₁₋₅ alkoxy or a halogen atom;

R_c is a halogen atom, C₁₋₁₀ alkyl optionally substituted with one or more halogen atoms, or a group -SF₅; and

R_d is cyano, hydroxy, a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, 5- to 10-membered heterocycloalkyl optionally substituted with one or more substituents R¹⁴, C₆₋₁₀ aryl optionally substituted with one or more substituents R¹⁴, or

5- to 10-membered heteroaryl optionally substituted with one or more substituents R^{14} .

4. The compound according to any one of claims 1 to 3 or a salt thereof, or a solvate of the compound or the salt, wherein

each R_a is independently selected from a halogen atom, nitro, cyano, $(C_{1-4}$ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyloxy (the heterocycloalkyloxy group is optionally substituted with optionally C_{1-4} alkoxy-substituted C_{1-4} alkyl), C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} , a group $-(O(CH_2)_{q1})_{q2}-NR^{41}R^{42}$, a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43}R^{44}$, a group $-(O(CH_2)_{s1})_{s2}-NR^{45}-C(O)R^{46}$, a group $-C(O)NR^{47}R^{48}$, pyridinyl, pyrrolyl, a group $-NR^{49}R^{50}$, and a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-NR^{51}R^{52}$;

R^{10} is carboxy, 3- to 6-membered heterocycloalkyl, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, or a group $-(O(CH_2)_o)_p-OH$;

R^{11} is hydroxy, carboxy, 5- or 6-membered heterocycloalkyl optionally substituted with one or more oxo groups, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, a group $-(O(CH_2)_o)_p-OH$, C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups, or a group $-NR^{39}R^{40}$;

R^{12} is a halogen atom, hydroxy, carboxy, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-4}$ alkoxy) C_{1-6} alkoxy, $(C_{1-3}$ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyl, 5- or 6-membered heteroaryl, or a group $-NR^{39}R^{40}$, wherein the 3- to 6-membered heterocycloalkyl group is optionally substituted with one or more substituents selected from oxo, a halogen atom, C_{1-4} alkyl (wherein the alkyl group is optionally substituted with one or more hydroxy groups), $(C_{1-4}$ alkoxy) C_{1-4} alkyl (wherein the alkoxy moiety is optionally substituted with one or more hydroxy groups), C_{1-4} alkoxy, $(C_{1-4}$ alkoxy)carbonyl, C_{1-4} alkylthio, morpholino, $(C_{1-3}$ alkyl)sulfonyl, and $-C(O)NR^{53}R^{54}$;

R^{13} is 5- or 6-membered heterocycloalkyl; and

o and p are each independently an integer selected from 2 to 4.

5. The compound according to any one of claims 1 to 4 or a salt thereof, or a solvate of the compound or the salt, wherein

each R_a is independently selected from a halogen atom, nitro, cyano, $(C_{1-4}$ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyloxy, C_{1-10} alkyl optionally substituted with one or more substituents R^{10} , C_{2-10} alkynyl optionally substituted with one or more substituents R^{11} , C_{1-8} alkoxy optionally substituted with one or more substituents R^{12} , and C_{1-4} alkylthio optionally substituted with one or more substituents R^{13} ;

each R^{10} is independently selected from carboxy, 3- to 6-membered heterocycloalkyl, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, and a group $-(O(CH_2)_o)_p-OH$;

each R^{11} is independently selected from hydroxy, carboxy, 5- or 6-membered heterocycloalkyl optionally substituted with one or more oxo groups, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, a group $-(O(CH_2)_o)_p-OH$, and C_{3-6} cycloalkyl optionally substituted with one or more hydroxy groups;

each R^{12} is independently selected from a halogen atom, hydroxy, C_{1-6} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-3}$ alkoxy)carbonyl, 3- to 6-membered heterocycloalkyl, 5- or 6-membered heteroaryl, and a group $-NR^{39}R^{40}$;

R^{13} is 5- or 6-membered heterocycloalkyl; and

o and p are each independently an integer selected from 2 to 4.

6. The compound according to any one of claims 1 to 4 or a salt thereof, or a solvate of the compound or the salt, wherein

each R_a is selected from a halogen atom, hydroxy, nitro, cyano, C_{1-6} alkyl, (carboxy) C_{1-8} alkyl, (morpholino) C_{1-4} alkyl, $[HO-((CH_2)_oO)]_pC_{1-6}$ alkyl, $(C_{1-6}$ alkoxy) C_{1-8} alkyl optionally substituted with one or more hydroxy groups, C_{1-6} alkoxy optionally substituted with one or more halogen atoms, $[N-((C_{1-3}$ alkoxy) C_{1-4} alkyl)- $N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, C_{1-6} alkoxy substituted with one or more hydroxy groups, $((C_{1-3}$ alkoxy)carbonyl) C_{1-3} alkoxy, $(C_{1-6}$ alkoxy) C_{1-8} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-3}$ alkoxy(C_{1-4} alkoxy)) C_{1-4} alkoxy, (carboxy) C_{1-8} alkoxy, (3- to 6-

membered heterocycloalkyl)C₁₋₆ alkoxy (the heterocycloalkyl moiety is optionally substituted with one or more substituents selected from oxo, a halogen atom, C₁₋₄alkyl (the alkyl is optionally substituted with one or more hydroxy groups), C₁₋₄ alkoxyC₁₋₄ alkyl (the alkoxy moiety is optionally substituted with one or more hydroxyl groups), C₁₋₄ alkoxy, (C₁₋₄ alkoxy)carbonyl, C₁₋₄ alkylthio, morpholino, (C₁₋₃ alkyl)sulfonyl, and (di(C₁₋₃ alkyl)amino)carbonyl), C₁₋₄ alkoxy substituted with a group -NH-CH((CH₂)_{v1}COOR⁵⁷)-(CH₂)_{v2}-COOR⁵⁷, [N-(3- to 6-membered oxacycloalkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di((hydroxy)C₁₋₄ alkyl)-amino]C₁₋₄ alkoxy, [N-((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (pyridinyl)C₁₋₄ alkoxy, (pyrimidinyl)C₁₋₄ alkoxy, (1,2,4-triazolyl)C₁₋₄ alkoxy, [N-(hydroxy)C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di((C₁₋₃ alkoxy)C₁₋₄ alkyl)amino]C₁₋₆ alkoxy, [N,N-di(C₁₋₃ alkyl)amino]C₁₋₆ alkoxy, [N-[N-(C₁₋₄alkyl)carbonyl-N-(C₁₋₃alkyl)amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino]C₁₋₄ alkoxy, [N-[N-(C₁₋₄alkyl)carbonyl-amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl) amino]C₁₋₄ alkoxy, a group -(O(CH₂)_{r1})_{r2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_w-SO₃H, C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_{x1}-CH(COOH)-(CH₂)_{x2}-SO₃H, a group -(O(CH₂)_{s1})_{s2}-NR⁴⁵-C(O)R⁴⁶, a group -C(O)NR⁴⁷R⁴⁸, pyridinyl, a group -NR⁴⁹R⁵⁰, a group -(O(CH₂)_{y1})_{y2}-O-CH₂-C(O)NR⁵¹R⁵², (carboxy)C₂₋₆ alkynyl, (3- to 6-membered heterocycloalkyl)C₂₋₆ alkynyl optionally substituted with one or more oxo groups, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)_oO)_p]C₂₋₈ alkynyl, (C₁₋₆ alkoxy)C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, (C₃₋₆ cycloalkyl)C₂₋₆ alkynyl optionally substituted with one or more hydroxy groups, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₂₋₆ alkynyl, (C₁₋₃ alkoxy)carbonyl, (morpholino)C₁₋₄ alkylthio optionally substituted with one or more oxo groups, 3- to 6-membered oxacycloalkyloxy, and 4- to 6-membered nitrogen containing heterocycloalkyloxy (the nitrogen containing heterocycloalkyl moiety is optionally substituted with one substituent selected from (C₁₋₃ alkoxy)C₁₋₄ alkyl and C₁₋₃ alkyl).

7. The compound according to any one of claims 1 to 6 or a salt thereof, or a solvate of the compound or the salt, wherein

each Ra is independently selected from a halogen atom, hydroxy, cyano, C₁₋₆ alkyl, (carboxy)C₁₋₈ alkyl, (morpholino)C₁₋₄ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl substituted with one or more hydroxy groups, C₁₋₆ alkoxy optionally substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (morpholino)C₁₋₆ alkoxy (the morpholino moiety may be substituted with one or two substituents selected from oxo and C₁₋₃ alkyl), (C₁₋₆ alkoxy)C₁₋₈ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy(C₁₋₄ alkoxy))C₁₋₄ alkoxy, (carboxy)C₁₋₈ alkoxy, (pyrrolidinyl)C₁₋₄ alkoxy (the pyrrolidinyl moiety is optionally substituted with (C₁₋₃ alkyl)C₁₋₄ alkoxy), C₁₋₄ alkoxy substituted with a group -NH-CH((CH₂)_{v1}COOR⁵⁷)-(CH₂)_{v2}-COOR⁵⁷, [N-(oxetanyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di(C₁₋₃ alkyl)amino]C₁₋₆ alkoxy, a group -(O(CH₂)_{t1})_{t2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_w-SO₃H, C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_{x1}-CH(COOH)-(CH₂)_{x2}-SO₃H, (carboxy)C₂₋₆ alkynyl, (morpholino)C₂₋₆ alkynyl, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)_oO)_p]C₂₋₈ alkynyl, (C₁₋₆ alkoxy)C₂₋₈ alkynyl substituted with one or more hydroxy groups, and (C₁₋₃ alkoxy)carbonyl.

8. The compound according to any one of claims 1 to 7 or a salt thereof, or a solvate of the compound or the salt, wherein Ra is independently selected from a halogen atom, hydroxy, cyano, C₁₋₃ alkyl, (carboxy)C₁₋₈ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl substituted with one or more hydroxy groups, C₁₋₃ alkoxy optionally substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (morpholino)C₁₋₆ alkoxy, (C₁₋₆ alkoxy)C₁₋₈ alkoxy substituted with one or more hydroxy groups, (carboxy)C₂₋₆ alkynyl, (morpholino)C₂₋₆ alkynyl, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)_oO)_p]C₂₋₈ alkynyl, (C₁₋₆ alkoxy)C₂₋₈ alkynyl substituted with one or more hydroxy groups, and (C₁₋₃ alkoxy)carbonyl.

9. The compound according to any one of claims 1 to 8 or a salt thereof, or a solvate of the compound or the salt, wherein R³ is methyl substituted with Re.

10. The compound according to any one of claims 1 to 9 or a salt thereof, or a solvate of the compound or the salt, wherein R³ is benzyl optionally substituted with one to three

substituents Ra on the benzene ring.

11. The compound according to any one of claims 1 to 5, 9, and 10 or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri is a halogen atom or C₁₋₃ alkoxy;

Rj is a halogen atom, nitro, or cyano; and

Rk is hydroxy, a halogen atom, (C₁₋₄ alkoxy)carbonyl, 5- to 10-membered heterocycloalkyloxy (the heterocycloalkyloxy group is optionally substituted with optionally C₁₋₄ alkoxy-substituted C₁₋₄ alkyl), C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰, C₂₋₁₀ alkenyl optionally substituted with one or more substituents R¹⁵, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², C₁₋₄ alkylthio optionally substituted with one or more substituents R¹³, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², a group -(O(CH₂)_{r1})_{r2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{s1})_{s2}-NR⁴⁵-C(O)R⁴⁶, a group -C(O)NR⁴⁷R⁴⁸, pyridinyl, pyrrolyl, a group -NR⁴⁹R⁵⁰, or a group -(O(CH₂)_{y1})_{y2}-O-CH₂-C(O)NR⁵¹R⁵².

12. The compound according to any one of claims 1 to 5, and 9 to 11 or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri is a halogen atom or C₁₋₃ alkoxy;

Rj is a halogen atom, nitro, or cyano; and

Rk is hydroxy, a halogen atom, (C₁₋₄ alkoxy)carbonyl, 5- to 10-membered heterocycloalkyloxy, C₁₋₁₀ alkyl optionally substituted with one or more substituents R¹⁰,

C₂₋₁₀ alkenyl optionally substituted with one or more substituents R¹⁵, C₂₋₁₀ alkynyl optionally substituted with one or more substituents R¹¹, C₁₋₈ alkoxy optionally substituted with one or more substituents R¹², or C₁₋₄ alkylthio optionally substituted with one or more substituents R¹³.

13. The compound according to claim 11 or a salt thereof, or a solvate of the compound or the salt, wherein

Rk is a halogen atom, hydroxy, C₁₋₆ alkyl, (carboxy)C₁₋₈ alkyl, (morpholino)C₁₋₄ alkyl, [HO-((CH₂)₆O)_p]C₁₋₆ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl optionally substituted with one or more hydroxy groups, C₁₋₆ alkoxy optionally substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, C₁₋₆ alkoxy substituted with one or more hydroxy groups, ((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkoxy, (C₁₋₆ alkoxy)C₁₋₈ alkoxy optionally substituted with one or more hydroxy groups, (C₁₋₃ alkoxy(C₁₋₄ alkoxy))C₁₋₄ alkoxy, (carboxy)C₁₋₈ alkoxy, (3- to 6-membered heterocycloalkyl)C₁₋₆ alkoxy (the heterocycloalkyl moiety contains one to three heteroatoms selected from O and N and is optionally substituted with one or more substituents selected from oxo, a halogen atom, C₁₋₄ alkyl (the alkyl is optionally substituted with one or more hydroxy groups), (C₁₋₄ alkoxy)C₁₋₄ alkyl (the alkoxy moiety is optionally substituted with one or more hydroxy groups), C₁₋₄ alkoxy, (C₁₋₄ alkoxy)carbonyl, C₁₋₄ alkylthio, morpholino, (C₁₋₃ alkyl)sulfonyl and (di(C₁₋₃ alkyl)amino)carbonyl), C₁₋₄ alkoxy substituted with a group -NH-CH((CH₂)_{v1}COOR⁵⁷)-(CH₂)_{v2}-COOR⁵⁷, [N-(3- to 6-membered oxacycloalkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di((hydroxy)C₁₋₄ alkyl)-amino]C₁₋₄ alkoxy, [N-((C₁₋₃ alkoxy)carbonyl)C₁₋₃ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (pyridinyl)C₁₋₄ alkoxy, (pyrimidinyl)C₁₋₄alkoxy, (1,2,4-triazolyl)C₁₋₄ alkoxy, [N-(hydroxy)C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N,N-di(C₁₋₃ alkoxy(C₁₋₃ alkyl))amino]C₁₋₆ alkoxy, [N,N-di(C₁₋₃ alkyl)amino]C₁₋₆ alkoxy, [N-[N-(C₁₋₄ alkyl)carbonyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, [N-[N-(C₁₋₄ alkyl)carbonyl-amino]C₁₋₄ alkyl-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, a group -(O(CH₂)_{r1})_{r2}-C(O)NR⁴³R⁴⁴, a group -(O(CH₂)_{q1})_{q2}-NR⁴¹R⁴², C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_w-SO₃H, C₁₋₄ alkoxy substituted with a group -NH-(CH₂)_{x1}-CH(COOH)-(CH₂)_{x2}-SO₃H,

a group $-(O(CH_2)_{s1})_{s2}-NR^{45}-C(O)R^{46}$, a group $-C(O)NR^{47}R^{48}$, pyridinyl, a group $-NR^{49}R^{50}$, a group $-(O(CH_2)_{y1})_{y2}-O-CH_2-C(O)NR^{51}R^{52}$, (carboxy) C_{2-8} alkynyl, (3- to 6-membered heterocycloalkyl) C_{2-6} alkynyl optionally substituted with one or more oxo groups, C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, $[HO-((CH_2)_oO)_p]C_{2-8}$ alkynyl, $(C_{1-6}$ alkoxy) C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, $(C_{3-6}$ cycloalkyl) C_{2-6} alkynyl optionally substituted with one or more hydroxy groups, $[N-((C_{1-3}$ alkoxy) C_{1-4} alkyl)-N-(C_{1-3} alkyl)amino] C_{2-6} alkynyl, $(C_{1-3}$ alkoxy)carbonyl, (morpholino) C_{1-4} alkylthio, 3- to 6-membered oxacycloalkyloxy, or 3- to 6-membered nitrogen containing heterocycloalkyloxy (the heterocycloalkyl moiety is optionally substituted with one substituent selected from $(C_{1-3}$ alkoxy) C_{1-4} alkyl and C_{1-3} alkyl).

14. The compound according to claim 11 or 13 or a salt thereof, or a solvate of the compound or the salt, wherein

Rk is hydroxy, a halogen atom, C_{1-6} alkyl, (carboxy) C_{1-8} alkyl, (morpholino) C_{1-4} alkyl, $(C_{1-6}$ alkoxy) C_{1-8} alkyl substituted with one or more hydroxy groups, C_{1-6} alkoxy optionally substituted with one or more halogen atoms, $[N-((C_{1-3}$ alkoxy) C_{1-4} alkyl)-N-(C_{1-3} alkyl)amino] C_{1-4} alkoxy, (morpholino) C_{1-6} alkoxy, $(C_{1-6}$ alkoxy) C_{1-8} alkoxy optionally substituted with one or more hydroxy groups, $(C_{1-3}$ alkoxy(C_{1-4} alkoxy)) C_{1-4} alkoxy, (carboxy) C_{1-8} alkoxy, (pyrrolidinyl) C_{1-4} alkoxy (the pyrrolidinyl moiety is substituted with $(C_{1-3}$ alkyl) C_{1-4} alkoxy), C_{1-4} alkoxy substituted with a group $-NH-CH((CH_2)_{v1}COOR^{57})-(CH_2)_{v2}-COOR^{57}$, $[N-(oxetanyl)-N-(C_{1-3}$ alkyl)amino] C_{1-4} alkoxy, $[N,N-di(C_{1-3}$ alkyl)amino] C_{1-6} alkoxy, a group $-(O(CH_2)_{r1})_{r2}-C(O)NR^{43}R^{44}$, a group $-(O(CH_2)_{q1})_{q2}-NR^{41}R^{42}$, C_{1-4} alkoxy substituted with a group $-NH-(CH_2)_w-SO_3H$, C_{1-4} alkoxy substituted with a group $-NH-(CH_2)_{x1}-CH(COOH)-(CH_2)_{x2}-SO_3H$, (carboxy) C_{2-8} alkynyl, (morpholino) C_{2-6} alkynyl, C_{2-8} alkynyl optionally substituted with one or more hydroxy groups, $[HO-((CH_2)_oO)_p]C_{2-8}$ alkynyl, $(C_{1-6}$ alkoxy) C_{2-8} alkynyl substituted with one or more hydroxy groups, or $(C_{1-3}$ alkoxy)carbonyl.

15. The compound according to any one of claims 11 to 14 or a salt thereof, or a solvate of the compound or the salt, wherein

Rk is hydroxy, a halogen atom, C₁₋₆ alkyl, (carboxy)C₁₋₈ alkyl, (C₁₋₆ alkoxy)C₁₋₈ alkyl substituted with one or more hydroxy groups, C₁₋₃ alkoxy optionally substituted with one or more halogen atoms, [N-((C₁₋₃ alkoxy)C₁₋₄ alkyl)-N-(C₁₋₃ alkyl)amino]C₁₋₄ alkoxy, (morpholino)C₁₋₆ alkoxy, (C₁₋₆ alkoxy)C₁₋₈ alkoxy substituted with one or more hydroxy groups, (carboxy)C₂₋₈ alkynyl, (morpholino)C₂₋₆ alkynyl, C₂₋₈ alkynyl optionally substituted with one or more hydroxy groups, [HO-((CH₂)₆O)_p]C₂₋₈ alkynyl, (C₁₋₆ alkoxy)C₂₋₈ alkynyl substituted with one or more hydroxy groups, or (C₁₋₃ alkoxy)carbonyl.

16. The compound according to any one of claims 1 to 15 or a salt thereof, or a solvate of the compound or the salt, wherein

R¹, R⁴, and R⁵ are as defined in any one of the following (1) to (3):

(1) R¹ is a hydrogen atom or C₁₋₆ alkyl;

R⁴ is C₁₋₄ alkyl optionally substituted with one or more halogen atom(s), or phenyl;

and

R⁵ is a hydrogen atom or C₁₋₄ alkyl;

(2) R¹ and R⁵ together with the carbon atom to which they are attached form a C₃₋₆ saturated carbocyclic ring; and

R⁴ is as defined above; and

(3) R¹ is a hydrogen atom or linear C₁₋₆ alkyl; and

R⁴ and R⁵ together with the carbon atom and the nitrogen atom to which they are attached form a 5- to 8-membered saturated heterocyclic ring.

17. The compound according to any one of claims 1, 2, 4, 6, 7, 9, 10 and 16 or a salt thereof, or a solvate of the compound or the salt, wherein

R¹, R⁴, and R⁵ are as defined in the following (1) or (2):

(1) R¹ is C₁₋₆ alkyl;

R⁴ is optionally halogen atom-substituted C₁₋₄ alkyl or phenyl; and

R⁵ is a hydrogen atom or C₁₋₄ alkyl; or

(2) R¹ and R⁵ together with the carbon atom to which they are attached form a C₃₋₆ saturated carbocyclic ring; and

R⁴ is as defined above;

R³ is C₁₋₄ alkyl substituted with Re;

Re is phenyl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom, optionally R¹¹-substituted C₂₋₆ alkynyl, and optionally R¹²-substituted C₁₋₆ alkoxy;

R¹¹ and R¹² are each independently selected from 5- or 6-membered heterocycloalkyl and -NR³⁹R⁴⁰;

R³⁹ and R⁴⁰ are each independently selected from hydrogen atom and optionally C₁₋₆ alkoxy-substituted C₁₋₆ alkyl;

Ar¹ is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, phenyl optionally substituted with one or more substituents R¹⁴, and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R¹⁴; and

each R¹⁴ is independently selected from a halogen atom, cyano, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, C₁₋₄ alkoxy, a group -SO₂NR³⁵R³⁶ (wherein R³⁵ and R³⁶ are each independently selected from C₁₋₄ alkyl), and C₁₋₄ alkylthio.

18. The compound according to any one of claims 1 to 17 or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is a halogen atom;

Rc is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, phenyl optionally substituted with one or more substituents R¹⁴, or 5- or 6-membered heteroaryl optionally substituted with one or more substituents R¹⁴.

19. The compound according to any one of claims 1 to 4, 9, 10, and 16 to 18 or a salt

thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one or more substituents Ra;

R¹¹ and R¹² are each independently selected from morpholinyl and a group -NR³⁹R⁴⁰;

R⁴ is optionally halogen atom-substituted C₁₋₄ alkyl or phenyl;

Ar¹ is phenyl, pyridinyl, or pyrimidinyl, wherein the phenyl, pyridinyl, and pyrimidinyl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb is a halogen atom;

Rc is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, phenyl, pyridinyl, or pyrimidinyl, wherein the phenyl, pyridinyl, and pyrimidinyl groups are each optionally substituted with one or more substituents R¹⁴.

20. The compound according to any one of claims 1 to 4, 9 to 12 and 16 to 19 or a salt thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri is a halogen atom;

Rj is a halogen atom; and

Rk is hydroxy, C₂₋₆ alkynyl optionally substituted with a substituent R¹¹, or C₁₋₆ alkoxy optionally substituted with a substituent R¹².

21. The compound according to any one of claims 17 to 20, or a salt thereof, or a solvate of the compound or the salt, wherein R¹¹ and R¹² are each independently selected from morpholinyl, and [N-((methoxy)ethyl)-N-(methyl)amino].

22. The compound according to any one of claims 1, 2, 4, 5, 9, 10 and 16 or a salt

thereof, or a solvate of the compound or the salt, wherein

R¹ and R⁵ together with the carbon atom to which they are attached form a C₃₋₆ saturated carbocyclic ring;

R⁴ is C₁₋₄ alkyl;

R³ is C₁₋₄ alkyl substituted with Re;

Re is phenyl optionally substituted with one or more substituents Ra;

each Ra is independently selected from a halogen atom and C₁₋₄ alkoxy optionally substituted with one or more substituents R¹²;

each R¹² is independently selected from 5- or 6-membered heterocycloalkyl and -NR³⁹R⁴⁰;

R³⁹ and R⁴⁰ are each independently selected from a hydrogen atom, and optionally C₁₋₄ alkoxy-substituted C₁₋₄ alkyl;

Ar¹ is phenyl or 5- or 6-membered heteroaryl, wherein the phenyl and heteroaryl groups are each optionally substituted with one to three substituents selected from Rb, Rc, and Rd;

Rb, Rc, and Rd are each independently selected from a halogen atom, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, and 5- or 6-membered heteroaryl optionally substituted with one or more substituents R¹⁴; and

each R¹⁴ is independently selected from a halogen atom, cyano, C₁₋₄ alkyl optionally substituted with one or more halogen atoms, and C₁₋₄ alkylthio.

23. The compound according to any one of claims 1 to 22 or a salt thereof, or a solvate of the compound or the salt, wherein

Rb is a halogen atom;

Rc is a halogen atom or C₁₋₄ alkyl optionally substituted with one or more halogen atoms; and

Rd is a halogen atom or 5- or 6-membered heteroaryl optionally substituted with one or more substituents R¹⁴.

24. The compound according to any one of claims 1 to 3, 9 to 12, 16, 22 and 23 or a salt

thereof, or a solvate of the compound or the salt, wherein

Re is phenyl optionally substituted with one to three substituents Ra;

the one to three substituents Ra are one substituent selected from Ri, Rj, and Rk, two substituents selected from combinations of Ri and Rj, Ri and Rk, and Rj and Rk, or three substituents Ri, Rj, and Rk;

Ri and Rj are each independently a halogen atom; and

Rk is C₁₋₄ alkoxy optionally substituted with one or more substituents R¹².

25. The compound according to any one of claims 1 to 5, 9 to 12, 16 to 18, 20 and 22 to 24 or a salt thereof, or a solvate of the compound or the salt, wherein

Ar¹ is phenyl or pyridinyl;

Rd is pyridinyl or pyrimidinyl;

R¹² is selected from morpholinyl, [N-((methoxy)ethyl)-N-(methyl)amiono], and [N,N-dimethylamino].

26. The compound according to any one of claims 1 to 25 or a salt thereof, or a solvate of the compound or the salt, wherein

Ar¹ is 4-(trifluoromethyl)-2-(6-methylthiopyridin-3-yl)phenyl, 4-(trifluoromethyl)-2-(6-trifluoromethylpyridin-3-yl)phenyl, 4-(trifluoromethyl)-2-(4-trifluoromethylpyrimidin-5-yl)phenyl, 4-(trifluoromethyl)-2-(6-trifluoromethylpyrimidin-4-yl)phenyl, 4-(trifluoromethyl)-2-(6-cyano-5-methylpyrimidin-4-yl)phenyl, 4-(trifluoromethyl)-2-(2-cyanopyridin-4-yl)phenyl, 4-chloro-2-(6-methylthiopyridin-3-yl)phenyl, 4-chloro-2-(6-trifluoromethylpyridin-3-yl)phenyl, 4-chloro-2-(4-trifluoromethylpyrimidin-5-yl)phenyl, 4-chloro-2-(6-cyano-5-methylpyrimidin-4-yl)phenyl, 4-chloro-2-(6-trifluoromethylpyrimidin-4-yl)phenyl, or 4-chloro-2-(2-cyanopyridin-4-yl)phenyl.

27. A compound selected from:

(4aR)-1-[(2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-[2-(6-cyano-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-

difluoro-4-(2-morpholin-4-ylethoxy)phenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-[2-(2-cyanopyridin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

6-[4-[[4-(4aR)-4-hydroxy-4a-methyl-2-oxo-3-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazin-1-yl)methyl]-2,3-difluorophenyl]hex-5-ynoic acid;

(4aR)-1-[[2,3-difluoro-4-(3-morpholin-4-ylprop-1-ynyl)phenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[3-[(2R)-2,3-dihydroxypropoxy]propoxy]-2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[4-[(2R)-2,3-dihydroxypropoxy]butoxy]-2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[4-[6-[(2R)-2,3-dihydroxypropoxy]hexoxy]-2,3-difluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[2-(trifluoromethyl)-4-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-5-yl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-1-[[2,3-difluoro-4-(morpholin-4-ylmethyl)phenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(4aR)-N-(4-bromo-3,5-difluorophenyl)-1-[(3-chloro-2-fluorophenyl)methyl]-4-hydroxy-4a-methyl-2-oxo-6,7-dihydro-5H-pyrrolo[1,2-b]pyridazine-3-carboxamide;

(3S)-3-tert-butyl-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-[6-(trifluoromethyl)pyridin-3-yl]phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[4-chloro-2-(6-methylsulfanylpurin-3-yl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

(3S)-3-tert-butyl-N-[2-(6-cyano-5-methylpyrimidin-4-yl)-4-(trifluoromethyl)phenyl]-1-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-4-hydroxy-2-methyl-6-oxo-3H-pyridazine-5-carboxamide;

6-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-(2-morpholin-4-ylethoxy)phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

6-[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-[2-[2-methoxyethyl(methyl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

4-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[4-(trifluoromethyl)-2-[6-

(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]butanoic acid;

5-[2,3-[difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]pentanoic acid;

6-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]hexanoic acid;

7-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]heptanoic acid;

7-[[2,3-difluoro-4-[2-[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

(2S)-2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]butanedioic acid;

3-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]pentanedioic acid;

6-(2,3-difluoro-4-(2-(methyl(oxetan-3-yl)amino)ethoxy)benzyl)-9-hydroxy-5-methyl-7-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-[2-[methyl(oxetan-3-yl)amino]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

7-(2,3-difluoro-4-(2-(methyl(oxetan-3-yl)amino)ethoxy)benzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-

diazaspiro[4.5]dec-9-ene-9-carboxamide;

7-(4-(3-(dimethylamino)-2,2-dimethylpropoxy)-2,3-difluorobenzyl)-10-hydroxy-6-methyl-8-oxo-N-(4-(trifluoromethyl)-2-(6-(trifluoromethyl)pyrimidin-4-yl)phenyl)-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

6-[[2,3-difluoro-4-[1-(2-methoxyethyl)azetidin-3-yl]oxyphenyl]methyl]-9-hydroxy-5-methyl-7-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-5,6-diazaspiro[3.5]non-8-ene-8-carboxamide;

7-[[2,3-difluoro-4-[4-[methyl-[(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]-4-oxobutoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

7-[[2,3-difluoro-4-[2-[2-[methyl-[(2S,3R,4R,5R)-2,3,4,5,6-pentahydroxyhexyl]amino]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

7-[[2,3-difluoro-4-[2-[2-[2-[2-[2-methoxyethyl(methyl)amino]ethoxy]ethoxy]ethoxy]ethoxy]phenyl]methyl]-10-hydroxy-6-methyl-8-oxo-N-[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]-6,7-diazaspiro[4.5]dec-9-ene-9-carboxamide;

2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]ethanesulfonic acid; and

2-[2-[2,3-difluoro-4-[[10-hydroxy-6-methyl-8-oxo-9-[[4-(trifluoromethyl)-2-[6-(trifluoromethyl)pyrimidin-4-yl]phenyl]carbamoyl]-6,7-diazaspiro[4.5]dec-9-en-7-yl]methyl]phenoxy]ethylamino]ethanesulfonic acid

or a salt thereof, or a solvate of the compound or the salt.

28. The compound according to any one of claims 1 to 27 or the salt thereof, or the solvate of the compound or the salt for use in the prevention and/or treatment of a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure.

29. A NaPi-IIb inhibitor comprising a compound according to any one of claims 1 to 27 or a salt thereof, or a solvate of the compound or the salt.
30. A PiT-1 inhibitor comprising a compound according to any one of claims 1 to 27 or a salt thereof, or a solvate of the compound or the salt.
31. A PiT-2 inhibitor comprising a compound according to any one of claims 1 to 27 or a salt thereof, or a solvate of the compound or the salt.
32. A preventive and/or therapeutic agent for a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, the agent comprising a compound according to any one of claims 1 to 27 or a salt thereof, or a solvate of the compound or the salt as an active ingredient.
33. A method for preventing and/or treating a disease selected from hyperphosphatemia, secondary hyperparathyroidism, and chronic renal failure, comprising administering a therapeutically effective amount of a compound according to any one of claims 1 to 27 or a salt thereof, or a solvate of the compound or the salt to a patient.