


Description

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a novel 4-tert.-butyl imidazole derivative having the formula [I]:



wherein R¹ is a hydrogen atom, a C₁-C₃ alkyl group, a C₁-C₃ alkylthio group or a C₂-C₃ alkoxyalkyl group; R² is a halogen atom, a nitro group or a trifluoromethyl group; R³ is a C₁-C₃ haloalkyl group, a C₁-C₃ haloalkoxy group; X is a nitrogen atom or a group represented by the formula,



in which Y is a halogen atom or a nitro group, a process for producing the same and insecticides containing the same as an active ingredient.


2. Description of the Related Art

It is described in U.S. patent No. 3,868,458, U.S. patent No. 3,940,484 and U.S. patent No. 3,996,366 that a certain imidazole derivative is useful as an active ingredient of insecticide.


As a result of extensive investigations on compounds having an excellent insecticidal effect, the present inventors have found a 4-tert.-butylimidazole derivative having the formula [I] exhibit an extremely high insecticidal effect, and thus have accomplished the present invention.

SUMMARY OF THE INVENTION

According to the present invention, there is provided a 4-tert.-butylimidazole derivative having the formula [I]:



wherein R¹ is a hydrogen atom, a C₁-C₃ alkyl group, a C₁-C₃ alkylthio group or a C₂-C₃ alkoxyalkyl group; R² is a halogen atom, a nitro group or a trifluoromethyl group; R³ is a C₁-C₃ haloalkyl group, a C₁-C₃ haloalkoxy group; X is a nitrogen atom or a group represented by the formula,



in which Y is a halogen atom or a nitro group, a process for producing the same and insecticides containing the same as an active ingredient.

In the formula [I], examples of the halogen atom and the same as the substituent include a fluorine atom, a chlorine atom or a bromine atom.

5

DESCRIPTION OF THE PREFERRED EMBODIMENTS

Hereinafter, the present invention is explained in detail.

Among the 4-tert.-butylimidazole derivative of the present invention, one wherein R¹ is a hydrogen atom, a methyl group, a methylthio group or a methoxymethyl group; R² is a halogen atom which comprises a fluorine atom, a chlorine atom or a bromine atom, a nitro group or a trifluoromethyl group; R³ is a trifluoromethyl group; X is a nitrogen atom or a group represented by the formula,

15



in which Y is a fluorine atom or a chlorine atom, is preferred.

More preferred is one wherein R¹ is a hydrogen atom or a methyl group; R² is a fluorine atom, a chlorine atom or a nitro group; R³ is a trifluoromethyl group; X is a group represented by the formula,

20




25

in which Y is a fluorine atom or a chlorine atom.

The compounds of the present invention can be produced according to the following reaction scheme.

30



35

40

45

50

wherein R¹, R², R³ and X are each as defined above and A is a halogen atom.

The compound of the present invention can be produced by reacting halide compounds having the formula [III] with imidazole derivatives having the formula [II]. The reaction may be carried out at about -5°C to about 150°C for about 1 to 24 hours in a solvent in the presence of a reagent for removing a hydrogen halide.

55

The amounts of the reagents used in the reaction may be 1-2 equivalents of the halide compounds having the formula [III] and 1-4 equivalents of the reagent for removing a hydrogen halide to one equivalent of the imidazole derivatives having the formula [II].

Examples of the solvent which is used for the both reactions described above include aliphatic hydrocarbons such as hexane, heptane, ligroin, petroleum ether, etc.; aromatic hydrocarbons such as benzene, toluene, xylene, etc.; halogenated hydrocarbons such as chloroform, carbon tetrachloride, dichloroethane, chlorobenzene, dichlorobenzene, etc.; ethers such as diethyl ether, diisopropyl ether, dioxan, tetrahydrofuran, ethylene glycol dimethyl ether, etc.; ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone, isophorone, cyclohexanone, etc.; esters such as ethyl acetate, butyl acetate, etc.; nitro compounds such as nitroethane, nitrobenzene, etc.; nitriles such as acetonitrile, isobutyronitrile, etc.; tertiary amines such as pyridine, triethylamine, N,N-diethylaniline, tributylamine, N-methylmorpholine, etc.; acid amides such as formamide, N,N-dimethylformamide, N,N-dimethylacetamide, etc.; sulfur compounds such as dimethylsulfoxide, sulfolane, etc.; or mixtures thereof.

Examples of the reagent of removing hydrogen halide include organic bases such as pyridine, triethylamine, N,N-diethylaniline, etc.; inorganic bases such as sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydrogen carbonate, calcium carbonate, sodium hydride, etc.; alkali metal alkoxides such as sodium methoxide, sodium ethoxide, etc.

After completion of the reaction, post-treatment follows in a conventional manner. If necessary and desired, the product may further be purified by chromatography, distillation, recrystallization, etc.

The imidazole derivatives having the formula [II] and the halide compounds having the formula [III] which are used as raw materials for the compounds of the present invention are prepared by the methods described in U.S. Patent No. 3,868,458, U.S. Patent No. 3,940,484, U.S. Patent No. 3,996,366 and U.S. Patent No. 3,888,932, U.S. Patent No. 3,928,416, European Patent No. 23,100, European Patent No. 34,402, West German Patent No. 2,606,393, West German Patent No. 3,545,570, U.S. Patent No. 4,184,041, J. Org. Chem., 25, 1710 (1960), British Patent No. 2,002,368, British Patent No. 1,121,211, Japan Patent (laid open) No. 84-20,269, respectively, or in a manner similar to the methods.

Examples of the compounds of the present invention are shown in Table 1 below.

TABLE 1

R^1	R^2	R^3	X
H	F	CF ₃	CF
H	F	CF ₃	CCl
H	F	CF ₃	CBr
H	F	CF ₃	CNO ₂
H	F	CF ₃	N
H	F	CF ₂ CF ₃	CF
H	F	CF ₂ CF ₃	CCl
H	F	CF ₂ CF ₃	CBr
H	F	CF ₂ CF ₃	CNO ₂
H	F	CF ₂ CF ₃	CF
H	F	CF ₂ CF ₂ CF ₃	CCl
H	F	CF ₂ CF ₂ CF ₃	CBr
H	F	CF ₂ CF ₂ CF ₃	CNO ₂
H	F	CF ₂ CF ₂ CF ₃	CF
H	F	CF ₂ CF ₂ CF ₃	CCl
H	F	CF ₂ CF ₂ CF ₃	CBr
H	F	CF ₂ CF ₂ CF ₃	CNO ₂
H	F	OCF ₃	CF
H	F	OCF ₃	CCl
H	F	OCF ₃	CBr
H	F	OCF ₃	CNO ₂

R^1	R^2	R^3	X
H	Cl	CF ₃	CF
H	Cl	CF ₃	CCl
H	Cl	CF ₃	CBr
H	Cl	CF ₃	CNO ₂
H	Cl	CF ₃	N
H	Cl	CF ₂ CF ₃	CF
H	Cl	CF ₂ CF ₃	CCl
H	Cl	CF ₂ CF ₃	CBr
H	Cl	CF ₂ CF ₃	CNO ₂
H	Cl	CF ₂ CF ₃	CF
H	Cl	CF ₂ CF ₂ CF ₃	CCl
H	Cl	CF ₂ CF ₂ CF ₃	CBr
H	Cl	CF ₂ CF ₂ CF ₃	CNO ₂
H	Cl	CF ₂ CF ₂ CF ₃	CF
H	Cl	CF ₂ CF ₂ CF ₃	CCl
H	Cl	CF ₂ CF ₂ CF ₃	CBr
H	Cl	CF ₂ CF ₂ CF ₃	CNO ₂
H	Cl	OCF ₃	CF
H	Cl	OCF ₃	CCl
H	Cl	OCF ₃	CBr
H	Cl	OCF ₃	CNO ₂

TABLE 1

R^1	R^2	R^3	X
H	Br	CF ₃	CF
H	Br	CF ₃	CCl
H	Br	CF ₃	CBr
H	Br	CF ₃	CNO ₂
H	Br	CF ₃	N
H	Br	CF ₂ CF ₃	CF
H	Br	CF ₂ CF ₃	CCl
H	Br	CF ₂ CF ₃	CBr
H	Br	CF ₂ CF ₃	CNO ₂
H	Br	CF ₂ CF ₃	CF
H	Br	CF ₂ CF ₂ CF ₃	CCl
H	Br	CF ₂ CF ₂ CF ₃	CBr
H	Br	CF ₂ CF ₂ CF ₃	CNO ₂
H	Br	CF ₂ CF ₂ CF ₃	CF
H	Br	CF ₂ CF ₂ CF ₃	CCl
H	Br	CF ₂ CF ₂ CF ₃	CBr
H	Br	CF ₂ CF ₂ CF ₃	CNO ₂
H	Br	OCF ₃	CF
H	Br	OCF ₃	CCl
H	Br	OCF ₃	CBr
H	Br	OCF ₃	CNO ₂

R^1	R^2	R^3	X
H	NO ₂	CF ₃	CF
H	NO ₂	CF ₃	CCl
H	NO ₂	CF ₃	CBr
H	NO ₂	CF ₃	CNO ₂
H	NO ₂	CF ₃	N
H	NO ₂	CF ₂ CF ₃	CF
H	NO ₂	CF ₂ CF ₃	CCl
H	NO ₂	CF ₂ CF ₃	CBr
H	NO ₂	CF ₂ CF ₃	CNO ₂
H	NO ₂	CF ₂ CF ₃	CF
H	NO ₂	CF ₂ CF ₂ CF ₃	CCl
H	NO ₂	CF ₂ CF ₂ CF ₃	CBr
H	NO ₂	CF ₂ CF ₂ CF ₃	CNO ₂
H	NO ₂	CF ₂ CF ₂ CF ₃	CF
H	NO ₂	OCF ₃	CCl
H	NO ₂	OCF ₃	CBr
H	NO ₂	OCF ₃	CNO ₂

TABLE 1

R^1	R^2	R^3	X
CH_3	F	CF_3	CF
CH_3	F	CF_3	CCl
CH_3	F	CF_3	CBr
CH_3	F	CF_3	CNO_2
CH_3	F	CF_3	N
CH_3	F	CF_2CF_3	CF
CH_3	F	CF_2CF_3	CCl
CH_3	F	CF_2CF_3	CBr
CH_3	F	CF_2CF_3	CNO_2
CH_3	F	CF_2CF_3	CF
CH_3	F	$CF_2CF_2CF_3$	CCl
CH_3	F	$CF_2CF_2CF_3$	CBr
CH_3	F	$CF_2CF_2CF_3$	CNO_2
CH_3	F	$CF_2CF_2CF_3$	CF
CH_3	F	OCF_3	CCl
CH_3	F	OCF_3	CBr
CH_3	F	OCF_3	CNO_2
CH_3	F	OCF_3	CF
CH_3	F	OCF_3	CCl
CH_3	F	OCF_3	CBr
CH_3	F	OCF_3	CNO_2

TABLE 1

R ¹	R ²	R ³	X
CH ₃	Br	CF ₃	CF
CH ₃	Br	CF ₃	CCl
CH ₃	Br	CF ₃	CBr
CH ₃	Br	CF ₃	CNO ₂
CH ₃	Br	CF ₃	N
CH ₃	Br	CF ₂ CF ₃	CF
CH ₃	Br	CF ₂ CF ₃	CCl
CH ₃	Br	CF ₂ CF ₃	CBr
CH ₃	Br	CF ₂ CF ₃	CNO ₂
CH ₃	Br	CF ₂ CF ₂ CF ₃	CF
CH ₃	Br	CF ₂ CF ₂ CF ₃	CCl
CH ₃	Br	CF ₂ CF ₂ CF ₃	CBr
CH ₃	Br	CF ₂ CF ₂ CF ₃	CNO ₂
CH ₃	Br	CF ₂ CF ₂ CF ₃	CF
CH ₃	Br	CF ₂ CF ₂ CF ₃	CCl
CH ₃	Br	CF ₂ CF ₂ CF ₃	CBr
CH ₃	Br	CF ₂ CF ₂ CF ₃	CNO ₂
CH ₃	Br	OCF ₃	CF
CH ₃	Br	OCF ₃	CCl
CH ₃	Br	OCF ₃	CBr
CH ₃	Br	OCF ₃	CNO ₂
CH ₃	Br	OCF ₃	CF
CH ₃	Br	OCF ₃	CCl
CH ₃	Br	OCF ₃	CBr
CH ₃	Br	OCF ₃	CNO ₂

TABLE I

R ¹	R ²	R ³	X
C ₂ H ₅	F	CF ₃	CF
C ₂ H ₅	F	CF ₃	CCl
C ₂ H ₅	F	CF ₃	CBr
C ₂ H ₅	F	CF ₃	CNO ₂
C ₂ H ₅	F	CF ₃	N
C ₂ H ₅	F	CF ₃	CF
C ₂ H ₅	F	CF ₂ CF ₃	CCl
C ₂ H ₅	F	CF ₂ CF ₃	CBr
C ₂ H ₅	F	CF ₂ CF ₃	CNO ₂
C ₂ H ₅	F	CF ₂ CF ₃	CF
C ₂ H ₅	F	CF ₂ CF ₂ CF ₃	CCl
C ₂ H ₅	F	CF ₂ CF ₂ CF ₃	CBr
C ₂ H ₅	F	CF ₂ CF ₂ CF ₃	CNO ₂
C ₂ H ₅	F	CF ₂ CF ₂ CF ₃	CF
C ₂ H ₅	F	CF ₂ CF ₂ CF ₃	CCl
C ₂ H ₅	F	CF ₂ CF ₂ CF ₃	CBr
C ₂ H ₅	F	CF ₂ CF ₂ CF ₃	CNO ₂
C ₂ H ₅	F	OCF ₃	CF
C ₂ H ₅	F	OCF ₃	CCl
C ₂ H ₅	F	OCF ₃	CBr
C ₂ H ₅	F	OCF ₃	CNO ₂
C ₂ H ₅	F	OCF ₃	CF
C ₂ H ₅	F	OCF ₃	CCl
C ₂ H ₅	F	OCF ₃	CBr
C ₂ H ₅	F	OCF ₃	CNO ₂

5
10
15
20
25
30
35
40
45
50
55

5
10
15
20
25
30
35
40
45
50
55

TABLE I

R^1	R^2	R^3	X
C_3H_7-n	F	CF_3	CF
C_3H_7-n	F	CF_3	CCl
C_3H_7-n	F	CF_3	CBr
C_3H_7-n	F	CF_3	CNO_2
C_3H_7-n	F	CF_3	N
C_3H_7-n	F	CF_2CF_3	CF
C_3H_7-n	F	CF_2CF_3	CCl
C_3H_7-n	F	CF_2CF_3	CBr
C_3H_7-n	F	CF_2CF_3	CNO_2
C_3H_7-n	F	CF_2CF_3	CF
C_3H_7-n	F	$CF_2CF_2CF_3$	CCl
C_3H_7-n	F	$CF_2CF_2CF_3$	CBr
C_3H_7-n	F	$CF_2CF_2CF_3$	CNO_2
C_3H_7-n	F	$CF_2CF_2CF_3$	CF
C_3H_7-n	F	$CF_2CF_2CF_3$	CCl
C_3H_7-n	F	$CF_2CF_2CF_3$	CBr
C_3H_7-n	F	$CF_2CF_2CF_3$	CNO_2
C_3H_7-n	F	OCF_3	CF
C_3H_7-n	F	OCF_3	CCl
C_3H_7-n	F	OCF_3	CBr
C_3H_7-n	F	OCF_3	CNO_2

R^1	R^2	R^3	X
C_3H_7-n	Cl	CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_2CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_2CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_2CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_2CF_3	C_3H_7-n
C_3H_7-n	Cl	CF_2CF_3	C_3H_7-n
C_3H_7-n	Cl	$CF_2CF_2CF_3$	C_3H_7-n
C_3H_7-n	Cl	$CF_2CF_2CF_3$	C_3H_7-n
C_3H_7-n	Cl	$CF_2CF_2CF_3$	C_3H_7-n
C_3H_7-n	Cl	$CF_2CF_2CF_3$	C_3H_7-n
C_3H_7-n	Cl	OCF_3	C_3H_7-n
C_3H_7-n	Cl	OCF_3	C_3H_7-n
C_3H_7-n	Cl	OCF_3	C_3H_7-n
C_3H_7-n	Cl	OCF_3	C_3H_7-n

TABLE 1

R^1	R^2	R^3	X
C_3H_7-n	Br	CF_3	CF
C_3H_7-n	Br	CF_3	CCl
C_3H_7-n	Br	CF_3	CBr
C_3H_7-n	Br	CF_3	CNO_2
C_3H_7-n	Br	CF_3	N
C_3H_7-n	Br	CF_2CF_3	CF
C_3H_7-n	Br	CF_2CF_3	CCl
C_3H_7-n	Br	CF_2CF_3	CBr
C_3H_7-n	Br	CF_2CF_3	CNO_2
C_3H_7-n	Br	$CF_2CF_2CF_3$	CF
C_3H_7-n	Br	$CF_2CF_2CF_3$	CCl
C_3H_7-n	Br	$CF_2CF_2CF_3$	CBr
C_3H_7-n	Br	$CF_2CF_2CF_3$	CNO_2
C_3H_7-n	Br	OCF_3	CF
C_3H_7-n	Br	OCF_3	CCl
C_3H_7-n	Br	OCF_3	CBr
C_3H_7-n	Br	OCF_3	CNO_2

R^1	R^2	R^3	X
C_3H_7-n	NO_2	CF_3	CF
C_3H_7-n	NO_2	CF_3	CCl
C_3H_7-n	NO_2	CF_3	CBr
C_3H_7-n	NO_2	CF_3	CNO_2
C_3H_7-n	NO_2	CF_3	N
C_3H_7-n	NO_2	CF_2CF_3	CF
C_3H_7-n	NO_2	CF_2CF_3	CCl
C_3H_7-n	NO_2	CF_2CF_3	CBr
C_3H_7-n	NO_2	CF_2CF_3	CNO_2
C_3H_7-n	NO_2	$CF_2CF_2CF_3$	CF
C_3H_7-n	NO_2	$CF_2CF_2CF_3$	CCl
C_3H_7-n	NO_2	$CF_2CF_2CF_3$	CBr
C_3H_7-n	NO_2	$CF_2CF_2CF_3$	CNO_2
C_3H_7-n	NO_2	OCF_3	CF
C_3H_7-n	NO_2	OCF_3	CCl
C_3H_7-n	NO_2	OCF_3	CBr
C_3H_7-n	NO_2	OCF_3	CNO_2

TABLE 1

R ¹	R ²	R ³	X
C ₃ H ₇ -iso	F	CF ₃	CF
C ₃ H ₇ -iso	F	CF ₃	CCl
C ₃ H ₇ -iso	F	CF ₃	CBr
C ₃ H ₇ -iso	F	CF ₃	CNO ₂
C ₃ H ₇ -iso	F	CF ₃	N
C ₃ H ₇ -iso	F	CF ₂ CF ₃	CF
C ₃ H ₇ -iso	F	CF ₂ CF ₃	CCl
C ₃ H ₇ -iso	F	CF ₂ CF ₃	CBr
C ₃ H ₇ -iso	F	CF ₂ CF ₃	CNO ₂
C ₃ H ₇ -iso	F	CF ₂ CF ₂ CF ₃	CF
C ₃ H ₇ -iso	F	CF ₂ CF ₂ CF ₃	CCl
C ₃ H ₇ -iso	F	CF ₂ CF ₂ CF ₃	CBr
C ₃ H ₇ -iso	F	CF ₂ CF ₂ CF ₃	CNO ₂
C ₃ H ₇ -iso	F	OCF ₃	CF
C ₃ H ₇ -iso	F	OCF ₃	CCl
C ₃ H ₇ -iso	F	OCF ₃	CBr
C ₃ H ₇ -iso	F	OCF ₃	CNO ₂
C ₃ H ₇ -iso	F	OCF ₃	CF
C ₃ H ₇ -iso	F	OCF ₃	CCl
C ₃ H ₇ -iso	F	OCF ₃	CBr
C ₃ H ₇ -iso	F	OCF ₃	CNO ₂

TABLE 1

R ¹	R ²	R ³	X
C ₃ H ₇ -iso	NO ₂	CF ₃	CF
C ₃ H ₇ -iso	NO ₂	CF ₃	CCl
C ₃ H ₇ -iso	NO ₂	CF ₃	CBr
C ₃ H ₇ -iso	NO ₂	CF ₃	CNO ₂
C ₃ H ₇ -iso	NO ₂	CF ₃	N
C ₃ H ₇ -iso	NO ₂	CF ₂ CF ₃	CF
C ₃ H ₇ -iso	NO ₂	CF ₂ CF ₃	CCl
C ₃ H ₇ -iso	NO ₂	CF ₂ CF ₃	CBr
C ₃ H ₇ -iso	NO ₂	CF ₂ CF ₃	CNO ₂
C ₃ H ₇ -iso	NO ₂	CF ₂ CF ₂ CF ₃	CF
C ₃ H ₇ -iso	NO ₂	CF ₂ CF ₂ CF ₃	CCl
C ₃ H ₇ -iso	NO ₂	CF ₂ CF ₂ CF ₃	CBr
C ₃ H ₇ -iso	NO ₂	CF ₂ CF ₂ CF ₃	CNO ₂
C ₃ H ₇ -iso	NO ₂	OCF ₃	CF
C ₃ H ₇ -iso	NO ₂	OCF ₃	CCl
C ₃ H ₇ -iso	NO ₂	OCF ₃	CBr
C ₃ H ₇ -iso	NO ₂	OCF ₃	CNO ₂
C ₃ H ₇ -iso	Br	CF ₃	CF
C ₃ H ₇ -iso	Br	CF ₃	CCl
C ₃ H ₇ -iso	Br	CF ₃	CBr
C ₃ H ₇ -iso	Br	CF ₃	CNO ₂
C ₃ H ₇ -iso	Br	CF ₃	N
C ₃ H ₇ -iso	Br	CF ₂ CF ₃	CF
C ₃ H ₇ -iso	Br	CF ₂ CF ₃	CCl
C ₃ H ₇ -iso	Br	CF ₂ CF ₃	CBr
C ₃ H ₇ -iso	Br	CF ₂ CF ₃	CNO ₂
C ₃ H ₇ -iso	Br	CF ₂ CF ₂ CF ₃	CF
C ₃ H ₇ -iso	Br	CF ₂ CF ₂ CF ₃	CCl
C ₃ H ₇ -iso	Br	CF ₂ CF ₂ CF ₃	CBr
C ₃ H ₇ -iso	Br	CF ₂ CF ₂ CF ₃	CNO ₂
C ₃ H ₇ -iso	Br	OCF ₃	CF
C ₃ H ₇ -iso	Br	OCF ₃	CCl
C ₃ H ₇ -iso	Br	OCF ₃	CBr
C ₃ H ₇ -iso	Br	OCF ₃	CNO ₂

TABLE I

R ¹	R ²	R ³	X
SCH ₃	F	CF ₃	CF
SCH ₃	F	CF ₃	CCl
SCH ₃	F	CF ₃	CBr
SCH ₃	F	CF ₃	CNO ₂
SCH ₃	F	CF ₃	N
SCH ₃	F	CF ₂ CF ₃	CF
SCH ₃	F	CF ₂ CF ₃	CCl
SCH ₃	F	CF ₂ CF ₃	CBr
SCH ₃	F	CF ₂ CF ₃	CNO ₂
SCH ₃	F	CF ₂ CF ₃	CF
SCH ₃	F	CF ₂ CF ₂ CF ₃	CCl
SCH ₃	F	CF ₂ CF ₂ CF ₃	CBr
SCH ₃	F	CF ₂ CF ₂ CF ₃	CNO ₂
SCH ₃	F	CF ₂ CF ₂ CF ₃	CF
SCH ₃	F	CF ₂ CF ₂ CF ₃	CCl
SCH ₃	F	CF ₂ CF ₂ CF ₃	CBr
SCH ₃	F	CF ₂ CF ₂ CF ₃	CNO ₂
SCH ₃	F	OCF ₃	CF
SCH ₃	F	OCF ₃	CCl
SCH ₃	F	OCF ₃	CBr
SCH ₃	F	OCF ₃	CNO ₂

R^1	R^2	R^3	X
SCH_3	Br	CF_3	CF
SCH_3	Br	CF_3	CCl
SCH_3	Br	CF_3	CBr
SCH_3	Br	CF_3	CNO_2
SCH_3	Br	CF_3	N
SCH_3	Br	CF_2CF_3	CF
SCH_3	Br	CF_2CF_3	CCl
SCH_3	Br	CF_2CF_3	CBr
SCH_3	Br	CF_2CF_3	CNO_2
SCH_3	Br	$CF_2CF_2CF_3$	CF
SCH_3	Br	$CF_2CF_2CF_3$	CCl
SCH_3	Br	$CF_2CF_2CF_3$	CBr
SCH_3	Br	$CF_2CF_2CF_3$	CNO_2
SCH_3	Br	OCF_3	CF
SCH_3	Br	OCF_3	CCl
SCH_3	Br	OCF_3	CBr
SCH_3	Br	OCF_3	CNO_2

TABLE 1

R^1	R^2	R^3	X
SC_2H_5	F	CF_3	CF
SC_2H_5	F	CF_3	CCl
SC_2H_5	F	CF_3	CBr
SC_2H_5	F	CF_3	CNO_2
SC_2H_5	F	CF_3	N
SC_2H_5	F	CF_2CF_3	CF
SC_2H_5	F	CF_2CF_3	CCl
SC_2H_5	F	CF_2CF_3	CBr
SC_2H_5	F	CF_2CF_3	CNO_2
SC_2H_5	F	CF_2CF_3	CF
SC_2H_5	F	$CF_2CF_2CF_3$	CCl
SC_2H_5	F	$CF_2CF_2CF_3$	CBr
SC_2H_5	F	$CF_2CF_2CF_3$	CNO_2
SC_2H_5	F	$CF_2CF_2CF_3$	CF
SC_2H_5	F	$CF_2CF_2CF_3$	CCl
SC_2H_5	F	$CF_2CF_2CF_3$	CBr
SC_2H_5	F	$CF_2CF_2CF_3$	CNO_2
SC_2H_5	F	OCF_3	CF
SC_2H_5	F	OCF_3	CCl
SC_2H_5	F	OCF_3	CBr
SC_2H_5	F	OCF_3	CNO_2

TABLE 1

R^1	R^2	R^3	X
SC_2H_5	Br	CF_3	CF
SC_2H_5	Br	CF_3	CCl
SC_2H_5	Br	CF_3	CBr
SC_2H_5	Br	CF_3	CNO_2
SC_2H_5	Br	CF_3	N
SC_2H_5	Br	CF_2CF_3	CF
SC_2H_5	Br	CF_2CF_3	CCl
SC_2H_5	Br	CF_2CF_3	CBr
SC_2H_5	Br	CF_2CF_3	CNO_2
SC_2H_5	Br	CF_2CF_3	CF
SC_2H_5	Br	$CF_2CF_2CF_3$	CCl
SC_2H_5	Br	$CF_2CF_2CF_3$	CBr
SC_2H_5	Br	$CF_2CF_2CF_3$	CNO_2
SC_2H_5	Br	$CF_2CF_2CF_3$	CF
SC_2H_5	Br	OCF_3	CCl
SC_2H_5	Br	OCF_3	CBr
SC_2H_5	Br	OCF_3	CNO_2
SC_2H_5	Br	OCF_3	CF
SC_2H_5	Br	OCF_3	CCl
SC_2H_5	Br	OCF_3	CBr
SC_2H_5	Br	OCF_3	CNO_2

TABLE I

R^1	R^2	R^3	X
SC_3H_7-n	F	CF_3	CF
SC_3H_7-n	F	CF_3	CCl
SC_3H_7-n	F	CF_3	CBr
SC_3H_7-n	F	CF_3	CNO_2
SC_3H_7-n	F	CF_3	N
SC_3H_7-n	F	CF_3	CF
SC_3H_7-n	F	CF_2CF_3	CCl
SC_3H_7-n	F	CF_2CF_3	CBr
SC_3H_7-n	F	CF_2CF_3	CNO_2
SC_3H_7-n	F	CF_2CF_3	CF
SC_3H_7-n	F	$CF_2CF_2CF_3$	CCl
SC_3H_7-n	F	$CF_2CF_2CF_3$	CBr
SC_3H_7-n	F	$CF_2CF_2CF_3$	CNO_2
SC_3H_7-n	F	$CF_2CF_2CF_3$	CF
SC_3H_7-n	F	OCF_3	CCl
SC_3H_7-n	F	OCF_3	CBr
SC_3H_7-n	F	OCF_3	CNO_2
SC_3H_7-n	F	OCF_3	CF
SC_3H_7-n	F	OCF_3	CCl
SC_3H_7-n	F	OCF_3	CBr
SC_3H_7-n	F	OCF_3	CNO_2

TABLE 1

R^1	R^2	R^3	X
SC_3H_7-n	Br	CF_3	CF
SC_3H_7-n	Br	CF_3	CCl
SC_3H_7-n	Br	CF_3	CBr
SC_3H_7-n	Br	CF_3	CNO_2
SC_3H_7-n	Br	CF_3	N
SC_3H_7-n	Br	CF_2CF_3	CF
SC_3H_7-n	Br	CF_2CF_3	CCl
SC_3H_7-n	Br	CF_2CF_3	CBr
SC_3H_7-n	Br	CF_2CF_3	CNO_2
SC_3H_7-n	Br	CF_2CF_3	CF
SC_3H_7-n	Br	$CF_2CF_2CF_3$	CCl
SC_3H_7-n	Br	$CF_2CF_2CF_3$	CBr
SC_3H_7-n	Br	$CF_2CF_2CF_3$	CNO_2
SC_3H_7-n	Br	$CF_2CF_2CF_3$	CF
SC_3H_7-n	Br	OCF_3	CCl
SC_3H_7-n	Br	OCF_3	CBr
SC_3H_7-n	Br	OCF_3	CNO_2
SC_3H_7-n	Br	OCF_3	CF
SC_3H_7-n	Br	OCF_3	CCl
SC_3H_7-n	Br	OCF_3	CBr
SC_3H_7-n	Br	OCF_3	CNO_2

TABLE 1

R ¹	R ²	R ³	X
SC ₃ H ₇ -iso	F	CF ₃	CF
SC ₃ H ₇ -iso	F	CF ₃	CCl
SC ₃ H ₇ -iso	F	CF ₃	CBr
SC ₃ H ₇ -iso	F	CF ₃	CNO ₂
SC ₃ H ₇ -iso	F	CF ₃	N
SC ₃ H ₇ -iso	F	CF ₃	CF
SC ₃ H ₇ -iso	F	CF ₂ CF ₃	CCl
SC ₃ H ₇ -iso	F	CF ₂ CF ₃	CBr
SC ₃ H ₇ -iso	F	CF ₂ CF ₃	CNO ₂
SC ₃ H ₇ -iso	F	CF ₂ CF ₃	CF
SC ₃ H ₇ -iso	F	CF ₂ CF ₂ CF ₃	CCl
SC ₃ H ₇ -iso	F	CF ₂ CF ₂ CF ₃	CBr
SC ₃ H ₇ -iso	F	CF ₂ CF ₂ CF ₃	CNO ₂
SC ₃ H ₇ -iso	F	CF ₂ CF ₂ CF ₃	CF
SC ₃ H ₇ -iso	F	OCF ₃	CCl
SC ₃ H ₇ -iso	F	OCF ₃	CBr
SC ₃ H ₇ -iso	F	OCF ₃	CNO ₂
SC ₃ H ₇ -iso	F	OCF ₃	CF
SC ₃ H ₇ -iso	F	OCF ₃	CCl
SC ₃ H ₇ -iso	F	OCF ₃	CBr
SC ₃ H ₇ -iso	F	OCF ₃	CNO ₂

TABLE 1

R ¹	R ²	R ³	X
SC ₃ H ₇ -iso	Br	CF ₃	CF
SC ₃ H ₇ -iso	Br	CF ₃	CCl
SC ₃ H ₇ -iso	Br	CF ₃	CBr
SC ₃ H ₇ -iso	Br	CF ₃	CNO ₂
SC ₃ H ₇ -iso	Br	CF ₃	N
SC ₃ H ₇ -iso	Br	CF ₂ CF ₃	CF
SC ₃ H ₇ -iso	Br	CF ₂ CF ₃	CCl
SC ₃ H ₇ -iso	Br	CF ₂ CF ₃	CBr
SC ₃ H ₇ -iso	Br	CF ₂ CF ₃	CNO ₂
SC ₃ H ₇ -iso	Br	CF ₂ CF ₂ CF ₃	CF
SC ₃ H ₇ -iso	Br	CF ₂ CF ₂ CF ₃	CCl
SC ₃ H ₇ -iso	Br	CF ₂ CF ₂ CF ₃	CBr
SC ₃ H ₇ -iso	Br	CF ₂ CF ₂ CF ₃	CNO ₂
SC ₃ H ₇ -iso	Br	OCF ₃	CF
SC ₃ H ₇ -iso	Br	OCF ₃	CCl
SC ₃ H ₇ -iso	Br	OCF ₃	CBr
SC ₃ H ₇ -iso	Br	OCF ₃	CNO ₂
SC ₃ H ₇ -iso	Br	OCF ₃	CF
SC ₃ H ₇ -iso	Br	OCF ₃	CCl
SC ₃ H ₇ -iso	Br	OCF ₃	CBr
SC ₃ H ₇ -iso	Br	OCF ₃	CNO ₂

TABLE 1

R^1	R^2	R^3	X
CH_2OCH_3	F	CF_3	CF
CH_2OCH_3	F	CF_3	CCl
CH_2OCH_3	F	CF_3	CBr
CH_2OCH_3	F	CF_3	CNO_2
CH_2OCH_3	F	CF_3	N
CH_2OCH_3	F	CF_2CF_3	CF
CH_2OCH_3	F	CF_2CF_3	CCl
CH_2OCH_3	F	CF_2CF_3	CBr
CH_2OCH_3	F	CF_2CF_3	CNO_2
CH_2OCH_3	F	$CF_2CF_2CF_3$	CF
CH_2OCH_3	F	$CF_2CF_2CF_3$	CCl
CH_2OCH_3	F	$CF_2CF_2CF_3$	CBr
CH_2OCH_3	F	$CF_2CF_2CF_3$	CNO_2
CH_2OCH_3	F	OCF_3	CF
CH_2OCH_3	F	OCF_3	CCl
CH_2OCH_3	F	OCF_3	CBr
CH_2OCH_3	F	OCF_3	CNO_2
CH_2OCH_3	F	OCF_3	CF
CH_2OCH_3	F	OCF_3	CCl
CH_2OCH_3	F	OCF_3	CBr
CH_2OCH_3	F	OCF_3	CNO_2

TABLE 1

R^1	R^2	R^3	X
CH_2OCH_3	Br	CF_3	CF
CH_2OCH_3	Br	CF_3	CCl
CH_2OCH_3	Br	CF_3	CBr
CH_2OCH_3	Br	CF_3	CNO_2
CH_2OCH_3	Br	CF_3	N
CH_2OCH_3	Br	CF_2CF_3	CF
CH_2OCH_3	Br	CF_2CF_3	CCl
CH_2OCH_3	Br	CF_2CF_3	CBr
CH_2OCH_3	Br	CF_2CF_3	CNO_2
CH_2OCH_3	Br	CF_2CF_3	CF
CH_2OCH_3	Br	$CF_2CF_2CF_3$	CCl
CH_2OCH_3	Br	$CF_2CF_2CF_3$	CBr
CH_2OCH_3	Br	$CF_2CF_2CF_3$	CNO_2
CH_2OCH_3	Br	$CF_2CF_2CF_3$	CF
CH_2OCH_3	Br	OCF_3	CCl
CH_2OCH_3	Br	OCF_3	CBr
CH_2OCH_3	Br	OCF_3	CNO_2
CH_2OCH_3	Br	OCF_3	CF
CH_2OCH_3	Br	OCF_3	CCl
CH_2OCH_3	Br	OCF_3	CBr
CH_2OCH_3	Br	OCF_3	CNO_2

TABLE 1

R^1	R^2	R^3	X
$CH_2OC_2H_5$	F	CF_3	CF
$CH_2OC_2H_5$	F	CF_3	CCl
$CH_2OC_2H_5$	F	CF_3	CBr
$CH_2OC_2H_5$	F	CF_3	CNO_2
$CH_2OC_2H_5$	F	CF_3	N
$CH_2OC_2H_5$	F	CF_3	CF
$CH_2OC_2H_5$	F	CF_2CF_3	CCl
$CH_2OC_2H_5$	F	CF_2CF_3	CBr
$CH_2OC_2H_5$	F	CF_2CF_3	CNO_2
$CH_2OC_2H_5$	F	CF_2CF_3	CF
$CH_2OC_2H_5$	F	$CF_2CF_2CF_3$	CCl
$CH_2OC_2H_5$	F	$CF_2CF_2CF_3$	CBr
$CH_2OC_2H_5$	F	$CF_2CF_2CF_3$	CNO_2
$CH_2OC_2H_5$	F	$CF_2CF_2CF_3$	CF
$CH_2OC_2H_5$	F	OCF_3	CCl
$CH_2OC_2H_5$	F	OCF_3	CBr
$CH_2OC_2H_5$	F	OCF_3	CNO_2
$CH_2OC_2H_5$	F	OCF_3	CF
$CH_2OC_2H_5$	F	OCF_3	CCl
$CH_2OC_2H_5$	F	OCF_3	CBr
$CH_2OC_2H_5$	F	OCF_3	CNO_2

R^1	R^2	R^3	X
$CH_2OC_2H_5$	Cl	CF_3	CF
$CH_2OC_2H_5$	Cl	CF_3	CCl
$CH_2OC_2H_5$	Cl	CF_3	CBr
$CH_2OC_2H_5$	Cl	CF_3	CNO_2
$CH_2OC_2H_5$	Cl	CF_3	N
$CH_2OC_2H_5$	Cl	CF_3	CF
$CH_2OC_2H_5$	Cl	CF_2CF_3	CCl
$CH_2OC_2H_5$	Cl	CF_2CF_3	CBr
$CH_2OC_2H_5$	Cl	CF_2CF_3	CNO_2
$CH_2OC_2H_5$	Cl	CF_2CF_3	CF
$CH_2OC_2H_5$	Cl	$CF_2CF_2CF_3$	CCl
$CH_2OC_2H_5$	Cl	$CF_2CF_2CF_3$	CBr
$CH_2OC_2H_5$	Cl	$CF_2CF_2CF_3$	CNO_2
$CH_2OC_2H_5$	Cl	$CF_2CF_2CF_3$	CF
$CH_2OC_2H_5$	Cl	OCF_3	CCl
$CH_2OC_2H_5$	Cl	OCF_3	CBr
$CH_2OC_2H_5$	Cl	OCF_3	CNO_2
$CH_2OC_2H_5$	Cl	OCF_3	CF

TABLE 1

R^1	R^2	R^3	X
$CH_2OC_2H_5$	Br	CF_3	CF
$CH_2OC_2H_5$	Br	CF_3	CCl
$CH_2OC_2H_5$	Br	CF_3	CBr
$CH_2OC_2H_5$	Br	CF_3	CNO_2
$CH_2OC_2H_5$	Br	CF_3	N
$CH_2OC_2H_5$	Br	CF_3	CF
$CH_2OC_2H_5$	Br	CF_2CF_3	CCl
$CH_2OC_2H_5$	Br	CF_2CF_3	CBr
$CH_2OC_2H_5$	Br	CF_2CF_3	CNO_2
$CH_2OC_2H_5$	Br	CF_2CF_3	CF
$CH_2OC_2H_5$	Br	$CF_2CF_2CF_3$	CCl
$CH_2OC_2H_5$	Br	$CF_2CF_2CF_3$	CBr
$CH_2OC_2H_5$	Br	$CF_2CF_2CF_3$	CNO_2
$CH_2OC_2H_5$	Br	$CF_2CF_2CF_3$	CF
$CH_2OC_2H_5$	Br	OCF_3	CCl
$CH_2OC_2H_5$	Br	OCF_3	CBr
$CH_2OC_2H_5$	Br	OCF_3	CNO_2
$CH_2OC_2H_5$	Br	OCF_3	CF
$CH_2OC_2H_5$	Br	OCF_3	CCl
$CH_2OC_2H_5$	Br	OCF_3	CBr
$CH_2OC_2H_5$	Br	OCF_3	CNO_2

R^1	R^2	R^3	X
CH_2OC_2	NO_2	CF_3	CF
CH_2OC_2	NO_2	CF_3	CCl
CH_2OC_2	NO_2	CF_3	CBr
CH_2OC_2	NO_2	CF_3	CNO_2
CH_2OC_2	NO_2	CF_3	N
CH_2OC_2	NO_2	CF_3	CF
$CH_2OC_2H_5$	NO_2	CF_2CF_3	CCl
$CH_2OC_2H_5$	NO_2	CF_2CF_3	CBr
$CH_2OC_2H_5$	NO_2	CF_2CF_3	CNO_2
$CH_2OC_2H_5$	NO_2	CF_2CF_3	CF
$CH_2OC_2H_5$	NO_2	$CF_2CF_2CF_3$	CCl
$CH_2OC_2H_5$	NO_2	$CF_2CF_2CF_3$	CBr
$CH_2OC_2H_5$	NO_2	$CF_2CF_2CF_3$	CNO_2
$CH_2OC_2H_5$	NO_2	$CF_2CF_2CF_3$	CF
$CH_2OC_2H_5$	NO_2	OCF_3	CCl
$CH_2OC_2H_5$	NO_2	OCF_3	CBr
$CH_2OC_2H_5$	NO_2	OCF_3	CNO_2
$CH_2OC_2H_5$	NO_2	OCF_3	CF

TABLE I

R ¹	R ²	R ³	X
H CH ₃ C ₂ H ₅ C ₃ H ₇ -n C ₃ H ₇ -iso SCH ₃ SC ₂ H ₅ SC ₃ H ₇ -n SC ₃ H ₇ -iso CH ₂ OCH ₃ CH ₂ OC ₂ H ₅	CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃	CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃ CF ₃	N N N N N N N N N N N N

Examples of harmful insects against which the compounds of the present invention exhibit remarkable effects include the following:

Harmful insects belonging to Hemiptera:

Planthoppers such as small brown planthopper (Laodelphax striatellus), brown planthopper (Nilaparvata lugens), white-backed rice planthopper (Sogatella furcifera), etc.; leafhoppers such as green rice leafhopper (Nephotettix cincticeps), (Nephotettix virescens), etc.; aphids, bugs, whiteflies, scales, lace bugs, psyllids, etc.

Lepidoptera:

Pyrallid moths such as rice stem borer (Chilo suppressalis), rice leafroller (Chaphalocrocis medinalis), Indian meal moth (Plodia interpunctella), etc.; moths such as tobacco cutworm (Spodoptera litura), rice armyworm (Pseudaletia separata), cabbage armyworm (Mamestra brassicae), etc.; Pieridae such as common cabbageworm (Pieris rapae crucivora), etc.; Tortricidae or tortricid moths such as Adoxophyes spp.; Grapholita spp., etc.; Carposinidae, Lyonetiid moths (Lyonetiidae), tussock moths (Lymantriidae), beet semi-loopers (Autographa nigrisigna); harmful insects belonging to Agrothis spp. such as turnip cutworm (Agrothis segetum), black cutworm (Agrothis ipsilon); harmful insects belonging to Hiliothis spp.; diamondback moth (Plutella xylostella), clothes moths (Tineidae), casemaking clothes moth (Tinea translucens), webbing clothes moth (Tineola bisselliella); etc.

Harmful insects belonging to Diptera:

Mosquitoes such as common mosquito (Culex pipiens pallens), Culex tritaeniorhynchus, etc.; Aedes spp. such as Aedes aegypti, Aedes albopictus, etc.; Anopheles spp. such as Anopheles sinensis, etc.; midges (Chironomidae); Muscidae such as housefly (Musca domestica), false stablefly (Muscina stabulans), etc.; Calliphoridae; Sarcophagidae; lesser housefly (Fannia canicularis); Anthomyiidae or anthomyiid flies such as seedcorn

maggot (Delia platura), onion maggot (Delia antiqua), etc.; fruit flies (Tephritidae); small fruit flies (Drosophilidae); moth flies (Psychodidae); black flies (Simuliidae); Tabanidae; stable flies (Stomoxysidae); etc. Harmful insects belonging to Coleoptera:

5 Corn root worms such as western corn rootworm (Diabrotica virgifera), southern corn root worm (Diabrotica undecimpunctata), etc.; scarabs (Scarabaeidae) such as cupreous chafer (Anomala cuprea), soybean beetle (Anomala rufocuprea), etc.; weevils such as maize weevil (Sitophilus zeamais), rice water weevil (Lissorhoptrus oryzophilus), adzuki bean weevil (Callosobruchys chinensis), etc.; darkling beetles (Tenebrionidae) such as yellow mealworm (Tenebrio molitor), red flour beetle (Tribolium castaneum), etc.; leaf beetles (Chrysomelidae) such as cucurbit leaf beetle (Aulacophora femoralis), striped flea beetles (Phyllotreta striolata), etc.; Anobiidae; Epilachna spp. such as twenty-eight-spotted ladybirds (Epilachna vigintioctopunctata), etc.; powderpost beetles (Lyctidae); false powderpost beetles (Bostrychidae), Cerambycidae; robe beetle (Paderus fuscipes), etc.

Harmful insects belonging to Dictyoptera:

15

German cockroach (Blattella germanica), smokybrown cockroach (Periplaneta fuliginosa), American cockroach (Periplaneta americana), brown cockroach (Periplaneta brunnea), oriental cockroach (Blatta orientalis), etc.

20 Harmful insects belonging to Thysanoptera:

Thrips palmi, flower thrips (Thrips hawaiiensis), etc.

Harmful insects belonging to Hymenoptera:

25

ants (Formicidae); hornets (Vespididae); bethylid wasps (Bethyidae); sawflies (Tenthredinidae) such as cabbage sawfly (Athalia rosae ruficornis), etc.

Harmful insects belonging to Orthoptera:

30

mole crickets (Gryllotalpidae); grasshoppers (Acrididae), etc.;

Harmful insects belonging to Aphaniptera:

35

Purex irritans, etc.

Harmful insects belonging to Anoplura:

40

Pediculus humanus capitis, phthirus pubis, etc.

Harmful insects belonging to Isoptera:

Reticulitermes speratus, Formosan subterranean termite (Coptotermes formosanus), etc.

45 Moreover, the compounds of the present invention are very effective to the insects which develop the resistance against conventional insecticides.

In the case that the compounds of the present invention are used as the active ingredient of insecticidal compositions, the compounds may be used as they are, without adding any other components but in general, the compounds are mixed with a solid carrier, a liquid carrier, a gaseous carrier, a feed, etc. and, if necessary and desired, the mixture is further supplemented with a surfactant and other adjuvants used to prepare insecticidal preparations and prepared into forms such as oil sprays, emulsifiable concentrates, wettable powders, flowable concentrates, granules, dusts, aerosol, fumigants (fogging, etc.), poison bait, etc.

50 These formulations may contain any effective amount of insecticide, generally 0.01 to 95% by weight of the compounds of the present invention as the active ingredient, together with an inert carrier.

55 Examples of the solid carrier used for making formulations include fine powders or granulates, etc. of clays (kaolin clay, diatomaceous earth, synthetic hydrated silicon dioxide, bentonite, Fubasami clay, terra alba, etc.), talc, ceramics, other inorganic minerals (sericite, quartz, sulfur, activated carbon, calcium carbonate, hydrated silica, etc.), chemical fertilizers (ammonium sulfate, ammonium phosphate, ammonium nitrate, urea, ammonium chloride, etc.), etc. Examples of the liquid carrier include water, alcohols (methanol, ethanol, etc.), ketones

(acetone, methyl ethyl ketone, etc.), aromatic hydrocarbons (benzene, toluene, xylene, ethylbenzene, methyl-naphthalene, etc.), aliphatic hydrocarbons (hexane, cyclohexane, kerosene, gas oil, etc.), esters (ethyl acetate, butyl acetate, etc.), nitriles (acetonitrile, isobutyronitrile, etc.), ethers (diisopropyl ether, dioxane, etc.), acid amides (N,N-dimethylformamide, N,N-dimethylacetamide, etc.), halogenated hydrocarbons (dichloromethane, trichloroethane, carbon tetrachloride, etc.), dimethylsulfoxide, vegetable oils such as soybean oil, cotton seed oil, etc. Examples of the gaseous carrier, i.e., propellant, include freon gas, butane gas, LPG (liquefied petroleum gas), dimethyl ether, carbon dioxide, etc.

Examples of the surfactant include alkyl sulfates, alkyl sulfonic acid salts, alkylaryl sulfonic acid salts, alkyl aryl ethers and polyoxyethylene derivatives thereof, polyethylene glycol ether, polyvalent alcohol esters, sugar alcohol derivatives, etc.

Examples of the adjuvants such as binders, dispersing agents, etc. for formulations include casein, gelatin, polysaccharides (starch powders, gum arabic, cellulose derivatives, alginic acid, etc.), lignin derivatives, bentonite, sugars, synthetic water-soluble high molecular substances (polyvinyl alcohol, polyvinyl-pyrrolidone, polyacrylic acid, etc.). Examples of the stabilizer include PAP (acidic isopropyl phosphate), BHT (2,6-di-tert-butyl-4-methylphenol), BHA (mixture of 2-tert-butyl-4-methoxyphenol and 3-tert-butyl-4-methoxyphenol), vegetable oils, mineral oils, surfactants, fatty acids or esters thereof, and the like.

As a base material for the poison baits, there are, for example, feed components such as crop powders, essential vegetable oil, sugars, crystalline cellulose etc.; antioxidants such as dibutylhydroxytoluene, nordihydroguaiaretic acid, etc.; preservatives such as dehydroacetic acid, etc.; feeding error preventing agents such as red pepper powders, etc.; incentive flavor such as cheese flavor, onion flavor, etc.

The thus obtained formulations may be used as they are or after diluting with water, etc. Alternatively, the formulations may be used as admixture with other insecticides, nematocides, acaricides, bactericides, herbicides, plant growth regulators, synergistic agents, fertilizers, soil conditioners, animal feed, etc., or may also be used simultaneously with them, without mixing therewith.

Where the compounds of the present invention are used as insecticides for agricultural use, the dose is generally 0.1 g to 100 g per 10 areas; when emulsifiable concentrates, wettable powders, flowable concentrates, etc. are used after diluting them with water, the concentration is 0.1 ppm to 500 ppm. Granules, dusts, etc. may be used as they are, without diluting them. For purposes of household and public hygiene, emulsifiable concentrates, wettable powders, flowable concentrates, etc. are diluted with water in a concentration of 0.1 ppm to 500 ppm; oils, aerosol, fumigants, poison baits, etc. may be used as they are.

These doses and concentrations may vary depending upon kind of formulations, timing for application, place applied, method for application, kinds of insect, condition of damages, etc. and may be increased or decreased, irrespective of the ranges set forth above.

Hereafter the present invention is described in more detail, by referring to synthesis examples, formulation examples and test examples.

Synthesis Example 1 (Synthesis of Compound No. (5))


To a solution of 0.13 g (1 mmol) of 2-methyl-4(5)-tert-butylimidazole in 5 ml of N,N-dimethylformamide was added 40 mg (1 mmol) of an oily sodium hydride (60%) while cooling with ice, followed by stirring at the same temperature for 15 minutes. After the reaction was completed, to the reaction mixture was added dropwise 0.22 g (1 mmol) of 3-chloro-4,5-difluorobenzotrifluoride, followed by stirring at 80-90°C for 4 hours. After the reaction was completed, the reaction mixture was poured into water, and extracted with ethyl acetate. Further, then the residue was washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The obtained crude product was subjected to silica gel column chromatography to give 0.17 g of 1-(2-chloro-6-fluoro-4-trifluoromethylphenyl)-2-methyl-4-tert.-butylimidazole.
m.p. 60-63°C

Synthesis Example 2 (Synthesis of Compound No. (10))

To a solution of 0.26 g (2 mmol) of 2-methyl-4(5)-tert-butylimidazole in 5 ml of N,N-dimethylformamide was added both of 0.41 g (3 mmol) of anhydrous potassium carbonate, and 0.49 g (2 mmol) of 3-chloro-4-fluoro-5-nitro-benzotrifluoride, followed by stirring at 80-100°C for 6 hours. After cooling, the reaction mixture was poured into water, and extracted with ethyl acetate. Further, then the residue was washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The obtained crude product was subjected to silica gel chromatography to give 0.3 g of 1-(2-chloro-6-nitro-4-trifluoromethylphenyl)-2-methyl-4-tert.-butylimidazole.
m.p. 115.7°C

Examples of the present invention prepared in the same manner as above are shown in Table 2.

Table 2



Compound No.	R ¹	R ²	R ³	X	Physical constant
(1)	H	Cl	CF ₃	CF	$n_D^{24.0}$ 1.4883
(2)	H	Cl	CF ₃	CCl	m.p. 94.6°C
(3)	H	Cl	CF ₃	N	m.p. 79.3°C
(4)	CH ₃	F	CF ₃	CF	$n_D^{24.0}$ 1.4735
(5)	CH ₃	Cl	CF ₃	CF	m.p. 60-63°C
(6)	CH ₃	Cl	CF ₃	CCl	m.p. 60.7°C

- cont'd -

Table 2 (cont'd)

(7)	CH ₃	Cl	CF ₃	N	$n_D^{25.2}$ 1.4999
(8)	CH ₃	Cl	OCF ₃	CCl	m.p. 152.0°C
(9)	CH ₃	Cl	CF ₂ CF ₂ CF ₃	CCl	m.p. 114.4°C
(10)	CH ₃	NO ₂	CF ₃	CCl	m.p. 115.7°C
(11)	CH ₃	NO ₂	CF ₃	CNO ₂	NMR δ (ppm (CDCl ₃ , TMS) 1.25(s, 9H), 2.18(s, 3H), 6.55(s, 1H), 8.46(s, 2H)
(12)	C ₂ H ₅	Cl	CF ₃	CCl	$n_D^{23.8}$ 1.5013
(13)	C ₂ H ₅	NO ₂	CF ₃	CNO ₂	$n_D^{24.9}$ 1.4938
(14)	n-C ₃ H ₇	NO ₂	CF ₃	CNO ₂	$n_D^{25.2}$ 1.4983
(15)	SCH ₃	Cl	CF ₃	CF	$n_D^{25.2}$ 1.5104

- cont'd -

Table 2 (cont'd)

(16)	SCH ₃	Cl	CF ₃	CCl	NMR δ (ppm (CDCl ₃ , TMS) 1.34(s, 9H), 2.51(s, 3H), 6.33(s, 1H), 7.74(s, 2H)
(17)	SCl ₃	Cl	CF ₃	N	^{25.2} _D 1.5281
(18)	SCH ₃	NO ₂	CF ₃	CCl	^{25.3} _D 1.5301
(19)	SCH ₃	NO ₂	CF ₃	CNO ₂	^{25.3} _D 1.5310
(20)	CH ₂ OCH ₃	Cl	CF ₃	CF	^{23.7} _D 1.4840
(21)	CH ₂ OCH ₃	Cl	CF ₃	CCl	m.p. 87.2°C
(22)	CH ₂ OCH ₃	NO ₂	CF ₃	CCl	^{23.7} _D 1.5037

Next, Formulation examples are shown, wherein parts are all by weight and the compounds of the present invention are designated by the compound numbers shown in Table 2.

Formulation Example 1. Emulsifiable concentrate

After 10 parts each of Compounds (1) through (22) of the present invention are dissolved in 35 parts of xylene and 35 parts of dimethylformamide, 14 parts of polyoxyethylene styrylphenyl ether and 6 parts of calcium dodecylbenzenesulfonate are added to the solutions. The resulting mixtures are thoroughly mixed stirred to give 10% emulsifiable concentrate, respectively.

Formulation Example 2. Wettable powder

After 20 parts of Compound (1) through (22) of the present invention are added to a mixture of 4 parts of sodium laurylsulfate, 2 parts of calcium ligninsulfonate, 20 parts of synthetic hydrated silicon dioxide fine powders and 54 parts of diatomaceous earth, the mixture is mixed and stirred with a juice mixer to give 20% wettable powder.

Formulation Example 3. Granule

After 5 parts of synthetic hydrated silicon dioxide fine powders, 5 parts of sodium dodecylbenzenesulfonate, 30 parts of bentonite and 55 parts of clay are added to 5 parts of Compound (1) through (22) of the present invention, the mixture is thoroughly mixed and stirred. A suitable amount of water is further added to the mixture followed by stirring. The mixture is granulated with a granulator and air-dried to give 5% granule.

Formulation Example 4. Dust

After 1 part of Compound (16) of the present invention is dissolved in an appropriate amount of acetone, 5 parts of synthetic hydrated silicon dioxide fine powders, 0.3 part of PAP ant 93.7 parts of clay are added to the solution. The mixture is mixed and stirred with a juice mixer and acetone is evaporated off to give 1% dust.

Formulation Example 5. Flowable concentrate

After 20 parts of Compound (4) of the present invention and 1.5 part of sorbitan trioleate are mixed with 28.5 parts of an aqueous solution containing 2 parts of polyvinyl alcohol, the mixture is finely divided (less than 3 μ in particle diameter) with a sand grinder. Then, 40 parts of aqueous solution containing 0.05 part of xanthane gum and 0.1 part of aluminum magnesium silicate are added to the powders and 10 parts of propylene glycol are further added thereto. The mixture is thoroughly mixed and stirred to give 20% flowable concentrate for aqueous suspension.

Formulation Example 6. Oil spray

After 0.1 part of Compound (1) through (22) of the present invention is dissolved in 5 parts of xylene and 5 parts of trichloroethane, the solution is mixed with 89.9 parts of deodorized kerosene to give 0.1% oil spray.

Formulation Example 7. Oil-based aerosol

After 0.1 part of Compound (1) through (22) of the present invention, 0.2 part of tetramethrin, 0.1 part of d-phenothrin, 10 parts of trichloroethane and 59.6 parts of deodorized kerosene are mixed with each other and dissolved. The solution is filled in an aerosol container. After a valve is mounted to the container, 30 parts of propellant (liquefied petroleum gas) are filled under pressure through the valve to give oil-based aerosol.

Formulation Example 8. Water-based aerosol

After 0.2 part of Compound (5) of the present invention, 0.2 part of d-allethrin, 0.2 part of d-phenothrin, 5 parts of xylene, 3.4 parts of deodorized kerosene and 1 part of emulsifier [ATMOS 300 (registered trademark, Atlas Chemical Co., Ltd.)] are mixed with each other and dissolved. The solution and 50 parts of distilled water are filled in an aerosol container. After a valve is mounted to the container, 40 parts of propellant (liquefied petroleum gas) are filled under pressure through the valve to give water-based aerosol.

Formulation Example 9. Mosquito coil

After 0.3 g of d-allethrin is added to 0.3 g of Compound (5) of the present invention, the mixture is dissolved in 20 ml of acetone. The solution is then uniformly mixed with 99.4 g of carrier for mosquito-coil (tapa powder : sake lees powder : wood powder of 4 : 3 : 3) with stirring and 120 ml of water is then added to the mixture. The mixture is thoroughly kneaded molded and dried to give mosquito-coil.

Formulation Example 10. Electric mosquito mat

Acetone is added to 0.4 g of Compound (5) of the present invention, 0.4 g of d-allethrin and 0.4 g of piperonyl butoxide to dissolve and make the whole volume 10 ml. This solution, 0.5 ml, is uniformly impregnated with a base material for electric mat (a mixture of cotton linter and pulp solidified in a plate-like form) having 2.5 cm x 1.5 cm and a thickness of 0.3 cm to give an electric mosquito mat.

Formulation Example 11. Fumigant

After 100 mg of Compound (16) of the present invention is dissolved in an appropriate amount of acetone, the solution is impregnated with a porous ceramic plate having 4.0 cm x 4.0 cm and a thickness of 1.2 cm to give a fumigant.

Formulation Example 12. Poison bait

After 10 mg of Compound (1) through (22) of the present invention is dissolved in a 0.5 ml acetone, the solution is applied to 5 g of the powder of dry animal food. The powder is dried to give a 0.5% poison bait.

Next, effectiveness of the compounds of the present invention as the active ingredient of insecticidal compositions is described below, with reference to test examples, wherein the compounds of the present invention are designated by the compound numbers shown in Table 2 and compounds used for comparison and control are designated by the compound numbers shown in Table 3.

Table 3

5	Compound symbol	Chemical Structure	Note
10	(A)	<chem>CC1=CN(C(=O)N(C)C)C(C1)C(C)(C)C</chem>	Compound described in U.S. Patent Nos. 3,868,458 and 3,940,484
20	(B)	<chem>CC1=CN(C(=O)N(C)C)C(C1)SC</chem>	Compound described in U.S. Patent No. 3,996,366

Test Example 1 (Insecticidal test on nymphs of brown planthopper)

The emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was diluted with water (corresponding to 500 or 5 ppm) and a rice plant seedling (length of about 12 cm) was immersed in the dilution for a minute. After air-drying, the rice plant seedling was put in a test tube and about 30 nymphs of brown planthopper (*Nilaparvata lugens*) were released. Six days after, the nymphs were observed if they were alive or dead. Criterion for the judgement is as follows.

- a: no insect was alive.
 - b: alive insects were 5 or less.
 - c: alive insects were 6 or more.
- The results are shown in Table 4.

Table 4

5	<u>Test compound</u>	<u>Concentration (ppm)</u>	<u>Efficacy</u>
	(1)	500	a
		5	a
10	(2)	500	a
		5	a
	(3)	500	a
		5	a
	(4)	500	a
		5	a
15	(5)	500	a
		5	a
	(6)	500	a
		5	a
20	(7)	500	a
		5	a
	(8)	500	a
	(9)	500	a
25	(10)	500	a
	(11)	500	a
	(12)	500	a
30	(13)	500	a
	(14)	500	a
35	(15)	500	a
		5	a
	(16)	500	a
		5	a
	(17)	500	a
40	(18)	500	a
	(19)	500	a
45	(20)	500	a
		5	a
	(21)	500	a
		5	a
	(22)	500	a
50	Untreated		c

55

Test Example 2 (Insecticidal test on southern corn rootworm)

On the bottom of a polyethylene cup having a diameter of 5.5 cm, a filter paper which is of the same size was laid down and 1 ml of an aqueous dilution (500 or 50 ppm) of the emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was dropped onto the filter paper and one corn sprout was put as feed. About 30 eggs of southern corn rootworm (*Diabrotica undecimpunctata*) were put in the cup. Eight days after the cup was covered, dead or alive larvae hatched were examined. Criterion for the judgement is as follows.

- a: no insect was alive.
 - b: alive insects were 5 or less.
 - c: alive insects were 6 or more.
- The results are shown in Table 5.

15

20

25

30

35

40

45

50

55

Table 5

	<u>Test compound</u>	<u>Concentration (ppm)</u>	<u>Efficacy</u>
5	(1)	500	a
		50	a
10	(2)	500	a
		50	a
	(3)	500	a
		50	a
	(4)	500	a
		50	a
15	(5)	500	a
		50	a
	(6)	500	a
		50	a
20	(7)	500	a
		50	a
	(8)	500	a
		50	a
	(9)	500	a
		50	a
25	(10)	500	a
		50	a
	(11)	500	a
		50	a
	(12)	500	a
		50	a
30	(13)	500	a
		50	a
	(14)	500	a
		50	a
35	(15)	500	a
		50	a
	(16)	500	a
		50	a
	(17)	500	a
		50	a
40	(18)	500	a
		50	a
	(19)	500	a
		50	a
45	(20)	500	a
		50	a
	(21)	500	a
		50	a
50	(22)	500	a
		50	a
	Untreated		c

55

Test Example 3 (Insecticidal test on common mosquito)

The emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was di-

luted with water and 0.7 ml of the dilution was added to 100 ml of ion exchange water (concentration of the effective ingredient was 3.5 ppm). In the mixture were released 20 last instar larvae of common mosquito (Culex pipiens pallens). One day after the release, mortality was examined.

Criterion for the judgement is as follows.

a: 90% or more

b: not less than 10% but less than 90%

c: less than 10%

The results are shown in Table 6.

Table 6

<u>Test Compound</u>	<u>Efficacy</u>
(1)	a
(2)	a
(3)	a
(4)	a
(5)	a
(6)	a
(7)	a
(8)	a
(9)	a
(10)	a
(11)	a
(12)	a
(13)	a
(14)	a
(15)	a
(16)	a
(17)	a
(18)	a
(19)	a
(20)	a
(21)	a
(22)	a
Untreated	c

Test Example 4 (Insecticidal test on German cockroach)

On the bottom of a polyethylene cup having a diameter of 5.5 cm, a filter paper which is of the same size was laid down and 0.7 ml of an aqueous dilution (500 ppm) of the emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was dropped onto the filter paper. As feed, 30 mg of sucrose

was uniformly spread thereon. In the cup, 10 adult males of German cockroach (*Blattella germanica*) were released. Six days after the cup was covered, dead or alive insects were examined to determine mortality. The results are shown in Table 7.

Table 7

10	<u>Test Compound</u>	<u>Mortality (%)</u> <u>500 ppm</u>
	(1)	100
	(2)	100
15	(3)	100
	(4)	100
	(5)	100
20	(6)	100
	(7)	100
	(8)	100
	(9)	100
25	(10)	100
	(11)	100
	(12)	100
30	(13)	100
	(14)	100
	(15)	100
35	(16)	100
	(17)	100
	(18)	100
	(19)	100
40	(20)	100
	(21)	100
	(22)	100
45	(A)	0
	(B)	0
50	Untreated	0

Test Example 5 (Insecticidal test on housefly)

On the bottom of a polyethylene cup having a diameter of 5.5 cm, a filter paper which is of the same size was laid down and 0.7 ml of an aqueous dilution (500 ppm) of the emulsifiable concentrate of the test compound prepared according to Formulation Example 1 was dropped onto the filter paper. As feed, 30 mg of sucrose was uniformly spread thereon. In the cup, 10 adult females of housefly (*Musca domestica*) were released. Forty eight hours after the cup was covered, dead or alive insects were examined to determine mortality (2 replica-

tions). The results are shown in Table 8.

Table 8

5

	<u>Test Compound</u>	<u>Mortality (%)</u> <u>500 ppm</u>
10	(1)	100
	(2)	100
	(3)	100
15	(4)	100
	(5)	100
	(6)	100
	(7)	100
20	(8)	100
	(9)	100
	(10)	100
25	(11)	100
	(12)	100
	(13)	100
	(14)	100
30	(15)	100
	(16)	100
	(17)	100
35	(18)	100
	(19)	100
	(20)	100
40	(21)	100
	(22)	100
45	Untreated	0

Claims

50

1. A 4-tert.-butylimidazole derivative having the formula:

55

tions). The results are shown in Table 8.

Table 8

5

	<u>Test Compound</u>	<u>Mortality (%)</u> <u>500 ppm</u>
10	(1)	100
	(2)	100
	(3)	100
15	(4)	100
	(5)	100
	(6)	100
	(7)	100
20	(8)	100
	(9)	100
	(10)	100
25	(11)	100
	(12)	100
	(13)	100
	(14)	100
30	(15)	100
	(16)	100
	(17)	100
35	(18)	100
	(19)	100
	(20)	100
40	(21)	100
	(22)	100
45	Untreated	0


Claims

50

1. A 4-tert.-butylimidazole derivative having the formula:

55





wherein R¹ is a hydrogen atom, a C₁-C₃ alkyl group, a C₁-C₃ alkylthio group or a C₂-C₃ alkoxyalkyl group; R² is a halogen atom, a nitro group or a trifluoromethyl group; R³ is a C₁-C₃ haloalkyl group, a C₁-C₃ haloalkoxy group; X is a nitrogen atom or a group represented by the formula,



in which Y is a halogen atom or a nitro group.

2. A 4-tert.-butylimidazole derivative according to claim 1, wherein R¹ is a hydrogen atom, a methyl group, a methylthio group or a methoxymethyl group; R² is a halogen atom which comprises a fluorine atom, a chlorine atom or a bromine atom, a nitro group or a trifluoromethyl group; R₃ is a trifluoromethyl group; X is a nitrogen atom or a group represented by the formula,




in which Y is a fluorine atom or a chlorine atom.

3. A 4-tert.-butylimidazole derivative according to claim 1, wherein R¹ is a hydrogen atom or a methyl group; R² is a fluorine atom, a chlorine atom or a nitro group; R₃ is a trifluoromethyl group; X is a group represented by the formula,



in which Y is a fluorine atom or a chlorine atom.


4. A 4-tert.-butylimidazole derivative according to claim 1, which is 1-(2-chloro-6-fluoro-4-trifluoromethylphenyl)-4-tert.-butylimidazole,
1-(2,6-dichloro-4-trifluoromethylphenyl)-4-tert.-butylimidazole,
1-(2,6-difluoro-4-trifluoromethylphenyl)-2-methyl-4-tert.-butylimidazole,
1-(2-chloro-6-fluoro-4-trifluoromethylphenyl)-2-methyl-4-tert.-butylimidazole, or
1-(2,6-dichloro-4-trifluoromethylphenyl)-2-methyl-4-tert.-butylimidazole, or
1-(2-chloro-6-nitro-4-trifluoromethylphenyl)-2-methyl-4-tert.-butylimidazole.
5. A process for producing a 4-tert.-butylimidazole derivative according to claim 1, which comprises reacting a halide compound having the formula:



wherein R^2 is a halogen atom, a nitro group or a trifluoromethyl group; R^3 is a C_1 - C_3 haloalkyl group, a C_1 - C_3 haloalkoxy group; X is a nitrogen atom or a group represented by the formula,



in which Y is a halogen atom or a nitro group and A is a halogen atom, with an imidazole derivative having the formula:




wherein R^1 is a hydrogen atom, a C_1 - C_3 alkyl group, a C_1 - C_3 alkylthio group or a C_2 - C_3 alkoxyalkyl group.

6. An insecticidal composition which comprises a 4-tert.-butylimidazole derivative according to claim 1 and an inert carrier.
7. A method for controlling insect pests which comprises applying a 4-tert.-butylimidazole derivative according to claim 1 to the insect pests or to a locus where insect pests may propagate.
8. Use of a 4-tert.-butylimidazole derivative according to claim 1 as an insecticide.

Patentansprüche

1. 4-tert.-Butylimidazol-Derivat der Formel:



wobei R^1 ein Wasserstoffatom, einen C_1 - C_3 -Alkylrest, einen C_1 - C_3 -Alkylthioest, oder einen C_2 - C_3 -Alkoxyalkylrest darstellt; R^2 ein Halogenatom, eine Nitrogruppe oder eine Trifluormethylgruppe darstellt; R^3 einen C_1 - C_3 -Halogenalkylrest oder einen C_1 - C_3 -Halogenalkoxyrest darstellt; X ein Stickstoffatom oder einen Rest, der durch die Formel



beschrieben wird, wobei Y ein Halogenatom oder eine Nitrogruppe ist, darstellt.

2. 4-tert.-Butylimidazol-Derivat nach Anspruch 1, wobei R^1 ein Wasserstoffatom, eine Methylgruppe, eine Methylthiogruppe oder eine Methoxymethylgruppe darstellt; R^2 ein Halogenatom, welches ein Fluoratom, ein Chloratom oder ein Bromatom umfaßt, eine Nitrogruppe oder eine Trifluormethylgruppe darstellt; R^3



eine Trifluormethylgruppe darstellt; X ein Stickstoffatom oder einen Rest, der durch die Formel




beschrieben wird, wobei Y ein Fluoratom oder ein Chloratom ist, darstellt.

3. 4-tert.-Butylimidazol-Derivat nach Anspruch 1, wobei R¹ ein Wasserstoffatom oder eine Methylgruppe darstellt; R² ein Fluoratom, ein Chloratom oder eine Nitrogruppe darstellt; R³ eine Trifluormethylgruppe darstellt; X einen Rest, der durch die Formel



beschrieben wird, wobei Y ein Fluoratom oder ein Chloratom ist, darstellt.


4. 4-tert.-Butylimidazol-Derivat nach Anspruch 1, nämlich
 1-(2-Chlor-6-fluor-4-trifluormethylphenyl)-4-tert.-butylimidazol,
 1-(2,6-Dichlor-4-trifluormethylphenyl)-4-tert.-butylimidazol,
 1-(2,6-Difluor-4-trifluormethylphenyl)-2-methyl-4-tert.-butylimidazol,
 1-(2-Chlor-6-fluor-4-trifluormethylphenyl)-2-methyl-4-tert.-butylimidazol,
 1-(2,6-Dichlor-4-trifluormethylphenyl)-2-methyl-4-tert.-butylimidazol, oder
 1-(2-Chlor-6-nitro-4-trifluormethylphenyl)-2-methyl-4-tert.-butylimidazol.
5. Verfahren zur Herstellung eines 4-tert.-Butylimidazol-Derivates nach Anspruch 1, welches die Umsetzung einer Halogenverbindung der Formel:



wobei R² ein Halogenatom, eine Nitrogruppe oder eine Trifluormethylgruppe darstellt; R³ einen C₁-C₃-Halogenalkylrest oder einen C₁-C₃-Halogenalkoxyrest darstellt; X ein Stickstoffatom oder einen Rest, der durch die Formel



beschrieben wird, wobei Y ein Halogenatom oder eine Nitrogruppe ist, darstellt und A ein Halogenatom darstellt, mit einem Imidazol-Derivat der Formel:



wobei R¹ ein Wasserstoffatom, einen C₁-C₃-Alkylrest, einen C₁-C₃-Alkylthioest, oder einen C₂-C₃-Alkoxy-




alkylrest darstellt, umfaßt.

6. Insektizides Mittel, welches ein 4-tert.-Butylimidazol-Derivat nach Anspruch 1 und einen inerten Träger umfaßt.
7. Verfahren zur Bekämpfung von Schadinsekten, welches die Anwendung eines 4-tert.-Butylimidazol-Derivates nach Anspruch 1 auf die Schadinsekten oder einen Ort, wo sich die Schadinsekten fortpflanzen könnten, umfaßt.
8. Verwendung eines 4-tert.-Butylimidazol-Derivates nach Anspruch 1 als Insektizid.

Revendications

1. Dérivé de 4-tert.-butylimidazole ayant la formule:



dans laquelle R¹ est un atome d'hydrogène, un groupe alkyle en C₁-C₃, un groupe alkylthio en C₁-C₃ ou un groupe alcoxyalkyle en C₂-C₃; R² est un atome d'halogène, un groupe nitro ou un groupe trifluorométhyle; R³ est un groupe halogénoalkyle en C₁-C₃, un groupe halogénoalcoxy en C₁-C₃; X est un atome d'azote ou un groupe représenté par la formule



dans laquelle Y est un atome d'halogène ou un groupe nitro.

2. Dérivé de 4-tert.-butylimidazole selon la revendication 1, dans lequel R¹ est un atome d'hydrogène, un groupe méthyle, un groupe méthylthio ou un groupe méthoxyméthyle; R² est un atome d'halogène qui comprend un atome de fluor, un atome de chlore ou un atome de brome, un groupe nitro ou un groupe trifluorométhyle; R³ est un groupe trifluorométhyle; X est un atome d'azote ou un groupe représenté par la formule



dans laquelle Y est un atome de fluor ou un atome de chlore.

3. Dérivé de 4-tert.-butylimidazole selon la revendication 1, dans laquelle R¹ est un atome d'hydrogène ou un groupe méthyle; R² est un atome de fluor, un atome de chlore ou un groupe nitro; R³ est un groupe trifluorométhyle; X est un groupe représenté par la formule




dans laquelle Y est un atome de fluor ou un atome de chlore.

4. Dérivé de 4-tert.-butylimidazole selon la revendication 1, qui est le 1-(2-chloro-6-fluoro-4-trifluorométhylphényl)-4-tert.-butylimidazole, le 1-(2,6-dichloro-4-trifluorométhylphényl)-4-tert.-butylimidazole,



le 1-(2,6-difluoro-4-trifluorométhylphényl)-2-méthyl-4-tert.-butylimidazole,
 le 1-(2-chloro-6-fluoro-4-trifluorométhylphényl)-2-méthyl-4-tert.-butylimidazole,
 le 1-2,6-dichloro-4-trifluorométhylphényl)-2-méthyl-4-tert.-butylimidazole, ou
 le 1-(2-chloro-6-nitro-4-trifluorométhylphényl)-2-méthyl-4-tert.-butylimidazole.


5. Procédé pour préparer un dérivé de 4-tert.-butylimidazole selon la revendication 1, qui comprend la réaction d'un halogénure ayant la formule :



dans laquelle R² est un atome d'halogène, un groupe nitro ou un groupe trifluorométhyle; R³ est un groupe halogénoalkyle en C₁-C₃, un groupe halogénoalcoxy en C₁-C₃; X est un atome d'azote ou un groupe représenté par la formule



dans laquelle Y est un atome d'halogène ou un groupe nitro et A est un atome d'halogène, avec un dérivé d'imidazole ayant la formule:



dans laquelle R¹ est un atome d'hydrogène, un groupe alkyle en C₁-C₃, un groupe alkylthio en C₁-C₃ ou un groupe alcoxyalkyle en C₂-C₃.

6. Composition insecticide qui comprend un dérivé de 4-tert.-butylimidazole selon la revendication 1 et un support inerte.
7. Procédé pour combattre les insectes parasites qui comprend l'application d'un dérivé de 4-tert.-butylimidazole selon la revendication 1 aux insectes parasites ou sur un lieu de propagation des insectes parasites.
8. Utilisation d'un dérivé de 4-tert.-butylimidazole selon la revendication 1 comme insecticide.

