


NEW BICYCLIC COMPOUNDS AS DUAL ATX/CA INHIBITORS

The present invention relates to organic compounds useful for therapy or prophylaxis in a mammal, and in particular to dual autotaxin (ATX) / carbonic anhydrase inhibitors which are inhibitors of lysophosphatidic acid (LPA) production and thus modulators of LPA levels and associated signaling, for the treatment or prophylaxis of inflammatory conditions, conditions of the nervous system, vascular and cardiovascular conditions, cancer, and ocular conditions.

The present invention provides novel compounds of formula (I)



wherein

- 10 R^1 is substituted phenyl, substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted phenyl- C_{2-6} -alkynyl, substituted pyridinyl, substituted pyridinyl- C_{1-6} -alkyl, substituted pyridinyl- C_{2-6} -alkenyl, substituted pyridinyl- C_{2-6} -alkynyl, substituted thiophenyl, substituted thiophenyl- C_{1-6} -alkyl, substituted thiophenyl- C_{2-6} -alkenyl or substituted thiophenyl- C_{2-6} -alkynyl, wherein
- 15 substituted phenyl, substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted phenyl- C_{2-6} -alkynyl, substituted pyridinyl, substituted pyridinyl- C_{1-6} -alkyl, substituted pyridinyl- C_{2-6} -alkenyl, substituted pyridinyl- C_{2-6} -alkynyl, substituted thiophenyl, substituted thiophenyl- C_{1-6} -alkyl, substituted thiophenyl- C_{2-6} -alkenyl and substituted thiophenyl- C_{2-6} -alkynyl are
- 20 substituted by R^3 , R^4 and R^5 ;

Y is -OC(O)- or -C(O)-;

W is -C(O)-, -S(O)₂- or -CR⁶R⁷-;

R^2 is substituted phenyl, substituted pyridinyl or substituted thiophenyl, wherein substituted phenyl, substituted pyridinyl and substituted thiophenyl are substituted by R^6 , R^7 and R^8 ;

R^3 is halogen, hydroxy, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, halo- C_{1-6} -alkoxy, halo- C_{1-6} -alkyl, hydroxy- C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkoxy, C_{3-8} -cycloalkoxy, C_{3-8} -cycloalkoxy- C_{1-6} -alkyl, C_{1-6} -alkylamino, C_{1-6} -alkylcarbonylamino, C_{3-8} -cycloalkylcarbonylamino, C_{1-6} -alkyltetrazolyl, C_{1-6} -alkyltetrazolyl- C_{1-6} -alkyl or heterocycloalkyl- C_{1-6} -alkoxy;

R^4 and R^5 are independently selected from H, halogen, hydroxy, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, halo- C_{1-6} -alkoxy, halo- C_{1-6} -alkyl, hydroxy- C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkoxy, C_{3-8} -cycloalkoxy, C_{3-8} -cycloalkoxy- C_{1-6} -alkyl, C_{1-6} -alkylcarbonylamino, C_{3-8} -cycloalkylcarbonylamino, C_{1-6} -alkyltetrazolyl, C_{1-6} -alkyltetrazolyl- C_{1-6} -alkyl or heterocycloalkyl- C_{1-6} -alkoxy;

R^6 is aminosulfonyl;

R^7 and R^8 are independently selected from H, halogen, hydroxy, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, halo- C_{1-6} -alkoxy, halo- C_{1-6} -alkyl, hydroxy- C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkoxy, C_{3-8} -cycloalkoxy and C_{3-8} -cycloalkoxy- C_{1-6} -alkyl;

R^9 is C_{1-6} -alkyl, C_{1-6} -alkoxy or halogen;

m, n, p and q are independently selected from 1 or 2;

or pharmaceutically acceptable salts.

Autotaxin (ATX) is a secreted enzyme also called ectonucleotide pyrophosphatase / phosphodiesterase 2 or lysophospholipase D that is important for converting lysophosphatidyl choline (LPC) to the bioactive signaling molecule lysophosphatidic acid (LPA). It has been shown that plasma LPA levels are well correlated with ATX activity and hence ATX is believed to be an important source of extracellular LPA. Early experiments with a prototype ATX inhibitor have shown that such a compound is able to inhibit the LPA synthesizing activity in mouse plasma. Work conducted in the 1970s and early 1980s has demonstrated that LPA can

elicit a wide range of cellular responses; including smooth muscle cell contraction, platelet activation, cell proliferation, chemotaxis and others. LPA mediates its effects via signaling to several G protein coupled receptors (GPCRs); the first members were originally denoted Edg (endothelial cell differentiation gene) receptors or ventricular zone gene-1(vzg-1) but are now
5 called LPA receptors. The prototypic group now consists of LPA1/Edg-2/VZG-1, LPA2/Edg-4, and LPA3/Edg-7. Recently, three additional LPA receptors LPA4/p2y9/GPR23, LPA5/GPR92 and LPA6/p2Y5 have been described that are more closely related to nucleotide-selective purinergic receptors than to the prototypic LPA1-3 receptors. The ATX-LPA signaling axis is involved in a large range of physiological and pathophysiological functions, including, for
10 example, nervous system function, vascular development, cardiovascular physiology, reproduction, immune system function, chronic inflammation, tumor metastasis and progression, organ fibrosis as well as obesity and/or other metabolic diseases such as diabetes mellitus. Therefore, increased activity of ATX and/or increased levels of LPA, altered LPA receptor expression and altered responses to LPA may contribute to the initiation, progression and/or
15 outcome of a number of different pathophysiological conditions related to the ATX/LPA axis.

Carbonic anhydrases (CA) are a family of zinc-dependent enzymes, which catalyze the equilibration between carbon dioxide and water and hydrogencarbonate and a proton. The CA reaction is involved in many physiological and pathological processes. Carbonic anhydrase inhibition is useful for the treatment of ocular conditions, conditions of reduced blood flow,
20 cancer, edema and inflammatory conditions including bacterial infections.


Dual acting ATX/CA inhibitors are expected to lower intraocular pressure by facilitating two independent pathways, such as inhibition of aqueous humor (AH) production through CA inhibition at the ciliary body and facilitation of AH outflow by ATX inhibition within the AH drainage system. In conditions of vascular leakage in the eye such as diabetic retinopathy, age
25 related macular disease, or retinal vein occlusion, CA levels have been shown or are expected to increase in the eye and facilitate an increase in pH. This is expected to activate many hydrolytic enzymes that can contribute to disease progression including ATX suggesting additional ATX inhibition by shifting the pH optimum .

In accordance with the invention, the compounds of formula (I) or their pharmaceutically
30 acceptable salts and esters can be used for the treatment or prophylaxis of diseases, disorders or conditions that are associated with the activity of autotaxin and/or the biological activity of lysophosphatidic acid (LPA).

The compounds of formula (I) or their pharmaceutically acceptable salts and esters herein inhibit autotaxin activity and carbonic anhydrase activity therefore inhibit LPA production and modulate LPA levels and associated signaling. Dual ATX/CA-II inhibitors described herein are useful as agents for the treatment or prevention of diseases or conditions in which ATX activity and/or LPA signaling participates, is involved in the etiology or pathology of the disease, or is otherwise associated with at least one symptom of the disease. The ATX-LPA axis has been implicated for example in angiogenesis, chronic inflammation, autoimmune diseases, fibrotic diseases, cancer and tumor metastasis and progression, ocular conditions, metabolic conditions such as obesity and/or diabetes mellitus, conditions such as cholestatic or other forms of chronic pruritus as well as acute and chronic organ transplant rejection.

Objects of the embodiments disclosed herein are the compounds of formula (I) and their aforementioned salts and esters and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments containing the said compounds, their pharmaceutically acceptable salts or esters, the use of the said compounds, salts or esters for the treatment or prophylaxis of disorders or conditions that are associated with the activity of ATX and/or the biological activity of lysophosphatidic acid (LPA), particularly in the treatment or prophylaxis of inflammatory conditions, conditions of the nervous system, conditions of the respiratory system, vascular and cardiovascular conditions, fibrotic diseases, cancer, ocular conditions, metabolic conditions, cholestatic and other forms of chronic pruritus and acute and- chronic organ transplant rejection, and the use of the said compounds, salts or esters for the production of medicaments for the treatment or prophylaxis of inflammatory conditions, conditions of the nervous system, conditions of the respiratory system, vascular and cardiovascular conditions, fibrotic diseases, cancer, ocular conditions, metabolic conditions, cholestatic and other forms of chronic pruritus and acute and chronic organ transplant rejection. More particularly, the compounds of formula (I) and their aforementioned salts and esters and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments containing the said compounds, their pharmaceutically acceptable salts or esters, the use of the said compounds, salts or esters for the treatment or prophylaxis of ocular conditions, furthermore particularly glaucoma.

In one aspect, there is provided compounds of formula (I)



wherein

- R^1 is substituted phenyl, substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted phenyl- C_{2-6} -alkynyl, substituted pyridinyl, substituted pyridinyl- C_{1-6} -alkyl, substituted pyridinyl- C_{2-6} -alkenyl, substituted pyridinyl- C_{2-6} -alkynyl, substituted thiophenyl, substituted thiophenyl- C_{1-6} -alkyl, substituted thiophenyl- C_{2-6} -alkenyl or substituted thiophenyl- C_{2-6} -alkynyl, wherein substituted phenyl, substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted phenyl- C_{2-6} -alkynyl, substituted pyridinyl, substituted pyridinyl- C_{1-6} -alkyl, substituted pyridinyl- C_{2-6} -alkenyl, substituted pyridinyl- C_{2-6} -alkynyl, substituted thiophenyl, substituted thiophenyl- C_{1-6} -alkyl, substituted thiophenyl- C_{2-6} -alkenyl and substituted thiophenyl- C_{2-6} -alkynyl are substituted by R^3 , R^4 and R^5 ;

Y is $-OC(O)-$ or $-C(O)-$;

- W is $-C(O)-$, $-S(O)_2-$ or $-CR^6R^7-$;

R^2 is substituted phenyl, substituted pyridinyl or substituted thiophenyl, wherein substituted phenyl, substituted pyridinyl and substituted thiophenyl are substituted by R^6 , R^7 and R^8 ;

- R^3 is halogen, hydroxy, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, halo- C_{1-6} -alkoxy, halo- C_{1-6} -alkyl, hydroxy- C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkoxy, C_{3-8} -cycloalkoxy, C_{3-8} -cycloalkoxy- C_{1-6} -alkyl, C_{1-6} -alkylamino, C_{1-6} -alkylcarbonylamino, C_{3-8} -cycloalkylcarbonylamino, C_{1-6} -alkyltetrazolyl, C_{1-6} -alkyltetrazolyl- C_{1-6} -alkyl or heterocycloalkyl- C_{1-6} -alkoxy;

- R^4 and R^5 are independently selected from H, halogen, hydroxy, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, halo- C_{1-6} -alkoxy, halo- C_{1-6} -alkyl, hydroxy- C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkoxy, C_{3-8} -cycloalkoxy, C_{3-8} -cycloalkoxy- C_{1-6} -alkyl, C_{1-6} -alkylcarbonylamino, C_{3-8} -cycloalkylcarbonylamino, C_{1-6} -alkyltetrazolyl, C_{1-6} -alkyltetrazolyl- C_{1-6} -alkyl or heterocycloalkyl- C_{1-6} -alkoxy;

R^6 is aminosulfonyl;

R^7 and R^8 are independently selected from H, halogen, hydroxy, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, halo- C_{1-6} -alkoxy, halo- C_{1-6} -alkyl, hydroxy- C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkoxy, C_{3-8} -cycloalkoxy and C_{3-8} -cycloalkoxy- C_{1-6} -alkyl;


R^9 is C_{1-6} -alkyl, C_{1-6} -alkoxy or halogen;

m, n, p and q are independently selected from 1 or 2;

or pharmaceutically acceptable salts.

In another aspect, there is provided a process to prepare a compound as defined herein

comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III), wherein R^1 , R^2 , R^9 , m, n, p and q are as defined herein and W is -C(O)-



In another aspect, there is provided a compound as defined herein for use as therapeutically active substance.

In another aspect, there is provided a pharmaceutical composition comprising a compound as defined herein and a therapeutically inert carrier.

In another aspect, there is provided a compound as defined herein for the treatment or prophylaxis of ocular conditions.

In another aspect, there is provided the use of a compound as defined herein for the preparation of a medicament for the treatment or prophylaxis of ocular conditions.

In another aspect, there is provided a compound as defined herein, when manufactured according to a process as defined herein.

The term “C₁₋₆-alkoxy” denotes a group of the formula -O-R', wherein R' is an C₁₋₆-alkyl group. Examples of C₁₋₆-alkoxy group include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy and tert-butoxy. Particular example is methoxy.

The term “C₂₋₆-alkenyl” denotes a monovalent linear or branched hydrocarbon group of 2 to 6 carbon atoms with at least one double bond. Particular example is ethylenyl.

The term “C₁₋₆-alkoxy-C₁₋₆-alkyl” denotes a C₁₋₆-alkyl group wherein at least one of the hydrogen atoms of the C₁₋₆-alkyl group is replaced by a C₁₋₆-alkoxy group. Particular examples
5 are methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, iso-propoxymethyl and iso-propoxyethyl.

The term “C₁₋₆-alkyl” denotes a monovalent linear or branched saturated hydrocarbon group of 1 to 6 carbon atoms. Examples of C₁₋₆-alkyl include methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, tert-butyl and pentyl. Particular alkyl groups include methyl, ethyl,
10 isopropyl, n-butyl and sec-butyl.

The term “C₁₋₆-alkylamino” a group of the formula -NH- R', wherein R' is an C₁₋₆-alkyl group. Particular C₁₋₆-alkylamino is a group of the formula -NH- R', wherein R' is ter-butyl.

The term “C₁₋₆-alkylcarbonylamino” denotes a group of the formula -NH-C(O)-R', wherein R' is an C₁₋₆-alkyl group. Particular C₁₋₆-alkylcarbonylamino is a group of the formula
15 -NH-C(O)-R', wherein R' is ter-butyl.

The term “C₁₋₆-alkyltetrazolyl” denotes tetrazolyl group substituted with one C₁₋₆-alkyl group. Particular C₁₋₆-alkyltetrazolyl is methyltetrazolyl.

The term “C₁₋₆-alkyltetrazolyl-C₁₋₆-alkyl” denotes C₁₋₆-alkyl group wherein one of the hydrogen atoms of the C₁₋₆-alkyl group is replaced by a C₁₋₆-alkyltetrazolyl group. Particular
20 example is methyltetrazolylmethyl.

The term “C₂₋₆-alkynyl” denotes a monovalent linear or branched hydrocarbon group of 2 to 6 carbon atoms with at least one triple bond.

The term “amino” denotes the -NH₂ group.

The term “aminosulfonyl” denotes -S(O)₂-NH₂ group.

25 The term “cyano” denotes a -C≡N group.

The term “C₃₋₈-cycloalkoxy” denotes a group of the formula -O-R', wherein R' is a C₃₋₈-cycloalkyl.

The term “C₃₋₈-cycloalkoxy-C₁₋₆-alkyl” denotes a C₁₋₆-alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a C₃₋₈-cycloalkoxy group.

The term “C₃₋₈-cycloalkyl” denotes a monovalent saturated monocyclic or bicyclic hydrocarbon group of 3 to 8 ring carbon atoms. Bicyclic means a ring system consisting of two saturated carbocycles having two carbon atoms in common. Examples for monocyclic cycloalkyl are cyclopropyl, cyclobutanyl, cyclopentyl, cyclohexyl or cycloheptyl. Examples for bicyclic C₃₋₈-cycloalkyl are bicyclo[2.2.1]heptanyl or bicyclo[2.2.2]octanyl. Particular C₃₋₈-cycloalkyl group is cyclopropyl.

The term “C₃₋₈-cycloalkyl-C₁₋₆-alkoxy” denotes a C₁₋₆-alkoxy group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a C₃₋₈-cycloalkyl group.

The term “C₃₋₈-cycloalkyl-C₁₋₆-alkyl” denotes a C₁₋₆-alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a C₃₋₈-cycloalkyl group.

The term “C₃₋₈-cycloalkylcarbonylamino” denotes a group of the formula -NH-C(O)-R', wherein R' is a C₃₋₈-cycloalkyl group.

The term “halo-C₁₋₆-alkoxy” denotes a C₁₋₆-alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group has been replaced by the same or different halogen atoms. Particular examples are trifluoromethoxy.

The term “halogen” and “halo” are used interchangeably herein and denote fluoro, chloro, bromo or iodo. Particular halogens are chloro and fluoro.

The term “halo-C₁₋₆-alkyl” denotes a C₁₋₆-alkyl group wherein at least one of the hydrogen atoms of the C₁₋₆-alkyl group has been replaced by the same or different halogen atoms. Particular examples are trifluoromethyl.

The term “heterocycloalkyl” denotes a monovalent saturated or partly unsaturated mono- or bicyclic ring system of 4 to 9 ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. Bicyclic means consisting of two cycles having two ring atoms in common, i.e. the bridge separating the two rings is either a single bond or a chain of one or two ring atoms. Examples for monocyclic saturated heterocycloalkyl are 4,5-dihydro-oxazolyl, oxetanyl, azetidiny, pyrrolidiny, 2-oxo-pyrrolidin-3-yl, tetrahydrofuranyl, tetrahydro-thienyl, pyrazolidiny, imidazolidiny, oxazolidiny, isoxazolidiny, thiazolidiny,

piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, thiomorpholinyl, 1,1-dioxo-thiomorpholin-4-yl, azepanyl, diazepanyl, homopiperazinyl, or oxazepanyl. Examples for bicyclic saturated heterocycloalkyl are 8-aza-bicyclo[3.2.1]octyl, quinuclidinyl, 8-oxa-3-aza-bicyclo[3.2.1]octyl, 9-aza-bicyclo[3.3.1]nonyl, 3-oxa-9-aza-bicyclo[3.3.1]nonyl, or 3-thia-9-aza-bicyclo[3.3.1]nonyl. Examples for partly unsaturated heterocycloalkyl are dihydrofuryl, imidazoliny, dihydro-oxazolyl, tetrahydro-pyridinyl, or dihydropyranyl. Particular example of heterocycloalkyl group is tetrahydropyranyl.

The term “heterocycloalkyl-C₁₋₆-alkoxy” denotes a C₁₋₆-alkoxy group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a heterocycloalkyl group. Particular example of heterocycloalkyl-C₁₋₆-alkoxy is tetrahydropyranyl-C₁₋₆-alkoxy, more particularly tetrahydropyranylmethoxy.

The term “hydroxy” denotes a -OH group.

The term “hydroxy-C₁₋₆-alkyl” denotes a C₁₋₆-alkyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a hydroxy group. Particular examples are hydroxymethyl and hydroxyethyl.

The term “phenoxy” denotes a group of the formula -O-R', wherein R' is a phenyl group.

The term “phenoxy-C₁₋₆-alkyl” denotes a C₁₋₆-alkyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a phenoxy group.

The term “phenyl-C₂₋₆-alkenyl” denotes a C₂₋₆-alkenyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a phenyl group. Particular example of phenyl-C₂₋₆-alkenyl is phenylethenyl.

The term “phenyl-C₁₋₆-alkyl” denotes a C₁₋₆-alkyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a phenyl group. Particular examples of phenyl-C₁₋₆-alkyl are phenylmethyl and phenylethyl.

The term “phenyl-C₂₋₆-alkynyl” denotes a C₂₋₆-alkynyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a phenyl group.

The term “pyridinyl-C₂₋₆-alkenyl” denotes a C₂₋₆-alkenyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a pyridinyl group.

The term "pyridinyl-C₁₋₆-alkyl" denotes a C₁₋₆-alkyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a pyridinyl group. Particular example of pyridinyl-C₁₋₆-alkyl is pyridinylmethyl, more particularly 2-pyridinylmethyl.

5 The term "pyridinyl-C₂₋₆-alkynyl" denotes a C₂₋₆-alkynyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a pyridinyl group.

The term "thiophenyl-C₂₋₆-alkenyl" denotes a C₂₋₆-alkenyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a thiophenyl group.

The term "thiophenyl-C₁₋₆-alkyl" denotes a C₁₋₆-alkyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a thiophenyl group.

10 The term "thiophenyl-C₂₋₆-alkynyl" denotes a C₂₋₆-alkynyl group wherein one of the hydrogen atoms of the alkyl group is replaced by a thiophenyl group.

The term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. The salts are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, in particular
15 hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, N-acetylcystein and the like. In addition, these salts may be
20 prepared by addition of an inorganic base or an organic base to the free acid. Salts derived from an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium salts and the like. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as
25 isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, lysine, arginine, N-ethylpiperidine, piperidine, polyimine resins and the like. Particular pharmaceutically acceptable salts of compounds of formula (I) are the hydrochloride salts, methanesulfonic acid salts and citric acid salts.

"Pharmaceutically acceptable esters" means that compounds of general formula (I) may be
30 derivatised at functional groups to provide derivatives which are capable of conversion back to the parent compounds in vivo. Examples of such compounds include physiologically acceptable

and metabolically labile ester derivatives, such as methoxymethyl esters, methylthiomethyl esters and pivaloyloxymethyl esters. Additionally, any physiologically acceptable equivalents of the compounds of general formula (I), similar to the metabolically labile esters, which are capable of producing the parent compounds of general formula (I) in vivo, are within the scope of this invention.

The term "protecting group" (PG) denotes a group which selectively blocks a reactive site in a multifunctional compound such that a chemical reaction can be carried out selectively at another unprotected reactive site in the meaning conventionally associated with it in synthetic chemistry. Protecting groups can be removed at the appropriate point. Exemplary protecting groups are amino-protecting groups, carboxy-protecting groups or hydroxy-protecting groups. Particular protecting groups are the tert-butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), fluorenylmethoxycarbonyl (Fmoc) and benzyl (Bn) groups. Further particular protecting groups are the tert-butoxycarbonyl (Boc) and the fluorenylmethoxycarbonyl (Fmoc) groups. More particular protecting group is the tert-butoxycarbonyl (Boc) group.

The abbreviation uM means microMolar and is equivalent to the symbol μM .

The abbreviation uL means microliter and is equivalent to the symbol μL .

The abbreviation ug means microgram and is equivalent to the symbol μg .

The compounds of formula (I) can contain several asymmetric centers and can be present in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates.

According to the Cahn-Ingold-Prelog Convention the asymmetric carbon atom can be of the "R" or "S" configuration.

Also an embodiment of the present invention are compounds according to formula (I) as described herein and pharmaceutically acceptable salts or esters thereof, in particular compounds according to formula (I) as described herein and pharmaceutically acceptable salts thereof, more particularly compounds according to formula (I) as described herein.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein

- 5 R^1 is substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted pyridinyl or substituted pyridinyl- C_{1-6} -alkyl, wherein substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted pyridinyl and substituted pyridinyl- C_{1-6} -alkyl are substituted by R^3 , R^4 and R^5 ;
- Y is a -OC(O)- or -C(O)-;
- W is -C(O)-;
- 10 R^2 is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R^6 , R^7 and R^8 ;
- R^3 is halo- C_{1-6} -alkoxy, C_{1-6} -alkylcarbonylamino or tetrahydropyranyl- C_{1-6} -alkoxy;
- R^4 is H, C_{1-6} -alkyl or C_{3-8} -cycloalkyl;
- R^5 is H;
- R^6 is aminosulfonyl;
- R^7 and R^8 are independently selected from H and halogen;
- 15 R^9 is C_{1-6} -alkyl, C_{1-6} -alkoxy or halogen;
- m, n, p and q are 1;
- or pharmaceutically acceptable salts.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, 20 substituted phenyl- C_{2-6} -alkenyl, substituted pyridinyl or substituted pyridinyl- C_{1-6} -alkyl, wherein substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted pyridinyl and substituted pyridinyl- C_{1-6} -alkyl are substituted by R^3 , R^4 and R^5 .

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is substituted pyridinyl or substituted pyridinyl- C_{1-6} - 25 alkyl, wherein substituted pyridinyl and substituted pyridinyl- C_{1-6} -alkyl are substituted by R^3 , R^4 and R^5 .

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein Y is -C(O)-.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R² is substituted phenyl or substituted pyridinyl, wherein
5 substituted phenyl and substituted pyridinyl are substituted by R⁶, R⁷ and R⁸.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R³ is halo-C₁₋₆-alkoxy, C₁₋₆-alkylcarbonylamino or tetrahydropyranyl-C₁₋₆-alkoxy.

A further particular embodiment of the present invention are compounds according to
10 formula (I) as described herein, wherein R³ is tetrahydropyranyl-C₁₋₆-alkoxy.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁴ is H, C₁₋₆-alkyl or C₃₋₈-cycloalkyl.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁴ is C₃₋₈-cycloalkyl.

15 A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁵ is H.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁷ and R⁸ are independently selected from H and halogen.

A further particular embodiment of the present invention are compounds according to
20 formula (I) as described herein, wherein R⁷ is H or halogen.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁸ is H.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁹ is halogen or C₁₋₆-alkoxy.

25 A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein m, n, p and q are 1.

A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein

R^1 is substituted pyridinyl or substituted pyridinyl- C_{1-6} -alkyl, wherein substituted pyridinyl and substituted pyridinyl- C_{1-6} -alkyl are substituted by R^3 , R^4 and R^5 ;

5 Y is $-C(O)-$;

W is $-C(O)-$;

R^2 is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R^6 , R^7 and R^8 ;

R^3 is tetrahydropyranyl- C_{1-6} -alkoxy;

10 R^4 is C_{3-8} -cycloalkyl;

R^5 is H;

R^6 is aminosulfonyl;

R^7 is H or halogen;


R^8 is H;

15 R^9 is halogen or C_{1-6} -alkoxy;

m , n , p and q are 1

or pharmaceutically acceptable salts.

A particular embodiment of the present invention are compounds according to formula I(a) as described herein,



wherein

R^1 is substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted pyridinyl or substituted pyridinyl- C_{1-6} -alkyl, wherein substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted pyridinyl and substituted pyridinyl- C_{1-6} -alkyl are substituted by R^3 , R^4 and R^5 ;

Y is a $-OC(O)-$ or $-C(O)-$;

W is $-C(O)-$;

R^2 is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R^6 , R^7 and R^8 ;

R^3 is halo- C_{1-6} -alkoxy, C_{1-6} -alkylcarbonylamino or tetrahydropyranyl- C_{1-6} -alkoxy;

R^4 is H, C_{1-6} -alkyl or C_{3-8} -cycloalkyl;

R^5 is H;


R^7 and R^8 are independently selected from H and halogen;

R^9 is C_{1-6} -alkyl, C_{1-6} -alkoxy or halogen;

m, n, p and q are 1;

or pharmaceutically acceptable salts.

A further particular embodiment of the present invention are compounds according to formula I(b) as described herein,



wherein

R¹ is substituted pyridinyl or substituted pyridinyl-C₁₋₆-alkyl, wherein substituted pyridinyl and substituted pyridinyl-C₁₋₆-alkyl are substituted by R³, R⁴ and R⁵;

Y is -C(O)-;

5 W is -C(O)-;

R² is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R⁶, R⁷ and R⁸;

R³ is tetrahydropyranyl-C₁₋₆-alkoxy;

R⁴ is C₃₋₈-cycloalkyl;

10 R⁵ is H;

R⁷ is H or halogen;

R⁸ is H;

R⁹ is halogen or C₁₋₆-alkoxy;

m, n, p and q are 1

15 or pharmaceutically acceptable salts.

Particular examples of compounds of formula (I) as described herein are selected from

trans-5-(2-fluoro-4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester;

20 trans-4-[3a-methoxy-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide;

trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide;

trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester;

trans-5-(4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

5 trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

trans-3a-methyl-5-(4-sulfamoyl-benzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

10 trans-5-(5-sulfamoylpyridine-2-carbonyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester;

trans-6-[3a-methoxy-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]pyridine-3-sulfonamide;

15 trans-6-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]pyridine-3-sulfonamide;

trans-5-(2-fluoro-4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

trans-4-[3a-fluoro-2-[(E)-3-[4-(trifluoromethoxy)phenyl]prop-2-enoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide;

20 trans-4-[3a-fluoro-2-[2-[4-(trifluoromethoxy)phenoxy]acetyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide;

trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-fluoro-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide;

25 trans-4-[3a-fluoro-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide;

(+)-(3aR,6aR)-3a-methyl-5-(4-sulfamoyl-benzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(-)-trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester ;

(-)-trans-5-(4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester ;

5 (-)-trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester ;

(-)-(3aS,6aS)-3a-methyl-5-(4-sulfamoylbenzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

10 (+)-trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(+)-trans-5-(4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester ;

15 (+)-trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester;

and pharmaceutically acceptable salts thereof.

Further particular examples of compounds of formula (I) as described herein are selected from

20 trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide;

trans-6-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]pyridine-3-sulfonamide;


25 trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-fluoro-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide;

and pharmaceutically acceptable salts thereof.

Processes for the manufacture of compounds of formula (I) as described herein are an object of the invention.

The preparation of compounds of formula (I) of the present invention may be carried out in sequential or convergent synthetic routes. Syntheses of the invention are shown in the following general schemes. The skills required for carrying out the reactions and purifications of the resulting products are known to those persons skilled in the art. In case a mixture of enantiomers or diastereoisomers is produced during a reaction, these enantiomers or diastereoisomers can be separated by methods described herein or known to the man skilled in the art such as e.g. (chiral) chromatography or crystallization. The substituents and indices used in the following description of the processes have the significance given herein.


Compounds of general formula (I) can be synthesised from amine precursor **1** and appropriate reagents, using methods well known in the art.

**1**

For instance, amine **1** is reacted with a suitable carboxylic acid of formula $R^1\text{-COOH}$ (**2**) leading to a compound of formula (I), wherein Y is -C(O)- . The reaction is performed in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

Amine **1** can also be reacted with suitable acylating reagents such as acyl chlorides of formula $R^1\text{-COCl}$ (**3**) to lead to compounds of formula (I), wherein Y is -C(O)- . The reaction is performed in a solvent such as dichloromethane, tetrahydrofuran, or N,N-dimethylformamide, in the presence of a base such as triethylamine or 4-methylmorpholine, at temperatures between 0°C and 80°C .

Alternatively, amine **1** is reacted with a suitable chloroformate ester of formula $R^1-O-C(O)-Cl$ (**4**), or with an imidazole-1-carboxylate ester of formula (**3**), leading to a compound of formula (I) wherein Y is $-OC(O)-$.



5 The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g., triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.


10 Chloroformate esters **4** are commercially available or can be synthesised from the corresponding alcohol of formula R^1-OH , by reaction with phosgene or a phosgene equivalent (e. g., diphosgene, triphosgene), as described in the literature.

15 Imidazole-1-carboxylate esters **5** are synthesised from the corresponding alcohols of formula R^1-OH , by reaction with 1,1'-carbonyldiimidazole. The reaction is performed at room temperature, in a solvent such as dichloromethane, tetrahydrofuran or acetonitrile. The imidazole-1-carboxylate esters **5** are typically not isolated but directly reacted with amines **1** as described above.

Alcohols of formula R^1-OH are commercially available or can be produced by methods described herein or known in the art.

20 Carboxylic acids (**2**) and acyl halides (**3**) are commercially available or can be prepared as described herein or in the literature.

Amines of general formula **1** are synthesised from suitably protected precursors **6**.




25 Suitable protective groups (PG) are tert-butoxycarbonyl or benzyloxycarbonyl. The deprotection of intermediates **6** can be performed using methods and reagents known in the art.

For instance, in the case where PG is benzyloxycarbonyl, the deprotection may be performed by hydrogenation at pressures between 1 bar and 100 bar, in the presence of a suitable catalyst such as palladium on activated charcoal, at temperatures between 20°C and 150°C in solvents such as methanol or ethanol.

- 5 Alternatively, in the case where PG is tert-butoxycarbonyl, the deprotection may be performed in the presence of a suitable acid, e. g. hydrochloric acid or trifluoroacetic acid, in a solvent such as water, 2-propanol, dichloromethane, or 1,4-dioxane at temperatures between 0°C and 30°C.

Intermediates **6** can be produced from amine precursors of general formula **7** by reaction
10 with appropriate reagents, using methods known in the art.



For instance, **7** is reacted with alkylating agents of general formula $X-CR^7R^8-R^2$ (**8**) where X is a leaving group such as Cl, Br, I, or OSO_2CH_3 , leading to **6**, wherein W is $-CR^7R^8-$. This reaction is performed in a solvent such as tetrahydrofuran or N,N-dimethylformamide, in the presence of a base, e. g. triethylamine or potassium carbonate, at
15 temperatures between 0°C and 100°C.

Alternatively, for compounds of formula **6**, wherein W is $-CR^7R^8-$, R^7 is hydrogen, alkyl or cycloalkyl, and R^8 is H, amine **7** is reacted with aldehydes or ketones of general formula $R^7-C(O)-R^2$ (**9**) in a reductive amination reaction, leading to **6**. This reaction is performed in the
20 presence of a suitable reducing agent, e. g., sodium borohydride or sodium triacetoxyborohydride, in a solvent such as methanol, acetic acid, tetrahydrofuran, 1,2-dichloroethane or mixtures thereof, at temperatures between 0°C and 50°C.


Alternatively, amine **7** is reacted with a suitable carboxylic acid of formula R^2-COOH (**10**), leading to compounds of formula **6**, wherein W is $-C(O)-$. The reaction is performed in the
25 presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium

hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

- 5 Alternatively, amine **7** is reacted with a suitable sulfonyl chloride of formula R^2-SO_2Cl (**11**), leading to compounds of formula **6**, wherein W is $-S(O_2)-$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g. triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.

Amines **7**, alkylating agents **8**, aldehydes/ketones **9**, carboxylic acids **10**, and sulfonyl chlorides **11** are commercially available or can be synthesised as described herein or in the literature.

- 15 Compounds of formula (I), can be produced from amine precursors of general formula **12** by reaction with appropriate reagents, using methods known in the art.




- For instance, an amine of formula **12** is reacted with alkylating agents of general formula $X-CR^7R^8-R^2$ (**8**) where X is a leaving group such as Cl, Br, I, or OSO_2CH_3 , leading to compounds of formula (I), wherein W is $-CR^7R^8-$. This reaction is performed in a solvent such as tetrahydrofuran or N,N-dimethylformamide, in the presence of a base, e. g., triethylamine or potassium carbonate, at temperatures between 0°C and 100°C.

- Alternatively, an amine of formula **12** is reacted with aldehydes or ketones of general formula $R^7-C(O)-R^2$ (**9**) in a reductive amination reaction, leading to compounds of formula (I) wherein W is $-CR^7R^8-$, R^7 is hydrogen, alkyl or cycloalkyl, and R^8 is H. This reaction is performed in the presence of a suitable reducing agent, e. g. sodium borohydride or sodium triacetoxyborohydride, in a solvent such as methanol, acetic acid, tetrahydrofuran, 1,2-dichloroethane or mixtures thereof, at temperatures between 0°C and 50°C.


Alternatively, amine **12** is reacted with a suitable carboxylic acid of formula $R^2\text{--COOH}$ (**10**), leading to compounds of formula (I) wherein W is --C(O)-- . The reaction is performed in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-
 5 (benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine,
 10 diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

Alternatively, amine **12** is reacted with a suitable sulfonyl chloride of formula $R^2\text{--SO}_2\text{Cl}$ (**11**), leading to (I) wherein W is $\text{--S(O}_2\text{)--}$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g. triethylamine, diisopropylethylamine,
 15 pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.

Amines **12** can be synthesised from their tert-butyl carbamate derivatives of formula **13** by carbamate deprotection. The deprotection may be performed in the presence of a suitable acid, e. g., hydrochloric acid or trifluoroacetic acid, in a solvent such as water, 2-propanol,
 20 dichloromethane, or 1,4-dioxane, at temperatures between 0°C and 30°C .



tert-Butyl carbamates **13** can be synthesised from amine precursors of formula **14** and appropriate reagents, using methods well known in the art.



25 For instance, amine **14** is reacted with a suitable carboxylic acid of formula $R^1\text{--COOH}$ (**2**) leading to compounds of formula **13**, wherein Y is --C(O)-- . The reaction is performed in the


presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium

- 5 hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

- Amine **14** can also be reacted with suitable acylating reagents, such as acyl chlorides of formula R^1-COCl (**3**) to provide compounds of formula **13**, wherein Y is $-C(O)-$. The reaction is performed in a solvent such as dichloromethane, tetrahydrofuran, or N,N-dimethylformamide, in the presence of a base such as triethylamine or 4-methylmorpholine, at temperatures between 0°C and 80°C.


- Alternatively, amine **14** is reacted with a suitable chloroformate ester of formula $R^1-O-C(O)-Cl$ (**4**), or with an imidazole-1-carboxylate ester of formula **5**, leading to a compound of formula **13**, wherein Y is $-OC(O)-$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g., triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.

- Alternatively, amine **14** can be reacted with a phosgene or a phosgene equivalent (e. g., triphosgene) to the corresponding N-chlorocarbonylamine **14A**, in the presence of a base (e. g., pyridine) in a suitable solvent, e. g., dichloromethane, at temperatures between -78°C and +20°C. N-Chlorocarbonylamine **14A** is then reacted with alcohol of formula R^1-OH , leading to a compound of formula **13**, wherein Y is $-OC(O)-$. This reaction is performed in a suitable solvent (e. g., acetonitrile or dichloromethane) in the presence of a suitable base (e. g., sodium hydride, pyridine or polystyrene-bound 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,3,2-diazaphosphorine), at temperatures between 20°C and the boiling point of the solvent.



Amines of formula **22** are commercially available or can be produced as described herein or in the literature.

Also an embodiment of the present invention is a process to prepare a compound of formula (I) as defined above comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III);



wherein R^1 , R^2 , R^9 , m , n , p and q are as defined above and W is $-\text{C}(\text{O})-$.

In particular, in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N' -dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O -(benzotriazol-1-yl)- $\text{N},\text{N},\text{N}',\text{N}'$ -tetramethyluronium hexafluoro-phosphate, O -(7-azabenzotriazol-1-yl)- $\text{N},\text{N},\text{N}',\text{N}'$ -tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, particularly O -(7-azabenzotriazol-1-yl)- $\text{N},\text{N},\text{N}',\text{N}'$ -tetramethyluronium hexafluoro-phosphate, in an aprotic solvent such as dichloromethane, tetrahydrofuran, N,N -dimethylformamide, N -methylpyrrolidinone and mixtures thereof, particularly N,N -dimethylformamide, in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine, particularly in the presence of 4-methylmorpholine and at a temperature comprised between -78°C and reflux, particularly between -10°C and room temperature.

Also an object of the present invention is a compound according to formula (I) as described herein for use as a therapeutically active substance.

Likewise an object of the present invention is a pharmaceutical composition comprising a compound according to formula (I) as described herein and a therapeutically inert carrier.

A particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of ocular conditions, particularly glaucoma.

The present invention also relates to the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of ocular
5 conditions, particularly glaucoma.

Also an object of the invention is a method for the treatment or prophylaxis of ocular conditions, particularly glaucoma, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Inflammatory conditions include, but are not limited to, arthritis, osteoarthritis,
10 multiple sclerosis, systemic lupus erythematoses, inflammatory bowel disease, abnormal evacuation disorder and the like as well as inflammatory airways diseases such as idiopathic pulmonary fibrosis (IPF), chronic obstructive pulmonary disease (COPD) or chronic asthma bronchiale.

Further conditions of the respiratory system include, but are not limited to, other diffuse
15 parenchymal lung diseases of different etiologies including iatrogenic drug-induced fibrosis, occupational and/or environmental induced fibrosis, systemic diseases and vasculitides, granulomatous diseases (sarcoidosis, hypersensitivity pneumonia), collagen vascular disease, alveolar proteinosis, Langerhans cell granulomatosis, lymphangioleiomyomatosis, inherited diseases (Hermansky-Pudlak Syndrome, tuberous sclerosis, neurofibromatosis, metabolic storage
20 disorders, familial interstitial lung disease), radiation induced fibrosis, silicosis, asbestos induced pulmonary fibrosis or acute respiratory distress syndrome (ARDS).

Conditions of the nervous system include, but are not limited to, neuropathic pain, schizophrenia, neuro-inflammation (e.g. astrogliosis), peripheral and/or autonomic (diabetic) neuropathies and the like.

25 Vascular conditions include, but are not limited to, atherosclerosis, thrombotic vascular disease as well as thrombotic microangiopathies, proliferative arteriopathy (such as swollen myointimal cells surrounded by mucinous extracellular matrix and nodular thickening), atherosclerosis, decreased vascular compliance (such as stiffness, reduced ventricular compliance and reduced vascular compliance), endothelial dysfunction and the like.

Cardiovascular conditions include, but are not limited to, acute coronary syndrome, coronary heart disease, myocardial infarction, arterial and pulmonary hypertension, cardiac arrhythmia such as atrial fibrillation, stroke and other vascular damage.

Fibrotic diseases include, but are not limited to myocardial and vascular fibrosis,
5 pulmonary fibrosis, skin fibrosis, scleroderma and encapsulating peritonitis.

Cancer and cancer metastasis include, but are not limited to, breast cancer, ovarian cancer, lung cancer, prostate cancer, mesothelioma, glioma, gastrointestinal cancers and progression and metastatic aggressiveness thereof.

Ocular conditions include, but are not limited to, proliferative and non-proliferative
10 (diabetic) retinopathy, dry and wet age-related macular degeneration (AMD), macular edema, central arterial /venous occlusion, traumatic injury, glaucoma and the like. Particularly, the ocular condition is glaucoma.

Metabolic conditions include, but are not limited to, obesity and diabetes.

Also an embodiment of the present invention are compounds of formula (I) as described
15 herein, when manufactured according to any one of the described processes.

Assay procedures

PRODUCTION OF HUMAN FULL LENGTH ATX, WITH AND WITHOUT HIS TAG

Autotaxin (ATX - ENPP2) cloning: cDNA was prepared from commercial human hematopoietic cells total RNA and used as template in overlapping PCR to generate a full length
20 human ENPP2 ORF with or without a 3'-6xHis tag. These full length inserts were cloned into the pcDNA3.1V5-His TOPO (Invitrogen) vector. The DNA sequences of several single clones were verified. The DNA from a correct full length clone was used to transfect Hek293 cells for verification of protein expression. The sequence of the encoded ENPP2 conforms to Swissprot entry Q13822, with or without the additional C-terminal 6xHis tag.

25 **ATX Fermentation:** Recombinant protein was produced by large-scale transient transfection in 20 L controlled stirred tank bioreactors (Sartorius). During cell growth and transfection, temperature, stirrer speed, pH and dissolved oxygen concentration were maintained at 37°C, 120 rpm, 7.1 and 30% DO, respectively. FreeStyle 293-F cells (Invitrogen) were cultivated in suspension in FreeStyle 293 medium (Invitrogen) and transfected at ca. 1-1.5 x 10E6 cells/mL

with above plasmid DNAs using X-tremeGENE Ro-1539 (commercial product, Roche Diagnostics) as complexing agent. Cells were fed a concentrated nutrient solution (J Immunol Methods 194 (1996), 19, 1-199 (page 193)) and induced by sodium butyrate (2 mM) at 72 h post-transfection and harvested at 96 h post-transfection. Expression was analyzed by Western Blot, enzymatic assay and/or analytical IMAC chromatography. After cooling the cell suspension to 4°C in a flow-through heat exchanger, cell separation and sterile filtration of supernatant was performed by filtration through Zeta Plus 60M02 E16 (Cuno) and Sartopore 2 XLG (Sartorius) filter units. The supernatant was stored at 4°C prior to purification.

ATX Purification: 20 liter of culture supernatant were conditioned for ultrafiltration by adding Brij 35 to a final concentration of 0.02% and by adjusting the pH to 7.0 using 1 M HCl. Then the supernatant was first microfiltered through a 0.2 µm Ultran-Pilot Open Channel PES filter (Whatman) and afterwards concentrated to 1 liter through an Ultran-Pilot Screen Channel PES filter with 30 kDa MWCO (Whatman). Prior to IMAC chromatography, NiSO₄ was added to a final concentration of 1 mM. The cleared supernatant was then applied to a HisTrap column (GE Healthcare) previously equilibrated in 50 mM Na₂HPO₄ pH 7.0, 0.5 M NaCl, 10% glycerol, 0.3% CHAPS, 0.02% NaN₃. The column was washed stepwise with the same buffer containing 20 mM, 40 mM and 50 mM imidazole, respectively. The protein was subsequently eluted using a linear gradient to 0.5 M imidazole in 15 column volumes. ATX containing fractions were pooled and concentrated using an Amicon cell equipped with a 30 kDa PES filter membrane. The protein was further purified by size exclusion chromatography on Superdex S-200 prep grade (XK 26/100) (GE Healthcare) in 20 mM BICINE pH 8.5, 0.15 M NaCl, 10% glycerol, 0.3% CHAPS, 0.02% NaN₃. Final yield of protein after purification was 5-10 mg ATX per liter of culture supernatant. The protein was stored at -80°C.

HUMAN ATX ENZYME INHIBITION ASSAY

ATX inhibition was measured by a fluorescence quenching assay using a specifically labeled substrate analogue (MR121 substrate). To obtain this MR121 substrate, BOC and TBS protected 6-amino-hexanoic acid (R)-3-({2-[3-(2-{2-[2-(2-amino-ethoxy)-ethoxy]-ethoxy}-ethoxy)-propionylamino]-ethoxy}-hydroxy-phosphoryloxy)-2-hydroxy-propyl ester (Ferguson et al., Org Lett 2006, 8 (10), 2023) was labeled with MR121 fluorophore (CAS 185308-24-1, 1-(3-carboxypropyl)-11-ethyl-1,2,3,4,8,9,10,11-octahydro-dipyrido[3,2-b:2',3'-i]phenoxazin-13-ium) on the free amine of the ethanolamine side and then, after deprotection, subsequently with tryptophan on the side of the aminohexanoic acid.

Assay working solutions were made as follows:

Assay buffer (50 mM Tris-HCl, 140 mM NaCl, 5 mM KCl, 1 mM CaCl₂, 1 mM MgCl₂, 0.01% Triton-X-100, pH 8.0;

ATX solution: ATX (human His-tagged) stock solution (1.08 mg/mL in 20mM bicine, pH 8.5,

5 0.15 M NaCl, 10% glycerol, 0.3% CHAPS, 0.02% NaN₃), diluted to 1.4 – 2.5x final concentration in assay buffer;

MR121 substrate solution: MR121 substrate stock solution (800 µM MR121 substrate in DMSO), diluted to 2 – 5x final concentration in assay buffer.

Test compounds (10 mM stock in DMSO, 8 µL) were obtained in 384 well sample plates

10 (Corning Costar #3655) and diluted with 8 µL DMSO. Row-wise serial dilutions were made by transferring 8 µL cpd solution to the next row up to row O. The compound and control solutions were mixed five times and 2 µL were transferred to 384 well assay plates (Corning Costar # 3702). Then, 15 µL of 41.7 nM ATX solution was added (30 nM final concentration), mixed five times and then incubated for 15 minutes at 30°C. 10 µL of MR121 substrate solution was added

15 (1µM final concentration), mixed 30 times and then incubated for 15 minutes at 30 °C.

Fluorescence was then measured every 2 minutes for 1 hour (Perkin Elmer plate: vision multimode reader); light intensity: 2.5%; exp. time: 1.4 sec, Filter: Fluo_630/690 nm) and IC₅₀ values were calculated from these readouts.

Human Carbonic Anhydrase-II Inhibition assay

20 Human carbonic anhydrase II (hCA-II) inhibition was measured by an absorbance method using 4-nitrophenyl acetate (4-NPA) as its substrate. 4-NPA can be catalyzed by active hCA II via a zinc-hydroxide mechanism. The nitrophenolate in the products can be ionized to generate a bright yellow anion with high absorbance at 348 to 400 nm, as reported in the literature (Armstrong et al., J. Biol. Chem. 1966, 241, 5137-5149). OD340 nm was chosen for detecting

25 hCA II substrate conversion.

Assay working solutions were made as follows:

Assay buffer: 50mM MOPS, 33mM Na₂SO₄, 1mM EDTA, 0.5 mg/ml BSA, pH 7.5;

Enzyme solution: hCA-II (human, full length) stock solution (1.0 mg/mL in 20mM HEPES, 50 mM NaCl, pH 7.4), diluted to 2133x final concentration in assay buffer;

4-NPA substrate solution: 4-NPA substrate stock solution (250 mM in DMSO, stored at -20°C), diluted to 50x final concentration in deionized water.

Test compounds (10 mM stock in DMSO, 100 μ L) were obtained in 96-well sample plates (Corning Costar #3655) and diluted to 0.5 mM. Column-wise serial dilutions were made by transferring 20 μ L compound solutions to the next column, from column 3 up to 22. After this, 1.2 μ L were transferred to 384 well assay plates (Corning Costar # 3701). Then 30 μ L of 16 nM hCA II solution was added (8 nM final concentration), mixed five times. 30 μ L of 4-NPA substrate solution was added (2.5 mM final concentration), mixed five times. Absorbance at 340 nm was then measured immediately as time zero. The assay plates were incubated at room temperature for 1 hour and then measured as time 1 hour (Perkin Elmer EnVision 2103; Filter: Photometric 340; Light intensity 60%; Number of flashes: 10). IC₅₀ values and K_i values were calculated from these readouts.

Ex	ATX IC50 (μ M)	CA-II IC50 (μ M)
1.00	0.008	0.005
1.01	0.01	0.008
1.02	0.006	0.01
1.03	0.006	
1.04	0.003	
1.05	0.02	
1.06	0.006	
1.07	0.006	0.013

Ex	ATX IC50 (μ M)	CA-II IC50 (μ M)
1.08	0.017	0.018
1.09	0.01	0.01
2.00	0.007	0.006
3.00	0.005	0.009
3.01	0.007	0.006
3.02	0.008	0.006
3.03	0.004	0.006
4.00	0.006	0.095

Ex	ATX IC50 (μ M)	CA-II IC50 (μ M)
4.01	0.013	0.018
4.02	0.008	0.01
4.03	0.011	0.013
5.00	0.007	0.038
5.01	0.006	0.027
5.02	0.004	0.015
5.03	0.013	0.02

15

Compounds of formula (I) and their pharmaceutically acceptable salts or esters thereof as described herein have IC_{50} values between $0.00001\ \mu M$ and $1000\ \mu M$, particular compounds have IC_{50} values between $0.0005\ \mu M$ and $500\ \mu M$, further particular compounds have IC_{50} values between $0.0005\ \mu M$ and $50\ \mu M$, more particular compounds have IC_{50} values between $0.0005\ \mu M$ and $5\ \mu M$. These results have been obtained by using the enzymatic assay described above.

The compounds of formula (I) and their pharmaceutically acceptable salts can be used as medicaments (e.g. in the form of pharmaceutical preparations). The pharmaceutical preparations can be administered internally, such as orally (e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatin capsules, solutions, emulsions or suspensions), nasally (e.g. in the form of nasal sprays), rectally (e.g. in the form of suppositories) or topical ocularly (e.g. in the form of solutions, ointments, gels or water soluble polymeric inserts). However, the administration can also be effected parenterally, such as intramuscularly, intravenously, or intraocularly (e.g. in the form of sterile injection solutions).

The compounds of formula (I) and their pharmaceutically acceptable salts can be processed with pharmaceutically inert, inorganic or organic adjuvants for the production of tablets, coated tablets, dragées, hard gelatin capsules, injection solutions or topical formulations. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts etc. can be used, for example, as such adjuvants for tablets, dragées and hard gelatin capsules.

Suitable adjuvants for soft gelatin capsules, are, for example, vegetable oils, waxes, fats, semi-solid substances and liquid polyols, etc.

Suitable adjuvants for the production of solutions and syrups are, for example, water, polyols, saccharose, invert sugar, glucose, etc.

Suitable adjuvants for injection solutions are, for example, water, alcohols, polyols, glycerol, vegetable oils, etc.

Suitable adjuvants for suppositories are, for example, natural or hardened oils, waxes, fats, semi-solid or liquid polyols, etc.

Suitable adjuvants for topical ocular formulations are, for example, cyclodextrins, mannitol or many other carriers and excipients known in the art.

5 Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, viscosity-increasing substances, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

10 The dosage can vary in wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of about 0.1 mg to 20 mg per kg body weight, preferably about 0.5 mg to 4 mg per kg body weight (e.g. about 300 mg per person), divided into preferably 1-3 individual doses, which can consist, for example, of the same amounts, should it be appropriate. In the case of topical
15 administration, the formulation can contain 0.001% to 15% by weight of medicament and the required dose, which can be between 0.1 and 25 mg in can be administered either by single dose per day or per week, or by multiple doses (2 to 4) per day, or by multiple doses per week It will, however, be clear that the upper or lower limit given herein can be exceeded when this is shown to be indicated.

The invention is illustrated hereinafter by Examples, which have no limiting character.

20 In case the preparative examples are obtained as a mixture of enantiomers, the pure enantiomers can be obtained by methods described herein or by methods known to those skilled in the art, such as e.g. chiral chromatography or crystallization.


Examples

All examples and intermediates were prepared under nitrogen atmosphere if not specified otherwise.

Abbreviations: aq. = aqueous; CAS-RN = Chemical Abstracts Service Registry Number; HPLC = high performance liquid chromatography; MS = mass spectrum; PS-BEMP = polystyrene-bound 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,3,2-diazaphosphorine; sat. = saturated




Example 1





- 10 **trans-5-(2-Fluoro-4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester**





- To a solution of trans-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester dihydrochloride (intermediate 5; 96 mg, 186 μ mol), 4-methylmorpholine (113 mg, 1.11 mmol) and 2-fluoro-4-sulfamoylbenzoic acid (CAS-RN 714968-42-0; 42.8 mg, 186 μ mol) in N,N-dimethylformamide (3 mL) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (70.6 mg, 186 μ mol) at room temperature, then after 18 h the reaction mixture was partitioned sat. aq. sodium hydrogen carbonate solution and ethyl acetate/2-methyltetrahydrofuran 4:1. The organic layer was washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate, filtered and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25 afforded the title compound (107 mg, 80%). White solid, MS: 646.2 (M+H)⁺.

The following examples were produced in analogy to example 1, replacing trans-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester dihydrochloride by the appropriate amine and 2-fluoro-4-sulfamoylbenzoic acid by the appropriate carboxylic acid.


Ex.	Systematic Name	Amine / Carboxylic acid	MS, m/e
1.01	trans-4-[3a-methoxy-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide 	trans-1-(3a-methoxyhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one dihydrochloride (intermediate 6.01) / 2-fluoro-4-sulfamoylbenzoic acid (CAS-RN 714968-42-0)	560.2 (M+H) ⁺
1.02	trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide 	trans-(2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridin-4-yl)3a-methoxyhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 6) / 2-fluoro-4-sulfamoylbenzoic acid (CAS-RN 714968-42-0)	603.3 (M+H) ⁺
1.03	trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester 	trans-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester dihydrochloride (intermediate 5) / 4-sulfamoylbenzoic acid (CAS-RN 138-41-0)	628.2 (M+H) ⁺

Ex.	Systematic Name	Amine / Carboxylic acid	MS, m/e
1.04	trans-5-(4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	trans-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester (intermediate 4.01) / 4-sulfamoylbenzoic acid (CAS-RN 138-41-0)	530.3 (M-H) ⁻
1.05	trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	trans-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester (intermediate 4.01) / 4-sulfamoylbenzoic acid (CAS-RN 138-41-0)	544.2 (M+H) ⁺
1.06	trans-3a-methyl-5-(4-sulfamoyl-benzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	trans-3a-methyl-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester hydrochloride (intermediate 4) / 4-sulfamoylbenzoic acid (CAS-RN 138-41-0)	528.2 (M+H) ⁺
1.07	trans-5-(5-sulfamoylpyridine-2-carbonyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester 	trans-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester dihydrochloride (intermediate 5) / 5-sulfamoylpicolinic acid (CAS-RN 1308677-67-9)	629.2 (M+H) ⁺

Ex.	Systematic Name	Amine / Carboxylic acid	MS, m/e
1.08	trans-6-[3a-methoxy-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]pyridine-3-sulfonamide 	trans-1-(3a-methoxyhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one dihydrochloride (intermediate 6.01) / 5-sulfamoylpicolinic acid (CAS-RN 1308677-67-9)	543.1 (M+H) ⁺
1.09	trans-6-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]pyridine-3-sulfonamide 	trans-(2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridin-4-yl)3a-methoxyhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 6) / 5-sulfamoylpicolinic acid(CAS-RN 1308677-67-9)	586.2 (M+H) ⁺

Example 2

trans-5-(2-Fluoro-4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester




5

To a solution of (4-(trifluoromethoxy)phenyl)methanol (26.1 mg, 136 μ mol) in acetonitrile (5 mL) was added 1,1'-carbonyldiimidazole (22 mg, 136 μ mol) and the reaction mixture was heated at 50°C for 2 h, then triethylamine (68.8 mg 680 μ mol) and 3-fluoro-4-(trans-3a-fluorooctahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzenesulfonamide hydrochloride (50 mg,

136 μmol) were added and the reaction mixture was heated at reflux for another 15 h. After cooling the reaction mixture was partitioned between ethyl acetate and sat. aq. sodium hydrogen carbonate solution. The organic layer was washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate, filtered and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25 afforded the title compound (59 mg, 79%). White foam, MS: 550.1 (M+H)⁺.




Example 3

trans-4-[3a-Fluoro-2-[(E)-3-[4-(trifluoromethoxy)phenyl]prop-2-enoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide



To a solution of 3-fluoro-4-((3aR,6aS)-3a-fluorooctahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzenesulfonamide hydrochloride (intermediate 9; 50 mg, 122 μmol), 4-methylmorpholine (61.9 mg, 612 μmol) and (E)-3-(4-(trifluoromethoxy)phenyl)acrylic acid (29 mg, 122 μmol) in N,N-dimethylformamide (4 mL) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (46.5 mg, 122 μmol) at room temperature, then after 18 h the reaction mixture was partitioned between sat. aq. sodium hydrogen carbonate solution and ethyl acetate/2-methyltetrahydrofuran 4:1. The organic layer was washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate filtered and evaporated. Chromatography (silica gel; gradient dichloromethane/methanol/25% aq. ammonia solution 97.5:2.5:0.25 to 95:5:0.25) afforded the title compound (48 mg, 72%). White solid, MS: 546.1 (M+H)⁺.


The following examples were produced in analogy to example 3, replacing 3-fluoro-4-((3aR,6aS)-3a-fluorooctahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzenesulfonamide hydrochloride by the appropriate amine and (E)-3-(4-(trifluoromethoxy)phenyl)acrylic acid by the appropriate carboxylic acid.

Ex.	Systematic name	Amine / Carboxylic acid	MS, m/e
3.01	trans-4-[3a-fluoro-2-[2-[4-(trifluoromethoxy)phenoxy]acetyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide 	3-fluoro-4-(trans-3a-fluorooctahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzenesulfonamide hydrochloride (intermediate 9) / 2-(4-(trifluoromethoxy)phenoxy)acetic acid (CAS-RN 72220-50-9)	548.2 (M+H) ⁺
3.02	trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-fluoro-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide 	3-fluoro-4-(trans-3a-fluorooctahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzenesulfonamide hydrochloride (intermediate 9) / 2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinic acid (intermediate 7)	591.2 (M+H) ⁺
3.03	trans-4-[3a-fluoro-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide 	3-fluoro-4-(trans-3a-fluorooctahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzenesulfonamide hydrochloride (intermediate 9) / 3-(4-(trifluoromethoxy)phenyl)propanoic acid	546.2 (M-H) ⁻

Examples 4 and 5

(+)-(3aR,6aR)-3a-Methyl-5-(4-sulfamoyl-benzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester and (-)-(3aS,6aS)-3a-methyl-5-(4-

5 sulfamoylbzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester



Racemic trans-3a-methyl-5-(4-sulfamoyl-benzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester (example 1.06; 180 mg, 341 μ mol) was separated by preparative HPLC using Chiralpak AD as the stationary phase and

- 5 heptane/ethanol/ammonium acetate 60:40:0.004 as the eluent. This produced the faster eluting (+)-enantiomer (example 4; 67 mg, 37%; white foam, MS: 528.2 (M+H)⁺) and the slower eluting (–)-enantiomer (example 5; 63 mg, 35%; white foam, MS: 528.2 (M+H)⁺).

The following examples were produced in analogy to examples 4 and 5 by chiral HPLC

- 10 separation of their racemates, using stationary phase and eluent as indicated below.

Ex.	Systematic Name	Optical rotation sign	Stationary phase; eluent	MS, m/e
4.01	trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester	(–)	Reprosil Chiral-NR; heptane/ethanol/ammonium acetate 60:40:0.004	542.3 (M–H) [–]
5.01	(example 1.05)	(+)		542.3 (M–H) [–]
4.02	trans-5-(4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester	(–)	Reprosil Chiral-NR; heptane/ethanol/ammonium acetate 70:30:0.003	530.2 (M–H) [–]
5.02	(example 1.04)	(+)		530.2 (M–H) [–]
4.03	trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester	(–)	Reprosil Chiral-NR; heptane/ethanol 60:40	628.3 (M+H) ⁺
5.03	(example 1.03)	(+)		628.3 (M+H) ⁺

IntermediatesIntermediate 1**trans-tert-Butyl 3a-methylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate**5 Step 1: trans-Dimethyl 1-benzyl-3-methylpyrrolidine-3,4-dicarboxylate

A solution of N-benzyl-1-methoxy-N-((trimethylsilyl)methyl)methanamine (CAS-RN 93102-05-7; 1.74 g, 17.0 mmol) in dichloromethane (5 mL) was added to an ice-cooled mixture of dimethyl 2-methylfumarate (CAS-RN 617-53-8; 1.00 g, 6.32 mmol) and trifluoroacetic acid (79.3 mg, 696 μ mol) in dichloromethane (10 mL) at 0–5°C. The resulting yellow solution was

- 10 allowed to reach room temperature over 20 h, then partitioned between sat. aq. hydrogencarbonate solution and dichloromethane. The organic layer was washed with brine, dried over magnesium sulfate, filtered and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 95:5:0.25) afforded the title compound (1.61 g, 87%). Light yellow oil, MS: 292.2 (M+H)⁺.

15 Step 2: trans-1-tert-Butyl 3,4-dimethyl 3-methylpyrrolidine-1,3,4-tricarboxylate

A solution of trans-dimethyl 1-benzyl-3-methylpyrrolidine-3,4-dicarboxylate (1.56 g, 5.35 mmol) and di-tert-butyl dicarbonate (1.29 g, 5.89 mmol) in methanol (20 mL) was stirred at room temperature under a hydrogen atmosphere (1 bar) in the presence of palladium (10% on activated charcoal, 165 mg, 1.55 mmol). After 3 h insoluble material was removed by filtration through

20 diatomaceous earth and the filtrate was concentrated. Chromatography (silica gel; gradient heptane to ethyl acetate/heptane 1:1) afforded the title compound (1.47 g, 91%). Colourless oil, MS: 202.1 (M+H–Me₃COCO)⁺.

Step 3: trans-tert-Butyl 3,4-bis(hydroxymethyl)-3-methylpyrrolidine-1-carboxylate

- 25 A solution of trans-1-tert-butyl 3,4-dimethyl 3-methylpyrrolidine-1,3,4-tricarboxylate (1.47 g, 4.87 mmol) in tetrahydrofuran (12 mL) was cooled to 0°C and treated with lithium borohydride solution (2 M in tetrahydrofuran, 5.47 mL, 10.9 mmol), then after 30 min the ice bath was removed and the reaction mixture was stirred at room temperature. After 18 h excess reagent was destroyed by slow addition of 1 M aq. hydrochloric acid solution at 0°C to pH 1. The resulting clear solution was extracted with dichloromethane, the organic layer was washed with

water and brine, dried over magnesium sulfate, filtered and evaporated. Chromatography (silica gel, heptane–ethyl acetate gradient) afforded the title compound (1.05 g, 88%). Colourless viscous oil, MS: 190.1 (M+H–isobutene)⁺.

Step 4: trans-tert-Butyl 3-methyl-3,4-bis(((methylsulfonyl)oxy)methyl)pyrrolidine-1-carboxylate

- 5 Methanesulfonyl chloride (1.41 g, 12.4 mmol) was added dropwise at 0°C to a clear colourless solution of trans-tert-butyl 3,4-bis(hydroxymethyl)-3-methylpyrrolidine-1-carboxylate (1.01 g, 4.12 mmol) and N,N-diisopropylethylamine (3.19 g, 24.7 mmol) in dichloromethane (10 mL) was cooled to 0 °C. After 30 min the reaction mixture was partitioned between sat. aq. ammonium chloride solution and dichloromethane. The organic layer was washed with sat. aq.
- 10 sodium hydrogencarbonate-solution and brine, dried over magnesium sulfate, filtered and evaporated. Chromatography (silica gel, heptane–ethyl acetate gradient) afforded the title compound (1.53 g, 92%). Yellow viscous oil, MS: 346.1 (M+H–isobutene)⁺.

Step 5: trans-tert-Butyl 5-benzyl-3a-methylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

- To a solution of trans-tert-butyl 3-methyl-3,4-bis(((methylsulfonyl)oxy)methyl)pyrrolidine-1-carboxylate (1.52 g, 3.79 mmol) in toluene (20 mL) were added triethylamine (1.15 g, 11.4 mmol) and benzylamine (811 mg, 7.57 mmol). The reaction mixture was heated at reflux, then after 20 h another portion of triethylamine (1.15 g, 11.4 mmol) and benzylamine (811 mg, 7.57 mmol) was added, then after another 20 h at reflux the reaction mixture was washed with 1 M aq. sodium hydroxide solution and brine, dried over magnesium sulfate, filtered and evaporated.
- 20 Chromatography (silica gel, heptane–ethyl acetate gradient) afforded the title compound (875 mg, 73%). Light yellow solid, MS: 317.2 (M+H)⁺.

Step 6: trans-tert-Butyl 3a-methylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

- A solution of (3aR,6aR)-tert-butyl 5-benzyl-3a-methylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (870 mg, 2.75 mmol) in methanol (10 mL) was stirred at room temperature under a
- 25 hydrogen atmosphere (1 bar) in the presence of palladium (10% on activated charcoal, 146 mg, 1.37 mmol), then after 5 h insoluble material was removed by filtration through diatomaceous earth. The filtrate was evaporated to afford the title compound (640 mg, 93%) containing ca. 10% of methanol. Colourless viscous oil, MS: 227.2 (M+H)⁺.

trans-tert-butyl 3a-fluorohexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

The title compound was produced in analogy to intermediate 1, replacing dimethyl 2-methylfumarate by diethyl 2-fluorofumarate (CAS-RN 4495-77-6). Colourless viscous oil, MS: 231.2 (M+H)⁺.

5

Intermediate 3**trans-tert-Butyl 3a-methoxyhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate**

The title compound was produced in analogy to intermediate 1, replacing dimethyl 2-methylfumarate by dimethyl 2-methoxyfumarate (CAS-RN 2215-05-6). Colourless viscous oil, MS: 243.2 (M+H)⁺.

10

Intermediate 4**trans-3a-Methyl-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester hydrochloride**

15 Step 1: trans-2-tert-Butyl 5-(4-(trifluoromethoxy)benzyl) 3a-methyltetrahydropyrrolo[3,4-c]pyrrole-2,5(1H,3H)-dicarboxylate

To a solution of (4-(trifluoromethoxy)phenyl)methanol (474 mg, 2.47 mmol) in acetonitrile (20 mL) was added 1,1'-carbonyldiimidazole (413 mg, 2.47 mmol) and the reaction mixture was heated to 50°C, then after 3 h triethylamine (1.25 g, 12.3 mmol) and trans-tert-butyl 3a-methylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 1; 621 mg, 2.47 mmol) were added and the reaction mixture was heated at reflux. After 15 h the reaction mixture was partitioned between ethyl acetate sat. aq. sodium hydrogen carbonate solution. The organic layer was washed with aq. sat. ammonium chloride solution and brine, dried over magnesium sulfate, filtered and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 95:5:0.25) afforded the title compound (821 mg, 75%). Light yellow oil, MS: 389.2 (M+H-isobutene)⁺.

20

25

Step 2: trans-3a-methyl-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester hydrochloride

A solution of trans-2-tert-butyl 5-(4-(trifluoromethoxy)benzyl) 3a-methyltetrahydropyrrolo[3,4-c]pyrrole-2,5(1H,3H)-dicarboxylate (812 mg, 1.83 mmol) and hydrochloric acid solution (5–6 M in 2-propanol, 10.2 mL, 51.2 mmol) in 2-propanol (5 mL) was stirred for 15 h at room temperature, then concentrated to dryness. The residue was triturated in tert-butyl methyl ether and the precipitate collected by filtration to produce the title compound (662 mg, 95%). White solid, MS: 345.1 (M+H)⁺.

Intermediate 4.01

trans-3a-Fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester

The title compound was produced in analogy to intermediate 4, replacing trans-tert-butyl 3a-methylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate by trans-tert-butyl 3a-fluorohexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 2). White solid, MS: 349.1 (M+H)⁺.

Intermediate 5

trans-3a-Methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester dihydrochloride

Step 1: trans-tert-Butyl 5-(chlorocarbonyl)-3a-methoxyhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

To a solution of trans-tert-butyl 3a-methoxyhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 3; 440 mg, 1.82 mmol) and pyridine (646 mg, 8.17 mmol) in dichloromethane (5 mL) was added dropwise a solution of triphosgene (242 mg, 817 μmol) in dichloromethane (4 mL) at 0°C. After 30 min the ice bath was removed, then after 4 h the reaction mixture was partitioned between dichloromethane and 1 M aq. hydrochloric acid solution. The organic layer was washed with water and brine, dried over magnesium sulfate, filtered, and evaporated to afford the title compound (372 mg, 67%) as a light yellow foam.

Step 2: trans-2-tert-Butyl 5-((3-pivalamido-5-(trifluoromethyl)pyridin-2-yl)methyl) 3a-methoxytetrahydropyrrolo[3,4-c]pyrrole-2,5(1H,3H)-dicarboxylate

To a solution of trans-tert-butyl 5-(chlorocarbonyl)-3a-methoxyhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (370 mg, 1.21 mmol) in acetonitrile (25 mL) was added N-(2-
5 (hydroxymethyl)-5-(trifluoromethyl)pyridin-3-yl)pivalamide (intermediate 8; 335 mg, 1.21 mmol) and PS-BEMP (CAS-RN 1446424-86-7; 1.5 g, 1.21 mmol). The orange suspension was heated at reflux for 21 h, then insoluble material was removed by filtration. To the filtrate was added PS-Trisamine (CAS-RN 1226492-10-9; 315 mg, 1.21 mmol) and the reaction mixture was stirred at room temperature for 18 h, then insoluble material was removed by filtration and the
10 filtrate was evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia 95:5:0.25) produced the title compound (333 mg, 50%). White foam, MS: 545.3 (M+H)⁺.

Step 3: trans-3a-Methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester dihydrochloride

15 A solution of trans-2-tert-butyl 5-((3-pivalamido-5-(trifluoromethyl)pyridin-2-yl)methyl) 3a-methoxytetrahydropyrrolo[3,4-c]pyrrole-2,5(1H,3H)-dicarboxylate (325 mg, 597 μmol) and hydrochloric acid solution (5–6 M in 2-propanol, 3.34 mL, 16.7 mmol) in 2-propanol (2 mL) was stirred at room temperature for 18 h, then the reaction mixture was concentrated to dryness. The residue was triturated in tert-butyl methyl ether and the precipitate collected by filtration to
20 afford the title compound (291 mg, 94%). White solid, MS: 445.2 (M+H)⁺.

Intermediate 6

trans-(2-Cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridin-4-yl)3a-methoxyhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride

25 Step 1: trans-tert-Butyl 5-(2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinoyl)-3a-methoxyhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

To a solution of trans-tert-butyl 3a-methoxyhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (500 mg, 2.06 mmol) in N,N-dimethylformamide (40 mL) were added 2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinic acid (intermediate 7; 572 mg, 2.06 mmol) and

4-methylmorpholine (1.04 g, 10.3 mmol). The solution was cooled to 0°C, then . O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (824 mg, 2.17 mmol) was added. The ice bath was removed, then after 96 h the reaction mixture was partitioned between sodium hydrogencarbonate solution and ethyl acetate/2-methyltetrahydrofuran 4:1. The organic layer was washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25 afforded the title compound (977 mg; 94%). Yellow viscous oil, MS: 502.3 (M+H)⁺.

Step 2: trans-(2-Cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)pyridin-4-yl)3a-methoxyhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride

A solution of trans-tert-butyl 5-(2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinoyl)-3a-methoxyhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (977 mg, 1.95 mmol) and hydrochloric acid solution (5–6 M in 2-propanol, 8.57 mL, 42.8 mmol). in 2-propanol (8 mL) was stirred at room temperature for 16 h, then the reaction mixture was concentrated to dryness. The residue was triturated in tert-butyl methyl ether and the precipitate collected by filtration to afford the title compound (858 mg, 88%). White solid, MS: 402.3 (M+H)⁺.

Intermediate 6.01

trans-1-(3a-methoxyhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one dihydrochloride

The title compound was produced in analogy to example 6, replacing 2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinic acid by 3-(4-(trifluoromethoxy)phenyl)propanoic acid. Brown viscous oil, MS: 359.2 (M+H)⁺.

Intermediate 7

2-Cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinic acid

Step 1: Methyl 6-cyclopropyl-2-oxo-1,2-dihydropyridine-4-carboxylate

A suspension of 6-cyclopropyl-2-oxo-1,2-dihydropyridine-4-carboxylic acid (CAS-RN 150190-28-6; 400 mg, 2.23 mmol) in methanol (4 mL) and sulfuric acid (12 μ L) was added was heated at 70°C for 48 h, then concentrated in vacuo. The residue was suspended in dichloromethane (10 mL), then insoluble material was removed by filtration and the filtrate evaporated to produce the
5 title compound (427 mg, 99%). Light brown semisolid, MS: 194.1 (M+H)⁺.

Step 2: Methyl 2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinate

To a stirring suspension of methyl 6-cyclopropyl-2-oxo-1,2-dihydropyridine-4-carboxylate (212 mg, 1.1 mmol) in acetonitrile (5 mL) were added potassium carbonate (455 mg, 3.29 mmol) and 4-(iodomethyl)tetrahydro-2H-pyran (CAS-RN 101691-94-5; 744 mg, 3.29 mmol). The reaction
10 mixture was heated at 80°C for 16 h and then evaporated in vacuo. The residue was purified by chromatography (silica gel; heptane–ethyl acetate gradient) to produce the title compound (188 mg, 59%). Colourless oil, MS: 292.2 (M+H)⁺.

Step 3: 2-Cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinic acid

To a solution of methyl 2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinate (184
15 mg, 632 μ mol) in tetrahydrofuran (2 mL) and water (2 mL) was added lithium hydroxide monohydrate (53.0 mg, 1.26 mmol) and the resulting mixture stirred at room temperature for 16 h. The mixture was partially evaporated in order to remove the tetrahydrofuran. The aqueous phase was partitioned between 1 M aq. hydrochloric acid solution and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered and evaporated to give the
20 title compound (218 mg, quant.). Colourless oil, MS: 276.1 (M–H)[–].

Intermediate 8

N-(2-(Hydroxymethyl)-5-(trifluoromethyl)pyridin-3-yl)pivalamide

Step 1: Methyl 3-pivalamido-5-(trifluoromethyl)picolinate

25 To a brown solution of methyl 3-amino-5-(trifluoromethyl)picolinate (CAS-RN 866775-17-9; 2.00 g, 8.63 mmol) in pyridine (25 mL) was added pivaloyl chloride (2.08 g, 17.3 mmol) at 0°C. After 20 min the ice-bath was removed, then after 5 h the reaction mixture was partitioned between 1 M aq. hydrochloric acid solution and ethyl acetate. The organic layer was washed with water and brine, dried over magnesium sulfate, filtered and evaporated. Chromatography

(silica gel; heptane–ethyl acetate gradient) afforded the title compound (2.46 g, 92%). Light yellow solid, MS: 305.1 (M+H)⁺.

Step 2: N-(2-(Hydroxymethyl)-5-(trifluoromethyl)pyridin-3-yl)pivalamide

To a clear light yellow solution of methyl 3-pivalamido-5-(trifluoromethyl)picolinate (2.45 g, 8.05 mmol) in tetrahydrofuran (60 mL) was added a solution of calcium chloride (1.79 g, 16.1 mmol) in ethanol (60 mL), then sodium borohydride (914 mg, 24.2 mmol) was added in three portions over a period of 30 min. The white suspension was stirred for 90 min at room temperature, then partitioned between water and sat. aq. ammonium chloride solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane–ethyl acetate gradient) afforded the title compound (1.97 g; 89%). Light yellow viscous oil, MS: 277.1 (M+H)⁺.

Intermediate 9

3-Fluoro-4-(trans-3a-fluorooctahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzenesulfonamide hydrochloride

Step 1: trans-tert-butyl 3a-fluoro-5-(2-fluoro-4-sulfamoylbenzoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

The title compound was produced in analogy to intermediate 6, step 1, replacing trans-tert-butyl 3a-methoxyhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate by trans-tert-butyl 3a-fluorohexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate and 2-cyclopropyl-6-((tetrahydro-2H-pyran-4-yl)methoxy)isonicotinic acid by 2-fluoro-4-sulfamoylbenzoic acid (CAS-RN 714968-42-0). Light yellow foam, MS: 432.2 (M+H)⁺.

Step 2: 3-Fluoro-4-(trans-3a-fluorooctahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzenesulfonamide hydrochloride

The title compound was produced in analogy to intermediate 6, step 2 from trans-tert-butyl 3a-fluoro-5-(2-fluoro-4-sulfamoylbenzoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate. White solid, MS: 332.0 (M+H)⁺.

Example A

A compound of formula (I) can be used in a manner known per se as the active ingredient for the production of tablets of the following composition:

Per tablet

	Active ingredient	200 mg
5	Microcrystalline cellulose	155 mg
	Corn starch	25 mg
	Talc	25 mg
	Hydroxypropylmethylcellulose	<u>20 mg</u>
		425 mg

10

Example B


A compound of formula (I) can be used in a manner known per se as the active ingredient for the production of capsules of the following composition:

Per capsule

15	Active ingredient	100.0 mg
	Corn starch	20.0 mg
	Lactose	95.0 mg
	Talc	4.5 mg
	Magnesium stearate	<u>0.5 mg</u>
20		220.0 mg

Claims

1. Compounds of formula (I)



wherein

- 5 R¹ is substituted phenyl, substituted phenyl-C₁₋₆-alkyl, substituted phenoxy-C₁₋₆-alkyl, substituted phenyl-C₂₋₆-alkenyl, substituted phenyl-C₂₋₆-alkynyl, substituted pyridinyl, substituted pyridinyl-C₁₋₆-alkyl, substituted pyridinyl-C₂₋₆-alkenyl, substituted pyridinyl-C₂₋₆-alkynyl, substituted thiophenyl, substituted thiophenyl-C₁₋₆-alkyl, substituted thiophenyl-C₂₋₆-alkenyl or substituted thiophenyl-C₂₋₆-alkynyl, wherein
- 10 substituted phenyl, substituted phenyl-C₁₋₆-alkyl, substituted phenoxy-C₁₋₆-alkyl, substituted phenyl-C₂₋₆-alkenyl, substituted phenyl-C₂₋₆-alkynyl, substituted pyridinyl, substituted pyridinyl-C₁₋₆-alkyl, substituted pyridinyl-C₂₋₆-alkenyl, substituted pyridinyl-C₂₋₆-alkynyl, substituted thiophenyl, substituted thiophenyl-C₁₋₆-alkyl, substituted thiophenyl-C₂₋₆-alkenyl and substituted thiophenyl-C₂₋₆-alkynyl are
- 15 substituted by R³, R⁴ and R⁵;

Y is -OC(O)- or -C(O)-;

W is -C(O)-, -S(O)₂- or -CR⁶R⁷-;

- R² is substituted phenyl, substituted pyridinyl or substituted thiophenyl, wherein substituted phenyl, substituted pyridinyl and substituted thiophenyl are substituted by
- 20 R⁶, R⁷ and R⁸;

- R³ is halogen, hydroxy, cyano, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-alkoxy-C₁₋₆-alkyl, halo-C₁₋₆-alkoxy, halo-C₁₋₆-alkyl, hydroxy-C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₃₋₈-cycloalkyl-C₁₋₆-alkyl, C₃₋₈-cycloalkyl-C₁₋₆-alkoxy, C₃₋₈-cycloalkoxy, C₃₋₈-cycloalkoxy-C₁₋₆-alkyl, C₁₋₆-alkylamino, C₁₋₆-alkylcarbonylamino, C₃₋₈-cycloalkylcarbonylamino, C₁₋₆-alkyltetrazolyl, C₁₋₆-alkyltetrazolyl-C₁₋₆-alkyl or heterocycloalkyl-C₁₋₆-alkoxy;
- 25

R^4 and R^5 are independently selected from H, halogen, hydroxy, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, halo- C_{1-6} -alkoxy, halo- C_{1-6} -alkyl, hydroxy- C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkoxy, C_{3-8} -cycloalkoxy, C_{3-8} -cycloalkoxy- C_{1-6} -alkyl, C_{1-6} -alkylcarbonylamino, C_{3-8} -cycloalkylcarbonylamino, C_{1-6} -alkyltetrazolyl, C_{1-6} -alkyltetrazolyl- C_{1-6} -alkyl or heterocycloalkyl- C_{1-6} -alkoxy;

R^6 is aminosulfonyl;

R^7 and R^8 are independently selected from H, halogen, hydroxy, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{1-6} -alkoxy- C_{1-6} -alkyl, halo- C_{1-6} -alkoxy, halo- C_{1-6} -alkyl, hydroxy- C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkyl, C_{3-8} -cycloalkyl- C_{1-6} -alkoxy, C_{3-8} -cycloalkoxy and C_{3-8} -cycloalkoxy- C_{1-6} -alkyl;

R^9 is C_{1-6} -alkyl, C_{1-6} -alkoxy or halogen;

m, n, p and q are independently selected from 1 or 2;

or pharmaceutically acceptable salts.

2. A compound according to claim 1, wherein

R^1 is substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted pyridinyl or substituted pyridinyl- C_{1-6} -alkyl, wherein substituted phenyl- C_{1-6} -alkyl, substituted phenoxy- C_{1-6} -alkyl, substituted phenyl- C_{2-6} -alkenyl, substituted pyridinyl and substituted pyridinyl- C_{1-6} -alkyl are substituted by R^3 , R^4 and R^5 ;

Y is a -OC(O)- or -C(O)-;

W is -C(O)-;

R^2 is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R^6 , R^7 and R^8 ;

R^3 is halo- C_{1-6} -alkoxy, C_{1-6} -alkylcarbonylamino or tetrahydropyranyl- C_{1-6} -alkoxy;

R^4 is H, C_{1-6} -alkyl or C_{3-8} -cycloalkyl;

R^5 is H;

R⁶ is aminosulfonyl;

R⁷ and R⁸ are independently selected from H and halogen;


R⁹ is C₁₋₆-alkyl, C₁₋₆-alkoxy or halogen;

m,n, p and q are 1;

5 or pharmaceutically acceptable salts.

3. A compound according to claim 1 or 2, wherein R¹ is substituted phenyl-C₁₋₆-alkyl, substituted phenoxy-C₁₋₆-alkyl, substituted phenyl-C₂₋₆-alkenyl, substituted pyridinyl or substituted pyridinyl-C₁₋₆-alkyl, wherein substituted phenyl-C₁₋₆-alkyl, substituted phenoxy-C₁₋₆-alkyl, substituted phenyl-C₂₋₆-alkenyl, substituted pyridinyl and substituted pyridinyl-C₁₋₆-alkyl are substituted by R³, R⁴ and R⁵.
10
4. A compound according to any one of claims 1 to 3, wherein R¹ is substituted pyridinyl or substituted pyridinyl-C₁₋₆-alkyl, wherein substituted pyridinyl and substituted pyridinyl-C₁₋₆-alkyl are substituted by R³, R⁴ and R⁵.
5. A compound according to any one of claims 1 to 4, wherein Y is -C(O)-.
- 15 6. A compound according to any one of claims 1 to 2, wherein R² is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R⁶, R⁷ and R⁸.
7. A compound according to any one of claims 1 to 6, wherein R³ is halo-C₁₋₆-alkoxy, C₁₋₆-alkylcarbonylamino or tetrahydropyranyl-C₁₋₆-alkoxy.
- 20 8. A compound according to any one of claims 1 to 7, wherein R³ is tetrahydropyranyl-C₁₋₆-alkoxy.
9. A compound according to any one of claims 1 to 8, wherein R⁴ is H, C₁₋₆-alkyl or C₃₋₈-cycloalkyl.
10. A compound according to any one of claims 1 to 9, wherein R⁴ is C₃₋₈-cycloalkyl.
- 25 11. A compound according to any one of claims 1 to 10, wherein R⁵ is H.

12. A compound according to any one of claims 1 to 11, wherein R^7 and R^8 are independently selected from H and halogen.
13. A compound according to any one of claims 1 to 12, wherein R^7 is H or halogen.
14. A compound according to any one of claims 1 to 13, wherein R^8 is H.
- 5 15. A compound according to any one of claims 1 to 13, wherein R^9 is halogen or C_{1-6} -alkoxy.
16. A compound according to any one of claims 1 to 15, wherein m, n, p and q are 1.
17. A compound according to any one of claims 1 to 16, wherein
 - R^1 is substituted pyridinyl or substituted pyridinyl- C_{1-6} -alkyl, wherein substituted pyridinyl and substituted pyridinyl- C_{1-6} -alkyl are substituted by R^3 , R^4 and R^5 ;
 - 10 Y is -C(O)-;
 - W is -C(O)-;
 - R^2 is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R^6 , R^7 and R^8 ;
 - R^3 is tetrahydropyranyl- C_{1-6} -alkoxy;
 - 15 R^4 is C_{3-8} -cycloalkyl;
 - R^5 is H;
 - R^6 is aminosulfonyl;
 - R^7 is H or halogen;
 - R^8 is H;
 - 20 R^9 is halogen or C_{1-6} -alkoxy;
 - m, n, p and q are 1
 - or pharmaceutically acceptable salts.
18. A compound according to claim 1 and of formula I(a),



wherein

R¹ is substituted phenyl-C₁₋₆-alkyl, substituted phenoxy-C₁₋₆-alkyl, substituted phenyl-C₂₋₆-alkenyl, substituted pyridinyl or substituted pyridinyl-C₁₋₆-alkyl, wherein substituted phenyl-C₁₋₆-alkyl, substituted phenoxy-C₁₋₆-alkyl, substituted phenyl-C₂₋₆-alkenyl, substituted pyridinyl and substituted pyridinyl-C₁₋₆-alkyl are substituted by R³, R⁴ and R⁵;

Y is a -OC(O)- or -C(O)-;

W is -C(O)-;

R² is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R⁶, R⁷ and R⁸;

R³ is halo-C₁₋₆-alkoxy, C₁₋₆-alkylcarbonylamino or tetrahydropyranyl-C₁₋₆-alkoxy;

R⁴ is H, C₁₋₆-alkyl or C₃₋₈-cycloalkyl;

R⁵ is H;


R⁷ and R⁸ are independently selected from H and halogen;

R⁹ is C₁₋₆-alkyl, C₁₋₆-alkoxy or halogen;

m, n, p and q are 1;

or pharmaceutically acceptable salts.

19. A compound according to claim 1 and of formula I(b),



wherein

R^1 is substituted pyridinyl or substituted pyridinyl- C_{1-6} -alkyl, wherein substituted pyridinyl and substituted pyridinyl- C_{1-6} -alkyl are substituted by R^3 , R^4 and R^5 ;

5 Y is $-C(O)-$;

W is $-C(O)-$;

R^2 is substituted phenyl or substituted pyridinyl, wherein substituted phenyl and substituted pyridinyl are substituted by R^6 , R^7 and R^8 ;

R^3 is tetrahydropyranyl- C_{1-6} -alkoxy;

10 R^4 is C_{3-8} -cycloalkyl;

R^5 is H;

R^7 is H or halogen;

R^8 is H;

R^9 is halogen or C_{1-6} -alkoxy;

15 m, n, p and q are 1

or pharmaceutically acceptable salts.

20. A compound according to any one of claims 1 to 19, selected from

trans-5-(2-fluoro-4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester;

20

trans-4-[3a-methoxy-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide;

trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide;

5 trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester;

trans-5-(4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

10 trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

trans-3a-methyl-5-(4-sulfamoyl-benzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

15 trans-5-(5-sulfamoylpyridine-2-carbonyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester;

trans-6-[3a-methoxy-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]pyridine-3-sulfonamide;

trans-6-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]pyridine-3-sulfonamide;

20 trans-5-(2-fluoro-4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

trans-4-[3a-fluoro-2-[(E)-3-[4-(trifluoromethoxy)phenyl]prop-2-enoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide;

25 trans-4-[3a-fluoro-2-[2-[4-(trifluoromethoxy)phenoxy]acetyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide;

trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-fluoro-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide;

trans-4-[3a-fluoro-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl]-3-fluorobenzenesulfonamide;

(+)-(3aR,6aR)-3a-methyl-5-(4-sulfamoyl-benzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

5 (-)-trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester ;

(-)-trans-5-(4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester ;

10 (-)-trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester ;

(-)-(3aS,6aS)-3a-methyl-5-(4-sulfamoylbenzoyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

15 (+)-trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(+)-trans-5-(4-sulfamoyl-benzoyl)-3a-fluoro-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester ;

20 (+)-trans-5-(4-sulfamoyl-benzoyl)-3a-methoxy-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2-dimethyl-propionylamino)-5-trifluoromethyl-pyridin-2-ylmethyl ester;

and pharmaceutically acceptable salts thereof.

21. A compound according to any one of claims 1 to 20, selected from


trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide;

25 trans-6-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-methoxy-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]pyridine-3-sulfonamide;

trans-4-[5-[2-cyclopropyl-6-(oxan-4-ylmethoxy)pyridine-4-carbonyl]-3a-fluoro-3,4,6,6a-tetrahydro-1H-pyrrolo[3,4-c]pyrrole-2-carbonyl]-3-fluorobenzenesulfonamide;

and pharmaceutically acceptable salts thereof.

22. A process to prepare a compound according to any one of claims 1 to 21 comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III), wherein R^1 , R^2 , R^9 , m, n, p and q are as defined in any one of claim 1 to 21 and W is -C(O)-.



23. A compound according to any one of claims 1 to 21 for use as therapeutically active substance.
24. A pharmaceutical composition comprising a compound according to any one of claims 1 to 21 and a therapeutically inert carrier.
25. A compound according to any one of claims 1 to 21 for the treatment or prophylaxis of ocular conditions.
26. The use of a compound according to any one of claims 1 to 21 for the preparation of a medicament for the treatment or prophylaxis of ocular conditions.
