

Specification

Title

PHARMACEUTICAL COMPOSITION FOR PREVENTING OR TREATING NONALCOHOLIC FATTY LIVER DISEASE, CONTAINING GPR119 LIGAND AS ACTIVE INGREDIENT

Technical Field

The present invention relates to a pharmaceutical composition, a food composition, and a feed composition for preventing or treating non-alcoholic fatty liver disease, comprising G protein coupled receptor 119 (GPR119) ligands as an effective ingredient; a method for treating and ameliorating non-alcoholic fatty liver disease; or a use thereof in preventing or treating non-alcoholic fatty liver disease.

Background

Fatty liver is a pathological condition in which triglycerides are excessively accumulated in hepatocytes, and medically defined as a condition in which triglycerides account for 5% or more of a liver weight. Fatty liver is classified into alcoholic and non-alcoholic fatty livers depending on whether fatty liver is caused by excessive alcohol intake or not. Non-alcoholic fatty liver disease (NAFLD) is a group of diseases that encompass all aspects of the disease ranging from non-alcoholic fatty liver to steatohepatitis and cirrhosis. Simple non-alcoholic fatty liver, which has only fat deposits in the liver and shows an increase in fat deposits in liver tissues caused by insulin resistance, etc. without any findings of damage to hepatocytes and fibrosis, progresses into non-alcoholic steatohepatitis (NASH), which is accompanied by damage to hepatocytes caused by inflammatory responses due to oxidative stress and sometimes fibrosis, and may progress into liver cirrhosis, which is accompanied by irreversible liver damage if not treated properly.

Recently, a prevalence rate of non-alcoholic fatty liver disease has rapidly

increased worldwide, and a prevalence rate of simple fatty liver is estimated to be 6.3 to 33%, and a prevalence rate of non-alcoholic steatohepatitis accompanied by inflammation is reported to be 3 to 5% (Hepatology, 2012(55): 2005-2023). Simple fatty liver very slowly progresses to serious liver disease and does not increase a mortality rate associated with liver disease, but steatohepatitis may lead to cirrhosis and liver cancer and increases a mortality rate associated with liver disease as well as an overall mortality. In addition, simple fatty liver may be easily ameliorated if a weight loss is made by 3 to 5% through revision of general living habits, but it is recommended to make a weight loss by at least 10% for reduction of inflammation. However, it is not clear whether or not this weight loss may induce inflammation and fibrosis to be ameliorated in the liver (Korean J Gastroenterol, 2012(60): 64-66).

So far, there has not been a therapeutic agent for non-alcoholic steatohepatitis. If the disease worsens, a liver transplantation is performed. According to U.S. statistical data, hepatitis C has been considered as the most common cause of liver transplantation, followed by non-alcoholic steatohepatitis as of 2013. In addition, an increase rate of liver transplantation for 10 years from 2004 was 14% for hepatitis C, while 170% for non-alcoholic steatohepatitis showing rapid increase and NASH is also expected to become the first cause of liver transplantation since 2020, leaving hepatitis C behind, and thus there is an urgent need to develop an effective therapeutic agent for non-alcoholic fatty liver disease including non-alcoholic steatohepatitis.

Currently, a drug therapy for non-alcoholic steatohepatitis includes a method of taking in an insulin resistance enhancer (e.g., pioglitazone) or an antioxidant (e.g., vitamin E), in which findings of fat accumulation and inflammation amelioration have been confirmed through small-scale clinical trials in patients with non-alcoholic steatohepatitis. However, such method has failed to show a clear effect of ameliorating inflammation and fibrosis, and it is still impossible to provide those drugs as an

intensive therapeutic option due to safety concerns about a long-term use.

G protein-coupled receptor 119 (GPR119) is distributed in L and K cells of the small intestine. If GPR119 is activated by triglyceride metabolites (2-monoacylglycerol), etc., GPR119 increases secretion of incretin hormones such as glucagon-like peptide-1 (GLP-1) or glucose-dependent insulintropic polypeptide (GIP). GPR119 is also distributed in pancreatic beta cells and alpha cells and contributes to maintaining blood sugar levels by controlling secretion of insulin and glucagon in a glucose-dependent manner. In addition, it is reported that, if GPR119 is overexpressed in a human macrophage line, an expression of GLP-1 receptor is increased to raise ATP-binding cassette protein A1 (ABCA1), which is a transporter of taking cholesterol out of cells, thereby increasing an ApoA1-mediated reflux of cholesterol, improving lipid metabolism, and decreasing a concentration of inflammatory cytokines in blood (J Lipid Res, 2014(55): 681-697).

However, it has not yet been reported if GPR119 ligand directly inhibits an inflammatory response and thus inhibits the inflammatory response, which is the core of hepatocyte damage in non-alcoholic fatty liver disease. Also, it has not been reported if GPR119 ligand ameliorates inflammation and fibrosis after the induction of non-alcoholic fatty liver disease.

Detailed Description of the Invention

Technical Problem


An object of the present invention is to provide a pharmaceutical composition for preventing or treating non-alcoholic fatty liver disease comprising GPR119 ligand as an effective ingredient.

Technical Solution

The present invention may provide a pharmaceutical composition for preventing or treating non-alcoholic fatty liver disease, which comprises a compound represented

by a following chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof as an effective ingredient:

[Chemical Formula 1]



The pharmaceutical composition according to the present invention may be useful in preventing or treating non-alcoholic fatty liver disease by remarkably inhibiting triglyceride deposition, inflammation and fibrosis in liver tissues.

In the chemical formula 1,

A is oxadiazole, dihydrooxazole, thiazole, or thiadiazole, the A is independently unsubstituted or substituted with at least one substituent selected from the group consisting of halogen, C1-C6 straight or branched-chain alkyl and C1-C6 alcohol, and the alkyl or alcohol group is independently unsubstituted or substituted with halogen or C1-C6 alkoxy group;


B is pyridine, pyrimidine, pyrazine, or oxadiazole, the B is independently unsubstituted or substituted with at least one substituent selected from the group consisting of halogen, C1-C6 straight or branched-chain alkyl, C1-C6 alcohol, C1-C6 alkoxy and oxadiazole group, and the C1-C6 straight or branched-chain alkyl, C1-C6 alcohol, C1-C6 alkoxy or oxadiazole group is independently unsubstituted or substituted with halogen, C1-C6 alkyl or C1-C6 alkoxy group; and

X is each independently F, Cl, Br or I, but not limited thereto.

According to one embodiment of the present invention, in the chemical formula

1,


it may be provided that



R1 to R3, R5 and R6 are each independently one or more substituents selected from the group consisting of hydrogen, halogen, C1-C6 straight or branched-chain alkyl and C1-C6 alcohol, and the alkyl or alcohol group is independently unsubstituted or substituted with halogen or C1-C6 alkoxy group.

According to another embodiment of the present invention, in the chemical formula 1,

it may be provided that



the R7 to R11 are independently substituted with at least one substituent selected from the group consisting of hydrogen, halogen, C1-C6 straight or branched-chain alkyl, C1-C6 alcohol, C1-C6 alkoxy and oxadiazole group, and the alkyl, alcohol, alkoxy or oxadiazole group is independently unsubstituted or substituted with halogen, C1-C6 alkyl or C1-C6 alkoxy group.

According to one embodiment of the present invention, in the chemical formula 1, it may be provided that A is oxadiazole substituted with C1-C6 straight or branched-chain alkyl, B is pyrimidine substituted with C1-C6 straight or branched-chain alkyl, and X is F.

In the present invention, the term "halogen" may refer to fluorine, chlorine, bromine or iodine.

In the present invention, the term "alkyl" may refer to a straight-chain or branched-chain hydrocarbon residue unless otherwise noted. An example of the C1-C6 alkyl may include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, hexyl, etc.

In the present invention, the term "alkoxy" may include an alkyl-oxygen radical having alkyl as defined above unless otherwise noted. An example of the C1-C6 alkoxy may include methoxy, ethoxy, propoxy, butoxy, pentoxy, etc.

In the present invention, the term "heterocycle" or "heterocyclic" may refer to 5 to 13-membered hetero-aromatic or non-aromatic compound including one to three heteroatoms selected from the group consisting of N, O, and S unless otherwise noted.

In the present invention, the compound represented by the chemical formula 1 may be a compound specifically selected from the group consisting of the following compounds:

2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-4,5-dihydrooxazole,

(R)-2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluoroph

enyl)-4-methyl-4,5-dihydrooxazole,

(S)-2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-4-methyl-4,5-dihydrooxazole,

(S)-2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-methyl-4,5-dihydrooxazole,

(R)-2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-methyl-4,5-dihydrooxazole,

2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5,5-dimethyl-4,5-dihydrooxazole,

(R)-(2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-4,5-dihydrooxazole-5-yl)methanol,

(S)-(2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-4,5-dihydrooxazole-5-yl)methanol,

(R)-3-(2-(4-(3-(3,5-difluoro-4-(5-methyl-4,5-dihydrooxazole-2-yl)phenoxy)propyl)piperidine-1-yl)pyrimidine-5-yl)-5-isobutyl-1,2,4-oxadiazole,

(R)-5-(4-(3-(3,5-difluoro-4-(4-methyl-4,5-dihydrooxazole-2-yl)phenoxy)propyl)piperidine-1-yl)-3-isopropyl-1,2,4-oxadiazole,

(S)-5-(4-(3-(3,5-difluoro-4-(5-methyl-4,5-dihydrooxazole-2-yl)phenoxy)propyl)piperidine-1-yl)-3-isopropyl-1,2,4-oxadiazole,

5-(4-(3-(4-(5,5-dimethyl-4,5-dihydrooxazole-2-yl)-3,5-difluorophenoxy)propyl)piperidine-1-yl)-3-isopropyl-1,2,4-oxadiazole,

3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-methyl-1,2,4-oxadiazole,

3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-propyl-1,2,4-oxadiazole,

3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)

-5-isopropyl-1,2,4-oxadiazole,

5-(tert-butyl)-3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-1,2,4-oxadiazole,

(3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-1,2,4-oxadiazole-5-yl)methanol,

2-(3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-1,2,4-oxadiazole-5-yl)ethane-1-ol,

(S)-1-(3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-1,2,4-oxadiazole-5-yl)propane-1-ol,

(R)-1-(3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-1,2,4-oxadiazole-5-yl)propane-2-ol,

(S)-1-(3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-1,2,4-oxadiazole-5-yl)propane-2-ol,

2-(3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-1,2,4-oxadiazole-5-yl)-2-methylpropane-1-ol,

3-(2,6-difluoro-4-(3-(1-(5-propylpyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-pentylpyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-(trifluoromethyl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-methoxypyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-isopropoxypyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,2,4-oxadiazole,

3-(4-(3-(1-(5-chloropyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,2,4-oxadiazole,

l)-5-isopropyl-1,2,4-oxadiazole,

3-(4-(3-(1-(5-bromopyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-(5-isobutyl-1,2,4-oxadiazole-3-yl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-methyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-(5-isobutyl-1,2,4-oxadiazole-3-yl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-ethyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-(5-isobutyl-1,2,4-oxadiazole-3-yl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,2,4-oxadiazole,

5-(sec-butyl)-3-(2,6-difluoro-4-(3-(1-(5-(5-isobutyl-1,2,4-oxadiazole-3-yl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-(5-isobutyl-1,2,4-oxadiazole-3-yl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-(methoxymethyl)-1,2,4-oxadiazole,

(S)-1-(3-(2,6-difluoro-4-(3-(1-(5-(5-isobutyl-1,2,4-oxadiazole-3-yl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-1,2,4-oxadiazole-5-yl)propane-1-ol,

2-(3-(2,6-difluoro-4-(3-(1-(5-(5-isobutyl-1,2,4-oxadiazole-3-yl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-1,2,4-oxadiazole-5-yl)-2-methylpropane-1-ol,

3-(4-(3-(1-(5-chloropyrazine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(5-(trifluoromethyl)pyridine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(3-isopropyl-1,2,4-oxadiazole-5-yl)piperidine-4-yl)propoxy)phenyl)-5-methyl-1,2,4-oxadiazole,

3-(2,6-difluoro-4-(3-(1-(3-isopropyl-1,2,4-oxadiazole-5-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,2,4-oxadiazole,

(3-(2,6-difluoro-4-(3-(1-(3-isopropyl-1,2,4-oxadiazole-5-yl)piperidine-4-yl)prop

oxy)phenyl)-1,2,4-oxadiazole-5-yl)methanol,

2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-methyl-1,3,4-oxadiazole,

2-ethyl-5-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-1,3,4-oxadiazole,

2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,3,4-oxadiazole,

5-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-N-isopropyl-1,3,4-oxadiazole-2-amine,

2-(2,6-difluoro-4-(3-(1-(5-(trifluoromethyl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-methyl-1,3,4-oxadiazole,

2-(2,6-difluoro-4-(3-(1-(5-(trifluoromethyl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-ethyl-1,3,4-oxadiazole,

2-(2,6-difluoro-4-(3-(1-(5-(trifluoromethyl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,3,4-oxadiazole,

2-(4-(3-(1-(5-chloropyrazine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-methyl-1,3,4-oxadiazole,

2-(4-(3-(1-(5-chloropyrazine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-ethyl-1,3,4-oxadiazole,

2-(4-(3-(1-(5-chloropyrazine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,3,4-oxadiazole,

5-(4-(3-(3,5-difluoro-4-(5-methyl-1,3,4-oxadiazole-2-yl)phenoxy)propyl)piperidine-1-yl)-3-propyl-1,2,4-oxadiazole,

5-(4-(3-(3,5-difluoro-4-(5-ethyl-1,3,4-oxadiazole-2-yl)phenoxy)propyl)piperidine-1-yl)-3-propyl-1,2,4-oxadiazole,

5-(4-(3-(3,5-difluoro-4-(5-isopropyl-1,3,4-oxadiazole-2-yl)phenoxy)propyl)piper

idine-1-yl)-3-propyl-1,2,4-oxadiazole,

5-(4-(3-(3,5-difluoro-4-(5-methyl-1,3,4-oxadiazole-2-yl)phenoxy)propyl)piperidine-1-yl)-3-isopropyl-1,2,4-oxadiazole,

5-(4-(3-(4-(5-ethyl-1,3,4-oxadiazole-2-yl)-3,5-difluorophenoxy)propyl)piperidine-1-yl)-3-isopropyl-1,2,4-oxadiazole,

5-(4-(3-(3,5-difluoro-4-(5-isopropyl-1,3,4-oxadiazole-2-yl)phenoxy)propyl)piperidine-1-yl)-3-isopropyl-1,2,4-oxadiazole,

5-(4-(3-(3,5-difluoro-4-(5-methyl-1,3,4-oxadiazole-2-yl)phenoxy)propyl)piperidine-1-yl)-3-(2,2,2-trifluoroethyl)-1,2,4-oxadiazole,

3-(4-(3-(3,5-difluoro-4-(5-isopropyl-1,3,4-oxadiazole-2-yl)phenoxy)propyl)piperidine-1-yl)-5-isopropyl-1,2,4-oxadiazole,

2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,3,4-thiadiazole,

2-(2,6-difluoro-4-(3-(1-(5-propylpyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,3,4-thiadiazole,

2-(2,6-difluoro-4-(3-(1-(5-pentylpyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,3,4-thiadiazole,

2-(2,6-difluoro-4-(3-(1-(5-fluoropyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,3,4-thiadiazole,

2-(2,6-difluoro-4-(3-(1-(5-(trifluoromethyl)pyrimidine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,3,4-thiadiazole,

2-(2,6-difluoro-4-(3-(1-(5-(trifluoromethyl)pyridine-2-yl)piperidine-4-yl)propoxy)phenyl)-5-isopropyl-1,3,4-thiadiazole, and

4-ethyl-2-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)thiazole.

In an embodiment of the present invention, the compound represented by the

chemical formula 1 of the present invention may be specifically 3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,2,4-oxadiazole.

In the present invention, a non-limiting example of the pharmaceutically acceptable salt of the compound represented by the chemical formula 1 may include inorganic acid salts such as hydrochloric acid, bromic acid, phosphoric acid or sulfuric acid; organic carboxylic acid salts such as acetic acid, trifluoroacetic acid, citric acid, maleic acid, oxalic acid, succinic acid, benzoic acid, tartaric acid, fumaric acid, mandelic acid, ascorbic acid or malic acid, or sulfonic acid salts such as methanesulfonic acid or para-toluenesulfonic acid; alkali metal salts such as sodium, potassium or lithium; various acid salts known to be capable of forming other pharmaceutically acceptable salts; or the like.

In one specific embodiment of the present invention, it was confirmed through an experiment of conducting a histological examination, measuring the concentration of ALT and AST in blood, measuring the expression level of genes and proteins associated with inflammation and fibrosis and the like that damage to liver tissues, deposition of triglycerides in liver tissues, infiltration of inflammatory cells and fibrosis are remarkably inhibited in an ob/ob mouse model, in which non-alcoholic fatty liver disease has been induced by special diet, by administering the compound represented by the chemical formula 1 during the process of inducing the disease (Example 1).

In one specific embodiment of the present invention, it was confirmed for the first time that the compound represented by the chemical formula 1 inhibits a differentiation of human monocytes and an activation of differentiated macrophages by an inflammatory factor in a concentration-dependent manner through a mechanism for inhibiting inflammatory responses (Example 2).

In one specific embodiment of the present invention, it was confirmed that

damage to liver tissues, deposition of triglycerides in liver tissues, infiltration of inflammatory cells and fibrosis are remarkably ameliorated in a C57BL6 mouse model, in which non-alcoholic fatty liver disease has been induced by special diet, by administering the compound represented by the chemical formula 1 after inducing the disease (Example 3).

The pharmaceutical composition for preventing or treating non-alcoholic fatty liver disease according to the present invention may be used in the form of a general pharmaceutical preparation. The pharmaceutical preparation may be administered in various oral and parenteral dosage forms upon administration, and the dosage form may be variously determined according to the method of use.

If the pharmaceutical preparation is formulated into various oral and parenteral dosage forms, it may be possible to use the generally used excipients such as fillers, diluents, extenders, binders, humectants, disintegrants, surfactants, etc.

A solid preparation for oral administration may include tablets, pills, powders, granules, capsules, etc., and this solid preparation may be prepared by mixing at least one excipient, for example, starch, calcium carbonate, sucrose, lactose, gelatin or the like in the pharmaceutical composition. In addition, lubricants such as magnesium stearate, talc, etc., may be used in addition to simple excipients.

Furthermore, a liquid preparation for oral administration may include suspending agents, liquids for internal use, emulsions, syrups, etc., but may also include various excipients, for example, humectants, sweetening agents, flavoring agents, preservatives, etc. in addition to water and liquid paraffin, which are the frequently used simple diluents.

A preparation for parenteral administration may include sterile aqueous solutions, non-aqueous solvents, suspensions, emulsions, lyophilized preparations and suppositories. The non-aqueous solvent and the suspending agent may include

propylene glycol, polyethylene glycol, vegetable oil like olive oil, injectable ester like ethyl oleate, etc. A base of the suppository may include witepsol, macrogol, tween 61, cacao butter, laurinum, glycerogelatin, etc.

In addition, the pharmaceutical composition for preventing or treating non-alcoholic fatty liver disease according to the present invention may represent an effective amount in an administration range of about 1 to about 1,000 mg. An administered amount or an intake amount may be administered in a variety of administered doses and methods, such as once a day or several times a day by dividing the composition depending on the subject's body weight, age, gender, health condition, diet, administration time, administration method, excretion rate, and severity of the disease.

In the present invention, non-alcoholic fatty liver disease (NAFLD) may include both primary and secondary non-alcoholic fatty liver diseases. Specifically, in the present invention, non-alcoholic fatty liver disease (NAFLD) may include simple steatosis, non-alcoholic steatohepatitis (NASH) and liver fibrosis and liver cirrhosis caused by the progression of these diseases, but is not limited to.

The pharmaceutical composition of the present invention may comprise the compound represented by the chemical formula 1 or one or more effective ingredients which show a similar function thereto.

The present invention may also provide a method for preventing or treating non-alcoholic fatty liver disease, which includes administering a therapeutically effective amount of the compound represented by the chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof into a subject in need of treatment.

In the present invention, the term “subject in need of treatment” may refer to mammals including humans, and the term “administration” may refer to providing a

predetermined material to subjects by any appropriate method. The term “therapeutically effective amount” may refer to an amount of an active ingredient or a pharmaceutical composition which induces animals or humans to show the biological or medical responses considered by investigators, veterinarians, doctors or other clinicians, and such amount may include an amount thereof for inducing a relief of diseases or disorders to be treated. It is apparent to those skilled in the art that the therapeutically effective dosage and the number of administration for effective ingredients of the present invention may vary depending on a desired effect.

In the present invention, a route of administration of the pharmaceutical composition of the present invention may be administered through any general route as long as it can reach a target tissue.

Administration may be performed orally, intraperitoneally, intravenously, intramuscularly, subcutaneously, endothelially, intranasally, intrapulmonarily, rectally, intracavitarily, intraperitoneally and intrathecally, but is not limited thereto.

The pharmaceutical composition of the present invention may be administered once a day or at least twice a day at a certain interval of time.

The pharmaceutical composition of the present invention may be used alone or in combination with a surgery, endocrinotherapy, drug treatment and methods of using a biologic response modifier, in order to prevent and treat non-alcoholic fatty liver disease.

The present invention may also provide a food composition for preventing or ameliorating non-alcoholic fatty liver disease, which comprises the compound represented by the chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof as an effective ingredient.

In the present invention, the term “alleviation” may refer to all the acts, in which

a disease gets better or takes a favorable turn by administering the composition.

In the present invention, the term "food" may refer to meats, sausages, breads, chocolates, candies, snacks, confectioneries, pizzas, instant noodles, other noodles, chewing gums, dairy products including ice creams, various types of soup, beverages, teas, health drinks, alcohol beverages, vitamin complexes, health functional foods, health foods, health supplement foods and the like, and include all the foods in a conventional sense.

The term "health functional food" may be the same term as food for special health use (FoSHU), and refer to the food having a high medical and medicinal effect, which is processed to efficiently show a biological regulation function in addition to supplying nutrients. In this case, the term "function(ality)" may refer to controlling nutrients or obtaining a beneficial effect on health such as a physiological action, etc., with regard to the structure and function of the human body.

The "health food" may refer to food having an effect of actively maintaining or enhancing health compared to general food, and "health supplement food" may refer to food for a health supplement purpose. In some cases, the terms of health functional food, health food, and health supplement food may be used interchangeably.

The food of the present invention may be prepared by a method conventionally used in the art, and raw materials and ingredients conventionally added in the art may be added to prepare the food during the preparation. Specifically, proteins, carbohydrates, fats, nutrients, seasoning agents, and flavoring agents may be included, and an example of the carbohydrates may include glucose, fructose, maltose, sucrose, oligosaccharide, dextrin, cyclodextrin, xylitol, sorbitol, erythrol, saccharin or synthetic flavoring agents, but is not limited thereto. The food composition of the present invention may be prepared into various dosage forms without limitation, as long as the dosage form is recognized as food.

The present invention may also provide a method for preventing or ameliorating non-alcoholic fatty liver disease, which includes administering the food composition comprising the compound represented by the chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrate or solvates thereof, or mixtures thereof as an effective ingredient into a subject in need of amelioration.

The present invention may also provide a feed composition for preventing or ameliorating non-alcoholic fatty liver disease, which comprises the compound represented by the chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof as an effective ingredient.

In the present invention, the term "feed" may refer to any natural or artificial diet, one meal, etc., or an ingredient of the one meal, which is to be consumed or digested by livestock or appropriate thereto. The feed may comprise feed additives or auxiliary feeds.

The kind of the feed is not particularly limited, and the feed conventionally used in the art may be used. A non-limiting example of the feed may include vegetable feeds such as grains, root fruits, food processing by-products, algae, fibers, pharmaceutical by-products, oils and fats, starches, oil meals, grain by-products or the like; animal feeds such as proteins, inorganic matters, oils and fats, minerals, single-cell proteins, zooplanktons, foods or the like. The feed may be used alone or by mixing at least two thereof.

The present invention may also provide a use of the compound represented by the chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof in preventing or treating non-alcoholic fatty liver disease.

The present invention may also provide a use of the compound represented by

the chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof in preparing a pharmaceutical preparation for preventing or treating non-alcoholic fatty liver disease.

In the therapeutic method, the food composition, the alleviation method, the feed composition and the use thereof, the compound represented by the chemical formula 1 may be specifically 3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,2,4-oxadiazole.

Matters mentioned in the pharmaceutical composition, the therapeutic method, the food composition, the alleviation method, the feed composition and the use thereof according to the present invention are applied the same, if not contradictory to each other.

Advantageous Effects

A pharmaceutical composition according to the present invention shows an excellent effect of alleviating lipid metabolism, reducing fat accumulation in liver tissues, and preventing histological damage caused by inflammation and fibrosis of liver tissues, and thus can be useful in preventing or treating non-alcoholic fatty liver disease.

Brief Description of the Drawings

FIG. 1 is a picture of a tissue sample showing an effect of reducing triglycerides and inhibiting infiltration of inflammatory cells by the compound of the present invention in a mouse model with induced non-alcoholic steatohepatitis.

FIG. 2 is an analysis graph showing an effect of inhibiting infiltration of inflammatory cells and inhibiting fibrosis by the compound of the present invention in a mouse model with induced non-alcoholic steatohepatitis (#, $p < 0.05$ vs. Normal; * & **, $p < 0.05$ & $p < 0.01$ vs. Ob-NASH).

FIG. 3 is a picture of a tissue sample showing an effect of inhibiting fibrosis by

the compound of the present invention in a mouse model with induced non-alcoholic steatohepatitis.

FIG. 4 is a view showing an effect of reducing AST and ALT by the compound of the present invention in a mouse model with induced non-alcoholic steatohepatitis (#, $p < 0.05$ vs. Normal; *, $p < 0.05$ vs. Ob-NASH).

FIG. 5 is a view showing results of measuring a concentration of proteins associated with inflammation and fibrosis in a mouse model with induced non-alcoholic steatohepatitis (###, $p < 0.001$ vs. Normal; ** & ***, $p < 0.01$ & $p < 0.001$ vs. Ob-NASH).

FIG. 6 is a view showing results of measuring an expression level of genes associated with inflammation and fibrosis in a mouse model with induced non-alcoholic steatohepatitis.

FIG. 7 is a view showing results of evaluating a gene expression of GPR119 in cells at a time point before differentiating human monocytes into macrophages and after differentiation for 24 and 48 hours.

FIG. 8 is a view showing results of evaluating an effect on differentiation of human monocytes into macrophages according to treatment with the compound of the present invention.

FIG. 9 is a view showing results of evaluating an effect on activation of differentiated macrophages according to treatment with the compound of the present invention.

FIG. 10 is a picture of a tissue sample showing an effect of reducing triglycerides and inhibiting infiltration of inflammatory cells in liver tissues by the compound of the present invention in a mouse model with induced non-alcoholic steatohepatitis.

FIG. 11 is a view showing results of measuring a content of triglycerides in liver tissues in a mouse model with induced non-alcoholic steatohepatitis (** & ***, $p < 0.01$ & $p < 0.001$ vs. DIO-NASH).

FIG. 12 is a view showing a change in NAFLD activity score (NAS) before and after administration into the same individual in a mouse model with induced non-alcoholic steatohepatitis.

FIG. 13 is a picture of a tissue sample showing an effect of inhibiting fibrosis by the compound of the present invention in a mouse model with induced non-alcoholic steatohepatitis.

FIG. 14 is a view showing results of measuring a content of type I collagen in liver tissues in a mouse model with induced non-alcoholic steatohepatitis (* & ***, $p < 0.05$ & $p < 0.001$ vs. DIO-NASH).

FIG. 15 is a view showing results of measuring a concentration of AST and ALT in blood in a mouse model with induced non-alcoholic steatohepatitis (** & ***, $p < 0.01$ & $p < 0.001$ vs. DIO-NASH).

FIG. 16 is a view showing results of measuring a gene expression level of monocyte attractant, macrophage markers and fibrosis markers in liver tissues of a mouse model with induced non-alcoholic steatohepatitis (*, ** & ***, $p < 0.05$, $p < 0.01$ & $p < 0.001$ vs. DIO-NASH).

Mode for Invention

The features and advantages of the present invention as well as methods for achieving them will be apparent with reference to exemplary embodiments described in detail hereinafter. However, the present invention is not limited to the Examples disclosed hereinafter, but will be implemented in various different forms. Hereinafter, the following Examples will be suggested for better understanding of the present invention and are provided only for the purpose of completely illustrating the scope of the present invention to those skilled in the art, and thus the present invention will be defined only by the scope of the claims thereto.

<Example 1> Confirmation of efficacy of GPR119 ligand in mouse

model with induced steatohepatitis due to supply of special diet

In order to confirm an effect of a GPR119 ligand compound according to the present invention on preventing non-alcoholic fatty liver disease, a following experiment was performed.

Preparation of mouse model with non-alcoholic steatohepatitis

A six-week-old male ob/ob mouse with leptin deficiency was fed with a special diet containing high fat, high fructose, and high cholesterol for 10 weeks, and thus was induced to develop non-alcoholic steatohepatitis. A mixed form of diet was prepared so that a daily dose of (3-(4-(3-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,2,4-oxadiazole (hereinafter, "compound 1") reaches 100 mg/kg/day in the compound represented by the chemical formula 1, and supplied to the mouse from a time point of starting the supply of the special diet for 10 weeks.

Histological examination

For a histological examination, the prepared mouse model with induced non-alcoholic steatohepatitis was subjected to autopsy to fix an isolated liver tissue in 10% formalin and prepare a paraffin block, thereby obtaining a 2 µm-thick tissue section. To confirm infiltration of inflammatory cells, Hematoxylin and Eosin (HE) stain was performed using an automatic staining machine (Autostainer XL, Leica), and the results thereof are shown in FIGS. 1 and 2.

As understood from FIGS. 1 and 2, the ob/ob mouse (Ob-NASH) fed with the special diet for 10 weeks showed a remarkable increase in triglycerides lipid droplets in liver tissues compared to C57BL/6 mouse fed with a normal diet (Normal), and the infiltration of inflammatory cells was also confirmed. In contrast, it was confirmed that the infiltration of inflammatory cells is significantly inhibited when the compound 1 of the present invention is simultaneously supplied with the special diet (Ob-NASH +

Compd 1).

Then, for evaluation of fibrosis, fibers in liver tissue were specifically stained by using Masson's Trichrome stain or Sirius Res stain, and the results thereof are shown in FIGS. 2 and 3.

As understood from FIGS. 2 and 3, it was confirmed that the ob/ob mouse fed with the special diet for 10 weeks shows purple-colored liver tissues as a whole with a remarkable progress of fibrosis unlike the C57BL/6 mouse fed with the normal diet, but fibrosis of liver tissues is significantly inhibited when the compound 1 of the present invention is simultaneously supplied with the special diet.

AST and ALT measurement in Blood

The mouse model with induced non-alcoholic steatohepatitis was subjected to autopsy, after which plasma was separated therefrom, so as to quantify aspartate aminotransferase (AST) and alanine aminotransferase (ALT) by using an automatic blood analyzer (Konelab 20i), and the results thereof are shown in FIG. 4.

As shown in FIG. 4, it was confirmed that the ob/ob mouse fed with the special diet for 10 weeks shows a significant increase in ALT and AST levels in blood, which is a marker of liver damage, due to damage to liver tissues unlike the C57BL/6 mouse fed with the normal diet. On the other hand, it was confirmed that ob/ob mouse is administered compound 1 of the present invention together with the special diet shows a significant inhibition of increase in ALT and AST, which means that hepatocytes damage is alleviated.

Confirmation of concentration of proteins associated with inflammation and fibrosis

To evaluate an effect on the expression of inflammation-related genes and fibrosis-related proteins in liver tissues, the liver tissues obtained from the mouse model were put into RIPA buffer (Cell Signaling) and pulverized with TissueLyser II™

(Quiagen), after which a total protein concentration was quantified with Pierce BCA Protein Assay Kit (Thermo Fisher Scientific) and a protein concentration of Mouse Ccl2 (=Mcp1; R&D Systems, MJE00) and Timp1 (R&D Systems, MTM100) in the liver tissues was quantified by using a commercially available ELISA kit, so that the results thereof are shown in FIG. 5.

As shown in FIG. 5, it was confirmed that the mouse fed with the special diet shows an increase in protein concentrations of Ccl2 (=Mcp1), in the liver tissues with regard to Ccl2 (=Mcp1), which is a factor for inducing inflammatory cells to the liver tissues, and Timp1, an endogenous inhibitor for enzymes degrading fibers deposited in the liver tissues, while protein levels of Ccl2 and Timp1 are significantly inhibited when the compound 1 of the present invention is administered together with the special diet.

Confirmation of expression of genes associated with inflammation and fibrosis

To measure an expression level of inflammation-related genes and fibrosis-related genes in liver tissues, Trizol® (Invitrogen) was added to the liver tissues collected from the mouse model, after which a total RNA was extracted from the tissues according to the method provided by the manufacturer, and then cDNA was synthesized by using a reverse transcriptase. Then, a real-time polymerase chain reaction was performed in LightCycler®480 Instrument II (RocheLifeScience) equipment by using a reagent of SYBR Green I Master Mix (Roche, 04707516001), and the results thereof are shown in FIG. 6.

As shown in FIG. 6, it was confirmed that the mouse fed with the special diet for 10 weeks shows an increase in expression of inflammation-related mouse genes (*Ccl2*, *Lgals3*, *Tnfa*) and fibrosis-related genes (*Tgfb1*, *Col1a*, *Timp1*, *Timp2*, *Acta2*), while the increase in expression of genes associated with inflammation and fibrosis is significantly inhibited when the compound 1 of the present invention is administered together with the special diet.

The above results suggest that the compound of the present invention has an excellent effect on non-alcoholic fatty liver disease by inhibiting damage, inflammation and fibrosis of liver tissues.

<Example 2> Confirmation of inhibitory effect on differentiation of human monocytes and activation of differentiated macrophages

To evaluate a direct anti-inflammatory effect of the compound according to the present invention, the effect on the differentiation of human monocytes and the activation thereof was confirmed.

First, as a result of evaluating a gene expression of GPR119 in cells at a time point before differentiating human monocytes (THP-1, ATCC® TIB-202™) into macrophages and after differentiation for 24 and 48 hours by using Phorbol 12-myristate 13-acetate (PMA), it was confirmed that an expression of GPR119 receptor increases with time as monocytes are differentiated into macrophages (FIG. 7).

Then, THP-1 monocytes were treated with PMA (50 ng/ml) for 48 hours to be differentiated into macrophages, while the compound 1 was co-treated, then replaced with serum-free medium, and then treated with lipopolysaccharides (LPS, 0.5 ng/ml) for four hours so as to quantify interleukin-1 β (IL-1 β , R&D Systems, DY201) secreted into the medium by using a commercialized ELISKA kit and evaluate an effect on the differentiation of monocytes, so that the results thereof are shown in FIG. 8.

As shown in FIG. 8, it was confirmed that IL-1 β secretion caused by activation of macrophages is decreased in a concentration-dependent manner when treated with the compound of the present invention in a process of differentiating monocytes into macrophages.

In addition, after treating the differentiated macrophages with the compound 1 for 48 hours, the secreted IL-1 β was quantified by the same method as above with regard to the activation of immune cells by LPS, so as to evaluate an effect on the

activation of differentiated macrophages. Thus, the results thereof are shown in FIG. 9.

As shown in FIG. 9, it was confirmed that IL-1 β secretion caused by activation of macrophages is decreased in a concentration-dependent manner even when the differentiated macrophages are treated with the compound of the present invention.

The above results suggest that the compound of the present invention has an excellent effect on preventing and treating non-alcoholic fatty liver disease by directly inhibiting the differentiation and activation of immune cells.

<Example 3> Confirmation of efficacy of GPR119 ligand in mouse model with induced steatohepatitis due to supply of special diet

In order to confirm an effect of a GPR119 ligand compound according to the present invention on treating non-alcoholic fatty liver disease, a following experiment was performed.

Preparation of a mouse model with non-alcoholic steatohepatitis and study design

A six-week-old male C57BL/6J mouse was fed with a special diet comprising high fat, high fructose, and high cholesterol for at least 26 weeks, and thus was induced to develop non-alcoholic steatohepatitis. It was confirmed that fatty liver and inflammatory fibrosis are induced through a biopsy of liver tissues three weeks before drug supply, and mice were evenly distributed into each group based on the area of collagen stained in the liver tissues. A feed was prepared by mixing the compounds according to the present invention in a special diet and supplied for additional eight weeks. According to the diet supplied to the mice, each of the groups was classified as a control group fed with a normal diet (Normal), a positive control group in which mice with induced non-alcoholic steatohepatitis due to the special diet were not supplied with a drug (DIO-NASH), and a group in which mice with induced non-alcoholic steatohepatitis were supplied with the compound 1 in an amount of 30 mg/kg/day (L)

and 100 mg/mg/day (H) (Cmpd1 (L) and Cmpd1 (H), respectively).

Histological examination

For a histological examination, the prepared C57BL/6J mouse model was subjected to autopsy to fix an isolated liver tissue in 10% formalin and prepare a paraffin block, thereby obtaining a 2 µm-thick tissue section. Then, Hematoxylin and Eosin (HE) stain was performed using an automatic staining machine (Autostainner XL, Leica), and the results thereof are shown in FIG. 10. In addition, the liver tissue isolated from the C57BL/6J mouse model was used to measure a content of triglycerides in the liver tissue by using a Triglyceride reagent (Roche Diagnostics, #22-045-795), and the results are shown in FIG. 11.

As understood from FIGS. 10 and 11, the mice fed with a special diet only (DIO-NASH) showed a remarkable increase in fat deposits and infiltration of inflammatory cells in liver tissues compared to the mice fed with a normal diet (Normal), while the DIO-NASH_Compd1 (H) group fed with the compound 1 of the present invention together with the special diet showed a significant decrease in fat deposits and infiltration of inflammatory cells.

In addition, as a result of calculating an NAFLD activity score (NAS) reflecting a ratio of fat cells, infiltration of inflammatory cells and damage to hepatocytes based on the above histological examination, before and after administration, the disease worsened in 50% of the DIO-NASH group, while no individual was deteriorated in case of administering the compound, and the disease was ameliorated in about 50% of the DIO-NASH_Compd1 group (FIG. 12).

Then, for the evaluation of fibrosis, the fibers of the liver tissues were specifically stained with Sirius Res stain in the mice of each group, and a content of type I collagen in the liver tissues was measured by an imaging analysis method after immunostaining, so that the results thereof are shown in FIGS. 13 and 14.

As shown in FIGS. 13 and 14, the mice fed with the special diet only (DIO-NASH) showed a remarkable progress of fibrosis in liver tissues compared to the mice fed with the normal diet (Normal), while the group fed with the compound 1 of the present invention together with the special diet (DIO-NASH_Compd1) showed a remarkable decrease in a cross-linking formation in fibers of the liver tissues and also showed a significant decrease in type I collagen.

AST and ALT measurement in Blood

Each of the mouse groups was subjected to autopsy, after which plasma was separated therefrom, so as to quantify aspartate aminotransferase (AST) and alanine aminotransferase (ALT) by using an automatic blood analyzer (Konelab 20i). Thus, the results thereof are shown in FIG. 15.

As shown in FIG. 15, the group of mice fed with the special diet only (DIO-NASH) showed a significant increase in a concentration of ALT and AST in blood compared to the mice fed with the normal diet, while the group of mice fed with the compound 1 of the present invention together with the special diet (DIO-NASH_Compd1) showed a significantly inhibited increase in ALT and AST.

Confirmation of change in gene expression

To analyze a change in total gene expression in liver tissues isolated from the above mouse model in eight weeks after drug administration, the isolated RNA fraction was used to generate a library by using NeoPrep (Illumina), and subjected to RNAseq profiling through NexSeq 500 (Illumina), after which bioinformatics analysis was performed and changes in major gene groups were classified for each function, so that the results thereof are shown in FIG. 16.

As shown in FIG. 16, the mice fed with the special diet only (DIO-NASH) showed an increase in the expression of monocyte attractants, macrophage markers and fibrosis markers in liver tissues compared to the mice fed with the normal diet (Normal), while


the DIO-NASH_Compd1 group of mice fed with the compound 1 of the present invention together with the special diet showed a remarkable decrease in the above markers.

The above results suggest that the compound of the present invention has an effect on non-alcoholic fatty liver disease by inhibiting the damage, inflammation and fibrosis of liver tissues, and has a more excellent therapeutic effect by co-administering a DPPIV inhibitor.

Claims

1. A pharmaceutical composition for preventing or treating non-alcoholic fatty liver disease, comprising a compound represented by a following chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof as an effective ingredient:

[Chemical Formula 1]




wherein A is oxadiazole, dihydrooxazole, thiazole, or thiadiazole, the A is independently unsubstituted or substituted with at least one substituent selected from the group consisting of halogen, C1-C6 straight or branched-chain alkyl and C1-C6 alcohol, and the alkyl or alcohol group is independently unsubstituted or substituted with halogen or C1-C6 alkoxy group;

B is pyridine, pyrimidine, pyrazine, or oxadiazole, the B is independently unsubstituted or substituted with at least one substituent selected from the group consisting of halogen, C1-C6 straight or branched-chain alkyl, C1-C6 alcohol, C1-C6 alkoxy and oxadiazole group, and the C1-C6 straight or branched-chain alkyl, C1-C6 alcohol, C1-C6 alkoxy or oxadiazole group is independently unsubstituted or substituted with halogen, C1-C6 alkyl or C1-C6 alkoxy group; and


X is independently F, Cl, Br or I.

2. The pharmaceutical composition according to claim 1, wherein, in the above chemical formula 1, A is



R1 to R3, R5 and R6 are each independently one or more substituents selected from the group consisting of hydrogen, halogen, C1-C6 straight or branched-chain alkyl and C1-C6 alcohol, and the alkyl or alcohol group is independently unsubstituted or substituted with halogen or C1-C6 alkoxy group.

3. The pharmaceutical composition according to claim 1, wherein, in the chemical formula 1, B is



the R7 to R11 are independently substituted with one or more substituents selected from the group consisting of hydrogen, halogen, C1-C6 straight or branched-chain alkyl, C1-C6 alcohol, C1-C6 alkoxy and oxadiazole group, and the alkyl, alcohol, alkoxy or oxadiazole group is independently unsubstituted or substituted with

halogen, C1-C6 alkyl or C1-C6 alkoxy group.

4. The pharmaceutical composition according to claim 1, wherein, in the chemical formula 1, X is F.

5. The pharmaceutical composition according to claim 1, wherein, in the chemical formula 1, A is oxadiazole substituted with C1-C6 straight or branched-chain alkyl, B is pyrimidine substituted with C1-C6 straight or branched-chain alkyl, and X is F.

6. The pharmaceutical composition according to claim 1, wherein the compound represented by the chemical formula 1 is 3-(4-(3-(1-(5-ethylpyrimidine-2-yl)piperidine-4-yl)propoxy)-2,6-difluorophenyl)-5-isopropyl-1,2,4-oxadiazole.

7. The pharmaceutical composition according to claim 1, wherein the non-alcoholic fatty liver disease is selected from the group consisting of simple fatty liver, non-alcoholic steatohepatitis, liver fibrosis and cirrhosis.

8. The pharmaceutical composition according to claim 1, wherein the pharmaceutical composition inhibits deposition of triglycerides in liver tissues.


9. The pharmaceutical composition according to claim 1, wherein the pharmaceutical composition inhibits infiltration of inflammatory cells in liver tissues.

10. The pharmaceutical composition according to claim 1, wherein

the pharmaceutical composition inhibits fibrosis of liver tissues.

11. A method for preventing or treating non-alcoholic fatty liver disease, the method comprising administering a therapeutically effective amount of a compound represented by a following chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof into a subject in need of treatment:


[Chemical Formula 1]



wherein A, B and X are the same as shown in claim 1.

12. A food composition for preventing or alleviating non-alcoholic fatty liver disease, which comprises a compound represented by a following chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof as an effective ingredient:


[Chemical Formula 1]



wherein A, B, and X are the same as in claim 1.

13. A method for preventing or alleviating non-alcoholic fatty liver disease, the method comprising administering a food composition comprising a compound represented by a following chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof into a subject in need of amelioration:


[Chemical Formula 1]



wherein A, B, and X are the same as in claim 1.

14. A feed composition for preventing or alleviating non-alcoholic fatty liver disease, which comprises a compound represented by a following chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof as an effective ingredient:


[Chemical Formula 1]



wherein A, B, and X are the same as in claim 1.

15. A use of a compound represented by a following chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof in preventing or treating non-alcoholic fatty liver disease:


[Chemical Formula 1]



wherein A, B, and X are the same as in claim 1.

16. Use of a compound represented by a following chemical formula 1, pharmaceutically acceptable salts thereof, optical isomers thereof, hydrates or solvates thereof, or mixtures thereof in the manufacture of a medicament for preventing or treating non-alcoholic fatty liver disease:

[Chemical Formula 1]



wherein A, B, and X are the same as in claim 1.