

**CD3/CD38 T cell retargeting hetero-dimeric immunoglobulins and methods of their
production**

Field of the Invention

5 The present invention relates to hetero-dimeric immunoglobulins that target both a component of the human CD3 antigen and a component of the human CD38 antigen and methods of making the same. The present invention also relates to antibodies which bind to the human CD38 antigen and derivatives thereof for use as therapeutic or diagnostic reagents and methods of making the same.

Background of the invention

10 The *in vivo* use of monoclonal antibodies (mAbs) has become a mainstay of routine clinical practice in the treatment of various human diseases and targeted immunotherapy with mAbs has become critical for the successful treatment of many forms of cancer. This is exemplified
15 by rituximab, a chimeric CD20 Antibody, which has revolutionized the treatment of several B cell malignancies such as follicular lymphoma. B cells, however, lose CD20 expression upon terminal differentiation into plasma cells, and rituximab consequently has conveyed very limited benefit to the treatment of Multiple Myeloma amongst other cancers.

20 The CD38 molecule is an attractive target because of its pattern of expression and its twin role as receptor and ectoenzyme. The targeting of CD38 has been proposed particularly as a means of treating multiple myeloma and chronic lymphocytic leukemia (CLL).

25 T cell redirected killing is a desirable mode of action in many therapeutic areas. Various bispecific antibody formats have been shown to mediate T cell redirection both in pre-clinical and clinical investigations (May C *et al.*, (2012) *Biochem Pharmacol*, 84(9): 1105-12; Frankel SR & Baeuerle PA, (2013) *Curr Opin Chem Biol*, 17(3): 385-92). All T cell retargeting bispecific antibodies or fragments thereof are engineered to have at least two antigen binding sites wherein one site binds a surface antigen on a target cell and the other site binds a T cell
30 surface antigen. Amongst T cell surface antigens, the human CD3 epsilon subunit from the TCR protein complex has been the most targeted to redirect T cell killing.

Many bispecific antibody formats have been used to redirect T cell killing, these mainly include tandem of scFv fragments and diabody based formats with only few examples of Fc-based bispecific antibody formats reported (Moore PA *et al.*, (2011) Blood, 117(17): 4542-51; May C *et al.*, (2012) *supra*; Frankel SR & Baeuerle PA, (2013) *supra*) . Bispecific formats that will encompass a human Fc region will have longer circulation half-lives which may result in enhanced efficacy and/or less frequent dosing regimens. Among possible Fc-based bispecific formats, one preferred format to redirect T cell killing is the so-called heavy chain hetero-dimer format. This format is of particular interest as it does not allow aggregation of multiple copies of human CD3 molecules at the T cell surface thereby preventing any T cell inactivation (Klein C *et al.*, (2012) MAbs, 4(6): 653-63).

The first described method to engineer heavy chain hetero-dimers is a method known as the “knob-into-hole” method (PCT Publication No: WO199627011; Merchant AM *et al.*, (1998) Nat Biotechnol, 16(7): 677-81). Recently a chemical method known as the FAB-arm exchange method wherein two antibodies are combined into one bispecific antibody via reduction and *in vitro* reshuffling of half-immunoglobulins has been reported (PCT Publication Nos: WO2008119353 (Schuurman J *et al.*) and WO2013060867 (Gramer M *et al.*); Labrijn AF *et al.*, (2013) Proc Natl Acad Sci USA, 110(13): 5145-50).

Summary of the Invention

The present invention provides an antibody or fragment thereof that binds to human CD38 comprising a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 25 or SEQ ID NO: 208, and/or a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 26 or SEQ ID NO: 209, and/or a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 27 or SEQ ID NO: 210; and/or comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 28 or SEQ ID NO: 211 or SEQ ID NO: 214, and/or a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 29 or SEQ ID NO: 212, and/or a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 30 or SEQ ID NO: 213; or

wherein the epitope binding region that binds the CD38 protein complex comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 31, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 32 and a heavy chain CDR3 comprising

the amino acid sequence of SEQ ID NO: 33, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 34, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 35 and a light chain CDR3 comprising the amino acid sequences of SEQ ID NO: 36.

- 5 In accordance with this aspect of the present invention the anti CD38 antibody or fragment thereof, is a murine antibody, chimeric antibody, humanized antibody or a fully human antibody.

In particular the antibody or fragment thereof comprises a heavy chain variable region sequence comprising the amino acid sequence consisting of SEQ ID NO: 52, SEQ ID NO: 58, SEQ ID NO: 60 or a sequence at least 80% identical to the non-CDR region of either of
10 said heavy chain variable region sequences and/or wherein the antibody or fragment thereof comprises a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 55, SEQ ID NO: 59, SEQ ID NO: 61 or a sequence at least 80% identical to the non-CDR region of any one of said light chain variable region sequences.

15 In particular the antibody or fragment thereof comprises a heavy chain sequence comprising the amino acid sequence consisting of SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 76, SEQ ID NO: 145, SEQ ID NO: 147, SEQ ID NO: 149 or a sequence at least 80% identical to the non-CDR region of either of said heavy chain variable region sequences and/or wherein the antibody or fragment thereof comprises a light chain sequence comprising the amino acid
20 sequence of SEQ ID NO: 74, SEQ ID NO: 77, SEQ ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150 or a sequence at least 80% identical to the non-CDR region of any one of said light chain variable region sequences.

In particular the antibody or fragment comprises human heavy and/or light chain constant regions and wherein the human heavy constant region is selected from the group of human
25 immunoglobulins consisting of IGHG1, non fucosylated IGHG1 and IGHG4.

The present invention also provides new anti-human CD3/CD38 bispecific antibodies.

In accordance with the present invention the target binding portion of the anti-human CD3 binding arm preferably comprises a single chain variable fragment (scFv) of SP34 or OKT3.

30 In particular wherein the anti-human CD3 binding arm is SP34 it has a better expression

profile in comparison to a scFv-Fc comprising the heavy and light variable regions encoded by SEQ ID NOs: 1 and 2, whilst maintaining its CD3 binding properties.

In particular when expressed as a scFv-Fc it has at least a twofold improvement in expression
5 in comparison to a SP34 chimera formatted as a scFv-Fc comprising the heavy and light variable regions encoded by SEQ ID NOs: 1 and 2. Preferably the improved SP34 scFv has at least a sixfold improvement in expression in comparison to a SP34 chimera formatted as an scFv comprising the heavy and light variable regions encoded by SEQ ID NOs: 1 and 2 and most preferably a twelvefold improvement in expression in comparison to a SP34 chimera
10 formatted as an scFv comprising the heavy and light variable regions encoded by SEQ ID NOs: 1 and 2.

In accordance with another aspect of the present invention when expressed as a scFv in a bispecific antibody comprising a FAB arm in the BEAT format, it has at least a twofold
15 improvement in expression in comparison to a SP34 chimera formatted as an scFv comprising the heavy and light variable regions encoded by SEQ ID NOs: 1 and 2. Preferably the improved SP34 scFv as a component of a BEAT bispecific antibody has at least a fivefold improvement in expression in comparison to a SP34 chimera formatted as an scFv, comprising the heavy and light variable regions encoded by SEQ ID NOs: 1 and 2, as a
20 component of a BEAT bispecific antibody.

In accordance with the present invention the two binding arms of the anti-human CD3 bispecific antibody each comprise an immunoglobulin constant region and wherein the first
25 arm or polypeptide binds to protein A and the second arm or polypeptide does not bind to protein A.

According to a further aspect of the present invention the epitope binding region of the polypeptide which binds the CD3 protein complex comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 3, a heavy chain CDR2 comprising the
30 amino acid sequence of SEQ ID NO: 4 and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 5 or SEQ ID NO: 9, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 6, a light chain CDR2 comprising the amino acid sequence of

SEQ ID NO: 7 or SEQ ID NO: 11 and a light chain CDR3 comprising the amino acid sequences of: SEQ ID NO: 8, SEQ ID NO: 10 or SEQ ID NO: 12; or

wherein the epitope binding region that binds the CD3 protein complex comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 163, and a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 164, and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 165; and comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 166, and a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 167, and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 168.

According to a further aspect of the present invention the epitope binding region of the polypeptide binds the CD38 protein comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 13, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 14 and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 15, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 16, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 17 and a light chain CDR3 comprising the amino acid sequences of: SEQ ID NO: 18; or

wherein the epitope binding region that binds the CD38 protein complex comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 19, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 20 and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 21, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 22, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 23 and a light chain CDR3 comprising the amino acid sequences of SEQ ID NO: 24; or

the epitope binding region of the polypeptide binds the CD38 protein comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 25, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 26 and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 27, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 28, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 29 and a light chain CDR3 comprising the amino acid sequences of: SEQ ID NO: 30; or

wherein the epitope binding region that binds the CD38 protein complex comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 31, a heavy chain CDR2

comprising the amino acid sequence of SEQ ID NO: 32 and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 33, and a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 34, a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 35 and a light chain CDR3 comprising the amino acid sequences of SEQ ID NO: 36.

Use of these new anti-human CD3/CD38 bispecific antibodies is not limited to but includes the treatments of various human cancers, autoimmune and inflammatory diseases. The specific destruction of cancer cells over healthy cells and tissues represents a primary objective in oncology. Therapeutics that could safely redirect T cell killing against tumour associated cell surface antigens may offer improved clinical efficacy. Known areas of clinical unmet needs in oncology include but are not limited to breast cancer, metastatic breast cancer, ovarian cancer, pancreatic cancer, lung cancer, lymphomas and multiple myeloma.

In accordance with a further aspect of the present invention the constant region of the second polypeptide of the hetero-dimeric immunoglobulin or fragment thereof, comprises an IgG3 CH3 region.

In accordance with a further aspect of the present invention the constant region of the second polypeptide of the hetero-dimeric immunoglobulin or fragment thereof, comprises a CH3 region other than that from IgG, and the non-IgG3 CH3 region comprises at least one substitution so as to decrease/abolish protein A binding.

According to a further aspect of the present invention the epitope binding region of second polypeptide of the hetero-dimeric immunoglobulin or fragment thereof comprises a VH3 region comprising at least one modification that reduces protein A binding.

The present invention also provides a method to produce anti-human CD3/CD38 bispecific heavy chain hetero-dimers having at least one VH3 based antigen binding site from a recombinant mammalian host cell line wherein the bispecific antibody product is readily isolated after a single Protein A chromatography step with a high degree of purity.

In particular the modified VH3 region comprises an amino acid substitution selected from the group consisting of: 57, 65, 81, 82a and combination 19/57/59 (Kabat numbering) and even more preferably wherein the modified VH3 region comprises an amino acid substitution selected from the group consisting of: 57A, 57E, 65S, 81E, 82aS and combination 19G/57A/59A (Kabat numbering).

According to a further aspect of the present invention the hetero-dimeric immunoglobulin or fragment thereof, may comprise further substitutions wherein the heavy chain variable framework region comprises an amino acid substitution selected from the group consisting of: I34M, V48I, A49G, R58N/Y, I69L, A71T and T73K (Kabat numbering) and the light chain variable framework region comprises an amino acid substitution selected from the group consisting of: M4L, V33M, A34N, L46R, L47W, T51A, R66G, F71Y and P96F (Kabat numbering); or wherein the heavy chain variable framework region comprises the amino acid substitutions I34M, A49G and A71T (Kabat numbering) and the light chain variable framework region comprises the amino acid substitutions M4L, L46R, L47W and F71Y (Kabat numbering).

In a further embodiment, the epitope binding region that binds to the CD3 protein complex comprises a heavy chain variable framework region that is the product of or derived from the human VH3 subclass. Preferably the heavy chain variable framework region is the product of or derived from human IGHV3-23. More preferably, the heavy chain variable framework region is the product of or derived from human IGHV3-23*04 (SEQ ID NO: 37). The heavy chain variable framework region comprises at least one amino acid modification from the corresponding framework region of the heavy chain variable region of the corresponding murine antibody comprising the amino acid sequence of SEQ ID NO: 38 or SEQ ID NO: 1.

In a preferred embodiment, the epitope binding region of the first polypeptide that binds to the CD3 protein complex comprises a light chain variable framework region that is the product of or derived from the human VK1 subclass or the human VK3 subclass. Preferably the light chain variable framework region is the product of or derived from human VK1-39 or VK3-20. More preferably the light chain variable framework region is the product of or derived from human IGKV1-39*01 (SEQ ID NO: 39) or IGKV3-20*01 (SEQ ID NO: 40). The light chain variable framework region comprises at least one amino acid modification from the

corresponding framework region of the light chain variable region of the corresponding murine antibody comprising the amino acid sequence of SEQ ID NO: 41 or SEQ ID NO: 2.

In a preferred embodiment, the epitope binding region that binds to the CD3 protein complex comprises a humanized heavy chain variable domain having the back mutations selected from the group consisting of: I34M, V48I, A49G, R58N/Y, I69L, A71T and T73K (Kabat numbering) and a humanized light chain variable domain having the back mutations selected from the group consisting of: M4L, V33M, A34N, L46R, L47W, R66G, F71Y and P96F (Kabat numbering). More preferably, the epitope binding region that binds to the CD3 protein complex comprises a humanized heavy chain variable domain having the back mutations I34M, A49G and A71T (Kabat numbering) and a humanized light chain variable domain having the back mutations M4L, L46R, L47W and F71Y (Kabat numbering).

According to a further aspect of the present invention the epitope binding region that binds the CD3 protein complex of the hetero-dimeric immunoglobulin or fragment thereof, wherein the epitope binding region that binds the CD3 protein complex comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 42, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 46; or wherein the epitope binding region that binds the CD3 protein complex comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 44, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 47; or wherein the epitope binding region that binds the CD3 protein complex comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 45, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 47; or wherein the epitope binding region that binds the CD3 protein complex comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 45, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 48; or wherein the epitope binding region that binds the CD3 protein complex comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 45, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 49.

The CD3 protein complex comprises a number of subunits, for example, delta, epsilon and gamma. In a preferred embodiment, the epitope binding region that binds to the CD3 protein complex binds to the CD3 epsilon subunit.

5 An epitope binding region as described herein includes the combination of one or more heavy chain variable domains and one or more complementary light chain variable domains which together form a binding site which permits the specific binding of the hetero-dimeric immunoglobulin or fragment thereof to one or more epitopes. In an embodiment of the present invention, the epitope binding region of the first poly peptide comprises a FAB and
10 the epitope binding region of the second polypeptide comprises a scFv. Alternatively, the epitope binding region of the first poly peptide comprises a scFv and the epitope binding region of the second polypeptide comprises a FAB.

. The epitope binding region that binds to CD38 comprises a heavy chain variable framework
15 region that is the product of or derived from the human VH3 subclass, preferably human VH3-23, more preferably human IGHV3-23*04 (SEQ ID NO: 37). The heavy chain variable framework region comprises at least one amino acid modification from the corresponding framework region of the heavy chain variable region of the corresponding murine antibody comprising the amino acid sequence of SEQ ID NO: 50 or 51 or 52. The epitope binding
20 region further comprises a light chain variable framework region that is the product of or derived from the human VK1 subclass, preferably human VK1-39, more preferably human IGKV1-39*01 (SEQ ID NO: 39). The light chain variable framework region comprises at least one amino acid modification from the corresponding framework region of the light chain variable region of the corresponding murine antibody comprising the amino acid sequence of
25 SEQ ID NO: 53 or 54 or 55.

In particular the CD38 binding polypeptide comprises variable heavy chain domain and variable light chain domain pair encoded by SEQ ID NOs: 56/57, 58/59, 60/61 and 62/63.

30 Anti-CD3 antibodies have been found to trigger toxicity by both direct and indirect mechanisms. Indirect mechanisms are mediated by the Fc region of the CD3 antibody which acts with the Fc receptor expressing immune cells and lead to transient T cell activation and

cytokine release. Therefore in order to improve the safety of the hetero-dimeric immunoglobulins or fragment thereof as described herein, the immunoglobulin constant region of the first and/or second polypeptide has reduced or no binding for effector immune cells and/or complement C1q. Preferably, the immunoglobulin constant region is engineered to abrogate Fc receptor binding in the lower hinge region. More preferably the immunoglobulin constant region of the first and/or second polypeptide comprises the substitution(s) L234A and/or L235A (EU numbering). Most preferably, the immunoglobulin constant region of the first and/or second polypeptide comprises the substitutions L234A and L235A (EU numbering).

In another aspect, the disclosure of the present invention also describes a hetero-dimeric immunoglobulin or fragment thereof wherein the epitope binding region binds to the CD3 epsilon subunit of the CD3 protein complex and comprises a FAB having a FAB thermo-stability superior to the FAB thermo-stability of the SP34 chimera comprising a heavy chain variable domain of amino acid sequence of SEQ ID NO: 1 and a light chain variable domain of amino acid sequence of SEQ ID NO: 2, as measured by Differential Scanning Calorimetry (DSC) as in Table 1. This increased thermostability will mean that these improved SP34 binding arms will have increased *in vivo* and *in vitro* stability, meaning better performance as a therapeutic and also in terms of its stability/storage/shelf life.

In a preferred embodiment, the present invention provides hetero-dimeric immunoglobulin or fragment thereof binding to:

i) the CD3 protein complex and CD38, wherein the first polypeptide has an amino acid sequence of SEQ ID NO: 65 and is assembled with a cognate light chain of amino acid sequence of SEQ ID NO: 66 and binds CD38, and wherein the second polypeptide has an amino acid sequence of SEQ ID NO: 67, SEQ ID NO: 68, SEQ ID NO: 69 and SEQ ID NO: 70 and binds CD3 epsilon;

ii) the CD3 protein complex and CD38, wherein the first polypeptide has an amino acid sequence of SEQ ID NO: 71 and is assembled with a cognate light chain of amino acid sequence of SEQ ID NO: 72 and binds CD38, and wherein the second polypeptide has an amino acid sequence of SEQ ID NO: 67, SEQ ID NO: 68, SEQ ID NO: 69 and SEQ ID NO: 70 and binds CD3 epsilon;

iii) the CD3 protein complex and CD38, wherein the first polypeptide has an amino acid sequence of SEQ ID NO: 73 and is assembled with a cognate light chain of amino acid sequence of SEQ ID NO: 74 and binds CD38, and wherein the second polypeptide has an amino acid sequence of SEQ ID NO: 67, SEQ ID NO: 68, SEQ ID NO: 69 and SEQ ID NO: 70 and binds CD3 epsilon;

In a further embodiment, the present invention provides hetero-dimeric immunoglobulin or fragment thereof binding to:

the CD3 protein complex and CD38, wherein the first polypeptide has an amino acid sequence of SEQ ID NO: 75 or 76 and is assembled with a light chain of amino acid sequence of SEQ ID NO: 77 and binds CD38, and wherein the second polypeptide has an amino acid sequence selected from the group comprising of SEQ ID NO: 67, SEQ ID NO: 68, SEQ ID NO: 69 and SEQ ID NO: 70 and binds CD3 epsilon;

the CD3 protein complex and CD38, wherein the first polypeptide has an amino acid sequence of SEQ ID NO: 78 and is assembled with a light chain of amino acid sequence of SEQ ID NO: 66 and binds CD38, and wherein the second polypeptide has an amino acid sequence selected from the group comprising of SEQ ID NO: 67, SEQ ID NO: 68, SEQ ID NO: 69 and SEQ ID NO: 70 and binds CD3 epsilon;

In accordance with a further aspect of the present invention there is provided a hetero-dimeric immunoglobulin or fragment thereof, comprising (a) a first polypeptide comprising an immunoglobulin constant region with a CH2 domain and a first engineered CH3 domain;

(b) a second polypeptide comprising an immunoglobulin constant region with a CH2 domain and a second engineered CH3 domain, wherein the two polypeptide heterodimerize through their engineered CH3 domains; and wherein the first polypeptide does not bind protein A or protein G and the second polypeptide only binds protein A and/or protein G through its immunoglobulin constant region; wherein the first engineered CH3 domain has substitutions at one or more of the following residues 20, 22, 26, 79, 85.1, 86, 88, 90 and the second engineered CH3 domain has substitutions in at least one residue selected from the group 3, 5, 20, 22, 26, 27, 81, 84, 85.1, 86, 88 and wherein said second chain also comprises a substitution at residue 84.4 according to the IMGT numbering.

According to a further aspect of the present invention the substitution at position 84.4 is selected from the group 84.4Q, 84.4V, 84.4A, 84.4S, 84.4T, 84.4N, 84.4G, 84.4L, 84.4L, 84.4Y, 84.4F, 84.4M.

- 5 In accordance with a further aspect of the present invention the first engineered CH3 domain has substitutions at least at the following residues 22, 86, 88 and the second engineered CH3 domain has substitutions at least at the following residues 7, 84.4, 85.1, 86 according to the IMGT numbering.
- 10 In accordance with a further aspect of the present invention the first engineered CH3 domain has substitutions at least at the following residues 20, 22, 79, 86, 88 and the second engineered CH3 domain has substitutions at least at the following residues 7, 26, 84, 84.4, 85.1, 86 according to the IMGT numbering.
- 15 In accordance with a further aspect of the present invention the first engineered CH3 domain has substitutions at least at the following residues 20, 22, 26, 79, 85.1, 86, 88, 90 and the second engineered CH3 domain has substitutions at least at the following residues 3, 5, 7, 20, 22, 26, 81, 84, 84.2, 84.4, 85.1, 86, 88, 90.
- 20 In accordance with a further aspect of the present invention the first engineered CH3 domain has substitutions at least at the following residues 3, 20, 22, 26, 79, 85.1, 86, 88, 90 and the second engineered CH3 domain has substitutions at least at the following residues 3, 5, 7, 20, 22, 26, 81, 84, 84.2, 84.4, 85.1, 86, 88, 90
- 25 In accordance with a further aspect of the present invention there is provided a heterodimeric immunoglobulin has a first chain encompassing a Fc region of the IgG3 isotype that will include a BTA CH3 domain and a non-VH3 variable domain or a VH3 based variable domain abrogated for protein A binding (using the G65S or N82aS substitutions for example) or no variable domain and therefore has no binding to protein A, and a second chain that
- 30 binds protein A encompassing a BTB D401Q CH3 domain (originating from a human IgG1 isotype for example) and either a non-VH3 variable domain or a VH3 variable domain abrogated for protein A binding (using the G65S or N82aS substitutions for example), or no variable domain.

Brief Description of the Figures

FIG. 1A-F: These figures all relate to OKT3 humanization on stable human frameworks.

FIG. 1A-C: Summary of humanized candidates formatted as human IgG1 antibodies. HPB-ALL staining relative to the chimeric OKT3 antibody: (-) indicates no binding, (+) weaker binding, (++) moderate binding and (+++) similar binding. **FIG. 1D:** DSC profiles of selected antibodies of candidates. **FIG. 1E:** Summary of humanized candidates formatted as scFv-Fc fusions. HPB-ALL staining relative to the chimeric OKT3 antibody: (-) indicates no binding, (+) weaker binding, (++) moderate binding and (+++) similar binding. **FIG. 1F:** DSC profiles of selected scFv-Fc candidates.

FIG. 2A-B: These figures all relate to SP34 humanization on stable human frameworks. **FIG. 2A:** Summary of humanized candidates formatted as human IgG1 antibodies. **FIG. 2B:** Summary of humanized candidates formatted as scFv-Fc fusion proteins (Fc of human IgG1 isotype). SPR data relative to the chimeric SP34 antibody for human and cynomolgus monkey CD3 epsilon 1-26_Fc fusion proteins: (-) indicates no binding, (+) weaker binding, (++) moderate binding, strong but not similar binding (+++), and (++++) similar binding.

FIG. 3 Shows the relative expression levels after reformatting from IgG1 to scFv-Fc for the SP34 chimera as well as SP34 H1L21, wherein a dramatic loss of expression was observed.

FIG. 4 Shows the effects on expression level of an SP34 H1L21 ScFv-Fc by Alanine scan in positions: T27, G27a, V27c, T28, T29, S30, N31, Y32, N52, K53, R54, P56, L90, Y92, S93, N94, and L95.

FIG. 5A Shows the effects on expression level of random mutation at position 29 of a SP34 H3L23 ScFv-Fc; **FIG. 5B** Shows the effects on expression level of random mutation at position 30 of a SP34 H3L23 ScFv-Fc; **FIG. 5C** Shows the effects on expression level of random mutation at position 95 of a SP34 H5L23 ScFv-Fc.

FIG. 6 shows the normalised expression level for several humanized SP34.

FIG. 7A: Antibody-antigen interaction measured by SPR between the chimeric HB-7 antibody and the human CD38 antigen. A CM5 sensor chip was covalently coupled with protein G and 200 RUs of chimeric HB-7 antibody were captured. Human CD38 protein (human CD38 extracellular domain with a poly-histidine tag) was injected at 125, 31, 7.8, 3.9, 1.9, 1 and 0.5 nM at a flow rate of 30 μ l/min in HBS-P. **FIG. 7B:** Antibody-antigen interaction measured by SPR between the humanized HB-7 best-fit antibody and the human CD38 antigen. A CM5 sensor chip was covalently coupled with protein G and 200 RUs of humanized HB-7 best-fit antibody were captured. Human CD38 protein (human CD38 extracellular domain with a poly-histidine tag) was injected at 50, 25, 12.5, 6.25 and 0.39 nM at a flow rate of 30 μ l/min in HBS-P. **FIG. 7C:** Antibody-antigen interaction measured by SPR between the humanized 9G7 best-fit antibody and the human CD38 antigen. A CM5 sensor chip was covalently coupled with protein G and 200 RUs of humanized 9G7 best-fit antibody were captured. Human CD38 protein (human CD38 extracellular domain with a poly-histidine tag) was injected at 25, 12.5, 6.25, 3.12, 1.56, 0.78, 0.39, 0.19, and 0.1 nM at a flow rate of 30 μ l/min in HBS-P. **FIG. 7D:** Antibody-antigen interaction measured by SPR between the humanized 9G7 best-framework antibody and the human CD38 antigen. A CM5 sensor chip was covalently coupled with protein G and 200 RUs of humanized 9G7 best-framework antibody were captured. Human CD38 protein (human CD38 extracellular domain with a poly-histidine tag) was injected at 50, 25, 12.5, 6.25, 3.12, 1.56, 0.78, 0.39, 0.19, and 0.1 nM at a flow rate of 30 μ l/min in HBS-P. **FIG. 7E:** Antibody-antigen interaction measured by SPR between the human 767 antibody and the human CD38 antigen. A CM5 sensor chip was covalently coupled with protein G and 200 RUs of human 767 antibody were captured. Human CD38 protein (human CD38 extracellular domain with a poly-histidine tag) was injected at 500, 250, 125, 62.5, 31.25, and 15.6 nM at a flow rate of 30 μ l/min in HBS-P. Affinity was obtained from a plot of the equilibrium response (Req) vs. analyte concentration (C) according to the following equation: $Req = KA * C * Rmax / (KA * C * n + 1)$, concentration at 50% saturation is KD. All SPR data are expressed as number of response units (abbreviated RU; Y axis) vs. time (X axis). **FIG. 7F:** DSC profiles of chimeric HB-7 and humanized HB-7 best-fit antibodies. **FIG. 7G:** DSC profiles of chimeric 9G7 and humanized 9G7 best-fit antibodies. **FIG. 7H:** DSC profiles of humanized 9G7 best-framework antibody. **FIG. 7I:** DSC profiles of human clone 767 antibody. **FIG. 7J:** summary table for the 9G7 humanized antibodies.

FIG. 8: Schematic diagram of the BEAT CD38-HB7bestfit/CD3 (format A) and BEAT CD38-767/CD3 (format B) antibodies. [(A+)] means functional Protein A binding site. [(A-)] means nonfunctional Protein A binding site.

5

FIG. 9A: Antibody-antigen interaction measured by SPR between the BEAT CD38-HB7bestfit/CD3 antibody and the human CD38 antigen. A CM5 sensor chip was covalently coupled with protein G and 200 RUs of BEAT CD38-HB7bestfit/CD3 antibody were captured. Human CD38 protein (poly-histidine tagged protein) was injected at 50, 25, 12.5, 6.25 and 0.39 nM at a flow rate of 30µl/min in HBS-P. Data are expressed as number of response units (abbreviated RU; Y axis) vs. time (X axis). **FIG. 9B:** BEAT CD38-HB7bestfit/CD3 antibody DSC profile.

10

FIG. 10: Example of T cell redirected killing by the BEAT CD38-HB7bestfit/CD3 antibody.

15

Readout: RDL-FACS method. Effector cells: purified human T cells. Effector cells-to-targeted cells ratio of 10:1. Mean of two donors with 48h incubation. Target cells: RPMI 8226. Antibody concentration is shown in nM.

FIG. 11: Example of T cell redirected killing by the BEAT CD38-767/CD3(SP34) antibody.

20

Readout: RDL-FACS method. Effector cells: human PBMCs. Effector cells-to-targeted cells ratio of 10:1. Mean of three donors with 24h incubation. Target cells: Daudi. Antibody concentration is shown in nM.

FIG. 12: Schematic diagram of the BEAT CD38-HB7bestfit/CD3(SP34) (format A) and

25

BEAT CD38-9G7bestfit/CD3(SP34-Kappa2) (format B) antibodies. [(A+)] means functional Protein A binding site.

FIG. 13: Example of T cell redirected killing by the BEAT CD38-HB7bestfit/CD3(SP34)

30

antibody. Readout: RDL-FACS method. Effector cells: Human PBMCs. Effector cells-to-targeted cells ratio of 10:1. Mean of three donors with 24h incubation. Target cells: Daudi cells. Antibody concentration is shown in nM.

FIG. 14: Antibody-antigen interaction measured by SPR between the BEAT CD38-9G7bestfit/CD3(SP34-Kappa2) antibody and the human CD3 epsilon 1-26_Fc fusion protein. A CM5 sensor chip was covalently coupled with 500 RUs of the human CD3 epsilon 1-26_Fc fusion protein. BEAT CD38-9G7bestfit/CD3(SP34-Kappa2) antibody was injected at 50, 25, 12.5, 6.2, 3.1, 0.8 and 0.4 nM at a flow rate of 30 µl/min in HBS-P. Data are expressed as number of response units (abbreviated RU; Y axis) vs. time (X axis).

FIG. 15: Example of T cell redirected killing by the BEAT CD38/CD3(SP34-Kappa2) antibody. Readout: RDL-FACS method. Effector cells: Human PBMCs. Effector cells-to-targeted cells ratio of 10:1. Mean of three donors with 24h incubation. Target cells: Daudi cells. Antibody concentration is shown in nM.

FIG. 16 shows the performance in an RDL assay of several CD38/CD3 bispecific antibodies, in which the CD3 binding arm comprises the original mouse SP34 reformatted as an scFv (SEQ ID NO: 137), or modified humanised SP34 scFv's comprising the heavy/light chain combinations H1/L21 (SEQ ID NO: 67), H5/L32 (SEQ ID NO: 68), H5/L65 (SEQ ID NO: 69) and H5/L67 (SEQ ID NO: 70).

FIG. 17: Non reducing SDS-PAGE analysis of BTB homodimers with (lanes group 1) and without protein A binding scFv moiety (lanes group 2). Purification using affinity chromatography: eluted and neutralized fractions were analyzed after protein A chromatography (A) or protein G chromatography (G). For group 2, flow-through or unbound fractions were collected and showed no binding to protein A or G (abbreviated A ft and G ft, respectively). Elutions were performed at pH 3.0. [(A+)] means a Protein A binding site. Molecular weight markers as indicated (kDa).

FIG. 18A: Non reducing SDS-PAGE analysis of the BEAT heterodimer containing only a single protein A site in the BTB chain carrying the D401Q substitution. Purification using affinity chromatography: eluted and neutralized fractions were analyzed after protein A chromatography (A) or protein G chromatography (G). Elutions were performed at pH 3.0. [(A+)] means a Protein A binding site. Molecular weight markers as indicated (kDa); **18B:** SDS-PAGE analysis of the BEAT heterodimer containing a single protein A site in the BTB chain lacking the D401Q substitution. Purification using affinity chromatography: eluted and

neutralized fractions were analyzed after protein A chromatography (A) or protein G chromatography (G). Elutions were performed at pH 3.0. [(A+)] means a Protein A binding site. Molecular weight markers as indicated (kDa).

FIG. 19: The graphs show the count per μl of blood for the CD4⁺ population (plain line) and CD8⁺ population (dotted line), for both animals that were injected with CD3/CD38 BEAT. The upper graphs show the counts for the animals dosed with 1 $\mu\text{g/kg}$. The bottom graphs show the counts for the animals dosed with 10 $\mu\text{g/kg}$. The counts at time points zero correspond to samples harvested 1 day prior to dosing.

FIG. 20: The graphs show the count per μl of blood for the CD14⁺CD38⁺ monocyte population, for both animals that were injected with CD3/CD38 BEAT. The upper graphs show the counts for the animals dosed with 1 $\mu\text{g/kg}$. The bottom graph shows the counts for the animals dosed with 10 $\mu\text{g/kg}$. The counts at time points zero correspond to samples harvested 1 day prior to dosing.

FIG. 21: The graphs show the mean fluorescence intensity (MFI) of the fluo-4 dye as a function of time in seconds. The isotype control condition is included in each graph. After the baseline acquisition during 40 s, the indicated antibodies were added into the samples and the acquisition was resumed. The isotype control was a human IgG1 antibody. Increase in MFI indicates a calcium mobilization into the cytoplasm of the cells.

FIG. 22: Epitope mapping of the humanized 9G7 antibody by SPR using peptide-Fc fusions derived from the extracellular domain of human CD38. Data are expressed as number of response units (abbreviated RU; Y axis) vs. time (X axis).

FIG. 23: 3D rendering of the extracellular domain of human CD38. The F6 peptide sequence is colored grey, positions M110 and T148 are colored black.

FIG. 24: Epitope mapping of the humanized 9G7 and SAR650984 antibodies by SPR using the extracellular domains of human and cynomolgus monkey CD38. Data are expressed as number of response units (abbreviated RU; Y axis) vs. time (X axis).

FIG. 25: Epitope mapping of the human 767 antibody by SPR using peptide-Fc fusions derived from the extracellular domain of human CD38. Data are expressed as number of response units (abbreviated RU; Y axis) vs. time (X axis).

5 **FIG. 26:** 3D rendering of the extracellular domain of human CD38. The F3 peptide sequence is colored grey, positions E76 and H79 are colored black.

10 **FIG. 27:** Epitope mapping of the humanized SAR650984 antibody by SPR using peptide-Fc fusions derived from the extracellular domain of human CD38. Data are expressed as number of response units (abbreviated RU; Y axis) vs. time (X axis).

Detailed description of the invention

The present invention relates generally to novel hetero-dimeric immunoglobulins that bind to the CD3 protein complex and a CD38 antigen.

15 Furthermore, these hetero-dimeric immunoglobulins have reduced or eliminated binding to protein A and therefore can be purified to a very high degree of purity using affinity chromatography.

20 For the purposes of interpreting this specification, the following definitions will apply and whenever appropriate, terms used in the singular will also include the plural and vice versa. It is to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

25 The terms “polypeptide” and “protein” refer to a polymer of amino acid residues wherein amino acids are combined via peptide bonds to form a chain of amino acids that have been linked together via dehydration synthesis. Polypeptides and proteins can be synthesized through chemical synthesis or recombinant expression and are not limited to a minimum amino acid length.

30 In accordance with the invention, the group of polypeptides comprises “proteins” as long as the proteins consist of a single polypeptide chain. Polypeptides may further form multimers such as dimers, trimers and higher oligomers, i.e. consisting of more than one polypeptide

molecule. Polypeptide molecules forming such dimers, trimers etc. may be identical or non-identical. The corresponding higher order structures of such multimers are, consequently, termed homo- or hetero-dimers, homo- or hetero-trimers etc. An example for a hetero-multimer is an antibody molecule, which, in its naturally occurring form, consists of two identical light polypeptide chains and two identical heavy polypeptide chains. The terms “polypeptide” and “protein” also refer to naturally modified polypeptides/proteins wherein the modification is effected e.g. by post-translational modifications like glycosylation, acetylation, phosphorylation and the like. Such modifications are well known in the art. Furthermore, for purposes of the present invention, a “polypeptide” refers to a protein which includes modifications, such as deletions, additions and substitutions (which can be conservative in nature) to the native sequence. These modifications may be deliberate, as through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the proteins or errors due to PCR amplification.

The term “CD3 complex” as used herein refers to the protein complex known as the CD3 (cluster of differentiation 3) T-cell co-receptor (Wucherpfennig KW *et al.*, (2010) Cold Spring Harb Perspect Biol, 2(4): a005140). The CD3 protein complex is composed of four distinct chains. In mammals, the complex contains a CD3 γ chain, a CD3 δ chain, and two CD3 ϵ chains. These chains associate with a molecule known as the T-cell receptor (TCR) and the ζ -chain to generate an activation signal in T lymphocytes (van der Merwe PA & Dushek O (2011) Nat Rev Immunol, 11(1): 47-55). The TCR, ζ -chain, and CD3 molecules together comprise the TCR complex. The CD3 γ , CD3 δ , and CD3 ϵ chains are highly related cell-surface proteins of the immunoglobulin superfamily containing a single extracellular immunoglobulin domain. The intracellular tails of the CD3 molecules contain a single conserved motif known as an immunoreceptor tyrosine-based activation motif or ITAM for short, which is essential for the signalling capacity of the TCR. Since CD3 is required for T-cell activation, drugs (often monoclonal antibodies) that target CD3 have and are being investigated as immunosuppressant therapies.

The term “immunoglobulin” as referred to herein can be used interchangeably with the term “antibody”. Immunoglobulin includes full-length antibodies and any antigen binding fragment or single chains thereof. Immunoglobulins can be homo-dimeric or hetero-dimeric. Immunoglobulins and specifically naturally occurring antibodies are glycoproteins which

exist as one or more copies of a Y-shaped unit, composed of four polypeptide chains. Each “Y” shape contains two identical copies of a heavy (H) chain and two identical copies of a light (L) chain, named as such by their relative molecular weights. Each light chain pairs with a heavy chain and each heavy chain pairs with another heavy chain. Covalent interchain

5 disulfide bonds and non-covalent interactions link the chains together. Immunoglobulins and specifically naturally occurring antibodies contain variable regions, which are the two copies of the antigen binding site. Papain, a proteolytic enzyme splits the “Y” shape into three separate molecules, two so called “Fab” or “FAB” fragments (Fab = fragment antigen binding) and one so called “Fc” fragment or “Fc region” (Fc = fragment crystallizable). A Fab

10 fragment consists of the entire light chain and part of the heavy chain. The heavy chain contains one variable region (VH) and either three or four constant regions (CH1, CH2, CH3 and CH4, depending on the antibody class or isotype). The region between the CH1 and CH2 regions is called the hinge region and permits flexibility between the two Fab arms of the Y-shaped antibody molecule, allowing them to open and close to accommodate binding to two

15 antigenic determinants separated by a fixed distance. The “hinge region” as referred to herein is a sequence region of 6-62 amino acids in length, only present in IgA, IgD and IgG, which encompasses the cysteine residues that bridge the two heavy chains. The heavy chains of IgA, IgD and IgG each have four regions, i.e. one variable region (VH) and three constant regions (CH1-3). IgE and IgM have one variable and four constant regions (CH1-4) on the heavy

20 chain. The constant regions of the immunoglobulins may mediate the binding to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the complement system classical pathway. Each light chain is usually linked to a heavy chain by one covalent disulfide bond. Each light chain contains one variable region (VL) and one light chain constant region. The light chain constant region is a kappa

25 light chain constant region designated herein as IGKC or is a lambda light chain constant region designated herein as IGLC. IGKC is used herein equivalently to C κ or CK and has the same meaning. IGLC is used herein equivalently to C λ or CL and has the same meaning. The term “an IGLC region” as used herein refer to all lambda light chain constant regions e.g. to all lambda light chain constant regions selected from the group consisting of IGLC1, IGLC2,

30 IGLC3, IGLC6 and IGLC7. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR or FW). Each VH and VL is composed of three CDRs and four FRs, arranged from amino- terminus to carboxy-terminus

in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain an epitope-binding region that interacts with an antigen. Engineered immunoglobulins can encompass different epitope binding region formats such as scFv, FAB or dAb fragments. These fragments are usually assembled in an antibody-like structure by genetic fusion to a IgG Fc region. Engineered immunoglobulins can be constructed as homo or hetero-dimers with or without the use of hetero-dimerization enhancing techniques, and can have mono- or bispecific binding properties.

The term “full length antibody” as used herein includes the structure that constitutes the natural biological form of an antibody, including variable and constant regions. For example, in most mammals, including humans and mice, the full length antibody of the IgG class is a tetramer and consists of two identical pairs of two immunoglobulin chains, each pair having one light and one heavy chain, each light chain comprising immunoglobulin regions VL and a light chain constant region, and each heavy chain comprising immunoglobulin regions VH, CH1 (C γ 1), CH2 (C γ 2), CH3 (C γ 3) and CH4 (C γ 4), depending on the antibody class or isotype). In some mammals, for example in camels and llamas, IgG antibodies may consist of only two heavy chains, each heavy chain comprising a variable region attached to the Fc region.

Antibodies are grouped into classes, also referred to as isotypes, as determined genetically by the constant region. Human constant light chains are classified as kappa (CK) and lambda (C λ) light chains. Heavy chains are classified as mu (μ), delta (δ), gamma (γ), alpha (α), or epsilon (ϵ) and define the antibody's isotype as IgM, IgD, IgG, IgA and IgE, respectively. Thus, “isotype” as used herein is meant any of the classes and/or subclasses of immunoglobulins defined by the chemical and antigenic characteristics of their constant regions. The known human immunoglobulin isotypes are IGHG1 (IgG1), IGHG2 (IgG2), IGHG3 (IgG3), IGHG4 (IgG4), IGHA1 (IgA1), IGHA2 (IgA2), IGHM (IgM), IGHD (IgD) and IGHE (IgE). The so-called human immunoglobulin pseudo-gamma IGHGP gene represents an additional human immunoglobulin heavy constant region gene which has been sequenced but does not encode a protein due to an altered switch region (Bensmana M *et al.*, (1988) Nucleic Acids Res, 16(7): 3108). In spite of having an altered switch region, the human immunoglobulin pseudo-gamma IGHGP gene has open reading frames for all heavy constant regions (CH1-CH3) and hinge. All open reading frames for its heavy constant

regions encode protein regions which align well with all human immunoglobulin constant regions with the predicted structural features. This additional pseudo-gamma isotype is referred herein as IgGP or IGHGP. Other pseudo immunoglobulin genes have been reported such as the human immunoglobulin heavy constant region epsilon P1 and P2 pseudo-genes (IGHEP1 and IGHEP2). The IgG class is the most commonly used for therapeutic purposes. In humans this class comprises subclasses IgG1, IgG2, IgG3 and IgG4. In mice this class comprises subclasses IgG1, IgG2a, IgG2b, IgG2c and IgG3.

The term “Immunoglobulin fragments” as used herein include, but is not limited to, (i) a region including for example a CH1, a CH2 or a CH3 region, (ii) the Fab fragment consisting of VL, VH, CL or CK and CH1 regions, including Fab' and Fab'-SH, (iii) the Fd fragment consisting of the VH and CH1 regions, (iv) the dAb fragment (Ward ES *et al.*, (1989) Nature, 341(6242): 544-6) which consists of a single variable region (v) F(ab')₂ fragments, a bivalent fragment comprising two linked Fab fragments (vi) single chain Fv fragments (scFv), wherein a VH region and a VL region are linked by a peptide linker which allows the two regions to associate to form an antigen binding site (Bird RE *et al.*, (1988) Science, 242(4877): 423-6; Huston JS *et al.*, (1988) Proc Natl Acad Sci U S A, 85(16): 5879-83), (vii) “diabodies” or “triabodies”, multivalent or multispecific fragments constructed by gene fusion (Holliger P *et al.*, (1993) Proc Natl Acad Sci U S A, 90(14): 6444-8; Tomlinson I & Holliger P, (2000) Methods Enzymol, 326:461-79), (viii) scFv, diabody or region antibody fused to an Fc region and (ix) scFv fused to the same or a different antibody.

The term “variable region” refers to the regions or domains that mediates antigen-binding and defines specificity of a particular antibody for a particular antigen. In naturally occurring antibodies, the antigen-binding site consists of two variable regions that define specificity: one located in the heavy chain, referred herein as heavy chain variable region (VH) and the other located in the light chain, referred herein as light chain variable region (VL). In humans, the heavy chain variable region (VH) can be divided into seven subgroups or subclasses: VH1, VH2, VH3, VH4, VH5, VH6 and VH7. In some cases, specificity may exclusively reside in only one of the two regions as in single-domain antibodies from heavy-chain antibodies found in camelids. The V regions are usually about 110 amino acids long and consist of relatively invariant stretches of amino acid sequence called framework regions (FRs or “non-CDR regions”) of 15-30 amino acids separated by shorter regions of extreme

variability called “hypervariable regions” that are 7-17 amino acids long. The variable domains of native heavy and light chains comprise four FRs, largely adopting a beta-sheet configuration, connected by three hypervariable regions, which form loops. The hypervariable regions in each chain are held together in close proximity by FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen binding site of antibodies (see Kabat EA *et al.*, *supra.*). The term “hypervariable region” as used herein refers to the amino acid residues of an antibody which are responsible for antigen binding. The hypervariable region generally comprises amino acid residues from a “complementary determining region” or “CDR”, the latter being of highest sequence variability and/or involved in antigen recognition. For all variable regions numbering is according to Kabat (Kabat EA *et al.*, *supra.*).

A number of CDR definitions are in use and are encompassed herein. The Kabat definition is based on sequence variability and is the most commonly used (Kabat EA *et al.*, *supra.*). Chothia refers instead to the location of the structural loops (Chothia & Lesk J. (1987) *Mol Biol*, 196: 901-917). The AbM definition is a compromise between the Kabat and the Chothia definitions and is used by Oxford Molecular's AbM antibody modelling software (Martin ACR *et al.*, (1989) *Proc Natl Acad Sci USA* 86:9268–9272; Martin ACR *et al.*, (1991) *Methods Enzymol*, 203: 121–153; Pedersen JT *et al.*, (1992) *Immunomethods*, 1: 126–136; Rees AR *et al.*, (1996) In Sternberg M.J.E. (ed.), *Protein Structure Prediction*. Oxford University Press, Oxford, 141–172). The contact definition has been recently introduced (MacCallum RM *et al.*, (1996) *J Mol Biol*, 262: 732-745) and is based on an analysis of the available complex structures available in the Protein Databank. The definition of the CDR by IMGT®, the international ImMunoGeneTics information system® (<http://www.imgt.org>) is based on the IMGT numbering for all immunoglobulin and T cell receptor V-REGIONS of all species (IMGT®, the international ImMunoGeneTics information system®; Lefranc MP *et al.*, (1999) *Nucleic Acids Res*, 27(1): 209-12; Ruiz M *et al.*, (2000) *Nucleic Acids Res*, 28(1): 219-21; Lefranc MP (2001) *Nucleic Acids Res*, 29(1): 207-9; Lefranc MP (2003) *Nucleic Acids Res*, 31(1): 307-10; Lefranc MP *et al.*, (2005) *Dev Comp Immunol*, 29(3): 185-203; Kaas Q *et al.*, (2007) *Briefings in Functional Genomics & Proteomics*, 6(4): 253-64). All Complementarity Determining Regions (CDRs) as referred to in the present invention, are defined preferably as follows (numbering according to Kabat EA *et al.*, *supra.*): LCDR1: 24-34, LCDR2: 50-56, LCDR3: 89-98, HCDR1: 26-35, HCDR2: 50-65, HCDR3:

95-102.

The “non-CDR regions” of the variable domain are known as framework regions (FR). The “non-CDR regions” of the VL region as used herein comprise the amino acid sequences: 1-23 (FR1), 35-49 (FR2), 57-88 (FR3) and 99-107 (FR4). The “non-CDR regions” of the VH region as used herein comprise the amino acid sequences: 1-25 (FR1), 36-49 (FR2), 66-94 (FR3) and 103-113 (FR4).

The CDRs of the present invention may comprise "extended CDRs" which are based on the aforementioned definitions and have variable domain residues as follows: LCDR1: 24-36, LCDR2: 46-56, LCDR3:89-97, HCDR1: 26-35, HCDR2:47-65, HCDR3: 93-102. These extended CDRs are numbered as well according to Kabat *et al., supra*. The “non-extended CDR region” of the VL region as used herein comprise the amino acid sequences: 1-23 (FR1), 37-45 (FR2), 57-88 (FR3) and 98- approximately 107 (FR4). The “non-extended CDR region” of the VH region as used herein comprise the amino acid sequences: 1-25 (FR1), 37-46 (FR2), 66-92 (FR3) and 103- approximately 113 (FR4).

The term “Fab” or “FAB” or “Fab region” or “FAB region” as used herein includes the polypeptides that comprise the VH, CH1, VL and light chain constant immunoglobulin regions. Fab may refer to this region in isolation, or this region in the context of a full length antibody or antibody fragment.

The term “Fc” or “Fc region”, as used herein includes the polypeptide comprising the constant region of an antibody heavy chain excluding the first constant region immunoglobulin region. Thus Fc refers to the last two constant region immunoglobulin regions of IgA, IgD and IgG or the last three constant region immunoglobulin regions of IgE and IgM, and the flexible hinge N-terminal to these regions. For IgA and IgM, Fc may include the J chain. For IgG, Fc comprises immunoglobulin regions Cgamma2 and Cgamma3 (Cγ2 and Cγ3) and the hinge between Cgamma1 (Cγ1) and Cgamma2 (Cγ2). Although the boundaries of the Fc region may vary, the human IgG heavy chain Fc region is usually defined to comprise residues C226 or P230 to its carboxyl-terminus, wherein the numbering is according to the EU index. Fc may refer to this region in isolation or this region in the context of an Fc polypeptide, for example an antibody.

The term “immunoglobulin constant region” as used herein refers to immunoglobulin or antibody heavy chain constant regions from human or animal species and encompasses all isotypes. Preferably, immunoglobulin constant regions are of human origin and are selected from the group consisting of, but not limited to: IGHG1 CH1, IGHG2 CH1, IGHG3 CH1, IGHG4 CH1, IGHA1 CH1, IGHA2 CH1, IGHE CH1, IGHEP1 CH1, IGHM CH1, IGHD CH1, IGHGP CH1, IGHG1 CH2, IGHG2 CH2, IGHG3 CH2, IGHG4 CH2, IGHA1 CH2, IGHA2 CH2, IGHE CH2, IGHEP1 CH2, IGHM CH2, IGHD CH2, IGHGP CH2, IGHG1 CH3, IGHG2 CH3, IGHG3 CH3, IGHG4 CH3, IGHA1 CH3, IGHA2 CH3, IGHE CH3, IGHEP1 CH3, IGHM CH3, IGHD CH3, IGHGP CH3, IGHE CH4 and IGHM CH4. Preferred “immunoglobulin constant regions” are selected from the group consisting of human IGHE CH2, IGHM CH2, IGHG1 CH3, IGHG2 CH3, IGHG3 CH3, IGHG4 CH3, IGHA1 CH3, IGHA2 CH3, IGHE CH3, IGHM CH3, IGHD CH3 and IGHGP CH3. More preferred “immunoglobulin constant regions” are selected from the group consisting of human IGHG1 CH3, IGHG2 CH3, IGHG3 CH3, IGHG4 CH3, IGHA1 CH3, IGHA2 CH3, IGHE CH3, IGHM CH3, IGHD CH3 and IGHGP CH3.

The term “epitope binding region” includes a polypeptide or a fragment thereof having minimal amino acid sequence to permit the specific binding of the immunoglobulin molecule to one or more epitopes. Naturally occurring antibodies have two epitope binding regions which are also known as antigen binding or combining sites or paratopes. Epitope binding regions in naturally occurring antibodies are confined within the CDR regions of the VH and/or VL domains wherein the amino acid mediating epitope binding are found. In addition to naturally occurring antibodies, artificial VH domains or VL domains or fragments thereof and combinations thereof can be engineered to provide epitope binding regions (Holt LJ *et al.*, (2003) Trends Biotechnol, 21(11): 484-490; Polonelli L *et al.*, (2008) PLoS ONE, 3(6): e2371). Examples of non-immunoglobulin based epitope binding regions can be found in artificial protein domains used as “scaffold” for engineering epitope binding regions (Binz HK *et al.*, (2005) Nat Biotechnol, 23(10): 1257-1268) or peptide mimetics (Murali R & Greene MI (2012) Pharmaceuticals, 5(2): 209-235). Preferably the term 'epitope binding region' includes the combination of one or more heavy chain variable domains and one or more complementary light chain variable domains which together forms a binding site which permits the specific binding of the immunoglobulin molecule to one or more epitopes.

Examples of an epitope binding region as exemplified in the present invention include scFv and FAB.

As used herein, the term “epitope” includes a fragment of a polypeptide or protein or a non-protein molecule having antigenic or immunogenic activity in an animal, preferably in a mammal and most preferably in a human. An epitope having immunogenic activity is a fragment of a polypeptide or protein that elicits an antibody response in an animal. An epitope having antigenic activity is a fragment of a polypeptide or protein to which an antibody or polypeptide specifically binds as determined by any method well-known to one of skill in the art, for example by immunoassays. Antigenic epitopes need not necessarily be immunogenic. Preferably, the term “epitope” as used herein refers to a polypeptide sequence of at least about 3 to 5, preferably about 5 to 10 or 15 and not more than about 1,000 amino acids (or any integer there between), which define a sequence that by itself or as part of a larger sequence, binds to an antibody generated in response to such sequence. There is no critical upper limit to the length of the fragment, which may comprise nearly the full-length of the protein sequence, or even a fusion protein comprising one or more epitopes. An epitope for use in the subject invention is not limited to a polypeptide having the exact sequence of the portion of the parent protein from which it is derived. Thus the term “epitope” encompasses sequences identical to the native sequence, as well as modifications to the native sequence, such as deletions, additions and substitutions (generally conservative in nature). The epitopes of protein antigens are divided into two categories, conformational epitopes and linear epitopes, based on their structure and interaction with the epitope binding site (Goldsby R *et al.*, (2003) “Antigens (Chapter 3)” Immunology (Fifth edition ed.), New York: W. H. Freeman and Company. pp. 57-75, ISBN 0-7167-4947-5). A conformational epitope is composed of discontinuous sections of the antigen's amino acid sequence. These epitopes interact with the paratope based on the 3-D surface features and shape or tertiary structure of the antigen. Most epitopes are conformational. By contrast, linear epitopes interact with the paratope based on their primary structure. A linear epitope is formed by a continuous sequence of amino acids from the antigen.

The term “hetero-dimeric immunoglobulin” or “hetero-dimeric fragment” or “hetero-dimer” or “hetero-dimer of heavy chains” as used herein includes an immunoglobulin molecule or part of comprising at least a first and a second polypeptide, like a first and a second region,

wherein the second polypeptide differs in amino acid sequence from the first polypeptide. Preferably, a hetero-dimeric immunoglobulin comprises two polypeptide chains, wherein the first chain has at least one non-identical region to the second chain, and wherein both chains assemble, i.e. interact through their non-identical regions. More preferably the hetero-dimeric immunoglobulin, has binding specificity for at least two different ligands, antigens or binding sites, i.e. is bispecific. Hetero-dimeric immunoglobulin as used herein includes but is not limited to full length bispecific antibodies, bispecific Fab, bispecific F(ab')₂, bispecific scFv fused to an Fc region, diabody fused to an Fc region and domain antibody fused to an Fc region.

The term "homo-dimeric immunoglobulin" or "homo-dimeric fragment" or "homo-dimer" or "homo-dimer of heavy chains" as used herein includes an immunoglobulin molecule or part of comprising at least a first and a second polypeptide, like a first and a second region, wherein the second polypeptide is identical in amino acid sequence to the first polypeptide. Preferably, a homo-dimeric immunoglobulin comprises two polypeptide chains, wherein the first chain has at least one identical region to the second chain, and wherein both chains assemble, i.e. interact through their identical regions. Preferably, a homo-dimeric immunoglobulin fragment comprises at least two regions, wherein the first region is identical to the second region, and wherein both regions assemble, i.e. interact through their protein-protein interfaces.

For all immunoglobulin constant regions included in the present invention, numbering can be according to the IMGT[®] (IMGT[®]; *supra*).

For all human CH1, CH2, CH3 immunoglobulin heavy chain constant regions selected from the group consisting of IGHG1, IGHG2, IGHG3 and IGHG4, numbering can be according to the "EU numbering system" (Edelman GM *et al.*, (1969) Proc Natl Acad Sci USA, 63(1): 78-85). A complete correspondence for the human CH1, hinge, CH2 and CH3 constant regions of IGHG1 can be found at the IMGT database (IMGT[®]; *supra*).

For the human kappa immunoglobulin light chain constant region (IGKC), numbering can be according to the "EU numbering system" (Edelman GM *et al.*, *supra*). A complete correspondence for the human CK region can be found at IMGT database (IMGT[®]; *supra*).

For the human lambda immunoglobulin light chain constant regions (IGLC1, IGLC2, IGLC3, IGLC6 and IGLC7), numbering can be according to the “Kabat numbering system” (Kabat EA *et al.*, *supra*). A complete correspondence for human IGLC regions can be found at the IMGT database (IMGT[®]; *supra*).

The human IGHG1 immunoglobulin heavy chain constant regions as referred to herein have the following region boundaries: CH1 region (EU numbering: 118-215), Hinge γ 1 region (EU numbering: 216-230), CH2 region (EU numbering: 231-340) and CH3 region (EU numbering: 341-447). The human CK region referred herein spans residues 108 to 214 (EU numbering). The human IGLC1, IGLC2, IGLC3, IGLC6 and IGLC7 regions referred herein span residues 108-215 (Kabat numbering).

The terms “amino acid” or “amino acid residue” as used herein includes natural amino acids as well as non-natural amino acids. Preferably natural amino acids are included.

The term “modification” or “amino acid modification” herein includes an amino acid substitution, insertion and/or deletion in a polypeptide sequence. The terms “substitution” or “amino acid substitution” or “amino acid residue substitution” as used herein refers to a substitution of a first amino acid residue in an amino acid sequence with a second amino acid residue, whereas the first amino acid residue is different from the second amino acid residue i.e. the substituted amino acid residue is different from the amino acid which has been substituted. For example, the substitution R94K refers to a variant polypeptide, in which the arginine at position 94 is replaced with a lysine. For example 94K indicates the substitution of position 94 with a lysine. For the purposes herein, multiple substitutions are typically separated by a slash or a comma. For example, “R94K/L78V” or “R94K, L78V” refers to a double variant comprising the substitutions R94K and L78V. By “amino acid insertion” or “insertion” as used herein is meant the addition of an amino acid at a particular position in a parent polypeptide sequence. For example, insert -94 designates an insertion at position 94. By “amino acid deletion” or “deletion” as used herein is meant the removal of an amino acid at a particular position in a parent polypeptide sequence. For example, R94- designates the deletion of arginine at position 94.

In certain embodiments, the terms “decrease”, “reduce”, or “reduction” in binding to Protein A refers to an overall decrease of at least 25%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, or 99% up to 100% (elimination) in the binding of a modified immunoglobulin or fragment thereof to Protein A detected by standard art known methods such as those described herein, as compared to a parental i.e. unmodified immunoglobulin or wild-type IgG or an IgG having the wild-type human IgG Fc region. In certain embodiments these terms alternatively may refer to an overall decrease of 10-fold (i.e. 1 log), 100-fold (2 logs), 1,000-fold (or 3 logs), 10,000-fold (or 4 logs), or 100,000-fold (or 5 logs).

The terms “eliminate”, “abrogate”, “elimination” or “abrogation” of binding to Protein A refers to an overall decrease of 100% in the binding of a modified immunoglobulin or fragment thereof to Protein A i.e. a complete loss of the binding of a modified immunoglobulin or fragment thereof to Protein A, detected by standard art known methods such as those described herein, as compared to a parental i.e. unmodified immunoglobulin or wild-type IgG or an IgG having the wild-type human IgG Fc region.

Similarly, the terms “decrease”, “reduce”, or “reduction” in binding to an affinity reagent refers to an overall decrease of at least 25%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, or 99% up to 100% (elimination) in the binding of a modified immunoglobulin or fragment thereof to the affinity reagent detected by standard art known methods such as those described herein, as compared to a parental, i.e. unmodified immunoglobulin or wild-type IgG or an IgG having the wild-type human IgG Fc region. In certain embodiments these terms alternatively may refer to an overall decrease of 10-fold (i.e. 1 log), 100-fold (2 logs), 1,000-fold (or 3 logs), 10,000-fold (or 4 logs), or 100,000-fold (or 5 logs).

The terms “eliminate” , “abrogate”, “elimination” or “abrogation” of binding to an affinity reagent refers to an overall decrease of 100% in the binding of a modified immunoglobulin or fragment thereof to the affinity reagent i.e. a complete loss of the binding of a modified immunoglobulin or fragment thereof to the affinity reagent detected by standard art known methods such as those described herein, as compared to a parental, i.e. unmodified immunoglobulin or wild-type IgG or an IgG having the wild-type human IgG Fc region.

“Bispecific antibodies” are monoclonal antibodies that have binding specificities for at least two different antigens. In certain embodiments, the bispecific antibodies are bispecific antibodies with one or more amino acid modifications in the VH region relative to the parental antibody. In certain embodiments, bispecific antibodies may be human or humanized antibodies. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express a target antigen. These antibodies possess a target-antigen-binding arm and an arm which binds a cytotoxic agent, such as, e.g., saporin, anti-interferon- α , vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten. Bispecific antibodies can be prepared as full length antibodies or antibody fragments. Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, (1983) *Nature*, 305: 537-40). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome and the product yields are low. Similar procedures are disclosed in WO1993/08829 and in Traunecker *et al.*, (1991) *EMBO J*, 10: 3655-9. According to a different approach, antibody variable regions with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant region sequences. The fusion, for example, is with an immunoglobulin heavy chain constant region, comprising at least part of the hinge, CH2 and CH3 regions. In certain embodiments, the first heavy-chain constant region (CH1), containing the site necessary for light chain binding, is present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors and are co-transfected into a suitable host organism. This provides for flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

Bispecific antibodies include cross-linked or “heteroconjugate” antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US4,676,980) and for treatment of HIV infection (WO1991/00360, WO1992/00373 and EP03089). Heteroconjugate antibodies may be made using any convenient cross-linking method. Suitable cross-linking agents are well known in the art (see US4,676,980), along with a number of cross-linking techniques. Antibodies with more than two valencies are also contemplated. For example, trispecific antibodies can be prepared (see Tutt A *et al.* (1991) J. Immunol. 147: 60-9).

Protein A: Protein A is a cell wall component produced by several strains of *Staphylococcus aureus* which consists of a single polypeptide chain. The Protein A gene product consists of five homologous repeats attached in a tandem fashion to the pathogen’s cell wall. The five domains are approximately 58 amino acids in length and denoted EDABC, each exhibiting immunoglobulin binding activity (Tashiro M & Montelione GT (1995) Curr. Opin. Struct. Biol., 5(4): 471-481). The five homologous immunoglobulin binding domains fold into a three-helix bundle. Each domain is able to bind proteins from many mammalian species, most notably IgGs (Hober S *et al.*, (2007) J. Chromatogr. B Analyt. Technol. Biomed. Life Sci., 848(1): 40-47). Protein A binds the heavy chain of most immunoglobulins within the Fc region but also within the Fab region in the case of the human VH3 family (Jansson B *et al.*, (1998) FEMS Immunol. Med. Microbiol., 20(1): 69-78). Protein A binds IgG from various species including human, mouse, rabbit and guinea pig but does not bind human IgG3 (Hober S *et al.*, (2007) *supra*). The inability of human IgG3 to bind Protein A can be explained by the H435R and Y436F substitutions in the human IgG3 Fc region (EU numbering, Jendeberg *et al.*, (1997) J. Immunol. Methods, 201(1): 25-34). Besides IgG, Protein A also interacts with IgM and IgA.

Amongst human VH subclasses, VH3 is the only subclass to bind Protein A (Graille M *et al.*, (2000) Proc. Natl. Acad. Sci. USA 97(10): 5399-5404), and all five domains of Protein A are known to bind this variable domain subclass (Jansson B *et al.*, (1998) FEMS Immunol. Med. Microbiol., 20(1): 69-78. VH3 based immunoglobulins or fragments thereof are of major importance to the biotechnology industry. VH3 based molecules have been extensively developed since their ability to bind Protein A facilitates their functional pre-screening, and as

such many synthetic or donor based phage display libraries or transgenic animal technologies used for antibody discovery are based on the VH3 subclass. In addition VH3 based molecules are often selected for their good expression and stability over other known heavy chain variable domain subclasses.

5

The capacity of Protein A to bind antibodies with such high affinity is the driving motivation for its industrial scale use in biologic pharmaceuticals. Protein A used for production of antibodies in bio-pharmaceuticals is usually produced recombinantly in *E. coli* and functions essentially the same as native Protein A (Liu HF *et al.*, (2010) MABs, 2(5): 480-499).

10

Most commonly, recombinant Protein A is bound to a stationary phase chromatography resin for purification of antibodies. Optimal binding occurs at pH8.2, although binding is also good at neutral or physiological conditions (pH 7.0-7.6). Elution is usually achieved through pH shift towards acidic pH (glycine-HCl, pH2.5-3.0). This effectively dissociates most protein-protein and antibody-antigen binding interactions without permanently affecting protein structure. Nevertheless, some antibodies and proteins are damaged by low pH and it is best to neutralize immediately after recovery by addition of 1/10th volume of alkaline buffer such as 1 M Tris-HCl, pH 8.0 to minimize the duration of time in the low-pH condition.

15

20

There are various commercially available Protein A chromatography resins. The main differences between these media are the support matrix type, Protein A ligand modification, pore size and particle size. The differences in these factors give rise to differences in compressibility, chemical and physical robustness, diffusion resistance and binding capacity of the adsorbents (Hober S *et al.*, (2007), *supra*). Examples of Protein A chromatography resins include but are not limited to the MabSelect SuRe™ Protein A resin and MabSelect™ Protein A resin from GE Healthcare as used in examples.

25

30

The term “chromatography” refers to protein liquid chromatography and includes fast protein liquid chromatography (FPLC) which is a form of liquid chromatography that is often used to analyze or purify mixtures of proteins. As in other forms of chromatography, separation is possible because the different components of a mixture have different affinities for two materials, a moving fluid (the mobile phase) which passes through a porous solid (the stationary phase). In FPLC, the mobile phase is an aqueous solution, or “buffer”. The buffer flow rate can be operated under gravity flow or controlled by a positive-displacement pump

which is normally kept at a constant rate, while the composition of the buffer can be varied by drawing fluids in different proportions from two or more external reservoirs. The stationary phase is a resin composed of beads, usually of cross-linked agarose, packed into a cylindrical glass or plastic column. FPLC resins are available in a wide range of bead sizes and surface ligands depending on the application.

The process of “affinity chromatography” involves the use of an affinity reagent as ligands which are cross-linked to the stationary phase and that have binding affinity to specific molecules or a class of molecules. Ligands can be bio-molecules, like protein ligands or can be synthetic molecules. Both types of ligand tend to have good specificity. The most commonly used protein ligand in production is the affinity reagent Protein A. In affinity chromatography when the solution (for example a crude cell supernatant containing a protein of interest) is loaded onto the column the target protein is usually adsorbed while allowing contaminants (other proteins, lipids, carbohydrates, DNA, pigments, etc.) to pass through the column. The adsorbent itself is normally packed in a chromatography column; though the adsorption stage can be performed by using the adsorbent as a stirred slurry in batch binding mode. The next stage after adsorption is the wash stage, in which the adsorbent is washed to remove residual contaminants. The bound protein is then eluted in a semi-pure or pure form. Elution is normally achieved by changing the buffer or salt composition so that the protein can no longer interact with the immobilized ligand and is released. In some instances the protein of interest may not bind the affinity resin and affinity chromatography is directed at binding unwanted contaminants and the unbound fraction is therefore collected to isolate the protein of interest. Affinity chromatography can be performed in a fixed bed or a fluidised bed.

The term “gradient mode chromatography” refers to a chromatography method wherein the proportion of the “elution” buffer (buffer B) is increased from 0% to 100% in a gradual or stepwise manner.

The terms “capture-elution mode chromatography” or “capture-elution purification mode” or “capture-elution purification” refers to a chromatography method wherein the proportion of the “elution” buffer (buffer B) is not increased from 0% to 100% in a gradual or stepwise

manner but rather directly applied at a 100% after capture and optionally a wash step with running buffer (buffer A).

Development of hetero-dimeric immunoglobulins targeting CD3 and CD38

5 The present invention provides an epitope binding region that binds the CD3 protein complex comprising the heavy and light chain CDRs as described *supra* and further comprising a heavy chain variable framework region that is the product of or derived from human gene IGHV3-23*04 (SEQ ID NO: 37). The heavy chain variable framework region comprises at least one amino acid modification from the corresponding framework region of the heavy
10 chain variable region of the corresponding murine antibody OKT3 comprising the amino acid sequence of SEQ ID NO: 38. Preferably the amino acid modification is an amino acid substitution. Typically, no more than seven, preferably no more than six, preferably no more than five, preferably no more than four, more preferably no more than three, even more preferably no more than two, most preferably no more than one amino acid modifications are
15 performed within a framework region. In some embodiments the present disclosure provides an epitope binding region that binds to the CD3 protein complex, wherein the amino acid modification of the framework regions of the heavy chain variable region comprise an amino acid substitution at amino acid position selected from the group consisting of: 34, 48, 49, 58, 69, 71 and 73 and wherein the amino acid position of each group member is indicated
20 according to the Kabat numbering. Preferably, amino acid substitutions of the framework regions of the heavy chain variable region are selected from the group consisting of: I34M, V48I, A49G, R58N, R58Y, I69L, A71T and T73K. Preferred amino acid substitution of the framework regions of the heavy chain variable region are at amino acid positions selected from the group consisting of 34, 49 and 71. More preferred amino acid substitutions of the
25 framework regions of the heavy chain variable region are selected from the group consisting of I34M, A49G and A71T.

In a further aspect, the epitope binding region of the first polypeptide that binds the CD3 protein complex comprises a light chain variable framework region that is the product of or
30 derived from a human gene selected from the group consisting of: IGKV1-39*01 (SEQ ID NO: 39) and IGKV3-20*01 (SEQ ID NO: 40). The light chain variable framework region comprises at least one amino acid modification from the corresponding framework region of the light chain variable region of the corresponding murine antibody OKT3 comprising the

amino acid sequence of SEQ ID NO: 41. Preferably the amino acid modification is an amino acid substitution. Typically, no more than eight, preferably no more than seven, preferably no more than six, preferably no more than five, preferably no more than four, more preferably no more than three, even more preferably no more than two, most preferably no more than one amino acid modifications are performed within a framework region. In some embodiments the present disclosure provides an epitope binding region that binds to the CD3 protein complex, wherein the amino acid modification of the framework regions of the light chain variable region sequence comprises an amino acid substitution at amino acid position selected from the group consisting of: 4, 33, 34, 46, 47, 66, 71 and 96. Preferably, amino acid substitutions of the framework regions of the light chain variable region are selected from the group consisting of: M4L, V33M, A34N, L46R, L47W, R66G, F71Y and P96F. Preferred amino acid substitution of the framework regions of the light chain variable region are at amino acid positions selected from the group consisting of 4, 46 and 47. More preferred amino acid substitutions of the framework regions of the light chain variable region are selected from the group consisting of M4L, L46R, L47W and F71Y. In some embodiments the epitope binding region of the first polypeptide that binds to the CD3 protein complex may comprise amino acid modifications of the framework regions of the heavy chain variable region sequence as set out above and amino acid modifications of the framework regions of the light chain variable region sequence as set out above.

The present disclosure also provides an antibody or fragment thereof that binds to the CD3 protein complex that comprises a heavy chain sequence selected from the group consisting of SEQ ID NOs: 79 to 90, 91-95 and 64, preferably selected consisting of SEQ ID NO: 64. The present disclosure also provides an antibody or fragment thereof that binds to the CD3 protein complex that comprises a light chain sequence selected from the group consisting of SEQ ID NOs: 96 to 104, 105 to 126, 127, 128 and 129 preferably consisting of SEQ ID NO: 127.

Given that each of these heavy and light chain variable region sequences can bind to the CD3 protein complex, the heavy and light chain variable region sequences can be “mixed and matched” to create anti-CD3 binding molecules of the invention. CD3 binding of such “mixed and matched” antibodies can be tested using the binding assays described e.g. in the Examples.

Engineering of the immunoglobulin constant region to promote hetero-dimer formation over homo-dimer formation

Methods to produce hetero-dimeric immunoglobulins are known in the art and one of the simplest methods relies on expressing the two distinct immunoglobulin chains in a single cell (WO95/33844, Lindhofer H & Thierfelder S). Without engineering, this straightforward method is limited by the formation of homo-dimeric species over the hetero-dimer of interest (Kufer P *et al.*, (2004) Trends Biotechnol., 22(5): 238-244). When using complementary technologies that will enhance heavy chain hetero-dimerization (Merchant AM *et al.*, (1998) Nat. Biotechnol., 16(7): 677-681), greater hetero-dimer production can be achieved but still results in the production of a significant amount of undesirable homo-dimers (Jackman J *et al.*, (2010) J Biol Chem., 285(27):20850-9, Klein C *et al.*, (2012) MAbs, 4(6):653-63). The present invention therefore utilises the BEAT[®] technology described method (PCT publication No: WO2012/131555), which is based on a unique concept of bio-mimicry that exhibit superior hetero-dimerisation over prior art methods. The BEAT technology is based on an interface exchange between naturally occurring homo or hetero-dimeric immunoglobulin domain pairs to create new hetero-dimers which can be used as building blocks for Fc-based bispecific antibodies.

In one aspect, the present invention provides a hetero-dimeric immunoglobulin or fragment thereof comprising first and second polypeptides comprising an engineered immunoglobulin constant region with a modified CH3 domain having a protein-protein interface, wherein the protein-protein interface of the first polypeptide comprises an amino acid substitution at a position selected from the group consisting of: 3, 5, 7, 20, 22, 26, 27, 79, 81, 84, 84.2, 85.1, 86, 88 and 90 (IMGT[®] numbering), and wherein the protein-protein interface of the second polypeptide comprises an amino acid substitution at position 84.4 and at a position selected from the group consisting of 3, 5, 7, 20, 22, 26, 27, 79, 81, 84, 84.2, 85.1, 86, 88 and 90 (IMGT[®] numbering).

In a further embodiment, the present invention provides a hetero-dimeric immunoglobulin or fragment thereof, wherein the first and second polypeptides comprise an engineered immunoglobulin constant region with a modified CH3 domain having a protein-protein interface, wherein the protein-protein interface of the first polypeptide comprises an amino acid substitution at position 88 and at a position selected from the group consisting of: 3, 5, 7,

20, 22, 26, 27, 79, 81, 84, 84.2, 85.1, 86 and 90 (IMGT[®] numbering), and wherein the protein-protein interface of the second polypeptide comprises an amino acid substitution at position 85.1 and/or 86 and at a position selected from the group consisting of 3, 5, 7, 20, 22, 26, 27, 79, 81, 84, 84.2, 84.4, 88 and 90 (IMGT[®] numbering), wherein the amino acid residue substituted at position 88 in the first engineered immunoglobulin constant region is interacting with the amino acid residue substituted at position 85.1 and/or 86 in the second engineered immunoglobulin constant region, wherein the amino acid position of each group member is indicated according to the IMGT[®] numbering.

Preferably the amino acid residue which is substituted in the protein-protein interface of the first engineered immunoglobulin constant region at position 88 is 88W and conservative amino acid substitutions thereof, wherein the amino acid position is indicated according to IMGT[®] numbering. More preferably, the amino acid residue which is substituted in the protein-protein interface of the first engineered immunoglobulin constant region at position 88 is 88W and wherein the further amino acid residue substituted in the protein-protein interface of the first engineered immunoglobulin constant region is selected from the group consisting of: 3A, 20V, 20T, 20A, 20N, 20Q, 20E, 20S, 20K, 20W, 22A, 22G, 22T, 22L, 22I, 22V, 26R, 26Q, 26T, 26K, 26V, 26S, 26N, 26E, 79Y, 85.1T, 85.1M, 85.1A, 85.1S, 85.1R, 85.1H, 85.1K, 85.1F, 85.1C, 85.1N, 85.1W, 86S, 86I, 86T, 86H, 86Q, 86V, 86W, 86Y, 86F and 90N, wherein the amino acid position is indicated according to the IMGT[®] numbering.

Preferably the amino acid residue which is substituted at position 85 and 86 in the protein-protein interface of the second engineered immunoglobulin constant region is selected from the group consisting of: 85.1A, 85.1S, 85.1C and 86S and conservative amino acid substitutions thereof (IMGT[®] numbering). More preferably the amino acid residue which is substituted in the protein-protein interface of the second engineered immunoglobulin constant region is selected from the group consisting of: 85.1A, 85.1S, 85.1C and 86S and wherein the further amino acid residue substituted in the protein-protein interface of the second engineered immunoglobulin constant region is selected from the group consisting of: 3E, 5A, 7F, 20T, 22V, 26T, 81D, 84L, 84.2E, 88R and 90R and conservative amino acid substitutions thereof (IMGT[®] numbering).

In a preferred embodiment the amino acid residue which is substituted in the protein-protein interface of the first engineered immunoglobulin constant region at position 88 is 88W and wherein the further amino acid residue substituted in the protein-protein interface of the first engineered immunoglobulin constant region is: 3A, 20K, 22V, 26T, 79Y, 85.1S, 86V and
5 90N and, wherein the amino acid residues which are substituted in the protein-protein interface of the second engineered immunoglobulin constant region at positions 85.1 and 86 are 85.1A, 85.1S or 85.1A and 86S and wherein the further amino acid residue substituted in the protein-protein interface of the second engineered immunoglobulin constant region is: 3E, 5A, 7F, 20T, 22V, 26T, 81D, 84L, 84.2E, 84.4Q, 88R and 90R (IMGT[®] numbering).

10 In an alternative embodiment, the present invention provides a hetero-dimeric immunoglobulin or fragment thereof, wherein the first and second polypeptides comprise an engineered immunoglobulin constant region with a modified CH3 domain having a protein-protein interface, wherein the protein-protein interface of the first polypeptide comprises an
15 amino acid substitution at position 20, and at a position selected from the group consisting of: 3, 5, 7, 22, 26, 27, 79, 81, 84, 84.2, 85.1, 86, 88 and 90 and, wherein the protein-protein interface of the second polypeptide comprises an amino acid substitution at position 26 and at a position selected from the group consisting of: 3, 22, 27, 79, 81, 84, 85.1, 86, and 88, wherein the amino acid residue substituted at position 20 in the first engineered
20 immunoglobulin constant region is interacting with the amino acid residue substituted at position 26 in the second engineered immunoglobulin constant region, wherein the amino acid position of each group member is indicated according to the IMGT[®] numbering.

25 Preferably the amino acid residues which are substituted in the protein-protein interface of the first engineered immunoglobulin chain comprise the amino acid residues at positions 20 and 22, and optionally a further amino acid residue at a position selected from the group consisting of: 3, 5, 7, 26, 27, 79, 81, 84, 84.2, 84.4, 85.1, 86, 88 and 90 and, wherein the amino acid residues which are substituted in the protein-protein interface of the second
30 engineered immunoglobulin chain comprise the amino acid residues at positions 26 and at a further position selected from the group consisting of: 3, 5, 7, 20, 22, 27, 79, 81, 84, 84.2, 84.4, 85.1, 86, 88 and 90, wherein the amino acid position of each group member is indicated according to the IMGT[®] numbering. Preferably the amino acid residues which are substituted

in the protein-protein interface of the first engineered immunoglobulin chain comprise the amino acid residues at positions 20 and 22, and optionally a further amino acid residue at a position selected from the group consisting of: 3, 5, 7, 26, 27, 79, 81, 84, 84.2, 84.4, 85.1, 86, 88 and 90 and, wherein the amino acid residues which are substituted in the protein-protein interface of the second engineered immunoglobulin chain comprise the amino acid residues at positions 26 and 86 and optionally at a further position selected from the group consisting of 3, 5, 7, 20, 22, 27, 79, 81, 84, 84.2, 84.4, 85.1, 88 and 90, wherein the amino acid position of each group member is indicated according to the IMGT[®] numbering.

More preferably the amino acid residue which is substituted at position 20 in the protein-protein interface of the first engineered immunoglobulin constant region is selected from the group consisting of 20V, 20T, 20A, 20N, 20Q, 20K, 20S, 20W and 20E and wherein the further amino acid residue substituted in the protein-protein interface of the first engineered immunoglobulin constant region is selected from the group consisting of 3A, 22A, 22G, 22L, 22I, 22V, 22T, 26K, 26R, 26Q, 26T, 26V, 26S, 26N, 26E, 79Y, 85.1W, 85.1F, 85.1T, 85.1M, 85.1A, 85.1S, 85.1R, 85.1H, 85.1K, 85.1C, 85.1N, 86W, 86Y, 86S, 86I, 86H, 86Q, 86V, 86T, 86F, 88Q, 88L, 88V, 88R, 88E, 88T, 88I, 88Y, 88K, 88W and 90N, and wherein the amino acid residue which is substituted at position 26 in the protein-protein interface of the second engineered immunoglobulin constant region is selected from the group consisting of 26T and 26E and conservative amino acid substitutions thereof, wherein the amino acid position is indicated according to the IMGT[®] numbering.

In a most preferred embodiment the amino acid residue which is substituted in the protein-protein interface of the first engineered immunoglobulin constant region at position 20 is 20K and wherein the further amino acid residue substituted in the protein-protein interface of the first engineered immunoglobulin constant region is 3A, 22V, 26T, 79Y, 85.1S, 86V, 88W and 90N and, wherein the amino acid residues which are substituted in the protein-protein interface of the second engineered immunoglobulin constant region at position 26 is 26T and wherein the further amino acid residue substituted in the protein-protein interface of the second engineered immunoglobulin constant region is 3E, 5A, 7F, 20T, 22V, 81D, 84L, 84.2E, 84.4Q, 85.1C/S/A, 86S, 88R and 90R (IMGT[®] numbering).

Examples

Example 1. Materials and Methods***Construction of expression vectors for transient mammalian cell expression***

cDNAs encoding the different polypeptide chains in part or in full were first gene synthesized
5 by GENEART AG (Regensburg, Germany) and modified using standard molecular biology
techniques. PCR products were digested with appropriate DNA restriction enzymes, purified
and ligated in a modified pcDNA3.1 plasmid (Invitrogen AG, Zug, Switzerland) carrying a
CMV promoter and a bovine hormone poly-adenylation (poly(A)) previously digested with
10 the same DNA restriction enzymes. All polypeptide chains were independently ligated in this
expression vector where secretion was driven by the murine VJ2C leader peptide.

Expression of recombinant proteins

Antibodies, ScFv-Fc fusion proteins, BEAT antibodies and antigens were expressed as
described below unless otherwise indicated. For transient expression, equal quantities of each
15 engineered chains vectors were co-transfected into suspension-adapted HEK293-EBNA cells
(ATCC-LGL standards, Teddington, UK; Cat. No: CRL-10852) using Polyethyleneimine
(PEI; Sigma, Buchs, Switzerland). Typically, 100ml of cells in suspension at a density of 0.8-
1.2 million cells per ml is transfected with a DNA-PEI mixture. When recombinant
expression vectors encoding each engineered chain genes are introduced into the host cells,
20 the immunoglobulin construct is produced by further culturing the cells for a period of 4 to 5
days to allow for secretion into the culture medium (EX-CELL 293, HEK293-serum-free
medium (Sigma), supplemented with 0.1% pluronic acid, 4mM glutamine and 0.25µg/ml
geneticin). Cell-free culture supernatants containing the secreted immunoglobulins were
prepared by centrifugation followed by sterile filtration and used for further analysis.

Differential Protein A affinity chromatography

Post production, cell-free supernatants were loaded onto a 1ml HiTrap™ MabSelect SuRe™
Protein A column pre-equilibrated in 0.2M phosphate citrate buffer pH 6.0 and operated on an
30 ÄKTApurifier™ chromatography system (both from GE Healthcare Europe GmbH; column
Cat. No: 11-0034-93) at a flow rate of 1ml/min. Running buffer was 0.2 M phosphate citrate
buffer pH 6. Elution of the hetero-dimer of interest was performed using 20 mM sodium
citrate buffer pH 4 whilst homo-dimeric species were eluted with 0.1 M glycine, pH3.0.

Elution was followed by OD reading at 280 nm; fraction containing the hetero-dimer of interest were pooled and neutralized with 0.1 volume of 1M Tris pH 8.0 (Sigma).

Supernatant, flow through and elution fractions were analysed under non-reduced conditions by SDS-PAGE (NuPAGE Bis-Tris 4-12% acrylamide, Invitrogen AG, Basel, Switzerland).

5

Differential Scanning Calorimetry (DSC)

The thermal stabilities of antibodies were compared using calorimetric measurements. Calorimetric measurements were carried out on a VP-DSC differential scanning microcalorimeter (MicroCal-GE Healthcare Europe GmbH, Glattbrugg, Switzerland). The cell volume was 0.128 ml, the heating rate was 1°C/min and the excess pressure was kept at 64 p.s.i. All protein fragments were used at a concentration of 1-0.5 mg/ml in PBS (pH 7.4). The molar heat capacity of each protein was estimated by comparison with duplicate samples containing identical buffer from which the protein had been omitted. The partial molar heat capacities and melting curves were analysed using standard procedures. Thermograms were baseline corrected and concentration normalized before being further analysed using a Non-Two State model in the software Origin v7.0.

10

15

20

25

The expected melting profiles for the human IgG subclasses are known (Garber E & Demarest SJ (2007) Biochem Biophys Res Commun, 355(3): 751-7) and all profiles have been shown to contain three unfolding transitions corresponding to the independent unfolding of the CH2, CH3 and FAB domains. Of the four human IgG subclasses, IGHG1 has the most stable CH3 domain (~85°C); while other subclasses CH3 domains are less stable, although none are known to melt below 70°C. Similarly, all subclasses are known to have a melting temperature of ~70°C for the CH2 domain.

Purity assessment by capillary gel electrophoresis

Non-reduced sample preparation

40 µg of desalted protein sample was buffered in SDS sample buffer (Beckman Coulter International S.A., Nyon, Switzerland; IgG Purity Kit, Cat. No: A10663) containing 5 mM Iodoacetamide (Sigma). A 10-kDa internal standard was added to the samples. The sample-mixtures were heated at 70°C for 10 min.

30

Capillary gel electrophoresis

Following sample preparation, samples were run on a ProteomeLab PA 800 (Beckman Coulter International S.A., Nyon, Switzerland) fitted with a photodiode array detector (DAD) set at 220 nm. Bare-fused silica capillaries of 50 μm ID $\times$ 30.2 cm (20.2 cm effective length to detector) were used as separation medium. Sample injection and separation were performed at constant voltages of 5 and 15 kV, respectively, with reverse polarity in SDS-molecular weight gel buffer. The data were recorded at a rate of 2 Hz and current was stable during separation. Capillary and samples were thermo-stated at 25°C.

Affinity measurements by SPR

SPR analysis was used to measure the association and dissociation rate constants for the binding kinetics of the different antibodies (murine and humanized antibodies). The binding kinetics of antibodies were measured on a BIAcore 2000 instrument (BIAcore-GE Healthcare Europe GmbH, Glattbrugg, Switzerland) at room temperature and analysed with the BiaEvaluation software (version 4.1, BIAcore-GE Healthcare Europe GmbH).

Measurements were performed on CM5 sensor chips (GE Healthcare Europe GmbH, Cat. No: BR-1000-14) individually coupled with the ligand of interest using a commercial amine coupling kit (GE Healthcare Europe GmbH, Cat. No: BR-1000-50). Protein G ligand was from Pierce (Thermo Fisher Scientific-Perbio Science S.A., Lausanne, Switzerland, Cat. No: 21193).

Data (sensorgram: fc2-fc1) were fitted with a 1:1 Langmuir model with or without mass transfer as indicated. In capture experiments, to account for the experimental variations in at the beginning of each measurement, the Rmax value was set to local in all fits. Dissociation times were of at least 350 seconds. Measurements were performed in triplicate and included zero-concentration samples for referencing. Both Chi2 and residual values were used to evaluate the quality of a fit between the experimental data and individual binding models.

Affinity measurements on HPB-ALL cells by FACS

HPB-ALL cells (DSMZ, Braunschweig, Germany, Cat. No: ACC483) were used as CD3 positive cell line for FACS staining. HPB-ALL were maintained in RPMI 1640 supplemented with 10% FCS and 100 U/ml Penicillin and 100ug/ml streptomycin. 100 μl dilution series of the chimeric OKT3 antibody and humanized variants were incubated with 4×10^5 HPB-all

cells in PBS supplemented with 1% BSA and 0.1% Sodium Azide (referred as FACS buffer) for 45 min on ice. An irrelevant human IgG1 was used as isotype control and the chimeric OKT3 antibody as positive control. After washing, cells were incubated with a 1/200 dilution of anti-Human Fc-PE (EBioscience, Vienna, Austria) for 45 min on ice. Cells were then washed again and resuspended in 200ul FACS buffer. The relative mean fluorescence of each sample was measured on FACSCalibur (BD Biosciences, Allschwil, Switzerland) Results are summarized in FIG. 9 as the relative staining of HBP-ALL compared to the chimeric OKT3 antibody.

Cell-lines for in vitro assays

Human CD38 positive cell lines

Human cells expressing CD38 antigen have been described in PCT Publication Nos: WO2005103083, WO2008047242, WO2011154453 and WO2012092612. CD38 positive human cell lines as used herein were as follows:

Stable recombinant CHO[CD38] cells

A gene coding for human CD38 was ordered at Source Biosciences (Berlin, Germany, Cat.-No.: IRAU37D11, 4309086). Human CD38 was amplified using primers adding a kozak sequence, a start codon followed by a signal peptide (murine V leader) to the 5' end and a NheI restriction site to the 3' end. The amplicon was cut using NheI and HindIII and cloned into the expression cassette of pT1, a pcDNA3.1 (Invitrogen AG) derived vector developed in-house. The expression cassette of pT1 links the expression of the gene of interest with expression of GFP and PAC (the gene for puromycin resistance) using two IRES (internal ribosome entry sites) on a polycistronic mRNA. A midiprep of the plasmid was prepared and the cloned CD38 open reading frame was confirmed by DNA sequencing. Suspension CHO-S cells (Invitrogen AG) were transfected using polyethyleneimine (JetPEI[®], Polyplus-transfection, Illkirch, France) in 50 ml bioreactor format (TubeSpin 50 bioreactors, TPP, Trasadingen, Switzerland). For this purpose, exponential growing cells were seeded in OptiMEM medium (Invitrogen AG, Cat. No.: 31985-047). A JetPEI[®]:DNA complex was added to the cells. After 5 h incubation of the cells with the JetPEI[®]:DNA complex at 37°C under shaking (200 RPM) for endocytosis, one volume of culture medium PowerCHO2 (Lonza, distributor RUWAG Lifescience, Bettlach, Switzerland, Cat. No:BE12-771Q) supplemented with 4 mM Gln was added to the cell suspension. The cells were then incubated

on a shaken platform at 37°C, 5% CO₂ and 80% humidity. One day after transfection the cells were seeded in 96 well plates at different concentrations in selective medium containing puromycin (Sigma, Cat. No: P8833-25mg). After approximately 14 days of selection under static conditions, 46 high GFP expressing cell pools were expanded as suspension cultures using TubeSpin 50 bioreactors. Once successfully adapted to suspension, the cells were analysed for CD38 by FACS. Stable CHO[CD38] clones with a homogenous CD38 staining profile were selected and used herein.

Other CD38 positive cell lines included:

NCI-H929 (ATCC-LGL standards; Cat. No: CRL-9068).

Namalwa (ATCC-LGL standards; Cat. No: CRL-1432)

U266 (ATCC-LGL standards; Cat. No: TIB-196)

RPMI 8226 (ATCC-LGL standards; Cat. No: CCL-155)

Culture conditions: RPMI 1640 medium supplemented with 10% heat-inactivated FBS, 1% penicillin-streptomycin (Invitrogen AG) and 1% GlutaMAX-1 (Invitrogen AG)

Raji (ATCC-LGL standards; Cat. No: CCL-86)

Daudi (ATCC-LGL standards; Cat. No: CCL-213)

Recombinant target antigens

Human CD3 gamma-epsilon-Fc fusion protein

A cDNA encoding the human CD3 gamma extracellular region (UniProt accession No: P09693 residues 23-103 (SEQ ID NO: 184); UniProt Consortium (2013) Nucleic Acids Res., 41(Database issue): D43-7; <http://www.uniprot.org/>) fused to the human CD3 epsilon extracellular region (UniProt accession No: P07766, residues 22-118 (SEQ ID NO: 185)) by a 26-residue peptide linker (sequence: GSADDAKKDAAKKDDAKKDDAKKDGS; SEQ ID NO: 169) was first synthesized by GENEART AG (Regensburg, Germany). This synthetic gene was fused to a human IgG1 Fc portion using standard overlap PCR techniques and a human IgG1 Fc cDNA template also obtained from Geneart AG. The resulting cDNA was cloned in the modified pcDNA3.1 plasmid mentioned above.

For transient expression of the CD3 gamma-epsilon-Fc protein (SEQ ID NO: 170), the recombinant vector was transfected into suspension-adapted HEK-EBNA cells (ATCC-CRL-10852) using Polyethyleneimine (PEI) as described above. The CD3 gamma-epsilon-Fc construct was then purified from cell-free supernatant using recombinant Streamline rProtein

A media (GE Healthcare Europe GmbH, Glattbrugg, Switzerland) and used for further analysis.

Human and cynomolgus monkey CD3 epsilon 1-26_Fc fusion proteins

5 A cDNA encoding the human CD3 epsilon peptide 1-26 (UniProt accession No: P07766, amino acids 23-48, SEQ ID NO: 171) and a cDNA encoding the cynomolgus CD3 epsilon peptide 1-26 (UniProt accession No: Q95LI5, amino acids 22-47, SEQ ID NO: 172) were PCR amplified from synthetic cDNAs obtained from GENEART A.G. for the human and cynomolgus monkey CD3 epsilon extracellular regions, respectively. The amplified products
10 were subsequently fused to a human IgG1 Fc portion using standard overlap PCR techniques. The human IgG1 Fc cDNA template was obtained from Geneart AG. The resulting cDNA were cloned in the modified pcDNA3.1 plasmid mentioned above.

For transient expression of human and cynomolgus CD3 epsilon constructs (SEQ ID NO: 173
15 and 174, respectively), the recombinant vectors were transfected into suspension-adapted HEK-EBNA cells (ATCC-CRL-10852) using Polyethyleneimine (PEI) as described above. The CD3 epsilon fusion constructs were then purified from cell-free supernatant using recombinant Streamline rProtein A media (GE Healthcare Europe GmbH, Glattbrugg, Switzerland) and used for further analysis. These two fusion proteins are referred herein as the
20 human and cynomolgus monkey CD3 epsilon 1-26_Fc fusion proteins.

Human and cynomolgus monkey CD38 extracellular regions

A cDNA for human CD38 was obtained from Source Biosciences (Erwin-Negelein-Haus, Germany, Cat. No.: IRAU37D11, 4309086), its extracellular region (UniProt accession No:
25 P28907 residues 43-300) was PCR amplified and cloned into an in-house expression vector derived from pcDNA3.1 (Invitrogen AG). This expression vector encompassed a kozak sequence and a start codon followed by the murine VJ2C leader peptide to the 5' end and a 6-His-tag to the 3' end of its multiple cloning site. The soluble extracellular region of human CD38 fused to a 6-His-tag (SEQ ID NO: 175) was expressed and purified as follows: one
30 volume of RPMI 1640 medium (PAA Laboratories, Cat. No: E15-039) containing HEK cells, 0.1% pluronic acid (Invitrogen AG), expression vector and polyethylenimine (JetPEI®, Polyplus-transfection, Illkirch, France) was incubated in a shaker flask at 37°C, 5% CO₂ and 80% humidity. One volume of ExCell293 medium supplemented with 6 mM glutamine was

added to the mixture after 4 hours and incubation continued further for a total of 5 days. Post production, cell-free supernatant was prepared by centrifugation and filtrated using 0.2 µm filters, pH was adjusted at 7.4 (4°C) using Tris 1 M pH 8.7. Ni-Sepharose Excell beads (GE Healthcare, Cat. No: 17-3712-03) were added to the solution and incubated overnight at 4°C under agitation. The solution was loaded on an Econo-Column (Bio-Rad Laboratories AG, Reinach, Switzerland, Cat. No: 737-4252) for gravity-flow purification. The beads were washed in PBS (2x), 20 mM imidazole and the protein was eluted in PBS, 500 mM Imidazole. Eluted fractions were pooled and buffer exchanged for PBS with two dialysis steps at 4°C. The purified human CD38 extracellular region was filtrated using 0.22 µm syringe filters.

Using the methods as described above the soluble extracellular region of cynomolgus monkey CD38 antigen fused to a 6-His-tag (SEQ ID NO: 176) was cloned, expressed and purified.

In vitro T cell redirection killing assay

Preparation of peripheral blood mononuclear cells

To produce peripheral blood mononuclear cells (PBMCs), blood filters containing human leukocytes were collected from the Blood Collection Centre in La Chaux-de-Fonds, Switzerland (Centre de Transfusion Sanguine et Laboratoire de Sérologie, rue Sophie-Mairet 29, CH-2300). Cells were removed from the filters by back-flushing with 60 ml of PBS containing 10 U/ml of liquemin (Drossapharm AG, Lucern, Switzerland). PBMCs were then purified with 50 mL Blood-Sep-Filter Tubes (Brunschwig, Basel, Switzerland) following manufacturer's instructions. Tubes were centrifuged for 20 min at 800g at room temperature (without brake) and the cells were collected from the interface. Cells were washed 3x with Roswell Park Memorial Institute (RPMI, PAA Laboratories, Pasching, Austria) medium without FBS or phosphate buffered Saline (PBS). PBMCs were resuspended at 10e6 cells/mL in RDL medium (RPMI supplemented with 10% heat inactivated Fetal bovine serum (FBS) and penicillin/streptomycin) and were cultured overnight at 37°C in a 5% CO₂ incubator prior to the assay.

T cell preparations

T cell purification was performed directly after the PBMC isolation using pan-T cell isolation kit II (Mytenyi Biotec GmbH, Bergisch Gladbach, Germany, Cat. No: 130-091-156)

following manufacturer's instructions. After purification, T cells were resuspended at 10e6 cells/mL in RDL medium and cultured overnight at 37°C in a 5% CO₂ incubator prior assay.

Assay readouts

- 5 Two different readouts which gave highly comparable results were used to quantify the redirected killing.

A flow cytometry method, referred herein as RDL-FACS method, based on fluorescence-cytometry as described in Schlereth B *et al.* ((2005) Cancer Res, 65: 2882-2889), Moore PA *et al.* ((2011) Blood, 117(17): 4542-51) and Friedrich M *et al.* ((2012) Mol Cancer Ther, 11: 2664-2673). Target cells were harvested, counted, washed once and resuspended at 5x10e6 cells/mL in PBS+1 µM Carboxyfluorescein succinimidyl ester (CFSE, Sigma). Cells were incubated 15 min at 37°C with gentle agitation every 5 min. CFSE loaded cells were washed 3x with RDL medium and resuspended at 2x10e5 cells/mL in RDL medium. PBMCs were harvested, counted and resuspended at 2x10e6 cells/mL in RDL medium. Antibodies serial dilutions (3x solutions) were prepared in RDL medium. Target cells (50 µl/well), T cells (50 µl/well) and 3x antibody solutions (50 µl/well) were distributed in flat-bottom 96-well plate (TPP, Trasadingen, Switzerland). The effector: target ratio was 10:1. The plates were incubated for 48h in a 5% CO₂ incubator at 37°C. After incubation the plates were centrifuged for 3 min at 300g, the supernatants were discarded by flicking the plates. The plates were washed once with 200 µl of PBS, centrifuged again and the PBS was discarded. A pre-warmed solution of accutase (Invitrogen AG) was added and the plates were incubated 10 min at 37°C. The detached adherent cells were resuspended by pipetting up and down after addition of 100 µL of RDL medium. The solution was transferred into a U-bottom 96-well plate (TPP). The U-bottom plates were centrifuged for 3 min at 300 g, the supernatants were discarded and the cells were resuspended in 200µl of cold FACS buffer (PBS + 2% FBS + 10% Versene) supplemented with 7-AAD (Becton Dickinson AG, Allschwil, Switzerland) at a 1/40 dilution. The plates were immediately acquired on a Guava easyCyte™ Flow Cytometer (Millipore AG, Zug, Switzerland). For each well, the absolute number of living target cells was determined by gating on CFSE positive 7ADD negative population using Flowjo® software (Miltenyi Biotec GmbH, Bergisch Gladbach, Germany). The percentage of specific cytotoxicity for each sample was determined using the condition in which only target cells were incubated as baseline. The EC₅₀ values were determined using nonlinear variable

slope regression method with Prism software (GraphPad software, La Jolla, CA, USA). The percentage of specific re-directed lysis (RDL) was calculated by subtracting the percentage of specific cytotoxicity of the condition without antibody to the conditions where a test antibody was added.

5

A cell viability method, referred herein as RDL-MTS method based on a colorimetric method to assess cell viability as described in in Bühler P *et al.* ((2008) Cancer Immunol Immunother, 57: 43–52, Labrijn AF *et al.* ((2013) Proc Natl Acad Sci USA, 110(13): 5145-50) and PCT Publication No: WO2012143524. Target cells were harvested, counted, washed once and resuspended at 2×10^5 cells/ml in RDL medium. PBMCs were harvested, counted and resuspended at 2×10^6 cells/mL in RDL medium. Antibodies serial dilutions (3x solutions) were prepared in RDL medium. Target cells (50 μ l/well), T cells (50 μ l/well) and 3x antibody solutions (50 μ l/well) were distributed in flat-bottom 96-well plate (TPP). The effector: target ratio was 10:1. The plates were incubated for 48 h in a 5% CO₂ incubator at 37°C. After incubation the supernatants were discarded and the plates were washed 3 times with 200 μ L of PBS to remove the PBMCs and 100 μ l of RDL medium was then added to each well. The readout was done using CellTiter 96[®] kit (Promega AG, Dübendorf, Switzerland) according to manufacturer's instructions. Briefly, 10-20 μ l of MTS reagent was added to each well and the plates were incubated 2-6h in a 5% CO₂ incubator at 37°C. The 490 nm absorbance was then read on a BioTek synergy plate reader (BioTek AG, Luzern, Switzerland). The percentage of specific killing was calculated using this formula: Specific killing = $100 \times [(SD - Sp)/(SD - MD)]$. SD is the absorbance measured in spontaneous death condition where target cells were incubated alone. Sp is the absorbance measured in each test condition (target cells + PBMCs + antibody). MD is the absorbance measured in the maximum death condition in which target cells were lysed by 3 freeze and thaw cycles. The percentage of specific redirected lysis (RDL) was calculated by subtracting the percentage specific cytotoxicity of the condition without antibody to the conditions where a test antibody was added. The EC₅₀ values were determined using nonlinear variable slope regression method with Prism software (GraphPad software).

30

Example 2: Antigen binding sites that target the human CD3 antigen, and the CD38 antigen

2.1 Antigen binding sites against the human CD3 antigen

The human CD3 epsilon subunit was selected to drive T cell redirect killing via bispecific engagement.

5 2.1A *Humanized variants of the mouse OKT3 antibody*

The anti-human CD3 epsilon antigen binding site used herein was derived from the mouse OKT3 antibody (Muromonab-CD3, trade name Orthoclone OKT3, marketed by Janssen-Cilag and subsequently discontinued; murine variable heavy chain and light chain domains with SEQ ID NO: 38 and 41, respectively). OKT3 murine variable domains were humanized
10 and formatted as scFv and FAB fragments.

Humanization followed the method described by Jung S & Plückthun A (1997, Protein Eng, 10(8): 959-66) to produce a highly stable humanized variant that would be suitable for both FAB and scFv formatting. The method makes use of the highly stable pair of VH/VL domains found in the Herceptin[®] antibody (rhUMAbHER2, huMAB4D5-8, trastuzumab or trade name Herceptin[®]; US Patent publication No.5,821,337; variable heavy chain and light chain domains with SEQ ID NO: 20 and 21, respectively) and follows the workflow of a humanization process onto fixed frameworks (Almagro JC & Fransson J (2008), Front Biosci, 13: 1619-33). Since the Herceptin[®] antibody is originally derived from the highly stable
15 human families of germline framework VH3 and VK1, germline frameworks from these two families can be equally used as a source of fixed frameworks. Alternatively, the human VK3 germline light chain framework family can be used instead of VK1 as it also has good stability properties (Ewert S *et al.*, (2003) J Mol Biol, 325: 531-553). In addition to mouse antibodies, human antibodies can be engineered using this fixed framework method to
20 improve stability. Preferred is the use of the human germline framework IGHV3-23*04, IGKV1-39*01 and IGKV3-20*01 having SEQ ID NO: 37, 39 and 40, respectively (referenced according to IMGT[®] (the international ImMunoGeneTics information system (Lefranc MP *et al.* (1999) Nucleic Acids Res, 27(1): 209-12; Ruiz M *et al.* (2000) Nucleic Acids Res, 28(1): 219-21; Lefranc MP (2001) Nucleic Acids Res, 29(1): 207-9; Lefranc MP (2003) Nucleic Acids Res, 31(1): 307-10; Lefranc MP *et al.*, (2005) Dev Comp Immunol, 29(3): 185-203; Kaas Q *et al.*, (2007) Briefings in Functional Genomics & Proteomics, 6(4): 253-64; <http://www.imgt.org>).
25
30

To this aim a first humanized antibody was constructed wherein the CDRs in the variable domains of the Herceptin[®] antibody were respectively replaced with the CDRs from the mouse OKT3 antibody and benchmarked against a chimera of the mouse OKT3 antibody (variable heavy chain and light chain with SEQ ID NO: 130 and 131, and referred herein as the chimeric OKT3 antibody).

The prototype antibody (variable heavy chain and light chain with SEQ ID NO: 79 and 96, and abbreviated VH/VL) had increased production levels in transient expression tests and increased FAB stability as measured by differential scanning calorimetry but had no binding to HPB-ALL cells (assessed by median fluorescence intensity in FACS experiments, see Materials and Methods section), a human CD3 epsilon positive T cell tumour line (FIG. 1A).

Based on a 3D model of the first prototype pair of variable domains, a subset of back mutations (from CDR grafted Herceptin[®] prototype to mouse OKT3 sequence) were selected and tested: I34M, V48I, A49G, R58N, R58Y, I69L, A71T and T73K in the variable heavy chain domain and M4L, V33M, A34N, L46R, L47W, R66G, F71Y and P96F in the variable light chain (Kabat numbering). Note that the R58N substitution corresponds to a CDR grafted Herceptin[®] prototype-to-mouse OKT3 mutation while the R58Y substitution corresponds to a CDR grafted Herceptin[®] prototype-to-human IGHV3-23*04 germline substitution. The engineering strategy with regard to the combination of substitutions was based on the complementarity of the different substitutions in terms of their putative influence on CDR regions and/or variable domain packing and/or immunogenicity.

In a first approach, all candidates were formatted as human IgG1 antibodies. Best variants were selected according to expression levels, FAB fragment thermo-stability and ability to bind HPB-ALL cells by FACS. Best humanized variants had the Protein A binding site present within their VH domain abrogated using the G65S or N82aS substitution. This engineering step was needed to further produce safe T cell retargeting BEAT antibodies free of anti-CD3 homo-dimers.

Back mutations in the VH of: I34M, A49G and A71T along with back mutations in the VL of: M4L, L46R, L47W and F71Y restored affinity. Best combinations of variable domains were VH8/VL4, VH8/VL8, VH11/VL4 and VH11/VL8 as these retained most of parental cell

binding (FIG. 1A-C). In addition, combinations VH8/VL8 (variable domains with SEQ ID NO: 131 and 132, respectively) and VH11/VL8 (variable domains with SEQ ID NO: 133 and 132, respectively) had enhanced FAB stability and (+2.8°C and +1.6°C, respectively, FIG. 1D).

5

Finally, best humanized variants were also reformatted as scFv-Fc fusions and transiently expressed. Variants were ranked in terms of their relative affinity, stability, expression levels in transient transfection in this format (FIG. 1E). Best combinations of variable domains in a scFv-Fc fusion format were similar to the combinations identified in an antibody format: VH8-VL4 (scFv fragment with SEQ ID NO: 135) and VH8-VL8 (scFv fragment with SEQ ID NO: 136). Both scFv fragments had good thermal stability with the scFv-Fc fusion format (FIG. 1F).

10

2.1B Humanized variants of the mouse SP34 antibody

15

The mouse antibody SP34 was first described in 1985 (Pessano S *et al.*, (1985) EMBO J, 4(2):337-44). It was produced by a hybridoma obtained from mice immunised with denatured protein extracts from HPB-ALL cells, the antibody has human specificity and cross-reactivity to cynomolgus monkey.

20

Following the methods and work flow described in this example *supra*, humanized VH and VL domains for the murine SP34 antibody having a VH domain with SEQ ID NO: 1 and a VL domain with SEQ ID NO: 2 were engineered via CDR grafting onto the VH3-23 and VK3 germline frameworks, respectively. The resulting VH3 based variable domains can be further abrogated for Protein A binding using the G65S or N82aS substitutions (Kabat numbering) depending on their usage in a BEAT antibody format.

25

To this aim a first humanized antibody was constructed wherein the CDRs in the variable domains of a human antibody having a germline VH3 heavy chain domain and a germline VK3 light chain domain were respectively replaced with the CDRs from the mouse SP34 antibody. The resulting humanized antibody was used a starting point for further affinity improvement and benchmarked against a chimera of the SP34 antibody (heavy chain and light chain with SEQ ID NO: 137 and 138, respectively, and referred herein as the chimeric SP34 antibody).

30

The prototype antibody (variable heavy chain and light chain with SEQ ID NO: 91 and 105, and abbreviated VH1/VL1) had a low binding to human CD3 epsilon 1-26_Fc fusion protein (assessed by SPR, see Materials and Methods section and FIG. 2A).

5

Based on a 3D model of the first prototype pair of variable domains, a subset of substitutions that corresponded to either back mutations between the CDR grafted human germline VH3/VK3 prototype and mouse SP34 sequence (human-to-mouse or mouse-to-human substitutions) or rationally designed amino acid changes was selected. The following changes were made and tested in various combinations: W100eF, and W100eY in the variable heavy chain domain and A2I, S25A, T27A, G27aA, V27cA, T28A, T29A, S30A, N31A, Y32A, E38Q, F44P, G46L, T51A, N52A, K53A, R54A, P56A, L66G, D69T, F87Y, Q89A, W91F, Y92A, S93A, N94A, and Q100G in the variable light chain (Kabat numbering; see FIG. 2A). The engineering strategy with regard to the combination of substitutions was based on the complementarity of the different substitutions in terms of their putative influence on CDR regions and/or variable domain packing and/or immunogenicity and/or impact on transient expression in mammalian cells.

10

15

20

In a first approach, all candidates were formatted as human IgG1 antibodies and later further tested in a scFv-Fc fusion protein format (FIG. 2B) with some variants having the Protein A binding site present within their VH domain abrogated using the G65S. Best humanized candidates were selected according to expression levels and ability to bind the human and cynomolgus monkey CD3 epsilon 1-26_Fc fusion proteins by SPR.

25

Surprisingly, when reformatted as scFv, SP34 chimera and H1L21 lead to a dramatic loss of expression compare to the IgG1 format, (Figure 3). However, scFv-Fc expression was enhanced by the combination of substitutions W100eY in the VH and W91F in the VL (FIG.2B).

30

To further improve the manufacturability of CD3 based bispecifics, humanized SP34 scFv was therefore further engineered. To identify key positions impacting SP34 scFv-Fc expression, an Alanine scan was performed in VL21 CDR. The following changes were made: T27A, G27aA, V27cA, T28A, T29A, S30A, N31A, Y32A, N52A, K53A, R54A,

P56A, L90A, Y92A, S93A, N94A, and L95A. Interestingly, only substitutions at position 29, 30 (VH1/VL26 SEQ NO: 139) and 95 (VH1/VL34 SEQ NO: 140) lead to significant increase in scFv-Fc expression (Figure 4), moreover these mutants were the only ones to retain full binding to human and Cynomologus monkey CD3 epsilon.

5

Based on these results, more substitutions were tested at position 29, 30 and 95 in the light chain.

10

As position 29 and 30 display hyper variability in all the different variable light chain families, these two positions were randomly mutated by site directed PCR. From all the mutants produced, only substitutions T29A, T29E, T29S, S30A and S30D improved significantly transient expression level Figures 5a and 5b while maintaining binding to human CD3 epsilon.

15

In a different approach, L95, which is known in the art as a canonical structure residue, was substituted with residues most frequently found at this very same position in the variable lambda families. The following changes were made L95A, L95G, L95T, L95S, L95D, and L95N. Unexpectedly, only L95G and L95T significantly improved expression while maintaining binding to the target (Figure 5c).

20

From this work, several humanized SP34 based BEAT were designed and tested for expression and binding to CD3 epsilon. Among all these constructs, bispecific containing H5L65 (SEQ ID NO: 69) and H5L67 (SEQ ID NO: 70) scFv, showed a 2 fold increase in expression over the H5L32 scFv based BEAT control, Figure 6.

25

H5L65 and H5L67 were further characterized by DSC either as IgG1 or scFv-Fc and compare to H5L32, H1L21 and SP34 chimera. H5L65 showed superior thermostability as IgG1 but more interestingly an increase of 5 to 2°C compared to other humanized variants in a scFv-Fc format. (Table 1) The *in vivo* stability of the improved antibodies is therefore increased as it also is *in vitro*.

30

	Fab	IgG (Fab)	scFvFc
SP34 Versions	Tm	Tm	Tm
	(°C)	(°C)	(°C)

SP34 chimera	66.9	66.7	N/A
hSP34 H1/L21	75.1	73.1	56.7
hSP34 H5/L32	77.5	75.9	59.7
hSP34 H5/L65	78.1	77.3	61.6
hSP34 H5/L67	77.4	76.5	59.9

Table 1

Preferred combinations of heavy chain and light chain variable domains with regard to antigen binding and recombinant expression were as follows: VH1 (SEQ ID NO: 42) or VH2 (SEQ ID NO: 43) or VH3 (SEQ ID NO: 44) or VH5 (SEQ ID NO: 45) paired with light chains domains VL21 (SEQ ID NO: 46), VL32 (SEQ ID NO: 47), VL65 (SEQ ID NO: 48) and VL67 (SEQ ID NO: 49).

2.2 Antigen binding sites against the human CD38 antigen

CD38 is a type II transmembrane glycoprotein which is normally found on hemopoietic cells and in solid tissues. CD38 is also expressed in a variety of malignant hematological diseases. Bispecific antibodies that would redirect T cells to kill CD38 positive cancer cells will be useful to treat a variety of malignant hematological diseases, including multiple myeloma, B-cell chronic lymphocytic leukaemia, B-cell acute lymphocytic leukaemia, Waldenström's macroglobulinemia, primary systemic amyloidosis, mantle-cell lymphoma, pro-lymphocytic/myelocytic leukaemia, acute myeloid leukaemia, chronic myeloid leukaemia, follicular lymphoma, NK-cell leukaemia and plasma-cell leukaemia. Several anti-CD38 antibodies have been described as research reagents or therapeutic candidates (PCT Publication No: WO2006099875). Amongst the best characterized anti-human CD38 antibodies are OKT-10 and HB-7 mouse hybridomas (Hoshino S *et al.*, (1997) J Immunol, 158(2): 741-7).

In a first approach, anti-human CD38 antigen binding sites can be derived from mouse hybridomas OKT10 (variable heavy chain and light chain with SEQ ID NO: 50 and 53, respectively) or HB-7 (variable heavy chain and light chain with SEQ ID NO: 51 and 54, respectively) and humanized versions thereof which can be further formatted as a FAB or scFv fragments. Following the methods and work flow described in Example 2.1, humanized VH and VL domains for the HB-7 hybridoma are can engineered via CDR grafting onto the VH3-23 and VK1 germline frameworks, respectively.

In a second approach, following the so-called best-fit humanization method described by Almagro JC & Fransson J (Front Biosci, (2008) 13: 1619-33), best-fit humanized VH and VL domains for the HB-7 hybridoma were engineered via CDR grafting onto the human IGHV4-59*03 and IGKV1-NL1*01 germline frameworks, respectively (referenced according to IMGT[®] *supra*). Humanized VH and VL variants with different degree of back mutations were investigated *in silico* and one preferred selection of humanized VH and VL was transiently expressed as a human IgG1 format and referred herein as humanized HB-7 best-fit VH (SEQ ID NO: 56) and VL (SEQ ID NO: 57) domains. The following mouse back mutations were introduced: (VH) S35H, I37V, I48L, V67L, V71K, T73N, F78V, Y91F and (VL): M4L, L48I, Y49S, T69K (Kabat numbering).

The humanized HB-7 best-fit antibody (heavy chain with SEQ ID NO: 141 and light chain with SEQ ID NO: 142) stained CHO[CD38] recombinant cells by FACS (data not shown). The humanized HB-7 best-fit antibody had a binding affinity for the CD38 extracellular region similar to that of the chimeric HB-7 antibody (heavy chain with SEQ ID NO: 143 and light chain with SEQ ID NO: 144) when assayed by SPR (KDs of 3.6 and 2.5 nM, respectively; FIG. 7A (chimeric) and FIG. 7B (humanized)). Surprisingly, the humanized HB-7 best-fit antibody displayed a significant enhancement (+14.6°C) in FAB fragment stability compared to the chimeric HB-7 antibody as judged from calorimetry profiles (76.4°C (chimeric) vs 91.0°C (humanized), FIG. 7F).

In a third approach, mice immunized with the human CD38 extracellular domain and human CD38⁺ cells were used to generate novel hybridoma candidates against human CD38. Methods to generate hybridomas are known and the methods used herein were similar to methods disclosed in PCT Publication No: WO2013008171. The 9G7 mouse antibody

candidate had a high affinity for both human and cynomolgus monkey CD38 (variable heavy chain and light chain with SEQ ID NO: 52 and 55, respectively). This mouse antibody was first humanized according to the methods described in this example *supra*. Using the best-fit approach, the germline VH framework IGHV2-5*09 and VK framework IGKV1-33*01 (referenced according to IMGT[®] *supra*) were selected as a starting point for the humanization process. Post CDR grafting, the first antibody prototype (formatted as a human IgG1 isotype, heavy chain SEQ ID NO: 145 and light chain with SEQ ID NO: 146) exhibited a strong binding to human CD38 only three fold lower than the mouse parental antibody as judged by SPR (chimeric 9G7 antibody with heavy chain SEQ ID NO: 147 and light chain with SEQ ID NO: 148; KD of 0.3 nM and 1 nM for the chimeric 9G7 antibody (data not shown) and first humanized prototype (data not shown), respectively). Affinity improved by two fold upon introduction of the F36Y back mutation in the variable light chain of the antibody (Kabat numbering) (the resulting antibody is referred herein as the humanized 9G7 best-fit antibody with heavy chain SEQ ID NO: 145 and light chain with SEQ ID NO: 74; KD of 0.5 nM for human CD38, FIG. 7C). The humanized 9G7 best-fit antibody also exhibited a high affinity for the cynomolgus monkey CD38 antigen (KD of 3.2 nM, data not shown), and an enhanced FAB thermo-stability (FAB T_m from DSC scans) over the chimeric 9G7 antibody (94°C vs. 82.2°C for the humanized 9G7 best-fit antibody and the chimeric 9G7 antibody, respectively; see FIG. 7G). The humanized 9G7 best-fit antibody has heavy chain variable domain with SEQ ID NO: 58 and light chain variable domain with SEQ ID NO: 59.

In addition, the 9G7 mouse antibody was humanized following the best-framework approach via CDR grafting onto the VH3-23 and VK1 germline frameworks. Humanized VH and VL variants with different degree of back mutations were investigated *in silico* and one preferred selection of humanized VH and VL combination was transiently expressed as a human IgG1 antibody (the resulting antibody is referred herein as the humanized 9G7 best-framework antibody with heavy chain SEQ ID NO: 149 and light chain with SEQ ID NO: 150). The following mouse back mutations were introduced: (VH) A24F, V37I, V48L, S49A, F67L, R71K, N73T, L78V, and K94R, and (VL) F36Y (Kabat numbering). This antibody exhibited a strong binding to human CD38 and cynomolgus monkey CD38 with affinity constants similar to that of the humanized 9G7 best-fit antibody (KD of 0.4 and 1 nM for human and cynomolgus monkey CD38, respectively; FIG. 7D). FAB thermo-stability (FAB T_m from DSC scans) was also very similar to that of the 9G7 best-fit F36Y humanized variant (89.2°C,

see FIG. 7H). FIG. 7J summarizes the different humanized 9G7 antibodies described above. The humanized 9G7 best-framework antibody has heavy chain variable domain with SEQ ID NO: 60 and light chain variable domain with SEQ ID NO: 61.

5 In a fourth approach, an antibody phage library was screened to generate additional scFv fragments against human CD38. The library had a diversity based on the naturally occurring human V genes. This donor derived antibody phage display library used cDNAs amplified from blood lymphocytes originating from 48 human donors of which 70% had an autoimmune disease (vasculitis, systemic lupus erythematosus, spondyloarthritis, rheumatoid arthritis and scleroderma). Library construction followed the protocol described
10 by Schofield et al. (2007, *Genome Biol.*, 8(11): R254) with a total diversity of 2.53×10^{10} clones. ScFv fragments recognizing human and/or cynomolgus monkey CD38 were isolated from this donor derived phage display library as follows. ScFv fragments were isolated in a series of repeated selection cycles on recombinantly derived human and/or cynomolgus monkey CD38 antigens (see Materials and Methods section). Methods to screen antibody
15 phage display libraries are known (Viti F et al., (2000) *Methods Enzymol*, 326: 480-505). Briefly, following incubation with the library, the immobilised antigen which had been previously coated on a plastic immunotube (overnight in PBS at a concentration of 20 $\mu\text{g/ml}$) or captured on streptavidin beads (when using a biotin labelled form of the antigen, antigen
20 captured at a concentration of 50 nM throughout the selection process), bound phages were recovered whilst unbound phages were washed away. Bound phages were rescued as described by Marks et al (Marks JD et al., (1991) *J Mol Biol*, 222(3): 581-97) and the selection process repeated three times. Over one thousand clones from the second and third round of panning were expressed and analysed by ELISA against the human and cynomolgus monkey CD38
25 antigens. Positive clones were subjected to DNA sequencing and some of the unique clones were further analysed for their ability to bind cell lines expressing human CD38. Following a first round of panning on a biotin labelled version of the human CD38 antigen immobilized on streptavidin beads and a second round of panning on a biotin labelled version of the cynomolgus monkey CD38 antigen immobilized on streptavidin beads, one preferred scFv
30 fragment (clone No 767) having a variable heavy chain sequence with SEQ ID NO: 62 and a variable light chain with SEQ ID NO: 63 was selected for its ability to bind both human and cynomolgus monkey CD38. When formatted as a human IgG1 antibody, clone 767 had a KD of about 300 nM for human CD38 (FIG. 7E) and about 1.2 μM for cynomolgus monkey

CD38 (data not shown) (clone 767 IgG1 antibody is referred herein as human 767 antibody with heavy chain SEQ ID NO: 151 and light chain with SEQ ID NO: 152). FAB thermostability (FAB T_m from DSC scans) was 70.2°C (FIG. 7I). Clone 767 VH domain belongs to the VH3 domain subclass.

5

EXAMPLE 3: CD38/CD3 targeting BEAT antibodies

Anti-CD38 and anti-CD3 epsilon arms can be formatted either as a scFv-Fc type of heavy chains consisting of a scFv fragment fused to a BEAT chain or as a heavy chain consisting of a FAB fragment fused to a BEAT chain similar to that of a naturally occurring antibody. The FAB based heavy chain requires its association with its cognate light chain to assemble into a functional antigen binding site.

10

15

L234A and L235A substitutions were introduced in CH2 regions and residual Protein A binding was abrogated within using the G65S or N82aS substitutions (Kabat numbering) when appropriate. Examples of BEAT antibodies targeting both human CD38 antigen and human CD3 epsilon were formatted as follows:

20

25

30

A first example of BEAT antibodies targeting both human CD38 antigen and human CD3 epsilon using the humanized HB7 bestfit VH and VL sequences was formatted as follows:

A BEAT CD38/CD3 antibody was engineered using a combination of antigen binding sites described in Example 2.1 and 2.2 for the anti-human CD3 epsilon and the anti-human CD38 arms, respectively. The anti-human CD38 arm of the hetero-dimeric immunoglobulin consisted of a BEAT heavy chain (SEQ ID NO: 153) encompassing a variable heavy chain region, a CH1 γ 1 region, a γ 1 hinge region, a γ 1 CH2 region with L234A and L235A substitutions (EU numbering), and a γ 1 based BEAT CH3 domain assembled with its cognate light chain (SEQ ID NO: 72). The anti-human CD3 epsilon arm of the hetero-dimeric immunoglobulin consisted of a BEAT heavy chain (SEQ ID NO: 154) encompassing a scFv fragment, a CH1 γ 1 region, a γ 1 hinge region, a γ 3 CH2 region with L234A and L235A substitutions (EU numbering), and a γ 3 based BEAT CH3 domain. This heavy chain encompassed part of a human IgG3 Fc region and therefore had no binding to Protein A but since the heavy chain used herein had its heavy chain variable domain originating from a VH3 framework, the VH domain was mutated to include the N82aS substitution thereby removing

any additional Protein A binding sites within the heavy chain.. The bispecific antibody is referred herein as BEAT CD38-HB7bestfit/CD3 antibody (FIG. 8 format A).

The BEAT CD38-HB7bestfit/CD3 antibody was expressed transiently, purified and tested *in vitro* for its affinity towards the CD38 and CD3 epsilon antigens, its stability and its ability to redirect T cell killing. The KD value was 3.2 nM for the human CD38 antigen (measured by SPR; FIG. 9A). DSC profiles for the bispecific antibody showed good thermo-stability profiles with a Tm of approximately 68°C for the scFv portion. The FAB portion had a Tm of approximately 91°C (FIG. 9B).

CD38 expressing cell lines (see Materials and Methods section) were used to assess redirected T cell killing in assays. FIG. 10 shows T cell redirected killing of RPMI 8226 myeloma cells using the BEAT CD38-HB7bestfit/CD3 antibody. Note that the assay used purified T cells as effector cells with an effector cells to target cells ratio of 10 to 1. When measured with the RDL-FACS method, the BEAT CD38-HB7bestfit/CD3 antibody had an EC₅₀ of 2.2 pM (mean of 2 donors, 48h incubation).

A second example of BEAT antibodies targeting both human CD38 antigen and human CD3 epsilon using the human clone 767 VH and VL sequences was formatted as follows: a BEAT CD38/CD3 antibody was engineered using a combination of antigen binding sites described in Example 2.1 and 2.2 for the anti-human CD3 epsilon and the anti-human CD38 arms, respectively. The anti-human CD38 arm of the hetero-dimeric immunoglobulin consisted of a BEAT heavy chain (SEQ ID NO: 65) encompassing a variable heavy chain region, a CH1 γ 1 region, a γ 1 hinge region, a γ 1 CH2 region with L234A and L235A substitutions (EU numbering), and a γ 1 based BEAT CH3 domain assembled with its cognate light chain (SEQ ID NO: 138). The anti-human CD3 epsilon arm of the hetero-dimeric immunoglobulin consisted of a BEAT heavy chain (SEQ ID NO: 155) encompassing a scFv fragment, a CH1 γ 1 region, a γ 1 hinge region, a γ 3 CH2 region with L234A and L235A substitutions (EU numbering), and a γ 3 based BEAT CH3 domain. This heavy chain encompassed part of a human IgG3 Fc region and therefore had no binding to Protein A but since the heavy chain used herein had its heavy chain variable domain originating from a VH3 framework, the VH domain was mutated to include the G65S substitution thereby removing any additional

Protein A binding sites within the heavy chain. This bispecific antibody is referred herein as BEAT CD38-767/CD3 antibody (FIG. 8 format B).

The BEAT CD38-767/CD3 antibody was expressed transiently, purified and tested *in vitro* for its affinity towards the CD38 and CD3 epsilon antigens, its stability and its ability to redirect T cell killing. CD38 expressing cell lines (see Materials and Methods section) were used to assess redirected T cell killing in assays similar to that of described in Example 3.2.1. FIG. 11 shows T cell redirected killing of Daudi cells using the BEAT CD38-767/CD3 antibody. Note that the assay used human PBMCs as effector cells with an effector cells to target cells ratio of 10:1. When measured with the RDL-FACS method, the BEAT CD38-767/CD3 antibody had an EC₅₀ of 244 pM (mean of 3 donors, 24h incubation).

Another example of BEAT antibodies targeting both human CD38 antigen and human CD3 epsilon using the humanized 9G7 best-framework VH and VL sequences is formatted as follows: a BEAT CD38/CD3 is engineered using a combination of antigen binding sites described in Example 2.1 and 2.2 for the anti-human CD3 epsilon and the anti-human CD38 antigen binding sites, respectively.

The anti-human CD38 arm of the hetero-dimeric immunoglobulin consists of a BEAT heavy chain (SEQ ID NO: 75 or 155) encompassing a variable heavy chain region, a CH1 γ 1 region, a γ 1 hinge region, a γ 3 CH2 region with L234A and L235A substitutions (EU numbering), and a γ 3 based BEAT CH3 domain assembled with its cognate light chain (SEQ ID NO: 77). This heavy chain encompasses part of a human IgG3 Fc region and therefore has no binding to Protein A but since the heavy chain used herein has its heavy chain variable domain originating from a VH3 framework, the VH domain is mutated to include the G65S substitution thereby removing any additional Protein A binding sites within the heavy chain.

The anti-human CD3 epsilon arm of the hetero-dimeric immunoglobulin consists of a BEAT heavy chain (SEQ ID NO: 68) encompassing a scFv fragment, a CH1 γ 1 region, a γ 1 hinge region, a γ 1 CH2 region with L234A and L235A substitutions (EU numbering), and a γ 1 based BEAT CH3 domain. This bispecific antibody is referred herein as BEAT CD38-9G7bestframework/CD3(SP34-Kappa2) antibody.

Another example of BEAT antibodies targeting both human CD38 antigen and human CD3 epsilon using the human clone 767 VH and VL sequences is formatted as follows: a BEAT CD38/CD3 is engineered using a combination of antigen binding sites described in Example 2.1 and 2.2 for the anti-human CD3 epsilon and the anti-human CD38 antigen binding sites, respectively.

The anti-human CD38 arm of the hetero-dimeric immunoglobulin consists of a BEAT heavy chain (SEQ ID NO: 78) encompassing a variable heavy chain region, a CH1 $\gamma 1$ region, a $\gamma 1$ hinge region, a $\gamma 3$ CH2 region with L234A and L235A substitutions (EU numbering), and a $\gamma 3$ based BEAT CH3 domain assembled with its cognate light chain (SEQ ID NO: 66). This heavy chain encompasses part of a human IgG3 Fc region and therefore has no binding to Protein A but since the heavy chain used herein has its heavy chain variable domain originating from a VH3 framework, the VH domain is mutated to include the G65S substitution thereby removing any additional Protein A binding sites within the heavy chain.

The anti-human CD3 epsilon arm of the hetero-dimeric immunoglobulin consists of a BEAT heavy chain (SEQ ID NO: 68) encompassing a scFv fragment, a CH1 $\gamma 1$ region, a $\gamma 1$ hinge region, a $\gamma 1$ CH2 region with L234A and L235A substitutions (EU numbering), and a $\gamma 1$ based BEAT CH3 domain. This bispecific antibody is referred herein as BEAT CD38-767 /CD3(SP34-Kappa2) antibody.

In a first set of experiments BEAT 9G7/SP34 showed the activation of T cells when in combination with PBMCs but not when isolated.

Example 4:CD3/CD38 BEAT antibodies encompassing only one VH3 domain

An example of BEAT antibodies targeting both human CD38 antigen and human CD3 epsilon using the humanized HB7/bestfit VH and VL sequences was formatted as follows: a BEAT CD38/CD3 was engineered using a combination of antigen binding sites described in Example 2.1 and 2.2 for the anti-human CD3 epsilon and the anti-human CD38 arms, respectively. The anti-human CD38 arm of the hetero-dimeric immunoglobulin consisted of a BEAT heavy chain (SEQ ID NO: 71) encompassing a variable heavy chain domain, a CH1 $\gamma 1$ region, a $\gamma 1$ hinge region, a $\gamma 3$ CH2 region with L234A and L235A substitutions (EU

numbering), and a $\gamma 3$ based BEAT CH3 domain assembled with its cognate light chain (SEQ ID NO: 72). This heavy chain had no binding to Protein A as it encompassed part of a human IgG3 Fc region and had its heavy chain variable domain originating from a non-VH3 domain subclass. The anti-human CD3 epsilon arm of the hetero-dimeric immunoglobulin consisted of a BEAT heavy chain (SEQ ID NO: 157) encompassing a scFv fragment, a CH1 $\gamma 1$ region, a $\gamma 1$ hinge region, a $\gamma 1$ CH2 region with L234A and L235A substitutions (EU numbering), and a $\gamma 1$ based BEAT CH3 domain. This heavy chain and light assembly encompassed a humanized version of the anti-human CD3 epsilon antibody (SP34) as described in PCT Publication No: WO2008119565. This BEAT antibody format is referred herein as BEAT CD38-HB7bestfit/CD3(SP34) antibody (FIG. 12 format A).

The ability of the BEAT CD38-HB7bestfit/CD3(SP34) antibody to redirect T cell killing towards CD38+ cells was investigated *in -vitro*. The CD38+ B lymphoblast cell line Daudi was used in killing assays. FIG. 13 show T cell redirected killing of Daudi cells by the BEAT CD38-HB7bestfit/CD3(SP34) antibody. The assays used human PBMCs as effector cells with an effector cells to target cells ratio of 10 to 1, and the RDL-FACS readout method after a 24h incubation period (see Materials and Methods section). The results show that the BEAT CD38-HB7bestfit/CD3(SP34) antibody was highly potent at redirecting T cell killing against the Daudi CD38+ cell line with an EC₅₀ of 1.8 pM (mean of 3 donors).

A second example of BEAT antibodies targeting both human CD38 antigen and human CD3 epsilon using the humanized 9G7 best-fit VH and VL sequences (SEQ ID NO: 58 and 59, respectively) was formatted as follows: a BEAT CD38/CD3 was engineered using a combination of antigen binding sites described in Example 2.1 and 2.2 for the anti-human CD3 epsilon and the anti-human CD38 arms, respectively. The anti-human CD38 arm of the hetero-dimeric immunoglobulin consisted of a BEAT heavy chain (SEQ ID NO: 73) encompassing a variable heavy chain domain, a CH1 $\gamma 1$ region, a $\gamma 1$ hinge region, a $\gamma 3$ CH2 region with L234A and L235A substitutions (EU numbering), and a $\gamma 3$ based BEAT CH3 domain assembled with its cognate light chain (SEQ ID NO: 74). This heavy chain had no binding to Protein A as it encompassed part of a human IgG3 Fc region and had its heavy chain variable domain originating from a non-VH3 domain subclass. The anti-human CD3 epsilon arm of the hetero-dimeric immunoglobulin consisted of a BEAT heavy chain (SEQ ID NO: 158) encompassing a scFv fragment, a CH1 $\gamma 1$ region, a $\gamma 1$ hinge region, a $\gamma 1$ CH2

region with L234A and L235A substitutions (EU numbering), and a $\gamma 1$ based BEAT CH3 domain. This arm of the bispecific antibody encompassed the variable domains of the humanized SP34 VH5/VL32 antibody described in Example 2.1. This BEAT antibody format is referred herein as BEAT CD38-9G7best-fit/CD3(SP34-Kappa2) antibody (FIG. 12 format B). CD38-9G7best-fit/CD3(SP34-Kappa2) antibody had a KD value of 18 nM for the human CD3 1-26_Fc fusion protein (FIG. 14).

The ability of the BEAT CD38-9G7best-fit/CD3(SP34-Kappa2) antibody to redirect T cell killing towards CD38+ cells was investigated *in vitro*. The CD38+ B lymphoblast cell line Daudi was used in killing assays. FIG. 15 show T cell redirected killing of Daudi cells by the BEAT CD38-9G7best-fit/CD3(SP34-Kappa2) antibody. The assays used human PBMCs as effector cells with an effector cells to target cells ratio of 10 to 1, and the RDL-FACS readout method after a 24h incubation period (see Materials and Methods section). The results show that the BEAT CD38-9G7best-fit/CD3(SP34-Kappa2) antibody was highly potent at redirecting T cell killing against the Daudi CD38+ cell line with an EC₅₀ of 2 pM (mean of 3 donors).

Example 5: Functional equivalence of improved SP34 in scFv format

In order to determine whether of the modifications made to the various SP34 scFv so as to improve their expression also affected the functional properties, namely the binding of CD3, they were tested in the context of a CD38xCD3 bispecific. The CD38 binding arm present as a FAB, comprises the heavy chain variable region encoded by SEQ ID NO: 60 and the light chain variable region encoded by SEQ ID NO: 61. The CD3 binding arm comprises the original mouse SP34 reformatted as an scFv (SEQ ID NO: 207), or modified humanised SP34 scFv's comprising the heavy/light chain combinations H1/L21 (SEQ ID NO: 67), H5/L32 (SEQ ID NO: 68), H5/L65 (SEQ ID NO: 69) and H5/L67 (SEQ ID NO: 70).

Each of the CD3/CD38 BEAT using the different version of SP34 were transiently expressed and purified. They were tested *in vitro* to compare their ability to redirect T cell killing. Raji CD38 expressing cell line (see Materials and Methods section) was used to assess redirected T cell killing. The assay used human PBMCs as effector cells with an effector cells to target cells ratio of 10:1.

When measured with the RDL-FACS method, all BEAT showed comparable EC₅₀ between 6 and 10pM (the mean of 2 donors, 24h incubation) Figure 16.

5 **Example 6: D401Q substitution abrogates protein A and G binding in one of the two homodimer species**

For analytical purposes, we produced the BTA and BTB homodimers of a FAB x scFv BEAT antibody wherein the FAB arm (BTA homodimer) had no binding to protein A since it encompassed a non-VH3 variable domain and part of a human IgG3 Fc region similarly to
10 that described above, and the scFv arm (BTB homodimer) bound protein A using both its VH3 based variable domain and its human IgG1 Fc region. Upon expression and purification (as described above), we found that while the BTA homodimer could readily be purified by protein G (heavy chain SEQ ID NO: 156 and light chain SEQ ID NO: 74), the BTB
15 homodimer could only be isolated by protein A and did not bind protein G, suggesting, for example, misfolding or conformational changes in the protein G binding site in the Fc region carrying the BTB CH3 domain (SEQ ID NO: 159). Considering that protein G binds to almost the same part of the Fc region as protein A does (at the CH2-CH3 interface), it is surprising that the BTB homodimers bound protein A but not protein G. It was then hypothesized that
20 protein A binding by the BTB homodimer was solely mediated by the protein A binding site located in the scFv moiety which as mentioned above encompassed a VH3 based variable domain. To assess the contribution of the scFv binding to protein A, a construct lacking the VH3 based scFv was produced and found to no longer bind protein A (SEQ ID NO: 160, see FIG. 17). It was concluded that in a BTB homodimer, the protein A and G binding sites in the
25 Fc region are both non-functional.

This later result suggested that the BTB homodimer was indeed misfolded or structurally altered in the protein A/G binding region. However, previous data wherein the BTB CH3 domain was used in the context of a heterodimeric immunoglobulin suggested otherwise. To
30 further investigate the functionality of the BTB CH3 domain in a heterodimeric context with regard to protein A binding, we produced a FAB x scFv BEAT antibody consisting of a FAB arm (BTA arm) with no binding to protein A since it encompassed a non-VH3 variable domain, a human IgG1 hinge, a human IgG3 CH2 domain and an engineered BTA CH3

domain originating from a human IgG3 isotype (FAB arm heavy chain with SEQ ID NO: 156 and light chain SEQ ID NO: 74), and a scFv arm (BTB arm) encompassing a VH3 based scFv fragment abrogated for protein A binding, a human IgG1 hinge, a human IgG1 CH2 domain and an engineered BTB CH3 domain originating from a human IgG1 isotype (scFv arm heavy chain with SEQ ID NO: 161) – therefore a heterodimer that was solely expected to bind protein A in the Fc region of the scFv arm. Upon expression and purification (as described above), it was found that indeed this molecule bound protein A and G (FIG.18A) and had no trace of BTB homodimers. Thus it was concluded that the protein A and G sites are intact in the context of a heterodimeric immunoglobulin and that only when the BTA and BTB chains are associated is a functional protein A/G binding site formed.

To understand the molecular basis of the protein A/G abrogation within the BTB homodimer, previously engineered BTB CH3 domains originating from a human IgG1 isotype were investigated. Substitution D401Q, EU numbering, a comparison between EU and IMGT numbering is provided at http://www.imgt.org/IMGTScientificChart/Numbering/Hu_IGHGnber.html, in the BTB CH3 domain was identified from modelling studies, and was engineered to abrogate a putative electrostatic contact between Arg411 of the first chain of the BTB homodimer with Asp401 of the second chain of the BTB homodimer with the view that this interaction could stabilize the homodimeric complex. Since this substitution was only implemented in the last step of the BTB CH3 domain engineering, a FAB x scFv BEAT antibody as described above was produced without the said D401Q substitution in the BTB CH3 domain (FAB arm with heavy chain and light chain having SEQ ID NOs: 156 and 74, respectively; scFv arm with heavy chain SEQ ID NO: 162). Upon expression and purification (as described above), it was found that this molecule bound protein A and G (FIG.18B) but showed the presence of BTB homodimers, thus indicating that the D401Q substitution was responsible for the lack of protein A/G binding observed in the BTB homodimers, thereby making a significant contribution within a heterodimeric immunoglobulin at reducing the said BTB contaminants. Importantly, residue 401 itself is not part of the protein A/G binding site, but is located oppositely at the CH3-CH3 interface. Thus, protein A/G abrogation is not a result of directly disrupting interactions between protein A/G and the Fc region. Rather it is suggested that, unexpectedly, D401Q leads to long range conformational changes at the CH2-CH3 interface as a result of the poor pairing capabilities of BTB chains with each other. In the

case of the heterodimer, the interfaces fit perfectly together and thus a proper conformation at the CH2/CH3 interface renders the protein A/G binding site functional.

This surprising result can be exploited to design a new strategy to efficiently isolate heterodimeric immunoglobulins without homodimer contaminants using the specific design that follows: the heterodimeric immunoglobulin has a first chain encompassing a Fc region of the IgG3 isotype that will include a BTA CH3 domain and a non-VH3 variable domain or a VH3 based variable domain abrogated for protein A binding (using the G65S or N82aS substitutions for example) or no variable domain and therefore has no binding to protein A, and a second chain that binds protein A encompassing a BTB D401Q CH3 domain (originating from a human IgG1 isotype for example) and either a non-VH3 variable domain or a VH3 variable domain abrogated for protein A binding (using the G65S or N82aS substitutions for example), or no variable domain. In this setting, both homodimers contain no protein A binding sites. The first homodimer is a dimer of the first chain and does not bind protein A as there is no protein A binding site while the second homodimer is a dimer of the second chain with a protein A binding site that is non-functional. The resulting heterodimer has only one protein A binding site found in the second chain (the BTB chain protein A binding site being functional only when paired with the BTA chain). Hence the heterodimer will be the only species binding to the protein A chromatography resin while unwanted homodimers of BTA and BTB chains will be washed away.

Example 7: CD3/CD38 BEAT activity *in vivo* in cynomolgus monkey.

Cynomolgus monkeys (macaca fascicularis, 1 male, 1 female per dose) were dosed by intravenous bolus injection of CD3/CD38 (comprising the CD3 heavy/light chain combination SP34 H5/L65 formatted as a scFv SEQ ID NO: 69 and the CD38 heavy chain variable region encoded by SEQ ID NO: 60 and the light chain variable region encoded by SEQ ID NO: 61 formatted as a FAB), referred to in this example as CD3/CD38 BEAT.

Immunophenotyping was performed on all animals from blood samples (EDTA) taken pre-treatment and on different time points following each administration of CD3/CD38 BEAT. Assessment of lymphocyte subsets was performed by flow cytometry using the following markers: Total T lymphocytes (CD4 and CD8), Monocytes (CD14) and CD38. The samples were acquired on a FACSCanto™ II flow cytometer. Data were analyzed using FACSDiva™

analysis software as follows: First, an electronic gate based on forward and side scatter characteristics was drawn to select the lymphocyte or monocyte populations (parent population). The absolute counts per μL of whole blood of the T lymphocyte population (CD4+ T cells and CD8+ T cells) and of the CD38 positive monocyte population (CD14+CD38+) was calculated from their relative percentages as derived from the parent population gate and the total parent population counts from a validated hematology analyzer according to the formula: Absolute population count ($\times 10^3/\mu\text{L}$) = (population relative % $\times$ total parent population count) / 100.

There was a clear early decrease of the T cell counts, both CD4 and CD8 positives, following the dose of 1 and 10 $\mu\text{g/kg}$ (Figure 19). Compared with cell counts obtained 1 day prior to dosing, there was a 48 and 78% decrease in CD4 T cell counts after the 1 $\mu\text{g/kg}$ dose and more than 90% after the 10 $\mu\text{g/kg}$ dose at the 4 hour time point. The CD8 T cell counts were even more drastically affected, with a drop of 86% after the 1 $\mu\text{g/kg}$ dose. The CD4 T cell counts rebounded rapidly and were overall similar to pre dose counts after 48 hours. However the CD8 T cell counts were slower to recover and reached similar levels than pre-dose after 1 week. This transient decrease of peripheral blood T cells was expected and reflects the activation of these populations by CD3/CD38 BEAT. These observations suggest that the T lymphocytes, and in particular the cytotoxic CD8 T cells are engaged into killing activities in the tissues.

There was also a clear decrease in the circulating CD38+ monocyte population (Figure 20). A very fast disappearance was observed, followed by a rebound between 24 and 48h. This rebound was generally followed by a second substantial decrease in the CD38+ monocytes counts indicating that this population is likely targeted and depleted by the cytotoxic action of the T lymphocytes.

Example 8: 9G7-based BEAT antibodies display enhanced killing activity than an OKT10xOKT3 BEAT

The potencies of two different 9G7-based BEAT antibodies vs. an OKT10-based BEAT antibody were investigated in RDL assays.

The bispecific antibody referred herein as BEAT CD38-9G7Mouse/CD3 is a BEAT CD38/CD3 antibody based on a combination of antigen binding sites described in Example 2 (OKT3 VH11/VL8 scFv for the anti-human CD3 epsilon arm and mouse-human chimeric 9G7 FAB for the anti-human CD38 arm). The anti-human CD38 arm of the bispecific antibody consisted of a BEAT heavy chain (SEQ ID NO: 177) assembled with its cognate light chain (SEQ ID NO: 178). The anti-human CD3 epsilon arm of the bispecific antibody consisted of a BEAT heavy chain (SEQ ID NO: 179) encompassing the anti-human CD3 epsilon scFv fragment. The bispecific antibody was expressed transiently and purified as described above.

The bispecific antibody referred herein as BEAT CD38-OKT10Mouse/CD3 is a BEAT CD38/CD3 antibody based on a combination of antigen binding sites described in Example 2 (OKT3 VH11/VL8 scFv for the anti-human CD3 epsilon arm and mouse-human chimeric OKT10 FAB for the anti-human CD38 arm). The anti-human CD38 arm of the bispecific antibody consisted of a BEAT heavy chain (SEQ ID NO: 180) assembled with its cognate light chain (SEQ ID NO: 181). The anti-human CD3 epsilon arm of the bispecific antibody consisted of a BEAT heavy chain (SEQ ID NO: 179) encompassing the anti-human CD3 epsilon scFv fragment. The bispecific antibody was expressed transiently and purified as described above.

The BEAT antibody referred herein as the BEAT CD38-9G7best-fit/CD3(SP34-Kappa2) antibody was described above in Example 4.

The cytotoxic potential of different BEAT antibodies based on the 9G7 or the OKT10 anti CD38 mAbs was assessed in a killing assay against the CD38⁺ target Daudi cells (ATCC). A flow cytometry-based readout was performed to quantify target cell death. Similar flow cytometry-based readouts were used in Moore et al. Blood. 2011 Apr 28;117(17):4542-51; Friedrich et al. Mol Cancer Ther 2012;11:2664-2673; Schlereth et al. Cancer Res 2005;65:2882-2889. Effector cells were non stimulated PBMC from healthy donors. The effector:target ratio was typically 10:1 and the incubation time was 48h. Briefly, target cells (Daudi) were labeled with a fluorescent cytoplasmic dye such as CFSE and were distributed in 96-well plates (TPP). Antibodies serial dilutions (3x solutions) and PBMCs (50 μ L/well) were distributed then distributing in corresponding wells. The Plates were incubated for 48 h

in a 5 % CO₂ incubator at 37°C. The cells were resuspended by pipetting and were transferred into U-bottom 96-well plates (TPP). The U-bottom plates were centrifuged 3 min at 300 g, the supernatants were discarded and the cells were resuspended in 200µL of cold FACS buffer (PBS + 2% FBS + 10% Versene) supplemented with 7-AAD (Becton Dickinson) at 1/40 dilution. The plates were immediately acquired on a guava easyCyte™ Flow Cytometer (Millipore). For each well, the absolute number of living target cells was determined by gating on Dye (CFSE) positive 7ADD negative population using flowjo software (Treestar). The % of specific cytotoxicity for each sample was determined using the condition in which only target cells were incubated as baseline or using the condition in which Target cells were mixed with PBMCs (no antibody condition). The EC₅₀ values were determined using nonlinear variable slope regression method with Prism software (GraphPad software).

The data in Table 2 indicates the EC₅₀ values obtained with the different BEAT constructs. The 9G7-based BEAT antibodies (BEAT CD38-9G7Mouse/CD3 and BEAT CD38-9G7best-fit/CD3(SP34-Kappa2) displayed a killing activity characterized by a clearly lower EC₅₀ than the BEAT CD38-OKT10Mouse/CD3 antibody (3.2 and 1.9 pM for 9G7 based BEAT antibodies versus 125.6 pM for the OKT10 based BEAT antibody), indicating a higher cytotoxic potential of the 9G7-based anti-CD38/CD3 BEAT antibodies.

	BEAT CD38-9G7Mouse/CD3	BEAT CD38-9G7best-fit/CD3(SP34-Kappa2)	BEAT CD38-OKT10Mouse/CD3
EC ₅₀ (pM)*	3.2	1.9	126

*The EC₅₀ indicated are average values of killing assays performed with different PBMC donors. N= 8 for 9G7xOKT3, N= 2 for OKT10xOKT3 and n=1 for 9G7xSP34(GV2)

Table 2: 9G7-based BEAT antibodies display enhanced killing activity than an OKT10xOKT3 BEAT

Example 9: 9G7 and 767 binders do not display agonism, in contrast to HB7 antibody

CD38 being competent for signaling, the agonist property of 9G7 and 767 antibodies were tested, in comparison to the HB7 clone, in a calcium flux assay on Jurkat human T cell lymphoma cell line which expresses naturally CD38. The Jurkat cells were loaded with Fluo-4 Dye at 2 μ M (Invitrogen) during 1 hour at 37°C. The cells were washed and resuspended in PBS supplemented with 1 mM Ca²⁺ and 1 mM Mg²⁺. The samples were acquired on a FACS Calibur flow cytometer (Becton Dickinson) for baseline during 40 s. Then test antibodies were added to the samples at the concentration of 10 μ g/ml and the acquisition was resumed until 7 minutes. The figure G shows the mean fluorescence intensity (MFI) of the Fluo-4 dye (FL-1 channel), which is a readout of calcium mobilization into the cytoplasm of the cells, as a function of time. While the isotype control (IgG1 human antibody) did not trigger any calcium flux, the HB7 anti-CD38 antibody did induce a strong signal. Strikingly neither the 9G7 nor the 767 antibody were able to trigger a calcium mobilization signal indicating that these antibodies, in contrast to the HB7 antibody do not have CD38 agonist properties.

During the same series of experiments it was also shown for 9G7 and 767 that they specifically bind to cells expressing CD38 in a specific manner.

Example 10: Epitope mapping

h9G7 epitope mapping

Linear peptide mapping was used to determine the epitope of humanized 9G7 (best-fit version). The extracellular domain of human CD38 (Residues 43-300) was divided into 19 consecutive peptides, 13 amino acids in length and a final peptide of 11 amino acids in length. Peptides were fused to a human IgG1 Fc region (CH2-CH3) with a two-amino-acid linker (Ala-Ser) between Fc and peptide sequence. These constructs were termed F1-F20 (SEQ ID NOs: 182-201) and their binding analyzed by SPR. The humanized 9G7 antibody (heavy chain SEQ ID NO: 145 and light chain SEQ ID: 146) was covalently immobilized on a CM5 sensor chip (about 1800 RUs). Peptide Fc-fusions were injected at 15 nM for 240 seconds. Human CD38 extracellular domain was injected as control. Only F6 and the control gave a binding signal (FIG. 22). All fragments were re-injected at 200 nM and F6 gave a stronger signal (FIG. 23) while the remaining fragments still showed no binding (data not shown). As F6 is part of the epitope of the SAR650984 antibody disclosed in PCT publication NO:

WO2008047242 (heavy chain with SEQ ID NO: 202 and light chain with SEQ ID NO: 203) their interaction was verified by SPR and confirmed (data not shown). Thus h9G7 and SAR650984 compete for binding on human CD38. Furthermore when the F6 sequence is mapped onto the 3D surface of the extracellular domain of human CD38, it overlaps with the SAR650984 epitope. However SAR650984 has been shown not to be cross-reactive with cynomolgus CD38 while humanized 9G7 binds to cynomolgus CD38 with high affinity (FIG. 24) suggesting that the two antibodies have overlapping but different epitopes.

Within the F6 sequence, residue M110 is a Valine in cynomolgus CD38. Thus in the context of the entire extracellular domain of human CD38, we mutated M110 to Valine (SEQ ID NO: 204) and compared binding of h9G7 and SAR650984. SAR650984 showed reduced binding compared to its binding on wild-type human CD38 while binding of humanized 9G7 was not impacted (data not shown). Based on the crystal structure of SAR650984 in complex with human CD38 extracellular domain (PDB accession code 4CMH, Deckert et al., 2014 Clin.Cancer Res. 20: 4574), we identified an additional residue that is potentially important for SAR650984 binding and that differs between human and cynomolgus CD38. This residue was T148 and is a methionine in cynomolgus CD38. Mutation of T148 to Methionine in the extracellular domain of human CD38 (SEQ ID NO: 205) significantly reduced the binding of SAR650984 while humanized 9G7 binding was unaffected (data not shown). A double mutant of the extracellular domain of human CD38 containing mutations M110V and T148M (SEQ ID NO: 206, see FIG. 23) abolished binding of SAR650984 while humanized 9G7 binding was not affected (FIG. 24). Thus it can be concluded that while humanized 9G7 and SAR650984 compete for binding on human CD38, the residues that are bound by these antibodies are not identical.

767 epitope mapping

The epitope of the 767 antibody was mapped using the same set of linear peptide-Fc fusions used for the epitope mapping experiments described above. The 767 antibody (heavy chain SEQ ID NO: 151 and light chain SEQ ID NO: 152) was covalently immobilized on a CM5 sensor chip (about 700 RUs). Peptide Fc-fusions (F1-F20, SEQ ID NO: 182-201) were injected at 1000 nM for 240 seconds. The extracellular domain of human CD38 extracellular was injected as control. Only F3 and the control gave a binding signal (FIG. 25). When the

F3 sequence is mapped onto the 3D surface of the extracellular domain of human CD38 (**FIG. 26**), it partially overlaps with the SAR650984 epitope, more specifically with two residues: E76 and H79. However 767 is cross-reactive to cynomolgus CD38 (with lower affinity) while SAR650984 is not, suggesting that the two antibodies have overlapping but different epitopes.

5 Furthermore SAR650984 did not bind the F3 construct by SPR (**FIG. 27**). Thus it can be concluded that while 767 and SAR650984 compete for binding on human CD38, the residues that are bound by these antibodies are not identical.

Claims

1. An antibody or fragment thereof that binds to human CD38 comprising a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 25 or SEQ ID NO: 208,
5 and/or a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 26 or SEQ ID NO: 209, and/or a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 27 or SEQ ID NO: 210; and/or comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 28 or SEQ ID NO: 211 or SEQ ID NO: 214, and/or a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 29 or
10 SEQ ID NO: 212, and/or a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 30 or SEQ ID NO: 213.
2. The antibody or fragment thereof of claim 1, wherein the antibody or fragment thereof is a murine antibody, chimeric antibody or a humanized antibody.
- 15 3. The antibody or fragment thereof of claim 1 or 2, wherein the antibody or fragment thereof comprises a heavy chain variable region sequence comprising the amino acid sequence consisting of SEQ ID NO: 52, SEQ ID NO: 58, SEQ ID NO: 60 or a sequence at least 80% identical to the non-CDR region of either of said heavy chain variable region sequences and/or wherein the antibody or fragment thereof comprises a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 55, SEQ ID NO: 59, SEQ ID NO: 61 or a sequence at least 80% identical to the non-CDR region of any one of said light chain variable region sequences.
- 20 4. The antibody or fragment thereof of claim 1 or 2, wherein the antibody or fragment thereof comprises a heavy chain sequence comprising the amino acid sequence consisting of SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 76, SEQ ID NO: 145, SEQ ID NO: 147, SEQ ID NO: 149 or a sequence at least 80% identical to the non-CDR region of either of said heavy chain variable region sequences and/or wherein the
25 antibody or fragment thereof comprises a light chain variable region sequence comprising the amino acid sequence of SEQ ID NO: 74, SEQ ID NO: 77, SEQ ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150 or a sequence at least 80% identical to the non-CDR region of any one of said light chain variable region sequences.
- 30

5. The antibody or fragment thereof of claim 1 to 3, wherein said antibody or fragment comprises human heavy and/or light chain constant regions and wherein the human heavy constant region is selected from the group of human immunoglobulins consisting of IGHG1, non fucosylated IGHG1 and IGHG4.
6. A bispecific antibody comprising an anti-CD38 antibody fragment according to any one of claims 1 to 3.
7. The bispecific antibody according to claim 6, wherein said anti-CD38 antibody fragment is selected from the group consisting of Fab, Fab', Fab'-SH, Fd, Fv, dAb, F(ab')₂, scFv, bispecific single chain Fv dimers, diabodies, triabodies and scFv.
8. The bispecific antibody according to any one of claims 6 to 7, further comprising an anti-CD3 antibody fragment selected from the group consisting of Fab, Fab', Fab'-SH, Fd, Fv, dAb, F(ab')₂, scFv, bispecific single chain Fv dimers, diabodies, triabodies and scFv.
9. The bispecific antibody according to claim 8, wherein said anti-CD3 antibody fragment comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 3, and a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 4, and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 5 or SEQ ID NO: 9; and comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 6, and a light chain CDR2 comprising the amino acid sequence of SEQ ID NO: 7, SEQ ID NO: 11, and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12; or wherein said anti-CD3 antibody fragment comprises a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 163, and a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 164, and a heavy chain CDR3 comprising the amino acid sequence of SEQ ID NO: 165; and comprising a light chain CDR1 comprising the amino acid sequence of SEQ ID NO: 166, and a light chain CDR2

comprising the amino acid sequence of SEQ ID NO: 167, and a light chain CDR3 comprising the amino acid sequence of SEQ ID NO: 168.

- 5 10. An antibody or fragment thereof that binds to human and cynomolgus CD38 and which binds to the same epitope as said antibody or fragment thereof according to any one of claims 1 to 5 or said bispecific antibody according to any one of claims 6 to 9.
- 10 11. An epitope on soluble human CD38 which is bound by said antibody or fragment thereof according to any one of claims 1 to 5 or said bispecific antibody according to any one of claims 6 to 9.
- 15 12. A composition comprising the antibody or fragment thereof of any one of claims 1 to 5 or said bispecific antibody according to any one of claims 6 to 9 and a pharmaceutically acceptable carrier.
- 20 13. An immunoconjugate comprising the antibody or fragment thereof of any one of claims 1 to 5 or said bispecific antibody according to any one of claims 6 to 9, linked to a therapeutic agent.
- 25 14. The antibody or fragment thereof according to any one of claims 1 to 5 or said bispecific antibody according to any one of claims 6 to 9, for use as a medicament.
- 30 15. The antibody or fragment thereof according to any one of claims 1 to 5 or said bispecific antibody according to any one of claims 6 to 9, for use as a medicament in treating a disease selected from the group consisting of malignant hematological diseases, including multiple myeloma, B-cell chronic lymphocytic leukaemia, B-cell acute lymphocytic leukaemia, Waldenström's macroglobulinemia, primary systemic amyloidosis, mantle-cell lymphoma, pro-lymphocytic/myelocytic leukaemia, acute myeloid leukaemia, chronic myeloid leukaemia, follicular lymphoma, NK-cell leukaemia and plasma-cell leukaemia.