

THIADIAZOLE COMPOUNDS USEFUL AS INHIBITORS OF CYSTEINE ACTIVITY DEPENDENT ENZYMES

FIELD OF THE INVENTION

This invention relates to novel compounds and their pharmaceutically acceptable acid addition salts and base addition salts for use in the treatment of acne, common cold, inflammatory joint disease by inhibition of cysteine proteases and cysteine activity dependent enzyme. In particular, it relates to novel compounds having pharmaceutical utility, to processes for their preparation, to compositions and uses in the treatment of various diseases by inhibition of cysteine proteases and cysteine activity dependent enzymes.

BACKGROUND OF THE INVENTION AND PRIOR ART

Transglutaminase, rhinovirus 3C protease, calpain, Interleukin beta converting enzyme, cathepsin B are examples of cysteine activity-dependent enzymes which are involved in the progression of various disorders and/or diseases such as acne, common cold and arthritis. These enzymes include in their chemical structure cysteine residues. It is believed that the thiol group of a cysteine residue in the enzyme acts as a nucleophile and causes the hydrolysis of the substrate thus permitting the progression of the disorder and/or disease. Accordingly, attempts have been made to develop thiol trapping agents to inhibit the catalytic activity of such enzymes in order to prevent the progression of the disease. Haghware reported the use of 1,2,4-thiadiazolines as inhibitors of alcohol dehydrogenase, a cysteine activity dependent enzyme.

EP-A-0 389 901, EP-A-0 473 980 and EP-A-0 548 650 disclosed the preparation of 1,2,4-thiadiazole-substituted acrylic acids and their use as pesticides. EP-A-0 473 980 also described the preparation of N-(1,2,4-thiadiazol-5-yl)-N-methylglycine ester and their use as pesticides.

US Patent 5,618,792 disclosed certain 3-substituted oxadiazole and 3-substituted thiadiazole peptoids which are serine protease inhibitors. US Patent 4,207,090 disclosed amino ester derivatives of 3-trihalomethyl-[1,2,4]-thiadiazoles as pesticides. US Patent 5,677,302 discloses the use of 1,2,4-thiadiazole [4,5-a] benzimidazoles and imidazo [1,2-d]-1,2,4-thiadiazoles as inhibitors of the enzyme H⁺/K⁺-ATPase, also known as the proton pump,



166166



G00002

another cysteine activity dependent enzyme. Condensed thiadiazole derivatives having a sulfonylimino group have been disclosed in US Patent 5,550,138 as being cathepsin B inhibitors. The various peptidyl inhibitors of cysteine proteases have been reviewed in protein profile, 1995, Vol. 2, Issue 14, p. 1587-1591. Unfortunately, most of the efforts have been largely frustrated by the reactivity of potential inhibitors with other nucleophiles such as alcohols and amines which are abundant in physiological systems.


The development of compounds which exhibits selective reactivity towards the thiol group of a cysteine of the cysteine activity dependent enzyme residue will represent an enormous advance in this field.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide novel pharmaceutical compounds, and composition containing such compounds which are active as cysteine activity dependent enzyme inhibitors and hence useful in the treatment of disorder and/or disease caused by the activity of such enzymes, and in particular transglutaminase human, rhinovirus 3C protease and cathepsin B.



It is a further object of the invention to provide processes for the synthesis of such compounds.

In a first aspect, the invention provides 3-5 disubstituted 1,2,4-thiadiazole compounds, having the general formula (II):



or their pharmaceutically acceptable salts thereof,

wherein A is an amino acid residue, or a peptide containing 2 to 3 amino acid residues with the N terminal of A directly attached to the C5 of the 1,2,4-thiadiazole ring or an isosteric form thereof and W represents a group of formula $-N(R^1)_2$ or $-OR^1$ with R^1 being independently hydrogen, lower alkyl, lower alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N or O atom, with the proviso that -A-W is not a group of formulas:



wherein R^e represents hydrogen or lower alkyl having a straight or branched chain of 1 to 4 carbon atoms;


R^f represents lower alkyl having a straight or branched chain of 1 to 4 carbon atoms;

R^i represents lower alkyl having a straight or branched chain of 1 to 4 carbon atoms;

R_j represents lower alkyl having a straight or branched chain of 1 to 4 carbon atoms, or benzyl which is unsubstituted or monosubstituted to trisubstituted by identical or different substituents selected from the group fluorine, chlorine, bromine, nitro, cyano, lower alkyl, methylthio, lower alkoxy, trifluoromethyl, trifluoromethoxy, difluoromethoxy, trifluoromethylthio, methoxycarbonyl, ethoxycarbonyl, methoximino-methyl, methoxyiminoethyl, ethoximinomethyl cyclopentyl, cyclohexyl, 1,3-propanediyl, but-2-ene-di-1,4-yl, 1,4-butanediyl, and phenyl, benzyl, phenoxy or benzyloxy in each case unsubstituted or monosubstituted to trisubstituted in the phenyl moiety by identical or different substituents selected from the group consisting of fluorine, chlorine, bromine, lower alkoxy, lower alkyl, methylthio, trifluoromethyl, trifluoromethoxy, difluoromethoxy, and trifluoromethylthio;


and Y is selected from:

- (1) lower alkoxy, lower cycloalkoxy, lower arylalkoxy, heterocycloxy, and lower heterocyclalkoxy wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;
- (2) lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocycl, aryl, lower arylalkyl, lower arylalkenyl, lower heterocyclalkenyl wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, and dialkylamino;
- (3) lower alkoxycarbonyl, carboxyl;
- (4) a ketone group of formula:



in which R² represents lower alkyl, lower cycloalkyl, lower heterocyclylalkyl, heterocyclyl, aryl, lower arylalkyl wherein the alkyl or aromatic ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;


- (5) a carbamoyl group of formula:



with R² being as defined above;


- (6) amino, lower alkylamino, lower dialkylamino;

- (7) amide of formula:




with R² being as defined above;

- (8) a group of formula:




wherein A is as defined above and the carboxyl terminal of A is directly attached to the nitrogen of the 3-amino-1,2,4-thiadiazole, R represents hydrogen, lower alkanoyl, lower cycloalkylcarbonyl, lower alkoxy carbonyl, lower arylalkyloxycarbonyl or N protecting group and R¹ is hydrogen, lower alkyl, lower alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N atom;

- (9) alcohol of formula:




with R^2 being as defined above;

- (10) sulfone of formula:




with R^2 being as defined above;

- (11) sulfoxide of formula:



with R^2 being as defined above;


- (12) sulfonamide of formula:



with R^2 being as defined above;


- (13) lower alkylthio, lower arylalkylthio, arylthio;

- (14) a group of formula:




with A as defined above and the N-terminal of A is directly attached to the methylene and W being as defined above;

- (15) a group of formula:



in which R^3 and R^4 are independently alkyl, aralkyl, heterocyclyl, heterocyclalkyl; R^3 and R^4 when taken together form with the N-atom a five or a six membered ring selected from the group piperidinyl, pyrrolidinyl, piperazinyl with the N-4 position of piperazine optionally substituted with pyridyl, heterocyclyl, alkyl, aralkyl and aryl.

In a second aspect, the invention provides 3-5 disubstituted 1,2,4-thiadiazole compounds, having the general formula (II):



or their pharmaceutically acceptable salts thereof,

wherein A-W is glycynamide, leucyl-proline, isoleucyl-proline, leucyl-octahydro-1*H*-indole-2-carboxylic acid, isoleucyl-octahydro-1*H*-indole-2-carboxylic acid, leucyl-octahydrocyclopenta[*b*]pyrrole-2-carboxylic acid, isoleucyl-octahydrocyclopenta[*b*]pyrrole-2-carboxylic acid or A is an amino acid residue A_1 or a dipeptide A_1 - A_2 with the N-terminal of A directly attached to the C5 of the 1,2,4-thiadiazole ring or an isomeric form thereof and W is -OH, -NH₂, lower alkoxy, lower alkylamino, the amino acid A_1 is selected from the group alanyl, β -alanyl, arginyl, aspartyl, glutamyl, glutaminyl, histidyl, isoleucyl, leucyl, lysyl, methionyl, norleucyl, norvalyl, ornithyl, phenylalanyl, prolyl, seryl, tryptophyl, tyrosyl, valyl, 4-aminobutanoyl, 2,6-diaminohexanoyl, 2,3-diaminopropanoyl, 2,4-diaminobutanoyl and 2,5-diaminopentanoyl, when A_1 is 2,6-diaminohexanoyl, 2,3-diaminopropanoyl, 2,4-diaminobutanoyl and 2,5-diaminopentanoyl, the N-attachment point is either at the α -NH₂ or the amino acid side-chain NH₂ group, A_2 is selected from the group alanyl, β -alanyl, arginyl, aspartyl, glycyl, glutamyl, glutaminyl, histidyl, isoleucyl, leucyl, lysyl, methionyl, norleucyl, norvalyl, ornithyl, phenylalanyl, prolyl, seryl, tryptophyl,

7a

tyrosyl, valyl, 2-indolinecarbonyl, octahydro-1*H*-indole-2-carbonyl, octahydrocyclopenta[*b*]pyrrole-2-carbonyl, octahydro-1*H*-isoindole-1-carbonyl, octahydrocyclopenta[*c*]pyrrole-1-carbonyl, 1,2,3,4-tetrahydro-3-isoquinolinecarbonyl, decahydro-3-isoquinolinecarbonyl;

5

and Y is selected from:

(1) lower alkoxy, lower cycloalkoxy, lower arylalkoxy, heterocycloxy, and lower heterocyclalkoxy wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;

10


(2) lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocyclyl, aryl, lower arylalkyl, lower arylalkenyl, lower heterocyclalkenyl wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, and dialkylamino;

15

(3) lower alkoxycarbonyl, carboxyl;

20

(4) a ketone group of formula:




25

in which R² represents lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocyclyl, aryl, lower arylalkyl wherein the alkyl or aromatic ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;

30

(5) a carbamoyl group of formula:



35

with R² being as defined above;

7b

(6) amino, lower alkylamino, lower dialkylamino;


(7) amide of formula:



with R^2 being as defined above;

10

(8) a group of formula:



15

wherein A is as defined above and the carboxyl terminal of A is directly attached to the nitrogen of the 3-amino-1,2,4-thiadiazole, R represents hydrogen, lower alkanoyl, lower cycloalkylcarbonyl, lower alkoxy carbonyl, lower arylalkyloxycarbonyl or N protecting group and R^1 is hydrogen, lower alkyl, lower alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N atom;

20


(9) alcohol of formula:



with R^2 being as defined above;

30

(10) sulfone of formula:




35

with R^2 being as defined above;


(11) sulfoxide of formula:

7c



with R^2 being as defined above;


(12) sulfonamide of formula:



with R^2 being as defined above;


(13) lower alkylthio, lower arylalkylthio, arylthio;

(14) a group of formula:




with A as defined above and the N-terminal of A is directly attached to the methylene and W being as defined above;

(15) a group of formula:



in which R^3 and R^4 are independently alkyl, aralkyl, heterocyclyl, heterocyclalkyl; R^3 and R^4 when taken together form with the N-atom a five or a six membered ring selected from the group piperidiny, pyrrolidiny, piperaziny with the N-4 position of piperazine optionally substituted with pyridyl, heterocyclyl, alkyl, aralkyl and aryl.


In a third aspect, the invention provides 3-5 disubstituted 1,2,4-thiadiazole compounds, having the general formula (III):



7d

or their pharmaceutically acceptable salts thereof,

wherein A is an amino acid residue, or a peptide containing 2 to 3 amino acid residues with the N-terminal of A attached by the means of a spacer X to the C5 of the 1,2,4-thiadiazole ring, or an isosteric form thereof, X is a spacer selected from the group of formula



$\text{—N(R}^1\text{)}_2$ or —OR^1 with R^1 being independently hydrogen, lower alkyl, lower alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N or O atom;


and Y is selected from:

(1) lower alkoxy, lower cycloalkoxy, lower arylalkoxy, heterocycloxy, and lower heterocyclalkoxy wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;

(2) lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocycl, aryl, lower arylalkyl, lower arylalkenyl, lower heterocyclalkenyl wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, and dialkylamino;

(3) lower alkoxycarbonyl, carboxyl;


(4) a ketone group of formula:



7e

in which R^2 represents lower alkyl, lower cycloalkyl,
lower heterocyclylalkyl, heterocyclyl, aryl, lower arylalkyl
wherein the alkyl or aromatic ring is optionally
substituted with 1 to 2 substituents selected from the
group amino, alkoxy, hydroxy, halo, amino, alkylamino,
dialkylamino;


- (5) a carbamoyl group of formula:



with R^2 being as defined above;


- (6) amino, lower alkylamino, lower dialkylamino;

- (7) amide of formula:




with R^2 being as defined above;

- (8) a group of formula:



wherein A is as defined above and the carboxyl
terminal of A is directly attached to the nitrogen of the
3-amino-1,2,4-thiadiazole, R represents hydrogen,
lower alkanoyl, lower cycloalkylcarbonyl, lower
alkoxycarbonyl, lower arylalkyloxycarbonyl or N
protecting group and R^1 being as defined as above;

- (9) alcohol of formula:




7f

with R² being as defined above;

(10) sulfone of formula:


5



with R² being as defined above;

10


(11) sulfoxide of formula:



with R² being as defined above;

15

(12) sulfonamide of formula:




20

with R² being as defined above;

(13) lower alkylthio, lower arylalkylthio, arylthio;

25


(14) a group of formula:



with A as defined above and the N-terminal of A is directly attached to the methylene and W being as defined above;

30

(15) a group of formula:




in which R³ and R⁴ are independently alkyl, aralkyl, heterocyclyl, heterocyclalkyl; R³ and R⁴ when taken together form with the N-atom a five or a six membered ring selected from the group piperidinyl, pyrrolidinyl, piperazinyl with the N-4 position of piperazine optionally

35

7g

substituted with pyridyl, heterocyclyl, alkyl, aralkyl and aryl.

5 In a fourth aspect, the invention provides 3-5 disubstituted 1,2,4-thiadiazole compounds, having the general formula (IV):



or their pharmaceutically acceptable salts thereof,

15 wherein A is an amino acid residue, or a peptide containing 2 to 3 amino acid residues, the carboxyl terminal of A is directly attached to the nitrogen of the 5-amino 1,2,4-thiadiazole and R represents hydrogen, lower alkanoyl, lower cycloalkylcarbonyl, lower alkoxy carbonyl, lower arylalkyloxycarbonyl or N protecting group and

20 R¹ is hydrogen, lower alkyl, lower alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N atom;

and Y is selected from:

- 25
- (1) lower alkoxy, lower cycloalkoxy, lower arylalkoxy, heterocycloxy, and lower heterocyclalkoxy wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;

30

 - (2) lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocyclyl, aryl, lower arylalkyl, lower arylalkenyl, lower heterocyclalkenyl wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, and dialkylamino;


35

7h

(3) lower alkoxy carbonyl, carboxyl;

(4) a ketone group of formula:

5




10

in which R^2 represents lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocycl, aryl, lower arylalkyl wherein the alkyl or aromatic ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;

15

(5) a carbamoyl group of formula:




20

with R^2 being as defined above;

(6) amino, lower alkylamino, lower dialkylamino;

(7) amide of formula:


25



with R^2 being as defined above;

(8) a group of formula:

30




35

wherein A is as defined above and the carboxyl terminal of A is directly attached to the nitrogen of the 3-amino-1, 2, 4-thiadiazole, R represents hydrogen, lower alkanoyl, lower cycloalkylcarbonyl, lower alkoxy carbonyl, lower arylalkyloxycarbonyl or N protecting group and R^1 is hydrogen, lower alkyl, lower


alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N atom;

- 5 (9) alcohol of formula:




10 with R² being as defined above;

- (10) sulfone of formula:




15 with R² being as defined above;

- (11) sulfoxide of formula.:



20 with R² being as defined above;

- 25 (12) sulfonamide of formula:



30 with R² being as defined above;


- (13) lower alkylthio, lower arylalkylthio, arylthio;

- (14) a group of formula:

35 -CH₂-A-W

with A as defined above and the N-terminal of A is directly attached to the methylene and W being as defined above;

(15) a group of formula:



in which R^3 and R^4 are independently alkyl, aralkyl, heterocyclyl, heterocyclalkyl; R^3 and R^4 when taken together form with the N-atom a five or a six membered ring selected from the group piperidinyl, pyrrolidinyl, piperazinyl with the N-4 position of piperazine optionally substituted with pyridyl, heterocyclyl, alkyl, aralkyl and aryl.

In a fifth aspect, the invention provides a pharmaceutical composition comprising a compound as described above or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable diluent or carrier thereof.

In a sixth aspect, the invention provides use of a compound or a pharmaceutically acceptable salt thereof as described above for the manufacture of a medicament for treating a mammal with a disease or disorder.

7k

5

BRIEF REFERENCE TO THE DRAWINGS

Fig. 1 is a diagrammatic representation of the chemical interaction between compounds (I) of the present invention and N-acetyl cysteine.


10

Fig. 2 is a diagrammatic representation of the reduction of IL-1-induced degradation of newly synthesized proteoglycan by a compound of formula (I).

DESCRIPTION OF THE PREFERRED EMBODIMENTS

15

One class of preferred compounds according to the invention are compounds corresponding to the following formula (II):



20


or their pharmaceutically acceptable salts thereof,

wherein:

A, W and Y are as previously defined.

25

A second class of preferred compounds according to the present invention are compounds having the general formula (III):



30

or their pharmaceutically acceptable salts thereof,


WO 99/45027

PCT/CA99/00160

8

wherein X, A, W and Y are as previously defined.

5 A further class of preferred compounds according to the present invention are compounds having the general formula (IV):



10

or their pharmaceutically acceptable salts thereof,

wherein:

15 A, R, R₁ and Y are as previously defined.

While the mechanism of action for all these compounds has not been confirmed, it is believed that the 3,5 disubstituted -1,2,4-thiadiazoles of the present invention react with the cysteine residue of the enzyme to form a disulfide bond thus inhibiting the activity of the enzyme. The S-N bond in the 3,5 disubstituted -1,2,4-thiadiazoles of the present invention has a high energy content which originates, at least in part, from non-bonded electron repulsion between sulfur atom d orbitals and nitrogen atom p orbitals. 3,5 disubstituted -1,2,4-thiadiazoles are therefore likely to be susceptible to nucleophilic attack. S-N bond cleavage of 1,2,4-thiadiazoles with reducing agents was reported over forty years ago (Gordeler, Chem. Ber., 1954, 87, 57). The thiol groups of cysteine dependent enzymes appear to act as reducing agents (nucleophiles), thereby becoming chemically modified with resulting inhibition of the enzymatic activity.

30

WO 99/45027

PCT/CA99/00160

9

Group Z at the C-5 position of the 1,2,4-thiadiazoles can be designed to incorporate a recognition sequence specific to the active site of a given cysteine activity dependent enzyme and group Y at the C-3 position may be
5 designed to tune the reactivity of the 5-substituted 1,2,4-thiadiazoles towards the enzyme by activating the adjacent bonds.

The reactivity of the compounds of the present invention towards cysteine activity dependent enzymes can be evaluated by their reactivity with N-acetyl
10 cysteine. Fig. 1 illustrates the reaction between compounds of formula (I) of the present invention with N-acetyl cysteine. The first step forms a disulfide compound by cleavage of the S-N bond of the 1,2,4-thiadiazoles. The disulfide IX reacts with a second thiol to produce a compound of formula X.

15 The compounds of the present invention are those having amino acid or peptide residue side chains. The side chains can be attached at position C3 or C5 positions of 1,2,4-thiadiazole. The use of amino acid or peptide residues as side chains in the monocyclic compounds used in the present invention, particularly when they are attached to the nucleus at position 5,
20 allows selection of an appropriate group having binding affinity for the enzyme which is to be inhibited by the compound. Furthermore, the binding affinity of the inhibitor can be tuned so that it binds to the enzyme at a close proximity of the active site cysteine residue with which it ultimately forms a disulfide bond.

25

The presence of an appropriately chosen enzyme binding or recognition group as a side chain on the compound at a position remote from the -S-N=C- group allows the compound to seek out and bind to the selected
enzyme, to enhance the chemical attack of the thiol group of the enzyme.

30 Due to the presence of recognition side group, compounds of this nature, are highly selective in their attack upon a specific, chosen enzyme, and are much less reactive towards other thiols which they might encounter in a biological system.

WO 99/45027

PCT/CA99/00160

10

Preferred compounds of formula (II) according to the invention are those in which A is glycyl, leucyl-prolyl and isoleucyl-prolyl; W is -NH₂ or -OH; and Y is lower alkyl (methyl), lower alkoxy (methoxy, n butoxy), lower aryl (phenyl),
5 cinnamyl, 1-H-Indol-3-yl-methyl, -CH₂-NR³R⁴ where R³ and R⁴ are as previously defined, -CH₂-A-W where A and W are as previously defined.

Preferred compounds of formula (III) according to the invention are those in which A is leucyl, leucyl-prolyl or isoleucyl-prolyl and Z and Y are as defined
10 above. Particularly preferred compounds of formula (III) are those in which Y is a lower alkoxy.

Preferred compounds of formula (IV) according to the invention are those in which A is phenyl alanyl, glycyl, R₁ is H and R is as previously defined.
15

The preferred compounds according to the present invention exhibit specificity to a particular cysteine activity dependent enzyme and thus are unreactive to other potential nucleophiles such as alcohols or amines.

20 As used herein:

The term "lower", as applied for example to lower alkyl, means 1 to 8 carbon atoms.

25 The term "aryl", alone or in combination, means a phenyl or naphthyl radical which optionally carries one or more substituents selected from alkyl, alkoxy, halogen, hydroxy, amino and the like, such as phenyl, *p*-tolyl, 4-methoxy-phenyl, 4-(*tert*-butoxy)phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-hydroxy-phenyl, 1-naphthyl, 2-naphthyl and the like.

30 The term "arylalkoxy carbonyl", alone or in combination, means a radical of the formula -C(O)-O-arylalkyl, in which the term "arylalkyl" has the significance given above. An example of an arylalkoxy-carbonyl radical is benzyloxycarbonyl.

The term "arylalkyl" means an alkyl radical in which one hydrogen atom is replaced by an aryl radical, such as benzyl, phenylethyl and the like.

- 5 The term "arylalkenyl" means an alkenyl radical in which one hydrogen atom is replaced by an aryl radical such as 3-phenylallyl, 2-phenylallyl, 1-phenylallyl and the like.

- 10 The term "cycloalkylcarbonyl" means an acyl group derived from a monocyclic or bridged cycloalkanecarboxylic acid such as cyclopropanecarbonyl, cyclohexanecarbonyl, adamantanecarbonyl, and the like, or from a benz-fused monocyclic cycloalkanecarboxylic acid which is optionally substituted by, for example, alkylamino, such as 1,2,3,4-tetrahydro-2-naphthoyl, 2-acetamido-1,2,3,4-tetrahydro-2-naphthoyl.

- 15 The term "arylalkanoyl" means an acyl radical derived from an aryl-substituted alkanecarboxylic acid such as phenylacetyl, 3-phenylpropionyl, hydrocinnamoyl, 4-phenylbutyryl, 2-naphthyl-acetyl, 4-chlorohydrocinnamoyl, 4-aminohydrocinnamoyl, 4-methoxyhydrocinnamoyl, and the like.

- 20 The term "aroyl" means an acyl radical derived from an aromatic carboxylic acid. Examples of such radicals include aromatic carboxylic acid, an optionally substituted benzoic or naphthoic acids such as benzoyl, 4-chloro-benzoyl, 4-carboxybenzoyl, 4-[(benzyloxy-carbonyl)] benzoyl, 1-naphthoyl, 2-naphthoyl, 6-carboxy-2-naphthoyl, 6-[(benzyloxy)formamido]-2-naphthoyl, and the like.

- 30 The term "heterocyclyl", as used herein except where noted, represents a stable 5- to 7-membered mono or bicyclic or stable 7- to 10-membered bicyclic heterocyclic ring which is either saturated or unsaturated, and which consists of carbon atoms, and from one to three heteroatoms selected from the group consisting of N, O, S and wherein the nitrogen and sulfur

WO 99/45027

PCT/CA99/00160

12

heteroatoms may be optionally oxidized, and the nitrogen atom may optionally be quaternized, and including any bicyclic group in which any of the above defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic elements, commonly known as heterocyclyl include piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 2-oxoazepinyl, azepinyl, pyrrolyl, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzothieryl, tetrahydroquinolinyl (e.g. 1,2,3,4-tetrahydro-2-quinolinyl, etc), 1,2,3,4-tetrahydro-isoquinolinyl (e.g. 1,2,3,4-tetrahydro-1-oxo-isoquinolinyl, etc.), quinoxaliny, beta-carboliny, 2-benzofurancarbonyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, oxadiazolyl and the like. The heterocycle may be substituted on one or more carbon atoms or heteroatom which results in the creation of a stable structure.

"Amino acid residue" means any of the naturally occurring alpha-, beta-, and gamma-amino carboxylic acids, including their D and L optical isomers and racemic mixtures thereof, and the N-lower alkyl- and N-phenyl lower alkyl-derivatives of these amino acids. The amino acid residue is either bonded through a nitrogen of the amino acid or the carboxyl carbon of the amino acid. The naturally occurring amino acids which can be incorporated into the present invention include, but are not limited to, alanine, arginine, asparagine, aspartic acid, cysteine, cystine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, ornithine, phenylalanine, proline, serine, threonine, thyroxine, tryptophan, tyrosine, valine, beta-alanine, and gamma-aminobutyric acid. Preferred amino acid residues include proline,


WO 99/45027

PCT/CA99/00160

13

lysine, leucine, phenylalanine, tyrosine, isoleucine, alanine, gamma-amino butyric acid, valine, glycine and phenylglycine.

- 5 "Amino acid residues" further includes commonly known synthetic unnatural amino acids and any logically designed peptidomimetic in which the H-bonding elements of the amino acid residue is taken into consideration. Synthetic unnatural amino acids include amino acids such as 4-hydroxy-
 10 proline, O-benzylthreonine, 4-cyclohexylproline, 3,4,5-trimethyl proline, 3,4-dimethylproline, 4,5-dimethylproline, 4-chlorophenylalanine, octahydro-indole-2-carboxylic acid, octahydro-isoquinoline-3-carboxylic acid, piperidiny-2-carboxylic acid, piperaziny-2-carboxylic acid, 4-phenylproline, 3-phenylproline, 4 cyclohexylproline, 3-cyclohexylproline, 4-aminoproline, octahydro-cyclopentane[b]pyrrole-2-carboxylic acid, statone, statine,
 15 norstatine derivatives, 4- amino-3-hydroxy-5-phenyl-pentanoic acid, 4-amino-3-oxo-5-phenyl-pentoic acid, 3-amino-2-hydroxy-4-phenylbutanoic acid, 2,2-difluorostatine, cyclohexylalanine, and the following amino acids:




WO 99/45027

PCT/CA99/00160

14

wherein a stands for 1 or 2. R_a stands for hydrogen, alkyl, hydroxy, alkoxy, carbonylamino.

5 Representative examples of logically designed peptidomimetics are illustrated below:




WO 99/45027

PCT/CA99/00160


15

where R_b , R_c and R_d independently represents hydrogen, and an alkyl.

5



10



15

All alpha-amino acids except glycine contain at least one asymmetric carbon atom. As a result, they are optically active, existing in either D or L form or as a racemic mixture. Accordingly, some of the compounds of the present invention may be prepared in optically active form, or as racemic mixtures of the compounds claimed herein.

A peptide residue contains a peptide bond which may be formed between the carbonyl function of an amino acid and another amino compound which may be another amino acid or an amine. Compound of the present invention may be in isosteric form e.g., $-\text{CH}_2\text{NH}-$ (reduced), $-\text{COCH}_2-$ (keto), $-\text{CH}(\text{OH})(\text{CH}_2-$ (hydroxy), $-\text{CH}(\text{NH}_2)\text{CH}_2-$ (amino), $-\text{CH}_2\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2\text{CH}_2-$ (hydrocarbon). Preferably a compound of the present invention has no peptidic carbamoyl in isosteric form. When it has peptidic carbamoyl groups in isosteric form, it has one or two, preferably one peptidic bond in isosteric form.

WO 99/45027

PCT/CA99/00160

16

A peptide residue consists of amino acid residues, preferably in their natural configuration. When there are amino acids in unnatural configuration there preferably is only one such amino acid in unnatural configuration. Amino acid
5 residue as used herein includes amino acid residues such as octahydro-indole-2-carboxylic acid and hydroxyproline.

The term "R", when referred as an "N-protecting group", is an amino protecting group. Examples of protecting groups as R are for example
10 disclosed in "Protective Groups in Organic Synthesis", T.V. Greene, J. Wiley & Sons NY (1981) 219-287. These include but are not limited to acyl such as acetyl, methoxysuccinyl, hydroxysuccinyl or benzoyl optionally substituted on the phenyl ring with for example p-methoxycarbonyl, p-methoxy or p-nitro; alkoxycarbonyl such as t-butyloxycarbonyl; arylmethoxycarbonyl such as
15 9-fluorenylmethoxycarbonyl or benzyloxy carbonyl optionally substituted on the phenyl ring with p-methoxy, p-nitro, p-chloro or m-phenyl; arylmethyl such as benzyl optionally substituted on the aromatic ring with p-methoxy, p-nitro or p-chloro; or arylsulfonyl such as phenylsulfonyl optionally substituted with p-methyl or p-methoxyl or naphthylsulfonyl optionally substituted on the
20 aromatic ring with example amino or dialkylamino.

The term "aryloxyalkanoyl" means an acyl radical of the formula aryl-O-alkanoyl and the term "heterocyclyloxycarbonyl" means an acyl group derived from heterocyclyl-O-CO- wherein heterocyclyl is defined above.
25

The term "heterocyclylalkanoyl" means an acyl radical derived from a heterocyclyl-substituted alkane carboxylic acid wherein heterocyclyl has the same meaning given above.
30

WO 99/45027

PCT/CA99/00160

17

The term "heterocyclyalkoxycarbonyl" means an acyl radical derived from a heterocyclyl-substituted alkyl-O-CO wherein heterocyclyl has the same significance given above.

5

The term "aminoalkanoyl" means an acyl radical derived from an amino-substituted alkanecarboxylic acid wherein the amino group can be a primary, secondary or tertiary amino group containing substituents selected from hydrogen, and alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl radicals and the like.

10

"Optional" or "optionally" means that the subsequently described event or circumstances may or may not occur, and that the description includes instances where the said event or circumstance occurs and instances in which it does not. For example, "phenyl ... optionally substituted" means that the phenyl may or may not be substituted and that the description includes both unsubstituted phenyl and phenyl wherein there is substitution.

15

Certain of the compounds of the invention have chiral centers and exist as optical antipodes. The invention described and claimed herein includes each of the individual enantiomers as well as their racemic modifications and the racemic mixture.

20

"Pharmaceutically acceptable, non-toxic salts" refers to pharmaceutically acceptable salts of the compounds of this invention which retain the biological activity of the parent compounds and are not biologically or otherwise undesirable (e.g. the salts are stable). Salts of the two types may be formed from the compounds of this invention: (1) Salts of inorganic and organic bases from compounds of formulae I, II, III and IV which have a carboxylic acid functional group and (2) Acid addition salts may be formed at the amine functional group of many of the compounds of this invention.

25

30

WO 99/45027

PCT/CA99/00160

18

Pharmaceutically acceptable salts derived from inorganic bases include sodium, potassium, lithium, ammonium, calcium, magnesium, ferrous, zinc, copper, manganous, aluminum, ferric, manganic salts and the like.

5 Particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Pharmaceutically acceptable, non-toxic salts derived from organic bases include salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins. Such salts are exemplified by, for
10 example isopropylamine, trimethyl-amine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, tromethamine, dicyclohexamine, lysine, arginine, histidine, caffeine, procaine, hydrabramine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, *N*-ethylpiperidine, polyamine
15 resins and the like. Particularly preferred organic non-toxic bases are isopropylamine, diethylamine, ethanolamine, piperidine, tromethamine, dicyclohexylamine, choline and caffeine.

Pharmaceutically acceptable acid addition salts are formed with inorganic
20 acids such as halo acids, sulfuric acid, nitric acid, phosphoric acid and the like and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid, salicylic
25 acid and the like.

The term "animals" refers to humans as well as all other animal species, particularly mammals (e.g. dogs, cats, horses, cattle, pigs, etc.), reptiles, fish, insects and helminths.

30

WO 99/45027

PCT/CA99/00160

19

The specific, most preferred compounds according to the present invention are the following:

- 5 2-(3-methoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide
 2-(3-n-Butoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide
 2-[3-(3-Phenylallyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide
 2-[3-(1*H*-Indol-3-ylmethyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide


- 10 *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline
 N-(3-Butoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline
 N-(3-Phenyl-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline
 N-(3-Methyl-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline
 N-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-isoleucyl-L-proline

15

N,N'-[3-methylene-[1,2,4]thiadiazol-5-yl]-di-(L-leucyl-L-proline methyl ester),

which has the following chemical formula:


20



25

· 3-[4-(2-pyridyl)piperazinylmethyl]-1,2,4-thiadiazol-5-yl-leucyl-proline,
 which has the following chemical formula:

30



WO 99/45027

PCT/CA99/00160

20

5-{3-methoxy-1,2,4-thiadiazolyl}carbamoyl-leucyl isoamylamide


5-{3-methoxy-1,2,4-thiadiazolyl}carbamoyl-isoleucyl isoamylamide

{3-methoxy-[1,2,4]-thiadiazol-5-yl}carbamoyl-L-leucyl-L-proline.

3-methoxy-5-carbobenzyloxy-phenylalaninamido-[1,2,4]-thiadiazole.


3,5-di-(*N*-carbobenzyloxy-L-phenylalaninamido)-[1,2,4]thiadiazole,

which has the following chemical formula:



The present invention provides synthetic methods for preparing compounds according to the invention. The choice of method depends largely upon the selected Z and Y group that is the substituent on C3 and C5 positions in the final compound.

The compounds of formula II are prepared by reaction of a compound of formula 3 with a primary or secondary amine. Examples of those amines are 2-pyridylpiperazine and leucyl-proline methyl ester. This method is appropriate for compounds in which Y is lower alkyl, lower alkoxy, heterocyclyl, 1-haloalkyl, aryl, dialkylamino:



WO 99/45027

PCT/CA99/00160


21

The reaction is normally carried out in an inert solvent such as N,N-dimethylformamide, tetrahydrofuran, dichloromethane, acetonitrile, dimethylsulfoxide in the presence or absence of a base such as 1 to 3 mole of triethylamine per mole of compound 3. These solvents may be used singularly or in combination of any ratio as necessary. Reaction temperature can be chosen over the range of from 0°C to 150°C, being preferably about 10 to 65°C. Reaction time is normally about 1 to 50 hours, preferably 1 to 8 hours. The amount of amine used is 1 to 3 mole per mole of compound 3.

The compounds of formula 3 can be produced by treating the corresponding amidine derivative with perchloromethyl mercaptan in a two phase mixture of dichloromethane and sodium hydroxide at 0 to 25°C for 2 to 6 hours.

Preparative methods for compound 1 include those reported in US patent 3,324,141; J. Org. Chem., 1962, 27, 2589; Chem. Ber., 1957, 90, 182.

Compounds of formula III wherein the spacer X is -NH-CO-CH₂ are prepared by N-alkylating a compound of formula (4) with an amine:




The compounds are isolated by conventional means. Compounds (4) are in turn prepared by reacting compound (5) with bromoacetyl bromide in an inert solvent such as methylene chloride or tetrahydrofuran in the presence of a base such as triethylamine. Compounds (5) are prepared from the amidine (1) with potassium thiocyanate (KSCN) in the presence of base such as sodium hydroxide in an inert solvent such as acetone and water.

WO 99/45027


PCT/CA99/00160

22

Compounds of formula III in which the spacer X is -NH-CO- are prepared by reaction of a compound (5) with H-A-W in the presence of 1,1-carbonyl diimidazole in an inert solvent such as THF or DMF.



The compounds of formula (IV) are produced by peptide coupling between compound (5) with N-protected amino acid or N-protected peptide carboxylic acid R-A-OH using conventional peptide coupling reagent such as 1,1-carbonyl di-imidazole or diphenylphosphoryl azide in an inert solvent such as dimethylformamide or tetrahydrofuran:



20

25 The compounds R-A-OH are generally commercially available from Sigma-Aldrich Inc., for example, carbobenzyloxy-L-phenylalanine, carbobenzyloxy-L-phenylalanyl-L-alanine (R-A'-OH as a N-protected amino acid), N-t-Boc-3-amino-2-hydroxy-5-methyl-hexanoic acid (R-A'-OH as a protected dipeptide with the peptide bond in carbamoyl form), N-t-Boc-3-amino-2-hydroxy-4-phenylbutyric acid (R-A'-OH as a N-protected peptide in isosteric form). If not


30 commercially available such compounds can be prepared according to the method disclosed in Synthetic Peptides, vol. 1 by George R. Pettit, Van Nostrand Reinhold, 1970.

WO 99/45027

PCT/CA99/00160

23


In an alternative route for the preparation of compounds of formula (IVA) within the scope of this invention, a compound of formula (6) is reacted with N-protected amino acid or peptide acid R-A-OH using conventional peptide coupling reagent such as 1,1-carbonyl di-imidazole or diphenylphosphoryl azide in an inert solvent such as dimethylformamide or tetrahydrofuran.



The preparation of pseudopeptides within the scope of the present invention involves the use of the bis-chloro derivative 7 as a starting material. The amines H-A-W reacts with compound 7 in the presence of a phase transfer catalyst such as tetra-N-butylammonium bromide in an inert solvent such as dimethylformamide at room temperature over a period of 20 to 30 hours to give compound 8. This reaction is temperature dependent. At elevated temperature, preferably 70-90°C, the disubstituted product compound II_A is formed. Compound 8 also reacts with other amines such as H-A-W or R³R⁴NH to give the compounds of formula II_A and II_B respectively.

WO 99/45027

PCT/CA99/00160



Certain compounds of this invention may be converted to their corresponding pharmaceutically acceptable acid addition salts by virtue of the presence of a basic amine nitrogen. These compounds may be converted from the free base form to various acid addition salts by treating with a stoichiometric excess of the appropriate organic or inorganic acid, such as, for example, phosphoric, pyruvic, hydrochloric or sulfuric acid and the like. Typically, the free base is dissolved in a polar organic solvent such as *p*-dioxane or dimethoxyethane, and the acid added thereto. The temperature is maintained between 0°C and 50°C. The resulting acid addition salt precipitates spontaneously or may be precipitated out of solution with a less polar solvent. These acid addition salts may be decomposed to the corresponding free base by treating with a stoichiometric amount of a suitable base, such as potassium carbonate or sodium hydroxide, typically in the presence of aqueous solvent, and at a temperature of between about 0°C and 50°. The free base form is isolated by conventional means, such as extraction with an organic solvent. Acid addition salts of the compounds of the present invention may be interchanged by taking advantage of differential solubilities of the salts, volatilities or acidities of the acids, or by treating with an appropriately loaded ion exchange resin. For example, the interchange is effected by the reaction of a salt of the compounds of formula I with a slight stoichiometric excess of an acid of a lower pKa than the acid component of the starting salt. This is carried out at a temperature between about 0°C and the boiling point of the solvent being used.

- For the treatment of diseases and/or disorders herein above referred to, the compounds of the present invention may be used orally, or parenterally in formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injection or infusion techniques. In addition to the treatment of warm-blooded animals such as mice, rats, horses, cattle, sheep, dogs, cats, etc., the compounds of the invention are effective in the treatment of humans.
- For compositions, conventional non-toxic solid carriers include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like may be used. The active compound as defined above may be formulated as liquid pharmaceutically administrable compositions can, for example, be prepared by mixing, dissolving, dispersing, etc. the active compounds as defined above the optional pharmaceutically adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, to hereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain a minor amount of non-toxic auxiliary substances such as wetting or emulsifying agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, etc. Actual methods of preparing such dosage forms are known, or will be apparent to those skilled in this art: for example, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa, 15th Edition, 1975. The composition of formulation to be administered will, in any event, contain a quantity of the active compounds in an amount effective to alleviate the symptoms of the subject being treated.
- The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard and

WO 99/45027

PCT/CA99/00160

26

soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions contain one or more
5 agents from the group consisting of sweetening agents, flavouring agents, colouring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations.

10 Tablets contain the active ingredient in admixture with the non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. The excipients may be for example, inert diluents, such as calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding
15 agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be coated by known techniques to delay the disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over long period.

20 Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredients are mixed with inert solid diluent, for example, calcium phosphate or kaolin, or as soft gelating capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

25 Aqueous suspensions contain the active materials in admixture with the excipient suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethyl-cellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum and gum acacia; dispersing or wetting agents may
30 be a naturally-occurring phosphate, for example lecithin, or condensation products of an alkene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecathyl-eneoxycetanol, or condensation

products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, P-hydroxy-benzoate, one or more colouring agents, such as sucrose or saccharin. Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with the dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional recipients, for example sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical composition of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphates, esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial ester with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavouring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavouring and colouring agents. The pharmaceutical compositions may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol.

Among the acceptable vehicles and solvents that may be employed are water, Ringer's solutions and isotonic sodium chloride solution. In addition, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Parenteral administration is generally characterized by injection, either subcutaneously, intramuscularly or intravenously. Injectables can be prepared in conventional forms, either as liquid solutions or suspension in liquid prior to injection, or as emulsions. Suitable excipients are for example, water, saline, dextrose, glycerol, ethanol or the like. In addition, if desired, the pharmaceutical compositions to be administered may also contain minor amounts of non-toxic auxiliary substance such as wetting or emulsifying agents, pH buffering agents and the like, such as for example, sodium acetate, sorbitan monolaurate, triethanolamine oleate, etc.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration of humans may contain from 0.5 mg to 5 mg of active agent compounded with an appropriate and convent amount of carrier material which may vary from about 5 to about 95% of the total composition. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet,
5 time of administration, drug combination and the severity of the particular disease undergoing therapy.

The invention is further described and illustrated in the following specific examples.

10

Example 1

A. Preparation of Amidines, compounds of formula I.

Selected amidines were from commercial sources (Aldrich and/or
15 Lancaster). Those which were not commercially available were prepared according to published procedures. [Sandler, S. R. and Karo, W. in Organic Functional Group Preparations, 2nd Ed., Academic Press, Inc., Toronto, Volume II, Chapter 7, 1986 and Volume III, Chapter 6, 1989 and references cited therein; MacLeod, A. M. et al. in J. Med. Chem. 1990, 33, 2052-2059].

20

B. Preparation of 2-(3-Chlorophenyl)-2-(tetrahydropyran-2-yloxy)acetamidine hydrochloride.

To a solution of sodium bisulfite (27.66 g, 0.266 mol) in water (120 mL) was added 3-chlorobenzaldehyde (25.0 g, 0.178 mol). After stirring for 20
25 min, a solution of sodium cyanide (12.4 g, 0.2525 mol) in water (80 mL) was added dropwise. Ethyl acetate (50 mL) was added and the resulting mixture was stirred for another 3 h. The organic layer was collected, washed with brine, dried (sodium sulfate), filtered and evaporated to dryness. The residue was dissolved in dichloromethane (200 mL) and cooled in ice. Dihydropyran
30 (20.62 g, 0.2451 mol) and pyridinium p-toluene sulfonate (6.5 g, 25.9 mmol) were then added successively and the mixture was stirred at room temperature for 16 h. The organic layer was successively washed with water

WO 99/45027

PCT/CA99/00160

30

(50 mL), 5% sodium bisulfite solution (2x50 mL) and brine (50 mL). The organic fraction was dried (sodium sulfate), filtered and evaporated to give an oil. Purification by column chromatography on silica gel using a mixture of hexane and ethyl acetate (95/5) as eluant afforded 31.5 g (70%) of 2-(3-chlorophenyl)-2-[(tetrahydropyran-2-yloxy)]acetonitrile (mixture of diastereoisomers) as a clear oil which was contaminated with <5% of 3-chlorobenzaldehyde. ¹H-NMR (CDCl₃) δ 7.30-7.60 (m, 4H, Ar-H), 5.59 (s, 0.3H, αH) and 5.41 (s, 0.7H, αH), 5.12 (t, J=2.9 Hz, 0.3H, OCHO) and 4.77 (t, J=2.9 Hz, 0.7H, OCHO), 3.60-4.05 (m, 2H, CH₂O), 1.50-2.06 (m, 6H, 3 CH₂). Small pieces of sodium (192 mg, 7.9 mmol) was added to anhydrous ethanol (200 mL) under a positive pressure of nitrogen at room temperature. After all the sodium had dissolved, 2-(3-chlorophenyl)-2-[(tetrahydropyran-2-yloxy)]acetonitrile (19.54 g, 77.6 mmol) was added neat and the contents of the flask was rinsed with ethanol (20 mL) and added to the reaction mixture. The progress of the reaction can be monitored by TLC using a mixture of hexane and ethyl acetate (1/1) as solvent system. After stirring for 16 h, the reaction mixture was cooled to ca. -40°C (dry ice-methanol-water) and an ethanolic solution of ammonia (77.6 mL of a 2M solution, 0.1552 mol) was quickly added followed by solid ammonium chloride (4.02 g, 75.15 mmol). The reaction mixture was allowed to warm to room temperature overnight. The reaction mixture was filtered over a pad of celite. The filtrate was collected and evaporated to dryness and then pumped under high vacuum to afford 2-(3-chlorophenyl)-2-(tetrahydropyran-2-yloxy)acetamidine hydrochloride (23 g) as a white foam in almost quantitative yield. ¹H-NMR (CDCl₃) δ 9.00-9.80 (br., 1H, NH), 8.00-8.80 (br., 1H, NH), 7.30-7.64 (m, 3H, Ar-H), 5.85 (s, 0.4H, αH) and 5.75 (s, 0.6H, αH), 4.84 (t, J=3.3 Hz, 0.4H, OCHO) and 4.57 (t, J=3.3 Hz, 0.6H, OCHO), 3.41-3.83 (m, 2H, CH₂O), 2.20-2.60 (br., 1H, NH), 1.46-1.86 (m, 6H, 3 CH₂).

C. In a similar manner, 2-pyridyl-2-(tetrahydropyran-2-yloxy)acetamidine hydrochloride salt was prepared as an orange foam (crude yield 96.0%). ¹H-NMR (CDCl₃) δ 9.00-9.80 (br., 1H, NH), 8.00-8.80 (br., 1H, NH), 8.56-8.61 (d, J=4.4 Hz, 1H, Py-H), 7.28-7.84 (m, 3H), 5.75 (s, 0.4H, αH) and 5.67 (s, 0.6H, αH), 4.93 (t, J=2.5 Hz, 0.4H, OCHO) and 4.71 (t, J=2.5 Hz, 0.6H, OCHO), 3.49-3.93 (m, 2H, CH₂O), 1.53-1.92 (m, 6H, 3 CH₂).

10 Example 2

A. Preparation of 5-Chloro-3-methoxy-[1,2,4]thiadiazole.

To an ice-cooled mixture of O-methylisourea hydrochloride (11.06 g, 0.1mol) in chloroform (150 mL) was added a solution of perchloromethyl mercaptan (11.0 mL, 0.1 mol) in chloroform (50 mL) dropwise over a period of 45 min. Then, a cold solution of sodium hydroxide (16 g, 0.4 mol in 30 ml water) was added dropwise while maintaining the reaction temperature to <5°C. The progress of the reaction was monitored by TLC using a mixture of hexane and ethyl acetate (9/1) as eluent. After 3 h at room temperature, the organic layer was collected, dried (sodium sulfate), filtered and then concentrated in vacuo. Purification by column chromatography on silica gel using a solvent gradient of a mixture of hexane and ethyl acetate (100/0, 95/5 then 9/1) afforded 5-chloro-3-methoxy-[1,2,4]thiadiazole (6.01 g, 39.9%) as a light yellow oil. ¹H-NMR (CDCl₃) δ 4.02 (s, OMe, 3H); ¹³C-NMR (CDCl₃) δ 173.56 (C3), 169.68 (C5), 57.07 (OMe); MS (APCI) m/z 151.1 (M⁺+1), 108.0, 94.0, 73.0, 58.0.

B. In a similar fashion, the following compounds were prepared:
3-Butoxy-5-chloro-[1,2,4]thiadiazole, light yellow oil, yield (50.9%). ¹H-NMR (CDCl₃) δ 4.35 (t, J=6.6 Hz, 2H, CH₂), 1.75 (m, 2H, CH₂), 1.44 (m, 2H, CH₂), 0.93 (t, J=7.1Hz, 3H, CH₃); ¹³C-NMR (CDCl₃) δ 173.27 (C3), 169.37 (C5), 70.17 (C1'), 30.70 (C2'), 18.96 (C3'), 13.67 (C4').

WO 99/45027

PCT/CA99/00160

32

5 *N*-(5-Chloro-[1,2,4]thiadiazol-3-yl)-acetamide or 3-acetamido-5-chloro-1,2,4-thiadiazole, light yellow solid on standing, yield (34.8%). ¹H-NMR (CDCl₃) δ 2.50 (s, Me); ¹³C-NMR (CDCl₃) δ 173.97 (C3), 171.86 (CO), 164.07 (C5), 22.54 (CH₃).

10 3-(5-Chloro-[1,2,4]thiadiazol-3-ylmethyl)-1*H*-indole or 5-chloro-3-{1*H*-indol-3-yl-methyl}-1,2,4-thiadiazole, light brown solid, yield (28%). ¹H-NMR (CDCl₃) δ 8.29 (br. s, 1H, NH), 7.68 (d, J=7.8 Hz, 1H, CHNH), 7.11-7.33 (m, 4H, Ar-H), 4.48 (s, 2H, CH₂); ¹³C-NMR (CDCl₃) δ 175.50 (C3), 173.00 (C5), 136.31, 127.13, 123.19, 122.26, 119.70, 119.05, 111.41, 110.55, 29.97 (CH₂); MS *m/z* 249.7 (M⁺), 130.1, 117.1.

15

Example 3

A. Preparation of 5-Chloro-3-[3-chlorophenyl-1-(tetrahydropyran-2-yloxy)methyl]-[1,2,4]thiadiazole.

20 To an ice-cooled solution of 2-(3-chlorophenyl)-2-(tetrahydropyran-2-yloxy)acetamidine hydrochloride (12.0 g, 39.51 mmol) in dichloromethane (20 mL) and sodium hydroxide (9.49 g, 0.237 mol, dissolved in 60 mL water) was added a solution of perchloromethyl mercaptan (9.18 g, 49.46 mmol) in dichloromethane (50 mL) over a period of 35 min. The reaction mixture was stirred at ice-cold temperature for a further 1h and the organic layer was
25 collected, dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography on silica gel using a mixture of hexane and ethyl acetate (94/6) afforded the title compound as a light yellow oil (8.4 g, 61.5%). ¹H-NMR (CDCl₃) δ 7.56 (d, J=6.9 Hz, 1H, Ar-H), 7.39-7.44 (m, 1H, Ar-H), 7.28-7.32 (m, 2H, Ar-H), 6.08 and 6.03 (s, 0.5H each, α-H),
30 4.84 and 4.72 (t, J=3.1 Hz, 0.5H each, OCHO), 3.49-3.92 (m, 2H, OCH₂), 1.55-1.94 (m, 6H, 3CH₂).

WO 99/45027

PCT/CA99/00160

33

B. In a similar manner, the following compounds were prepared:

- 5 3-[(5-Chloro-[1,2,4]thiadiazol-3-yl)-(tetrahydropyran-2-yloxy)methyl]-pyridine, brown oil, 28.7% yield (mixture of diastereoisomers). ¹H-NMR (CDCl₃) δ 8.55 and 8.52 (d, J=4.5 Hz, 0.67 and 0.33 H each), 7.69-7.81 (m, 2H), 7.18-7.28 (m, 1H), 6.20 and 6.15 (s, 0.33 and 0.67 H each, ArCHO), 4.81-4.87 (m, 1H, OCHO), 3.84-3.90 (m, 1H, OCH), 3.44-3.54 (m, 1H, OCH), 1.90-1.96 (m, 1H),
10 1.75-1.80 (m, 2H), 1.51-1.60 (m, 3H); MS m/z 312.1 (M⁺+1), 228, 210, 85. 5-Chloro-3-chloromethyl-[1,2,4]thiadiazole, brown oil, 83.6% yield. ¹H-NMR (CDCl₃) δ 4.70 (s); ¹³C-NMR (CDCl₃) δ 174.33 (C3), 170.56 (C5), 40.17 (CH₂).

Example 4

- 15 Preparation of 2-(3-methoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide.

To a suspension of glycine hydrochloride (0.55 g, 5 mmol) in DMF (10 mL) was added triethylamine (1.7 mL, 12 mmol) and the resulting mixture was stirred for 15 min at room temperature. This was added to a solution of 5-chloro-3-methoxy-[1,2,4]thiadiazole (0.38 g, 2.5 mmol, prepared according to
20 the procedure by Goerdeler, J. et al. in Chem. Ber. 1955, 88, 843) in DMF (15 mL). Then, tetra-butylammonium bromide (50 mg) was added and the resulting suspension was heated at 80-85°C for 1.5 h. On cooling to room temperature, the reaction mixture was diluted with water and ethyl acetate. The organic layer was collected, washed with brine, sat. Sodium bicarbonate
25 solution, brine, dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography on silica gel using a solvent mixture of dichloromethane and methanol (95/5 and 9/1) afforded
2-(3-methoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide (75 mg, 16%) as a white solid. ¹H-NMR (MeOD) δ 4.03 (s, 2H, NCH₂), 3.94 (s, 3H, OMe); ¹³C-NMR (MeOD) δ 182.0, 173.3, 170.0, 56.7 (MeO), 47.9; MS (m/z) 189 (M⁺+1), 172,
30 144, 87.

WO 99/45027

PCT/CA99/00160

34

Example 5

5 A. Preparation of (3-methoxy-[1,2,4]thiadiazol-5-ylamino)-acetic acid methyl ester or N-(3-methoxy-[1,2,4]thiadiazol-5-yl)-glycine methyl ester.

In a similar manner as described in example 4, a suspension of glycine methyl ester hydrochloride (0.63 g, 5 mmol) in DMF (10 mL) was added triethylamine (1.7 mL, 12 mmol) and the resulting mixture was stirred for 15 min at room temperature. This was added to a solution of 5-chloro-3-methoxy-[1,2,4]thiadiazole (0.38 g, 2.5 mmol) in DMF (15 mL). Then, tetra-butylammonium bromide (50 mg) was added and the resulting suspension was heated at 80-85°C for 1.5 h. On cooling to room temperature, the reaction mixture was worked up as usual and the product was purified by column chromatography on silica gel using a solvent mixture of hexane and ethyl acetate (6/4) thereby affording (3-methoxy-[1,2,4]thiadiazol-5-ylamino)-acetic acid methyl ester (310 mg, 60.1%) as a white solid. ¹H-NMR (CDCl₃) δ 7.20 (br.s, 1H, NH), 4.12 (br. s, 2H, NCH₂), 3.91 (s, 3H, OMe), 3.74 (s, 3H, OMe); ¹³C-NMR (CDCl₃) δ 182.5 (C3), 170.0 (C=O), 168.1 (C5), 56.2 (OMe), 52.6 (OMe), 46.1 (CH₂).

20

B. Proceeding in a similar manner as described in example 5A, the following compounds were prepared:

(3-n-Butoxy-[1,2,4]thiadiazol-5-ylamino)-acetic acid methyl ester or N-(3-n-butoxy-[1,2,4]thiadiazol-5-yl)-glycine methyl ester, off-white solid on standing, 65% yield. ¹H-NMR (CDCl₃) δ 6.98 (br. s, 1H, NH), 4.29 (t, J=5.0 Hz, 2H, OCH₂), 4.18 (br. s, 2H, NCH₂), 3.80 (s, 3H, OMe), 1.75 (m, 2H, OCH₂ CH₂), 1.46 (m, 2H, CH₂ CH₃), 0.94 (t, J=7.1 Hz, 3H, CH₃); ¹³C-NMR (CDCl₃) δ 182.1 (C3), 170.0 (C=O), 167.8 (C5), 69.0 (OCH₂), 52.7 (OMe), 46.1 (NCH₂), 30.9, 19.0, 13.7.

30

WO 99/45027

PCT/CA99/00160

35

- 5 [3-(3-Phenyl-allyl)-[1,2,4]thiadiazol-5-ylamino]-acetic acid methyl ester or N-{3-cinnamyl-[1,2,4]thiadiazol-5-yl}-glycine methyl ester; off-white solid, 68% yield. ¹H-NMR (CDCl₃) δ 7.18-7.40 (m, 6H, Ar-H and NH), 6.42-6.55 (m, 2H, HC=CH), 4.12 (br. s, 2H, NCH₂), 3.75 (s, 3H, OMe), 3.66 (d, J=6.0 Hz, 2H, C=CCH₂); ¹³C-NMR (CDCl₃) δ 183.6 (C3), 172.0 (C5), 170.0 (C=O), 137.2, 132.3, 128.5 (C-ortho), 127.3, 126.3 (C-meta), 125.2, 52.6 (OMe), 46.7 (NCH₂), 36.8 (C=CCH₂).
- 10 [3-(1*H*-Indol-3-ylmethyl)-[1,2,4]thiadiazol-5-ylamino]-acetic acid methyl ester or N-{3-(1*H*-Indol-3-ylmethyl)-[1,2,4]thiadiazol-5-yl}-glycine methyl ester; light brown solid, 71.4% yield. ¹H-NMR (CDCl₃) δ (rotamers) 8.36 (br. s, 1H, NH), 7.72 and 7.64 (d, J=7.7 Hz, 0.25H and 0.75H each, C=CH), 6.93-7.28 (m, 5H, Ar-H and NH), 4.22 and 4.19 (s, 0.6H and 1.4H each, In-CH₂), 3.87 (s, 2H, NCH₂), 3.71 (s, 3H, OMe); ¹³C-NMR (CDCl₃) δ 183.6 (C3), 172.0 (C5), 169.8 (C=O), 136.3, 127.3, 123.0, 121.9, 119.3, 119.2, 111.4, 111.3, 52.6 (OMe), 46.5 (NCH₂), 29.7 (InCH₂).
- 15 [3-[1-Phenyl-1-(tetrahydropyran-2-yloxy)-methyl]-[1,2,4]thiadiazol-5-ylamino]-acetic acid methyl ester or N-{3-[1-Phenyl-1-(tetrahydropyran-2-yloxy)-methyl]-[1,2,4]thiadiazol-5-yl}glycine methyl ester; colorless oil, 89% yield. ¹H-NMR (CDCl₃) δ (diastereoisomers) 7.45-7.55 (m, 2H, Ar-H), 7.23-7.40 (m, 3H, Ar-H), 7.00 and 6.65 (br. s, 0.5H and 0.5H each, NH), 5.91 and 5.87 (s, 0.5H and 0.5H each, aH), 4.85 and 4.66 (t, 0.5H and 0.5H each, OCHO), 4.06 (s, 2H, NCH₂), 3.78 (s, 3H, OMe), 3.52-4.05 (m, 2H, OCH₂), 1.56-1.92 (m, 6H, 3CH₂).
- 20
- 25

30

C. Proceeding in a similar manner as described in example 5A, the following compounds are prepared:

- 5 N-(3-methoxy-[1,2,4]-thiadiazol-5-yl)-leucine methyl ester
N-(3-methoxy-[1,2,4]-thiadiazol-5-yl)-isoleucine methyl ester
N-(3-methoxy-[1,2,4]-thiadiazol-5-yl)-alanine methyl ester
N-(3-methoxy-[1,2,4]-thiadiazol-5-yl)-proline methyl ester
N-(3-methoxy-[1,2,4]-thiadiazol-5-yl)-phenylalanine methyl ester
10 N-(3-methoxy-[1,2,4]-thiadiazol-5-yl)-tyrosine methyl ester
N-(3-methoxy-[1,2,4]-thiadiazol-5-yl)-serine methyl ester
N-(3-methoxy-[1,2,4]-thiadiazol-5-yl)-threonine methyl ester

- D. Preparation of *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-proline methyl ester.
15

To a suspension of L-proline methyl ester hydrochloride (331 mg, 2 mmol) in DMF (15 mL) was added triethylamine (0.56 mL, 4 mmol) and the resulting mixture was stirred for 10 min at room temperature. Then, a solution of 5-chloro-3-methoxy-[1,2,4]thiadiazole (151 mg, 1 mmol) in DMF (5 mL)
20 was added dropwise. The resulting reaction mixture was stirred at room temperature for 16h. The mixture was diluted with water and ethyl acetate. The organic layer was collected, washed with brine, dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography on silica gel using a solvent mixture of hexane and ethyl acetate (1/1) afforded
25 *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-proline methyl ester (210 mg, 86.4%) as a light yellow oil. ¹H-NMR (CDCl₃) δ 4.50 (m, 1H, CHCO₂), 3.96 (s, 3H, OMe), 3.75 (s, 3H, OMe), 3.55 (m, 1H, NCHpro), 3.40 (m, 1H, NCHpro), 2.11-2.35 (m, 4H, 2CH₂pro).

Example 6

A. Preparation of 2-(3-methoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide or
5 b-N-(3-methoxy-[1,2,4]thiadiazol-5-yl)-glycinamide.

The ester (101 mg, 0.5 mmol) was stirred with aq. ammonia (0.5 mL of
a 28-30% solution) for 30 min. Acetonitrile (15 mL) was added and volatile
materials were removed in vacuo. The residue was purified by column
chromatography on silica gel using a solvent mixture of dichloromethane and
10 methanol (9/1) thereby affording
2-(3-methoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide (92 mg, 98%) as a white
solid.

B. In a similar manner as described in example 6A, the following
15 compounds were prepared:

2-(3-n-Butoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide or β -N-(3-n-butoxy-
[1,2,4]thiadiazol-5-yl)-glycinamide; white solid, 95.7% yield. M.p. 148.5-
149.5°C; $^1\text{H-NMR}$ (DMSO) δ 8.51 (br. s, 1H, NH), 7.51 (br. s, 1H, NH), 7.14
(br. s, 1H, NH), 4.18 (t, J=6.6 Hz, 2H, OCH_2), 3.88 (br. d, J=2.8 Hz, 2H, NCH_2),
20 1.63 (m, 2H, OCH_2CH_2), 1.38 (m, 2H, CH_2CH_3), 0.90 (t, J=7.2 Hz, 3H, CH_3);
 $^{13}\text{C-NMR}$ (DMSO) δ 181.6 (C3), 169.9 (C=O), 167.1 (C5), 67.9 (OCH_2), 46.8
(NCH_2), 30.5, 18.6, 13.6; MS (m/z) 231 ($\text{M}^+ + 1$), 253, 231, 202, 175, 99, 83,
61.

25 2-[3-(3-Phenylallyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide or β -N-(3-
cinnamyl-[1,2,4]thiadiazol-5-yl)-glycinamide; white solid, 85% yield. M.p.
150.0-152.5°C; $^1\text{H-NMR}$ (MeOD) δ 7.37-7.39 (m, 2H, Ar-H), 7.27-7.31 (m,
2H, Ar-H); 7.18-7.22 (m, 1H, Ar-H), 6.42-6.55 (m, 2H, HC=CH), 4.09 (s, 2H,
 NCH_2), 3.59 (d, J=6.2 Hz, 2H, C=CCH_2); MS (m/z) 275 ($\text{M}^+ + 1$), 258, 230, 189,
30 152, 117, 91.

WO 99/45027

PCT/CA99/00160

38

2-[3-(1*H*-Indol-3-ylmethyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide or β -N-[3-(1*H*-Indol-3-ylmethyl)-[1,2,4]thiadiazol-5-yl]-glycinamide; off-white solid, 94.0% yield. M.p. 200.0-200.5°C; ¹H-NMR (DMSO) δ 8.46 (t, J=5.4 Hz, 1H, NH), 7.58 (d, J=7.9 Hz, 1H), 7.53 (br. s, 1H, NH), 7.35 (d, J=8.1 Hz, 1H), 7.20 (br. s, 1H), 7.07 (t, J=7.3 Hz, 1H), 6.97 (t, J=7.5 Hz, 1H), 4.04 (s, 2H, In-CH₂), 3.94 (d, J=5.4 Hz, 2H, NCH₂); ¹³C-NMR (DMSO) δ 182.9 (C3), 171.7 (C5), 170.2 (C=O), 136.2, 127.2, 123.4, 120.9, 118.9, 118.3, 111.3, 110.5, 47.3 (NCH₂), 29.3 (InCH₂); MS (m/z) 288 (M⁺+1), 271, 243, 184, 155, 130.

2-[3-[Phenyl-(tetrahydropyran-2-yloxy)-methyl]-[1,2,4]thiadiazol-5-ylamino]-acetamide or β -N-[3-[Phenyl-(tetrahydropyran-2-yloxy)-methyl]-[1,2,4]thiadiazol-5-yl]-glycinamide; white solid, 80.2% yield. M.p. 147.5-148.5°C; ¹H-NMR (CDCl₃) δ (diastereoisomers) 7.50-7.52 (m, 3H, Ar-H), 7.28-7.38 (m, 2H, Ar-H), 6.82 and 6.68 (br. t, 0.5H and 0.5H each, NHCH₂), 6.08 and 5.98 (br. s, 0.5H and 0.5H each, NH), 5.92 and 5.87 (s, 0.5H and 0.5H each, α H), 5.68 (br. s, 1H, NH), 4.86 and 4.65 (t, J=3.7 Hz, 0.5H and 0.5H each, OCHO), 3.99 (t, J=6.5 Hz, 2H, NCH₂), 3.52-4.05 (m, 2H, OCH₂), 1.52-1.80 (m, 6H, 3CH₂).

C. In a similar manner as described in example 6A, the following compounds are prepared:

β -N-[3- n-butoxy-[1,2,4]-thiadiazol-5-yl]-alaninamide

β -N-[3- n-butoxy-[1,2,4]-thiadiazol-5-yl]-phenylalaninamide

β -N-[3- n-butoxy-[1,2,4]-thiadiazol-5-yl]-leucinamide

β -N-[3- n-butoxy-[1,2,4]-thiadiazol-5-yl]-isoleucinamide

β -N-[3- n-butoxy-[1,2,4]-thiadiazol-5-yl]-tyrosinamide

WO 99/45027

PCT/CA99/00160

39

Example 7

5 A. Preparation of *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline methyl ester.

 A solution of *N*-L-leucyl-L-proline methyl ester hydrochloride (5.6 g, 20 mmol, prepared according to the procedure by Jones, J. B. et al. in *J. Med. Chem.* 1995, 38, 3078) and triethylamine (5.6 mL, 40 mmol) in DMF (60 mL) was stirred at room temperature for 15 min. Then, a solution of 5-chloro-3-methoxy-[1,2,4]thiadiazole (1.51 g, 10 mmol) in DMF (10 mL) and tetra-butylammonium bromide (150 mg) were added and the resulting mixture was stirred at room temperature for 24 h. The reaction mixture was quenched with water and extracted into ethyl acetate. The organic layer was washed with a saturated solution of sodium bicarbonate, brine, dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography on silica gel using a mixture of hexane and ethyl acetate (7/3 and 6/4) afforded *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline methyl ester (1.92 g, 53.9%) as an off-white foam. ¹H-NMR (CDCl₃) δ 7.10 (d, J=8.5 Hz, 1H, NHCH), 4.74 (m, 1H, NHCH), 4.57 (dd, J=8.6, 4.6 Hz, 1H, CHCO₂), 3.92 (s, 3H, OMe), 3.85 (m, 1H), 3.69 (s, 3H, OMe), 3.62 (m, 1H), 2.17-2.23 (m, 1H), 1.95-2.07 (m, 4H), 1.77-1.82 (m, 1H), 1.59-1.66 (m, 1H), 0.97 (d, J=6.5 Hz, 3H, Me), 0.94 (d, J=6.7 Hz, 3H, Me); MS (m/z) 357 (M⁺+1), 228, 200, 130, 101, 70.

25 B. In a similar manner, the following compounds were prepared:
N-(3-Butoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline methyl ester, light yellow oil, 20.3% yield. ¹H-NMR (CDCl₃) δ 7.25 (d, J=8.8 Hz, 1H, NHCH), 4.75 (m, 1H, NHCH), 4.58 (dd, J=8.6, 4.6 Hz, 1H, CHCO₂), 4.23 (t, J=6.8 Hz, 2H, OCH₂), 3.92 (m, 1H), 3.61-3.72 (m, 1H), 3.69 (s, 3H, OMe), 2.20-2.25 (m, 1H), 1.98-2.06 (m, 3H), 1.63-1.82 (m, 5H), 1.39-1.45 (m, 2H), 0.90-0.97 (m, 9H, 3Me).

WO 99/45027

PCT/CA99/00160

40

- N*-(3-Phenyl-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline methyl ester, white solid, 85% yield. M.p. 154.0-154.5°C; ¹H-NMR (CDCl₃) δ 8.14-8.16 (m, 2H, Ar-H), 7.83 (d, J=8.7 Hz, 1H, NHCH), 7.40-7.42 (m, 3H, Ar-H), 5.02 (m, 1H, NHCH), 4.68 (dd, J=8.6, 4.6 Hz, 1H, CHCO₂), 4.09-4.14 (m, 1H), 3.77-3.82 (m, 1H), 3.76 (s, 3H, OMe), 1.72-2.29 (m, 7H), 1.09 (d, J=6.5 Hz, 3H, Me), 0.98 (d, J=6.6 Hz, 3H, Me).
- 10 *N*-(3-Methyl-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline methyl ester, white solid, 30% yield. M.p. 88-89°C; ¹H-NMR (CDCl₃) δ 7.80 (d, J=8.6 Hz, 1H, NHCH), 4.68 (m, 1H, NHCH), 4.58 (dd, J=8.6, 4.6 Hz, 1H, CHCO₂), 3.94-3.98 (m, 1H), 3.62-3.68 (m, 1H), 3.68 (s, 3H, OMe), 2.31 (s, 3H, Me), 2.07-2.22 (m, 1H), 1.96-2.07 (m, 3H), 1.69-1.79 (m, 4H), 0.96 (d, J=6.6 Hz, 3H, Me), 0.94 (d, J=6.7 Hz, 3H, Me).

C. Proceeding in a similar manner as described in example 7A, the following compounds are prepared:

- N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-octahydro-(3aS, 6aS)-cyclopentan[b]pyrrole- 2(S)-carboxylic acid methyl ester
- 20 *N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-(4-tert-butoxycarbonyl)piperazine - 2(R or S)-carboxylic acid methyl ester
- N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-decahydro-(4aR,8aS)-isoquinoline-3(S)-carboxic acid methyl ester
- 25 *N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-octahydroindole- 2-carboxylic acid methyl ester
- N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-indoline-2-carboxylic acid methyl ester
- N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-isoleucyl-L-octahydroindole- 2-carboxylic acid methyl ester
- 30 *N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-isoleucyl-L- L-octahydro-(3aS, 6aS)-cyclopentan[b]pyrrole- 2(S)-carboxylic acid methyl ester.

WO 99/45027

PCT/CA99/00160

41

Example 8

Preparation of *N*-(3-chloromethyl-[1,2,4]thiadiazol-5-yl)-*L*-leucyl-*L*-proline methyl ester.

5 A solution of *N*-*L*-leucyl-*L*-proline methyl ester hydrochloride (2.22 g, 8 mmol) and triethylamine (2.8 mL, 20 mmol) in DMF (50 mL) was stirred at room temperature for 15 min. Then, a solution of 3-chloromethyl-5-chloro-[1,2,4]thiadiazole (0.34 g, 2 mmol) in DMF (10 mL) and tetra-butylammonium
10 bromide (100 mg) were added and the resulting mixture was stirred at room temperature for 24 h. The reaction mixture was quenched with water and extracted into ethyl acetate. The organic layer was washed with a saturated ammonium chloride solution, brine, dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography on silica gel
15 using a mixture of hexane and ethyl acetate (7/3, 6/4 and 1/1) afforded *N*-(3-chloromethyl-[1,2,4]thiadiazol-5-yl)-*L*-leucyl-*L*-proline methyl ester as a white foam, 0.49 g, 65.3% yield. ¹H-NMR (CDCl₃) δ 7.56 (d, J=8.2 Hz, 1H, NHCH), 4.78 (m, 1H, NHCH), 4.60 (dd, J=8.5, 4.6 Hz, 1H, CHCO₂), 4.46 (s, 2H, CH₂Cl), 4.09 (m, 1H), 3.69-3.73 (m, 1H), 3.74 (s, 3H, OMe), 2.25-2.30 (m, 1H), 1.68-2.11 (m, 6H), 0.99-1.03 (2d, 6H, 2Me); MS (*m/z*) 375 (*M*⁺+1), 357, 305, 246, 218, 162, 150, 130.

Example 9

Preparation of *N*-{3-[1-[4-(2-pyridyl)piperazinyl]-methyl]-[1,2,4]thiadiazol-5-yl]-*L*-leucyl-*L*-proline methyl ester.

25 A mixture of *N*-(3-chloromethyl-[1,2,4]thiadiazol-5-yl)-*L*-leucyl-*L*-proline methyl ester (0.45 g, 1.2 mmol), 1-(2-pyridyl)piperazine (0.62 mL, 4 mmol), triethylamine (1.2 mL, 8 mmol) and tetra-butylammonium bromide (100 mg) in DMF (20 mL) was stirred at room temperature for 24 h. The reaction mixture
30 was diluted with water and ethyl acetate. The organic layer was collected, washed with water, dried (sodium sulfate), filtered and concentrated in vacuo

to give a light yellow oil. Purification by column chromatography on silica gel using a solvent mixture of 5% methanol in dichloromethane afforded *N*-{3-[1-
5 [4-(2-pyridyl)piperazinyl]-methyl]-[1,2,4]thiadiazol-5-yl]-L-leucyl-L-proline methyl ester as a white foam (0.55g, 91.7 %). ¹H-NMR (CDCl₃) δ 8.19 (m, 1H), 7.48-7.52 (m, 1H), 7.14 (br. m, 1H, NH), 6.64-6.67 (m, 2H), 4.67 (m, 1H, CHCO), 4.57 (dd, J= 8.4 and 4.6 Hz, 1H, CHCO₂), 3.92 (m, 1H, NCHpro), 3.62-3.78 (m, 10H, OMe, NCHpro, N=CCH₂, 2NCH₂), 2.83 (m, 4H, 2NCH₂),
10 1.62-2.30 (m, 6H, 2CH₂pro and CH₂), 0.90-1.03 (2d, J=6.5 and 6.6 Hz, 7H, 2Me and 1CH); MS (m/z) 502 (M⁺+1), 459, 408, 382, 300, 253, 225, 176, 147, 121, 95.

Example 10

15 Preparation of *N, N'*-{3-methylene-[1,2,4]thiadiazol-5-yl}-di-[L-leucyl-L-proline methyl ester].

A solution of *N*-L-leucyl-L-proline methyl ester hydrochloride (2.22 g, 8 mmol) and triethylamine (2.8 mL, 20 mmol) in DMF (50 mL) was stirred at room temperature for 15 min. Then, a solution of 3-chloromethyl-5-chloro-
20 [1,2,4]thiadiazole (0.34 g, 2 mmol) in DMF (10 mL) and tetra-butylammonium bromide (100 mg) were added and the resulting mixture was heated at 75-78°C for 19 h. The reaction mixture was quenched with water and extracted into ethyl acetate. The organic layer was washed with water, dried (sodium sulfate), filtered and concentrated in vacuo to a brown solid. Purification by
25 column chromatography on silica gel using a mixture of dichloromethane and ethyl acetate (1/1) followed by a mixture of dichloromethane and methanol (95/5) afforded *N, N'*-{3-methylene-[1,2,4]thiadiazol-5-yl}-di-[L-leucyl-L-proline methyl ester] as a light yellow semi-solid (0.43 g, 37.1%). ¹H-NMR (CDCl₃) δ 6.83 (d, J=6.8 Hz, 1H, SCNH), 4.78 (m, 1H, CHCO), 4.61 (dd, J= 8.7 and 4.8
30 Hz, 1H, CHCO₂), 4.59 (m, 1H, CHCO), 4.42 (dd, J= 8.3 and 4.1 Hz, 1H, CHCO₂), 3.73 (s, 3H, OMe), 3.71 (s, 3H, OMe), 3.68 (m, 2H, N=CCH₂), 3.38-

3.70 (m, 5H, 2NCH₂pro and NH), 1.38-2.40 (m, 12H, 4CH₂pro and 2CH₂),
0.85-1.05 (4d, J=6.5, 6.7, 6.7 and 6.4 Hz, 14H, 4Me and 2CH); MS (m/z) 581
5 (M⁺+1), 549, 452, 424, 392, 295, 267, 211, 130.

Example 11

A. Preparation of *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline.

To an ice-cooled solution of *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-
10 leucyl-L-proline methyl ester (0.68 g, 1.91 mmol) in methanol (10 mL) was
added a solution of 1N sodium hydroxide (2.4 mL, 2.4 mmol). The resulting
mixture was stirred in ice for 3 h, then at room temperature for 16 h. Volatile
materials were removed in vacuo and the residue was dissolved in water (10
mL) and washed with ethyl acetate. The aqueous layer was collected and
15 acidified with 1N HCl solution (pH ca. 5.6) as a voluminous white precipitate
separated. The mixture was extracted into ethyl acetate (6x25 mL) and the
combined organic layers was dried (sodium sulfate), filtered and concentrated
to a light yellow foam. Trituration with diethyl ether gave *N*-(3-methoxy-
[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline as a white solid (0.58 g, 89%). M.p.
20 foamed at 78°C and melted at 89.0-92.0°C; ¹H-NMR (CDCl₃) δ 7.65 (br. s,
1H, NHCH), 4.70 (br. t, 1H, NHCH), 4.49 (t, J=5.3 Hz, 1H, CHCO₂), 3.97 (s,
3H, OMe), 3.56-3.58 (m, 2H), 2.03-2.21 (m, 3H), 1.60-1.76 (m, 3H), 0.76-0.99
(m, 1H), 0.99 (d, J=6.5 Hz, 3H, Me), 0.96 (d, J=6.2 Hz, 3H, Me); MS (m/z) 343
(M⁺+1), 228, 200, 102, 83.

25

B. In a similar manner, the following compounds were prepared:

N-(3-Butoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline, white solid, 95.8% yield.
M.p. foamed at 78°C and melted at 110.0-113.0°C; ¹H-NMR (CDCl₃) δ 7.90 (br.
s, 1H, NHCH), 4.50-4.70 (m, 1H, NHCH), 4.45 (dd, J=7.1, 3.4 Hz, 1H, CHCO₂),
30 4.26 (t, J=6.7 Hz, 2H, OCH₂), 3.92 (m, 1H), 3.53-3.61 (m, 1H), 1.85-2.35 (m, 5H),
1.67-1.76 (m, 3H), 1.55-1.62 (m, 1H), 1.39-1.48 (m, 2H), 0.85-0.97 (m, 10H, 3Me
and 1CH); MS (m/z) 385 (M⁺+1), 329, 270, 242, 214, 116, 70.

WO 99/45027

PCT/CA99/00160

44

N-(3-Phenyl-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline, white solid, 91.8% yield. M.p. foamed at 112°C and melted at 136.0-138.0°C; ¹H-NMR (CDCl₃) δ 8.08-8.12 (m, 2H, Ar-H), 7.58 (br. s, 1H, NHCH), 7.38-7.42 (m, 3H, Ar-H), 4.90 (m, 1H, NHCH), 4.53 (m, 1H, CHCO₂), 4.09-4.14 (m, 1H), 3.52-3.62 (m, 1H), 2.00-2.32 (m, 4H), 1.75-1.95 (m, 2H), 1.55-1.64 (m, 1H), 1.04 (d, J=5.9 Hz, 3H, Me), 0.99 (d, J=6.0 Hz, 3H, Me), 0.85-0.92 (m, 1H); MS (m/z) 389 (M⁺+1), 333, 274, 246, 190, 116, 70.

N-(3-Methyl-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline, white solid, 80% yield. M.p. foamed at 81°C and melted at 112-115°C; ¹H-NMR (CDCl₃) δ 7.30 (br. s, NHCH), 4.42-4.60 (m, 1H, NHCH), 3.80-4.00 (m, 1H, CHCO₂), 3.54-3.65 (m, 1H), 2.37 (s, 3H, Me), 2.07-2.42 (m, 5H), 1.50-1.90 (m, 4H), 0.96-1.10 (m, 6H, 2Me), 0.80-0.95 (m, 1H); MS (m/z) 327 (M⁺+1), 271, 212, 184, 116, 70.

N-{3-[1-(4-(2-Pyridyl)piperazin-1-yl)-methyl]-[1,2,4]thiadiazol-5-yl}-L-leucyl-L-proline, light yellow solid, 70.7% yield. M.p. 163.5-166.0°C; ¹H-NMR (CDCl₃) δ 8.21 (m, 1H), 7.46-7.49 (m, 1H), 6.60-6.65 (m, 2H), 4.20-4.70 (m, 2H, CHCO and CHCO₂), 3.80 (m, 1H, NCHpro), 3.40-3.75 (m, 7H, NCHpro, N=CCH₂, 2NCH₂), 2.72 (m, 4H, 2NCH₂), 1.60-2.05 (m, 6H, 2CH₂pro and CH₂), 0.90-1.03 (2d, J=5.5 and 5.7 Hz, 7H, 2Me and 1CH); MS (m/z) 488 (M⁺+1), 442, 368, 345, 297, 253, 225, 176, 147, 121.

N, N'-{3-methylene-[1,2,4]thiadiazol-5-yl}-di-[L-leucyl-L-proline], light yellow solid (84.2%). M.p.; ¹H-NMR (CDCl₃) δ 6.83 (d, J=6.8 Hz, 1H, SCNH), 4.78 (m, 1H, CHCO), 4.61 (dd, J= 8.7 and 4.8 Hz, 1H, CHCO₂), 4.59 (m, 1H, CHCO), 4.42 (dd, J= 8.3 and 4.1 Hz, 1H, CHCO₂), 3.68 (m, 2H, N=CCH₂), 3.38-3.70 (m, 5H, 2NCH₂pro and NH), 1.38-2.40 (m, 12H, 4CH₂pro and 2CH₂), 0.85-1.05 (4d, J=6.5, 6.7, 6.7 and 6.4 Hz, 14H, 4Me and 2CH); MS (m/z) 553 (M⁺+1).

C. Proceeding in a similar manner, the following compounds are made:

- N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-octahydro-(3aS, 6aS)-
 5 cyclopentan[b]pyrrole- 2(S)-carboxylic acid
N-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-{4-tert-butoxycarbonyl}piperazine
 - 2(R or S)-carboxylic acid
N-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-decahydro-(4aR,8aS)-
 isoquinoline-3(S)-carboxic acid
 10 *N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-octahydroindole- 2-carboxylic
 acid
N-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-indoline-2-carboxylic acid
N-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-isoleucyl-L-octahydroindole- 2-
 carboxylic acid
 15 *N*-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-L-isoleucyl-L-L-octahydro-(3aS, 6aS)-
 cyclopentan[b]pyrrole- 2(S)-carboxylic acid.

Example 12

- A. Preparation of *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-iso-leucyl-L-proline
 20 methyl ester.

To a solution of *N*-L-iso-leucyl-L-proline methyl ester hydrochloride (5.58
 g, 20 mmol, prepared according to a procedure reported by Jones, J. B. et al.
 in J. Med. Chem. 1995, 38, 3078) in DMF (50 mL) was added triethylamine (5.6
 mL, 40 mmol), tetra-butylammonium bromide (0.4 g) and 5-chloro-3-methoxy-
 25 1,2,4-thiadiazole (1.51 g, 10 mmol). The resulting mixture was stirred at room
 temperature for 16 h. The reaction mixture was diluted with water and ethyl
 acetate. The organic layer was collected, dried (sodium sulfate), filtered and
 concentrated in vacuo. The residue was purified by column chromatography on
 silica gel using a solvent mixture of hexane and ethyl acetate (7/3 and 6/4)
 30 thereby affording the title compound as a light yellow oil. Trituration with hexane
 and filtration gave *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-iso-leucyl-L-proline

WO 99/45027

PCT/CA99/00160

46

5 methyl ester as a white solid (3.1 g, 87.1%). M.p. 129-130°C; ¹H-NMR (CDCl₃) δ 7.90 (d, J=9.2 Hz, 1H, NHCH), 4.61-4.65 (m, 2H, NHCH and CHCO₂), 3.96-4.00 (m, 1H), 3.92 (s, 3H, OMe), 3.69-3.72 (m, 1H), 3.72 (s, 3H, OMe), 2.21-2.24 (m, 1H), 1.82-2.05 (m, 4H), 1.62-1.66 (m, 1H), 1.18-1.24 (m, 1H), 1.03 (d, J=6.8 Hz, 3H, Me), 0.88 (t, J=7.2 Hz, 3H, Me).

10 B. In a similar fashion, *N*-[3-(1-phenyl-1-tetrahydropyran-2-yloxy)methyl-[1,2,4]thiadiazol-5-yl]-L-iso-leucyl-L-proline methyl ester was prepared as a white foam in 54.7% yield. ¹H-NMR (CDCl₃) δ 7.46-7.50 (m, 2H), 7.26-7.36 (m, 4H, Ar-H and NH), 5.79-5.85 (m, 1H, αH), 4.58-4.95 (m, 1H, OCHO), 4.38-4.51 (m, 2H), 3.81-4.02 (m, 2H), 3.74 (s, 3H, OMe), 3.60-3.70 (m, 1H), 3.45-3.55 (m, 1H), 1.45-2.10 (m, 12H), 1.04 (d, J=6.6 Hz, 3H, Me), 0.90 (t, J=7.3 Hz, 3H, Me).

15

Example 13

Preparation of *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-iso-leucyl-L-proline

In a similar manner as described in example 18A, base saponification of *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-iso-leucyl-L-proline methyl ester followed by acidification, afforded *N*-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-iso-leucyl-L-proline (93.3% yield) as a white solid. M.p. foamed at 85°C and melted at 112-114°C; ¹H-NMR (CDCl₃) δ 7.40-7.90 (br, 1H, NHCH), 4.50-4.60 (m, 2H, NHCH and CHCO₂), 3.90-4.10 (m, 1H), 3.96 (s, 3H, OMe), 3.69-3.72 (m, 1H), 1.82-2.25 (m, 4H), 1.62-1.75 (m, 1H), 1.15-1.32 (m, 1H), 1.02 (d, J=6.7 Hz, 3H, Me), 0.88 (t, J=6.6 Hz, 3H, Me); MS (m/z) 343 (M⁺+1), 287, 228, 200, 116, 70.

25

Example 14

A. Preparation of *N*-[3-(1-Phenyl-1-hydroxy)methyl-[1,2,4]thiadiazol-5-yl]-L-iso-leucyl-L-proline methyl ester.

30 A solution of *N*-[3-(1-Phenyl-1-(tetrahydropyran-2-yloxy))methyl-[1,2,4]thiadiazol-5-yl]-L-iso-leucyl-L-proline methyl ester (0.9 g, 1.74 mmol) in

WO 99/45027

PCT/CA99/00160

47

methanol (5 mL) and 3N HCl (10 mL) was stirred at room temperature for 5 min. Methanol was reduced under reduced pressure. The reaction mixture was
5 diluted with ethyl acetate and basified with 3N sodium hydroxide solution (pH ca. 10). The organic layer was collected, washed with water, dried (sodium sulfate), filtered and concentrated in vacuo. The material was purified by column chromatography (60% EtOAc : hexane) to give the titled compound (0.72g, 95.6% yield) as a white foam, 95.6% yield. ¹H-NMR (CDCl₃) δ 7.28-7.46 (m, 6H, 5Ar-H and NH), 5.75-5.85 (m, 1H, αH), 4.40-4.60 (m, 2H), 3.58-3.95 (m, 3H),
10 3.72 (s, 3H, OMe), 1.85-2.20 (m, 4H), 1.62-1.82 (m, 2H), 1.14-1.32 (m, 1H), 1.04 (d, J=6.3 Hz, 3H, Me), 0.91 (t, J=7.0 Hz, 3H, Me).

B. In a similar manner, the following compounds was prepared:
15 *N*-[3-(1-phenyl-1-hydroxy)methyl-[1,2,4]thiadiazol-5-yl]-L-iso-leucyl-L-proline as a white solid in 76% yield. M.p. foamed at 100°C and melted at 163-166°C; ¹H-NMR (CDCl₃) δ 7.79 (br, 1H, NH), 7.38-7.48 (m, 2H), 7.26-7.36 (m, 3H, Ar-H), 5.69-5.74 (m, 1H, αH), 3.50-4.50 (m, 3H), 1.80-2.20 (m, 6H), 1.60-1.780 (m, 1H), 1.10-1.30 (m, 1H), 0.91 (d, J=5.7 Hz, 3H, Me), 0.87 (t, J=7.5 Hz, 3H, Me); MS
20 (m/z) 419 (M⁺+1), 401, 349, 304, 242, 218, 163, 107, 83.

Example 15

A. Preparation of *N*-[3-Benzoyl-[1,2,4]thiadiazol-5-yl]-L-iso-leucyl-L-proline methyl ester.

25 To an ice-cooled suspension the compound from example 14A (0.65 g, 1.25 mmol) in acetone (10 mL) was added dropwise over a period of ca. 10 min a solution of chromium trioxide (0.15 g, 1.5 mmol) dissolved in water (10 mL) and conc. sulfuric acid (0.13 mL). The resulting mixture was allowed to warm to room temperature and stirred for 16 h. The reaction mixture was diluted
30 with ethyl acetate (100 mL), then made basic (pH ca. 10) by the addition of 3N sodium hydroxide solution. The organic layer was collected, washed with

WO 99/45027

PCT/CA99/00160

48

water, dried (sodium sulfate), filtered and concentrated in vacuo. Purification
by column chromatography on silica gel using a solvent gradient of a mixture
of hexane and ethyl acetate (6/4 and 4/6) afforded the title compound (0.63g
5 mg, 92.3%) as a colorless oil. ¹H-NMR (CDCl₃) δ 8.16 (d, J=7.5 Hz, 2H), 8.08
(d, J=8.4 Hz, 1H, NH), 7.59 (t, J=7.2 Hz, 1H), 7.44 (t, J=7.6 Hz, 2H), 4.62-
4.66 (m, 2H), 4.30-4.32 (m, 1H), 3.76 (s, 3H, OMe), 3.75-3.80 (m, 1H), 2.24-
2.30 (m, 1H), 1.95-2.10 (m, 5H), 1.76-1.78 (m, 1H), 1.28-1.34 (m, 1H), 1.13
10 (d, J=6.6 Hz, 3H, Me), 0.96 (t, J=7.4 Hz, 3H, Me).

B. In a similar fashion, the following compounds were prepared:
2-(3-Benzoyl-[1,2,4]thiadiazol-5-ylamino)-acetamide or β-N-{3-benzoyl-1,2,4-
thiadiazol-5-yl}-glycinamide, white solid, 46.2% yield. M.p. 227-228°C; ¹H-
15 NMR (DMSO) δ 8.84 (br. s, 1H, NH), 8.03-8.05 (m, 2H), 7.69 (t, J=6.8 Hz, 1H,
NHCH₂), 7.54-7.57 (m, 3H), 7.19 (br. s, 1H, NH), 4.02 (d, J=5.3 Hz, 2H,
NCH₃); MS m/z 263 (M⁺+1), 246, 218, 185, 158, 140, 105.

Example 16

20 Preparation of *N*-[3-benzoyl-[1,2,4]thiadiazol-5-yl]-L-iso-leucyl-L-proline.

Proceeding in a similar manner as described in example 11,
saponification of the ester from example 15A followed by acidification
afforded *N*-[3-benzoyl-[1,2,4]thiadiazol-5-yl]-L-iso-leucyl-L-proline as a white
solid in 93.8% yield. M.p. foamed at 103°C and melted at 132-134°C; ¹H-NMR
25 (CDCl₃) δ 8.15 (br. s, 1H, NH), 8.13 (d, J=7.6 Hz, 2H), 7.55 (t, J=7.0 Hz, 1H),
7.42 (t, J=7.8 Hz, 2H), 4.58-4.62 (m, 1H), 4.55 (dd, J=7.6, 4.0 Hz, 1H), 4.18-
4.22 (m, 1H), 3.58-3.63 (m, 1H), 1.92-2.2 (m, 6H), 1.65-1.75 (m, 1H), 1.21-
1.30 (m, 1H), 0.98 (d, J=6.8 Hz, 3H, Me), 0.90 (t, J=7.4 Hz, 3H, Me); MS m/z
417 (M⁺+1), 371, 302, 274, 206, 116, 70.

30

WO 99/45027

PCT/CA99/00160

49

Example 17

Preparation of {3-methoxy-[1,2,4]-thiadiazol-5-yl}carbamoyl-L-leucine-L-proline methyl ester.

To 5-amino-3-methoxy-[1,2,4]thiadiazole (1.31 g, 10 mmol) in dichloromethane (15 mL) at room temperature was added 1,1'-carbonyldiimidazole, CDI, (1.78 g, 11 mmol) followed by triethylamine (2.1 mL, 15 mmol). The resulting suspension was stirred under nitrogen for 2.5 h. Volatile materials were removed in vacuo and the residue was dissolved in DMF (20 mL). The latter was then added to a solution of L-leucine-L-proline methyl ester hydrochloride (4.18 g, 15 mmol) and triethyl amine (2.1 mL, 15.1 mmol) in DMF (25 mL). The resulting mixture was heated at 120 °C for 2 h and then allowed to cool to room temperature. Volatile materials were removed in vacuo and the residue was diluted with ethyl acetate (250 mL) and water (100 mL). The aqueous layer was extracted with ethyl acetate (2x100 mL). The combined organic layers was dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography on silica gel using a solvent mixture of hexane and ethyl acetate (1/1 and 2/3) afforded {3-methoxy-[1,2,4]-thiadiazol-5-yl}carbamoyl-L-leucine-L-proline methyl ester as a light yellow foam (3.2 g, 76.6%). M.p. 88-90°C; ¹H-NMR (CDCl₃) δ 12.75 (s, 1H, NH), 6.70 (d, J=12.0 Hz, 1H, NH), 4.79-4.87 (m, 1H, NCHpro), 4.54-4.59 (dd, J=11.8, 6.0 Hz, 1H, NCH), 4.12 (s, 3H, OMe), 3.64-3.90 (m, 2H, NCH₂pro), 1.56-2.26 (m, 6H, 3CH₂), 0.94-1.12 (two d, J=9.7 Hz, 7H, 2CH₃ + CH); ¹³C-NMR (CDCl₃) δ 178.3 (C3), 172.3, 171.1 (CO₂), 165.9 (C5), 153.7 (NHC=ONH), 58.8, 56.6, 52.2, 49.8, 47.0 (CH₂), 42.0 (CH₂), 29.0 (CH₂), 24.8 (CH₂), 24.5, 23.2, 22.0; MS (m/z) 400 (M⁺+1), 326, 271, 243, 209, 158, 130, 87.

30

WO 99/45027

PCT/CA99/00160

50

Example 18

Preparation of {3-methoxy-[1,2,4]-thiadiazol-5-yl}carbamoyl-L-leucine-L-proline.

5

In a similar manner as shown in example 18, base saponification of 3-methoxy-5-(N- L-leucine-L-proline)carbamoyl-[1,2,4]-thiadiazole methyl ester using a solution of 1N sodium hydroxide followed by acidification using a solution of 1N HCl {3-methoxy-[1,2,4]-thiadiazol-5-yl}carbamoyl-L-leucine-L-proline as a white solid (55%). M.p. 209-211°C; ¹H-NMR (MeOD) δ 4.64-4.67 (dd, J=8.1, 2.3 Hz, 1H, NCH), 4.44-4.45 (m, 1H, NCHpro), 3.94 (s, 3H, OMe), 3.60-3.86 (m, 2H, NCH₂pro), 1.54-2.27 (m, 6H, 3CH₂), 0.94-1.01 (two d, J=6.4 Hz, 7H, 2CH₃ + CH); MS (m/z) 386 (M⁺+1), 271, 229, 158, 116, 86.

10

Example 19

Preparation of 3-methoxy-5-(N-carbobenzyloxy-L-phenylalanyl-L-alaninamido)-[1,2,4]thiadiazole

To a solution of 5-amino-3-methoxy-[1,2,4]thiadiazole (0.37 g, 2.8 mmol) and N-carbobenzyloxy-L-phenylalanyl-L-alanine, N-Cbz-Phe-Ala-OH, (1.0 g, 2.7 mmol) in DMF (25 mL) was added 1,3-dicyclohexylcarbodiimide, DCC, (0.56 g, 2.7 mmol). After stirring for 30 min, 1-hydroxybenzotriazole hydrate, HOBT, (0.36 g, 2.7 mmol) was added and the resulting mixture was stirred at room temperature for 20 h. Volatile materials were removed in vacuo and the residue was purified by column chromatography on silica gel using a solvent mixture of dichloromethane and methanol (96/4) thereby affording the title compound as a white solid (1.2 g). Further purification by cristallization and chromatography on silica gel gave 3-methoxy-5-(N-carbobenzyloxy-L-phenylalanyl-L-alaninamido)-[1,2,4]thiadiazole (0.87 g, 67% yield). M.p. 161-162°C; ¹H-NMR (CDCl₃) δ 7.14-7.38 (m, 12H, 10Ar-H and 2 NH), 5.92 (br. s, 1H, NH), 5.06 (s, 2H, OCH₂), 4.38 (m, 1H, CHCH₂Ar), 4.00 (m, 1H, CHCH₂), 3.97 (s, 3H, OMe), 3.00-3.08 (m, 2H, CHCH₂Ar), 1.34 (d, J=8.9 Hz, 3H, Me); MS (m/z) 484 (M⁺+1), 416, 361, 316, 285, 185, 132,

20

25

30

75. The latter compound was also prepared by using diphenylphosphoryl azide and triethylamine instead of DCC/HOBT in 27.7% yield.

5

Example 20

Preparation of 3-methoxy-5-(*N*-carbobenzyloxy-L-phenylalaninamido)-[1,2,4]thiadiazole.

To an ice-cooled solution of 5-amino-3-methoxy-[1,2,4]thiadiazole (1.31 g, 10 mmol) and *N*-carbobenzyloxy-L-phenylalanine, (Cbz-L-phe-OH 2.99 g, 10 mmol) in DMF (100 mL) was added 1,3-dicyclohexylcarbodiimide (2.06 g, 10 mmol) followed by 1-hydroxybenzotriazole hydrate (1.35 g, 10 mmol). The resulting mixture was stirred at room temperature for 16 h. The mixture was diluted with ethyl acetate (250 mL) and water (200 mL). The organic layer was collected and successively washed with a solution of 1N HCl (15 mL), water, a saturated solution of sodium bicarbonate, and water. The organic layer was dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography using a solvent mixture of hexane and ethyl acetate (3/1) afforded 3-methoxy-5-(*N*-carbobenzyloxy-L-phenylalaninamido)-[1,2,4]thiadiazole as a colorless oil. Crystallization from ether and hexane gave the product as a white solid (1.1 g, 26.7%). ¹H-NMR (CDCl₃) δ 12.50 (s, 1H, NH), 7.06-7.35 (m, 10H, Ar-H), 6.02 (br. d, J=7.6 Hz, 1H, NH), 5.19-5.22 (m, 1H, CH), 5.11 (s, 2H, OCH₂), 4.05 (s, 3H, OMe), 3.11-3.20 (m, 2H, CHCH₂Ar); MS (m/z) 413 (M⁺+1), 369, 222, 210, 132, 91.

25

Example 21

Preparation of 3,5-di-(*N*-carbobenzyloxy-L-phenylalaninamido)-[1,2,4]thiadiazole.

To an ice-cooled solution of 3,5-diamino-[1,2,4]thiadiazole (1.16 g, 10 mmol, prepared according to the procedure reported by Kurzer, F. In J. Chem. Soc. 1955, 1) and *N*-carbobenzyloxy-L-phenylalanine, (Cbz-L-phe-OH, 4.98 g, 20 mmol) in DMF (100 mL) was added 1,3-

30

WO 99/45027

PCT/CA99/00160

52

dicyclohexylcarbodiimide (4.12 g, 20 mmol) followed by 1-hydroxybenzotriazole hydrate (2.70 g, 20 mmol). The resulting mixture was stirred at room temperature for 16 h. The mixture was diluted with ethyl acetate and water. The organic layer was collected and successively washed with a solution of 1N HCl, brine, a saturated solution of sodium bicarbonate and brine. The organic layer was dried (sodium sulfate), filtered and concentrated in vacuo to give a light yellow solid. Purification by column chromatography using a solvent mixture of hexane and ethyl acetate (7/3 and 6/4) afforded 3,5-di-(*N*-carbobenzyloxy-L-phenylalanylamido)-[1,2,4]thiadiazole as a white solid. Recrystallization from ethyl acetate and hexane gave the product as a white solid (3.5 g, 51.5%). M.p. 192-194°C; ¹H-NMR (CDCl₃) δ 13.3 (s, 1H, NH), 11.1 (s, 1H, NH), 7.95 (d, J=7.5 Hz, 1H, NHCH), 7.68 (d, J=8.4 Hz, 1H, NHCH), 7.19-7.39 (m, 20H, Ar-H), 4.98 (s, 4H, 2OCH₂), 4.58-4.64 (m, 2H, 2CHCH₂), 3.03-3.12 (m, 2H, PhCH₂CH), 2.77-2.92 (m, 2H, PhCH₂CH); MS (m/z) 679.5 (M⁺+1), 635, 591, 488, 445, 398, 354, 297, 210, 117.

Example 22

Preparation of 5-bromoacetamido-3-methoxy-[1,2,4]thiadiazole.

To an ice-cooled solution of 5-amino-3-methoxy-[1,2,4]thiadiazole (1.31 g, 10 mmol) in THF (25 mL) was added triethylamine (2.22 g, 11 mmol) followed by dropwise addition of bromoacetyl bromide (1.52 g, 15 mmol). The resulting mixture was stirred at room temperature for 16 h. The mixture was diluted with water (25 mL) and ethyl acetate (50 mL). The aqueous layer was extracted with ethyl acetate (3x50 mL). The combined organic layers was dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography on silica gel using a solvent gradient of a mixture of dichloromethane and methanol (98/2, 97/3 and 96/4) afforded 3-methoxy-5-bromoacetamido-[1,2,4]thiadiazole as a white solid (1.26 g, 50%). M.p. 198-199°C; ¹H-NMR (CDCl₃) δ 13.10 (br. s, 1H, NH), 4.28 (s, 2H, CH₂), 4.11 (s,

WO 99/45027

PCT/CA99/00160

53

3H, OMe); ^{13}C -NMR (CDCl_3) δ 176.8 (C3), 166.9 (C5), 166.6 (C=O), 56.9 (OMe), 25.9 (CH_2).

5

B. Similarly, when chloroacetyl chloride was used, 5-chloroacetamido-3-methoxy-[1,2,4]thiadiazole was obtained as a white solid (52%). M.p. 207-208°C; ^1H -NMR (CDCl_3) δ 12.04 (br. s, 1H, NH), 4.45 (s, 2H, CH_2), 4.11 (s, 3H, OMe).

10

Example 23

Preparation of N-[(3-methoxy-1,2,4-thiadiazol-5-yl)carbamoyl]methyl]-L-leucyl-L-proline methyl ester.

To a solution of 5-bromoacetamido-3-methoxy-[1,2,4]thiadiazole (0.76 g, 3 mmol) and L-leucine-L-proline methyl ester (0.97 g, 4 mmol) was added solid sodium iodide (45 mg, 0.3 mmol), tetrabutylammonium bromide (97 mg, 0.3 mmol) and triethylamine (0.7 mL, 5 mmol). The resulting mixture was stirred for 3 h, then quenched with brine and extracted into ethyl acetate. The organic layer was washed with a saturated solution of sodium bicarbonate, brine, dried (sodium sulfate), filtered and concentrated in vacuo. Purification by column chromatography on silica gel using a solvent mixture of hexane and ethyl acetate (1/1) followed by a mixture of dichloromethane and methanol (96/4) gave N-[(3-methoxy-1,2,4-thiadiazol-5-yl)carbamoyl]methyl]-L-leucyl-L-proline methyl ester as an off-white solid (1.09 g, 88%). M.p. 154-155°C; MS (m/z) 414 ($\text{M}^+ + 1$), 382, 340, 257, 199, 130, 75.

Example 24

Preparation of N-[(3-methoxy-1,2,4-thiadiazol-5-yl)carbamoyl]methyl]-L-leucyl-L-proline.

30

Base saponification N-[(3-methoxy-1,2,4-thiadiazol-5-yl)carbamoyl]methyl]-L-leucyl-L-proline methyl ester with a 1N sodium

hydroxide solution followed by acidification with a 1N HCl solution similar to example 3h (B) afforded N-(((3-methoxy-1,2,4-thiadiazol-5-yl)carbamoyl)methyl)-L-leucyl-L-proline methyl ester (80% yield). M.p. 195-197°C with decomposition; ¹H-NMR (MeOD) δ 4.63 (d, J=16.9 Hz, 1H, COCH₂), 4.41 (br. dd, 1H, CHCO₂H), 4.17 (d, J=17.1 Hz, 1H, COCH₂), 4.05 (m, 1H, CHCH₂), 4.00 (s, 3H, OMe), 3.48-3.70 (m, 2H, CH₂pro), 1.72-2.40 (m, 6H, 3CH₂), 0.92-1.03 (m, 7H, 2CH₃ + CH); MS (m/z) 400 (M⁺+1).

10

Example 25

A. Preparation of acid addition salt of a compound of formula I.

To an ice-cooled suspension of β-N-{3-(α-hydroxybenzyl)-1,2,4-thiadiazol-5-yl}-glycinamide (60 mg, 0.22 mmol) in methanol (10 mL) was bubbled HCl gas for ca. 2 min as a light yellow solution resulted. The volume of the reaction mixture was reduced to ca. 2 mL by rotary evaporation and diethyl ether was added. The voluminous yellow precipitate of the hydrochloride salt was collected by filtration and dried at 50°C under vacuum for 3 h (67 mg, 99.1%). 2-[3-(1-Hydroxy-1-phenylmethyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide hydrochloride salt or β-N-{3-(α-hydroxybenzyl)-1,2,4-thiadiazol-5-yl}-glycinamide hydrochloride salt, white solid. ¹H-NMR (MeOD) δ 7.47-7.51 (m, 2H), 7.30-7.40 (m, 3H), 5.81 (s, 1H, αH), 4.19 (s, 2H, NCH₂).

20

B. In a similar manner, the following compounds were made:

25 2-[3-(3-Phenylallyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide hydrochloride salt or β-N-{3-cinnamyl-1,2,4-thiadiazol-5-yl}glycinamide hydrochloride salt, white solid, 99% yield, m.p. 209.0-210.5°C with decomposition.

2-[3-(1-Hydroxy-1-phenylmethyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide hydrochloride salt or β-N-{3-(α-hydroxybenzyl)-1,2,4-thiadiazol-5-yl}-glycinamide hydrochloride salt, white solid, 99.1% yield. ¹H-NMR (MeOD) δ 7.47-7.51 (m, 2H), 7.30-7.40 (m, 3H), 5.81 (s, 1H, αH), 4.19 (s, 2H, NCH₂).

30

WO 99/45027

PCT/CA99/00160

55

2-(3-n-Butoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide hydrochloride salt or β -N-(3-n-butoxy-1,2,4-thiadiazol-5-yl)-glycinamide hydrochloride salt, white solid, 86.5% yield, m.p. foamed at 124°C and melted at 178-190°C with decomposition.

2-[3-(1H-Indol-3-ylmethyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide hydrochloride salt, white solid, 97.8% yield, m.p. foamed at 125°C and melted at 209.0-210.5°C with decomposition.

10

Example 26

Inhibition of human rhinovirus protease (3C P) by compounds of formula I.

Human Rhinovirus protease (3Cp) was dissolved in 50 mM of potassium Phosphate pH 7.5, 0.25 mM EDTA, 10% glycerol at 0.13 mg/mL. The total assay volume was 500 micro liters and the assays were run at room temperature (RT) which was 23°C. Assay buffer contained 50 mM TRIS/HCl, pH 7.0, 1 mM EDTA, 100 mM NaCl, 0.005mM DTT. Human Rhinovirus 3Cp substrate (3CpS: Bachem M-2075, Abz-Glu-Thr-Leu-Phe-Gln-Gly-Pro-Val-p-nitro-Phe-NH₂) was dissolved in 80 mM NaHCO₃, 15% DMSO. All inhibitors were dissolved in DMSO to 20mM. All inhibitors were dissolved immediately prior to their assays. The enzyme 3Cp was added to the assay buffer to 0.007 mg/mL or 0.3 microM. The mixture was incubated at RT for 1 0-15 minutes. The reaction was initiated by the addition of 3Cp to 0.03 mM. When the inhibitors were assayed, they were added immediately after the addition of 3Cp. Fluorescence readings were recorded at 1 second intervals for 600 sec. The data were downloaded to a disk, uploaded as text file, and converted to dF vs time(s) values. The data were transferred to the program MacCurveFit and, fit to the linear equation $y = mx + b$, where y - dF, x = time and m = rate(dF/s). $R^2 > 0.97$. V_o = steady-state rate (dF/s) without inhibitors. V_i = steady-state rate (dF/s) with inhibitor. % Inhibition = $1 - V_i/V_o$.

30

N-(3-methoxy-1,2,4-thiadiazol-5-yl)-glycinamide; IC₅₀ = 62 μ M.

WO 99/45027

PCT/CA99/00160

56

Example 27

Inhibition of cathepsin B by 1,2,4-thiadiazoles. Enzyme assays and kinetic measurements.

Conditions for the above experiments can be found in the following references: Menard R. et al., 30 Biochemistry 1990, 29, 6706-6713; Fox T. et al., Biochemistry 1992, 31, 12571-12576; Cannona E. et al. Biochemistry 1996, 35, 8149-8157. A typical experiment consists in choosing an inhibitor concentration such that maximum inhibition could be achieved in less than two hours, monitoring the complete progress curve (i.e. fluorescence vs time), and analyzing the data. The analysis yield two parameters: the % inhibition once steady state was reached, and a rate constant which represents the rate at which this steady state is reached. Typically, the enzyme activity decreases with time until the maximum level of inhibition is reached (i.e. steady state) where the enzyme activity remains constant. Since significant levels of activity could still be detected at steady state (i.e. inhibition is not complete), the data was fitted to equation (1), which is normally used for slow-binding reversible inhibitors.

$$[P] = v_o \cdot t + \{(v_i - v_o) [1 - e^{-k_{obs} t}]\} / k_{obs} \quad (1)$$


$$\% \text{ inhibition} = (1 - v_i / v_o) \cdot 100 \quad (2)$$

In this equation, [P] represents the concentration of product (obtained from the fluorescence readings), k_{obs} is the first order rate constant to reach steady state, v_o is the initial rate which corresponds to the rate in the absence of the inhibitor, and v_i is the rate of the inhibited enzyme at steady state. The % inhibition was obtained by using equation (2), where the rate measured in the absence of the inhibitor was used for v_o . An example of a simple mechanism for such a process is given below:

WO 99/45027

PCT/CA99/00160

57



For this mechanism, the value of k_{obs} determined experimentally would correspond to:

10

$$k_{\text{obs}} = k_{\text{on}} [\text{inh}] + k_{\text{off}}$$

The value of k_{off} varies with inhibitor concentration. Often k_{off} is much slower than $k_{\text{on}}[\text{inh}]$ and $k_{\text{obs}} / [\text{inh}]$ can be used to approximate k_{on} . This approach is used in this study and the results are shown in below:

15 3-methoxy-5-(carbobenzyloxy-L-phenylalanyl-L-alaninamido)-1,2,4-thiadiazole, 65% inhibition at 400 mM, $k_{\text{obs}} / [\text{Inh}] = 38 \text{ M}^{-1} \text{ s}^{-1}$.
 N-[3-methoxy-{1,2,4}-thiadiazol-5-yl]-(L-leucyl-L-proline), 85% inhibition at 60 mM, $k_{\text{obs}} / [\text{Inh}] = 170 \text{ M}^{-1} \text{ s}^{-1}$.

20 **Example 28**

Inhibition of transglutaminase by compounds of formula I (Enzyme Assay).

Transglutaminase activity is measured by the colorimetric hydroxamate procedure (J. Biol. Chem., 1971, 246, 1093). The tested compounds were dissolved in methanol and added to a buffered solution of purified
 25 Transglutaminase to give a final concentration of 0.1 mM (10% MeOH). After on hour of incubation, the solution was assayed for residual activity using the standard activity assay. 3-Methoxy-5-(carbobenzyloxy-L-phenylalanyl-L-alaninamido)-1,2,4-thiadiazole completely inhibits the enzyme at 100 μM .

30

Example 29**Preparation of Pharmaceutically Acceptable Salt.**

5 A solution of tris(hydroxymethyl)methylamine (61 mg, 0.5 mmol) in 2 mL water was added dropwise to a solution of *N*-{(3-methoxy)-[1,2,4]thiadiazol-5-yl}-L-leucyl-L-proline (171 mg, 0.5 mmol) in 10 mL methanol. The resulting clear solution was stirred for 1 h at room temperature. Volatile materials were removed in vacuo and the residual white foam was
10 dried under vacuum at room temperature for 16 h (225 mg, yield 97.1%). M.p. foamed at 49 °C and melted at 127-128 °C; ¹H-NMR (MeOD) δ 4.80 (m, 1H, CH/N), 4.30 (m, 1H, CHCO₂), 3.93 (s, 2H), 3.91 (s, 2H), 3.90 (s, 2H), 3.62 (s, 3H, OMe), 3.50-3.60 (m, 2H), 2.00-2.20 (m, 3H), 1.60-1.85 (m, 3H), 0.94-1.05 (m, 7H, 2Me and 2CH).

15

Example 30

The effect of *N*-{(3-methoxy)-[1,2,4]thiadiazol-5-yl}-L-leucyl-L-proline. tris(hydroxymethyl)methylamine salt (Apo501) on IL-1-Induced Proteoglycan Degradation.

20 The effect of Apo501 on IL-1 (interleukin) induced proteoglycan degradation of articular cartilage from normal calf joints was investigated *in-vitro*. Articular cartilage was sliced aseptically from the distal forelimb joints of less than 6-month old calves. The joint showed no clinical sign of skeletal disease. The cartilage was cut out with a sterile biopsy punch to the diameter
25 of 3.5mm with comparable thickness. The pooled, punched cartilage was washed and incubated in 20 ml of Ham's F12 medium with 5% FBS and 3% antibiotics for 72 hr at 37°C for recovery.

Revoered explants were washed and replaced with a fresh medium. 20 μl (10 μCi/ml) of sodium [³⁵S] sulfate was added and cartilage explants
30 incubated for 72 h at 37°C to radiolabel proteoglycans. The explants was washed thoroughly (5 times) to remove free radioisotope and incubated in serum free, F-12 medium (3% Ab) overnight. Explants were then incubated in

WO 99/45027

PCT/CA99/00160


59

quadruplicate in fresh, serum free mediums containing varied concentrations of Apo-501 (10^{-7} , 10^{-6} , 10^{-5} , and 10^{-4}) for 24 h. On the next day, each cartilage was washed and transferred to a well in a 24-well plate, to which 1 ml of freshly prepared medium (serum-free, 3% Ab) that contains corresponding Apo-501 concentration and 50 ng of IL- β was added. The explants were incubated for 72 h at 37°C. The extent of proteoglycan degradation in each cartilage treated with different concentrations of Apo501 was determined by measuring the radioactivity of [^{35} S] sulfate-glycosaminoglycans by a liquid scintillation counter. Sample of 100 μ l medium was added to 5 ml of scintillation cocktail (Ready Safe, Beckmann) and counted. Proteoglycan degradation was expressed as radiolabelled glycosaminoglycans released into the media (counts per min per ml) per dry weight (mg) of cartilage.

As shown in Fig. 2, Apo501 demonstrated an inhibitory effect on the IL-1 induced proteoglycan degradation. Apo501 resulted in a significant reduction (60%) of IL-1 (50 ng/ml)-induced degradation of newly synthesized proteoglycan ($p = 0.023$) at 1×10^{-6} M.

CLAIMS


1. 3-5 disubstituted 1,2,4-thiadiazole compounds, having the
 5 general formula (II):




10

- or their pharmaceutically acceptable salts thereof,
 wherein A is an amino acid residue, or a peptide containing 2 to 3
 amino acid residues with the N terminal of A directly attached to the C5
 of the 1,2,4-thiadiazole ring or an isosteric form thereof and W
 15 represents a group of formula $-N(R^1)_2$ or $-OR^1$ with R^1 being
 independently hydrogen, lower alkyl, lower alkenyl, lower alkynyl in
 which the unsaturated bond is at least one carbon removed from the N
 or O atom, with the proviso that -A-W is not a group of formulas:

20



25



30

wherein R^e represents hydrogen or lower alkyl having a straight or
 branched chain of 1 to 4 carbon atoms;

R^f represents lower alkyl having a straight or branched chain
 of 1 to 4 carbon atoms;


35

R^i represents lower alkyl having a straight or branched chain of 1 to 4 carbon atoms;

R_j represents lower alkyl having a straight or branched chain of 1 to 4 carbon atoms, or benzyl which is unsubstituted or monosubstituted to trisubstituted by identical or different substituents selected from the group fluorine, chlorine, bromine, nitro, cyano, lower alkyl, methylthio, lower alkoxy, trifluoromethyl, trifluoromethoxy, difluoromethoxy, trifluoromethylthio, methoxycarbonyl, ethoxycarbonyl, methoximino-methyl, methoxyiminoethyl, ethoximinomethyl cyclopentyl, cyclohexyl, 1,3-propanediyl, but-2-ene-di-1,4-yl, 1,4-butanediyl, and phenyl, benzyl, phenoxy or benzyloxy in each case unsubstituted or monosubstituted to trisubstituted in the phenyl moiety by identical or different substituents selected from the group consisting of fluorine, chlorine, bromine, lower alkoxy, lower alkyl, methylthio, trifluoromethyl, trifluoromethoxy, difluoromethoxy, and trifluoromethylthio;


and Y is selected from:

- (1) lower alkoxy, lower cycloalkoxy, lower arylalkoxy, heterocycloxy, and lower heterocyclalkoxy wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;
- (2) lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocycl, aryl, lower arylalkyl, lower arylalkenyl, lower heterocyclalkenyl wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, and dialkylamino;
- (3) lower alkoxycarbonyl, carboxyl;
- (4) a ketone group of formula:



in which R² represents lower alkyl, lower cycloalkyl, lower heterocyclylalkyl, heterocyclyl, aryl, lower arylalkyl wherein the alkyl or aromatic ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;


- (5) a carbamoyl group of formula:



with R² being as defined above;


- (6) amino, lower alkylamino, lower dialkylamino;

- (7) amide of formula:



with R² being as defined above;


- (8) a group of formula:



wherein A is as defined above and the carboxyl terminal of A is directly attached to the nitrogen of the 3-amino-1,2,4-thiadiazole, R represents hydrogen, lower alkanoyl, lower cycloalkylcarbonyl, lower alkoxy carbonyl, lower arylalkyloxycarbonyl or N protecting group and R¹ is hydrogen, lower alkyl, lower alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N atom;


63

- (9) alcohol of formula:




with R^2 being as defined above;

- (10) sulfone of formula:




with R^2 being as defined above;

- (11) sulfoxide of formula:



with R^2 being as defined above;


- (12) sulfonamide of formula:



with R^2 being as defined above;


- (13) lower alkylthio, lower arylalkylthio, arylthio;

- (14) a group of formula:



with A as defined above and the N-terminal of A is directly attached to the methylene and W being as defined above;

- (15) a group of formula:




in which R^3 and R^4 are independently alkyl, aralkyl, heterocyclyl, heterocyclalkyl; R^3 and R^4 when taken together form with the N-atom a five or a six membered ring selected from the group piperidinyl, pyrrolidinyl, piperazinyl with the N-4 position of piperazine optionally substituted with pyridyl, heterocyclyl, alkyl, aralkyl and aryl.

5

2. 3-5 disubstituted 1,2,4-thiadiazole compounds, having the general formula (II):

10



15

or their pharmaceutically acceptable salts thereof,

20

wherein A-W is glycylamide, leucyl-proline, isoleucyl-proline, leucyl-octahydro-1*H*-indole-2-carboxylic acid, isoleucyl-octahydro-1*H*-indole-2-carboxylic acid, leucyl-octahydrocyclopenta[*b*]pyrrole-2-carboxylic acid, isoleucyl-octahydrocyclopenta[*b*]pyrrole-2-carboxylic acid or A is an amino acid residue A_1 or a dipeptide A_1-A_2 with the N-terminal of A directly attached to the C5 of the 1,2,4-thiadiazole ring or an isomeric form thereof and W is -OH, -NH₂, lower alkoxy, lower alkylamino, the amino acid A_1 is selected from the group alanyl, β -alanyl, arginyl, aspartyl, glutamyl, glutaminyl, histidyl, isoleucyl, leucyl, lysyl, methionyl, norleucyl, norvalyl, ornithyl, phenylalanyl, prolyl, seryl, tryptophyl, tyrosyl, valyl, 4-aminobutanoyl, 2,6-diaminohexanoyl, 2,3-diaminopropanoyl, 2,4-diaminobutanoyl and 2,5-diaminopentanoyl, when A_1 is 2,6-diaminohexanoyl, 2,3-diaminopropanoyl, 2,4-diaminobutanoyl and 2,5-diaminopentanoyl, the N-attachment point is either at the α -NH₂ or the amino acid side-chain NH₂ group, A_2 is selected from the group alanyl, β -alanyl, arginyl, aspartyl, glycyl, glutamyl, glutaminyl, histidyl, isoleucyl, leucyl, lysyl, methionyl, norleucyl, norvalyl, ornithyl, phenylalanyl, prolyl, seryl, tryptophyl,

35

tyrosyl, valyl, 2-indolinecarbonyl, octahydro-1*H*-indole-2-carbonyl, octahydrocyclopenta[*b*]pyrrole-2-carbonyl, octahydro-1*H*-isoindole-1-carbonyl, octahydrocyclopenta[*c*]pyrrole-1-carbonyl, 1,2,3,4-tetrahydro-3-isoquinolinecarbonyl, decahydro-3-isoquinolinecarbonyl;

5

and Y is selected from:

(1) lower alkoxy, lower cycloalkoxy, lower arylalkoxy, heterocyclyloxy, and lower heterocyclylalkoxy wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;

10


(2) lower alkyl, lower cycloalkyl, lower heterocyclylalkyl, heterocyclyl, aryl, lower arylalkyl, lower arylalkenyl, lower heterocyclylalkenyl wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, and dialkylamino;

15

(3) lower alkoxycarbonyl, carboxyl;

20

(4) a ketone group of formula:




25

in which R² represents lower alkyl, lower cycloalkyl, lower heterocyclylalkyl, heterocyclyl, aryl, lower arylalkyl wherein the alkyl or aromatic ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;

30

(5) a carbamoyl group of formula:



35

with R² being as defined above;

(6) amino, lower alkylamino, lower dialkylamino;


(7) amide of formula:



with R^2 being as defined above;

10

(8) a group of formula:



15

wherein A is as defined above and the carboxyl terminal of A is directly attached to the nitrogen of the 3-amino-1,2,4-thiadiazole, R represents hydrogen, lower alkanoyl, lower cycloalkylcarbonyl, lower alkoxy carbonyl, lower arylalkyloxycarbonyl or N protecting group and R^1 is hydrogen, lower alkyl, lower alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N atom;

20


(9) alcohol of formula:



with R^2 being as defined above;

30


(10) sulfone of formula:



35


with R^2 being as defined above;

(11) sulfoxide of formula:



with R² being as defined above;


(12) sulfonamide of formula:



with R² being as defined above;


(13) lower alkylthio, lower arylalkylthio, arylthio;

(14) a group of formula:




with A as defined above and the N-terminal of A is directly attached to the methylene and W being as defined above;

(15) a group of formula:




in which R³ and R⁴ are independently alkyl, aralkyl, heterocyclyl, heterocyclalkyl; R³ and R⁴ when taken together form with the N-atom a five or a six membered ring selected from the group piperidinyl, pyrrolidinyl, piperazinyl with the N-4 position of piperazine optionally substituted with pyridyl, heterocyclyl, alkyl, aralkyl and aryl.

3. 3-5 disubstituted 1,2,4-thiadiazole compounds, having the general formula (III):



68

or their pharmaceutically acceptable salts thereof,
 wherein A is an amino acid residue, or a peptide containing 2 to 3
 amino acid residues with the N-terminal of A attached by the means
 of a spacer X to the C5 of the 1,2,4-thiadiazole ring, or an isosteric
 form thereof, X is a spacer selected from the group of formula



—N(R¹)₂ or —OR¹ with R¹ being independently hydrogen, lower alkyl,
 lower alkenyl, lower alkynyl in which the unsaturated bond is at least
 one carbon removed from the N or O atom;


and Y is selected from:

(1) lower alkoxy, lower cycloalkoxy, lower arylalkoxy,
 heterocycloxy, and lower heterocyclalkoxy wherein
 the alkyl or aryl ring is optionally substituted with 1 to 2
 substituents selected from the group amino, alkoxy,
 hydroxy, halo, amino, alkylamino, dialkylamino;

(2) lower alkyl, lower cycloalkyl, lower heterocyclalkyl,
 heterocycl, aryl, lower arylalkyl, lower arylalkenyl,
 lower heterocyclalkenyl wherein the alkyl or aryl ring
 is optionally substituted with 1 to 2 substituents
 selected from the group amino, alkoxy, hydroxy, halo,
 amino, alkylamino, and dialkylamino;

(3) lower alkoxycarbonyl, carboxyl;


(4) a ketone group of formula:



69

in which R^2 represents lower alkyl, lower cycloalkyl,
lower heterocyclylalkyl, heterocyclyl, aryl, lower arylalkyl
wherein the alkyl or aromatic ring is optionally
substituted with 1 to 2 substituents selected from the
group amino, alkoxy, hydroxy, halo, amino, alkylamino,
dialkylamino;


- (5) a carbamoyl group of formula:



with R^2 being as defined above;


- (6) amino, lower alkylamino, lower dialkylamino;

- (7) amide of formula:




with R^2 being as defined above;

- (8) a group of formula:



wherein A is as defined above and the carboxyl
terminal of A is directly attached to the nitrogen of the
3-amino-1,2,4-thiadiazole, R represents hydrogen,
lower alkanoyl, lower cycloalkylcarbonyl, lower
alkoxycarbonyl, lower arylalkoxycarbonyl or N
protecting group and R^1 being as defined as above;


- (9) alcohol of formula:



with R² being as defined above;

(10) sulfone of formula:


5



with R² being as defined above;

10


(11) sulfoxide of formula:



with R² being as defined above;

15

(12) sulfonamide of formula:




20

with R² being as defined above;

(13) lower alkylthio, lower arylalkylthio, arylthio;

25


(14) a group of formula:



with A as defined above and the N-terminal of A is directly attached to the methylene and W being as defined above;

30

(15) a group of formula:




in which R³ and R⁴ are independently alkyl, aralkyl, heterocyclyl, heterocyclalkyl; R³ and R⁴ when taken together form with the N-atom a five or a six membered ring selected from the group piperidinyl, pyrrolidinyl, piperazinyl with the N-4 position of piperazine optionally

35

substituted with pyridyl, heterocyclyl, alkyl, aralkyl and aryl.

- 5 4. 3-5 disubstituted 1,2,4-thiadiazole compounds, having the general formula (IV):



or their pharmaceutically acceptable salts thereof,

- 15 wherein A is an amino acid residue, or a peptide containing 2 to 3 amino acid residues, the carboxyl terminal of A is directly attached to the nitrogen of the 5-amino 1,2,4-thiadiazole and R represents hydrogen, lower alkanoyl, lower cycloalkylcarbonyl, lower alkoxy carbonyl, lower arylalkyloxycarbonyl or N protecting group and
- 20 R¹ is hydrogen, lower alkyl, lower alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N atom;


and Y is selected from:

- 25 (1) lower alkoxy, lower cycloalkoxy, lower arylalkoxy, heterocycloxy, and lower heterocyclalkoxy wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;
- 30 (2) lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocyclyl, aryl, lower arylalkyl, lower arylalkenyl, lower heterocyclalkenyl wherein the alkyl or aryl ring is optionally substituted with 1 to 2 substituents
- 35 selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, and dialkylamino;

(3) lower alkoxy carbonyl, carboxyl;

(4) a ketone group of formula:

5




in which R² represents lower alkyl, lower cycloalkyl, lower heterocyclalkyl, heterocycl, aryl, lower arylalkyl wherein the alkyl or aromatic ring is optionally substituted with 1 to 2 substituents selected from the group amino, alkoxy, hydroxy, halo, amino, alkylamino, dialkylamino;

10

(5) a carbamoyl group of formula:

15




with R² being as defined above;

20

(6) amino, lower alkylamino, lower dialkylamino;

(7) amide of formula:


25



with R² being as defined above;

(8) a group of formula:

30




wherein A is as defined above and the carboxyl terminal of A is directly attached to the nitrogen of the 3-amino-1, 2, 4-thiadiazole, R represents hydrogen, lower alkanoyl, lower cycloalkylcarbonyl, lower alkoxy carbonyl, lower arylalkyloxycarbonyl or N protecting group and R¹ is hydrogen, lower alkyl, lower

35


alkenyl, lower alkynyl in which the unsaturated bond is at least one carbon removed from the N atom;

- 5 (9) alcohol of formula:



- 10 with R^2 being as defined above;


- (10) sulfone of formula:



15

with R^2 being as defined above;


- (11) sulfoxide of formula.:



20

with R^2 being as defined above;

- 25 (12) sulfonamide of formula:




30

with R^2 being as defined above;

- (13) lower alkylthio, lower arylalkylthio, arylthio;


- (14) a group of formula:

35



with A as defined above and the N-terminal of A is directly attached to the methylene and W being as defined above;

(15) a group of formula:



in which R^3 and R^4 are independently alkyl, aralkyl, heterocyclyl, heterocyclalkyl; R^3 and R^4 when taken together form with the N-atom a five or a six membered ring selected from the group piperidiny, pyrrolidiny, piperaziny with the N-4 position of piperazine optionally substituted with pyridyl, heterocyclyl, alkyl, aralkyl and aryl.

5. 3-5 disubstituted 1,2,4-thiadiazole compounds according to claim 1 or 2 wherein A is glycol, W is $-NH_2$ and Y is as defined in claim 1 or 2.

6. A compound according to claim 5 wherein Y is methoxy, which is N-(3-methoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide.

7. A compound according to claim 5 wherein Y is n-butoxyl N-(3-n-Butoxy-[1,2,4]thiadiazol-5-ylamino)-acetamide.

8. A compound according to claim 5 wherein Y is cinnamyl N-(3-Phenylallyl)-[1,2,4]thiadiazol-5-ylamino]-acetamide.

9. A compound according to claim 5 wherein Y is 1-H-Indol-3-ylmethyl N-{3-(1-H-Indol-3-ylmethyl)-[1,2,4]thiadiazol-5-ylamino}-acetamide.

10. 3-5 disubstituted 1,2,4-thiadiazole compounds according to claim 1 or 2 wherein A is either leucyl-propyl or isoleucyl-propyl, W is $-OH$ and Y is as defined in claim 1 or 2.

11. Compounds according to claim 10 wherein A is leucyl-propyl, W is $-OH$ and Y is as defined in claim 1 or 2.

12. Compounds according to claim 10 wherein A is isoleucyl-propyl, W is $-OH$ and Y is as defined in claim 1 or 2.

13. A compound according to claim 11 wherein Y is methoxy, which is N-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline.

14. A compound according to claim 11 wherein Y is n-butoxy, which is N-(3-Butoxy-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline.

5 15. A compound according to claim 11 wherein Y is methyl, which is N-(3-Methyl-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline.

16. A compound according to claim 11 wherein Y is phenyl, which is N-(3-Phenyl-[1,2,4]thiadiazol-5-yl)-L-leucyl-L-proline.

10 17. A compound according claim 12 wherein Y is methoxy, which is N-(3-methoxy-[1,2,4]thiadiazol-5-yl)-L-isoleucyl-L-proline.

18. A compound according to claim 10 which is N,N'-{3-methylene-[1,2,4]-
15 thiadiazol-5-yl}-di-{L-leucyl-L-proline}.

19. A compound according to claim 10 which is {3-[4-(2-pyridyl)piperazinyl-methyl]-1,2,4-thiadiazol-5-yl}-leucyl-proline.

20 20. A compound according to claim 10 which is N,N'-{3-methylene-[1,2,4]-thiadiazol-5-yl}-di-{L-leucyl-L-proline methyl ester}.

21. Compounds according to claim 3 wherein X is -NH-C- , A, W and Y are as
25 defined in claim 3.



22. Compounds according to claim 3 wherein X is $\text{-NH-C-CH}_2\text{-}$, A, W and Y are
as defined in claim 3.



23. Compounds according to claim 21 wherein A is leucyl and W and Y are as
30 defined in claim 3.

24. Compounds according to claim 23 wherein W is NH-isoamyl and Y is as defined in claim 3.

35 25. A compound according to claim 24 wherein Y is methoxy, which is 5-{3-methoxy-1,2,4-thiadiazolyl}carbonyl-isoleucyl isoamylamide.

26. Compounds according to claim 22 wherein A is leucyl-prolyl and W and Y are as defined in claim 3.
- 5 27. A compound according to claim 26 wherein W is OH and Y is as defined in claim 3.
28. A compound according to claim 27 wherein W is methoxy, which is N-((3-methoxy-[1,2,4]-thiadiazol-5-yl)carbamoylmethyl)-L-leucine-L-proline.
- 10 29. Compounds according to claim 4 wherein A is phenylalanyl, R¹ is hydrogen, R is carbobenzyloxy and Y is as defined in claim 4.
30. A compound according to claim 29 wherein Y is methoxy, which is 5-(N-carbobenzyloxy-L-phenylalaninamido)-3-methoxy-[1,2,4]thiadiazole.
- 15 31. A compound according to claim 29 wherein Y is carbobenzyloxy-phenylalaninamido, which is N,N'-{3,5-di-(N-carbobenzyloxy-L-phenylalaninamido)}-[1,2,4]thiadiazole.
- 20 32. A pharmaceutical composition comprising a compound according to claim 1 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable diluent or carrier thereof.
- 25 33. A pharmaceutical composition comprising a compound according to claim 2 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable diluent or carrier thereof.
- 30 34. A pharmaceutical composition comprising a compound according to claim 3 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable diluent or carrier thereof.
- 35 35. A pharmaceutical composition comprising a compound according to claim 4 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable diluent or carrier thereof.

36 The compound of any one of claims 1 to 31 or the composition of any one of claims 32 to 35 when used in the treatment of a mammal with a disease or disorder.

5 37 The compound or composition of claim 36 wherein the disorder is acne.

38 The compound or composition of claim 36 wherein the disease is a common cold or an inflammatory joint disease

- 39 Use of a compound or a pharmaceutically acceptable salt thereof according to any one of claims 1 to 31 for the manufacture of a medicament for treating a mammal with a disease or disorder.
- 40 The use of claim 39 wherein the disease or disorder is ameliorated by inhibiting at least a cysteine activity dependent enzyme.
- 41 Use of a composition according to any one of claims 32 to 35 for the manufacture of a medicament for treating a mammal with a disease or disorder.
- 42 The use of claim 41 wherein the disease or disorder is ameliorated by inhibiting at least a cysteine activity dependent enzyme.
- 43 The use according to any one of claims 39 to 42 wherein the disorder is acne.
- 44 The use according to any one of claims 39 to 42 wherein the disease is common cold.
- 45 The use according to any one of claims 39 to 42 wherein the disease is inflammatory joint disease.