

COMPOSITIONS COMPRISING EXTRACTS FROM SANGUINARIA OR MACLEAYA

The present invention relates to novel compositions containing anthocyanosides and/or procyanidins in combination with bactericidal isoquinoline alkaloids and optionally with a lipophilic extract of *Zanthoxylum bungeanum* and/or *Echinacea angustifolia*.

TECHNOLOGICAL BACKGROUND

Throat redness and inflammation with formation of plaques are a frequent aspect of common influenza, cold and other cold diseases. Common cold and influenza, which affect on the average up to three times a year both children and adults, are related to mild viral infections caused by rhinovirus (40%), coronavirus (10%) and to a less extent adenovirus and parainfluenza virus. Although no specific treatments for these pathologies exist, antihistamines and decongestants are considered useful, as the oedema reduction alleviates pain and shortens the duration of the disease underlying inflammation. Said conditions can sometimes involve complications due to secondary bacterial infections, as nasal sinuses are often obstructed by the congestion of mucous membranes wherein pathogenic germs can easily proliferate. In this event, it is necessary to undergo antibiotic treatment.

DISCLOSURE OF THE INVENTION

The present invention relates to novel compositions containing

- a. Anthocyanosides, and/or
- b. Procyanidins,
- c. Isoquinoline alkaloids extracted from *Sanguinaria canadensis*, *Macleaya cordata* or *Macleaya microcarpa*,
and optionally
- d. a lipophilic extract of *Zanthoxylum bungeanum* and/or of *Echinacea*

angustifolia.

According to the present invention, "anthocyanosides" comprises both anthocyanosides *per se* and their aglycones (anthocyanidins).

Anthocyanosides preferably derive from extracts of *Vaccinium*
5 *myrtillus*. The extract of bilberry (*Vaccinium myrtillus*) is known to have marked, particularly topical, anti-inflammatory activity, thanks to its action on capillary permeability and fragility. The preparation of anthocyanosides-containing bilberry extracts is known. Moreover, both bilberry anthocyanosides and procyanindins have bacteriostatic action which prevents
10 bacterial and fungal adhesion, for example at the dental, paradental and mucous membrane levels.

According to the present invention, procyanindins preferably derive from *Vitis vinifera* extracts, obtained as described in GB 1541469 or from *Camellia sinensis* extracts, as disclosed in EP 0814823, or other plants
15 containing them, preferably edible plants.

The content in anthocyanosides- and/or procyanosides- containing extracts can range from 20 to 80 mg per unit dose.

According to the present invention, "isoquinoline alkaloids" mean isoquinoline alkaloids having antibacterial activity, particularly sanguinarines
20 and chelitrines, preferably obtainable from water-alcoholic extracts of *Sanguinaria canadensis*, *Macleaya cordata* or *Macleaya microcarpa*. They have broad-spectrum antimicrobial activity. 0.2% Water-alcoholic solutions have shown therapeutic activity in a number of conditions of the upper respiratory tract, acting on pathogenic gram-positive and gram-negative bacteria, mycetes,
25 fungi and protozoa. The extracts containing these alkaloids have shown strong anti-inflammatory topical activity. The extracts containing the isoquinoline alkaloids are typically present in amounts from 2 to 20 mg per unit dose.

Preferably, the compositions of the invention also comprise lipophilic

extracts of *Zanthoxylum bungeanum* and/or *Echinacea angustifolia* enriched in isobutylamides. Said extracts have useful analgesic topical activity through inhibition of nervous conduction. Each one of said extracts can be added to the compositions of the invention in amounts ranging from 0.02 to 0.05 mg
5 per unit dose.

Therefore, according to a preferred aspect, the compositions of the invention will contain, in addition to components 1 and/or 2, and 3 indicated above, also an isobutylamide-enriched lipophilic extract of *Zanthoxylum bungeanum* and/or *Echinacea angustifolia*. The *Zanthoxylum bungeanum*
10 extract can be prepared e.g. according to WO 00/02570. The *Echinacea* extract can be prepared e.g. according to EP 0 464 298. The compositions of the invention are capable of preventing the formation of purulent plaques infected with various saprophytes of the oral cavity, thus avoiding the use of antibiotics, while reducing the progress of the infection. In particular, the
15 compositions of the invention proved to exert a synergistic effect mainly on the duration of the condition. Furthermore, the compositions of the invention are capable of exerting favourable actions, cleaning the oral cavity and removing dental plaque, thanks to the bacterial adhesiveness-reducing effect, as already mentioned, of both bilberry extracts and procyanindins, and to the
20 high activity of isoquinoline alkaloids on anaerobic bacterial strains.

The pharmaceutical compositions will preferably be formulated as tablets for the slow dissolution in the oral cavity or as chewable forms, particularly gums, which provide the slow release of the active principles. These compositions are used in both prophylactic and curative treatments, as
25 well as for the hygiene of the oral cavity.

Therefore, the present invention also relates to the use of

- a. Anthocyanosides, and/or
- b. Procyanidins,

c. Isoquinoline alkaloids extracted from *Sanguinaria canadensis*,
Macleaya cordata or *Macleaya microcarpa*,
 and optionally

d. *Zanthoxylum bungeanum* and/or *Echinacea angustifolia* lipophilic
 5 extract,

for the preparation of compositions for the prevention and the treatment
 of bacterial infections of the oral cavity and for the hygiene of the oral cavity.

According to a preferred aspect, the compositions of the present
 invention will also contain essential oils.

10 Said compositions will be prepared according to well-known conventional
 methods, such as those described in "Remington's Pharmaceutical Handbook",
 Mack Publishing Co., N.Y., USA, together with suitable excipients.

The following examples illustrate the invention in detail.

EXAMPLE I - 1000 mg chewable tablets containing:

15

1.	<i>Vaccinium myrtillus</i> alcoholic extract.....	40 mg
2.	<i>Sanguinaria canadensis</i> alcoholic extract.....	2 mg
3.	<i>Zanthoxylum bungeanum</i> extracted purified.....	0.025 mg
4.	Soy lecithin.....	30 mg
5.	Anhydrous citric acid.....	5 mg
6.	L-Cysteine.....	5 mg
7.	Lactose.....	200 mg
8.	Mannitol.....	552.475 mg
9.	Methylcellulose.....	40 mg
10.	Glycerol palmitostearate.....	50 mg
11.	Berry flavour.....	40 mg
12.	potassium acesulfame.....	0.5 mg
13.	Talc.....	10 mg
14.	Sodium bicarbonate.....	25 mg

EXAMPLE II - 1000 mg chewable tablets containing:

1.	<i>Vitis vinifera</i> alcoholic extract.....	80 mg
2.	<i>Sanguinaria canadensis</i> alcoholic extract.....	2 mg
3.	<i>Zanthoxylum bungeanum</i> extracted purified.....	0.025 mg
4.	Soy lecithin.....	30 mg
5.	Anhydrous citric acid.....	5 mg
6.	L-Cysteine.....	5 mg
7.	Lactose.....	200 mg
8.	Mannitol.....	512.475 mg
9.	Methylcellulose.....	40 mg
10.	Glycerol palmitostearate.....	50 mg
11.	Berry flavours.....	40 mg
12.	Potassium acesulfame.....	0.5 mg
13.	Talc.....	10 mg
14.	Sodium bicarbonate.....	25 mg

EXAMPLE III - 1000 mg chewable tablets containing

1.	<i>Vaccinium myrtillus</i> alcoholic extract.....	40 mg
2.	<i>Sanguinaria canadensis</i> alcoholic extract.....	2 mg
3.	<i>Echinacea angustifolia</i> lipophylic extract.....	5 mg
4.	Soy lecithin.....	30 mg
5.	Anhydrous citric acid.....	5 mg
6.	L-Cysteine.....	5 mg
7.	Lactose.....	200 mg
8.	Mannitol.....	547.5 mg
9.	Methylcellulose.....	40 mg
10.	Glyceryl palmitostearate.....	50 mg
11.	Berry flavours.....	40 mg
12.	Potassium acesulfame.....	0.5 mg
13.	Talc.....	10 mg
14.	Sodium bicarbonate.....	25 mg

EXAMPLE IV - 1000 mg containing chewable tablets

1.	<i>Vitis vinifera</i> alcoholic extract.....	80 mg
2.	<i>Sanguinaria canadensis</i> alcoholic extract.....	2 mg
3.	<i>Echinacea angustifolia</i> lipophylic extract.....	5 mg
4.	Soy lecithin.....	30 mg
5.	Anhydrous citric acid.....	5 mg
6.	L-Cysteine.....	5 mg
7.	Lactose.....	200 mg
8.	Mannitol.....	507.5 mg
9.	Methylcellulose.....	40 mg
10.	Glycerol palmitostearate.....	50 mg
11.	Berry flavours.....	40 mg
12.	Potassium acesulfame.....	0.5 mg
13.	Talc.....	10 mg
14.	Sodium bicarbonate.....	25 mg

CLAIMS

1. Compositions containing:
 - a. Anthocyanosides extracted from *Vaccinium myrtillus*, and/or
 - 5 b. Procyanidins extracted from *Vitis vinifera*,
 - c. Sanguinarines and chelitrines extracted from *Sanguinaria canadensis*, *Macleaya cordata* or *Macleaya microcarpa*, and optionally
 - d. a *Zanthoxylum bungeanum* and/or *Echinacea angustifolia* lipophilic
10 extract.
2. Compositions as claimed in claim 1 further containing a *Zanthoxylum bungeanum* and/or *Echinacea angustifolia* lipophilic extract.
3. Compositions as claimed in claim 1 or 2 in the form of tablets or chewing gums.
- 15 4. The use of:
 - a. Anthocyanosides extracted from *Vaccinium myrtillus*, and/or
 - b. Procyanidins extracted from *Vitis vinifera*,
 - c. Sanguinarines and chelitrines extracted from *Sanguinaria canadensis*, *Macleaya cordata* or *Macleaya microcarpa*,
20 and optionally
 - d. a *Zanthoxylum bungeanum* and/or *Echinacea angustifolia* lipophilic extract,for the preparation of compositions for the prevention and the treatment of bacterial infections of the oral cavity.
- 25 5. The use of:
 - a. Anthocyanosides extracted from *Vaccinium myrtillus*, and/or

- b. Procyanidins extracted from *Vitis vinifera*,
 - c. Sanguinarines and chelerytrines extracted from *Sanguinaria canadensis*, *Macleaya cordata* or *Macleaya microcarpa*,
and optionally
 - 5 d. a *Zanthoxylum bungeanum* and/or *Echinacea angustifolia* lipophilic
extract,
- for the preparation of compositions for the hygiene of the oral cavity.