Figure 1. Representative Plot of the Concentration-Effect Relationship of Compound (Ia) on Short-Circuit Current by Canine Bronchial Epithelial Cells




Figure 2. Dose-Response of Compound (Ia) on Sheep MCC at 4h Post-dose




Figure 3. Effect of Compound (Ia) and HS on Sheep MCC at 4h post-dose




Figure 4. Effect Compound (Ia) and HS on Sheep MCC at 8h post-dose






Figure 5. Effect of Sodium Channel Block of Compound (Ia) on Surface Liquid Retention 0-8 h in the *in vitro* CBE model.



Results are reported as the mean ± SD; * indicates significance ($p < 0.05$) from vehicle. † Indicates significance ($p < 0.05$) from amiloride

Figure 6. Effect of Compound (Ia) on Surface Liquid Retention at 24 hours in the *in vitro* CBE model



Results are reported as the mean ± SD; * indicates significance ($p < 0.05$) from control

Figure 7. Effect of ENaC Blockers I and Comparative Example I on Sheep MCC at 8 hrs




Figure 8. Effect of ENaC Blockers I and Comparative Example 1 on Sheep Plasma Potassium Levels




Figure 9. Comparison of Comparative Example 4 and Compound 1 on Sheep MCC at 4h Post-dose




Figure 10 . Comparative Example 4 Raises Plasma K^+ at an Effective Dose Compound 1 Has No Effect on Plasma K^+ at 1000x an Effective Dose In Sheep

