


Accordingly, one aspect relates to compounds of formula (I)



(I)

wherein

5 R_1 is selected from the group consisting of H and C_1 to C_3 alkyl;

R_2 is selected from the group consisting of H and C_1 - C_3 alkyl, wherein the C_1 - C_3 alkyl optionally is substituted one or more times with one or more substituents selected from the group consisting of phenyl and C_3 - C_6 cycloalkyl,

R_3 is selected from the group consisting of H, methyl and ethyl,

10 R_4 is selected from the group consisting of H, hydroxy, methoxy and ethoxy,

R_5 , R_6 and R_7 are H,

R_8 is selected from the group consisting of H, methyl, ethyl and cyclopropyl


R_9 is selected from the group consisting of H, methyl and ethyl

and pharmaceutically acceptable acid addition salts of Compound I, racemic
15 mixtures of Compound I, or the corresponding enantiomer and/or optical isomer of Compound I. Another aspect relates to the compound as disclosed herein for use as a medicament.

In another aspect, there is provided the use of the compound as disclosed herein in the manufacture of a medicament for the treatment of ADHD, schizophrenia or
20 cognitive impairment associated with schizophrenia.

Claims

1. A compound having the structure



Compound (I)

5 wherein

R_1 is selected from the group consisting of H and C_1 - C_3 alkyl;

R_2 is selected from the group consisting of H and C_1 - C_3 alkyl

 wherein the C_1 - C_3 alkyl optionally is substituted with phenyl or C_3 - C_6 cycloalkyl;

10 R_3 is selected from the group consisting of H, methyl and ethyl

R_4 is selected from the group consisting of H, , hydroxyl, methoxy and ethoxy and;

R_5 , R_6 and R_7 are H

R_8 is selected from the group consisting of H, methyl, ethyl and cyclopropyl

15 R_9 is selected from the group consisting of H, methyl and ethyl

and pharmaceutically acceptable acid addition salts of Compound I, racemic mixtures of Compound I, or the corresponding enantiomer and/or optical isomer of Compound I.

2. The compound according to claim 1, wherein R_2 is H.